

Lorenzo Foscolo · Mark Haskins · Johannes Nordström

Infinitely many new families of complete cohomogeneity one G_2 -manifolds: G_2 analogues of the Taub–NUT and Eguchi–Hanson spaces

Received June 28, 2018

Abstract. We construct infinitely many new 1-parameter families of simply connected complete non-compact G_2 -manifolds with controlled geometry at infinity. The generic member of each family has so-called asymptotically locally conical (ALC) geometry. However, the nature of the asymptotic geometry changes at two special parameter values: at one special value we obtain a unique member of each family with asymptotically conical (AC) geometry; on approach to the other special parameter value the family of metrics collapses to an AC Calabi–Yau 3-fold. Our infinitely many new diffeomorphism types of AC G_2 -manifolds are particularly noteworthy: previously the three examples constructed by Bryant and Salamon in 1989 furnished the only known simply connected AC G_2 -manifolds.

We also construct a closely related conically singular G_2 -holonomy space: away from a single isolated conical singularity, where the geometry becomes asymptotic to the G_2 -cone over the standard nearly Kähler structure on the product of a pair of 3-spheres, the metric is smooth and it has ALC geometry at infinity. We argue that this conically singular ALC G_2 -space is the natural G_2 analogue of the Taub–NUT metric in 4-dimensional hyperKähler geometry and that our new AC G_2 -metrics are all analogues of the Eguchi–Hanson metric, the simplest ALE hyperKähler manifold. Like the Taub–NUT and Eguchi–Hanson metrics, all our examples are cohomogeneity one, i.e. they admit an isometric Lie group action whose generic orbit has codimension one.

Keywords. Differential geometry, Einstein and Ricci-flat metrics, special and exceptional holonomy, non-compact G_2 holonomy manifolds, cohomogeneity one metrics, collapsed Riemannian manifolds

1. Introduction

Over the past 40 years cohomogeneity one Riemannian metrics, i.e. metrics admitting an isometric Lie group action with generic orbit of codimension one, have played a distinguished role in the construction of complete Ricci-flat or Einstein metrics, particularly in the cases of metrics with special or exceptional holonomy. Throughout its history the subject has attracted considerable interest from both mathematicians and theoretical physi-

L. Foscolo: Department of Mathematics, University College London, London, UK;
e-mail: l.foscolo@ucl.ac.uk

M. Haskins: Mathematics Department, Duke University, Durham, NC, USA;
e-mail: mhaskins@math.duke.edu

J. Nordström: Department of Mathematical Sciences, University of Bath, Bath, UK;
e-mail: j.nordstrom@bath.ac.uk

Mathematics Subject Classification (2020): Primary 53C25, 53C29; Secondary 53C80

cists, with key contributions and important new insights from both communities [5, 10, 14, 17–20, 26, 31, 32, 37, 53]. Moreover, until very recently [35], the *only* known complete non-compact G_2 -manifolds, i.e. Riemannian 7-manifolds admitting metrics with holonomy group the compact exceptional Lie group G_2 , were of cohomogeneity one.

The cohomogeneity one property affords a reduction of the system of non-linear partial differential equations that characterises an Einstein metric or a holonomy reduction, to a system of non-linear ordinary differential equations (ODEs). In some cases these ODEs can be integrated explicitly and completeness of such metrics can then be approached directly [5, 10, 17–20, 26, 27, 29, 31, 53]. In more complicated cases qualitative methods from the theory of ODEs are needed to prove the existence and to establish qualitative properties of solutions; it is usually then a significant challenge to understand when such (non-explicit) solutions give rise to complete metrics [8, 13–15, 34].

Prior to this paper a very limited number of complete cohomogeneity one G_2 -metrics had been constructed: the three rigid *asymptotically conical* (AC) examples constructed by Bryant–Salamon in 1989 [18] and a 1-parameter family of so-called *asymptotically locally conical* (ALC) examples constructed in 2013 by Bogoyavlenskaya [13]. The asymptotic geometry of ALC spaces will be described a little later in this introduction. The existence of the latter family, denoted \mathbb{B}_7 in the physics literature, was first predicted in 2001 by Brandhuber–Gomis–Gubser–Gukov based on an informal analysis of deforming away from a single explicit ALC G_2 -metric that they constructed [17]. In addition to these rigorously constructed examples, numerical analysis of the relevant ODE systems by Brandhuber, Cvetič–Gibbons–Lü–Pope and later Hori–Hosomichi–Page–Rabadán–Walcher, suggested the existence of a further three 1-parameter families of ALC G_2 -metrics, denoted \mathbb{A}_7 [40], \mathbb{C}_7 [16, 28] and \mathbb{D}_7 [16, 25], in the physics literature. Up to discrete symmetries, in the first case the group G acting is $SU(2) \times SU(2)$, while the latter two cases have the enhanced symmetry group $G = SU(2) \times SU(2) \times U(1)$.

Main results. In the current paper we revisit the theory of non-compact G_2 -manifolds with a cohomogeneity one action of $G = SU(2) \times SU(2)$, with a particular focus on the case where the symmetry enhances to $G = SU(2) \times SU(2) \times U(1)$. Our main results are stated in Theorems A–E later in this introduction. Theorem B proves the existence of the previously predicted 1-parameter family of cohomogeneity one G_2 -metrics \mathbb{D}_7 , and also gives a new proof of the existence of the \mathbb{B}_7 family. Theorems C and D construct *infinitely* many new 1-parameter families of complete simply connected cohomogeneity one G_2 -manifolds, all with $G = SU(2) \times SU(2) \times U(1)$. Theorem C is particularly noteworthy because it constructs infinitely many new asymptotically conical G_2 -metrics; previously only the three classical examples due to Bryant–Salamon, dating back to 1989, were known. The metrics constructed in Theorems C and D include the previously predicted 1-parameter family of G_2 -metrics \mathbb{C}_7 as a special case. Theorem E states that our existence results recover all complete simply connected $SU(2) \times SU(2) \times U(1)$ -invariant G_2 -manifolds.

The general qualitative features of all these 1-parameter families turn out to be the same. We will give an explanation for this, which relies on the existence of a new *singular* cohomogeneity one G_2 -metric φ_{cs} on $(0, \infty) \times S^3 \times S^3$ which is forward complete with

ALC geometry as $t \rightarrow \infty$ but which as $t \rightarrow 0$ has a conically singular (CS) end modelled on the G_2 -cone C over the standard homogeneous nearly Kähler structure on $S^3 \times S^3$: see Theorem A.

Motivation from highly collapsed G_2 -metrics. Recently, in [35], we developed a new analytic method for the construction of complete non-compact G_2 -metrics with ALC geometry. This method is very powerful: it gives the first constructions of complete non-compact G_2 -metrics with very little symmetry; easily yields G_2 -metrics on infinitely many different simply connected 7-manifolds and also produces high-dimensional families of G_2 -metrics. The method of construction necessarily produces G_2 -metrics that are highly collapsed, that is, these G_2 -metrics are Gromov–Hausdorff close to a complete non-compact Calabi–Yau 3-fold B . Since smooth complete G_2 -metrics typically deform in a smooth finite-dimensional moduli space, it is natural to try to understand deformations of these highly collapsed G_2 -metrics, both in a local and in a global sense.

The local deformation theory of G_2 -metrics is already well established in other settings: in the smooth compact case by Joyce [41], in the asymptotically cylindrical case by the third author [47] and in the asymptotically conical and conically singular cases by Karigiannis–Lotay [43]. This local deformation theory can be adapted to the ALC setting once a Fredholm theory for elliptic operators on suitable weighted spaces on ALC spaces is developed. We have developed such analytic tools, the details of which, together with the local deformation theory, will appear elsewhere.

However, currently it seems (far) out of reach to hope to understand *large* deformations of our highly collapsed G_2 -metrics in any generality. In this paper we focus on some particular cases where this large deformation question turns out to be tractable. Although the general G_2 -metric we construct using the methods of [35] admits only a circle symmetry, the symmetries are enhanced if the base Calabi–Yau B admits symmetries. In particular, if the limiting Calabi–Yau 3-fold B has cohomogeneity one, then so do our highly collapsed G_2 -metrics. It is then not difficult to deduce that there are infinitely many topological types of simply connected 7-manifolds that admit complete non-compact G_2 -metrics of cohomogeneity one. The goal of this paper is to understand all these complete non-compact G_2 -metrics including those *far from the highly collapsed regime*. We achieve this by using cohomogeneity one methods. It turns out that the behaviour of these moduli spaces of complete cohomogeneity one G_2 -metrics on each of the 7-manifolds in question is qualitatively the same. However, despite the qualitative similarities to the already-understood \mathbb{B}_7 family, constructing these new families of complete cohomogeneity one solutions will require various new ideas.

Geometry of the \mathbb{B}_7 family. We now turn to a description of the key properties of the \mathbb{B}_7 family, since these will be shared by all the cohomogeneity one families we construct and moreover motivate our approach to constructing these new families. The \mathbb{B}_7 family is a 1-parameter family of complete cohomogeneity one G_2 -metrics g_α on $S^3 \times \mathbb{R}^4$, parametrised by a finite interval, say $(0, 1]$. The group acting is $SU(2) \times SU(2) \times U(1)$ with principal orbits diffeomorphic to $S^3 \times S^3$ and singular orbit $S^3 \times \{0\}$. $U(1)$ acts trivially on S^3 and as the standard Hopf action on \mathbb{R}^4 . The quotient space by the (non-fixed-point-free) $U(1)$ -action is nevertheless a manifold, homeomorphic to $S^3 \times \mathbb{R}^3$. For

$\alpha \in (0, 1)$, all the metrics g_α share the same basic asymptotic behaviour: they are complete with (submaximal) volume growth of large geodesic balls of order r^6 .

ALF and ALC spaces. To model finer asymptotic behaviour of n -dimensional Ricci-flat spaces with volume growth of order r^{n-1} it is natural to consider a metric g_∞ on a circle bundle M over a Ricci-flat cone (C, g_C) of dimension $n - 1$ of the form

$$g_\infty = g_C + \ell^2 \theta^2,$$

where $\ell > 0$ is some constant and θ is some fixed connection on the circle bundle M . Thus the model metric g_∞ is a Riemannian submersion over the cone C with circle fibres of constant length $2\pi\ell$. Under appropriate conditions on curvature decay, we might then expect that an n -dimensional Ricci-flat manifold with volume growth of order r^{n-1} must, outside a compact set, become asymptotic to (an exterior domain in) such a model end. If $n = 4$, then (up to a possible \mathbb{Z}_2 -quotient) necessarily $C = \mathbb{R}^3$. Such metrics were therefore termed *asymptotically locally flat* (ALF) spaces by physicists. The higher-dimensional analogues of ALF spaces were subsequently termed *asymptotically locally conical* (ALC) spaces [26]. For $\alpha \in (0, 1)$, any metric g_α in the \mathbb{B}_7 family is ALC. For any ALC G_2 -manifold the cone C should be a 3-dimensional Calabi–Yau cone. Many such cones are now known to exist, e.g. see the discussion in [35, §9] and references therein. Hence, unlike the very rigid situation for ALF hyperKähler 4-manifolds, there are many different asymptotic models for ALC G_2 -metrics. For all ALC metrics in the \mathbb{B}_7 family the Calabi–Yau cone C is the conifold, i.e. the cone over the standard homogeneous Sasaki–Einstein metric on $S^2 \times S^3$, viewed as the homogeneous space $SU(2) \times SU(2)/\Delta U(1)$.

Transitions in the asymptotic geometry of the \mathbb{B}_7 family. As we approach either endpoint of the interval $(0, 1]$, the asymptotic geometry of the metrics g_α degenerates. Fixing the scale of g_α by requiring the size of the singular orbit $S^3 \times \{0\} \subset S^3 \times \mathbb{R}^4$ to be constant, as $\alpha \rightarrow 0$ we find that $\ell \rightarrow 0$ and the G_2 -metrics collapse to the Stenzel metric [20, 53] on the smoothing of the conifold, a well-known example of a cohomogeneity one AC Calabi–Yau 3-fold. Collapse occurs with bounded curvature, except close to the singular S^3 , which is fixed by the $U(1)$ -action.

As $\alpha \rightarrow 1$, instead $\ell \rightarrow \infty$ and the metric g_1 has a different asymptotic behaviour: it is still complete but the volume growth jumps from submaximal r^6 up to maximal growth r^7 . The geometry at infinity of g_1 is *asymptotically conical* (AC), i.e. it is modelled by a G_2 -holonomy cone over a smooth nearly Kähler 6-manifold: in this case $S^3 \times S^3$ endowed with its homogeneous nearly Kähler structure. In fact, the metric g_1 is the classical Bryant–Salamon G_2 -metric on $S^3 \times \mathbb{R}^4$ [18].

When the ALC/AC transition occurs at $\alpha = 1$ nothing catastrophic happens to the *local geometry*, in the sense that there is an extended 1-parameter family g_α , $\alpha \in (0, \infty)$, of local cohomogeneity one G_2 -metrics that continues to close smoothly in the neighbourhood of the singular orbit $S^3 \times \{0\}$. However, for $\alpha > 1$, these local solutions g_α are incomplete. So there are two “phases” of locally well-behaved solutions: one consisting entirely of complete ALC metrics and the other entirely of incomplete metrics. The AC solution appears at the transition between these two phases.

A conically singular ALC G_2 -space, its desingularisations and the G_2 -flop. Given the freedom to rescale any G_2 -metric, when describing the moduli space of G_2 -metrics it is sometimes natural to impose a choice that breaks the scale invariance, and so describes solutions up to scale. Different ways to fix this scale invariance can however lead to different behaviour of solutions in various limiting regimes. In the above description of the \mathbb{B}_7 family, scale invariance was broken by fixing the size of the singular orbit S^3 . For a family of ALC metrics there is another geometrically natural way to break scale invariance: keep the length ℓ of the asymptotic circle of all the metrics fixed. Adopting this alternative scale fixing forces the size of the singular orbit to vary within the \mathbb{B}_7 family and the size shrinks to zero as α approaches 1, where the ALC/AC transition occurred.

If we let \tilde{g}_α denote this rescaling of the \mathbb{B}_7 metrics, then there is an obvious guess for the behaviour of the limit of \tilde{g}_α as $\alpha \rightarrow 1$. There should exist a cohomogeneity one G_2 -metric on $(0, \infty) \times S^3 \times S^3$ with the following properties: at the end where $t \rightarrow \infty$ the metric is forward complete with ALC asymptotics; at the end where $t \rightarrow 0$ we have a conically singular (CS) end, i.e. the metric completion over $t = 0$ has an isolated conical singularity, modelled on the G_2 -cone C over the homogeneous nearly Kähler structure on $S^3 \times S^3$.

Our first main result confirms the existence of this CS ALC G_2 -space: its existence and significance does not seem to have been anticipated in the physics literature.

Theorem A. *Let C be the G_2 -cone over the homogeneous nearly Kähler structure on $S^3 \times S^3$.*

- (i) *There exists a 1-parameter family of $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structures φ_{cs}^c , $c \in \mathbb{R}$, defined on $(0, \epsilon) \times S^3 \times S^3$ for some $\epsilon > 0$ and such that the metric $g_{\varphi_{cs}^c}$ has a conical singularity as $t \rightarrow 0$ asymptotic to the cone C . If $c = 0$ the solution coincides with the cone C .*
- (ii) *If $c > 0$ then the local conically singular solution from (i) is forward complete and extends to a torsion-free G_2 -structure on $(0, \infty) \times SU(2) \times SU(2)$ with a conically singular end as $t \rightarrow 0$ and an ALC end as $t \rightarrow \infty$. All solutions with $c > 0$ differ only by rescaling.*
- (iii) *If $c < 0$ then the local conically singular solution from (i) is forward incomplete.*

Desingularising CS G_2 -metrics. The existence of the CS ALC G_2 -metric constructed in Theorem A immediately suggests the possibility of obtaining families of smooth ALC G_2 -metrics from it, by a desingularisation procedure. Assuming the existence of a compact CS G_2 -space (X, g_{cs}) , Karigiannis [42] used a gluing method to find a 1-parameter family of G_2 -metrics g_t on a smooth compact 7-manifold M that degenerates to (X, g_{cs}) as $t \rightarrow 0$. More specifically, he assumed that the G_2 -cone C that models the singularity admits a G_2 -desingularisation, i.e. a complete AC G_2 -manifold asymptotic at infinity to the given cone C . The smooth compact manifold M is obtained from X by first replacing a neighbourhood of the conical singularity with a rescaled copy of the AC manifold and then correcting to G_2 -holonomy by analytic methods. In the compact case there are two significant obstacles to turning Karigiannis' desingularisation method into a practical way to construct smooth G_2 -metrics: firstly, it is still unknown how to produce any compact

CS G_2 -spaces; secondly, topological obstructions to smoothing do occur in the compact setting.

One can also consider the same G_2 -desingularisation procedure in the setting of non-compact CS G_2 -spaces. In the specific case of the CS ALC G_2 -space constructed in Theorem A the classical Bryant–Salamon AC G_2 -metric on $S^3 \times \mathbb{R}^4$ provides a G_2 -desingularisation of the cone over the standard nearly Kähler structure on $S^3 \times S^3$. For instance, for $\delta > 0$ small and $\alpha \in (1 - \delta, 1)$, the \mathbb{B}_7 metric \tilde{g}_α above should arise this way as a G_2 -desingularisation of the CS ALC G_2 -metric of Theorem A.

To adapt Karigiannis' approach to the non-compact ALC setting requires non-trivial additional analytic work to be undertaken. However, the CS ALC setting has two notable advantages: (i) we have already proven the existence of at least one CS ALC G_2 -space; (ii) the obstructions to smoothing present in the compact setting no longer arise, essentially because of the extra freedom to vary the asymptotic geometry. We will give the details of this ALC G_2 -desingularisation method elsewhere, since this also entails developing the requisite weighted analysis on ALC spaces.

In fact, there are three variants of the AC Bryant–Salamon G_2 -metric on $S^3 \times \mathbb{R}^4$, that are equivalent up to diffeomorphism, but not up to $SU(2) \times SU(2) \times U(1)$ -equivariant diffeomorphism. Depending on which variant is used to perform the desingularisation one obtains three different 1-parameter families of smooth complete ALC G_2 -metrics. Naturally, one family is the \mathbb{B}_7 family, whose degeneration behaviour motivated us to seek the CS ALC solution in the first place. Both the other families are versions of the so-called \mathbb{D}_7 family: families of G_2 -metrics that in the collapsed limit arise from the two small resolutions of the conifold (whereas the \mathbb{B}_7 family collapses to the smoothing of the conifold). The transition between the \mathbb{B}_7 and \mathbb{D}_7 family through the CS ALC space of Theorem A gives a metric version of the so-called G_2 -flop, whose physical significance was emphasised by Acharya and Atiyah–Maldacena–Vafa in the context of large N duality [1, 6].

Since both the CS ALC G_2 -metric of Theorem A and the Bryant–Salamon AC metric have cohomogeneity one, one would hope to be able to prove the existence of these particular G_2 -desingularisations more directly by ODE methods. This is indeed the case.

Theorem B.

- (i) *There exists a 1-parameter family (up to scale) of $SU(2) \times SU(2) \times U(1)$ -invariant complete G_2 -metrics on $S^3 \times \mathbb{R}^4$ with ALC geometry where $U(1)$ acts trivially on S^3 and via the standard Hopf action on \mathbb{R}^4 .*
- (ii) *There exists a 1-parameter family (up to scale) of $SU(2) \times SU(2) \times U(1)$ -invariant complete G_2 -metrics on $S^3 \times \mathbb{R}^4$ with ALC geometry where $U(1)$ acts via the standard Hopf action on S^3 and trivially on \mathbb{R}^4 .*

The family constructed in (i) is the \mathbb{B}_7 family whose existence was first proven in 2013 by Bogoyavlenskaya [13], following earlier work by Brandhuber–Gomis–Gubser–Gukov [17] and Cvetiĉ–Gibbons–Lü–Pope [24]. The family constructed in (ii) is the \mathbb{D}_7 family. Numerical evidence for its existence was given by Cvetiĉ–Gibbons–Lü–Pope [25] and Brandhuber [16, §3.2] We proved the existence of sufficiently collapsed members of this

family in [35]. The advantage of the cohomogeneity one methods of this paper is that they allow us to construct the full 1-parameter space of solutions, not only those that are sufficiently collapsed or close to the CS ALC solution.

Highly collapsed compact CS G_2 -spaces. Singular exceptional holonomy spaces play a crucial role in physics: M theory compactified on a smooth G_2 -manifold has low energy behaviour that is too simple to model known features of the Standard Model of Particle Physics, but M theory on singular G_2 -spaces with a combination of codimension 4 orbifold singularities and codimension 7 singularities can correct this problem [1–3, 11, 55]. Understanding compact singular spaces with G_2 -holonomy is therefore an important, but currently open, problem. The existence of our CS ALC G_2 -space suggests a possible approach (that we are currently pursuing) to construct highly collapsed compact CS G_2 -spaces that are close to a compact CS Calabi–Yau 3-fold.

Finite quotients and infinitely many new AC G_2 -manifolds. We have just seen the importance of AC G_2 -metrics for constructing smooth G_2 -manifolds by desingularising CS G_2 -spaces. AC G_2 -manifolds have also been studied from a physics perspective, e.g. the local physics associated with the three Bryant–Salamon AC G_2 -manifolds was studied in detail by Atiyah–Witten [7]. However, these three classical examples remained the only known AC G_2 -metrics. As we now explain, our previous work on highly collapsed G_2 -metrics also suggests the existence of infinitely many new cohomogeneity one AC G_2 -metrics.

Earlier, we explained that specialising our analytic construction of highly collapsed G_2 -metrics to the case where the collapsed AC Calabi–Yau limit is also of cohomogeneity one, yielded the existence of infinitely many simply connected cohomogeneity one G_2 -manifolds. In each of these cases (up to scale) there is a 1-parameter family of ALC G_2 -metrics close to the highly collapsed limit; it is then natural to try to understand this 1-parameter family away from this highly collapsed regime. Motivated by the geometry of the \mathbb{B}_7 family of G_2 -metrics, it is natural to conjecture that, as we deform *any* of these infinitely many 1-parameter families away from the collapsed limit, eventually an isolated conical singularity develops while maintaining ALC asymptotics at infinity.

CS ALC G_2 -spaces consistent with this conjecture exist: they are quotients of the CS ALC space constructed in Theorem A by particular finite cyclic subgroups of its group of G_2 -isometries. However, in order for such a CS ALC G_2 -space to be a limit of smooth ALC G_2 -metrics we must have previously observed an AC G_2 -metric with prescribed topology and asymptotic geometry bubbling off. Thus continuing these 1-parameter families of highly collapsed cohomogeneity one G_2 -metrics far from the collapsed regime predicts the existence of infinitely many new cohomogeneity one AC G_2 -metrics, asymptotic to particular finite quotients of the standard G_2 -cone over $S^3 \times S^3$.

Infinitely many new cohomogeneity one AC G_2 -metrics. For each $m, n \in \mathbb{Z}$, let $M_{m,n}$ denote the total space of the circle bundle over $K_{\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1}$ whose restriction to the zero section has first Chern class $c_1 = m[\omega] - n[\omega']$, where ω and ω' denote the standard Kähler–Einstein metrics on the two factors. We describe $M_{m,n}$ as a cohomogeneity one space as follows. Denote by $K_{m,n} \subset T^2 \subset \text{SU}(2) \times \text{SU}(2)$ the kernel of the homomorphism $\rho_{m,n} : T^2 \rightarrow \text{U}(1)$ defined by $(e^{i\theta_1}, e^{i\theta_2}) \mapsto e^{i(m\theta_1 + n\theta_2)}$. Then $K_{m,n}$ is isomorphic

to $U(1) \times \mathbb{Z}_d$ where $d = \gcd(m, n)$. If m and n are coprime then $M_{m,n}$ is a simply connected cohomogeneity one 7-manifold with $G = SU(2) \times SU(2)$: it has singular orbit type $G/K_{m,n} \simeq S^2 \times S^3$ and principal orbit type G/Γ_{m+n} where $\Gamma_{m+n} := K_{m,n} \cap K_{2,-2}$ is isomorphic to a cyclic group of order $2|m+n|$ whose generator ζ is embedded in $T^2 \subset SU(2) \times SU(2)$ via $\zeta \mapsto (\zeta^n, \zeta^{-m})$ (up to the action of the outer automorphism the image in fact depends only on $m+n$). In fact, each $M_{m,n}$ also admits a cohomogeneity one action of $SU(2) \times SU(2) \times U(1)$.

Theorem C. *Suppose that m and n are coprime positive integers.*

- (i) *There exists a 1-parameter family φ_α , $\alpha > 0$, of smooth $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structures defined in a tubular neighbourhood of the singular orbit in $M_{m,n}$.*
- (ii) *There exists a unique $\alpha_{ac} > 0$ such that $\varphi_{\alpha_{ac}}$ extends to a complete torsion-free AC G_2 -structure on $M_{m,n}$ asymptotic to the Γ_{m+n} -quotient of the homogeneous nearly Kähler structure on $S^3 \times S^3$.*

These new AC G_2 -metrics are rigid up to scaling. This is consistent with our conjecture that they arise as limits of 1-parameter families of ALC metrics: otherwise the desingularisation construction would yield a larger parameter family of smooth ALC G_2 -metrics close to the CS ALC limit.

Note also that the tangent cone at infinity of our AC G_2 -metrics depends only on the sum $m+n$, not on m and n separately, whereas we prove (see Remark 7.2) that metrics with different coprime pairs (m, n) satisfying $0 < m \leq n$ are not isometric. Hence by considering all such pairs with fixed sum we obtain finitely many different AC G_2 -metrics asymptotic to the same G_2 -cone. This gives rise to infinitely many new geometric transitions in G_2 -geometry.

Cohomogeneity one ALC metrics from AC ones. The existence of these new AC G_2 -metrics asymptotic to the cone over $(S^3 \times S^3)/\Gamma_{m+n}$ allows us to use the ALC version of the G_2 -desingularisation technique to construct new smooth ALC G_2 -metrics close to CS ALC limits (and therefore far from the collapsed limit). However, the cohomogeneity one methods we developed to prove the existence of the new AC G_2 -metrics also enable us to construct such smooth ALC G_2 -metrics more directly. Moreover, these methods allow us to construct the whole 1-parameter family of G_2 -metrics interpolating between the highly collapsed regime and the CS ALC/AC limit, whereas analytic methods can at present produce only solutions close to one of those two limiting regimes. Under the assumptions of Theorem C and adopting its notation we have the following.

Theorem D.

- (i) *If $0 < \alpha < \alpha_{ac}$ then the local solution φ_α constructed in Theorem C extends to a complete torsion-free ALC G_2 -structure on $M_{m,n}$, asymptotic to a circle bundle over a \mathbb{Z}_2 -quotient of the conifold.*
- (ii) *If $\alpha > \alpha_{ac}$ then the local solution φ_α constructed in Theorem C cannot be extended to a complete invariant G_2 -metric.*

Theorems C and D specialised to the case $m = n = 1$ establish the existence of the whole \mathbb{C}_7 family of metrics previously conjectured to exist in [28].

Dihedral ALC G_2 -spaces. When $m = n = 1$, the desingularisation procedure also suggests the existence of another family of complete cohomogeneity one ALC G_2 -metrics. Indeed we can quotient the CS ALC space of Theorem A by a “dihedral” group \mathbb{Z}_4 that does not preserve the circle action, in which case the quotient inherits only a cohomogeneity one action of $SU(2)^2$. However, since the tangent cone at the conical singularity has enhanced $SU(2)^3$ -symmetry, the local singularity model is in fact isomorphic to the cone over $S^3 \times S^3 / \Gamma_2$. Hence the new AC G_2 -metric given by Theorem C can be used to desingularise this singularity. The resulting smooth complete cohomogeneity one ALC G_2 -manifolds have only $SU(2)^2$ -symmetry and belong to the \mathbb{A}_7 family predicted by Hori–Hosomichi–Page–Rabadán–Walcher [40]; because of its smaller symmetry group at present we are unable to recover the \mathbb{A}_7 family by cohomogeneity one methods. We defer the analytic construction of the \mathbb{A}_7 family close to either the CS ALC or the highly collapsed limits to elsewhere.

Classification of $SU(2)^2 \times U(1)$ -invariant complete G_2 -metrics. While we therefore expect the existence of at least another family of complete cohomogeneity one $SU(2)^2$ -invariant G_2 -manifolds, we show that our constructions recover all complete G_2 -manifolds with enhanced $SU(2)^2 \times U(1)$ -symmetry.

Theorem E. *Any complete simply connected $SU(2)^2 \times U(1)$ -invariant G_2 -manifold is isometric to one of the complete metrics of Theorems B, C and D.*

Analogies with 4-dimensional hyperKähler geometry. There are various analogies between the geometry of hyperKähler 4-manifolds and the geometry of G_2 -manifolds. Here we pursue the very strong analogy between the CS ALC G_2 -metric constructed in Theorem A and the Taub–NUT metric on \mathbb{C}^2 . The reader is encouraged to skip to the plan of the paper if analogies with hyperKähler geometry seem unlikely to enlighten.

The Taub–NUT family. Recall that the Gibbons–Hawking ansatz [36] provides a local description of any 4-dimensional hyperKähler metric admitting a triholomorphic circle action in terms of a positive harmonic function h on some domain in \mathbb{R}^3 . We will refer to the family of hyperKähler metrics obtained by choosing $h_m(x) = m + \frac{1}{2|x|}$ for any choice of $m \in \mathbb{R}$ as the *Taub–NUT family* of metrics g_{in}^m . By scaling we can reduce to the three cases: $m = 0$, $m = +1$ or $m = -1$. For $m = 0$ we obtain the flat metric on \mathbb{R}^4 . For $m = 1$ we obtain the (Euclidean) Taub–NUT metric on \mathbb{R}^4 : a complete cohomogeneity one ALF space discovered by Hawking [38]. Many other ALF hyperKähler 4-manifolds can be derived from Taub–NUT as we describe below. For $m = -1$, h is no longer everywhere positive, and the resulting metric is incomplete. However, even this incomplete cohomogeneity one metric is not without interest. For r sufficiently large, h is strictly negative and one therefore obtains a (negative) definite hyperKähler ALF end from it. The asymptotic geometry of the Atiyah–Hitchin metric [5], another more complicated ALF hyperKähler 4-manifold that does not arise from the Gibbons–Hawking ansatz, nevertheless, turns out to be exponentially well approximated by (a finite quotient of) such a negative mass Taub–NUT metric g_{in}^m .

The conically singular family of G_2 -metrics as the G_2 analogue of the Taub–NUT family. It is natural to view the family of CS torsion-free G_2 -structures φ_{cs}^c we constructed in Theorem A as the G_2 analogue of the Taub–NUT family of metrics g_{tn}^m . Certain similarities are immediately apparent: we have a 1-parameter family of cohomogeneity one special holonomy metrics for which the singular orbit is a point; up to scale there are only three different solution types $c = 0$, $c > 0$ or $c < 0$; when $c = 0$ we obtain a Ricci-flat cone C with special holonomy which is rigid and that has enhanced symmetry compared to the solutions with $c \neq 0$; for $c > 0$ the CS solution φ_{cs}^c is forward complete and has an ALC end as $t \rightarrow \infty$; for $c < 0$ the CS solution φ_{cs}^c is forward incomplete.

ALF hyperKähler metrics from quotients of Taub–NUT. Further similarities between the CS family of G_2 -metrics φ_{cs}^c and the Taub–NUT family of metrics g_{tn}^m are connected with the infinitely many new families of cohomogeneity one G_2 -metrics we constructed in Theorems C and D. First we need to recall how to obtain other ALF hyperKähler 4-manifolds by combining the (positive mass) Taub–NUT metric with ALE hyperKähler 4-manifolds, following Biquard–Minerbe [12].

If we identify \mathbb{R}^4 with the quaternions \mathbb{H} then left multiplication by any unit quaternion and right multiplication by unit quaternions in the normaliser $U(1) \rtimes \mathbb{Z}_2$ of the maximal torus $U(1)$ of $SU(2)$ all give isometries of the Taub–NUT metric; the left $SU(2)$ action rotates the 2-sphere of compatible complex structures, while $U(1) \rtimes \mathbb{Z}_2$ acts triholomorphically. On flat \mathbb{R}^4 the group of triholomorphic isometries fixing the origin enhances to $SU(2)$. For any ADE subgroup $\Gamma \subset SU(2)$ the minimal resolution of \mathbb{C}^2/Γ admits a family of complete ALE hyperKähler metrics asymptotic to the Euclidean orbifold metric on \mathbb{C}^2/Γ [44]. In the simplest case, $\Gamma = \mathbb{Z}_2$ is the centre of $SU(2)$ and the minimal resolution of $\mathbb{C}^2/\mathbb{Z}_2$, biholomorphic to $T^*\mathbb{C}P^1$, admits a unique (up to scale) cohomogeneity one $SU(2)$ -invariant ALE hyperKähler metric, the Eguchi–Hanson metric [31]. For any cyclic or dihedral subgroup $\Gamma \subset SU(2)$, but *not* any of the exceptional ones, since these are not contained in $U(1) \rtimes \mathbb{Z}_2$, we may consider the ALF hyperKähler orbifold, Taub–NUT quotiented by Γ . We may resolve its isolated orbifold singularity by gluing in an ALE hyperKähler metric on the minimal resolution of \mathbb{C}^2/Γ . Biquard–Minerbe [12] showed that one can glue in such ALE hyperKähler spaces without destroying the ALF geometry at infinity. In the simplest case $\Gamma = \mathbb{Z}_2$, one obtains a 1-parameter family of cohomogeneity one ALF hyperKähler metrics on $T^*\mathbb{C}P^1$. In fact, whenever Γ is cyclic, both the ALE metric on the minimal resolution and the Taub–NUT metric admit triholomorphic circle actions, and the result of gluing arises more explicitly from the Gibbons–Hawking ansatz. Gluing in a dihedral ALE space will however destroy the triholomorphic circle symmetry of Taub–NUT and the main point of [12] was to construct dihedral ALF spaces.

In the analogy with hyperKähler geometry, quotients of the CS ALC G_2 -metric constructed in Theorem A by finite subgroups of its group of G_2 -structure-preserving isometries acting freely on $S^3 \times S^3$, are the G_2 analogues of orbifold quotients of Taub–NUT. The AC G_2 -metrics constructed in Theorem C, being rigid (up to scale) and of cohomogeneity one, can all be regarded as analogues of the Eguchi–Hanson metric. Finally the ALC version of G_2 -desingularisation, applied to finite quotients of the CS ALC

G_2 -metric of Theorem A, and desingularised by appropriate AC G_2 -metrics, e.g. using the new AC G_2 -metrics we construct, is the G_2 analogue of the Biquard–Minerbe construction.

Plan of the paper. In the rest of the Introduction we give the plan of the paper, which also serves as a detailed outline of the proof of our four main theorems.

In Section 2 we describe an infinite family of simply connected non-compact 7-manifolds admitting cohomogeneity one actions of $SU(2) \times SU(2) \times U(1)$ which, except in one case, arise as the total space of a non-trivial circle bundle over a cohomogeneity one AC Calabi–Yau 3-fold B . There are three main examples of the latter: the small resolutions and the smoothing of the conifold and the canonical bundle of $\mathbb{C}P^1 \times \mathbb{C}P^1$. The small resolutions and the smoothing of the conifold give rise to a single simply connected cohomogeneity one 7-manifold each. Both are diffeomorphic to $S^3 \times \mathbb{R}^4$, with principal orbit $S^3 \times S^3$, but they have inequivalent singular orbit types and are therefore not equivariantly diffeomorphic. $K_{\mathbb{C}P^1 \times \mathbb{C}P^1}$, because it has 2-dimensional second cohomology, provides by far the richest source of simply connected cohomogeneity one 7-manifolds: there are infinitely many such circle bundles $M_{m,n}$ over $K_{\mathbb{C}P^1 \times \mathbb{C}P^1}$, parametrised as described above by a pair of coprime integers m and n . Provided m and n have the same sign, the circle bundles $M_{m,n}$ all satisfy the hypotheses in our recent construction of complete highly collapsed G_2 -manifolds with ALC geometry from circle bundles over an AC Calabi–Yau 3-fold B [35]. Hence we know that each $M_{m,n}$ admits a 1-parameter family of highly collapsed ALC G_2 -metrics, which must be invariant under the action of $SU(2) \times SU(2) \times U(1)$. The principal orbits in $M_{m,n}$ are diffeomorphic to quotients of $S^3 \times S^3$ by a cyclic group Γ_{m+n} of order $2|m + n|$.

The goal of the rest of the paper is to use cohomogeneity one techniques to understand the whole 1-parameter family of complete cohomogeneity one $SU(2) \times SU(2) \times U(1)$ -invariant G_2 -metrics—not only the highly collapsed regime accessible via our analytic methods—for each of the 7-manifolds described above.

In Section 3 we describe the system of non-linear ODEs that governs the local behaviour of $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structures φ on $I \times (S^3 \times S^3)/\Gamma$, where I is some interval and Γ is either trivial or one of the finite cyclic subgroups $\Gamma_k \simeq \mathbb{Z}_{2k}$. The case where Γ is non-trivial is essential to this paper. Both Hamiltonian and Lagrangian formulations of the ODE system arise from Hitchin’s work on stable forms [39]; our paper makes use of both. Two real parameters p and q determine the cohomology class of the fundamental 3-form φ . For each choice of parameters p and q , local $SU(2) \times SU(2) \times U(1)$ -invariant G_2 -metrics on $I \times (S^3 \times S^3)$ depend on two parameters. The main task is to understand which of these local solutions extend to complete metrics, or in the case of the CS ALC solution of Theorem A, to a metric that is forward complete with ALC geometry as $t \rightarrow \infty$ and has prescribed singular behaviour as $t \rightarrow 0$.

In Sections 4 and 5 we understand solutions that extend smoothly over a singular orbit and solutions that have either conically singular or asymptotically conical end behaviour. Proposition 4.5, the main result of Section 4, characterises which members of the 2-parameter family of local $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structures

extend smoothly over the different classes of singular orbits that we need to consider. In each case there are strong restrictions on the values of the constants p and q compatible with smooth extension over a given singular orbit type and in all cases there is (up to scale) a 1-parameter family of smooth solutions defined in a neighbourhood of each type of singular orbit. To prove these results we adapt to our first-order ODE systems the representation-theoretic approach to singular initial value problems for cohomogeneity one Einstein metrics developed by Eschenburg–Wang [32]. The first two authors used the same framework in the course of proving the existence of complete cohomogeneity one nearly Kähler 6-manifolds [34, §4]. The main result in Section 5 is Proposition 5.3, which proves the existence of 1-parameter families of $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structures on either a CS or AC end; in the CS case necessarily $p = q = 0$, while in the AC case we get a 1-parameter family of AC ends φ_{ac}^c for each fixed p, q . Describing solutions with either type of end behaviour leads to a class of singular initial value problems not widely studied in the previous extensive work on cohomogeneity one Einstein metrics. We state a general existence result, Theorem 5.1, which yields convergent *generalised* power series solutions to a wide class of first-order singular initial value problems, including ours, and for which solutions depend real analytically on a finite number of real parameters.

The most novel arguments in the paper appear beginning in Section 6 where we develop criteria that guarantee that a locally-defined cohomogeneity one torsion-free G_2 -structure extends to a forward complete solution and then establish additional conditions under which we can bootstrap from forward completeness to finer asymptotic metric behaviour—in our case ALC asymptotics.

Proposition 6.1 gives a necessary and sufficient condition for forward completeness in terms of the positivity of the mean curvature of all principal orbits. The fact that (non-flat) principal orbits cannot be minimal in *complete* cohomogeneity one Ricci-flat manifolds was observed previously by Böhm [15]. The sufficiency of the positivity of mean curvature of all principal orbits for forward completeness, however, uses the fact that we have a *first-order* ODE system for the metric g . It is a pleasure to thank Burkhard Wilking for suggesting this idea. Close to a singular orbit, or to an isolated conical singularity, the mean curvature of any principal orbit is necessarily large and positive. The next step is therefore to determine conditions under which the positivity of the mean curvature of the principal orbits persists for all time.

Most of the results of Sections 2 to 5 are stated more generally for $SU(2) \times SU(2)$ -invariant torsion-free G_2 -structures that do not necessarily enjoy an enhanced symmetry group $SU(2) \times SU(2) \times U(1)$. At this stage, to make further progress, it is crucial that we restrict to the case where there is an additional $U(1)$ -symmetry. Once the values of p and q have been fixed, two coefficient functions a and b determine any closed $SU(2) \times SU(2) \times U(1)$ -invariant G_2 -structure φ . The rest of our analysis of the global behaviour of solutions is based on the Lagrangian formulation of the problem. Under the additional symmetry assumption this leads to the single second-order non-linear ODE, equation (3.16), which we write schematically as $G_{p,q}(a, b, a', b', a'', b'') = 0$. In Lemma 6.4 we prove that solutions to equation (3.16) that begin in a certain open subset \mathcal{O}_{fc} of the set of principal orbits must remain in \mathcal{O}_{fc} , at least while they continue to exist with the fundamental

3-form φ remaining well-defined. Moreover, it turns out that the mean curvature of any principal orbit in \mathcal{O}_{fc} is strictly positive. Therefore, by Proposition 6.1, any solution that ever enters \mathcal{O}_{fc} cannot blow up in finite time and is therefore forward complete.

Significantly more work is needed to pass from forward completeness to a statement about the complete end necessarily having ALC geometry. The first step is to clarify what asymptotic behaviour the coefficients a and b should exhibit along an ALC end. On the total space of the circle bundle $\mathbb{R}^+ \times \text{SU}(2) \times \text{SU}(2) \rightarrow \mathbb{R}^+ \times (\text{SU}(2) \times \text{SU}(2))/\Delta\text{U}(1) = C$ over the conifold C there is an explicit 1-parameter family of closed $\text{SU}(2) \times \text{SU}(2) \times \text{U}(1)$ -invariant G_2 -structures φ_∞^ℓ , where the parameter $\ell > 0$ is the asymptotic length of the circle fibre. These closed G_2 -structures φ_∞^ℓ serve as models for the possible asymptotic behaviour of ALC ends of invariant G_2 -metrics. Convergence of φ to the asymptotic model φ_∞^ℓ implies that, in appropriate parametrisations, a and b behave asymptotically like s^3 and s^2 respectively.

For a solution to (3.16) that enters the open set \mathcal{O}_{fc} above, forward completeness already forces both a and b to go to infinity. However, without further assumptions it does not force the asymptotic behaviour needed for an ALC end. To guarantee this we need to restrict to a smaller open subset $\mathcal{O}_{alc} \subset \mathcal{O}_{fc}$ of the space of principal orbits. Proposition 6.11 establishes that if a solution of (3.16) enters \mathcal{O}_{alc} then it stays in \mathcal{O}_{alc} for all future times and has an ALC end modelled on φ_∞^ℓ for some $\ell > 0$. Proposition 6.11 is our main tool for establishing the existence of forward complete invariant torsion-free G_2 -structures with an ALC end. On the other hand, Proposition 6.13 establishes the existence of an open subset \mathcal{O}_{in} of the space of principal orbits with the property that any solution of (3.16) that enters \mathcal{O}_{in} is forward *incomplete*.

We can now easily prove the existence of the conically singular ALC G_2 -metric described in Theorem A and also the two 1-parameter families \mathbb{B}_7 and \mathbb{D}_7 of ALC G_2 -metrics described in Theorem B, by combining Proposition 6.11 with information about the local solutions extending smoothly on singular orbits constructed in Section 4, or the local solutions with CS ends constructed in Section 5. The point is that for all these local solutions we have expansions for the coefficients a and b of the 3-form φ in a neighbourhood of the singular orbit. Using these we can identify for exactly which parameter values local solutions enter \mathcal{O}_{alc} or \mathcal{O}_{in} . For instance, the local CS solutions φ_{cs}^c enter into \mathcal{O}_{alc} (respectively, \mathcal{O}_{in}) precisely when $c > 0$ ($c < 0$).

One also expects a similar behaviour for the infinitely many 1-parameter (up to scale) families of local solutions defined in a neighbourhood of the singular orbit in $M_{m,n}$: there are two phases, one corresponding to solutions that enter \mathcal{O}_{alc} and one corresponding to solutions that enter \mathcal{O}_{in} ; the two phases are separated by a unique (up to scale) solution corresponding to a smooth AC metric on $M_{m,n}$. However, unlike the previous cases, it is impossible to determine which phase each solution belongs to simply by looking at its behaviour near the singular orbit. The key to proving Theorems C and D turns out to be first to focus attention on the single member of each 1-parameter family that has AC geometry. Once we prove the existence of the unique AC member of each 1-parameter family, we will be able to compare the behaviour of our other local solutions with it. We will use this to conclude that, for all parameter values on one side of the AC solution, local solutions are eventually forced to enter \mathcal{O}_{alc} , while on the other side of the AC

solution, local solutions that exist for a sufficiently long time must eventually enter \mathcal{O}_{in} ; see Sections 7.2 and 7.3.

It remains therefore to describe the approach we take in Section 7.1 to the construction of a new rigid complete AC G_2 -metric on each of the infinitely many spaces $M_{m,n}$. To illustrate the somewhat delicate nature of the problem consider the following facts. From Proposition 4.5(iii, iv), for each choice of m and n there is (up to scale) a 1-parameter family of local $SU(2) \times SU(2) \times U(1)$ -invariant solutions that extend smoothly over the singular orbit $SU(2) \times SU(2)/K_{m,n}$. Moreover, the constants p and q that specify the cohomology class of the G_2 -structure φ are determined (up to scale) by the choice of m and n . Also from Proposition 5.3(ii), for each choice of p and q , there exists a 1-parameter family of $SU(2) \times SU(2) \times U(1)$ -invariant solutions with AC ends. Finally, recall also that the analysis of the ODEs on the principal orbits showed that, for each fixed p and q , there is a 2-parameter family of local $SU(2) \times SU(2) \times U(1)$ -invariant solutions. Putting this together, a complete AC G_2 -metric corresponds to an intersection point of two curves in a 2-dimensional manifold. For dimensional reasons we might hope that such matching happens a (non-zero!) finite number of times.

We consider the problem of “shooting from infinity”, i.e. we consider the Γ_{m+n} quotient of the 1-parameter family φ_{ac}^c , $c \in \mathbb{R}$, of invariant AC end solutions φ_{ac}^c we constructed on $(T, \infty) \times S^3 \times S^3$ and investigate extending these solutions backwards. Our aim is to show that as we vary the parameter $c \in \mathbb{R}$, for exactly one value of c the maximal backward extension of the AC end satisfies the conditions to close smoothly on the singular orbit $SU(2) \times SU(2)/K_{m,n}$ at its backward extinction time. Proposition 7.6 provides our main criterion for backward extension of invariant AC ends: if a principal orbit belongs to a certain open subset $\mathcal{O}_{\text{ac}}^+$ of the set of principal orbits then the solution can be continued backwards in time remaining in $\mathcal{O}_{\text{ac}}^+$, provided that the fundamental 3-form φ remains well-defined. The invariant AC ends φ_{ac}^c constructed in Proposition 5.3(ii) belong to $\mathcal{O}_{\text{ac}}^+$ precisely when the parameter $c \in \mathbb{R}$ is positive. The fundamental 3-form φ fails to be well-defined whenever the curve (a, b) hits the zero-locus of an explicit quartic polynomial F (depending on m and n). The level set $F = 0$ contains a unique singular point. In Proposition 7.12 we show that there exists a unique parameter value $c_{\text{ac}} > 0$ such that the solution φ_{ac}^c extends backward until it hits this distinguished point. Finally, in Proposition 7.13 we show that this unique solution in fact defines a smooth torsion-free G_2 -structure on $M_{m,n}$ which by construction is AC.

The classification result Theorem E is also proved in Section 7 by showing that, up to a finite quotient, any $SU(2)^2 \times U(1)$ -invariant G_2 -manifold closing smoothly on a singular orbit must be equivariantly diffeomorphic to one of the manifolds we consider in the paper; cf. Theorem 7.3.

2. Cohomogeneity one ALC G_2 -manifolds

In this section we describe the 7-manifolds with a cohomogeneity one action of $SU(2) \times SU(2)$ known to admit complete G_2 -metrics. We start by describing the \mathbb{B}_7 family [13, 17] and then discuss our analytic construction of highly collapsed ALC G_2 -metrics [35], specialised to the cohomogeneity one setting.

A *cohomogeneity one manifold* X is a Riemannian manifold acted upon isometrically by a Lie group G with generic orbits of codimension one. The cohomogeneity one manifolds we consider in this paper are complete non-compact irreducible Ricci-flat manifolds and G will be compact. The Splitting Theorem implies that they have only one end and therefore the orbit space X/G is a half-line $[0, \infty)$. The cohomogeneity one structure of X is encoded by a pair of closed subgroups $K_0 \subset K \subset G$ called its *group diagram*. Orbits corresponding to points in $(0, \infty)$ are hypersurfaces all diffeomorphic to G/K_0 and are called *principal orbits*. The orbit over 0 is the only lower-dimensional orbit G/K and it is called the *singular orbit*. X has the topology of a disc bundle over the singular orbit. The complement of the zero section of this disc bundle is foliated by the principal orbits G/K_0 , which themselves are sphere bundles over G/K . In particular, K/K_0 must be diffeomorphic to a sphere. All the manifolds we consider in this paper admit a cohomogeneity one action of $G = \text{SU}(2) \times \text{SU}(2)$ or of $G = \text{SU}(2) \times \text{SU}(2) \times \text{U}(1)$.

2.1. The \mathbb{B}_7 family and its symmetries

The \mathbb{B}_7 family is a 1-parameter family (up to scale) of G_2 -metrics on $S^3 \times \mathbb{R}^4$ admitting a cohomogeneity one action of $\text{SU}(2) \times \text{SU}(2)$, in fact of $\text{SU}(2) \times \text{SU}(2) \times \text{U}(1)$. As explained in the Introduction, the generic member of the family has ALC asymptotic geometry, while at a special parameter value we have the Bryant–Salamon AC metric [18]. One special explicit ALC member of the family was found by Brandhuber–Gomis–Gubser–Gukov [17], while the existence of the full 1-parameter family of ALC metrics was later established by Bogoyavlenskaya [13]. We now carefully describe the cohomogeneity one structure of these metrics and their full group of symmetries.

We identify $S^3 = \text{SU}(2)$ with the unit quaternions. $S^3 \times \mathbb{R}^4$ admits a cohomogeneity one action of $\text{SU}(2) \times \text{SU}(2)$ described by the group diagram

$$\{1\} \subset \Delta\text{SU}(2) \subset \text{SU}(2) \times \text{SU}(2). \tag{2.1}$$

More explicitly, we can identify $[0, \infty) \times S^3 \times S^3/\sim$ with $S^3 \times \mathbb{R}^4$ via $(t, q_1, q_2) \mapsto (q_1\bar{q}_2, tq_1)$, where $\text{SU}(2) \times \text{SU}(2)$ acts by left multiplication on $S^3 \times S^3$ and $(t, q_1, q_2) \sim (t', q'_1, q'_2)$ if and only if $t = 0 = t'$ and $q'_1 = q_1q, q'_2 = q_2q$ for some $q \in \text{SU}(2)$.

The full symmetry group of any G_2 -metric in the \mathbb{B}_7 family is however larger than $\text{SU}(2) \times \text{SU}(2)$; we describe it first for the most symmetric member of the family, the AC Bryant–Salamon metric. In this case the group of continuous G_2 -isometries is $G^{\text{bs}} = \text{SU}(2)^2 \times \text{SU}(2)/\Delta\mathbb{Z}_2$, where the first factor $\text{SU}(2)^2$ acts by left multiplication, the third $\text{SU}(2)$ acts diagonally by right multiplication and the quotient by $\Delta\mathbb{Z}_2$ appears because to get an effective action we must quotient by the centre \mathbb{Z}_2 of $\text{SU}(2)$ embedded diagonally in the three factors. The asymptotic cone C of the AC Bryant–Salamon metric is the cone over the standard homogeneous nearly Kähler structure on $S^3 \times S^3$. The induced metric on $S^3 \times S^3$ has additional discrete isometries arising from outer automorphisms of $\text{SU}(2)^3$. More concretely there is an action of the symmetric group S_3 on $S^3 \times S^3$ generated by the pair of involutions

$$(q_1, q_2) \mapsto (\bar{q}_1, q_2\bar{q}_1), \quad (q_1, q_2) \mapsto (q_1\bar{q}_2, \bar{q}_2),$$

whose composition has order 3. However this action of S_3 does not extend to the AC Bryant–Salamon metric. Instead the Bryant–Salamon metric can be realised in three inequivalent ways as a cohomogeneity one manifold with an action of $SU(2) \times SU(2)$, corresponding to the group diagrams (2.1), (2.4) and its image under the outer automorphism of $SU(2)^2$.

The symmetry group of the ALC members of the \mathbb{B}_7 family is smaller than G^{bs} . The standard Hopf action of S^1 on \mathbb{R}^4 induces a circle action on $S^3 \times \mathbb{R}^4$. The action fixes $S^3 \times \{0\}$, but the quotient $S^3 \times \mathbb{R}^3$ is still a manifold. Under our identification of $S^3 \times \mathbb{R}^4$ with $[0, \infty) \times S^3 \times S^3/\sim$, the projection $\pi : [0, \infty) \times S^3 \times S^3/\sim \rightarrow S^3 \times \mathbb{R}^3$ is given by

$$(t, q_1, q_2) \mapsto (q_1\bar{q}_2, t^2q_1i\bar{q}_1),$$

where we identify \mathbb{R}^3 with $\text{Im } \mathbb{H}$. The group of continuous isometries that preserves the generic G_2 -structure in the \mathbb{B}_7 family is the subgroup G_π of G^{bs} that preserves this projection π . Note that since, away from $t = 0$, π is a principal circle bundle endowed with a natural connection θ (the left-invariant 1-form on $SU(2)^2$ dual to the vector field generating the diagonal $U(1)$ subgroup), there is an exact sequence

$$1 \rightarrow G_\pi^+ \rightarrow G_\pi \rightarrow \mathbb{Z}_2 \rightarrow 1,$$

where G_π^+ is the subgroup that preserves the connection θ (whereas arbitrary elements of G_π are allowed to send θ to $\pm\theta$). We have $G_\pi^+ = SU(2)^2 \times U(1)/\Delta\mathbb{Z}_2$, where $SU(2)^2$ is the left-acting group, $U(1)$ acts diagonally on the right and $\Delta\mathbb{Z}_2$ is the centre of $SU(2)$ embedded diagonally in each factor, and $G_\pi = SU(2)^2 \times N/\Delta\mathbb{Z}_2$, where $N \simeq U(1) \rtimes \mathbb{Z}_2$ is the normaliser of $U(1)$ in the right-acting $SU(2)$. There is a further \mathbb{Z}_2 -action generated by the outer automorphism of $SU(2)^2$, i.e. the involution of $S^3 \times S^3$ that exchanges the two factors. This automorphism is an isometry of all members of the \mathbb{B}_7 family but it always acts non-trivially on the G_2 -structure.

By considering quotients of the \mathbb{B}_7 family by finite subgroups $\Gamma \subset G_\pi$ that act freely on $S^3 \times S^3$ we can obtain either non-simply-connected smooth ALC G_2 -manifolds or ALC G_2 -orbifolds with singularities contained in a compact set. Finite subgroups of G_π acting freely on $S^3 \times S^3$ exist in abundance: since the subgroup $SU(2)^2$ acts by left translations on $S^3 \times S^3$, any finite subgroup $\Gamma \subset SU(2)^2 \subset G_\pi$ necessarily acts freely. (More generally, a partial classification of subgroups of $SU(2)^3$ acting freely on $S^3 \times S^3$ was obtained recently in [22], though the complete classification remains open.) In general the resulting quotient space $(S^3 \times \mathbb{R}^4)/\Gamma$ will no longer admit a cohomogeneity one action. However, if one chooses Γ to be contained in the subgroup $\mathbb{Z}_2^2 \times N/\Delta\mathbb{Z}_2$ of G_π that commutes with the left action of $SU(2)^2$ then the quotient space does still admit a cohomogeneity one action of $SU(2) \times SU(2)$. Note that $\mathbb{Z}_2^2 \times N/\Delta\mathbb{Z}_2 \simeq \mathbb{Z}_2 \times N$ and the action on the two factors of $S^3 \times \mathbb{R}^4$ is given precisely by this isomorphism. There are three possibilities for Γ :

- (i) $\Gamma = \Gamma_0$ where Γ_0 is a cyclic or binary dihedral subgroup of $SU(2)$;
- (ii) $\Gamma = \mathbb{Z}_2 \times \Gamma_0$ with Γ_0 as above;
- (iii) Γ is a cyclic group of even order or a binary dihedral group embedded in $\mathbb{Z}_2 \times N$ via the homomorphism $\phi \times \iota$, where $\phi : \Gamma \rightarrow \mathbb{Z}_2$ is a non-trivial homomorphism and ι is the standard embedding of Γ in $N \subset SU(2)$.

The resulting orbifolds are $S^3 \times (\mathbb{R}^4/\Gamma_0)$, $\mathbb{R}P^3 \times (\mathbb{R}^4/\Gamma_0)$ and $(S^3 \times (\mathbb{R}^4/\Gamma_0))/\mathbb{Z}_2$, respectively, where in the latter case $\Gamma_0 := \phi^{-1}(1)$. In this paper the case where Γ is a cyclic group of even order embedded in $\mathbb{Z}_2 \times U(1) \subset \mathbb{Z}_2 \times N$ as in (iii) plays a distinguished role.

We introduce the following notation that we use throughout the rest of the paper: for each $m, n \in \mathbb{Z}$ let $K_{m,n}$ be the subgroup of the maximal torus $T^2 \subset SU(2) \times SU(2)$ defined by

$$K_{m,n} = \{(e^{i\theta_1}, e^{i\theta_2}) \in T^2 \mid e^{i(m\theta_1+n\theta_2)} = 1\}, \tag{2.2}$$

i.e. $K_{m,n}$ is the kernel of the group homomorphism $(e^{i\theta_1}, e^{i\theta_2}) \mapsto e^{i(m\theta_1+n\theta_2)}$ from T^2 to $U(1)$. In particular, $K_{1,-1}$ is the diagonally embedded circle $\Delta U(1)$ in $SU(2) \times SU(2)$ and the subgroup $\mathbb{Z}_2 \times U(1)$ of elements in G_π^+ that commute with the left $SU(2)^2$ -action on $S^3 \times S^3$ can be identified with the right-acting $K_{2,-2}$. Indeed, note that $K_{m,n}$ is isomorphic to the direct product $U(1) \times \mathbb{Z}_{\gcd(m,n)}$ via the map $(e^{i\theta}, \zeta) \mapsto (e^{ik\theta} \zeta^r, e^{-ih\theta} \zeta^s)$, where $m = \gcd(m, n)h, n = \gcd(m, n)k$ and $rh + sk = 1$. We also note that conjugation by $j \in SU(2)$ (or any other unit quaternion orthogonal to 1 and i) on the first [second] factor identifies $K_{m,n}$ with $K_{-m,n}$ [$K_{m,-n}$] since $-je^{i\theta}j = e^{-i\theta}$.

Now, if $\Gamma_k \simeq \mathbb{Z}_{2k}$ is the subgroup of $K_{2,-2} = \mathbb{Z}_2 \times U(1)$ defined by the embedding $\zeta \mapsto (\zeta^k, \zeta)$, then the orbifold $(S^3 \times \mathbb{R}^4)/\Gamma_k$ admits a 1-parameter family of cohomogeneity one G_2 -metrics (the quotients of the \mathbb{B}_7 family of metrics) with group diagram

$$\Gamma_k \subset \Delta SU(2) \subset SU(2) \times SU(2).$$

We can think of this orbifold as a *partial* resolution of the cone over $(S^3 \times S^3)/\Gamma_k$. In the next section we will see that there are *smooth* 7-manifolds that topologically resolve the same cone and that are known to carry complete cohomogeneity one ALC G_2 -metrics.

Remark 2.3. When k is odd, the isomorphism $\mathbb{Z}_{2k} \rightarrow \mathbb{Z}_2 \times \mathbb{Z}_k$ defined by $\zeta \mapsto (\zeta^k, \zeta^2)$ and the outer automorphism of $SU(2) \times SU(2)$ that exchanges the two factors yield an isomorphism between Γ_k and the subgroup $\mathbb{Z}_2 \times \mathbb{Z}_{2k} \subset \mathbb{Z}_2 \times U(1)$ of type (ii).

2.2. Highly collapsed ALC G_2 -metrics

The starting point for this paper is our construction of complete G_2 -manifolds from AC Calabi–Yau 3-folds [35] and its specialisation to the cohomogeneity one case. Our analytic construction yields the existence of infinitely many new families of cohomogeneity one G_2 -metrics, but recovers only metrics close to a certain degenerate (collapsed) limit. In this section we describe these cohomogeneity one manifolds. The main goal of the paper is to describe the full moduli space of invariant G_2 -metrics on each of the 7-manifolds in question.

Cohomogeneity one AC Calabi–Yau 3-folds. The cross-sections of Calabi–Yau cones are called *Sasaki–Einstein manifolds*. The only homogeneous Sasaki–Einstein 5-manifolds are the round 5-sphere and $S^2 \times S^3 = SU(2) \times SU(2)/\Delta U(1)$, or finite quotients thereof [52, §2.2]. The case of the round 5-sphere and its quotients is not relevant for this paper and therefore we consider only AC Calabi–Yau 3-folds B asymptotic to (quotients of) the

conifold, i.e. the Calabi–Yau cone $C(\Sigma)$ over $\Sigma = S^2 \times S^3$ endowed with its $SU(2) \times SU(2)$ -invariant Sasaki–Einstein structure. If we insist that the Calabi–Yau 3-fold B has a cohomogeneity one action then there are three known possibilities. In the following we use the notation $K_{m,n}$ introduced in (2.2).

- (i) B is one of the two small resolutions of the conifold [20]. They have group diagrams

$$\begin{aligned} K_{1,-1} &\subset U(1) \times SU(2) \subset SU(2) \times SU(2), \\ K_{1,-1} &\subset SU(2) \times U(1) \subset SU(2) \times SU(2). \end{aligned}$$

The two resolutions are in fact isomorphic under the outer automorphism of $SU(2) \times SU(2)$ that exchanges the two factors. Without loss of generality we can therefore concentrate only on the first case, which from now on we will refer to as *the small resolution of the conifold*.

- (ii) B is the canonical line bundle of $\mathbb{C}P^1 \times \mathbb{C}P^1$. Its group diagram is

$$K_{2,-2} \subset T^2 \subset SU(2) \times SU(2).$$

In this case the tangent cone at infinity is a (free) \mathbb{Z}_2 -quotient of the conifold. Moreover, while the AC Calabi–Yau metric on the small resolution of the conifold is unique (up to scale), in this case there is a 2-parameter family of AC Calabi–Yau metrics on B , parametrised by the Kähler class of the corresponding Kähler form. The metrics corresponding to the unique (up to scale) compactly supported Kähler class are due to Calabi [19]; the full 2-parameter family of AC Calabi–Yau metrics was first considered in [49].

- (iii) $B = T^*S^3$ is the smoothing of the conifold [20, 53]. Its group diagram is

$$K_{1,-1} \subset \Delta SU(2) \subset SU(2) \times SU(2).$$

Remark. In (ii), it would perhaps be more natural to define $S^2 \times S^3/\mathbb{Z}_2$ as $SU(2) \times SU(2)/K_{2,2}$. Our choice is motivated by the desire to use the same conventions adopted in the physics literature, in particular those of [16]. The main confusing consequence of our choice is that the complex structures on the two factors in $\mathbb{C}P^1 \times \mathbb{C}P^1 = SU(2) \times SU(2)/T^2$ are conjugate to each other. In particular, with our choices Kähler classes on $\mathbb{C}P^1 \times \mathbb{C}P^1$ are parametrised by pairs $(\alpha, -\beta)$ with $\alpha, \beta > 0$.

Highly collapsed G_2 -metrics on circle bundles over AC Calabi–Yau 3-folds. Let $(B, g_0, \omega_0, \Omega_0)$ be one of the three simply connected cohomogeneity one AC Calabi–Yau 3-folds just described. Here ω_0 is the Kähler form of the AC Calabi–Yau metric g_0 and Ω_0 is a holomorphic complex volume form normalised so that $\omega_0^3 = \frac{3}{2} \operatorname{Re} \Omega_0 \wedge \operatorname{Im} \Omega_0$. We now consider a *non-trivial* circle bundle M over B : therefore we consider only cases (i) and (ii), since the second cohomology vanishes in case (iii).

- (i) When B is the small resolution of the conifold, up to finite quotients and a change of orientation, there is only one possible choice of line bundle. The total space M is $S^3 \times \mathbb{R}^4$, endowed with an $SU(2) \times SU(2)$ -action with group diagram

$$\{1\} \subset \{1\} \times SU(2) \subset SU(2) \times SU(2). \tag{2.4}$$

- (ii) When $B = K_{\mathbb{C}P^1 \times \mathbb{C}P^1}$ we have an infinite family of examples: for a pair (m, n) of coprime integers we consider the simply connected manifold $M_{m,n}$ described by the group diagram

$$K_{m,n} \cap K_{2,-2} \subset K_{m,n} \subset \text{SU}(2) \times \text{SU}(2). \tag{2.5}$$

The subgroup $K_{m,n} \cap K_{2,-2}$ is isomorphic to $\mathbb{Z}_{2|n+m|}$ embedded in $T^2 \subset \text{SU}(2) \times \text{SU}(2)$ via $\zeta \mapsto (\zeta^n, \zeta^{-m}) = (\zeta^n, \zeta^{n+m}\zeta^n)$. If n is odd then $\text{gcd}(n, 2|n+m|) = 1$ since n and m are coprime and therefore $\zeta \mapsto \zeta^n$ is an isomorphism; $K_{m,n} \cap K_{2,-2}$ is then isomorphic to the group $\Gamma_{|n+m|}$ defined in the previous section. If $n = 2n'$ is even, then $\text{gcd}(n', |n+m|) = 1$ and $K_{m,n} \cap K_{2,-2}$ is isomorphic to the subgroup $\{(\xi, \epsilon \xi) \in T^2 \mid \xi \in \mathbb{Z}_{|n+m|}, \epsilon \in \mathbb{Z}_2\}$ via the composition of the isomorphism $\mathbb{Z}_{2|n+m|} \simeq \mathbb{Z}_{|n+m|} \times \mathbb{Z}_2$ given by $\zeta \mapsto (\zeta^2, \zeta^{|n+m|})$ and the automorphism of $\mathbb{Z}_{|n+m|}$ defined by $\xi \mapsto \xi^{n'}$. As explained in Remark 2.3 the outer automorphism that exchanges the two factors of $\text{SU}(2) \times \text{SU}(2)$ identifies this group with $\Gamma_{|n+m|}$ (in fact, note that if n is even then m must be odd).

In all cases there exists a unique $\text{SU}(2) \times \text{SU}(2)$ -invariant connection θ on the circle bundle $M \rightarrow B$ (the left-invariant 1-form dual to the vector field that generates the diagonal $\text{U}(1)$ subgroup) and therefore a 1-parameter family of $\text{SU}(2)^2 \times \text{U}(1)$ -invariant metrics

$$\bar{g}_\epsilon = g_0 + \epsilon^2 \theta^2.$$

Note that the model metric \bar{g}_ϵ has ALC asymptotic geometry. Furthermore, for $\epsilon > 0$ sufficiently small, \bar{g}_ϵ is approximately G_2 . In [35] we use analytic methods to show that, under the necessary topological condition $c_1(M) \cup [\omega_0] = 0 \in H^4(B)$, for all $\epsilon > 0$ sufficiently small, we can perturb \bar{g}_ϵ to an $\text{SU}(2)^2 \times \text{U}(1)$ -invariant ALC G_2 -metric g_ϵ . The metric g_ϵ is *highly collapsed*: g_ϵ is arbitrarily close (in certain weighted Hölder spaces) to the model metric \bar{g}_ϵ and therefore (M, g_ϵ) collapses to (B, g_0) as $\epsilon \rightarrow 0$ (with globally bounded curvature). The specialisation of the main result of [35] to the cohomogeneity one setting therefore yields the following theorem.

Theorem 2.6.

- (i) *The manifold $M = S^3 \times \mathbb{R}^4$ described by the group diagram (2.4) carries a 1-parameter family (up to scale) of highly collapsed $\text{SU}(2)^2 \times \text{U}(1)$ -invariant ALC G_2 -metrics.*
- (ii) *For every pair of coprime positive integers m, n the manifold $M_{m,n}$ described by the group diagram (2.5) carries a 1-parameter family (up to scale) of highly collapsed $\text{SU}(2)^2 \times \text{U}(1)$ -invariant ALC G_2 -metrics.*

Proof. Given the main result of [35], we need only explain why the necessary topological constraint $c_1(M) \cup [\omega_0] = 0 \in H^4(B)$ is satisfied. The uniqueness (modulo diffeomorphism) of metrics in the construction of [35] implies that the continuous isometries $\text{SU}(2) \times \text{SU}(2)$ of the Calabi–Yau base B extend to M , and rotation in the circle fibres provides the additional $\text{U}(1)$. In case (i) the constraint is automatically satisfied since $H^4(B) = 0$. In case (ii) note that $B = K_{\mathbb{C}P^1 \times \mathbb{C}P^1}$ retracts onto its singular orbit $D = \mathbb{C}P^1 \times \mathbb{C}P^1$. Hence the 7-manifold $M_{m,n}$ retracts onto its singular orbit $M_{m,n}|_D$, a principal circle bundle over D . We can then understand the constraint $c_1(M_{m,n}) \cup [\omega_0] = 0$ by restriction to this singular orbit.

Fix a basis of left-invariant 1-forms $e_1, e_2, e_3, e'_1, e'_2, e'_3$ (the dual vector fields will be denoted by E_i, E'_i) on $SU(2) \times SU(2)$ with the property that

$$de_i = -e_j \wedge e_k, \quad de'_i = -e'_j \wedge e'_k \tag{2.7}$$

for (ijk) any cyclic permutation of (123) . When identifying \mathfrak{su}_2 with $\text{Im } \mathbb{H}$ we adopt the convention that $E_1 = i/2, E_2 = j/2, E_3 = k/2$; indeed by the first Maurer–Cartan structure equation, we must have $[E_i, E_j] = E_k$ in order for (2.7) to be satisfied. In particular, note that E_i, E'_i have period 4π . Without loss of generality we can assume that the maximal torus T^2 in $SU(2) \times SU(2)$ is generated by E_3 and E'_3 . The area forms of the two factors of $\mathbb{C}P^1 \times \mathbb{C}P^1 = SU(2) \times SU(2)/T^2$ are

$$\omega_1 = -\frac{1}{2}de_3 = \frac{1}{2}e_1 \wedge e_2, \quad \omega_2 = -\frac{1}{2}de'_3 = \frac{1}{2}e'_1 \wedge e'_2.$$

Now, $c_1(M_{m,n}|_D) = m[\omega_1] + n[\omega_2], [\omega_0|_D] = \alpha[\omega_1] - \beta[\omega_2]$ for some $\alpha, \beta > 0$ and we must have $m\beta - n\alpha = 0$, i.e. $(\alpha, \beta) = a(m, n)$ for some $a \neq 0$. In particular, m and n must have the same sign and so up to changing the circle bundle to its dual we can assume that both are positive. □

3. Invariant half-flat structures and Hitchin’s flow

Let M be a cohomogeneity one 7-manifold acted upon by $SU(2) \times SU(2)$ and described by one of the group diagrams (2.1), (2.4) or (2.5). The first step of our analysis is to describe torsion-free G_2 -structures on the open dense subset of principal orbits, i.e. on a cylinder of the form $(0, \infty) \times SU(2) \times SU(2)/K_0$, where K_0 is either trivial or the finite cyclic subgroup $\mathbb{Z}_{2|m+n|} = K_{m,n} \cap K_{2,-2}$ for coprime integers m, n . Any cohomogeneity one G_2 -structure on $(0, \infty) \times SU(2) \times SU(2)/K_0$ can be thought of as a 1-parameter family of invariant $SU(3)$ -structures on the principal orbit $SU(2) \times SU(2)/K_0$. The condition that the G_2 -structure be torsion-free can then be written as “static” and “evolution” equations for the corresponding family of $SU(3)$ -structures. The “static” equations constrain the torsion of the $SU(3)$ -structures; solutions are called *half-flat* structures. The “evolution” equations form a system of first-order ODEs known as *Hitchin’s flow*. The notions of stable forms and volume functionals introduced by Hitchin [39] allow one to interpret the ODE system as a Hamiltonian system on the space of invariant half-flat structures. We also give an alternative description of cohomogeneity one torsion-free G_2 -structures based on Hitchin’s description of torsion-free G_2 -structures as critical points of a volume functional on the space of closed stable 3-forms on a 7-manifold in a fixed cohomology class. Palais’ Principle of Symmetric Criticality (which holds in our context since the symmetry group is compact [48, Theorem 5.4]) allows us to give an alternative Lagrangian formulation of Hitchin’s flow.

3.1. Invariant half-flat structures on $SU(2) \times SU(2)$

The holonomy reduction of a Riemannian 7-manifold to G_2 is conveniently expressed as the existence of a closed and coclosed (in fact, parallel) 3-form φ with special algebraic properties at each point. The natural action of $GL(7, \mathbb{R})$ on $\Lambda^3(\mathbb{R}^7)^*$ has two open orbits;

one of these is isomorphic to $GL(7, \mathbb{R})/G_2$ and we say that a 3-form φ on a 7-manifold M is *positive* if φ_x lies in this orbit for every $x \in M$. Since the stabiliser of a positive 3-form is conjugate to G_2 , the existence of φ is equivalent to the reduction of the frame bundle of M to G_2 . Moreover, since G_2 is a subgroup of $SO(7)$, every positive 3-form φ defines a Riemannian metric g_φ and volume form dv_φ on M .

A G_2 -structure on a family of parallel hypersurfaces such as the principal orbits in a cohomogeneity one manifold is described by a 1-parameter family of *half-flat* $SU(3)$ -structures.

Definition 3.1. An $SU(3)$ -structure on a 6-manifold is a pair of smooth differential forms (ω, Ω) , where ω is a non-degenerate 2-form and Ω is a complex volume form, satisfying the algebraic constraints

$$\omega \wedge \operatorname{Re} \Omega = 0, \quad \frac{1}{6}\omega^3 = \frac{1}{4} \operatorname{Re} \Omega \wedge \operatorname{Im} \Omega. \tag{3.2}$$

A *half-flat structure* is an $SU(3)$ -structure (ω, Ω) such that

$$d\omega \wedge \omega = 0 = d \operatorname{Re} \Omega. \tag{3.3}$$

Invariant half-flat structures on $SU(2) \times SU(2)$ have been studied by Schulte-Hengesbach [51, Chapter 5] and Madsen–Salamon [45]. We now briefly summarise their results. It is useful to recall first the formal geometric set-up introduced by Hitchin [39, §6]. Let N be a compact 6-manifold. Let \mathcal{U} and \mathcal{V} be the space of closed *stable* 4-forms and 3-forms representing a fixed pair of cohomology classes on N of degree 4 and degree 3 respectively. Here a 4-form on N is stable if it can be written as $\frac{1}{2}\omega^2$ for a non-degenerate 2-form ω (uniquely determined up to sign) and a 3-form is stable if it is the real part of a holomorphic volume form Ω (in this case the imaginary part of Ω is uniquely determined by its real part). The tangent space of $\mathcal{U} \times \mathcal{V}$ at any point $(\frac{1}{2}\omega^2, \operatorname{Re} \Omega)$ is the product of affine spaces $\Omega_{\text{exact}}^4 \oplus \Omega_{\text{exact}}^3$. There is a non-degenerate pairing

$$\langle \sigma, \rho \rangle = \int_N \alpha \wedge \rho = - \int_N \sigma \wedge \beta,$$

where $\sigma = d\alpha$ is an exact 4-form and $\rho = d\beta$ an exact 3-form, which can be used to define a symplectic form Ψ on $\mathcal{U} \times \mathcal{V}$.

The diffeomorphism group of N acts naturally on $\mathcal{U} \times \mathcal{V}$ preserving Ψ . Given $(\frac{1}{2}\omega^2, \operatorname{Re} \Omega) \in \mathcal{U} \times \mathcal{V}$ and a vector field X on N , the infinitesimal action v_X of X on $\mathcal{U} \times \mathcal{V}$ is by Lie derivative and therefore

$$(v_X \lrcorner \Psi)(\sigma, \rho) = \int_N (X \lrcorner \frac{1}{2}\omega^2) \wedge \rho - \sigma \wedge (X \lrcorner \operatorname{Re} \Omega).$$

We claim that v_X is a Hamiltonian vector field on $\mathcal{U} \times \mathcal{V}$. Indeed, it is clear that the function μ_X on $\mathcal{U} \times \mathcal{V}$ defined by

$$\mu_X(\frac{1}{2}\omega^2, \operatorname{Re} \Omega) = \int_N (X \lrcorner \frac{1}{2}\omega^2) \wedge \operatorname{Re} \Omega = - \int_N \frac{1}{2}\omega^2 \wedge (X \lrcorner \operatorname{Re} \Omega)$$

satisfies $d\mu_X = v_X \lrcorner \Psi$. Moreover, since ω is non-degenerate, $X \mapsto X \lrcorner \omega$ is an isomorphism: rewriting

$$\mu_X(\frac{1}{2}\omega^2, \text{Re } \Omega) = \int_N (X \lrcorner \omega) \wedge \omega \wedge \text{Re } \Omega$$

we see that the vanishing of the moment map μ_X for all X is equivalent to $\omega \wedge \text{Re } \Omega = 0$.

Hitchin defines volume functionals $V(\frac{1}{2}\omega^2)$ and $V(\text{Re } \Omega)$ and a diffeomorphism-invariant functional H by taking a certain linear combination of these:

$$H(\frac{1}{2}\omega^2, \text{Re } \Omega) = \int_N 2\omega^3 - 3 \text{Re } \Omega \wedge \text{Im } \Omega.$$

By diffeomorphism invariance H descends to the symplectic quotient

$$\mathcal{M} = \mu^{-1}(0)/\text{Diff}_0(N).$$

The zero-level set of H in \mathcal{M} can almost be identified with the moduli space of half-flat structures on N in the given cohomology classes: the second constraint in (3.2) is only satisfied in an integral sense. However, if N is a homogeneous space and we restrict to invariant forms (and invariant diffeomorphisms) then the zero-level set of H in \mathcal{M} does indeed parametrise invariant half-flat structures on N with fixed cohomology classes.

We now specialise this general framework to the case where $N = \text{SU}(2) \times \text{SU}(2)$ (or, later, a finite free quotient of this). The group of Lie algebra inner automorphisms of $\mathfrak{su}_2 \oplus \mathfrak{su}_2$ (i.e. the group of invariant diffeomorphisms of N isotopic to the identity) is $\text{Aut} = \text{SO}(3) \times \text{SO}(3)$. Fix $(p, q) \in \mathbb{R}^2$. We consider the space \mathcal{U} of invariant non-degenerate 2-forms ω such that $d\omega \wedge \omega = 0$ (since $H^4(N) = 0$ the closed 4-form ω^2 is necessarily exact) and the space \mathcal{V} of invariant closed stable 3-forms $\text{Re } \Omega$ of the form

$$\text{Re } \Omega = p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d\eta$$

for some invariant 2-form η . Here e_i, e'_i denote the left-invariant 1-forms defined in (2.7). By [51, Chapter 5, Lemmas 1.1 and 1.3], \mathcal{U} and \mathcal{V} are each identified with open subsets of the space $M_{3 \times 3}$ of real 3×3 matrices via

$$\omega = \sum_{i,j} A_{ij} e_i \wedge e'_j, \quad \eta = \sum_{i,j} B_{ij} e_i \wedge e'_j.$$

Via the double cover $\text{SO}(4) \rightarrow \text{SO}(3) \times \text{SO}(3)$, we identify the $\text{SO}(3) \times \text{SO}(3)$ -representation $M_{3 \times 3}$ with the $\text{SO}(4)$ -representation $\text{Sym}_0^2(\mathbb{R}^4)$ [45, Lemma 1]. Then $(\mathcal{U} \times \mathcal{V}, \Psi)$ can be identified with $T^*\text{Sym}_0^2(\mathbb{R}^4)$ endowed with its canonical symplectic form. The vanishing of the moment map μ for the action of $\text{SO}(4)$ guarantees that the two matrices A and B , thought of as traceless symmetric 4×4 matrices, commute [45, Theorem 1]. By singular symplectic reduction, $\mu^{-1}(0)/\text{SO}(4)$ is the cotangent space of $\text{Sym}_0^2(\mathbb{R}^4)/\text{SO}(4)$ and is identified with $\mathbb{R}^3 \times \mathbb{R}^3/W$, where W is the symmetric group on three elements acting diagonally on $\mathbb{R}^3 \times \mathbb{R}^3$ [45, Corollary 1].

Concretely [51, Chapter 5, Theorem 1.4], up to the action of Aut we can assume that

$$\begin{aligned} \omega &= \alpha_1 e_1 \wedge e'_1 + \alpha_2 e_2 \wedge e'_2 + \alpha_3 e_3 \wedge e'_3, \\ \text{Re } \Omega &= p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a_1 e_1 \wedge e'_1 + a_2 e_2 \wedge e'_2 + a_3 e_3 \wedge e'_3). \end{aligned} \tag{3.4}$$

The 2-form ω is non-degenerate and $\text{Re } \Omega$ is stable if and only if, respectively, $\alpha_1, \alpha_2, \alpha_3 > 0$ and

$$\begin{aligned} \Lambda(a_1, a_2, a_3) &= a_1^4 + a_2^4 + a_3^4 - 2a_1^2 a_2^2 - 2a_2^2 a_3^2 - 2a_3^2 a_1^2 \\ &\quad + 4(p - q)a_1 a_2 a_3 + 2pq(a_1^2 + a_2^2 + a_3^2) + p^2 q^2 < 0. \end{aligned} \tag{3.5}$$

Moreover, the second constraint in (3.2) forces

$$2\alpha_1 \alpha_2 \alpha_3 = \sqrt{-\Lambda(a_1, a_2, a_3)}.$$

Note that there is a residual ambiguity in the parametrisation (3.4): the Weyl group W acts permuting $(\alpha_1, \alpha_2, \alpha_3)$ and (a_1, a_2, a_3) simultaneously.

Remark. Due to different choices of basis of \mathfrak{su}_2 , the definition of Λ in (3.5) is different from the one given by Schulte-Hengesbach [51, Chapter 5, equation (1.7)]. The two formulas are related by $a_i \mapsto -a_i$ for all $i = 1, 2, 3$.

Remark 3.6. In addition to the (left) $\text{SU}(2) \times \text{SU}(2)$ -invariance, when $\alpha_1 = \alpha_2$ and $a_1 = a_2$, the half-flat structure (3.4) is invariant under the right action of $\Delta\text{U}(1)$ (in fact, of the normaliser N of $\text{U}(1)$ in $\text{SU}(2)$ if we do not restrict to continuous symmetries) on $\text{SU}(2) \times \text{SU}(2)$. Similarly, if $\alpha_1 = \alpha_2 = \alpha_3$ and $a_1 = a_2 = a_3$ then the half-flat structure is invariant under the right action of $\Delta\text{SU}(2)$. The Bryant–Salamon AC G_2 -metric on the spinor bundle of S^3 [18] has the additional $\Delta\text{SU}(2)$ -symmetry; cf. Example 3.14. All the global examples we construct in this paper will have the additional $\Delta\text{U}(1)$ -symmetry.

The case of non-trivial stabiliser of the principal orbits. We now extend the previous discussion to the case where the stabiliser K_0 of the principal orbits is non-trivial. In the situation of interest, the isotropy representation of $K_0 = K_{m,n} \cap K_{2,-2} \simeq \mathbb{Z}_{2|m+n|}$ on $\mathfrak{su}_2 \oplus \mathfrak{su}_2$ is generated by the automorphism

$$T = \text{diag}(\zeta^{2|m|}, 1, \zeta^{2|m|}, 1), \tag{3.7}$$

where we identify $\text{span}(E_1, E_2)$ and $\text{span}(E'_1, E'_2)$ with \mathbb{C} , and ζ is a generator of $\mathbb{Z}_{2|m+n|}$. The subgroup Aut_T of $\text{Aut} = \text{SO}(3) \times \text{SO}(3)$ of inner automorphisms that commute with T is $\text{Aut}_T = \text{O}(2) \times \text{O}(2)$, where $\text{O}(2)$ is the subgroup of $\text{SO}(3)$ that fixes E_3 . Similarly, an invariant 2-form

$$\sum_{i,j} C_{ij} e_i \wedge e'_j$$

on $\text{SU}(2) \times \text{SU}(2)$ determined by a 3×3 matrix C descends to $\text{SU}(2) \times \text{SU}(2)/K_0$ if and only if $T^t \begin{pmatrix} 0 & C \\ -C^t & 0 \end{pmatrix} T = \begin{pmatrix} 0 & C \\ -C^t & 0 \end{pmatrix}$. A computation shows that

$$C = \begin{pmatrix} c_{11} & c_{12} & 0 \\ c_{21} & c_{22} & 0 \\ 0 & 0 & c_{33} \end{pmatrix},$$

with the upper-left 2×2 block commuting with the rotation of \mathbb{R}^2 of angle $\frac{2\pi|n|}{|n+m|}$. The latter condition forces $c_{11} = c_{22}$ and $c_{12} = -c_{21}$ unless $e^{\frac{2\pi i|n|}{|n+m|}} = \pm 1$. This can only happen if there exists $d \in \mathbb{Z}$ such that $(d + 1)m + (d - 1)n = 0$.

The constraints that K_0 -invariant forms must satisfy and the smaller group of automorphisms Aut_T play off against each other and, as in the case of trivial stabiliser of the principal orbit, we deduce the following proposition.

Proposition 3.8. *Up to the action of Aut_T any invariant half-flat structure (ω, Ω) on $\text{SU}(2) \times \text{SU}(2)/K_{m,n} \cap K_{2,-2}$ can be put in the normal form*

$$\begin{aligned} \omega &= \alpha_1 e_1 \wedge e'_1 + \alpha_2 e_2 \wedge e'_2 + \alpha_3 e_3 \wedge e'_3, \\ \text{Re } \Omega &= p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a_1 e_1 \wedge e'_1 + a_2 e_2 \wedge e'_2 + a_3 e_3 \wedge e'_3) \end{aligned}$$

with $\alpha_1, \alpha_2, \alpha_3, -\Lambda(a_1, a_2, a_3) > 0$ and $2\alpha_1\alpha_2\alpha_3 = \sqrt{-\Lambda(a_1, a_2, a_3)}$. Furthermore, $\alpha_1 = \alpha_2$ and $a_1 = a_2$ unless there exists $d \in \mathbb{Z}$ such that $(d + 1)m + (d - 1)n = 0$.

3.2. The fundamental ODE system

We now introduce the evolution equations for a 1-parameter family of invariant half-flat structures on $\text{SU}(2) \times \text{SU}(2)/K_0$ to define a torsion-free G_2 -structure.

The Hamiltonian formulation: Hitchin’s flow. Consider a G_2 -structure $\varphi = dt \wedge \omega + \text{Re } \Omega$ on a cylinder $(0, t_0) \times N$. Here (ω, Ω) is a 1-parameter family of $\text{SU}(3)$ -structures on the 6-manifold N . Then we have $*\varphi = -dt \wedge \text{Im } \Omega + \frac{1}{2}\omega^2$ and the condition that φ be closed and coclosed is equivalent to the half-flat equations (3.3) for (ω, Ω) together with the evolution equations

$$\partial_t \text{Re } \Omega = d\omega, \quad \partial_t(\omega^2) = -2d \text{Im } \Omega. \tag{3.9}$$

We specialise now to the case of a cohomogeneity one G_2 -structure on $M = (0, t_0) \times N$ with $N = \text{SU}(2) \times \text{SU}(2)/K_0$. Fix $p, q \in \mathbb{R}$ and consider a closed $\text{SU}(2) \times \text{SU}(2)$ -invariant 3-form

$$\varphi = p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a_1 e_1 \wedge e'_1 + a_2 e_2 \wedge e'_2 + a_3 e_3 \wedge e'_3), \tag{3.10}$$

where d denotes the differential in 7 dimensions. We rewrite φ as $\varphi = dt \wedge \omega + \text{Re } \Omega$ for a 1-parameter family of pairs of differential forms

$$\begin{aligned} \omega &= \dot{a}_1 e_1 \wedge e'_1 + \dot{a}_2 e_2 \wedge e'_2 + \dot{a}_3 e_3 \wedge e'_3, \\ \text{Re } \Omega &= p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a_1 e_1 \wedge e'_1 + a_2 e_2 \wedge e'_2 + a_3 e_3 \wedge e'_3), \end{aligned}$$

where now d is the differential on the 6-manifold $\text{SU}(2) \times \text{SU}(2)/K_0$ and $\dot{a}_i = \frac{da_i}{dt}$. Assuming that t is the arc-length parameter along a geodesic meeting all principal orbits orthogonally, the 3-form φ is a G_2 -structure if and only if the pair (ω, Ω) defines an $\text{SU}(3)$ -structure for all t , i.e.

$$\dot{a}_i > 0, \quad \Lambda(a_1, a_2, a_3) < 0, \quad 2\dot{a}_1\dot{a}_2\dot{a}_3 = \sqrt{-\Lambda(a_1, a_2, a_3)}.$$

Remark 3.11. The last condition is equivalent to the requirement that t be the arc-length parameter along a geodesic meeting all principal orbits orthogonally. In the following it will sometimes be convenient to drop this constraint. In that case, we will only require $\dot{a}_i > 0$ and $\Lambda(a_1, a_2, a_3) < 0$.

Set $y_i = a_i$ and $x_i = \dot{a}_i \dot{a}_k$. Then we have

$$\begin{aligned} \frac{1}{2}\omega^2 &= x_1 e_2 \wedge e'_2 \wedge e_3 \wedge e'_3 + x_2 e_3 \wedge e'_3 \wedge e_1 \wedge e'_1 + x_3 e_1 \wedge e'_1 \wedge e_2 \wedge e'_2, \\ \text{Re } \Omega &= p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(y_1 e_1 \wedge e'_1 + y_2 e_2 \wedge e'_2 + y_3 e_3 \wedge e'_3). \end{aligned}$$

We are going to rewrite (3.9) as an ODE system for the pair $(x, y) \in \mathbb{R}^3 \times \mathbb{R}^3$. In Hitchin’s formalism, the evolution equations (3.9) are interpreted as the Hamiltonian flow of the Hamiltonian function H on the space $\mathcal{U} \times \mathcal{V}$ of half-flat structures in fixed cohomology classes [39, Theorem 8]. Specialising to the $SU(2) \times SU(2)$ -invariant setting, the Hamiltonian H is given by

$$H(x, y) = \sqrt{-\Lambda(y_1, y_2, y_3)} - 2\sqrt{x_1 x_2 x_3}, \tag{3.12}$$

where Λ was defined in (3.5). Then (3.9) is equivalent to the Hamiltonian system

$$\begin{aligned} \dot{x}_i &= \frac{\partial H}{\partial y_i} = \frac{2}{\sqrt{-\Lambda(y_1, y_2, y_3)}} (y_i(-y_i^2 + y_j^2 + y_k^2 - pq) - (p - q)y_j y_k), \\ \dot{y}_i &= -\frac{\partial H}{\partial x_i} = \frac{x_j x_k}{\sqrt{x_1 x_2 x_3}}. \end{aligned} \tag{3.13}$$

In particular, H is constant along the flow. In fact $H = 0$ along the flow since we require the normalisation $2\omega^3 = 3 \text{Re } \Omega \wedge \text{Im } \Omega$.

Remark. Alternatively, one can use [51, Corollary 1.5] (with the usual change of sign due to our different choice of basis of \mathfrak{su}_2) to write

$$\begin{aligned} \sqrt{-\Lambda} \text{Im } \Omega &= (2y_1 y_2 y_3 - p(y_1^2 + y_2^2 + y_3^2 + pq)) e_1 \wedge e_2 \wedge e_3 \\ &\quad + (2y_1 y_2 y_3 + q(y_1^2 + y_2^2 + y_3^2 + pq)) e'_1 \wedge e'_2 \wedge e'_3 \\ &\quad + (y_i(y_i^2 - y_j^2 - y_k^2 + pq) - 2qy_j y_k) e_i \wedge e'_j \wedge e'_k \\ &\quad + (y_i(y_i^2 - y_j^2 - y_k^2 + pq) + 2py_j y_k) e'_i \wedge e_j \wedge e_k, \end{aligned}$$

with the convention that we sum over cyclic permutations (ijk) of (123) . The equivalence between (3.9) and (3.13) is then immediate.

Example 3.14. By Remark 3.6, solutions of (3.13) satisfying $x_1 = x_2 = x_3 = x$ and $y_1 = y_2 = y_3 = y$ correspond to $SU(2)^3$ -invariant torsion-free G_2 -structures. In this highly symmetric setting, (3.13) reduces to algebraic equations. Indeed, the vanishing of the Hamiltonian function H describes a curve $4x^3 = 3y^4 - 4(p - q)y^3 - 6pqy^2 - p^2q^2$ in the (x, y) -plane. The solution with $p = 0 = q$ induces a conical metric: the G_2 -cone \mathbb{C} over the homogeneous nearly Kähler structure on $S^3 \times S^3$. Since the asymptotic behaviour

of the curve $H(x, y) = 0$ for large y is independent of p and q , solutions for arbitrary p and q have one AC end asymptotic to C. As observed in [16, §2.1], only when $(p, q) = (r_0^3, -r_0^3), (-r_0^3, 0)$ or $(0, r_0^3)$ for some $r_0 > 0$, do solutions close smoothly on a singular orbit. Up to scale and (not necessarily $SU(2) \times SU(2)$ -equivariant) diffeomorphisms there exists a unique complete AC metric asymptotic to the cone C: the Bryant–Salamon metric on the spinor bundle of S^3 [18, §3].

The Lagrangian formulation. Later in the paper it will be useful to have a different formulation of (3.13), first introduced by Brandhuber [16].

Let φ be a closed G_2 -structure on a 7-manifold M and denote by g_φ the induced Riemannian metric. Hitchin [39, Theorem 1] showed that the equation $d^*\varphi = 0$ is the Euler–Lagrange equation for the volume functional $\varphi \mapsto \text{Vol}(M, g_\varphi)$ restricted to variations of φ amongst closed 3-forms with fixed cohomology class. Assume that $M = (0, s_0) \times SU(2) \times SU(2)/K_0$, with coordinate s on the first factor, and let φ be a closed $SU(2) \times SU(2)$ -invariant G_2 -structure of the form (3.10), i.e.

$$\varphi = p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a_1 e_1 \wedge e'_1 + a_2 e_2 \wedge e'_2 + a_3 e_3 \wedge e'_3).$$

Then up to a constant we have

$$\text{Vol}(M, g_\varphi) = \int_0^{s_0} L(a, a') ds, \quad L(a, a') = (-a'_1 a'_2 a'_3 \Lambda(a_1, a_2, a_3))^{1/3}. \quad (3.15)$$

Here $a'_i = \frac{da_i}{ds}$. In order to prove (3.15) observe that the fixed parameter s and the arc-length parameter t are related by

$$2 \left(\frac{ds}{dt} \right)^3 a'_1 a'_2 a'_3 = \sqrt{-\Lambda(a_1, a_2, a_3)}$$

and that the volume form of g_φ is

$$dv_{g_\varphi} = \frac{1}{6} dt \wedge \omega^3 = \dot{a}_1 \dot{a}_2 \dot{a}_3 dt \wedge e_1 \wedge e'_1 \wedge e_2 \wedge e'_2 \wedge e_3 \wedge e'_3.$$

By the Principle of Symmetric Criticality, we conclude that φ is coclosed if and only if the triple $a = (a_1, a_2, a_3)$ satisfies the second-order Lagrangian system

$$(\partial_{a'_i} L(a, a'))' - \partial_{a_i} L(a, a') = 0.$$

However, since Hitchin’s volume functional is invariant under diffeomorphisms, we know *a priori* that $(\partial_{a'_i} L(a, a'))' - \partial_{a_i} L(a, a')$ is orthogonal to the vector field (a'_1, a'_2, a'_3) . We therefore reduce to an ODE system of two second-order equations in three variables. This reformulation makes sense since we have the freedom to change the parametrisation s .

We carry out the relevant calculations explicitly in the case where $a_1 = a_2$, i.e. when there is an additional $U(1)$ -symmetry; cf. Remark 3.6. All the global solutions we find in this paper will admit this additional $U(1)$ -symmetry. Set $a := a_1 = a_2, b := a_3$ and

$$F(a, b) := -\Lambda(a, a, b) = 4a^2(b - p)(b + q) - (b^2 + pq)^2.$$

Let F_a and F_b denote the partial derivatives of F . Then we calculate

$$9L^5 \left(\frac{d}{ds} \left(\frac{\partial L}{\partial a'} \right) - \frac{\partial L}{\partial a} \right) = (a')^2 b' F(2F(a'b'' - b'a'') - a'b'(a'F_a - 2b'F_b)),$$

$$9L^5 \left(\frac{d}{ds} \left(\frac{\partial L}{\partial b'} \right) - \frac{\partial L}{\partial b} \right) = (a')^3 F(-2F(a'b'' - b'a'') + a'b'(a'F_a - 2b'F_b)).$$

Thus the pair of functions (a, b) yields a torsion-free $SU(2) \times SU(2) \times U(1)$ -invariant G_2 -structure if and only if

$$2F(a'b'' - b'a'') - a'b'(a'F_a - 2b'F_b) = 0. \tag{3.16}$$

As a sanity check, note that (3.16) can be immediately derived from (3.13). Indeed, in the presence of the additional $U(1)$ -symmetry, (3.13) becomes the ODE system

$$\begin{aligned} \dot{x}_1 &= \frac{F_a(y_1, y_2)}{4\sqrt{F(y_1, y_2)}}, & \dot{x}_2 &= \frac{F_b(y_1, y_2)}{2\sqrt{F(y_1, y_2)}}, \\ \dot{y}_1 &= \frac{x_1 x_2}{\sqrt{x_1^2 x_2}}, & \dot{y}_2 &= \frac{x_1^2}{\sqrt{x_1^2 x_2}}, \end{aligned} \tag{3.17}$$

for the four functions $x_1 = \dot{a}b, x_2 = \dot{a}^2, y_1 = a, y_2 = b$. Then (3.16) is an immediate consequence of (3.17). Moreover, the variable t in (3.17) is the arc-length parameter along a geodesic meeting all principal orbits orthogonally and therefore we have the further normalisation $2\dot{a}^2 b = \sqrt{F(a, b)}$; cf. Remark 3.11.

3.3. The induced metric

For later use, we briefly discuss properties of the map $\varphi \mapsto g_\varphi$ in the $SU(2) \times SU(2)$ -invariant setting. Lemma 3.19 below shows that Hitchin’s flow (3.13) can be regarded as an evolution equation for the family of Riemannian metrics induced by the 1-parameter family of half-flat structures. In fact, cohomogeneity one $SU(2) \times SU(2)$ -invariant G_2 -manifolds are described by a first-order ODE system for the metric coefficients in the work of Brandhuber–Gomis–Gubser–Gukov [17], Cvetič–Gibbons–Lü–Pope [23–25, 28], Hori–Hosomichi–Page–Rabadán–Walcher [40] and Bazaikin–Bogoyavlenskaya [9, 13].

Consider the $SU(2) \times SU(2)$ -invariant closed G_2 -structure

$$\varphi = p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a_1 e_1 \wedge e'_1 + a_2 e_2 \wedge e'_2 + a_3 e_3 \wedge e'_3).$$

The induced metric g_φ takes the form $g_\varphi = dt^2 + g_t$, where t is the arc-length parameter along a geodesic meeting all $SU(2) \times SU(2)$ -orbits orthogonally and g_t is a 1-parameter family of $SU(2) \times SU(2)$ -invariant metrics on the principal orbits. By [51, Corollary 1.5] (with the usual change of signs)

$$\frac{1}{2}\sqrt{-\Lambda} g_t = \dot{a}_i(a_j a_k - p a_i) e_i \otimes e_i + \dot{a}_i(a_j a_k + q a_i) e'_i \otimes e'_i + \dot{a}_i(a_i^2 - a_j^2 - a_k^2 - pq) e_i \otimes e'_i, \tag{3.18}$$

where (ijk) runs over cyclic permutations of (123) .

Remark. When $p + q = 0$ the metric g_t is invariant under the involution generated by the outer automorphism of $SU(2) \times SU(2)$ that exchanges the two factors. This additional symmetry however does not preserve the G_2 -structure φ .

Lemma 3.19. Fix $p, q \in \mathbb{R}$ and assume that $\dot{a}_1, \dot{a}_2, \dot{a}_3$ and $2\dot{a}_1\dot{a}_2\dot{a}_3 = \sqrt{-\Lambda(a_1, a_2, a_3)}$ are all positive. Then $(\dot{a}_1, \dot{a}_2, \dot{a}_3, a_1, a_2, a_3)$ is uniquely determined by the metric g_t (up to discrete symmetries when $p + q = 0$). In other words, the map that associates to each invariant half-flat structure in a given cohomology class (p, q) its induced metric is a (local) diffeomorphism.

Proof. By (3.18), up to the action of the automorphism group of $\mathfrak{su}_2 \oplus \mathfrak{su}_2$, the $SU(2) \times SU(2)$ -invariant metrics g that could possibly be induced by an invariant half-flat structure must be of the form

$$g = A_i e_i \otimes e_i + B_i e'_i \otimes e'_i + C_i e_i \otimes e'_i.$$

Furthermore, $A_i = B_i$ if $p + q = 0$. If the coefficients A_i, B_i, C_i are given by (3.18) then a straightforward computation shows that

$$4\dot{a}_j\dot{a}_k = \sqrt{(4A_j B_j - C_j^2)(4A_k B_k - C_k^2)}.$$

Thus $\dot{a}_1, \dot{a}_2, \dot{a}_3$ are uniquely determined by the metric g . Furthermore, $(p + q)a_i = \dot{a}_j\dot{a}_k(B_i - A_i)$ and therefore a_1, a_2, a_3 are also uniquely determined whenever $p + q \neq 0$.

If $p + q = 0$ we calculate instead

$$\begin{aligned} \dot{a}_j\dot{a}_k(A_i + B_i + C_i) &= (a_j - a_k - a_i + p)(a_k - a_j - a_i + p), \\ -\dot{a}_j\dot{a}_k(A_i + B_i - C_i) &= (a_1 + a_2 + a_3 + p)(a_i - a_j - a_k + p), \end{aligned}$$

from which it follows, using $A_i = B_i$, that

$$\Lambda(a_1, a_2, a_3) = (a_1 + a_2 + a_3 + p)(a_1 - a_2 - a_3 + p)(a_2 - a_3 - a_1 + p)(a_3 - a_1 - a_2 + p).$$

Together with $2\dot{a}_1\dot{a}_2\dot{a}_3 = \sqrt{-\Lambda(a_1, a_2, a_3)}$, these relations allow us to determine $a_i - a_j - a_k + p$ up to sign:

$$4(a_i - a_j - a_k + p)^2 = (A_i + B_i - C_i)(A_j + B_j + C_j)(A_k + B_k + C_k). \quad \square$$

A Riemannian quantity that will play an important role later in the paper is the mean curvature of the principal orbits in an $SU(2) \times SU(2)$ -invariant G_2 -manifold. Consider a 1-parameter family of invariant half-flat structures satisfying Hitchin’s flow (3.13). The shape operator L of the hypersurface $\{t = \text{const}\}$ is $L = \frac{1}{2}g_t^{-1}\dot{g}_t$, where the dot stands for time differentiation. The mean curvature is then $l = \text{tr}(L)$ or alternatively the time derivative of the logarithm of the orbital volume function $2\dot{a}_1\dot{a}_2\dot{a}_3 = \sqrt{-\Lambda(a_1, a_2, a_3)}$. We easily calculate

$$l = \frac{1}{2(\dot{a}_1\dot{a}_2\dot{a}_3)^2} \sum_{i=1}^3 \dot{a}_i (a_i(-a_i^2 + a_j^2 + a_k^2 - pq) - (p - q)a_j a_k). \quad (3.20)$$

4. Local solutions in a neighbourhood of the singular orbit

We are interested in 7-manifolds M with a cohomogeneity one action of $G = \text{SU}(2) \times \text{SU}(2)$ and one singular orbit $Q = G/K$, where up to automorphisms of G the stabiliser K can be taken as one of the following:

$$\Delta\text{SU}(2), \quad \{1\} \times \text{SU}(2), \quad K_{m,n}$$

for two coprime integers m, n , with $K_{m,n}$ as defined in (2.2). Note that $Q \simeq S^3$ in the first two cases and $Q \simeq S^2 \times S^3$ when $K = K_{m,n}$. The only reason we restrict to coprime m, n is that the resulting 7-manifold M is simply connected in this case. We recover the general case by taking finite quotients of the simply connected manifolds.

A tubular neighbourhood of the singular orbit Q in M is equivariantly diffeomorphic to a neighbourhood of the zero section in the vector bundle

$$G \times_K V \rightarrow Q,$$

where V is an orthogonal representation of K of dimension $d = 1 + \dim(K)$: V is the standard representation of $\text{SU}(2)$ when $K = \Delta\text{SU}(2)$ or $\{1\} \times \text{SU}(2)$, while V is the irreducible 2-dimensional real representation with weight $2|m + n|$ when $K = K_{m,n} \simeq \text{SO}(2)$. Fix a vector v_0 in the unit sphere in V and denote by K_0 the stabiliser of v_0 in K . Then K_0 is the principal orbit stabiliser; K_0 is trivial when $K = \Delta\text{SU}(2)$ and $K = \{1\} \times \text{SU}(2)$, and $K_0 \simeq \mathbb{Z}_{2|m+n|}$ when $K = K_{m,n}$.

In this section we derive explicit conditions on p, q and the a_i so that the invariant 3-form

$$\varphi = p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a_1 e_1 \wedge e'_1 + a_2 e_2 \wedge e'_2 + a_3 e_3 \wedge e'_3)$$

defines a smooth invariant closed positive 3-form in a neighbourhood of the zero section in $G \times_K V$. We then use this analysis to set up and study singular initial value problems for the ODE system (3.13) that correspond to local cohomogeneity one torsion-free G_2 -structures defined in a neighbourhood of the three types of singular orbit. The main result of the section is Proposition 4.5 where we parametrise the space of cohomogeneity one torsion-free G_2 -structures defined in a neighbourhood of each of the three types of singular orbit.

4.1. Smooth extension over the singular orbit

An efficient way of understanding conditions for the smooth extension of tensors along a singular orbit $Q = G/K$ in a cohomogeneity one manifold $M = G \times_K V$ has been given by Eschenburg–Wang [32, §1]. We briefly recall their approach. Write $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ where \mathfrak{g} and \mathfrak{k} are the Lie algebras of G and K respectively. Given a point $q \in Q$ we can identify $T_q Q$ with \mathfrak{p} and $T_q M$ with $\mathfrak{p} \oplus V$. By G -invariance, the 3-form φ is uniquely determined by its restriction to the fibre of $G \times_K V$ over q . We can therefore interpret φ as a K -equivariant map $\varphi: V \rightarrow \Lambda^3(\mathfrak{p} \oplus V)^*$. Furthermore, if we fix a point v_0 in the unit sphere \mathbb{S}^{d-1} in V and denote by K_0 its stabiliser in K , by K -equivariance $\varphi: V \rightarrow \Lambda^3(\mathfrak{p} \oplus V)^*$

is uniquely determined by the curve $t \mapsto \varphi(tv_0)$, which must lie in the subspace of K_0 -invariant 3-forms on $\mathfrak{p} \oplus V$. Similarly, every K -equivariant map from the unit sphere in V to $\Lambda^3(\mathfrak{p} \oplus V)^*$ is uniquely determined by its value at v_0 . Eschenburg and Wang show that $\varphi: V \rightarrow \Lambda^3(V \oplus \mathfrak{p})^*$ defines a smooth G -invariant 3-form if and only if for all $p \geq 0$ there exists a homogeneous degree p polynomial $\varphi_p: \mathbb{S}^{d-1} \rightarrow \Lambda^3(V \oplus \mathfrak{p})^*$ with values in the subspace of K_0 -invariant forms such that $t \mapsto \varphi(tv_0)$ has Taylor series $\sum_{p \geq 0} \varphi_p(v_0) t^p$.

Proposition 4.1. *Set $G = \text{SU}(2) \times \text{SU}(2)$ and (K, V) to be one of $(\Delta\text{SU}(2), \mathbb{C}^2)$, $(\{1\} \times \text{SU}(2), \mathbb{C}^2)$ or $(K_{m,n}, \mathbb{R}^2_{2|m+n|})$, where \mathbb{C}^2 denotes the standard representation of $\text{SU}(2)$ and $\mathbb{R}^2_{2|m+n|}$ is the irreducible real 2-dimensional representation of $K_{m+n} \simeq \text{SO}(2)$ with weight $2|m+n|$. Let*

$$\varphi = p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a_1 e_1 \wedge e'_1 + a_2 e_2 \wedge e'_2 + a_3 e_3 \wedge e'_3)$$

be an invariant closed 3-form defined on the complement of the zero section in a disc subbundle of the vector bundle $G \times_K V \rightarrow G/K$.

- (i) *If $K = \Delta\text{SU}(2)$ then φ extends as a smooth positive 3-form over the zero section of $G \times_K V$ if and only if*
 - (a) $p + q = 0$;
 - (b) for $i = 1, 2, 3$, a_i is an even function of t with $a_i(t) = p + \frac{1}{2}\alpha t^2 + O(t^4)$ for some $\alpha \in \mathbb{R}$;
 - (c) $8\alpha^3 = p > 0$.
- (ii) *If $K = \{1\} \times \text{SU}(2)$ then φ extends as a smooth positive 3-form over the zero section of $G \times_K V$ if and only if*
 - (a) $q = 0$;
 - (b) for $i = 1, 2, 3$, a_i is an even function of t vanishing at the origin with $a_i = \frac{1}{2}\alpha_i t^2 + O(t^4)$ for some $\alpha_i > 0$;
 - (c) $8\alpha_1\alpha_2\alpha_3 = -p > 0$.
- (iii) *If $K = K_{m,n}$ then φ extends as a smooth positive 3-form over the zero section of $G \times_K V$ if and only if*
 - (a) $mn > 0$;
 - (b) $p = -m^2r_0^3$ and $q = n^2r_0^3$ for some $r_0 \neq 0$;
 - (c) a_3 is an even function of t with $a_3(0) = mnr_0^3$ and $\ddot{a}_3(0) > 0$;
 - (d) $a_1 + a_2$ is an odd function of t and $\dot{a}_1(0) + \dot{a}_2(0) > 0$;
 - (e) $a_1 = a_2$ unless $m = n = \pm 1$; in the latter case $a_1 - a_2$ is an even function of t and $\alpha = \frac{1}{2}(a_1(0) - a_2(0))$ satisfies $\alpha^2 < r_0^6$.

Remark. In Theorem 7.3 we will prove that, up to finite quotients, the proposition covers all possible singular orbits of smooth G_2 -metrics with enhanced $\text{SU}(2) \times \text{SU}(2) \times \text{U}(1)$ -symmetry.

We now prove the three cases of the proposition.

The case $K = \Delta\text{SU}(2)$. The singular orbit is $Q = \text{SU}(2) \times \text{SU}(2)/\Delta\text{SU}(2) \simeq S^3$ and the stabiliser of points on principal orbits is trivial. A tubular neighbourhood of the singular orbit S^3 is equivariantly diffeomorphic to $S^3 \times \mathbb{R}^4$, where $\text{SU}(2) \times \text{SU}(2)$ acts on $S^3 \times \mathbb{R}^4 \subset \mathbb{H} \oplus \mathbb{H}$ by $(q_1, q_2) \cdot (x, y) = (q_2 x q_1^*, q_1 y)$. Therefore along the ray $\gamma(t) = (1, t) \in S^3 \times \mathbb{H} \subset \mathbb{H}^2$ we have

$$\begin{aligned} E_1 &= \frac{1}{2}(-i, it), & E_2 &= \frac{1}{2}(-j, jt), & E_3 &= \frac{1}{2}(-k, kt), \\ E'_1 &= \frac{1}{2}(i, 0), & E'_2 &= \frac{1}{2}(j, 0), & E'_3 &= \frac{1}{2}(k, 0). \end{aligned}$$

In particular, if we define $e_i^\pm = \frac{1}{2}(e_i \pm e'_i)$ and let t, x_1, x_2, x_3 be Euclidean coordinates on \mathbb{R}^4 we have

$$e_1^+ = 2t^{-1}dx_1, \quad e_2^+ = 2t^{-1}dx_2, \quad e_3^+ = 2t^{-1}dx_3.$$

For $t \neq 0$ the closed positive 3-form φ can then be written as

$$\begin{aligned} \frac{1}{2}\varphi &= V e_1^- \wedge e_2^- \wedge e_3^- + \sum_{i=1}^3 e_i^- \wedge \omega_i + 8(p+q)t^{-3}dx_1 \wedge dx_2 \wedge dx_3 \\ &\quad + 2(p+q)t^{-1} \sum_{i=1}^3 dx_i \wedge e_j^- \wedge e_k^-. \end{aligned}$$

Here $\omega_i = -2\dot{a}_i t^{-1}dt \wedge dx_i + 4V_i t^{-2}dx_j \wedge dx_k$ and

$$V = a_1 + a_2 + a_3 + \frac{p-q}{2}, \quad V_i = a_i - a_j - a_k + \frac{p-q}{2}. \tag{4.2}$$

If φ extends smoothly at $t = 0$ then clearly $p + q = 0$, since otherwise the coefficient of $dx_1 \wedge dx_2 \wedge dx_3$ would blow up. In order to detect subtler conditions for the smooth extension along the singular orbit, following Eschenburg–Wang’s analysis we regard φ as a map $\varphi: \mathbb{H} \rightarrow \Lambda^3(\mathbb{H} \oplus \text{Im } \mathbb{H})$ identifying vector spaces with their duals using their standard metrics. As an $\text{SU}(2)$ -representation we have

$$\begin{aligned} \Lambda^3(\mathbb{H} \oplus \text{Im } \mathbb{H}) &= \Lambda^3\mathbb{H} \oplus (\Lambda^+\mathbb{H} \otimes \text{Im } \mathbb{H}) \oplus (\Lambda^-\mathbb{H} \otimes \text{Im } \mathbb{H}) \oplus (\mathbb{H} \otimes \Lambda^2 \text{Im } \mathbb{H}) \oplus (\Lambda^3 \text{Im } \mathbb{H}) \\ &\simeq \mathbb{H} \oplus (\text{Im } \mathbb{H} \otimes \text{Im } \mathbb{H}) \oplus (\mathbb{R}^3 \otimes \text{Im } \mathbb{H}) \oplus (\mathbb{H} \otimes \text{Im } \mathbb{H}) \oplus \mathbb{R}, \end{aligned}$$

where \mathbb{H} and $\text{Im } \mathbb{H}$ are the standard and adjoint representations of $\text{SU}(2)$, respectively, and we used the fact that the induced action of $\text{SU}(2)$ on $\Lambda^2\mathbb{H}$ acts trivially on anti-self-dual forms and acts via the adjoint representation on the space of self-dual forms (with respect to the volume form $dt \wedge dx_1 \wedge dx_2 \wedge dx_3$). By applying Eschenburg–Wang’s analysis we deduce conditions for the extension of φ over the singular orbit as a smooth 3-form:

- (i) $e_1^- \wedge e_2^- \wedge e_3^-$ is $\text{SU}(2)$ -invariant and therefore V must be even.
- (ii) $e_i^- \wedge \omega_i^- \in \mathbb{R}^3 \times \text{Im } \mathbb{H}$ corresponds to a degree 2 polynomial of the form $q \mapsto quq^*$ for $q \in \mathbb{H}$ and some fixed $u \in \text{Im } \mathbb{H}$. Here ω_i^- is the anti-self-dual part of ω_i . We deduce that

$$\frac{t\dot{a}_i + 2V_i}{t^2}$$

must be even and vanish at $t = 0$.

(iii) The triple $(e_1^- \wedge \omega_1^+, e_2^- \wedge \omega_2^+, e_3^- \wedge \omega_3^+)$ represents a map $\text{Im } \mathbb{H} \rightarrow \text{Im } \mathbb{H}$ which is diagonal in the standard basis of $\text{Im } \mathbb{H}$ and therefore symmetric. We decompose this map into a multiple of the identity and a traceless part. The equivariant polynomial $\mathbb{H} \rightarrow \text{Sym}(\text{Im } \mathbb{H})$ which corresponds to the identity by evaluation at 1 is clearly the constant polynomial $q \mapsto [u \mapsto u]$. On the other hand, an equivariant map $\mathbb{H} \rightarrow \text{Sym}_0(\text{Im } \mathbb{H})$ which has value $A \in \text{Sym}_0(\text{Im } \mathbb{H})$ at $q = 1$ must correspond to the degree 4 polynomial $q \mapsto \text{Ad}_q \circ A \circ \text{Ad}_{q^*}$. We therefore conclude that

$$\frac{t\dot{a}_i - 2V_i}{t^2}$$

must have an even Taylor series expansion with 0th and 2nd order coefficients independent of i .

In summary we must have

$$p + q = 0, \quad a_i = p + \frac{1}{2}\alpha t^2 + O(t^4)$$

for some $\alpha \in \mathbb{R}$.

Finally, the requirements that φ be a positive 3-form and t be the arc-length parameter along a geodesic meeting orthogonally all principal orbits impose further constraints. As in the proof of Lemma 3.19, it is useful to observe that when $q = -p$ we can factor $\Lambda(a_1, a_2, a_3)$ as $V V_1 V_2 V_3$, where V, V_i are defined in (4.2). Since $V \approx 4p + \frac{3}{2}\alpha t^2$ and $V_i \approx -\frac{1}{2}\alpha t^2$ as $t \rightarrow 0$, at leading order in t the conditions $\dot{a}_i, -\Lambda(a_1, a_2, a_3) > 0$ and $2\dot{a}_1\dot{a}_2\dot{a}_3 = \sqrt{-\Lambda(a_1, a_2, a_3)}$ are equivalent to the requirement

$$8\alpha^3 = p > 0.$$

The case $K = \{1\} \times \text{SU}(2)$. In this case we have $Q = S^3$ and $M = S^3 \times \mathbb{R}^4$ with action of $\text{SU}(2) \times \text{SU}(2)$ given by the left multiplication by unit quaternions on $S^3 \subset \mathbb{H}$ and $\mathbb{R}^4 \simeq \mathbb{H}$. Hence as K -representations we have $V = \mathbb{H}$ and $\mathfrak{p} = \text{Im } \mathbb{H}$. The 1-forms e_1, e_2, e_3 define a coframe on \mathfrak{p} . If t, x_1, x_2, x_3 are Euclidean coordinates on V , then along the ray $t \in \mathbb{H} \simeq V$ we have

$$e'_i = 2t^{-1}dx_i, \quad i = 1, 2, 3.$$

Hence we can write

$$\begin{aligned} \varphi &= p e_1 \wedge e_2 \wedge e_3 + 8qt^{-3} dx_1 \wedge dx_2 \wedge dx_3 \\ &+ \sum_{i=1}^3 t^{-1}(-\dot{a}_i + 2t^{-1}a_i)e_i \wedge (dt \wedge dx_i + dx_j \wedge dx_k) \\ &- t^{-1}(\dot{a}_i + 2t^{-1}a_i)e_i \wedge (dt \wedge dx_i - dx_j \wedge dx_k) - 2t^{-1}a_i dx_i \wedge e_j \wedge e_k, \end{aligned}$$

where for each $i = 1, 2, 3$ we have $\epsilon_{ijk} = 1$. Now, for φ to be smooth we must certainly have $q = 0$. Moreover, Eschenburg–Wang’s analysis implies that necessary and sufficient conditions for the smoothness of φ are that for all $i = 1, 2, 3$,

- (i) $2t^{-1}(\dot{a}_i \pm 2t^{-1}a_i)$ has Taylor series involving only even powers of t and also $2t^{-1}(\dot{a}_i - 2t^{-1}a_i)$ vanishes at the origin;
- (ii) $2t^{-1}a_i$ has Taylor series only involving odd powers of t .

These conditions are satisfied if and only if a_1, a_2, a_3 are smooth even functions vanishing at $t = 0$. Finally, at leading order in t the requirement that φ be a positive 3-form and t be the arc-length parameter along a geodesic meeting all orbits imposes the further constraints

$$\alpha_i > 0, \quad \delta\alpha_1\alpha_2\alpha_3 = -p > 0,$$

where we write $a_i(t) = \frac{1}{2}\alpha_i t^2 + O(t^4)$.

The case $K = K_{m,n}$. In this case the singular orbit is a circle bundle over

$$D = S^2 \times S^2 = \text{SU}(2) \times \text{SU}(2)/T^2$$

and the 7-manifold M is the total space of a $\mathbb{C} \times \mathbb{S}^1$ -bundle over D . More precisely, $Q = G \times_{T^2} S^1$ and $M = G \times_{T^2} (S^1 \times \mathbb{C})$, where T^2 acts on S^1 and \mathbb{C} with weights (m, n) and $(2, -2)$ respectively. The $K_{m,n}$ -representations V and \mathfrak{p} are naturally induced by the T^2 -representations $V = \mathbb{C}_{2,-2}$ and $\mathfrak{p} = \mathbb{R} \oplus \mathbb{C}_{2,0} \oplus \mathbb{C}_{0,2}$. Since m and n are coprime, $K_{m,n} \simeq \text{U}(1)$ is embedded in T^2 via $e^{i\theta} \mapsto (e^{in\theta}, e^{-im\theta})$ and as real $K_{m,n}$ -representations we have $V = \mathbb{R}^2_{2|n+m|}$ and $\mathfrak{p} = \mathbb{R} \oplus \mathbb{R}^2_{2|n|} \oplus \mathbb{R}^2_{2|m|}$.

It is convenient to introduce a new \mathbb{Z} -basis for the Lie algebra of the maximal torus T^2 in $\text{SU}(2) \times \text{SU}(2)$: fix integers (r, s) with $mr + ns = 1$ and consider the basis elements

$$nE_3 - mE'_3, \quad rE_3 + sE'_3.$$

The dual basis of left-invariant 1-forms is $se_3 - re'_3, me_3 + ne'_3$. Note that

$$e_3 = n(se_3 - re'_3) + r(me_3 + ne'_3), \quad e'_3 = -m(se_3 - re'_3) + s(me_3 + ne'_3).$$

Let t, x be coordinates on $V \simeq \mathbb{R}^2$. Since $nE_3 - mE'_3$ has period $\frac{2\pi}{|m+n|}$ in $\text{SU}(2) \times \text{SU}(2)/K_0$, along the ray t in V we must have (up to changing x into $-x$)

$$(n + m)(se_3 - re'_3) = t^{-1} dx.$$

On the other hand e_1, e_2 and e'_1, e'_2 are coframes on $\mathfrak{n} = \mathbb{R}^2_{2|n|}$ and $\mathfrak{n}' = \mathbb{R}^2_{2|m|}$ respectively and $me_3 + ne'_3$ generates the trivial real factor in \mathfrak{p} .

Although we work with real representations of $K_{m,n}$, in order to apply Eschenburg–Wang’s analysis it is convenient to complexify $V \oplus \mathfrak{p}$ and work with complex $\text{U}(1)$ -representations instead. We have

$$V \otimes \mathbb{C} = \mathbb{C}_{2(n+m)} \oplus \mathbb{C}_{-2(n+m)}, \quad \mathbb{R} \otimes \mathbb{C} = \mathbb{C}_0, \quad \mathfrak{n} \otimes \mathbb{C} = \mathbb{C}_{2n} \oplus \mathbb{C}_{-2n}, \quad \mathfrak{n}' = \mathbb{C}_{-2m} \oplus \mathbb{C}_{2m}.$$

Introduce the following basis of $(V \oplus \mathfrak{p}) \otimes \mathbb{C}$ (identified with its dual) adapted to the decomposition of $(V \oplus \mathfrak{p}) \otimes \mathbb{C}$ into $K_{m,n}$ -representations:

$$\begin{aligned} dz_V &= dt + idx = dt + i(n + m)(se_3 - re'_3), \\ d\bar{z}_V &= dt - idx = dt - i(n + m)(se_3 - re'_3), \\ dz_{\mathbb{R}} &= me_3 + ne'_3, \\ dz_{\mathfrak{n}} &= e_1 + ie_2, \quad d\bar{z}_{\mathfrak{n}} = e_1 - ie_2, \quad dz_{\mathfrak{n}'} = e'_1 + ie'_2, \quad d\bar{z}_{\mathfrak{n}'} = e'_1 - ie'_2. \end{aligned}$$

We can then rewrite φ as

$$\begin{aligned} \varphi = & \frac{i}{2}(pr - sa_3)dz_{\mathbb{R}} \wedge dz_n \wedge d\bar{z}_n + \frac{i}{2}(a_3r + sq)dz_{\mathbb{R}} \wedge dz_{n'} \wedge d\bar{z}_{n'} \\ & + \frac{i}{2} \frac{pn + ma_3}{(m+n)t} \operatorname{Im}(dz_V) \wedge dz_n \wedge d\bar{z}_n + \frac{i}{2} \frac{a_3n - mq}{(m+n)t} \operatorname{Im}(dz_V) \wedge dz_{n'} \wedge d\bar{z}_{n'} \\ & + \frac{i}{2} \frac{\dot{a}_3}{(m+n)t} dz_{\mathbb{R}} \wedge dz_V \wedge d\bar{z}_V \\ & + \frac{r+s}{2}(a_1 - a_2)dz_{\mathbb{R}} \wedge \operatorname{Im}(dz_n \wedge dz_{n'}) - \frac{r-s}{2}(a_1 + a_2)dz_{\mathbb{R}} \wedge \operatorname{Im}(dz_n \wedge d\bar{z}_{n'}) \\ & + \frac{t(\dot{a}_1 + \dot{a}_2) - (a_1 + a_2)}{4t} \operatorname{Re}(dz_V \wedge dz_n \wedge d\bar{z}_{n'}) \\ & + \frac{t(\dot{a}_1 + \dot{a}_2) + (a_1 + a_2)}{4t} \operatorname{Re}(d\bar{z}_V \wedge dz_n \wedge d\bar{z}_{n'}) \\ & + \frac{t(\dot{a}_1 - \dot{a}_2) + \frac{m-n}{m+n}(a_1 - a_2)}{4t} \operatorname{Re}(dz_V \wedge dz_n \wedge dz_{n'}) \\ & + \frac{t(\dot{a}_1 - \dot{a}_2) - \frac{m-n}{m+n}(a_1 - a_2)}{4t} \operatorname{Re}(d\bar{z}_V \wedge dz_n \wedge dz_{n'}). \end{aligned}$$

Working with complex representations makes it very easy to understand which $U(1)$ -representation each component of φ in this decomposition belongs to. We collect the weights of the corresponding real representations in the following table.

$dz_{\mathbb{R}} \wedge dz_n \wedge d\bar{z}_n$	0	$dz_{\mathbb{R}} \wedge dz_{n'} \wedge d\bar{z}_{n'}$	0
$\operatorname{Im}(dz_V) \wedge dz_n \wedge d\bar{z}_n$	$2 m+n $	$\operatorname{Im}(dz_V) \wedge dz_{n'} \wedge d\bar{z}_{n'}$	$2 m+n $
$dz_{\mathbb{R}} \wedge dz_V \wedge d\bar{z}_V$	0	$dz_{\mathbb{R}} \wedge \operatorname{Im}(dz_n \wedge dz_{n'})$	$2 m-n $
$dz_{\mathbb{R}} \wedge \operatorname{Im}(dz_n \wedge d\bar{z}_{n'})$	$2 m+n $	$\operatorname{Re}(dz_V \wedge dz_n \wedge d\bar{z}_{n'})$	$4 m+n $
$\operatorname{Re}(d\bar{z}_V \wedge dz_n \wedge d\bar{z}_{n'})$	0	$\operatorname{Re}(dz_V \wedge dz_n \wedge dz_{n'})$	$4 n $
$\operatorname{Re}(d\bar{z}_V \wedge dz_n \wedge dz_{n'})$	$4 m $		

The principal orbits have non-trivial stabiliser $K_0 = \mathbb{Z}_{2|m+n|}$ and all terms that are not K_0 -invariant must vanish. Note that $\mathbb{R}_{2|m-n|}^2$, $\mathbb{R}_{4|m|}^2$ and $\mathbb{R}_{4|n|}^2$ are trivial K_0 -representations if and only if there exists $d \in \mathbb{Z}$ such that $(d+1)m + (d-1)n = 0$. If this is not the case we must have $a_1 = a_2$, as we already know from Proposition 3.8.

From the table above we read off the necessary and sufficient conditions for φ to extend smoothly along the principal orbit:

- (i) a_3 is even and there exists $r_0 \in \mathbb{R}$ such that

$$p = -m^2r_0^3, \quad q = n^2r_0^3, \quad a_3(0) = mnr_0^3;$$

- (ii) $a_1 + a_2$ is odd;
- (iii) if there exists $d \in \mathbb{Z}$ such that $(d+1)m + (d-1)n = 0$ then the Taylor series of $a_1 - a_2$ only involves monomials of the form $t^{|d|+2l}$, $l \geq 0$ (in fact $l \geq 1$ if $d \neq 0$); otherwise $a_1 = a_2$.

Indeed, note that if $(d + 1)m + (d - 1)n = 0$ then $m - n = -d(m + n)$, $2m = -(d - 1)(m + n)$ and $2n = (d + 1)(m + n)$.

Finally, we determine constraints imposed by the requirement that φ defines a G_2 -structure for small t . Note that the conditions (i), (ii) and (iii) above allow us to write

$$a_1 = \alpha + \beta t + \gamma t^2 + O(t^3), \quad a_2 = -\alpha + \beta t - \gamma t^2 + O(t^3), \quad a_3 = mn r_0^3 + \delta t^2 + O(t^4),$$

with $\alpha = 0 = \gamma$ unless $m = n = \pm 1$ (i.e. $d = 0$). Thus $\beta, \delta > 0$ guarantee that $\dot{a}_i > 0$ for $i = 1, 2, 3$. Moreover, we calculate that the first three coefficients of the Taylor series of $\Lambda(a_1, a_2, a_3)$ at $t = 0$ are

$$4mnr_0^6(m - n)^2\alpha^2, \quad 0, \quad -4r_0^6\beta^2mn(m + n)^2 + 16\alpha^2\beta^2.$$

Note that the first coefficient, i.e. the value of $\Lambda(a_1, a_2, a_3)$ at $t = 0$, always vanishes, in agreement with the requirement $2\dot{a}_1\dot{a}_2\dot{a}_3 = \sqrt{-\Lambda(a_1, a_2, a_3)}$. Hence $\Lambda(a_1, a_2, a_3) < 0$ for small $t > 0$ if and only if $mn > 0$ and $\alpha^2 < r_0^6$ when $n = m = \pm 1$. Note also that once we know that m and n have the same sign, then the only way that $(d + 1)m + (d - 1)n = 0$ for some $d \in \mathbb{Z}$ is if $d = 0$ and $m = n$ (and therefore $m = n = \pm 1$ since we assume that $\gcd(m, n) = 1$).

4.2. The singular IVP

We now aim to construct and parametrise solutions to the fundamental ODE system (3.13) satisfying the smoothness conditions of Proposition 4.1. The main tool is the following existence result for a special type of singular initial value problems; cf. [32, §5; 33, §4; 46, Theorem 7.1].

Theorem 4.3. *Consider the singular initial value problem*

$$\dot{y} = \frac{1}{t}M_{-1}(y) + M(t, y), \quad y(0) = y_0, \tag{4.4}$$

where y takes values in \mathbb{R}^k , $M_{-1}: \mathbb{R}^k \rightarrow \mathbb{R}^k$ is a smooth function of y in a neighbourhood of y_0 and $M: \mathbb{R} \times \mathbb{R}^k \rightarrow \mathbb{R}^k$ is smooth in t, y in a neighbourhood of $(0, y_0)$. Assume that

- (i) $M_{-1}(y_0) = 0$;
- (ii) $h \text{Id} - d_{y_0}M_{-1}$ is invertible for all $h \in \mathbb{N}, h \geq 1$.

Then there exists a unique solution $y(t)$ of (4.4). Furthermore y depends continuously on y_0 satisfying (i) and (ii).

The main results of the section are summarised in the following proposition. In the statement, t is the arc-length parameter along a geodesic meeting all principal orbits orthogonally.

Proposition 4.5. *Let M be one of the cohomogeneity one 7-manifolds considered in Proposition 4.1, labelled by the stabiliser K of the singular orbit. Consider $SU(2) \times SU(2)$ -invariant G_2 -structures on M of the form*

$$\varphi = p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a_1 e_1 \wedge e'_1 + a_2 e_2 \wedge e'_2 + a_3 e_3 \wedge e'_3).$$

- (i) If $K = \Delta\text{SU}(2)$ then there exists a 3-parameter family of torsion-free G_2 -structures φ defined in a neighbourhood of the singular orbit. The family is parametrised by $r_0 > 0$ and $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ such that

$$64r_0(\alpha_1 + \alpha_2 + \alpha_3) = 1, \quad p = -q = r_0^3, \quad a_i(t) = r_0^3 + \frac{1}{4}r_0t^2 + \alpha_i t^4 + O(t^6).$$

- (ii) If $K = \{1\} \times \text{SU}(2)$ then there exists a 3-parameter family of torsion-free G_2 -structures φ defined in a neighbourhood of the singular orbit. The family is parametrised by $r_0 > 0$ and $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ such that

$$\alpha_1\alpha_2\alpha_3 = 1, \quad p = -r_0^3, \quad q = 0, \quad a_i(t) = \frac{1}{4}r_0\alpha_i t^2 + O(t^4).$$

- (iii) If $K = K_{1,1}$ then there exists a 3-parameter family of torsion-free G_2 -structures φ defined in a neighbourhood of the singular orbit. The family is parametrised by $r_0, \alpha, \beta \in \mathbb{R}$ with $\beta > 0$ via $p = -q = -r_0^3$ and

$$a_1 = r_0^3\alpha + r_0^2\beta t + O(t^2), \quad a_2 = -r_0^3\alpha + r_0^2\beta t + O(t^2), \quad a_3 = r_0^3 + O(t^2).$$

- (iv) If $K = K_{m,n}$ for coprime integers with $mn > 1$ then there exists a 2-parameter family of torsion-free G_2 -structures φ defined in a neighbourhood of the singular orbit. The family is parametrised by $r_0 \in \mathbb{R}$ and $\beta > 0$ via $p = -m^2r_0^3, q = n^2r_0^3$ and

$$a_1 = a_2 = r_0^2\beta t + O(t^3), \quad a_3 = mnr_0^3 + O(t^2).$$

In the rest of the section we show how to apply Theorem 4.3 to prove the four cases of the proposition.

The case $K = \Delta\text{SU}(2)$. Fix $r_0 > 0$. According to Proposition 4.1(i) we must have $p = -q = r_0^3$ and

$$x_i = \dot{a}_j \dot{a}_k = \frac{r_0^2}{4}t^2 + 2r_0(\alpha_j + \alpha_k)t^4 + O(t^6), \quad y_i = a_i = r_0^3 + \frac{r_0}{4}t^2 + \alpha_i t^4 + O(t^6).$$

The constraint $H(x, y) = 0$ is already satisfied up to fourth order at $t = 0$. The vanishing of the fifth t -derivative of $H(x, y)$ at $t = 0$ imposes the further constraint

$$64r_0(\alpha_1 + \alpha_2 + \alpha_3) = 1.$$

Fix any triple $(\alpha_1, \alpha_2, \alpha_3)$ satisfying this constraint and write

$$x_i = \frac{r_0^2}{4}t^2 + t^4 X_i, \quad y_i = r_0^3 + \frac{r_0}{4}t^2 + t^4 Y_i,$$

where $X_i(0) = 2r_0(\alpha_j + \alpha_k)$ and $Y_i(0) = \alpha_i$. Writing $O(t^k)$ for any (convergent) series $\sum_{i \geq k} c_i t^i$ with coefficients c_i depending on X_1, \dots, Y_3, r_0 , we find that (X, Y) satisfies an ODE system

$$\begin{aligned} \dot{X}_i &= t^{-1}(-4X_i + r_0(-5Y_i + 3Y_j + 3Y_k) + \frac{5}{64}) + O(1), \\ \dot{Y}_i &= t^{-1}(\frac{1}{r_0}(-X_i + X_j + X_k) - 4Y_i) + O(1) \end{aligned}$$

with the same structure as the one considered in Theorem 4.3. The condition $64r_0(\alpha_1 + \alpha_2 + \alpha_3) = 1$ guarantees that the initial condition y_0 for (X, Y) satisfies $M_{-1}(y_0) = 0$.

Moreover, we have

$$d_{y_0}M_{-1} = \begin{pmatrix} -4 & 0 & 0 & -5r_0 & 3r_0 & 3r_0 \\ 0 & -4 & 0 & 3r_0 & -5r_0 & 3r_0 \\ 0 & 0 & -4 & 3r_0 & 3r_0 & -5r_0 \\ -\frac{1}{r_0} & \frac{1}{r_0} & \frac{1}{r_0} & -4 & 0 & 0 \\ \frac{1}{r_0} & -\frac{1}{r_0} & \frac{1}{r_0} & 0 & -4 & 0 \\ \frac{1}{r_0} & \frac{1}{r_0} & -\frac{1}{r_0} & 0 & 0 & -4 \end{pmatrix}$$

and therefore for all $h \geq 1$,

$$\det(h \text{ Id} - d_{y_0}M_{-1}) = h^2(h + 3)(h + 5)(h + 8)^2 > 0.$$

The case $K = \{1\} \times \text{SU}(2)$. Fix $r_0 > 0$. According to Proposition 4.1(ii) we must have $p = -r_0^3, q = 0$ and

$$x_i = \frac{r_0^2}{4} \alpha_j \alpha_k t^2 + O(t^4), \quad y_i = \frac{r_0}{4} \alpha_i t^2 + O(t^4)$$

for $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ satisfying $\alpha_1 \alpha_2 \alpha_3 = 1$.

The pair (X, Y) defined by $x_i = t^2 X_i$ and $y_i = t^2 Y_i$ satisfies an ODE system of the form of Theorem 4.3:

$$\dot{X}_i = t^{-1} \left(-2X_i + \sqrt{\frac{r_0^3 Y_j Y_k}{Y_i}} \right) + O(1), \quad \dot{Y}_i = t^{-1} \left(-2Y_i + \sqrt{\frac{X_j X_k}{X_i}} \right) + O(1).$$

The initial condition $y_0 = \left(\frac{r_0^2}{4} \alpha_j \alpha_k, \frac{r_0}{4} \alpha_i \right)$ for (X, Y) satisfies $M_{-1}(y_0) = 0$ precisely when $\alpha_1 \alpha_2 \alpha_3 = 1$. Moreover, for any such triple $(\alpha_1, \alpha_2, \alpha_3)$ we have

$$d_{y_0}M_{-1} = \begin{pmatrix} -2 & 0 & 0 & -\alpha_2^2 \alpha_3^2 r_0 & -\alpha_3 r_0 & -\alpha_2 r_0 \\ 0 & -2 & 0 & -\alpha_3 r_0 & -\alpha_3^2 \alpha_1^2 r_0 & -\alpha_1 r_0 \\ 0 & 0 & -2 & -\alpha_2 r_0 & -\alpha_1 r_0 & -\alpha_1^2 \alpha_2^2 r_0 \\ -\frac{\alpha_1^2}{r_0} & -\frac{\alpha_1 \alpha_2}{r_0} & -\frac{\alpha_3 \alpha_1}{r_0} & -2 & 0 & 0 \\ -\frac{\alpha_1 \alpha_2}{r_0} & -\frac{\alpha_2^2}{r_0} & -\frac{\alpha_2 \alpha_3}{r_0} & 0 & -2 & 0 \\ -\frac{\alpha_3 \alpha_1}{r_0} & -\frac{\alpha_2 \alpha_3}{r_0} & -\frac{\alpha_3^2}{r_0} & 0 & 0 & -2 \end{pmatrix}$$

and therefore for all $h \geq 1$,

$$\det(h \text{ Id} - d_{y_0}M_{-1}) = h^2(h + 4)^2(h + 1)(h + 3) > 0.$$

Remark 4.6. For future use we record higher-order expansions of a_1, a_2, a_3 in the special case where there is an additional $U(1)$ -symmetry:

$$a_1(t) = a_2(t) = \frac{1}{4} r_0 \alpha_1 t^2 + \frac{8 - 5\alpha_3^3}{576 \alpha_1 r_0} t^4 + O(t^6),$$

$$a_3(t) = \frac{1}{4} r_0 \alpha_3 t^2 - \frac{(4 - 7\alpha_3^3) \alpha_3}{576 \alpha_1^2 r_0} t^4 + O(t^6).$$

The case $K = K_{1,1}$. Fix $r_0 \in \mathbb{R}$. According to Proposition 4.1(iii) we must have $p = -r_0^3, q = r_0^3$ and

$$y_1 = r_0^3\alpha + r_0^2\beta t + O(t^2), \quad y_2 = -r_0^3\alpha + r_0^2\beta t + O(t^2), \quad y_3 = r_0^3 + O(t^2),$$

for some $\alpha \in \mathbb{R}$ and $\beta > 0$ (in particular we can assume that $y_1 + y_2 > 0$ for small $t > 0$; we will use this condition freely below to simplify square roots).

We now set

$$\begin{aligned} x_1 &= tX_1, & x_2 &= tX_2, & x_3 &= r_0^4\beta^2 + t^2X_3, \\ y_1 &= r_0^3\alpha + tY_1, & y_2 &= -r_0^3\alpha + tY_2, & y_3 &= r_0^3 + t^2Y_3. \end{aligned}$$

Then (X, Y) satisfies an ODE system of the form of Theorem 4.3:

$$\begin{aligned} \dot{X}_1 &= \frac{1}{t}(2\epsilon r_0^3\sqrt{1-\alpha^2} - X_1) + O(1), \\ \dot{X}_2 &= \frac{1}{t}(2\epsilon r_0^3\sqrt{1-\alpha^2} - X_2) + O(1), \\ \dot{X}_3 &= \frac{1}{t}\left(\frac{(Y_1 + Y_2)^2 - 2Y_3r_0^3(1-\alpha^2)}{(Y_1 + Y_2)\epsilon\sqrt{1-\alpha^2}} - 2X_3\right) + O(1), \\ \dot{Y}_1 &= \frac{1}{t}\left(\frac{r_0^2\beta X_2}{\sqrt{X_1X_2}} - Y_1\right) + O(1), \\ \dot{Y}_2 &= \frac{1}{t}\left(\frac{r_0^2\beta X_1}{\sqrt{X_1X_2}} - Y_2\right) + O(1), \\ \dot{Y}_3 &= \frac{1}{t}\left(\frac{X_1X_2}{r_0^2\beta\sqrt{X_1X_2}} - 2Y_3\right) + O(1), \end{aligned}$$

where ϵ is the sign of r_0 .

There is a unique solution

$$y_0 = \left(2\epsilon r_0^3\sqrt{1-\alpha^2}, 2\epsilon r_0^3\sqrt{1-\alpha^2}, \frac{\epsilon\beta r_0^2}{\sqrt{1-\alpha^2}} - \frac{r_0^2(1-\alpha^2)}{2\beta^2}, r_0^2\beta, r_0^2\beta, \frac{\epsilon r_0\sqrt{1-\alpha^2}}{\beta}\right)$$

to the equation $M_{-1}(y_0) = 0$. Moreover, $d_{y_0}M_{-1}$ is given by

$$\begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2\frac{2\epsilon\beta^3+(1-\alpha^2)\sqrt{1-\alpha^2}}{2\beta^3\sqrt{1-\alpha^2}} & \frac{2\epsilon\beta^3+(1-\alpha^2)\sqrt{1-\alpha^2}}{2\beta^3\sqrt{1-\alpha^2}} & -\frac{\epsilon r_0\sqrt{1-\alpha^2}}{\beta} & 0 \\ -\frac{\epsilon\beta}{4r_0\sqrt{1-\alpha^2}} & \frac{\epsilon\beta}{4r_0\sqrt{1-\alpha^2}} & 0 & -1 & 0 & 0 \\ \frac{\epsilon\beta}{4r_0\sqrt{1-\alpha^2}} & -\frac{\epsilon\beta}{4r_0\sqrt{1-\alpha^2}} & 0 & 0 & -1 & 0 \\ \frac{1}{2\beta r_0^2} & \frac{1}{2\beta r_0^2} & 0 & 0 & 0 & -2 \end{pmatrix}.$$

and

$$\det(h \text{Id} - d_{y_0}M_{-1}) = (h + 1)^4(h + 2)^2 > 0$$

for all $h \geq 1$.

The case $K = K_{m,n}$, $mn > 1$. This case is very similar to the previous one. Fix $r_0 \in \mathbb{R}$. According to Proposition 4.1(iii) we must have $p = -m^2r_0^3$, $q = n^2r_0^3$ and

$$y_1 = y_2 = r_0^2\beta t + O(t^2), \quad y_3 = mn r_0^3 + O(t^2),$$

for some $\beta > 0$. Using $y_1 = y_2$ and $x_1 = x_2$, we now set

$$x_1 = tX_1, \quad x_3 = r_0^4\beta^2 + t^2X_3, \quad y_1 = tY_1, \quad y_3 = mn r_0^3 + t^2Y_3.$$

Then (X_1, X_3, Y_1, Y_3) satisfies an ODE system of the form of Theorem 4.3:

$$\begin{aligned} \dot{X}_1 &= \frac{1}{t}(\sqrt{mn}(m+n)\epsilon r_0^3 - X_1) + O(1), \\ \dot{X}_3 &= \frac{1}{t}\left(\frac{(m+n)^2Y_1^2 - 2m^2n^2Y_3r_0^3}{\sqrt{mn}(m+n)\epsilon Y_1} - 2X_3\right) + O(1), \\ \dot{Y}_1 &= \frac{1}{t}(r_0^2\beta X_1 - Y_1) + O(1), \\ \dot{Y}_3 &= \frac{1}{t}\left(\frac{X_1}{r_0^2\beta} - 2Y_3\right) + O(1), \end{aligned}$$

where ϵ is the sign of r_0 .

There is a unique solution

$$y_0 = \left(\epsilon\sqrt{mn}(m+n)r_0^3, \frac{\epsilon(m+n)\beta r_0^2}{2\sqrt{mn}} - \frac{r_0^2m^2n^2}{2\beta^2}, r_0^2\beta, \frac{\epsilon\sqrt{mn}(m+n)r_0}{2\beta} \right)$$

to the equation $M_{-1}(y_0) = 0$. Moreover,

$$d_{y_0}M_{-1} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -2 & \frac{\epsilon(m+n)\beta^3 + m^2n^2\sqrt{mn}}{\sqrt{mn}\beta^3} & -\frac{2\epsilon r_0 mn\sqrt{mn}}{(m+n)\beta} \\ 0 & 0 & -1 & 0 \\ \frac{1}{2\beta r_0^2} & 0 & 0 & -2 \end{pmatrix}$$

and

$$\det(h \text{Id} - d_{y_0}M_{-1}) = (h + 1)^2(h + 2)^2 > 0$$

for all $h \geq 1$.

Remark 4.7. For future use note that the proof yields

$$y_1 = r_0^2\beta t + O(t^3), \quad y_2 = mn r_0^3 + \frac{\sqrt{mn}(m+n)|r_0|}{2\beta}t^2 + O(t^4)$$

as $t \rightarrow 0$.

Remark. A more geometric interpretation of the parameters in Proposition 4.5 is given by studying the leading order behaviour of the induced metric (3.18) as $t \rightarrow 0$. For example, in the case of Proposition 4.5(ii) with $\alpha_1 = \alpha_2$ (so that we have the enhanced $U(1)$ -symmetry) we have

$$g_t \approx r_0^2\alpha_1^2(e_1 \otimes e_1 + e_2 \otimes e_2) + r_0^2\alpha_3^2 e_3 \otimes e_3 + \frac{1}{4}t^2(e'_1 \otimes e'_1 + e'_2 \otimes e'_2 + e'_3 \otimes e'_3).$$

The metric on the singular orbit is a Berger metric on the 3-sphere. If we fix $r_0 = 1$ by scaling, the parameters $\alpha_1 = \alpha_2$ and α_3 determine, respectively, the size of the base S^2 and of the Hopf fibres. Similarly, in the case of Proposition 4.5 (iv) we calculate

$$g_t \approx \frac{r_0^2 \beta}{\sqrt{mn}} (m(e_1 \otimes e_1 + e_2 \otimes e_2) + (e'_1 \otimes e'_1 + e'_2 \otimes e'_2)) \\ + \frac{mnr_0^2}{\beta^2} (me_3 + ne'_3)^2 + t^2(m+n)^2(se_3 - re'_3)^2$$

as $t \rightarrow 0$. Fixing $r_0 = 1$ by scaling, we see that the metric on the singular orbit is a squashed metric on the principal circle bundle $S^2 \times S^3 \rightarrow S^2 \times S^2$: the metric on the base $S^2 \times S^2$ is a product of the round metrics with ratio m/n between the areas of the two factors; the parameter β determines the respective sizes of the base and the circle fibres.

Remark. As an aside, we note that Alekseevsky–Dotti–Ferraris have classified invariant Einstein metrics on $SU(2) \times SU(2)/K_{m,n}$ [4, Theorem 4.1]. When $mn > 1$ there exists a unique invariant Einstein metric, which coincides with the Einstein metric obtained by Wang–Ziller on any circle bundle over a product of Kähler–Einstein manifolds [54, Theorem 1.4]; when $m = n = 1$ there are two Einstein metrics, the product of round metrics and the Sasaki–Einstein metric. The restriction of g_t to the singular orbit is never Einstein unless $m = n = 1$ and $2\beta^6 = 1$.

5. Conically singular and asymptotically conical ends

In addition to the solutions of (3.13) just constructed in Proposition 4.5 and which are defined in the neighbourhood of various classes of singular orbits we will need two further classes of local solutions to (3.13): (i) solutions with an isolated conical singularity modelled on the cone over the homogeneous nearly Kähler structure on $S^3 \times S^3$ and (ii) solutions with an asymptotically conical end asymptotic to the same cone. Describing solutions with either type of end behaviour leads to a class of singular initial value problems not widely studied in the previous extensive work on cohomogeneity one Einstein metrics. The closest work we are aware of is Dancer–Wang’s work [30] on Painlevé-type expansions of singular solutions to the cohomogeneity one Ricci-flat equations. In many cases their expansions construct continuous families of either AC or ALC Ricci-flat ends, subsets of which often have special or exceptional holonomy. However, because they consider only “rational resonances”, their approach often does not capture the full space of ends, e.g. in [30, Theorem 5.6] the only G_2 -holonomy AC ends obtained arise from the original Bryant–Salamon AC metric.

In order to construct solutions with prescribed singular/asymptotic behaviour we will use the following extension of Theorem 4.3. The theorem is proved in Chapter I of Volume 3 of Picard’s treatise [50] using the method of majorants; cf. in particular [50, Chapter I, §13].

Theorem 5.1. *Consider the singular initial value problem*

$$t\dot{y} = \Phi(y, t), \quad y(0) = y_0, \tag{5.2}$$

where y takes values in \mathbb{R}^k and $\Phi: \mathbb{R}^k \times \mathbb{R} \rightarrow \mathbb{R}^k$ is a real analytic function in a neighbourhood of $(y_0, 0)$ with $\Phi(y_0, 0) = 0$. Assume also that $\partial_t \Phi(y_0, 0)$ lies in the image of $\text{Id} - d_{y_0} \Phi(\cdot, 0)$ and fix a preimage $y_1 \in \mathbb{R}^k$. After possibly a change of basis, assume that $d_{y_0} \Phi(\cdot, 0)$ contains a diagonal block $\text{diag}(\lambda_1, \dots, \lambda_m)$ in the upper-left corner. Furthermore assume that the eigenvalues $\lambda_1, \dots, \lambda_m$ satisfy:

- (i) $\lambda_1, \dots, \lambda_m > 0$;
- (ii) for every $\mathbf{h} = (h_0, \dots, h_m) \in \mathbb{Z}_{\geq 0}^{m+1}$ with $|\mathbf{h}| = h_0 + \dots + h_m \geq 2$ the matrix

$$(\mathbf{h} \cdot \boldsymbol{\lambda}) \text{Id} - d_{y_0} \Phi(\cdot, 0)$$

is invertible. Here $\boldsymbol{\lambda} = (1, \lambda_1, \dots, \lambda_m)$ and $\mathbf{h} \cdot \boldsymbol{\lambda} = \sum_{i=0}^m h_i \lambda_i$.

Then for every $(u_1, \dots, u_m) \in \mathbb{R}^m$ there exists a unique solution $y(t)$ of (5.2) given as a convergent generalised power series

$$y(t) = y_0 + y_1 t + (u_1 t^{\lambda_1}, \dots, u_m t^{\lambda_m}, 0, \dots, 0) + \sum_{|\mathbf{h}| \geq 2} y_{\mathbf{h}} t^{\mathbf{h} \cdot \boldsymbol{\lambda}}.$$

Furthermore, the solutions depend real analytically on u_1, \dots, u_m .

Remark. It is clear that if Φ does not depend on t then the solution y has a generalised power series expansion in powers of $t^{\lambda_1}, \dots, t^{\lambda_m}$ only and we should take $h_0 = 0$ in condition (ii).

Remark. The existence theorem can be extended to the case of smooth (rather than real analytic) Φ by truncation of a formal generalised power series solution to sufficiently high order and a contraction mapping argument.

We now use this existence result to construct 1-parameter families of conically singular and asymptotically conical ends.

Proposition 5.3. *Let C be the G_2 -holonomy cone over the homogeneous nearly Kähler structure on $S^3 \times S^3$ and set $v_0 = \frac{\sqrt{145}-7}{2} \approx 2.5$ and $v_\infty = \frac{\sqrt{145}+7}{2} \approx 9.5$.*

- (i) *For every $c \in \mathbb{R}$ there exists a unique $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structure*

$$\varphi = d(a(e_1 \wedge e'_1 + e_2 \wedge e'_2) + b e_3 \wedge e'_3)$$

defined on $(0, \epsilon) \times S^3 \times S^3$ for some $\epsilon > 0$, where the functions $t^{-3}a$ and $t^{-3}b$ admit convergent generalised power series expansions in powers of t^{v_0} satisfying

$$\frac{54}{\sqrt{3}} t^{-3} a(t) = 1 + \frac{1}{2} c t^{v_0} + O(t^{2v_0}), \quad \frac{54}{\sqrt{3}} t^{-3} b(t) = 1 - c t^{v_0} + O(t^{2v_0}).$$

In particular the associated metric g_φ has a conical singularity as $t \rightarrow 0$ asymptotic to the cone C with rate v_0 .

(ii) Fix $p, q \in \mathbb{R}$. For every $c \in \mathbb{R}$ there exists a unique $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structure

$$\varphi = p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a(e_1 \wedge e'_1 + e_2 \wedge e'_2) + b e_3 \wedge e'_3)$$

on $(T, \infty) \times S^3 \times S^3$ for some $T > 0$, where the functions $t^{-3}a$ and $t^{-3}b$ admit convergent generalised power series expansions in powers of t^{-3} and $t^{-\nu_\infty}$ satisfying

$$\frac{54}{\sqrt{3}}t^{-3}a = 1 + O(t^{-3}), \quad \frac{54}{\sqrt{3}}t^{-3}b = 1 + O(t^{-3}), \quad \frac{54}{\sqrt{3}}t^{-3}(b-a) = ct^{-\nu_\infty} + O(t^{-12}).$$

In fact if $p = 0 = q$ then $t^{-3}a$ and $t^{-3}b$ admit convergent generalised power series expansions in powers of $t^{-\nu_\infty}$ only. In particular the associated metric g_φ has a complete asymptotically conical end as $t \rightarrow \infty$ asymptotic to the cone C with rate $-\nu_\infty$ if $p = 0 = q$ and rate -3 otherwise.

Proof. First of all, note that, because of its invariance under $SU(2)^3$ and scaling, the solution to the fundamental ODE system (3.13) corresponding to the G_2 -cone over the homogeneous nearly Kähler structure on $S^3 \times S^3$ must satisfy $a = b = Ct^3$. It is then immediate to check that $C = \frac{\sqrt{3}}{54}$.

The proof of the existence of CS solutions is a straightforward application of Theorem 5.1. We write $a = \frac{\sqrt{3}}{54}t^3(1 + Y_1)$ and $b = \frac{\sqrt{3}}{54}t^3(1 + Y_2)$. Define functions X_1 and X_2 by $\dot{a}\dot{b} = \frac{1}{108}t^4(1 + X_1)$ and $\dot{a}^2 = \frac{1}{108}t^4(1 + X_2)$. Then the $U(1)$ -enhanced ODE system (3.17) for $(\dot{a}\dot{b}, \dot{a}^2, a, b)$ becomes an ODE system for (X_1, X_2, Y_1, Y_2) of the form (5.2):

$$\begin{aligned} t\dot{X}_1 &= -4X_1 + \frac{4\sqrt{3}(1 + Y_1)(1 + Y_2)^2}{\sqrt{4(1 + Y_1)^2(1 + Y_2)^2 - (1 + Y_2)^4}} - 4, \\ t\dot{X}_2 &= -4X_2 + \frac{4\sqrt{3}(1 + Y_2)(2(1 + Y_1)^2 - (1 + Y_2)^2)}{\sqrt{4(1 + Y_1)^2(1 + Y_2)^2 - (1 + Y_2)^4}} - 4, \\ t\dot{Y}_1 &= -3Y_1 + \frac{3(1 + X_1)(1 + X_2)}{\sqrt{(1 + X_1)^2(1 + X_2)}} - 3, \\ t\dot{Y}_2 &= -3Y_2 + \frac{3(1 + X_1)^2}{\sqrt{(1 + X_1)^2(1 + X_2)}} - 3. \end{aligned}$$

The linearisation of Φ at $y_0 = (0, 0, 0, 0)$ is

$$L = \begin{pmatrix} -4 & 0 & -\frac{4}{3} & \frac{16}{3} \\ 0 & -4 & \frac{32}{3} & -\frac{20}{3} \\ 0 & \frac{3}{2} & -3 & 0 \\ 3 & -\frac{3}{2} & 0 & -3 \end{pmatrix}. \tag{5.4}$$

L has four distinct eigenvalues $-1, -6, -\nu_\infty = -\nu_0 - 7, \nu_0$. The corresponding eigenvectors are, respectively,

$$(4, 4, 3, 3), \quad (2, 2, -1, -1), \quad (3 + \nu_0, -6 - 2\nu_0, -3, 6), \quad (4 + \nu_0, -8 - 2\nu_0, 3, -6).$$

The proof of part (i) now follows immediately from Theorem 5.1.

Constructing AC ends is more involved: Theorem 5.1 cannot be applied immediately because the non-resonance condition (ii) fails. As above we work with the system (3.17) for the 4-tuple $(x_1 = x_2 = \dot{a}\dot{b}, x_3 = \dot{a}^2, y_1 = y_2 = a, y_3 = b)$ but it is now convenient to change variable $s = 1/t$. Define functions (X_1, X_2, Y_1, Y_2) by $s^3y_1 = \frac{\sqrt{3}}{54}(1+Y_1), s^3y_3 = \frac{\sqrt{3}}{54}(1+Y_2), s^4x_1 = \frac{1}{108}(1+X_1)$ and $s^4x_3 = \frac{1}{108}(1+X_2)$. The 4-tuple (X_1, X_2, Y_1, Y_2) satisfies a system of the form $s\dot{y} = \Phi(y, s^3)$ where Φ does not depend on s^3 if $p = 0 = q$. The linearisation of $\Phi(\cdot, 0)$ at 0 is $-L$, with L as in (5.4).

Now, the presence of 1 in the spectrum of $-L$ is explained by the fact that the original system (3.17) is t -invariant. By a translation $t \mapsto t + t_0$ we can therefore always reduce to the case when the eigenvalue 1 is never excited. The eigenvalue 6 should not be excited either. Indeed, we are interested in solutions of (3.17) satisfying the conservation law $H(x, y) = 0$. Rewriting $H = H(s^3, X_1, X_2, Y_1, Y_2)$, a calculation shows that H vanishes at leading order if and only if $3(2X_1 + X_2) - 4(2Y_1 + Y_2) = 0$.

The discussion above motivates us to look for a 1-parameter family of AC solutions to the ODE system (3.17) given by convergent generalised power series in powers of s^3, s^{ν_∞} and parametrised by the coefficient of s^{ν_∞} . A further change of variable $s \mapsto s^3$ justifies the fact that powers of s^3 and not s should be considered.

Given that $\partial_s \Phi(0, 0) = 0$, the conditions to be satisfied in order to apply Theorem 5.1 are

$$3h_0 + \nu_\infty h_1 \neq 1, 6, \nu_\infty, -\nu_0$$

for every $h_0, h_1 \in \mathbb{Z}_{\geq 0}$ with $h_0 + h_1 \geq 2$. The non-resonance condition fails only when $(h_0, h_1) = (2, 0)$. There is therefore a potential obstruction to solve the system: the coefficient $y_{2,0}$ of s^6 must satisfy an equation

$$(L + 6)y_{2,0} = Q_{2,0}(y_0, y_{1,0}), \tag{5.5}$$

where $Q_{2,0}$ is a real analytic function of the initial condition $y_0 = 0$ for (X, Y) and the coefficient $y_{1,0}$ of s^3 , which is uniquely determined by the equation. We know that $L + 6$ has a 1-dimensional kernel and cokernel. Only if $Q_{2,0}(y_0, y_{1,0})$ lies in the hyperplane $\text{im}(L + 6)$ can we solve (5.5). Assuming this is the case, we fix the choice of a solution $y_{2,0}$ to (5.5) by imposing the vanishing of the coefficient of order s^6 of the Hamiltonian H . Since the linearisation of H does not annihilate the eigenvector of $-L$ of eigenvalue 6, this requirement fixes a unique choice for $y_{2,0}$. Once $y_{2,0}$ is uniquely determined, the iteration procedure to find a formal generalised power series solution to (3.17) can be continued without further obstructions and the majorisation argument in the proof of Theorem 5.1 still guarantees that the generalised power series converges.

The key observation now is that the potential obstruction to solve (5.5) does in fact vanish. Instead of showing this by computation, we observe that for every $p, q \in \mathbb{R}$ there exists an AC solution of the ODE system (3.17) with an enhanced $SU(2)$ -symmetry, i.e. with $a = b$. Using the conserved quantity $H = 0$ we can describe such a solution by the curve $(x, y) \in \mathbb{R}^2$ defined by the equation $4x^3 = 3y^4 - 4(p - q)y^3 - 6pqy^2 - p^2q^2$. Recalling that $y = a = b$ and $x = \dot{a}^2$ we can rewrite this equation as the ODE

$$4\dot{a}^6 = 3a^4 - 4(p - q)a^3 - 6pqa^2 - p^2q^2$$

for the function a . Taking the sixth root of both sides of the equation ($\dot{a} > 0$ with our conventions), changing variable $s = 1/t$ and writing $a = \frac{\sqrt{3}}{54}s^{-3}(1 + w)$, the equation can be written in the form $s\dot{w} = w + O(w^2 + s^3)$. Theorem 5.1 then guarantees that the solution admits a convergent power series expansion in $s^3 = t^{-3}$, unique up to a time translation $t \mapsto t - t_0$. We conclude that all the coefficients $y_{h,0}$ in a formal generalised power series expansion of y (in particular the coefficient $y_{2,0}$) are uniquely determined by the equation up to a time translation. \square

Remark. In order to relate the proposition to the general deformation theory of CS and AC G_2 -manifolds developed by Karigiannis–Lotay [43], one can check that $\sigma = e_1 \wedge e'_1 + e_2 \wedge e'_2 - 2e_3 \wedge e'_3$ is a coclosed primitive (1, 1)-form with

$$\Delta\sigma = (v_0 + 3)(v_0 + 4)\sigma = (-v_\infty + 3)(-v_\infty + 4)\sigma = 36\sigma$$

on $S^3 \times S^3$ endowed with its homogeneous nearly Kähler structure. It follows that $d(t^{v_0+3}\sigma)$ and $d(t^{-v_\infty+3}\sigma)$ are closed and coclosed 3-forms of type 27 on the G_2 -cone C and therefore infinitesimal deformations of C as a G_2 -manifold. On the other hand, differentiating our solutions with respect to the parameter c in the construction yields solutions $d(t^{v_0+3}\sigma)$ and $d(t^{-v_\infty+3}\sigma)$ of the linearisation of (3.17) at the conical solution $a = b = \frac{\sqrt{3}}{54}t^3$.

6. Existence of ALC metrics

In this section we obtain our first main (global) existence results, Theorems A and B in the Introduction: we prove the existence of two 1-parameter families of complete ALC G_2 -metrics and the existence of an ALC G_2 -space with an isolated conical singularity. One of the two families of ALC G_2 -metrics we obtain, named \mathbb{B}_7 in the physics literature, is already known to exist thanks to work of Bogoyavlenskaya [13] (one member of this family is explicit and was found earlier by Brandhuber–Gomis–Gubser–Gukov [17]); the other family, named \mathbb{D}_7 in the physics literature, is currently only known numerically or in the collapsed limit of [35]. The existence of the CS ALC space seems not to have been anticipated in the physics literature: as explained in the Introduction it can be used, together with the Bryant–Salamon AC metric on $S^3 \times \mathbb{R}^4$, to explain the existence of both the \mathbb{B}_7 and \mathbb{D}_7 families of ALC manifolds by a gluing construction.

6.1. A criterion for forward completeness

In the following proposition we relate the forward completeness of a cohomogeneity one torsion-free G_2 -structure to the sign of the mean curvature l of the principal orbits. One direction is very closely related to a result of Böhm [15, Proposition 3.2], while the other implication, which plays a crucial role in our analysis, was suggested to us by Wilking.

Proposition 6.1. *Let (M, φ) be a cohomogeneity one $SU(2) \times SU(2)$ -invariant G_2 -manifold (not necessarily complete). Assume that $l(t_0) > 0$ for some initial time t_0 corresponding to a principal orbit. Then the solution blows up in finite time if and only if there*

exists $t_* > t_0$ such that $l(t_*) = 0$, i.e. if and only if there exists a principal orbit that is a minimal hypersurface.

Proof. Since the 7-dimensional metric $dt^2 + g_t$ induced by φ is Ricci-flat, the mean curvature l of the principal orbits satisfies $0 = l' + |L|^2 = l' + \frac{1}{6}l^2 + |\mathring{L}|^2$, where \mathring{L} denotes the traceless part of the second fundamental form L of the principal orbits; cf. for example [32, Proposition 2.1].

Note that M cannot contain a totally geodesic principal orbit \mathcal{O} . To see this, let ψ denote the parallel spinor on M and recall that using Clifford multiplication the covariant derivative of $\psi|_{\mathcal{O}}$ can be identified with the second fundamental form of \mathcal{O} . If \mathcal{O} were totally geodesic, it would therefore carry a parallel spinor and hence a homogeneous Calabi–Yau (and therefore Ricci-flat) metric. This is impossible since homogeneous Ricci-flat metrics must be flat, while the principal orbits of M , which are finite quotients of $S^3 \times S^3$, do not carry flat metrics. In particular, $l(t_*) = 0$ implies $l(t) < 0$ for $t > t_*$. Then comparison with the solution of $u' + \frac{1}{6}u^2 = 0$ shows that l must blow up in finite time.

Conversely, assume that the solution exists only on a finite interval $[t_0, T)$. By Lemma 3.19 we can regard Hitchin’s flow as a *first-order* ODE system for the metric g . Since it satisfies a first-order equation, the metric must degenerate as we approach the maximal existence time T (otherwise we could use local existence to extend the solution beyond $t = T$). Hence as $t \rightarrow T$ the norm of a Jacobi field J , i.e. a vector field satisfying $J' = LJ$, converges either to zero or to infinity. In either case, we deduce that $\int_{t_0}^T |L| \geq \lim_{t \rightarrow T} \log |J|(t) - \log |J|(t_0) = \infty$. Since $T < \infty$, by Hölder’s inequality we also have $\int_{t_0}^T |L|^2 = \infty$. Integration of $l' + |L|^2 = 0$ shows that $l(t) \rightarrow -\infty$ as $t \rightarrow T$. Since $l(t_0) > 0$ we deduce there must exist $t_* \in (t_0, T)$ such that $l(t_*) = 0$. \square

Remark. If M closes smoothly on a singular orbit or has an isolated conical singularity at $t = 0$ then $\lim_{t \rightarrow 0} l(t) = +\infty$ so the assumption about the positivity of the mean curvature at an initial time is certainly satisfied in these cases.

Remark. Finite time blow-up when there exists a minimal principal orbit can be deduced as in the proposition from any condition that would exclude the existence of a totally geodesic principal orbit. For example, in [15, Proposition 3.2] Böhm shows that in a complete cohomogeneity one Ricci-flat manifold that does not contain a line and is not flat principal, orbits cannot be minimal.

6.2. The $U(1)$ -enhanced symmetric system

Let us now specialise to the setting where we assume an additional $U(1)$ -symmetry. In the rest of the paper we will give a detailed qualitative analysis of the single second-order equation (3.16) arising from the Lagrangian formulation of the problem. Hence in the rest of the paper $\dot{}$ denotes differentiation with respect to an arbitrary parameter s . We reserve the freedom to change parametrisation (compatible with a fixed orientation) and will specify the choice of a parametrisation when additional properties are needed. We

consider pairs of functions (a, b) satisfying

$$2F(\dot{a}\ddot{b} - \dot{b}\ddot{a}) = -\dot{a}\dot{b}(2\dot{b}F_b - \dot{a}F_a),$$

where $F = 4a^2(b - p)(b + q) - (b^2 + pq)^2$ and F_a, F_b denote its partial derivatives, and the constraints

$$\dot{a}, \dot{b} > 0, \quad F(a, b) > 0.$$

Note that the latter condition forces a to have a definite sign and $b - p, b + q$ to have the same definite sign.

The following formulas will play an important role in our analysis:

$$2F_b - F_a = 8(a - b)(a(2b + q - p) + b^2 + pq), \tag{6.2a}$$

$$2bF_b - aF_a = 8(a - b)(a + b)(b^2 + pq). \tag{6.2b}$$

Furthermore the formula (3.20) for the mean curvature l of the principal orbits reads

$$l = \frac{\dot{a}F_a + \dot{b}F_b}{2F}. \tag{6.3}$$

We also observe that in all cases we are interested in, p and q satisfy $pq \leq 0$. By Theorem 7.3 below this is no accident: every $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structure closing smoothly on a singular orbit must satisfy this condition.

Lemma 6.4. *Assume that (a, b) is a solution to (3.16) with $pq \leq 0$. Then the conditions*

$$\dot{a} > \dot{b}, \quad a > b > \max(p, -q, \sqrt{-pq}) \geq 0 \tag{6.5}$$

are preserved as long as the solution exists with $\dot{a}, \dot{b}, F > 0$.

Proof. Since $\dot{a}, \dot{b} > 0$ the inequalities $a > 0$ and $b > \max(p, -q, \sqrt{-pq}) \geq 0$ are certainly preserved. If $\dot{a} > \dot{b}$ then also $a > b$ is preserved. We therefore must show that the condition $\dot{a} - \dot{b} > 0$ is preserved as long as the solution satisfies the open constraints to define a G_2 -structure. Now, at a point where $\dot{a} = \dot{b}$, (3.16) yields

$$2F(\ddot{a} - \ddot{b}) = \dot{a}^2(2F_b - F_a).$$

Hence by (6.2a), $\ddot{a} - \ddot{b} > 0$ as long as $a > b > \max(p, -q, \sqrt{-pq}) \geq 0$. □

Proposition 6.6. *If the conditions (6.5) are satisfied then the solution (a, b) is forward complete. In particular, $a \rightarrow \infty$ as we approach the complete end.*

Proof. By (6.3) the mean curvature of the principal orbit has the same sign as $2(\dot{a}F_a + \dot{b}F_b) = \dot{b}(2F_b - F_a) + (2\dot{a} + \dot{b})F_a > 0$ if the conditions (6.5) are satisfied (note that $F_a = 8a(b - p)(b + q)$ is strictly positive). By Proposition 6.1 the solution cannot blow up in finite time.

In order to prove that $a \rightarrow \infty$, parametrise with respect to the arc-length parameter t along a geodesic meeting all principal orbits orthogonally. Note that since $\dot{a} > \dot{b}$ we have $2\dot{a}^3 > 2\dot{a}^2\dot{b}$. Up to a positive multiplicative constant, the right-hand side is the orbital volume function $\text{Vol}(t)$. Since the mean curvature l is positive, $\text{Vol}(t) \geq \text{Vol}(t_0) > 0$ for all $t \geq t_0$. Hence \dot{a} is bounded below by a positive constant. Since t is unbounded along the complete end, so is a . □

Remark. In the course of proving ALC asymptotics in Proposition 6.11 we will show that the assumptions (6.5) also force $b \rightarrow \infty$ along the complete end.

6.3. ALC asymptotics

We are now going to show that under an additional assumption the complete ends of Proposition 6.6 are ALC. First note that the conical Calabi–Yau structure (ω_C, Ω_C) on the conifold $C = C(\Sigma)$, $\Sigma = \text{SU}(2) \times \text{SU}(2)/K_{1,-1}$, is given by

$$\text{Re } \Omega_C = d\left(\frac{1}{18}t^3(e_1 \wedge e'_1 + e_2 \wedge e'_2)\right), \quad \omega_C = -d\left(\frac{1}{6}t^2(e_3 - e'_3)\right).$$

Moreover, $\theta = \frac{1}{2}(e_3 + e'_3)$ is the (unique up to gauge transformations) Hermitian Yang–Mills connection on the circle bundle $\mathbb{R}_+ \times \text{SU}(2) \times \text{SU}(2) \rightarrow C$, i.e. $d\theta \wedge \omega_C^2 = 0 = d\theta \wedge \Omega_C$. Fix $\ell > 0$ and consider the closed G_2 -structure φ_∞ on the total space of this circle bundle,

$$\varphi_\infty = d\left(\frac{1}{18}t^3(e_1 \wedge e'_1 + e_2 \wedge e'_2)\right) + \frac{1}{6}\ell t^2 e_3 \wedge e'_3. \tag{6.7}$$

Since θ is Hermitian Yang–Mills, $d *_{\varphi_\infty} \varphi_\infty = O(t^{-2})$. Moreover, up to terms that decay as t^{-1} the metric induced by φ_∞ is

$$g_{\varphi_\infty} = dt^2 + t^2 g_{\text{se}} + \ell^2 \theta^2 + O(t^{-1}),$$

where g_{se} is the (pull-back to the total space of the circle bundle of the) Sasaki–Einstein metric on $\Sigma = \text{SU}(2) \times \text{SU}(2)/\Delta\text{U}(1)$. (In particular here t is the arc-length parameter along a geodesic meeting all principal orbits orthogonally.) We therefore regard φ_∞ as the asymptotic model for an $\text{SU}(2) \times \text{SU}(2)$ -invariant torsion-free G_2 -structure inducing an ALC metric.

Lemma 6.8. *Let*

$$\varphi = p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a(e_1 \wedge e'_1 + e_2 \wedge e'_2) + b e_3 \wedge e'_3)$$

be an $\text{SU}(2) \times \text{SU}(2) \times \text{U}(1)$ -invariant torsion-free G_2 -structure defined on $(t_0, \infty) \times \text{SU}(2) \times \text{SU}(2)$. Assume that a, b are positive increasing functions with $\lim_{t \rightarrow \infty} a = \infty$, $\lim_{t \rightarrow \infty} \frac{a^2}{b^3} = \frac{2}{3\ell^3}$ for some $\ell > 0$ and $\lim_{t \rightarrow \infty} \frac{a}{b} \frac{db}{da} = \frac{2}{3}$. Then the functions $\tilde{a}(t) = 18t^{-3}a(t) - 1$ and $\tilde{b}(t) = 6\ell^{-1}t^{-2}b(t) - 1$ satisfy $\tilde{a}^{(k)}(t) = O(t^{-k-1})$ and $\tilde{b}^{(k)}(t) = O(t^{-k-1})$ as $t \rightarrow \infty$ for all $k \geq 0$. In particular, φ converges to the closed G_2 -structure φ_∞ of (6.7) in C^∞ as $t \rightarrow \infty$.

Proof. Since a is a positive increasing function of t we can introduce a parameter s such that $a = \frac{1}{18}s^3$. Then by our assumptions on a and b , $b \approx \frac{1}{6}\ell s^2$ in C^0 . Hence, $\frac{2}{\ell s} \frac{db}{ds} \approx \frac{a}{b} \frac{db}{da} \approx \frac{2}{3}$ and therefore $b - \frac{1}{6}\ell s^2$ converges to zero in C^1 . Now, the arc-length parameter t is related to s by

$$2\left(\frac{ds}{dt}\right)^3 \left(\frac{da}{ds}\right)^2 \frac{db}{ds} = \sqrt{F(a, b)}.$$

As $s \rightarrow \infty$ we therefore have $\frac{ds}{dt} \approx 1$ and by integration $s \approx t$ in C^1 . Thus $a(t) - \frac{1}{18}t^3$ and $b(t) - \frac{1}{6}\ell t^2$ converge to zero in C^1 .

Now consider the system (3.17) for the four functions $x_1 = \dot{a}b$, $x_2 = \dot{a}^2$, $y_1 = a$, $y_2 = b$. Write $x_1 = \frac{\ell}{18}t^3(1 + X_1)$, $x_2 = \frac{1}{36}t^4(1 + X_2)$, $y_1 = \frac{1}{18}t^3(1 + Y_1)$ and $y_2 = \frac{\ell}{6}t^2(1 + Y_2)$. After changing variable $e^\tau = t$, one can check that (X_1, X_2, Y_1, Y_2) is a solution to an initial value problem of the form

$$\begin{aligned} \dot{X}_1 &= -3(1 + X_1) + 3 \frac{(1 + Y_1)(1 + Y_2)^2 + O(e^{-2\tau})}{\sqrt{(1 + Y_1)^2(1 + Y_2)^2 + O(e^{-2\tau})}}, \\ \dot{X}_2 &= -4(1 + X_2) + 4 \frac{(1 + Y_1)^2(1 + Y_2) + O(e^{-2\tau})}{\sqrt{(1 + Y_1)^2(1 + Y_2)^2 + O(e^{-2\tau})}}, \\ \dot{Y}_1 &= -3(1 + Y_1) + 3\sqrt{1 + X_2}, \\ \dot{Y}_2 &= -2(1 + Y_2) + 2 \frac{1 + X_1}{\sqrt{1 + X_2}}, \end{aligned} \tag{6.9}$$

where $O(e^{-2\tau})$ indicates a real analytic function of (X_1, X_2, Y_1, Y_2) with coefficients depending real analytically on $e^{-\tau}$ and vanishing at $e^{-\tau} = 0$ at least with order 2.

Define $Z = (X_1, X_2, Y_1, Y_2, e^{-\tau})$ so that (6.9) can be rewritten as an autonomous system $\frac{dZ}{ds} = \Phi(Z)$ for a real analytic map Φ . The linearisation $d_0\Phi$ has a 1-dimensional kernel and four negative eigenvalues -1 (with multiplicity 2), -5 and -6 . The presence of a 1-dimensional kernel is explained by the freedom to change ℓ . In fact $\{(c, 0, 0, c, 0) \mid c \in \mathbb{R}\}$ is the centre manifold of the system $\frac{dZ}{ds} = \Phi(Z)$. Hence standard centre manifold theory [21, §2.4, Theorem 2(b)] implies that any solution Z that stays in a neighbourhood of 0 must satisfy $Z = (c, 0, 0, c, 0) + O(e^{-\tau})$ as $\tau \rightarrow \infty$ for some $c \in \mathbb{R}$. In particular, if (x_1, x_2, y_1, y_2) is a solution to (3.17) asymptotic to $(\frac{\ell}{18}t^3, \frac{1}{36}t^4, \frac{1}{18}t^3, \frac{\ell}{6}t^2)$ for some $\ell > 0$ then

$$(t^{-3}x_1, t^{-4}x_2, t^{-3}y_1, t^{-2}y_2) = \left(\frac{\ell}{18}, \frac{1}{36}, \frac{1}{18}, \frac{\ell}{6}\right) + O(t^{-1}).$$

The statement about the decay of derivatives of $t^{-3}a$ and $t^{-2}b$ then follows from a bootstrap argument. □

Remark. In terms of a new independent variable $s = \frac{1}{\log \tau}$, the system (6.9) takes the form $s\dot{z} = \Phi(z, s^2)$ of Theorem 5.1. The linearisation of $\Phi(\cdot, 0)$ at the origin has four distinct eigenvalues 0, 1, 5, 6. We have already observed that the 1-dimensional kernel is due to the freedom of choosing $\ell > 0$. The eigenvector with eigenvalue 1 is due to the t -invariance of the original system. The eigenvectors with eigenvalue 5, 6 are, respectively, $(-3, 0, 0, 2)$ and $(-1, 2, -1, 1)$. Moreover, the Hamiltonian constraint $H(x, y) = 0$ is satisfied at leading order as $(s, z) \rightarrow 0$ if and only if $2X_1 + X_2 = 2(Y_1 + Y_2)$. The eigenvector with eigenvalue 5 does not satisfy this constraint. One could then try to use Theorem 5.1 to show that, up to translations in t and the scaling freedom to fix the asymptotic length ℓ of the circle fibre, there exists a 1-parameter family of $SU(2) \times SU(2) \times U(1)$ -invariant ALC G_2 -holonomy ends. However, since the non-resonance condition (ii) in

Theorem 5.1 is not satisfied, this does not follow immediately from that theorem. We do not pursue the matter further here since it is not necessary in our analysis.

Thanks to Lemma 6.8, in order to prove that a complete end is ALC it is enough to control the quantities $\frac{a^2}{b^3}$ and $\frac{a}{b} \frac{db}{da}$.

Let us fix the notation that will be used throughout the rest of the section. Let $\lambda = \dot{a}/\dot{b}$. Then (3.16) can be rewritten as

$$2F\dot{\lambda} = \lambda\dot{b}(2F_b - \lambda F_a).$$

We fix a parameter s that satisfies $s \rightarrow \infty$ along the complete end (for example we can take $s = a$ or $s = t$, the arc-length parameter along a geodesic meeting all principal orbits orthogonally).

For $\alpha \in \mathbb{R}$ we introduce the following pair of (positive) ratios:

$$P_\alpha = \frac{b^{1+\alpha}}{a}, \quad Q_\alpha = \frac{b^\alpha}{\lambda}.$$

Using (3.16) as rewritten above we find that these two quantities satisfy

$$a^2 \dot{P}_\alpha = b^\alpha \dot{b} R_\alpha, \quad 2F \dot{Q}_\alpha = b^{-1} \dot{b} Q_\alpha (S_\alpha - F_a R_\alpha), \tag{6.10a}$$

where

$$R_\alpha = (1 + \alpha)a - b\lambda, \quad S_\alpha = \alpha(2F + aF_a) - (2bF_b - aF_a).$$

Moreover,

$$2F \dot{R}_\alpha = \lambda \dot{b} (S_\alpha - F_a R_\alpha). \tag{6.10b}$$

Proposition 6.11. *Assume that either $q \geq p$ or $q = -p \leq 0$. If (a, b) is a solution to (3.16) satisfying (6.5) and additionally*

$$a\dot{b} - \dot{a}b < 0 \tag{6.12}$$

holds at some initial time, then along the complete end the ratio $\frac{a^2}{b^3}$ converges to a constant and $\frac{a}{b} \frac{db}{da}$ converges to $\frac{2}{3}$.

Proof. The proof is based on the study of the behaviour of the positive ratios P_α and Q_α introduced above for $\alpha \geq 0$.

We begin by analysing (6.10b) for α sufficiently large. Since $a > 0$, for all α sufficiently large R_α is positive at the initial time s_0 . We claim that, assuming α even larger if necessary, R_α remains positive for all $s \geq s_0$. Indeed, using $2\alpha F > 0$ we find

$$\begin{aligned} S_\alpha &> (1 + \alpha)aF_a - 2bF_b \\ &= 8a^2((1 + \alpha)(b - p)(b + q) - (b - p) - (b + q)) + 8b^2(b^2 + pq) \\ &> 8a^2((1 + \alpha)(b - p)(b + q) - (b - p) - (b + q)). \end{aligned}$$

If $\alpha > \frac{1}{2b-p+q}$ then $(1 + \alpha)(b - p)(b + q) - (b - p) - (b + q)$ is an increasing function of b . Since $\dot{b} > 0$ we conclude that $S_\alpha > 0$ provided we choose α so that

$$(1 + \alpha)(b - p)(b + q) - (b - p) - (b + q) > 0, \quad \alpha > \frac{1}{2b - p + q}$$

are satisfied at the initial time s_0 . Then (6.10b) shows that for large enough α , $\dot{R}_\alpha > 0$ whenever $R_\alpha = 0$. We conclude that $R_\alpha(s) > 0$ for all $s \geq s_0$ and all sufficiently large α as claimed. Hence by (6.10a), for all sufficiently large α , P_α is an increasing function of s , and in particular it is bounded away from 0. In particular b , as well as a , is unbounded along the complete end. For otherwise $P_\alpha = b^{1+\alpha}/a \leq c/a \rightarrow 0$. Note that we have not yet made use of assumption (6.12), only of assumptions (6.5).

We now consider the equation (6.10b) for small α . Note that our assumption (6.12) is equivalent to the assumption that R_0 is negative at s_0 . Hence by continuity R_α is also negative at s_0 for $\alpha > 0$ sufficiently small. We want to show that for all $\alpha \geq 0$ sufficiently small R_α remains negative for all $s \geq s_0$. First consider the case $\alpha = 0$. We have $S_0 = -(2bF_b - aF_a) < 0$ by (6.2b) and (6.5). Hence (6.10b) shows that at a point where $R_0 = 0$ we must have $\dot{R}_0 < 0$. We therefore conclude that $R_0(s) < 0$ for all $s \geq s_0$. In particular, P_0 is strictly decreasing by (6.10a).

Now, since $S_0 < 0$, for all $s \geq s_0$ there exists α_s such that $S_\alpha(s) < 0$ for all $0 \leq \alpha < \alpha_s$. In order to show that we can choose α_s independent of s , using $a, b \rightarrow \infty$ we calculate

$$\lim_{s \rightarrow \infty} S_\alpha = \lim_{s \rightarrow \infty} \alpha(16a^2b^2 - 2b^4) - 8a^2b^2 + 8b^4.$$

Thus $\lim_{s \rightarrow \infty} S_\alpha < 0$ if and only if

$$\alpha < \lim_{s \rightarrow \infty} \frac{4(a^2 - b^2)}{8a^2 - b^2} = \lim_{s \rightarrow \infty} \frac{4(1 - P_0^2)}{8 - P_0^2}.$$

Note that this rational function of P_0 is decreasing for P_0 in $(0, 1)$ with range $(0, 1/2)$. Since P_0 takes values in $(0, 1)$ and is decreasing in s we conclude that $\lim_{s \rightarrow \infty} \alpha_s > 0$ and therefore α_s is bounded below.

As in the case $\alpha = 0$, we now conclude that $R_\alpha < 0$ for all $s \geq s_0$ and any $\alpha \geq 0$ sufficiently small. Then P_α is decreasing and therefore bounded above. We conclude that $b/a \rightarrow 0$ as $s \rightarrow \infty$ since otherwise $P_\alpha = \frac{b}{a}b^\alpha \geq cb^\alpha$ could not be bounded.

We can now study the behaviour of R_α for all $\alpha \in \mathbb{R}$. Using the fact that $a, b \rightarrow \infty$ and a dominates b to show that $2F \approx 8a^2b^2$ and $S_\alpha \approx 8(2\alpha - 1)a^2b^2$ along the complete end, we now conclude that for every $\epsilon > 0$ there exists $s_\epsilon > 0$ such that

$$\frac{dR_\alpha}{da} \leq 2\alpha - 1 + \epsilon \text{ if } R_\alpha \geq 0, \quad \frac{dR_\alpha}{da} \geq 2\alpha - 1 - \epsilon \text{ if } R_\alpha \leq 0$$

for all $s > s_\epsilon$. We conclude that R_α eventually becomes strictly negative if $\alpha < 1/2$ and strictly positive if $\alpha > 1/2$. In other words, for all $\delta > 0$ there exists $s_\delta > 0$ such that

$$\pm(\frac{3}{2}a - \lambda b \pm \delta a) > 0$$

for all $s > s_\delta$. Rearranging and writing $\lambda^{-1} = \frac{db}{da}$, we therefore have $\lim_{t \rightarrow \infty} \frac{a}{b} \frac{db}{da} = \frac{2}{3}$. Note that so far we have not used the assumption that either $q \geq p$ or $q = -p \leq 0$.

In order to show that $P_{1/2}$ converges we now consider the two quantities $P_{1/2}$ and $Q_{1/2}$ at the same time. First note that along the end $S_{1/2} \approx 7b^4 + 8(q - p)a^2b$.

Assume now that $q \geq p$ so that $S_{1/2}$ is eventually positive. In particular, $R_{1/2}$ has a definite sign along the complete end, since if it becomes positive for large enough s then it must remain positive by (6.10b).

If $R_{1/2}$ is eventually negative, then by (6.10a), $P_{1/2}$ is eventually decreasing, and in particular bounded above, and $Q_{1/2}$ is eventually increasing and therefore bounded below. Moreover, since $R_{1/2} < 0$ can be rewritten as $3Q_{1/2} \leq 2P_{1/2}$, we conclude that $P_{1/2}$ and $Q_{1/2}$ are monotone and bounded and therefore convergent. If $R_{1/2}$ is eventually positive, then eventually $3Q_{1/2} \geq 2P_{1/2}$, $P_{1/2}$ is increasing, while $Q_{1/2}$ satisfies

$$\frac{d \log Q_{1/2}}{db} \leq \frac{S_{1/2}}{2bF} \approx \frac{7b}{8a^2} + \frac{q - p}{b^2}.$$

Now, since we know that b^γ/a converges to zero along the end for all $\gamma < 3/2$, we conclude that for b sufficiently large,

$$\frac{d \log Q_{1/2}}{db} \leq c_\gamma(b^{1-2\gamma} + b^{-2})$$

for some constant $c_\gamma > 0$. Choosing $2\gamma \in (2, 3)$ makes the right-hand side integrable in b as $b \rightarrow \infty$. Thus $Q_{1/2}$ is bounded along the end. We conclude that $P_{1/2}$ is bounded above and increasing and therefore convergent. The proof in the case $q \geq p$ is now complete.

When $q = -p < 0$ the final part of the argument, i.e. the convergence of $P_{1/2}$, breaks down because $S_{1/2}$ does not necessarily have a definite sign. We modify the argument as follows. For $\alpha \geq 0$ we now consider functions

$$P_\alpha = \frac{(b - p)^{1+\alpha}}{a - p}, \quad Q_\alpha = \frac{(b - p)^\alpha}{\lambda}.$$

The analogues of (6.10) are

$$(a - p)^2 \dot{P}_\alpha = (b - p)^\alpha \dot{b} R_\alpha, \quad 2F \dot{Q}_\alpha = (b - p)^{-1} \dot{b} Q_\alpha (S_\alpha - F_a R_\alpha),$$

but now we must define

$$R_\alpha = (1 + \alpha)(a - p) - (b - p)\lambda, \quad S_\alpha = \alpha(2F + aF_a - pF_a) - (2bF_b - aF_a) + p(2F_b - Fa).$$

With these new definitions we also have

$$2F \dot{R}_\alpha = \lambda \dot{b} (S_\alpha - F_a R_\alpha).$$

Using $q = -p$ we now calculate

$$\frac{S_\alpha}{2(b - p)^2} = 4(2\alpha - 1)a^2 - 4(1 + \alpha)pa + (4 - \alpha)(b + p)^2 - 4p(b + p).$$

Indeed, if $q = -p$ then $F = (b - p)^2(2a - b - p)(2a + b + p)$ and the formulas in (6.2) become

$$2F_b - F_a = 8(a - b)(2a + b + p)(b - p), \quad 2bF_b - aF_a = 8(a - b)(a + b)(b - p)(b + p).$$

Since $a, b, a/b \rightarrow \infty$ as $s \rightarrow \infty$, we see that $S_\alpha \approx 8(2\alpha - 1)a^2b^2 + 2(4 - \alpha)b^4$ for large s . In particular S_α is eventually positive for all $\alpha \geq 1/2$. The proof now proceeds exactly as before. \square

6.4. Incompleteness

The tools we have developed to prove forward completeness and ALC asymptotics can also be used to prove that certain solutions yield incomplete metrics.

Proposition 6.13. *Let (a, b) be a solution to (3.16) with $a > 0, b > \max(p, -q, \sqrt{-pq})$ and $\dot{a}, \dot{b}, F(a, b) > 0$. If there exists a time such that*

$$0 < \frac{\dot{a}}{\dot{b}} < \frac{a}{b} < 1 \tag{6.14}$$

then the solution cannot be forward complete.

Proof. Recall the functions P_α, R_α and S_α introduced just before Proposition 6.11 and the evolution equations (6.10) for P_α and R_α . We will assume that the solution is complete and derive a contradiction by studying the behaviour of the function P_α for small enough $\alpha < 0$.

We first show that the conditions (6.14) are preserved for all time the solution exists and satisfies $\dot{a}, \dot{b}, F(a, b) > 0$. The fact that the conditions $a < b$ and $\dot{a} < \dot{b}$ persist is proved as in Lemma 6.4 exploiting the fact that $2F_b - F_a < 0$ whenever $0 < a < b$ and $b > \max(p, -q, \sqrt{-pq})$ by (6.2a). In order to prove the persistence of the condition $\dot{a}/\dot{b} < a/b$ consider the quantity R_0 and note that $\dot{a}/\dot{b} < a/b$ is equivalent to $R_0 > 0$. It is enough to observe that $\dot{R}_0 > 0$ at any point where $R_0 = 0$. Indeed, by (6.10b) at a point where $R_0 = 0, \dot{R}_0$ has the same sign as $S_0 = aF_a - 2bF_b$. By (6.2b), $S_0 > 0$ whenever $0 < a < b$ and $b > \max(p, -q, \sqrt{-pq})$. We conclude that $R_0 > 0$ for all time. Note that in particular $P_0 = b/a$ is strictly increasing by (6.10a).

Assume now for a contradiction that the solution is complete. Using $\dot{a} < \dot{b}$ we conclude that $b \rightarrow \infty$ along the complete end in the same way as we proved that a was unbounded in Proposition 6.6. Since $F(a, b) > 0$ for all time, we must also have

$$4\left(\frac{a}{b}\right)^2 > \frac{(b^2 + pq)^2}{b^2(b - p)(b + q)} \rightarrow 1 \tag{6.15}$$

as $b \rightarrow \infty$. Hence a is also unbounded along the complete end. Moreover, since $P_0 = b/a$ is increasing and initially $P_0 > 1$ by (6.14), $a \approx kb$ for some $1/2 \leq k < 1$ as $b \rightarrow \infty$.

We are now going to derive a contradiction to the completeness assumption by studying the behaviour of the function $P_\alpha = b^\alpha/a$ for small enough $\alpha < 0$. Since $S_0 > 0$ as observed earlier, there exists $\alpha_s > 0$ such that $S_\alpha(s) > 0$ for all $|\alpha| < \alpha_s$. In order to show that α_s can be chosen independent of s , using $a \approx kb \rightarrow \infty$ we calculate

$$\lim_{s \rightarrow \infty} \frac{S_\alpha}{b^4} = \lim_{s \rightarrow \infty} \frac{\alpha(16a^2b^2 - 2b^4) - 8a^2b^2 + 8b^4}{b^4} = 2\alpha(8k^2 - 1) + 8(1 - k^2).$$

Since $1/2 \leq k < 1$, both of these coefficients are positive and we conclude that $\lim_{s \rightarrow \infty} S_\alpha > 0$ for all $\alpha > -\frac{4(1-k^2)}{8k^2-1}$. We conclude that $S_\alpha > 0$ for all α sufficiently small as claimed.

Now, the condition $R_0(s_0) > 0$ for some initial time s_0 , which holds by assumption (6.14), forces $R_\alpha(s_0) > 0$ for all small enough α by continuity. Since $S_\alpha > 0$ we can now conclude that $R_\alpha > 0$ for all time and any α sufficiently small as we did above in the case $\alpha = 0$. In particular P_α is positive and increasing for α sufficiently small. If $\alpha < 0$ we now reach a contradiction since $P_\alpha = \frac{b}{a}b^\alpha \approx \frac{1}{k}b^\alpha \rightarrow 0$. \square

6.5. Global behaviour

We now use Propositions 6.11 and 6.13 to describe the global behaviour of the local cohomogeneity one torsion-free G_2 -structures of Proposition 4.5(i, ii) and of Proposition 5.3(i) under an enhanced $U(1)$ -symmetry assumption.

Theorem 6.16. *Let (ijk) be a cyclic permutation of (123) . Consider the local cohomogeneity one torsion-free G_2 -structures of Proposition 4.5(i) and the subfamily defined by $\alpha_j = \alpha_k$.*

- (i) *If $\alpha_i < \alpha_j = \alpha_k$ then the torsion-free G_2 -structure extends to a complete $SU(2) \times SU(2) \times U(1)$ -invariant ALC G_2 -metric on $S^3 \times \mathbb{R}^4$. Here the $U(1)$ -action is the Hopf circle action on the second factor.*
- (ii) *If $\alpha_i = \alpha_j = \alpha_k$ then the torsion-free G_2 -structure is the Bryant–Salamon complete $SU(2)^3$ -invariant AC G_2 -metric on $S^3 \times \mathbb{R}^4$.*
- (iii) *If $\alpha_i > \alpha_j = \alpha_k$ then the torsion-free G_2 -structure is incomplete.*

Theorem 6.17. *Let (ijk) be a cyclic permutation of (123) . Consider the local cohomogeneity one torsion-free G_2 -structures of Proposition 4.5(ii) and the subfamily defined by $\alpha_j = \alpha_k$.*

- (i) *If $\alpha_i < \alpha_j = \alpha_k$ then the torsion-free G_2 -structure extends to a complete $SU(2) \times SU(2) \times U(1)$ -invariant ALC G_2 -metric on $S^3 \times \mathbb{R}^4$. Here the $U(1)$ -action is the Hopf circle action on the first factor.*
- (ii) *If $\alpha_i = \alpha_j = \alpha_k$ then the torsion-free G_2 -structure is the Bryant–Salamon complete $SU(2)^3$ -invariant AC G_2 -metric on $S^3 \times \mathbb{R}^4$.*
- (iii) *If $\alpha_i > \alpha_j = \alpha_k$ then the torsion-free G_2 -structure is incomplete.*

Theorem 6.18. *Consider the conically singular $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structure of Proposition 5.3(i) parametrised by $c \in \mathbb{R}$.*

- (i) *If $c > 0$ then the solution extends to a torsion-free G_2 -structure on $(0, \infty) \times S^3 \times S^3$ with a CS end as $t \rightarrow 0$ and an ALC end as $t \rightarrow \infty$.*
- (ii) *If $c = 0$ then the solution is the G_2 -cone over the $SU(2)^3$ -invariant nearly Kähler structure over $S^3 \times S^3$.*
- (iii) *If $c < 0$ then the solution is incomplete.*

Remark. In all three cases one parameter can be fixed by scaling, hence we have found two 1-parameter families of complete ALC metrics and a unique CS ALC manifold up to scale. The ALC family of Theorem 6.16 is the \mathbb{B}_7 family of Brandhuber–Gomis–Gubser–Gukov [17] and Bogoyavlenskaya [13]. The ALC family of Theorem 6.17 is the conjectured \mathbb{D}_7 family in the physics literature.

Case (ii) in Theorems 6.16–6.18 is characterised by an enhanced $SU(2)$ -symmetry and is therefore dealt with in Example 3.14. We therefore concentrate on proving cases (i) and (iii) in each theorem. Up to a change of basis, we can assume that the additional $U(1)$ -action is generated by $E_3 + E'_3$ in all cases, i.e. $(ijk) = (312)$ in Theorems 6.16 and 6.17.

Proposition 4.5(i, ii) and Proposition 5.3(i) provide the leading-order behaviour of $a = a_1 = a_2$ and $b = a_3$ as $t \rightarrow 0$. We need to check that the hypotheses of Propositions 6.11 and 6.13 are satisfied, i.e. the conditions on p and q together with the inequalities (6.5) and (6.12) or (6.14).

- In the case of Theorem 6.16, $q = -p < 0$ so $b - p, b + q, b - \sqrt{-pq} > 0$. Moreover for small t the signs of $a - b, \dot{a} - \dot{b}$ and $\dot{a}b - a\dot{b}$ are all the same as the sign of $\alpha_1 - \alpha_3$.
- In the case of Theorem 6.17, $q = 0$ and $p = -r_0^3 < 0$. Moreover, $\alpha_3 > 0$ since $\alpha_1\alpha_2\alpha_3 = 1$. Then $b - p, b + q, b - \sqrt{-pq} > 0$. The signs of $a - b, \dot{a} - \dot{b}$ are the same as the sign of $\alpha_1 - \alpha_3$. In order to control the sign of $\dot{a}b - a\dot{b}$ for small t we have to use the higher-order expansions of a, b in Remark 4.6. We find

$$\dot{a}b - a\dot{b} = \frac{(1 - \alpha_3^3)\alpha_3}{96\alpha_1}t^5 + O(t^7).$$

Since $\alpha_1^2\alpha_3 = 1$, if $\alpha_3 < \alpha_1$ then $0 < \alpha_3 < 1 < \alpha_1$, while $0 < \alpha_1 < 1 < \alpha_3$ if $\alpha_3 > \alpha_1$.

- In the case of Theorem 6.18, $p = 0 = q$ and for small $t > 0$ we have $b > 0$. The signs of $a - b, \dot{a} - \dot{b}$ are the same as the sign of c . In order to control the sign of $\dot{a}b - a\dot{b}$ we calculate

$$\frac{54^2(\dot{a}b - a\dot{b})}{3t^5} \approx \frac{3}{2}c\nu_0t^{\nu_0}$$

as $t \rightarrow 0$.

Remark. Theorems 6.16 and 6.17 do not address the issue of what happens to solutions that do not enjoy an enhanced $U(1)$ -symmetry. Our expectation is that all of these local cohomogeneity one metrics are incomplete.

7. Existence of AC metrics

In Proposition 4.5 we constructed four different families of local solutions to (3.13) that close smoothly over various different singular orbits. In the previous section we have shown that a subset of the local solutions closing smoothly on a singular orbit S^3 con-

structured in Proposition 4.5(i, ii) extend to complete ALC metrics. None of the local solutions constructed in Proposition 4.5(iii, iv) is covered by these results: no choice of parameters there allows us to satisfy all the hypotheses of Propositions 6.11 or 6.13 for small positive t . In this section we use a different approach to study $SU(2) \times SU(2) \times U(1)$ -invariant G_2 -manifolds with singular orbit $SU(2) \times SU(2)/K_{m,n}$ and thus prove the following theorem.

Theorem 7.1. *Fix coprime positive integers m, n and a real number $r_0 > 0$. For $\beta > 0$, let*

$$\varphi_\beta = -m^2 r_0^3 e_1 \wedge e_2 \wedge e_3 + n^2 r_0^3 e'_1 \wedge e'_2 \wedge e'_3 + d(a(e_1 \wedge e'_1 + e_2 \wedge e'_2) + b e_3 \wedge e'_3)$$

be the (locally defined) $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structure closing smoothly on $SU(2) \times SU(2)/K_{m,n}$ defined in Proposition 4.5(iii) (when $m = n = 1$) or (iv) satisfying

$$a = r_0^2 \beta t + O(t^3), \quad b = mnr_0^3 + O(t^2)$$

as $t \rightarrow 0$. There exists $\beta_{ac} > 0$ such that the following holds.

- (i) If $\beta > \beta_{ac}$ then φ_β extends to a complete torsion-free ALC G_2 -structure asymptotic to a circle bundle over a \mathbb{Z}_2 -quotient conifold.
- (ii) If $\beta = \beta_{ac}$ then φ_β extends to a complete torsion-free AC G_2 -structure asymptotic to the cone over the $\mathbb{Z}_{2(m+n)}$ -quotient of the homogeneous nearly Kähler structure on $S^3 \times S^3$ with rate -3 .
- (iii) If $\beta < \beta_{ac}$ then φ_β does not extend to a complete torsion-free G_2 -structure.

Figure 1 illustrates a small number of solution curves (a, b) to the ODE system (3.16) illustrating all three cases of Theorem 7.1 in the case where $m = 1$ and $n = 2$.

Remark. The existence of the ALC metrics in part (i) of the theorem when β is sufficiently large is guaranteed by our analytic construction of highly collapsed ALC G_2 -metrics on circle bundles over AC Calabi–Yau 3-folds [35, Theorem 9.7].

Remark. The case $m = n = 1$ has also been considered by Bazaikin–Bogoyavlenskaya [9] and Cvetič–Gibbons–Lü–Pope [28, §3]. In [9] the existence of ALC metrics for every positive value of β was claimed. However, there appear to be mistakes in the proof of [9, Lemma 9]. In [28, §3] numerical experiments suggested the existence of a full 3-parameter family of ALC G_2 -metrics closing smoothly on the singular orbit $SU(2) \times SU(2)/K_{1,1}$. In the collapsed limit this contradicts our analysis in [35, Theorem 9.7].

The most interesting part of Theorem 7.1 is part (ii). Only three simply connected AC G_2 -metrics (up to symmetries and scaling) are currently known [18]. Part (ii) of the theorem provides infinitely many new AC G_2 -metrics. It is essential that we consider AC manifolds asymptotic to a non-trivial quotient of the G_2 -cone over $S^3 \times S^3$ since, by Karigiannis–Lotay [43, Corollary 6.10], the Bryant–Salamon metric on the spinor bundle of S^3 is the unique (up to scale) AC G_2 -metric asymptotic to the cone over $S^3 \times S^3$.




Fig. 1. Numerical solutions of (3.16) with $m = 1, n = 2$ satisfying the initial conditions of Proposition 4.5(iv): There are two incomplete solutions approaching the set $F = 0$, the unique AC solution asymptotic to the diagonal $a = b$ and two complete solutions crossing the diagonal transversally.

Remark 7.2. The smooth 7-manifold $M_{m,n}$ underlying the G_2 -metrics constructed in the theorem depends only on the sum $m+n$. In fact, $M_{m,n}$ can be identified with $H^{2(n+m)} \times S^3$, where $H^{2(n+m)}$ is the total space of the \mathbb{R}^2 -bundle on S^2 with Euler class $2(n+m)$. However, metrics for different choices of (m, n) can never be isometric. Indeed, an isometry would have to respect the $SU(2) \times SU(2) \times U(1)$ -orbit structure: otherwise the tangent space at a point would be spanned by Killing vectors and the metric would be homogeneous; the latter is impossible since the metric is Ricci-flat but cannot be flat. The group of $SU(2) \times SU(2) \times U(1)$ -equivariant diffeomorphisms of the principal orbit $SU(2) \times SU(2)/\mathbb{Z}_{2(m+n)}$ is $SU(2) \times SU(2) \times N \rtimes \mathbb{Z}_2$, where N is the normaliser of $U(1)$ in $SU(2)$ acting on the right on $SU(2) \times SU(2)$ and \mathbb{Z}_2 is generated by the outer automorphism of $SU(2) \times SU(2)$ that exchanges the two factors. The induced action of $SU(2) \times SU(2) \times N \rtimes \mathbb{Z}_2$ on cohomology is generated by the involution that exchanges $e_1 \wedge e_2 \wedge e_3$ and $e'_1 \wedge e'_2 \wedge e'_3$. Since the image of the cohomology class of φ in the cohomology of the principal orbits depends on the pair (m, n) , we conclude that different choices of (m, n) with $0 < m \leq n$ and $\gcd(m, n) = 1$ give rise to non-isometric metrics. In fact this argument also shows that, modulo the outer automorphism of $SU(2) \times SU(2)$, there is no diffeomorphism between $M_{m,n}$ and $M_{m',n'}$ asymptotic to an isometry of the asymptotic cone. In particular, considering pairs (m, n) with $0 < m \leq n$ and fixed (sufficiently large) $m+n$, part (ii) of the theorem yields different AC G_2 -metrics asymptotic

to the same G_2 -cone and therefore gives rise to infinitely many new geometric transitions in G_2 -geometry.

Before proving Theorem 7.1 we show that the complete solutions obtained in Theorems 6.16, 6.17 and 7.1(i, ii) are the only complete simply connected $SU(2) \times SU(2) \times U(1)$ -invariant G_2 -manifolds.

Theorem 7.3. *Let (M, g) be a complete $SU(2) \times SU(2) \times U(1)$ -invariant G_2 -metric with M simply connected. Then up to symmetries (M, g) is isometric to one of the complete metrics of Theorems 6.16, 6.17 and 7.1.*

Proof. Given the completeness and incompleteness statements in Theorems 6.16, 6.17 and 7.1, it only remains to prove that if (M, g) is an $SU(2) \times SU(2) \times U(1)$ -invariant G_2 -metric closing smoothly on a singular orbit Q , then, up to a finite cover and the action of the outer automorphism of $SU(2)^2$, M is described by one of the group diagrams (2.4), (2.5) and (2.1).

Let

$$\varphi = p e_1 \wedge e_2 \wedge e_3 + q e'_1 \wedge e'_2 \wedge e'_3 + d(a(e_1 \wedge e'_1 + e_2 \wedge e'_2) + b e_3 \wedge e'_3)$$

be an $SU(2) \times SU(2) \times U(1)$ -invariant torsion-free G_2 -structure defined in a neighbourhood of a singular orbit Q . Let t be the arc-length parameter along a geodesic meeting all orbits orthogonally and assume that the point $t = 0$ lies on the singular orbit Q . In particular, a, b are smooth functions defined on $[0, t_0)$ for some $t_0 > 0$.

In order to determine the behaviour of the functions a and b as $t \rightarrow 0$, observe that $F \rightarrow 0$ as $t \rightarrow 0$ since $\sqrt{F(a, b)} = 2\dot{a}^2\dot{b}$ is the orbital volume function. Moreover, the evolution equations for $x_1 = \dot{a}\dot{b}$ and $x_2 = \dot{a}^2$ in (3.17) show that $(F_a, F_b) \rightarrow 0$ as $t \rightarrow 0$, i.e. the point $(a_0, b_0) = (a, b)|_{t=0}$ must be a critical point of F on the level set $F = 0$. As an aside, note that we must have $pq \leq 0$ since $F = 0 = F_a$ force $b^2 + pq = 0$. Since the coefficients of the autonomous ODE system (3.17) depend real analytically on a and b , we conclude that there exist a critical point (a_0, b_0) of F with $F(a_0, b_0) = 0$, positive integers h, k and $a_1, b_1 \neq 0$ such that $a = a_0 + a_1 t^h + O(t^{h+1})$ and $b = b_0 + b_1 t^k + O(t^{k+1})$. In fact, since with our conventions $\dot{a}, \dot{b} > 0$ for $t > 0$, we must have $a_1, b_1 > 0$.

We now consider the metric $g_\varphi = dt^2 + g_t$ induced by φ . Regard M as a cohomogeneity one manifold with group diagram

$$K_0 \subset K \subset SU(2) \times SU(2),$$

where $Q = SU(2)^2/K$, K_0 is a finite subgroup of $SU(2)^2$ and K/K_0 is a sphere. As $t \rightarrow 0$, g_t converges to a smooth metric g_0 on the singular orbit Q . Moreover, thinking of g_0 as a symmetric endomorphism of $\mathfrak{su}_2 \oplus \mathfrak{su}_2$, the kernel of g_0 coincides with the Lie algebra \mathfrak{k} of K . By studying the behaviour of g_t as $t \rightarrow 0$ we can therefore determine the possibilities for \mathfrak{k} and therefore the group diagram of M up to finite quotients.

Denote by $\mathfrak{n}, \mathfrak{n}'$ and \mathfrak{t} the subspaces of $\mathfrak{su}_2 \oplus \mathfrak{su}_2$ defined by $\text{span}(E_1, E_2)$, $\text{span}(E'_1, E'_2)$ and $\text{span}(E_3, E'_3)$ respectively. Since K/K_0 is a sphere and K_0 is finite, we

deduce that \mathfrak{k} cannot contain $\mathfrak{n} \oplus \mathfrak{n}'$ or \mathfrak{t} . Indeed, if $\mathfrak{n} \oplus \mathfrak{n}' \subseteq \mathfrak{k}$ then $K = \text{SU}(2) \times \text{SU}(2)$ (since $\mathfrak{n} \oplus \mathfrak{n}'$ generates $\mathfrak{su}_2 \oplus \mathfrak{su}_2$ as a Lie algebra) and K/K_0 cannot be a sphere; similarly, if $\mathfrak{t} \subseteq \mathfrak{k}$ then K is diffeomorphic to a 2-torus, $S^3 \times S^1$ or $S^3 \times S^3$, none of which finitely covers a sphere.

Now, in order to study the behaviour of g_t for small $t \geq 0$, regard it as a symmetric endomorphism of $\mathfrak{su}_2 \oplus \mathfrak{su}_2$ and note that the decomposition $\mathfrak{su}_2 \oplus \mathfrak{su}_2 = (\mathfrak{n} \oplus \mathfrak{n}') \oplus \mathfrak{t}$ is g_t -orthogonal. By (3.18) the restriction of g_t to the first factor $\mathfrak{n} \oplus \mathfrak{n}'$ is the block matrix

$$\frac{2\dot{a}}{\sqrt{F}} \begin{pmatrix} a(b-p) & -\frac{1}{2}(b^2 + pq) \\ -\frac{1}{2}(b^2 + pq) & a(b+q) \end{pmatrix},$$

and the restriction of g_t to \mathfrak{t} is

$$\frac{2\dot{b}}{\sqrt{F}} \begin{pmatrix} a^2 - pb & -\frac{1}{2}(2a^2 - b^2 + pq) \\ -\frac{1}{2}(2a^2 - b^2 + pq) & a^2 + qb \end{pmatrix}.$$

Now, consider first the case where $pq(p+q) \neq 0$. In this case F has only two critical points contained in the level set $F = 0$, $(0, 0, \pm\sqrt{-pq})$. In fact, since $\dot{a}, \dot{b} > 0$ for $t > 0$ we must have $b_0 = \sqrt{-pq}$, $p < 0$ and $q > 0$. Indeed, since $a(0) = 0$, the sign constraint $\dot{a} > 0$ for $t > 0$ forces the same sign constraint for a . Then the positive definiteness of g_t for $t > 0$ forces $-pb_0, qb_0, b_0 - p, b_0 + q > 0$.

Using $a = a_1 t^h + O(t^{h+1})$ and $b = \sqrt{-pq} + b_1 t^k + O(t^{k+1})$, we now calculate $F = O(t^m)$ where $m \geq 2h$. Indeed, $h \leq k$ since $F > 0$ for $t > 0$. Moreover, $m > 2h$ if and only if $h = k$ and a_1, b_1 are appropriately chosen so that the coefficient of t^{2h} in F vanishes.

At leading order in t as $t \rightarrow 0$, the restriction of g_t to \mathfrak{t} takes the form

$$ct^{k-1-m/2} \begin{pmatrix} |p| & \sqrt{-pq} \\ \sqrt{-pq} & |q| \end{pmatrix}$$

for some $c > 0$, while the restriction of g_t to $\mathfrak{n} \oplus \mathfrak{n}'$ is of the form

$$\begin{pmatrix} O(t^{2h-1-m/2}) & O(t^{h+k-1-m/2}) \\ O(t^{h+k-1-m/2}) & O(t^{2h-1-m/2}) \end{pmatrix}.$$

Since the kernel \mathfrak{k} of g_0 cannot contain \mathfrak{t} or $\mathfrak{n} \oplus \mathfrak{n}'$, we deduce that $k - 1 - m/2 = 2h - 1 - m/2 = 0$, i.e. $h = 1, k = 2$. Then \mathfrak{k} is 1-dimensional, spanned by $\sqrt{|q|}E_3 - \sqrt{|p|}E'_3$. The orbit of this vector field in $\text{SU}(2)^2$ is closed if and only if $\sqrt{-p/q} \in \mathbb{Q}$. If this is the case there exist relatively prime positive integers m, n and $r_0 > 0$ such that $K = K_{m,n}$ up to finite quotients and $p = -m^2 r_0^3, q = n^2 r_0^3$.

Finally, since $K = K_{m,n}$ is a circle we must also argue that the principal orbit stabiliser K_0 is $K_{m,n} \cap K_{2,-2}$. This is a consequence of the proof of Proposition 4.5(iv). Indeed, the proposition parametrises all smooth solutions (x_1, x_2, y_1, y_2) to (3.17) with $p = -m^2 r_0^3$ and $q = n^2 r_0^3$ satisfying $x_1 = O(t), x_2 = r_0^4 \beta + O(t^2), y_1 = O(t)$ and $y_2 = mn r_0^3 + O(t^2)$. We have already established that $a = a_1 t + O(t^2)$ and

$b = mn r_0^3 + b_1 t^2 + O(t^3)$ for some $a_1, b_1 > 0$. Hence $(x_1 = \dot{a}b, x_2 = \dot{a}^2, y_1 = a, y_2 = b)$ coincides with one of the solutions of Proposition 4.5(iv).

We now briefly indicate the changes to the proof in the case where $pq(p + q) = 0$. If $p + q = 0$ the critical locus of F contained in the zero-level set is $\{(a_0, p) \mid a_0 \in \mathbb{R}\} \cup \{(0, -p)\}$. Consider first the 1-dimensional component. Boundedness of the restriction of g_t to \mathfrak{t} as $t \rightarrow 0$ forces $a_0 = \pm p$. If $p \neq 0$, consideration of the behaviour of the restriction of g_t to $\mathfrak{n} \oplus \mathfrak{n}'$ as $t \rightarrow 0$ implies that $p > 0$ and $\mathfrak{k} = \Delta \mathfrak{su}_2$. If $p = 0$, one shows instead that it is impossible to find $h, k \geq 1$ so that g_t remains bounded as $t \rightarrow 0$ and \mathfrak{k} contains neither $\mathfrak{n} \oplus \mathfrak{n}'$ nor \mathfrak{t} . The case $p \neq 0$ and $(a_0, b_0) = (0, -p)$ is analysed exactly as in the case $pq(p + q) \neq 0$. When $p \neq 0$ and $q = 0$ (the case $p = 0, q \neq 0$ can be reduced to this by acting with the outer automorphism of $SU(2)^2$), F has a unique critical point, $(0, 0)$, on its zero-level set. Analysis of the behaviour of the restriction of g_t to $\mathfrak{n} \oplus \mathfrak{n}'$ as $t \rightarrow 0$ forces $\mathfrak{n} \oplus \{0\} \subseteq \mathfrak{k}$. The only possibility for \mathfrak{k} is then $\mathfrak{su}_2 \oplus \{0\}$. Finally, since K is 3-dimensional in all these cases, K_0 is automatically trivial. \square

Remark 7.4. In particular, we deduce a strong rigidity and uniqueness result for the AC G_2 -metrics of Theorem 7.1(ii). Indeed, by [43, Propositions 6.3 and 6.8] any complete AC G_2 -manifold (M, g) asymptotic to the cone over $S^3 \times S^3/\mathbb{Z}_{2(n+m)}$ must be $SU(2) \times SU(2) \times U(1)$ -invariant. By Theorem 7.3, up to a finite quotient, any such metric has group diagram either (2.4), (2.5) and (2.1). The proof of Theorems 6.16, 6.17 and 7.1 then shows that (M, g) is either a finite quotient of the Bryant–Salamon AC metric on $S^3 \times \mathbb{R}^4$ (if $\mathbb{Z}_{2(m+n)}$ acts freely on $S^3 \times \mathbb{R}^4$) or one of the AC metrics of Theorem 7.1(ii).

In the rest of the section we prove Theorem 7.1. We first establish part (ii) of the theorem and the existence of the critical value β_{ac} , which is not explicit. Our strategy is to consider the AC ends constructed in Proposition 5.3(ii) and study which of these extend *backward* to close smoothly on the singular orbit $SU(2) \times SU(2)/K_{m,n}$. Once part (ii) of Theorem 7.1 is established, a comparison argument with the AC solution will let us obtain the existence of the ALC G_2 -metrics in part (i) and the incompleteness result in part (iii).

7.1. Extending AC ends backwards

Fix a pair of positive coprime integers m, n and $r_0 \in \mathbb{R}$ and set $p = -m^2 r_0^3, q = n^2 r_0^3$. We consider pairs of functions (a, b) satisfying the ODE (3.16), i.e.

$$2F(\ddot{a}b - \dot{a}\ddot{b}) = -\dot{a}\dot{b}(2\dot{b}F_b - \dot{a}F_a),$$

where

$$F = 4a^2(b + m^2 r_0^3)(b + n^2 r_0^3) - (b^2 - m^2 n^2 r_0^6)^2. \tag{7.5}$$

By Proposition 5.3(ii) for each $c \in \mathbb{R}$ there exists a solution (a, b) corresponding to an AC end asymptotic to the cone over the homogeneous nearly Kähler structure on $S^3 \times S^3$. We now consider the problem of extending these AC ends backwards, i.e. to decreasing values of the parameter t .

Proposition 7.6. *Suppose that $r_0 \geq 0$ and (a, b) is a solution to (3.16) satisfying*

$$\dot{a}, \dot{b} > 0, \quad a > 0, \quad b > \max(-m^2 r_0^3, -n^2 r_0^3), \quad F > 0 \tag{7.7a}$$

and

$$b > a, \quad \dot{a} > \dot{b} > 0 \tag{7.7b}$$

at some time t_0 (or equivalently, since (7.7) are open conditions, on an open interval (t_1, t_2) of existence). Then the solution extends backwards in time, with the conditions (7.7) persisting, until $F(a, b) \rightarrow 0$, i.e. until

$$2a - \frac{|b^2 - m^2 n^2 r_0^6|}{\sqrt{(b + m^2 r_0^3)(b + n^2 r_0^3)}} \rightarrow 0.$$

Proof. Set $\mu := \dot{b}/\dot{a}$. As at the beginning of the proof of Proposition 6.11, we rewrite (3.16) in the form

$$2F\dot{\mu} = \mu\dot{a}(F_a - 2\mu F_b). \tag{7.8}$$

The coefficient of $\dot{\mu}$ on the left-hand side of the equation is positive. On the other hand, thanks to our hypotheses (7.7b), μ takes values in the interval $(0, 1)$ and therefore $F_a - 2\mu F_b$ is greater than the minimum of F_a and $F_a - 2F_b$. Using (7.7a) one can check that the former is always positive and the latter is a concave function of a which is non-negative for both $a = 0$ and $a = b$. Thus $F_a - 2\mu F_b$ is positive whenever $b > a > 0$. We conclude that $\dot{\mu} > 0$.

Now, since $\dot{\mu} > 0$, the inequality $\mu < 1$ is preserved as we evolve backwards, and hence $b > a$ is also preserved. It remains to prove that we can extend backwards until $F(a, b) \rightarrow 0$.

Since (3.16) is equivalent to Hitchin’s flow (3.17) for the 4-tuple $(\dot{a}\dot{b}, \dot{a}^2, a, b)$, it is clear that solutions fail to extend only when one of a, b, \dot{a}, \dot{b} diverges or when one of the inequalities $\dot{a}, \dot{b}, F(a, b) > 0$ fails to be satisfied. Now, for any $M > 0$, the curves $\{a = b\}$, $\{F(a, b) = 0\}$ and $\{a^2 + b^2 = M^2\}$ bound a compact region $\mathcal{R} \subset \{F \geq 0\}$ in the first quadrant in the (a, b) -plane. By assumption, the curve (a, b) lies in \mathcal{R} for some $M > 0$ and therefore the solution can be extended backward until $F(a, b) \rightarrow 0$ provided we control \dot{a}, \dot{b} . Since $2\dot{a}^2\dot{b} = \sqrt{F(a, b)}$ and $\dot{b} < \dot{a}$ it is enough to prove that \dot{b} is bounded away from zero until $F(a, b) \rightarrow 0$.

Now, since $\dot{a} > 0$, we can reparametrise so that $\dot{a} = 1$. Then (7.8) becomes

$$\frac{d \log \dot{b}}{da} = \frac{F_a - 2\mu F_b}{2F}$$

The right-hand side can only blow up as $F(a, b) \rightarrow 0$, so until then $\log \dot{b}$ remains bounded. Thus \dot{b} is bounded away from zero until $F(a, b) \rightarrow 0$. □

The solutions (a, b) of (3.16) constructed in Proposition 5.3(ii) always satisfy (7.7a) in the interior of a maximal interval of existence. Moreover, the solutions of Proposition 5.3(ii) satisfy

$$b - a \approx \frac{\sqrt{3}}{54} ct^{3-\nu_\infty}$$

as $t \rightarrow \infty$ for some $c \in \mathbb{R}$. When c is positive, (7.7b) will then also hold for t sufficiently large. Hence we can apply Proposition 7.6 to conclude that the solutions constructed in Proposition 5.3(ii) with $c > 0$ extend backward until $F(a, b) \rightarrow 0$. In the limiting case $c = 0$, the uniqueness statement in Proposition 5.3(ii) implies that $a = b$.

We will prove that there exists $c_{ac} > 0$ such that, after taking the quotient of the principal orbits by $\mathbb{Z}_{2|m+n|}$, the AC G_2 -metric corresponding to the solution (a_{ac}, b_{ac}) constructed in Proposition 5.3 (ii) with $c = c_{ac}$ extends smoothly over a singular orbit $SU(2) \times SU(2)/K_{m,n}$. The following lemma will be used to show that there exists a unique such value c_{ac} .

Lemma 7.9. *Fix a pair of positive coprime integers m and n and suppose that $r_0 \geq 0$. Let (a_1, b_1) and (a_2, b_2) be solutions of (3.16) satisfying*

$$b > \max(a, mnr_0^3), \quad F(a, b) > 0, \quad \dot{a}, \dot{b} > 0. \tag{7.10}$$

Parametrise the curves (a_1, b_1) and (a_2, b_2) so that $a_1(s) = s = a_2(s)$.

- (i) *The inequalities $b_1 > b_2, \dot{b}_1 < \dot{b}_2$ are preserved evolving (a_1, b_1) and (a_2, b_2) backwards until either solution hits the boundary of the region defined by (7.10).*
- (ii) *The inequalities $b_1 < b_2, \dot{b}_1 < \dot{b}_2$ are preserved evolving (a_1, b_1) and (a_2, b_2) forwards until either solution hits the the boundary of the region defined by (7.10).*

Proof. Let (a, b) be a solution (3.16) parametrised so that $\dot{a} = 1$. Then (3.16) can be rewritten as

$$2\ddot{b} = \left(\frac{F_a}{F} - 2\dot{b} \frac{F_b}{F} \right) b\dot{b}. \tag{7.11}$$

In order to compare two solutions (a_1, b_1) and (a_2, b_2) we now observe that, for each fixed $a > 0$, F_a/F and $-F_b/F$ are strictly increasing functions of b on the range defined by the inequalities $b > \max(a, mnr_0^3)$ and $F(a, b) > 0$. Indeed, the function

$$\frac{F_a}{F} = \frac{8a}{4a^2 - \frac{(b^2 - m^2n^2r_0^6)^2}{(b+m^2r_0^3)(b+n^2r_0^3)}}$$

is increasing in b if and only if

$$\frac{(b^2 - m^2n^2r_0^6)^2}{(b + m^2r_0^3)(b + n^2r_0^3)} = (b - mnr_0^3)^2 \frac{(b + mnr_0^3)^2}{(b + mnr_0^3)^2 + (m - n)^2r_0^3b}$$

is. Each factor on the right-hand side is increasing precisely when $b > mnr_0^3$. Meanwhile $F_{bb} = 4(2a^2 - 3b^2 + m^2n^2r_0^6) < 0$ in the given range, while F is a priori positive. Thus

$$-\frac{d}{db} \left(\frac{F_b}{F} \right) = \frac{-F_{bb}F + F_b^2}{F^2} > 0.$$

Going back to (7.11), we now conclude that at every point where $\dot{b}_1 = \dot{b}_2, \ddot{b}_1 - \ddot{b}_2$ has the same sign as $b_1 - b_2$. Hence the inequalities $b_1 > b_2$ and $\dot{b}_1 < \dot{b}_2$ [$b_1 < b_2$ and $\dot{b}_1 < \dot{b}_2$] are preserved as we evolve backwards [forwards] as long as both solutions remain in the region defined by the inequalities (7.10). □

Proposition 7.12. *For each $r_0 > 0$ there exists a unique $c_{ac} > 0$ such that the solution of (3.16) constructed in Proposition 5.3(ii) with $c = c_{ac}$ extends backwards until $a \rightarrow 0$ and $b \rightarrow mnr_0^3$. Moreover, for any $k \in (1, 2)$ the solution satisfies $ka > \frac{|b^2 - m^2n^2r_0^6|}{\sqrt{(b+m^2r_0^3)(b+n^2r_0^3)}}$ and $b > mnr_0^3$ whenever $(a, b) \neq (0, mnr_0^3)$.*

Proof. Fix $k \in (1, 2)$. For any $c \geq 0$, evolve the AC end solution of Proposition 5.3(ii) backwards until we hit the curve γ which is the union of the curves

$$\gamma_1 = \{b = mnr_0^3\} \quad \text{and} \quad \gamma_2 = \left\{ ka = \frac{|b^2 - m^2n^2r_0^6|}{\sqrt{(b + m^2r_0^3)(b + n^2r_0^3)}} \right\}.$$

Since γ is smooth except at the intersection point $(0, mnr_0^3)$ of γ_1 and γ_2 , the intersection point of the trajectory (a, b) with γ depends continuously on c unless (a, b) approaches $(0, mnr_0^3)$.

The uniqueness statement in Proposition 5.3(ii) implies that the solution with $c = 0$ satisfies $a = b$ and therefore it must hit γ along the segment γ_1 . In order to analyse the behaviour of the intersection point as $c \rightarrow \infty$, we can rescale so that $r_0 \rightarrow 0$ and $c > 0$ is fixed. This solution must hit the half-line $ka = b, b > 0$ away from $(0, 0)$ since $a - b$ is strictly increasing for all time and strictly negative along the AC end.

We conclude that for each $r_0 > 0$ the set S of $c \geq 0$ for which we hit the segment γ_1 is non-empty and bounded. Set $c_{ac} = \sup S$. The solution corresponding to $c = c_{ac}$ can hit neither γ_1 nor γ_2 , so by Proposition 7.6 the only possibility is that it approaches $(0, mnr_0^3)$.

The uniqueness part of the claim follows from Lemma 7.9(i). Indeed, if (a, b) is a solution of (3.16) given by Proposition 5.3(ii) then

$$a \approx \sqrt[3]{\frac{3}{54}}t^3, \quad b - a \approx \sqrt[3]{\frac{3}{54}}ct^{3-\nu_\infty}$$

for some $c \in \mathbb{R}$. Hence if we parametrise (a, b) so that $a(s) = s$ we have

$$b \approx s + \left(\frac{\sqrt{3}}{54}\right)^{\nu_\infty/3} cs^{(3-\nu_\infty)/3}$$

as $s \rightarrow \infty$. Since $3 - \nu_\infty \approx -6.5$, if (a_1, b_1) and (a_2, b_2) are two solutions corresponding to $c_1 > c_2 > 0$ then

$$b_1 > b_2, \quad \dot{b}_1 < \dot{b}_2$$

for large s . □

In order to conclude the proof of Theorem 7.1(ii) we must show that after a $\mathbb{Z}_{2(m+n)}$ -quotient the AC solution with $c = c_{ac}$ singled out by Proposition 7.12 extends smoothly across the singular orbit $SU(2) \times SU(2)/K_{m,n}$.

Proposition 7.13. Fix $k \in (1, 2)$. Consider a solution (a, b) satisfying (7.7), $b > mnr_0^3$ and

$$ka > \frac{b^2 - m^2n^2r_0^6}{\sqrt{(b + m^2r_0^3)(b + n^2r_0^3)}}. \tag{7.14}$$

Assume that $(a, b) \rightarrow (0, mnr_0^3)$. Then (a, b) defines a cohomogeneity one torsion-free G_2 -structure with principal orbits $SU(2) \times SU(2)/\mathbb{Z}_{2(m+n)}$ and extending smoothly on the singular orbit $SU(2) \times SU(2)/K_{m,n}$.

Proof. We must show that (a, b) satisfies the boundary conditions of Proposition 4.1(iii).

First of all note that (7.14) implies that the function F in (7.5) satisfies

$$(4 - k^2)(b + m^2r_0^3)(b + n^2r_0^3)a^2 \leq F(a, b) \leq 4(b + m^2r_0^3)(b + n^2r_0^3)a^2. \tag{7.15a}$$

Differentiating the expression in (7.5) and using (7.14) and (7.15a) we also obtain

$$\sqrt{(b + m^2r_0^3)(b + n^2r_0^3)} \leq \frac{F_a}{4\sqrt{F}} \leq \frac{2}{\sqrt{4 - k^2}} \sqrt{(b + m^2r_0^3)(b + n^2r_0^3)} \tag{7.15b}$$

and

$$-2kb(b + mnr_0^2) \leq \frac{F_b}{2\sqrt{F}} \leq \frac{2}{\sqrt{4 - k^2}} \frac{(2b + (m^2 + n^2)r_0^3)a}{\sqrt{(b + m^2r_0^3)(b + n^2r_0^3)}}. \tag{7.15c}$$

Reparametrise so that $\dot{a} = 1$ and consider the first-order equation (7.8) for $\mu = \frac{db}{da}$. Since $\frac{dF}{da} = F_a + \mu F_b$, we find

$$\frac{d}{da} \left(\frac{\sqrt{F}}{\mu} \right) = \frac{\sqrt{F}}{\mu} \frac{3\mu F_b}{2F} \leq C \frac{\sqrt{F}}{\mu}$$

for some $C > 0$. Indeed, F_b/F is bounded above by (7.15a) and (7.15c) and $0 < \mu < 1$. Thus the logarithm of \sqrt{F}/μ is a function of a whose derivative is bounded above. It follows that \sqrt{F}/μ is bounded below away from zero as $a \rightarrow 0$.

Now change parametrisation and introduce the arc-length parameter t along a geodesic meeting all principal orbits orthogonally. Recall that t is defined by the normalisation $2\dot{a}^2\dot{b} = \sqrt{F(a, b)}$. By (7.15a) and the hypothesis $\dot{b} < \dot{a}$ in (7.7b) we have

$$Ca \leq \sqrt{F(a, b)} = 2\dot{a}^2\dot{b} < 2\dot{a}^3$$

for some $C > 0$. Since the function $a \mapsto a^{-1/3}$ is integrable near $a = 0$, t has a finite limit as $(a, b) \rightarrow (0, mnr_0^3)$. By a time translation we can assume that $t \rightarrow 0$ as $(a, b) \rightarrow (0, mnr_0^3)$. Moreover, since

$$\frac{\sqrt{F}}{\mu} = \frac{2\dot{a}^2\dot{b}}{\mu} = 2\dot{a}^3$$

and \sqrt{F}/μ is bounded below away from zero, so is \dot{a} .

Consider now the ODE system (3.17) for $x_1 = \dot{a}\dot{b}$, $x_2 = \dot{a}^2$, $y_1 = a$, $y_2 = b$. By (7.15b) and (7.15c), \dot{x}_1 and \dot{x}_2 remain bounded and therefore x_1 and x_2 have a well-defined limit as $t \rightarrow 0$. Since $2\dot{b}^3 \leq 2\dot{a}^2\dot{b} = \sqrt{F} \rightarrow 0$ as $t \rightarrow 0$ and \dot{a} is bounded away from zero, $(x_1, x_2) \rightarrow (0, r_0^4\beta_{ac}^2)$ as $t \rightarrow 0$ for some $\beta_{ac} > 0$. Then the right-hand side of (3.17) is bounded as $t \rightarrow 0$ and a bootstrap argument shows that x_1, x_2, y_1, y_2 are smooth functions of t up to $t = 0$. Uniqueness of the solutions in Proposition 4.5(iii, iv) then forces (a, b) to satisfy the conditions of Proposition 4.1(iii). \square

7.2. Existence of ALC metrics

We now prove Theorem 7.1(i). Theorem 7.1(ii) guarantees the existence of $\beta_{ac} > 0$ such that the $SU(2) \times SU(2) \times U(1)$ -invariant local solution (a_{ac}, b_{ac}) of Proposition 4.5(iii) (when $m = n = 1$) or (iv) with

$$a_{ac} = r_0^2\beta_{ac}t + O(t^3), \quad b_{ac} = mnr_0^3 + O(t^2)$$

exists for all time $t \geq 0$ and gives rise to a complete AC metric. From the proof of Theorem 7.1(ii) we also know that

$$b_{ac} > \max(a_{ac}, mnr_0^3), \quad \dot{a}_{ac} > \dot{b}_{ac} > 0$$

for all $t > 0$.

Consider now one of the local solutions (a, b) of Proposition 4.5(iii, iv) with $\beta > \beta_{ac}$. We want to show that (a, b) eventually satisfies the constraints (6.5) and (6.12), i.e.

$$a > b > mnr_0^3, \quad \dot{a} > \dot{b}, \quad a\dot{b} - \dot{a}b < 0,$$

so that Propositions 6.6 and 6.11 can be applied to guarantee that the solution (a, b) is immortal and gives rise to an ALC end as $t \rightarrow \infty$. In fact, it is enough to show that the solution (a, b) , initially contained in the region

$$a < b, \quad \dot{a} > \dot{b},$$

will intersect the line $a = b$ with the condition $\dot{a} > \dot{b}$ preserved, for (6.5) and (6.12) will then be satisfied immediately after the intersection time.

First of all, note that $a > 0$ and $b > mnr_0^3$ are preserved as long as the solution exists. From Remark 4.7 we know that $a = r_0^2\beta t + O(t^3)$ and $b = mnr_0^3 + \frac{\sqrt{mn}(m+n)r_0}{2\beta}t^2 + O(t^4)$ as $t \rightarrow 0$. In order to compare (a, b) and (a_{ac}, b_{ac}) , we now reparametrise both solutions so that $a(s) = s = a_{ac}(s)$. Then

$$b = mnr_0^3 + \frac{\sqrt{mn}(m+n)}{2\beta^3r_0^3}s^2 + O(s^4), \quad b_{ac} = mnr_0^3 + \frac{\sqrt{mn}(m+n)}{2\beta_{ac}^3r_0^3}s^2 + O(s^4) \tag{7.16}$$

as $s \rightarrow 0$. In particular, if $\beta > \beta_{ac}$ we have

$$b < b_{ac}, \quad \dot{b} < \dot{b}_{ac}$$

for $s > 0$ sufficiently small. By Lemma 7.9(ii) these conditions are preserved as long as $b > \max(a, mnr_0^3)$.

Now, on the one hand the solution (a, b) certainly exists as long as $b > a > 0$, because then (a, b) is bounded to stay in the region $\{b_{ac} > b > a, b > mnr_0^3\}$ where (3.16) cannot blow up. On the other hand, since $b_{ac} \approx a_{ac} = s = a(s)$ for large s , for all $\epsilon > 0$ there exists s_0 such that $b_{ac} < a + \epsilon$ for all $s \geq s_0$. Moreover, as long as $b > \max(a, mnr_0^3)$, $b - b_{ac}$ is strictly decreasing and therefore bounded above by a definite constant $-\delta_0 < 0$ depending only on $\beta - \beta_{ac}$. Hence if $b > a$ for all $s \in [0, 2s_0]$ we would have

$$b - a - \epsilon < b - b_{ac} < -\delta_0 < 0$$

for all $s \in [s_0, 2s_0]$. If ϵ is chosen small enough we reach a contradiction. We conclude that (a, b) must intersect the boundary of the region $\{b_{ac} > b > a, b > mnr_0^3\}$. By Lemma 7.9(ii) the only possibility is that (a, b) intersects the line $a = b$. Moreover since we must have

$$\dot{b} \leq \dot{b}_{ac} < 1$$

up to and including the intersection time, at the intersection time we also have $\dot{b} < \dot{a}$.

An application of Propositions 6.6 and 6.11 now concludes the proof of Theorem 7.1(i).

7.3. Incompleteness results

We now conclude the proof of Theorem 7.1 establishing part (iii).

Let (a, b) be one of the local solutions constructed in Proposition 4.5(iii, iv) with $\beta < \beta_{ac}$. Assume for a contradiction that (a, b) yields a complete G_2 -metric. We want to show that the constraints (6.14), i.e.

$$0 < \frac{\dot{a}}{b} < \frac{a}{b} < 1,$$

are eventually satisfied, for Proposition 6.13 would otherwise give a contradiction.

We compare (a, b) and (a_{ac}, b_{ac}) : parametrise both solutions so that $a(s) = s = a_{ac}(s)$ and use (7.16) to conclude that since $\beta < \beta_{ac}$ we have

$$b > b_{ac}, \quad \dot{b} > \dot{b}_{ac}$$

by Lemma 7.9 (ii). Since $b_{ac} > a_{ac} = s$ we already conclude that $b > a = s$ for all time.

We will also need to know that b is unbounded and exists for all $s \geq 0$. For this note that $2\dot{b} = \sqrt{F}$ is the orbital volume function, which must be bounded below if (a, b) is complete. Hence b is unbounded. By (6.15), the condition $F > 0$ forces the ratio $a/b = s/b$ to be bounded below and therefore $s \rightarrow \infty$ as $b \rightarrow \infty$.

Now consider the quantities $R = s\dot{b} - b$ and $R_{ac} = s\dot{b}_{ac} - b_{ac}$ and, taking into account the parametrisation $a(s) = s$, note that $R > 0$ is equivalent to

$$\frac{\dot{a}}{b} < \frac{a}{b}.$$

By (7.16), R and R_{ac} are both initially strictly negative, but $R_{ac} \rightarrow 0$ along the AC end. We make two further crucial observations about R and R_{ac} :

- (i) $R - R_{ac}$ is strictly increasing: indeed, $\dot{R} - \dot{R}_{ac} = s(\ddot{b} - \ddot{b}_{ac})$ and $\ddot{b} - \ddot{b}_{ac} > 0$ for $b, b_{ac} > s$ is proved in Lemma 7.9;
- (ii) $R - R_{ac} > 0$ for small $s > 0$, by (7.16).

Hence by (i) and (ii), $R - R_{ac}$ has a strictly positive lower bound. Since R_{ac} converges to zero as $s \rightarrow \infty$, we conclude that R must eventually be positive.

Acknowledgments. LF would like to thank the Royal Society for the support of his research under a University Research Fellowship and NSF for the partial support of his work under grant DMS-1608143. MH and JN would like to thank the Simons Foundation for its support of their research under the Simons Collaboration on Special Holonomy in Geometry, Analysis and Physics (grants #488620, Mark Haskins, and #488631, Johannes Nordström).

All three authors would like to thank MSRI and the program organisers for providing an extremely productive environment during the Differential Geometry Program at MSRI in Spring 2016. Research at MSRI was partially supported by NSF grant DMS-1440140.

LF and MH would like to thank the Riemann Centre for Geometry and Physics at Leibniz Universität Hannover for supporting LF under a Riemann Research Fellowship and for hosting a research visit of MH in Autumn 2015. They would also like to thank the Newton Institute in Cambridge for hosting them during the Metric and Analytic Aspects of Moduli Spaces workshop in Summer 2015. Discussions about this work took place at both institutions.

MH would like to thank Christoph Böhm and Burkhard Wilking for very useful discussions during his visit to Münster in November 2015, that led to Proposition 6.1. MH would also like to thank Robert Bryant for several discussions about singular initial value problems for ODEs and PDEs, and for suggesting classical references that eventually led us to consult Picard's book [50].

The authors would also like to thank Harvey Tollick for help with producing Figure 1.

References

- [1] Acharya, B. S.: On realising $N = 1$ Super Yang–Mills in M theory. [hep-th/0011089](#) (2000)
- [2] Acharya, B. S., Gukov, S.: M theory and singularities of exceptional holonomy manifolds. *Phys. Rep.* **392**, 121–189 (2004) [Zbl 1162.53036](#) [MR 2039166](#)
- [3] Acharya, B. S., Witten, E.: Chiral fermions from manifolds of G_2 holonomy. [hep-th/0109152](#) (2001)
- [4] Alekseevsky, D., Dotti, I., Ferraris, C.: Homogeneous Ricci positive 5-manifolds, *Pacific J. Math.* **175**, 1–12 (1996) [Zbl 0865.53041](#) [MR 1419469](#)
- [5] Atiyah, M., Hitchin, N.: *The Geometry and Dynamics of Magnetic Monopoles*. Princeton Univ. Press, Princeton, NJ (1988) [Zbl 0671.53001](#) [MR 934202](#)
- [6] Atiyah, M., Maldacena, J., Vafa, C.: An M-theory flop as a large N duality. *J. Math. Phys.* **42**, 3209–3220 (2001) [Zbl 1061.81056](#) [MR 1840340](#)
- [7] Atiyah, M., Witten, E.: M -theory dynamics on a manifold of G_2 holonomy. *Adv. Theor. Math. Phys.* **6**, 1–106 (2002) [Zbl 1033.81065](#) [MR 1992874](#)
- [8] Bazaikin, Y. V.: On new examples of complete noncompact metrics with Spin(7) holonomy group. *Sibirsk. Mat. Zh.* **48**, 11–32 (2007) (in Russian) [Zbl 1164.53360](#) [MR 2304875](#)
- [9] Bazaikin, Y. V., Bogoyavlenskaya, O. A.: Complete Riemannian metrics with holonomy group G_2 on deformations of cones over $S^3 \times S^3$. *Math. Notes* **93**, 643–653 (2013); transl. from *Mat. Zametki* **93**, 645–657 (2013) [Zbl 1281.53052](#) [MR 3206014](#)

- [10] Belinskiĭ, V. A., Gibbons, G. W., Page, D. N., Pope, C. N.: Asymptotically Euclidean Bianchi IX metrics in quantum gravity. *Phys. Lett. B* **76**, 433–435 (1978) [MR 0496343](#)
- [11] Berglund, P., Brandhuber, A.: Matter from G_2 manifolds. *Nuclear Phys. B* **641**, 351–375 (2002) [Zbl 0998.81069](#) [MR 1928190](#)
- [12] Biquard, O., Minerbe, V.: A Kummer construction for gravitational instantons. *Comm. Math. Phys.* **308**, 773–794 (2011) [Zbl 1234.53020](#) [MR 2855540](#)
- [13] Bogoyavlenskaya, O. A.: On a new family of complete Riemannian metrics on $S^3 \times \mathbb{R}^4$ with holonomy group G_2 . *Sibirsk. Mat. Zh.* **54**, 551–562 (2013) (in Russian) [Zbl 1278.53046](#) [MR 3112613](#)
- [14] Böhm, C.: Inhomogeneous Einstein metrics on low-dimensional spheres and other low-dimensional spaces. *Invent. Math.* **134**, 145–176 (1998) [Zbl 0965.53033](#) [MR 1646591](#)
- [15] Böhm, C.: Non-compact cohomogeneity one Einstein manifold. *Bull. Soc. Math. France* **127**, 135–177 (1999) [Zbl 0935.53021](#) [MR 1700472](#)
- [16] Brandhuber, A.: G_2 holonomy spaces from invariant three-forms. *Nuclear Phys. B* **629**, 393–416 (2002) [Zbl 1039.83036](#) [MR 1903163](#)
- [17] Brandhuber, A., Gomis, J., Gubser, S. S., Gukov, S.: Gauge theory at large N and new G_2 holonomy metrics. *Nuclear Phys. B* **611**, 179–204 (2001) [Zbl 0971.81071](#) [MR 1857379](#)
- [18] Bryant, R. L., Salamon, S. M.: On the construction of some complete metrics with exceptional holonomy. *Duke Math. J.* **58**, 829–850 (1989) [Zbl 0681.53021](#) [MR 1016448](#)
- [19] Calabi, E.: Métriques kählériennes et fibrés holomorphes. *Ann. Sci. École Norm. Sup. (4)* **12**, 269–294 (1979) [Zbl 0431.53056](#) [MR 543218](#)
- [20] Candelas, P., de la Ossa, X. C.: Comments on conifolds. *Nuclear Phys. B* **342**, 246–268 (1990) [MR 1068113](#)
- [21] Carr, J.: *Applications of Centre Manifold Theory*. *Appl. Math. Sci.* 35, Springer, New York (1981) [Zbl 0464.58001](#) [MR 635782](#)
- [22] Cortés, V., Vázquez, J. J.: Locally homogeneous nearly Kähler manifolds. *Ann. Global Anal. Geom.* **48**, 269–294 (2015) [Zbl 1326.53104](#) [MR 3396468](#)
- [23] Cvetič, M., Gibbons, G. W., Lü, H., Pope, C. N.: A G_2 unification of the deformed and resolved conifolds. *Phys. Lett. B* **534**, 172–180 (2002) [Zbl 0994.83045](#) [MR 1913118](#)
- [24] Cvetič, M., Gibbons, G. W., Lü, H., Pope, C. N.: Cohomogeneity one manifolds of Spin(7) and G_2 holonomy. *Phys. Rev. D* **65**, art. 106004, 29 pp. (2002) [MR 1919035](#)
- [25] Cvetič, M., Gibbons, G. W., Lü, H., Pope, C. N.: M-theory conifolds. *Phys. Rev. Lett.* **88**, 121602 (2002) [MR 1901726](#)
- [26] Cvetič, M., Gibbons, G. W., Lü, H., Pope, C. N.: New complete noncompact Spin(7) manifolds. *Nuclear Phys. B* **620**, 29–54 (2002) [Zbl 1037.53032](#) [MR 1873145](#)
- [27] Cvetič, M., Gibbons, G. W., Lü, H., Pope, C. N.: New cohomogeneity one metrics with Spin(7) holonomy. *J. Geom. Phys.* **49**, 350–365 (2004) [Zbl 1092.53024](#) [MR 2076750](#)
- [28] Cvetič, M., Gibbons, G. W., Lü, H., Pope, C. N.: Orientifolds and slumps in G_2 and Spin(7) metrics. *Ann. Physics* **310**, 265–301 (2004) [MR 2044740](#)
- [29] Dancer, A., Wang, M. Y.: Kähler–Einstein metrics of cohomogeneity one. *Math. Ann.* **312**, 503–526 (1998) [Zbl 0965.53034](#) [MR 1654828](#)
- [30] Dancer, A., Wang, M. Y.: Painlevé expansions, cohomogeneity one metrics and exceptional holonomy. *Comm. Anal. Geom.* **12**, 887–926 (2004) [Zbl 1092.53039](#) [MR 2104080](#)
- [31] Eguchi, T., Hanson, A. J.: Self-dual solutions to Euclidean gravity. *Ann. Physics* **120**, 82–106 (1979) [Zbl 0409.53020](#) [MR 540896](#)
- [32] Eschenburg, J.-H., Wang, M. Y.: The initial value problem for cohomogeneity one Einstein metrics. *J. Geom. Anal.* **10**, 109–137 (2000) [Zbl 0992.53033](#) [MR 1758585](#)
- [33] Ferus, D., Karcher, H.: Nonrotational minimal spheres and minimizing cones. *Comment. Math. Helv.* **60**, 247–269 (1985) [Zbl 0566.53052](#) [MR 800005](#)

- [34] Foscolo, L., Haskins, M.: New G_2 -holonomy cones and exotic nearly Kähler structures on S^6 and $S^3 \times S^3$. *Ann. of Math. (2)* **185**, 59–130 (2017) [Zbl 1381.53086](#) [MR 3583352](#)
- [35] Foscolo, L., Haskins, M., Nordström, J.: Complete non-compact G_2 -manifolds from asymptotically conical Calabi–Yau 3-folds. [arXiv:1709.04904](#) (2017)
- [36] Gibbons, G. W., Hawking, S. W.: Gravitational multi-instantons. *Phys. Lett. B* **78**, 430–432 (1978)
- [37] Gibbons, G. W., Page, D. N., Pope, C. N.: Einstein metrics on S^3 , \mathbf{R}^3 and \mathbf{R}^4 bundles. *Comm. Math. Phys.* **127**, 529–553 (1990) [Zbl 0699.53053](#) [MR 1040893](#)
- [38] Hawking, S. W.: Gravitational instantons. *Phys. Lett. A* **60**, 81–83 (1977) [MR 0465052](#)
- [39] Hitchin, N.: Stable forms and special metrics. In: *Global Differential Geometry: the Mathematical Legacy of Alfred Gray* (Bilbao, 2000), *Contemp. Math.* 288, Amer. Math. Soc., Providence, RI, 70–89 (2001) [Zbl 004.53034](#) [MR 1871001](#)
- [40] Hori, K., Hosomichi, K., Page, D. C., Rabadán, R., Walcher, J.: Non-perturbative orientifold transitions at the conifold. *J. High Energy Phys.* **2005**, art. 026, 61 pp. [MR 2181684](#)
- [41] Joyce, D. D.: *Compact Manifolds with Special Holonomy*. Oxford Math. Monogr., Oxford Univ. Press, Oxford (2000) [Zbl 1027.53052](#) [MR 1787733](#)
- [42] Karigiannis, S.: Desingularization of G_2 manifolds with isolated conical singularities. *Geom. Topol.* **13**, 1583–1655 (2009) [Zbl 1161.53348](#) [MR 2496053](#)
- [43] Karigiannis, S., Lotay, J.: Deformation theory of G_2 conifolds. *Comm. Anal. Geom.* **28**, 1057–1210 (2020). [Zbl 07268919](#) [MR 4165315](#)
- [44] Kronheimer, P. B.: The construction of ALE spaces as hyper-Kähler quotients. *J. Differential Geom.* **29**, 665–683 (1989) [Zbl 0671.53045](#) [MR 992334](#)
- [45] Madsen, T. B., Salamon, S.: Half-flat structures on $S^3 \times S^3$. *Ann. Global Anal. Geom.* **44**, 369–390 (2013) [Zbl 1285.53038](#) [MR 3132080](#)
- [46] Malgrange, B.: Sur les points singuliers des équations différentielles. *Enseign. Math. (2)* **20**, 147–176 (1974) [Zbl 0299.34011](#) [MR 0368074](#)
- [47] Nordström, J.: Deformations and gluing of asymptotically cylindrical manifolds with exceptional holonomy. http://people.bath.ac.uk/jlpn20/thesis_final_twoside.pdf (2008)
- [48] Palais, R. S.: The principle of symmetric criticality. *Comm. Math. Phys.* **69**, 19–30 (1979) [Zbl 0417.58007](#) [MR 547524](#)
- [49] Pando Zayas, L. A., Tseytlin, A. A.: 3-branes on spaces with $\mathbf{R} \times S^2 \times S^3$ topology. *Phys. Rev. D (3)* **63**, art. 086006, 11 pp. (2001) [MR 1831169](#)
- [50] Picard, E.: *Traité d’analyse. Tome III. Les Grands Classiques Gauthier-Villars, Éditions Jacques Gabay, Sceaux* (1991) [MR 1188644](#)
- [51] Schulte-Hengesbach, F.: Half-flat structures on Lie groups. Ph.D. thesis, Hamburg, <http://www.math.uni-hamburg.de/home/schulte-hengesbach/diss.pdf> (2010)
- [52] Sparks, J.: Sasaki–Einstein manifolds. In: *Geometry of Special Holonomy and Related Topics*, *Surv. Differ. Geom.* 16, Int. Press, Somerville, MA, 265–324 (2011) [Zbl 1256.53037](#) [MR 2893680](#)
- [53] Stenzel, M. B.: Ricci-flat metrics on the complexification of a compact rank one symmetric space. *Manuscripta Math.* **80**, 151–163 (1993) [Zbl 0811.53049](#) [MR 1233478](#)
- [54] Wang, M. Y., Ziller, W.: Einstein metrics on principal torus bundles. *J. Differential Geom.* **31**, 215–248 (1990) [Zbl 0691.53036](#) [MR 1030671](#)
- [55] Witten, E.: Anomaly cancellation on G_2 manifolds. [hep-th/0108165](#) (2001)