



Saugata Basu · Deepam Patel

VC density of definable families over valued fields

Received May 4, 2018

Abstract. We prove a tight bound on the number of realized 0/1 patterns (or equivalently on the Vapnik–Chervonenkis codensity) of definable families in models of the theory of algebraically closed valued fields with a non-archimedean valuation. Our result improves the best known result in this direction proved by Aschenbrenner, Dolich, Haskell, Macpherson and Starchenko, who proved a weaker bound in the restricted case where the characteristics of the field K and its residue field are both assumed to be 0. The bound obtained here is optimal and without any restriction on the characteristics.

We obtain the aforementioned bound as a consequence of another result on bounding the Betti numbers of semi-algebraic subsets of certain Berkovich analytic spaces, mirroring similar results known already in the case of o-minimal structure and for real closed as well as algebraically closed fields. The latter result is the first result in this direction and is possibly of independent interest. Its proof relies heavily on recent results of Hrushovski and Loeser on the topology of semi-algebraic subsets of Berkovich analytic spaces.

Keywords. VC-density, Berkovich space, algebraically closed valued field, o-minimal, Betti numbers

Contents

1. Introduction	2362
1.1. Combinatorial definitions	2362
1.2. Brief history	2363
2. Main results	2365
2.1. Model theory of algebraically closed valued fields	2365
2.2. New results	2366
3. Proofs of the main results	2367
3.1. Methods used	2367
3.2. Proof of Theorem 2	2369
3.3. Proof of Theorem 1	2393
3.4. Proof of Corollary 1	2396

S. Basu: Department of Mathematics, Purdue University, 150 N. University Street, West Lafayette, IN 47907, U.S.A.; e-mail: sbasu@math.purdue.edu

D. Patel: Department of Mathematics, Purdue University, 150 N. University Street, West Lafayette, IN 47907, U.S.A.; e-mail: patel471@purdue.edu

Mathematics Subject Classification (2020): Primary 03C98; Secondary 14F45

Appendix 2396
 A.1. Review of singular cohomology 2397
 A.2. Recollections from Hrushovski–Loeser 2398
 References 2402

1. Introduction

In this article, we prove a tight bound on the number of realized 0/1 patterns (or equivalently on the Vapnik–Chervonenkis codensity) of definable families in models of the theory of algebraically closed valued fields with a non-archimedean valuation (henceforth referred to just as ACVF). This result improves on the best known upper bound on this quantity previously obtained by Aschenbrenner et al. [AD⁺16]. Our result is a consequence of a topological result giving an upper bound on the Betti numbers of certain semi-algebraic sets obtained as Berkovich analytifications of definable sets in certain models of ACVF which we will recall more precisely in the next section.

In order to state our main combinatorial result we need to introduce some preliminary notation and definitions.

1.1. Combinatorial definitions

Suppose V and W are sets, and $X \subset V \times W$ is a subset. Let $\pi_V : X \rightarrow V, \pi_W : X \rightarrow W$ denote the restrictions to X of the natural projection maps. For any $v \in V, w \in W$, we set $X_v := \pi_W(\pi_V^{-1}(v))$, and $X_w := \pi_V(\pi_W^{-1}(w))$.

Notation 1.1.1. For each $n > 0$, we define a function

$$\chi_{X,V,W;n} : V \times W^n \rightarrow \{0, 1\}^n$$

as follows. For $\bar{w} := (w_1, \dots, w_n) \in W^n$ and $v \in V$, we set

$$(\chi_{X,V,W;n}(v, \bar{w}))_i := \begin{cases} 0 & \text{if } v \notin X_{w_i}, \\ 1 & \text{otherwise.} \end{cases}$$

(Note that in the special case when $n = 1, \chi_{X,V,W;1}$ is just the usual characteristic function of the subset $X \subset V \times W$.)

For $\bar{w} \in W^n$ and $\sigma \in \{0, 1\}^n$, we will say that σ is *realized by the tuple* $(X_{w_1}, \dots, X_{w_n})$ *of subsets of* V *if there exists* $v \in V$ *such that* $\chi_{X,V,W;n}(v, \bar{w}) = \sigma$. We will often refer to elements of $\{0, 1\}^n$ colloquially as ‘0/1 patterns’.

Finally, we define the function

$$\chi_{X,V,W} : \mathbb{N} \rightarrow \mathbb{N} \quad \text{by} \quad \chi_{X,V,W}(n) := \max_{\bar{w} \in W^n} \text{card}(\chi_{X,V,W;n}(V, \bar{w})).$$

The function $\chi_{X,V,W}$ is closely related to the notion of VC-codensity of a set system. Since some of the prior results (for example, those in [AD⁺16]) have been stated in terms of VC-codensity it is useful to recall its definition here.

Definition 1.1.2. Let X be a set and $\mathcal{S} \subset 2^X$. The *shatter function* of \mathcal{S} , $\pi_{\mathcal{S}} : \mathbb{N} \rightarrow \mathbb{N}$, is defined by setting

$$\pi_{\mathcal{S}}(n) := \max_{A \subset X, \text{card}(A)=n} \text{card}(\{A \cap Y \mid Y \in \mathcal{S}\}).$$

We denote

$$\text{vcd}_{\mathcal{S}} := \limsup_{n \rightarrow \infty} \frac{\log(\pi_{\mathcal{S}}(n))}{\log(n)}.$$

Given a definable subset $X \subset V \times W$ in some structure, we will denote

$$\text{vcd}(X, V, W) := \text{vcd}_{\mathcal{S}},$$

where $\mathcal{S} = \{X_v \mid v \in V\} \subset 2^W$. We will call (following the convention in [AD⁺16]), $\text{vcd}(X, V, W)$, the *VC-codensity* of the family of subsets, $\{X_w \mid w \in W\}$, of V . More generally, if $\phi(\bar{X}, \bar{Y})$ is a first-order formula (with parameters) in the theory of some structure M , we set

$$\text{vcd}(\phi) := \text{vcd}(S, M^{|\bar{X}|}, M^{|\bar{Y}|}),$$

where $S \subset M^{|\bar{X}|} \times M^{|\bar{Y}|}$ is the set defined by ϕ . (Here and elsewhere in the paper, $|\bar{X}|$ denotes the length of the finite tuple of variables \bar{X} .) Note also that if M is an NIP structure (see for example [Sim15, Chapter 2] for definition), then $\text{vcd}(\phi) < \infty$ for every (parted) formula ϕ .

The problem of proving upper bounds on $\text{vcd}(X, V, W)$ of a definable family can be reduced to proving upper bounds on the function $\chi_{X, V, W}$ (see Proposition 3.4.1 below). We will henceforth concentrate on the problem of obtaining tight upper bounds on the function $\chi_{X, V, W}$.

1.2. Brief history

For definable families of hypersurfaces in \mathbb{F}^k of fixed degree over a field \mathbb{F} , Babai, Rónyai, and Ganapathy [RBG01] gave an elegant argument using linear algebra to show that the number of 0/1 patterns (see Notation 1.1.1) realized by n such hypersurfaces in \mathbb{F}^k is bounded by $C \cdot n^k$, where C is a constant that depends on the family (but independent of n). This bound is easily seen to be optimal. A more refined topological estimate on these realized 0/1 patterns (in terms of the sums of the Betti numbers) is given in [BPR09], where the methods are more in line with the methods in the current paper.

A similar result was proved in [BPR05] for definable families of semi-algebraic sets in \mathbb{R}^k , where \mathbb{R} is an arbitrary real closed field. For definable families in M^k , where M is an arbitrary o-minimal expansion of a real closed field, the first author [Bas10] adapted the methods of [BPR05] to prove a bound of $C \cdot n^k$ on the number of 0/1 patterns for such families where C is a constant that depends on the family (see also [JL10]). These bounds were obtained as a consequence of more general results bounding the individual Betti numbers of definable sets defined in terms of the members of the family, and more sophisticated homological techniques (as opposed to just linear algebra) played an important role in obtaining these bounds.

If K is an algebraically closed valued field, then the problem of obtaining tight bounds on $\text{vcd}(\phi)$ for parted formulas, $\phi(\bar{X}, \bar{Y})$, in the one-sorted language of valued fields with parameters in K was considered by Aschenbrenner et al. [AD⁺16]. They obtained the non-trivial bound of $2|\bar{X}|$ on $\text{vcd}(\phi)$ in the case when the characteristic pair of K (consisting of the characteristic of the field K and that of its residue field) is $(0, 0)$ [AD⁺16, Corollary 6.3]. In terms of 0/1 patterns (Proposition 3.4.1) their result can be restated as saying that for each $k > 0$ and any fixed definable family of subsets of K^k , there exists $C > 0$ (depending on the family) such that for all $n > 0$ the number of 0/1 patterns realized by any n sets of the family is bounded from above by $C \cdot n^{2k}$.

Given that the model-theoretic/algebraic techniques used thus far do not immediately yield the tight upper bound of $|\bar{X}|$ on $\text{vcd}(\phi(\bar{X}, \bar{Y}))$ for valued fields, it is natural to consider a more topological approach as in [Bas10]. However, for definable families over a (complete) valued field, it is not a priori clear that there exists an appropriate well-behaved cohomology theory (i.e. with the required finiteness/cohomological dimension properties) that makes the approach in [Bas10] feasible in this situation. For example, ordinary sheaf cohomology with respect to the Zariski or étale site for schemes is clearly unsuitable. Fortunately, the recent break-through results of Hrushovski and Loeser [HL16] give us an opening in this direction. Instead of considering the original definable subset of an affine variety V defined over K , we can consider the corresponding *semi-algebraic* subset of the Berkovich analytification $B_{\mathbf{F}}(V)$ of V (see §A.2 below for the definitions). These semi-algebraic subsets have certain key topological tameness properties which are analogous to those used in the case of o-minimal structures, and moreover crucially they are homotopy equivalent to a simplicial complex of dimension at most $\dim(V)$. Therefore, their cohomological dimension is at most $\dim(V)$. In particular, the singular cohomology of the underlying topological spaces satisfies the required properties. Thus, in order to bound the number of realizable 0/1 patterns of a finite set of definable subsets of V , we can first replace the latter set by the corresponding semi-algebraic subsets of $B_{\mathbf{F}}(V)$, and then try to make use of their tame topological properties to obtain a bound on the number of 0/1 patterns realized by these subsets. An upper bound on the latter quantity will also be an upper bound on the number of 0/1 patterns realized by the definable subsets we started with (this fact is elucidated later in Observation 3.3.1).

Using the results of Hrushovski and Loeser, one can then hope to proceed with the o-minimal case as the guiding principle. While the arguments are somewhat similar in spirit, there are several technical challenges that need to be overcome—for example, an appropriate definition of “tubular neighborhoods” with the required properties (see §3.1 below for a more detailed description of these challenges). The bounds on the sum of the Betti numbers of the semi-algebraic subsets of Berkovich spaces that we obtain in this way are exactly analogous to the ones in the algebraic, semi-algebraic, as well as o-minimal cases. The fact that the cohomological dimension of the semi-algebraic subsets of $B_{\mathbf{F}}(V)$ is bounded by $\dim(V)$ is one key ingredient in obtaining these tight bounds.

Our results on bounding the Betti numbers of semi-algebraic subsets of Berkovich spaces are of independent interest, and they seem to suggest a more general formalism of cohomology associated to NIP structures. For example, one obtains bounds (on the Betti numbers) of the exact same shape and having the same exponents for definable families in

the case of algebraic, semi-algebraic, o-minimal and valued field structures. Moreover, in each of these cases, these bounds are obtained as a consequence of general bounds on the dimension of certain cohomology groups. Therefore, it is perhaps reasonable to hope for some general cohomology theory (say for NIP structures which are fields) which would in turn give a uniform method of obtaining tight bounds on VC-density via cohomological methods. More generally, it shows that cohomological methods can play an important role in model theory in general.

As a consequence of the bound on the Betti numbers (in fact using the bound only on the 0-th Betti number) we prove that $\text{vcd}(\phi(\bar{X}, \bar{Y}))$ over an arbitrary algebraically closed valued field is bounded by $|\bar{X}|$. One consequence of our methods (unlike the techniques used in [AD⁺16]) is that there are no restrictions on the characteristic pair of the valued field K .

Finally, note that in [AD⁺16] the authors also obtain a bound of $2|\bar{X}| - 1$ on $\text{vcd}(\phi(\bar{X}, \bar{Y}))$, over \mathbb{Q}_p , where ϕ is a formula in Macintyre’s language [Mac76]. However, our methods right now do not yield results in this case.

Outline of the paper. In §2 we first introduce the necessary technical background (in §2.1), and then state the main results of the paper, namely Theorems 1 and 2, and Corollary 1 (in §2.2). The proofs of the main results appear in §3. We first give an outline of the proofs in §3.1. We next prove the main topological result of the paper (Theorem 2) in in §3.2, and prove Theorem 1 and Corollary 1 in §3.3 and §3.4 respectively.

In order to make the paper self-contained and for the benefit of the readers, we include in an appendix a review of some very classical results about singular cohomology (in §A.1), as well as much more recent ones related to semi-algebraic sets associated to definable sets in models of ACVF proved by Hrushovski and Loeser [HL16] (in §A.2). These results are used heavily in the proofs of the main theorems.

2. Main results

2.1. Model theory of algebraically closed valued fields

In this section, K will always denote an algebraically closed non-archimedean valued field, and the value group of K will be denoted by Γ . Let $R := K[X_1, \dots, X_N]$ and $\mathbb{A}_K^N = \text{Spec}(R)$. Given a closed affine subvariety $V = \text{Spec}(A)$ of $\mathbb{A}_K^N = \text{Spec}(R)$ and an extension K' of K , we will denote by $V(K') \subset \mathbb{A}_K^N(K')$ the set of K' points of V .

We denote by \mathcal{L} the two-sorted language

$$(0_K, 1_K, +_K, \times_K, |\cdot| : K \rightarrow \Gamma \cup \{0_\Gamma\}, \leq_\Gamma, \times_\Gamma),$$

where the subscript K denotes constants, functions, relations etc. of the field sort and the subscript Γ denotes the same for the value group sort. When the context is clear we will drop the subscripts. The constant 0_Γ is interpreted as the valuation of 0 (and does not technically belong to the value group).

Now suppose that $\phi(X_1, \dots, X_n)$ is a quantifier-free formula with parameters in $(K; \Gamma \cup \{0_\Gamma\})$ in the language \mathcal{L} with free variables only of the field sort. Then ϕ is

a quantifier-free formula with atoms of the form $|F| \leq \lambda \cdot |G|$ where $F, G \in R$ and $\lambda \in \Gamma \cup \{0_\Gamma\}$. The formula ϕ gives rise to a definable subset of \mathbb{A}_K^N , and in particular ϕ defines a subset of $\mathbb{A}_K^N(K')$ for every valued extension K' of K . We will denote the intersection of this subset with V by $\mathcal{R}(\phi, V)$, and by $\mathcal{R}(\phi, V)(K')$ the corresponding subset of $V(K')$.

Let ϕ be a formula with parameters in $(K; \Gamma \cup \{0_\Gamma\})$ in the language \mathcal{L} with free variables only of the field sort. Note that every such formula is equivalent modulo the two-sorted theory of $(K; \Gamma \cup \{0_\Gamma\})$ to a quantifier-free formula (see for example [HHM08, Theorem 7.1(ii)]). Because of this fact, we can assume without loss of generality in what follows that ϕ is a quantifier-free formula, and is thus a quantifier-free formula with atoms of the form $|F| \leq \lambda \cdot |G|$ where $F, G \in R$ and $\lambda \in \Gamma \cup \{0_\Gamma\}$.

2.2. New results

Our main result is the following.

Theorem 1 (Bound on the number of 0/1 patterns). *Let K be an algebraically closed valued field with value group Γ . Suppose that $V \subset \mathbb{A}_K^N$ and $W \subset \mathbb{A}_K^M$ are closed affine subvarieties and let*

$$\phi(X_1, \dots, X_N; Y_1, \dots, Y_M)$$

be a formula with parameters in $(K; \Gamma \cup \{0_\Gamma\})$ in the language \mathcal{L} (with free variables only of the field sort). Then there exists a constant $C = C_{\phi, V, W}$ such that for all $n > 0$,

$$\chi_{\mathcal{R}(\phi, (V \times W)(K), V(K), W(K))}(n) \leq C \cdot n^k, \quad \text{where } k = \dim V.$$

As an immediate corollary of Theorem 1 we obtain the following bound on the VC-codensity for definable families over algebraically closed valued fields.

Corollary 1 (Bound on the VC-codensity for definable families over ACVF). *Let K be an algebraically closed valued field with value group Γ . Let $\phi(\overline{X}, \overline{Y})$ be a formula with parameters in $(K; \Gamma \cup \{0_\Gamma\})$ in the language \mathcal{L} . Then*

$$\text{vcd}(\phi) \leq |\overline{X}|.$$

Theorem 1 will follow from a more general topological theorem which we will now state. First, we recall some more notation.

We assume that K is an algebraically closed complete valued field with a non-archimedean valuation whose value group Γ is a subgroup of the multiplicative group $\mathbb{R}_{>0}$.

Given an affine variety V as before, Hrushovski–Loeser [HL16] associate to V a locally compact Hausdorff topological space, denoted by $B_{\mathbf{F}}(V)$. More generally, they associate a locally compact Hausdorff topological space $B_{\mathbf{F}}(X)$ to any definable subset $X \subset V$ which is functorial in definable maps. In the present setting, $B_{\mathbf{F}}(V)$ can be identified with the Berkovich analytic space associated to V and has an explicit description in terms of valuations. We refer the reader to §A.2 for a brief review of this construction and its main properties.

Notation 2.2.1. If $V \subset \mathbb{A}_K^N$ is an affine closed subvariety, and ϕ a formula in the language with parameters in $(K; \Gamma \cup \{0_\Gamma\})$ in the language \mathcal{L} with free variables only of the field sort, we will denote by $\tilde{\mathcal{R}}(\phi, V)$ the *semi-algebraic* subset $B_{\mathbf{F}}(\tilde{\mathcal{R}}(\phi, V))$ of $B_{\mathbf{F}}(V)$.

Suppose now that $V \subset \mathbb{A}_K^N$ and $W \subset \mathbb{A}_K^M$ are closed affine subvarieties and let $\phi(\cdot; \cdot)$ be a formula in disjunctive normal form without negations and with atoms of the form $|F| \leq \lambda \cdot |G|$, $F, G \in K[X_1, \dots, X_N, Y_1, \dots, Y_M]$, $\lambda \in \Gamma \cup \{0_\Gamma\}$. Then for each $w \in W(K)$, $\tilde{\mathcal{R}}(\phi(\cdot, w), V)$ is a semi-algebraic subset of $B_{\mathbf{F}}(V)$.

For $\bar{w} = (w_1, \dots, w_n) \in W(K)^n$ and $\sigma \in \{0, 1\}^n$, we set

$$(2.2.2) \quad \tilde{\mathcal{R}}(\sigma, \bar{w}) := \tilde{\mathcal{R}}(\phi_\sigma(\bar{w}), V),$$

where

$$\phi_\sigma(\bar{w}) := \bigwedge_{i, \sigma(i)=1} \phi(\cdot, w_i) \wedge \bigwedge_{i, \sigma(i)=0} \neg\phi(\cdot, w_i).$$

Given a topological space Z , we denote by $H^i(Z)$ the corresponding i -th singular cohomology group of X with rational coefficients. We refer the reader to §A.1 for a brief recollection of the main properties of these groups. We note that for $Z = \tilde{\mathcal{R}}(\sigma, \bar{w})$ these groups are finite-dimensional \mathbb{Q} -vector spaces. Let

$$b_i(\tilde{\mathcal{R}}(\sigma, \bar{w})) = \dim_{\mathbb{Q}} H^i(\tilde{\mathcal{R}}(\sigma, \bar{w}))$$

denote the corresponding i -th Betti number.

The following theorem, mirroring a similar theorem in the o-minimal case [Bas10], is the main technical result of this paper.

Theorem 2 (Bound on the Betti numbers). *Let K be an algebraically closed complete valued field with a non-archimedean valuation whose value group Γ is a subgroup of the multiplicative group $\mathbb{R}_{>0}$. Suppose that $V \subset \mathbb{A}_K^N$ and $W \subset \mathbb{A}_K^M$ are closed affine subvarieties and let $\phi(\cdot; \cdot)$ be a formula in disjunctive normal form without negations and with atoms of the form $|F| \leq \lambda \cdot |G|$, $F, G \in K[X_1, \dots, X_N, Y_1, \dots, Y_M]$, $\lambda \in \Gamma \cup \{0_\Gamma\}$. Let $\dim(V) = k$. Then there exists a constant $C = C_{\phi, V, W} > 0$ such that for all $\bar{w} \in W(K)^n$ and $0 \leq i \leq k$,*

$$\sum_{\sigma \in \{0, 1\}^n} b_i(\tilde{\mathcal{R}}(\sigma, \bar{w})) \leq C \cdot n^{k-i}.$$

3. Proofs of the main results

In this section we prove our main results. Before starting the formal proof we first give a brief outline of our methods.

3.1. Methods used

Our main technical result, Theorem 2, gives a bound, for each $i, 0 \leq i \leq k$, and $\bar{w} \in W(K)^n$, on the sum over $\sigma \in \{0, 1\}^n$ of the i -th Betti numbers of $\tilde{\mathcal{R}}(\sigma, \bar{w})$. The technique

for achieving this is an adaptation of the topological methods used to prove a similar result in the o-minimal category in [Bas10, Theorem 2.1]. We recall here the main steps of the proof in [Bas10].

We assume that $V = \mathbb{R}^N$, $W = \mathbb{R}^M$, where \mathbb{R} is a real closed field and $X \subset V \times W$ is a closed definable subset in an o-minimal expansion of \mathbb{R} .

- Step 1. One constructs definable infinitesimal tubes around the fibers X_{w_1}, \dots, X_{w_n} .
 Step 2. Let $\sigma \in \{0, 1\}^n$, and C be a connected component of

$$\bigcap_{\sigma(i)=1} X_{w_i} \cap \bigcap_{\sigma(i)=0} (V \setminus X_{w_i}).$$

One proves that there exists a unique connected component D of the complement of the boundaries of the tubes constructed in Step 1 such that C is homotopy equivalent to D . The homotopy equivalence is proved using the local conical structure theorem for o-minimal structures.

- Step 3. As a consequence of Step 2, in order to bound $\sum_{\sigma} b_i(\mathbb{R}(\sigma, \bar{w}))$, it suffices (using Alexander duality) to bound the Betti numbers of the union of the boundaries of the tubes constructed in Step 1.
 Step 4. Bounding those Betti numbers is achieved using certain inequalities which follow from the Mayer–Vietoris exact sequence (see Properties A.1.1(5)). In these inequalities only the Betti numbers of at most k -ary intersections of the boundaries play a role.
 Step 5. One then uses Hardt’s triviality theorem for o-minimal structures to get a uniform bound on each of these Betti numbers that depends only on the definable family under consideration, i.e. on X , V , and W . Thus, the only part of the bound that grows with n comes from certain binomial coefficients counting the number of different possible intersections one needs to consider.

The method we use to prove Theorem 2 is close in spirit to that in [Bas10, proof of Theorem 2.1] as outlined above but different in many important details. For each of the steps above we give the corresponding step in the proof of Theorem 2.

- Step 1'. We again construct certain tubes around the fibers and give explicit descriptions of the tubes in terms of the formula ϕ defining the given semi-algebraic set $\tilde{\mathcal{R}}(\sigma, \bar{w})$. The definition of these tubes is somewhat more complicated than in the o-minimal case (see Notation 3.2.2). The use of two different infinitesimals to define these tubes is necessitated by the singular behavior of the semi-algebraic set defined by $|F| \leq \lambda \cdot |G|$ near the common zeros of F and G .
 Step 2'. The homotopy equivalence property analogous to Step 2 above is proved in Proposition 3.2.6, and the role of the local conical structure theorem in the o-minimal case is now played by a corresponding result of Hrushovski and Loeser (see Theorem A.3).
 Step 3'. We avoid the use of Alexander duality by directly using a Mayer–Vietoris type inequality giving a bound on the Betti numbers of intersections of open sets in terms of the Betti numbers of up to k -fold unions (see Proposition 3.2.47).

Step 4'. This step is subsumed by Step 3'.

Step 5'. Finally, instead of using Hardt's triviality to obtain a constant bound on the Betti numbers of these 'small' unions, we use a theorem of Hrushovski and Loeser which states that the number of homotopy types amongst the fibers of any fixed map in the analytic category that we consider is finite (see Theorem A.4).

We apply Theorem 2 directly to obtain the VC-codensity bound in the case of the theory of ACVF (using Observation 3.3.1). One extra subtlety here is in removing the assumption on the formula ϕ (which occurs in the hypothesis of Theorem 2). Actually, in order to prove Corollary 1 in general it suffices only to consider ϕ of the special form having just one atom of the form $|F| \leq \lambda \cdot |G|$ or $|F| = \lambda \cdot |G|$. This reduction from the general case to the special case is encapsulated in a combinatorial result (Proposition 3.3.2). With the help of Proposition 3.3.2, Corollary 1 becomes a consequence of Theorem 2 and Observation 3.3.1.

We now give the proofs in full detail. In the next subsection (§3.2) we give the proof of Theorem 2. In §3.3, we show how to deduce Theorem 1 from Theorem 2. Finally, in §3.4 we show how to deduce Corollary 1 from Theorem 2.

3.2. Proof of Theorem 2

In the following, K will be a fixed algebraically closed non-archimedean (complete real-valued) field and V is an affine variety over K . We shall freely use the results of Hrushovski and Loeser [HL16] on the spaces $B_{\mathbf{F}}(X)$ associated to definable subsets $X \subset V$. For the reader's convenience, an exposition (with references) of the results we require below is provided in §A.2. We shall also make use of some standard facts about singular cohomology of topological spaces; we refer the reader to §A.1 for a review of these facts.

Notation 3.2.1 (Closed cube). For $R \in \mathbb{R}$, $R > 0$, and $N > 0$, we denote by $\text{Cube}_N(R)$ the semi-algebraic subset $\tilde{\mathcal{R}}(\psi, \mathbb{A}_K^N)$, where

$$\psi = \bigwedge_{1 \leq i \leq N} |X_i| \leq R,$$

and $\mathbb{A}_K^N = \text{Spec}(K[X_1, \dots, X_N])$ is the usual affine space. Notice that $\text{Cube}_N(R)$ is a closed topological space since the $|X_i|$ are continuous functions (see A.2.2(4, 5)). Moreover, it is compact (see A.2.2(6)). If $V = \text{Spec}(A) \subset \mathbb{A}_K^N$ is a closed subvariety, then we set $\text{Cube}_V(R) := \text{Cube}_N(R) \cap B_{\mathbf{F}}(V)$. Note that this a closed semi-algebraic subset of $B_{\mathbf{F}}(V)$.

Notation 3.2.2 (Open, closed $(\varepsilon, \varepsilon')$ -tubes). Suppose $\phi(\cdot)$ is a formula in disjunctive normal form without negations and with atoms of the form $|F| \leq \lambda \cdot |G|$ with $F, G \in K[X_1, \dots, X_N]$ and $\lambda \in \mathbb{R}_+ := \mathbb{R}_{\geq 0}$. We denote by

$$\phi^o(\cdot; T, T')$$

the formula obtained from ϕ by replacing each atom $|F| \leq \lambda \cdot |G|$ with $\lambda, G \neq 0$ by the formula

$$(|F| < (\lambda \cdot T) \cdot |G|) \vee ((|F| < T') \wedge (|G| < T')),$$

and each atom $|F| \leq \lambda \cdot |G|$ with $\lambda = 0$ or $G = 0$ by the formula

$$|F| < T',$$

where T, T' are new variables of the value sort. Similarly, we denote by

$$\phi^c(\cdot; T, T')$$

the formula obtained from ϕ by replacing each atom $|F| \leq \lambda \cdot |G|$ by the formula

$$(|F| \leq (\lambda \cdot T) \cdot |G|) \vee ((|F| \leq T') \wedge (|G| \leq T'))$$

if $\lambda, G \neq 0$, and by the formula

$$|F| \leq T'$$

if $\lambda = 0$ or $G = 0$. Here again T, T' are new variables of the value sort.

For $\varepsilon > 1, \varepsilon' > 0$, and V a closed subvariety of \mathbb{A}_K^N we set

$$\text{Tube}_{V,\phi}^o(\varepsilon, \varepsilon') := \tilde{\mathcal{R}}(\phi^o(\cdot; \varepsilon, \varepsilon'), V), \quad \text{Tube}_{V,\phi}^c(\varepsilon, \varepsilon') := \tilde{\mathcal{R}}(\phi^c(\cdot; \varepsilon, \varepsilon'), V).$$

For each $R > 0$, we set

$$(3.2.3) \quad \text{Tube}_{V,\phi}^o(\varepsilon, \varepsilon', R) := \text{Cube}_V(R) \cap \text{Tube}_{V,\phi}^o(\varepsilon, \varepsilon'),$$

$$(3.2.4) \quad \text{Tube}_{V,\phi}^c(\varepsilon, \varepsilon', R) := \text{Cube}_V(R) \cap \text{Tube}_{V,\phi}^c(\varepsilon, \varepsilon').$$

We set

$$\text{TubeCompl}_{V,\phi}^c(\varepsilon, \varepsilon', R) := \text{Cube}_V(R) \setminus \text{Tube}_{V,\phi}^o(\varepsilon, \varepsilon', R).$$

Notice that by definition, $\text{Tube}_{V,\phi}^o(\varepsilon, \varepsilon', R)$ (resp. $\text{TubeCompl}_{V,\phi}^c(\varepsilon, \varepsilon', R)$) is an open (resp. closed) subset of $\text{Cube}_V(R)$. Moreover, both are semi-algebraic as subsets of $B_{\mathbf{F}}(V)$.

Finally, we set

$$\text{TubeBoundary}_{V,\phi}^c(\varepsilon, \varepsilon', R) := \text{Tube}_{V,\phi}^c(\varepsilon, \varepsilon', R) \cap \text{TubeCompl}_{V,\phi}^c(\varepsilon, \varepsilon', R).$$

Remark 3.2.5. Note that in our notation for the ‘tubes’ above, a superscript o (resp. c) indicates that the corresponding tube is open (resp. closed).

The next proposition is the key ingredient for the proof of Theorem 2.

Proposition 3.2.6. *Let $V \subset \mathbb{A}_K^N$ and $W \subset \mathbb{A}_K^M$ be closed affine subvarieties. Let $\phi(\cdot, \cdot)$ be a formula in disjunctive normal form without negations and with atoms of the form $|F| \leq \lambda \cdot |G|$ where $F, G \in K[X_1, \dots, X_N, Y_1, \dots, Y_M]$. For each $\bar{w} \in W(K)^n$, $\sigma \in \{0, 1\}^n$, and for all sufficiently large $R > 0$ and $\delta, \delta', \varepsilon, \varepsilon' \in \mathbb{R}_+$ satisfying $0 < \delta - 1 \ll \delta' \ll \varepsilon - 1 \ll \varepsilon' \ll 1$,*

$$H^*(\tilde{\mathcal{R}}(\sigma, \bar{w})) \cong H^*(S_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)),$$

where

$$S_\sigma(\delta, \delta', \varepsilon, \varepsilon', R) := \bigcap_{i, \sigma(i)=1} \text{Tube}_{V, \phi(\cdot, w_i)}^o(\delta, \delta', R) \cap \bigcap_{i, \sigma(i)=0} \text{TubeCompl}_{V, \phi(\cdot, w_i)}^c(\varepsilon, \varepsilon', R),$$

and $\tilde{\mathcal{R}}(\sigma, \bar{w})$ is as in (2.2.2).

The proof of Proposition 3.2.6 will use the following lemma.

Lemma 3.2.7. *With notation as in Proposition 3.2.6:*

- (1) *For every fixed $\delta', \varepsilon, \varepsilon', R \in \mathbb{R}_+$, there exists $\delta_0 = \delta_0(\delta', \varepsilon, \varepsilon', R) > 1$ such that for all $1 < t_1 \leq t_2 \leq \delta_0$, the inclusion map $S_\sigma(t_1, \delta', \varepsilon, \varepsilon', R) \hookrightarrow S_\sigma(t_2, \delta', \varepsilon, \varepsilon', R)$ is a homotopy equivalence.*
- (2) *For every fixed $\varepsilon, \varepsilon', R \in \mathbb{R}_+$, there exists $\delta'_0 = \delta'_0(\varepsilon, \varepsilon', R) > 0$ such that for all $0 < t'_1 \leq t'_2 \leq \delta'_0$, the inclusion map*

$$\bigcap_{t>1} S_\sigma(t, t'_1, \varepsilon, \varepsilon', R) \hookrightarrow \bigcap_{t>1} S_\sigma(t, t'_2, \varepsilon, \varepsilon', R)$$

is a homotopy equivalence.

- (3) *Let*

$$S'_\sigma(\varepsilon, \varepsilon', R) := \bigcap_{t>1, t'>0} S_\sigma(t, t', \varepsilon, \varepsilon', R).$$

For every fixed $\varepsilon', R \in \mathbb{R}_+$, there exists $\varepsilon_0 = \varepsilon_0(\varepsilon', R) > 1$ such that for all $1 < s_1 \leq s_2 \leq \varepsilon_0$, the natural inclusion

$$S'_\sigma(s_2, \varepsilon', R) \hookrightarrow S'_\sigma(s_1, \varepsilon', R)$$

is a homotopy equivalence.

- (4) *For every fixed $R \in \mathbb{R}_+$, there exists $\varepsilon'_0 = \varepsilon'_0(R) > 0$ such that for all $0 < s'_1 \leq s'_2 \leq \varepsilon'_0$, the natural inclusion*

$$\bigcup_{s>1} S'_\sigma(s, s'_2, R) \hookrightarrow \bigcup_{s>1} S'_\sigma(s, s'_1, R)$$

is a homotopy equivalence.

(5) The following equality holds:

$$\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R) = \bigcup_{s>1, s'>0} S'_\sigma(s, s', R).$$

(6) There exists $R_0 > 0$ such that for all $R > R_0$, the natural inclusion

$$\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R) \hookrightarrow \tilde{\mathcal{R}}(\sigma, \bar{w})$$

is a homotopy equivalence.

Remark 3.2.8. (i) The subsets $S_\sigma(t, \delta', \varepsilon, \varepsilon', R)$ form an *increasing* sequence in t , i.e. if $t_1 < t_2$, then $S_\sigma(t_1, \delta', \varepsilon, \varepsilon', R) \subset S_\sigma(t_2, \delta', \varepsilon, \varepsilon', R)$. The analogous assertion also holds for $S_\sigma(\delta, t', \varepsilon, \varepsilon', R)$ (with t' replacing t).

(ii) The subsets $S_\sigma(\delta, \delta', s, \varepsilon', R)$ form a *decreasing* sequence in s , i.e. if $s_1 < s_2$, then $S_\sigma(\delta, \delta', s_2, \varepsilon', R) \subset S_\sigma(\delta, \delta', s_1, \varepsilon', R)$. The analogous assertion also holds for $S_\sigma(\delta, \delta', \varepsilon, s', R)$.

(iii) The sequence of subsets $S_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)$ is increasing in R .

Proof of Lemma 3.2.7. We prove each part separately.

Proof of (1). Let

$$S_\sigma^1(\delta', \varepsilon, \varepsilon', R) := \bigcup_{t>1} S_\sigma(t, \delta', \varepsilon, \varepsilon', R).$$

First observe that $S_\sigma^1(\delta', \varepsilon, \varepsilon', R)$ is a semi-algebraic subset of $B_{\mathbf{F}}(V)$. To see this let

$$\begin{aligned} \Phi_{\sigma, \delta', \varepsilon, \varepsilon'}(\cdot; T) &:= \bigwedge_{i, \sigma(i)=1} \phi^o(\cdot, w_i; T, \delta') \wedge \bigwedge_{i, \sigma(i)=0} \neg(\phi^o(\cdot, w_i; \varepsilon, \varepsilon')) \\ &\quad \wedge \bigwedge_{1 \leq i \leq N} (|X_i| \leq R), \\ \Phi_{\sigma, \delta', \varepsilon, \varepsilon'}^1(\cdot) &:= (\exists T) (T > 1) \wedge \Phi_{\sigma, \delta', \varepsilon, \varepsilon'}(\cdot; T). \end{aligned}$$

By A.2.2(7),

$$S_\sigma^1(\delta', \varepsilon, \varepsilon', R) = \tilde{\mathcal{R}}(\Phi_{\sigma, \delta', \varepsilon, \varepsilon'}^1, V).$$

It follows that $S_\sigma^1(\delta', \varepsilon, \varepsilon', R)$ is a semi-algebraic subset of $B_{\mathbf{F}}(V)$.

Now consider the function $f : \mathcal{R}(\Phi_{\sigma, \delta', \varepsilon, \varepsilon'}^1, V) \rightarrow \mathbb{R}_+$ defined by

$$f(x) := \inf_{\{(x, t) \mid \Phi_{\sigma, \delta', \varepsilon, \varepsilon'}(x; t)\}} t.$$

It is clear that f is definable. Note that $S_\sigma(t, \delta', \varepsilon, \varepsilon', R) = \tilde{\mathcal{R}}(\Phi_{\sigma, \delta', \varepsilon, \varepsilon'}^1 \wedge f \geq t, V)$. The claim now follows directly from Theorem A.3. □

Proof of (2). Let

$$S_\sigma^2(\varepsilon, \varepsilon', R) := \bigcup_{t'>0} \bigcap_{t>1} S_\sigma(t, t', \varepsilon, \varepsilon', R).$$

Then $S_\sigma^2(\varepsilon, \varepsilon', R)$ is a semi-algebraic subset of $B_{\mathbf{F}}(V)$. To see this let

$$\begin{aligned} \Phi_{\sigma, \varepsilon, \varepsilon'}^2(\cdot; T') &:= \bigwedge_{\sigma(i)=1} \phi^c(\cdot, w_i; 1, T') \wedge \bigwedge_{\sigma(i)=0} \neg(\phi^o(\cdot, w_i; \varepsilon, \varepsilon')) \wedge \bigwedge_{1 \leq i \leq N} (|X_i| \leq R), \\ \Phi_{\sigma, \varepsilon, \varepsilon'}^3(\cdot) &:= (\exists T') (T' > 0) \wedge \Phi_{\sigma, \varepsilon, \varepsilon'}^2(\cdot; T'). \end{aligned}$$

As in the previous part, $S_\sigma^2(\delta', \varepsilon, \varepsilon', R) = \widetilde{\mathcal{R}}(\Phi_{\sigma, \varepsilon, \varepsilon'}^3, V)$. In particular, $S_\sigma^2(\delta', \varepsilon, \varepsilon', R)$ is semi-algebraic.

Moreover, let $g : \mathcal{R}(\Phi_{\sigma, \varepsilon, \varepsilon'}^3, V) \rightarrow \mathbb{R}_+$ be defined by

$$g(x) := \inf_{\{(x, t') \mid \Phi_{\sigma, \varepsilon, \varepsilon'}^3(x; t')\}} t'.$$

Clearly, g is definable and $S_\sigma^2(t', \varepsilon, \varepsilon', R) = \widetilde{\mathcal{R}}(\Phi_{\sigma, \varepsilon, \varepsilon'}^3 \wedge g \geq t', V)$. As in the previous part, the result follows by applying Theorem A.3 to g . \square

Proof of (3). First note that $S_\sigma^3(\varepsilon', R) = \bigcup_{s>1} S'_\sigma(s, \varepsilon', R)$ is a semi-algebraic subset of $B_{\mathbf{F}}(V)$. To see this let

$$\begin{aligned} \Phi_{\sigma, \varepsilon'}^4(\cdot; S) &:= \bigwedge_{\sigma(i)=1} \phi^c(\cdot, w_i; 1, 0) \wedge \bigwedge_{\sigma(i)=0} \neg(\phi^o(\cdot, w_i; S, \varepsilon')) \wedge \bigwedge_{1 \leq i \leq N} (|X_i| \leq R), \\ \Phi_{\sigma, \varepsilon'}^5(\cdot) &:= (\exists S) (S > 1) \wedge \Phi_{\sigma, \varepsilon'}^4(\cdot; S). \end{aligned}$$

Then $S_\sigma^3(\varepsilon', R) = \widetilde{\mathcal{R}}(\Phi_{\sigma, \varepsilon'}^5, V)$. In particular, $S_\sigma^3(\varepsilon', R)$ is semi-algebraic.

Let $h : \mathcal{R}(\Phi_{\sigma, \varepsilon'}^5, V) \rightarrow \mathbb{R}_+$ be given by

$$h(x) := \sup_{\{(x, s) \mid \Phi_{\sigma, \varepsilon'}^5(x; s)\}} s.$$

Clearly, h is definable. Moreover, $S'_\sigma(s, \varepsilon', R)$ equals $\widetilde{\mathcal{R}}(\Phi_{\sigma, \varepsilon'}^5 \wedge h \geq s, V)$ and therefore is also semi-algebraic. Now apply Theorem A.3. \square

Proof of (4). Let $S_\sigma^4(R) := \bigcup_{s'>0} S_\sigma^3(s', R)$, and consider

$$\Phi_\sigma^6(\cdot) := (\exists S') (S' > 0) \wedge \Phi_{\sigma, S'}^5(\cdot).$$

Then $S_\sigma^4(R) = \widetilde{\mathcal{R}}(\Phi_\sigma^6, V)$. In particular, $S_\sigma^4(R)$ is semi-algebraic.

We can now consider the function $h : \mathcal{R}(\Phi_\sigma^6, V) \rightarrow \mathbb{R}_+$ given by

$$h(x) := \sup_{\{(x, s') \mid \Phi_{\sigma, s'}^5(x)\}} s'.$$

One can now argue as in (3). \square

Proof of (5). This follows from the definition of $S'_\sigma(s, s', R)$. \square

Proof of (6). This part follows immediately from Theorem A.3. For example, consider the definable function h on $\tilde{\mathcal{R}}(\sigma, \bar{w})$ given by

$$h(x) := \frac{1}{\max_i(\max(1, |x_i|))},$$

where x_i 's are the coordinates. Then $h(x) \geq 0$ for all $x \in V$, and for all $0 < \varepsilon \leq 1$,

$$h(x) \geq \varepsilon \iff x \in \text{Cube}_V(1/\varepsilon).$$

Then there exists $0 < \varepsilon_0 < 1$ such that for all $0 < \varepsilon \leq \varepsilon_0$ the natural inclusions

$$\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(1/\varepsilon) \hookrightarrow \tilde{\mathcal{R}}(\sigma, \bar{w}) = \tilde{\mathcal{R}}(\sigma, \bar{w}) \cap B_{\mathbb{F}}(h \geq 0)$$

are homotopy equivalences. Now we set $R_0 := 1/\varepsilon_0 > 0$, and for any $R \geq R_0$, we consider $\varepsilon(R) := 1/R$ to obtain the desired conclusion. \square

This completes the proof of Lemma 3.2.7. \square

We now prove Proposition 3.2.6. Since the proof is long and technical, we begin by giving a general outline. Because of the nature of the argument, the steps enumerated do not actually occur in the same order as in the list below.

Step 1. By Lemma 3.2.7(6), there exists an $R_0 > 0$ such that for all $R > R_0$ the natural inclusion

$$\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R) \hookrightarrow \tilde{\mathcal{R}}(\sigma, \bar{w})$$

induces an isomorphism

$$H^*(\tilde{\mathcal{R}}(\sigma, \bar{w})) \xrightarrow{\cong} H^*(\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)).$$

So we fix some $R > 0$ large enough and consider only the semi-algebraic set $\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)$.

Step 2. By Lemma 3.2.7(5), we have natural inclusions

$$S'_\sigma(s, s', R) \hookrightarrow \bigcup_{s>1, s'>0} S'_\sigma(s, s', R) = \tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R).$$

We shall see in Claim 4 below that this induces an isomorphism

$$H^*(\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)) \cong \varprojlim_{s'} \varprojlim_s H^*(S'_\sigma(s, s', R)).$$

Step 3. We shall see in Claim 1 below that the natural inclusions

$$S'_\sigma(\varepsilon, \varepsilon', R) \hookrightarrow S_\sigma(t, t', \varepsilon, \varepsilon', R)$$

induce an isomorphism

$$\varinjlim_{t'} \varinjlim_t H^*(S_\sigma(t, t', \varepsilon, \varepsilon', R)) \cong H^*(S'_\sigma(\varepsilon, \varepsilon', R)).$$

Step 4. In order to conclude, we shall show that the direct and inverse limits appearing in Step 2 (proved in Claim 6) and Step 3 (proved in Claim 3) ‘stabilize’. This will be a consequence of the homotopy equivalences proved in Lemma 3.2.7, and it is proved in two intermediate steps (Claims 4 and 5 for Step 2, and Claims 2 and 3 for Step 3).

The proofs involving commutation of the limit (or colimit) functors with cohomology in Steps 2 and 3 all rely on proving that a certain increasing family of compact subspaces S_λ , of a semi-algebraic set T , indexed by a real parameter λ , are cofinal in the family of all compact subspaces of $S := \bigcup_\lambda S_\lambda$ (the families are different for different steps). One then uses Lemma A.1.2 to obtain the desired commutation of various limits (or colimits) with cohomology. The proofs of all these cofinality statements rely on the following basic lemma that we extract out for clarity.

Lemma 3.2.9. *Let T be a compact Hausdorff space, Λ a partially ordered set, $(C_\lambda)_{\lambda \in \Lambda}$ an increasing sequence of compact subsets of T , and $S := \bigcup_\lambda C_\lambda$. Suppose that there is a continuous function $\theta : S \rightarrow \mathbb{R}_{>0} \cup \{\infty\}$ such that the following property holds:*

$$(3.2.10) \text{ For each } \theta_0 \in \mathbb{R}_{>0}, \text{ there exists a } \lambda(\theta_0) \in J \text{ such that } x \in C_{\lambda(\theta_0)} \text{ if } \theta(x) \geq \theta_0.$$

Then $(C_\lambda)_{\lambda \in \Lambda}$ is cofinal in the family of compact subsets of S .

Proof. Let C be a compact subset of S in T . We need to show that there is a λ such that $C \subset C_\lambda$. Since C is compact, $F|_C$ attains its minimum $\theta_0 > 0$ on C . Let $\lambda(\theta_0)$ be as in the proposition. Clearly,

$$x \in C \implies \theta(x) \geq \theta_0 \implies x \in C_{\lambda(\theta_0)}.$$

It follows that $C \subset C_{\lambda(\theta_0)}$, and so $(C_\lambda)_{\lambda \in \Lambda}$ is cofinal in the family of compact subsets of S . □

Proof of Proposition 3.2.6.

Claim 1. *The natural inclusions*

$$(3.2.11) \quad S'_\sigma(\varepsilon, \varepsilon', R) := \bigcap_{t>1, t'>0} S_\sigma(t, t', \varepsilon, \varepsilon', R) \hookrightarrow S_\sigma(t, t', \varepsilon, \varepsilon', R)$$

induce an isomorphism

$$(3.2.12) \quad H^*(S'_\sigma(\varepsilon, \varepsilon', R)) \cong \varinjlim_{t, t'} H^*(S_\sigma(t, t', \varepsilon, \varepsilon', R)).$$

As an immediate consequence,

$$(3.2.13) \quad H^*(S'_\sigma(\varepsilon, \varepsilon', R)) \cong \varinjlim_{t'} \varinjlim_t H^*(S_\sigma(t, t', \varepsilon, \varepsilon', R)).$$

(Here the inductive limit in (3.2.12) is taken over the poset $\mathbb{R}_{>1} \times \mathbb{R}_{>0}$ partially ordered by

$$(t_1, t'_1) \preceq (t_2, t'_2) \iff t_2 \leq t_1 \text{ and } t'_2 \leq t'_1,$$

and for $(t_1, t'_1) \preceq (t_2, t'_2)$, the morphism

$$H^*(S_\sigma(t_1, t'_1, \varepsilon, \varepsilon', R)) \rightarrow H^*(S_\sigma(t_2, t'_2, \varepsilon, \varepsilon', R))$$

is induced by the inclusion $S_\sigma(t_2, t'_2, \varepsilon, \varepsilon', R) \hookrightarrow S_\sigma(t_1, t'_1, \varepsilon, \varepsilon', R)$.

Proof of Claim 1. First note that the isomorphism (3.2.13) is an immediate consequence of the isomorphism (3.2.12) and the fact that

$$\lim_{\substack{\longrightarrow \\ t'}} \lim_{\substack{\longrightarrow \\ t}} H^*(S_\sigma(t, t', \varepsilon, \varepsilon', R)) \cong \lim_{\substack{\longrightarrow \\ t, t'}} H^*(S_\sigma(t, t', \varepsilon, \varepsilon', R))$$

(see for example [SGA72, Exp. 1, p. 13] for the last isomorphism).

We now proceed to prove the isomorphism (3.2.12). Let

$$T := \bigcap_{i, \sigma(i)=0} \text{TubeCompl}_{V, \phi(\cdot, w_i)}^c(\varepsilon, \varepsilon', R).$$

Since each $\text{TubeCompl}_{V, \phi(\cdot, w_i)}^c(\varepsilon, \varepsilon', R)$ is compact, T is a compact Hausdorff space. Notice that for each $t > 1, t' > 0, S_\sigma(t, t', \varepsilon, \varepsilon', R) \subset T$.

We will now show that for fixed $\varepsilon, \varepsilon', R$, the family of semi-algebraic sets

$$(3.2.14) \quad (S_\sigma(t, t', \varepsilon, \varepsilon', R))_{t>1, t'>0}$$

is a cofinal system of open neighborhoods of

$$\bigcap_{t>1, t'>0} S_\sigma(t, t', \varepsilon, \varepsilon', R)$$

in T . Assuming this fact, the claim follows from Lemma A.1.2(1).

We first prove the following cofinality statement from which the cofinality of (3.2.14) will follow.

Suppose that I is a finite set, and for each $i \in I$ let $F_i, G_i \in K[X_1, \dots, X_N]$ and $\lambda_i \in \mathbb{R}_+$. Let V be as before, $R > 0$, and $T^{(1)}$ a compact semi-algebraic subset of $\text{Cube}_V(R)$. We define

$$S^{(1)}(t, t', R) := T^{(1)} \cap \bigcap_{i \in I} \text{Tube}_{V, |F_i| \leq \lambda_i \cdot |G_i|}^o(t, t', R).$$

Notice that for each $t > 1, t' > 0, S^{(1)}(t, t', R) \subset T^{(1)}$, and hence

$$\bigcap_{t>1, t'>0} S^{(1)}(t, t', R) \subset T^{(1)}$$

as well.

Claim 1a. *The family of semi-algebraic sets $(S^{(1)}(t, t', R))_{t>1, t'>0}$ is a cofinal system of open neighborhoods of*

$$\bigcap_{t>1, t'>0} S^{(1)}(t, t', R)$$

in $T^{(1)}$.

Proof of Claim 1a. Proving the claim is equivalent to proving the cofinality of the family of compact subsets

$$(T^{(1)} \setminus S^{(1)}(t, t', R))_{t>1, t'>0}$$

in the partially ordered family of compact subsets of $T^{(1)} \setminus \bigcap_{t>1, t'>0} S^{(1)}(t, t', R)$. To prove the latter we use Lemma 3.2.9 with $\Lambda := \mathbb{R}_{>1} \times \mathbb{R}_{>0}$, and the family $(C_\lambda)_{\lambda \in \Lambda} := (T^{(1)} \setminus S^{(1)}(t, t', R))_{(t, t') \in \Lambda}$ of compact semi-algebraic subsets of the compact set $T^{(1)}$.

We now define a continuous function $\theta : T^{(1)} \setminus \bigcap_{t>1, t'>0} S^{(1)}(t, t', R) \rightarrow \mathbb{R}_{\geq 0}$. We first introduce the following auxiliary functions. For $\lambda \geq 0$, let $H_\lambda(u, v) : \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be defined as follows. If $\lambda = 0$, then

$$H_0(u, v) := u,$$

and if $\lambda > 0$,

$$(3.2.15) \quad H_\lambda(u, v) := \begin{cases} \min\left(\max(u, v), \max\left(0, \frac{u}{\lambda v} - 1\right)\right) & \text{if } v \neq 0, \\ u & \text{else.} \end{cases}$$

It is easy to check that the functions $H_\lambda(u, v)$ are continuous.

For each $i \in I$, let $\theta_i : T^{(1)} \setminus \bigcap_{t>1, t'>0} S^{(1)}(t, t', R) \rightarrow \mathbb{R}_{\geq 0}$ be defined by

$$\theta_i(x) := H_{\lambda_i}(|F_i(x)|, |G_i(x)|),$$

and let $\theta : T^{(1)} \setminus \bigcap_{t>1, t'>0} S^{(1)}(t, t', R) \rightarrow \mathbb{R}_{\geq 0}$ be defined by

$$\theta(x) := \max_{i \in I} \theta_i(x).$$

Notice that each θ_i (and hence also θ) are continuous, since they are compositions of continuous functions.

In order to apply Lemma 3.2.9 it remains to check that with λ defined by $\lambda(\theta_0) = (1 + \theta_0, \theta_0)$:

- (1) $\theta(x) > 0$ for each $x \in T^{(1)} \setminus \bigcap_{t>1, t'>0} S^{(1)}(t, t', R)$.
- (2) θ satisfies (3.2.10) with λ defined by $\lambda(\theta_0) = (1 + \theta_0, \theta_0)$.

To prove (1), suppose that $\theta(x) = 0$. Then $\theta_i(x) = 0$ for each $i \in I$.

If $\lambda_i = 0$, then $\theta_i(x) = 0$ implies that $|F_i(x)| = 0$. If $\lambda_i > 0$, then $\theta_i(x) = 0$ implies that either $|F_i(x)| = |G_i(x)| = 0$ or $|F_i(x)|/(\lambda_i \cdot |G_i(x)|) \leq 1$ or equivalently $|F_i(x)| \leq \lambda_i \cdot |G_i(x)|$. Together they imply that $x \in \bigcap_{t>1, t'>0} S^{(1)}(t, t', R)$, which is a contradiction.

To prove (2), suppose $\theta(x) \geq \theta_0$. First note that

$$\begin{aligned} T^{(1)} \setminus S^{(1)}(1 + \theta_0, \theta_0, R) &= T^{(1)} \setminus \bigcap_{i \in I} \text{Tube}_{V, |F_i| \leq \lambda_i \cdot |G_i|}^o(1 + \theta_0, \theta_0, R) \\ &= T^{(1)} \cap \bigcup_{i \in I} \text{TubeComp}^c_{V, |F_i| \leq \lambda_i \cdot |G_i|}(1 + \theta_0, \theta_0, R), \end{aligned}$$

which is equal to the set

$$T^{(1)} \cap \bigcup_{i \in I} \widetilde{\mathcal{R}}(|F_i| \geq \lambda_i \cdot (1 + \theta_0) \cdot |G_i|) \wedge ((|F_i| \geq \theta_0) \vee (|G_i| \geq \theta_0)), V).$$

Since $\theta(x) \geq \theta_0$, there exists an i such that $\theta(x) = \theta_i(x) = \theta_0$. Hence $|F_i(x)|$ and $|G_i(x)|$ are not simultaneously 0. We have two cases. If $\lambda_i = 0$, then

$$|F_i(x)| = \theta_i(x) \geq \theta_0,$$

which implies that

$$(3.2.16) \quad x \in \widetilde{\mathcal{R}}(|F_i| \geq (\lambda_i \cdot (1 + \theta_0) \cdot |G_i|) \wedge ((|F_i| \geq \theta_0) \vee (|G_i| \geq \theta_0)), V).$$

Otherwise, $\lambda_i > 0$. If $|G_i(x)| \neq 0$, we have

$$\max(|F_i(x)|, |G_i(x)|) \geq \theta_i(x) \geq \theta_0,$$

and

$$\max\left(0, \frac{|F_i(x)|}{\lambda_i |G_i(x)|} - 1\right) \geq \theta_i(x) \geq \theta_0,$$

which again implies (3.2.16). If $|G_i(x)| = 0$, then $|F_i(x)| = \theta_0$, and again (3.2.16) holds. This yields (2), thus completing the proof of Claim 1a. □

Now we return to the proof of Claim 1. Let $\phi = \bigvee_{h \in H} \phi^{(h)}$, where each $\phi^{(h)}$ is a conjunction of weak inequalities, $|F_{jh}| \leq \lambda_{jh} \cdot |G_{jh}|$, $j \in J_h$, and H, J_h are finite sets.

Let $I_\sigma := \{i \in [1, n] \mid \sigma_i = 1\}$ and let H^{I_σ} denote the set of maps $\psi : I_\sigma \rightarrow H$. Note that

$$S_\sigma(t, t', \varepsilon, \varepsilon', R) = \bigcap_{I_\sigma} \left(\bigcup_{h \in H} \bigcap_{j \in J_h} \text{Tube}_{V, |F_{jh}(\cdot, w_i)| \leq \lambda_{jh} \cdot |G_{jh}(\cdot, w_i)|}^o(t, t', R) \right) \cap T.$$

(Recall that $T = \bigcap_{i, \sigma_i=0} \text{TubeComp}^c_{V, \phi(\cdot, w_i)}(\varepsilon, \varepsilon', R)$ is a compact semi-algebraic set.) Then

$$S_\sigma(t, t', \varepsilon, \varepsilon', R) = \bigcup_{\psi \in H^{I_\sigma}} S_\sigma^{(\psi)}(t, t', \varepsilon, \varepsilon', R),$$

where for $\psi \in H^{I_\sigma}$,

$$S_\sigma^{(\psi)}(t, t', \varepsilon, \varepsilon', R) := T \cap \bigcap_{i, \sigma_i=1} \text{Tube}_{V, \phi^{(\psi(i))}(\cdot, w_i)}^o(t, t', R).$$

An open neighborhood U of $\bigcap_{t>1, t'>0} S_\sigma(t, t', \varepsilon, \varepsilon', R)$ in T is clearly also an open neighborhood of $\bigcap_{t>1, t'>0} S_\sigma^{(\psi)}(t, t', \varepsilon, \varepsilon', R)$ for each $\psi \in H^{I_\sigma}$.

Fixing a $\psi \in H^{I_\sigma}$, we apply Claim 1a with

$$T^{(1)} := T, \quad I := \{(j, \psi(i)) \mid i \in I_\sigma, j \in J_{\psi(i)}\},$$

and for $i_0 = (j, \psi(i)) \in I$,

$$F_{i_0} = F_{j, \psi(i)}, \quad G_{i_0} = G_{j, \psi(i)}, \quad \lambda_{i_0} = \lambda_{j, \psi(i)}.$$

We deduce that for each $\psi \in H^{I_\sigma}$, there exists $\theta_0^{(\psi)} > 0$ such that

$$S_\sigma^{(\psi)}(1 + \theta_0^{(\psi)}, \theta_0^{(\psi)}, \varepsilon, \varepsilon', R) \subset U.$$

Now take $\theta_0 := \min_{\psi \in H^{I_\sigma}} \theta_0^{(\psi)}$. Then

$$S_\sigma(1 + \theta_0, \theta_0, \varepsilon, \varepsilon', R) = \bigcup_{\psi \in H^{I_\sigma}} S_\sigma^{(\psi)}(1 + \theta_0, \theta_0, \varepsilon, \varepsilon', R) \subset U.$$

This proves (3.2.12) and concludes the proof of Claim 1. □

Claim 2. *The natural inclusions*

$$\bigcap_{t>1} S_\sigma(t, t', \varepsilon, \varepsilon', R) \hookrightarrow S_\sigma(t, t', \varepsilon, \varepsilon', R)$$

induce for each fixed $t' > 0, \varepsilon > 1, \varepsilon' > 0, R > 0$ an isomorphism

$$(3.2.17) \quad H^*\left(\bigcap_{t>1} S_\sigma(t, t', \varepsilon, \varepsilon', R)\right) \cong \varinjlim_t H^*(S_\sigma(t, t', \varepsilon, \varepsilon', R)).$$

Proof of Claim 2. The proof is structurally similar to the proof of Claim 1. Let

$$T := \bigcap_{i, \sigma(i)=0} \text{TubeCompl}_{V, \phi(\cdot, w_i)}^c(\varepsilon, \varepsilon', R).$$

Then T is compact. We will show for fixed $t', \varepsilon, \varepsilon', R$, the family of semi-algebraic sets

$$(3.2.18) \quad (S_\sigma(t, t', \varepsilon, \varepsilon', R))_{t>1}$$

is a cofinal system of open neighborhoods of $\bigcap_{t>1} S_\sigma(t, t', \varepsilon, \varepsilon', R)$ in T . Assuming this fact, the claim follows from Lemma A.1.2(1).

We first prove the following cofinality statement from which the cofinality of (3.2.18) will follow.

Suppose that I is a finite set, and for each $i \in I$ let $F_i, G_i \in K[X_1, \dots, X_N]$ and $\lambda_i \in \mathbb{R}_+$. Let V be as before, $R > 0$, and $T^{(2)}$ a compact semi-algebraic subset of $\text{Cube}_V(R)$. We define

$$S^{(2)}(t, t', R) := T^{(2)} \cap \bigcap_{i \in I} \text{Tube}_{V, |F_i| \leq \lambda_i \cdot |G_i|}^o(t, t', R).$$

Claim 2a. *The family of semi-algebraic sets $(S^{(2)}(t, t', R))_{t>1}$ is a cofinal system of open neighborhoods of*

$$\bigcap_{t>1} S^{(2)}(t, t', R)$$

in $T^{(2)}$.

Proof of Claim 2a. Proving the claim is equivalent to proving that the family of compact semi-algebraic sets

$$(T^{(2)} \setminus S^{(2)}(t, t', R))_{t>1}$$

is cofinal in the family of compact subsets of $T^{(2)} \setminus \bigcap_{t>1} S^{(2)}(t, t', R)$.

Let

$$\begin{aligned} S_i^{(2)}(t, t', R)^c &:= T^{(2)} \cap \text{TubeComp}^c_{V, |F_i| \leq \lambda_i \cdot |G_i|}(t, t', R) \\ &= \begin{cases} T^{(2)} \cap \tilde{\mathcal{R}}(|F_i| \geq t \cdot \lambda_i \cdot |G_i|) \\ \wedge ((|F_i| \geq t') \vee (|G_i| \geq t')), V & \text{if } \lambda_i > 0, \\ T^{(2)} \cap \tilde{\mathcal{R}}(|F_i| \geq t', V) & \text{if } \lambda_i = 0. \end{cases} \end{aligned}$$

Note that

$$T^{(2)} \setminus S^{(2)}(t, t', R) = \bigcup_{i \in I} S_i^{(2)}(t, t', R)^c,$$

and

$$T^{(2)} \setminus \bigcap_{t>1} S^{(2)}(t, t', R) = \bigcup_{i \in I} \bigcup_{t>1} S_i^{(2)}(t, t', R)^c.$$

The last cofinality statement would follow if for each i we can show that the family of compact semi-algebraic sets $(S_i^{(2)}(t, t', R)^c)_{t>1}$ is cofinal in the family of compact subspaces of $\bigcup_{t>1} S_i^{(2)}(t, t', R)^c$. This is because if for each compact subspace

$$C \subset T^{(2)} \setminus \bigcap_{t>1} S^{(2)}(t, t', R) = \bigcup_{i \in I} \bigcup_{t>1} S_i^{(2)}(t, t', R)^c$$

and $i \in I$, there exists $t_{0,i} > 1$ such that $C \cap \bigcup_{t>1} S_i^{(2)}(t, t', R)^c \subset S_i^{(2)}(t_{0,i}, t', R)^c$, then $C \subset T^{(2)} \setminus S^{(2)}(t_0, t', R)$ with $t_0 := \min_i t_{0,i}$.

We now proceed to show the cofinality of $(S_i^{(2)}(t, t', R)^c)_{t>1}$ in the family of compact subspaces of $\bigcup_{t>1} S_i^{(2)}(t, t', R)^c$ using Lemma 3.2.9.

For each $i \in I$, define the continuous function $\theta_i : \bigcup_{t>1} S_i^{(2)}(t, t', R)^c \rightarrow \mathbb{R}_+ \cup \{\infty\}$ by

$$(3.2.19) \quad \theta_i(x) := \begin{cases} |F_i(x)| & \text{if } \lambda_i = 0, \\ \frac{|F_i(x)|}{\lambda_i |G_i(x)|} & \text{if } \lambda_i > 0. \end{cases}$$

It is an easy exercise to check that the functions θ_i are positive and satisfy (3.2.10) with the map λ defined by

$$\lambda(\theta_0) := \begin{cases} t' & \text{if } \lambda_i = 0, \\ \theta_0 & \text{if } \lambda_i > 0. \end{cases}$$

This finishes the proof of Claim 2a. □

Claim 2 follows from Claim 2a in exactly the same manner as in the proof of Claim 1 from Claim 1a. □

Claim 3. For every fixed $\varepsilon > 1, \varepsilon' > 0$ and $R > 0$, there exists $\delta'_0 > 0$ such that for each $0 < \delta' \leq \delta'_0$, there exists $\delta_0(\delta') > 1$ such that the inclusion

$$S'_\sigma(\varepsilon, \varepsilon', R) \hookrightarrow S_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)$$

induces an isomorphism

$$(3.2.20) \quad H^*(S'_\sigma(\varepsilon, \varepsilon', R)) \cong H^*(S_\sigma(\delta, \delta', \varepsilon, \varepsilon', R))$$

for all $1 < \delta \leq \delta_0(\delta')$.

Proof of Claim 3. We fix $\varepsilon > 1, \varepsilon' > 0$ and $R > 0$. First, note that (3.2.13) yields

$$(3.2.21) \quad H^*(S'_\sigma(\varepsilon, \varepsilon', R)) \cong \varinjlim_{t'} \varinjlim_t H^*(S_\sigma(t, t', \varepsilon, \varepsilon', R)).$$

By Lemma 3.2.7(2) there exists δ'_0 such that for all $0 < t'_2 \leq t'_1 \leq \delta'_0$, the inclusion map

$$\bigcap_{t>1} S_\sigma(t, t'_2, \varepsilon, \varepsilon', R) \hookrightarrow \bigcap_{t>1} S_\sigma(t, t'_1, \varepsilon, \varepsilon', R)$$

induces an isomorphism

$$H^*\left(\bigcap_{t>1} S_\sigma(t, t'_1, \varepsilon, \varepsilon', R)\right) \rightarrow H^*\left(\bigcap_{t>1} S_\sigma(t, t'_2, \varepsilon, \varepsilon', R)\right).$$

It follows that, for any $0 < \delta' \leq \delta'_0$,

$$(3.2.22) \quad \varinjlim_{t'} H^*\left(\bigcap_{t>1} S_\sigma(t, t', \varepsilon, \varepsilon', R)\right) \cong H^*\left(\bigcap_{t>1} S_\sigma(t, \delta', \varepsilon, \varepsilon', R)\right).$$

Moreover, it follows from (3.2.17) that

$$(3.2.23) \quad H^*\left(\bigcap_{t>1} S_\sigma(t, t', \varepsilon, \varepsilon', R)\right) \cong \varinjlim_t H^*(S_\sigma(t, t', \varepsilon, \varepsilon', R))$$

for each fixed $t' > 0, \varepsilon > 1, \varepsilon' > 0$ and $R > 0$. Hence, from (3.2.21)–(3.2.23) we get

$$(3.2.24) \quad H^*(S'_\sigma(\varepsilon, \varepsilon', R)) \cong \varinjlim_t H^*(S_\sigma(t, \delta', \varepsilon, \varepsilon', R))$$

It again follows from Lemma 3.2.7(1) that for each fixed δ' , there exists $\delta_0(\delta')$ such that for all $1 < t_2 \leq t_1 \leq \delta_0(\delta')$ the inclusion $S_\sigma(t_2, \delta', \varepsilon, \varepsilon', R) \hookrightarrow S_\sigma(t_1, \delta', \varepsilon, \varepsilon', R)$ induces an isomorphism

$$H^*(S_\sigma(t_1, \delta', \varepsilon, \varepsilon', R)) \rightarrow H^*(S_\sigma(t_2, \delta', \varepsilon, \varepsilon', R)),$$

which implies that

$$(3.2.25) \quad \varprojlim_{t'} H^*(S_\sigma(t, \delta', \varepsilon, \varepsilon', R)) \cong H^*(S_\sigma(t_0, \delta', \varepsilon, \varepsilon', R))$$

for all $1 < t_0 \leq \delta_0(\delta')$. Claim 3 follows from (3.2.24) and (3.2.25), after taking δ'_0 and $\delta_0(\delta')$ as above. □

Claim 4. *The inclusion*

$$\bigcup_{s>1, s'>0} S'_\sigma(s, s', R) \hookrightarrow \widetilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)$$

induces an isomorphism

$$(3.2.26) \quad H^*(\widetilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)) \cong \varprojlim_{s', s} H^*(S'_\sigma(s, s', R)).$$

As an immediate consequence,

$$(3.2.27) \quad H^*(\widetilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)) \cong \varprojlim_{s'} \varprojlim_s H^*(S'_\sigma(s, s', R)).$$

(Here the projective limit is taken over the poset $\mathbb{R}_{>1} \times \mathbb{R}_{>0}$ partially ordered by

$$(s_1, s'_1) \leq (s_2, s'_2) \iff s_2 \leq s_1 \text{ and } s'_2 \leq s'_1,$$

and for $(s_1, s'_1) \leq (s_2, s'_2)$, the morphism

$$H^*(S'_\sigma(s_2, s'_2, R)) \rightarrow H^*(S'_\sigma(s_1, s'_1, R))$$

is induced by inclusion.)

Proof of Claim 4. First note that (3.2.27) is an immediate consequence of (3.2.26) and the fact that

$$\varprojlim_{s'} \varprojlim_s H^*(S'_\sigma(s, s', R)) \cong \varprojlim_{s, s'} H^*(S'_\sigma(s, s', R))$$

(see for example [SGA72, Exp. 1, p. 13] for the last isomorphism). Note that the semi-algebraic sets $S'_\sigma(s, s', R)$ are compact for each choice of $s > 1, s' > 0$ and $R > 0$. In order to see this, recall that by definition (see (3.2.11)) $S'_\sigma(s, s', R)$ is the intersection of $\bigcap_{i, \sigma(i)=1} \bigcap_{t>1, t'>0} \text{Tube}_{V, \phi(\cdot, w_i)}^O(t, t', R)$ with the compact semi-algebraic set $\bigcap_{i, \sigma(i)=0} \bigcap_{t>1, t'>0} \text{TubeCompl}_{V, \phi(\cdot, w_i)}^C(s, s', R)$. Therefore, it suffices to prove that the semi-algebraic set

$$\bigcap_{t>1, t'>0} \text{Tube}_{V, \phi(\cdot, w_i)}^O(t, t', R)$$

is compact for each i . In general, $\phi = \bigvee_{h \in H} \phi^{(h)}$ where each $\phi^{(h)}$ is a conjunction of weak inequalities $|F_{jh}| < \lambda_{jh} \cdot |G_{jh}|$, $j \in J_h$, where H and J_h are finite sets. It follows that the semi-algebraic set $\bigcap_{t>1, t'>0} \text{Tube}_{V, \phi(\cdot, w_i)}^o(t, t', R)$ is the union over H of the intersection over J_h of the semi-algebraic sets

$$\bigcap_{t>1, t'>0} \text{Tube}_{V, |F_{jh}(\cdot, w_i)| \leq \lambda_{jh} \cdot |G_{jh}(\cdot, w_i)|}^o(t, t', R).$$

We claim that

$$(3.2.28) \quad \bigcap_{t>1, t'>0} \text{Tube}_{V, |F_{jh}(\cdot, w_i)| \leq \lambda_{jh} \cdot |G_{jh}(\cdot, w_i)|}^o = \text{Cube}_V(R) \cap \tilde{\mathcal{R}}(|F_{jh}(\cdot, w_i)| \leq \lambda_{jh} \cdot |G_{jh}(\cdot, w_i)|, V),$$

and the latter set is easily seen to be compact. Verifying (3.2.28) is an easy exercise starting from the definition in (3.2.3). It follows that

$$\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R) = \bigcup_{s>1, s'>0} S'_\sigma(s, s', R),$$

where each $S'_\sigma(s, s', R)$ is a compact subset of $\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)$. We now prove that the family

$$(3.2.29) \quad (S'_\sigma(s, s', R))_{s>1, s'>0}$$

is cofinal in the family of compact subspaces of

$$\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R) = \bigcup_{s>1, s'>0} S'_\sigma(s, s', R).$$

Then the isomorphism (3.2.26) will follow from Lemma A.1.2(2).

We first prove the following cofinality statement from which the cofinality of (3.2.29) will follow.

Suppose that I is a finite set, and for each $i \in I$, let $F_i, G_i \in K[X_1, \dots, X_N]$ and $\lambda_i \in \mathbb{R}_+$. Let V and $R > 0$ be as before. We define

$$S^{(3)}(s, s', R) := \bigcup_{i \in I} \text{TubeComp}^c_{V, |F_i| \leq \lambda_i \cdot |G_i|}(s, s', R) = \begin{cases} \text{Cube}_V(R) \cap \bigcup_{i \in I} \tilde{\mathcal{R}}(|F_i| \geq s', V) & \text{if } \lambda_i = 0, \\ \text{Cube}_V(R) \cap \bigcup_{i \in I} \tilde{\mathcal{R}}(|F_i| \geq s \cdot \lambda_i \cdot |G_i|) \wedge (|F_i| \geq s' \vee |G_i| \geq s'), V & \text{if } \lambda_i > 0. \end{cases}$$

Claim 4a. *The family of semi-algebraic sets $(S^{(3)}(s, s', R))_{s>1, s'>0}$ is cofinal in the directed family of compact subspaces of*

$$\bigcup_{s>1, s'>0} S''(s, s', R).$$

Proof of Claim 4a. One can deduce this formally from Claim 1a by taking complements and setting $T^{(1)} = \text{Cube}_V(R)$. On the other hand, one can also proceed via Lemma 3.2.9 using the function

$$\theta : \bigcup_{s>1, s'>0} S^{(3)}(s, s', R) \rightarrow \mathbb{R}_{\geq 0}$$

defined as follows. For each $i \in I$, let $\theta_i : \bigcup_{s>1, s'>0} S^{(3)}(s, s', R) \rightarrow \mathbb{R}_{\geq 0}$ be defined by

$$\theta_i(x) := H_{\lambda_i}(|F_i(x)|, |G_i(x)|)$$

(see (3.2.15) for the definition of $H_{\lambda_i}(\cdot, \cdot)$), and let $\theta : \bigcup_{s>1, s'>0} S^{(3)}(s, s', R) \rightarrow \mathbb{R}_{\geq 0}$ be defined by

$$\theta(x) := \max_{i \in I} \theta_i(x).$$

One can now directly verify that θ is positive and satisfies (3.2.10) with the map λ defined by $\lambda(\theta_0) = (1 + \theta_0, \theta_0)$. We leave the details to the reader. This concludes the proof of Claim 4a. □

The proof of Claim 4 from Claim 4a is formally analogous to the similar derivation of Claim 1 from Claim 1a and is omitted. □

Claim 5. For each fixed $s' > 0$ and $R > 0$ the natural inclusion

$$S'_\sigma(s, s', R) \hookrightarrow \bigcup_{s>1} S'_\sigma(s, s', R)$$

induces an isomorphism

$$(3.2.30) \quad H^*\left(\bigcup_{s>1} S'_\sigma(s, s', R)\right) \cong \varprojlim_s H^*(S'_\sigma(s, s', R)).$$

Proof of Claim 5. The proof is structurally similar to the proof of Claim 4.

We will now show that for fixed s', R , the family of semi-algebraic sets

$$(3.2.31) \quad (S_\sigma(s, s', R))_{s>1}$$

is a cofinal system of compact subsets of $\bigcup_{s>1} S_\sigma(s, s', R)$. Assuming this fact, the claim follows from Lemma A.1.2(2).

We first prove the following cofinality statement from which the cofinality of (3.2.31) will follow.

Suppose that I is a finite set, and for each $i \in I$, let $F_i, G_i \in K[X_1, \dots, X_N]$ and $\lambda_i \in \mathbb{R}_+$. Let V and $R > 0$ be as before. We define

$$S^{(4)}(s, s', R) := \bigcup_{i \in I} \text{TubeComp}^c_{V, |F_i| \leq \lambda_i, |G_i|}(s, s', R).$$

Claim 5a. The family of semi-algebraic sets $(S^{(4)}(s, s', R))_{s>1}$ is a cofinal system of compact semi-algebraic subsets of

$$\bigcup_{s>1} S^{(4)}(s, s', R).$$

Proof of Claim 5a. One can deduce this formally from Claim 2a by taking complements and $T^{(2)} := \text{Cube}_V(R)$. Alternatively, one can argue directly as follows.

For each $i \in I$, let

$$S_i^{(4)}(s, s', R) := \text{TubeCompl}_{V, |F_i| \leq \lambda_i |G_i|}^c(s, s', R) \\ = \begin{cases} \text{Cube}_V(R) \cap \tilde{\mathcal{R}}(|F_i| \geq s', V) & \text{if } \lambda_i = 0, \\ \text{Cube}_V(R) \cap \tilde{\mathcal{R}}((|F_i| \geq s \cdot \lambda_i \cdot |G_i|) \\ \quad \wedge ((|F_i| \geq s') \vee (|G_i| \geq s')), V) & \text{if } \lambda_i > 0. \end{cases}$$

Note that

$$S^{(4)}(s, s', R) = \bigcup_{i \in I} S_i^{(4)}(s, s', R),$$

and

$$\bigcup_{s>1} S^{(4)}(s, s', R) = \bigcup_{i \in I} \bigcup_{s>1} S_i^{(4)}(s, s', R).$$

Note that the cofinality statement in our claim would follow if for each i we can show that the family of compact semi-algebraic sets $(S_i^{(4)}(s, s', R))_{s>1}$ is cofinal in the family of compact subspaces of $\bigcup_{s>1} S_i^{(4)}(s, s', R)$. To see this, suppose that we have proven the latter cofinality statement (for each i). Let $C \subset \bigcup_{s>1} S^{(4)}(s, s', R)$ be a compact subspace. Then $C_i := C \cap \bigcup_{s>1} S_i^{(4)}(s, s', R)$ is a compact subspace and by hypothesis for each $i \in I$, there exists $s_{0,i} > 1$ such that $C_i \subset S_i^{(4)}(s_{0,i}, s', R)$. It follows that $C \subset S^{(4)}(s_0, s', R)$ with $s_0 := \min_i s_{0,i}$.

We now proceed to show the cofinality of $(S_i^{(4)}(s, s', R))_{s>1}$ in the family of compact subspaces of $\bigcup_{s>1} S_i^{(4)}(s, s', R)$ using Lemma 3.2.9. For each $i \in I$, consider the continuous function $\theta_i : \bigcup_{s>1} S_i^{(4)}(s, s', R) \rightarrow \mathbb{R}_+ \cup \{\infty\}$ defined by

$$\theta_i(x) := \begin{cases} |F_i(x)| & \text{if } \lambda_i = 0, \\ \frac{|F_i(x)|}{\lambda_i |G_i(x)|} & \text{if } \lambda_i > 0. \end{cases}$$

It is an easy exercise to check that the functions θ_i are positive and satisfy (3.2.10) with the map λ defined by $\lambda(\theta_0) = \theta_0$. This completes the proof of Claim 5a. □

Claim 5 follows from Claim 5a in exactly the same manner as in the proof of Claim 1 from Claim 1a. □

Claim 6. Let $R > 0$. Then there exists $\varepsilon'_0(R) > 0$ such that for each $0 < \varepsilon' \leq \varepsilon'_0(R)$, there exists $\varepsilon_0(\varepsilon') > 1$ such that

$$(3.2.32) \quad H^*(\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)) \cong H^*(S'_\sigma(\varepsilon, \varepsilon', R))$$

for all $1 < \varepsilon \leq \varepsilon_0(\varepsilon')$.

Proof of Claim 6. It follows from (3.2.27) that

$$(3.2.33) \quad H^*(\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)) \cong \varprojlim_{s'} \varprojlim_s H^*(S'_\sigma(s, s', R)).$$

It follows from Lemma 3.2.7(2) that there exists $\varepsilon'_0(R)$ such that for all $0 < s'_2 \leq s'_1 \leq \varepsilon'_0(R)$, the inclusion map

$$\bigcup_{s>1} S'_\sigma(s, s'_1, R) \hookrightarrow \bigcup_{s>1} S'_\sigma(s, s'_2, R)$$

induces an isomorphism

$$H^*\left(\bigcup_{s>1} S'_\sigma(s, s'_2, R)\right) \rightarrow H^*\left(\bigcup_{s>1} S'_\sigma(s, s'_1, R)\right).$$

Hence

$$(3.2.34) \quad \varprojlim_{s'} H^*\left(\bigcup_{s>1} S'_\sigma(s, s', R)\right) \cong H^*\left(\bigcup_{s>1} S'_\sigma(s, \varepsilon', R)\right)$$

for all $0 < \varepsilon' \leq \varepsilon'_0(R)$. Moreover, it follows from (3.2.30) that

$$(3.2.35) \quad H^*\left(\bigcup_{s>1} S'_\sigma(s, \varepsilon', R)\right) \cong \varprojlim_s H^*(S'_\sigma(s, \varepsilon', R)).$$

Hence, from (3.2.33)–(3.2.35) we get an isomorphism

$$(3.2.36) \quad H^*(\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)) \cong \varprojlim_s H^*(S'_\sigma(s, \varepsilon', R))$$

It again follows from Lemma 3.2.7(1) that for each fixed s' , and hence for $s' = \varepsilon'$, there exists $\varepsilon_0(\varepsilon') > 1$ such that for all $1 < s_2 \leq s_1 \leq \varepsilon_0(\varepsilon')$, the inclusion map $S'_\sigma(s_1, \varepsilon', R) \hookrightarrow S'_\sigma(s_2, \varepsilon', R)$ induces an isomorphism

$$H^*(S'_\sigma(s_2, \varepsilon', R)) \rightarrow H^*(S'_\sigma(s_1, \varepsilon', R)),$$

which implies that

$$(3.2.37) \quad \varprojlim_s H^*(S'_\sigma(s, \varepsilon', R)) \cong H^*(S'_\sigma(\varepsilon, \varepsilon', R))$$

for all $1 < \varepsilon \leq \varepsilon_0(\varepsilon')$. Claim 6 follows from (3.2.36) and (3.2.37). □

We now return to the proof of Proposition 3.2.6. Using Lemma 3.2.7(6), we find that there exists $R_0 > 0$ such that for all $R \geq R_0$,

$$(3.2.38) \quad H^*(\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)) \cong H^*(\tilde{\mathcal{R}}(\sigma, \bar{w})).$$

Fix $R \geq R_0$. It follows from (3.2.32) that there exists $\varepsilon'_0(R) > 0$ such that for each $0 < \varepsilon' \leq \varepsilon'_0(R)$, there exists $\varepsilon_0(\varepsilon') > 1$ such that for all $1 < \varepsilon \leq \varepsilon_0(\varepsilon')$,

$$(3.2.39) \quad H^*(\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)) \cong H^*(S_\sigma(\varepsilon, \varepsilon', R)).$$

Fix ε' and ε satisfying $0 < \varepsilon' \leq \varepsilon'_0(R)$ and $1 < \varepsilon \leq \varepsilon_0(\varepsilon')$.

Now it follows from (3.2.20) that there exists $\delta'_0(\varepsilon, \varepsilon', R) > 0$ such that for each $0 < \delta' \leq \delta'_0(\varepsilon, \varepsilon', R)$, there exists $\delta_0(\delta') > 1$ such that for all $1 < \delta \leq \delta_0(\delta')$,

$$H^*(S'_\sigma(\varepsilon, \varepsilon', R)) \cong H^*(S_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)).$$

Choose δ', δ satisfying $0 < \delta' \leq \delta'_0(\varepsilon, \varepsilon', R)$ and $1 < \delta \leq \delta_0(\delta')$. It is now clear that with the above choices of $R, \varepsilon', \varepsilon, \delta', \delta$, we have

$$H^*(\tilde{\mathcal{R}}(\sigma, \bar{w})) \cong H^*(S_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)).$$

This concludes the proof of Proposition 3.2.6. □

Before stating the next proposition we introduce some notation. As in the hypothesis of Proposition 3.2.6, let $V \subset \mathbb{A}_K^N$ and $W \subset \mathbb{A}_K^M$ be closed affine subvarieties and $\phi(\cdot, \cdot)$ a formula in disjunctive normal form without negations and with atoms of the form $|F| \leq \lambda \cdot |G|$ where $F, G \in K[X_1, \dots, X_N, Y_1, \dots, Y_M]$.

For $\delta, \varepsilon > 1$ and $\delta', \varepsilon' > 0$ let

$$S''_\sigma(\delta, \delta', \varepsilon, \varepsilon', R) := \bigcap_{i, \sigma(i)=1} \text{Tube}^o_{V, \phi(\cdot, w_i)}(\delta, \delta', R) - \bigcup_{i, \sigma(i)=0} \text{Tube}^c_{V, \phi(\cdot, w_i)}(\varepsilon, \varepsilon').$$

It follows from the definition that for all $\delta, \varepsilon > 1$ and $\delta', \varepsilon' > 0$,

$$S''_\sigma(\delta, \delta', \varepsilon, \varepsilon', R) \subset S_\sigma(\delta, \delta', \varepsilon, \varepsilon', R).$$

Note that the sets $S''_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)$ and $S_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)$ shrink as δ, δ' decreases, and they grow with decreasing $\varepsilon, \varepsilon'$. More precisely, for all $\delta_i, \delta'_i, \varepsilon_i, \varepsilon'_i, i = 1, 2$, satisfying $1 < \delta_1 < \delta_2, 0 < \delta'_1 < \delta'_2, 1 < \varepsilon_2 < \varepsilon_1, 0 < \varepsilon'_2 < \varepsilon'_1$, we have

$$\begin{aligned} S_\sigma(\delta_1, \delta'_1, \varepsilon_1, \varepsilon'_1, R) &\subset S_\sigma(\delta_2, \delta'_2, \varepsilon_2, \varepsilon'_2, R), \\ S''_\sigma(\delta_1, \delta'_1, \varepsilon_1, \varepsilon'_1, R) &\subset S''_\sigma(\delta_2, \delta'_2, \varepsilon_2, \varepsilon'_2, R). \end{aligned}$$

Proposition 3.2.40. *With notation as above, for all $\delta, \delta', \varepsilon, \varepsilon' \in \mathbb{R}_+$ satisfying $0 < \delta - 1 < \delta' < \varepsilon - 1 < \varepsilon'$, every connected component of $S''_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)$ is a connected component of the semi-algebraic set*

$$(3.2.41) \quad U_{\phi, \delta, \delta', \varepsilon, \varepsilon', R} := \bigcap_{1 \leq i \leq n} (U_{i, \varepsilon, \varepsilon', R} \cap U_{i, \delta, \delta', R}),$$

where for $1 \leq i \leq n$ and $t > 1, t' > 0$,

$$U_{i, t, t', R} := \text{Cube}_V(R) \setminus \text{TubeBoundary}^c_{V, \phi(\cdot, w_i)}(t, t', R).$$

Before proving Proposition 3.2.40, we note that Propositions 3.2.40 and 3.2.6 imply

Proposition 3.2.42. *For each $\bar{w} \in W(K)^n$, there exist $\delta > 1, \delta' > 0, \varepsilon > 1, \varepsilon' > 0$, and $R > 0$ such that for each $\sigma \in \{0, 1\}^n$ and $0 \leq i < k$,*

$$(3.2.43) \quad \sum_{\sigma \in \{0, 1\}^n} b_i(\tilde{\mathcal{R}}(\sigma, \bar{w})) \leq b_i(U_{\phi, \delta, \delta', \varepsilon, \varepsilon', R}).$$

Proof. By Proposition 3.2.6 and using the notation of its proof there exist an $R > 0$, an $\varepsilon'(R) > 0$, and for each $0 < \varepsilon' < \varepsilon'(R)$, there exists an $\varepsilon_0(\varepsilon') > 1$ such that

$$(3.2.44) \quad H^*(\tilde{\mathcal{R}}(\sigma, \bar{w}) \cap \text{Cube}_V(R)) \cong H^*(S'_\sigma(\varepsilon, \varepsilon', R))$$

for all $1 < \varepsilon \leq \varepsilon_0(\varepsilon')$. Fix ε'_i and ε_i ($i = 1, 2$) satisfying $0 < \varepsilon'_1 < \varepsilon'_2 \leq \varepsilon'_0(R)$ and $1 < \varepsilon_1 < \varepsilon_2 \leq \min(\varepsilon_0(\varepsilon'_1), \varepsilon_0(\varepsilon'_2))$. Recall that it follows from (3.2.20) that there exists $\delta'_0(\varepsilon_i, \varepsilon'_i, R) > 0$ such that for each $0 < \delta' \leq \delta'_0(\varepsilon_i, \varepsilon'_i, R)$, there exists $\delta_0^{(i)}(\delta') > 1$ (depending on δ' and $\delta'_0(\varepsilon_i, \varepsilon'_i, R)$) such that for all $1 < \delta \leq \delta_0^{(i)}(\delta')$,

$$H^*(S'_\sigma(\varepsilon_i, \varepsilon'_i, R)) \cong H^*(S_\sigma(\delta, \delta', \varepsilon_i, \varepsilon'_i, R)).$$

Let

$$0 < \delta' \leq \min(\delta'_0(\varepsilon_1, \varepsilon'_1, R), \delta'_0(\varepsilon_2, \varepsilon'_2, R)), \quad 1 < \delta \leq \min(\delta_0^{(1)}(\delta'), \delta_0^{(2)}(\delta')).$$

With the above choices of $R, \varepsilon'_i, \varepsilon_i, \delta', \delta$, we have

$$H^*(\tilde{\mathcal{R}}(\sigma, \bar{w})) \cong H^*(S_\sigma(\delta, \delta', \varepsilon_i, \varepsilon'_i, R)).$$

On the other hand, let $T_i := S_\sigma(\delta, \delta', \varepsilon_i, \varepsilon'_i, R)$ and $T''_i := S''_\sigma(\delta, \delta', \varepsilon_i, \varepsilon'_i, R)$. Then $T_2 \subset T''_1 \subset T_1$, and by the previous remarks the natural map

$$H^i(T_1) \rightarrow H^i(T_2)$$

is an isomorphism. On the other hand, this map factors through $H^i(T''_1)$ and therefore the natural map

$$H^i(T''_1) \rightarrow H^i(T_1)$$

is surjective. It follows that $b_i(T_1) \leq b_i(T''_1)$. Since the connected components of the T''_1 (as σ varies) are connected components of $U_{\phi, \delta, \delta', \varepsilon, \varepsilon', R}$ (by Proposition 3.2.40), the inequality (3.2.43) follows immediately. \square

Proof of Proposition 3.2.40. Recall that ϕ is a disjunction of the formulas $\phi_h, h \in H$, where H is a finite set, and each ϕ_h is a conjunction of weak inequalities $|F_{hj}| \leq \lambda_{hj} \cdot |G_{hj}|, j \in J_h$, where J_h is a finite set. As before for each i we let $F_{ihj} := F_{hj}(\cdot, w_i), G_{ihj} := G_{hj}(\cdot, w_i)$.

We first observe that $S''_\sigma(\delta, \delta', \varepsilon, \varepsilon', R) \subset U_{\phi, \delta, \delta', \varepsilon, \varepsilon', R}$. To see this, for $t' > 0, t > 1$, and $i \in [1, n]$, let $\theta_{i,t,t'} : B_{\mathbb{F}}(V) \rightarrow \mathbb{R}$ be the continuous function defined by

$$(3.2.45) \quad \theta_{i,t,t'}(x) := \max_{h \in H} \min_{j \in J_h} \mu_{i,h,j,t,t'}(x),$$

where

$$\mu_{i,h,j,t,t'}(x) := \begin{cases} t' - |F_{ihj}(x)| & \text{if } \lambda_{hj} = 0, \\ \max(\lambda_j \cdot t \cdot |G_{ihj}(x)| - |F_{ihj}(x)|, \min(t' - |F_{ihj}(x)|, t' - |G_{ihj}(x)|)) & \text{if } \lambda_{hj} > 0. \end{cases}$$

The formula defining $\theta_{i,t,t'}$ might seem formidable at first glance, but becomes easier to understand with the observation that each occurrence of \max and \min in (3.2.45) corresponds to an occurrence of respectively \bigvee and \bigwedge in the formula $\phi^o(\cdot; T, T')$ (see Notation 3.2.2). With this observation, and the obvious facts that for any $A \subset \mathbb{R}$,

$$\bigvee_{a \in A} (a > 0) \iff \max_{a \in A} a > 0, \quad \bigwedge_{a \in A} (a > 0) \iff \min_{a \in A} a > 0,$$

it is easy to verify that

$$\begin{aligned} x \in \text{Tube}_{V,\phi(\cdot,w_i)}^o(\delta, \delta') &\iff \theta_{i,\delta,\delta'}(x) > 0, \\ x \in \text{Tube}_{V,\phi(\cdot,w_i)}^c(\delta, \delta') &\iff \theta_{i,\delta,\delta'}(x) \geq 0, \end{aligned}$$

and finally that for any $R > 0$,

(3.2.46)

$$x \in \text{TubeBoundary}_{V,\phi(\cdot,w_i)}^c(\delta, \delta', R) \iff x \in \text{Cube}_V(R) \wedge (\theta_{i,\delta,\delta'}(x) = 0).$$

Now let $x \in S''_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)$. Then, for each i with $\sigma(i) = 1$, we have $x \in \text{Tube}_{V,\phi(\cdot,w_i)}^o(\delta, \delta', R)$, and hence $x \notin \text{TubeBoundary}_{V,\phi(\cdot,w_i)}^c(\delta, \delta', R)$.

Using the fact that $\delta' < \varepsilon'$ and $\delta < \varepsilon$, one can also check that $\theta_{i,\delta,\delta'}(x) > 0$ implies $\theta_{i,\varepsilon,\varepsilon'}(x) > 0$ as well. This in turn implies that

$$x \in \text{Tube}_{V,\phi(\cdot,w_i)}^o(\delta, \delta', R) \implies x \notin \text{TubeBoundary}_{V,\phi(\cdot,w_i)}^c(\varepsilon, \varepsilon', R).$$

Hence,

$$x \notin \text{TubeBoundary}_{V,\phi(\cdot,w_i)}^c(\delta, \delta', R) \cup \text{TubeBoundary}_{V,\phi(\cdot,w_i)}^c(\varepsilon, \varepsilon', R)$$

for all i with $\sigma(i) = 1$. In particular, $x \in U_{i,\varepsilon,\varepsilon',R} \cap U_{i,\delta,\delta',R}$.

We now consider the case of i such that $\sigma(i) = 0$. Then $x \in \text{Cube}_V(R) \setminus \text{Tube}_{V,\phi(\cdot,w_i)}^c(\varepsilon, \varepsilon', R)$, and hence $x \notin \text{TubeBoundary}_{V,\phi(\cdot,w_i)}^c(\varepsilon, \varepsilon', R)$.

Also, if $x \notin \text{Tube}_{V,\phi(\cdot,w_i)}^c(\varepsilon, \varepsilon', R)$, then $x \notin \text{TubeBoundary}_{V,\phi(\cdot,w_i)}^c(\delta, \delta', R)$, since clearly

$$\text{TubeBoundary}_{V,\phi(\cdot,w_i)}^c(\delta, \delta', R) \subset \text{Tube}_{V,\phi(\cdot,w_i)}^c(\varepsilon, \varepsilon', R).$$

Hence,

$$x \notin \text{TubeBoundary}_{V,\phi(\cdot,w_i)}^c(\delta, \delta', R) \cup \text{TubeBoundary}_{V,\phi(\cdot,w_i)}^c(\varepsilon, \varepsilon', R)$$

for all i with $\sigma(i) = 0$. Combining everything, we have $x \in U_{\phi,\delta,\delta',\varepsilon,\varepsilon',R}$.

Now let C be a connected component of $S''_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)$, and D be the connected component of $U_{\phi,\delta,\delta',\varepsilon,\varepsilon',R}$ containing C . We claim that $D = C$. Let $x \in D$, and let y be any point of C . Since $y \in D$ and D is path connected, there exists a path $\gamma : [0, 1] \rightarrow D$ with $\gamma(0) = y$, $\gamma(1) = x$, and $\gamma([0, 1]) \subset D$. We claim that $\gamma([0, 1]) \subset S''_\sigma(\delta, \delta', \varepsilon, \varepsilon', R)$, which immediately implies $D = C$.

We first show that for each i with $\sigma(i) = 1$, $\gamma([0, 1]) \subset \text{Tube}_{V, \phi(\cdot, w_i)}^o(\delta, \delta', R)$. For each i with $\sigma(i) = 1$, consider the continuous function $\theta_i : [0, 1] \rightarrow \mathbb{R}$ defined by

$$\theta_i(t) := \theta_{i, \delta, \delta'}(\gamma(t)).$$

It follows from (3.2.46) that $\theta_i(t) = 0$ implies that

$$\gamma(t) \in \text{TubeBoundary}_{V, \phi(\cdot, w_i)}^c(\delta, \delta', R).$$

Consequently, since

$$\gamma([0, 1]) \subset \text{Cube}_V(R) \setminus \text{TubeBoundary}_{V, \phi(\cdot, w_i)}^c(\delta, \delta', R)$$

for each i , θ_i cannot vanish anywhere on $[0, 1]$. Also $\theta_i(t) > 0$ if and only if $\gamma(t) \in \text{Tube}_{V, \phi(\cdot, w_i)}^o(\delta, \delta', R)$. Since $\gamma(0) = y \in S''_{\sigma, \delta, \delta', \varepsilon, \varepsilon', R}$, this implies that $\theta_i(0) > 0$, hence $\theta_i(t) > 0$ for each $t \in [0, 1]$, and

$$\gamma([0, 1]) \subset \bigcap_{i, \sigma(i)=1} \text{Tube}_{V, \phi(\cdot, w_i)}^o(\delta, \delta', R).$$

Finally, we show that

$$\gamma([0, 1]) \subset \bigcap_{i, \sigma(i)=0} (\text{Cube}_V(R) \setminus \text{Tube}_{V, \phi(\cdot, w_i)}^c(\varepsilon, \varepsilon', R)).$$

For each i with $\sigma(i) = 0$, consider the continuous function $\mu_i : [0, 1] \rightarrow \mathbb{R}$ defined by

$$\mu_i(t) := -\theta_{i, \varepsilon, \varepsilon'}(\gamma(t)).$$

Notice that $\mu_i(t) = 0$ implies $\gamma(t) \in \text{TubeBoundary}_{V, \phi(\cdot, w_i)}^c(\varepsilon, \varepsilon', R)$, and since $\gamma([0, 1]) \subset \text{Cube}_V(R) \setminus \text{TubeBoundary}_{V, \phi(\cdot, w_i)}^c(\varepsilon, \varepsilon', R)$ for each i , μ_i cannot vanish anywhere on $[0, 1]$. Moreover, $\mu_i(t) > 0$ if and only if $\gamma(t) \in \text{Cube}_V(R) \setminus \text{Tube}_{V, \phi(\cdot, w_i)}^c(\varepsilon, \varepsilon', R)$. Since $\gamma(0) = y \in S''_{\sigma}(\delta, \delta', \varepsilon, \varepsilon', R)$, this implies that $\mu_i(0) > 0$, hence $\mu_i(t) > 0$ for each $t \in [0, 1]$, and so

$$\gamma([0, 1]) \subset \bigcap_{i, \sigma(i)=0} (\text{Cube}_V(R) \setminus \text{Tube}_{V, \phi(\cdot, w_i)}^c(\varepsilon, \varepsilon', R)).$$

This proves that $D = C$. □

Let $X \subset V$ be a definable subset where V is an affine variety of dimension k , and let U_1, \dots, U_n be open semi-algebraic subsets of $B_{\mathbb{F}}(X)$. For $J \subset [1, n]$, we denote $U^J := \bigcup_{j \in J} U_j$ and $U_J := \bigcap_{j \in J} U_j$. We have the following proposition, which is very similar to [BPR06, Proposition 7.33(ii)].

Proposition 3.2.47. *With notation as above, for each $0 \leq i \leq k = \dim(V)$,*

$$b_i(U_{[1, n]}) \leq \sum_{j=1}^{k-i} \sum_{J \subset [1, n], \text{card}(J)=j} b_{i+j-1}(U^J) + \binom{n}{k-i} b_k(B_{\mathbb{F}}(V)).$$

Proof. We first prove the claim when $n = 1$. If $0 \leq i \leq k - 1$, the claim is

$$b_i(U_1) \leq b_i(U_1) + b_k(B_{\mathbf{F}}(V)),$$

which is clear. If $i = k$, the claim is $b_k(U_1) \leq b_k(B_{\mathbf{F}}(V))$, which is true by Corollary A.6(d).

The claim is now proved by induction on n . Assume that the induction hypothesis holds for all $n - 1$ open semi-algebraic subsets of $B_{\mathbf{F}}(V)$, and for all $0 \leq i \leq k$.

It follows from the standard Mayer–Vietoris sequence (see Properties A.1.1(5)) that

$$(3.2.48) \quad b_i(U_{[1,n]}) \leq b_i(U_{[1,n-1]}) + b_i(U_n) + b_{i+1}(U_{[1,n-1]} \cup U_n).$$

Applying the induction hypothesis to the set $U_{[1,n-1]}$, we deduce that

$$(3.2.49) \quad \begin{aligned} b_i(U_{[1,n-1]}) &\leq \sum_{j=1}^{k-i} \sum_{J \subset [1,n-1], \text{card}(J)=j} b_{i+j-1}(U^J) \\ &\quad + \binom{n-1}{k-i} b_k(B_{\mathbf{F}}(V)). \end{aligned}$$

Next, applying the induction hypothesis to the set

$$U_{[1,n-1]} \cup U_n = \bigcap_{1 \leq j \leq n-1} (U_j \cup U_n),$$

we get

$$(3.2.50) \quad \begin{aligned} b_{i+1}(U_{[1,n-1]} \cup U_n) &\leq \sum_{j=1}^{k-i-1} \sum_{J \subset [1,n-1], \text{card}(J)=j} b_{i+j}(U^{J \cup \{n\}}) \\ &\quad + \binom{n-1}{k-i-1} b_k(B_{\mathbf{F}}(V)). \end{aligned}$$

From the inequalities (3.2.48)–(3.2.50) we obtain

$$b_i(U_{[1,n]}) \leq \sum_{j=1}^{k-i} \sum_{J \subset [1,n], \text{card}(J)=j} b_{i+j-1}(U^J) + \binom{n}{k-i} b_k(B_{\mathbf{F}}(V)),$$

which finishes the induction. □

Proof of Theorem 2. Using Proposition 3.2.42 we find that there exist $\delta > 1$, $\delta' > 0$, $\varepsilon > 1$, $\varepsilon' > 0$, $R > 0$ (which we fix for the remainder of the proof) such that for each i , $0 \leq i \leq k$,

$$(3.2.51) \quad \sum_{\sigma \in \{0,1\}^n} b_i(\tilde{\mathcal{R}}(\sigma, \bar{w})) \leq b_i(U_{\phi, \delta, \delta', \varepsilon, \varepsilon', R}).$$

From the definition of $U_{\phi, \delta, \delta', \varepsilon, \varepsilon', R}$ in (3.2.41), we see that $U_{\phi, \delta, \delta', \varepsilon, \varepsilon', R}$ is an intersection of the sets

$$\begin{aligned} & \text{Cube}_V(R) \setminus \text{TubeBoundary}_{V, \phi(\cdot, w_j)}^c(\varepsilon, \varepsilon', R), \\ & \text{Cube}_V(R) \setminus \text{TubeBoundary}_{V, \phi(\cdot, w_j)}^c(\delta, \delta', R), \end{aligned}$$

for $1 \leq j \leq n$.

Now for each $m \geq 1$ and $m', m'' \geq 0$ with $m' + m'' = m$, let

$$\Phi_{m', m''}(\bar{X}, \bar{Y}^{(1)}, \dots, \bar{Y}^{(m)}; s, s', t, t', R) := (\Psi_1 \vee \Psi_2) \wedge (\Psi_3 \wedge \Psi_4),$$

where

$$\begin{aligned} \Psi_1 &:= \bigvee_{1 \leq j \leq m'} (-\phi^c(\bar{X}, \bar{Y}^{(j)}; s, s') \vee \phi^o(\bar{X}, \bar{Y}^{(j)}; s, s')), \\ \Psi_2 &:= \bigvee_{m'+1 \leq j \leq m} (-\phi^c(\bar{X}, \bar{Y}^{(j)}; t, t') \vee \phi^o(\bar{X}, \bar{Y}^{(j)}; t, t')), \\ \Psi_3 &:= \Phi_V(\bar{X}; R), \\ \Psi_4 &:= \bigwedge_{1 \leq j \leq m} \Phi_W(\bar{Y}^{(j)}). \end{aligned}$$

Here $\Phi_{V,R}(\bar{X}; R)$ is a formula such that $\text{Cube}_V(R) = \widetilde{\mathcal{R}}(\Phi_{V,R}, V)$, and $\Phi_W(\bar{Y})$ is a formula such that $B_{\mathbb{F}}(W) = \widetilde{\mathcal{R}}(\Phi_W, V)$.

Denote by $X_{m', m''}$ the definable subset of $V \times \underbrace{W \times \dots \times W}_m \times \mathbb{R}^5$ defined by the formula

$$\Phi_{m', m''}(\bar{X}, \bar{Y}^{(1)}, \dots, \bar{Y}^{(m)}; s, s', t, t', R),$$

and let

$$\pi_{m', m''} : X_{m', m''} \rightarrow \underbrace{W \times \dots \times W}_m \times \mathbb{R}^5$$

denote the projection map. It follows from Theorem A.4 (with $Y = \underbrace{W \times \dots \times W}_m, V$

viewed as a quasi-projective variety in \mathbb{P}^N and $X_{m', m''}$ as above) that the number of homotopy types amongst the semi-algebraic sets

$$B_{\mathbb{F}}(\pi_{m', m''}^{-1}(w'_1, \dots, w'_m, s, s', t, t', R))$$

is finite, and moreover since each such fiber is homotopy equivalent to a finite simplicial complex by Theorem A.5, there exists a finite bound $C_{i, m', m''} \in \mathbb{Z}_{\geq 0}$ such that

$$b_i(B_{\mathbb{F}}(\pi_{m', m''}^{-1}(w'_1, \dots, w'_m, s, s', t, t', R))) \leq C_{i, m', m''}$$

for all $(w'_1, \dots, w'_m) \in W(K)^m$ and $s, s', t, t', R \in \mathbb{R}$.

Let

$$(3.2.52) \quad C_{i, m} := \max_{\substack{m', m'' \geq 0 \\ m' + m'' = m}} C_{i, m', m''}.$$

Note that $C_{i, m}$ depends only on V and ϕ .

It follows from Notation 3.2.2 that for each $1 \leq j \leq n$, the semi-algebraic set

$$\tilde{\mathcal{R}}((\neg(\phi^c(\bar{X}, w_j; \cdot, \cdot) \vee \phi^o(\bar{X}, w_j; \cdot, \cdot)), V) \cap \text{Cube}_V(R)$$

is equal to

$$\text{Cube}_V(R) \setminus \text{TubeBoundary}_{V, \phi(\cdot, w_j)}^c(\cdot, \cdot, R).$$

It follows that for any

$$J' = (j'_1, \dots, j'_{\text{card}(J')}), \quad J'' = (j''_1, \dots, j''_{\text{card}(J'')}) \subset [1, n]$$

with $J' \cap J'' = \emptyset$, the semi-algebraic set

$$\tilde{\mathcal{R}}(\Phi_{\text{card}(J'), \text{card}(J'')}(\cdot, w_{j'_1}, \dots, w_{j'_{\text{card}(J')}}), w_{j''_1}, \dots, w_{j''_{\text{card}(J'')}}; \varepsilon, \varepsilon', \delta, \delta', R)$$

is the union of

$$\bigcup_{j \in J'} (\text{Cube}_V(R) \setminus \text{TubeBoundary}_{V, \phi(\cdot, w_j)}^c(\varepsilon, \varepsilon', R))$$

and

$$\bigcup_{j \in J''} (\text{Cube}_V(R) \setminus \text{TubeBoundary}_{V, \phi(\cdot, w_j)}^c(\delta, \delta', R)).$$

Also, since each m -ary union amongst the semi-algebraic sets

$$\text{Cube}_V(R) \setminus \text{TubeBoundary}_{V, \phi(\cdot, w_j)}^c(\varepsilon, \varepsilon', R),$$

$$\text{Cube}_V(R) \setminus \text{TubeBoundary}_{V, \phi(\cdot, w_j)}^c(\delta, \delta', R)$$

is clearly homeomorphic to one of the sets $B_{\mathbf{F}}(\pi_{m', m''}^{-1}(w'_1, \dots, w'_m, s, s', t, t', R))$, $m' + m'' = m$, $(w'_1, \dots, w'_m) \in W(K)^m$, $s, s', t, t', R \in \mathbb{R}$, the i -th Betti number of every such union is bounded by $C_{i, m}$.

It now follows from (3.2.52) and Proposition 3.2.47 that

$$\sum_{\sigma \in \{0, 1\}^n} b_i(\tilde{\mathcal{R}}(\sigma, \bar{w})) \leq \sum_{j=1}^{k-i} \binom{2n}{j} C_{i+j-1, j} + \binom{2n}{k-i} b_k(B_{\mathbf{F}}(V)).$$

The theorem follows after noticing that $\binom{2n}{j} \leq (2n)^j$ for all $n, j \geq 0$. □

3.3. Proof of Theorem 1

We need a couple of preliminary results of a set-theoretic nature, starting with the following observation.

Observation 3.3.1. *Let Y, Y', V, V', W, W' be sets such that $Y \subset V \times W, Y' \subset V' \times W', V \subset V', W \subset W',$ and $Y' \cap (V \times W) = Y.$ Then, for every $n > 0,$*

$$\chi_{Y, V, W}(n) \leq \chi_{Y', V', W}(n).$$

Proof. Note that a 0/1 pattern is realized by the tuple $(Y_{w_1}, \dots, Y_{w_n})$ in V only if it is realized by the tuple $(Y'_{w_1}, \dots, Y'_{w_n})$ in $V'.$ This follows from the fact that $Y' \cap (V \times W) = Y,$ and therefore for all $w \in W, Y'_w \cap V = Y_w.$ □

Let V, W be sets, I a finite set, and for each $\alpha \in I,$ let X_α be a subset of $V \times W.$ Let $i_\alpha : X_\alpha \hookrightarrow V \times W$ denote the inclusion map. Suppose that X is a subset of $V \times W$ obtained as a Boolean combination of the X_α 's. Let $W' := \coprod_{\alpha \in I} W,$ and for $\alpha \in I$ let $j_\alpha : W \hookrightarrow W'$ denote the canonical inclusion. Let $X' := \bigcup_{\alpha \in I} \text{Im}((1_V \times j_\alpha) \circ i_\alpha) \subset V \times W'.$ With this notation we have the following proposition.

Proposition 3.3.2.

$$\chi_{X, V, W}(n) \leq \chi_{X', V, W'}(\text{card}(I) \cdot n).$$

Proof. For $v \in V$ and $S \subset W$ (resp. $S' \subset W'$) we set $S_v := S \cap X_v$ (resp. $S'_v := S' \cap X'_v$). Let $\bar{w} \in W^n.$ We claim that for $v, v' \in V,$

$$\begin{aligned} \chi_{X, V, W; n}(v, \bar{w}) \neq \chi_{X, V, W; n}(v', \bar{w}) \\ \implies \chi_{X', V, W'; \text{card}(I) \cdot n}(v, j_n(\bar{w})) \neq \chi_{X', V, W'; \text{card}(I) \cdot n}(v', j_n(\bar{w})), \end{aligned}$$

where $j_n : W^{[1, n]} \rightarrow W'^{I \times [1, n]}$ is defined by

$$j_n(w_1, \dots, w_n)_{(\alpha, i)} := j_\alpha(w_i).$$

To prove the claim first observe that since $\chi_{X, V, W; n}(v, \bar{w}) \neq \chi_{X, V, W; n}(v', \bar{w}),$ there exists $i \in [1, n]$ such that $v \in X_{w_i} \Leftrightarrow v' \notin X_{w_i}.$

Since X is a Boolean combination of the $X_\alpha, \alpha \in I,$ there must exist $\alpha \in I$ such that $v \in (X_\alpha)_{w_i} \Leftrightarrow v' \notin (X_\alpha)_{w_i}.$ It now follows from the definition of X', W' that $\chi_{X', V, W'; \text{card}(I) \cdot n}(v, j_n(\bar{w})) \neq \chi_{X', V, W'; \text{card}(I) \cdot n}(v', j_n(\bar{w})).$ This implies that

$$\text{card}(\chi_{X, V, W; n}(V, \bar{w})) \leq \text{card}(\chi_{X', V, W'; \text{card}(I) \cdot n}(V, j_n(\bar{w}))).$$

It follows immediately that $\chi_{X, V, W}(n) \leq \chi_{X', V, W'}(\text{card}(I) \cdot n).$ □

Proof of Theorem 1. We make two reductions. We first claim that it suffices to prove the theorem in the case of an algebraically closed complete valued field of rank 1, i.e. the value group is a subgroup of the multiplicative group $\mathbb{R}_+.$ Secondly, we claim that we can assume without loss of generality that the formula ϕ is in disjunctive normal form without negations and with atoms of the form $|F| \leq \lambda \cdot |G|.$

Reduction to complete algebraically closed field of rank 1. The theory of algebraically closed valued fields in the two-sorted language \mathcal{L} becomes complete once we fix the characteristic of the field and that of the residue field. Moreover, for each such characteristic pair $(0, 0), (0, p),$ or (p, p) (p a prime) there exists a model $(K; \Gamma)$ of the theory of algebraically closed valued field such that the value group is a multiplicative subgroup

of \mathbb{R}_+ (i.e. of rank 1) and K is complete. It follows by a standard transfer argument that it suffices to prove the theorem for such a model.

Reduction to the case of disjunctive normal form without negations and with atoms of the form $|F| \leq \lambda \cdot |G|$. We now observe that it suffices to prove the theorem in the case when the formula ϕ is equivalent to a formula in disjunctive normal form without negations with atoms of the form $|F| \leq \lambda \cdot |G|$. Furthermore, using the first reduction, we may assume that the value group is \mathbb{R}_+ and K is an algebraically closed complete valued field. In particular, we assume that the atoms of ϕ are of the form $|F| \leq \lambda \cdot |G|$ with $\lambda \in \mathbb{R}_+$ and $F, G, \in K[\bar{X}, \bar{Y}]$. Let $(\phi_\alpha)_{\alpha \in I}$ be the finite tuple of atomic formulas appearing in ϕ . Denote

$$\phi'' := \left(\bigvee_{\alpha \in I} (\phi_\alpha(\bar{X}, \bar{Y}^{(\alpha)}) \wedge (|Z_\alpha - 1| = 0)) \right) \wedge \bigvee_{\alpha \in I} \theta_\alpha((Z_\alpha)_{\alpha \in I}),$$

where $\theta_\alpha((Z_\alpha)_{\alpha \in I})$ is the closed formula

$$(|Z_\alpha - 1| = 0) \wedge \bigwedge_{\beta \neq \alpha} (|Z_\beta| = 0).$$

Note that ϕ'' is equivalent to a formula in disjunctive normal form without negations and with atoms of the form $|F| \leq \lambda \cdot |G|$.


Let $X_\alpha := \mathcal{R}(\phi_\alpha, V \times W)(K)$ and $X := \mathcal{R}(\phi, V \times W)(K)$. Then X is a Boolean combination of the X_α 's and we can consider $X' \subset V(K) \times W(K)'$ where X' and $W(K)'$ are defined as in Proposition 3.3.2. In particular, we let $\pi_1 : X' \rightarrow V(K)$ and $\pi'_1 : X' \rightarrow W(K)'$ denote the natural projection maps. Similarly, we let

$$\pi''_2 : \mathcal{R}(\phi'', V \times W \times \mathbb{A}^{|I|})(K) \rightarrow W(K) \times \mathbb{A}^{|I|}(K)$$


and

$$\pi''_1 : \mathcal{R}(\phi'', V \times W \times \mathbb{A}^{|I|})(K) \rightarrow V(K)$$

denote the natural projection maps. Note that the diagram



is isomorphic to the diagram



By isomorphism, we mean that there are natural bijections $\mathcal{R}(\phi'', V \times W \times \mathbb{A}^{|I|})(K) \rightarrow X'$ and $\text{Im}(\pi''_2) \rightarrow \text{Im}(\pi'_2)$ making the resulting morphism of diagrams above commute (with the identity as the map on $V(K)$).

Using Proposition 3.3.2, we get

$$\chi_{\mathcal{R}(\phi, (V \times W))(K), V(K), W(K)}(n) \leq \chi_{X', V(K), (W(K))'}(\text{card}(I) \cdot n),$$

and the right hand side clearly equals

$$\chi_{\mathcal{R}(\phi'', (V \times W \times \mathbb{A}^{|I|}))(K), V(K), W(K) \times \mathbb{A}^{|I|}(K)}(\text{card}(I) \cdot n).$$

So it suffices to prove that there exists a constant C (depending only on V and ϕ) such that for all n ,

$$\chi_{\mathcal{R}(\phi'', (V \times W \times \mathbb{A}^{|I|}))(K), V(K), W(K) \times \mathbb{A}^{|I|}(K)}(n) \leq C \cdot n^{\dim(V)}.$$

This shows that we can assume that ϕ is equivalent to a formula in disjunctive normal form without negations and with atoms of the form $|F| \leq \lambda \cdot |G|$.

We now use the special case of Theorem 2 obtained by setting $i = 0$. In that case, $b_0(\tilde{\mathcal{R}}(\sigma, \bar{w}))$ is the number of connected components, which is at least 1 as soon as $\tilde{\mathcal{R}}(\sigma, \bar{w})$ is non-empty. Now use Observation 3.3.1 with $V' := B_{\mathbf{F}}(V)$, $Y' := \bigcup_{w \in W(K)} (\tilde{\mathcal{R}}(\phi(\cdot, w), V) \times \{w\})$ and $Y := \mathcal{R}(\phi, (V \times W))(K)$, noting that there exists a canonical injective map $\iota : V(K) \hookrightarrow B_{\mathbf{F}}(V)$ such that for each $w \in W(K)$ the following diagram of injective maps commutes:

$$\begin{array}{ccc} V(K) & \xrightarrow{\iota_V} & B_{\mathbf{F}}(V) \\ \uparrow & & \uparrow \\ \mathcal{R}(\phi(\cdot, w), V)(K) & \longrightarrow & \tilde{\mathcal{R}}(\phi(\cdot, w), V) \end{array}$$

This finishes the proof. □

3.4. Proof of Corollary 1

Corollary 1 follows immediately from Theorem 1 and the following proposition, which is well known, but whose proof we include for the sake of completeness.

Proposition 3.4.1. *Suppose that there exists a constant $C > 0$ such that for all $n > 0$, $\chi_{X, V, W}(n) \leq C \cdot n^k$. Then $\text{vcd}(X, V, W) \leq k$.*

Proof. Notice that for $v \in V$ and $w \in W$, $w \in X_v \Leftrightarrow v \in X_w$. Let $\mathcal{S} = \{X_v \mid v \in V\}$, $A = \{w_1, \dots, w_n\} \subset W$, and $I \subset [1, n]$. For $v \in V$, we have $w_i \in X_v$ for all $i \in I$; $w_i \notin X_v$ for all $i \in [1, n] \setminus I$ if and only if $v \in X_{w_i}$ for all $i \in I$; $v \notin X_{w_i}$ for all $i \in [1, n] \setminus I$. This implies that

$$\text{card}(\{A \cap Y \mid Y \in \mathcal{S}\}) = \chi_{X, V, W; n}(V, \bar{w}) \leq C \cdot n^k.$$

The proposition now follows from Definition 1.1.2. □

Appendix

A.1. Review of singular cohomology

In this section we recall some basic statements about singular cohomology groups which are used throughout this article. These facts are all standard and we refer the reader to [Spa66] for their proofs.

Given any topological space X , one can associate to X the singular cohomology groups $H^i(X, \mathbb{Q})$ (for $i \geq 0$) which satisfy the following general properties (see for example [Spa66, pp. 238–240]):

- Properties A.1.1.** (1) *The $H^i(X, \mathbb{Q})$ are \mathbb{Q} -vector spaces. If X is a finite simplicial complex of dimension n , then each $H^i(X, \mathbb{Q})$ is finite-dimensional, and moreover $H^i(X, \mathbb{Q}) = 0$ for all $i > n$.*
- (2) *The singular cohomology groups are contravariant and homotopy invariant, i.e. a continuous morphism $f : X \rightarrow Y$ induces a linear map $f^* : H^i(Y, \mathbb{Q}) \rightarrow H^i(X, \mathbb{Q})$, and if f is a homotopy equivalence, then the induced map f^* is an isomorphism.*
- (3) (Connected components) *The dimension of $H^0(X, \mathbb{Q})$ equals the number of connected components of X .*
- (4) *For any subspace $Y \subset X$, one can define relative cohomology groups $H^i(X, Y; \mathbb{Q})$ which fit into a long exact sequence*

$$\dots \rightarrow H^i(X, Y; \mathbb{Q}) \rightarrow H^i(X, \mathbb{Q}) \rightarrow H^i(Y, \mathbb{Q}) \rightarrow H^{i+1}(X, Y; \mathbb{Q}) \rightarrow \dots .$$

- (5) (Mayer–Vietoris) *If $U, V \subset X$ are open subsets such that $U \cup V = X$, then there is a long exact sequence of cohomology groups*

$$\dots \rightarrow H^i(X, \mathbb{Q}) \rightarrow H^i(U, \mathbb{Q}) \oplus H^i(V, \mathbb{Q}) \rightarrow H^i(U \cap V, \mathbb{Q}) \rightarrow H^{i+1}(X, \mathbb{Q}) \rightarrow \dots .$$

Note that this implies immediately that

$$b_i(U \cap V) \leq b_i(U) + b_i(V) + b_{i+1}(X).$$

Finally, we recall some properties of singular cohomology with regard to projective and injective limits. These properties are used in the proof of Proposition 3.2.6. Below, we drop the coefficients \mathbb{Q} from the notation of singular cohomology groups.

Let I be a directed set, $(U_i)_{i \in I}$ be a directed system of topological spaces, and

$$U = \varinjlim_i U_i$$

denote the corresponding direct limit. In particular, for all $i \leq j$ ($i, j \in I$), we have continuous maps $f_{ij} : U_i \rightarrow U_j$ which induce morphisms $f_{ij}^* : H^k(U_j) \rightarrow H^k(U_i)$. The latter cohomology groups form an inverse system, and the natural continuous maps $U_i \rightarrow U$ induce a morphism

$$H^k(U) \rightarrow \varprojlim_i H^k(U_i).$$

Similarly, an inverse system $(U_i)_{i \in I}$ of topological spaces gives rise to a direct system of corresponding cohomology groups and a natural morphism

$$\varinjlim_i H^k(U_i) \rightarrow H^k(U), \quad \text{where } U = \varprojlim_i U_i.$$

In this article, we only consider direct systems U_i given by an increasing sequence of subspaces of a space X or inverse systems U_i given by a decreasing sequence of subspaces. In the former case, the direct limit U is given by the union of these spaces, and in the latter case the inverse limit is given by the intersection of these subspaces. The following lemma is our main tool for understanding the corresponding cohomology groups.

Lemma A.1.2. *Let X be a paracompact Hausdorff space having the homotopy type of a finite simplicial complex, and I a directed set.*

- (1) *Let $\{U_i\}_{i \in I}$ be a decreasing sequence of open subspaces of X , and $S := \bigcap_i U_i$. Suppose that the family U_i is cofinal in the family of open neighborhoods of S in X . Then the natural map*

$$\varinjlim_i H^k(U_i) \rightarrow H^k(S)$$

is an isomorphism.

- (2) *Let $\{C_i\}_{i \in I}$ be an increasing sequence of compact subspaces of S , and $S := \bigcup_i C_i$. Suppose that the family C_i is cofinal in the family of compact subspaces of S . Then the natural map*

$$H^k(S) \rightarrow \varprojlim_i H^k(C_i)$$

is an isomorphism.

Proof of (1). This is Theorem 5 in [LR68]. □

Proof of (2). The statement follows from the fact that the singular homology of any space is isomorphic to the direct limit of the singular homology of its compact subspaces [Spa66, Theorem 4.4.6], the fact that the singular cohomology group $H^*(S, \mathbb{Q})$ is canonically isomorphic to $\text{Hom}(H_*(S, \mathbb{Q}), \mathbb{Q})$ since \mathbb{Q} is a field, and that the dual of a direct limit of finite-dimensional vector spaces is the inverse limit of the duals of those spaces. □

Remark A.1.3. Note that a compact Hausdorff space is paracompact Hausdorff. In the applications considered in this paper, the previous lemma is applied in the setting of compact Hausdorff spaces.

A.2. Recollections from Hrushovski–Loeser

In this section we recall some results from the theory of non-archimedean tame topology due to Hrushovski and Loeser [HL16]. The main reference for this section is [HL16, Chapter 14], but we refer the reader to [Duc16] for an excellent survey. In particular, we

will deal with the model theory of valued fields. We denote by K a complete valued field with values in the ordered multiplicative group of positive real numbers.

We consider a two-sorted language with the two sorts corresponding to valued fields and the value group. The signature of this two-sorted language will be

$$(0, 1, +_K, \times_K, |\cdot| : K \rightarrow \mathbb{R}_+, \leq_{\mathbb{R}_+}, \times_{\mathbb{R}}),$$

where the subscript K denotes constants, functions, relations etc. of the field sort and the subscript \mathbb{R}_+ denotes the same for the value group sort. When the context is clear we will drop the subscripts.

We denote by $|\cdot|$ the valuation written multiplicatively. The valuation $|\cdot|$ satisfies

$$|x + y| \leq \max(|x|, |y|), \quad |x \cdot y| = |x| |y|, \quad |0| = 0.$$

Remark A.2.1. Note that we follow Berkovich’s convention and write our valuations multiplicatively. In particular, the terminology ‘valuation’ is somewhat abusive, and here we really mean a non-archimedean absolute value. In [HL16], all valuations are written additively.

Following [HL16, §14.1], we will denote by \mathbf{F} the two-sorted structure $(K; \mathbb{R}_+)$ viewed as a substructure of a model of ACVF (with value group \mathbb{R}_+). Given a quasi-projective variety V defined over K and an \mathbf{F} -definable subset X of $V \times \mathbb{R}_+^n$, Hrushovski and Loeser [HL16] associate to X (functorially) a topological space $B_{\mathbf{F}}(X)$. By definition, this is the space of types, in X , defined over \mathbf{F} which are almost orthogonal to the definable set \mathbb{R}_+ . Given a variety V as above, we say that a subset $Z \subset B_{\mathbf{F}}(V)$ is *semi-algebraic* if it is of the form $B_{\mathbf{F}}(X)$ for an \mathbf{F} -definable subset $X \subset V$. We note that X itself can be identified in $B_{\mathbf{F}}(X)$ as the set of realized types, and hence there is a canonically defined injection $X \hookrightarrow B_{\mathbf{F}}(X)$.

We now recall a description of the spaces $B_{\mathbf{F}}(X)$ in some special cases and some of their properties; these are the only properties which are used in this article.

- Properties A.2.2.** (1) ([HL16, 14.4.1]) *For every \mathbf{F} -definable set X , $B_{\mathbf{F}}(X)$ is a Hausdorff topological space which is locally path connected. This construction is functorial in definable maps, i.e. a definable map $f : X \rightarrow Y$ induces a continuous map of the corresponding topological spaces.*
- (2) ([HL16, 14.1, p. 194]) *If V is an affine variety and $X \subset V$ a definable subset, then $B_{\mathbf{F}}(X)$ is a subspace of $B_{\mathbf{F}}(V)$. In fact, it is a semi-algebraic subset (in the sense of Berkovich spaces, see property (3) below).*
- (3) ([HL16, 14.1, p. 194]) *Suppose X is an affine variety $\text{Spec}(A)$. In this case, $B_{\mathbf{F}}(X)$ can be identified with the Berkovich analytic space associated to X . Its points can be described in terms of multiplicative semi-norms as follows. A point of $B_{\mathbf{F}}(X)$ is a multiplicative map $\phi : A \rightarrow \mathbb{R}_+$ such that $\phi(a + b) \leq \max(\phi(a), \phi(b))$.*
- (4) *With $X = \text{Spec}(A)$, the topology on $B_{\mathbf{F}}(X)$ is the one inherited from viewing it as a natural subset of \mathbb{R}_+^A . If $f \in A$, then f gives rise to a continuous function*

$$f : B_{\mathbf{F}}(X) \rightarrow \mathbb{R}_+$$

defined as follows:

$$f(\phi) = \phi(f) \in \mathbb{R}_+.$$

This follows from the previous observation and the definition of the topology on Berkovich analytic spaces.

- (5) ([HL16, 14.1, p. 194]) Let $V = \text{Spec}(A)$. Then any formula ϕ of the form $f \bowtie \lambda g$, where $f, g \in A$, $\lambda \in \mathbb{R}_+$ and $\bowtie \in \{\leq, <, \geq, >\}$, gives a definable subset X of V , and therefore a semi-algebraic subset $B_{\mathbf{F}}(X)$ of $B_{\mathbf{F}}(V)$. It can be described in the language of valuations as the set $\{x \in B_{\mathbf{F}}(V) \mid f(x) \bowtie \lambda g(x)\}$. In general, the semi-algebraic subset associated to a Boolean combination of such formulas is the corresponding Boolean combination of the semi-algebraic subsets associated to each formula. Moreover, a subset of $B_{\mathbf{F}}(V)$ is semi-algebraic if and only if it is a Boolean combination of subsets of the form $\{x \in B_{\mathbf{F}}(X) \mid f(x) \bowtie \lambda g(x)\}$, where $f, g \in A$, $\lambda \in \mathbb{R}_+$ and $\bowtie \in \{\leq, <, \geq, >\}$.
- (6) ([HL16, 14.1.2]) If X is an \mathbf{F} -definable subset of an algebraic variety V , then $B_{\mathbf{F}}(X)$ is compact if and only if $B_{\mathbf{F}}(X)$ is closed in $B_{\mathbf{F}}(V')$ where V' is a complete algebraic variety containing V .
- (7) Suppose that K is algebraically closed, $V = \text{Spec}(A) \subset \mathbb{A}_K^N$ is an affine subvariety, and $\phi(X; T)$ (with $X = (X_1, \dots, X_N)$) a formula with parameters in \mathbf{F} . Here the X_i are free variables of the field sort and T is a free variable of the value sort. Suppose $a \in \mathbb{R}_+$ is such that for all t, t' satisfying $a < t < t'$, $(K; \mathbb{R}_+) \models \phi(X; t') \rightarrow \phi(X, t)$. Let $\psi(X)$ be the formula

$$\exists T (T > a) \wedge \phi(X, T).$$

Then

$$\tilde{\mathcal{R}}(\psi, V) = \bigcup_{a < t} \tilde{\mathcal{R}}(\phi(\cdot; t), V).$$

Proof of (7). The inclusion $\bigcup_{a < t} \tilde{\mathcal{R}}(\phi(\cdot; t), V) \subset \tilde{\mathcal{R}}(\psi, V)$ is obvious, since for each $t > a$, $(K; \mathbb{R}_+) \models \phi(X, t) \rightarrow \psi(X)$, which implies that $\tilde{\mathcal{R}}(\phi(\cdot; t), V) \subset \tilde{\mathcal{R}}(\psi(\cdot), V)$.

To prove the reverse inclusion, let $p \in \tilde{\mathcal{R}}(\psi, V)$. Then, by definition p is a type which is almost orthogonal to the value group, and moreover there exists $x \in \mathcal{R}(\psi, V)(K')$ such that $x \models p$ and $(K'; \mathbb{R}_+)$ is an elementary extension of $(K; \mathbb{R}_+)$ (since types which are orthogonal to \mathbb{R}_+ can always be realized in such a model). Hence, there exists $t_0 > a$, $t_0 \in \mathbb{R}_+$, such that $(K'; \mathbb{R}_+) \models \phi(x, t_0)$, and so $p \in \tilde{\mathcal{R}}(\phi(\cdot, t_0), V)$. This proves that

$$\tilde{\mathcal{R}}(\psi, V) \subset \bigcup_{a < t} \tilde{\mathcal{R}}(\phi(\cdot; t), V). \quad \square$$

Given an \mathbf{F} -definable map $f : X \rightarrow \mathbb{R}_+$, we will denote by $B_{\mathbf{F}}(f) : B_{\mathbf{F}}(X) \rightarrow B_{\mathbf{F}}(\mathbb{R}_+) = \mathbb{R}_+$ the induced map. We will say that $B_{\mathbf{F}}(f)$ is a semi-algebraic map.

The following theorems which are easily deduced from the main theorems in [HL16, Chapter 14] will play a key role in the results of this paper. We will use the same notation as above.

Theorem A.3 ([HL16, Theorem 14.4.4]). *Let V be a quasi-projective variety over K , $X \subset V$ be an \mathbf{F} -definable subset and $f : X \rightarrow \mathbb{R}_+$ be an \mathbf{F} -definable map. For $t \in \mathbb{R}_+$, let $B_{\mathbf{F}}(X)_{\geq t}$ denote the semi-algebraic subset $B_{\mathbf{F}}(X \cap (f \geq t)) = B_{\mathbf{F}}(X) \cap B_{\mathbf{F}}(f \geq t)$ of $B_{\mathbf{F}}(V)$. Then there exists a finite partition \mathcal{P} of \mathbb{R}_+ into intervals such that for each $I \in \mathcal{P}$ and for all $\varepsilon \leq \varepsilon' \in I$, the inclusion $B_{\mathbf{F}}(X)_{\geq \varepsilon'} \hookrightarrow B_{\mathbf{F}}(X)_{\geq \varepsilon}$ is a homotopy equivalence.*

Theorem A.4 ([HL16, Theorem 14.3.1(1)]). *Let Y be a variety and $X \subset Y \times \mathbb{R}_+^r \times \mathbb{P}^m$ be an \mathbf{F} -definable set. Let $\pi : X \rightarrow Y \times \mathbb{R}_+^r$ be the projection map. Then there are finitely many homotopy types amongst the fibers $(B_{\mathbf{F}}(\pi^{-1}(y, t)))_{(y,t) \in Y \times \mathbb{R}_+^r}$.*

Theorem A.5 ([HL16, Theorem 14.2.4]). *Let V be a quasi-projective variety defined over K , and X an \mathbf{F} -definable subset of V such that $B_{\mathbf{F}}(X)$ is compact. Then there exists a family of finite simplicial complexes $(X_i)_{i \in I}$ (where I is a directed partially ordered set) embedded in $B_{\mathbf{F}}(X)$ of dimension $\leq \dim(V)$, deformation retractions $\pi_{i,j} : X_i \rightarrow X_j$, $j < i$, and deformation retractions $\pi_i : B_{\mathbf{F}}(X) \rightarrow X_i$, such that $\pi_{i,j} \circ \pi_i = \pi_j$ and the canonical map $B_{\mathbf{F}}(X) \rightarrow \varprojlim_i X_i$ is a homeomorphism.*

As an immediate consequence of Theorem A.5 we have, using the same notation:

Corollary A.6. *Let $V \subset \mathbb{A}_K^N$ be a closed affine subvariety, and let $B_{\mathbf{F}}(X)$ be a semi-algebraic subset of V .*

- (a) *Every connected component of $B_{\mathbf{F}}(X)$ is path connected.*
- (b) *$H^i(B_{\mathbf{F}}(X)) = 0$ for $i > \dim(V)$.*
- (c) *$\dim H^*(B_{\mathbf{F}}(X)) < \infty$.*
- (d) *The restriction homomorphism $H^{\dim(V)}(B_{\mathbf{F}}(V)) \rightarrow H^{\dim(V)}(B_{\mathbf{F}}(X))$ is surjective.*

Proof. Recall the definition of $\text{Cube}_V(R)$ (see Notation 3.2.1) and that $\text{Cube}_V(R)$ is a compact topological space. Similar remarks apply to $\text{Cube}_V(R) \cap B_{\mathbf{F}}(X)$. Moreover, arguing as for part (6) of Lemma 3.2.7, for sufficiently large R the natural inclusions $\text{Cube}_V(R) \cap X \hookrightarrow B_{\mathbf{F}}(X)$ and $\text{Cube}_V(R) \hookrightarrow B_{\mathbf{F}}(V)$ induce homotopy equivalences. In the following, we fix an R large enough such that both inclusions are homotopy equivalences. Note that (a)–(c) now follow directly from Theorem A.5. We shall now prove (d).

By the previous remarks, it is sufficient to prove that the natural induced morphism

$$H^{\dim(V)}(\text{Cube}_V(R)) \rightarrow H^{\dim(V)}(\text{Cube}_V(R) \cap B_{\mathbf{F}}(X))$$

is surjective.

By Theorem A.5, $\text{Cube}_V(R)$ has the homotopy type of a finite simplicial polyhedron of dimension at most $\dim(V)$. Since $\text{Cube}_V(R)$ is compact, it follows that the cohomological dimension (in the sense of [Ive86, p. 196, Definition 9.4]) of $\text{Cube}_V(R)$ is $\leq \dim(V)$.

It follows again from Theorem A.5 that there exists a compact polyhedron $Z \subset \text{Cube}_V(R) \cap X$ such that Z is a deformation retract of $\text{Cube}_V(R) \cap B_{\mathbf{F}}(X)$. Let $\iota : Z \hookrightarrow \text{Cube}_V(R) \cap B_{\mathbf{F}}(X)$ be the inclusion map. Since the inclusion of Z in $\text{Cube}_V(R)$ factors through ι , and ι induces isomorphisms in cohomology, it follows (using the long exact sequence of cohomology for pairs) that

$$H^*(\text{Cube}_V(R), \text{Cube}_V(R) \cap B_{\mathbf{F}}(X)) \cong H^*(\text{Cube}_V(R), Z).$$

We now prove that

$$H^{\dim(V)+1}(\text{Cube}_V(R), \text{Cube}_V(R) \cap B_{\mathbf{F}}(X)) \cong H^{\dim(V)+1}(\text{Cube}_V(R), Z) = 0.$$

This gives the desired result by an application of the long exact sequence in cohomology associated to the pair $(\text{Cube}_V(R), \text{Cube}_V(R) \cap B_{\mathbf{F}}(X))$.

Recall that $\text{Cube}_V(R)$ is a Hausdorff space, and so Z is a closed subspace of $\text{Cube}_V(R)$. By [Ive86, p. 198, Proposition 9.7] it follows that the cohomological dimension of $U := \text{Cube}_V(R) \setminus Z$ is also $\leq \dim(V)$. This implies that $H_c^{\dim(V)+1}(U) \cong H^{\dim(V)+1}(\text{Cube}_V(R), Z) = 0$, which finishes the proof. \square

Acknowledgments. S.B. thanks Institut Henri Poincaré, Paris, for hosting him during the trimester on “Model Theory, Combinatorics and Valued Fields” where part of this work was done. Both authors thank the anonymous referees for very detailed reports which helped to substantially improve the paper.

S.B. would like to acknowledge support from the National Science Foundation awards and DMS-1620271, CCF-1618918 and CCF-1910441. D.P. would like to acknowledge support from the National Science Foundation award DMS-1502296.

References

- [AD⁺16] Aschenbrenner, M., Dolich, A., Haskell, D., Macpherson, D., Starchenko, S.: Vapnik–Chervonenkis density in some theories without the independence property, I. *Trans. Amer. Math. Soc.* **368**, 5889–5949 (2016) [Zbl 1423.03119](#) [MR 3458402](#)
- [Bas10] Basu, S.: Combinatorial complexity in o-minimal geometry. *Proc. London Math. Soc.* (3) **100**, 405–428 (2010) [Zbl 1186.52017](#) [MR 2595744](#)
- [BPR05] Basu, S., Pollack, R., Roy, M.-F.: On the Betti numbers of sign conditions. *Proc. Amer. Math. Soc.* **133**, 965–974 (2005) [Zbl 1080.14068](#) [MR 2117195](#)
- [BPR06] Basu, S., Pollack, R., Roy, M.-F.: *Algorithms in Real Algebraic Geometry*. 2nd ed., *Algorithms Comput. Math.* 10, Springer, Berlin (2006) [Zbl 1102.14041](#) [MR 2248869](#)
- [BPR09] Basu, S., Pollack, R., Roy, M.-F.: An asymptotically tight bound on the number of connected components of realizable sign conditions. *Combinatorica* **29**, 523–546 (2009) [Zbl 1212.14005](#) [MR 2604321](#)
- [Duc16] Ducros, A.: About Hrushovski and Loeser’s work on the homotopy type of Berkovich spaces. In: *Nonarchimedean and Tropical Geometry*, *Simons Sympos.*, Springer, 99–131 (2016) [Zbl 1349.14098](#) [MR 3702309](#)
- [HHM08] Haskell, D., Hrushovski, E., Macpherson, D.: *Stable Domination and Independence in Algebraically Closed Valued Fields*. *Lecture Notes in Logic* 30, Assoc. Symbolic Logic, Chicago, IL, and Cambridge Univ. Press, Cambridge (2008) [Zbl 1217.03026](#) [MR 2369946](#)
- [HL16] Hrushovski, E., Loeser, F.: Non-Archimedean Tame Topology and Stably Dominated Types. *Ann. of Math. Stud.* 192, Princeton Univ. Press, Princeton, NJ (2016) [Zbl 1365.14033](#) [MR 3445772](#)
- [Ive86] Iversen, B.: *Cohomology of Sheaves*. Universitext, Springer, Berlin (1986) [Zbl 1272.55001](#) [MR 842190](#)
- [JL10] Johnson, H. R., Laskowski, M. C.: Compression schemes, stable definable families, and o-minimal structures. *Discrete Comput. Geom.* **43**, 914–926 (2010) [Zbl 1201.03022](#) [MR 2610477](#)

-
- [LR68] Lee, C. N., Raymond, F.: Čech extensions of contravariant functors. *Trans. Amer. Math. Soc.* **133**, 415–434 (1968) [Zbl 0162.55001](#) [MR 0234450](#)
- [Mac76] Macintyre, A.: On definable subsets of p -adic fields. *J. Symbolic Logic* **41**, 605–610 (1976) [Zbl 0362.02046](#) [MR 0485335](#)
- [RBG01] Rónyai, L., Babai, L., Ganapathy, M.: On the number of zero-patterns of a sequence of polynomials. *J. Amer. Math. Soc.* **14**, 717–735 (2001) [Zbl 0978.12001](#) [MR 1824986](#)
- [SGA72] Théorie des topos et cohomologie étale des schémas. Tome 1: Théorie des topos. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), *Lecture Notes in Math.* 269, Springer, Berlin (1972) [Zbl 0245.00002](#) [MR 0354652](#)
- [Sim15] Simon, P.: *A Guide to NIP Theories*. *Lecture Notes in Logic* 44, Assoc. Symbolic Logic, Chicago, IL, and Cambridge Sci. Publ., Cambridge (2015) [Zbl 1332.03001](#) [MR 3560428](#)
- [Spa66] Spanier, E. H.: *Algebraic Topology*. McGraw-Hill, New York (1966) [Zbl 0145.43303](#) [MR 0210112](#)