



Peter Hästö · Jihoon Ok

Maximal regularity for local minimizers of non-autonomous functionals

Received April 2, 2019

Abstract. We establish local $C^{1,\alpha}$ -regularity for some $\alpha \in (0, 1)$ and C^α -regularity for any $\alpha \in (0, 1)$ of local minimizers of the functional

$$v \mapsto \int_{\Omega} \varphi(x, |Dv|) dx,$$

where φ satisfies a (p, q) -growth condition. Establishing such a regularity theory with sharp, general conditions has been an open problem since the 1980s. In contrast to previous results, we formulate the continuity requirement on φ in terms of a single condition for the map $(x, t) \mapsto \varphi(x, t)$, rather than separately in the x - and t -directions. Thus we can obtain regularity results for functionals without assuming that the gap q/p between the upper and lower growth bounds is close to 1. Moreover, for $\varphi(x, t)$ with particular structure, including p -, Orlicz-, $p(x)$ - and double phase-growth, our single condition implies known, essentially optimal, regularity conditions. Hence, we handle regularity theory for the above functional in a universal way.

Keywords. Maximal regularity, non-autonomous functional, variable exponent, double phase, non-standard growth, minimizer, Hölder continuity, generalized Orlicz space, Musielak–Orlicz space

1. Introduction

The calculus of variations is a classical and still active topic in mathematics which is connected not only to other mathematical fields (partial differential equations, geometry, ...) and but also to applications (physics, engineering, economy, ...). Research on regularity of minimizers of the functional

$$v \mapsto \mathcal{F}(v, \Omega) := \int_{\Omega} F(x, Dv) dx$$

Peter Hästö: Department of Mathematics and Statistics, FI-20014 University of Turku, Finland, and Department of Mathematics, FI-90014 University of Oulu, Finland;
peter.hasto@oulu.fi / peter.hasto@utu.fi

Jihoon Ok: Department of Mathematics, Sogang University, Seoul 04107, Republic of Korea;
jihoonok@sogang.ac.kr

Mathematics Subject Classification (2020): Primary 49N60; Secondary 35A15, 35B65, 35J62, 46E35

has been a major topic in the calculus of variations and PDEs. If F depends only on the gradient, i.e. $F(x, z) \equiv F(z)$, \mathcal{F} is called an *autonomous* functional. The simplest non-linear model case is the p -power function

$$F(z) = |z|^p, \quad 1 < p < \infty.$$

The corresponding Euler–Lagrange equation is the p -Laplace equation $\operatorname{div}(|Du|^{p-2}Du) = 0$, and the maximal regularity of weak solutions of p -Laplace equations is $C^{1,\alpha}$ for some $\alpha \in (0, 1)$ depending only on p and the dimension n . We refer to [1, 32, 40, 58, 62, 78–80, 82, 83] for classical results on $C^{1,\alpha}$ -regularity for equations and systems of p -Laplacian type.

On the other hand, if F depends on both the space variable and the gradient, \mathcal{F} is called a *non-autonomous* functional, and this has been a central topic in contemporary regularity theory. The main approach to such minimization problems is due to Giaquinta and Giusti [47, 48]. It is based on the following p -type growth conditions:

$$\begin{cases} z \mapsto F(x, z) \text{ is } C^2, \\ \nu|z|^p \leq F(x, z) \leq L(1 + |z|^p), \\ \nu(\mu^2 + |z|)^{(p-2)/2}|\lambda|^2 \leq F_{zz}(x, z)\lambda \cdot \lambda \leq L(\mu^2 + |z|^2)^{(p-2)/2}|\lambda|^2, \\ |F(x, z) - F(y, z)| \leq \omega(|x - y|)(1 + |z|^p). \end{cases}$$

This essentially corresponds to the perturbed case $a(x)|z|^p$ with the same p -type growth assumed at all points. Lieberman [60] extended this to the case where $|z|^p$ is replaced by $\varphi(|z|)$. However, such structure conditions fail to accommodate many kinds of energy functionals since the variability in the x - and z -directions are treated separately.

The need to treat the x - and z -directions separately leads Mingione to conclude in his influential survey that “regularity results should be chased [in more general cases] by looking at special classes of functionals and thinking of relevant model examples, thereby limiting the degree of generality one wants to achieve” [71, p. 405]. In this spirit, the most significant non-autonomous functionals in the literature have so-called Uhlenbeck structure, i.e. F depends on $t := |z|$ instead of z ,

$$F(x, z) = \varphi(x, |z|) = \varphi(x, t),$$

and are the following:

- I. Perturbed Orlicz: $a(x)\psi(t)$, where $0 < \nu \leq a(\cdot) \leq L$ and $\psi'(t) \approx t\psi''(t)$.
- II. Variable exponent: $t^{p(x)}$, where $1 < p^- \leq p(\cdot) \leq p^+ < \infty$.
- III. Double phase: $t^p + a(x)t^q$, where $1 < p \leq q$ and $a(\cdot) \geq 0$.

These models were first studied by Zhikov [85, 86] in the 1980’s in relation to Lavrent’ev’s phenomenon and have been considered in hundreds of papers since [71, 75]. In keeping with Mingione’s thesis, regularity results for these cases have been established in independent, idiosyncratic ways (cf. Section 2). Moreover, various variants and borderline cases have been investigated, such as:

- IV. Perturbed variable exponent: $t^{p(x)} \log(e + t)$, e.g. [44, 59, 72, 74].

- V. Orlicz variable exponent: $[\psi(t)]^{p(x)}$ or $\psi(t^{p(x)})$, e.g. [21, 45].
- VI. Degenerate double phase: $t^p + a(x)t^p \log(e + t)$, e.g. [9, 16].
- VII. Orlicz double phase: $\psi(t) + a(x)\xi(t)$, e.g. [17].
- VIII. Triple phase: $t^p + a(x)t^q + b(x)t^r$, e.g. [31, 43].
- IX. Double variable exponent: $t^{p(x)} + t^{q(x)}$, e.g. [22, 77, 84].
- X. Variable exponent double phase $t^{p(x)} + a(x)t^{q(x)}$, e.g. [61, 76].

In this paper, we establish a general regularity theory for non-autonomous functionals with Uhlenbeck structure based on a single condition involving both the x - and t -directions. Specifically, we prove maximal local regularity properties, i.e. $C^{1,\alpha}$ -regularity for some $\alpha \in (0, 1)$ and C^α -regularity for any $\alpha \in (0, 1)$. We consider a convex function $\varphi : \Omega \times [0, \infty) \rightarrow [0, \infty)$ satisfying the following “vanishing A1” variant of (A1) (see Definitions 3.4 and 4.1 below):

(VA1) There exists a non-decreasing continuous function $\omega : [0, \infty) \rightarrow [0, 1]$ with $\omega(0) = 0$ such that for any small ball $B_r \Subset \Omega$,

$$\varphi_{B_r}^+(t) \leq (1 + \omega(r))\varphi_{B_r}^-(t) \quad \text{for all } t > 0 \text{ satisfying } \varphi_{B_r}^-(t) \in [\omega(r), |B_r|^{-1}],$$

where $\varphi_{B_r}^+(t)$ and $\varphi_{B_r}^-(t)$ are the supremum and infimum of $\varphi(\cdot, t)$ in B_r , respectively. Let us point out that (VA1) is optimal for Theorem 1.1 in the following sense: For any $\theta < 1$ assume that (VA1) is replaced by

$$\varphi_{B_r}^+(t) \leq (1 + \omega(r))\varphi_{B_r}^-(t) \quad \text{for all } t > 0 \text{ satisfying } \varphi_{B_r}^-(t) \in [\omega(r), |B_r|^{-\theta}].$$

Then the conclusions of the theorem do not hold, as is shown by examples in [71] already in the double phase case (cf. Corollary 8.6); see also [6, 13]. Furthermore, $1 + \omega(r)$ in the inequality from (VA1) ensures the continuity of the function, which is necessary already in the perturbed linear case (cf. Corollary 8.1 and Remarks 1.3 and 1.4).

Theorem 1.1. *Let $\varphi \in \Phi_w(\Omega)$, $\varphi(x, \cdot) \in C^1([0, \infty))$ for every $x \in \Omega$ with $\partial_t \varphi$ satisfying (A0), (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for some $1 < p \leq q$ (see Definition 3.1) and let $u \in W_{\text{loc}}^{1,\varphi}(\Omega)$ be a local minimizer of the φ -energy*

$$\int_{\Omega} \varphi(x, |\nabla u|) dx. \quad (1.2)$$

- (1) *If φ satisfies (VA1), then $u \in C_{\text{loc}}^\alpha(\Omega)$ for any $\alpha \in (0, 1)$.*
- (2) *If φ satisfies (VA1) and $\omega(r) \leq cr^\beta$ for some $c, \beta > 0$, then $u \in C_{\text{loc}}^{1,\alpha}(\Omega)$ for some $\alpha \in (0, 1)$. Here α depends only on n, p, q, L and β , where $L \geq 1$ is from (A0).*

Remark 1.3. In this paper, we consider $\varphi(x, t)$ continuous in x . It is clear that we cannot remove the assumption $\lim_{r \rightarrow 0} \omega(r) = 0$ from (VA1) and still obtain C^α -regularity for all $\alpha \in (0, 1)$. However, continuity is not strictly speaking necessary, as it is known for $\varphi(x, t) = a(x)\psi(t)$ with a locally VMO (vanishing mean oscillation) that the corresponding minimizer is in C_{loc}^α for any $\alpha \in (0, 1)$, in fact, in $W_{\text{loc}}^{1,p}$ for any $p > 1$. It seems that for this result the special multiplicative structure is important.

Remark 1.4. If we consider solutions of the general linear elliptic equation $\operatorname{div}(A(x)Du) = 0$, where $A(x)$ is a bounded and uniformly elliptic $n \times n$ matrix, then the continuity of A does not imply that the function is Lipschitz or its derivative is continuous [56, Propositions 1.5 and 1.6]. Therefore, we cannot expect to remove the assumption $\omega(r) \leq cr^\beta$ from (VA1) and still obtain $C^{1,\alpha}$ -regularity.

We shall introduce notation, assumptions and properties of generalized Φ -functions and related spaces later in Section 3. Recall that *local minimizer* means that u satisfies

$$\int_{\Omega'} \varphi(x, |\nabla u|) dx \leq \int_{\Omega'} \varphi(x, |\nabla v|) dx$$

for every $v \in W^{1,\varphi}(\Omega')$ with $u - v \in W_0^{1,\varphi}(\Omega')$ and $\Omega' \Subset \Omega$.

In fact, we will generalize (VA1) to a weaker version, (wVA1), which covers not only (VA1) but its borderline cases (see Remark 4.2) as well as the PDE case (see Remark 4.3), and under this condition we will prove C^α - and $C^{1,\alpha}$ -regularity (see Theorems 7.2 and 7.4). As far as we know, these theorems cover all previously known results (and several new ones) of C^α - or $C^{1,\alpha}$ -regularity for the functionals I–X (see Section 8) with the exception of VMO coefficients (Remark 1.3).

Even in the case of autonomous functionals (i.e. $\varphi(x, t) \equiv \varphi(t)$), our results provide slight extensions to the state-of-the-art. Up to now, maximal regularity for autonomous functionals has been established assuming $\varphi \in C^1([0, \infty)) \cap C^2((0, \infty))$. However, in this paper we only assume $\varphi \in C^1([0, \infty))$, that is, we do not assume that φ is twice differentiable. For instance, $\varphi(t) := \int_0^t \min\{s, s^2\} ds$ (cf. [5]) is covered by our result but is not C^2 .

Let us conclude the introduction by outlining the approach of the paper and pointing out the main difficulties and innovations.

The first difficulty for a reasonable regularity theory is to find a well-designed condition for general φ . The regularity conditions on φ for the types I–III seem unconnected to one another, since in these cases, the behaviors of φ with respect to x and t can be investigated separately. Recently, on the other hand, the C^α -continuity with some small $\alpha > 0$ for (quasi-)minimizers of the general non-autonomous functional has been established under the so-called (A1) condition [13, 54, 55]:

$$\varphi_{B_r}^+(t) \leq L\varphi_{B_r}^-(t) \quad \text{for all } t > 0 \text{ satisfying } \varphi_{B_r}^-(t) \in [1, |B_r|^{-1}]. \quad (1.5)$$

From this, it is natural to require $L \rightarrow 1$ as $r \rightarrow 0$ for higher regularity. Additionally, small values $t \leq 1$ were previously lumped into an additive constant using decay at infinity. A more precise estimate, on the other hand, requires the previous condition to be extended from $[1, |B_r|^{-1}]$ to $[\omega(r), |B_r|^{-1}]$.

The main difficulty is to find a suitably regular auxiliary autonomous function $\tilde{\varphi}(t)$ for the perturbation technique in which one approximates the minimizer with the solution to a related but simpler minimization problem. In order for the perturbation argument to work under the assumption (VA1), the autonomous function $\tilde{\varphi}(t)$ should satisfy the following requirements:

- (1) $\tilde{\varphi} \in C^1([0, \infty)) \cap C^2((0, \infty))$ and $t\tilde{\varphi}''(t) \approx \tilde{\varphi}'(t)$.
- (2) For a given B_r with small $r \in (0, 1)$, $\tilde{\varphi}(t)$ is sufficiently close in some sense to $\varphi(x, t)$ for all $(x, t) \in B_r \times [t_1, t_2]$, where $t_1 := (\varphi_{B_r}^-)^{-1}(\omega(r))$ and $t_2 := (\varphi_{B_r}^-)^{-1}(|B_r|^{-1})$.
- (3) $\theta_0(x, t) := \varphi(x, \tilde{\varphi}^{-1}(t))$ satisfies (A0), (aInc)₁, (aDec)_{q/p} and (A1).

The construction of such $\tilde{\varphi}$ is quite non-trivial, since the property (3) is not satisfied in general for either $\tilde{\varphi}(t) = \varphi(y, t)$ with any choice of $y \in \overline{B_r}$ or $\tilde{\varphi}(t) = \varphi_{B_r}^-(t)$ (the expected choices based on previous research). Note that for type II (variable exponent) or type III (double phase), one can simply take $\tilde{\varphi}(t) = t^{p_r}$ or $\tilde{\varphi}(t) = t^p + a_r t^q$, where $p_r := \inf_{B_r} p(\cdot)$ and $a_r := \inf_{B_r} a(\cdot)$, so this provides no guidance for the general case: in these special cases $t \mapsto \varphi(x, \tilde{\varphi}^{-1}(t))$ satisfies (aInc)₁ since a single point captures the slowest growth for all values of t , whereas in general the slowest growth may occur at different locations for different t .

The requirements (1)–(3) above are crucially used in our comparison step. Let v be a minimizer of an autonomous functional with $\tilde{\varphi}$ -energy in B_r satisfying $v = u$ on ∂B_r . Then by (1) and known regularity results for Orlicz growth, we find that v is locally C^{1, α_0} for some $\alpha_0 \in (0, 1)$ (Lemma 4.12). Moreover, from (3) we can deduce a global non-linear Calderón–Zygmund type estimate in the generalized Orlicz space L^θ with $\theta = \theta_0^{1+\alpha_0}$ for some $\alpha_0 > 0$ (Lemma 4.15), which implies that $Dv \in L^\varphi(B_r)$ and so, with this v , we can use the minimizing property of u . Note that this approach is new even for the double phase problem, type III.

The Calderón–Zygmund type estimates (Lemma 4.15) in generalized Orlicz space L^θ for the norm will be obtained by an extrapolation argument [29] and in this process (A1) for θ suffices. However, we need a mean integral version of Calderón–Zygmund type estimate that is stable under change of size of the underlying domain and here (A1) for θ is not enough. We overcome this problem by replacing $\theta(x, t)$ with $\theta(x, t) + t^{p_1}$ for suitable $p_1 > 1$ along with delicate analysis. Note that $\theta(x, t) + t^{p_1}$ satisfies a stronger assumption than (A1). As a consequence, there is “+1” in the mean integral version of estimate (4.17).

We construct our approximation $\tilde{\varphi}$ and derive the comparison estimate for φ and $\tilde{\varphi}$ in Section 5. In Proposition 5.12 we show that our approximation satisfies the assumptions in (3) above, and in this step a new framework for generalized Orlicz spaces from [51] is rather crucial. Then a comparison argument along with (2) and a higher integrability result for Du imply that Du is sufficiently close to Dv in the mean oscillation sense (Corollary 6.3).

We present proofs of some regularity results for autonomous problems in Appendices A and B. We start this article with an overview of regularity theory in the (p, q) -growth case (Section 2) and with notation and background (Section 3).

Remark 1.6. Constructing a suitable $\tilde{\varphi}$ is the main problem also in extending this approach to the case without Uhlenbeck structure, i.e. energy functionals depending on the derivative Du , not just its norm. Namely, an approximation $\tilde{\varphi} : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}$ affords us much less room to operate in than $\tilde{\varphi} : \Omega \times [0, \infty) \rightarrow \mathbb{R}$. Indeed, it is not even clear how to state the appropriate assumptions in this case. In addition, the main tools from [51]

concern only the isotropic case $\varphi(x, |Du|)$. Therefore, the regularity of the anisotropic minimization problem $\int \varphi(x, Du) dx$ remains a question for future research.

Remark 1.7. The vectorial case, i.e. $u : \Omega \rightarrow \mathbb{R}^N$ with $N > 1$, is also an interesting issue. The main difficulty in this case is the following: in order that the local minimizer of the regular autonomous functional with Orlicz function $\tilde{\varphi} = \tilde{\varphi}(t)$ have $C^{1,\alpha}$ -regularity, $\tilde{\varphi}$ should apparently satisfy not only $t\tilde{\varphi}''(t) \approx \tilde{\varphi}'(t)$ but also a Hölder type vanishing condition on $\tilde{\varphi}''$ (see [35, Assumption 2.2]). It is unclear whether (VA1) or some modification implies the additional condition on $\tilde{\varphi}$. This is also a future research topic.

2. Overview of regularity for (p, q) -growth and special cases

An alternative extension to the approach of Giaquinta and Giusti is to consider different upper and lower growth rates, and replace the exponent on the right-hand side by $q > p$. This leads to so-called (p, q) -growth functionals, for instance with assumptions

$$\begin{cases} z \mapsto F(x, z) \text{ is } C^2, \\ v|z|^p \leq F(x, z) \leq L(1 + |z|^q), \\ v(1 + |z|)^{(p-2)/2}|\lambda|^2 \leq F_{zz}(x, z)\lambda \cdot \lambda \leq L(1 + |z|^2)^{(q-2)/2}|\lambda|^2, \\ |F(x, z) - F(y, z)| \leq \omega(|x - y|)(1 + |z|^q). \end{cases}$$

This case was introduced and systematically studied by Marcellini [63–67]. Several other researchers also contributed to the theory [11, 38, 71]. For instance, Marcellini [64] started by showing that every minimizer in $W_{\text{loc}}^{1,q}(\Omega)$ has locally bounded gradient provided $2 \leq p \leq q$ and

$$\frac{q}{p} \leq 1 + \frac{2}{n-2} \quad \text{when } n > 2$$

(the proof uses PDE techniques and entails several additional assumptions, which are not presented here; see also a recent improvement in [12]). Note, however, that $W_{\text{loc}}^{1,q}(\Omega)$ is already higher integrability, so this is not a natural assumption in this context and was addressed in [64, Section 3]. Later, Esposito, Leonetti and Mingione [39] showed that every minimizer in $W_{\text{loc}}^{1,p}(\Omega)$ also belongs to $W_{\text{loc}}^{1,q}(\Omega)$, but only when

$$q/p \leq 1 + \beta/n \quad \text{for } \omega \in C^\beta.$$

Furthermore, they provide an example showing that if the latter condition does not hold, then a minimizer in $W_{\text{loc}}^{1,p}(\Omega)$ need not belong to $W_{\text{loc}}^{1,q}(\Omega)$ so the Lavrent'ev phenomenon occurs.

It seems that (p, q) -growth is the most general class of non-autonomous functionals in the calculus of variations. Regularity theory, including C^α - and $C^{1,\alpha}$ -regularity, in this general class is not easily obtained from classical regularity theory for functionals with standard p -growth (see for instance [71]). Furthermore, there are no general results in the (p, q) -case which cover the special cases I–X, so in that sense the theory is incomplete.

We note that some recent papers [13, 23, 24, 54, 55, 81] deal with the calculus of variations in generalized Orlicz spaces, but these papers do not cover higher regularity.

Indeed, the C^α - and $C^{1,\alpha}$ -regularity theories for type I–III functionals have been proved in independent ways. For I, φ is nothing but an autonomous functional with coefficient, and so regularity results can be obtained by using a standard perturbation argument. On the other hand, II and III are quite different from I, since they are potentially non-uniformly elliptic problems. Formally, we can rewrite the energy functions as

$$\text{II: } |Du|^{p(x)-p^-} |Du|^{p^-} \quad \text{and} \quad \text{III: } (1 + a(x)|Du|^{q-p}) |Du|^p.$$

Here, $|Du|^{p(x)-p^-}$ and $1 + a(x)|Du|^{q-p}$ blow up or vanish when $|Du|$ does. Therefore, by identifying $a(x)$ in I with $|Du|^{p(x)-p^-}$ or $1 + a(x)|Du|^{q-p}$, we see that a is neither bounded nor far away from the zero. Let us briefly introduce regularity results for the above types. Let u be a minimizer of the φ -energy (1.2) with φ being one of I–III. Then the following is known:

For type I, i.e. $\varphi(x, t) = a(x)\psi(t)$, suppose a is continuous with modulus of continuity ω_a . Then

$$\begin{aligned} \lim_{r \rightarrow 0^+} \omega_a(r) = 0 &\implies u \in C^\alpha \text{ for any } \alpha \in (0, 1), \\ \omega_a(r) \lesssim r^\beta \text{ for some } \beta > 0 &\implies u \in C^{1,\alpha} \text{ for some } \alpha \in (0, 1) \end{aligned} \quad (2.1)$$

(see for instance [71] and references therein).

For type II, i.e. $\varphi(x, t) = t^{p(x)}$, suppose p is continuous with modulus of continuity ω_p . Then

$$\begin{aligned} \lim_{r \rightarrow 0^+} \omega_p(r) \ln \frac{1}{r} = 0 &\implies u \in C^\alpha \text{ for any } \alpha \in (0, 1), \\ \omega_p(r) \lesssim r^\beta \text{ for some } \beta > 0 &\implies u \in C^{1,\alpha} \text{ for some } \alpha \in (0, 1). \end{aligned} \quad (2.2)$$

For these results, we refer to a series of papers of Acerbi, Coscia and Mingione [2, 3, 28]; see also [4, 18, 41, 42].

For type III, i.e. $\varphi(x, t) = t^p + a(x)t^q$, suppose $a \in C^{0,\beta}$ for some $\beta \in (0, 1]$. Then

$$q/p \leq 1 + \beta/n \implies u \in C^{1,\alpha} \text{ for some } \alpha \in (0, 1). \quad (2.3)$$

For this result, we refer to a series of papers of Baroni, Colombo and Mingione [10, 26]; see also [8, 15, 25, 27, 73]. Note that no independent condition implies C^α -regularity. In other words, we cannot ensure even C^α -regularity for u if $q/p > 1 + \beta/n$. We also mention that the $C^{1,\alpha}$ -regularity for type III was first proved under the following condition instead of (2.3):

$$q/p < 1 + \beta/n \implies u \in C^{1,\alpha} \text{ for some } \alpha \in (0, 1) \quad (\text{see [26]}), \quad (2.4)$$

and later it was extended to the borderline case $q/p = 1 + \beta/n$ in [10] (see also [30]).

As mentioned in the introduction, our general results cover all of these special cases. Specifically, Theorem 1.1(1) implies (2.1)₁ and (2.2)₁ and Theorem 1.1(2) implies (2.1)₂,

(2.2)₂ and (2.4). We notice that Theorem 1.1(2) does not imply (2.3). In fact, (VA1) holds when $\varphi(x, t) = t^p + a(x)t^q$ with $a(\cdot) \in C^{0,\beta}$ if and only if the strict inequality $q/p < 1 + \beta/n$ holds. This gap will be filled by Theorem 7.4; this is one main reason why we consider the slightly weaker assumption (wVA1).

Furthermore, many other, previously unstudied cases can also be covered: see, e.g. Corollary 8.3, and Section 8 more generally. Originally, the double phase model was introduced to model the situation when two phases (the p -growth and the q -growth phases) mix. Since only the larger exponent affects the nature of the problem, this was simplified in the form $t^p + a(x)t^q$ that we have seen. However, we can also consider a variant which is more closely related to the original motivation:

$$\varphi(x, t) = (1 - a(x))t^p + a(x)t^q, \quad \text{where } 1 < p \leq q, \quad a(\cdot) : \Omega \rightarrow [0, 1]. \quad (2.5)$$

Now a indicates the relative amount of material at a point from the q -phase. Such functionals have been treated by Eleuteri–Marcellini–Mascolo [36–38]. More generally, we can also deal with general double phase problems of the type

$$\varphi(x, t) = a(x)\psi(t) + b(x)\xi(t),$$

where $a(\cdot), b(\cdot) \geq 0$ satisfy $v \leq a(\cdot) + b(\cdot) \leq L$ and ψ', ξ' satisfy (A0), (Inc) _{$p-1$} and (Dec) _{$q-1$} , which includes the following examples:

$$t^p + a(x)t^q, \quad a(x)t^p + t^q, \quad a(x)t^p + b(x)t^q, \quad \text{and} \quad \psi(t) + a(x)\psi(t) \ln(e + t).$$

We present conditions for the above functions to satisfy (wVA1) or (VA1) in Corollaries 8.4 and 8.6, so that C^α - and $C^{1,\alpha}$ -regularity results for (2.5) are obtained as special cases. We note that the second example $a(x)t^p + t^q$ can be understood as a functional with standard q -growth and hence q/p has no upper bound to obtain the regularity results. Here, we explain the regularity results for this functional as a special case of double phase problems. In addition, in the same spirit, one could consider functionals with infinitely many phases such that

$$\varphi(x, t) = \sum_{i=1}^{\infty} a_i(x)t^{p_i}, \quad \text{where } 1 < p \leq p_i \leq q, \quad a_i(\cdot) \geq 0 \quad \text{and} \quad 0 < v \leq \sum_{i=1}^{\infty} a_i(\cdot) \leq L,$$

which satisfies the fundamental assumption of Theorem 1.1.

3. Generalized Orlicz spaces

Notation and assumptions

For $x_0 \in \mathbb{R}^n$ and $r > 0$, $B_r(x_0)$ is the ball in \mathbb{R}^n with radius r and center x_0 . We write $B_r = B_r(x_0)$ when the center is clear or unimportant. For an integrable function f in $U \subset \mathbb{R}^n$, we define $(f)_U$ to be the average of f in U in the integral sense, that is, $(f)_U := \int_U f \, dx := |U|^{-1} \int_U f \, dx$.

We say that $f : [0, \infty) \rightarrow [0, \infty)$ is *almost increasing* or *almost decreasing* if there exists $L \geq 1$ such that for any $0 < t < s < \infty$, $f(t) \leq Lf(s)$ or $f(s) \leq Lf(t)$, respectively. In particular, if $L = 1$ we say f is *non-decreasing* or *non-increasing*.

We refer to [51] for more details about basics of Φ -functions and generalized Orlicz spaces. For $\varphi : \Omega \times [0, \infty) \rightarrow [0, \infty)$ and $B_r \subset \Omega$, we write

$$\varphi_{B_r}^+(t) := \sup_{x \in B_r} \varphi(x, t) \quad \text{and} \quad \varphi_{B_r}^-(t) := \inf_{x \in B_r} \varphi(x, t).$$

If the map $t \mapsto \varphi(x, t)$ is non-decreasing for every $x \in \Omega$, then the (left-continuous) inverse function with respect to t is defined by

$$\varphi^{-1}(x, t) := \inf \{ \tau \geq 0 : \varphi(x, \tau) \geq t \}.$$

If φ is strictly increasing and continuous in t , then this is just the normal inverse function.

Definition 3.1. Let $\varphi : \Omega \times [0, \infty) \rightarrow [0, \infty)$ and $\gamma > 0$. We define some conditions related to regularity with respect to the t -variable.

- (aInc) $_\gamma$ The map $t \mapsto \varphi(x, t)/t^\gamma$ is almost increasing with constant $L \geq 1$ uniformly in $x \in \Omega$.
- (Inc) $_\gamma$ The map $t \mapsto \varphi(x, t)/t^\gamma$ is non-decreasing for every $x \in \Omega$.
- (aDec) $_\gamma$ The map $t \mapsto \varphi(x, t)/t^\gamma$ is almost decreasing with constant $L \geq 1$ uniformly in $x \in \Omega$.
- (Dec) $_\gamma$ The map $t \mapsto \varphi(x, t)/t^\gamma$ is non-increasing for every $x \in \Omega$.
- (A0) There exists $L \geq 1$ such that $L^{-1} \leq \varphi(x, 1) \leq L$ for every $x \in \Omega$.

Note that this version of (A0) is slightly stronger than the one used in [51], but they are equivalent under the doubling assumption (aDec). Let $0 < c \leq 1 \leq C < \infty$. If φ satisfies (aInc) $_\gamma$ with constant $L \geq 1$, then

$$\varphi(x, ct) \leq Lc^\gamma \varphi(x, t) \quad \text{and} \quad L^{-1}C^\gamma \varphi(x, t) \leq \varphi(x, Ct) \quad \text{for all } (x, t) \in \Omega \times [0, \infty).$$

On the other hand, if φ satisfies (aDec) $_\gamma$ with constant $L \geq 1$, then

$$L^{-1}c^\gamma \varphi(x, t) \leq \varphi(x, ct) \quad \text{and} \quad \varphi(x, Ct) \leq LC^\gamma \varphi(x, t) \quad \text{for all } (x, t) \in \Omega \times [0, \infty).$$

Remark 3.2. If φ satisfies (aInc) $_\gamma$ or (aDec) $_\gamma$ for some $\gamma > 0$, then so do $\varphi_{B_r}^-$ and $\varphi_{B_r}^+$ for any $B_r \subset \Omega$.

Remark 3.3. Suppose that $\varphi(x, \cdot) \in C^1([0, \infty))$ for each $x \in \Omega$ and that $\gamma > 0$. Then

- φ satisfies (Inc) $_\gamma$ if and only if $\gamma\varphi(x, t) \leq t\varphi'(x, t)$ for all $x \in \Omega$ and $t \in [0, \infty)$;
- φ satisfies (Dec) $_\gamma$ if and only if $\gamma\varphi(x, t) \geq t\varphi'(x, t)$ for all $x \in \Omega$ and $t \in [0, \infty)$.

These conclusions are obtained by differentiating the function $t \mapsto \varphi(x, t)/t^\gamma$.

For functions $f, g : U \rightarrow \mathbb{R}$ with $U \subset \mathbb{R}^n$, $f \lesssim g$ or $f \approx g$ (in U) mean that there exists $C \geq 1$ such that $f(y) \leq Cg(y)$ or $C^{-1}f(y) \leq g(y) \leq Cf(y)$, respectively, for all $y \in U$. In particular, in this paper we shall use these symbols when the relevant constants C depend only on n and the constants from the fundamental conditions $(\text{aInc})_\gamma$, $(\text{aDec})_\gamma$, $(\text{Inc})_\gamma$, $(\text{Dec})_\gamma$ and (A0) . By following this, for instance, (A0) can be written as $\varphi(\cdot, 1) \approx 1$ in Ω . We use some results from papers with a weaker notion of equivalence, $f \simeq g$ (in U), which means that there exists $C \geq 1$ such that $f(C^{-1}y) \leq g(y) \leq f(Cy)$ for all $y \in U$. However, if (aDec) holds, then \simeq and \approx are equivalent and furthermore the constants can be moved inside and outside of φ as observed above.

Basic properties of generalized φ -functions and related function spaces

We next introduce classes of Φ -functions. Let $L^0(\Omega)$ be the set of measurable functions on Ω .

Definition 3.4. Let $\varphi : \Omega \times [0, \infty) \rightarrow [0, \infty]$. We call φ a (generalized) Φ -prefunction if $x \mapsto \varphi(x, |f(x)|)$ is measurable for every $f \in L^0(\Omega)$, and $t \mapsto \varphi(x, t)$ is non-decreasing for every $x \in \Omega$ and satisfies $\varphi(x, 0) = \lim_{t \rightarrow 0^+} \varphi(x, t) = 0$ and $\lim_{t \rightarrow \infty} \varphi(x, t) = \infty$ for every $x \in \Omega$. A prefunction φ is a

- (1) (generalized weak) Φ -function, denoted $\varphi \in \Phi_w(\Omega)$, if it satisfies $(\text{aInc})_1$;
- (2) (generalized) convex Φ -function, denoted $\varphi \in \Phi_c(\Omega)$, if $t \mapsto \varphi(x, t)$ is left-continuous and convex for every $x \in \Omega$.

If φ is independent of x , then we write $\varphi \in \Phi_w$ or $\varphi \in \Phi_c$ without “ (Ω) ”. In what follows we omit the words “generalized” and “weak” from the parentheses.

We note that convexity implies $(\text{Inc})_1$ so that $\Phi_c(\Omega) \subset \Phi_w(\Omega)$. For $\varphi \in \Phi_w(\Omega)$, the generalized Orlicz space (also known as the Musielak–Orlicz space) is defined by

$$L^\varphi(\Omega) := \{f \in L^0(\Omega) : \|f\|_{L^\varphi(\Omega)} < \infty\}$$

with the (Luxemburg) norm

$$\|f\|_{L^\varphi(\Omega)} := \inf \{\lambda > 0 : \varrho_\varphi(f/\lambda) \leq 1\}, \quad \text{where} \quad \varrho_\varphi(f) := \int_\Omega \varphi(x, |f(x)|) dx.$$

We denote by $W^{1,\varphi}(\Omega)$ the set of $f \in L^\varphi(\Omega)$ such that $\partial_1 f, \dots, \partial_n f \in L^\varphi(\Omega)$, where $\partial_i f$ is the weak derivative of f in the x_i -direction, with the norm $\|f\|_{W^{1,\varphi}(\Omega)} := \|f\|_{L^\varphi(\Omega)} + \sum_i \|\partial_i f\|_{L^\varphi(\Omega)}$. Note that if φ satisfies $(\text{aDec})_q$ for some $q \geq 1$, then $f \in L^\varphi(\Omega)$ if and only if $\varrho_\varphi(f) < \infty$, and if φ satisfies (A0) , $(\text{aInc})_p$ and $(\text{aDec})_q$ for some $1 < p \leq q$, then $L^\varphi(\Omega)$ and $W^{1,\varphi}(\Omega)$ are reflexive Banach spaces. In addition we denote by $W_0^{1,\varphi}(\Omega)$ the closure of $C_0^\infty(\Omega)$ in $W^{1,\varphi}(\Omega)$. For more information about the generalized Orlicz and Orlicz–Sobolev spaces, we refer to the monographs [51, 69] and also [34, Chapter 2].

For $\varphi : [0, \infty) \rightarrow [0, \infty)$, we define the conjugate function by

$$\varphi^*(x, t) := \sup_{s \geq 0} (st - \varphi(x, s)).$$

By definition, we have the following *Young inequality*:

$$ts \leq \varphi(x, t) + \varphi^*(x, s) \quad \text{for all } s, t \geq 0.$$

If $\varphi \in \Phi_c(\Omega)$, then $(\varphi^*)^* = \varphi$ [34, Theorem 2.2.6].

We state some properties of Φ -functions, for which we refer to [51, Chapter 2].

Proposition 3.5. *Let φ be a Φ -prefunction.*

- (1) *If φ satisfies (aInc)₁, then there exists $\psi \in \Phi_c(\Omega)$ such that $\varphi \simeq \psi$.*
- (2) *If φ satisfies (aDec)₁, then there exists $\psi \in \Phi_c(\Omega)$ such that $\varphi \approx \psi^{-1}$. Note that $\psi^{-1}(x, \cdot)$ is concave.*
- (3) *Let $p, q \in (1, \infty)$. Then φ satisfies (aInc)_p or (aDec)_q if and only if φ^* satisfies (aDec)_{p/(p-1)} or (aInc)_{q/(q-1)}, respectively.*
- (4) *Let $\varphi \in \Phi_w(\Omega)$ and $\gamma \geq 1$. Then φ satisfies (aInc)_γ or (aDec)_γ if and only if φ^{-1} satisfies (aDec)_{1/γ} or (aInc)_{1/γ}, respectively.*
- (5) *If φ satisfies (aInc)_p and (aDec)_q, then for any $s, t \geq 0$ and $\kappa \in (0, 1)$,*

$$ts \leq \varphi(x, \kappa^{1/p}t) + \varphi^*(x, \kappa^{-1/p}s) \lesssim \kappa \varphi(x, t) + \kappa^{-1/p-1} \varphi^*(x, s)$$

and

$$ts \leq \varphi(x, \kappa^{-1/q'}t) + \varphi^*(x, \kappa^{1/q'}s) \lesssim \kappa^{-(q-1)} \varphi(x, t) + \kappa \varphi^*(x, s).$$

If $\varphi \in \Phi_c(\Omega)$, then there exists $\varphi' = \varphi'(x, t)$, which is non-decreasing and right-continuous, such that

$$\varphi(x, t) = \int_0^t \varphi'(x, s) ds.$$

Such φ' is called the *right-derivative* of φ . Note that this derivative was denoted by $\partial_t \varphi$ in the introduction.

We next collect some results about the derivative φ' . For (4), we give a simple direct proof, since earlier proofs of the inequality used additional assumptions.

Proposition 3.6. *Let $\gamma > 0$ and suppose that $\varphi \in \Phi_c(\Omega)$ with derivative φ' .*

- (1) *If φ' satisfies (aInc)_γ, (aDec)_γ, (Inc)_γ or (Dec)_γ, then φ satisfies (aInc)_{γ+1}, (aDec)_{γ+1}, (Inc)_{γ+1} or (Dec)_{γ+1}, respectively, with the same constant $L \geq 1$.*
- (2) *If φ' satisfies (aDec)_γ with constant L , then $\varphi(x, t) \approx t\varphi'(x, t)$, more precisely*

$$\frac{t\varphi'(x, t)}{2^{\gamma+1}L} \leq \varphi(x, t) \leq t\varphi'(x, t) \quad \text{for } (x, t) \in \Omega \times [0, \infty).$$

- (3) *If φ' satisfies (A0) and (aDec)_γ with constant $L \geq 1$, then φ also satisfies (A0), with constant depending on L and γ .*
- (4) *$\varphi^*(x, \varphi'(x, t)) \leq t\varphi'(x, t)$.*

Proof. We start with (1) and suppose that φ' satisfies $(\text{aInc})_\gamma$. Fix $0 < t < s < \infty$ and set $a := s/t > 1$. Then $(\text{aInc})_\gamma$ of φ' implies that

$$\begin{aligned} \frac{\varphi(x, t)}{t^{\gamma+1}} &= \frac{1}{t^{\gamma+1}} \int_0^t \varphi'(x, \tau) d\tau \\ &\leq \frac{L}{t^{\gamma+1}} \int_0^t \frac{\varphi'(x, a\tau)}{a^\gamma} d\tau \stackrel{\tilde{\tau}=a\tau}{=} \frac{L}{(at)^{\gamma+1}} \int_0^{at} \varphi'(x, \tilde{\tau}) d\tilde{\tau} = L \frac{\varphi(x, s)}{s^{\gamma+1}}, \end{aligned}$$

which means φ satisfies $(\text{aInc})_{\gamma+1}$. In the same way we can also prove that $(\text{aDec})_\gamma$ of φ' implies $(\text{aDec})_{\gamma+1}$ of φ . The claims regarding (Inc) and (Dec) follow when $L = 1$.

We next prove (2). Since φ' is non-decreasing, it follows that

$$\frac{t}{2} \varphi'(x, \frac{t}{2}) \leq \underbrace{\int_0^t \varphi'(x, \tau) d\tau}_{=\varphi(x, t)} \leq t \varphi'(x, t).$$

By the $(\text{aDec})_\gamma$ condition of φ' , we have $\varphi'(x, t/2) \geq L^{-1} 2^{-\gamma} \varphi'(x, t)$, which implies $\varphi(x, t) \approx t \varphi'(x, t)$.

Then, we prove (3). By (2) and (A0) of φ' it follows that $\varphi(\cdot, 1) \approx 1 \cdot \varphi'(\cdot, 1) \approx 1$, so φ satisfies (A0) .

Finally, we prove (4). Since φ is convex, $\varphi(x, s) \geq \varphi(x, t) + k(s - t)$, where $k := \varphi'(x, t)$ is the slope. Then from the definition of the conjugate function we have

$$\begin{aligned} \varphi^*(x, \varphi'(x, t)) &= \sup_{s \geq 0} (sk - \varphi(x, s)) \leq \sup_{s \geq 0} (sk - \varphi(x, t) - k(s - t)) \\ &= tk - \varphi(x, t) \leq t \varphi'(x, t). \end{aligned} \quad \blacksquare$$

We end this subsection with some properties of C^1 -regular Φ -functions. Note that Proposition 3.8(2) below is proved for C^2 -functions in [33, Lemma 3] – here we provide a more elementary proof which is based on a reduction to the same claim for the function t^p , that is,

$$(|x|^{p-2}x - |y|^{p-2}y) \cdot (x - y) \approx (|x| + |y|)^{p-2} |x - y|^2 \quad \text{for } p > 1. \quad (3.7)$$

While versions of this claim are commonly known, we have not found this precise formulation in the literature. Rather than providing a proof of (3.7), we just invoke [33, Lemma 3], since t^p is certainly a C^2 -function.

Proposition 3.8. *Let $\varphi \in \Phi_c \cap C^1([0, \infty))$ with φ' satisfying $(\text{Inc})_{p-1}$ and $(\text{Dec})_{q-1}$ for some $1 < p \leq q$. Then for $\kappa \in (0, \infty)$ and $x, y \in \mathbb{R}^n$ the following hold:*

- (1) $\frac{\varphi'(|x| + |y|)}{|x| + |y|} |x - y|^2 \approx \left(\frac{\varphi'(|x|)}{|x|} x - \frac{\varphi'(|y|)}{|y|} y \right) \cdot (x - y);$
- (2) $\frac{\varphi'(|x| + |y|)}{|x| + |y|} |x - y|^2 \lesssim \varphi(|x|) - \varphi(|y|) - \frac{\varphi'(|y|)}{|y|} y \cdot (x - y);$

$$(3) \quad \varphi(|x - y|) \lesssim \kappa[\varphi(|x|) + \varphi(|y|)] + \kappa^{-1} \frac{\varphi'(|x| + |y|)}{|x| + |y|} |x - y|^2.$$

If additionally $\varphi \in C^2((0, \infty))$, then $t\varphi''(t) \approx \varphi'(t)$ and $\frac{\varphi'(|x|+|y|)}{|x|+|y|}$ can be replaced by $\varphi''(|x| + |y|)$.

Proof. When $\varphi \in C^2((0, \infty))$, the inequalities $t\varphi''(t) \approx \varphi'(t)$ are direct consequences of Remark 3.3 and Proposition 3.6(1). This also implies $\frac{\varphi'(|x|+|y|)}{|x|+|y|} \approx \varphi''(|x| + |y|)$.

For (1), we may assume without loss of generality that $|x| \geq |y|$. By (Inc)_{p-1} and (Dec)_{q-1},

$$\left(\frac{|y|}{|x|}\right)^{q-1} \varphi'(|x|) \leq \varphi'(|y|) \leq \left(\frac{|y|}{|x|}\right)^{p-1} \varphi'(|x|).$$

Thus there exists $\gamma \in [p-1, q-1]$ such that $\varphi'(|y|) = \left(\frac{|y|}{|x|}\right)^\gamma \varphi'(|x|)$. Hence

$$\left(\frac{\varphi'(|x|)}{|x|}x - \frac{\varphi'(|y|)}{|y|}y\right) \cdot (x - y) = \frac{\varphi'(|x|)}{|x|^\gamma} (|x|^{\gamma-1}x - |y|^{\gamma-1}y) \cdot (x - y).$$

We use (3.7) with $\gamma + 1$ in place of p . Furthermore, $|x| \geq |y|$ implies $|x| + |y| \approx |x|$, and so

$$\left(\frac{\varphi'(|x|)}{|x|}x - \frac{\varphi'(|y|)}{|y|}y\right) \cdot (x - y) \approx \frac{\varphi'(|x|)}{|x|^\gamma} (|x| + |y|)^{\gamma-1} |x - y|^2 \approx \frac{\varphi'(|x| + |y|)}{|x| + |y|} |x - y|^2.$$

We next prove (2). Denote $\eta := \frac{x-y}{|x-y|}$ and $z_s := y + \eta s$. Then

$$\varphi(|x|) - \varphi(|y|) = \int_0^{|x-y|} \varphi'(|z_s|) \frac{z_s}{|z_s|} \cdot \eta \, ds.$$

Furthermore, since $x - y = \eta|x - y|$, we have

$$\begin{aligned} \varphi(|x|) - \varphi(|y|) - \frac{\varphi'(|y|)}{|y|}y \cdot (x - y) &= \int_0^{|x-y|} \left(\frac{\varphi'(|z_s|)}{|z_s|}z_s - \frac{\varphi'(|y|)}{|y|}y \right) \cdot (x - y) \, ds \\ &\approx \int_0^{|x-y|} \frac{\varphi'(|z_s| + |y|)}{|z_s| + |y|} |x - y| s \, ds, \end{aligned}$$

where the second step follows from (1) since $x - y = \frac{|x-y|}{s}(z_s - y)$. When $s \geq \frac{3}{4}|x - y|$, $|z_s| + |x| \approx |x| + |y|$ and (2) follows.

We finally prove (3). By the inequality $ab \leq \frac{1}{2}(a^2 + b^2)$, we find that

$$|x - y| \leq \frac{1}{2}\kappa(|x| + |y|) + \frac{1}{2}\kappa^{-1}(|x| + |y|)^{-1}|x - y|^2.$$

Therefore, since φ' is non-decreasing and $|x - y| \leq |x| + |y|$, we find by $t\varphi'(t) \approx \varphi(t)$ that

$$\begin{aligned} \varphi(|x - y|) &\lesssim \varphi'(|x| + |y|)|x - y| \\ &\leq \kappa\varphi'(|x| + |y|)(|x| + |y|) + \kappa^{-1}\varphi'(|x| + |y|)(|x| + |y|)^{-1}|x - y|^2 \\ &\approx \kappa[\varphi(|x|) + \varphi(|y|)] + \kappa^{-1} \frac{\varphi'(|x| + |y|)}{|x| + |y|} |x - y|^2. \end{aligned} \quad \blacksquare$$

4. Preliminary regularity results

Assumptions for higher regularity

Here we introduce the new assumptions that are used to obtain C^α -regularity for any $\alpha \in (0, 1)$ or $C^{1,\alpha}$ -regularity for some $\alpha \in (0, 1)$ of local minimizers of (1.2). We also restate the definition of (VA1) from the introduction, so that it can be more easily compared with its weaker variant, (wVA1).

In the next definition, we have several conditions which are assumed to hold “for any small ball”; this means that the condition holds for all $r < r_0$ for some $r_0 > 0$.

Definition 4.1. Let $\varphi \in \Phi_w(\Omega)$. We define some conditions related to regularity with respect to the x -variable.

(A1) There exists $L \geq 1$ such that for any $B_r \Subset \Omega$ with $|B_r| < 1$,

$$\varphi_{B_r}^+(t) \leq L\varphi_{B_r}^-(t) \quad \text{for all } t > 0 \text{ with } \varphi_{B_r}^-(t) \in [1, |B_r|^{-1}].$$

(VA1) There exists a non-decreasing continuous function $\omega : [0, \infty) \rightarrow [0, 1]$ with $\omega(0) = 0$ such that for any small $B_r \Subset \Omega$,

$$\varphi_{B_r}^+(t) \leq (1 + \omega(r))\varphi_{B_r}^-(t) \quad \text{for all } t > 0 \text{ with } \varphi_{B_r}^-(t) \in [\omega(r), |B_r|^{-1}].$$

(wVA1) For any $\varepsilon > 0$, there exists a non-decreasing continuous function $\omega = \omega_\varepsilon : [0, \infty) \rightarrow [0, 1]$ with $\omega(0) = 0$ such that for any small ball $B_r \Subset \Omega$,

$$\varphi_{B_r}^+(t) \leq (1 + \omega(r))\varphi_{B_r}^-(t) + \omega(r) \quad \text{for all } t > 0 \text{ with } \varphi_{B_r}^-(t) \in [\omega(r), |B_r|^{-1+\varepsilon}].$$

Intuitively, (A1) is a jump condition that restricts the amount that φ can jump between nearby points, whereas (VA1) and (wVA1) are continuity conditions that imply continuity with respect to the x -variable.

Remark 4.2. We see that (VA1) implies (wVA1), which in turn implies (A1). Assumption (VA1) is easier to understand but we emphasize that (wVA1) covers an interesting borderline case which has arisen in the double phase case: see Corollary 8.6.

Remark 4.3. Finally, we would like to explain why we adapt the methodology of the calculus of variations, instead of one of partial differential equations, since indeed u is a minimizer of (1.2) if and only if it is a weak solution to

$$\operatorname{div} \left(\frac{\varphi'(x, |Du|)}{|Du|} Du \right) = 0 \quad \text{in } \Omega$$

(see [53]). In the comparison step in our approach, we take advantage of the minimizing property of u . If we would instead use the PDE approach, to the best of our understanding, the main assumption (VA1) would be replaced by the assumption

$$(\varphi')_{B_r}^+(t) \leq (1 + \omega(r))(\varphi')_{B_r}^-(t) \quad \text{for all } t > 0 \text{ satisfying } \varphi_{B_r}^-(t) \in [\omega(r), |B_r|^{-1}].$$

Compared with (VA1), φ is replaced by φ' in the inequality. Since small values are not covered in this assumption or (VA1), these two assumptions are not comparable, i.e. one may hold but not the other, in either direction. However, if φ satisfies the basic assumption in Theorem 1.1 (this is always assumed in our main theorems), we show that (wVA1) is implied by this assumption: for any $\varepsilon > 0$, any small $B_r \Subset \Omega$, any $t > 0$ satisfying $\varphi_{B_r}^-(t) \in [\omega(r), |B_r|^{-1+\varepsilon}] \subset [\omega(r), |B_r|^{-1}]$ and any $x, y \in B_r$,

$$\begin{aligned} \varphi(x, t) &= \int_0^t \varphi'(x, s) ds \\ &\leq (1 + \omega(r)) \int_{(\varphi_{B_r}^-)^{-1}(\omega(r))}^t (\varphi')_{B_r}^-(s) ds + \int_0^{(\varphi_{B_r}^-)^{-1}(\omega(r))} \varphi'(x, s) ds \\ &\leq (1 + \omega(r)) \int_0^t \varphi'(y, s) ds + \varphi(x, (\varphi_{B_r}^-)^{-1}(\omega(r))) \\ &\leq (1 + \omega(r)) \varphi(y, t) + c\omega(r)^{p/q}. \end{aligned}$$

Thus (wVA1) holds with the function $c\omega(r)^{p/q}$. Moreover, we could also consider a (wVA1)-type assumption with φ' instead of φ , but the same argument shows that this also implies (wVA1).

We note that such difference between regularity assumptions for the minimizer and the PDE problem does not appear in types I–III. This also shows that regularity theory for general $\varphi(x, t)$ cannot be understood easily by just mixing the ones for types I–III.

Higher integrability and reverse Hölder type inequality

We prove higher integrability of minimizers of (1.2) and, as a corollary, a reverse Hölder type inequality. In this subsection we assume (A1).

The following higher integrability result appears as [52, Theorem 1.1] in the case $\delta = 1$. From the proof in that article, one can derive the stated dependence on δ with the help of the (aDec) $_q$ assumption; alternatively, one can use that result and a covering argument.

Lemma 4.4 (Higher integrability). *Let $\varphi \in \Phi_w(\Omega)$ satisfy (A0), (A1), (aInc) $_p$ and (aDec) $_q$ with constant $L \geq 1$ and $1 < p \leq q$. If $u \in W_{\text{loc}}^{1,\varphi}(\Omega)$ is a local minimizer of (1.2), then there exist $\sigma_0 = \sigma_0(n, p, q, L) > 0$, $c_1 = c_1(n, p, q, L) \geq 1$ and $\sigma_1 = \sigma_1(\sigma_0, n, q)$ such that*

$$\left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1}{1+\sigma_0}} \leq c_1 \delta^{-\sigma_1} \left(\int_{B_{(1+\delta)r}} \varphi(x, |Du|) dx + 1 \right) \quad (4.5)$$

for any $B_{2r} \Subset \Omega$ with $\|Du\|_{L^\varphi(B_{2r})} \leq 1$ and $\delta \in (0, 1]$.

Remark 4.6. Fix $\Omega' \Subset \Omega$. Since $\int_{\Omega'} \varphi(x, |Du|) dx < \infty$, there exists $R > 0$ such that

$$\int_{B_r} \varphi(x, |Du|) dx \leq 1 \quad (\text{or equivalently } \|Du\|_{L^\varphi(B_r)} \leq 1)$$

for $B_r \subset \Omega'$ with $r \leq R$. In view of the previous lemma, this means that $\varphi(\cdot, |Du|) \in L_{\text{loc}}^{1+\sigma_0}(\Omega)$.

The next lemma contains reverse Hölder type estimates for Du .

Lemma 4.7. *Let $\varphi \in \Phi_w(\Omega)$ satisfy (A0), (A1), (aInc) $_p$ and (aDec) $_q$ with constant $L \geq 1$ and $1 < p \leq q$. Suppose that $u \in W_{\text{loc}}^{1,\varphi}(\Omega)$ is a local minimizer of (1.2) and $B_{2r} \Subset \Omega$ with $\|Du\|_{L^\varphi(B_{2r})} \leq 1$. There exist $\sigma_0 = \sigma_0(n, p, q, L)$ and, for every $t \in (0, 1]$, $c_t = c(n, p, q, L, t) > 0$ such that*

$$\left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1}{1+\sigma_0}} \leq c_t \left(\left(\int_{B_{2r}} \varphi(x, |Du|)^t dx \right)^{1/t} + 1 \right) \quad (4.8)$$

and $c = c(n, p, q, L) \geq 1$ such that

$$\int_{B_r} \varphi(x, |Du|) dx \leq \left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1}{1+\sigma_0}} \leq c \left(\varphi_{B_{2r}}^- \left(\int_{B_{2r}} |Du| dx \right) + 1 \right).$$

Proof. We start with the first inequality. In (4.5) we split $\varphi = \varphi^\theta \varphi^{1-\theta}$ with $\theta \in (0, 1)$ and use Hölder's inequality with exponents $\frac{1+\sigma_0}{\theta}$ and $\frac{1+\sigma_0}{1+\sigma_0-\theta}$ and Young's inequality with exponents $\frac{1}{\theta}$ and $\frac{1}{1-\theta}$:

$$\begin{aligned} & \left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1}{1+\sigma_0}} \\ & \leq c_1 \left[\delta^{-\sigma_1} \left(\int_{B_{(1+\delta)r}} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{\theta}{1+\sigma_0}} \left(\int_{B_{2r}} \varphi(x, |Du|)^t dx \right)^{\frac{1-\theta}{t}} + 1 \right] \\ & \leq \frac{1}{2} \left(\int_{B_{(1+\delta)r}} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1}{1+\sigma_0}} + c_2 \delta^{-\frac{\sigma_1}{1-\theta}} \left(\int_{B_{2r}} \varphi(x, |Du|)^t dx \right)^{1/t} + c_1 \end{aligned}$$

where we have set $t := \frac{(1+\sigma_0)(1-\theta)}{1+\sigma_0-\theta}$. Now we see from a standard iteration lemma (e.g. [55, Lemma 4.2]) that the first claim holds.

We move on to the second claim. The first inequality directly follows from Hölder's inequality, hence we prove the second inequality. Taking $t = 1/q$ in (4.8), we see that

$$\left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1}{1+\sigma_0}} \leq c_{1/q} \left[\left(\int_{B_{2r}} \varphi_{B_{2r}}^+ (|Du|)^{1/q} dx \right)^q + 1 \right].$$

We notice that the map $t \mapsto [\varphi_{B_{2r}}^+(t)]^{1/q}$ satisfies (aDec) $_1$, since $\varphi_{B_{2r}}^+$ satisfies (aDec) $_q$. Therefore, by Jensen's inequality and Proposition 3.5(2), we have

$$\left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1}{1+\sigma_0}} \leq c \left[\varphi_{B_{2r}}^+ \left(\int_{B_{2r}} |Du| dx \right) + 1 \right] \quad (4.9)$$

for some $c = c(c_1, q, L) \geq 1$. In addition, since

$$\int_{B_{2r}} \varphi_{B_{2r}}^-(|Du|) dx \leq \int_{B_{2r}} \varphi(x, |Du|) dx \leq 1 \quad (\Longleftrightarrow \|Du\|_{L^\varphi(B_{2r})} \leq 1),$$

it follows by Jensen's inequality that $\varphi_{B_{2r}}^-(\int_{B_{2r}} |Du| dx) \lesssim |B_{2r}|^{-1}$. If also the inequality $\varphi_{B_{2r}}^-(\int_{B_{2r}} |Du| dx) \geq 1$ holds, then (A1) implies that

$$\varphi_{B_{2r}}^+\left(\int_{B_{2r}} |Du| dx\right) \leq L \varphi_{B_{2r}}^-\left(\int_{B_{2r}} |Du| dx\right),$$

whereas in the case $\varphi_{B_{2r}}^-(\int_{B_{2r}} |Du| dx) \leq 1$, (A0) gives an upper bound of c for the right-hand side of (4.9). ■

Regularity results for the autonomous case

In this subsection, we consider $\varphi \in \Phi_c \cap C^1([0, \infty)) \cap C^2((0, \infty))$ with φ' satisfying (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for some $1 < p \leq q$. Fix $v_0 \in W^{1,\varphi}(B_r)$ and let $v \in v_0 + W_0^{1,\varphi}(B_r)$ be a solution of the minimization problem

$$\min_{w \in v_0 + W_0^{1,\varphi}(B_r)} \int_{B_r} \varphi(|Dw|) dx, \quad (4.10)$$

or equivalently a weak solution to

$$\begin{cases} \operatorname{div}\left(\frac{\varphi'(|Dv|)}{|Dv|} Dv\right) = 0 & \text{in } B_r, \\ v = v_0 & \text{on } \partial B_r. \end{cases} \quad (4.11)$$

We start with the $C^{1,\alpha}$ -regularity in the autonomous case, with appropriate estimates.

Lemma 4.12. *Let $\varphi \in \Phi_c \cap C^1([0, \infty)) \cap C^2((0, \infty))$ with φ' satisfying (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for some $1 < p \leq q$. If $v \in W^{1,\varphi}(B_r)$ is a minimizer of (4.10) or a weak solution to (4.11), then $Dv \in C_{\text{loc}}^{\alpha_0}(B_r, \mathbb{R}^n)$ for some $\alpha_0 \in (0, 1)$ with the following estimates: for any $B_\rho(x_0) \subset B_r$,*

$$\sup_{B_{\rho/2}(x_0)} |Dv| \leq c \int_{B_\rho(x_0)} |Dv| dx, \quad (4.13)$$

and for any $\tau \in (0, 1)$,

$$\int_{B_{\tau\rho}(x_0)} |Dv - (Dv)_{B_{\tau\rho}(x_0)}| dx \leq c \tau^{\alpha_0} \int_{B_\rho(x_0)} |Dv| dx. \quad (4.14)$$

Here $\alpha_0 \in (0, 1)$ and $c > 0$ depend only on n , p and q .

The previous lemma is expected from [60]. In particular, we refer to [7] for the case $p \geq 2$. However, we cannot find any result treating the case $p < 2$ with the above estimates in the literature. Hence, we give a proof of the above lemma in Appendix A. We also note that (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for φ' are equivalent to $t\varphi''(t) \approx \varphi'(t)$ by Remark 3.3, since we assume $\varphi \in C^2((0, \infty))$.

We next state Calderón–Zygmund type estimates in B_r with non-zero boundary data.

Lemma 4.15 (Calderón–Zygmund estimates). *Let $\varphi \in \Phi_c \cap C^1([0, \infty)) \cap C^2((0, \infty))$ with φ' satisfying (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for some $1 < p \leq q$, and $|B_r| \leq 1$. If $v \in W^{1,\varphi}(B_r)$ is a minimizer of (4.10) or a weak solution to (4.11), then there exists $c = c(n, p, q, p_1, q_1, L) > 0$ such that*

$$\|\varphi(|Dv|)\|_{L^\theta(B_r)} \leq c \|\varphi(|Dv_0|)\|_{L^\theta(B_r)} \quad (4.16)$$

for any $\theta \in \Phi_w(B_r)$ satisfying (A0), (A1), (aInc) $_{p_1}$ and (aDec) $_{q_1}$ with constant $L \geq 1$ and $1 < p_1 \leq q_1$.

Moreover, fix $\kappa > 0$ and assume that $\int_{B_r} \theta(x, \varphi(|Dv_0|)) dx \leq \kappa$. Then there exists $c = c(n, p, q, p_1, q_1, L) > 0$ such that

$$\int_{B_r} \theta(x, \varphi(|Dv|)) dx \leq c(\kappa^{q_1/p_1-1} + 1) \left(\int_{B_r} \theta(x, \varphi(|Dv_0|)) dx + 1 \right). \quad (4.17)$$

Proof. In view of known results about gradient estimates for equations of p -Laplacian type or (4.11) (see for instance [14, 20, 70]), it is expected that for any $1 < s < \infty$ and any Muckenhoupt weight $w \in A_s$,

$$\int_{B_r} \varphi(|Dv|)^s w(x) dx \leq c \int_{B_r} \varphi(|Dv_0|)^s w(x) dx, \quad (4.18)$$

where $c > 0$ depends only on n, p, q, s and $[w]_{A_s}$ (see Appendix B for the definition of the Muckenhoupt class A_s). We outline the proof of (4.18) in Appendix B.

We may assume that $\|\varphi(|Dv_0|)\|_{L^\theta(B_r)} < \infty$, since otherwise (4.16) is trivial. Then $\|\varphi(|Dv_0|)\|_{L^{p_1}(B_r)} < \infty$ by (aInc) $_{p_1}$ of θ and so $\|\varphi(|Dv|)\|_{L^{p_1}(B_r)} < \infty$ by (4.18) with $s = p_1$. We define $\theta_j(x, t) := \min\{\theta(x, t), jt^{p_1}\}$, $j > 0$, and conclude that $\|\varphi(|Dv|)\|_{L^{\theta_j}(B_r)} < \infty$. Since $\varphi(|Dv|) \in L^{\theta_j}(B_r)$, extrapolation for the generalized Orlicz functions (see [51, Corollary 5.3.4]) gives

$$\|\varphi(|Dv|)\|_{L^{\theta_j}(B_r)} \lesssim \|\varphi(|Dv_0|)\|_{L^{\theta_j}(B_r)} \leq \|\varphi(|Dv_0|)\|_{L^\theta(B_r)}.$$

We note that in the statement of [51, Corollary 5.3.4], φ is also assumed to satisfy the so-called (A2) condition, which is however not needed if the domain Ω is bounded [51, Lemma 4.2.3], and in our case, $\Omega = B_r$. Finally, (4.16) follows from this by monotone convergence: $\|\varphi(|Dv|)\|_{L^\theta(B_r)} = \lim_{j \rightarrow \infty} \|\varphi(|Dv|)\|_{L^{\theta_j}(B_r)}$ [51, Lemma 3.1.4].

We next prove the second claim, inequality (4.17). If $\int_{B_r} \theta(x, \varphi(|Dv_0|)) dx \geq 1$, then it follows from (4.16) by [51, Lemma 3.2.10] that

$$\begin{aligned} \int_{B_r} \theta(x, \varphi(|Dv|)) dx &\leq c \left(\int_{B_r} \theta(x, \varphi(|Dv_0|)) dx \right)^{q_1/p_1} \\ &\leq c\kappa^{q_1/p_1-1} \int_{B_r} \theta(x, \varphi(|Dv_0|)) dx, \end{aligned}$$

which implies (4.17).

Now, we suppose that $\int_{B_r} \theta(x, \varphi(|Dv_0|)) dx < 1$. We assume first that the (A1) inequality holds also in $[0, 1]$, i.e., for some $L_1 \geq 1$,

$$\theta_{B_\rho}^+(t) \leq L_1 \theta_{B_\rho}^-(t) \quad \text{for all } t > 0 \text{ satisfying } \theta_{B_\rho}^-(t) \in [0, |B_\rho|^{-1}], \quad (4.19)$$

whenever $B_\rho \subset B_r$. Define $\theta^\pm(t) := \theta_{B_r}^\pm(t)$,

$$M := (\theta^-)^{-1} \left(\int_{B_r} \theta(x, \varphi(|Dv_0|)) dx \right) \quad \text{and} \quad \bar{\theta}(x, t) := \frac{\theta(x, Mt)}{\theta^-(M)}.$$

Note that $\theta^-(M) \in [0, 1]$. Then $\bar{\theta}$ also satisfies (aInc) $_{p_1}$ and (aDec) $_{q_1}$, with the same constants as θ . We next prove that $\bar{\theta}$ satisfies (A0). It is clear that $\bar{\theta}^-(1) = 1$. On the other hand, since $\theta^-(M) \in [0, 1]$, we see by (4.19) with $B_\rho = B_r$ that $\bar{\theta}^+(1) = \theta^+(M)/\theta^-(M) \leq L_1$. Finally, we show that $\bar{\theta}$ satisfies (A1). Let $B_\rho \subset B_r$ and consider $t > 0$ such that $\bar{\theta}_{B_\rho}^-(t) \in [1, |B_\rho|^{-1}]$. Then $\theta_{B_\rho}^-(Mt) = \bar{\theta}_{B_\rho}^-(t)\theta^-(M) \leq |B_\rho|^{-1}$. Therefore, in view of (4.19), we have

$$\bar{\theta}_{B_\rho}^+(t) = \frac{\theta_{B_\rho}^+(Mt)}{\theta^-(M)} \leq \frac{L_1 \theta_{B_\rho}^-(Mt)}{\theta^-(M)} = L_1 \bar{\theta}_{B_\rho}^-(t)$$

so that $\bar{\theta}$ satisfies the (A1) condition with constant L_1 .

Let $m := \varphi^{-1}(M)$ and set

$$\bar{v} := \frac{v}{m}, \quad \bar{v}_0 := \frac{v_0}{m}, \quad \text{and} \quad \bar{\varphi}(t) := \frac{\varphi(mt)}{M}.$$

Then $\bar{\varphi}'(t) = \frac{\varphi'(mt)m}{M}$ and $\bar{v} \in W^{1,\bar{\varphi}}(B_r)$ is a weak solution to

$$\operatorname{div} \left(\frac{\bar{\varphi}'(|D\bar{v}|)}{|D\bar{v}|} D\bar{v} \right) = 0 \quad \text{in } B_r \quad \text{with} \quad \bar{v} = \bar{v}_0 \quad \text{on } \partial B_r.$$

Note that $\bar{\varphi}'$ also satisfies (Inc) $_{p-1}$ and (Dec) $_{q-1}$ with the same constant as φ' . In addition, by the definitions of $\bar{\theta}$, $\bar{\varphi}$ and M ,

$$\begin{aligned} \int_{B_r} \bar{\theta}(x, \bar{\varphi}(|D\bar{v}_0|)) dx &= \frac{1}{\theta^-(M)} \int_{B_r} \theta(x, \varphi(|Dv_0|)) dx \leq 1 \\ &\implies \|\bar{\varphi}(|D\bar{v}_0|)\|_{L^{\bar{\theta}}(B_r)} \leq 1. \end{aligned}$$

Therefore, applying (4.16) to $(\theta, \varphi, v, v_0) = (\bar{\theta}, \bar{\varphi}, \bar{v}, \bar{v}_0)$, we have

$$\|\bar{\varphi}(|D\bar{v}|)\|_{L^{\bar{\theta}}(B_r)} \leq c \|\bar{\varphi}(|D\bar{v}_0|)\|_{L^{\bar{\theta}}(B_r)} \leq c$$

for some $c = c(n, p, q, p_1, q_1, L_1) \geq 1$. Finally, this implies that

$$\frac{1}{\theta^-(M)} \int_{B_r} \theta(x, \varphi(|Dv|)) dx = \int_{B_r} \bar{\theta}(x, \bar{\varphi}(|D\bar{v}|)) dx \leq c$$

for some $c = c(n, p, q, p_1, q_1, L_1) \geq 1$. In view of the definition of M , we have

$$\int_{B_r} \theta(x, \varphi(|Dv|)) dx \leq c \int_{B_r} \theta(x, \varphi(|Dv_0|)) dx \quad (4.20)$$

in the case when (4.19) holds. Note that (4.20) is stronger than (4.17), but requires the stronger assumption (4.19).

We return to the case that θ satisfies (A1) with normal range and define

$$\tilde{\theta}(x, t) := \theta(x, t) + t^{p_1}.$$

It is easy to check that $\tilde{\theta}$ satisfies (A0), (aInc) $_{p_1}$, (aDec) $_{q_1}$ and $\theta(x, t) \leq \tilde{\theta}(x, t) \lesssim \theta(x, t) + 1$. Let us show that $\tilde{\theta}$ satisfies (4.19). Fix $B_\rho \subset B_r$ and $t \geq 0$ satisfying $\tilde{\theta}_{B_\rho}^-(t) \in [0, |B_\rho|^{-1}]$. Then

$$\theta_{B_\rho}^-(t) = \tilde{\theta}_{B_\rho}^-(t) - t^{p_1} \leq |B_\rho|^{-1}.$$

If $\theta_{B_\rho}^-(t) \geq 1$, then (A1) for θ implies that

$$\tilde{\theta}_{B_\rho}^+(t) = \theta_{B_\rho}^+(t) + t^{p_1} \leq L\theta_{B_\rho}^-(t) + t^{p_1} \leq L(\theta_{B_\rho}^-(t) + t^{p_1}) = L\tilde{\theta}_{B_\rho}^-(t).$$

On the other hand, if $\theta_{B_\rho}^-(t) \leq 1$, by (A0), (aInc) $_{p_1}$ and (aDec) $_{q_1}$ for θ we have $t \lesssim 1$ and then $\theta_{B_\rho}^+(t) \approx t^{p_1} \leq \theta_{B_\rho}^-(t)$. Hence $\tilde{\theta}$ satisfies (4.19). Finally, since $\int_{B_r} \tilde{\theta}(x, \varphi(|Dv_0|)) dx \leq c(1 + |B_r|) \leq \tilde{c}$, applying the result (4.20) for the function $\tilde{c}^{-1}\tilde{\theta}(x, t)$, we obtain

$$\begin{aligned} \int_{B_r} \theta(x, \varphi(|Dv|)) dx &\leq \int_{B_r} \tilde{\theta}(x, \varphi(|Dv|)) dx \leq c \int_{B_r} \tilde{\theta}(x, \varphi(|Dv_0|)) dx \\ &\leq c \int_{B_r} [\theta(x, \varphi(|Dv_0|)) + 1] dx, \end{aligned}$$

which completes the proof of (4.17). ■

5. Comparison results without continuity assumption

Assume that $\varphi \in \Phi_c(\Omega) \cap C^1([0, \infty))$ satisfies a stronger version of (A1): there exists $L \geq 1$ and a non-decreasing continuous function $\omega : [0, \infty) \rightarrow [0, 1]$ with $\omega(0) = 0$ such that for any small $B_r \Subset \Omega$,

$$\varphi_{B_r}^+(t) \leq L\varphi_{B_r}^-(t) \quad \text{for all } t > 0 \text{ with } \varphi_{B_r}^-(t) \in [\omega(r), |B_r|^{-1}]. \quad (5.1)$$

Note that this condition is implied by (wVA1) with $L = 3$ and $\omega = \omega_\varepsilon$ for any fixed ε . Further, we assume that φ' satisfies (A0) with the same constant $L \geq 1$, as well as (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for some $1 < p \leq q$.

We fix $\Omega' \Subset \Omega$ and consider $B_{2r} = B_{2r}(x_0) \subset \Omega'$ with $r > 0$ satisfying

$$r \leq \frac{1}{2}, \quad \omega(2r) \leq \frac{1}{L}, \quad |B_{2r}| \leq \min \left\{ \frac{1}{2L}, 2^{-\frac{2(1+\sigma_0)}{\sigma_0}} \left(\int_{\Omega'} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{-\frac{2+\sigma_0}{\sigma_0}} \right\}, \quad (5.2)$$

where $\sigma_0 \in (0, 1)$ is given in Lemma 4.4. Note that $\varphi(\cdot, |Du|) \in L_{\text{loc}}^{1+\sigma_0}(\Omega)$ (see Remark 4.6). Hence from Hölder's inequality and (5.2) we have

$$\int_{B_{2r}} \varphi(x, |Du|)^{1+\sigma_0/2} dx \leq |B_{2r}| \left(\int_{B_{2r}} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1+\sigma_0/2}{1+\sigma_0}} \leq \frac{1}{2}, \quad (5.3)$$

so that

$$\int_{B_{2r}} \varphi(x, |Du|) dx \leq \int_{B_{2r}} \varphi(x, |Du|)^{1+\sigma_0/2} dx + |B_{2r}| \leq \frac{1}{2} + \frac{1}{2} = 1. \quad (5.4)$$

Therefore, we can take advantage of the results in Lemmas 4.4 and 4.7. For convenience, we write $\varphi^\pm(t) := \varphi_{B_{2r}}^\pm(t)$.

Construction of a regularized Orlicz function

We construct a regularized function $\tilde{\varphi} \in C^1([0, \infty)) \cap C^2((0, \infty))$ with $t\tilde{\varphi}''(t) \approx \tilde{\varphi}'(t)$, which is independent of the x variable and sufficiently close to $\varphi(x_0, t)$ in a suitable range of t . This procedure is quite delicate since we want improved differentiability, and moreover we want to find $\tilde{\varphi}$ satisfying in particular the assumptions of Proposition 5.12 below. The challenge lies in ensuring that $\varphi(x, \tilde{\varphi}^{-1}(t))$ satisfies (aInc)₁ and (aDec) _{γ} with some $\gamma > 1$ for small and large values of t , as we only have the comparison property when t is in some range $[t_1, t_2]$. We approach this problem by requiring p -growth for small and large values of t . This is counter-intuitive, because it means that the resulting function is neither a lower nor an upper bound of the original function, in contrast to estimates used in previous articles.

We first define

$$B := B_{2r} = B_{2r}(x_0), \quad t_1 := (\varphi^-)^{-1}(\omega(2r)) \quad \text{and} \quad t_2 := (\varphi^-)^{-1}(|B|^{-1}). \quad (5.5)$$

Note that it follows from $\omega(2r), |B_{2r}| \leq L^{-1}$ in (5.2) and (A0) of φ that $t_1 \leq 1 \leq t_2$. Let

$$\psi_B(t) := \begin{cases} a_1(t/t_1)^{p-1} & \text{if } 0 \leq t < t_1, \\ \varphi'(x_0, t) & \text{if } t_1 \leq t \leq t_2, \\ a_2(t/t_2)^{p-1} & \text{if } t_2 < t < \infty, \end{cases} \quad (5.6)$$

where the constants $a_1 := \varphi'(x_0, t_1)$ and $a_2 := \varphi'(x_0, t_2)$ are chosen so that ψ_B is continuous. We then define

$$\varphi_B(t) := \int_0^t \psi_B(s) ds.$$

Note that these functions depend on B via the center point x_0 as well as the values t_1 and t_2 .

When $t \in [t_1, t_2]$, the coincidence of derivatives implies that

$$\varphi_B(t) - \varphi(x_0, t) = \varphi_B(t_1) - \varphi(x_0, t_1) = \frac{1}{p} t_1 \varphi'(x_0, t_1) - \varphi(x_0, t_1)$$

and so, using the facts that $\frac{1}{q} t_1 \varphi'(x_0, t_1) \leq \varphi(x_0, t_1) \leq \frac{1}{p} t_1 \varphi'(x_0, t_1)$ by [\(Inc\)_p](#) and [\(Dec\)_q](#) as well as [\(5.1\)](#), we find that

$$0 \leq \varphi_B(t) - \varphi(x_0, t) \leq (q/p - 1) \varphi(x_0, t_1) \approx \varphi^-(t_1) = \omega(2r) \quad \text{for all } t \in [t_1, t_2]. \quad (5.7)$$

Fix $\eta \in C_0^\infty(\mathbb{R})$ with $\eta \geq 0$, $\text{supp } \eta \subset (0, 1)$ and $\|\eta\|_1 = 1$. We define

$$\tilde{\varphi}(t) := \int_0^\infty \varphi_B(t\sigma) \eta_r(\sigma - 1) d\sigma = \int_0^\infty \varphi_B(s) \eta_{rt}(s - t) ds \quad \text{where} \quad \eta_r(t) := \frac{1}{r} \eta\left(\frac{t}{r}\right); \quad (5.8)$$

the second expression is valid for $t > 0$. From the second formula, we see that $\tilde{\varphi} \in C^\infty((0, \infty))$.

For the next proof, we recall the following elementary inequalities which follow from the mean value theorem for $s \mapsto (1 + s)^\gamma$ on $[0, 1]$: for $\gamma > 0$ and $0 < s \leq 1$,

$$1 + \min\{1, 2^{\gamma-1}\} \gamma s \leq (1 + s)^\gamma \leq 1 + \max\{1, 2^{\gamma-1}\} \gamma s. \quad (5.9)$$

For the functions defined above, we have the following properties.

Proposition 5.10. *Let $\tilde{\varphi}$ be from [\(5.8\)](#). Then:*

- (1) $\varphi_B(t) \leq \tilde{\varphi}(t) \leq (1 + cr)\varphi_B(t) \leq c\varphi_B(t)$ for all $t > 0$ with $c > 0$ depending only on q . Furthermore,

$$0 \leq \tilde{\varphi}(t) - \varphi(x_0, t) \leq c(r\varphi^-(t) + \omega(2r)) \leq c\varphi^-(t) \quad \text{for all } t \in [t_1, t_2].$$

- (2) $\tilde{\varphi} \in C^1([0, \infty))$ and it satisfies [\(A0\)](#), [\(Inc\)_p](#) and [\(Dec\)_q](#) while $\tilde{\varphi}'$ satisfies [\(A0\)](#), [\(Inc\)_{p-1}](#) and [\(Dec\)_{q-1}](#). In particular, $\tilde{\varphi}'(t) \approx t\tilde{\varphi}''(t)$ for all $t > 0$.

- (3) $\tilde{\varphi}(t) \leq c\varphi(x, t)$ for all $(x, t) \in B \times [1, \infty)$, and so $\tilde{\varphi}(t) \leq c(\varphi(x, t) + 1)$ for all $(x, t) \in B \times [0, \infty)$.

Here, the constants $c > 0$ depend only on n , p , q and L .

Proof. It follows from the construction that ψ_B satisfies [\(Inc\)_{p-1}](#), [\(Dec\)_{q-1}](#) and [\(A0\)](#). By Proposition 3.6, φ and φ_B satisfy [\(Inc\)_p](#), [\(Dec\)_q](#) and [\(A0\)](#).

(1) We note that $\eta_r(\sigma - 1)$ is only non-zero when $\sigma - 1 \in [0, r]$. As φ_B is increasing and $\eta \geq 0$, we obtain

$$\tilde{\varphi}(t) = \int_0^\infty \varphi_B(t\sigma) \eta_r(\sigma - 1) d\sigma \leq \int_1^{1+r} \varphi_B((1+r)t) \eta_r(\sigma - 1) d\sigma = \varphi_B((1+r)t)$$

since $\|\eta_r\|_1 = 1$. Similarly, we obtain $\tilde{\varphi}(t) \geq \varphi_B(t)$. In addition, by (Dec)_q for φ_B and (5.9), we have

$$\tilde{\varphi}(t) \leq \varphi_B((1+r)t) \leq (1+r)^q \varphi_B(t) \leq (1+2^{q-1}qr) \varphi_B(t) \quad \text{for all } t \geq 0.$$

By this inequality and (5.7), we estimate

$$\begin{aligned} \tilde{\varphi}(t) - \varphi(x_0, t) &\leq (1+2^{q-1}qr) \varphi_B(t) - \varphi_B(t) + c\omega(2r) = 2^{q-1}qr \varphi_B(t) + c\omega(2r) \\ &\lesssim r\varphi^-(t) + \omega(2r) \end{aligned}$$

for all $t \in [t_1, t_2]$, where we have also used (5.7) and (5.1) to estimate φ_B in the last step. In addition, we know that $\omega(2r) = \varphi^-(t_1) \leq \varphi^-(t)$ for all $t \in [t_1, t_2]$. The lower bound follows from $\tilde{\varphi}(t) \geq \varphi_B(t) \geq \varphi(x_0, t)$.

(2) The claims for $\tilde{\varphi}'$ will be derived from the equality

$$\tilde{\varphi}'(t) = \int_0^\infty \sigma \psi_B(t\sigma) \eta_r(\sigma-1) d\sigma \quad \text{for } t \geq 0, \quad (5.11)$$

which is obtained by differentiating under the integral sign. The continuity of ψ_B implies that $\tilde{\varphi}' \in C([0, \infty))$ and so $\tilde{\varphi} \in C^1([0, \infty))$. As in (1), since the support of η_r is in $[0, r]$, $\|\eta_r\|_1 = 1$, and since ψ_B is increasing and satisfies (Dec)_{q-1}, we see that

$$\psi_B(t) \leq \tilde{\varphi}'(t) \leq (1+r)\psi_B((1+r)t) \leq (1+r)^q \psi_B(t) \leq 2^q \psi_B(t) \quad \text{for all } t > 0,$$

that is, $\psi_B \approx \tilde{\varphi}'$. Hence $\tilde{\varphi}'(1) \approx \psi_B(1) = \varphi'(x_0, 1)$, which implies that $\tilde{\varphi}'$ satisfies (A0). From the expression for $\tilde{\varphi}'$, we also see, since ψ_B satisfies (Inc)_{p-1}, that

$$\tilde{\varphi}'(\lambda t) = \int_0^\infty \sigma \psi_B(\lambda t \sigma) \eta_r(\sigma-1) d\sigma \geq \lambda^{p-1} \int_0^\infty \sigma \psi_B(t\sigma) \eta_r(\sigma-1) d\sigma = \lambda^{p-1} \tilde{\varphi}'(t)$$

for $\lambda \geq 1$ and $t > 0$. This yields (Inc)_{p-1} for $\tilde{\varphi}'$. Similarly we prove that $\tilde{\varphi}'$ satisfies (Dec)_{q-1}. The properties for $\tilde{\varphi}$ follow by Proposition 3.6.

(3) Fix $x \in B$. When $t \in [1, t_2]$, we see by (1) and (5.1) that $\tilde{\varphi}(t) \lesssim \varphi^-(t) \leq \varphi(x, t)$. For $t \geq t_2$, we observe that

$$\psi_B(t) = \varphi'(x_0, t_2) \left(\frac{t}{t_2} \right)^{p-1} \approx \varphi'(x, t_2) \left(\frac{t}{t_2} \right)^{p-1} \leq \varphi'(x, t),$$

since $\varphi'(x_0, t_2) \approx \varphi(x_0, t_2)/t_2 \approx \varphi(x, t_2)/t_2 \approx \varphi'(x, t_2)$ by (5.1) and φ' satisfies (Inc)_{p-1}. Then

$$\varphi_B(t) = \varphi_B(t_2) + \int_{t_2}^t \psi_B(\tau) d\tau \lesssim \varphi(x, t_2) + \int_{t_2}^t \varphi'(x, \tau) d\tau = \varphi(x, t). \quad \blacksquare$$

The next result shows the strength of the approach with (aInc) and (aDec), since it would be difficult to construct an approximating function to guarantee (Inc) and (Dec).

Proposition 5.12. *For $\tilde{\varphi}$ defined in (5.8) and any $\sigma \in (0, 1)$, set*

$$\theta(x, t) := [\varphi(x, \tilde{\varphi}^{-1}(t))]^{1+\sigma}.$$

Then $\theta \in \Phi_w(B_r)$ satisfies (A0), (aInc) $_{1+\sigma}$, (aDec) $_{q(1+\sigma)/p}$ and (A1). Here the constants depend only on n, p, q and L (from the assumptions on φ) and are independent of σ .

Proof. That $\theta \in \Phi_w(B_r)$ is clear once we show (aInc) $_1$. As φ and $\tilde{\varphi}$ satisfy (A0), so does θ . Now we prove that θ satisfies (aInc) $_{1+\sigma}$, which holds if $t \mapsto \varphi(x, \tilde{\varphi}^{-1}(t))$ satisfies (aInc) $_1$. Let $t > s > 0$, $\tilde{\varphi}(\tau) = t$ and $\tilde{\varphi}(\sigma) = s$. Then

$$L \frac{\varphi(x, \tilde{\varphi}^{-1}(t))}{t} \geq \frac{\varphi(x, \tilde{\varphi}^{-1}(s))}{s} \iff L \frac{\varphi(x, \tau)}{\tilde{\varphi}(\tau)} \geq \frac{\varphi(x, \varsigma)}{\tilde{\varphi}(\varsigma)}.$$

By Proposition 5.10 and Remark 3.3, we have $\tilde{\varphi}(t) \approx \varphi_B(t)$, $\varphi(x, t) \approx t\varphi'(x, t)$ and $\varphi_B(t) \approx t\psi_B(t)$ for all $(x, t) \in B \times (0, \infty)$. Therefore, it suffices to show that the function $t \mapsto \varphi'(x, t)/\psi_B(t)$ is almost increasing. Let t_1 and t_2 be from (5.5). Then by the definition of ψ_B in (5.6) we see that $t \mapsto \varphi'(x, t)/\psi_B(t)$ is non-decreasing on $(0, t_1] \cup [t_2, \infty)$, since φ' satisfies (Inc) $_{p-1}$. By (5.1) together with $t\varphi'(x, t) \approx \varphi(x, t)$, we have $\varphi'(x, t)/\psi_B(t) \approx 1$ in $[t_1, t_2]$. Therefore, we see that $t \mapsto \varphi'(x, t)/\psi_B(t)$ is almost increasing. The property (aDec) $_{q(1+\sigma)/p}$ is proved analogously.

Finally, we show that θ satisfies (A1). Let $B_\rho \subset B_r$, and assume $\theta_{B_\rho}^-(t) \in [1, |B_\rho|^{-1}]$. Then

$$\varphi_{B_\rho}^-(\tilde{\varphi}^{-1}(t)) = \theta_{B_\rho}^-(t)^{1/(1+\sigma)} \in [1, |B_\rho|^{-1/(1+\sigma)}] \subset [1, |B_\rho|^{-1}].$$

Therefore, (A1) for φ implies that

$$\theta_{B_\rho}^+(t) = [\varphi_{B_\rho}^+(\tilde{\varphi}^{-1}(t))]^{1+\sigma} \leq [L\varphi_{B_\rho}^-(\tilde{\varphi}^{-1}(t))]^{1+\sigma} \leq L^2\theta_{B_\rho}^-(t)$$

and so θ satisfies (A1). ■

Comparison estimates

Let $\tilde{\varphi} : [0, \infty) \rightarrow [0, \infty)$ be the function constructed in the previous subsection. We then consider the minimizer $v \in W^{1, \tilde{\varphi}}(B_r)$ of

$$\int_{B_r} \tilde{\varphi}(|Dv|) dx \quad \text{with} \quad v = u \text{ on } \partial B_r, \quad (5.13)$$

where $u \in W_{\text{loc}}^{1, \varphi}(\Omega)$ is a minimizer of (1.2), and derive a comparison estimate between the gradients of u and v . We note from Proposition 5.10(3) that $u \in W^{1, \tilde{\varphi}}(B_r)$, so it is an appropriate boundary-value function and thus there exists a unique minimizer of (5.13). The minimizer v is also a weak solution to

$$\operatorname{div} \left(\frac{\tilde{\varphi}'(|Dv|)}{|Dv|} Dv \right) = 0 \quad \text{in } B_r \quad \text{with} \quad v = u \quad \text{on } \partial B_r. \quad (5.14)$$

Before stating the main comparison result, we observe the following reverse Hölder type estimate for Du and Calderón–Zygmund type estimate for the problem (5.13).

Lemma 5.15. *Let $u \in W_{\text{loc}}^{1,\varphi}(\Omega)$ be a local minimizer of (1.2) and $v \in W^{1,\tilde{\varphi}}(B_r)$ be the minimizer of (5.13), where $B_{2r} \Subset \Omega$ with $r > 0$ satisfying (5.2) and $\tilde{\varphi}$ is defined in (5.8). Then*

$$\left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1}{1+\sigma_0}} \leq c \left[\tilde{\varphi} \left(\int_{B_{2r}} |Du| dx \right) + 1 \right] \quad (5.16)$$

and

$$\begin{aligned} \int_{B_r} \varphi(x, |Dv|) dx &\leq \left(\int_{B_r} \varphi(x, |Dv|)^{1+\sigma_0/2} dx \right)^{\frac{1}{1+\sigma_0/2}} \\ &\leq c \left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma_0/2} dx + 1 \right)^{\frac{1}{1+\sigma_0/2}}. \end{aligned} \quad (5.17)$$

Moreover,

$$\int_{B_r} |Dv| dx \leq c \left(\int_{B_{2r}} |Du| dx + 1 \right). \quad (5.18)$$

Here the constants $c \geq 1$ depend on n, p, q and L .

Proof. We first prove (5.16). We note that u satisfies the reverse Hölder inequality (4.5) for some $c_1 = c_1(n, p, q, L) \geq 1$. Then by Lemma 4.7, we have

$$\begin{aligned} \left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1}{1+\sigma_0}} &\lesssim \varphi^- \left(\int_{B_{2r}} |Du| dx \right) + 1 \\ &\leq \varphi(x_0, t_0) + 1, \end{aligned}$$

where $t_0 := \int_{B_{2r}} |Du| dx$. This and (A0) imply that (5.16) holds when $t_0 \leq 1$; we therefore assume that $t_0 \geq 1$. By Jensen's inequality and (5.4),

$$1 \leq t_0 \lesssim (\varphi^-)^{-1} \left(\int_{B_{2r}} \varphi^- (|Du|) dx \right) \leq (\varphi^-)^{-1} (|B_{2r}|^{-1}) = t_2,$$

where t_2 is defined in (5.5). Therefore, it follows from Proposition 5.10(1) that

$$\left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma_0} dx \right)^{\frac{1}{1+\sigma_0}} \lesssim \varphi(x_0, t_0) \lesssim \tilde{\varphi}(t_0) = \tilde{\varphi} \left(\int_{B_{2r}} |Du| dx \right).$$

As for (5.17), we only prove the second inequality, since the first inequality directly follows from Hölder's inequality. Let $\theta \in \Phi_w(B_r)$ be from Proposition 5.12 with $\sigma = \sigma_0/2$. By the proposition, we may apply Lemma 4.15 with $v_0 = u$ and $\kappa = 1$ (see (5.3)) to conclude that

$$\begin{aligned} \int_{B_r} \varphi(x, |Dv|)^{1+\sigma_0/2} dx &= \int_{B_r} \theta(x, \tilde{\varphi}(|Dv|)) dx \lesssim \int_{B_r} [\theta(x, \tilde{\varphi}(|Du|)) + 1] dx \\ &\approx \int_{B_r} [\varphi(x, |Du|) + 1]^{1+\sigma_0/2} dx. \end{aligned}$$

By Jensen's inequality, Proposition 5.10(3), (5.17) and (5.16),

$$\tilde{\varphi}\left(\int_{B_r} |Dv| dx\right) \leq \int_{B_r} \tilde{\varphi}(|Dv|) dx \lesssim \int_{B_r} [\varphi(x, |Dv|) + 1] dx \lesssim \tilde{\varphi}\left(\int_{B_{2r}} |Du| dx\right) + 1.$$

Then (5.18) follows when we apply $\tilde{\varphi}^{-1}$ to both sides and use (aDec) $_q$ to move “+1” inside $\tilde{\varphi}$. ■

6. Comparison results with continuity assumption

Assume that $\varphi \in \Phi_c(\Omega) \cap C^1([0, \infty))$ satisfies (wVA1), in addition to the assumptions at the beginning of the previous section, i.e. (A0) with constant $L \geq 1$, as well as (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for some $1 < p \leq q$. At this stage, we fix

$$\varepsilon_0 := \frac{\sigma_0}{2(2 + \sigma_0)} \quad (6.1)$$

where $\sigma_0 \in (0, 1)$ is determined in Lemma 4.4. We will use (wVA1) for $\varepsilon = \varepsilon_0$, which fixes ω in that condition. We take r so small that (5.2) holds for this ω . Now we derive a gradient comparison estimate between u and v .

Lemma 6.2. *Let $u \in W_{\text{loc}}^{1,\varphi}(\Omega)$ be a local minimizer of (1.2) and $v \in W^{1,\tilde{\varphi}}(B_r)$ be a minimizer of (5.13), where $B_{2r} \Subset \Omega$ with $r > 0$ satisfying (5.2) and $\tilde{\varphi}$ is defined in (5.8). Then there exist $\gamma = \gamma(n, p, q, L) \in (0, 1)$ and $c = c(n, p, q, L) \geq 1$ such that*

$$\int_{B_r} \tilde{\varphi}''(|Du| + |Dv|) |Du - Dv|^2 dx \leq c(\omega(2r)^{p/q} + r^\gamma) \left(\tilde{\varphi}\left(\int_{B_{2r}} |Du| dx\right) + 1 \right).$$

Proof. By Proposition 5.10(3) and Lemma 5.15, we see that $u \in W^{1,\tilde{\varphi}}(B_r)$ and $v \in W^{1,\varphi}(B_r)$. By Proposition 3.8(2),

$$\tilde{\varphi}''(|Du| + |Dv|) |Du - Dv|^2 \lesssim \tilde{\varphi}(|Du|) - \tilde{\varphi}(|Dv|) - \frac{\tilde{\varphi}'(|Dv|)}{|Dv|} Dv \cdot (Du - Dv).$$

Since $u - v \in W_0^{1,\tilde{\varphi}}(B_r)$ and v is a weak solution to (5.14),

$$\begin{aligned} & \int_{B_r} \tilde{\varphi}''(|Du| + |Dv|) |Du - Dv|^2 dx \\ & \lesssim \int_{B_r} [\tilde{\varphi}(|Du|) - \tilde{\varphi}(|Dv|)] dx - \underbrace{\int_{B_r} \tilde{\varphi}'(|Dv|) \frac{Dv}{|Dv|} \cdot (Du - Dv) dx}_{=0 \text{ by (5.14)}} \\ & = \int_{B_r} \{ [\tilde{\varphi}(|Du|) - \varphi(x, |Du|)] + [\varphi(x, |Du|) - \varphi(x, |Dv|)] + [\varphi(x, |Dv|) - \tilde{\varphi}(|Dv|)] \} dx \\ & \leq \underbrace{\int_{B_r} [\tilde{\varphi}(|Du|) - \varphi(x, |Du|)] dx}_{=: I_1} + \underbrace{\int_{B_r} [\varphi(x, |Dv|) - \tilde{\varphi}(|Dv|)] dx}_{=: I_2}; \end{aligned}$$

in the last step we have used $\int_{B_r} [\varphi(x, |Du|) - \varphi(x, |Dv|)] dx \leq 0$ since u is a minimizer of (1.2). We shall estimate I_2 . We split B_r into three regions E_1, E_2 and E_3 defined by

$$\begin{aligned} E_1 &:= B_r \cap \{\varphi^-(|Dv|) \leq \omega(2r)\}, \\ E_2 &:= B_r \cap \{\omega(2r) < \varphi^-(|Dv|) \leq |B_{2r}|^{-1+\varepsilon_0}\}, \\ E_3 &:= B_r \cap \{|B_{2r}|^{-1+\varepsilon_0} < \varphi^-(|Dv|)\}. \end{aligned}$$

In the set E_1 , (Dec) $_q$ and (A0) for φ imply that $|Dv| \lesssim \omega(2r)^{1/q}$. Therefore by (Inc) $_p$ and (A0) for φ and $\tilde{\varphi}$,

$$\int_{B_r} |\varphi(x, |Dv|) - \tilde{\varphi}(|Dv|)| \chi_{E_1} dx \lesssim \omega(2r)^{p/q} \int_{B_r} \chi_{E_1} dx \leq \omega(2r)^{p/q}.$$

In the set E_3 , Proposition 5.10(3) and the fact that $1 < |B_{2r}|^{1-\varepsilon_0} \varphi^-(|Dv|)$ imply that

$$\begin{aligned} |\varphi(x, |Dv|) - \tilde{\varphi}(|Dv|)| &\lesssim \varphi(x, |Dv|) + 1 \lesssim \varphi(x, |Dv|) \\ &\leq [|B_{2r}|^{1-\varepsilon_0} \varphi^-(|Dv|)]^{\sigma_0/2} \varphi(x, |Dv|) \\ &\lesssim r^{n(1-\varepsilon_0)\sigma_0/2} \varphi(x, |Dv|)^{1+\sigma_0/2}. \end{aligned}$$

Integrating this inequality over E_3 and using (6.1), we find that

$$\begin{aligned} \int_{B_r} |\varphi(x, |Dv|) - \tilde{\varphi}(|Dv|)| \chi_{E_3} dx \\ \lesssim r^{\frac{n(4+\sigma_0)\sigma_0}{4(2+\sigma_0)}} \left(\int_{B_r} \varphi(x, |Dv|)^{1+\sigma_0/2} dx \right)^{\sigma_0/2+\sigma_0+\frac{2}{2+\sigma_0}}. \end{aligned}$$

On the one hand, by (5.17) and (5.16), we have

$$\left(\int_{B_r} \varphi(x, |Dv|)^{1+\sigma_0/2} dx \right)^{\frac{2}{2+\sigma_0}} \lesssim \tilde{\varphi} \left(\int_{B_{2r}} |Du| dx \right) + 1.$$

On the other hand, by (5.3),

$$\left(\int_{B_r} \varphi(x, |Dv|)^{1+\sigma_0/2} dx \right)^{\frac{\sigma_0}{2+\sigma_0}} \leq |B_r|^{-\frac{\sigma_0}{2+\sigma_0}} \lesssim r^{-\frac{4n\sigma_0}{4(2+\sigma_0)}}.$$

Therefore, combining the previous three inequalities, we have

$$\int_{B_r} |\varphi(x, |Dv|) - \tilde{\varphi}(|Dv|)| \chi_{E_3} dx \lesssim r^{\frac{n\sigma_0^2}{4(2+\sigma_0)}} \left(\tilde{\varphi} \left(\int_{B_{2r}} |Du| dx \right) + 1 \right).$$

Recall that t_1 and t_2 are defined in (5.5). In the set E_2 , we observe that

$$\omega(2r) < \varphi^-(|Dv|) \leq |B_{2r}|^{-1+\varepsilon_0} < |B_{2r}|^{-1} \quad \text{and so} \quad t_1 < |Dv| < t_2.$$

Hence it follows from (wVA1) and Proposition 5.10(1) that

$$\begin{aligned} |\varphi(x, |Dv|) - \tilde{\varphi}(|Dv|)| &\leq |\varphi(x, |Dv|) - \varphi(x_0, |Dv|)| + |\varphi(x_0, |Dv|) - \tilde{\varphi}(|Dv|)| \\ &\lesssim (r + \omega(r))\varphi^-(|Dv|) + \omega(2r). \end{aligned}$$

Therefore, applying (5.17) and (5.16), we have

$$\int_{B_r} |\varphi(x, |Dv|) - \tilde{\varphi}(|Dv|)| \chi_{E_2} dx \lesssim (r + \omega(2r)) \left(\tilde{\varphi} \left(\int_{B_{2r}} |Du| dx \right) + 1 \right).$$

We have shown that

$$\begin{aligned} |I_2| &\leq \int_{B_r} |\varphi(x, |Dv|) - \tilde{\varphi}(|Dv|)| dx \\ &\lesssim (\omega(2r)^{p/q} + r^{\frac{n\sigma_0^2}{4(2+\sigma_0)}} + r) \left(\tilde{\varphi} \left(\int_{B_{2r}} |Du| dx \right) + 1 \right). \end{aligned}$$

The estimate for I_1 is analogous, with sets E_i defined with Du instead of Dv . \blacksquare

The following corollary is the key to the regularity results in the next section. Indeed, once we have the estimate from the corollary, the main results follow using standard methods.

Corollary 6.3. *Under the assumptions of Lemma 6.2, we have*

$$\int_{B_r} |Du - Dv| dx \leq c(\omega(r)^{\frac{p}{2q^2}} + r^{\gamma_1}) \left(\int_{B_{2r}} |Du| dx + 1 \right)$$

for some $\gamma_1 = \gamma_1(n, p, q, L) \in (0, 1)$ and $c = c(n, p, q, L) \geq 1$.

Proof. Set $\tilde{\omega}(r) := \omega(r)^{p/q} + r^\gamma$ with γ from Lemma 6.2 and note that $\tilde{\omega}(\cdot) \leq 2$. Applying Proposition 3.8(3) with $\kappa = \tilde{\omega}(r)^{1/2}$, Proposition 5.10(3) and Lemmas 6.2 and 5.15, we find that

$$\begin{aligned} &\int_{B_r} \tilde{\varphi}(|Du - Dv|) dx \\ &\lesssim \tilde{\omega}(r)^{1/2} \int_{B_r} [\tilde{\varphi}(|Du|) + \tilde{\varphi}(|Dv|)] dx + \tilde{\omega}(r)^{-1/2} \int_{B_r} \tilde{\varphi}''(|Du| + |Dv|) |Du - Dv|^2 dx \\ &\lesssim \tilde{\omega}(r)^{1/2} \int_{B_r} [\varphi(x, |Du|) + \varphi(x, |Dv|) + 1] dx + \tilde{\omega}(r)^{1/2} \left(\tilde{\varphi} \left(\int_{B_{2r}} |Du| dx \right) + 1 \right) \\ &\lesssim \tilde{\omega}(r)^{1/2} \left(\tilde{\varphi} \left(\int_{B_{2r}} |Du| dx \right) + 1 \right). \end{aligned}$$

Therefore, by Jensen's inequality and (Dec) $_q$ for $\tilde{\varphi}$, we have

$$\tilde{\varphi} \left(\int_{B_r} |Du - Dv| dx \right) \leq \int_{B_r} \tilde{\varphi}(|Du - Dv|) dx \lesssim \tilde{\varphi} \left(\tilde{\omega}(r)^{\frac{1}{2q}} \left(\int_{B_{2r}} |Du| dx + 1 \right) \right).$$

The claim follows, since $\tilde{\varphi}$ is strictly increasing. \blacksquare

7. Proofs of main results

In this section, we prove the main theorems. Before starting the proof we introduce a basic iteration lemma. We refer to [46, Lemma 2.1 in Chapter III].

Lemma 7.1. *Let $f : [0, r_0] \rightarrow [0, \infty)$ be a non-decreasing function. Suppose that for all $0 < \rho \leq r \leq r_0$,*

$$f(\rho) \leq C((\rho/r)^n + \varepsilon)f(r) + Cr^n$$

with a positive constant C . Then for any $\tau \in (0, n)$, there exist $\varepsilon_1, c > 0$ depending only on n, C and τ such that if $\varepsilon < \varepsilon_1$, then

$$f(\rho) \leq c(\rho/r)^{n-\tau}(f(r) + r^{n-\tau}).$$

In the next results, we denote by ω the function from (wVA1) for $\varepsilon = \varepsilon_0$ (see the beginning of Section 6). Likewise, we denote by $L \geq 1$ the constant from (A0). Now let us state and prove our main results. For iteration techniques used in the proof, we refer to [2, 10].

Theorem 7.2. *Let $\varphi \in \Phi_c(\Omega) \cap C^1([0, \infty))$ with φ' satisfying (A0), (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for some $1 < p \leq q$ and let $u \in W_{\text{loc}}^{1,\varphi}(\Omega)$ be a local minimizer of (1.2). If φ satisfies (wVA1), then $u \in C_{\text{loc}}^\alpha(\Omega)$ for any $\alpha \in (0, 1)$.*

Proof. Let $r_0 \in (0, 1)$ be a sufficiently small positive number which will be determined later. We fix $\Omega' \Subset \Omega$ and assume that (5.2) holds $r = r_0 > 0$. For any $B_{2r} \subset \Omega'$ with $0 < 2r \leq r_0$, let $v \in W^{1,\tilde{\varphi}}(B_r)$ be the minimizer of (5.13) with $\tilde{\varphi}$ determined in (5.8). When $\rho \in (0, r/2)$, applying Corollary 6.3 with $\omega_0(r) := \omega(r)^{p/(2q^2)} + r^{\gamma_1}$, (4.13) and (5.18), we have

$$\begin{aligned} \int_{B_\rho} |Du| dx &\leq \int_{B_r} |Du - Dv| dx + \int_{B_\rho} |Dv| dx \\ &\lesssim \omega_0(2r) \int_{B_{2r}} [|Du| + 1] dx + \rho^n \sup_{B_{r/2}} |Dv| \\ &\leq \omega_0(r_0) \int_{B_{2r}} [|Du| + 1] dx + \rho^n \int_{B_r} |Dv| dx \\ &\lesssim \left(\left(\frac{\rho}{r} \right)^n + \omega_0(r_0) \right) \int_{B_{2r}} |Du| dx + r^n. \end{aligned}$$

Moreover, if $r/2 \leq \rho \leq 2r$, the above estimate is obvious since then $1/2 \leq \rho/r$.

Let $f(\rho) := \int_{B_\rho} |Du| dx$, fix $\tau \in (0, n)$ and choose r_0 so small that

$$\omega_0(r_0) = \omega(r_0)^{\frac{1}{2q}} + r_0^{\gamma_1} \leq \varepsilon_1,$$

where ε_1 is from Lemma 7.1. Then, applying the lemma and using (5.4) with $2r = r_0$, we have the following Morrey type estimate:

$$\int_{B_\rho} |Du| dx \lesssim \left(\frac{\rho}{r_0} \right)^{n-\tau} \left(\int_{B_{r_0}} |Du| dx + r_0^{n-\tau} \right) \quad \text{for all } B_\rho \subset \Omega' \text{ with } \rho \in (0, r_0] \quad (7.3)$$

with implicit constant $c = c(n, p, q, L, \tau) \geq 1$. We note that in (7.3), $\rho \in (0, r_0]$ and $B_\rho \subset \Omega'$ are arbitrary and the implicit constant is universal. Therefore, by taking $\tau = 1 - \alpha$ for each $\alpha \in (0, 1)$ in (7.3), we have $u \in C_{\text{loc}}^\alpha(\Omega')$ by a Morrey type embedding (see for instance [46, Chapter 3, Theorem 1.1]). More precisely, we obtain

$$[u]_{C^\alpha(B_{r_0/2})} \lesssim r_0^{1-\alpha} \int_{B_{r_0}} |Du| dx + 1. \quad \blacksquare$$

Next we prove the second main theorem, $C^{1,\alpha}$ -regularity.

Theorem 7.4. *Let $\varphi \in \Phi_c(\Omega) \cap C^1([0, \infty))$ with φ' satisfying (A0), (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for some $1 < p \leq q$ and let $u \in W_{\text{loc}}^{1,\varphi}(\Omega)$ be a minimizer of (1.2). If φ satisfies (wVA1) with*

$$\omega(r) \lesssim r^\beta \quad \text{for all } r \in (0, 1] \text{ and for some } \beta \in (0, 1),$$

then $u \in C_{\text{loc}}^{1,\alpha}(\Omega)$ for some $\alpha \in (0, 1)$. Here α depends only on n, p, q, L and β .

Proof. Fix $\Omega' \Subset \Omega$. We first notice from (7.3) that for any $\tau \in (0, n)$,

$$\int_{B_{2r}} |Du| dx \leq c_\tau r^{-\tau} \quad \text{for all } B_{2r} \subset \Omega' \text{ with } 2r \in (0, r_0],$$

where $r_0 > 0$ is from the proof of Theorem 7.2 and $c_\tau \geq 1$ depends on n, p, q, L, r_0 and τ . Consider sufficiently small $2r < r_0$, which will be determined later. Let $v \in W^{1,\tilde{\varphi}}(B_r)$ be a minimizer of (5.13) with $\tilde{\varphi}$ determined in (5.8). Then for $0 < \rho < r/2$, applying (4.14) with $B_\rho(x_0) = B_{r/2}$ and $\tau = 2\rho/r$, Corollary 6.3 with $\omega(r) \lesssim r^\beta$ and (5.18), we have

$$\begin{aligned} \int_{B_\rho} |Du - (Du)_\rho| dx &\leq 2 \int_{B_\rho} |Du - (Dv)_\rho| dx \\ &\leq 2 \int_{B_\rho} |Du - Dv| dx + 2 \int_{B_\rho} |Dv - (Dv)_\rho| dx \\ &\lesssim \left(\frac{r}{\rho}\right)^n \int_{B_r} |Du - Dv| dx + \left(\frac{\rho}{r}\right)^{\alpha_0} \int_{B_{r/2}} |Dv| dx \\ &\lesssim \left(r^{\gamma_0} \left(\frac{r}{\rho}\right)^n + \left(\frac{\rho}{r}\right)^{\alpha_0}\right) \left(\int_{B_{2r}} |Du| dx + 1\right) \\ &\leq c_\tau r^{-\tau} \left(r^{\gamma_0} \left(\frac{r}{\rho}\right)^n + \left(\frac{\rho}{r}\right)^{\alpha_0}\right), \end{aligned}$$

where $\gamma_0 := \min\{\frac{p\beta}{2q^2}, \gamma_1\}$. Finally, we choose $\rho := r^{1+\gamma_0/(2n)}$ and $\tau := \frac{\alpha_0\gamma_0}{4n}$. Suppose that $2r \leq \min\{r_0, 4^{-2n/\gamma_0}\}$. Then $\rho < r/2$ and for the concentric balls $B_\rho \subset B_{2r} \subset \Omega'$ we have

$$\int_{B_\rho} |Du - (Du)_\rho| dx \lesssim c_\tau r^{\gamma_0/2-\tau} + r^\tau \leq 2c_\tau r^\tau = 2c_\tau \rho^{\frac{\tau}{1+\gamma_0/(2n)}}.$$

This implies that $Du \in C_{\text{loc}}^\alpha(\Omega')$ with $\alpha = \frac{\alpha_0 \gamma_0}{4n+2\gamma_0}$ by a Campanato type embedding (see for instance [46, Chapter 3]). Since $\Omega' \Subset \Omega$ is arbitrary, we have the conclusion. The last inequality also yields an estimate for the seminorm $[Du]_{C^\alpha(B_{r_0/2})}$; however, once we unravel the dependence on c_τ , it is a somewhat complicated formula. ■

Remark 7.5. By following the proofs, one can see that we use the condition (wVA1) only for fixed $\varepsilon = \varepsilon_0$ determined in (6.1). Therefore, in Theorems 7.2 and 7.4, the condition (wVA1) can be replaced with the combination of (A1) and (wVA1) with fixed $\varepsilon > 0$, where ε is sufficiently small and depends on n, p, q and L .

8. Examples of special structures

In this section, we show that our results include previous regularity results for special structures presented in the introduction. We provide details only for some cases, as the remaining ones can be handled by similar techniques. We denote by C^ω continuous functions with modulus of continuity ω .

Corollary 8.1 (Perturbed autonomous case). *Let $a : \Omega \rightarrow [v, \Lambda]$ for some $0 < v \leq \Lambda$, and let $\psi \in \Phi_c \cap C^1([0, \infty))$ with ψ' satisfying (Inc) $_{p-1}$ and (aDec) $_{q-1}$ for some $1 < p \leq q$. Define $\varphi(x, t) := a(x)\psi(t)$. Then φ satisfies (VA1), with $\omega(r) \approx \omega_a(2r)$, if and only if $a \in C^{\omega_a}$.*

Proof. For any $B_r \subset \Omega$,

$$\varphi_{B_r}^+(t) = a_{B_r}^+ \psi(t) = \left(1 + \frac{a_{B_r}^+ - a_{B_r}^-}{a_{B_r}^-}\right) \varphi_{B_r}^-(t).$$

Since $a_{B_r}^- \in [v, \Lambda]$, we obtain

$$\frac{\varphi_{B_r}^+(t) - \varphi_{B_r}^-(t)}{\varphi_{B_r}^-(t)} \approx a_{B_r}^+ - a_{B_r}^-,$$

and so the claim follows. ■

Corollary 8.2 (Variable exponent case). *Let $p : \Omega \rightarrow [p_1, p_2]$ for some $1 < p_1 \leq p_2$. Define $\varphi(x, t) := t^{p(x)}$. Then φ satisfies (VA1) if and only if there exists ω_p with*

$$\lim_{r \rightarrow 0} \omega_p(r) \ln \frac{1}{r} = 0 \quad \text{and} \quad p \in C^{\omega_p}.$$

Moreover, φ satisfies (VA1) with $\omega(r) \lesssim r^\beta$ for some $\beta > 0$ if and only if

$$\omega_p(r) \lesssim r^{\tilde{\beta}} \quad \text{for some } \tilde{\beta} > 0.$$

Proof. Fix $B_r \subset \Omega$ with $|B_r| \leq 1$ and set $p^\pm = p_{B_r}^\pm$. Then we have $\varphi_{B_r}^-(t) = t^{p^-}$ and $\varphi_{B_r}^+(t) = t^{p^+}$ for $t \geq 1$ as well as $\varphi_{B_r}^-(t) = t^{p^-}$ and $\varphi_{B_r}^+(t) = t^{p^+}$ for $t < 1$.

Let us derive an equivalent form of the inequality in condition (VA1). We may consider the range $[|B_r|, |B_r|^{-1}]$ in the condition, since it turns out that this choice of lower bound entails no additional restrictions in the variable exponent case. When $t \geq 1$, we have

$$\varphi_{B_r}^+(t) - \varphi_{B_r}^-(t) = (t^{p^+ - p^-} - 1)t^{p^-} = (t^{p^+ - p^-} - 1)\varphi_{B_r}^-(t).$$

When $t \leq 1$, the exponents p^+ and p^- are interchanged. Since we consider the range

$$t \in [(\varphi^-)_{B_r}^{-1}(|B_r|), (\varphi^-)_{B_r}^{-1}(|B_r|^{-1})] = [|B_r|^{1/p^+}, |B_r|^{-1/p^-}],$$

we obtain

$$\sup_{t \in [|B_r|^{1/p^+}, |B_r|^{-1/p^-}]} \frac{\varphi_{B_r}^+(t) - \varphi_{B_r}^-(t)}{\varphi_{B_r}^-(t)} = |B_r|^{\frac{p^- - p^+}{p^-}} - 1.$$

Suppose that $p \in C^{\omega_p}$. By the mean value theorem, $e^x - 1 \leq e^x x$. Thus

$$\begin{aligned} |B_r|^{\frac{p^- - p^+}{p^-}} - 1 &\leq |B_r|^{-\omega_p(2r)} - 1 \leq e^{n\omega_p(2r)\ln(1/r)} - 1 \\ &\leq n e^{n\omega_p(2r)\ln(1/r)} \omega_p(2r) \ln \frac{1}{r} =: \omega(r) \end{aligned}$$

and the inequality from (VA1) holds with this ω . Moreover, if $\lim_{r \rightarrow 0} \omega_p(r) \ln \frac{1}{r} = 0$, then ω tends to zero, hence we obtain (VA1). If $\omega_p(r) \lesssim t^\beta$, then ω is of order $\beta - \varepsilon$ for any ε .

Suppose next that φ satisfies (VA1) with a function ω . Then, for $r \leq 1/2$,

$$|B_r|^{\frac{p^- - p^+}{p^-}} - 1 \leq \omega(r)$$

hence

$$p^+ - p^- \leq -\frac{p^- \log(1 + \omega(r))}{\log |B_r|} \lesssim \frac{p_2 \log(1 + \omega(r))}{n \log \frac{1}{r}} =: \omega_p(2r).$$

Then $\omega_p(2r) \log \frac{1}{r} \approx \log(1 + \omega(r)) \rightarrow 0$. If $\omega(r) \leq r^{\tilde{\beta}}$, then $\omega_p(2r) \approx \log(1 + r^{\tilde{\beta}})/\log \frac{1}{r} \lesssim r^{\tilde{\beta}}$. \blacksquare

Rădulescu and colleagues [22, 77, 84] have considered a functional with model case $\varphi(x, t) = t^{p(x)} + t^{q(x)}$, which they call “double phase” (it is different from the double phase functional of Zhikov, considered below). To the best of our knowledge, this is the first regularity result for this functional.

Corollary 8.3 (Rădulescu’s double phase). *Let $p, q : \Omega \rightarrow [p_1, p_2]$ for some $1 < p_1 \leq p_2$ and $\varphi(x, t) = t^{p(x)} + t^{q(x)}$. Then φ satisfies (VA1) if there exist ω_m and ω_M with*

$$\lim_{r \rightarrow 0} \omega_m(r) = 0, \quad \min\{p, q\} \in C^{\omega_m}, \quad \lim_{r \rightarrow 0} \omega_M(r) \ln \frac{1}{r} = 0 \quad \text{and} \quad \max\{p, q\} \in C^{\omega_M}.$$

In addition, φ satisfies (VA1) with $\omega(r) \leq r^\beta$ for some $\beta > 0$ if

$$\lim_{r \rightarrow 0} \omega_m(r) r^{-\tilde{\beta}} = 0 \quad \text{and} \quad \lim_{r \rightarrow 0} \omega_M(r) r^{-\tilde{\beta}} = 0 \quad \text{for some } \tilde{\beta} > 0.$$

This result can be proved with the same methods as Corollary 8.2; the details are left to the interested reader. Note that the regularity required of the minimum is lower than the regularity required of the maximum. This is due to the fact that we only require the inequality of (VA1) in the range $[\omega(r), 1]$ where the minimum determines φ , whereas the maximum is used in the range $[1, |B_r|]$.

We now consider double phase problems in the sense of Zhikov and Mingione.

Corollary 8.4 (Double phase case). *Let $a \in C^{\omega_a}(\Omega)$ and $b \in C^{\omega_b}(\Omega)$ be non-negative with $0 < \nu \leq a(\cdot) + b(\cdot) \leq \Lambda$ for some $0 < \nu \leq \Lambda$, and $\psi, \xi \in \Phi_c \cap C^1([0, \infty))$ with ψ', ξ' satisfying (A0), and (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for some $1 < p \leq q$. Suppose that ξ/ψ is almost increasing. Define*

$$\varphi(x, t) := a(x)\psi(t) + b(x)\xi(t)$$

and, for $\varepsilon \in [0, 1)$,

$$\omega_\varepsilon(r) := \omega_a(r) + \omega_b(r)r^{n(1-\varepsilon)}\xi(\psi^{-1}(r^{-n(1-\varepsilon)})).$$

If ω_ε is bounded with $\lim_{r \rightarrow 0} \omega_\varepsilon(r) = 0$ when $\varepsilon > 0$, then φ satisfies (wVA1) with $\omega \approx \omega_\varepsilon$.

Proof. Fix $B_r \subset \Omega$ so small that $\omega_a(2r), \omega_b(2r) \leq \nu/2$. Set $a^\pm := a_{B_r}^\pm$, $b^\pm := b_{B_r}^\pm$ and $\varphi^\pm(t) := \varphi_{B_r}^\pm(t)$. For $0 \leq \varepsilon < 1$, suppose $t \in (0, t_2)$ with $t_2 := (\varphi^-)^{-1}(|B_r|^{-1+\varepsilon})$.

We first consider $t > 1$. Assume first that $b^- \geq \nu/4$. Then $\varphi^-(t) \gtrsim \xi(t) \gtrsim \psi(t)$ and so

$$\begin{aligned} \varphi^+(t) - \varphi^-(t) &\leq (a^+ - a^-)\psi(t) + (b^+ - b^-)\xi(t) \\ &\lesssim (\omega_a(r) + \omega_b(r))\varphi^-(t). \end{aligned}$$

We note that the case $b^- < \nu/4$ and $a^- < \nu/4$ cannot occur, since $a + b \geq \nu$ and $\omega_a(2r), \omega_b(2r) \leq \nu/2$.

Next, we consider $t > 1$ and $a^- \geq \nu/4$. Then $\varphi^-(t) \gtrsim \psi(t)$. Note that $\varphi(x, t) \approx \max\{a(x)\psi(t), b(x)\xi(t)\}$ and that by the continuity of the functions a and b , there exists $x_t \in B_r$ such that $\varphi^-(t) = \varphi(x_t, t)$. Using these and that ξ/ψ is almost increasing, we have

$$\frac{\xi(t)}{\varphi^-(t)} \approx \min \left\{ \frac{\xi(t)}{a(x_t)\psi(t)}, \frac{1}{b(x_t)} \right\} \lesssim \min \left\{ \frac{\xi(t_2)}{a(x_t)\psi(t_2)}, \frac{\xi(t_2)}{b(x_t)\xi(t_2)} \right\} \lesssim \frac{\xi(t_2)}{\varphi^-(t_2)}.$$

Since $\frac{\psi(t)}{\varphi^-(t)} \leq \frac{4}{\nu}$ and $\psi(t_2) \lesssim a(x_{t_2})\psi(t_2) \lesssim \varphi^-(t_2) \approx r^{-n(1-\varepsilon)}$, we conclude that

$$\begin{aligned} \frac{\varphi^+(t) - \varphi^-(t)}{\varphi^-(t)} &\leq (a^+ - a^-) \frac{\psi(t)}{\varphi^-(t)} + (b^+ - b^-) \frac{\xi(t)}{\varphi^-(t)} \\ &\lesssim \omega_a(2r) + \omega_b(2r) \frac{\xi(t_2)}{\varphi^-(t_2)} \\ &\lesssim \omega_a(r) + \omega_b(r)r^{n(1-\varepsilon)}\xi(\psi^{-1}(r^{-n(1-\varepsilon)})). \end{aligned}$$

We note that the factor multiplying $\omega_b(r)$ in the last expression is greater than $c > 0$ depending on the parameters, so it can absorb the $+\omega_b(r)$ from the other cases to give ω_ε in the statement of the result.

The necessary inequality has been established for all cases when $t > 1$. We next consider $t \leq 1$. By (A0) for ψ and ξ ,

$$\varphi^+(t) - \varphi^-(t) \leq (a^+ - a^-)\psi(t) + (b^+ - b^-)\xi(t) \lesssim \omega_a(r) + \omega_b(r).$$

We use this as the additive term “ $+\omega(r)$ ” in the definition of (wVA1) to cover small t . This concludes the proof of (wVA1). \blacksquare

Remark 8.5. In the previous proof, we used the additive error “ $+\omega(r)$ ” for (wVA1) to handle the case $t \leq 1$. If $a \equiv 1$, then this is not needed, and we also have the following conclusion: if ω_0 is bounded with $\lim_{r \rightarrow 0} \omega_0(r) = 0$, then φ satisfies (VA1).

Suppose that $\xi(t) = \psi(t) \ln(e + t)$ and $a \equiv 1$ in Corollary 8.4. Then we have

$$\xi(\psi^{-1}(r^{-n(1-\varepsilon)})) = r^{-n(1-\varepsilon)} \ln(e + \psi^{-1}(r^{-n(1-\varepsilon)})) \approx r^{-n(1-\varepsilon)} \ln(e + 1/r)$$

since ψ satisfies (Inc) $_p$ and (Dec) $_q$. We see that the degenerate double phase functional satisfies (VA1) if b is vanishing log-Hölder continuous.

From Corollary 8.4, we obtain sharp regularity conditions for φ satisfying particular structures of double phase with power functions.

Corollary 8.6. *Let $1 < p \leq q$, $\beta \in (0, 1]$, and $a \in C^{\omega_a}$ and $b \in C^{0,\beta}$ be non-negative. Define $\gamma_\varepsilon := \beta - n(q - p)(1 - \varepsilon)/p$, $\varepsilon \geq 0$.*

- (1) *Let $\varphi(x, t) = t^p + b(x)t^q$.
If $q/p < 1 + \beta/n$, then φ satisfies (VA1) with $\omega(r) \approx r^{\gamma_0}$.
If $q/p \leq 1 + \beta/n$, then φ satisfies (wVA1) with $\omega(r) \approx r^{\gamma_\varepsilon}$.*
- (2) *Let $\varphi(x, t) = a(x)t^p + t^q$. Then φ satisfies (wVA1) with $\omega(r) \approx \omega_a(r)$.*
- (3) *Let $a(x)t^p + b(x)t^q$ with $v \leq a + b \leq \Lambda$. If $q/p \leq 1 + \beta/n$, then φ satisfies (wVA1) with $\omega(r) \approx r^{\gamma_\varepsilon} + \omega_a(r)$.*

Appendix A. $C^{1,\alpha}$ -Regularity for autonomous problems

In this section, we prove Lemma 4.12. We follow the ideas in [35, 58]. In fact, it is almost enough to replace the map $t \mapsto t^p$ by the map $t \mapsto \varphi(t)$ in the proof in [58]. However, for completeness, we present the proof. Suppose $\varphi \in C^1([0, \infty)) \cap C^2((0, \infty))$ and φ' satisfies (Inc) $_{p-1}$ and (Dec) $_{q-1}$ for some $1 < p \leq q$. We first consider the following non-degenerate problem for $\varepsilon > 0$:

$$\operatorname{div} \left(\frac{\varphi_\varepsilon(|Du_\varepsilon|)}{|Du_\varepsilon|} Du_\varepsilon \right) = 0 \quad \text{in } \Omega, \quad \text{where} \quad \varphi_\varepsilon(t) := \int_0^t \frac{\varphi'(\varepsilon + s)s}{\varepsilon + s} ds, \quad (\text{A.1})$$

which is the Euler–Lagrange equation of the minimization problem

$$\min_w \int_{\Omega} \varphi_{\varepsilon}(|Dw|) dx. \quad (\text{A.2})$$

(In Lemma 4.12, $\Omega = B_r$.) By the definition of φ_{ε} we have

$$\frac{\varphi'_{\varepsilon}(t)}{t} = \frac{\varphi'(\varepsilon + t)}{\varepsilon + t}, \quad \text{so that} \quad \lim_{t \rightarrow 0^+} \frac{\varphi'_{\varepsilon}(t)}{t} = \frac{\varphi'(\varepsilon)}{\varepsilon} > 0.$$

Hence (A.1) is non-degenerate. We emphasize that all hidden constants in \approx and \lesssim in this appendix depend only on n , p and q , but are independent of ε . We observe by the first equality above and (Inc) $_{p-1}$ and (Dec) $_{q-1}$ of φ' that

$$\begin{aligned} \varphi''_{\varepsilon}(t) &= \frac{\varphi'(\varepsilon + t)}{\varepsilon + t} \left(1 + \left(\frac{\varphi''(\varepsilon + t)}{(\varepsilon + t)\varphi'(\varepsilon + t)} - 1 \right) \frac{t}{\varepsilon + t} \right) \\ &\geq \frac{\varphi'_{\varepsilon}(t)}{t} \left(1 + (p-2) \frac{t}{\varepsilon + t} \right) \geq \min\{1, p-1\} \frac{\varphi'_{\varepsilon}(t)}{t} \end{aligned} \quad (\text{A.3})$$

and

$$\varphi''_{\varepsilon}(t) \leq \frac{\varphi'_{\varepsilon}(t)}{t} \left(1 + (q-2) \frac{t}{\varepsilon + t} \right) \leq \max\{1, q-1\} \frac{\varphi'_{\varepsilon}(t)}{t}. \quad (\text{A.4})$$

Therefore, φ'_{ε} satisfies (Inc) $_{\min\{1, p-1\}}$ and (Dec) $_{\max\{1, q-1\}}$, which implies that

$$t\varphi''_{\varepsilon}(t) \approx \varphi'_{\varepsilon}(t), \quad t\varphi'_{\varepsilon}(t) \approx \varphi_{\varepsilon}(t), \quad \frac{\varphi_{\varepsilon}(t)}{t^2} \approx \frac{\varphi'_{\varepsilon}(t)}{t} = \frac{\varphi'(\varepsilon + t)}{\varepsilon + t}. \quad (\text{A.5})$$

In view of [35], in particular Lemmas 5.7 and 5.8, we find that $u_{\varepsilon} \in W_{\text{loc}}^{2,2}(\Omega)$ and $\varphi_{\varepsilon}(|Du_{\varepsilon}|) \in W_{\text{loc}}^{1,2}(\Omega)$ if $\varepsilon > 0$ and that for any $B_{2\rho} \Subset \Omega$ and $\varepsilon \geq 0$,

$$\sup_{B_{\rho}} \varphi_{\varepsilon}(|Du_{\varepsilon}|) \lesssim \int_{B_{2\rho}} \varphi_{\varepsilon}(|Du_{\varepsilon}|) dx. \quad (\text{A.6})$$

Here $u_0 = u$ and $\varphi_0 = \varphi$.

Fix $\varepsilon > 0$ and $B_{2\rho} \Subset \Omega$. From now on, for convenience, we shall simply write

$$u = u_{\varepsilon} \quad \text{and} \quad v = \varphi_{\varepsilon}(|Du|) = \varphi_{\varepsilon}(|Du_{\varepsilon}|). \quad (\text{A.7})$$

We first notice from (A.1) and $u \in W_{\text{loc}}^{2,2}(\Omega)$ that

$$\operatorname{div} \left(\frac{\varphi'_{\varepsilon}(|Du|)}{|Du|} Du \right) = \operatorname{div} \left(\frac{\varphi'(\varepsilon + |Du|)}{\varepsilon + |Du|} Du \right) = \sum_{i,j=1}^n a_{ij} u_{x_i} x_j = 0 \quad \text{a.e. in } \Omega,$$

where $a_{ij} = a_{ij}(Du)$, $b_{ij} = b_{ij}(Du)$,

$$a_{ij}(z) := \frac{\varphi'(\varepsilon + |z|)}{\varepsilon + |z|} b_{ij}(z) = \frac{\varphi'_{\varepsilon}(|z|)}{|z|} b_{ij}(z)$$

and

$$b_{ij}(z) := \left(\frac{\varphi''(\varepsilon + |z|)(\varepsilon + |z|)}{\varphi'(\varepsilon + |z|)} - 1 \right) \frac{z_i z_j}{(\varepsilon + |z|)|z|} + \delta_{ij}$$

for $z \in \mathbb{R}^n$ (δ_{ij} is the Kronecker delta). As in (A.3) and (A.4) along with the fact that $\sum_{i,j} z_i z_j \eta_i \eta_j = (z \cdot \eta)^2$, we conclude that

$$\min \{1, p-1\} |\eta|^2 \leq \sum_{i,j=1}^n b_{ij}(z) \eta_i \eta_j \leq \max \{1, q-1\} |\eta|^2 \quad \text{for all } z, \eta \in \mathbb{R}^n. \quad (\text{A.8})$$

Consider the weak form of (A.1) and a unit vector $v \in S^{n-1}$. We see that

$$\begin{aligned} 0 &= - \int_{\Omega} \frac{\varphi'_{\varepsilon}(|Du|)}{|Du|} Du \cdot D(\zeta_v) dx = \int_{\Omega} \sum_{i=1}^{\infty} \left(\frac{\varphi'_{\varepsilon}(|Du|)}{|Du|} u_{x_i} \right)_v \zeta_{x_i} dx \\ &= \int_{\Omega} \sum_{i,j=1}^{\infty} a_{ij} u_{x_j} v \zeta_{x_i} dx \end{aligned}$$

for any $\zeta \in C_0^{\infty}(\Omega)$, where the subscript v indicates directional derivatives. Thus we have shown that

$$\sum_{i,j=1}^n (a_{ij} u_{v x_j})_{x_i} = 0 \quad (\text{A.9})$$

in the weak sense. In addition, by the definition of v in (A.7), we have

$$v_{x_j} = \frac{\varphi'_{\varepsilon}(|Du|)}{|Du|} \sum_{k=1}^n u_{x_k} u_{x_k x_j},$$

so that $b_{ij} v_{x_j} = \sum_{k=1}^n a_{ij} u_{x_k x_j} u_{x_k}$. We conclude, with (A.9) for the second equality, that

$$\begin{aligned} - \int_{B_{\rho}} \sum_{i,j=1}^n b_{ij} v_{x_j} \zeta_{x_i} dx &= - \sum_{k=1}^n \int_{B_{\rho}} \sum_{i,j=1}^n a_{ij} u_{x_k x_j} (u_{x_k} \zeta)_{x_i} dx \\ &\quad + \int_{B_{\rho}} \sum_{i,j,k=1}^n a_{ij} u_{x_k x_j} u_{x_k x_i} \zeta dx \\ &= \int_{B_{\rho}} \sum_{i,j,k=1}^n a_{ij} u_{x_k x_j} u_{x_k x_i} \zeta dx \end{aligned}$$

for all $\zeta \in C_0^{\infty}(B_{\rho})$ with $B_{\rho} \Subset \Omega$. Therefore, we have

$$\mathcal{L}v := \sum_{i,j=1}^n (b_{ij} v_{x_i})_{x_j} = \sum_{i,j,k=1}^n a_{ij} u_{x_k x_j} u_{x_k x_i} =: g \quad (\text{A.10})$$

in the weak sense. Moreover, by (A.8) with $\eta = D(u_{x_k})$ and (A.5), we have

$$g \approx \frac{\varphi_\varepsilon(|Du|)}{|Du|^2} |D^2u|^2, \quad (\text{A.11})$$

where $|D^2u|^2 := \sum_{i,j} (u_{x_i x_j})^2$. In the same way as in [58, Lemma 1] with $v = \varphi_\varepsilon(|Du|)$, and $v^{-1/p}$ replaced by $[\varphi_\varepsilon^{-1}(v)]^{-1} = |Du|^{-1}$, we deduce that for any $B_{4\rho} \Subset \Omega_r$,

$$\int_{B_\rho} g^{1+\delta} dx \lesssim \left(\int_{B_{4\rho}} g dx \right)^{1+\delta}. \quad (\text{A.12})$$

Next, set

$$v_0 := v = \varphi_\varepsilon(|Du|) \quad \text{and} \quad v_k := \frac{v}{\varphi_\varepsilon^{-1}(v)} u_{x_k} = \frac{\varphi_\varepsilon(|Du|)}{|Du|} u_{x_k}, \quad k = 1, \dots, n.$$

Then

$$v_{0,x_j} = v_{x_j} = \frac{\varphi'_\varepsilon(|Du|) Du_{x_j} \cdot Du}{|Du|}, \quad (\text{A.13})$$

and, for $k = 1, \dots, n$,

$$\begin{aligned} v_{k,x_j} &= \frac{v}{\varphi_\varepsilon^{-1}(v)} u_{x_k x_j} + \left(1 - \frac{v}{\varphi_\varepsilon^{-1}(v) \varphi'_\varepsilon(\varphi_\varepsilon^{-1}(v))} \right) \frac{v_{x_j}}{\varphi_\varepsilon^{-1}(v)} u_{x_k} \\ &= \frac{\varphi_\varepsilon(|Du|)}{|Du|} u_{x_k x_j} + \left(1 - \frac{\varphi_\varepsilon(|Du|)}{|Du| \varphi'_\varepsilon(|Du|)} \right) \frac{\varphi'_\varepsilon(|Du|) Du_{x_j} \cdot Du}{|Du|^2} u_{x_k}. \end{aligned}$$

Here we note from (Inc)_{min{2,p}} and (Dec)_{max{2,q}} for φ_ε that

$$0 < 1 - \frac{1}{\min\{2, p\}} \leq \underbrace{1 - \frac{\varphi_\varepsilon(|Du|)}{|Du| \varphi'_\varepsilon(|Du|)}}_{=: A(|Du|)} \leq 1 - \frac{1}{\max\{2, q\}} < 1.$$

Then the previous expression for the partial derivatives implies that $v_k \in W_{\text{loc}}^{1,2}(\Omega)$ since $u \in W_{\text{loc}}^{2,2}(\Omega)$, $v = \varphi_\varepsilon(|Du|) \in W_{\text{loc}}^{1,2}(\Omega)$ and $Du \in L_{\text{loc}}^\infty(\Omega)$. Moreover,

$$\begin{aligned} \sum_{k=1}^n |Dv_k|^2 &= \sum_{k,j=1}^n v_{k,x_j}^2 \\ &= \frac{\varphi_\varepsilon(|Du|)^2}{|Du|^2} |D^2u|^2 + A(|Du|)^2 \frac{\varphi'_\varepsilon(|Du|)^2}{|Du|^4} \sum_{k,j=1}^n (Du_{x_j} \cdot Du)^2 u_{x_k}^2 \\ &\quad + 2A(|Du|) \frac{\varphi_\varepsilon(|Du|) \varphi'_\varepsilon(|Du|)}{|Du|^3} \sum_{j=1}^n \left[Du_{x_j} \cdot Du \sum_{k=1}^n u_{x_k x_j} u_{x_k} \right]. \end{aligned}$$

Since $A(|Du|) \approx 1$ and $\sum_{k,j} (Du_{x_j} \cdot Du)^2 u_{x_k}^2 \leq |Du|^4 |D^2u|^2$, and since we have $Du_{x_j} \cdot Du \sum_k u_{x_k x_j} u_{x_k} = (Du_{x_j} \cdot Du)^2 \geq 0$, we obtain

$$gv \approx \frac{\varphi_\varepsilon(|Du|)^2}{|Du|^2} |D^2u|^2 \leq \sum_{k=1}^n |Dv_k|^2 \lesssim \frac{\varphi_\varepsilon(|Du|)^2}{|Du|^2} |D^2u|^2 \approx gv. \quad (\text{A.14})$$

Using the expression for the partial derivative v_{k,x_j} and (A.9), we see that for $k = 1, \dots, n$,

$$\begin{aligned} \mathcal{L}v_k &= \sum_{i,j=1}^n (b_{ij} v_{k,x_j})_{x_i} \\ &= \sum_{i,j,l=1}^n \left(a_{ij} u_{x_l x_j} \left[\frac{|Du| v \delta_{lk}}{\varphi'_\varepsilon(|Du|)|Du|} + \frac{u_{x_k} u_{x_l}}{|Du|} - \frac{|Du| v u_{x_k} u_{x_l}}{\varphi'_\varepsilon(|Du|)|Du|^3} \right] \right)_{x_i} \\ &= \frac{\varphi'_\varepsilon(|Du|)}{|Du|} \sum_{i,j,l=1}^n b_{ij} u_{x_l x_j} \left(\frac{u_{x_k} u_{x_l}}{|Du|} + \frac{v \delta_{lk}}{\varphi'_\varepsilon(|Du|)} - \frac{v u_{x_k} u_{x_l}}{\varphi'_\varepsilon(|Du|)|Du|^2} \right)_{x_i} \\ &=: g_k \end{aligned} \quad (\text{A.15})$$

in the weak sense for test functions in $C_0^\infty(B_\rho)$. Note that g_k depends on first and second partial derivatives of u . We use the estimates $|u_{x_j}| \leq |Du|$ and $|u_{x_l x_j}| \leq |D^2u|$ to conclude that

$$\left| \left(\frac{u_{x_k} u_{x_l}}{|Du|} \right)_{x_i} \right| = \left| \frac{u_{x_k x_i} u_{x_l} + u_{x_k} u_{x_l x_i}}{|Du|} - \frac{u_{x_k} u_{x_l} \sum_{m=1}^n u_{x_m x_i}}{|Du|^2} \right| \lesssim |D^2u|.$$

Similarly, using also $t\varphi'_\varepsilon(t) \approx \varphi_\varepsilon(t)$ from (A.5), we estimate the other multipliers of $b_{ij} u_{x_l x_j}$ by $|D^2u|$ as well. Since $b_{ij} \approx 1$ by (A.8), we conclude by (A.11) that

$$|g_k| \lesssim \left| \frac{\varphi'_\varepsilon(|Du|)}{|Du|} \sum_{i,j,l=1}^n b_{ij} u_{x_l x_j} |D^2u| \right| \lesssim \frac{\varphi'_\varepsilon(|Du|)}{|Du|} |D^2u|^2 \approx g. \quad (\text{A.16})$$

From now on, fix $B_{32s} \Subset \Omega$. For $k = 0, 1, \dots, n$, let $h_k \in W^{1,2}(B_s)$ be a weak solution to

$$\mathcal{L}h_k = (b_{ij} h_{k,x_j})_{x_i} = 0 \quad \text{in } B_s, \quad h_k = v_k \quad \text{on } \partial B_s,$$

and let

$$w_k = h_k - v_k \in W_0^{1,2}(B_s).$$

Then, by De Giorgi's theory for linear equations (see for instance [49, Theorem 7.7]), for any concentric balls $B_\rho \subset B_s$ with $0 < \rho \leq s$ we have

$$\int_{B_\rho} |Dh_k|^2 dx \lesssim \left(\frac{\rho}{s} \right)^{\beta-2} \int_{B_s} |Dh_k|^2 dx \lesssim \left(\frac{\rho}{s} \right)^{\beta-2} \int_{B_s} |Dv_k|^2 dx \quad (\text{A.17})$$

for some $\beta = \beta(n, p, q) \in (0, 2)$. In addition, by (A.8) and $\mathcal{L}h_k = 0$, and by (A.10), (A.15) and (A.16),

$$\begin{aligned} \int_{B_s} |Dw_k|^2 dx &\lesssim \int_{B_s} \sum_{i,j=1}^n b_{ij} (h_k - v_k)_{x_j} w_{k,x_i} dx \\ &= - \int_{B_s} \sum_{i,j=1}^n b_{ij} v_{k,x_j} w_{k,x_i} dx = \int_{B_s} g_k w_k dx \lesssim \int_{B_s} g |w_k| dx. \end{aligned}$$

Here we interpret $g_0 := g$. (Note that $w_k \notin C_0^\infty(B_\rho)$, but we can use w_k as a test function by an approximation argument.)

Hence applying Hölder's inequality and (A.12) we find that

$$\int_{B_s} |Dw_k|^2 dx \lesssim \left(\int_{B_{4s}} g dx \right) \left(\int_{B_s} |w_k|^{1+1/\delta} dx \right)^{\frac{1}{1+1/\delta}}.$$

Furthermore, the same arguments used to prove [58, (3.8) and (3.13)] (here we need (A.13) and (A.14)) yield

$$\int_{B_{4s}} g dx \lesssim \frac{1}{M(4s)} \sum_{k=0}^n \int_{B_{8s}} |Dv_k|^2 dx, \quad \text{where} \quad M(\rho) := \sup_{B_\rho} v,$$

and

$$\frac{1}{M(4s)} \left(\int_{B_s} |w_k|^{1+1/\delta} dx \right)^{\frac{1}{1+1/\delta}} \leq c \left(1 - \frac{M(\rho)}{M(8s)} \right)^{\frac{1}{1+1/\delta}} \quad \text{for all } \rho \in (0, s].$$

Therefore, combining the last three estimates and (A.17) we see that, for $\rho \in (0, s]$,

$$\sum_{k=0}^n \int_{B_\rho} |Dv_k|^2 dx \lesssim \left(\left(\frac{\rho}{s} \right)^{n-2+\beta} + \left(1 - \frac{M(\rho)}{M(8s)} \right)^{\frac{1}{1+1/\delta}} \right) \sum_{k=0}^n \int_{B_{8s}} |Dv_k|^2 dx.$$

Finally, by a standard iteration argument as in [58, p. 857] and by Poincaré's inequality, we can find $\beta_1 \in (0, \beta)$ such that

$$\sum_{k=0}^n \int_{B_\rho} |v_k - (v_k)_{B_\rho}|^2 dx \lesssim \rho^2 \sum_{k=0}^n \int_{B_\rho} |Dv_k|^2 dx \lesssim \left(\frac{\rho}{s} \right)^{\beta_1} M(4s)^2$$

for any $\rho \in (0, s]$. With the definition of v_k , this implies that, for any $x, y \in B_s$,

$$\begin{aligned} \left| \frac{\varphi_\varepsilon(|Du(x)|)}{|Du(x)|} u_{x_k}(x) - \frac{\varphi_\varepsilon(|Du(y)|)}{|Du(y)|} u_{x_k}(y) \right| &= |v_k(x) - v_k(y)| \\ &\lesssim \left(\frac{|x - y|}{s} \right)^{\beta_1/2} M(8s). \end{aligned}$$

We use Proposition 3.8(1) with φ_ε in place of φ' to conclude that

$$\left| \frac{\varphi_\varepsilon(|z_1|)}{|z_1|} z_1 - \frac{\varphi_\varepsilon(|z_2|)}{|z_2|} z_2 \right| \gtrsim \frac{\varphi_\varepsilon(|z_1| + |z_2|)}{|z_1| + |z_2|} |z_1 - z_2| \geq \varphi_\varepsilon(|z_1 - z_2|)$$

where we have used $|z_1| + |z_2| \geq |z_1 - z_2|$ and (Inc)₁ for φ_ε in the last step. Applying this in the previous estimate with $z_1 = Du(x)$ and $z_2 = Du(y)$, we find that

$$\varphi_\varepsilon(|Du(x) - Du(y)|) \lesssim \left(\frac{|x - y|}{s} \right)^{\beta_1/2} M(8s).$$

We now undo the convention of omitting ε from (A.7) for the final part. Inserting (A.6) with the definition of $M(\rho)$ into the above estimate, we deduce that

$$\varphi_\varepsilon(|Du_\varepsilon(x) - Du_\varepsilon(y)|) \lesssim \left(\frac{|x - y|}{s} \right)^{\beta_1/2} \int_{B_{16s}} \varphi_\varepsilon(|Du_\varepsilon|) dx.$$

At this point, we restrict our attention to the case $\Omega = B_{32s}$ and consider minimizers u_ε of (A.2) with the boundary value restriction $w \in u + W_0^{1,\varphi_\varepsilon}(B_{32s})$. We apply φ_ε^{-1} to both sides and use (Dec)_{max{2,q}} for φ_ε to get

$$|Du_\varepsilon(x) - Du_\varepsilon(y)| \lesssim \left(\frac{|x - y|}{s} \right)^{\alpha_0} \varphi_\varepsilon^{-1} \left(\int_{B_{16s}} \varphi_\varepsilon(|Du|) dx \right)$$

for some $\alpha_0 \in (0, 1)$. Letting $\varepsilon \rightarrow 0$, we can remove ε in the above estimate as in [35, proof of Lemma 4.9]. Finally, by the same argument as in the proof of Lemma 4.7 with (A.6) and Jensen's inequality for the concave function equivalent to $\varphi^{1/q}$ (see Proposition 3.5(2)) we also see that

$$\int_{B_{16s}} \varphi(|Du|) dx \lesssim \left(\int_{B_{32s}} \varphi(|Du|)^{1/q} dx \right)^q \lesssim \varphi \left(\int_{B_{32s}} |Du| dx \right). \quad (\text{A.18})$$

These imply, for any $x, y \in B_s$ and $B_{32s} \Subset \Omega$, that

$$|Du(x) - Du(y)| \leq c \left(\frac{|x - y|}{s} \right)^{\alpha_0} \int_{B_{32s}} |Du| dx,$$

which shows (4.14). In addition, from (A.6) and (A.18), we also have (4.13).

Appendix B. Weighted estimate for autonomous problems

In this appendix, we discuss the global weighted estimate (4.18). For global regularity estimates, the regularity of the boundary of the domain is a delicate issue. In particular, the Reifenberg flat condition is considered sharp for Calderón–Zygmund type estimates for problems in divergence form. Hence we shall give a result for domains satisfying this condition. We say that a bounded domain Ω is (δ, R) -Reifenberg flat for some small

$\delta \in (0, 1)$ and $R > 0$ if for any $y \in \partial\Omega$ and $r \in (0, R]$ there exists an isometric coordinate system with origin at y , say (x_1, \dots, x_n) , such that in this coordinate system,

$$B_r(0) \cap \{x_n > \delta r\} \subset \Omega \cap B_r(0) \subset B_r(0) \cap \{x_n > -\delta r\}.$$

Note that a domain with Lipschitz boundary with Lipschitz seminorm $\delta \in (0, 1)$ is (δ, R) -Reifenberg flat for some $R > 0$ and that the ball B_r is $(\delta, 2\delta r)$ -Reifenberg flat for any $\delta \in (0, 1/2)$.

For $1 \leq s \leq \infty$, let A_s be the Muckenhoupt class. In particular, for $1 < s < \infty$, a weight w (i.e., $w \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $w \geq 0$) is an A_s -weight, $w \in A_s$, if

$$[w]_{A_s} := \sup_{B \subset \mathbb{R}^n} \left(\int_B w \, dx \right) \left(\int_B w^{-\frac{1}{s-1}} \, dx \right)^{s-1} < \infty.$$

For the properties of the A_s class, we refer to [50].

Theorem B.1. *Let $\varphi \in \Phi_c \cap C^1([0, \infty)) \cap C^2((0, \infty))$ with φ' satisfying $(\text{Inc})_{p-1}$ and $(\text{Dec})_{q-1}$ for some $1 < p \leq q < \infty$, and let $w \in A_s$ for some $s \in (1, \infty)$. There exists a small $\delta = \delta(n, p, q, s, [w]_{A_s}) \in (0, 1)$ such that if Ω is (δ, R) -Reifenberg flat for some $R > 0$, $v_0 \in W^{1,\varphi}(\Omega)$ satisfies $\varphi(|Dv_0|) \in L^s_w(\Omega)$ and $v \in W^{1,\varphi}(\Omega)$ is a weak solution to*

$$\operatorname{div} \left(\frac{\varphi'(|Dv|)}{|Dv|} Dv \right) = 0 \quad \text{in } \Omega \quad \text{with} \quad v = v_0 \quad \text{on } \partial\Omega, \quad (\text{B.2})$$

then

$$\int_{\Omega} \varphi(|Dv|)^s w \, dx \leq c \int_{\Omega} \varphi(|Dv_0|)^s w \, dx$$

for some $c = c(n, p, q, s, [w]_{A_s}, \operatorname{diam}(\Omega)/R) > 0$. In particular, letting $\Omega = B_r$ we have (4.18), since B_r is $(\delta, 2\delta r)$ -Reifenberg flat.

Remark B.3. In Theorem B.1, δ is decreasing as a function of $[w]_{A_s}$ [70, Remark 2.2]. Moreover, we can also see by analyzing the proof that the constant c is increasing in $[w]_{A_s}$ and in $\operatorname{diam}(\Omega)/R$ when the other is constant. Therefore, when $\Omega = B_r$, the constant c is increasing in $[w]_{A_s}$, since $\operatorname{diam}(\Omega)/R = 1/(2\delta)$.

Sketch of the proof of Theorem B.1. For the p -Laplacian case, that is, $\varphi(t) = t^p$, the weighted estimate has been proved in [70] (see also [20]) for the equation

$$\operatorname{div}(|Dv|^{p-2} Dv) = \operatorname{div}(|F|^{p-2} F) \quad \text{in } \Omega \quad \text{with} \quad v = 0 \quad \text{on } \partial\Omega.$$

Specifically, in [70], it has been shown that for the above equation,

$$\int_{\Omega} |Dv|^{ps} w \, dx \leq c \int_{\Omega} |F|^{ps} w \, dx$$

for any $w \in A_s$ and any $F \in L^{ps}_w(\Omega, \mathbb{R}^n)$. Moreover, it turns out that this result without a weight (i.e., $w \equiv 1$) is naturally extended [14] to the equation involving a general function φ ,

$$\operatorname{div} \left(\frac{\varphi'(|Dv|)}{|Dv|} Dv \right) = \operatorname{div} \left(\frac{\varphi'(|F|)}{|F|} F \right) \quad \text{in } \Omega \quad \text{with} \quad v = 0 \quad \text{on } \partial\Omega. \quad (\text{B.4})$$

Therefore, proceeding as in [70] with minor modification, one can prove that for the equation (B.4),

$$\int_{\Omega} \varphi(|Dv|)^s w \, dx \leq c \int_{\Omega} \varphi(|F|)^s w \, dx$$

for any $w \in A_s$ and any $F \in L^\varphi(\Omega, \mathbb{R}^n)$ satisfying $\varphi(|F|) \in L_w^s(\Omega)$.

In this theorem, we consider non-zero boundary data v_0 . However, the gradient of v_0 can be handled in much the same way as for F in the results mentioned above. Hence, by the same argument as in [70], replacing t^p by $\varphi(t)$ and changing boundary comparison estimates from [70, Lemma 4.6] to Lemmas B.5 and B.11 below, we obtain the desired estimate. \blacksquare

For the rest of the paper, we suppose the assumptions of Theorem B.1 hold. We consider our problem (B.2) on a local region near the boundary of Ω . Define $\Omega_r(x) := \Omega \cap B_r(x)$, $B_r := B_r(0)$, $\Omega_r = \Omega_r(0)$, $B_r^+ := B_r \cap \{x_n > 0\}$ and $B_r^- := B_r \cap \{x_n < 0\}$. Then we consider our equation in the region Ω_{5r} with $5r < R$ and

$$B_{5r}^+ \subset \Omega_{5r} \subset B_{5r} \cap \{x_n > -10\delta r\}.$$

Here, $\delta \in (0, 1)$ and $R > 0$ come from the (δ, R) -Reifenberg flat condition of Ω and so δ has to be determined later and R is given. Note that in view of the scaling invariance property of (B.2) (see for instance the proof of Lemma 4.15), we may let $r = 1$ and consider assumption (B.7) below.

We first compare our equation (B.2) with an equation having zero boundary values on $\partial\Omega$ in a local region near the boundary.

Lemma B.5. *For $v_0 \in W^{1,\varphi}(\Omega)$ let $v \in W_0^{1,\varphi}(\Omega)$ be a weak solution to (B.2). For any $\varepsilon > 0$ there exists a small $\delta \in (0, 1)$ depending on n, p, q and ε such that if Ω is $(\delta, 5)$ -Reifenberg flat and*

$$B_5^+ \subset \Omega_5 \subset B_5 \cap \{x_n > -10\delta\}, \quad (\text{B.6})$$

$$\int_{\Omega_5} \varphi(|Dv|) \, dx \leq 1 \quad \text{and} \quad \int_{\Omega_5} \varphi(|Dv_0|) \, dx \leq \delta, \quad (\text{B.7})$$

then for the weak solution $w \in W^{1,\varphi}(\Omega_5)$ to

$$\operatorname{div} \left(\frac{\varphi'(|Dw|)}{|Dw|} Dw \right) = 0 \quad \text{in } \Omega_5 \quad \text{and} \quad w = v - v_0 \quad \text{on } \partial\Omega_5, \quad (\text{B.8})$$

we have

$$\int_{\Omega_5} \varphi(|Dw|) \, dx \leq c \quad \text{and} \quad \int_{\Omega_5} \varphi(|Dv - Dw|) \, dx \leq \varepsilon. \quad (\text{B.9})$$

Here, $c > 0$ depends on p and q , but is independent of ε .

Proof. Since $w - v + v_0 \in W_0^{1,\varphi}(\Omega_5)$, by (B.8) and (B.2) we have

$$\int_{\Omega_5} \frac{\varphi'(|Dw|)}{|Dw|} Dw \cdot D(w - v + v_0) \, dx = 0 = \int_{\Omega_5} \frac{\varphi'(|Dv|)}{|Dv|} Dv \cdot D(w - v + v_0) \, dx. \quad (\text{B.10})$$

In view of $\varphi(t) \leq t\varphi'(t)$ and Propositions 3.5(5) and 3.6(4), the first equality above implies that

$$\begin{aligned} \int_{\Omega_5} \varphi(|Dw|) dx &\leq \int_{\Omega_5} \varphi'(|Dw|)|Dv - Dv_0| dx \\ &\leq \int_{\Omega_5} \left[\frac{1}{2}\varphi(|Dw|) dx + c\varphi(|Dv - Dv_0|) \right] dx \end{aligned}$$

for some $c = c(p, q) > 0$. By (B.7), we obtain the first estimate in (B.9).

We next prove the second estimate in (B.9). By Proposition 3.8(1), (B.10) and Propositions 3.5(5) and 3.6(4) we find that for $\kappa_1 \in (0, 1)$,

$$\begin{aligned} \int_{\Omega_5} \frac{\varphi'(|Dw| + |Dv|)}{|Dw| + |Dv|} |Dw - Dv|^2 dx \\ \lesssim \int_{\Omega_5} \left(\frac{\varphi'(|Dw|)}{|Dw|} Dw - \frac{\varphi'(|Dv|)}{|Dv|} Dv \right) \cdot D(w - v) dx \\ \lesssim \int_{\Omega_5} (\varphi'(|Dw|) + \varphi'(|Dv|)) |Dv_0| dx \\ \lesssim \kappa_1 \int_{\Omega_5} [\varphi(|Dw|) + \varphi(|Dv|)] dx + \frac{1}{\kappa_1^{q-1}} \int_{\Omega_5} \varphi(|Dv_0|) dx. \end{aligned}$$

Moreover, by Proposition 3.8(3), for any $\kappa_2 \in (0, 1)$,

$$\varphi(|Dw - Dv|) \leq \kappa_2(\varphi(Dw) + \varphi(Dv)) + \kappa_2^{-1} \frac{\varphi'(|Dw| + |Dv|)}{|Dw| + |Dv|} |Dw - Dv|^2.$$

Combining the above two estimates we have

$$\begin{aligned} \int_{\Omega_5} \varphi(|Dw - Dv|) dx &\lesssim \kappa_2 \int_{\Omega_5} [\varphi(Dw) + \varphi(Dv)] dx \\ &\quad + \frac{\kappa_1}{\kappa_2} \int_{\Omega_5} [\varphi(|Dw|) + \varphi(|Dv|)] dx \\ &\quad + \frac{1}{\kappa_2 \kappa_1^{q-1}} \int_{\Omega_5} \varphi(|Dv_0|) dx. \end{aligned}$$

Finally, applying (B.7) and the first estimate in (B.9) and choosing sufficiently small κ_1 , κ_2 and δ depending n, p, q on ε , we get the second estimate in (B.9). ■

We also notice that the weak solution w to (B.8) has value zero on $\partial\Omega_5 \cap B_5$. We next compare (B.8), which assumes zero boundary values on $\partial\Omega_5 \cap B_5$, with an equation defined in B_2^+ with zero boundary values on $B_2 \cap \{x_n = 0\}$. A similar result can be found in [14, Lemma 3.6]. The proof of that lemma employs a compactness argument. Here we give a more direct approach which clearly shows the dependence on δ .

Lemma B.11. *Let $\eta = \eta(x_n) \in C^\infty(\mathbb{R})$ with $\eta = 0$ if $x_n \leq 0$, $\eta = 1$ if $x_n \geq \delta$ and $|\eta'| \leq 2/\delta$. For any $\varepsilon > 0$ there exists a small $\delta > 0$ depending on n, p, q and ε such that, under the assumptions of the above lemma, if w_0 is the weak solution to*

$$\operatorname{div}\left(\frac{\varphi'(|Dw_0|)}{|Dw_0|}Dw_0\right) = 0 \quad \text{in } B_2^+ \quad \text{and} \quad w_0 = \eta w \quad \text{on } \partial B_2^+,$$

then

$$\int_{B_2^+} \varphi(|Dw_0|) dx \leq c \quad \text{and} \quad \int_{B_2^+} \varphi(|Dw - Dw_0|) dx \leq \varepsilon. \quad (\text{B.12})$$

Moreover,

$$\|\varphi(|Dw_0|)\|_{L^\infty(\Omega_1)} = \|\varphi(|Dw_0|)\|_{L^\infty(B_1^+)} \leq c \int_{B_2^+} \varphi(|Dw_0|) dx \leq c, \quad (\text{B.13})$$

where we extend w_0 by zero to B_2^- . Here the constants c depend on n, p and q , but are independent of ε .

Proof. We follow the technique of [57, Lemma 2.5] (see also [19, Lemma 2.5]). Clearly, (δ, R) -Reifenberg flat domains with $\delta \in (0, 1/2)$ satisfy the measure density condition $|B_r|4^{-n} \leq |\Omega_r(x)| \leq |B_r|$ and $4^{-n}|B_r| \leq |B_r(x) \setminus \Omega_r(x)|$ for all $x \in \partial\Omega$ and $r \in (0, R]$. One can show as in [72, Theorem 3.9] that, for equation (B.8), there exists $\sigma = \sigma(n, p, q) \in (0, 1)$ such that $\varphi(|Dw|) \in L_{\text{loc}}^{1+\sigma}(B_5)$ (we extend w by 0 in $B_5 \setminus \Omega_5$) and

$$\left(\int_{\Omega_3} \varphi(|Dw|)^{1+\sigma} dx\right)^{\frac{1}{1+\sigma}} \lesssim \int_{\Omega_4} \varphi(|Dw|) dx.$$

Then by Hölder's inequality with (B.6), we observe that

$$\int_{\Omega_3 \cap \{x_n \leq \delta\}} \varphi(|Dw|) dx \lesssim \delta^{\frac{\sigma}{1+\sigma}} \left(\int_{\Omega_3} \varphi(|Dw|)^{1+\sigma} dx\right)^{\frac{1}{1+\sigma}} \lesssim \delta^{\frac{\sigma}{1+\sigma}} \int_{\Omega_4} \varphi(|Dw|) dx. \quad (\text{B.14})$$

In addition, using the fact that $w \equiv 0$ in $B_4 \setminus \Omega_4$ and w is absolutely continuous on almost all lines parallel to the coordinate axes, as well as Jensen's inequality, we find that

$$\begin{aligned} \int_{\Omega_2 \cap \{x_n \leq \delta\}} \varphi(|D\eta| |w|) dx &\lesssim \int_{\Omega_2 \cap \{x_n \leq \delta\}} \varphi\left(\frac{1}{\delta} \left|\int_{-8\delta}^{x_n} D_n w(x', y) dy\right|\right) dx \\ &\lesssim \int_{\Omega_2 \cap \{x_n \leq \delta\}} \varphi\left(\int_{-8\delta}^{\delta} |D_n w(x', y)| dy\right) dx \\ &\lesssim \int_{\{|x'| \leq 2\} \times \{-10\delta < x_n \leq \delta\}} \int_{-8\delta}^{\delta} \varphi(|D_n w(x', y)|) dy dx' dx_n \\ &\lesssim \int_{\Omega_3 \cap \{(x', y): y \leq \delta\}} \varphi(|Dw(x', y)|) dx' dy. \end{aligned} \quad (\text{B.15})$$

In the last inequality above, we have used the facts that $\{|x'| \leq 2\} \cap \{|x_n| \leq 2\} \subset B_3$ and $Dw \equiv 0$ in $B_3 \setminus \Omega_3$.

Since $w_0 - \eta w \in W_0^{1,\varphi}(B_2^+)$ and w_0 is a minimizer, using also (B.15) we find that

$$\begin{aligned} \int_{B_2^+} \varphi(|Dw_0|) dx &\leq \int_{B_2^+} \varphi(|D(\eta w)|) dx \\ &\lesssim \int_{B_2^+} \varphi(|Dw|) dx + \int_{B_2^+ \cap \{x_n \leq \delta\}} \varphi(|D\eta||w|) dx \\ &\lesssim \int_{\Omega_3} \varphi(|Dw|) dx, \end{aligned} \quad (\text{B.16})$$

which together with (B.6) and (B.9) yields the first estimate in (B.12).

We next derive the second estimate in (B.12). Since $w_0 - \eta w \in W_0^{1,\varphi}(B_2^+) \cap W_0^{1,\varphi}(\Omega_2)$ we have

$$\int_{B_2^+} \left(\frac{\varphi'(|Dw_0|)}{|Dw_0|} Dw_0 - \frac{\varphi'(|Dw|)}{|Dw|} Dw \right) \cdot D(w_0 - \eta w) dx = 0,$$

which together with Propositions 3.5(5) and 3.6(4) implies that for any $\kappa_1 \in (0, 1)$,

$$\begin{aligned} \int_{B_2^+} \left(\frac{\varphi'(|Dw_0|)}{|Dw_0|} Dw_0 - \frac{\varphi'(|Dw|)}{|Dw|} Dw \right) \cdot D(w_0 - w) dx \\ = \int_{B_2^+ \cap \{x_n \leq \delta\}} \left(\frac{\varphi'(|Dw_0|)}{|Dw_0|} Dw_0 - \frac{\varphi'(|Dw|)}{|Dw|} Dw \right) \cdot D(\eta w - w) dx \\ \leq \int_{B_2^+ \cap \{x_n \leq \delta\}} (\varphi'(|Dw_0|) + \varphi'(|Dw|))(|D\eta||w| + |Dw|) dx \\ \lesssim \kappa_1 \int_{B_2^+ \cap \{x_n \leq \delta\}} [\varphi(|Dw_0|) + \varphi(|Dw|)] dx \\ + \frac{1}{\kappa_1^{q-1}} \int_{B_2^+ \cap \{x_n \leq \delta\}} [\varphi(|D\eta||w|) + \varphi(|Dw|)] dx. \end{aligned}$$

Applying Proposition 3.8(3) and (B.14)–(B.16), we see that for any $\kappa_2 \in (0, 1)$,

$$\int_{B_2^+} \varphi(|Dw_0 - Dw|) dx \lesssim \left(\kappa_2 + \frac{\kappa_1}{\kappa_2} \right) \int_{\Omega_3} \varphi(|Dw|) dx + \frac{\delta^{\frac{q}{1+\sigma}}}{\kappa_2 \kappa_1^{q-1}} \int_{\Omega_4} \varphi(|Dw|) dx.$$

Therefore, using the first estimate of (B.9) and taking sufficiently small κ_1 , κ_2 and δ depending on n , p , q and ε , we have the second estimate in (B.12).

Let $\tilde{w}_0 \in W^{1,\varphi}(B_2)$ be an even extension of w_0 so that $\tilde{w}_0(x) = w_0(x)$ if $x \in B_2^+$ and $\tilde{w}_0(x_1, \dots, x_{n-1}, x_n) = w_0(x_1, \dots, x_{n-1}, -x_n)$ if $(x_1, \dots, x_n) \in B_2^-$. Note that \tilde{w}_0 is well defined since $w_0 = 0$ on $B_r \cap \{x_n = 0\}$. Moreover \tilde{w}_0 is a weak solution to

$$\operatorname{div} \left(\frac{\varphi'(|D\tilde{w}_0|)}{|D\tilde{w}_0|} D\tilde{w}_0 \right) = 0 \quad \text{in } B_2$$

(see for instance [68, Theorem 3.4]). Therefore, (B.13) follows from Lemma 4.12. \blacksquare

Acknowledgments. We thank the referee for useful comments.

Funding. J. Ok was supported by the National Research Foundation of Korea funded by the Korean Government (NRF-2017R1C1B2010328) and P. Hästö was supported in part by the Magnus Ehrnrooth Foundation.

References

- [1] Acerbi, E., Fusco, N.: Regularity for minimizers of nonquadratic functionals: the case $1 < p < 2$. *J. Math. Anal. Appl.* **140**, 115–135 (1989) Zbl [0686.49004](#) MR [997847](#)
- [2] Acerbi, E., Mingione, G.: Regularity results for a class of functionals with non-standard growth. *Arch. Ration. Mech. Anal.* **156**, 121–140 (2001) Zbl [0984.49020](#) MR [1814973](#)
- [3] Acerbi, E., Mingione, G.: Regularity results for stationary electro-rheological fluids. *Arch. Ration. Mech. Anal.* **164**, 213–259 (2002) Zbl [1038.76058](#) MR [1930392](#)
- [4] Acerbi, E., Mingione, G.: Gradient estimates for the $p(x)$ -Laplacean system. *J. Reine Angew. Math.* **584**, 117–148 (2005) Zbl [1093.76003](#) MR [2155087](#)
- [5] Azzollini, A., d’Avenia, P., Pomponio, A.: Quasilinear elliptic equations in \mathbb{R}^N via variational methods and Orlicz–Sobolev embeddings. *Calc. Var. Partial Differential Equations* **49**, 197–213 (2014) Zbl [1285.35039](#) MR [3148112](#)
- [6] Balci, A. K., Diening, L., Surnachev, M.: New examples on Lavrentiev gap using fractals. *Calc. Var. Partial Differential Equations* **59**, art. 180, 34 pp. (2020) Zbl [1453.35082](#) MR [4153906](#)
- [7] Baroni, P.: Riesz potential estimates for a general class of quasilinear equations. *Calc. Var. Partial Differential Equations* **53**, 803–846 (2015) Zbl [1318.35041](#) MR [3347481](#)
- [8] Baroni, P., Colombo, M., Mingione, G.: Harnack inequalities for double phase functionals. *Nonlinear Anal.* **121**, 206–222 (2015) Zbl [1321.49059](#) MR [3348922](#)
- [9] Baroni, P., Colombo, M., Mingione, G.: Nonautonomous functionals, borderline cases and related function classes. *Algebra i Analiz* **27**, 6–50 (2015) Zbl [1335.49057](#) MR [3570955](#)
- [10] Baroni, P., Colombo, M., Mingione, G.: Regularity for general functionals with double phase. *Calc. Var. Partial Differential Equations* **57**, art. 62, 48 pp. (2018) Zbl [1394.49034](#) MR [3775180](#)
- [11] Beck, L., Mingione, G.: Lipschitz bounds and nonuniform ellipticity. *Comm. Pure Appl. Math.* **73**, 944–1034 (2020) Zbl [1445.35140](#) MR [4078712](#)
- [12] Bella, P., Schäffner, M.: On the regularity of minimizers for scalar integral functionals with (p, q) -growth. *Anal. PDE* **13**, 2241–2257 (2020) Zbl [07324216](#) MR [4175825](#)
- [13] Benyaiche, A., Harjulehto, P., Hästö, P., Karppinen, A.: The weak Harnack inequality for unbounded supersolutions of equations with generalized Orlicz growth. *J. Differential Equations* **275**, 790–814 (2021) Zbl [1455.35088](#) MR [4191341](#)
- [14] Byun, S.-S., Cho, Y.: Nonlinear gradient estimates for generalized elliptic equations with non-standard growth in nonsmooth domains. *Nonlinear Anal.* **140**, 145–165 (2016) Zbl [1338.35167](#) MR [3492733](#)
- [15] Byun, S.-S., Oh, J.: Global gradient estimates for non-uniformly elliptic equations. *Calc. Var. Partial Differential Equations* **56**, art. 46, 36 pp. (2017) Zbl [1378.35139](#) MR [3624942](#)
- [16] Byun, S.-S., Oh, J.: Global gradient estimates for the borderline case of double phase problems with BMO coefficients in nonsmooth domains. *J. Differential Equations* **263**, 1643–1693 (2017) Zbl [1372.35115](#) MR [3632230](#)

- [17] Byun, S.-S., Oh, J.: Regularity results for generalized double phase functionals. *Anal. PDE* **13**, 1269–1300 (2020) Zbl [07271830](#) MR [4149062](#)
- [18] Byun, S.-S., Ok, J.: On $W^{1,q(\cdot)}$ -estimates for elliptic equations of $p(x)$ -Laplacian type. *J. Math. Pures Appl.* (9) **106**, 512–545 (2016) Zbl [1344.35058](#) MR [3520446](#)
- [19] Byun, S.-S., Ok, J., Youn, Y.: Global gradient estimates for spherical quasi-minimizers of integral functionals with $p(x)$ -growth. *Nonlinear Anal.* **177**, 186–208 (2018) Zbl [1402.35095](#) MR [3865194](#)
- [20] Byun, S.-S., Ryu, S.: Global weighted estimates for the gradient of solutions to nonlinear elliptic equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **30**, 291–313 (2013) Zbl [1292.35127](#) MR [3035978](#)
- [21] Capone, C., Cruz-Uribe, D., Fiorenza, A.: A modular variable Orlicz inequality for the local maximal operator. *Georgian Math. J.* **25**, 201–206 (2018) Zbl [1392.42019](#) MR [3808281](#)
- [22] Cencelj, M., Rădulescu, V. D., Repovš, D. D.: Double phase problems with variable growth. *Nonlinear Anal.* **177**, 270–287 (2018) Zbl [1421.35235](#) MR [3865198](#)
- [23] Chlebicka, I., Gwiazda, P., Zatorska-Goldstein, A.: Well-posedness of parabolic equations in the non-reflexive and anisotropic Musielak–Orlicz spaces in the class of renormalized solutions. *J. Differential Equations* **265**, 5716–5766 (2018) Zbl [1397.35126](#) MR [3857497](#)
- [24] Chlebicka, I., Gwiazda, P., Zatorska-Goldstein, A.: Parabolic equation in time and space dependent anisotropic Musielak–Orlicz spaces in absence of Lavrentiev’s phenomenon. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **36**, 1431–1465 (2019) Zbl [1419.35092](#) MR [3985549](#)
- [25] Colombo, M., Mingione, G.: Bounded minimisers of double phase variational integrals. *Arch. Ration. Mech. Anal.* **218**, 219–273 (2015) Zbl [1325.49042](#) MR [3360738](#)
- [26] Colombo, M., Mingione, G.: Regularity for double phase variational problems. *Arch. Ration. Mech. Anal.* **215**, 443–496 (2015) Zbl [1322.49065](#) MR [3294408](#)
- [27] Colombo, M., Mingione, G.: Calderón–Zygmund estimates and non-uniformly elliptic operators. *J. Funct. Anal.* **270**, 1416–1478 (2016) Zbl [06535749](#) MR [3447716](#)
- [28] Coscia, A., Mingione, G.: Hölder continuity of the gradient of $p(x)$ -harmonic mappings. *C. R. Acad. Sci. Paris Sér. I Math.* **328**, 363–368 (1999) Zbl [0920.49020](#) MR [1675954](#)
- [29] Cruz-Uribe, D., Hästö, P.: Extrapolation and interpolation in generalized Orlicz spaces. *Trans. Amer. Math. Soc.* **370**, 4323–4349 (2018) Zbl [1391.46037](#) MR [3811530](#)
- [30] De Filippis, C., Mingione, G.: A borderline case of Calderón–Zygmund estimates for nonuniformly elliptic problems. *Algebra i Analiz* **31**, 82–115 (2019) Zbl [1435.35127](#) MR [3985927](#)
- [31] De Filippis, C., Oh, J.: Regularity for multi-phase variational problems. *J. Differential Equations* **267**, 1631–1670 (2019) Zbl [1422.49037](#) MR [3945612](#)
- [32] DiBenedetto, E.: $C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations. *Nonlinear Anal.* **7**, 827–850 (1983) Zbl [0539.35027](#) MR [709038](#)
- [33] Diening, L., Ettwein, F.: Fractional estimates for non-differentiable elliptic systems with general growth. *Forum Math.* **20**, 523–556 (2008) Zbl [1188.35069](#) MR [2418205](#)
- [34] Diening, L., Harjulehto, P., Hästö, P., Růžička, M.: *Lebesgue and Sobolev Spaces with Variable Exponents*. Lecture Notes in Math. 2017, Springer, Heidelberg (2011) Zbl [1222.46002](#) MR [2790542](#)
- [35] Diening, L., Stroffolini, B., Verde, A.: Everywhere regularity of functionals with φ -growth. *Manuscripta Math.* **129**, 449–481 (2009) Zbl [1168.49035](#) MR [2520895](#)

- [36] Eleuteri, M., Marcellini, P., Mascolo, E.: Lipschitz continuity for energy integrals with variable exponents. *Rend. Lincei Mat. Appl.* **27**, 61–87 (2016) Zbl [1338.35169](#) MR [3470676](#)
- [37] Eleuteri, M., Marcellini, P., Mascolo, E.: Lipschitz estimates for systems with ellipticity conditions at infinity. *Ann. Mat. Pura Appl.* (4) **195**, 1575–1603 (2016) Zbl [1354.35035](#) MR [3537963](#)
- [38] Eleuteri, M., Marcellini, P., Mascolo, E.: Regularity for scalar integrals without structure conditions. *Adv. Calc. Var.* **13**, 279–300 (2020) Zbl [1445.35159](#) MR [4116617](#)
- [39] Esposito, L., Leonetti, F., Mingione, G.: Sharp regularity for functionals with (p, q) growth. *J. Differential Equations* **204**, 5–55 (2004) Zbl [1072.49024](#) MR [2076158](#)
- [40] Evans, L. C.: A new proof of local $C^{1,\alpha}$ regularity for solutions of certain degenerate elliptic p.d.e. *J. Differential Equations* **45**, 356–373 (1982) Zbl [0508.35036](#) MR [672713](#)
- [41] Fan, X.: Global $C^{1,\alpha}$ regularity for variable exponent elliptic equations in divergence form. *J. Differential Equations* **235**, 397–417 (2007) Zbl [1143.35040](#) MR [2317489](#)
- [42] Fan, X., Zhao, D.: A class of De Giorgi type and Hölder continuity. *Nonlinear Anal.* **36**, 295–318 (1999) Zbl [0927.46022](#) MR [1688232](#)
- [43] Fang, Y., Zhang, C., Zhang, X.: Estimates for multi-phase problems in Campanato space. Preprint (2019).
- [44] Giannetti, F., Passarelli di Napoli, A.: Regularity results for a new class of functionals with non-standard growth conditions. *J. Differential Equations* **254**, 1280–1305 (2013) Zbl [1255.49064](#) MR [2997371](#)
- [45] Giannetti, F., Passarelli di Napoli, A., Ragusa, M. A., Tachikawa, A.: Partial regularity for minimizers of a class of non autonomous functionals with nonstandard growth. *Calc. Var. Partial Differential Equations* **56**, art. 153, 29 pp. (2017) Zbl [1383.49047](#) MR [3709297](#)
- [46] Giaquinta, M.: *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*. Ann. of Math. Stud. 105, Princeton Univ. Press, Princeton, NJ (1983) Zbl [0516.49003](#) MR [717034](#)
- [47] Giaquinta, M., Giusti, E.: Differentiability of minima of nondifferentiable functionals. *Invent. Math.* **72**, 285–298 (1983) Zbl [0513.49003](#) MR [700772](#)
- [48] Giaquinta, M., Giusti, E.: Global $C^{1,\alpha}$ -regularity for second order quasilinear elliptic equations in divergence form. *J. Reine Angew. Math.* **351**, 55–65 (1984) Zbl [0528.35014](#) MR [749677](#)
- [49] Giusti, E.: *Direct Methods in the Calculus of Variations*. World Sci., River Edge, NJ (2003) Zbl [1028.49001](#) MR [1962933](#)
- [50] Grafakos, L.: *Classical Fourier Analysis*. 3rd ed., Grad. Texts in Math. 249, Springer, New York (2014) Zbl [1304.42001](#) MR [3243734](#)
- [51] Harjulehto, P., Hästö, P.: *Orlicz Spaces and Generalized Orlicz Spaces*. Lecture Notes in Math. 2236, Springer, Cham (2019) Zbl [1436.46002](#) MR [3931352](#)
- [52] Harjulehto, P., Hästö, P., Karppinen, A.: Local higher integrability of the gradient of a quasiminimizer under generalized Orlicz growth conditions. *Nonlinear Anal.* **177**, 543–552 (2018) Zbl [1403.49034](#) MR [3886589](#)
- [53] Harjulehto, P., Hästö, P., Klén, R.: Generalized Orlicz spaces and related PDE. *Nonlinear Anal.* **143**, 155–173 (2016) Zbl [1360.46029](#) MR [3516828](#)
- [54] Harjulehto, P., Hästö, P., Lee, M.: Hölder continuity of quasiminimizers and ω -minimizers with generalized Orlicz growth. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **22**, 549–582 (2021)

- [55] Harjulehto, P., Hästö, P., Toivanen, O.: Hölder regularity of quasiminimizers under generalized growth conditions. *Calc. Var. Partial Differential Equations* **56**, art. 22, 26 pp. (2017) Zbl [1366.35036](#) MR [3606780](#)
- [56] Jin, T., Maz'ya, V., Van Schaftingen, J.: Pathological solutions to elliptic problems in divergence form with continuous coefficients. *C. R. Math. Acad. Sci. Paris* **347**, 773–778 (2009) Zbl [1198.35055](#) MR [2543981](#)
- [57] Kim, Y., Ryu, S.: Global gradient estimates for parabolic equations with measurable nonlinearities. *Nonlinear Anal.* **164**, 77–99 (2017) Zbl [1373.35165](#) MR [3712021](#)
- [58] Lewis, J. L.: Regularity of the derivatives of solutions to certain degenerate elliptic equations. *Indiana Univ. Math. J.* **32**, 849–858 (1983) Zbl [0554.35048](#) MR [721568](#)
- [59] Liang, S., Cai, M., Zheng, S.: Global regularity in Lorentz spaces for nonlinear elliptic equations with $L^{p(\cdot)} \log L$ -growth. *J. Math. Anal. Appl.* **467**, 67–94 (2018) Zbl [1397.35052](#) MR [3834795](#)
- [60] Lieberman, G. M.: The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva for elliptic equations. *Comm. Partial Differential Equations* **16**, 311–361 (1991) Zbl [0742.35028](#) MR [1104103](#)
- [61] Maeda, F.-Y., Mizuta, Y., Ohno, T., Shimomura, T.: Sobolev's inequality for double phase functionals with variable exponents. *Forum Math.* **31**, 517–527 (2019) Zbl [1423.46049](#) MR [3918454](#)
- [62] Manfredi, J. J.: Regularity of the gradient for a class of nonlinear possibly degenerate elliptic equations. PhD thesis, Washington Univ. in St. Louis (1986) MR [2635642](#)
- [63] Marcellini, P.: Regularity of minimizers of integrals of the calculus of variations with non-standard growth conditions. *Arch. Ration. Mech. Anal.* **105**, 267–284 (1989) Zbl [0667.49032](#) MR [969900](#)
- [64] Marcellini, P.: Regularity and existence of solutions of elliptic equations with p, q -growth conditions. *J. Differential Equations* **90**, 1–30 (1991) Zbl [0724.35043](#) MR [1094446](#)
- [65] Marcellini, P.: Regularity for elliptic equations with general growth conditions. *J. Differential Equations* **105**, 296–333 (1993) Zbl [0812.35042](#) MR [1240398](#)
- [66] Marcellini, P.: Everywhere regularity for a class of elliptic systems without growth conditions. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **23**, 1–25 (1996) Zbl [0922.35031](#) MR [1401415](#)
- [67] Marcellini, P.: Regularity for some scalar variational problems under general growth conditions. *J. Optim. Theory Appl.* **90**, 161–181 (1996) Zbl [0901.49030](#) MR [1397651](#)
- [68] Martio, O.: Reflection principle for solutions of elliptic partial differential equations and quasiregular mappings. *Ann. Acad. Sci. Fenn. Ser. A I Math.* **6**, 179–187 (1981) Zbl [0473.30016](#) MR [639975](#)
- [69] Méndez, O., Lang, J.: *Analysis on Function Spaces of Musielak–Orlicz Type*. CRC Press, Boca Raton, FL (2019) Zbl [06993277](#) MR [3889985](#)
- [70] Mengesha, T., Phuc, N. C.: Global estimates for quasilinear elliptic equations on Reifenberg flat domains. *Arch. Ration. Mech. Anal.* **203**, 189–216 (2012) Zbl [1255.35113](#) MR [2864410](#)
- [71] Mingione, G.: Regularity of minima: an invitation to the dark side of the calculus of variations. *Appl. Math.* **51**, 355–426 (2006) Zbl [1164.49324](#) MR [2291779](#)
- [72] Ok, J.: Gradient estimates for elliptic equations with $L^{p(\cdot)} \log L$ growth. *Calc. Var. Partial Differential Equations* **55**, art. 26, 30 pp. (2016) Zbl [1342.35090](#) MR [3465442](#)

- [73] Ok, J.: Regularity of ω -minimizers for a class of functionals with non-standard growth. *Calc. Var. Partial Differential Equations* **56**, art. 48, 31 pp. (2017) Zbl [1369.49050](#) MR [3626319](#)
- [74] Ok, J.: Harnack inequality for a class of functionals with non-standard growth via De Giorgi's method. *Adv. Nonlinear Anal.* **7**, 167–182 (2018) Zbl [1397.35115](#) MR [3794882](#)
- [75] Rădulescu, V. D.: Nonlinear elliptic equations with variable exponent: old and new. *Nonlinear Anal.* **121**, 336–369 (2015) Zbl [1321.35030](#) MR [3348928](#)
- [76] Ragusa, M. A., Tachikawa, A.: Regularity for minimizers for functionals of double phase with variable exponents. *Adv. Nonlinear Anal.* **9**, 710–728 (2020) Zbl [1420.35145](#) MR [3985000](#)
- [77] Shi, X., Rădulescu, V. D., Repovš, D. D., Zhang, Q.: Multiple solutions of double phase variational problems with variable exponent. *Adv. Calc. Var.* **13**, 385–401 (2020) Zbl [1454.49006](#) MR [4156782](#)
- [78] Tolksdorf, P.: Regularity for a more general class of quasilinear elliptic equations. *J. Differential Equations* **51**, 126–150 (1984) Zbl [0488.35017](#) MR [727034](#)
- [79] Uhlenbeck, K.: Regularity for a class of non-linear elliptic systems. *Acta Math.* **138**, 219–240 (1977) Zbl [0372.35030](#) MR [474389](#)
- [80] Ural'tseva, N. N.: Degenerate quasilinear elliptic systems. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklova* **7**, 184–222 (1968) (in Russian) Zbl [0196.12501](#) MR [0244628](#)
- [81] Wang, B., Liu, D., Zhao, P.: Hölder continuity for nonlinear elliptic problem in Musielak–Orlicz–Sobolev space. *J. Differential Equations* **266**, 4835–4863 (2019) Zbl [1420.35099](#) MR [3912735](#)
- [82] Wang, L.: Compactness methods for certain degenerate elliptic systems. *Manuscripta Math.* **78**, 273–285 (1993) Zbl [0819.35067](#) MR [1206157](#)
- [83] Wang, L.: Compactness methods for certain degenerate elliptic equations. *J. Differential Equations* **107**, 341–350 (1994) Zbl [0792.35067](#) MR [1264526](#)
- [84] Zhang, Q., Rădulescu, V. D.: Double phase anisotropic variational problems and combined effects of reaction and absorption terms. *J. Math. Pures Appl.* (9) **118**, 159–203 (2018) Zbl [1404.35191](#) MR [3852472](#)
- [85] Zhikov, V. V.: Averaging of functionals of the calculus of variations and elasticity theory. *Izv. Akad. Nauk SSSR Ser. Mat.* **50**, 675–710, 877 (1986) (in Russian) Zbl [0599.49031](#) MR [0864171](#)
- [86] Zhikov, V. V.: On Lavrentiev's phenomenon. *Russian J. Math. Phys.* **3**, 249–269 (1995) Zbl [0910.49020](#) MR [1350506](#)