

François Hamel · Nikolai Nadirashvili

Circular flows for the Euler equations in two-dimensional annular domains, and related free boundary problems

Received September 3, 2019

Abstract. In this paper, we first consider steady Euler flows in two-dimensional bounded annuli, as well as in complements of disks, in punctured disks and in the punctured plane. We prove that if the flow does not have any stagnation point and satisfies rigid wall boundary conditions together with further conditions at infinity in the case of unbounded domains and at the center in the case of punctured domains, then the flow is circular, in the sense that the streamlines are concentric circles. In other words, the flow then inherits the radial symmetry of the domain. We then show two classification results for the steady Euler equations in simply or doubly connected bounded domains with free boundaries. Here, the flows are further assumed to have constant norm on each connected component of the boundary, and the domains are then proved to be disks or annuli. On the one hand, the proofs use ODE and PDE arguments to establish some geometric properties of the trajectories of the flow and the orthogonal trajectories of the gradient of the stream function. On the other hand, we also show some comparison results of independent interest for a derived semilinear elliptic equation satisfied by the stream function. These last results, which are based on the method of moving planes, adapted here to some almost circular domains located between some streamlines of the flow, lead with a limiting argument to the radial symmetry of the stream function and the streamlines of the flow.

Keywords. Euler equations, circular flows, semilinear elliptic equations, free boundary problems, method of moving planes

1. Introduction and main results

Throughout this paper, $|\cdot|$ denotes the Euclidean norm in \mathbb{R}^2 and, for $0 \leq a < b \leq \infty$, $\Omega_{a,b}$ denotes the two-dimensional domain defined by

$$\Omega_{a,b} = \{x \in \mathbb{R}^2 : a < |x| < b\}.$$

François Hamel: Aix Marseille Univ, CNRS, I2M, Marseille, France; francois.hamel@univ-amu.fr

Nikolai Nadirashvili: Aix Marseille Univ, CNRS, I2M, Marseille, France;
nikolay.nadirashvili@univ-amu.fr

Mathematics Subject Classification (2020): 35B06, 35B50, 35B53, 35J61, 76B03

When $a < b$ are two positive real numbers, $\Omega_{a,b}$ is a bounded smooth annulus. When $0 < a < b = \infty$, $\Omega_{a,\infty}$ is an exterior domain which is the complement of a closed disk. When $0 = a < b < \infty$, $\Omega_{0,b}$ is a punctured disk. When $0 = a < b = \infty$, $\Omega_{0,\infty}$ is the punctured plane $\mathbb{R}^2 \setminus \{0\}$, where we denote $0 = (0, 0)$ with a slight abuse of notation.

We also denote

$$e_r(x) = \frac{x}{|x|} \quad \text{and} \quad e_\theta(x) = e_r(x)^\perp = \left(-\frac{x_2}{|x|}, \frac{x_1}{|x|} \right)$$

for $x = (x_1, x_2) \in \mathbb{R}^2 \setminus \{0\}$. Moreover, for $x \in \mathbb{R}^2$ and $r > 0$,

$$B(x, r) = \{y \in \mathbb{R}^2 : |y - x| < r\}$$

denotes the open Euclidean disk with center x and radius r . We also write $B_r = B(0, r)$ and

$$C_r = \partial B_r = \{x \in \mathbb{R}^2 : |x| = r\}.$$

1.1. Liouville-type radial symmetry results for steady Euler flows in $\Omega_{a,b}$

In $\Omega_{a,b}$, we consider steady flows $v = (v_1, v_2)$ of an inviscid fluid, solving the system of the Euler equations:

$$\begin{cases} v \cdot \nabla v + \nabla p = 0 & \text{in } \Omega_{a,b}, \\ \operatorname{div} v = 0 & \text{in } \Omega_{a,b}, \end{cases} \tag{1.1}$$

where the solutions v and p are always understood in the classical sense, that is, they are (at least) of class C^1 in $\Omega_{a,b}$ and therefore satisfy (1.1) everywhere in $\Omega_{a,b}$. We always assume rigid wall boundary conditions, that is, v is (at least) continuous up to the regular parts of $\partial\Omega_{a,b}$ and tangential there:

$$\begin{cases} v \cdot e_r = 0 \text{ on } C_a & \text{if } a > 0, \\ v \cdot e_r = 0 \text{ on } C_b & \text{if } b < \infty. \end{cases} \tag{1.2}$$

The Euler equations are an old but still very active research field. The search for qualitative properties of steady solutions is an important aspect of the study of the Euler flows, and the first main motivation of our paper is to understand the effect of the geometry of the underlying domain $\Omega_{a,b}$ on the properties of steady flows, and more precisely to get some conditions on the flow which guarantee its radial symmetry (see the precise definition below). In other words, how does the solution inherit the geometric radial symmetry properties of the domain? Our primary goal is thus to establish some Liouville-type results for the Euler equations. But the paper is also concerned with related Serrin-type free boundary problems, for which the underlying domain, simply or doubly connected, is free, but is eventually proved to be circular due to additional boundary conditions. Lastly, the paper contains some new comparison results of independent interest on semilinear elliptic equations in doubly connected domains, which are used to show the rigidity results for the Euler equations in given domains and for the related free boundary problems.

A flow v in $\Omega_{a,b}$ is called a *circular flow* if $v(x)$ is parallel to the vector $e_\theta(x)$ at every point $x \in \Omega_{a,b}$, that is, $v \cdot e_r = 0$ in $\Omega_{a,b}$. The main goal of the results of this subsection is to show that, under some conditions, the flow is circular. We obtain such results in the four cases $0 < a < b < \infty$, $0 < a < b = \infty$, $0 = a < b < \infty$, and $0 = a < b = \infty$.

The case of bounded smooth annuli $\Omega_{a,b}$ with $0 < a < b < \infty$. The first result is concerned with flows having no stagnation point in the closed annulus $\overline{\Omega_{a,b}}$. Throughout the paper, the stagnation points of a flow v are the points x for which $|v(x)| = 0$.

Theorem 1.1. *Assume $0 < a < b < \infty$. Let v be a $C^2(\overline{\Omega_{a,b}})$ flow solving (1.1)–(1.2) and such that $|v| > 0$ in $\overline{\Omega_{a,b}}$. Then v is a circular flow, and there is a $C^2([a, b])$ function V with constant strict sign such that*

$$v(x) = V(|x|)e_\theta(x) \quad \text{for all } x \in \overline{\Omega_{a,b}}.$$

It actually turns out that the assumption $|v| > 0$ in $\overline{\Omega_{a,b}}$ can be slightly relaxed. Namely, if $|v| > 0$ in the open annulus $\Omega_{a,b}$ and if the set of stagnation points is assumed to be properly included in one of the connected components of $\partial\Omega_{a,b}$, then the same conclusion holds, and then in fact $|v| > 0$ in $\overline{\Omega_{a,b}}$. This is the purpose of the following result.

Theorem 1.2. *Assume $0 < a < b < \infty$. Let v be a $C^2(\overline{\Omega_{a,b}})$ flow solving (1.1)–(1.2) and such that*

$$\{x \in \overline{\Omega_{a,b}} : |v(x)| = 0\} \not\subset C_a \quad \text{or} \quad \{x \in \overline{\Omega_{a,b}} : |v(x)| = 0\} \not\subset C_b. \quad (1.3)$$

Then $|v| > 0$ in $\overline{\Omega_{a,b}}$ and the conclusion of Theorem 1.1 holds.

Theorem 1.2 is clearly stronger than Theorem 1.1, but we preferred to state Theorem 1.1 separately since the assumption is simpler.

Several further comments are in order. First of all, despite the fact that $\Omega_{a,b}$ is not simply connected, the flow v has a stream function $u : \overline{\Omega_{a,b}} \rightarrow \mathbb{R}$ of class $C^3(\overline{\Omega_{a,b}})$ defined by

$$\nabla^\perp u = v, \quad \text{that is,} \quad \frac{\partial u}{\partial x_1} = v_2 \quad \text{and} \quad \frac{\partial u}{\partial x_2} = -v_1 \quad (1.4)$$

in $\overline{\Omega_{a,b}}$, since v is divergence free and tangential on C_a . Notice that the stream function u is uniquely defined in $\overline{\Omega_{a,b}}$ up to an additive constant. Theorems 1.1 and 1.2 can then be viewed as Liouville-type symmetry results since their conclusion means that the stream function u is radially symmetric (and strictly monotone with respect to $|x|$ in $\overline{\Omega_{a,b}}$). Furthermore, if for x in $\overline{\Omega_{a,b}}$ one denotes by ξ_x the solution of

$$\begin{cases} \dot{\xi}_x(t) = v(\xi_x(t)), \\ \xi_x(0) = x, \end{cases} \quad (1.5)$$

the conclusion of Theorems 1.1 and 1.2 implies that each function ξ_x is defined in \mathbb{R} and periodic, and that the streamlines $\Xi_x = \xi_x(\mathbb{R})$ of the flow are concentric circles.

Theorems 1.1 and 1.2 also mean equivalently that any $C^2(\overline{\Omega_{a,b}})$ noncircular flow for (1.1)–(1.2) must have a stagnation point either in the open annulus $\Omega_{a,b}$, or in both circles C_a and C_b , or the set of stagnation points is equal to C_a , or it is equal to C_b .

Without the assumption $|v| > 0$ in $\overline{\Omega_{a,b}}$ or the weaker one (1.3), the conclusion of Theorems 1.1 and 1.2 obviously does not hold in general, in the sense that there are noncircular flows which do not fulfill (1.3). To construct such flows explicitly, we first point out that, for any continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ and any nonradial $C^2(\overline{\Omega_{a,b}})$ solution u of

$$\Delta u + f(u) = 0 \tag{1.6}$$

in $\Omega_{a,b}$ which is constant on C_a and on C_b and which has a critical point in $\overline{\Omega_{a,b}}$, the $C^1(\overline{\Omega_{a,b}})$ field $v = \nabla^\perp u$ is a noncircular solution of (1.1)–(1.2) with a stagnation point in $\overline{\Omega_{a,b}}$: notice indeed that $v = \nabla^\perp u$ satisfies the boundary condition

$$v \cdot e_r = -\nabla u \cdot e_\theta = 0$$

on $\partial\Omega_{a,b}$ since u is constant on C_a and on C_b , and v solves (1.1) with pressure

$$p = -|v|^2/2 - F(u) = -|\nabla u|^2/2 - F(u),$$

where $F' = f$. As an example, let $\lambda \in \mathbb{R}$ and $\varphi \in C^\infty([a, b])$ be the principal eigenvalue and the principal eigenfunction of the eigenvalue problem

$$-\varphi''(r) - r^{-1}\varphi'(r) + r^{-2}\varphi(r) = \lambda\varphi(r) \quad \text{in } [a, b]$$

with $\varphi > 0$ in (a, b) and Dirichlet boundary condition $\varphi(a) = \varphi(b) = 0$ (the principal eigenvalue λ is unique and the principal eigenfunction φ is unique up to multiplication by positive constants). The $C^\infty(\overline{\Omega_{a,b}})$ function u defined by $u(x) = \varphi(|x|)x_1/|x|$ (that is, $u(x) = \varphi(r) \cos(\theta)$ in the usual polar coordinates) satisfies

$$\Delta u + \lambda u = 0 \quad \text{in } \overline{\Omega_{a,b}}$$

and it has some critical points in $\overline{\Omega_{a,b}}$ (since $\min_{\overline{\Omega_{a,b}}} u < 0 < \max_{\overline{\Omega_{a,b}}} u$ and $u = 0$ on $\partial\Omega_{a,b}$). Actually, it can easily be seen that φ has only one critical point in $[a, b]$ and that u has exactly six critical points in $\overline{\Omega_{a,b}}$ (two in $\Omega_{a,b}$, two on C_a , and two on C_b). Then the $C^\infty(\overline{\Omega_{a,b}})$ flow $v = \nabla^\perp u$ is a noncircular flow solving (1.1)–(1.2) and having two stagnation points in $\Omega_{a,b}$ and four on $\partial\Omega_{a,b}$.

However, we do not know whether the hypothesis (1.3) could be more relaxed for the conclusion of Theorems 1.1 and 1.2 to still hold. For instance, would it be sufficient to assume that v has no stagnation point in $\Omega_{a,b}$? We refer to the comments after the proof of Theorems 1.1 and 1.2 in Section 3.1 below for further details on this question.

On the other hand, we point out that the sufficient conditions $|v| > 0$ in $\overline{\Omega_{a,b}}$ or the more general one (1.3) are obviously not equivalent to being a circular flow, in the sense that there are circular flows for (1.1)–(1.2) which do not fulfill (1.3) (besides the trivial flow $v = (0, 0)$!). Actually, any $C^1(\overline{\Omega_{a,b}})$ circular flow $v(x) = V(|x|)e_\theta(x)$ solving (1.1)–(1.2) and for which $V \in C^1([a, b])$ does not have a constant strict sign, has a set of

stagnation points containing at least a circle. For instance, let $\mu \in \mathbb{R}$ and $\phi \in C^\infty([a, b])$ be the principal eigenvalue and the principal eigenfunction of the eigenvalue problem

$$-\phi''(r) - r^{-1}\phi'(r) = \mu\phi(r) \quad \text{in } [a, b],$$

with $\phi > 0$ in (a, b) and Dirichlet boundary condition $\phi(a) = \phi(b) = 0$, and let $u = \phi(|\cdot|)$. Then $v = \nabla^\perp u = \phi'(|\cdot|)e_\theta$ is a $C^\infty(\overline{\Omega_{a,b}})$ nontrivial circular flow solving (1.1)–(1.2) with pressure $p(x) = -\phi'(|x|)^2/2 - \mu\phi(|x|)^2/2$ and with a circle of stagnation points in $\overline{\Omega_{a,b}}$: more precisely, if $r^* \in (a, b)$ denotes a real number such that $\phi(r^*) = \max_{[a,b]}\phi$ (it is easy to see that r^* is the only critical point of ϕ in $[a, b]$), then the set of stagnation points of the flow v is equal to the whole circle C_{r^*} .

Lastly, the assumption on the $C^2(\overline{\Omega_{a,b}})$ smoothness of v is a technical assumption which is used in the proof. It warrants the C^1 smoothness of the vorticity

$$\omega = \frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2},$$

satisfying $v \cdot \nabla \omega = 0$ in $\Omega_{a,b}$, and the C^1 smoothness of the vorticity function f arising in the semilinear elliptic equation of the type (1.6) satisfied by the stream function u . We refer to the proofs of the preliminary results in Section 2 and especially Lemma 2.8 below for further details.

The case of exterior domains $\Omega_{a,\infty}$ with $0 < a < \infty$.

Theorem 1.3. *Assume $0 < a < \infty$ and $b = \infty$. Let v be a $C^2(\overline{\Omega_{a,\infty}})$ flow solving (1.1)–(1.2) and such that*

$$\{x \in \overline{\Omega_{a,\infty}} : |v(x)| = 0\} \subsetneq C_a \quad \text{and} \quad \liminf_{|x| \rightarrow +\infty} |v(x)| > 0. \quad (1.7)$$

Assume moreover that

$$v(x) \cdot e_r(x) = o(1/|x|) \quad \text{as } |x| \rightarrow +\infty. \quad (1.8)$$

Then $|v| > 0$ in $\overline{\Omega_{a,\infty}}$ and v is a circular flow, namely there is a $C^2([a, +\infty))$ function V with constant strict sign such that $v(x) = V(|x|)e_\theta(x)$ for all $x \in \overline{\Omega_{a,\infty}}$.

As for Theorems 1.1 and 1.2, the conclusion of Theorem 1.3 says that the stream function u is radially symmetric and strictly monotone with respect to $|x|$ in $\overline{\Omega_{a,\infty}}$, and that the streamlines of the flow v are concentric circles.

In the proof of Theorem 1.3 given in Section 3.2, the $o(1/|x|)$ behavior in (1.8) seems nearly optimal. Let us show that without (1.8) the conclusion of Theorem 1.3 does not hold in general. Consider the $C^\infty(\overline{\Omega_{a,\infty}})$ function u defined by $u(x) = 2(|x|^2/a^2 - 1) + (|x|/a - a/|x|)x_1/|x|$, that is,

$$u = 2\left(\frac{r^2}{a^2} - 1\right) + \left(\frac{r}{a} - \frac{a}{r}\right)\cos\theta$$

in the usual polar coordinates. The function u satisfies $\Delta u - 8/a^2 = 0$ in $\overline{\Omega_{a,\infty}}$ with Dirichlet boundary condition $u = 0$ on C_a , and the $C^\infty(\overline{\Omega_{a,\infty}})$ field $v = \nabla^\perp u$ satisfies (1.1)–(1.2) with pressure $p = -|v|^2/2 + 8u/a^2$. In the usual polar coordinates, the field v is given by

$$v = \left[\frac{4r}{a^2} + \left(\frac{1}{a} + \frac{a}{r^2} \right) \cos \theta \right] e_\theta + \left[\left(\frac{1}{a} - \frac{a}{r^2} \right) \sin \theta \right] e_r. \tag{1.9}$$

It satisfies condition (1.7) (and even $\inf_{\Omega_{a,\infty}} |v| \geq 2/a > 0$). But

$$v(x) \cdot e_r(x) = \left(\frac{1}{a} - \frac{a}{|x|^2} \right) \frac{x_2}{|x|} \neq o\left(\frac{1}{|x|} \right) \text{ as } |x| \rightarrow +\infty,$$

and v is not a circular flow. However, since $u(x) \rightarrow +\infty$ as $|x| \rightarrow +\infty$ and $u = 0$ on C_a and since u has no critical point, it is easily seen that all solutions ξ_x of (1.5) are defined in \mathbb{R} and periodic and that all streamlines $\Xi_x = \xi_x(\mathbb{R})$ (which are level sets of u) surround the origin.¹ Nevertheless, the streamlines do not converge to any family of circles at infinity since a calculation yields $\max_{y \in \Xi_x} |y| - \min_{y \in \Xi_x} |y| = \max_{\mathbb{R}} |\xi_x(\cdot)| - \min_{\mathbb{R}} |\xi_x(\cdot)| \rightarrow a/2 > 0$ as $|x| \rightarrow +\infty$. In this counterexample, one actually has $0 < \limsup_{|x| \rightarrow +\infty} |v(x) \cdot e_r(x)| < +\infty$. Thus, there may be another critical behavior than $o(1/|x|)$ in (1.8) for which the conclusion would still hold, although a different proof would be necessary. The question of the characterization of a critical behavior is left open.

We point out that, in Theorem 1.3, the flow v is not assumed to be bounded. Actually, there are unbounded circular flows satisfying all assumptions of Theorem 1.3: consider for instance the $C^\infty(\overline{\Omega_{a,\infty}})$ unbounded circular flow v defined by

$$v(x) = |x|e_\theta(x),$$

solving (1.1)–(1.2) with stream function $u(x) = |x|^2/2$ and pressure $p(x) = |x|^2/2$, and satisfying $\inf_{\Omega_{a,\infty}} |v| = a > 0$.

Notice lastly that condition (1.7) is fulfilled in particular when $\inf_{\Omega_{a,\infty}} |v| > 0$. Furthermore, as soon as $|v| > 0$ on C_a (which holds if $\inf_{\Omega_{a,\infty}} |v| > 0$), the boundary condition (1.2) and the continuity of v imply in particular that $v \cdot e_\theta$ has a constant strict sign on C_a . Under the condition $\inf_{\Omega_{a,\infty}} |v| > 0$, the following result then provides some estimates on the infimum or the supremum of the vorticity $\frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2}$ in $\Omega_{a,\infty}$, in terms of the sign of $v \cdot e_\theta$ on C_a .

Theorem 1.4. *Assume $0 < a < \infty$ and $b = \infty$. Let v be a $C^2(\overline{\Omega_{a,\infty}})$ flow solving (1.1)–(1.2) and such that $\inf_{\Omega_{a,\infty}} |v| > 0$. If $v \cdot e_\theta > 0$ on C_a (respectively $v \cdot e_\theta < 0$ on C_a), then*

$$\sup_{\Omega_{a,\infty}} \left(\frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2} \right) > 0 \quad (\text{respectively } \inf_{\Omega_{a,\infty}} \left(\frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2} \right) < 0).$$

¹Throughout the paper, we say that a Jordan curve \mathcal{C} surrounds the origin if the bounded connected component of $\mathbb{R}^2 \setminus \mathcal{C}$ contains the origin.

The flow v given by (1.9) is an example of a flow satisfying the assumptions of Theorem 1.4, with $v \cdot e_\theta > 0$ on C_a , and for which the vorticity (namely Δu) is actually equal to the positive constant $8/a^2$ everywhere in $\overline{\Omega_{a,\infty}}$.

Theorem 1.4 can also be viewed as a Liouville-type result. Namely, we show in its proof that if $\inf_{\Omega_{a,\infty}} |v| > 0$ and $v \cdot e_\theta > 0$ on C_a , and if the vorticity is nonpositive everywhere in $\Omega_{a,\infty}$, then v is a circular flow of the type $v = V(|\cdot|)e_\theta$ with $V : [a, +\infty) \rightarrow [\eta, +\infty)$ for some $\eta > 0$. Therefore, the vorticity $\frac{\partial v_2}{\partial x_1}(x) - \frac{\partial v_1}{\partial x_2}(x) = V'(|x|) + V(|x|)/|x|$ cannot be nonpositive everywhere (since otherwise the function $r \mapsto r V(r) (\geq \eta r)$ would be nonincreasing in $[a, +\infty)$, leading to a contradiction).²

Notice that Theorem 1.4 does not hold good if the assumption $\inf_{\Omega_{a,\infty}} |v| > 0$ is dropped. There are actually some circular flows v satisfying (1.1)–(1.2) such that $|v| > 0$ in $\overline{\Omega_{a,\infty}}$ and $v \cdot e_\theta > 0$ on C_a , but $\inf_{\Omega_{a,\infty}} |v| = 0$ and for which the vorticity is negative everywhere. Consider for instance the $C^\infty(\overline{\Omega_{a,\infty}})$ circular flow

$$v(x) = \frac{1}{|x|^2} e_\theta(x),$$

solving (1.1)–(1.2) with stream function $u(x) = -1/|x|$ and pressure $p(x) = -1/(4|x|^2)$: one has $|v| > 0$ in $\overline{\Omega_{a,\infty}}$ and $v \cdot e_\theta > 0$ on C_a , but $\inf_{\Omega_{a,\infty}} |v| = 0$ and $\frac{\partial v_2}{\partial x_1}(x) - \frac{\partial v_1}{\partial x_2}(x) = -1/|x|^3 < 0$ in $\overline{\Omega_{a,\infty}}$.

The case of punctured disks $\Omega_{0,b}$ with $0 < b < \infty$

Theorem 1.5. *Assume $a = 0$ and $0 < b < \infty$. Let v be a $C^2(\overline{\Omega_{0,b}} \setminus \{0\})$ flow solving (1.1)–(1.2) and such that*

$$\{x \in \overline{\Omega_{0,b}} \setminus \{0\} : |v(x)| = 0\} \subsetneq C_b \tag{1.10}$$

and

$$\int_{C_\varepsilon} |v \cdot e_r| \rightarrow 0 \quad \text{as } \varepsilon \xrightarrow{+} 0. \tag{1.11}$$

Then $|v| > 0$ in $\overline{\Omega_{0,b}} \setminus \{0\}$ and v is a circular flow, namely there is a $C^2((0, b])$ function V with constant strict sign such that $v(x) = V(|x|)e_\theta(x)$ for all $x \in \overline{\Omega_{0,b}} \setminus \{0\}$.

Notice that (1.11) is fulfilled in particular if $v(x) \cdot e_r(x) = o(1/|x|)$ as $|x| \xrightarrow{+} 0$. Let us show that without (1.11) the conclusion of Theorem 1.5 does not hold in general. To do so, let us give a counterexample similar to (1.9) above (which was there defined in $\overline{\Omega_{a,\infty}}$). More precisely, consider the $C^\infty(\overline{\Omega_{0,b}} \setminus \{0\})$ function u defined by $u(x) = (|x|/b - b/|x|)x_1/|x|$, that is,

$$u = \left(\frac{r}{b} - \frac{b}{r} \right) \cos \theta$$

in the usual polar coordinates. The function u satisfies $\Delta u = 0$ in $\overline{\Omega_{0,b}} \setminus \{0\}$ with Dirichlet boundary condition $u = 0$ on C_b , and the $C^\infty(\overline{\Omega_{0,b}} \setminus \{0\})$ field $v = \nabla^\perp u$ satisfies (1.1)–

²The same arguments do not lead to any contradiction in the case of bounded annuli $\Omega_{a,b}$ and $\Omega_{0,b}$ with $b < \infty$; see Remark 1.6.

(1.2) with pressure $p = -|v|^2/2$ (and vorticity equal to 0). In polar coordinates, the field v is given by

$$v = \left[\left(\frac{1}{b} + \frac{b}{r^2} \right) \cos \theta \right] e_\theta + \left[\left(\frac{1}{b} - \frac{b}{r^2} \right) \sin \theta \right] e_r. \tag{1.12}$$

It has only two stagnation points in $\overline{\Omega_{0,b}} \setminus \{0\}$ and they both lie on C_b . Hence, (1.10) is fulfilled. But $\int_{C_\varepsilon} |v \cdot e_r| = 4(b/\varepsilon - \varepsilon/b) \rightarrow 0$ as $\varepsilon \xrightarrow{+} 0$, and v is not a circular flow. In this counterexample, one actually has $\int_{C_\varepsilon} |v \cdot e_r| \sim 4b/\varepsilon$ as $\varepsilon \xrightarrow{+} 0$. Thus, there may be another critical behavior than $o(1)$ in condition (1.11) for which the conclusion would still hold, although a different proof would be necessary. The question of characterizing the critical behavior is still open.

Lastly, in Theorem 1.5, the flow v is not assumed to be bounded. Actually, there are unbounded circular flows satisfying all assumptions of Theorem 1.5: consider for instance the $C^\infty(\overline{\Omega_{0,b}} \setminus \{0\})$ unbounded circular flow v defined by

$$v(x) = \frac{1}{|x|} e_\theta(x) \tag{1.13}$$

solving (1.1)–(1.2) with stream function $u(x) = \ln|x|$ and pressure $p(x) = -1/(2|x|^2)$, and satisfying $|v| > 0$ in $\overline{\Omega_{0,b}} \setminus \{0\}$ and then (1.10)–(1.11).

Remark 1.6. A result similar to Theorem 1.4 does not hold in the punctured disk $\Omega_{0,b}$. For instance, the $C^\infty(\overline{\Omega_{0,b}} \setminus \{0\})$ flow (1.13) satisfies (1.1)–(1.2), $v \cdot e_\theta > 0$ on C_b , $\inf_{\Omega_{0,b}} |v| > 0$, but $\frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2} \equiv 0$ in $\Omega_{0,b}$. The same observation holds good in a smooth annulus $\Omega_{a,b}$ with $0 < a < b < \infty$.

Remark 1.7. Let us comment here the similar conditions (1.8) and (1.11). Condition (1.8) implies that $\int_{C_R} |v \cdot e_r| \rightarrow 0$ as $R \rightarrow +\infty$, which would be the dual of (1.11). But the stronger pointwise asymptotic behavior (1.8) is truly used in the proof of Theorem 1.3, and in particular in the proof of Lemma 3.1 below. Replacing (1.8) by the weaker integral condition $\lim_{R \rightarrow +\infty} \int_{C_R} |v \cdot e_r| = 0$ still yields some intermediary results (see the common preliminary results of Section 2 below), but it is an open question to decide whether the conclusion of Theorem 1.3 would still hold with the integral condition instead of the pointwise one.

The case of the punctured plane $\Omega_{0,\infty}$. The last geometric configuration considered in the paper is the punctured plane

$$\Omega_{0,\infty} = \mathbb{R}^2 \setminus \{0\}.$$

Theorem 1.8. *Let v be a $C^2(\Omega_{0,\infty})$ flow solving (1.1) and such that $|v| > 0$ in $\Omega_{0,\infty}$ and $\liminf_{|x| \rightarrow +\infty} |v(x)| > 0$. Assume moreover that*

$$v(x) \cdot e_r(x) = o(1/|x|) \text{ as } |x| \rightarrow +\infty \quad \text{and} \quad \int_{C_\varepsilon} |v \cdot e_r| \rightarrow 0 \text{ as } \varepsilon \xrightarrow{+} 0. \tag{1.14}$$

Then v is a circular flow. Furthermore, there is a $C^2((0, +\infty))$ function V with constant strict sign such that $v(x) = V(|x|)e_\theta(x)$ for all $x \in \Omega_{0,\infty}$.

The conclusion says that, under roughly speaking the absence of stagnation points in the punctured plane and at infinity, and under the same conditions as in Theorems 1.3 and 1.5 on the behavior of the radial component of v at infinity and at the origin, all streamlines are closed and are just concentric circles.

Remark 1.9. Let us mention here other rigidity results for the stationary solutions of (1.1) in various geometrical configurations. The analyticity of the streamlines under a condition of the type $v_1 > 0$ in the unit disk was shown in [19]. The local correspondence between the vorticities of the solutions of (1.1) and the co-adjoint orbits of the vorticities for the nonstationary version of (1.1) in more general annular domains was investigated in [8]. In a previous paper [14] (see also [15]), we considered the case of a two-dimensional strip with bounded section and the case of bounded flows in a half-plane, assuming in both cases that the flows v are tangential on the boundary and that $\inf |v| > 0$; all streamlines are then proved to be lines which are parallel to the boundary of the domain (in other words, the flow is a parallel flow). Compared to [14], the results of the present paper are concerned with different geometrical situations, and the cases of punctured disks or exterior domains involve specific difficulties. We here also include and prove some new comparison results of independent interest for solutions of semilinear elliptic equations in doubly connected domains (see Proposition 1.14 in Section 1.3 below), not to mention the Serrin-type free boundary problems considered in Section 1.2. These types of problems, as well as some methods used here such as the method of moving planes or the Kelvin transform, were not used in [14–16]. Earlier results by Kalisch [18] were concerned with flows in two-dimensional strips under the additional assumption $v \cdot e \neq 0$, where e is the main direction of the strip. In [16], we considered the case of the whole plane \mathbb{R}^2 and we showed that any $C^2(\mathbb{R}^2)$ bounded flow v is still a parallel flow under the condition $\inf_{\mathbb{R}^2} |v| > 0$, with completely different tools based on the study of the growth of the argument of the flow at infinity.

1.2. Serrin-type free boundary problems with overdetermined boundary conditions

The last main results on the solutions of the Euler equations (1.1) are two Serrin-type results in smooth simply or doubly connected bounded domains whose boundaries are free but on which the flow is assumed to satisfy an additional condition.

Theorem 1.10. *Let Ω be a C^2 nonempty simply connected bounded domain of \mathbb{R}^2 . Let $v \in C^2(\overline{\Omega})$ satisfy the Euler equations (1.1) and assume that $v \cdot n = 0$ and $|v|$ is constant on $\partial\Omega$, where n denotes the outward unit normal on $\partial\Omega$. Assume moreover that v has a unique stagnation point in $\overline{\Omega}$. Then, up to translation,*

$$\Omega = B_R$$

for some $R > 0$. Furthermore, the unique stagnation point of v is the center of the disk and v is a circular flow, that is, there is a $C^2([0, R])$ function $V : [0, R] \rightarrow \mathbb{R}$ such that $V \neq 0$ in $(0, R]$, $V(0) = 0$, and $v(x) = V(|x|)e_\theta(x)$ for all $x \in \overline{B_R} \setminus \{0\}$.

In the proof, we will show that the $C^3(\overline{\Omega})$ stream function u defined by (1.4) satisfies a semilinear elliptic equation $\Delta u + f(u) = 0$ in $\overline{\Omega}$. Furthermore, up to normalization, the function u vanishes on $\partial\Omega$ and is positive in Ω . Lastly, since $|v|$ is assumed to be constant along $\partial\Omega$, the normal derivative $\frac{\partial u}{\partial n}$ of u along $\partial\Omega$ is constant. This problem is therefore an elliptic equation with overdetermined boundary conditions. Since the celebrated paper by Serrin [25], it has been known that these overdetermined boundary conditions on $\partial\Omega$ determine the geometry of Ω , namely, Ω is then a ball and the function u is radially symmetric (hence, here, v would then be a circular flow). The proof is based on the method of moving planes developed in [3, 6, 11, 25] and on the maximum principle, and it relies on the Lipschitz continuity of the function f . In our case, the function f is given in terms of the function u itself and it is continuous in $[0, \max_{\overline{\Omega}} u]$, as will be seen in the proof of Theorem 1.10. But it can be non-Lipschitz-continuous on the whole range $[0, \max_{\overline{\Omega}} u]$. More precisely, it can be non-Lipschitz-continuous in any left neighborhood of the maximum value $\max_{\overline{\Omega}} u$.³ One therefore has to adapt the proof to this case by removing small neighborhoods of size ε around the maximum point of u (which is the unique stagnation point of v); one shows the symmetry of the domain in all directions up to ε and one concludes by passing to the limit as $\varepsilon \searrow 0$.

Remark 1.11. Other free boundary problems related to the Euler equations have been considered by Gómez-Serrano, Park, Shi and Yao [13]. They proved that if a solution v of the Euler equations (1.1) in \mathbb{R}^2 has a vorticity which is the indicator function of a bounded set (a patch) and if v is tangential on the boundary of this set, then v is circular (up to translation) and the patch is a disk (see also [10, Chapter 4] for an earlier result when the patch is assumed to be simply connected). It was also shown in [13] that smooth solutions of (1.1) in \mathbb{R}^2 with nonnegative compactly supported vorticity must be radially symmetric (up to translation). Other rigidity results of [13, 17] also deal with nonstationary uniformly rotating solutions.

In connection with Theorems 1.5 and 1.10, we state the following conjecture.

Conjecture 1.12. *Let D be an open nonempty disk and let $z \in D$. Let v be a $C^2(\overline{D} \setminus \{z\})$ and bounded flow solving (1.1) and $v \cdot n = 0$ on ∂D , where n denotes the outward unit normal on ∂D . Assume that $|v| > 0$ in $\overline{D} \setminus \{z\}$. Then z is the center of the disk and the flow is circular with respect to z .*

Up to translation, one can assume that $D = B_b$ for some $b \in (0, +\infty)$, hence $n = e_r$ on ∂D . If the point z is a priori assumed to be the center of the disk, i.e. the origin, then Theorem 1.5 implies that v is a circular flow. Up to rotation, assume now that $z = (\alpha, 0)$

³For instance, for any $R > 0$, the $C^\infty(\overline{B_R})$ flow $v(x) = -4|x|^2 x^\perp$ satisfies (1.1) in $\overline{B_R}$ with $v \cdot e_r = 0$ on C_R , and with pressure $p(x) = (8/3)|x|^6$ (up to an additive constant). Furthermore, $|v|$ is constant on C_R and the only stagnation point of v in $\overline{B_R}$ is the center of the disk. The stream function $u(x) = R^4 - |x|^4$ (up to an additive constant) satisfies the elliptic equation $\Delta u + f(u) = 0$ in $\overline{B_R}$ with $f(s) = 16\sqrt{R^4 - s}$, and the function f is not Lipschitz continuous in any left neighborhood of $\max_{\overline{B_R}} u = R^4$.

for some $\alpha \in (0, b)$ and, without loss of generality, that the stream function u is positive in $D \setminus \{z\}$ and vanishes on ∂D . The goal would be to reach a contradiction. As far as Theorem 1.10 is concerned, the method of proof described in the paragraph following the statement shows simultaneously the symmetry of the domain and the symmetry of the function u (which obeys an equation of the type $\Delta u + f(u) = 0$), thanks to the overdetermined boundary conditions satisfied by u . Here in Conjecture 1.12, the same techniques based on the method of moving planes implies for instance on the one hand that the function u is even in x_2 in $\overline{\Omega_{0,b}} \setminus \{z\}$, and on the other hand that $u(x_1, x_2) < u(2\alpha - x_1, x_2)$ for all $(x_1, x_2) \in \overline{\Omega_{0,b}}$ such that $x_1 > \alpha$. But, regarding the second property, the Hopf lemma might not apply to the function $(x_1, x_2) \mapsto u(x_1, x_2) - u(2\alpha - x_1, x_2)$ at the point $z = (\alpha, 0)$ since the vorticity function f might not be Lipschitz continuous around the limiting value of u at z (see also the comments after the proof of Theorems 1.1 and 1.2 in Section 3.1 below, and notice that u is not differentiable at z , unless one further assumes that $|v(x)| \rightarrow 0$ as $x \rightarrow z$). Therefore, the arguments as in the proof of Theorem 1.10 do not lead to an obvious contradiction if z is not the center of the disk. However, Conjecture 1.12 seems natural and will be the purpose of further investigation.

A related weaker conjecture (with stronger assumptions) can also be formulated: if D is an open nonempty disk and $z \in D$, and if $v \in C^2(\overline{D})$ solves (1.1) with $v \cdot n = 0$ on ∂D and z is the only stagnation point of v in \overline{D} , then z is the center of the disk and v is circular with respect to z . For the same reasons as in the previous paragraph (since the vorticity function f might not be Lipschitz continuous around $u(z)$), the proof of that second conjecture is not clear either.

The last main result related to the Euler equations is concerned with the case of doubly connected bounded domains.

Theorem 1.13. *Let ω_1 and ω_2 be two C^2 nonempty simply connected bounded domains of \mathbb{R}^2 such that $\overline{\omega_1} \subset \omega_2$, and denote*

$$\Omega = \omega_2 \setminus \overline{\omega_1}.$$

Let $v \in C^2(\overline{\Omega})$ satisfy the Euler equations (1.1). Assume that $v \cdot n = 0$ on $\partial\Omega = \partial\omega_1 \cup \partial\omega_2$, where n denotes the outward unit normal on $\partial\Omega$, and that $|v|$ is constant on $\partial\omega_1$ and on $\partial\omega_2$. Assume moreover that $|v| > 0$ in $\overline{\Omega}$. Then ω_1 and ω_2 are two concentric disks and, up to translation,

$$\Omega = \Omega_{a,b}$$

for some $0 < a < b < \infty$ and v is a circular flow satisfying the conclusion of Theorem 1.1 in $\overline{\Omega}$.

In this case, by using the arguments of Section 2 below (which also lead to the proof of Theorems 1.1 and 1.2 in $\Omega_{a,b}$ with $0 < a < b < \infty$), it follows that the stream function u of the flow v satisfies a semilinear elliptic equation $\Delta u + f(u) = 0$ in $\overline{\Omega}$, with $u = c_1$ on $\partial\omega_1$ and $u = c_2$ on $\partial\omega_2$, for some real numbers $c_1 \neq c_2$. Furthermore, $\min(c_1, c_2) < u < \max(c_1, c_2)$ in Ω and the normal derivative $\frac{\partial u}{\partial n}$ is constant along $\partial\omega_1$ and along $\partial\omega_2$. Since v has no stagnation point in $\overline{\Omega}$, the function f is then shown to be Lipschitz continuous

in the whole interval $[\min(c_1, c_2), \max(c_1, c_2)]$, and known results of Reichel [21] and Sirakov [27] imply that, up to translation, $\Omega = \Omega_{a,b}$ for some $0 < a < b < \infty$, and u is radially symmetric.

Further symmetry results have been obtained for nonlinear elliptic equations of the type $\Delta u + f(u) = 0$ or more general ones in exterior domains with overdetermined boundary conditions (see e.g. [1, 22, 27]), or in the whole space (see e.g. [12, 20, 26]), in both cases with further assumptions on the solution u at infinity and on the function f . Such conditions are in general not satisfied by the stream function u and the vorticity function f of a flow v that would be defined in the complement of a simply connected bounded domain or in the whole or punctured plane. Lastly, we refer to [5, 9, 23, 24] for further references on overdetermined boundary value elliptic problems in domains with more complex topology or in unbounded epigraphs.

1.3. *Directional comparison results for semilinear elliptic equations in some doubly connected domains*

As briefly mentioned after Theorems 1.10 and 1.13, the main strategy of these results, as well as the other ones in the fixed annular domains $\Omega_{a,b}$, is to show that the stream function u satisfies a semilinear elliptic equation $\Delta u + f(u) = 0$ in the domain under consideration and that this stream function is then radially symmetric (and the domain itself is circular if the boundary is free). The radial symmetry of the stream function u means that the flow v is circular. The proof of the radial symmetry of u follows from its even symmetry and monotonicity with respect to each direction. The proof of the symmetry and monotonicity relies on some directional comparison results which are themselves based on the method of moving planes in doubly connected domains trapped between two level sets of u . We point out that these level sets are not known a priori to be circles and their precise shape is not known. That is why we have to show in this framework a key proposition containing some new directional comparison results, which we think are of independent interest and which we state and will use for more general semilinear heterogeneous elliptic equations. Proposition 1.14 below will be used as a key step in the proof of Theorems 1.3–1.5, 1.8 and 1.10. We also point out that the heterogeneity in (1.17) below is not an artifice, since we will truly deal with heterogeneous equations obtained after a Kelvin transform of some original equations set in exterior domains.

To do so, let us first introduce a few notations. For $e \in \mathbb{S}^1 = C_1$ and $\lambda \in \mathbb{R}$, we denote

$$T_{e,\lambda} = \{x \in \mathbb{R}^2 : x \cdot e = \lambda\}, \quad H_{e,\lambda} = \{x \in \mathbb{R}^2 : x \cdot e > \lambda\}, \quad (1.15)$$

and, for $x \in \mathbb{R}^2$,

$$R_{e,\lambda}(x) = x_{e,\lambda} = x - 2(x \cdot e - \lambda)e. \quad (1.16)$$

In other words, $R_{e,\lambda}$ is the orthogonal reflection with respect to the line $T_{e,\lambda}$.

Proposition 1.14. *Let Ξ and Ξ' be two C^1 Jordan curves surrounding the origin, and let Ω and Ω' be the bounded connected components of $\mathbb{R}^2 \setminus \Xi$ and $\mathbb{R}^2 \setminus \Xi'$, respectively.*




Fig. 1. The sets Ω , Ω' , $\omega = \Omega \setminus \overline{\Omega'}$, $H_{e,\lambda} \cap \overline{\Omega}$ (light blue background), $R_{e,\lambda}(\Omega')$ (with dashed boundary), and $\omega_{e,\lambda}$ (dashed red).

Assume that $\overline{\Omega'} \subset \Omega$ and let

$$\omega = \Omega \setminus \overline{\Omega'}$$

be the nonempty and doubly connected domain located between Ξ and Ξ' , with boundary

$$\partial\omega = \Xi \cup \Xi'.$$

Write $R' = \min_{x \in \Xi'} |x| > 0$ and $R = \max_{x \in \Xi} |x| > R'$. Let $e \in \mathbb{S}^1$, let $\bar{\lambda} = \max_{x \in \Xi} x \cdot e > 0$ and let $\varepsilon \in [0, \bar{\lambda})$. Let $c_1 < c_2 \in \mathbb{R}$ and let $\varphi \in C^2(\overline{\omega})$ be a solution of

$$\begin{cases} \Delta\varphi + F(|x|, \varphi) = 0 & \text{in } \overline{\omega}, \\ c_1 < \varphi < c_2 & \text{in } \omega, \\ \varphi = c_1 \text{ on } \Xi, \quad \varphi = c_2 \text{ on } \Xi', \end{cases} \tag{1.17}$$

with a continuous function $F : [R', R] \times [c_1, c_2] \rightarrow \mathbb{R}$ that is nonincreasing with respect to its first variable and uniformly Lipschitz continuous with respect to its second variable. Assume that

$$R_{e,\lambda}(H_{e,\lambda} \cap \overline{\Omega}) \subset \Omega \quad \text{for all } \lambda > \varepsilon, \tag{1.18}$$

$$R_{e,\lambda}(H_{e,\lambda} \cap \Xi') \subset \Omega' \quad \text{for all } \lambda > \varepsilon \tag{1.19}$$

(see Fig. 1). Then, for every $\lambda \in [\varepsilon, \bar{\lambda})$,

$$\varphi(x) \leq \varphi_{e,\lambda}(x) = \varphi(x_{e,\lambda}) \quad \text{for all } x \in \overline{\omega_{e,\lambda}} \tag{1.20}$$

with

$$\omega_{e,\lambda} = (H_{e,\lambda} \cap \omega) \setminus R_{e,\lambda}(\overline{\Omega'}).^4$$

Organization of the paper. In Section 2, we give some common preliminaries for the proofs of Theorems 1.1–1.5, 1.8, 1.10 and 1.13 in fixed or free domains. Namely, we study the properties of the streamlines of the flow and we derive a semilinear elliptic equation

⁴Notice that $\omega_{e,\lambda}$ is open by definition, and it is nonempty for each $\lambda \in [0, \bar{\lambda})$: indeed, for such λ , the set $T_{e,\lambda} \cap \Xi$ is not empty and, for any $x \in T_{e,\lambda} \cap \Xi$ and $r > 0$, $\omega_{e,\lambda} \cap B(x, r) \neq \emptyset$. But $\omega_{e,\lambda}$ may not be connected, as can be seen in Fig. 1.

for the stream function. Section 3 is devoted to the proofs of Theorems 1.1–1.5 and 1.8, when the circular domain $\Omega_{a,b}$ is fixed. The cases of the exterior domains $\Omega_{a,\infty}$ and the punctured disks $\Omega_{0,b}$ and punctured plane $\Omega_{0,\infty}$ involve some additional difficulties and require specific additional assumptions. The proof of the Serrin-type Theorems 1.10 and 1.13 is carried out in Section 4. Lastly, Section 5 is concerned with the proof of Proposition 1.14. Proposition 1.14 and its subsequent limiting argument showing that the flow considered is circular are necessary for the proofs of Theorems 1.3–1.5 and 1.8 in the case of exterior or punctured domains, as well as for the proof of Theorem 1.10 about simply connected domains with free boundary.

2. Some common preliminaries

In this section, we state and prove some common properties which will be used in the proofs of the main results on the Euler equations (1.1) in a fixed annular domain $\Omega_{a,b}$ or in simply or doubly connected domains with free boundaries. To cover all possible cases, throughout this section we consider two C^1 Jordan curves \mathcal{C}_1 and \mathcal{C}_2 and we assume that they both surround the origin, that is, the bounded connected components ω_1 and ω_2 of $\mathbb{R}^2 \setminus \mathcal{C}_1$ and $\mathbb{R}^2 \setminus \mathcal{C}_2$ contain the origin 0. We further assume that

$$\overline{\omega_1} \subset \omega_2.$$

Our aim here is to study the properties of the stream function u and the streamlines of a divergence-free flow v in the open connected set Ω in one of the following four possible cases:

$$\Omega = \omega_2 \setminus \overline{\omega_1}, \quad \text{or} \quad \Omega = \mathbb{R}^2 \setminus \overline{\omega_1}, \quad \text{or} \quad \Omega = \omega_2 \setminus \{0\}, \quad \text{or} \quad \Omega = \mathbb{R}^2 \setminus \{0\}.$$

Notice that Ω is bounded if $\Omega = \omega_2 \setminus \overline{\omega_1}$ or $\omega_2 \setminus \{0\}$, and unbounded in the other two cases. We say that Ω is *not punctured* if $\Omega = \omega_2 \setminus \overline{\omega_1}$ or $\mathbb{R}^2 \setminus \overline{\omega_1}$, and *punctured* in the other two cases. We point out that the annular domains $\Omega_{a,b}$ with $0 \leq a < b \leq \infty$ fall within these four types of domains Ω . We also set

$$\begin{cases} D = \overline{\Omega} & \text{if } \Omega \text{ is not punctured,} \\ D = \overline{\Omega} \setminus \{0\} & \text{if } \Omega \text{ is punctured.} \end{cases} \tag{2.1}$$

Throughout this section, $v = (v_1, v_2)$ is a $C^1(D)$ vector field. We point out that the $C^2(D)$ regularity of v will be specified and used only in Lemma 2.8 below, as will also be the Euler equations $v \cdot \nabla v + \nabla p = 0$ themselves. We also point out that if Ω is punctured, then v is not assumed to be defined at 0. It is however always assumed that

$$\operatorname{div} v = 0 \quad \text{in } D, \quad |v| > 0 \quad \text{in } \Omega, \tag{2.2}$$

and

$$\begin{cases} v \cdot n = 0 \text{ on } \partial\omega_1 & \text{if } \Omega \text{ is not punctured,} \\ \int_{\mathcal{C}_\varepsilon} |v \cdot e_r| \rightarrow 0 \text{ as } \varepsilon \xrightarrow{\geq} 0 & \text{if } \Omega \text{ is punctured,} \end{cases} \tag{2.3}$$

where n denotes the outward unit normal to Ω (that is, on \mathcal{C}_1 and/or \mathcal{C}_2 as appropriate). The assumptions (2.2)–(2.3) are in force throughout Section 2, and are therefore not repeated in the statements.

The first common result is concerned with the existence and some elementary properties of the stream function u .

Lemma 2.1. (i) *There is a unique (up to additive constants) $C^2(D)$ scalar function u such that*

$$\nabla^\perp u = v \quad \text{in } D,$$

and there is a constant c_{in} such that

$$\begin{cases} u = c_{\text{in}} \in \mathbb{R} \text{ on } \partial\omega_1 & \text{if } \Omega \text{ is not punctured,} \\ u(x) \rightarrow c_{\text{in}} \in \overline{\mathbb{R}} \text{ as } |x| \xrightarrow{\geq} 0 & \text{if } \Omega \text{ is punctured.} \end{cases} \quad (2.4)$$

(ii) *If one further assumes that*

$$\begin{cases} v \cdot n = 0 \text{ on } \partial\omega_2 & \text{if } \Omega \text{ is bounded,} \\ \int_{C_R} |v \cdot e_r| \rightarrow 0 \text{ as } R \rightarrow +\infty & \text{if } \Omega \text{ is unbounded,} \end{cases} \quad (2.5)$$

then there is a constant c_{out} such that

$$\begin{cases} u = c_{\text{out}} \in \mathbb{R} \text{ on } \partial\omega_2 & \text{if } \Omega \text{ is bounded,} \\ u(x) \rightarrow c_{\text{out}} \in \overline{\mathbb{R}} \text{ as } |x| \rightarrow +\infty & \text{if } \Omega \text{ is unbounded.} \end{cases} \quad (2.6)$$

Furthermore,

$$c_{\text{in}} \neq c_{\text{out}} \quad \text{and} \quad \min(c_{\text{in}}, c_{\text{out}}) < u < \max(c_{\text{in}}, c_{\text{out}}) \quad \text{in } \Omega.^5 \quad (2.7)$$

Proof. (i) The existence and uniqueness (up to additive constants) of a stream function $u \in C^2(D)$ is a consequence of the fact that Ω is doubly connected, v is divergence free and satisfies conditions (2.3) on the inner boundary of Ω .

Assume now that Ω is not punctured, and consider a parametrization of its inner boundary \mathcal{C}_1 by a $C^1(\mathbb{R})$ periodic function ζ_1 such that $|\zeta_1'| > 0$ in \mathbb{R} . From the first condition in (2.3), the function $u \circ \zeta_1$ is constant in \mathbb{R} , that is, there is $c_{\text{in}} \in \mathbb{R}$ such that $u = c_{\text{in}}$ on $\mathcal{C}_1 = \partial\omega_1$.

Assume now that Ω is punctured, and let $\rho > 0$ be such that $\overline{B_r} \setminus \{0\} \subset \Omega$ for all $r \in (0, \rho)$. Since u has no critical point in Ω (because $|v| > 0$ in Ω by (2.2)), it follows that

$$u > \min \left(\min_{C_{r_1}} u, \min_{C_{r_3}} u \right) \quad \text{in } \Omega_{r_1, r_3}, \text{ for all } 0 < r_1 < r_3 < \rho.$$

⁵The subscript “in” in c_{in} refers to the “inner” boundary of Ω , which is $\partial\omega_1$ (resp. $\{0\}$) if Ω is not punctured (resp. punctured). Similarly, the subscript “out” in c_{out} refers to the “outer” boundary of Ω , which is $\partial\omega_2$ (resp. the infinity) if Ω is bounded (resp. unbounded).

In particular, $\min_{C_{r_2}} u > \min(\min_{C_{r_1}} u, \min_{C_{r_3}} u)$ for all $0 < r_1 < r_2 < r_3 < \rho$, hence the function $r \mapsto \min_{C_r} u$ is strictly monotone in a right neighborhood of 0 and has a limit $c_{in} \in \overline{\mathbb{R}}$ at 0. On the other hand, the integral condition in (2.3) means that $\int_{C_r} |\nabla u \cdot e_\theta| \rightarrow 0$ as $r \xrightarrow{\geq} 0$, hence the oscillation of u on C_r converges to 0 as $r \xrightarrow{\geq} 0$, that is,

$$\text{osc}_{C_r} u := \max_{C_r} u - \min_{C_r} u \rightarrow 0 \quad \text{as } r \xrightarrow{\geq} 0. \tag{2.8}$$

Finally, $\max_{C_r} u \rightarrow c_{in}$ as $r \xrightarrow{\geq} 0$, and $u(x) \rightarrow c_{in}$ as $|x| \xrightarrow{\geq} 0$.

(ii) Similarly, conditions (2.5), together with (2.2), imply the existence of $c_{out} \in \overline{\mathbb{R}}$ such that (2.6) holds in both the bounded and unbounded cases. In the unbounded case, the integral condition in (2.5) also implies, as in (2.8) above, that

$$\max_{C_R} u - \min_{C_R} u \rightarrow 0 \quad \text{as } R \rightarrow +\infty. \tag{2.9}$$

Finally, in all cases, since u has no critical point in Ω , the properties (2.4) and (2.6) immediately yield (2.7). ■

The second common result is concerned with the trajectories of the gradient flow ∇u , with u given by Lemma 2.1. Namely, for $x \in D$, with D as in (2.1), let σ_x be the solution of

$$\begin{cases} \dot{\sigma}_x(t) = \nabla u(\sigma_x(t)), \\ \sigma_x(0) = x. \end{cases} \tag{2.10}$$

For $y \in \mathbb{R}^2$ and $E \subset \mathbb{R}^2$, we set

$$\text{dist}(y, E) = \inf_{z \in E} |y - z|.$$

Lemma 2.2. (i) *If $x \in \Omega$, then there exist $-\infty \leq t_x^- < 0 < t_x^+ \leq +\infty$ such that $\sigma_x : (t_x^-, t_x^+) \rightarrow \Omega$ is of class C^1 with $(u \circ \sigma_x)' > 0$ in (t_x^-, t_x^+) and*

$$\begin{cases} \text{dist}(\sigma_x(t), \partial\Omega) \rightarrow 0 & \text{as } t \rightarrow t_x^\pm \text{ if } \Omega \text{ is bounded,} \\ \text{dist}(\sigma_x(t), \partial\Omega) \rightarrow 0 \text{ or } |\sigma_x(t)| \rightarrow +\infty & \text{as } t \rightarrow t_x^\pm \text{ if } \Omega \text{ is unbounded.} \end{cases} \tag{2.11}$$

(ii) *If Ω is not punctured and $x \in \partial\omega_1$, then either $\sigma_x(t) = x$ for all $t \in \mathbb{R}$, or there is $t_x^+ \in (0, +\infty]$ such that $\sigma_x : [0, t_x^+) \rightarrow \Omega \cup \partial\omega_1$ is of class C^1 with $(u \circ \sigma_x)' > 0$ in $[0, t_x^+)$, $\sigma_x((0, t_x^+)) \subset \Omega$, and*

$$\begin{aligned} \text{dist}(\sigma_x(t), \partial\omega_2) &\xrightarrow[t \rightarrow t_x^+]{} 0 \text{ if } \Omega \text{ is bounded} \\ &\text{(resp. } |\sigma_x(t)| \xrightarrow[t \rightarrow t_x^+]{} +\infty \text{ if } \Omega \text{ is unbounded),} \end{aligned} \tag{2.12}$$

or there is $t_x^- \in [-\infty, 0)$ such that $\sigma_x : (t_x^-, 0] \rightarrow \Omega \cup \partial\omega_1$ is of class C^1 with $(u \circ \sigma_x)' > 0$ in $(t_x^-, 0]$, $\sigma_x((t_x^-, 0)) \subset \Omega$, and (2.12) holds with t_x^- instead of t_x^+ .

(iii) If Ω is bounded, if (2.5) holds and if $x \in \partial\omega_2$, then either $\sigma_x(t) = x$ for all $t \in \mathbb{R}$, or there is $t_x^+ \in (0, +\infty)$ such that $\sigma_x : [0, t_x^+) \rightarrow \Omega \cup \partial\omega_2$ is of class C^1 with $(u \circ \sigma_x)' > 0$ in $[0, t_x^+)$, $\sigma_x((0, t_x^+)) \subset \Omega$, and

$$\begin{aligned} \text{dist}(\sigma_x(t), \partial\omega_1) &\xrightarrow[t \rightarrow t_x^+]{\quad} 0 \text{ if } \Omega \text{ is not punctured} \\ &\text{(resp. } |\sigma_x(t)| \xrightarrow[t \rightarrow t_x^+]{\quad} 0 \text{ if } \Omega \text{ is punctured),} \end{aligned} \quad (2.13)$$

or there is $t_x^- \in [-\infty, 0)$ such that $\sigma_x : (t_x^-, 0] \rightarrow \Omega \cup \partial\omega_2$ is of class C^1 with $(u \circ \sigma_x)' > 0$ in $(t_x^-, 0]$, $\sigma_x((t_x^-, 0)) \subset \Omega$, and (2.13) holds with t_x^- instead of t_x^+ .

Proof. (i) Consider any x in Ω . Since ∇u is of class $C^1(D)$, the solution σ_x of (2.10) is defined in a neighborhood of 0 and the quantities t_x^\pm defined by

$$\begin{cases} t_x^+ = \sup \{t > 0 : \sigma_x((0, t)) \subset \Omega\}, \\ t_x^- = \inf \{t < 0 : \sigma_x((t, 0)) \subset \Omega\} \end{cases} \quad (2.14)$$

are such that $-\infty \leq t_x^- < 0 < t_x^+ \leq +\infty$. The functions $\sigma_x : (t_x^-, t_x^+) \rightarrow \Omega$ and $u \circ \sigma_x : (t_x^-, t_x^+) \rightarrow \mathbb{R}$ are of class C^1 , with

$$(u \circ \sigma_x)'(t) = |\nabla u(\sigma_x(t))|^2 = |v(\sigma_x(t))|^2 > 0 \quad \text{for all } t \in (t_x^-, t_x^+),$$

since $|v| > 0$ in Ω by (2.2). Let us then show (2.11) as $t \rightarrow t_x^+$ (the limit as $t \rightarrow t_x^-$ can be treated similarly). Let us assume by way of contradiction that (2.11) (with $t \rightarrow t_x^+$) does not hold. So, in all configurations, there exist an increasing sequence $(t_n)_{n \in \mathbb{N}}$ in (t_x^-, t_x^+) converging to t_x^+ and a point $y \in \Omega$ such that $\sigma_x(t_n) \rightarrow y$ as $n \rightarrow +\infty$. Since the continuous field $|\nabla u| = |v|$ does not vanish in Ω by (2.2), there are real numbers $r > 0$, $\eta > 0$ and $\tau > 0$ such that

$$\begin{cases} \overline{B(y, r)} \subset \Omega, & |\nabla u| \geq \eta \text{ in } \overline{B(y, r)}, \\ \sigma_z(t) \in \overline{B(y, r)} & \text{for all } z \in \overline{B(y, r/2)} \text{ and } t \in [-\tau, \tau]. \end{cases}$$

Since $\sigma_x(t_n) \rightarrow y$ as $n \rightarrow +\infty$, one has $\sigma_x(t_n) \in \overline{B(y, r/2)}$ for all n large enough, hence σ_x is defined in $[t_n - \tau, t_n + \tau]$ with $\sigma_x(t) \in \overline{B(y, r)} \subset \Omega$ for all $t \in [t_n - \tau, t_n + \tau]$ and n large enough. This implies that $t_x^+ = +\infty$. Furthermore, for all n large enough, one has

$$(u \circ \sigma_x)'(t) = |\nabla u(\sigma_x(t))|^2 \geq \eta^2 \quad \text{for all } t \in [t_n - \tau, t_n + \tau],$$

hence

$$u(\sigma_x(t_n + \tau)) \geq u(\sigma_x(t_n - \tau)) + 2\eta^2\tau.$$

Since $u \circ \sigma_x$ is increasing in (t_x^-, t_x^+) and since $t_n \rightarrow t_x^+ = +\infty$ as $n \rightarrow +\infty$, one then gets $u(\sigma_x(t)) \rightarrow +\infty$ as $t \rightarrow t_x^+ = +\infty$, contradicting the fact that $\sigma_x(t_n) \rightarrow y$, $t_n \rightarrow t_x^+$ and the continuity of u at y . Therefore, (2.11) has been proved.

(ii) Assume now that Ω is not punctured and consider $x \in \partial\omega_1$. From (2.3), either $|\nabla u(x)| = 0$ (and then $\sigma_x(t) = x$ for all $t \in \mathbb{R}$), or $|\nabla u(x)| > 0$ and

$$\nabla u(x) = \pm |\nabla u(x)|n(x),$$

with $n(x)$ the outward normal vector to Ω at x . If $\nabla u(x) = -|\nabla u(x)|n(x)$ (resp. $\nabla u(x) = |\nabla u(x)|n(x)$) with $|\nabla u(x)| > 0$, then $\nabla u(x)$ points into (resp. outward from) Ω at x and the quantity t_x^+ (resp. t_x^-) given in (2.14) is well defined and satisfies $0 < t_x^+ \leq +\infty$ together with $\sigma_x((0, t_x^+)) \subset \Omega$ and $\sigma_x([0, t_x^+)) \subset \Omega \cup \partial\omega_1$ (resp. $-\infty \leq t_x^- < 0$, $\sigma_x((t_x^-, 0)) \subset \Omega$, and $\sigma_x((t_x^-, 0]) \subset \Omega \cup \partial\omega_1$). Furthermore, σ_x and $u \circ \sigma_x$ are of class $C^1([0, t_x^+))$ (resp. $C^1((t_x^-, 0])$) and $(u \circ \sigma_x)' > 0$ in $[0, t_x^+)$ (resp. in $(t_x^-, 0]$). Lastly, if $\nabla u(x) = -|\nabla u(x)|n(x)$ with $|\nabla u(x)| > 0$, then, as in (i), (2.11) still holds with $t \rightarrow t_x^+$, and since the continuous function u is equal to the constant $c_{in} \in \mathbb{R}$ on $\partial\omega_1$ by Lemma 2.1 (i), one gets (2.12). If $\nabla u(x) = |\nabla u(x)|n(x)$ with $|\nabla u(x)| > 0$, one similarly gets (2.12) with t_x^- instead of t_x^+ .

(iii) If Ω is bounded and (2.5) holds, and if $x \in \partial\omega_2$, then either $|\nabla u(x)| = 0$ and $\sigma_x(t) \equiv x$ in \mathbb{R} , or σ_x is defined in $(t_x^-, 0]$ or $[0, t_x^+)$ and ranges in $\Omega \cup \partial\omega_2$, and the conclusion follows as in (ii). ■

Remark 2.3. The extremal values t_x^\pm in Lemma 2.2 can be finite or infinite. Consider for instance the case $\Omega = \Omega_{a,b}$ with $0 \leq a < b \leq \infty$ and a circular flow $v(x) = V(|x|)e_\theta(x)$ with a $C^1((a, b))$ positive scalar function V (the function V can be assumed to be extended in a C^1 fashion at a and b when they are positive real numbers, and therefore v is of class $C^1(D)$ with D as in (2.1)). Assumptions (2.2)–(2.3) and (2.5) are fulfilled, and the stream function u is given by $u(x) = U(|x|)$ in D , with $U' = V$. For any $x \in \Omega_{a,b}$, the solution σ_x of (2.10) can then be written as

$$\sigma_x(t) = \zeta_x(t)e_r(x) \quad \text{with } \zeta_x'(t) = V(\zeta_x(t)) \text{ for all } t \in (t_x^-, t_x^+).$$

Therefore, the finiteness of t_x^+ (resp. of t_x^-) is equivalent to the integrability of the function $1/V$ at b (resp. at a).

The third common result provides the existence of a C^1 curve in D connecting the inner and outer boundaries of Ω , along which u is strictly monotone, and which will be as in Lemma 2.2 a trajectory of the gradient flow.

Lemma 2.4.

$$S = \{x \in D : |v(x)| = 0\}$$

the set of stagnation points of v in D , defined by (2.1) (notice that $S \subset \partial\Omega \cap D$ by (2.2)).

(i) Assume that

$$\begin{cases} S \subsetneq \mathcal{C}_1 \text{ or } S \subsetneq \mathcal{C}_2, \text{ and } v \cdot n = 0 \text{ on } \mathcal{C}_2 & \text{if } \Omega = \omega_2 \setminus \overline{\omega_1}, \\ S \subsetneq \mathcal{C}_1 & \text{if } \Omega = \mathbb{R}^2 \setminus \overline{\omega_1}, \\ S \subsetneq \mathcal{C}_2 \text{ and } v \cdot n = 0 \text{ on } \mathcal{C}_2 & \text{if } \Omega = \omega_2 \setminus \{0\}, \\ \int_{C_R} |v \cdot e_r| \rightarrow 0 \text{ as } R \rightarrow +\infty & \text{if } \Omega = \mathbb{R}^2 \setminus \{0\}. \end{cases} \tag{2.15}$$

Then there exist $-\infty \leq t_{\text{in}} < t_{\text{out}} \leq +\infty$, an interval $I \subset \mathbb{R}$ and a C^1 function $\sigma : I \rightarrow D$ such that $(u \circ \sigma)' \neq 0$ in I and

$$\begin{cases} I = [t_{\text{in}}, t_{\text{out}}], \sigma(t_{\text{in}}) \in \partial\omega_1, \sigma(t_{\text{out}}) \in \partial\omega_2 & \text{if } \Omega = \omega_2 \setminus \overline{\omega_1}, \\ I = [t_{\text{in}}, t_{\text{out}}), \sigma(t_{\text{in}}) \in \partial\omega_1, |\sigma(t)| \rightarrow +\infty \text{ as } t \nearrow t_{\text{out}} & \text{if } \Omega = \mathbb{R}^2 \setminus \overline{\omega_1}, \\ I = (t_{\text{in}}, t_{\text{out}}], |\sigma(t)| \rightarrow 0 \text{ as } t \searrow t_{\text{in}}, \sigma(t_{\text{out}}) \in \partial\omega_2 & \text{if } \Omega = \omega_2 \setminus \{0\}, \\ I = (t_{\text{in}}, t_{\text{out}}), |\sigma(t)| \rightarrow 0 \text{ as } t \searrow t_{\text{in}}, |\sigma(t)| \rightarrow +\infty \text{ as } t \nearrow t_{\text{out}} & \text{if } \Omega = \mathbb{R}^2 \setminus \{0\}. \end{cases} \tag{2.16}$$

(ii) If $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1}$ together with

$$S \not\subset \mathcal{C}_1 \quad \text{and} \quad \liminf_{|x| \rightarrow +\infty} |v(x)| > 0, \tag{2.17}$$

then there exist $-\infty < t_{\text{in}} < t_{\text{out}} \leq +\infty$ and a C^1 function $\sigma : [t_{\text{in}}, t_{\text{out}}) \rightarrow D = \overline{\Omega}$ such that $(u \circ \sigma)' \neq 0$ in $[t_{\text{in}}, t_{\text{out}})$ and

$$\sigma(t_{\text{in}}) \in \partial\omega_1, \quad |\sigma(t)| \rightarrow +\infty \text{ as } t \nearrow t_{\text{out}}, \quad \text{and} \quad |u(\sigma(t))| \rightarrow +\infty \text{ as } t \nearrow t_{\text{out}}.$$

Furthermore, one can take $t_{\text{in}} = 0$ without loss of generality.

Remark 2.5. Notice that condition (2.15) implies (2.5) in all possible configurations of Ω except when $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1}$: in that case, in (2.15) we do not assume that $\lim_{R \rightarrow +\infty} \int_{C_R} |v \cdot e_r| = 0$, whereas in (2.5) we do. We also point out that, again if $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1}$, condition (2.17) implies (2.15), but still does not imply (2.5).

Proof of Lemma 2.4. (i) Here we assume (2.15). Consider first the case $\Omega = \omega_2 \setminus \overline{\omega_1}$, and assume that $S \not\subset \mathcal{C}_1$ in (2.15) (the case $S \not\subset \mathcal{C}_2$ can be handled similarly). Then $|v| = |\nabla u| > 0$ on $\mathcal{C}_2 = \partial\omega_2$ and there exists a point

$$A \in \mathcal{C}_1 = \partial\omega_1$$

such that $|\nabla u(A)| = |v(A)| > 0$ and $\nabla u(A)$ is parallel to the normal vector $n(A)$. Assume first that $\nabla u(A)$ points into Ω at A , that is, $\nabla u(A) = -|\nabla u(A)|n(A)$, and let

$$\sigma = \sigma_A$$

be the solution of (2.10) with $x = A$. With the notations of Lemma 2.2 (ii), the function σ is of class $C^1([0, t_A^+))$ and $(u \circ \sigma)' > 0$ in $[0, t_A^+)$, with $u(\sigma(0)) = u(A) = c_{\text{in}}$ by Lemma 2.1 (i). Furthermore, (2.12) implies that $\text{dist}(\sigma(t), \partial\omega_2) \rightarrow 0$ as $t \rightarrow t_A^+$, hence $u(\sigma(t)) \rightarrow c_{\text{out}}$ as $t \rightarrow t_A^+$, where $c_{\text{out}} \in \mathbb{R}$ is given by Lemma 2.1 (ii) (since (2.5) is satisfied by (2.15), here with $\Omega = \omega_2 \setminus \overline{\omega_1}$). In particular, one has $c_{\text{in}} < c_{\text{out}}$ and $c_{\text{in}} < u < c_{\text{out}}$ in Ω . On the other hand, the function $|\nabla u \circ \sigma|$ is continuous in $[0, t_A^+)$, positive at 0 (since $|\nabla u(\sigma(0))| = |v(A)| > 0$), positive in $(0, t_A^+)$ (since $\sigma((0, t_A^+)) \subset \Omega$), and

$$\liminf_{t \rightarrow t_A^+} |\nabla u(\sigma(t))| = \liminf_{t \rightarrow t_A^+} |v(\sigma(t))| > 0$$

(since the continuous field $|v|$ is positive on the compact set $\mathcal{C}_2 = \partial\omega_2$ and $\text{dist}(\sigma(t), \partial\omega_2) \rightarrow 0$ as $t \rightarrow t_A^+$). As a consequence, there is $\eta > 0$ such that $|\dot{\sigma}(t)| = |\nabla u(\sigma(t))| \geq \eta$ for all $t \in [0, t_A^+)$. Therefore, $(u \circ \sigma)'(t) = |\nabla u(\sigma(t))|^2 \geq \eta^2$ for all $t \in [0, t_A^+)$ and t_A^+ is a positive real number, since u is bounded in the compact set $\overline{\Omega}$. Moreover, for every $t \in [0, t_A^+)$,

$$c_{\text{out}} - c_{\text{in}} \geq u(\sigma(t)) - u(A) = u(\sigma(t)) - u(\sigma(0)) = \int_0^t |\nabla u(\sigma(s))|^2 ds \geq \eta \int_0^t |\dot{\sigma}(s)| ds,$$

hence the length of the curve $\sigma([0, t_A^+))$ is finite. Finally, there is a point $A^+ \in \mathcal{C}_2 = \partial\omega_2$ such that $\sigma(t) \rightarrow A^+$ as $t \rightarrow t_A^+$. By setting $\sigma(t_A^+) = A^+$ and remembering that the field ∇u is (at least) continuous in $\overline{\Omega}$, it follows that the function $\sigma : [0, t_A^+] \rightarrow \overline{\Omega} = D$ is then of class $C^1([0, t_A^+])$ and $(u \circ \sigma)' > 0$ in $[0, t_A^+]$. To sum up, if $\nabla u(A)$ points into Ω at A , then (2.16) holds in the case $\Omega = \omega_2 \setminus \overline{\omega_1}$ with $t_{\text{in}} = 0$ and $t_{\text{out}} = t_A^+$. Similarly, if $\nabla u(A)$ points outward from Ω at A , then (2.16) still holds, with $t_{\text{in}} = 0$, $t_{\text{out}} = -t_A^- \in (0, +\infty)$, $\sigma(t) = \sigma_A(-t)$ for $t \in [0, -t_A^-]$, and σ_A of class $C^1([t_A^-, 0])$.

Consider now the case $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1}$. By (2.3) and (2.15), there is a point $A \in \mathcal{C}_1 = \partial\omega_1$ such that $\nabla u(A)$ is not zero and parallel to the normal vector $n(A)$. Assume that $\nabla u(A)$ points into Ω at A (the other case can be treated similarly) and let $\sigma = \sigma_A$. The function $\sigma : [0, t_A^+) \rightarrow \Omega \cup \partial\omega_1 = \overline{\Omega} = D$ is of class $C^1([0, t_A^+))$ and $(u \circ \sigma)' > 0$ in $[0, t_A^+)$, for some $t_A^+ \in (0, +\infty]$. It then follows from Lemma 2.2 (ii) (with $t \rightarrow t_A^+$) that $|\sigma(t)| \rightarrow +\infty$ as $t \rightarrow t_A^+$. This yields (2.16) with $t_{\text{in}} = 0$ and $t_{\text{out}} = t_A^+$.

Consider then the case $\Omega = \omega_2 \setminus \{0\}$. By (2.15), there is a point $A \in \mathcal{C}_2 = \partial\omega_2$ such that $\nabla u(A)$ is not zero and parallel to the normal vector $n(A)$. Assume that $\nabla u(A)$ points outward from Ω at A (the other case can be treated similarly) and let $\sigma = \sigma_A$. The function $\sigma : (t_A^-, 0] \rightarrow \Omega \cup \partial\omega_2 = D$ is of class $C^1((t_A^-, 0])$ and $(u \circ \sigma)' > 0$ in $(t_A^-, 0]$, for some $t_A^- \in [-\infty, 0)$. Since (2.5) is satisfied by (2.15) (here with $\Omega = \omega_2 \setminus \{0\}$), it then follows from Lemma 2.2 (iii) (with $t \rightarrow t_A^-$) that $|\sigma(t)| \rightarrow 0$ as $t \rightarrow t_A^-$. This yields (2.16) with $t_{\text{in}} = t_A^-$ and $t_{\text{out}} = 0$.

Lastly, consider the case $\Omega = \mathbb{R}^2 \setminus \{0\}$. By (2.15), condition (2.5) is fulfilled. By Lemma 2.1, there are then $c_{\text{in}} \neq c_{\text{out}}$ in $\overline{\mathbb{R}}$ such that $u(x) \rightarrow c_{\text{in}}$ as $|x| \xrightarrow{\sim} 0$ and $u(x) \rightarrow c_{\text{out}}$ as $|x| \rightarrow +\infty$. Pick any point $A \in \Omega$. By Lemma 2.2 (i), the function $\sigma_A : (t_A^-, t_A^+) \rightarrow \Omega = D$ is of class $C^1((t_A^-, t_A^+))$, with $-\infty \leq t_A^- < t_A^+ \leq +\infty$, and $(u \circ \sigma_A)' > 0$ in (t_A^-, t_A^+) . Using also (2.11), one finds that either $|\sigma_A(t)| \rightarrow 0$ as $t \rightarrow t_A^-$ and $|\sigma_A(t)| \rightarrow +\infty$ as $t \rightarrow t_A^+$, or $|\sigma_A(t)| \rightarrow +\infty$ as $t \rightarrow t_A^-$ and $|\sigma_A(t)| \rightarrow 0$ as $t \rightarrow t_A^+$. In the former case, (2.16) holds with $\sigma = \sigma_A$ and $(t_{\text{in}}, t_{\text{out}}) = (t_A^-, t_A^+)$, whereas in the latter case (2.16) holds with $\sigma = \sigma_A(-\cdot)$ and $(t_{\text{in}}, t_{\text{out}}) = (-t_A^+, -t_A^-)$.

(ii) Assume now that $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1}$ and that (2.17) is fulfilled. In particular, (2.15) holds and the previous part (i) yields the existence of $0 = t_{\text{in}} < t_{\text{out}} \leq +\infty$ and a C^1 function $\sigma : [t_{\text{in}}, t_{\text{out}}) \rightarrow \overline{\Omega}$ such that $(u \circ \sigma)' \neq 0$ in $[t_{\text{in}}, t_{\text{out}})$, $\sigma(t_{\text{in}}) = \sigma(0) \in \partial\omega_1$, and $|\sigma(t)| \rightarrow +\infty$ as $t \xrightarrow{\sim} t_{\text{out}}$. Furthermore, by construction, either $\sigma = \sigma_A$, with $A = \sigma(0) \in \partial\omega_1$ such that $|v(A)| > 0$, or $\sigma = \sigma_A(-\cdot)$. The only thing to prove is that $|u(\sigma(t))| \rightarrow +\infty$ as $t \xrightarrow{\sim} t_{\text{out}}$. By (2.17), there are $R > 0$ and $\eta > 0$ such that $\mathbb{R}^2 \setminus B_R \subset \Omega$ and $|v(x)| \geq \eta$ for

all $|x| \geq R$. Let $T \in (0, t_{\text{out}})$ be such that $|\sigma(s)| \geq R$ for all $s \in [T, t_{\text{out}})$. For all $t \in [T, t_{\text{out}})$, one has

$$|u(\sigma(t)) - u(\sigma(T))| = \int_T^t |\nabla u(\sigma(s))|^2 ds \geq \eta \int_T^t |\dot{\sigma}(s)| ds \geq \eta(|\sigma(t)| - |\sigma(T)|).$$

Consequently, $|u(\sigma(t))| \rightarrow +\infty$ as $t \xrightarrow{\sim} t_{\text{out}}$. The proof of Lemma 2.4 is complete. ■

The fourth common result states that the streamlines of the flow are C^1 Jordan curves surrounding the origin, and that they approach 0 or infinity as appropriate. For x in D defined by (2.1), ξ_x denotes the solution of (1.5), ranging in D and defined in a maximal interval I_x containing 0, and

$$\Xi_x = \xi_x(I_x)$$

denotes the streamline of the flow containing x . We recall that, by definition, the stream function u given in Lemma 2.1 is constant along each streamline of the flow.

Lemma 2.6. (i) *If (2.5) holds, then, for every $x \in \Omega$, the function ξ_x is defined in \mathbb{R} and periodic, and the streamline $\Xi_x = \xi_x(\mathbb{R})$ is a C^1 Jordan curve surrounding the origin. Furthermore,*

$$\begin{cases} \min_{\mathbb{R}} |\xi_x| \rightarrow +\infty & \text{as } |x| \rightarrow +\infty & \text{if } \Omega \text{ is unbounded,} \\ \max_{\mathbb{R}} |\xi_x| \rightarrow 0 & \text{as } |x| \xrightarrow{\sim} 0 & \text{if } \Omega \text{ is punctured.} \end{cases}$$

(ii) *If (2.5) holds with Ω unbounded and $\liminf_{|x| \rightarrow +\infty} |v(x) \cdot e_\theta(x)| > 0$, then*

$$\max_{\mathbb{R}} |\xi_x| - \min_{\mathbb{R}} |\xi_x| \rightarrow 0 \quad \text{as } |x| \rightarrow +\infty.$$

(iii) *If $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1}$ and $\inf_{\Omega} |v| > 0$, then, for every $x \in \overline{\Omega}$, the function ξ_x is defined in \mathbb{R} and periodic, and the streamline $\Xi_x = \xi_x(\mathbb{R})$ is a C^1 Jordan curve surrounding the origin. Furthermore, $\min_{\mathbb{R}} |\xi_x| \rightarrow +\infty$ and $|u(x)| \rightarrow +\infty$ as $|x| \rightarrow +\infty$, and $u - c_{\text{in}}$ has a constant strict sign in Ω , where $u = c_{\text{in}}$ on $\partial\omega_1$.*

Proof. (i) From Lemma 2.1, there are $c_{\text{in}} \neq c_{\text{out}}$ in $\overline{\mathbb{R}}$ such that the stream function u satisfies (2.4) and (2.6)–(2.7). Therefore, the continuity of u in D and the fact that u is constant along each streamline of the flow imply that, for each $x \in \Omega$, $\inf_{t \in I_x} \text{dist}(\xi_x(t), \partial\Omega) > 0$ and $\Xi_x = \xi_x(I_x)$ is bounded. Since $|\nabla u| = |v| > 0$ in Ω by (2.2), it is standard to conclude that, for each $x \in \Omega$, the function ξ_x is periodic and the streamline $\Xi_x = \xi_x(\mathbb{R})$ (here, $I_x = \mathbb{R}$) is a C^1 Jordan curve surrounding the origin.

If Ω is unbounded, then $u(x) \rightarrow c_{\text{out}} \in \overline{\mathbb{R}}$ as $|x| \rightarrow +\infty$ by (2.6). Applying also (2.4), (2.7), and the continuity of u in D , it easily follows that $\min_{t \in \mathbb{R}} |\xi_x(t)| \rightarrow +\infty$ as $|x| \rightarrow +\infty$. Similarly, if Ω is punctured, then $\max_{t \in \mathbb{R}} |\xi_x(t)| \rightarrow 0$ as $|x| \xrightarrow{\sim} 0$.

(ii) In the unbounded case, assume now that $\liminf_{|x| \rightarrow +\infty} |v(x) \cdot e_\theta(x)| > 0$, in addition to (2.5). From (i), one knows that, for each $x \in \Omega$, the streamline Ξ_x is a C^1 Jordan curve surrounding the origin, with $\min_{\mathbb{R}} |\xi_x| \rightarrow +\infty$ as $|x| \rightarrow +\infty$. Let us now show that $\max_{\mathbb{R}} |\xi_x| - \min_{\mathbb{R}} |\xi_x| \rightarrow 0$ as $|x| \rightarrow +\infty$. Let $R > 0$ and $\eta > 0$ be such that $\mathbb{R}^2 \setminus B_R \subset \Omega$

and $|\nabla u(x) \cdot e_r(x)| = |v(x) \cdot e_\theta(x)| \geq \eta$ for all $|x| \geq R$. In view of the continuity of ∇u , let us only consider the case

$$\nabla u(x) \cdot e_r(x) \geq \eta \quad \text{for all } |x| \geq R \tag{2.18}$$

(the case $\nabla u(x) \cdot e_r(x) \leq -\eta$ can be handled similarly). Consider any $\varepsilon > 0$. From (2.9) and (2.18), there is $R_\varepsilon \geq R + \varepsilon$ such that

$$\max_{C_{|x|-\varepsilon}} u < u(x) - \varepsilon \eta/2 \quad \text{and} \quad \min_{C_{|x|+\varepsilon}} u > u(x) + \varepsilon \eta/2, \quad \text{for all } |x| \geq R_\varepsilon.$$

Thus, for every x with $|x| \geq R_\varepsilon$, one has $\Xi_x \subset \Omega_{|x|-\varepsilon, |x|+\varepsilon}$ and $\max_{\mathbb{R}} |\xi_x| - \min_{\mathbb{R}} |\xi_x| < 2\varepsilon$.

(iii) Consider now the case $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1}$ and assume that

$$\eta := \inf_{\Omega} |v| > 0. \tag{2.19}$$

Notice that we do not assume condition (2.5), so Lemma 2.1 (ii) and parts (i)–(ii) of the present lemma cannot be applied. In particular, we do not know yet that u has a limit at infinity or that all streamlines of the flow surround the origin. However, Lemma 2.4 (ii) can be applied, since (2.17) is fulfilled here. Therefore, there exist $t_{\text{out}} \in (0, +\infty]$, a point $A \in \partial\omega_1$ (actually, A can here be arbitrarily chosen on $\partial\omega_1$ since $|v| > 0$ on $\partial\omega_1$), and a C^1 function $\sigma : [0, t_{\text{out}}] \rightarrow D$ such that $\sigma(0) = A$, $(u \circ \sigma)' \neq 0$ in $[0, t_{\text{out}})$, and $|\sigma(t)| \rightarrow +\infty$ and $|u(\sigma(t))| \rightarrow +\infty$ as $t \nearrow t_{\text{out}}$. Assume that $(u \circ \sigma)' > 0$ in $[0, t_{\text{out}})$ (the other case can be handled similarly), hence

$$u(\sigma(t)) \rightarrow +\infty \quad \text{as } t \nearrow t_{\text{out}}. \tag{2.20}$$

By construction of σ in Lemma 2.4 (ii), this case corresponds to the condition that $\nabla u(A) \cdot n(A) < 0$.

Denote

$$E = \{s \in [0, t_{\text{out}}] : \text{the streamline } \Xi_{\sigma(s)} \text{ is a } C^1 \text{ Jordan curve surrounding the origin}\},$$

and let us show that

$$E = [0, t_{\text{out}}]. \tag{2.21}$$

To do so, we prove that E is not empty (it contains 0), relatively open in $[0, t_{\text{out}})$ and that the largest interval containing 0 and contained in E is actually $[0, t_{\text{out}})$. Note first that since $v \cdot n = 0$ and $|v| > 0$ on $\mathcal{C}_1 = \partial\omega_1$, the streamline $\Xi_{\sigma(0)} = \Xi_A$ is equal to the C^1 Jordan curve \mathcal{C}_1 and it surrounds the origin by assumption. In other words, $0 \in E$.

Let us now show that E is relatively open in $[0, t_{\text{out}})$. Let $s_0 \in E$ and denote $x = \sigma(s_0) \in \overline{\Omega}$. By definition, the function ξ_x is periodic, with some period $T_x > 0$. Remember also that u is constant along each streamline of the flow. Therefore, since v is (at least) continuous and $|v(x)| = |\nabla u(x)| > 0$ in $\overline{\Omega}$, there are real numbers $r > 0$ and $\tau \in (0, T_x)$ such that, for every $y \in B(x, r) \cap \overline{\Omega}$, there are real numbers t_y^\pm such that

$$-\tau < t_y^- < 0 < t_y^+ < \tau \quad \text{and} \quad B(x, r) \cap \Xi_y = B(x, r) \cap u^{-1}(\{u(y)\}) = \xi_y((t_y^-, t_y^+)).$$

On the other hand, since $\xi_x(T_x) = \xi_x(0) = x$, the Cauchy–Lipschitz theorem provides the existence of $r' \in (0, r]$ such that, for every $z \in B(x, r') \cap \overline{\Omega}$, the function ξ_z is defined (and of class C^1) at least on the interval $[0, T_x]$ and $\xi_z(T_x) \in B(x, r) \cap \overline{\Omega}$. Furthermore, by continuity of σ , there is $\varepsilon > 0$ such that $s_0 + \varepsilon < t_{\text{out}}$ and

$$\sigma(s) \in B(\sigma(s_0), r') \cap \overline{\Omega} = B(x, r') \cap \overline{\Omega} \quad \text{for all } s \in [\max(0, s_0 - \varepsilon), s_0 + \varepsilon].$$

As a consequence, for every $s \in [\max(0, s_0 - \varepsilon), s_0 + \varepsilon]$, the points $z := \sigma(s) \in B(x, r') \cap \overline{\Omega}$ and $y := \xi_z(T_x) \in B(x, r) \cap \overline{\Omega}$ satisfy $u(z) = u(y)$, hence

$$z \in B(x, r') \cap u^{-1}(\{u(y)\}) \subset B(x, r) \cap u^{-1}(\{u(y)\})$$

and $z = \xi_y(t)$ for some $t \in (t_y^-, t_y^+) \subset (-\tau, \tau)$. Thus, $\xi_y(-T_x) = z = \xi_y(t)$ and since $|t| < \tau < T_x$, the function ξ_y is defined in \mathbb{R} and $(T_x + t)$ -periodic. So is ξ_z since $z \in \Xi_y$. In other words, for every $s \in [\max(0, s_0 - \varepsilon), s_0 + \varepsilon]$, the function $\xi_{\sigma(s)} = \xi_z$ is defined in \mathbb{R} and periodic. Since $|\nabla u| = |v| > 0$ in $\overline{\Omega}$, one then concludes that $\Xi_{\sigma(s)}$ is a C^1 Jordan curve surrounding the origin. Finally, the set E is relatively open in $[0, t_{\text{out}}]$.

Denote

$$T_* = \sup \{t \in [0, t_{\text{out}}] : [0, t] \subset E\}.$$

The previous paragraphs imply that $0 < T_* \leq t_{\text{out}}$. The proof of (2.21) will be complete once we show that $T_* = t_{\text{out}}$. Assume by way of contradiction that $T_* < t_{\text{out}}$ (in particular, T_* is then a positive real number). Consider any increasing sequence $(s_n)_{n \in \mathbb{N}}$ in $(0, T_*)$ and converging to T_* . Owing to the definition of T_* , each function $\xi_{\sigma(s_n)}$ is periodic and each streamline $\Xi_{\sigma(s_n)}$ surrounds the origin. Furthermore, since each s_n is positive and $u \circ \sigma$ is increasing in $[0, t_{\text{out}}]$ and u is constant on \mathcal{C}_1 ($\ni A = \sigma(0)$), each streamline $\Xi_{\sigma(s_n)}$ is included in the (open) set Ω . Consider now any $n \in \mathbb{N}$ and any point

$$x \in \Xi_{\sigma(s_n)}.$$

Notice that $u(x) = u(\sigma(s_n))$ and remember that $u \circ \sigma$ is increasing in $[0, t_{\text{out}}]$, hence

$$u(A) = u(\sigma(0)) < u(\sigma(s_n)) = u(x) < u(\sigma(T_*)).$$

Moreover, by Lemma 2.2 (i), there exist $-\infty \leq t_x^- < 0 < t_x^+ \leq +\infty$ such that $\sigma_x : (t_x^-, t_x^+) \rightarrow \Omega$ is of class C^1 with

$$(u \circ \sigma_x)' > 0 \text{ in } (t_x^-, t_x^+), \text{ and } \text{dist}(\sigma_x(t), \partial\omega_1) \rightarrow 0 \text{ or } |\sigma_x(t)| \rightarrow +\infty \text{ as } t \rightarrow t_x^\pm. \tag{2.22}$$

The nonzero vector $\dot{\sigma}_x(0) = \nabla u(\sigma_x(0)) = \nabla u(x)$ is orthogonal to $\Xi_{\sigma(s_n)}$ at x by definition of u . Since $u(\sigma_x(0)) = u(x) > u(A) = c_{\text{in}}$ with $u = c_{\text{in}}$ on $\mathcal{C}_1 = \partial\omega_1$, and since $\Xi_{\sigma(s_n)}$ is a C^1 Jordan curve surrounding the origin and meeting $\sigma_x((t_x^-, t_x^+))$ orthogonally at the only point x , it follows from (2.22) that $\text{dist}(\sigma_x(t), \partial\omega_1) \rightarrow 0$ as $t \rightarrow t_x^-$ and $u(\sigma_x(t)) > c_{\text{in}}$ for all $t \in (t_x^-, t_x^+)$. Then, for any $t \in (t_x^-, 0)$,

$$\begin{aligned} u(\sigma(T_*)) > u(x) &= u(\sigma_x(0)) = \int_t^0 |\nabla u(\sigma_x(s))|^2 ds + u(\sigma_x(t)) \\ &\geq \eta \int_t^0 |\dot{\sigma}_x(s)| ds + c_{\text{in}} \geq \eta(|x| - |\sigma_x(t)|) + c_{\text{in}} \end{aligned}$$

with $\eta > 0$ as in (2.19). In the limit as $t \rightarrow t_x^-$, one gets $|x| \leq R_1 + (u(\sigma(T_*)) - c_{in})/\eta$ with $R_1 := \max_{y \in \mathcal{C}_1} |y|$. This holds for any $n \in \mathbb{N}$ and any $x \in \Xi_{\sigma(s_n)}$, hence

$$\sup_{n \in \mathbb{N}} \left(\max_{\mathbb{R}} |\xi_{\sigma(s_n)}| \right) \leq R_1 + \frac{u(\sigma(T_*)) - c_{in}}{\eta} =: M.$$

Finally, consider the streamline $\Xi_{\sigma(T_*)}$ parametrized by the function $\xi_{\sigma(T_*)}$. If there is a real number t such that $|\xi_{\sigma(T_*)}(t)| > M$, then $|\xi_{\sigma(s_n)}(t)| > M$ for all n large enough, by the Cauchy-Lipschitz theorem. Therefore, $\Xi_{\sigma(T_*)} \subset \overline{B_M}$, and since $|v| > 0$ in $\overline{\Omega}$, it follows that $\xi_{\sigma(T_*)}$ is defined in \mathbb{R} and periodic, and $\Xi_{\sigma(T_*)}$ is a C^1 Jordan curve surrounding the origin. In other words, $T_* \in E$. Since E has been proved to be relatively open in $[0, t_{out})$, one is led to a contradiction with the definition of T_* if $T_* < t_{out}$. Eventually, $T_* = t_{out}$ and (2.21) is proved.

We now claim that

$$\min_{\mathbb{R}} |\xi_{\sigma(s)}| \rightarrow +\infty \quad \text{as } s \xrightarrow{\sim} t_{out}. \tag{2.23}$$

Indeed, for any $R > R_1$, let $C \in [0, +\infty)$ be such that $|u| \leq C$ in $\overline{\Omega} \cap \overline{B_R}$. By (2.20), there is then $\tau \in (0, t_{out})$ such that $u(\sigma(s)) > C$ for all $s \in (\tau, t_{out})$, hence $u(\xi_{\sigma(s)}(t)) = u(\sigma(s)) > C$ and $|\xi_{\sigma(s)}(t)| > R$ for all $s \in (\tau, t_{out})$ and $t \in \mathbb{R}$. Thus, $\min_{\mathbb{R}} |\xi_{\sigma(s)}| > R$ for all $s \in (\tau, t_{out})$. This yields (2.23).

Consider then any point $x \in \overline{\Omega}$ and let us deduce that Ξ_x surrounds the origin. From (2.20)–(2.21) and (2.23), there is $s \in (0, t_{out})$ such that $u(\sigma(s)) > u(x)$ and the streamline $\Xi_{\sigma(s)}$ is a C^1 Jordan curve surrounding both x and the origin. Therefore, the streamline Ξ_x is bounded (it belongs to the bounded connected component of $\mathbb{R}^2 \setminus \Xi_{\sigma(s)}$). Using again the fact that $|v| > 0$ in $\overline{\Omega}$, one concludes that the function ξ_x is periodic and Ξ_x is a C^1 Jordan curve surrounding the origin.

From the previous properties, we easily deduce that $\min_{\mathbb{R}} |\xi_x| \rightarrow +\infty$ as $|x| \rightarrow +\infty$. Indeed, for any fixed $R > R_1 = \max_{y \in \partial\omega_1} |y|$, there is $s \in (0, t_{out})$ such that $\min_{\mathbb{R}} |\xi_{\sigma(s)}| > R$, by (2.23). Then, for any x with $|x| > R' := \max_{\mathbb{R}} |\xi_{\sigma(s)}| > R$, the streamlines Ξ_x and $\Xi_{\sigma(s)}$ do not intersect, and both are C^1 Jordan curves surrounding the origin, hence $\min_{\mathbb{R}} |\xi_x| > \min_{\mathbb{R}} |\xi_{\sigma(s)}| > R$. This shows that $\min_{\mathbb{R}} |\xi_x| \rightarrow +\infty$ as $|x| \rightarrow +\infty$.

Finally, let us prove that $u(x) \rightarrow +\infty$ as $|x| \rightarrow +\infty$. Fix any $B > 0$, and, by (2.20), let $s \in (0, t_{out})$ be such that $u \circ \sigma > B$ in (s, t_{out}) . Let $R_B > R_1$ be such that $\min_{\mathbb{R}} |\xi_x| > \max_{t \in [0, s]} |\sigma(t)|$ for all $|x| \geq R_B$. For every x such that $|x| \geq R_B$, the streamline Ξ_x surrounds the origin and necessarily crosses $\sigma([0, t_{out}))$, at a point $\sigma(s_x)$ with $s_x \in (s, t_{out})$, hence $u(x) = u(\sigma(s_x)) > B$. This shows that $u(x) \rightarrow +\infty$ as $|x| \rightarrow +\infty$. Since $u = c_{in}$ on $\partial\omega_1$ and u has no critical point in Ω , one concludes that $u > c_{in}$ in Ω . The proof of Lemma 2.6 is thereby complete. ■

Remark 2.7. If $|v| > 0$ on $\mathcal{C}_1 = \partial\omega_1$ (resp. on $\mathcal{C}_2 = \partial\omega_2$), then the boundary conditions (2.3) (resp. (2.5)) imply that, for any $x \in \partial\omega_1$ with Ω not punctured (resp. $x \in \partial\omega_2$ with Ω bounded), ξ_x is still defined and periodic in \mathbb{R} with $\Xi_x = \partial\omega_1$ (resp. $\Xi_x = \partial\omega_2$). If $x \in \partial\Omega \cap D$ and $|v(x)| = 0$, then $\xi_x(t) = x$ for all $t \in \mathbb{R}$ and $\Xi_x = \{x\}$. If $x \in \partial\omega_1$

with Ω not punctured (resp. $x \in \partial\omega_2$ with Ω bounded) with $|v(x)| > 0$ and if v has some stagnation points on $\partial\omega_1$ (resp. $\partial\omega_2$), then ξ_x is still defined in \mathbb{R} , but it is not periodic anymore and Ξ_x is a proper arc of $\partial\omega_1$ (resp. $\partial\omega_2$) which is relatively open in $\partial\omega_1$ (resp. $\partial\omega_2$).

In the last common preliminary result, we derive a semilinear elliptic equation $\Delta u + f(u) = 0$ in D for some C^1 function f defined in the range of u . We recall that D is defined in (2.1).

Lemma 2.8. *Assume that v is of class $C^2(D)$, solves the Euler equations (1.1) in D , and satisfies (2.2)–(2.3). Let u in $C^3(D)$ be given by Lemma 2.1 and let J be its range $J = \{u(x) : x \in D\}$.*

(i) *If (2.5) and (2.15) are fulfilled, then there is a C^1 function $f : J \rightarrow \mathbb{R}$ such that*

$$\Delta u + f(u) = 0 \quad \text{in } D.$$

(ii) *If $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1}$ and $\inf_{\Omega} |v| > 0$, then the same conclusion holds.*

Proof. First of all, since v is of class $C^2(D)$, the stream function u given by Lemma 2.1 is now of class $C^3(D)$. By continuity of u and connectedness of D , the range J of u is an interval.

(i) Assume (2.5) and (2.15). It follows from Lemma 2.1 and its notations that the interior of J is $(\min(c_{\text{in}}, c_{\text{out}}), \max(c_{\text{in}}, c_{\text{out}}))$. Furthermore, J is open at c_{in} if and only if Ω is punctured, while J is open at c_{out} if and only if Ω is unbounded (for instance, if $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1}$, then $J = [c_{\text{in}}, c_{\text{out}})$ or $(c_{\text{out}}, c_{\text{in}}]$). It then follows from Lemmas 2.1 and 2.4 (i) that, with the same notations as there, the function $g := u \circ \sigma : I \rightarrow J$ is a C^1 diffeomorphism from I onto J . Let $g^{-1} : J \rightarrow I$ be its C^1 inverse diffeomorphism, and define

$$f(\tau) = -\Delta u(\sigma(g^{-1}(\tau))) \quad \text{for } \tau \in J. \quad (2.24)$$

By the chain rule, the function f is of class $C^1(J)$ (remember that Δu is now of class $C^1(D)$). The above formula means that the equation $\Delta u + f(u) = 0$ is satisfied along the curve $\sigma(I)$. Let us now check it in the whole set D . Consider first any point $x \in \Omega$. From Lemmas 2.4 (i) and 2.6 (i), the streamline Ξ_x surrounds the origin and meets the curve $\sigma(I)$. Hence, there is $s \in I$ such that $\sigma(s) \in \Xi_x$. On the one hand, the stream function u is constant along the streamline Ξ_x . On the other hand, the $C^1(D)$ vorticity $\frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2} = \Delta u$ satisfies $v \cdot \nabla(\Delta u) = 0$ in D from the Euler equations (1.1), hence Δu is constant along the streamline Ξ_x too. As a consequence, (2.24) yields

$$\Delta u(x) + f(u(x)) = \Delta u(\sigma(s)) + f(u(\sigma(s))) = \Delta u(\sigma(s)) + f(g(s)) = 0.$$

Therefore, $\Delta u + f(u) = 0$ in Ω . Finally, since both functions Δu and $f \circ u$ are (at least) continuous in D , one concludes that $\Delta u + f(u) = 0$ in D .

(ii) Assume that $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1}$ and $\inf_{\Omega} |v| > 0$. From Lemmas 2.1 (i) and 2.6 (iii), one has $J = [c_{\text{in}}, +\infty)$ or $(-\infty, c_{\text{in}}]$. By Lemma 2.4 (ii), there are t_{out} in $(0, +\infty)$ and a C^1 curve $\sigma : [0, t_{\text{out}}) \rightarrow D = \overline{\Omega}$ such that $g := u \circ \sigma$ is a C^1 diffeomorphism from $[0, t_{\text{out}})$

onto J . Let $g^{-1} : J \rightarrow [0, t_{\text{out}})$ be its C^1 inverse diffeomorphism, and define $f \in C^1(J)$ as in (2.24). Since, by Lemma 2.6 (iii), each streamline Ξ_x (for each $x \in D = \overline{\Omega}$) surrounds the origin, it follows as in (i) above that the equation $\Delta u + f(u) = 0$ holds, here directly in D . The proof of Lemma 2.8 is thereby complete. ■

3. Proof of the main results in fixed annular domains $\Omega_{a,b}$

This section is devoted to the proofs of Theorems 1.1–1.5 and 1.8 on the Euler flows in the fixed annular domains $\Omega_{a,b}$ with $0 \leq a < b \leq \infty$. The proofs rely on the common properties proved in Section 2, as well as on various applications of Proposition 1.14 and further specific arguments in the unbounded and punctured cases.

3.1. The case of bounded annuli $\Omega_{a,b}$: proofs of Theorems 1.1 and 1.2

This section is devoted to the proof of Theorem 1.2 (we recall that Theorem 1.1 is a particular case). Throughout this section, we consider real numbers $0 < a < b$ and a $C^2(\overline{\Omega_{a,b}})$ solution v of (1.1)–(1.2) satisfying (1.3), that is,

$$\{x \in \overline{\Omega_{a,b}} : |v(x)| = 0\} \not\subset C_a \quad \text{or} \quad \{x \in \overline{\Omega_{a,b}} : |v(x)| = 0\} \not\subset C_b.$$

This situation falls within the general framework of Section 2, with $\mathcal{C}_1 = C_a$, $\omega_1 = B_a$, $\mathcal{C}_2 = C_b$, $\omega_2 = B_b$, $\Omega = \omega_2 \setminus \overline{\omega_1} = \Omega_{a,b}$, and $D = \overline{\Omega_{a,b}}$. Notice also that conditions (2.2)–(2.3), (2.5), and (2.15) are fulfilled by assumption. Therefore, the flow has a $C^3(\overline{\Omega_{a,b}})$ stream function u and, by Lemmas 2.1 and 2.8 (i), there are real numbers $c_{\text{in}} \neq c_{\text{out}}$ and a $C^1([\min(c_{\text{in}}, c_{\text{out}}), \max(c_{\text{in}}, c_{\text{out}})])$ function f such that

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \overline{\Omega_{a,b}}, \\ \min(c_{\text{in}}, c_{\text{out}}) < u < \max(c_{\text{in}}, c_{\text{out}}) & \text{in } \Omega_{a,b}, \\ u = c_{\text{in}} \text{ on } C_a, \quad u = c_{\text{out}} \text{ on } C_b. \end{cases} \tag{3.1}$$

It then follows from [27, Theorem 5]⁶ that u is radially symmetric and strictly monotone with respect to $|x|$ in $\overline{\Omega_{a,b}}$. Therefore, there is a $C^3([a, b])$ strictly monotone function $U : [a, b] \rightarrow \mathbb{R}$ such that $u(x) = U(|x|)$ for all $x \in \overline{\Omega_{a,b}}$. The flow $v = \nabla^\perp u$ is then given by

$$v(x) = V(|x|)e_\theta(x)$$

for all $x \in \overline{\Omega_{a,b}}$, with $V = U' \in C^2([a, b])$. Lastly, since $|v|$ is continuous in $\overline{\Omega_{a,b}}$ and does not vanish in $\Omega_{a,b}$, nor in the whole circle C_a or in the whole circle C_b , the function V has a constant strict sign in $[a, b]$. The proof of Theorem 1.2 is thereby complete. ■

⁶Notice that this result holds in any dimension $n \geq 2$. It is similar to the classical radial symmetry property proved in [11] in the case where u is a positive solution of the equation $\Delta u + f(u) = 0$ in a ball, with Dirichlet condition $u = 0$ on the boundary.

A related open question. For a $C^2(\overline{\Omega_{a,b}})$ flow v solving (1.1)–(1.2), could the assumption (1.3) be slightly relaxed for v still to be necessarily a circular flow? As we mentioned in the introduction, the conclusion does not hold in general if v has stagnation points in $\Omega_{a,b}$. So a natural question is:

$$\text{if } |v| > 0 \text{ in } \Omega_{a,b}, \text{ is } v \text{ a circular flow?} \tag{3.2}$$

We first point out that Lemmas 2.1, 2.2 and 2.6 (i), with $\Omega = \Omega_{a,b}$, still hold since they do not use the whole assumption (2.15) (more precisely, they do not use (1.3)), but only $|v| > 0$ in $\Omega_{a,b}$. Consider then any point $y \in \Omega_{a,b}$. With the notations of Lemma 2.2, and assuming that $c_{in} < c_{out}$ (after possibly changing v to $-v$ and u to $-u$), there are some quantities t_y^\pm such that $-\infty \leq t_y^- < 0 < t_y^+ \leq +\infty$ and the solution σ_y of (2.10) with y in place of x is of class $C^1((t_y^-, t_y^+))$ and ranges in $\Omega_{a,b}$, with

$$\begin{cases} |\sigma_y(t)| \rightarrow a \text{ and } u(\sigma_y(t)) \rightarrow c_{in} & \text{as } t \rightarrow t_y^-, \\ |\sigma_y(t)| \rightarrow b \text{ and } u(\sigma_y(t)) \rightarrow c_{out} & \text{as } t \rightarrow t_y^+. \end{cases} \tag{3.3}$$

The $C^1((t_y^-, t_y^+))$ function $g := u \circ \sigma_y$ is increasing (we recall that $(u \circ \sigma_y)'(t) = |\nabla u(\sigma_y(t))|^2 = |v(\sigma_y(t))|^2 > 0$ for all $t \in (t_y^-, t_y^+)$), and so g is an increasing homeomorphism from (t_y^-, t_y^+) onto (c_{in}, c_{out}) . The function $f : (c_{in}, c_{out}) \rightarrow \mathbb{R}$ defined by

$$f(\tau) = -\Delta u(\sigma_y(g^{-1}(\tau))) \quad \text{for } \tau \in (c_{in}, c_{out}) \tag{3.4}$$

is of class $C^1((c_{in}, c_{out}))$ and, since for every $x \in \Omega_{a,b}$ the streamline Ξ_x intersects $\sigma_y((t_y^-, t_y^+))$ by Lemma 2.6 (i), the same arguments as in the proof of Lemma 2.8 (i) imply that

$$\Delta u + f(u) = 0 \quad \text{in } \Omega_{a,b}.$$

Furthermore, remembering from Lemma 2.6 (i) that, for each $x \in \Omega_{a,b}$, the C^1 solution ξ_x of (1.5) is periodic and ranges in $\Omega_{a,b}$, it follows from the continuity of u in the compact set $\overline{\Omega_{a,b}}$ and the facts that $u = c_{in}$ on C_a , $u = c_{out}$ on C_b and $c_{in} < u < c_{out}$ in $\Omega_{a,b}$, that

$$\max_{t \in \mathbb{R}} |\xi_x(t)| \rightarrow a \text{ as } |x| \xrightarrow{\geq} a \quad \text{and} \quad \min_{t \in \mathbb{R}} |\xi_x(t)| \rightarrow b \text{ as } |x| \xrightarrow{\leq} b.$$

Since the function Δu is constant along any streamline of the flow from the Euler equations (1.1) and since Δu is uniformly continuous in $\overline{\Omega_{a,b}}$, it follows from the previous observations that Δu is constant on C_a and constant on C_b . Let d_1 and d_2 be the values of Δu on C_a and C_b , respectively, and set $f(c_{in}) = -d_1$ and $f(c_{out}) = -d_2$. One then infers from (3.3)–(3.4) that $f : [c_{in}, c_{out}] \rightarrow \mathbb{R}$ is continuous in $[c_{in}, c_{out}]$ and the equation $\Delta u + f(u) = 0$ holds in the closed annulus $\overline{\Omega_{a,b}}$ (so u is a classical $C^2(\overline{\Omega_{a,b}})$ solution of (3.1)). However, since

$$f'(\tau) = -\frac{\nabla(\Delta u)(\sigma_y(g^{-1}(\tau))) \cdot \nabla u(\sigma_y(g^{-1}(\tau)))}{|\nabla u(\sigma_y(g^{-1}(\tau)))|^2} \quad \text{for all } \tau \in (c_{in}, c_{out})$$

and since $|\nabla u(\sigma_y(g^{-1}(\tau)))|$ may converge to 0 as $\tau \rightarrow c_{\text{in}}$ or c_{out} (this happens if $|v| = 0$ on C_a or if $|v| = 0$ on C_b), the function f' may be unbounded in $(c_{\text{in}}, c_{\text{out}})$.⁷ The argument used in the proof of Theorem 1.2 to conclude that the solution u of (3.1) is radially symmetric relies on [27, Theorem 5], which itself uses the Lipschitz continuity of f over the range of u . Thus, the same argument cannot be applied in general when v is just assumed to have no stagnation point in $\Omega_{a,b}$, without the assumption (1.3). Other arguments should then be used to prove that v is circular or to disprove this property in general. We leave this question for further work.

3.2. The case of unbounded annuli $\Omega_{a,\infty}$: proofs of Theorems 1.3 and 1.4

Throughout this section, we fix a positive real number a and we consider a $C^2(\overline{\Omega_{a,\infty}})$ flow v solving (1.1)–(1.2) and such that

$$\{x \in \overline{\Omega_{a,\infty}} : |v(x)| = 0\} \subsetneq C_a \quad \text{and} \quad |v| \geq \eta > 0 \text{ in } \overline{\Omega_{a+1,\infty}} \tag{3.5}$$

for some positive real number $\eta > 0$ (these conditions are fulfilled in both Theorems 1.3 and 1.4). We also assume that either $v(x) \cdot e_r(x) = o(1/|x|)$ as $|x| \rightarrow +\infty$ (that is, (1.8) holds, for Theorem 1.3) or $\inf_{\Omega_{a,\infty}} |v| > 0$ (for Theorem 1.4). This situation fits into the framework of Section 2, with $\mathcal{C}_1 = C_a$, $\omega_1 = B_a$, $\Omega = \mathbb{R}^2 \setminus \overline{\omega_1} = \Omega_{a,\infty}$, and $D = \overline{\Omega_{a,\infty}}$. Notice also that conditions (2.2)–(2.3) and (2.15) are fulfilled by assumption. So is (2.17), hence Lemma 2.4 (ii) can be applied. Furthermore, either (2.5) is fulfilled (from (1.8), for Theorem 1.3) and Lemmas 2.1 (ii) and 2.8 (i) can be applied, or the conditions of Lemmas 2.6 (iii) and 2.8 (ii) are fulfilled. Therefore, from Lemmas 2.1, 2.4 (ii), 2.6 (iii) and 2.8, the flow has a $C^3(\overline{\Omega_{a,\infty}})$ stream function u , and there exist a real number c_{in} (which can be taken to be 0, since u is unique up to additive constants) and $c_{\text{out}} = \pm\infty$ (we can assume that $c_{\text{out}} = +\infty$, even if it means changing v to $-v$ and u to $-u$), together with a $C^1([0, +\infty))$ function f such that

$$\begin{cases} \Delta u + f(u) = 0 \text{ in } \overline{\Omega_{a,\infty}}, \\ u > 0 \text{ in } \Omega_{a,\infty}, \\ u = 0 \text{ on } C_a, \quad u(x) \rightarrow +\infty \text{ as } |x| \rightarrow +\infty. \end{cases} \tag{3.6}$$

3.2.1. Proof of Theorem 1.3. In addition to (3.5), we further assume that

$$v(x) \cdot e_r(x) = o(1/|x|) \quad \text{as } |x| \rightarrow +\infty. \tag{3.7}$$

⁷For instance, the smooth flow $v(x) = (|x| - a)e_\theta(x)$ solves (1.1)–(1.2) with pressure $p(x) = |x|^2/2 - 2a|x| + a^2 \ln|x|$ and stream function $u(x) = (|x| - a)^2/2$ (up to additive constants), while $|v| = 0$ on C_a and $|v| > 0$ in $\Omega_{a,b}$. Here, $c_{\text{in}} = 0$, $c_{\text{out}} = (b - a)^2/2$ and $f(s) = -2 + a/(a + \sqrt{2s})$ for $s \in [c_{\text{in}}, c_{\text{out}}] = [0, (b - a)^2/2]$, hence f' is not bounded in $(c_{\text{in}}, c_{\text{out}})$. Notice that this example is a circular flow, which makes question (3.2) still relevant.

As a consequence, (2.5) holds, and there is $R_0 \geq a + 1$ such that

$$|\nabla u \cdot e_r| = |v \cdot e_\theta| \geq \eta/2 \quad \text{in } \overline{\Omega_{R_0, \infty}}. \tag{3.8}$$

Hence, Lemma 2.6 (i, ii) can be applied and the streamlines $\Xi_x = \xi_x(\mathbb{R})$ (with $x \in \Omega_{a, \infty}$) surround the origin and are such that

$$\min_{\mathbb{R}} |\xi_x| \rightarrow +\infty \quad \text{and} \quad \max_{\mathbb{R}} |\xi_x| - \min_{\mathbb{R}} |\xi_x| \rightarrow 0 \quad \text{as } |x| \rightarrow +\infty \tag{3.9}$$

(actually, the second conclusion is stronger than the first, since $\xi_x(0) = x$). Together with the normalization of u (that is, $u > 0$ in $\Omega_{a, \infty}$), properties (3.8)–(3.9) yield the existence of $R_1 \geq R_0 \geq a + 1$ such that

$$\nabla u(x) \cdot e_r(x) \geq \eta/2 \text{ for all } |x| \geq R_0, \quad \min_{\mathbb{R}} |\xi_x| \geq R_0 \text{ for all } |x| \geq R_1.$$

For any x with $|x| \geq R_1$, it then follows that, for every $\theta \in \mathbb{R}$, there is a unique $\varrho_x(\theta) \geq R_0$ such that $(\varrho_x(\theta) \cos \theta, \varrho_x(\theta) \sin \theta) \in \Xi_x$, and moreover

$$\Xi_x = \{(\varrho_x(\theta) \cos \theta, \varrho_x(\theta) \sin \theta) : \theta \in \mathbb{R}\}. \tag{3.10}$$

Notice also that the 2π -periodic function ϱ_x is of class $C^3(\mathbb{R})$ from the implicit function theorem.

If some streamlines were true circles centered at the origin, then [27, Theorem 5] would imply that the stream function u is radially symmetric in the bounded region between C_a and these streamlines. To circumvent the fact that the streamlines are not known to be true circles a priori, we use Lemma 3.1 below and Proposition 1.14 to compare the stream function u with its reflection with respect to some lines approximating any line containing the origin. We then proceed by passing to the limit as the approximation parameter goes to 0. With Proposition 1.14, it then easily follows that u is radially symmetric and that all streamlines are truly circular, thus completing the proof of Theorem 1.3.

To apply this strategy, let us now introduce some additional notations which will be used in this section, as well as in the proof of Theorems 1.4, 1.5, 1.8 and 1.10 in the following sections. For $x \in \Omega_{a, \infty}$, let Ω_x denote the bounded connected component of $\mathbb{R}^2 \setminus \Xi_x$. Notice that Ω_x is well defined and contains the origin, by Lemma 2.6 (i). Notice also that u is equal to the positive constant $u(x)$ along Ξ_x , while u vanishes along C_a and has no critical point in $\Omega_{a, \infty}$. Hence,

$$0 < u(y) < u(x) \quad \text{for all } y \in \Omega_x \cap \Omega_{a, \infty}, \tag{3.11}$$

where $\Omega_x \cap \Omega_{a, \infty}$ is the bounded domain located between Ξ_x and C_a . As a consequence, $\nabla u(z)$ points outwards from Ω_x at each point $z \in \Xi_x$.

Recalling that the sets $T_{e, \lambda}$ and $H_{e, \lambda}$ and the reflection $R_{e, \lambda}$ have been defined in (1.15)–(1.16), the following lemma says that, for any $\varepsilon > 0$, the set $\Omega_x \cap H_{e, \lambda}$ will be an admissible set for the method of moving planes for any $e \in \mathbb{S}^1$ and $\lambda > \varepsilon > 0$, provided $|x|$ is large enough.




Fig. 2. The sets Ω_x , $H_{e,\lambda} \cap \overline{\Omega_x}$ and $R_{e,\lambda}(H_{e,\lambda} \cap \overline{\Omega_x})$.

Lemma 3.1. For each $\varepsilon > 0$, there exists $R_\varepsilon > a$ such that

$$R_{e,\lambda}(H_{e,\lambda} \cap \overline{\Omega_x}) \subset \Omega_x$$

for all $e \in \mathbb{S}^1$, $\lambda > \varepsilon$ and $|x| \geq R_\varepsilon$ (see Fig. 2).

Proof. Fix $\varepsilon > 0$, and assume that the conclusion does not hold. Then there are sequences $(x_n)_{n \in \mathbb{N}}$ in $\Omega_{a,\infty}$, $(e_n)_{n \in \mathbb{N}}$ in \mathbb{S}^1 , $(\lambda_n)_{n \in \mathbb{N}}$ in $(\varepsilon, +\infty)$ and $(y_n)_{n \in \mathbb{N}}$ such that

$$\lim_{n \rightarrow +\infty} |x_n| = +\infty, \text{ and } y_n \in H_{e_n, \lambda_n} \cap \overline{\Omega_{x_n}} \text{ and } z_n := R_{e_n, \lambda_n}(y_n) \notin \Omega_{x_n} \text{ for all } n \in \mathbb{N}.$$

From (3.9), there is a sequence $(r_n)_{n \in \mathbb{N}}$ of positive real numbers converging to 0 such that $B_{|x_n| - r_n} \subset \Omega_{x_n} \subset B_{|x_n| + r_n}$ for all $n \in \mathbb{N}$, hence $|y_n| \leq |x_n| + r_n$. On the other hand, since $y_n \cdot e_n > \lambda_n > \varepsilon > 0$, one has

$$|y_n|^2 - |z_n|^2 = |y_n|^2 - |R_{e_n, \lambda_n}(y_n)|^2 = 4\lambda_n(y_n \cdot e_n - \lambda_n) > 0,$$

hence $|y_n| > |z_n| \geq |x_n| - r_n$ since $z_n \notin \Omega_{x_n}$. As a consequence, $|x_n| - r_n \leq |z_n| < |y_n| \leq |x_n| + r_n$ for all $n \in \mathbb{N}$, and $\lim_{n \rightarrow +\infty} (|y_n| - |x_n|) = \lim_{n \rightarrow +\infty} (|y_n| - |z_n|) = 0$. The inequality $|y_n|^2 - |z_n|^2 = 4\lambda_n(y_n \cdot e_n - \lambda_n) > 4\varepsilon(y_n \cdot e_n - \lambda_n) > 0$ yields $\lim_{n \rightarrow +\infty} (y_n \cdot e_n - \lambda_n) = 0$. Hence,

$$\text{dist}(y_n, \Xi_{x_n} \cap T_{e_n, \lambda_n}) \rightarrow 0 \text{ and } |y_n - z_n| \rightarrow 0 \text{ as } n \rightarrow +\infty.$$

For each $n \in \mathbb{N}$, let $\varphi_n \in \mathbb{R}$ be such that $e_n = (\cos \varphi_n, \sin \varphi_n)$. Since $y_n \cdot e_n > \lambda_n > \varepsilon > 0$, there is a unique $\theta_n \in (-\pi/2, \pi/2)$ such that

$$y_n/|y_n| = (\cos(\varphi_n + \theta_n), \sin(\varphi_n + \theta_n)).$$

Similarly, since $(z_n - y_n) \cdot e_n \rightarrow 0$ as $n \rightarrow +\infty$, one has $z_n \cdot e_n > \varepsilon/2$ for all large n and there is a unique $\theta'_n \in (-\pi/2, \pi/2)$ such that

$$z_n/|z_n| = (\cos(\varphi_n + \theta'_n), \sin(\varphi_n + \theta'_n)).$$




Fig. 3. The points y_n, y'_n, z_n, z'_n , and ζ_n (here $e_n = (1, 0)$ and $\varphi_n = 0$).

Since $\lim_{n \rightarrow +\infty} |y_n - z_n| = 0$ and $\lim_{n \rightarrow +\infty} |y_n| = \lim_{n \rightarrow +\infty} |z_n| = \lim_{n \rightarrow +\infty} |x_n| = +\infty$, one also infers that $\theta_n - \theta'_n \rightarrow 0$ as $n \rightarrow +\infty$. We also recall that (3.10) holds with x_n instead of x , for all n large enough. It then follows from Lemma 2.6 (i) and the assumptions on y_n and z_n that $|y_n| \leq \varrho_{x_n}(\varphi_n + \theta_n)$ and $|z_n| \geq \varrho_{x_n}(\varphi_n + \theta'_n)$ for all n large enough. Denote, for n large enough,

$$y'_n = (\varrho_{x_n}(\varphi_n + \theta_n) \cos(\varphi_n + \theta_n), \varrho_{x_n}(\varphi_n + \theta_n) \sin(\varphi_n + \theta_n)) \in \Xi_{x_n},$$

$$z'_n = (\varrho_{x_n}(\varphi_n + \theta'_n) \cos(\varphi_n + \theta'_n), \varrho_{x_n}(\varphi_n + \theta'_n) \sin(\varphi_n + \theta'_n)) \in \Xi_{x_n},$$

and observe that $y_n \in (0, y'_n]$ and $z'_n \in (0, z_n]$.

We now claim that $\theta'_n \neq \theta_n$ for all n large enough. Indeed, otherwise, up to extraction of a subsequence, $y'_n = z'_n$ and the four points $0, y_n, y'_n = z'_n$ and z_n would be aligned in that order. But since $y_n - z_n = 2(y_n \cdot e_n - \lambda_n)e_n$ with $y_n \cdot e_n - \lambda_n > 0$, the vectors y_n and z_n would be parallel to e_n . Hence, $y_n = (y_n \cdot e_n)e_n$ with $y_n \cdot e_n > \lambda_n > \varepsilon > 0$ and $z_n = (z_n \cdot e_n)e_n$ with $z_n \cdot e_n = 2\lambda_n - y_n \cdot e_n < \lambda_n < y_n \cdot e_n$. This contradicts the fact that $0, y_n$ and z_n lie on the half-line \mathbb{R}_+e_n in that order. Thus, $\theta'_n \neq \theta_n$ for all n large enough, so for all n without loss of generality. Notice that the same arguments also imply that $\theta_n \neq 0$ and $\theta'_n \neq 0$ for all n large enough (since otherwise in either case one would have $\theta_n = \theta'_n = 0$ up to extraction of a subsequence), thus for all n without loss of generality. In particular, either $0 < \theta_n < \pi/2$ or $-\pi/2 < \theta_n < 0$.

Assume first that, up to extraction of a subsequence, $0 < \theta_n < \pi/2$ for all n . One then infers from the definition of $z_n = R_{e_n, \lambda_n}(y_n)$ and the previous paragraph that

$$0 < \theta_n < \theta'_n < \pi/2.$$

Remember now that $0 < u(y) < u(x_n)$ for every y in the domain $\Omega_{x_n} \cap \Omega_{a, \infty}$ between Ξ_{x_n} and C_a , and $\nabla u(z)$ points outwards from Ω_{x_n} at each point $z \in \Xi_{x_n}$. For each $n \in \mathbb{N}$, since $y_n \in (0, y'_n], z'_n \in (0, z_n]$ and $y_n - z_n = \zeta_n e_n$ with $\zeta_n := 2(y_n \cdot e_n - \lambda_n) > 0$, there is an angle $\phi_n \in [\theta_n, \theta'_n] \subset (0, \pi/2)$ such that

$$\zeta_n := (\varrho_{x_n}(\varphi_n + \phi_n) \cos(\varphi_n + \phi_n), \varrho_{x_n}(\varphi_n + \phi_n) \sin(\varphi_n + \phi_n)) \in \Xi_{x_n} \cap [y_n, z_n]$$

and $\nabla u(\zeta_n) \cdot e_n \leq 0$ (see Fig. 3). The point ζ_n can be defined as the first point on $\Xi_{x_n} \cap [y_n, z_n]$ when going from y'_n to z'_n along Ξ_{x_n} with increasing angle. Notice that $|\zeta_n| \rightarrow +\infty$ since $|y_n| \rightarrow +\infty$ and $|y_n - z_n| \rightarrow 0$ as $n \rightarrow +\infty$. Let $v_{1,n} = v(\zeta_n) \cdot e_n$ and $v_{2,n} = v(\zeta_n) \cdot e_n^\perp$. The inequality $\nabla u(\zeta_n) \cdot e_n \leq 0$ means that $v_{2,n} \leq 0$. Therefore,

$$v(\zeta_n) \cdot e_r(\zeta_n) = v_{1,n} \cos \phi_n + v_{2,n} \sin \phi_n \leq v_{1,n} \cos \phi_n, \tag{3.12}$$

while $0 < \eta/2 \leq |v(\zeta_n) \cdot e_\theta(\zeta_n)| = |-v_{1,n} \sin \phi_n + v_{2,n} \cos \phi_n|$ for all n large enough, from (3.8) and $\lim_{n \rightarrow +\infty} |\zeta_n| = +\infty$. But since the continuous function $v \cdot e_\theta = \nabla u \cdot e_r$ has a constant strict positive sign at infinity, it follows that

$$\eta/2 \leq v(\zeta_n) \cdot e_\theta(\zeta_n) = -v_{1,n} \sin \phi_n + v_{2,n} \cos \phi_n$$

for all n large enough. Since $v_{2,n} \leq 0$ and $0 < \phi_n < \pi/2$, one gets $-v_{1,n} \sin \phi_n \geq \eta/2$, hence $v_{1,n} \leq -\eta/2$. Together with (3.12), it follows that $v(\zeta_n) \cdot e_r(\zeta_n) \leq -(\eta/2) \cos \phi_n$ for all n large enough. On the other hand, since $\zeta_n \in [y_n, z_n]$ and $\lim_{n \rightarrow +\infty} |z_n - y_n| = \lim_{n \rightarrow +\infty} (y_n \cdot e_n - \lambda_n) = 0$, we have $\zeta_n \cdot e_n - \lambda_n \rightarrow 0$, hence $\zeta_n \cdot e_n \geq \varepsilon/2$ for all n large enough (since $\lambda_n > \varepsilon > 0$ for all n). Finally,

$$\cos \phi_n = \frac{\zeta_n \cdot e_n}{|\zeta_n|} \geq \frac{\varepsilon}{2|\zeta_n|} \quad \text{and} \quad v(\zeta_n) \cdot e_r(\zeta_n) \leq -\frac{\eta\varepsilon}{4|\zeta_n|}$$

for all n large enough. That last inequality contradicts the assumption (3.7) and the limit $\lim_{n \rightarrow +\infty} |\zeta_n| = +\infty$.

The second case, for which, up to extraction of a subsequence, $-\pi/2 < \theta_n < 0$ for all n (and so $-\pi/2 < \theta'_n < \theta_n < 0$) can be handled similarly and leads to a contradiction as well. The proof of Lemma 3.1 is thereby complete. ■

Proof of Theorem 1.3. We shall show that the stream function u is radially symmetric in $\overline{\Omega_{a,\infty}}$. Notice that we already know that $u = 0$ on C_a . Let then $x \neq y \in \Omega_{a,\infty}$ be such that $|x| = |y| (> a)$, and set

$$e = \frac{y - x}{|y - x|} \in \mathbb{S}^1. \tag{3.13}$$

Pick $0 < \varepsilon < a$. Let $R_\varepsilon > a$ be as in Lemma 3.1. From (3.9), there is a point $x_\varepsilon \in \Omega_{a,\infty}$ such that $|x_\varepsilon| \geq R_\varepsilon$ and $\min_{\mathbb{R}} |\xi_{x_\varepsilon}| > |x| = |y|$. Lemma 3.1 then yields

$$R_{e,\lambda}(H_{e,\lambda} \cap \overline{\Omega_{x_\varepsilon}}) \subset \Omega_{x_\varepsilon} \quad \text{for all } \lambda > \varepsilon. \tag{3.14}$$

We are now going to apply Proposition 1.14 with

$$\left\{ \begin{array}{l} \Xi = \Xi_{x_\varepsilon}, \quad \Omega = \Omega_{x_\varepsilon}, \quad \Xi' = C_a, \quad \Omega' = B_a, \quad \omega = \Omega_{x_\varepsilon} \setminus \overline{B_a}, \\ R' = a, \quad R = \max_{\mathbb{R}} |\xi_{x_\varepsilon}| > a, \quad \bar{\lambda} = \max_{z \in \Xi_{x_\varepsilon}} z \cdot e > a > \varepsilon > 0, \\ \varphi = -u \in C^3(\overline{\omega}), \quad c_1 = -u(x_\varepsilon) = -u|_{\Xi_{x_\varepsilon}} < 0, \quad c_2 = 0 = -u|_{C_a}, \\ F(r, s) = F(s) = -f(-s) \quad \text{for } (r, s) \in [a, R] \times [-u(x_\varepsilon), 0]. \end{array} \right.$$

Notice immediately that assumption (1.19) is automatically satisfied. The function F clearly satisfies the assumptions of Proposition 1.14 since f is of class $C^1([0, +\infty))$. The function φ satisfies $\Delta\varphi + F(\varphi) = 0$ in $\bar{\omega}$, with $c_1 < \varphi < c_2$ in ω (since $0 < u < u(x_\varepsilon)$ in $\Omega_{x_\varepsilon} \cap \Omega_{a,\infty}$ by (3.11)). In view of (3.14), all assumptions of Proposition 1.14 are satisfied. Proposition 1.14 applied with $\lambda = \varepsilon$ implies that $\varphi \leq \varphi_{e,\varepsilon}$, so $u \geq u_{e,\varepsilon}$ in $\overline{\omega_{e,\varepsilon}}$ with

$$\omega_{e,\varepsilon} = (H_{e,\varepsilon} \cap \omega) \setminus R_{e,\varepsilon}(\overline{\Omega'}) = (H_{e,\varepsilon} \cap (\Omega_{x_\varepsilon} \setminus \overline{B_a})) \setminus R_{e,\varepsilon}(\overline{B_a}).$$

Observe now that $y \cdot e = (|y|^2 - x \cdot y)/|y - x| > 0$ since $|x| = |y|$ and $x \neq y$, and remember that $a < |y| < \min_{\mathbb{R}} |\xi_{x_\varepsilon}|$, hence $y \in \omega$. Therefore, $y \in \omega_{e,\varepsilon}$ for all $\varepsilon > 0$ small enough, and

$$u(y) \geq u_{e,\varepsilon}(y) = u(y_{e,\varepsilon}) = u(y - 2(y \cdot e - \varepsilon)e)$$

for all $\varepsilon > 0$ small enough. By passing to the limit $\varepsilon \xrightarrow{\geq} 0$ and using the definition of e and the assumption $|x| = |y|$, one infers that

$$u(y) \geq u(y - 2(y \cdot e)e) = u(x).$$

Since the last inequality holds for all $x \neq y \in \Omega_{a,\infty}$ such that $|x| = |y|$ (and also for all $x, y \in C_a$), the $C^3(\overline{\Omega_{a,\infty}})$ function u is radially symmetric in $\overline{\Omega_{a,\infty}}$. By (3.5)–(3.6), there is then a $C^3([a, +\infty))$ function U such that $U(a) = 0$, $U' > 0$ in $[a, +\infty)$ and $u(x) = U(|x|)$ for all $x \in \overline{\Omega_{a,\infty}}$. This means that $v(x) = V(|x|)e_\theta(x)$ for all $x \in \overline{\Omega_{a,\infty}}$ with $V = U' \in C^2([a, +\infty))$ and $V > 0$ in $[a, +\infty)$. The proof of Theorem 1.3 is thereby complete. ■

3.2.2. *Proof of Theorem 1.4.* In addition to (3.5), we actually assume the stronger condition

$$|v| = |\nabla u| \geq \eta > 0 \quad \text{in } \overline{\Omega_{a,\infty}} \tag{3.15}$$

for some $\eta > 0$. Since from our normalization the function u is positive in $\Omega_{a,\infty}$ and vanishes on C_a , conditions (1.2) and (3.15) imply that

$$v \cdot e_\theta = \nabla u \cdot e_r \geq \eta > 0 \quad \text{on } C_a. \tag{3.16}$$

To prove Theorem 1.4, we have to show that the supremum of the vorticity is positive:

$$\sup_{\Omega_{a,\infty}} \left(\frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2} \right) > 0. \tag{3.17}$$

Assume by way of contradiction that

$$\Delta u = \frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2} \leq 0 \quad \text{in } \overline{\Omega_{a,\infty}}. \tag{3.18}$$

We will show that u is radially symmetric, and this will easily lead to a contradiction. To prove the radial symmetry of u , let us use a Kelvin transform of the variables by setting

$$w(x) = u(x/|x|^2) \quad \text{for } x \in \overline{\Omega_{0,1/a}} \setminus \{0\},$$

and let us show that the $C^3(\overline{\Omega_{0,1/a}} \setminus \{0\})$ function w is radially symmetric in $\overline{\Omega_{0,1/a}} \setminus \{0\}$. From (3.6), one has

$$w = 0 \text{ on } C_{1/a}, \quad w > 0 \text{ in } \Omega_{0,1/a} \quad \text{and} \quad w(x) \rightarrow +\infty \text{ as } |x| \xrightarrow{\geq} 0,$$

and a straightforward calculation yields

$$\Delta w(x) + \frac{1}{|x|^4} f(w(x)) = 0 \quad \text{for all } x \in \overline{\Omega_{0,1/a}} \setminus \{0\},$$

that is, $\Delta w(x) + F(|x|, w(x)) = 0$ in $\overline{\Omega_{0,1/a}} \setminus \{0\}$ with

$$F : (0, 1/a] \times [0, +\infty) \rightarrow \mathbb{R}, \quad (r, s) \mapsto F(r, s) = r^{-4} f(s).$$

The function F is of class $C^1((0, 1/a] \times [0, +\infty))$. Furthermore, the range of u is the whole interval $[0, +\infty)$ by (3.6), and $f \geq 0$ in $[0, +\infty)$ by (3.6) and (3.18). Therefore, F is nonincreasing with respect to its first variable in $(0, 1/a] \times [0, +\infty)$.

Consider now any two points $x \neq y \in \overline{\Omega_{0,1/a}} \setminus \{0\}$ with $|x| = |y|$. As in the proof of Theorem 1.3, denote

$$e = \frac{y - x}{|y - x|} \in \mathbb{S}^1$$

and consider an arbitrary real number ε such that

$$0 < \varepsilon < |x| = |y| \leq 1/a.$$

By Lemma 2.6 (iii), there is a point $x_\varepsilon \in \Omega_{a,\infty}$ such that $\min_{\mathbb{R}} |\xi_{x_\varepsilon}| > 1/\varepsilon > a$. One knows that the streamline Ξ_{x_ε} surrounds the origin and that $u = u(x_\varepsilon) > 0$ along Ξ_{x_ε} . Furthermore, as in (3.11), one has $0 < u < u(x_\varepsilon)$ in the domain $\Omega_{x_\varepsilon} \cap \Omega_{a,\infty}$ between Ξ_{x_ε} and C_a , since $u = 0$ on C_a and u has no critical point in $\Omega_{a,\infty}$.

Denote $\Xi = C_{1/a}$ and

$$\Xi' = \{z \in \mathbb{R}^2 : z/|z|^2 \in \Xi_{x_\varepsilon}\}.$$

Notice that the Jordan curve Ξ' surrounds the origin and $\Xi' \subset B_\varepsilon (\subset B_{1/a})$ by definition of x_ε . Set $\Omega = B_{1/a}$, let Ω' be the bounded connected component of $\mathbb{R}^2 \setminus \Xi'$, and let

$$\omega = \Omega \setminus \overline{\Omega'} = B_{1/a} \setminus \overline{\Omega'} (\supset \Omega_{\varepsilon,1/a}).$$

Denote $R = 1/a$,

$$0 < R' = \min_{z \in \Xi'} |z| = \frac{1}{\max_{\mathbb{R}} |\xi_{x_\varepsilon}|} < \varepsilon < R,$$

and $\bar{\lambda} = \max_{z \in \Xi} z \cdot e = 1/a > 0$. One has $0 < \varepsilon < 1/a$, hence $\varepsilon \in [0, \bar{\lambda})$. The function $\varphi = w$ is of class $C^3(\bar{\omega})$ with

$$\varphi = c_1 = 0 \text{ on } \Xi = C_{1/a}, \quad \varphi = c_2 = u(x_\varepsilon) > 0 \text{ on } \Xi', \quad \text{and} \quad 0 < \varphi < u(x_\varepsilon) \text{ in } \omega$$

(since $0 < u < u(x_\varepsilon)$ in $\Omega_{x_\varepsilon} \cap \Omega_{a,\infty}$). Furthermore, φ satisfies $\Delta\varphi + F(|x|, \varphi) = 0$ in $\overline{\omega}$ with $F(r, s) = r^{-4}f(s)$ and $(r, s) \in [R', 1/a] \times [0, u(x_\varepsilon)]$. The function F then satisfies the conditions of Proposition 1.14. Lastly, condition (1.18) is immediately satisfied since $\Omega = B_{1/a}$, and (1.19) also holds since $H_{e,\lambda} \cap \Xi' = \emptyset$ for all $\lambda > \varepsilon$ (because $\Xi' \subset B_\varepsilon$). To sum up, all assumptions of Proposition 1.14 are fulfilled. Its conclusion with $\lambda = \varepsilon$ yields $w \leq w_{e,\varepsilon}$ in $\overline{\omega_{e,\varepsilon}}$ with

$$\omega_{e,\varepsilon} = (H_{e,\varepsilon} \cap (B_{1/a} \setminus \overline{\Omega'})) \setminus R_{e,\varepsilon}(\overline{\Omega'}).$$

Since $y \cdot e > 0$ and since $\overline{\Omega'} \subset B_\varepsilon$ and $R_{e,\varepsilon}(\overline{\Omega'}) \subset B_{3\varepsilon}$, it follows that $y \in \omega_{e,\varepsilon}$ for all $\varepsilon > 0$ small enough. As a consequence, $w(y) \leq w_{e,\varepsilon}(y) = w(y_{e,\varepsilon}) = w(y - 2(y \cdot e - \varepsilon)e)$ for all $\varepsilon > 0$ small enough and the passage to the limit as $\varepsilon \rightarrow 0$ yields

$$w(y) \leq w(y - 2(y \cdot e)e) = w(x)$$

by definition of e . Since this holds for all $x \neq y \in \overline{\Omega_{0,1/a}} \setminus \{0\}$ with $|x| = |y|$, this means that w is radially symmetric in $\overline{\Omega_{0,1/a}} \setminus \{0\}$, hence u is radially symmetric in $\overline{\Omega_{a,\infty}}$. In view of (3.16), there is then a $C^3([a, +\infty))$ function U such that $u(x) = U(|x|)$ and $U' \geq \eta > 0$ in $[a, +\infty)$. But $\Delta u \leq 0$ in $\overline{\Omega_{a,\infty}}$ by (3.18). Hence $U''(r) + r^{-1}U'(r) \leq 0$ in $[a, +\infty)$ and the function $r \mapsto rU'(r)$ is nonincreasing in $[a, +\infty)$, a contradiction with $U' \geq \eta > 0$.

As a conclusion, (3.18) cannot hold, that is, (3.17) holds and the proof of Theorem 1.4 is thereby complete. ■

3.3. The case of punctured disks $\Omega_{0,b}$: proof of Theorem 1.5

Throughout this section, we fix a positive real number b and we consider a $C^2(\overline{\Omega_{0,b}} \setminus \{0\})$ flow v solving (1.1)–(1.2) and such that

$$\{x \in \overline{\Omega_{0,b}} \setminus \{0\} : |v(x)| = 0\} \subsetneq C_b \quad \text{and} \quad \int_{C_\varepsilon} |v \cdot e_r| \rightarrow 0 \text{ as } \varepsilon \xrightarrow{\geq} 0. \quad (3.19)$$

This situation falls within the framework of Section 2, with $\mathcal{C}_2 = C_b$, $\omega_2 = B_b$, $\Omega = \omega_2 \setminus \{0\} = \Omega_{0,b}$, and $D = \overline{\Omega_{0,b}} \setminus \{0\}$, and conditions (2.2)–(2.3), (2.5), and (2.15) are fulfilled by assumption. Therefore, by Lemmas 2.1 and 2.8 (i), the flow has a $C^3(\overline{\Omega_{0,b}} \setminus \{0\})$ stream function u , and there exist a real number c_{out} (which can be taken to be 0, since u is unique up to additive constants) and $0 \neq c_{\text{in}} \in \overline{\mathbb{R}}$ (we can assume that $0 < c_{\text{in}} \leq +\infty$, even if it means changing v to $-v$ and u to $-u$), together with a $C^1([0, c_{\text{in}}))$ function f such that

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \overline{\Omega_{0,b}} \setminus \{0\}, \\ u > 0 & \text{in } \Omega_{0,b}, \\ u = 0 & \text{on } C_b, \quad u(x) \rightarrow c_{\text{in}} \quad \text{as } |x| \xrightarrow{\geq} 0. \end{cases} \quad (3.20)$$

One goal is to show that u is radially symmetric in $\overline{\Omega_{0,b}} \setminus \{0\}$. Consider any two points $x \neq y \in \overline{\Omega_{0,b}} \setminus \{0\}$ with $|x| = |y|$. As in the proofs of Theorems 1.3 and 1.4, denote

$$e = \frac{y - x}{|y - x|} \in \mathbb{S}^1$$

and consider an arbitrary real number ε such that

$$0 < \varepsilon < |x| = |y| \leq b.$$

By Lemma 2.6 (i), there is a point $x_\varepsilon \in \Omega_{0,b}$ such that $\max_{\mathbb{R}} |\xi_{x_\varepsilon}| < \varepsilon < b$, and the streamline Ξ_{x_ε} surrounds the origin with $u = u(x_\varepsilon) > 0$ along Ξ_{x_ε} . Since $u = 0$ on C_b and u has no critical point in $\Omega_{0,b}$, one infers that $0 < u < u(x_\varepsilon)$ in the domain ω between Ξ_{x_ε} and C_b . Denote $\Xi = C_b$, $\Xi' = \Xi_{x_\varepsilon}$, $\Omega = B_b$, let $\Omega' = \Omega_{x_\varepsilon}$ be the bounded connected component of $\mathbb{R}^2 \setminus \Xi_{x_\varepsilon}$, and notice that

$$\omega = \Omega \setminus \overline{\Omega'}.$$

Set

$$R = b, \quad 0 < R' = \min_{z \in \Xi'} |z| < \varepsilon < R \quad \text{and} \quad \bar{\lambda} = \max_{z \in \Xi} z \cdot e = b > 0.$$

One has $\varepsilon \in (0, \bar{\lambda})$. The function $\varphi = u$ is of class $C^3(\overline{\omega})$ with

$$\varphi = c_1 = 0 \text{ on } \Xi = C_b, \quad \varphi = c_2 = u(x_\varepsilon) > 0 \text{ on } \Xi', \quad \text{and} \quad 0 < \varphi < u(x_\varepsilon) \text{ in } \omega.$$

Furthermore, $\Delta\varphi + F(\varphi) = 0$ in $\overline{\omega}$, with $F : [R', b] \times [0, u(x_\varepsilon)] \ni (r, s) \mapsto F(r, s) = f(s)$ satisfying all conditions of Proposition 1.14. Lastly, condition (1.18) is immediately satisfied since $\Omega = B_b$, and (1.19) also holds since $H_{e,\lambda} \cap \Xi' = \emptyset$ for all $\lambda > \varepsilon$ (because $\Xi' \subset B_\varepsilon$). To sum up, all assumptions of Proposition 1.14 are fulfilled. Its conclusion with $\lambda = \varepsilon$ yields $u \leq u_{e,\varepsilon}$ in $\overline{\omega_{e,\varepsilon}}$, with

$$\omega_{e,\varepsilon} = (H_{e,\varepsilon} \cap (B_b \setminus \overline{\Omega'})) \setminus R_{e,\varepsilon}(\overline{\Omega'}).$$

Since $y \cdot e > 0$ and since $\overline{\Omega'} \subset B_\varepsilon$ and $R_{e,\varepsilon}(\overline{\Omega'}) \subset B_{3\varepsilon}$, it follows that $y \in \omega_{e,\varepsilon}$ for all $\varepsilon > 0$ small enough. As a consequence, $u(y) \leq u(y_{e,\varepsilon}) = u(y - 2(y \cdot e - \varepsilon)e)$ for all $\varepsilon > 0$ small enough and the passage to the limit as $\varepsilon \xrightarrow{\sim} 0$ yields

$$u(y) \leq u(y - 2(y \cdot e)e) = u(x)$$

by definition of e . Since this holds for all $x \neq y \in \overline{\Omega_{0,b}} \setminus \{0\}$ with $|x| = |y|$, this means that u is radially symmetric in $\overline{\Omega_{0,b}} \setminus \{0\}$. By (3.19)–(3.20), there is then a $C^3((0, b])$ function U such that $u(x) = U(|x|)$ and $U' < 0$ in $(0, b]$. Hence, $v(x) = U'(|x|)e_\theta(x)$ for all $x \in \overline{\Omega_{0,b}} \setminus \{0\}$, and the proof of Theorem 1.5 is thereby complete. ■

3.4. The case of the punctured plane $\Omega_{0,\infty} = \mathbb{R}^2 \setminus \{0\}$: proof of Theorem 1.8

Let v be a $C^2(\Omega_{0,\infty})$ flow solving (1.1) and such that $|v| > 0$ in $\Omega_{0,\infty}$, we have $\liminf_{|x| \rightarrow +\infty} |v(x)| > 0$, and (1.14) holds. This situation fits into the framework of Section 2, with $\Omega = D = \mathbb{R}^2 \setminus \{0\} = \Omega_{0,\infty}$, and conditions (2.2)–(2.3), (2.5), and (2.15) are fulfilled by assumption. Therefore, from Lemmas 2.1 and 2.8 (i), the flow has a $C^3(\Omega_{0,\infty})$ stream function u , and there exist $c_{\text{in}} \neq c_{\text{out}} \in \mathbb{R}$ (we can assume that $c_{\text{in}} > c_{\text{out}}$, even if

it means changing v to $-v$ and u to $-u$), together with a $C^1((c_{\text{out}}, c_{\text{in}}))$ function f such that

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \Omega_{0,\infty}, \\ c_{\text{out}} < u < c_{\text{in}} & \text{in } \Omega_{0,\infty}, \\ u(x) \rightarrow c_{\text{in}} \text{ as } |x| \xrightarrow{\geq} 0, & u(x) \rightarrow c_{\text{out}} \text{ as } |x| \rightarrow +\infty. \end{cases} \tag{3.21}$$

Moreover, (1.14) and $\liminf_{|x| \rightarrow +\infty} |v(x)| > 0$ yield $\liminf_{|x| \rightarrow +\infty} |v(x) \cdot e_\theta(x)| > 0$. It then follows from Lemma 2.6 (i, ii) that each streamline $\Xi_x = \xi_x(\mathbb{R})$ is a C^1 Jordan curve surrounding the origin, with

$$\max_{\mathbb{R}} |\xi_x| - \min_{\mathbb{R}} |\xi_x| \rightarrow 0 \text{ as } |x| \rightarrow +\infty \quad \text{and} \quad \max_{\mathbb{R}} |\xi_x| \rightarrow 0 \text{ as } |x| \xrightarrow{\geq} 0.$$

Since u is unique up to additive constants, one can also assume that $c_{\text{in}} > 0 > c_{\text{out}}$. Pick then a point $X \in \Omega_{0,\infty}$ such that $u(X) = 0$, and let Ω_X be the bounded connected component of $\mathbb{R}^2 \setminus \Xi_X$ (then $0 \in \Omega_X$). In view of the inequalities $c_{\text{in}} > 0 > c_{\text{out}}$ and the assumption $|\nabla u| > 0$ in $\Omega_{0,\infty}$, it follows that $u > 0$ in $\Omega_X \setminus \{0\}$ and $u < 0$ in $\mathbb{R}^2 \setminus \overline{\Omega_X}$. Furthermore, by definition of u , v is orthogonal to the normal vector to Ω_X . As a consequence, Lemma 2.6 (iii) can also be applied with $\omega_1 = \Omega_X$, hence $c_{\text{out}} = -\infty$ in (3.21).

Now, still using the notations (1.15)–(1.16), it follows as in Lemma 3.1, from the assumptions $\liminf_{|x| \rightarrow +\infty} |v(x)| > 0$ and $v(x) \cdot e_r(x) = o(1/|x|)$ as $|x| \rightarrow +\infty$, that, for every $\varepsilon > 0$, there is $R_\varepsilon > 0$ such that

$$R_{e,\lambda}(H_{e,\lambda} \cap \overline{\Omega_x}) \subset \Omega_x$$

for all $e \in \mathbb{S}^1, \lambda > \varepsilon$ and $|x| > R_\varepsilon$.

Lastly, let $x \neq y \in \Omega_{0,\infty}$ with $|x| = |y|$. Define $e \in \mathbb{S}^1$ as in (3.13) and let $0 < \varepsilon < |x| = |y|$. As in the proofs of Theorems 1.3 and 1.5, there are $x_\varepsilon \in \mathbb{R}^2 \setminus \overline{\Omega_X}$ and $x'_\varepsilon \in \Omega_X$ such that $\min_{\mathbb{R}} |\xi_{x_\varepsilon}| > |x| = |y| > \varepsilon$,

$$R_{e,\lambda}(H_{e,\lambda} \cap \overline{\Omega_{x_\varepsilon}}) \subset \Omega_{x_\varepsilon} \quad \text{for all } \lambda > \varepsilon, \tag{3.22}$$

and $\max_{\mathbb{R}} |\xi_{x'_\varepsilon}| < \varepsilon < |x| = |y|$. The streamlines $\Xi = \Xi_{x_\varepsilon}$ and $\Xi' = \Xi_{x'_\varepsilon}$ surround the origin, and u is equal to $c_1 = u(x_\varepsilon) < 0$ along Ξ and to $c_2 = u(x'_\varepsilon) > 0$ along Ξ' . Furthermore, $u(x_\varepsilon) < u < u(x'_\varepsilon)$ in the domain

$$\omega = \Omega_{x_\varepsilon} \setminus \overline{\Omega_{x'_\varepsilon}}$$

located between Ξ_{x_ε} and $\Xi_{x'_\varepsilon}$. Denote $R' = \min_{z \in \Xi'} |z| = \min_{\mathbb{R}} |\xi_{x'_\varepsilon}| \in (0, \varepsilon)$, $R = \max_{z \in \Xi} |z| = \max_{\mathbb{R}} |\xi_{x_\varepsilon}| > |x| = |y| > \varepsilon > R'$, and $\bar{\lambda} = \max_{z \in \Xi} z \cdot e > \min_{\mathbb{R}} |\xi_{x_\varepsilon}| > |x| = |y| > \varepsilon > 0$. The $C^3(\overline{\omega})$ function $\varphi = u$ satisfies (1.17) with $[R', R] \times [c_1, c_2] \ni (r, s) \mapsto F(r, s) = f(s)$ satisfying the assumptions of Proposition 1.14 since f is of class $C^1((-\infty, c_{\text{in}}))$. By (3.22) and the fact that $H_{e,\lambda} \cap \Xi' = \emptyset$ for all $\lambda > \varepsilon$ (since $\Xi' \subset B_\varepsilon$), assumptions (1.18)–(1.19) are satisfied. All assumptions of Proposition 1.14 are therefore fulfilled. Proposition 1.14 applied with $\lambda = \varepsilon$ implies that $u \leq u_{e,\varepsilon}$ in $\overline{\omega_{e,\varepsilon}}$ with

$$\omega_{e,\varepsilon} = (H_{e,\varepsilon} \cap (\Omega_{x_\varepsilon} \setminus \overline{\Omega_{x'_\varepsilon}})) \setminus R_{e,\varepsilon}(\overline{\Omega_{x'_\varepsilon}}).$$

As in the proof of Theorem 1.5, one has $y \in \omega_{e,\varepsilon}$ for all $\varepsilon > 0$ small enough, hence

$$u(y) \leq u_{e,\varepsilon}(y) = u(y_{e,\varepsilon}) = u(y - 2(y \cdot e - \varepsilon)e)$$

for all $\varepsilon > 0$ small enough. By passing to the limit as $\varepsilon \xrightarrow{\geq} 0$ and using the definition of e and the assumption $|x| = |y|$, one infers that $u(y) \leq u(y - 2(y \cdot e)e) = u(x)$. Since the last inequality holds for any $x \neq y \in \Omega_{0,\infty}$ such that $|x| = |y|$, the $C^3(\Omega_{0,\infty})$ function u is radially symmetric in $\Omega_{0,\infty}$. Since $|\nabla u| = |v| > 0$ in $\Omega_{0,\infty}$ and $c_{in} > c_{out} = -\infty$, there is then a $C^3((0, +\infty))$ function U such that $U' < 0$ in $(0, +\infty)$ and $u(x) = U(|x|)$ for all $x \in \Omega_{0,\infty}$. This means that $v(x) = V(|x|)e_\theta(x)$ for all $x \in \Omega_{0,\infty}$ with $V = U' \in C^2((0, +\infty))$ and $V < 0$ in $(0, +\infty)$. The proof of Theorem 1.8 is thereby complete. ■

4. Proofs of the Serrin-type Theorems 1.10 and 1.13

We start in Section 4.1 with the proof of Theorem 1.13 dealing with the case of doubly connected bounded domains, since the proof follows easily from the arguments used in the proof of Theorems 1.1–1.2 and from some known results of Reichel [21] and Sirakov [27] on elliptic overdetermined boundary value problems. Section 4.2 is then devoted to the proof of Theorem 1.10.

4.1. Proof of Theorem 1.13

Let $\omega_1, \omega_2, \Omega = \omega_2 \setminus \overline{\omega_1}$ and v be as in Theorem 1.13. Up to translation, one can assume that $0 \in \omega_1$, and the assumptions of Theorem 1.13 then fall within the framework of Section 2, with $\mathcal{C}_1 = \partial\omega_1, \mathcal{C}_2 = \partial\omega_2, \Omega = \omega_2 \setminus \overline{\omega_1}, D = \overline{\Omega} = \overline{\omega_2} \setminus \omega_1$, and conditions (2.2)–(2.3), (2.5), and (2.15) are fulfilled by assumption. Therefore, the flow has a $C^3(\overline{\Omega})$ stream function u , and by Lemmas 2.1 and 2.8(i) there are real numbers $c_{in} \neq c_{out}$ (one can again assume that $c_{in} > c_{out}$) and a $C^1([c_{out}, c_{in}])$ function f such that

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \overline{\Omega}, \\ c_{out} < u < c_{in} & \text{in } \Omega, \\ u = c_{in} \text{ on } \partial\omega_1, \quad u = c_{out} \text{ on } \partial\omega_2. \end{cases} \tag{4.1}$$

Since

$$\left| \frac{\partial u}{\partial n} \right| = |\nabla u| = |v| > 0 \quad \text{along } \partial\Omega,$$

where n denotes the outward unit normal on $\partial\Omega$, and since $|v|$ is constant along $\partial\omega_1$ and along $\partial\omega_2$, it follows that $\frac{\partial u}{\partial n}$ is also constant along $\partial\omega_1$ and along $\partial\omega_2$ (and $\frac{\partial u}{\partial n} > 0$ on $\partial\omega_1$ and $\frac{\partial u}{\partial n} < 0$ on $\partial\omega_2$). One concludes from [21, 27] (see also [2, 28]) that, up to translation, $\Omega = \Omega_{a,b}$ for some $0 < a < b < \infty$ and u is radially symmetric and decreasing with respect to $|x|$ in $\overline{\Omega} = \overline{\Omega_{a,b}}$. The assumptions and the conclusion of Theorem 1.1 are then satisfied, and the proof of Theorem 1.13 is complete. ■

4.2. Proof of Theorem 1.10

Let ω be a C^2 nonempty simply connected bounded domain of \mathbb{R}^2 (we here call this domain ω instead of Ω to distinguish it from the notations of Section 2, which are used below). Let $v \in C^2(\bar{\omega})$ satisfy the Euler equations (1.1) in $\bar{\omega}$. We assume that $v \cdot n = 0$ and $|v|$ is constant on $\partial\omega$, where n denotes the outward unit normal on $\partial\omega$, and v has a unique stagnation point in $\bar{\omega}$. Since ω is simply connected and v is divergence free, there is a $C^3(\bar{\omega})$ stream function u satisfying (1.4). Furthermore, u is constant along $\partial\omega$ since $v \cdot n = 0$ on $\partial\omega$. Up to normalization, one can assume that

$$u = 0 \quad \text{on } \partial\omega. \quad (4.2)$$

By assumption, u has a unique critical point in $\bar{\omega}$, and $|\nabla u| = |v|$ is constant along $\partial\omega$. Then $|\frac{\partial u}{\partial n}| = |\nabla u| = |v| > 0$ on $\partial\omega$. Up to changing v to $-v$ and u to $-u$, one can assume that

$$\frac{\partial u}{\partial n} = \gamma < 0 \quad \text{on } \partial\omega \quad (4.3)$$

for some negative constant γ . Hence, u has a unique maximum point in $\bar{\omega}$ (which is actually in ω) and this point is the unique critical point of u in $\bar{\omega}$. Up to translation, one can assume that this point is the origin 0. One also infers from the uniqueness of the critical point of u that

$$0 < u < u(0) \quad \text{for all } x \in \omega \setminus \{0\}. \quad (4.4)$$

This situation then falls within the framework of Section 2, with $\mathcal{C}_2 = \partial\omega$, $\omega_2 = \omega$, $\Omega = \omega \setminus \{0\}$, and $D = \bar{\omega} \setminus \{0\}$, and conditions (2.2)–(2.3), (2.5), and (2.15) are fulfilled by assumption (in particular, the integral condition (2.3) in the present punctured case is satisfied since here v is continuous at 0 and $|v(0)| = 0$). Therefore, from Lemma 2.8 (i) and the previous notations, there exists a $C^1([0, u(0)])$ function f such that $\Delta u + f(u) = 0$ in $\bar{\omega} \setminus \{0\}$. Furthermore, by setting $f(u(0)) = -\Delta u(0)$, it follows from (4.2), (4.4) and the continuity of u and Δu in $\bar{\omega}$ that f is continuous at $u(0)$, and hence in the whole interval $[0, u(0)]$. By the continuity of u and Δu at 0, the equation

$$\Delta u + f(u) = 0$$

holds in $\bar{\omega}$. Our goal is to show that ω is then a ball centered at the origin and that u is radially symmetric and decreasing with respect to $|x|$ in $\bar{\omega}$.

Remembering that u satisfies (4.2)–(4.4), it would then follow from [25] that $\omega = B_R$ for some $R > 0$ and u is radially symmetric and decreasing with respect to $|x|$ in $\bar{\omega}$, if the function f were known to be Lipschitz continuous in $[0, u(0)]$. However, f' is not bounded in a neighborhood of $u(0)$ in general (see the comments after Theorem 1.10 in Section 1.2). We will nevertheless still be able to show the desired symmetry of ω and of u by removing from ω small neighborhoods of 0 and applying Serrin's strategy and the method of moving planes in punctured domains. The images by u of the closures of these punctured domains are intervals of the type $[0, L]$, with $0 < L < u(0)$, and thus f is Lipschitz continuous in $[0, L]$.

More precisely, let first $\rho > 0$ be such that $\overline{B_\rho} \subset \omega$ and let e be any unit vector. Let η be any real number in $(0, \rho)$, and denote

$$\bar{\lambda}_e = \max_{x \in \partial\omega} x \cdot e > \rho > \eta.$$

Using the same notations $T_{e,\lambda}$, $H_{e,\lambda}$ and $R_{e,\lambda}$ as in (1.15)–(1.16), it follows from [4] that there is $\tilde{\lambda} \in (\rho, \bar{\lambda}_e)$ such that

$$R_{e,\lambda}(H_{e,\lambda} \cap \bar{\omega}) \subset \omega \quad \text{for all } \lambda \in (\tilde{\lambda}, \bar{\lambda}_e). \tag{4.5}$$

Since $\max_{\mathbb{R}} |\xi_x| \rightarrow 0$ as $|x| \xrightarrow{+} 0$ by Lemma 2.6 (i) (or here, more simply, because of (4.2), (4.4) and the continuity of u in $\bar{\omega}$), there is $x_\eta \in \omega \setminus \{0\}$ such that $\Xi_{x_\eta} \subset B_\eta$. Let then Ω' be the bounded connected component of $\mathbb{R}^2 \setminus \Xi_{x_\eta}$ (notice that $\overline{\Omega'} \subset B_\eta \subset \omega$) and let

$$\tilde{\omega} = \omega \setminus \overline{\Omega'}$$

be the doubly connected bounded domain located between Ξ_{x_η} and $\partial\omega$. Notice that $\partial\tilde{\omega} = \Xi_{x_\eta} \cup \partial\omega$, $0 \notin \tilde{\omega}$ and

$$0 < u < u(x_\eta) \quad \text{in } \tilde{\omega} \tag{4.6}$$

since u has no critical point in $\tilde{\omega}$.

From (4.5), two cases can occur: either

(a) we have

$$R_{e,\lambda}(H_{e,\lambda} \cap \bar{\omega}) \subset \omega \quad \text{for all } \lambda \in [\eta, \bar{\lambda}_e), \tag{4.7}$$

- (b) there is $\lambda^* \in [\eta, \tilde{\lambda}]$ such that $R_{e,\lambda}(H_{e,\lambda} \cap \bar{\omega}) \subset \omega$ for all $\lambda \in (\lambda^*, \bar{\lambda}_e)$, and either
- (internal tangency) there is a point $x^* \in H_{e,\lambda^*} \cap \partial\omega$ such that $x_{e,\lambda^*}^* = R_{e,\lambda^*}(x^*)$ is in $\partial\omega$, or
 - (orthogonality) T_{e,λ^*} meets $\partial\omega$ orthogonally, at some point p^* .

We will prove that only case (a) occurs.

Assume by way of contradiction that case (b) occurs. Denote

$$\begin{cases} \Xi = \partial\Omega, & \Xi' = \Xi_{x_\eta}, & R' = \min_{x \in \Xi'} |x| \in (0, \rho), & R = \max_{x \in \Xi} |x| > \rho > R', \\ \varepsilon = \lambda^* \in [\eta, \tilde{\lambda}] \subset [0, \bar{\lambda}_e), \\ c_1 = 0 = u|_{\partial\omega}, & c_2 = u(x_\eta) = u|_{\Xi_{x_\eta}} \in (0, u(0)). \end{cases}$$

The $C^3(\bar{\omega})$ function $\varphi = u$ satisfies $c_1 < \varphi < c_2$ in $\tilde{\omega}$ by (4.6) and $\Delta\varphi + F(\varphi) = 0$ in $\tilde{\omega}$, where $F(s) = f(s)$ for $s \in [c_1, c_2] \subset [0, u(0))$. The function F is therefore C^1 in $[c_1, c_2]$. Condition (1.18) holds by definition of ε , λ^* and $\bar{\lambda}_e$, and (1.19) is automatically fulfilled since $\Xi' \subset B_\eta$ and $\varepsilon = \lambda^* \geq \eta$. Therefore, all assumptions of Proposition 1.14 are satisfied and it follows from the conclusion applied with $\lambda = \varepsilon = \lambda^*$ that

$$u \leq u_{e,\lambda^*} \quad \text{in } \overline{\omega_{e,\lambda^*}} \quad \text{with } \omega_{e,\lambda^*} = (H_{e,\lambda^*} \cap \tilde{\omega}) \setminus R_{e,\lambda^*}(\overline{\Omega'}).$$

Denote

$$w = u_{e,\lambda^*} - u.$$

Since $F = f$ is of class C^1 in $[c_1, c_2] \subset [0, u(0))$, the nonnegative $C^3(\overline{\omega_{e,\lambda^*}})$ function w satisfies an equation of the type $\Delta w + cw = 0$ in $\overline{\omega_{e,\lambda^*}}$ for some $c \in C(\overline{\omega_{e,\lambda^*}})$. Thus, the strong maximum principle implies that, for each connected component ω' of ω_{e,λ^*} , either $w > 0$ in ω' , or $w \equiv 0$ in $\overline{\omega'}$. We shall now consider separately the internal tangency case and the orthogonality case.

Consider first the case of internal tangency: there is a point $x^* \in H_{e,\lambda^*} \cap \partial\omega$ such that $x_{e,\lambda^*}^* = R_{e,\lambda^*}(x^*) \in \partial\omega$, hence

$$u(x^*) = u(x_{e,\lambda^*}^*) = 0 \quad \text{and} \quad w(x^*) = 0.$$

Since $\overline{\Omega'} \cap \partial\omega = \emptyset$, one has $x^* \notin \overline{\Omega'} \cup R_{e,\lambda^*}(\overline{\Omega'})$. There is a connected component ω^* of ω_{e,λ^*} such that $x^* \in \partial\omega^*$, and $B(x^*, r) \cap \omega = B(x^*, r) \cap \omega^*$ for all $r > 0$ small enough (in particular, the interior sphere condition in ω^* is satisfied at $x^* \in \partial\omega^*$). Let $n(\zeta)$ be the generic notation for the outward normal to ω at a point $\zeta \in \partial\omega$. Owing to the definitions of λ^* and x^* , one has $R_{e,\lambda^*}(n(x^*)) = n(x_{e,\lambda^*}^*)$, while $\nabla u_{e,\lambda^*}(x^*) = R_{e,\lambda^*}(\nabla u(x_{e,\lambda^*}^*))$ owing to the definition of u_{e,λ^*} . Hence,

$$\begin{aligned} \nabla w(x^*) \cdot n(x^*) &= \nabla u_{e,\lambda^*}(x^*) \cdot n(x^*) - \nabla u(x^*) \cdot n(x^*) \\ &= R_{e,\lambda^*}(\nabla u(x_{e,\lambda^*}^*)) \cdot R_{e,\lambda^*}(n(x_{e,\lambda^*}^*)) - \nabla u(x^*) \cdot n(x^*) \\ &= \nabla u(x_{e,\lambda^*}^*) \cdot n(x_{e,\lambda^*}^*) - \nabla u(x^*) \cdot n(x^*) = 0 \end{aligned}$$

since $\nabla u \cdot n$ is equal to the constant γ on $\partial\omega$ by (4.3). It then follows from the Hopf lemma applied to the function w at the point x^* , together with the strong maximum principle, that

$$w \equiv 0 \text{ in } \overline{\omega^*}, \quad \text{that is,} \quad u \equiv u_{e,\lambda^*} \text{ in } \overline{\omega^*}.$$

On the other hand, as for (5.3) in the proof of Proposition 1.14, one has

$$\begin{aligned} \partial\omega^* \subset \partial\omega_{e,\lambda^*} \\ \subset \underbrace{((T_{e,\lambda^*} \cap \tilde{\omega}) \setminus R_{e,\lambda^*}(\Omega'))}_{=: \partial_1\omega_{e,\lambda^*}} \cup \underbrace{((H_{e,\lambda^*} \cap \partial\omega) \setminus R_{e,\lambda^*}(\Omega'))}_{=: \partial_2\omega_{e,\lambda^*}} \cup \underbrace{(H_{e,\lambda^*} \cap \tilde{\omega} \cap R_{e,\lambda^*}(\Xi_{x_\eta}))}_{=: \partial_3\omega_{e,\lambda^*}}. \end{aligned}$$

Since $u_{e,\lambda^*} = u(x_\eta)$ on $R_{e,\lambda^*}(\Xi_{x_\eta})$ and $u < u(x_\eta)$ in $\tilde{\omega}$, one has $w = u_{e,\lambda^*} - u > 0$ on $\partial_3\omega_{e,\lambda^*}$, hence $\partial_3\omega_{e,\lambda^*} \cap \partial\omega^* = \emptyset$ and

$$\begin{aligned} \partial\omega^* &\subset ((T_{e,\lambda^*} \cap \tilde{\omega}) \setminus R_{e,\lambda^*}(\Omega')) \cup ((H_{e,\lambda^*} \cap \partial\omega) \setminus R_{e,\lambda^*}(\Omega')) \\ &\subset (T_{e,\lambda^*} \cap \tilde{\omega}) \cup (H_{e,\lambda^*} \cap \partial\omega). \end{aligned}$$

Therefore, ω^* is a connected component of $H_{e,\lambda^*} \cap \omega$. Since $w \equiv 0$ in $\overline{\omega^*}$, the arguments of Reichel [22] (see also [1, 27]) imply that $\tilde{\omega} = \overline{\omega^* \cup R_{e,\lambda^*}(\omega^*)}$. Hence ω symmetric with respect to the line T_{e,λ^*} , and moreover u is itself symmetric with respect to T_{e,λ^*} , which is impossible since $0 \notin T_{e,\lambda^*}$ and 0 is the only maximum point of u . As a consequence, the case of internal tangency is ruled out.

Consider now the case of orthogonality, that is, T_{e,λ^*} meets $\partial\omega$ orthogonally, at some point p^* . By definition of u_{e,λ^*} , one has $u(p^*) = u_{e,\lambda^*}(p^*)$, thus $w(p^*) = 0$. Notice

also that, as in the case of internal tangency, $p^* \notin \overline{\Omega'} \cup R_{e,\lambda^*}(\overline{\Omega'})$. There is a connected component ω^* of ω_{e,λ^*} such that $p^* \in \partial\omega^*$, and $B(p^*, r) \cap \omega \cap H_{e,\lambda^*} = B(p^*, r) \cap \omega^*$ for all $r > 0$ small enough. Since u and $\frac{\partial u}{\partial n}$ are constant on $\partial\omega$ and since T_{e,λ^*} meets $\partial\omega$ orthogonally at p^* , it follows as in [22] that all first and second order derivatives of w vanish at p^* . Serrin's corner lemma [25] and the strong maximum principle then yield $w \equiv 0$ in $\overline{\omega^*}$. One is then led to a contradiction as in the previous paragraph.

As a consequence, only case (a) occurs. Thus, (4.7) holds. By arguing as at the beginning of the study of case (b) and applying Proposition 1.14 with $\varepsilon = \eta$ this time, one infers that

$$u \leq u_{e,\lambda} \quad \text{in } \overline{\omega_{e,\lambda}} \quad \text{for all } \lambda \in [\eta, \bar{\lambda}_e), \tag{4.8}$$

with $\omega_{e,\lambda} = (H_{e,\lambda} \cap \tilde{\omega}) \setminus R_{e,\lambda}(\overline{\Omega'})$. Since (4.7) and (4.8) hold for every direction $e \in \mathbb{S}^1$ and for every $\eta \in (0, \rho)$, one finally concludes that $\omega = B_R$ for some $R > 0$ and that, as in the proof of Theorem 1.5, u is radially symmetric in $\overline{\omega} = \overline{B_R}$. Since 0 is the unique critical point of the $C^3(\overline{B_R})$ function u and since $u = 0$ on ∂B_R with $u > 0$ in B_R , there is a $C^3([0, R])$ function $U : [0, R] \rightarrow \mathbb{R}$ such that $u(x) = U(|x|)$ in $\overline{B_R}$, with $U'(0) = 0$ and $U' < 0$ in $(0, R]$. Therefore,

$$v(x) = \nabla^\perp u(x) = U'(|x|)e_\theta(x) \quad \text{for all } x \in \overline{B_R} \setminus \{0\}$$

and the $C^2([0, R])$ function $V = U'$ satisfies the desired conclusion. The proof of Theorem 1.10 is thereby complete. ■

5. Proof of Proposition 1.14

The proof is based on the method of moving planes developed in [3, 6, 11, 25], adapted to our geometrical configuration. The idea is to compare the function φ with its reflection $\varphi_{e,\lambda}$ in $\overline{\omega_{e,\lambda}}$ by moving the lines $T_{e,\lambda}$ and decreasing λ from the value $\bar{\lambda}$ to ε . We recall that

$$\omega_{e,\lambda} = (H_{e,\lambda} \cap \omega) \setminus R_{e,\lambda}(\overline{\Omega'}).$$

Notice in particular that $R' \leq |x| \leq R$ for all $\lambda \in [\varepsilon, \bar{\lambda})$ and $x \in \overline{\omega_{e,\lambda}}$, since $\omega_{e,\lambda} \subset \omega = \Omega \setminus \overline{\Omega'}$.

Consider first any $\lambda \in (\varepsilon, \bar{\lambda})$. For each $x \in \omega_{e,\lambda}$,

$$x_{e,\lambda} = R_{e,\lambda}(x) \in R_{e,\lambda}(H_{e,\lambda} \cap \omega) \subset R_{e,\lambda}(H_{e,\lambda} \cap \overline{\Omega}) \subset \Omega$$

by (1.18), and $x_{e,\lambda} \notin \overline{\Omega'}$, hence $x_{e,\lambda} \in \omega$. Thus

$$R_{e,\lambda}(\overline{\omega_{e,\lambda}}) \subset \overline{\omega}$$

and the function $\varphi_{e,\lambda}$ given in (1.20) is well defined and of class C^2 in $\overline{\omega_{e,\lambda}}$. Furthermore, $\Delta\varphi_{e,\lambda} + F(|x_{e,\lambda}|, \varphi_{e,\lambda}) = 0$ in $\overline{\omega_{e,\lambda}}$. Since $|x| \geq |x_{e,\lambda}|$ for all $x \in \overline{\omega_{e,\lambda}}$ (remember that $\lambda > \varepsilon \geq 0$) and since F is nonincreasing with respect to its first variable, it follows that

$$\Delta\varphi_{e,\lambda} + F(|x|, \varphi_{e,\lambda}) \leq 0 \quad \text{in } \overline{\omega_{e,\lambda}}.$$

Let

$$\Phi_{e,\lambda} = \varphi_{e,\lambda} - \varphi,$$

which is well defined and of class C^2 in $\overline{\omega_{e,\lambda}}$. We have

$$\Delta \Phi_{e,\lambda} + c_{e,\lambda} \Phi_{e,\lambda} \leq 0 \quad \text{in } \overline{\omega_{e,\lambda}}, \tag{5.1}$$

where, say,

$$c_{e,\lambda}(x) = \begin{cases} \frac{F(|x|, \varphi_{e,\lambda}(x)) - F(|x|, \varphi(x))}{\varphi_{e,\lambda}(x) - \varphi(x)} & \text{if } \varphi_{e,\lambda}(x) \neq \varphi(x), \\ 0 & \text{if } \varphi_{e,\lambda}(x) = \varphi(x). \end{cases}$$

Since the function F is assumed to be Lipschitz continuous with respect to its second variable, uniformly with respect to the first one, the function $c_{e,\lambda}$ is in $L^\infty(\omega_{e,\lambda})$, and moreover there is a constant $M \geq 0$ such that

$$|c_{e,\lambda}(x)| \leq M \quad \text{for all } \lambda \in (\varepsilon, \bar{\lambda}) \text{ and all } x \in \overline{\omega_{e,\lambda}}. \tag{5.2}$$

Consider again any $\lambda \in (\varepsilon, \bar{\lambda})$ and decompose the boundary of $\omega_{e,\lambda}$ into three parts. More precisely, since

$$\partial(A \cap B \cap C) \subset (\partial A \cap \bar{B} \cap \bar{C}) \cup (A \cap \partial B \cap \bar{C}) \cup (A \cap B \cap \partial C)$$

for any three sets A, B and C , since $\partial\omega = \Xi \cup \Xi'$ and since $(H_{e,\lambda} \cap \Xi') \setminus R_{e,\lambda}(\Omega') = \emptyset$ by assumption (1.19), one has (with $A = H_{e,\lambda}$, $B = \omega$ and $C = \mathbb{R}^2 \setminus R_{e,\lambda}(\bar{\Omega}')$)

$$\partial\omega_{e,\lambda} \subset \underbrace{((T_{e,\lambda} \cap \bar{\omega}) \setminus R_{e,\lambda}(\Omega'))}_{=:\partial_1\omega_{e,\lambda}} \cup \underbrace{((H_{e,\lambda} \cap \Xi) \setminus R_{e,\lambda}(\Omega'))}_{=:\partial_2\omega_{e,\lambda}} \cup \underbrace{(H_{e,\lambda} \cap \omega \cap R_{e,\lambda}(\Xi'))}_{=:\partial_3\omega_{e,\lambda}} \tag{5.3}$$

(see Fig. 4). Notice that since $T_{e,\lambda} \cap \Xi = T_{e,\lambda} \cap \partial\Omega$ is nonempty (because $\lambda \in (\varepsilon, \bar{\lambda}) \subset [0, \bar{\lambda})$), both sets $\partial_1\omega_{e,\lambda}$ and $\partial_2\omega_{e,\lambda}$ are nonempty (however, $\partial_3\omega_{e,\lambda}$ may be empty). Furthermore, even if $\omega_{e,\lambda}$ may not be connected (as in Fig. 4), the boundary of each connected component of $\omega_{e,\lambda}$ intersects $\partial_2\omega_{e,\lambda} \cup \partial_3\omega_{e,\lambda}$.

Let us now study the sign of $\Phi_{e,\lambda}$ on $\partial\omega_{e,\lambda}$, for any $\lambda \in (\varepsilon, \bar{\lambda})$. First, on $\partial_1\omega_{e,\lambda}$ ($\subset T_{e,\lambda}$), one has $\varphi_{e,\lambda} = \varphi$, hence $\Phi_{e,\lambda} = 0$. Secondly, for each $x \in \partial_2\omega_{e,\lambda}$, one has

$$x_{e,\lambda} \in R_{e,\lambda}(H_{e,\lambda} \cap \Xi) \subset R_{e,\lambda}(H_{e,\lambda} \cap \bar{\Omega}) \subset \Omega$$

by (1.18), hence $x_{e,\lambda} \in \omega \cup \Xi'$ and $\varphi_{e,\lambda}(x) = \varphi(x_{e,\lambda}) > c_1$ by (1.17), while $x \in \Xi$ and $\varphi(x) = c_1$. Thus, $\Phi_{e,\lambda}(x) = \varphi_{e,\lambda}(x) - \varphi(x) > 0$ for each $x \in \partial_2\omega_{e,\lambda}$. Thirdly, for each $x \in \partial_3\omega_{e,\lambda}$, one has $x_{e,\lambda} \in \Xi'$ and $\varphi_{e,\lambda}(x) = \varphi(x_{e,\lambda}) = c_2$, while $x \in \omega$ and $\varphi(x) < c_2$, by (1.17). Thus, $\Phi_{e,\lambda}(x) = \varphi_{e,\lambda}(x) - \varphi(x) > 0$ for each $x \in \partial_3\omega_{e,\lambda}$. As a consequence, $\Phi_{e,\lambda} \geq 0$ on $\partial\omega_{e,\lambda}$ and even $\Phi_{e,\lambda} > 0$ on $\partial_2\omega_{e,\lambda} \cup \partial_3\omega_{e,\lambda}$ ($\neq \emptyset$), hence

$$0 \leq \Phi_{e,\lambda} \neq 0 \quad \text{on the boundary of each connected component of } \omega_{e,\lambda}. \tag{5.4}$$




Fig. 4. The three parts $\partial_1\omega_{e,\lambda}$, $\partial_2\omega_{e,\lambda}$ and $\partial_3\omega_{e,\lambda}$ of the boundary of $\omega_{e,\lambda}$ (dashed red).

Let us now consider $\lambda \simeq \bar{\lambda}$ with $\lambda < \bar{\lambda}$. Since the functions $\Phi_{e,\lambda}$ satisfy (5.1)–(5.4), and since the sets $\omega_{e,\lambda}$ are all included in the given bounded domain Ω and the Lebesgue measure $|\omega_{e,\lambda}|$ of $\omega_{e,\lambda}$ goes to 0 as $\lambda \rightarrow \bar{\lambda}$ owing to the definition of $\bar{\lambda}$, it follows for instance from the maximum principle in sets with bounded diameter and small Lebesgue measure and from the strong maximum principle [7] that there is $\lambda_0 \in (\varepsilon, \bar{\lambda})$ such that $\Phi_{e,\lambda} > 0$ in $\omega_{e,\lambda}$ for all $\lambda \in (\lambda_0, \bar{\lambda})$.

Let us finally define

$$\lambda_* = \inf \{ \lambda \in (\varepsilon, \bar{\lambda}) : \Phi_{e,\lambda'} > 0 \text{ in } \omega_{e,\lambda'} \text{ for all } \lambda' \in (\lambda, \bar{\lambda}) \},$$

and notice that $\varepsilon \leq \lambda_* \leq \lambda_0 < \bar{\lambda}$. Our goal is to show that $\lambda_* = \varepsilon$. Assume by way of contradiction that $\lambda_* > \varepsilon$. Notice that $\Phi_{e,\lambda_*} \geq 0$ in $\overline{\omega_{e,\lambda_*}}$ by continuity (indeed, for each $x \in \omega_{e,\lambda_*}$, we have $x \in \omega_{e,\lambda}$ for $\lambda - \lambda_* > 0$ small, hence $\varphi(x) < \varphi_{e,\lambda}(x)$ for $\lambda - \lambda_* > 0$ small, and $\varphi(x) \leq \varphi_{e,\lambda_*}(x)$ by passing to the limit $\lambda \rightarrow \lambda_*$ and by continuity of φ ; therefore, $\varphi \leq \varphi_{e,\lambda_*}$ in $\overline{\omega_{e,\lambda_*}}$ again by continuity of φ). On the other hand, $0 \leq \Phi_{e,\lambda_*} \not\equiv 0$ on the boundary of each connected component of ω_{e,λ_*} , because $\lambda_* \in (\varepsilon, \bar{\lambda})$. Hence, $\Phi_{e,\lambda_*} > 0$ in ω_{e,λ_*} from the strong maximum principle. As in the previous paragraph, from [7], there exists $\delta > 0$ such that the weak maximum principle holds in any open set $\omega' \subset \omega$ for the solutions $\Phi \in C^2(\omega') \cap C(\overline{\omega'})$ of $\Delta\Phi + c\Phi \leq 0$ in ω' with $\Phi \geq 0$ on $\partial\omega'$ and $\|c\|_{L^\infty(\omega')} \leq M$, as soon as $|\omega'| \leq \delta$. Let then K be a compact subset of ω_{e,λ_*} such that

$$|\omega_{e,\lambda_*} \setminus K| < \delta/2.$$

Since $\min_K \Phi_{e,\lambda_*} > 0$, it follows from the continuity of φ in $\bar{\omega}$ that there exists $\underline{\lambda} \in (\varepsilon, \lambda_*)$ such that, for all $\lambda \in [\underline{\lambda}, \lambda_*]$,

$$\min_K \Phi_{e,\lambda} > 0, \quad \partial(\omega_{e,\lambda} \setminus K) = \partial\omega_{e,\lambda} \cup \partial K \quad \text{and} \quad |\omega_{e,\lambda} \setminus K| < \delta.$$

For any such $\lambda \in [\underline{\lambda}, \lambda_*]$, one then has $0 \leq \Phi_{e,\lambda} \not\equiv 0$ on the boundary of each connected component of $\omega_{e,\lambda} \setminus K$ and one infers from the choice of δ and from the strong maximum principle that $\Phi_{e,\lambda} > 0$ in $\omega_{e,\lambda} \setminus K$, and finally $\Phi_{e,\lambda} > 0$ in $\omega_{e,\lambda}$. This last property contradicts the definition of λ_* .

As a conclusion, $\lambda_* = \varepsilon$. Therefore, for every $\lambda \in (\varepsilon, \bar{\lambda})$, one has $\Phi_{e,\lambda} > 0$ in $\omega_{e,\lambda}$, so $\varphi < \varphi_{e,\lambda}$ in $\omega_{e,\lambda}$ and $\varphi \leq \varphi_{e,\lambda}$ in $\overline{\omega_{e,\lambda}}$ by continuity of φ . As in the previous paragraph, it also follows by continuity that $\varphi \leq \varphi_{e,\varepsilon}$ in $\overline{\omega_{e,\varepsilon}}$. The proof of Proposition 1.14 is thereby complete. ■

Acknowledgements. The authors are grateful to the anonymous reviewers for pointing out the references [10, 13] and for very valuable comments which improved the readability and quality of the paper.

The authors are grateful to Jean-Michel Coron for suggesting this research topic and to Boyan Sirakov for pointing out his reference [27].

Funding. This work has been carried out in the framework of Archimède Labex of Aix-Marseille University. The project leading to this publication has received funding from Excellence Initiative of Aix-Marseille University – A*MIDEX, a French “Investissements d’Avenir” programme, from the European Research Council under the European Union’s Seventh Framework Programme (FP/2007-2013) ERC Grant Agreement n. 321186 – ReaDi – Reaction-Diffusion Equations, Propagation and Modelling and from the ANR NONLOCAL (ANR-14-CE25-0013) and RESISTE (ANR-18-CE45-0019) projects.

References

- [1] Aftalion, A., Busca, J.: Radial symmetry of overdetermined boundary-value problems in exterior domains. Arch. Ration. Mech. Anal. **143**, 195–206 (1998) Zbl [0911.35008](#) MR [1650014](#)
- [2] Alessandrini, G.: A symmetry theorem for condensers. Math. Methods Appl. Sci. **15**, 315–320 (1992) Zbl [0756.35118](#) MR [1170529](#)
- [3] Alexandrov, A. D.: A characteristic property of spheres. Ann. Mat. Pura Appl. (4) **58**, 303–315 (1962) Zbl [0107.15603](#) MR [143162](#)
- [4] Amick, C. J., Fraenkel, L. E.: The uniqueness of Hill’s spherical vortex. Arch. Ration. Mech. Anal. **92**, 91–119 (1986) Zbl [0609.76018](#) MR [816615](#)
- [5] Berestycki, H., Caffarelli, L. A., Nirenberg, L.: Monotonicity for elliptic equations in unbounded Lipschitz domains. Comm. Pure Appl. Math. **50**, 1089–1111 (1997) Zbl [0906.35035](#) MR [1470317](#)
- [6] Berestycki, H., Nirenberg, L.: On the method of moving planes and the sliding method. Bol. Soc. Brasil. Mat. (N.S.) **22**, 1–37 (1991) Zbl [0784.35025](#) MR [1159383](#)
- [7] Berestycki, H., Nirenberg, L., Varadhan, S. R. S.: The principal eigenvalue and maximum principle for second-order elliptic operators in general domains. Comm. Pure Appl. Math. **47**, 47–92 (1994) Zbl [0806.35129](#) MR [1258192](#)
- [8] Choffrut, A., Šverák, V.: Local structure of the set of steady-state solutions to the 2D incompressible Euler equations. Geom. Funct. Anal. **22**, 136–201 (2012) Zbl [1256.35076](#) MR [2899685](#)
- [9] Farina, A., Valdinoci, E.: Flattening results for elliptic PDEs in unbounded domains with applications to overdetermined problems. Arch. Ration. Mech. Anal. **195**, 1025–1058 (2010) Zbl [1236.35058](#) MR [2591980](#)
- [10] Fraenkel, L. E.: An Introduction to Maximum Principles and Symmetry in Elliptic Problems. Cambridge Tracts in Math. 128, Cambridge Univ. Press, Cambridge (2000) Zbl [1214.35001](#) MR [1751289](#)
- [11] Gidas, B., Ni, W. M., Nirenberg, L.: Symmetry and related properties via the maximum principle. Comm. Math. Phys. **68**, 209–243 (1979) Zbl [0425.35020](#) MR [544879](#)

- [12] Gidas, B., Ni, W. M., Nirenberg, L.: Symmetry of positive solutions of nonlinear elliptic equations in \mathbf{R}^n . In: *Mathematical Analysis and Applications, Part A*, Adv. Math. Suppl. Stud. 7, Academic Press, New York, 369–402 (1981) Zbl [0469.35052](#) MR [634248](#)
- [13] Gómez-Serrano, J., Park, J., Shi, J., Yao, Y.: Symmetry in stationary and uniformly rotating solutions of active scalar equations. *Duke Math. J.* **170**, 2957–3038 (2021) MR [4312192](#)
- [14] Hamel, F., Nadirashvili, N.: Shear flows of an ideal fluid and elliptic equations in unbounded domains. *Comm. Pure Appl. Math.* **70**, 590–608 (2017) Zbl [1360.35155](#) MR [3602531](#)
- [15] Hamel, F., Nadirashvili, N.: Parallel and circular flows for the two-dimensional Euler equations. In: *Séminaire Laurent Schwartz—Équations aux dérivées partielles et applications. Année 2017–2018*, Ed. Éc. Polytech., Palaiseau, exp. V, 13 pp. (2018) MR [3968022](#)
- [16] Hamel, F., Nadirashvili, N.: A Liouville theorem for the Euler equations in the plane. *Arch. Ration. Mech. Anal.* **233**, 599–642 (2019) Zbl [1421.35263](#) MR [3951689](#)
- [17] Hmidi, T.: On the trivial solutions for the rotating patch model. *J. Evol. Equations* **15**, 801–816 (2015) Zbl [1338.35365](#) MR [3427065](#)
- [18] Kalisch, H.: Nonexistence of coherent structures in two-dimensional inviscid channel flow. *Math. Model. Nat. Phenom.* **7**, 77–82 (2012) Zbl [1251.35084](#) MR [2892779](#)
- [19] Koch, H., Nadirashvili, N.: Partial analyticity and nodal sets for nonlinear elliptic systems. arXiv:1506.06224 (2015)
- [20] Li, C.: Monotonicity and symmetry of solutions of fully nonlinear elliptic equations on unbounded domains. *Comm. Partial Differential Equations* **16**, 585–615 (1991) Zbl [0741.35014](#) MR [1113099](#)
- [21] Reichel, W.: Radial symmetry by moving planes for semilinear elliptic BVPs on annuli and other non-convex domains. In: *Elliptic and Parabolic Problems (Pont-à-Mousson, 1994)*, Pitman Res. Notes Math. Ser. 325, Longman Sci. Tech., Harlow, 164–182 (1995) Zbl [0839.35047](#) MR [1416582](#)
- [22] Reichel, W.: Radial symmetry for elliptic boundary-value problems on exterior domains. *Arch. Ration. Mech. Anal.* **137**, 381–394 (1997) Zbl [0891.35006](#) MR [1463801](#)
- [23] Ros, A., Ruiz, D., Sicbaldi, P.: A rigidity result for overdetermined elliptic problems in the plane. *Comm. Pure Appl. Math.* **70**, 1223–1252 (2017) Zbl [1373.35207](#) MR [3666566](#)
- [24] Ros, A., Sicbaldi, P.: Geometry and topology of some overdetermined elliptic problems. *J. Differential Equations* **255**, 951–977 (2013) Zbl [1284.35297](#) MR [3062759](#)
- [25] Serrin, J.: A symmetry problem in potential theory. *Arch. Ration. Mech. Anal.* **43**, 304–318 (1971) Zbl [0222.31007](#) MR [333220](#)
- [26] Serrin, J., Zou, H.: Symmetry of ground states of quasilinear elliptic equations. *Arch. Ration. Mech. Anal.* **148**, 265–290 (1999) Zbl [0940.35079](#) MR [1716665](#)
- [27] Sirakov, B.: Symmetry for exterior elliptic problems and two conjectures in potential theory. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **18**, 135–156 (2001) Zbl [0997.35014](#) MR [1808026](#)
- [28] Willms, N. B., Gladwell, G. M. L., Siegel, D.: Symmetry theorems for some overdetermined boundary value problems on ring domains. *Z. Angew. Math. Phys.* **45**, 556–579 (1994) Zbl [0807.35099](#) MR [1289661](#)