© 2022 European Mathematical Society Published by EMS Press



David Loeffler · Sarah Livia Zerbes

# On the Bloch–Kato conjecture for the symmetric cube

Received May 13, 2020

**Abstract.** We prove one inclusion in the Iwasawa main conjecture, and the Bloch–Kato conjecture in analytic rank 0, for the symmetric cube of a level 1 modular form.

Keywords. L-values, Bloch-Kato conjecture, Siegel modular forms

# 1. Introduction

In this short note, we study the arithmetic of the *p*-adic symmetric cube Galois representation associated to a level 1 modular cusp form. We prove that the Bloch–Kato conjecture holds in analytic rank 0 for the critical twists of this representation (corresponding to the critical values of the *L*-function); and we prove one inclusion of the cyclotomic Iwasawa Main Conjecture for this representation, showing that the characteristic ideal of the Selmer group divides the *p*-adic *L*-function. We deduce these results from the results of [5] on the arithmetic of automorphic representations of GSp(4), using the existence of a symmetric cube lifting from GL(2) to GSp(4).

**Remark 1.1.** • The results of this paper were inspired by the striking work of Haining Wang [7], where he proves many cases of the Bloch–Kato conjecture for symmetric cubes of modular forms via arithmetic level-raising on triple products of Shimura curves. Our results are complementary to those of Wang, since we restrict to level 1 (and hence weight  $\geq$  12), while he considers weight 2 and general level.

• The converse implication of the Bloch–Kato conjecture in this setting – i.e. that the vanishing of the central critical *L*-value implies the existence of a non-zero Selmer class – will be treated in forthcoming work of Samuel Mundy, using a functorial lifting to the exceptional group  $G_2$ .

• The restriction to modular forms of level 1 is carried over from the results in *op. cit.*, which rely on forthcoming work of Barrera, Dimitrov and Williams regarding interpola-

Sarah Livia Zerbes: Department of Mathematics, ETH Zürich, Rämistrasse 101, 8092 Zürich, Switzerland; sarah.zerbes@math.ethz.ch

Mathematics Subject Classification (2020): Primary 11F67; Secondary 11F03, 11F46, 11F80

David Loeffler: Mathematics Institute, Zeeman Building, University of Warwick, Coventry CV4 7AL, UK; d.a.loeffler@warwick.ac.uk

tion of *p*-adic *L*-functions for GL(2n) in finite-slope families. At present these results are only available for level 1, but we are confident that it should be possible to relax this assumption. In this case, all of the results of this paper will generalize without change to modular forms of higher level.

In forthcoming work, we will prove the analogous results for the Galois representations attached to quadratic Hilbert modular forms, using the twisted Yoshida lift to GSp(4).

#### 2. Large Galois image

Let f be a cuspidal, new, non-CM eigenform of weight  $k \ge 2$  and level  $\Gamma_1(N)$  for some  $N \ge 1$ , and write F for the coefficient field of f. Throughout this section, let pbe a prime > 3. Let v be a prime of F above p, and denote by  $V_{f,v}$  the 2-dimensional  $F_v$ -linear Galois representation associated to f. We are interested in the representation Sym<sup>3</sup>  $V_{f,v}$ .

**Remark 2.1.** (1) Our conventions are such that the trace of *geometric* Frobenius at a prime  $\ell \nmid pN$  is the Hecke eigenvalue  $a_{\ell}(f)$ .

- (2) The Hodge–Tate weights of  $(\text{Sym}^3 V_{f,v})|_{G_{\mathbb{Q}p}}$  are 3 3k, 2 2k, 1 k, 0.
- (3) We have  $(\text{Sym}^3 V_{f,v})^* \cong (\text{Sym}^3 V_{f,v})(3k-3)$ .
- (4) The critical twists of Sym<sup>3</sup>  $V_{f,v}$  are  $(\text{Sym}^3 V_{f,v})(j)$  for  $k \le j \le 2k 2$ .

**Proposition 2.2.** If  $p \gg 0$ , then the *p*-adic representation  $W_{\Pi}^* = \text{Sym}^3 V_{f,v}^*$  has large image: there exists a  $G_{\mathbb{Q}}$ -invariant lattice  $\mathcal{T}_{\Pi}$  of  $W_{\Pi}^*$  with the following properties:

- $\mathcal{T}_{\Pi} \otimes k_v$  is an irreducible  $k_v[\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\mu_{p^{\infty}}))]$ -module, where  $k_v$  is the residue field of  $\mathcal{O}_{F_v}$ , and
- there exists  $\tau \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\mu_{p^{\infty}}))$  such that  $\text{rank}_{\mathcal{O}_{F_{\mathcal{V}}}} \mathcal{T}_{\Pi}/(\tau-1)\mathcal{T}_{\Pi} = 1$ .

In order to prove this proposition, we quote the following result due to Ribet [6]:

**Lemma 2.3.** If p is sufficiently large, then there exists a  $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -stable  $\mathcal{O}_{F_v}$ -lattice T of  $V_{f,v}^*$  such that the image of the homomorphism

$$\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\mu_{p^{\infty}})) \to \operatorname{GL}_2(T) \cong \operatorname{GL}_2(\mathcal{O}_{F_v})$$

contains  $SL_2(\mathbb{Z}_p)$ . Here, the last isomorphism is given by a basis of T.

Proof of Proposition 2.2. Let  $\tau \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\mu_{p^{\infty}}))$  be such that its image in  $\text{GL}_2(\mathcal{O}_{F_v})$  is equal to  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ . Then the matrix of  $\tau$  with respect to the natural basis of  $\mathcal{T}_{\Pi} = \text{Sym}^3 T$  is

$$\begin{pmatrix} 1 & 3 & 3 & 1 \\ & 1 & 2 & 1 \\ & & 1 & 1 \\ & & & 1 \end{pmatrix},$$

so since p > 3, we deduce that rank $_{\mathcal{O}_{F_v}} \mathcal{T}_{\Pi}/(\tau - 1)\mathcal{T}_{\Pi} = 1$ .

The irreducibility of  $\mathcal{T}_{\Pi} \otimes k_v$  as a  $k_v[\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\mu_{p^{\infty}}))]$ -module is immediate.

**Remark 2.4.** We can clearly choose  $\tau \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}^{ab})$ ). Hence

$$\operatorname{rank}_{\mathcal{O}_{F_n}} \mathcal{T}_{\Pi}(\chi)/(\tau-1)\mathcal{T}_{\Pi}(\chi) = 1$$

for every Dirichlet character  $\chi$  of prime-to-*p*-conductor.

# 3. Functoriality

**Notation 3.1.** Let  $\pi$  be the associated automorphic representation of  $GL_2(\mathbb{A}_f)$ , and write  $\operatorname{Sym}^3 \pi$  for the image of  $\pi$  under the Langlands functoriality transfer from  $GL_2$  to  $GL_4$  attached to  $\operatorname{Sym}^3 : \operatorname{GL}_2(\mathbb{C}) \to \operatorname{GL}_4(\mathbb{C})$ .

**Theorem 3.2.** There exists a globally generic cuspidal automorphic representation  $\Pi = \bigotimes_v \Pi_v$  of  $\operatorname{GSp}_4(\mathbb{A})$  of weight (2k - 1, k + 1) such that  $\operatorname{Sym}^3 \pi$  is the functorial lift of  $\Pi$  under the embedding  $\operatorname{GSp}_4(\mathbb{C}) \to \operatorname{GL}_4(\mathbb{C})$ . In particular,

(1)  $\Pi$  is unramified at the primes not dividing N,

(2)  $L(\Pi, s - w/2) = L(\text{Sym}^3 \pi, s)$ , where w = 3(k + 1), and

(3) the p-adic spin representation  $W_{\Pi}$  associated to  $\Pi$  is equal to Sym<sup>3</sup>  $V_{f,v}$ .

*Here, we normalise*  $L(\Pi, s)$  *so that the centre of the functional equation is at* s = 1/2*.* 

*Proof.* See [3] and [1, §10].

**Remark 3.3.** In more classical terms,  $\Pi$  (or more precisely its *L*-packet) corresponds to a holomorphic vector-valued Siegel modular form taking values in the representation det<sup>*k*+1</sup>  $\otimes$  Sym<sup>3*k*-3</sup> of GL<sub>2</sub>( $\mathbb{C}$ ).

#### 4. Application to the Bloch-Kato conjecture

**Theorem 4.1.** Assume that f is a cuspidal eigenform of weight k and level 1, and let  $V = (\text{Sym}^3 V_{f,v})(2k-2)$ . Let p be a prime > 3, and assume that f is ordinary at p. Let  $0 \le j \le k-2$ , and let  $\rho$  be a finite order character of  $\mathbb{Z}_p^{\times}$ . If  $L(\text{Sym}^3 \pi \otimes \rho, k+j) \ne 0$ , then  $H_f^1(\mathbb{Q}, V(-j-\rho)) = 0$ .

*Proof.* The assumption on f being ordinary at p implies that the representation  $\Pi$  is Borel ordinary at p. Moreover, Remark 2.4 implies that the big image assumption  $\text{Hyp}(\mathbb{Q}(\mu_{p^{\infty}}), \mathcal{T}_{\Pi}(\chi))$  (cf. [5, §2]) is satisfied for all Dirichlet characters  $\chi$  of prime-to-p conductor. We are hence in the situation where we can apply [5, Theorem D].

**Remark 4.2.** Observe that if  $j \neq (k-2)/2$  (the central value), then the *L*-value  $L(\text{Sym}^3 \pi \otimes \rho, k+j)$  is automatically non-zero, since  $L(\Pi, s)$  is non-vanishing for Re(s) > 1.

# 5. Application to the Iwasawa main conjecture

We similarly obtain applications to the cyclotomic Iwasawa main conjecture. We preserve the notations and hypotheses of the previous section. Let  $\Gamma = \text{Gal}(\mathbb{Q}(\mu_{p^{\infty}})/\mathbb{Q})$ , and write  $\Lambda = \mathcal{O}_{F_{v}}[[\Gamma]]$  for the Iwasawa algebra.

# 5.1. p-adic L-functions for the symmetric cube

We recall the following two known results:

**Theorem 5.1** (Algebraicity of Sym<sup>3</sup> *L*-values). There exist constants  $\Omega_{\Pi}^+, \Omega_{\Pi}^- \in \mathbb{C}$ , welldefined up to multiplication by  $F^{\times}$ , such that for every Dirichlet character  $\chi$  and every  $0 \le j \le k - 2$ , the quantity

$$L^{\mathrm{alg}}(\mathrm{Sym}^3 f \otimes \bar{\rho}, j+k) \coloneqq \frac{G(\rho)^2}{(2\pi i)^j \cdot \Omega_{\Pi}^{\varepsilon}} L(\mathrm{Sym}^3 f \otimes \bar{\rho}, j+k)$$

lies in  $F(\chi)$ , and depends  $\operatorname{Gal}(\overline{F}/F)$ -equivariantly on  $\chi$ . Here  $\varepsilon = (-1)^j \rho(-1)$ .

**Theorem 5.2** (Existence of the Sym<sup>3</sup> *p*-adic *L*-function). Let  $\alpha$  be the unit root of the Hecke polynomial of f at p. Then there exists an element  $L_{v,\alpha}(\text{Sym}^3 f) \in \Lambda$  with the following interpolation property: for any Dirichlet character  $\rho$  of p-power conductor, and any  $0 \le j \le k-2$ , we have

$$L_{\nu,\alpha}(\operatorname{Sym}^{3} f)(j+\rho) = j!(j+k-1)!R_{p}(\operatorname{Sym}^{2} f,\rho,j)L^{\operatorname{alg}}(\operatorname{Sym}^{3} f \otimes \bar{\rho},j+k),$$

where

$$R_p(\operatorname{Sym}^3 f, \rho, j) := \begin{cases} \left(\frac{p^{2(j+k-1)}}{\alpha^5 \beta}\right)^m & \text{if } \rho \text{ has conductor } p^m > 1, \\ \left(1 - \frac{p^{j+k-1}}{\alpha^3}\right)\left(1 - \frac{p^{j+k-1}}{\alpha^2 \beta}\right)\left(1 - \frac{\alpha\beta^2}{p^{j+k}}\right)\left(1 - \frac{\beta^3}{p^{j+k}}\right) & \text{if } \rho = 1. \end{cases}$$

These results are due to Dimitrov, Januszewski and Raghuram [2], as a special case of general theorems applying to any automorphic representation  $\Pi$  of GL(4) admitting a Shalika model. An alternative proof using coherent cohomology of Siegel Shimura varieties is given in [4, Theorem A].

#### 5.2. The Iwasawa main conjecture

Recall that V denotes the Galois representation  $(\text{Sym}^3 V_{f,v})(2k-2)$ .

**Definition 5.3.** Let  $\widetilde{R\Gamma}_{\text{Iw}}(\mathbb{Q}(\mu_{p^{\infty}}), V)$  denote the Nekovář Selmer complex with the unramified local condition at  $\ell \neq p$ , and the Greenberg local condition at p determined by the ordinarity of f.

We shall be interested in the degree 2 cohomology  $\tilde{H}^2_{Iw}(\mathbb{Q}(\mu_{p^{\infty}}), V)$ . Note that this can be described in more classical terms: if T is a Galois-stable lattice in V, and  $(-)^{\vee}$ 

denotes Pontryagin duality, then for any  $0 \le j \le k - 2$ , there is a canonical isomorphism

$$\widetilde{H}^{2}_{\mathrm{Iw}}(\mathbb{Q}_{\mu_{p^{\infty}}}, V) \cong \left( \varinjlim_{n} H^{1}_{\mathrm{f}}(\mathbb{Q}(\mu_{p^{n}}), T^{\vee}(1+j)) \right)^{\vee}(j) \otimes \mathbb{Q}_{p}.$$

**Theorem 5.4.** If the assumptions of Theorem 4.1 are satisfied, then  $\widetilde{H}^2_{Iw}(\mathbb{Q}(\mu_{p^{\infty}}), V)$  is  $\Lambda[1/p]$ -torsion, and its characteristic ideal divides the p-adic L-function  $L_{\nu,\alpha}(\text{Sym}^3 f)$ .

*Proof.* This follows by applying [5, Theorem C] to the representation  $\Pi$ .

### References

- Conti, A.: Galois level and congruence ideal for *p*-adic families of finite slope Siegel modular forms. Compos. Math. 155, 776–831 (2019) Zbl 1435.11084 MR 3932577
- [2] Dimitrov, M., Januszewski, F., Raghuram, A.: L-functions of GL<sub>2n</sub>: p-adic properties and nonvanishing of twists. Compos. Math. 156, 2437–2468 (2020) MR 4199213
- [3] Kim, H. H., Shahidi, F.: Functorial products for GL<sub>2</sub> × GL<sub>3</sub> and the symmetric cube for GL<sub>2</sub>. Ann. of Math. (2) 155, 837–893 (2002) Zbl 1040.11036 MR 1923967
- [4] Loeffler, D., Pilloni, V., Skinner, C., Zerbes, S. L.: Higher Hida theory and *p*-adic *L*-functions for GSp<sub>4</sub>. Duke Math. J. **170**, 4033–4121 Zbl 07467789 MR 4348233
- [5] Loeffler, D., Zerbes, S. L.: On the Bloch–Kato conjecture for GSp(4). arXiv:2003.05960 (2020)
- [6] Ribet, K. A.: On *l*-adic representations attached to modular forms. II. Glasgow Math. J. 27, 185–194 (1985) Zbl 0596.10027 MR 819838
- [7] Wang, H.: Arithmetic level raising on triple product of Shimura curves and Gross–Schoen diagonal cycles I: ramified case. Algebra Number Theory, to appear; arXiv:2004.00555