© 2022 European Mathematical Society Published by EMS Press



Bradley Dirks · Mircea Mustață

Minimal exponents of hyperplane sections: a conjecture of Teissier

Received September 3, 2020

Abstract. We prove a conjecture of Teissier asserting that if f has an isolated singularity at P and H is a smooth hypersurface through P, then $\tilde{\alpha}_P(f) \ge \tilde{\alpha}_P(f|_H) + \frac{1}{\theta_P(f)+1}$, where $\tilde{\alpha}_P(f)$ and $\tilde{\alpha}_P(f|_H)$ are the minimal exponents at P of f and $f|_H$, respectively, and $\theta_P(f)$ is an invariant obtained by comparing the integral closures of the powers of the Jacobian ideal of f and of the ideal defining P. The proof builds on the approaches of Loeser (1984) and Elduque–Mustață (2021). The new ingredients are a result concerning the behavior of Hodge ideals with respect to finite maps and a result about the behavior of certain Hodge ideals for families of isolated singularities with constant Milnor number. In the opposite direction, we show that for every f, if H is a general hypersurface through P, then $\tilde{\alpha}_P(f) \le \tilde{\alpha}_P(f|_H) + \frac{1}{\text{mult}_P(f)}$, extending a result of Loeser from the case of isolated singularities.

Keywords. Arnold exponent, minimal exponent, Hodge ideals, spectrum

1. Introduction

Let X be a smooth complex algebraic variety and $f \in \mathcal{O}_X(X)$ nonzero, defining a hypersurface Y. For a point $P \in Y$, the *minimal exponent* $\tilde{\alpha}_P(f)$ can be defined as the negative of the largest root of the reduced Bernstein–Sato polynomial of f at P. This is a very interesting invariant of singularities that refines the *log canonical threshold* lct_P(f); more precisely, by a result of Lichtin and Kollár, we have lct_P(f) = min { $\tilde{\alpha}_P(f)$, 1} (see [13, Section 10]). Moreover, by a result of Saito [27, Theorem 0.4] the hypersurface Y has rational singularities at P if and only if $\tilde{\alpha}_P(f) > 1$. In the setting where Y has an isolated singularity at P, the minimal exponent can be described via asymptotic expansion of integrals along vanishing cycles [16, 17]. In this incarnation, it has been extensively studied in [1] and is also known as the Arnold exponent of f at P.

Bradley Dirks: Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA; bdirks@umich.edu

Mircea Mustață: Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA; mmustata@umich.edu

Mathematics Subject Classification (2020): Primary 14F10; Secondary 14B05, 32S25

In this article we are interested in the behavior of the minimal exponent under restriction to a smooth hypersurface H in X, containing P. When Y has an isolated singularity at P, Teissier [33] introduced and studied the invariant $\theta_P(f)$ defined as

$$\theta_P(f) = \max_E \frac{\operatorname{ord}_E(J_f)}{\operatorname{ord}_E(\mathfrak{m}_P)},\tag{1}$$

where *E* runs over the prime divisors over *X* with center at *P*, \mathfrak{m}_P is the ideal defining *P* in *X*, and J_f is the Jacobian ideal of *f*. Using the description of the integral closure of an ideal in terms of divisorial valuations (see [14, Chapter 9.6.A]) one can see that the maximum in (1) is achieved by a divisor on the normalized blow-up of *X* along $\mathfrak{m}_P \cdot J_f$; moreover, $\theta_P(f)$ is the minimum of the positive rational numbers r/s with the property that \mathfrak{m}_P^r is contained in the integral closure $\overline{J_f^s}$ of J_f^s . The following is our main result, giving a positive answer to a conjecture of Teissier [34].

Theorem 1.1. Suppose that $n = \dim(X) \ge 2$ and the hypersurface defined by f in X has an isolated singularity at P. If H is a smooth hypersurface in X, with $f|_H \ne 0$ and $P \in H$, then

$$\widetilde{\alpha}_P(f) \ge \widetilde{\alpha}_P(f|_H) + \frac{1}{\theta_P(f) + 1}.$$

By successively applying the theorem for general hyperplane sections (which automatically have isolated singularities), we obtain the following:

Corollary 1.2. If the hypersurface Y defined by f in X has an isolated singularity at P and if H_1, \ldots, H_{n-1} are general smooth hypersurfaces in X, containing P, then

$$\widetilde{\alpha}_{P}(f) \geq \frac{1}{\theta_{P}(f) + 1} + \frac{1}{\theta_{P}(f|_{H_{1}}) + 1} + \dots + \frac{1}{\theta_{P}(f|_{H_{1}} \cap \dots \cap H_{n-1}) + 1}$$

The inequality in Theorem 1.1 was proved by Loeser [15], with $\theta_P(f)$ replaced by its round-up $\lceil \theta_P(f) \rceil$ (that is, the smallest integer that is $\geq \theta_P(f)$). Assuming that $f \in \mathbb{C}[x_1, \ldots, x_n]$, P = 0, and H is the hyperplane given by $x_n = 0$, the argument in [15] made use of the family of hypersurfaces

$$h_t(x_1,\ldots,x_n) = f(x_1,\ldots,x_{n-1},tx_n) + (1-t)x_n^{\theta+1},$$

where $\theta = \lceil \theta_P(f) \rceil$. Note that $\tilde{\alpha}_0(h_1) = \tilde{\alpha}_0(f)$ and $\tilde{\alpha}_0(h_0) = \tilde{\alpha}_0(f|_H) + \frac{1}{\theta+1}$ by the Thom–Sebastiani property of Arnold exponents (see for example [16, Example (6.8)]). The definition of $\theta_0(f)$ implies that the Milnor number of h_t at 0 is constant in a neighborhood *V* of 1, hence by a result of Varchenko [35] it follows that $\tilde{\alpha}_0(h_t)$ is constant on *V*. Finally, by the semicontinuity of the Arnold exponent [30, Theorem 2.11], it follows that $\tilde{\alpha}_0(h_t) \geq \tilde{\alpha}_0(h_0)$ for $t \in V$, and we conclude that $\tilde{\alpha}_0(f) \geq \tilde{\alpha}_0(f|_H) + \frac{1}{\theta+1}$.

This argument was modified in [6] to prove the weaker version of Theorem 1 for log canonical thresholds. The idea was to consider the same family $(h_t)_{t \in \mathbb{C}}$, with $\theta = \theta_P(f)$. In order to make sense of this, one pulls back this expression by the finite cover given by $\pi(x_1, \ldots, x_n) = (x_1, \ldots, x_{n-1}, x_n^d)$, where *d* is a divisible enough positive integer.

Recall that the log canonical threshold is characterized by the triviality of certain invariants associated to f, the *multiplier ideals* (see [14, Chapter 9]); due to the presence of the finite cover π , the argument in [6] relied on considering whether the equation x_n^{d-1} (defining the relative canonical divisor of π) lies in a suitable multiplier ideal of $h_t \circ \pi$, by making use of various properties of multiplier ideals.

In this note we follow the same approach. Since we deal with the minimal exponent, we need to make use of more refined invariants, the *Hodge ideals* $I_p(f^{\lambda})$ introduced and studied in [21]. The definition of these invariants (as well as the proofs of their basic properties) makes use of Saito's theory of mixed Hodge modules [26]. It was shown in [22,29] that in the same way that triviality of multiplier ideals characterizes log canonical thresholds, triviality of Hodge ideals characterizes minimal exponents. In order to extend the approach in [6] to the setting of Hodge ideals, we need two new properties of these invariants, that are of independent interest.

Theorem 1.3. Let $\pi: Y \to X$ be a finite surjective morphism between smooth varieties and let $K_{Y/X}$ be the effective divisor on Y defined by the determinant of the Jacobian matrix of π . If $0 \neq f \in \mathcal{O}_X(X)$ and $g = \pi^*(f)$ both define reduced divisors, then for every $h \in \mathcal{O}_X(X)$, every nonnegative integer p, and every $\lambda \in \mathbb{Q}_{>0}$, if $\pi^*(h) \cdot \mathcal{O}_Y(-K_{Y/X}) \subseteq I_p(g^{\lambda})$, then $h \in I_p(f^{\lambda})$.

For a more general statement, which does not assume that f and g define reduced divisors, see Theorem 3.5 below. We also give a partial converse of this result in the case of a Galois cover (see Theorem 3.6). At least for such covers, we thus have an extension of the formula in [14, Theorem 9.5.42] relating multiplier ideals under finite maps.

The next result is concerned with certain Hodge ideals associated to families of hypersurfaces with constant Milnor number. Let $\varphi: \mathcal{X} \to T$ be a smooth morphism of smooth complex algebraic varieties (in particular, T is connected), and let $s: T \to \mathcal{X}$ be a section of φ . Let $f \in \mathcal{O}_{\mathcal{X}}(\mathcal{X})$ be such that $f \circ s = 0$ and for every $t \in T$, the restriction f_t to $\mathcal{X}_t = \varphi^{-1}(t)$ is nonzero. We assume that for every $t \in T$, the hypersurface defined by f_t in \mathcal{X}_t is reduced, has at most one singular point at s(t), and furthermore the Milnor number of this hypersurface at s(t) is independent of $t \in T$ (note that the condition to be reduced is a consequence of isolated singularities as long as the relative dimension is at least 2). In this case, a result of Varchenko [35] says that the spectrum of f_t at s(t)is independent of $t \in T$; in particular, the minimal exponent $\tilde{\alpha}_{s(t)}(f_t)$ is independent of $t \in T$.

Theorem 1.4. With the above notation, if $\alpha = \tilde{\alpha}_{s(t)}(f_t)$ for $t \in T$, then for every nonnegative integer p and every $\gamma \in \mathbf{Q} \cap (0, 1]$ with $p + \gamma \leq \alpha + 1$, the subscheme of \mathcal{X} defined by $I_p(f^{\gamma})$ is finite and flat over T (possibly empty). Moreover, for every $t \in T$, we have

$$I_p(f_t^{\gamma}) = I_p(f^{\gamma}) \cdot \mathcal{O}_{\mathcal{X}_t}.$$

We deduce this result from the above-mentioned result of Varchenko on the constancy of the spectrum and a result due to Jung, Kim, Saito, and Yoon [9] that allows us to relate the Hodge ideals satisfying the condition in the theorem to the Hodge filtration in Steenbrink's mixed Hodge structure on the cohomology of the Milnor fiber. Finally, for the proof of Theorem 1.1 we also need the Restriction Theorem for Hodge ideals from [21], as well as a version of the Thom–Sebastiani property for certain Hodge ideals. Regarding the latter property, in our setting it is enough to use a Thom–Sebastiani type result for some related ideals, Saito's *microlocal multiplier ideals*; this property was proved by Maxim, Saito, and Schürmann [20].

Our main result gives a lower bound for the difference between the minimal exponent of f and the minimal exponent of a restriction of f to a smooth hypersurface, in terms of Teissier's invariant $\theta_P(f)$. Our last result is an upper bound for the same difference in terms of multiplicity, when we restrict to a *general* hypersurface. We note that in this result we do not require isolated singularities.

Theorem 1.5. Let X be a smooth complex algebraic variety with $\dim(X) \ge 2$, $f \in \mathcal{O}_X(X)$ nonzero, and $P \in X$ such that f(P) = 0. If H is a general hypersurface in X containing P, then

$$\widetilde{\alpha}_P(f|_H) \ge \widetilde{\alpha}_P(f) - \frac{1}{\operatorname{mult}_P(f)}$$

The case when f has an isolated singularity at P is a consequence of a stronger bound proved by Loeser [15]. We deduce the general case from this one by making use of a result from [22]. As a consequence of Theorem 1.5 we get conditions, in terms of the minimal exponent $\tilde{\alpha}_P(f)$, that guarantee that successive general hyperplane sections through P have rational singularities (see Corollary 6.1). Another application concerns a characterization of singular points with maximal minimal exponent (see Corollary 6.3).

The paper is structured as follows. In the next section we review briefly the Hodge ideals and the microlocal multiplier ideals, the connection between them, and the corresponding characterization of minimal exponents. In Section 3 we discuss the behavior of Hodge ideals under finite maps and prove Theorem 1.3, as well as its partial converse in the case of Galois finite covers. In Section 4 we recall the relevant result from [9] and use it to relate certain jumping numbers for Hodge ideals to the spectrum. In particular, we prove Theorem 1.4. We combine these results to give the proof of Theorem 1.1 in Section 5. The last section of the paper is devoted to the proof of the bound in the opposite direction in Theorem 1.5 above and of the above-mentioned applications.

2. Hodge ideals and minimal exponents

In this section we review some basic facts about Hodge ideals and their connection with minimal exponents and microlocal multiplier ideals, following [21,22]. Let X be a smooth *n*-dimensional complex algebraic variety and $f \in \mathcal{O}_X(X)$ a nonzero regular function. We denote by \mathcal{D}_X the sheaf of differential operators on X.

For every positive rational number α , we have a left \mathcal{D}_X -module

$$\mathcal{M}(f^{-\alpha}) := \mathcal{O}_X[1/f]f^{-\alpha}$$

This is a free module of rank 1 over the sheaf $\mathcal{O}_X[1/f]$, generated by the element $f^{-\alpha}$, with differential operators acting in the expected way: if D is a derivation on \mathcal{O}_X , then

$$D \cdot (hf^{-\alpha}) = D(h)f^{-\alpha} - \alpha \frac{hD(f)}{f}f^{-\alpha}.$$

Since $\mathcal{M}(f^{-\alpha})$ is a filtered direct summand of a mixed Hodge module in the sense of [26], it carries a canonical filtration $F_{\bullet}\mathcal{M}(f^{-\alpha})$, compatible with the filtration on \mathcal{D}_X given by order of differential operators. The Hodge ideals $(I_p(f^{\alpha}))_{p\geq 0}$ describe this filtration.

In what follows we will only be interested in the case when f defines a reduced divisor D. In this case, the Hodge ideals are given by

$$F_p \mathcal{M}(f^{-\alpha}) = I_p(f^{\alpha}) \cdot \mathcal{O}_X(pD) \cdot f^{-\alpha}$$

(we note that $I_p(f^{\alpha})$ was denoted by $I_p(\alpha D)$ in [21]).

It is sometimes convenient to also consider the right \mathcal{D}_X -module corresponding to $\mathcal{M}(f^{-\alpha})$. Recall that there is an equivalence of categories between left and right \mathcal{D}_X -modules such that if \mathcal{M} is a left \mathcal{D}_X -module, the \mathcal{O}_X -module underlying the right \mathcal{D}_X -module corresponding to \mathcal{M} is $\omega_X \otimes_{\mathcal{O}_X} \mathcal{M}$ (see [8, Section 1.2]). We denote by $\mathcal{M}_r(f^{-\alpha})$ the right \mathcal{D}_X -module corresponding to $\mathcal{M}(f^{-\alpha})$. The filtration on $\mathcal{M}(f^{-\alpha})$ induces a filtration on $\mathcal{M}_r(f^{-\alpha})$, with the convention

$$F_{p-n}\mathcal{M}_r(f^{-\alpha}) = \omega_X \otimes_{\mathcal{O}_X} F_p\mathcal{M}(f^{-\alpha}).$$

We next recall the V-filtration associated to f. Let $\iota: X \to X \times \mathbf{A}^1$ be the graph embedding given by $\iota(x) = (x, f(x))$. We denote the standard coordinate on \mathbf{A}^1 by t. The \mathcal{D} -module-theoretic push-forward $B_f := \iota_+ \mathcal{O}_X$ of \mathcal{O}_X can be described as

$$B_f = \mathcal{O}_X \otimes_{\mathbf{C}} \mathbf{C}[\partial_t]$$

(see [8, Example 1.3.5]). We thus have an \mathcal{O}_X -basis of B_f given by $\partial_t^j \delta$, for $j \ge 0$, where we put $\delta = 1 \otimes 1 \in B_f$. The action of *t* on the elements of this basis is given by

$$t \cdot \partial_t^j \delta = f \partial_t^j \delta - j \partial_t^{j-1} \delta.$$

The V-filtration on B_f is a rational filtration $(V^{\gamma}B_f)_{\gamma \in \mathbf{Q}}$ which is exhaustive, decreasing, left-continuous and discrete.¹ The filtration, constructed by Malgrange [18], is uniquely characterized by the following properties:

- (i) Every $V^{\gamma} B_f$ is a coherent module over $\mathcal{D}_X[t, \partial_t t]$.
- (ii) $t \cdot V^{\gamma} B_f \subseteq V^{\gamma+1} B_f$ for all $\gamma \in \mathbf{Q}$, with equality if $\gamma > 0$.
- (iii) $\partial_t \cdot V^{\gamma} B_f \subseteq V^{\gamma-1} B_f$ for all $\gamma \in \mathbf{Q}$.
- (iv) For every $\gamma \in \mathbf{Q}$, the operator $\partial_t t \gamma$ on $\operatorname{Gr}_V^{\gamma} := V^{\gamma} B_f / V^{>\gamma} B_f$ is nilpotent, where $V^{>\gamma} B_f = \bigcup_{\nu' > \nu} V^{\nu'} B_f$.

¹More precisely, the last two properties mean that there is a positive integer ℓ such that $V^{\gamma}B_f$ is constant for $\gamma \in (i/\ell, (i+1)/\ell]$, with $i \in \mathbb{Z}$.

Remark 2.1. It follows from property (iv) above that if $\alpha \neq 1$, then $t\partial_t$ is invertible on $\operatorname{Gr}_V^{\alpha}$; in particular, $\partial_t: \operatorname{Gr}_V^{\alpha} \to \operatorname{Gr}_V^{\alpha-1}$ is injective. This implies that if $u \in B_f$ is such that $\partial_t u \in V^{>0}B_f$, then $u \in V^{>1}B_f$.

In particular, the *V*-filtration on B_f induces a *V*-filtration $(V^{\gamma} \mathcal{O}_X)_{\gamma \in \mathbf{Q}}$ on \mathcal{O}_X via the inclusion $\mathcal{O}_X \hookrightarrow B_f$ that maps *h* to *h* δ . Saito [28] introduced a *microlocal* version of the *V*-filtration. This in turn induces the *microlocal V*-filtration $(\tilde{V}^{\gamma} \mathcal{O}_X)_{\gamma \in \mathbf{Q}}$ on \mathcal{O}_X , which we can describe via the usual *V*-filtration as follows. For $\gamma \leq 0$, we have $\tilde{V}^{\gamma} \mathcal{O}_X = \mathcal{O}_X$. If $\gamma > 0$, write $\gamma = p + \alpha$ for an integer *p* and $\alpha \in (0, 1]$ (hence $p = \lceil \alpha \rceil - 1$). With this notation, $\tilde{V}^{\gamma} \mathcal{O}_X$ consists of those regular functions $h \in \mathcal{O}_X$ with the property that there are regular functions $h_0, \ldots, h_{p-1} \in \mathcal{O}_X$ such that

$$h\partial_t^p \delta + h_{p-1}\partial_t^{p-1}\delta + \dots + h_0\delta \in V^{\alpha}B_f.$$

Whenever the function f is not clear from the context, we write $\tilde{V}^{\gamma} \mathcal{O}_X(f)$ for $\tilde{V}^{\gamma} \mathcal{O}_X$.

Remark 2.2. A basic fact is that $\delta \in V^{>0}B_f$. For example, this follows from Sabbah's description of the *V*-filtration in terms of *b*-functions (see [24]) and the fact, due to Kashiwara [11], that all the roots of the *b*-function of *f* are negative rational numbers.

Remark 2.3. The microlocal *V*-filtration on \mathcal{O}_X is a rational, decreasing, exhaustive, left-continuous and discrete filtration by coherent ideals. With the above definition, the only assertion that is not clear is that $\tilde{V}^{\gamma_1}\mathcal{O}_X \subseteq \tilde{V}^{\gamma_2}\mathcal{O}_X$ if $\gamma_1 > \gamma_2$. In order to check this, we can easily reduce to the case when $\gamma_2 = p$ is a positive integer and $\gamma_1 = p + \alpha$ for a rational number $\alpha \in (0, 1]$. In order to prove the inclusion, suppose that $h \in \tilde{V}^{p+\alpha}\mathcal{O}_X$, hence there are $h_0, \ldots, h_{p-1} \in \mathcal{O}_X$ such that

$$h\partial_t^p \delta + h_{p-1}\partial_t^{p-1}\delta + \dots + h_0\delta \in V^{\alpha}B_f.$$

Since $h_0 \delta \in V^{>0} B_f$ by Remark 2.2, it follows that if

$$u = h\partial_t^{p-1}\delta + h_{p-1}\partial_t^{p-2}\delta + \dots + h_1\delta,$$

then $\partial_t u \in V^{>0}B_f$, hence $u \in V^{>1}B_f \subseteq V^1B_f$ by Remark 2.1. This implies that $h \in \tilde{V}^p \mathcal{O}_X$.

Remark 2.4. The *microlocal multiplier ideals* of f are defined by

$$\widetilde{\mathcal{J}}(f^{\gamma}) := \widetilde{V}^{\gamma+\epsilon} \mathcal{O}_X \quad \text{for } 0 < \epsilon \ll 1$$

(see [20, 29]). The shift in the definition is convenient since it implies that for $\gamma < 1$, the microlocal multiplier ideal $\tilde{\mathcal{J}}(f^{\gamma})$ coincides with the usual multiplier ideal $\mathcal{J}(f^{\gamma})$ (this is a consequence of a theorem of Budur and Saito [4] relating multiplier ideals to the *V*-filtration on \mathcal{O}_X). In what follows we will not shift by ϵ since, as we will see shortly, this indexing matches the one for Hodge ideals, but we will still refer to the elements of the filtration $(\tilde{V}^{\gamma} \mathcal{O}_X)_{\gamma \in \mathbf{Q}}$ as *microlocal multiplier ideals*.

The following is the main result relating Hodge ideals and microlocal multiplier ideals. It was proved in [29, Theorem 1] for $\gamma \in \mathbb{Z}_{>0}$ and in [22, Theorem A'] in general.

Theorem 2.5. If f defines a reduced divisor, then for every $\gamma = p + \alpha$ with $p \in \mathbb{Z}_{\geq 0}$ and $\alpha \in \mathbb{Q} \cap (0, 1]$, we have

$$I_p(f^{\alpha}) + (f) = \tilde{V}^{\gamma} \mathcal{O}_X + (f).$$

In particular, it follows from the theorem that given $P \in X$ with f(P) = 0, we have $I_p(f^{\alpha}) = \mathcal{O}_X$ around P if and only if $\tilde{V}^{\gamma} \mathcal{O}_X = \mathcal{O}_X$ around P. It was shown by Saito [29] that if $\tilde{\alpha}_P(f)$ is the *minimal exponent* of f at P, then

$$V^{\gamma}\mathcal{O}_X = \mathcal{O}_X \text{ around } P \quad \text{if and only if } \gamma \leq \widetilde{\alpha}_P(f)$$
 (2)

(see also [22, Remark 6.13]). As a consequence, if f defines a reduced divisor, then

$$I_p(f^{\alpha}) = \mathcal{O}_X \text{ around } P \text{ if and only if } \widetilde{\alpha}_P(f) \ge p + \alpha$$
 (3)

(see [22, Corollary 6.1]).

Remark 2.6. We note that the definition of the minimal exponent that is used in the above results is in terms of the Bernstein–Sato polynomial of f. We will only need the above characterization and thus do not recall the precise definition. For more details and basic properties of the minimal exponent that follow from the above characterization in terms of Hodge ideals, we refer to [22]. The fact that for isolated singularities the minimal exponent coincides with the Arnold exponent follows from [17]. We will discuss in more detail the case of isolated singularities in Section 4.

For us it will be important that Hodge ideals are equal to microlocal multiplier ideals also in an interval of length 1 starting with the minimal exponent. More precisely, we have the following result. Recall that the *Jacobian ideal* J_f of f is defined as follows: if x_1, \ldots, x_n are algebraic coordinates in an open subset U of X, then $J_f|_U$ is generated by $\frac{\partial f}{\partial x_1}, \ldots, \frac{\partial f}{\partial x_n}$ (this definition is independent of the choice of coordinates and thus by gluing the local definitions we get a coherent ideal sheaf of \mathcal{O}_X).

Proposition 2.7. Let f be a nonzero regular function on the smooth complex algebraic variety X, defining a reduced divisor, and let $P \in X$ be such that f(P) = 0. Suppose that γ is a positive rational number and we write

$$\gamma = p + \alpha$$
 with $p = \lceil \gamma \rceil - 1$.

If $\gamma \leq \tilde{\alpha}_P(f) + 1$, then

$$I_p(f^{\alpha}) = \tilde{V}^{\gamma} \mathcal{O}_X$$

in a neighborhood of P; moreover, in such a neighborhood these ideals contain $(f) + J_f$.

Proof. Suppose first that $\gamma \leq 1$. In this case p = 0 and both ideals $I_0(f^{\gamma})$ and $\tilde{V}^{\gamma} \mathcal{O}_X$ are equal to the multiplier ideal $\mathcal{J}(f^{\gamma-\epsilon})$ for $0 < \epsilon \ll 1$ (for the Hodge ideal this follows from [21, Proposition 9.1], while for the microlocal multiplier ideal it follows from the result of Budur and Saito [4] relating multiplier ideals and the *V*-filtration). These ideals contain f since $(f) = \mathcal{J}(f) \subseteq \mathcal{J}(f^{\gamma-\epsilon})$ for every $\gamma \leq 1$. The fact that J_f is contained in $\mathcal{J}(f^{1-\epsilon})$ for $\epsilon > 0$ (which, in turn, is contained in $\mathcal{J}(f^{\gamma-\epsilon})$) is proved in [5, Theorem 4.2].

Suppose now that $\gamma > 1$, hence $p \ge 1$, and that $\gamma \le \tilde{\alpha}_P(f) + 1$. In this case we have

$$I_{p-1}(f^{\alpha}) = \mathcal{O}_X = \widetilde{V}^{\gamma-1}\mathcal{O}_X$$

in a neighborhood of P by (2) and (3). After possibly replacing X by this neighborhood of P, we may and will assume that these equalities hold on X. If we show that $f \in I_p(f^{\alpha}) \cap \tilde{V}^{\gamma} \mathcal{O}_X$, then the equality of the ideals in the proposition follows from Theorem 2.5. Then the last assertion follows as well if we show that $J_f \subseteq I_p(f^{\alpha})$.

The fact that $f \in I_p(f^{\alpha})$ has already been noticed in [22, Corollary 5.5]. The point is that since $I_{p-1}(f^{\alpha}) = \mathcal{O}_X$, we have

$$F_{p-1}\mathcal{M}(f^{-\alpha}) = \left(\mathcal{O}_X \frac{1}{f^{p-1}}\right) \cdot f^{-\alpha}.$$

The fact that $F_{p-1}\mathcal{M}(f^{-\alpha}) \subseteq F_p\mathcal{M}(f^{-\alpha})$ and the definition of $I_p(f^{\alpha})$ then give $f \in I_p(f^{\alpha})$. Moreover, since we have $F_1\mathcal{D}_X \cdot F_{p-1}\mathcal{M}(f^{-\alpha}) \subseteq F_p\mathcal{M}(f^{-\alpha})$, we see that if x_1, \ldots, x_n are local algebraic coordinates on X, then

$$-(p+\alpha-1)\frac{1}{f^p}\cdot\frac{\partial f}{\partial x_i}f^{-\alpha} = \partial_{x_i}\cdot\left(\frac{1}{f^{p-1}}f^{-\alpha}\right)\in F_p\mathcal{M}(f^{-\alpha}),$$

hence $J_f \subseteq I_p(f^{\alpha})$.

In order to complete the proof, it is enough to show that $f \in \tilde{V}^{\gamma} \mathcal{O}_X$. Since $\tilde{V}^{\gamma-1} \mathcal{O}_X = \mathcal{O}_X$, it follows that we have in $V^{\alpha} B_f$ an element of the form

 $u = \partial_t^{p-1} \delta$ + lower order terms.

In this case we have in $V^{\alpha}B_f$ also the element

$$\partial_t t u = f \partial_t^p \delta + \text{lower order terms},$$

hence $f \in \widetilde{V}^{\gamma} \mathcal{O}_X$.

3. Hodge ideals under finite maps

In this section we consider the behavior of Hodge ideals under finite surjective morphisms. Let us fix such a morphism $\pi: Y \to X$ between smooth complex *n*-dimensional algebraic varieties. Since we deal with push-forward of \mathcal{D} -modules, in this section we typically consider right \mathcal{D} -modules.

Recall that $\pi^*(\mathcal{D}_X)$ has a canonical structure of $(\mathcal{D}_Y, \pi^{-1}(\mathcal{D}_X))$ -bimodule; as such, it is denoted by $\mathcal{D}_{Y \to X}$. As for every proper morphism, we have an induced pushforward morphism between the corresponding derived categories of (quasi-coherent) right \mathcal{D} -modules given by $R\pi_*(-\otimes_{\mathcal{D}_Y}^L \mathcal{D}_{Y \to X})$ [8, Chapters 1.3 and 2.5]. The case of finite maps is easier: first, π_* is exact on quasi-coherent \mathcal{O}_Y -modules. Second, $\mathcal{D}_{Y \to X}$ is a flat left \mathcal{D}_Y -module (see [2, Theorem 2.11.10] or [10, Proposition 2.10]). This implies that the functor between derived categories is induced by the exact functor $\pi_+ = \pi_*(- \otimes_{\mathcal{D}_Y} \mathcal{D}_{Y \to X})$ between the corresponding Abelian categories of right $\mathcal{D}_$ modules. Note that if π is étale, then $\mathcal{D}_Y = \pi^*(\mathcal{D}_X)$ and the functor π_+ is equal to π_* on right \mathcal{D}_Y -modules.

Remark 3.1. We have a morphism of left \mathcal{D}_Y -modules $\mathcal{D}_Y \to \mathcal{D}_{Y \to X}$ that maps 1 to $\pi^*(1)$. This induces a canonical morphism of \mathcal{O}_X -modules $\pi_*(\mathcal{M}) \to \pi_+(\mathcal{M})$, which is an isomorphism if π is étale.

We also write i_+ for the push-forward functor between the Abelian categories of \mathcal{D} modules when $i: U \hookrightarrow X$ is the open immersion corresponding to the complement of an effective divisor in X. We thus have $i_+\mathcal{M} = i_*\mathcal{M}$ for every right \mathcal{D}_U -module \mathcal{M} . It is straightforward to see (and well-known) that if $j: V \hookrightarrow Y$ is the open immersion with $V = \pi^{-1}(U)$ and $\varphi: V \to U$ is the induced morphism, then we have a canonical isomorphism of functors $\pi_+ \circ j_+ \simeq i_+ \circ \varphi_+$.

Lemma 3.2. If $\pi: Y \to X$ is a finite surjective morphism between smooth complex algebraic varieties and if \mathcal{M} is a right \mathcal{D}_Y -module having no torsion as an \mathcal{O}_Y -module, then $\pi_+(\mathcal{M})$ has no torsion as an \mathcal{O}_X -module.

Proof. The assertion is local on X, hence we may assume that X (and thus also Y) is affine. By generic smoothness, we can find a nonzero $h \in \mathcal{O}_X(X)$ such that π is étale over the open subset $U = (h \neq 0)$. Let $V = \pi^{-1}(U)$ and $j: V \hookrightarrow Y$ and $i: U \hookrightarrow X$ be the corresponding open immersions. Note that we have a canonical morphism $\mathcal{M} \to j_+(\mathcal{M}|_V)$, which is injective since \mathcal{M} has no torsion as an \mathcal{O}_Y -module. By taking the direct image, we get an injective morphism

$$\pi_+ \mathcal{M} \to \pi_+ j_+ (\mathcal{M}|_V) \simeq i_+ \varphi_+ (\mathcal{M}|_V),$$

where $\varphi: V \to U$ is the induced morphism; therefore it is enough to show that the righthand side has no torsion. Note that as an \mathcal{O}_X -module, $i_+\varphi_+(\mathcal{M}|_V)$ is simply $i_*\varphi_*(\mathcal{M}|_V)$, since φ is étale. Since $\mathcal{M}|_V$ is an \mathcal{O}_V -module without torsion, it follows that $i_*\varphi_*(\mathcal{M}|_V)$ is an \mathcal{O}_X -module without torsion. This completes the proof.

In what follows, we will make use of Saito's theory of pure and mixed Hodge modules, for which we refer to [25, 26]. Recall that a mixed Hodge module has an underlying filtered (right) \mathcal{D} -module. For example, on a smooth *n*-dimensional variety *X*, we have the pure Hodge module $\mathbf{Q}_X^H[n]$, whose underlying \mathcal{D}_X -module is ω_X (the right \mathcal{D}_X module corresponding to \mathcal{O}_X), with the filtration given by $F_{p-n}\omega_X = \omega_X$ for $p \ge 0$, and $F_{p-n}\omega_X = 0$ otherwise.

Suppose now that $\varphi: W \to X$ is a morphism between smooth varieties which is either finite and surjective, or an open immersion given by the complement of an effective divisor. If $(\mathcal{M}, F_{\bullet})$ is a filtered \mathcal{D}_W -module on W that underlies a mixed Hodge module M, then we have by [26] a mixed Hodge module that we will denote by φ_+M and whose underlying filtered \mathcal{D}_X -module we will denote by $\varphi_+(\mathcal{M}, F_{\bullet})$. The corresponding \mathcal{D}_X module is just $\varphi_+(\mathcal{M})$, but the description of the filtration is rather subtle. When W is the complement of the hypersurface defined by $f \in \mathcal{O}_X(X)$ and $M = \mathbf{Q}_W^H[n]$, we get the filtration on $\mathcal{O}_X[1/f]$ described by the Hodge ideals of f. One easy case is when φ is finite and étale, in which case $F_p(\varphi_+\mathcal{M}) = \varphi_*(F_p\mathcal{M})$ for all p.

Given a finite surjective morphism $\pi: Y \to X$ of smooth *n*-dimensional algebraic varieties, we have a canonical morphism of mixed Hodge modules

$$\mathbf{Q}_X^H[n] \to \pi_+ \mathbf{Q}_Y[n] \tag{4}$$

that commutes with restriction to open subsets of X. At the level of \mathcal{D}_X -modules, this is given by the composition

$$\omega_X \hookrightarrow \pi_* \omega_Y \to \pi_+ \omega_Y,$$

where the first morphism maps a form $\eta \in \omega_X$ to its pull-back $\pi^*(\eta) \in \omega_Y$ and the second morphism is the one in Remark 3.1.

The next lemma provides the ingredient to relate Hodge ideals under finite maps. Suppose that $\pi: Y \to X$ is a finite surjective morphism between *n*-dimensional smooth varieties, $f \in \mathcal{O}_X(X)$ is nonzero, and $g = f \circ \pi \in \mathcal{O}_Y(Y)$. Recall that associated to f and g we have the filtered right \mathcal{D} -modules $\mathcal{M}_r(f^{-\alpha})$ and $\mathcal{M}_r(g^{-\alpha})$ on X and Y, respectively. Note that for every $\alpha \in \mathbf{Q}_{>0}$, we have a canonical morphism of \mathcal{O}_X -modules

$$\mathcal{M}_r(f^{-\alpha}) \to \pi_* \mathcal{M}_r(g^{-\alpha})$$
 (5)

that maps $u = f^{-\alpha} \frac{\eta}{f^m}$ with $\eta \in \omega_Y$ to $\pi^*(u) := g^{-\alpha} \frac{\pi^*(\eta)}{g^m}$.

Lemma 3.3. With the above notation, for every $\alpha \in \mathbf{Q}_{>0}$, the map τ given by the composition

$$\mathcal{M}_r(f^{-\alpha}) \to \pi_* \mathcal{M}_r(g^{-\alpha}) \to \pi_+ \mathcal{M}_r(g^{-\alpha}),$$

where the first map is the one in (5) and the second map is the one in Remark 3.1, is an injective strict morphism of filtered \mathcal{D}_X -modules.

Proof. For the proof, we will need to make use of the definition of the filtrations on $\mathcal{M}_r(f^{-\alpha})$ and $\mathcal{M}_r(g^{-\alpha})$ in [21, Section 2]. Let $X' = X \setminus V(f)$ and $Y' = Y \setminus V(g)$. Choose an integer $m \ge 2$ such that $m\alpha \in \mathbb{Z}$ and let

$$Y'' = \operatorname{Spec}(\mathcal{O}_{Y'}[y]/(y^m - g^{-m\alpha})) \quad \text{and} \quad X'' = \operatorname{Spec}(\mathcal{O}_{X'}[y]/(y^m - f^{-m\alpha})),$$

so that we have a diagram with Cartesian squares

$$\begin{array}{c|c} Y'' & \stackrel{q}{\longrightarrow} Y' & \stackrel{j}{\longrightarrow} Y \\ \psi & & \varphi & & & \\ \psi & & \varphi & & & \\ X'' & \stackrel{p}{\longrightarrow} X' & \stackrel{i}{\longrightarrow} X \end{array}$$

with i and j open immersions and p and q finite étale morphisms. We have by (4) a canonical morphism of mixed Hodge modules

$$\mathbf{Q}_{X''}^H[n] \to \psi_+ \mathbf{Q}_{Y''}^H[n],$$

that induces, after applying i_+p_+ and taking the underlying filtered \mathcal{D}_X -modules, a morphism of filtered \mathcal{D}_X -modules

$$\bigoplus_{k=0}^{m-1} \mathcal{M}_r(f^{-k\alpha}) \to \pi_+ \bigoplus_{k=0}^{m-1} \mathcal{M}_r(g^{-k\alpha}).$$

By taking the suitable eigenspace with respect to the $\mathbf{Z}/m\mathbf{Z}$ action on both sides, we obtain, for k = 1, a morphism of filtered \mathcal{D}_X -modules

$$\mathcal{M}_r(f^{-\alpha}) \to \pi_+ \mathcal{M}_r(g^{-\alpha})$$

Checking that it is given by the composition in the statement is an easy exercise. Strictness follows from the fact that for every morphism of mixed Hodge modules, the underlying morphism of filtered \mathcal{D} -modules is strict. The fact that τ is injective is clear if π is étale; the general case follows by restricting to an open subset U of X such that π is étale over U and using the fact that as an \mathcal{O}_X -module, $\mathcal{M}_r(f^{-\alpha})$ has no torsion.

In order to describe the filtration on $\pi_+ \mathcal{M}_r(g^{-\alpha})$, we take the usual approach, by factoring π as $q \circ \rho$, where $\rho: Y \to Y \times X$ is the graph embedding given by $\rho(y) = (y, \pi(y))$ and $q: Y \times X \to X$ is the projection onto the second component. Let us assume that we have algebraic coordinates x_1, \ldots, x_n defined on X (we can always reduce to this case by taking a suitable cover of X). Let $\pi_i = \pi^*(x_i) \in \mathcal{O}_Y(Y)$. If \mathcal{M} is a right \mathcal{D}_Y -module, then the \mathcal{D} -module push-forward via ρ is easy to compute: we have an isomorphism

$$\rho_{+}\mathcal{M} \simeq \mathcal{M} \otimes_{\mathbb{C}} \mathbb{C}[\partial_{x_{1}}, \dots, \partial_{x_{n}}], \tag{6}$$

where on the right-hand side a function $h \in \mathcal{O}_X$ acts via

$$(u \otimes 1)h = u\pi^*(h) \otimes 1$$
 for all $u \in \mathcal{M}$

and a derivation $D \in \text{Der}_{\mathbb{C}}(\mathcal{O}_Y)$ acts by

$$(u \otimes \partial_x^{\beta})D = uD \otimes \partial_x^{\beta} - \sum_{i=1}^n uD(\pi_i) \otimes \partial_x^{\beta+e_i} \quad \text{for } u \in \mathcal{M}, \ \beta \in \mathbb{Z}^n_{\geq 0},$$
(7)

where we use the multi-index notation and e_1, \ldots, e_n is the standard basis of \mathbb{Z}^n . Moreover, if $(\mathcal{M}, F_{\bullet})$ is a filtered \mathcal{D}_Y -module, then via the isomorphism (6) we have

$$F_k \rho_+ \mathcal{M} = \sum_{\beta \in \mathbb{Z}_{\geq 0}^n} F_{k-|\beta|} \mathcal{M} \otimes \partial_x^{\beta} \quad \text{for all } k \in \mathbb{Z},$$
(8)

where for $\beta = (\beta_1, \dots, \beta_n)$, we put $|\beta| = \sum_i \beta_i$.

On the other hand, the \mathcal{D} -module push-forward via q is computed by the relative Spencer complex. Given a right $\mathcal{D}_{Y \times X}$ -module \mathcal{N} , the *relative Spencer complex* of \mathcal{N} is the complex

$$C^{\bullet}(\mathcal{N}): 0 \to \mathcal{N} \otimes_{\mathcal{O}_{Y \times X}} \bigwedge^{n} p^{*}(\mathcal{T}_{Y}) \to \cdots \to \mathcal{N} \otimes_{\mathcal{O}_{Y \times X}} p^{*}(\mathcal{T}_{Y}) \xrightarrow{d_{1}} \mathcal{N} \to 0,$$

where \mathcal{T}_Y is the tangent sheaf of Y and $p: Y \times X \to Y$ is the projection onto the first component. The map d_1 , which is the only one we will need, is given by right multiplication.

If \mathcal{M} is a \mathcal{D}_Y -module, then we have a canonical isomorphism

$$\pi_+ \mathcal{M} \simeq \mathcal{H}^0(C^{\bullet}(\rho_+ \mathcal{M})) = \operatorname{coker}(\rho_+ \mathcal{M} \otimes p^*(\mathcal{T}_Y) \to \rho_+ \mathcal{M})$$

Moreover, if $(\mathcal{M}, F_{\bullet})$ is the filtered \mathcal{D}_Y -module underlying a mixed Hodge module, then via this isomorphism, the filtration on $\pi_+ \mathcal{M}$ is the quotient filtration induced by the filtration on $\rho_+ \mathcal{M}$ described in (8).

Remark 3.4. If we are in the setting of Lemma 3.3, then via the isomorphism

$$\pi_{+}\mathcal{M}_{r}(g^{-\alpha}) \simeq \operatorname{coker}\left(\rho_{+}\mathcal{M}_{r}(g^{-\alpha}) \otimes p^{*}(\mathcal{T}_{Y}) \to \rho_{+}\mathcal{M}_{r}(g^{-\alpha})\right),$$
(9)

the morphism τ maps $u \in \mathcal{M}_r(f^{-\alpha})$ to the class of $\pi^*(u) \otimes 1 \in \rho_+ \mathcal{M}_r(g^{-\alpha})$. Indeed, since as an \mathcal{O}_X -module $\pi_+ \mathcal{M}_r(g^{-\alpha})$ has no torsion by Lemma 3.2, in order to check the assertion we may restrict to an open subset U of X such that π is étale over U; in this case the assertion follows via an easy computation.

Theorem 3.5. Let $\pi: Y \to X$ be a finite surjective morphism between smooth *n*-dimensional complex algebraic varieties. If $f \in \mathcal{O}_X(X)$ is nonzero and $g = \pi^*(f) \in \mathcal{O}_Y(Y)$, then for every $\alpha \in \mathbf{Q}_{>0}$ and $k \in \mathbf{Z}$, we have the following inclusion:

$$\{u \in \mathcal{M}_r(f^{-\alpha}) \mid \pi^*(u) \in F_k \mathcal{M}_r(g^{-\alpha})\} \subseteq F_k \mathcal{M}_r(f^{-\alpha}).$$

Note that if we assume that f and g define reduced divisors, then by passing from right to left \mathcal{D} -modules and using the definition of Hodge ideals, we obtain the assertion in Theorem 1.3.

Proof of Theorem 3.5. After taking a suitable open cover of X, we may assume that we have algebraic coordinates defined on X. If $u \in \mathcal{M}_r(f^{-\alpha})$ is such that $\pi^*(u) \in F_k \mathcal{M}_r(g^{-\alpha})$, then it follows from the above description of the filtration on $\pi_+ \mathcal{M}_r(g^{-\alpha})$ that the class of $\pi^*(u) \otimes 1$ in $\operatorname{coker}(\rho_+ \mathcal{M}_r(g^{-\alpha}) \otimes p^*(\mathcal{T}_Y) \to \rho_+ \mathcal{M}_r(g^{-\alpha}))$ lies in $F_k(\pi_+ \mathcal{M}_r(g^{-\alpha}))$. Since this is equal to $\tau(u)$ and τ is strict by Lemma 3.3, we conclude that $u \in F_k \mathcal{M}_r(f^{-\alpha})$.

Our next goal is to understand how far the inclusion in Theorem 3.5 is from being an equality. We will do this under the extra assumption that π is Galois. This result will not be used in the following sections.

Let $\pi: Y \to X$ be a finite surjective morphism between smooth complex algebraic varieties. Recall that π is Galois if the corresponding field extension $\mathbf{C}(X) \hookrightarrow \mathbf{C}(Y)$ is normal (hence Galois, since we are in characteristic 0). Suppose that this is the case and let *G* be the Galois group of this field extension. Since *Y* is the integral closure of *X* in $\mathbf{C}(Y)$, we have an induced action of *G* on *Y* and π is the quotient morphism with respect to this action.

Theorem 3.6. Let $\pi: Y \to X$ be a Galois finite surjective morphism between smooth *n*-dimensional complex algebraic varieties. Suppose that $f \in \mathcal{O}_X(X)$ is nonzero and let $g = \pi^*(f) \in \mathcal{O}_Y(Y)$. For every $\alpha \in \mathbf{Q}_{>0}$ and $k \in \mathbf{Z}$, put

$$F'_k \mathcal{M}_r(f^{-\alpha}) := \{ u \in \mathcal{M}_r(f^{-\alpha}) \mid \pi^*(u) \in F_k \mathcal{M}_r(g^{-\alpha}) \}.$$

Then, for every $k \in \mathbb{Z}$,

$$F_k \mathcal{M}_r(f^{-\alpha}) = \sum_{i \ge 0} F'_{k-i} \mathcal{M}_r(f^{-\alpha}) \cdot F_i \mathcal{D}_X.$$
 (10)

In particular,

$$F_k \mathcal{M}_r(f^{-\alpha}) = F_{k-1} \mathcal{M}_r(f^{-\alpha}) \cdot F_1 \mathcal{D}_X + F'_k \mathcal{M}_r(f^{-\alpha}).$$

Proof. After taking a suitable open cover of X, we may and will assume that X (hence also Y) is affine and we have algebraic coordinates x_1, \ldots, x_n defined on X. We may thus use the description of the filtration on $\pi_+ \mathcal{M}_r(g^{-\alpha})$ that we have discussed.

Note that the action of G on Y induces a G-action on $\mathcal{O}_Y(Y)$. Moreover, it also induces a G-action on $\Gamma(Y, \omega_Y)$ that makes ω_Y a G-equivariant sheaf.² Moreover, its subspace of invariant sections is precisely the image of the map

$$\Gamma(X, \omega_X) \to \Gamma(Y, \omega_Y)$$

given by the pull-back of differential forms (see [3, Theorem 1]). Since g is a G-invariant section of \mathcal{O}_Y , it follows that the sheaf $\mathcal{M}_r(g^{-\alpha})$ has an induced structure of G-equivariant sheaf; moreover, its subspace of G-invariant sections is the image of the pull-back map

$$\Gamma(X, \mathcal{M}_r(f^{-\alpha})) \to \Gamma(Y, \mathcal{M}_r(g^{-\alpha})).$$

We also have an induced *G*-action on $\Gamma(X, \pi_+\mathcal{M}_r(g^{-\alpha}))$. To see this, note that we have a natural *G*-action on $\Gamma(Y \times X, \rho_+\mathcal{M}_r(g^{-\alpha})) = \Gamma(Y, \mathcal{M}_r(g^{-\alpha})) \otimes_{\mathbb{C}} \mathbb{C}[\partial_{x_1}, \dots, \partial_{x_n}]$, where *G* acts trivially on $\mathbb{C}[\partial_{x_1}, \dots, \partial_{x_n}]$. The *G*-action on *Y* induces an action of *G* on $\Gamma(Y, \mathcal{T}_Y)$ such that the multiplication map $\rho_+\mathcal{M}_r(g^{-\alpha}) \otimes \mathcal{T}_Y \to \rho_+\mathcal{M}_r(g^{-\alpha})$ is compatible with the *G*-actions. We thus get an induced *G*-action on the cokernel of this map and thus on $\Gamma(X, \pi_+\mathcal{M}_r(g^{-\alpha}))$ via the isomorphism (9). It also follows from the description of the canonical morphism

$$\tau: \Gamma(X, \mathcal{M}_r(f^{-\alpha})) \to \Gamma(X, \pi_+ \mathcal{M}_r(g^{-\alpha}))$$

in Remark 3.4 that the image of τ lands in the subspace of *G*-invariant sections. A key point is that due to the fact that *g* is *G*-invariant, the filtration on $\Gamma(Y, \mathcal{M}_r(g^{-\alpha}))$ is preserved by the *G*-action and therefore so is the filtration on $\Gamma(X, \pi_+ \mathcal{M}_r(g^{-\alpha}))$.

²Since *G* is a finite group and *Y* is affine, this simply means that the scalar multiplication $\mathcal{O}_Y(Y) \times \Gamma(Y, \omega_Y) \to \Gamma(Y, \omega_Y)$ is compatible with the *G*-actions.

Given a complex vector space V with a G-action, we consider the linear map $A = A_V: V \to V$ given by $A(v) = \frac{1}{|G|} \sum_{g \in G} gv$. Note that A(v) lies in the subspace V^G of invariant elements for every $v \in V$ and A(v) = v if $v \in V^G$.

The inclusion " \supseteq " in (10) is clear: it follows from Theorem 3.5 that for every j we have $F'_j \mathcal{M}_r(f^{-\alpha}) \subseteq F_j \mathcal{M}_r(f^{-\alpha})$ and thus

$$F'_{k-i}\mathcal{M}_r(f^{-\alpha})\cdot F_i\mathcal{D}_X\subseteq F_k\mathcal{M}_r(f^{-\alpha})$$

We now prove the reverse inclusion. Given a global section u of $F_k \mathcal{M}_r(f^{-\alpha})$, it follows from Lemma 3.3 that $\tau(u)$ is a global section of $F_k(\pi_+\mathcal{M}_r(g^{-\alpha}))$. This means that we can find global sections w_β of $F_{k-|\beta|}\mathcal{M}_r(g^{-\alpha})$ for $\beta \in \mathbb{Z}_{\geq 0}^n$, with $w_\beta = 0$ for all but finitely many β , such that

$$\pi^*(u) \otimes 1 - \sum_{\beta} w_{\beta} \otimes \partial_x^{\beta} \in \operatorname{Im}(\rho_+ \mathcal{M}_r(g^{-\alpha}) \otimes \mathcal{T}_Y \to \rho_+ \mathcal{M}_r(g^{-\alpha})).$$

If we put $w'_{\beta} = A(w_{\beta}) \in \Gamma(Y, F_{k-|\beta|}\mathcal{M}_r(g^{-\alpha}))^G$, we see that

$$\pi^*(u) \otimes 1 - \sum_{\beta} w'_{\beta} \otimes \partial_x^{\beta} \in \operatorname{Im}(\rho_+ \mathcal{M}_r(g^{-\alpha}) \otimes \mathcal{T}_Y \to \rho_+ \mathcal{M}_r(g^{-\alpha}))$$

Since each w'_{β} is *G*-invariant, it follows that we can write $w'_{\beta} = \pi^*(u_{\beta})$ for some global section u_{β} of $F'_{k-|\beta|}\mathcal{M}_r(f^{-\alpha})$. We now claim that for every β , $\pi^*(u_{\beta}) \otimes \partial_x^{\beta}$ and $\pi^*(u_{\beta}\partial_x^{\beta}) \otimes 1$ have the same image in $\Gamma(X, \pi_+\mathcal{M}_r(g^{-\alpha}))$. If this is the case, it follows that $\tau(u) = \tau(\sum_{\beta} u_{\beta}\partial_x^{\beta})$. Since τ is injective by Lemma 3.3, we conclude that $u \in \sum_{i\geq 0} F'_{k-i}\mathcal{M}_r(f^{-\alpha}) \cdot F_i\mathcal{D}_X$.

In order to prove the above claim, since $\pi_+ \mathcal{M}_r(g^{-\alpha})$ is an \mathcal{O}_X -module with no torsion by Lemma 3.2, it is enough to prove it on some nonempty open subset of X. We thus may and will assume that π is étale. In this case, if $\pi_i = x_i \circ \pi$, then π_1, \ldots, π_n give an algebraic system of coordinates on Y and we have a corresponding system of derivations $\partial_{\pi_1}, \ldots, \partial_{\pi_n}$. Of course, arguing by induction on $|\beta|$, it is enough to show that for every global section η of $\mathcal{M}_r(f^{-\alpha})$, the elements $\pi^*(\eta \partial_{x_i}) \otimes 1$ and $\pi^*(\eta) \otimes \partial_{x_i}$ in $\rho_+ \mathcal{M}_r(g^{-\alpha})$ have the same image in $\pi_+ \mathcal{M}_r(g^{-\alpha})$. This follows from the fact that by (7), the map $\rho_+ \mathcal{M}_r(g^{-\alpha}) \otimes \mathcal{T}_Y \to \rho_+ \mathcal{M}_r(g^{-\alpha})$ maps $\pi^*(\eta) \otimes \partial_{\pi_i}$ to

$$\pi^*(\eta)\partial_{\pi_i}\otimes 1-\pi^*(\eta)\otimes \partial_{x_i}=\pi^*(\eta\partial_{x_i})\otimes 1-\pi^*(\eta)\otimes \partial_{x_i}.$$

This completes the proof of (10) and the last assertion in the theorem is an immediate consequence.

We now give a consequence of the results in Theorems 1.3 and 3.6 for minimal exponents.

Corollary 3.7. Let $\pi: Y \to X$ be a finite surjective morphism between smooth complex algebraic varieties and let $K_{Y/X}$ be the effective divisor on Y locally defined by the determinant of a Jacobian matrix of π . If $0 \neq f \in \mathcal{O}_X(X)$ and $g = f \circ \pi$ both define reduced

divisors, then the following hold for every *P* in *X* with f(P) = 0, every nonnegative integer *p*, and every $\alpha \in \mathbf{Q} \cap (0, 1]$:

- (i) If $\mathcal{O}_Y(-K_{Y/X}) \subseteq I_p(g^{\alpha})$ in a neighborhood of the fiber $\pi^{-1}(P)$, then $\tilde{\alpha}_P(f) \ge p + \alpha$.
- (ii) If π is Galois and the hypersurface defined by f is not smooth at P, then the converse of (i) holds: if α̃_P(f) ≥ p + α, then 𝒪_Y(−K_{Y/X}) ⊆ I_p(g^α) in a neighborhood of the fiber π⁻¹(P).

Remark 3.8. Note that a similar assertion holds for log canonical thresholds of arbitrary regular functions in the setting of a finite surjective morphism between smooth varieties, as above: we have $\operatorname{lct}_P(f) > \lambda$ if and only if $\mathcal{O}_Y(-K_{Y/X})$ is contained in the multiplier ideal $\mathcal{J}(g^{\lambda})$ in a neighborhood of the fiber $\pi^{-1}(P)$. This follows from the fact that $\operatorname{lct}_P(f) > \lambda$ if and only if $\mathcal{J}(f^{\lambda}) = \mathcal{O}_X$ in a neighborhood of P and the theorem relating the multiplier ideals of f and g (see [14, Theorem 9.5.42]).

Proof of Corollary 3.7. The first assertion follows directly from Theorem 1.3 and the characterization of $\tilde{\alpha}_P(f)$ via the Hodge ideals of f in (3). Suppose now that we are in the setting of (ii). Using again the characterization of $\tilde{\alpha}_P(f)$ via the Hodge ideals of f, we conclude using the hypothesis that $I_p(f^{\alpha}) = \mathcal{O}_X$ in a neighborhood of P. Equivalently, we have

$$\mathcal{O}_X \cdot \frac{1}{f^p} f^{-\alpha} = F_p \mathcal{M}(f^{-\alpha}).$$
(11)

For every $k \in \mathbb{Z}$, let us put

$$F'_k\mathcal{M}(f^{-\alpha}) = \left\{ u = \frac{h}{f^p} f^{-\alpha} \in F_k\mathcal{M}(f^{-\alpha}) \ \middle| \ \frac{\pi^*(h)}{g^p} \mathcal{O}_Y(-K_{Y/X}) \cdot g^{-\alpha} \subseteq F_k\mathcal{M}(g^{-\alpha}) \right\}.$$

It follows from equation (10) in Theorem 3.6 (after passing from right to left \mathcal{D} -modules) that

$$F_p \mathcal{M}(f^{-\alpha}) = F'_p \mathcal{M}(f^{-\alpha}) + F_1 \mathcal{D}_X \cdot F_{p-1} \mathcal{M}(f^{-\alpha}).$$

We next show that since f has a singular point at P, we have

$$F_1 \mathcal{D}_X \cdot F_{p-1} \mathcal{M}(f^{-\alpha}) \subseteq \mathfrak{m}_P \cdot \frac{1}{f^p} f^{-\alpha}, \qquad (12)$$

where \mathfrak{m}_P is the ideal of functions vanishing at P. In order to see this, let us choose local coordinates x_1, \ldots, x_n centered at P. First, recall that

$$F_{p-1}\mathcal{M}(f^{-\alpha}) \subseteq \mathcal{O}_X \cdot \frac{1}{f^{p-1}}f^{-\alpha} \subseteq \mathfrak{m}_P \cdot \frac{1}{f^p}f^{-\alpha},$$

where the second inclusion follows from the fact that $f \in \mathfrak{m}_P$. Moreover, for every *i* and every $h \in \mathcal{O}_X$, we have

$$\partial_{x_i} \cdot \frac{h}{f^{p-1}} f^{-\alpha} = \frac{\partial h}{\partial x_i} \cdot \frac{f}{f^p} f^{-\alpha} - (\alpha + p - 1) \frac{\partial f}{\partial x_i} \cdot \frac{h}{f^p} f^{-\alpha} \in \mathfrak{m}_P \cdot \frac{1}{f^p} f^{-\alpha},$$

where we use the fact that $f, \frac{\partial f}{\partial x_i} \in \mathfrak{m}_P$. This proves (12).

We thus have the \mathcal{O}_X -submodule $F'_p \mathcal{M}(f^{-\alpha})$ of $\mathcal{O}_X \cdot \frac{1}{f^p} f^{-\alpha}$ and (11) and (12) give

$$\mathcal{O}_X \cdot \frac{1}{f^p} f^{-\alpha} \subseteq F'_p \mathcal{M}(f^{-\alpha}) + \mathfrak{m}_P \cdot \frac{1}{f^p} f^{-\alpha}.$$

We deduce using Nakayama's lemma that $\frac{1}{f^p}f^{-\alpha} \in F'_p\mathcal{M}(f^{-\alpha})$ around P, that is, $\mathcal{O}_Y(-K_{Y/X}) \subseteq I_p(g^{\alpha})$.

4. Hodge ideals for families with constant Milnor number

Let *X* be a smooth *n*-dimensional complex algebraic variety and $f \in \mathcal{O}_X(X)$ a nonzero regular function. Let $P \in X$ be a point with f(P) = 0; we assume that *P* is a singular point of *f* and that *f* has an isolated singularity at *P*. Recall that J_f denotes the Jacobian ideal of *f*, generated by $\frac{\partial f}{\partial x_1}, \ldots, \frac{\partial f}{\partial x_n}$, where x_1, \ldots, x_n are local coordinates on *X*. Our assumption on *f* implies that the dimension $\mu = \mu_P(f) := \dim_{\mathbb{C}}(\mathcal{O}_{X,P}/J_{f,P})$ is a finite positive number; this is the *Milnor number* of *f* at *P*. We note that a more natural context for the discussion in this section is that of holomorphic functions on complex manifolds; however, this is not really that different from the algebraic case since we deal with isolated singularities, so that we prefer to stick to the algebraic case, as in the rest of the article.

For basic facts on the spectrum of f and its connection to the *V*-filtration and the mixed Hodge structure on the cohomology of the Milnor fiber of f at P, we refer to [9, Section 1] and the references therein. We only recall that if $F_{f,P}$ denotes the Milnor fiber of f at P, then on its cohomology $H^{n-1}(F_{f,P}, \mathbf{Q})$ there is a mixed Hodge structure and a compatible action of the monodromy T (the *inverse* of the Milnor monodromy). Note that dim_{**Q**} $H^{n-1}(F_{f,P}, \mathbf{Q}) = \mu$. If T_s is the semisimple part of the monodromy and if λ is an eigenvalue of T_s on the above cohomology, then $H^{n-1}(F_{f,P}, \mathbf{C})_{\lambda}$ denotes the corresponding eigenspace.

The *spectrum* of f at P is a collection of μ positive rational numbers, not necessarily distinct and indexed nondecreasingly $\alpha_1, \ldots, \alpha_{\mu}$ such that for every $\beta \in \mathbf{Q}_{>0}$, we have

$$#\{i \mid \alpha_i = \beta\} = \dim_{\mathbf{C}} \operatorname{Gr}_F^q H^{n-1}(F_{f,P}, \mathbf{C})_{\lambda},$$
(13)

where $\lambda = \exp(-2\pi i\beta)$ and $q = n - \lceil\beta\rceil$. Note that α_1 appears with multiplicity 1 and it is equal to $\tilde{\alpha}_P(f)$.

Let us review now the connection between the Hodge filtration on $H^{n-1}(F_{f,P}, \mathbf{Q})$ and the *V*-filtration on the Brieskorn lattice. Recall that the *Brieskorn lattice* of *f* at *P* is

$$H_{f,P}'' := \Omega_{X,P}^n / df \wedge d \Omega_{X,P}^{n-2}$$

where the stalks of the sheaves of differential forms are the analytic ones (associated to the complex manifold X^{an}). This has a structure of free module of rank μ over C{{t}} and over C{{ d_t^{-1} }}, where the action of t is given by $t \cdot [\omega] = [f\omega]$ and the action of ∂_t^{-1} is given by $\partial_t^{-1} \cdot [\omega] = [df \wedge \eta]$, where $d\eta = \omega$. The *Gauss–Manin system* of f at P is

$$G_{f,P} := H_{f,P}''[\partial_t].$$

Note that we have an injective map $H''_{f,P} \hookrightarrow G_{f,P}$ and a surjective map

$$H_{f,P}'' \to \Omega_{f,P}^n := \Omega_{X,P}^n / df \wedge \Omega_{X,P}^{n-1}.$$

A choice of local coordinates x_1, \ldots, x_n at P gives an isomorphism

$$\Omega_{f,P}^n \simeq \mathcal{O}_{X,P}/J_{f,P}$$

On the Gauss–Manin system there is a *V*-filtration, similar to the one discussed in Section 2, such that $\partial_t t - \beta$ is nilpotent on $\operatorname{Gr}_V^\beta G_{f,P}$ for all $\beta \in \mathbf{Q}$. This *V*-filtration induces a *V*-filtration on the submodule $H_{f,P}^{"}$ and then a quotient filtration on $\Omega_{f,P}^n$. An important fact is that for every $\beta \in \mathbf{Q}$, we have an isomorphism

$$\operatorname{Gr}_{V}^{\beta}\Omega_{f,P}^{n} \simeq \operatorname{Gr}_{F}^{q} H^{n-1}(F_{f,P}, \mathbf{C})_{\lambda},$$
(14)

where $\lambda = \exp(-2\pi i\beta)$ and $q = n - \lceil \beta \rceil$ (see [9, (1.2.2)]).

The key result for us is that via the isomorphism $\Omega_{f,P}^n \simeq \mathcal{O}_{X,P}/J_{f,P}$, the quotient *V*-filtration on $\Omega_{f,P}^n$ coincides with the quotient filtration on $\mathcal{O}_{X,P}/J_{f,P}$ induced by the microlocal *V*-filtration: for every $\beta \in \mathbf{Q}$, the isomorphism identifies

$$V^{\beta}\Omega^{n}_{f,P}$$
 with $(\tilde{V}^{\beta}\mathcal{O}_{X,P} + J_{f,P})/J_{f,P}$ (15)

(see [9, Proposition 1.4]).

Recall now that if $\beta' \leq \tilde{\alpha}_P(f) + 1$, then $J_f \subseteq \tilde{V}^{\beta'} \mathcal{O}_X$ around *P* (see Proposition 2.7). We thus deduce from (14) and (15) that if $\beta < \tilde{\alpha}_P(f) + 1$, then we have an isomorphism

$$\widetilde{V}^{\beta}\mathcal{O}_{X,P}/\widetilde{V}^{>\beta}\mathcal{O}_{X,P}\simeq \mathrm{Gr}_{F}^{q}H^{n-1}(F_{f,P},\mathbf{C})_{\lambda},$$

where $\lambda = \exp(-2\pi i\beta)$ and $q = n - \lceil\beta\rceil$. We thus obtain the following:

Proposition 4.1. If f has an isolated singularity at P and $\beta < \tilde{\alpha}_P(f) + 1$, then β is a jumping number for the microlocal multiplier ideals of f at P (that is, $\tilde{V}^{\beta}\mathcal{O}_{X,P} \neq \tilde{V}^{>\beta}\mathcal{O}_{X,P}$) if and only if β is in the spectrum of f at P. More precisely, the multiplicity of β in this spectrum is equal to $\dim_{\mathbb{C}}(\tilde{V}^{\beta}\mathcal{O}_{X,P}/\tilde{V}^{>\beta}\mathcal{O}_{X,P})$.

We also obtain the result stated in the Introduction.

Proof of Theorem 1.4. If $\mu_{s(t)}(f_t) = 0$ (that is, if the hypersurface defined by f_t is smooth at s(t)), then the assertions in the theorem are trivial, since all Hodge ideals coincide with the corresponding structure sheaves. Hence from now on we assume that $\mu_{s(t)}(f_t) > 0$. By Varchenko's theorem [35], the constancy of the Milnor number implies that the spectrum of f_t is independent of t. Since by assumption $p + \gamma \le \alpha + 1$, it follows from Proposition 2.7 that

$$I_p(f_t^{\gamma}) = \tilde{V}^{p+\gamma} \mathcal{O}_{\mathcal{X}_t}(f_t) \supseteq J_{f_t}$$

(note that it is enough to check these at s(t), since the hypersurface defined by f is smooth away from this point). By the definition of the spectrum (see (13)) and using (14) and (15),

we conclude that

$$\dim_{\mathbb{C}}(\mathcal{O}_{\mathcal{X}_t}/I_p(f_t^{\gamma})) \quad \text{is independent of } t \in T.$$
(16)

Since each hypersurface defined by every f_t is reduced, it follows that the hypersurface defined by f is reduced as well. Let Z be the closed subscheme of X defined by $I_p(f^{\gamma})$ and $\tau: Z \to T$ the morphism induced by φ . Note that τ is finite, being proper, with finite fibers: in fact, the support of Z is a closed subset of s(T).

Using the Restriction Theorem for Hodge ideals (see [21, Theorem A (vi)]), we see that

$$I_p(f_t^{\gamma}) \subseteq I_p(f^{\gamma}) \cdot \mathcal{O}_{\mathfrak{X}_t} \quad \text{for all } t \in T,$$
(17)

with equality for general $t \in T$. For the finite morphism τ , we know that the map

$$T \ni t \mapsto \dim_{\mathbb{C}}(\mathcal{O}_{\mathfrak{X}_t}/I_p(f^{\gamma}) \cdot \mathcal{O}_{\mathfrak{X}_t})$$

is upper-semicontinuous, and it is constant if and only if τ is flat. We thus deduce from (16) and (17) that the inclusion in (17) is an equality for all $t \in T$ and that τ is flat. This completes the proof of the theorem.

5. Proof of Teissier's conjecture

Before giving the proof of Theorem 1.1, we make some preliminary remarks. We begin by reviewing some facts about semicontinuity of minimal exponents, which we will also use in the next section.

Suppose first that we have a smooth morphism $\varphi: \mathcal{X} \to T$ of complex algebraic varieties and $s: T \to \mathcal{X}$ is a section of φ . Suppose that $f \in \mathcal{O}_{\mathcal{X}}(\mathcal{X})$ is such that for every $t \in T$, the restriction f_t of f to $\mathcal{X}_t = \varphi^{-1}(t)$ is nonzero. We assume that $f \circ s = 0$, hence we may consider $\tilde{\alpha}_{s(t)}(f_t)$ for all $t \in T$. In this case, the function

$$T \ni t \mapsto \widetilde{\alpha}_{s(t)}(f_t)$$

is lower semicontinuous by [22, Theorem E (2)] (when each f_t has an isolated singularity at s(t), this result was also proved in [30, Theorem 2.11]). In fact, the proof in [22] shows something stronger: for every $\alpha > 0$, the set { $t \in T \mid \tilde{\alpha}_{s(t)}(f_t) \ge \alpha$ } is open in T. Since a countable intersection of nonempty open subsets is nonempty, it follows that the set { $\tilde{\alpha}_{s(t)}(f_t) \mid t \in T$ } has a maximum, which is achieved on an open subset of T.

We next make two remarks concerning the hypothesis in Theorem 1.1.

Remark 5.1. In the statement of Theorem 1.1, we may also assume that the hypersurface defined by $f|_H$ in H has an isolated singularity at P. For this, it is enough to show that there is a smooth hypersurface H' containing P, with the hypersurface defined by $f|_{H'}$ having an isolated singularity at P, and such that $\tilde{\alpha}_P(f|_{H'}) \ge \tilde{\alpha}_P(f|_H)$. It is clear that in the statement of the theorem we may assume that X is affine and that we have a system of algebraic coordinates x_1, \ldots, x_n on X, centered at P, such that H is generated by x_1 .

If H' is defined by $a_1x_1 + \cdots + a_nx_n$, where $a_1, \ldots, a_n \in \mathbb{C}$ are general, then it follows from the semicontinuity of minimal exponents that $\tilde{\alpha}_P(f|_{H'}) \ge \tilde{\alpha}_P(f|_H)$. On the other hand, since the hypersurface Y defined by f has an isolated singularity at P, it is easy to see that also the hypersurface in H' defined by $f|_{H'}$ has an isolated singularity at P. This proves our assertion.

Remark 5.2. In order to prove Theorem 1.1, it is enough to consider the case when $X = \mathbf{A}^n$, P = 0, and H is the hyperplane defined by $x_n = 0$. Indeed, we may assume that X is affine and we have a system of algebraic coordinates x_1, \ldots, x_n centered at P such that H is defined by x_n . In this case, the map $\sigma = (x_1, \ldots, x_n)$: $X \to \mathbf{A}^n$ is étale, $\sigma(P) = 0$, and H is the inverse image of the hyperplane H', defined by the vanishing of the last coordinate. By Remark 5.1, we may also assume that $f \mid_H$ has isolated singularities. For every $d \ge 1$, there is $f_d \in \mathcal{O}_{\mathbf{A}^n}(\mathbf{A}^n)$ such that $f - \sigma^*(f_d) \in (x_1, \ldots, x_n)^d$. Since f has an isolated singularity at P, it follows that f and $\sigma^*(f_d)$ differ by an analytic change of coordinates for $d \gg 0$ (see [7, Corollary 2.24]); since both the minimal exponent and Teissier's invariant $\theta_P(f)$ can be computed by passing to the local ring of the corresponding complex manifold, we have

$$\widetilde{\alpha}_P(f) = \widetilde{\alpha}_P(\sigma^*(f_d))$$
 and $\theta_P(f) = \theta_P(\sigma^*(f_d))$.

The same argument implies that $\tilde{\alpha}_P(f|_H) = \tilde{\alpha}_P(\sigma^*(f_d)|_H)$ for $d \gg 0$. On the other hand, since σ induces a biholomorphic map in a neighborhood of P, we also have

$$\widetilde{\alpha}_P(\sigma^*(f_d)) = \widetilde{\alpha}_0(f_d), \quad \widetilde{\alpha}_P(\sigma^*(f_d)|_H) = \widetilde{\alpha}_0(f_d|_{H'}), \quad \text{and} \quad \theta_P(\sigma^*(f_d)) = \theta_0(f_d).$$

This completes the proof of our assertion.

We can now give the proof of our main result.

Proof of Theorem 1.1. It follows from Remarks 5.1 and 5.2 that we may and will assume that $X = \mathbf{A}^n$, with coordinates x_1, \ldots, x_n , H is the hyperplane defined by $x_n = 0$, and P is the origin; moreover, $g = f(x_1, \ldots, x_{n-1}, 0)$ has an isolated singularity at 0 in \mathbf{A}^{n-1} . We choose a positive integer d such that $m := d(\theta_P(f) + 1)$ is an integer. The case when the hypersurface defined by f is smooth at P is trivial, since in this case $\tilde{\alpha}_P(f) = \infty$. Therefore from now on we assume that this hypersurface is singular at P. We thus have $J_f \subseteq (x_1, \ldots, x_n)$, hence $\theta_P(f) \ge 1$. Let $\lambda = \tilde{\alpha}_P(g) + \frac{1}{\theta_P(f)+1}$, so that we need to show that $\tilde{\alpha}_P(f) \ge \lambda$.

If n = 2, then $\tilde{\alpha}_P(f) = \operatorname{lct}_P(f)$ since by [28, Theorem (0.4)], we have $\tilde{\alpha}_P(f) \le n/2 = 1$. Therefore the assertion in the theorem follows from the main result in [6]. Hence from now on we assume $n \ge 3$. In this case, since g has an isolated singularity at 0, it follows that it is reduced in a neighborhood of 0.

As we have explained in the Introduction, we follow the approach in [6], replacing log canonical thresholds and multiplier ideals by minimal exponents and Hodge ideals, respectively. For technical reasons, in our setting it is important to consider the following two-parameter family of hypersurfaces: let

$$h(x_1,...,x_n,y,z) = f(x_1,...,x_{n-1},yx_n^d) + zx_n^m \in \mathbf{C}[x_1,...,x_n,y,z]$$

and for every $s, t \in \mathbb{C}$ we consider $h_{s,t} = h|_{y=s,z=t} \in \mathbb{C}[x_1, \dots, x_n]$. Note that for every s and t we have $h_{s,t}|_H = f|_H$. First, since $f|_H$ is reduced in a neighborhood of P, it follows that every $h_{s,t}$ is reduced in a neighborhood of P and h is reduced in a neighborhood of $\{P\} \times \mathbb{A}^2$. We can thus use the results in Section 2 for the Hodge ideals of $h_{s,t}$ and h around the respective subsets. Second, we have

$$\widetilde{\alpha}_P(h_{s,t}) \ge \widetilde{\alpha}_P(g) \quad \text{and} \quad \widetilde{\alpha}_{(P,s,t)}(h) \ge \widetilde{\alpha}_P(g) \quad \text{for all } s, t \in \mathbb{C}$$
 (18)

since the minimal exponent does not go up under restriction to a smooth hypersurface, see [22, Theorem E(1)]. Since $\lambda - \tilde{\alpha}_P(g) \le 1/2$, we deduce from Proposition 2.7 that if we write

$$\lambda = p + \gamma$$
 with $p = \lceil \lambda \rceil - 1$

then

$$I_{p}(h_{s,t}^{\gamma}) = \tilde{V}^{\lambda} \mathcal{O}_{\mathbf{A}^{n}}(h_{s,t}) \quad \text{around } P \text{ for all } s, t \in \mathbf{C},$$
$$I_{p}(h^{\gamma}) = \tilde{V}^{\lambda} \mathcal{O}_{\mathbf{A}^{n+2}}(h) \quad \text{around } \{P\} \times \mathbf{A}^{2}.$$

We put

$$I := I_p(h^{\gamma}) \subseteq S = \mathbf{C}[x_1, \dots, x_n, y, z].$$

We consider on S the grading such that $wt(x_n) = 1$, wt(y) = -d, wt(z) = -m, and $wt(x_i) = 0$ for $1 \le i \le n - 1$, so that h is homogeneous of weight 0. This implies that I is a graded ideal of S (since the hypersurface defined by h is preserved by the C^{*}-action corresponding to the grading of S, the same holds for the Hodge ideals of this hypersurface).

Step 1. We first show that $x_n^{d-1} \in I_p(h|_{y=0}^{\gamma})$ around $\{P\} \times \{0\} \times \mathbf{A}^1$. Note that $h|_{y=0} = g(x_1, \ldots, x_{n-1}) + zx_n^m$. We thus deduce from the Thom–Sebastiani theorem for microlocal multiplier ideals (see [20, Theorem 2.2]) that around $\{P\} \times \{0\} \times \mathbf{A}^1$ we have

$$\widetilde{V}^{\lambda}\mathcal{O}_{\mathbf{A}^{n+1}}(h|_{y=0}) = \sum_{\beta_1+\beta_2=\lambda} \left(\widetilde{V}^{\beta_1}\mathcal{O}_{\mathbf{A}^{n-1}}(g) \cdot \mathbf{C}[x_1,\ldots,x_n,z] \right) \cdot \left(\widetilde{V}^{\beta_2}\mathcal{O}_{\mathbf{A}^2}(zx_n^m) \cdot \mathbf{C}[x_1,\ldots,x_n,z] \right).$$
(19)

By the characterization of the minimal exponent in terms of microlocal multiplier ideals (see (2)), we have $\tilde{V}^{\beta_1} \mathcal{O}_{A^{n-1}}(g) = \mathbb{C}[x_1, \dots, x_{n-1}]$ around the origin for $\beta_1 = \tilde{\alpha}_0(g)$. On the other hand, if $\beta_2 = \frac{1}{\theta_P(f)+1}$, then $\beta_2 < 1$ and thus the corresponding microlocal multiplier ideal is a usual multiplier ideal for a simple normal crossing divisor, which is easy to compute:

$$\begin{split} \widetilde{V}^{\beta_2} \mathcal{O}_{\mathbf{A}^2}(z x_n^m) &= \mathcal{J}(\mathbf{A}^2, (z x_n^m)^{\beta_2 - \epsilon}) \quad \text{for } 0 < \epsilon \ll 1 \\ &= \mathcal{J}(\mathbf{A}^2, z^{\beta_2 - \epsilon} x_n^{m(\beta_2 - \epsilon)}) = (x_n)^{\lfloor m(\beta_2 - \epsilon) \rfloor} = (x_n^{d-1}) \end{split}$$

(recall that $m/(\theta_P(f)+1) = d$). We thus conclude from (19) that around $\{P\} \times \{0\} \times \mathbf{A}^1$, x_n^{d-1} lies in $\tilde{V}^{\lambda} \mathcal{O}_{\mathbf{A}^{n+1}}(h|_{y=0}) = I_p(h|_{y=0}^{\gamma})$.

Step 2. We next show that there is an open subset U of \mathbf{A}^2 such that for every $(s, t) \in U$, we have $x_n^{d-1} \in I_p(h_{s,t}^{\gamma})$ around P. Note first that we have

$$I_p(h|_{y=0}^{\gamma}) \subseteq I \cdot \mathcal{O}_{\mathbf{A}^{n+2}}|_{y=0} \text{ around } \{P\} \times \{0\} \times \mathbf{A}^1.$$

This is a consequence of the Restriction Theorem for Hodge ideals in [21, Theorem A (vi)]. We deduce from Step 1 that $x_n^{d-1} \in I \cdot \mathcal{O}_{\mathbf{A}^{n+2}}|_{y=0}$ around $\{P\} \times \{0\} \times \mathbf{A}^1$. This implies that there is a polynomial $q \in \mathbf{C}[x_1, \ldots, x_n, z]$ such that $q \cdot x_n^{d-1} \in I \cdot \mathcal{O}_{\mathbf{A}^{n+2}}|_{y=0}$ and $1 - q \in (x_1, \ldots, x_n, z]$. Indeed, for every $a \in \mathbf{C}$ we have a polynomial $q_a \in \mathbf{C}[x_1, \ldots, x_n, z]$ such that $q_a \cdot x_n^{d-1} \in I \cdot \mathcal{O}_{\mathbf{A}^{n+2}}|_{y=0}$ and $q_a(0, a) \neq 0$. If we write $q_a = q'_a + q''_a$ with

$$q'_a \in (x_1, \ldots, x_n) \mathbf{C}[x_1, \ldots, x_n, z]$$
 and $q''_a \in \mathbf{C}[z]$

we see that the gcd of the q_a'' is 1, hence they generate the unit ideal. This immediately implies the existence of q as asserted.

We can thus find polynomials $Q_1, Q_2 \in S$ with $Q_1 \in I$ such that

$$q \cdot x_n^{d-1} = Q_1 + y Q_2. \tag{20}$$

If we use the grading that we defined on S and for any polynomial $Q \in S$, we denote by Q_j the homogeneous component of Q of weight j, then we can write

$$q_0 = 1 + \sum_{i \ge 0} z^i x_n^{mi} u_i(x_1, \dots, x_{n-1}) \quad \text{with } u_0 \in (x_1, \dots, x_{n-1}) \mathbb{C}[x_1, \dots, x_{n-1}],$$
$$(yQ_2)_{d-1} = y \cdot \sum_{j,k \ge 0} z^j y^k x_n^{mj+d(k+2)-1} v_{j,k}(x_1, \dots, x_{n-1}).$$

Since *I* is graded, it follows from (20) that $q_0 \cdot x_n^{d-1} - (yQ_2)_{d-1} \in I$. We thus see that if

$$R = 1 + \sum_{i \ge 0} z^{i} x_{n}^{mi} u_{i}(x_{1}, \dots, x_{n-1}) - y \cdot \sum_{j,k \ge 0} z^{j} y^{k} x_{n}^{d(k+1)+mj} v_{j,k}(x_{1}, \dots, x_{n-1}),$$

then R(0, y, z) = 1 and $R \cdot x_n^{d-1} \in I$.

The second half of the Restriction Theorem for Hodge ideals (see [21, Theorem A (vi)]) says that there is an open subset U of A^2 such that for every $(s, t) \in U$, we have

$$I_p(h_{s,t}^{\gamma}) = I \cdot \mathcal{O}_{\mathbf{A}^{n+2}}|_{y=s, z=t}.$$

Since $R(P, s, t) \neq 0$, we conclude that for all such (s, t), we have $x_n^{d-1} \in I_p(h_{s,t}^{\gamma})$ around P.

Step 3. We now prove that $x_n^{d-1} \in I_p(h_{1,0}^{\gamma})$ around *P*. Note first that the condition that $m \ge d(\theta_P(f) + 1)$ implies that there is an open neighborhood *V* of $(1,0) \in \mathbf{A}^2$ and an integer μ such that for every $(s,t) \in V$, the hypersurface $h_{s,t}$ has an isolated singularity at *P*, with Milnor number μ . Indeed, for every $(s,t) \in \mathbf{A}^2$, with $s \ne 0$, an easy change of

variable implies that $h_{s^d,t}$ has the same Milnor number at P as

$$f(x_1, \dots, x_{n-1}, x_n^d) + \frac{t}{s^m} x_n^m.$$
 (21)

It follows from the proof of [6, Lemma 2.10] that the inequality $m \ge d(\theta_P(f) + 1)$ implies the existence of an open neighborhood V_0 of 0 in \mathbf{A}^1 such that the Milnor number at *P* of the hypersurface in (21) is finite and independent of $t/s^m \in V_0$. If *V* is the image of the open set $\{(s,t) \mid s \ne 0, t/s^m \in V_0\}$ via the map $\mathbf{A}^2 \rightarrow \mathbf{A}^2$ that maps (s,t) to (s^d,t) , then *V* satisfies the required property (note that this map is open, being flat).

It is then easy to see that there is an open neighborhood W of $\{P\} \times V$ in $\mathbf{A}^n \times V$ such that for every $(s, t) \in V$, the only singular point of $h_{s,t}$ in $W \cap (\mathbf{A}^n \times \{(s, t)\})$ is (P, s, t) (see for example [6, Proposition 2.9 (ii)]). Let Z be the closed subscheme of W defined by $I \cdot \mathcal{O}_W$ and $\tau: Z \to V$ the morphism induced by the projection $\mathbf{A}^n \times \mathbf{A}^2 \to \mathbf{A}^2$. It follows from our choice of W that we can apply Theorem 1.4 to the morphism τ and to the function $h|_W$ to conclude that τ is a finite flat morphism and we have

$$I_p(h_{s,t}^{\gamma}) = I \cdot \mathcal{O}_W|_{y=s, z=t} \quad \text{for all } (s,t) \in V.$$
(22)

Since *V* is an irreducible variety and τ is finite and flat, the fact that $x_n^{d-1} \in I_p(h_{s,t}^{\gamma})$ for all $(s,t) \in U \cap V$ gives $x_n^{d-1} \in I \cdot \mathcal{O}_W$ on *W*. Indeed, recall first that by Step 2, for every $(s,t) \in U \cap V$ we have $x_n^{d-1} \in I \cdot \mathcal{O}_W|_{y=s,z=t}$ (a priori, we only know this around *P*, but *P* is the only singular point of $h_{s,t}$ in $W \cap (\mathbf{A}^n \times \{(s,t)\})$). Since τ is finite and flat, this implies³ that $x_n^{d-1}|_{\tau^{-1}(U \cap V)} = 0$. Using the fact that *V* is a reduced scheme and τ is flat, we now deduce that $x_n^{d-1}|_Z = 0$. In particular, the equality in (22) for (s,t) = (1,0) gives $x_n^{d-1} \in I_p(h_{1,0}^{\gamma})$ around *P*.

We can now conclude. Note that $h_{1,0}(x_1, \ldots, x_n) = f(x_1, \ldots, x_{n-1}, x_n^d)$. Consider the finite morphism $\varphi: \mathbf{A}^n \to \mathbf{A}^n$ given by $\varphi(x_1, \ldots, x_n) = (x_1, \ldots, x_{n-1}, x_n^d)$. Since $\varphi^*(f) = h_{1,0}$ and P is the only point in the fiber of φ over P, it follows from Step 3 that we may apply Theorem 1.3 for the restriction of φ over a suitable neighborhood of P to conclude that $1 \in I_p(f^\gamma)$ around P. We thus have $\tilde{\alpha}_P(f) \ge p + \gamma = \lambda$ by the characterization of the minimal exponent in terms of Hodge ideals (see (3)). This completes the proof of the theorem.

Proof of Corollary 1.2. Since H_1, \ldots, H_{n-1} are general smooth hypersurfaces in X containing P, it follows that each $Z_i := H_1 \cap \cdots \cap H_i$ is smooth and $f|_{Z_i}$ has an isolated singularity at P, for $1 \le i \le n-1$. We can thus apply Theorem 1.1 to each of $f, f|_{Z_1}, \ldots, f|_{Z_{n-2}}$ to conclude that

$$\widetilde{\alpha}_{P}(f) \geq \frac{1}{\theta_{P}(f) + 1} + \frac{1}{\theta_{P}(f|_{Z_{1}}) + 1} + \dots + \frac{1}{\theta_{P}(f|_{Z_{n-2}}) + 1} + \widetilde{\alpha}_{P}(f|_{Z_{n-1}}).$$

Note now that Z_{n-1} is a smooth curve. If $m = \text{mult}_P(f|_{Z_{n-1}})$, then $\tilde{\alpha}(f|_{Z_{n-1}}) = 1/m = \frac{1}{\theta_P(f|_{Z_{n-1}})+1}$. We thus obtain the inequality in the statement of the corollary.

³Note that if A is a reduced algebra of finite type over C and B is a finite flat A-algebra, then the injective homomorphism $A \hookrightarrow \prod_{\mathfrak{m} \in Max(A)} A/\mathfrak{m}$ induces an injective homomorphism $B \hookrightarrow \prod_{\mathfrak{m} \in Max(A)} B/\mathfrak{m}B$.

6. A lower bound for the minimal exponent of general hyperplane sections

Our goal in this section is to prove Theorem 1.5. We begin with a few comments regarding what we mean by restriction to a *general hypersurface*. Let *P* be a point on a smooth complex algebraic variety *X*. We consider a system of regular functions y_1, \ldots, y_N defined on an open neighborhood of *P* and whose images in the local ring $\mathcal{O}_{X,P}$ generate the maximal ideal. If *H* is a hypersurface in *X* defined around *P* by a linear combination $\sum_{i=1}^{N} a_i y_i$, with $a_1, \ldots, a_N \in \mathbb{C}$ general, we refer to *H* as a general hypersurface in *X* containing *P*. It is clear that such a hypersurface is smooth at *P*.

If $f \in \mathcal{O}_X(X)$ is such that f(P) = 0, then it follows from the semicontinuity statement for minimal exponents discussed at the beginning of the previous section that for such a general H, the minimal exponent $\tilde{\alpha}_P(f|_H)$ is independent of H. Moreover, this is the largest of all minimal exponents $\tilde{\alpha}_P(f|_{H'})$, where H' is *any* hypersurface in X that is smooth at P and with $f|_{H'} \neq 0$ (this follows by enlarging the given system of generators of the maximal ideal of $\mathcal{O}_{X,P}$ with the germ of an equation defining H' around P).

We can now prove the lower bound for the minimal exponent for the restriction to a general hypersurface containing P.

Proof of Theorem 1.5. Let $d = \operatorname{mult}_P(f)$. The assertion in the theorem is clear if the hypersurface defined by f is smooth at P: indeed, in this case also the hypersurface defined by $f|_H$ in H is smooth, hence $\tilde{\alpha}_P(f|_H) = \infty = \tilde{\alpha}_P(f)$. Therefore from now on we may assume $d \ge 2$.

Suppose first that we know the assertion when f has an isolated singularity at P. In order to deduce the general case, we may and will assume that X is affine and we have an algebraic system of coordinates x_1, \ldots, x_n on X, centered at P. It follows from the discussion at the beginning of this section that it is enough to exhibit one smooth hypersurface H containing P such that $\tilde{\alpha}_P(f|_H) \ge \tilde{\alpha}_P(f) - 1/d$. For every m > d, consider n

$$f_m = f + \sum_{i=1}^n a_{i,m} x_i^m,$$

where $a_{1,m}, \ldots, a_{n,m} \in \mathbb{C}$ are general. Note that by the Kleiman-Bertini theorem the hypersurface defined by f_m has at most one singular point, namely P. Note also that $\operatorname{mult}_P(f_m) = d$. By the isolated singularity case, it follows that if H is general (depending on f_m), then $\widetilde{\alpha}_P(f_m) = \widetilde{\alpha}_P(f_m) = 1/d$

$$\widetilde{\alpha}_P(f_m|_H) \ge \widetilde{\alpha}_P(f_m) - 1/d$$

Since the intersection of a countable family of nonempty Zariski open subsets of an irreducible complex algebraic variety is nonempty, it follows that we can choose such a smooth hypersurface H that satisfies these conditions for all m. Since we have

$$\lim_{m \to \infty} \tilde{\alpha}_P(f_m) = \tilde{\alpha}_P(f) \quad \text{and} \quad \lim_{m \to \infty} \tilde{\alpha}_P(f_m|_H) = \tilde{\alpha}_P(f|_H)$$

by [22, Proposition 6.6(3)], we deduce that

$$\widetilde{\alpha}_P(f|_H) \ge \widetilde{\alpha}_P(f) - 1/d.$$

Therefore f satisfies the assertion in the theorem.

We thus see that it is enough to treat the case when f has an isolated singularity at P. This follows from a sharper inequality proved by Loeser [15]. For the benefit of the reader, we include a slightly modified version of his proof, explaining in detail how various arguments from [32] come in the picture.

Arguing as in Remark 5.2, we see that it is enough to consider the case when $X = \mathbf{A}^n$ and P = 0. After a suitable choice of coordinates x_1, \ldots, x_n , we may assume that H is the hyperplane defined by $x_n = 0$. Consider the following family of polynomials:

$$h_t(x_1, \dots, x_n) = f(x_1, \dots, x_{n-1}, tx_n) + (1-t)x_n^d$$
 for $t \in \mathbf{C}$.

Note that $h_0 = g + x_n^d$, where $g = f(x_1, ..., x_{n-1}, 0)$, hence the Thom–Sebastiani formula for the minimal exponent gives

$$\widetilde{\alpha}_0(h_0) = \widetilde{\alpha}_0(g) + 1/d$$

(see [16, Example (6.8)] or [22, Example 6.7]). On the other hand, we have $h_1 = f$, hence by the semicontinuity of the minimal exponent (see the discussion at the beginning of the previous section) there is a Zariski open neighborhood U of 1 such that $\tilde{\alpha}_0(h_t) \ge \tilde{\alpha}_0(f)$ for all $t \in U$. The key point is to show that there is an open neighborhood V of 0 such that for $t \in V$, h_t has an isolated singularity at 0 and the Milnor number is constant. Indeed, in this case Varchenko's theorem [35] implies that for $t \in V$, we have

$$\widetilde{\alpha}_0(h_t) = \widetilde{\alpha}_0(h_0) = \widetilde{\alpha}_0(g) + 1/d.$$

By taking $t \in U \cap V$, we thus obtain

$$\widetilde{\alpha}_0(f|_H) = \widetilde{\alpha}_0(g) \ge \widetilde{\alpha}_0(f) - 1/d.$$

Note that the Milnor number of h_0 at 0 is $\mu_0(g + x_n^{d-1}) = (d-1) \cdot \mu_0(g)$. Suppose now that $t \neq 0$ and we want to compute the Milnor number μ_t of h_t at 0 for general such *t*. Note that since h_0 has an isolated singularity at 0, the same holds for h_t with *t* general.

An easy change of variable gives

$$\mu_t = \dim_{\mathbf{C}}(\mathcal{O}_{\mathbf{A}^n,\mathbf{0}}/J_t), \quad \text{where } J_t = (\partial f/\partial x_1, \dots, \partial f/\partial x_{n-1}, dsx_n^{d-1} + \partial f/\partial x_n) \cdot \mathcal{O}_{\mathbf{A}^n,\mathbf{0}}$$

with $s = \frac{1-t}{t^d}$. We will make use of various facts about Hilbert–Samuel multiplicities, for which we refer to [19, Chapter 14]) and, more generally, of mixed multiplicities, for which we refer to [31,32]. Note that since h_t has an isolated singularity at 0, the elements $\frac{\partial f}{\partial x_1}, \ldots, \frac{\partial f}{\partial x_{n-1}}, dsx_n^{d-1} + \frac{\partial f}{\partial x_n}$ form a system of parameters, hence a regular sequence, in the Cohen–Macaulay ring $\mathcal{O}_{\mathbf{A}^n,0}$. It follows that if Γ is defined by $\frac{\partial f}{\partial x_1}, \ldots, \frac{\partial f}{\partial x_{n-1}}$, then its local ring $\mathcal{O}_{\Gamma,0}$ is Cohen–Macaulay and

$$\mu_t = e(J_t; \mathcal{O}_{\mathbf{A}^n, \mathbf{0}}) = e((dsx_n^{d-1} + \partial f/\partial x_n) \cdot \mathcal{O}_{\Gamma, \mathbf{0}}; \mathcal{O}_{\Gamma, \mathbf{0}})$$
(23)

(see [19, Theorem 14.11]). On the other hand, since t is general, s is general too, hence

$$e((sdx_n^{d-1} + \partial f/\partial x_n) \cdot \mathcal{O}_{\Gamma,0}; \mathcal{O}_{\Gamma,0}) = e((x_n^{d-1}, \partial f/\partial x_n) \cdot \mathcal{O}_{\Gamma,0}; \mathcal{O}_{\Gamma,0})$$
(24)

(see [19, Theorems 14.13 and 14.14]). Using again the fact that $\frac{\partial f}{\partial x_1}, \ldots, \frac{\partial f}{\partial x_{n-1}}$ form a regular sequence and the definition of the Milnor number, we see that

$$\mu_0(g) = \dim_{\mathbb{C}} \left(\mathcal{O}_{\mathbb{A}^n, 0} / (x_n, \partial f / \partial x_1, \dots, \partial f / \partial x_{n-1}) \right) = e((x_n) \cdot \mathcal{O}_{\Gamma, 0}; \mathcal{O}_{\Gamma, 0}).$$
(25)

The fact that *H* is general is used in two ways. First, the Jacobian $J(f|_H)$ of $f|_H$ and the restriction of J_f to $\mathcal{O}_{H,0}$ have the same integral closure (see [32, Proposition 2.7]). This implies that

$$\mu_0(f|_H) = e(J_f|_H; \mathcal{O}_{H,0}) = e(J_f \cdot \mathcal{O}_{H,0}; \mathcal{O}_{H,0}).$$
(26)

On the other hand, since H is general, a basic property of mixed multiplicities (see [32, Corollary 2.2] or [31, Theorem 2.5]) gives

$$e(J_f \cdot \mathcal{O}_{H,0}; \mathcal{O}_{H,0}) = e(J_f^{[n-1]}, \mathfrak{m}^{[1]}; \mathcal{O}_{\mathbf{A}^n, 0}) = e(\mathfrak{m} \cdot \mathcal{O}_{\Gamma,0}; \mathcal{O}_{\Gamma,0}),$$
(27)

where m is the maximal ideal in $\mathcal{O}_{\mathbf{A}^n,0}$ (for the last equality, note that since *H* is general, $\partial f/\partial x_1, \ldots, \partial f/\partial x_{n-1}$ are general linear combinations of a system of generators of J_f). By combining (26) and (27), we conclude that $\mu_0(g) = e(\mathfrak{m} \cdot \mathcal{O}_{\Gamma,0}; \mathcal{O}_{\Gamma,0})$, and using also (25), it follows that

$$e((x_n) \cdot \mathcal{O}_{\Gamma,0}; \mathcal{O}_{\Gamma,0}) = e(\mathfrak{m} \cdot \mathcal{O}_{\Gamma,0}; \mathcal{O}_{\Gamma,0}).$$

The completion of $\mathcal{O}_{\Gamma,0}$ has all its minimal primes of the same dimension (in fact, it is Cohen–Macaulay), hence we can apply a theorem of Rees [23] to conclude that the ideals $(x_n) \cdot \mathcal{O}_{\Gamma,0} \subseteq \mathfrak{m} \cdot \mathcal{O}_{\Gamma,0}$ have the same integral closure. Since $d = \operatorname{mult}_0(f)$, it follows that $\partial f / \partial x_n \in \mathfrak{m}^{d-1}$ and we see that the ideals $(x_n^{d-1}, \partial f / \partial x_n) \cdot \mathcal{O}_{\Gamma,0}$ and $\mathfrak{m}^{d-1} \cdot \mathcal{O}_{\Gamma,0}$ have the same integral closure. Therefore

$$e((x_n^{d-1}, \partial f/\partial x_n) \cdot \mathcal{O}_{\Gamma,0}; \mathcal{O}_{\Gamma,0}) = e(\mathfrak{m}^{d-1} \cdot \mathcal{O}_{\Gamma,0}; \mathcal{O}_{\Gamma,0}) = (d-1) \cdot e(\mathfrak{m} \cdot \mathcal{O}_{\Gamma,0}; \mathcal{O}_{\Gamma,0}).$$

Using now (23)–(25), we conclude that for t general, we have $\mu_t = (d - 1) \cdot \mu_0(g) = \mu_0(h_0)$. This completes the proof of the theorem.

Corollary 6.1. Let X be a smooth complex algebraic variety of dimension n and P a point in X. Let $f \in \mathcal{O}_X(X)$ be nonzero such that f(P) = 0 and let $d = \operatorname{mult}_P(f)$. If $\tilde{\alpha}_P(f) > 1 + r/d$ for some $r \leq n - 1$ and H_1, \ldots, H_r are general hypersurfaces in X containing P, then the hypersurface of $Y = H_1 \cap \cdots \cap H_r$ defined by $f|_Y$ has rational singularities at P.

Proof. A repeated application of Theorem 1.5 gives $\tilde{\alpha}_P(f|_Y) > 1$. This implies that the hypersurface in *Y* defined by $f|_Y$ has rational singularities at *P* by [27, Theorem 0.4].

Example 6.2. Let $n \ge 2$ and let $f = \det(x_{i,j})_{1\le i,j\le n}$ be the determinant of an $n \times n$ matrix of indeterminates. In this case the reduced Bernstein–Sato polynomial of f (at 0) is given by

$$\widetilde{b}_f(s) = \prod_{i=2}^n (s+i)$$

(see for example [12, Appendix]). We thus have $\tilde{\alpha}_0(f) = 2$. Since $\operatorname{mult}_0(f) = n$, it follows from Corollary 6.1 that if $L \subseteq \mathbf{A}^{n^2}$ is a general linear subspace containing 0, of codimension < n, then the restriction $f|_L$ defines a hypersurface with rational singularities (note that $f|_L$ is a homogeneous polynomial, hence having rational singularities at 0 implies rational singularities everywhere).

Corollary 6.3. Let X be a smooth complex algebraic variety of dimension n and P a point in X. If $f \in O_X(X)$ is nonzero and P is a singular point of the hypersurface Y defined by f, then the following are equivalent:

- (i) $\tilde{\alpha}_P(f) = n/2$.
- (ii) The tangent cone $C_P(f)$ of Y at P is a quadric cone of rank n.
- (iii) There are analytic local coordinates x_1, \ldots, x_n on X centered at P such that $f = \sum_{i=1}^n x_i^2$.

Proof. The equivalence between (ii) and (iii) is well-known: for example, it is a consequence of the Morse lemma. Note also that if $C_P(f)$ is a quadric cone of rank n, then it is well-known that $\tilde{\alpha}_P(f) = n/2$ (see, for example, [22, Theorem E(3)]). Therefore it is enough to prove the converse.

Recall that since *P* is a singular point of *Y*, we always have $\tilde{\alpha}_P(f) \le n/2$ by [28, Theorem (0.4)]. We prove by induction on $n \ge 1$ that if $\tilde{\alpha}_P(f) = n/2$, then *f* satisfies the condition in (ii). The case n = 1 is trivial. For the induction step, suppose that $n \ge 2$ and that we know the assertion for n - 1. If $\tilde{\alpha}_P(f) = n/2$ and $d = \text{mult}_P(f)$, then for a general hypersurface *H* in *X* containing *P*, it follows from Theorem 1.5 that

$$\widetilde{\alpha}_P(f|_H) \ge \widetilde{\alpha}_P(f) - \frac{1}{d} = \frac{n}{2} - \frac{1}{d} \ge \frac{n-1}{2}.$$

Since $\tilde{\alpha}_P(f|_H) \leq \frac{n-1}{2}$, we conclude that d = 2 and $\tilde{\alpha}_P(f|_H) = \frac{n-1}{2}$. By the induction hypothesis, it follows that $C_P(f|_H)$ is a quadric cone of rank n - 1. In particular, we deduce that $C_P(f)$ is a quadric cone of rank $\geq n - 1$. By the Morse lemma, there are local analytic coordinates x_1, \ldots, x_n on X centered at P such that $f = x_1^m + \sum_{i=2}^n x_i^2$ for some $m \geq 2$. In this case the Thom–Sebastiani formula gives $\tilde{\alpha}_P(f) = \frac{1}{m} + \frac{n-1}{2}$. Since $\tilde{\alpha}_P(f) = \frac{n}{2}$, we deduce that m = 2, completing the proof of the induction step.

Acknowledgments. We are grateful to Eva Elduque for many discussions related to this project. We would like to thank Sebastián Olano and Jakub Witaszek for a conversation that led to Corollary 6.3. We are also indebted to the anonymous referees for the careful reading of the paper and for many useful comments.

Funding. The authors were partially supported by NSF grant DMS-1701622.

References

 [1] Arnol'd, V. I., Guseĭn-Zade, S. M., Varchenko, A. N.: Singularities of Differentiable Maps. Vol. II. Monogr. Math. 83, Birkhäuser Boston, Boston, MA (1988) Zbl 0659.58002 MR 966191

- Björk, J.-E.: Analytic D-modules and Applications. Math. Appl. 247, Kluwer, Dordrecht (1993) Zbl 0805.32001 MR 1232191
- Brion, M.: Differential forms on quotients by reductive group actions. Proc. Amer. Math. Soc. 126, 2535–2539 (1998) Zbl 0901.14029 MR 1451789
- Budur, N., Saito, M.: Multiplier ideals, V-filtration, and spectrum. J. Algebraic Geom. 14, 269–282 (2005) Zbl 1086.14013 MR 2123230
- [5] Ein, L., Lazarsfeld, R., Smith, K. E., Varolin, D.: Jumping coefficients of multiplier ideals. Duke Math. J. 123, 469–506 (2004) Zbl 1061.14003 MR 2068967
- [6] Elduque, E., Mustață, M.: On a conjecture of Teissier: the case of log canonical thresholds. Sb. Math. 212, 433–448 (2021) Zbl 1470.14010 MR 4223977
- [7] Greuel, G.-M., Lossen, C., Shustin, E.: Introduction to Singularities and Deformations. Springer Monogr. Math., Springer, Berlin (2007) Zbl 1125.32013 MR 2290112
- [8] Hotta, R., Takeuchi, K., Tanisaki, T.: D-Modules, Perverse Sheaves, and Representation Theory. Progr. Math. 236, Birkhäuser Boston, Boston, MA (2008) Zbl 1136.14009 MR 2357361
- Jung, S.-J., Kim, I.-K., Saito, M., Yoon, Y.: Hodge ideals and spectrum of isolated hypersurface singularities. Ann. Inst. Fourier (Grenoble) 72, 465–510 (2022) Zbl 07554662 MR 4448602
- [10] Källström, R.: *D*-modules and finite maps. arXiv:1811.06796 (2018)
- [11] Kashiwara, M.: B-functions and holonomic systems. Rationality of roots of B-functions. Invent. Math. 38, 33–53 (1976/77) Zbl 0354.35082 MR 430304
- [12] Kimura, T.: Introduction to Prehomogeneous Vector Spaces. Transl. Math. Monogr. 215, Amer. Math. Soc., Providence, RI (2003) Zbl 1035.11060 MR 1944442
- [13] Kollár, J.: Singularities of pairs. In: Algebraic Geometry (Santa Cruz, CA, 1995), Proc. Sympos. Pure Math. 62, Amer. Math. Soc., Providence, RI, 221–287 (1997) Zbl 0905.14002 MR 1492525
- [14] Lazarsfeld, R.: Positivity in Algebraic Geometry. II. Ergeb. Math. Grenzgeb. (3) 49, Springer, Berlin (2004) Zbl 1093.14500 MR 2095472
- [15] Loeser, F.: Exposant d'Arnold et sections planes. C. R. Acad. Sci. Paris Sér. I Math. 298, 485–488 (1984) Zbl 0591.32010 MR 753895
- [16] Malgrange, B.: Intégrales asymptotiques et monodromie. Ann. Sci. École Norm. Sup. (4) 7, 405–430 (1974) Zbl 0305.32008 MR 372243
- [17] Malgrange, B.: Le polynôme de Bernstein d'une singularité isolée. In: Fourier Integral Operators and Partial Differential Equations (Nice, 1974), Lecture Notes in Math. 459, Springer, Berlin, 98–119 (1975) Zbl 0308.32007 MR 0419827
- [18] Malgrange, B.: Polynômes de Bernstein–Sato et cohomologie évanescente. In: Analysis and Topology on Singular Spaces, II, III (Luminy, 1981), Astérisque 101, 243–267 (1983) Zbl 0528.32007 MR 737934
- [19] Matsumura, H.: Commutative Ring Theory. 2nd ed., Cambridge Stud. Adv. Math. 8, Cambridge Univ. Press, Cambridge (1989) Zbl 0666.13002 MR 1011461
- [20] Maxim, L., Saito, M., Schürmann, J.: Thom–Sebastiani theorems for filtered D-modules and for multiplier ideals. Int. Math. Res. Notices 2020, 91–111 Zbl 1476.32003 MR 4050564
- [21] Mustață, M., Popa, M.: Hodge ideals for Q-divisors: birational approach. J. École Polytech. Math. 6, 283–328 (2019) Zbl 1427.14045 MR 3959075
- [22] Mustață, M., Popa, M.: Hodge ideals for \mathbb{Q} -divisors, V-filtration, and minimal exponent. Forum Math. Sigma **8**, art. e19, 41 pp. (2020) MR 4089396
- [23] Rees, D.: α-transforms of local rings and a theorem on multiplicities of ideals. Proc. Cambridge Philos. Soc. 57, 8–17 (1961) MR 118750
- [24] Sabbah, C.: D-modules et cycles évanescents (d'après B. Malgrange et M. Kashiwara). In: Géométrie algébrique et applications, III (La Rábida, 1984), Travaux en Cours 24, Hermann, Paris, 53–98 (1987) Zbl 0623.32013 MR 907935

- [25] Saito, M.: Modules de Hodge polarisables. Publ. RIMS Kyoto Univ. 24, 849–995 (1989) (1988) Zbl 0691.14007 MR 1000123
- [26] Saito, M.: Mixed Hodge modules. Publ. RIMS Kyoto Univ. 26, 221–333 (1990)
 Zbl 0727.14004 MR 1047415
- [27] Saito, M.: On *b*-function, spectrum and rational singularity. Math. Ann. 295, 51–74 (1993)
 Zbl 0788.32025 MR 1198841
- [28] Saito, M.: On microlocal *b*-function. Bull. Soc. Math. France **122**, 163–184 (1994)
 Zbl 0810.32004 MR 1273899
- [29] Saito, M.: Hodge ideals and microlocal V-filtration. arXiv:1612.08667 (2016)
- [30] Steenbrink, J. H. M.: Semicontinuity of the singularity spectrum. Invent. Math. 79, 557–565 (1985) Zbl 0568.14021 MR 782235
- [31] Swanson, I.: Multigraded Hilbert functions and mixed multiplicities. In: Syzygies and Hilbert Functions, Lecture Notes Pure Appl. Math. 254, Chapman & Hall/CRC, Boca Raton, FL, 267–280 (2007) Zbl 1145.13012 MR 2309934
- [32] Teissier, B.: Cycles évanescents, sections planes et conditions de Whitney. In: Singularités à Cargèse (Cargèse, 1972), Astérisque 7-8, 285–362 (1973) Zbl 0295.14003 MR 0374482
- [33] Teissier, B.: Variétés polaires. I. Invariants polaires des singularités d'hypersurfaces. Invent. Math. 40, 267–292 (1977) Zbl 0446.32002 MR 470246
- [34] Teissier, B.: Polyèdre de Newton jacobien et équisingularité. In: Seminar on Singularities (Paris, 1976/1977), Publ. Math. Univ. Paris VII 7, Univ. Paris VII, Paris, 193–221 (1980) MR 683624
- [35] Varchenko, A. N.: The complex singularity index does not change along the stratum $\mu = \text{const.}$ Funktsional. Anal. i Prilozhen. **16**, no. 1, 1–12, 96 (1982) (in Russian) Zbl 0498.32010 MR 648803