



Ioan Bejenaru · Sebastian Herr

## On global well-posedness and scattering for the massive Dirac–Klein–Gordon system

Received September 25, 2014

**Abstract.** We prove global well-posedness and scattering for the massive Dirac–Klein–Gordon system with small initial data of subcritical regularity in dimension three. To achieve this, we impose a nonresonance condition on the masses.

**Keywords.** Global well-posedness, scattering, Dirac–Klein–Gordon system, nonresonance condition, null structure

### 1. Introduction

The Dirac–Klein–Gordon system is a basic model of proton–proton interactions (one proton is scattered in a meson field produced by a second proton) or neutron–neutron interaction (see Bjorken and Drell [4]). In physics these are known as the strong interactions which are responsible for the forces which bind nuclei.

The mathematical formulation of the Dirac–Klein–Gordon system is as follows (see e.g. [8]):

$$\begin{cases} (-i\gamma^\mu \partial_\mu + M)\psi = \phi\psi, \\ (\square + m^2)\phi = \psi^\dagger \gamma^0 \psi. \end{cases} \quad (1.1)$$

Here,  $\square$  denotes the d'Alembertian  $\square = \partial_t^2 - \Delta_x$ ,  $\psi : \mathbb{R}^{1+3} \rightarrow \mathbb{C}^4$  is the spinor field (column vector), and  $\phi : \mathbb{R}^{1+3} \rightarrow \mathbb{R}$  is a scalar field. For  $\mu = 0, \dots, 3$ ,  $\gamma^\mu$  are the  $4 \times 4$  Dirac matrices given by

$$\gamma^0 = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix}, \quad \gamma^j = \begin{pmatrix} 0 & \sigma^j \\ -\sigma^j & 0 \end{pmatrix},$$

where for  $j = 1, 2, 3$  the Pauli matrices  $\sigma^j$  are

$$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

I. Bejenaru: Department of Mathematics, University of California, San Diego, La Jolla, CA 92093-0112, USA; e-mail: [ibejenaru@math.ucsd.edu](mailto:ibejenaru@math.ucsd.edu)

S. Herr: Fakultät für Mathematik, Universität Bielefeld, Postfach 10 01 31, 33501 Bielefeld, Germany; e-mail: [herr@math.uni-bielefeld.de](mailto:herr@math.uni-bielefeld.de)

*Mathematics Subject Classification (2010):* Primary 35Q40; Secondary 35Q41, 35L70

$\psi^\dagger$  denotes the conjugate transpose of  $\psi$ , i.e.  $\psi^\dagger = \overline{\psi}^t$ . The matrices  $\gamma^\mu$  satisfy

$$\gamma^\alpha \gamma^\beta + \gamma^\beta \gamma^\alpha = 2g^{\alpha\beta} I_4, \quad g^{\alpha\beta} = \text{diag}(1, -1, -1, -1).$$

We will study the Cauchy problem with initial condition

$$(\psi, \phi, \partial_t \phi)|_{t=0} = (\psi_0, \phi_0, \phi_1). \quad (1.2)$$

Before turning to the mathematical analysis of the Dirac–Klein–Gordon equations we highlight a key property of the physical model presented in Bjorken and Drell [4, Chapter 10.2]. The mass  $M$  is effectively  $938 \frac{\text{MeV}}{c^2}$  (proton) or  $939 \frac{\text{MeV}}{c^2}$  (neutron). There are many types of meson fields, but those believed to be major contributors to the nuclear force at large distances are the  $\pi$ -mesons (pions), and their masses are  $m = 140 \frac{\text{MeV}}{c^2}$  for  $\pi^\pm$ ,  $m = 135 \frac{\text{MeV}}{c^2}$  for  $\pi^0$ . Heavier mesons such as K mesons (kaons) may also play a role for small impact parameter collisions; the masses of kaons are  $m = 494 \frac{\text{MeV}}{c^2}$  for  $K^\pm$  and  $m = 498 \frac{\text{MeV}}{c^2}$  for  $K^0$ . It is then reasonable to assume that in the Dirac–Klein–Gordon equations

$$2M > m > 0.$$

We are not implying that all mesons are lighter than baryons (protons or neutrons in our context), but that this is a reasonable assumption in the context of our model. Higher energy (more massive) mesons were created momentarily in the Big Bang but are not thought to play a role in nature today. Such particles are also regularly created in experiments; for instance the heaviest meson created is the upsilon meson with mass  $9.46 \frac{\text{GeV}}{c^2}$  (roughly 10 times the mass of the proton/neutron). However, these heavy mesons do not play a role in the model described by Dirac–Klein–Gordon equations.

We now turn our attention to the mathematical aspects of (1.1). The fundamental question is that of global regularity of solutions. For smooth and small initial data endowed with additional algebraic structure, Chadam and Glassey [6] established global regularity for solutions of (1.1). The work of Klainerman [13] on nonlinear Klein–Gordon equations paved the way of establishing a more general result. Following those ideas and taking advantage of the null structure present in the system, Bachelot [1] established global regularity for (very) smooth and small initial data. The next direction of research was to obtain a local in time result for rough data as close as possible to the critical space which is

$$\psi_0 \in L^2, \quad (\phi_0, \phi_1) \in H^{1/2} \times H^{-1/2}.$$

Beals and Bézard [2] proved that for small initial data  $(\phi_0, \phi_1) \in H^2 \times H^1$  and  $\psi_0 \in H^1$  one has a local well-posedness theory for (1.1). Bournaveas [5] improved this local in time result to  $(\phi_0, \phi_1) \in H^{1+\epsilon} \times H^\epsilon$  and  $\psi_0 \in H^{1/2+\epsilon}$ , for any  $\epsilon > 0$ . D’Ancona, Foschi and Selberg [8] established local well-posedness of (1.1) for data  $(\phi_0, \phi_1) \in H^{1/2+\epsilon} \times H^{-1/2+\epsilon}$  and  $\psi_0 \in H^\epsilon$ , for any  $\epsilon > 0$ ; hence the last result covers the full subcritical regime.

Recently, Wang [20] proved a global in time result for small initial data in the critical Besov space  $(\phi_0, \phi_1) \in \dot{B}_{2,1}^{1/2} \times \dot{B}_{2,1}^{-1/2}$  and  $\psi_0 \in \dot{B}_{2,1}^0$  (for  $M = m = 0$ ), additionally

assuming that an angular derivative is bounded in the same space; the proof exploits the observation of Sterbenz [19] that angular regularity acts as a null structure. The result is then extended to nonzero masses under the condition  $2M > m > 0$ .

It is worth mentioning that in all of the above results the masses  $M, m$  are arbitrary; the result in [20] is an exception. In the context of a local in time result, the terms  $M\psi$ ,  $m^2\phi$  can be treated as perturbations, thus allowing an analysis of (1.1) as a system of wave equations. Obviously, this cannot be the case for a global in time theory which includes scattering.

In the context of the cubic Dirac system [3] we proposed a different approach that incorporates the terms  $M\psi$  and  $m^2\phi$  into the linear part of the operator, as they naturally appear. This will help us treat (1.1) as a system of (half) Klein–Gordon equations after using projectors which are adapted to our context from the work of D’Ancona, Foschi and Selberg [8]. Then we restrict our attention to the physically relevant case  $2M > m > 0$  and obtain a global (in time) result and scattering for small initial data in the subcritical regime. The resolution spaces used here have a simpler structure compared to [3]. Our main result is the following

**Theorem 1.1.** *Assume that  $\epsilon > 0$  and  $2M > m > 0$ . Then the Cauchy problem (1.1)–(1.2) is globally well-posed for small initial data*

$$\psi_0 \in H^\epsilon(\mathbb{R}^3; \mathbb{C}^4), \quad (\phi_0, \phi_1) \in H^{1/2+\epsilon}(\mathbb{R}^3; \mathbb{R}) \times H^{-1/2+\epsilon}(\mathbb{R}^3; \mathbb{R}),$$

and these solutions scatter to free solutions for  $t \rightarrow \pm\infty$ .

We refer to Subsection 4.2 for more details. Our result is at the same level of regularity as the one proved by D’Ancona, Foschi and Selberg [8]. Its strength lies in the global in time and scattering parts. In terms of Sobolev regularity it is slightly more restrictive than Wang’s result [20]. However, we do not assume additional angular regularity on the initial data (cf. also Remark 4.2).

A key observation is that under the assumption  $2M > m > 0$  the system (1.1) has no resonances. It was known from prior works on Klein–Gordon type systems with multiple speeds that, under certain conditions between the masses, resonant interactions do not occur and the well-posedness theory improves. We refer the reader to the works of Delort and Fang [9], Schottdorf [17] and Germain [10] and to the references therein. We will use this, together with some localized Strichartz estimates, to prove the key nonlinear estimates.

Note that unlike many of the previous works which dealt with power type nonlinearities for the Klein–Gordon equation, the Dirac–Klein–Gordon system contains derivatives. This is not apparent from our formulation of (1.1); however, if one wants to write (1.1) as a system of Klein–Gordon equations, one should apply  $-i\gamma^\mu\partial_\mu - M$  to the first equation, and then it is obvious that the right hand side contains derivatives.

We conclude this section with an overview of the paper. In Section 2 we introduce some basic notation and rewrite the original system (1.1) in the equivalent form (2.2) which has two advantages: it is first order in time and it unveils the null structure. The gains from the null structure are quantified in Subsection 2.3 in a manner that fits our analysis. In Section 3 we define the resolution space in which we iterate our system. Without

getting into technical details at this point, there is one particular aspect of this section that deserves to be highlighted. Proving Strichartz estimates has become a standard type argument due to the Christ–Kiselev Lemma [7]. However, proving localized versions of the Strichartz estimates using Christ–Kiselev type arguments is not straightforward. In Section 3 we provide an alternative argument for establishing (localized) Strichartz estimates using  $U^p, V^p$  spaces and we think that this part of the paper may be of independent interest. In Section 4 we prove the trilinear estimates based on which we prove our main result in Theorem 1.1.

## 2. Reductions

### 2.1. Notation

We write  $A \lesssim B$  if there is a harmless constant  $c > 0$  such that  $A \leq cB$ , and  $A \gtrsim B$  iff  $B \lesssim A$ . Further, we write  $A \approx B$  iff both  $A \lesssim B$  and  $B \lesssim A$ . Also, we write  $A \ll B$  if the constant  $c$  can be chosen such that  $c < 2^{-10}$ . Also,  $A \gg B$  iff  $B \ll A$ .

Similarly, we write  $A \leq B$  iff  $2^A \lesssim 2^B$ ,  $A \geq B$  iff  $2^A \gtrsim 2^B$ ,  $A \sim B$  iff  $2^A \approx 2^B$ ,  $A < B$  iff  $2^A \ll 2^B$ ,  $A > B$  iff  $2^A \gg 2^B$ .

Let  $\rho^0 \in C_c^\infty(-2, 2)$  be a fixed smooth, even, cutoff satisfying  $\rho^0(s) = 1$  for  $|s| \leq 1$  and  $0 \leq \rho \leq 1$ . For  $k \in \mathbb{Z}$  we define  $\rho_k : \mathbb{R}^3 \rightarrow \mathbb{R}$ ,  $\rho_k(y) := \rho^0(2^{-k}|y|) - \rho^0(2^{-k+1}|y|)$ , such that  $A_k := \text{supp}(\rho_k) \subset \{y \in \mathbb{R}^3 : 2^{k-1} \leq |y| \leq 2^{k+1}\}$ . Let  $\tilde{\rho}_k = \rho_{k-1} + \rho_k + \rho_{k+1}$  and  $\tilde{A}_k := \text{supp}(\tilde{\rho}_k)$ . For  $k \geq 1$ , let  $P_k$  be the Fourier multiplication operators with respect to  $\rho_k$ , and  $P_0 = I - \sum_{k \geq 1} P_k$ . For  $j \in \mathbb{Z}$  we define

$$\mathcal{F}[Q_j^{\pm, m} f](\tau, \xi) = \rho_j(\tau \pm \langle \xi \rangle_m) \mathcal{F}f(\tau, \xi).$$

Similarly, we define  $\tilde{P}_k$  and  $\tilde{Q}_j^{\pm, m}$ .

We also define  $P_{\leq k} = \sum_{0 \leq k' \leq k} P_{k'}$ ,  $P_{< k} = \sum_{0 \leq k' < k} P_{k'}$ ,  $P_{> k} = I - P_{\leq k}$ ,  $P_{\geq k} = I - P_{< k}$ , and similarly  $Q_{\leq j}^{\pm, m}$ ,  $Q_{< j}^{\pm, m}$ ,  $Q_{\geq j}^{\pm, m}$ , and  $Q_{j \in J}^{\pm, m}$  for an interval  $J$ . In the obvious way we also define the analogous operators based on  $\tilde{P}_k$  and  $\tilde{Q}_j^{\pm, m}$ .

In the case  $m = 1$  we suppress the superscripts, e.g.  $Q_j^{\pm, 1} = Q_j^\pm$ .

Further, for  $l \in \mathbb{N}$  let  $\mathcal{K}_l$  denote a set of spherical caps of radius  $2^{-l}$  which is a covering of  $\mathbb{S}^2$  with finite overlap. For a cap  $\kappa \in \mathcal{K}_l$  we denote its center in  $\mathbb{S}^2$  by  $\omega(\kappa)$ . Let  $\Gamma_\kappa$  be the cone generated by  $\kappa \in \mathcal{K}_l$  and  $(\eta_\kappa)_{\kappa \in \mathcal{K}_l}$  be a smooth partition of unity subordinate to  $(\Gamma_\kappa)_{\kappa \in \mathcal{K}_l}$ . Let  $P_\kappa$  denote the Fourier multiplication operator with symbol  $\eta_\kappa$  such that  $I = \sum_{\kappa \in \mathcal{K}_l} P_\kappa$ . Further, let  $\tilde{P}_\kappa$  be with doubled support such that  $P_\kappa = \tilde{P}_\kappa P_\kappa = P_\kappa \tilde{P}_\kappa$ . For notational convenience, we also define  $\mathcal{K}_0 = \{\mathbb{S}^2\}$  and  $P_\kappa = I$  if  $\kappa \in \mathcal{K}_0$ .

### 2.2. Setup of the system and null structure

As written in (1.1), the cubic Dirac–Klein–Gordon system has a linear part whose coefficients are matrices, and it is technically easier to work with scalar equations. To do so, we adapt the setup introduced in [8, Section 2 and 3] to take into account the mass terms,

similarly to our prior work on the cubic Dirac equation [3] (however, the sign convention is in accordance with [8]). We repeat here the essential steps for the convenience of the reader. As highlighted in [8], the new setup is able to identify a null structure in the non-linearity, although the presence of mass terms alters the effectiveness of this structure at very small scales.

For  $j = 1, 2, 3$  the matrices  $\alpha^j := \gamma^0 \gamma^j$ ,  $\beta := \gamma^0$  have the properties

$$\alpha^j \beta + \beta \alpha^j = 0, \quad \alpha^j \alpha^k + \alpha^k \alpha^j = 2\delta^{jk} I_4;$$

see [8, p. 878] for more details.

We introduce the Fourier multiplication operators  $\Pi_{\pm}^M(D)$  with symbol

$$\Pi_{\pm}^M(\xi) = \frac{1}{2} \left[ I \pm \frac{1}{\langle \xi \rangle_M} (\xi \cdot \alpha + M\beta) \right].$$

In the case  $M = 1$  we suppress the superscript, i.e.  $\Pi_{\pm}(D) = \Pi_{\pm}^1(D)$ .

We then define  $\psi_{\pm} = \Pi_{\pm}^M(D)\psi$  and split  $\psi = \psi_+ + \psi_-$ . Also, define  $\langle D \rangle = \sqrt{1 - \Delta}$ . By applying the operators  $\Pi_{\pm}^M(D)$  to the system (1.1) we obtain the following system of equations:

$$\begin{cases} (-i\partial_t + \langle D \rangle_M)\psi_+ = \Pi_+^M(D)(\phi\beta\psi), \\ (-i\partial_t - \langle D \rangle_M)\psi_- = \Pi_-^M(D)(\phi\beta\psi), \\ (\square + m^2)\phi = \langle \psi, \beta\psi \rangle. \end{cases} \tag{2.1}$$

In order to have a fully first order system, we define  $\phi_{\pm} = \phi \pm i\langle D \rangle_m^{-1}\partial_t\phi$ , and thus

$$(-i\partial_t + \langle D \rangle_m)\phi_+ = \langle D \rangle_m^{-1}\langle \psi, \beta\psi \rangle.$$

Note that  $\phi = \Re\phi_{\pm}$  and  $\phi_- = \overline{\phi_+}$  since  $\phi$  is real-valued. The system which we will study is

$$\begin{cases} (-i\partial_t + \langle D \rangle_M)\psi_+ = \Pi_+^M(D)(\Re\phi_+\beta\psi), \\ (-i\partial_t - \langle D \rangle_M)\psi_- = \Pi_-^M(D)(\Re\phi_+\beta\psi), \\ (-i\partial_t + \langle D \rangle_m)\phi_+ = \langle D \rangle_m^{-1}\langle \psi, \beta\psi \rangle. \end{cases} \tag{2.2}$$

We aim to provide a global theory for this system for initial data  $(\psi_{\pm,0}, \phi_{+,0}) \in H^\epsilon \times H^{1/2+\epsilon}$ . It is an easy exercise that this translates back into a global theory for the original system with  $(\psi_0, \phi_0, \phi_1) \in H^\epsilon \times H^{1/2+\epsilon} \times H^{-1/2+\epsilon}$ .

There is a null structure in the system (2.2), which we describe next. This is again inspired by the work in [8] and was adapted to the current setup in [3]. For more details, we refer the reader to [8, 3].

We decompose  $\langle \psi, \beta\psi \rangle$  as

$$\begin{aligned} \langle \psi, \beta\psi \rangle &= \langle \Pi_+^M(D)\psi_+, \beta\Pi_+^M(D)\psi_+ \rangle + \langle \Pi_-^M(D)\psi_-, \beta\Pi_-^M(D)\psi_- \rangle \\ &\quad + \langle \Pi_+^M(D)\psi_+, \beta\Pi_-^M(D)\psi_- \rangle + \langle \Pi_-^M(D)\psi_-, \beta\Pi_+^M(D)\psi_+ \rangle. \end{aligned}$$

We have

$$\Pi_{\pm}^M(D)\beta = \beta\Pi_{\mp}^M(D) \pm M\langle D \rangle_M^{-1}\beta. \tag{2.3}$$

The following lemma, which corresponds to [3, Lemma 3.1] and [8, Lemma 2], analyzes the symbols of the bilinear operators above.

**Lemma 2.1.** *For fixed  $M \geq 0$ ,*

$$\begin{aligned} \Pi_{\pm}^M(\xi)\Pi_{\mp}^M(\eta) &= \mathcal{O}(\angle(\xi, \eta)) + \mathcal{O}(\langle \xi \rangle^{-1} + \langle \eta \rangle^{-1}), \\ \Pi_{\pm}^M(\xi)\Pi_{\pm}^M(\eta) &= \mathcal{O}(\angle(-\xi, \eta)) + \mathcal{O}(\langle \xi \rangle^{-1} + \langle \eta \rangle^{-1}). \end{aligned} \tag{2.4}$$

We now explain heuristically why this is useful here; see Lemma 3.3 for the technical result which will be used in the nonlinear analysis. By (2.3), for  $s_1, s_2 \in \{+, -\}$ ,

$$\begin{aligned} \mathcal{F}_x \langle \Pi_{s_1} \psi_1, \beta \Pi_{s_2} \psi_2 \rangle(\xi) &= \int_{\xi=\xi_1-\xi_2} \langle \Pi_{s_1}(\xi_1) \widehat{\psi}_1(\xi_1), \beta \Pi_{s_2}(\xi_2) \widehat{\psi}_2(\xi_2) \rangle d\xi_1 d\xi_2 \\ &= \int_{\xi=\xi_1-\xi_2} \langle \beta \Pi_{-s_2}(\xi_2) \Pi_{s_1}(\xi_1) \widehat{\psi}_1(\xi_1), \widehat{\psi}_2(\xi_2) \rangle d\xi_1 d\xi_2 \\ &\quad + s_2 M \int_{\xi=\xi_1-\xi_2} \langle \xi_1 \rangle_M^{-1} \langle \beta \Pi_{s_1}(\xi_1) \widehat{\psi}_1(\xi_1), \widehat{\psi}_2(\xi_2) \rangle d\xi_1 d\xi_2. \end{aligned}$$

Hence, smallness of the angle  $\angle(s_1\xi_1, s_2\xi_2)$  can be exploited as long as it exceeds  $\max(\langle \xi_1 \rangle_M^{-1}, \langle \xi_2 \rangle_M^{-1})$ . See [8, p. 885] for the analogue of this in the massless case, where we have  $\Pi_{-}^0(\xi_1)\Pi_{+}^0(\xi_2) = 0$  if  $\angle(\xi_1, \xi_2) = 0$ , which makes the null structure effective at all angular scales. In the massive case  $M > 0$  the null structure does not bring gains beyond  $\max(\langle \xi_1 \rangle_M^{-1}, \langle \xi_2 \rangle_M^{-1})$ . To compensate for this we need to use the fact that there are no resonances present in (2.2).

In fact, as observed in [8], there is a second and similar null structure in the nonlinearities present in the equations for  $\psi_{\pm}$ , which will be exploited by duality in Section 4.

### 2.3. Modulation analysis

A key aspect in the nonlinear analysis is the lack of resonant terms. Arguments of similar nature are contained in [17, Lemma 2] (see also [9, 10]). Additionally, we will prove that smallness of the maximal modulation induces angular constraints. In the context of the cubic Dirac equation a similar result is contained in [3, Lemma 6.5]. We first provide lower bounds for the resonance function.

**Lemma 2.2.** *Fix  $0 < m < 2M$ . For  $s_1, s_2 \in \{+, -\}$  define the resonance function*

$$\mu^{s_1, s_2}(\xi_1, \xi_2) := \langle \xi_1 - \xi_2 \rangle_m + s_1 \langle \xi_1 \rangle_M - s_2 \langle \xi_2 \rangle_M. \tag{2.5}$$

*Then we have the following bounds:*

- *Case 1: If*
  - (a)  $s_1 = +, s_2 = -$  or

(b)  $s_1 = -, s_2 = +$  and  $\langle \xi_1 - \xi_2 \rangle_m \ll \min(\langle \xi_1 \rangle_M, \langle \xi_2 \rangle_M)$ ,

then

$$|\mu^{s_1, s_2}(\xi_1, \xi_2)| \gtrsim \max(\langle \xi_1 - \xi_2 \rangle, \langle \xi_1 \rangle, \langle \xi_2 \rangle). \quad (2.6)$$

• *Case 2: If*

(a)  $s_1 = s_2$  or

(b)  $s_1 = -, s_2 = +$  and  $\langle \xi_1 - \xi_2 \rangle_m \gtrsim \min(\langle \xi_1 \rangle_M, \langle \xi_2 \rangle_M)$ ,

then

$$|\mu^{s_1, s_2}(\xi_1, \xi_2)| \gtrsim_{m, M} \frac{\langle \xi_1 \rangle \cdot \langle \xi_2 \rangle}{\langle \xi_1 - \xi_2 \rangle} \angle (s_1 \xi_1, s_2 \xi_2)^2. \quad (2.7)$$

With any choice of signs, we have both

$$|\mu^{s_1, s_2}(\xi_1, \xi_2)| \gtrsim_{m, M} \min(\langle \xi_1 \rangle, \langle \xi_2 \rangle) \angle (s_1 \xi_1, s_2 \xi_2)^2, \quad (2.8)$$

and the non-resonance bound

$$|\mu^{s_1, s_2}(\xi_1, \xi_2)| \gtrsim_{m, M} \max(\langle \xi_1 - \xi_2 \rangle^{-1}, \langle \xi_1 \rangle^{-1}, \langle \xi_2 \rangle^{-1}). \quad (2.9)$$

*Proof.* In Case 1 the lower bound (2.6) is obvious, which implies all other claims.

Suppose now that we are in Case 2(a). Then

$$\begin{aligned} & (\langle \xi_1 - \xi_2 \rangle_m - |\langle \xi_1 \rangle_M - \langle \xi_2 \rangle_M|)(\langle \xi_1 - \xi_2 \rangle_m + |\langle \xi_1 \rangle_M - \langle \xi_2 \rangle_M|) \\ &= 2(|\xi_1| |\xi_2| - \xi_1 \cdot \xi_2) + m^2 + 2(\langle \xi_1 \rangle_M \langle \xi_2 \rangle_M - |\xi_1| |\xi_2| - M^2). \end{aligned}$$

Now, we compute

$$\langle \xi_1 \rangle_M \langle \xi_2 \rangle_M - (|\xi_1| |\xi_2| + M^2) = M^2 \frac{(|\xi_1| - |\xi_2|)^2}{\langle \xi_1 \rangle_M \langle \xi_2 \rangle_M + |\xi_1| |\xi_2| + M^2}. \quad (2.10)$$

Since this is nonnegative, we conclude that

$$\begin{aligned} & (\langle \xi_1 - \xi_2 \rangle_m - |\langle \xi_1 \rangle_M - \langle \xi_2 \rangle_M|)(\langle \xi_1 - \xi_2 \rangle_m + |\langle \xi_1 \rangle_M - \langle \xi_2 \rangle_M|) \\ & \geq 2|\xi_1| |\xi_2| (1 - \cos \angle(\xi_1, \xi_2)) + m^2 \gtrsim |\xi_1| |\xi_2| \angle(\xi_1, \xi_2)^2 + m^2. \end{aligned}$$

Now, because of  $m > 0$  and  $\langle \xi_1 - \xi_2 \rangle_m + |\langle \xi_1 \rangle_M - \langle \xi_2 \rangle_M| \lesssim \langle \xi_1 - \xi_2 \rangle_m$  the estimates (2.8) and (2.7) follow. Also, (2.9) follows if  $\langle \xi_1 - \xi_2 \rangle \lesssim \min(\langle \xi_1 \rangle, \langle \xi_2 \rangle)$ . Otherwise, we have  $\max(\langle \xi_1 \rangle, \langle \xi_2 \rangle) \gg \min(\langle \xi_1 \rangle, \langle \xi_2 \rangle)$ , and the estimate (2.9) follows from

$$\begin{aligned} & (\langle \xi_1 - \xi_2 \rangle_m - |\langle \xi_1 \rangle_M - \langle \xi_2 \rangle_M|)(\langle \xi_1 - \xi_2 \rangle_m + |\langle \xi_1 \rangle_M - \langle \xi_2 \rangle_M|) \\ & \geq M^2 \frac{(|\xi_1| - |\xi_2|)^2}{\langle \xi_1 \rangle_M \langle \xi_2 \rangle_M + |\xi_1| |\xi_2| + M^2}, \end{aligned}$$

where we have used (2.10) again.

Suppose now that we are in Case 2(b). A computation similar to the above yields

$$\begin{aligned} & (\langle \xi_1 \rangle_M + \langle \xi_2 \rangle_M - \langle \xi_1 - \xi_2 \rangle_m)(\langle \xi_1 \rangle_M + \langle \xi_2 \rangle_M + \langle \xi_1 - \xi_2 \rangle_m) \\ &= 2(|\xi_1| |\xi_2| + \xi_1 \cdot \xi_2) + 2M^2 - m^2 + 2(\langle \xi_1 \rangle_M \langle \xi_2 \rangle_M - |\xi_1| |\xi_2|) \\ &\geq |\xi_1| |\xi_2| \angle(-\xi_1, \xi_2)^2 + 4M^2 - m^2. \end{aligned}$$

By assumption  $4M^2 - m^2 > 0$ , so the estimate (2.8) is proved, and due to  $\langle \xi_1 - \xi_2 \rangle \approx \max(\langle \xi_1 \rangle, \langle \xi_2 \rangle)$  the claim (2.7) follows, too. Also, if  $|\xi_1| \approx |\xi_2|$ , then (2.9) follows. Otherwise, we use the lower bound provided by (2.10) to obtain (2.9).  $\square$

**Remark 2.3.** From now on we fix  $M = m = 1$  in order to simplify the exposition. In view of Lemma 2.2 it will be obvious that all arguments carry over to the case  $2M > m > 0$  with modified (implicit) constants depending on  $m, M$ .

**Lemma 2.4.** Let  $s_1, s_2 \in \{+, -\}$ . Consider  $k, k_1, k_2 \in \mathbb{N}_0, j, j_1, j_2 \in \mathbb{Z}$ , and  $\phi = \tilde{P}_k \tilde{Q}_j^+ \phi, u_i = \tilde{P}_{k_i} \tilde{Q}_{j_i}^{s_i} u_i$ .

(i) If  $\max(j, j_1, j_2) < -\min(k, k_1, k_2)$ , we have

$$\int_{\mathbb{R}^{1+3}} \phi \cdot u_1 \overline{u_2} \, dt \, dx = 0. \tag{2.11}$$

(ii) Case 1: Suppose that either

- $s_1 = +, s_2 = -, \text{ or}$
- $s_1 = -, s_2 = + \text{ and } k < \min(k_1, k_2)$ .

If  $\max(j, j_1, j_2) < \max(k, k_1, k_2)$ , then (2.11) holds true.

Case 2: Suppose that either

- $s_1 = s_2, \text{ or}$
- $s_1 = -, s_2 = + \text{ and } k \geq \min(k_1, k_2)$ .

If  $l \geq 1, \kappa_1, \kappa_2 \in \mathcal{K}_l$  with  $d(s_1 \kappa_1, s_2 \kappa_2) \geq 2^{-l}$  and  $\max(j, j_1, j_2) < k_1 + k_2 - k - 2l$ , then

$$\int_{\mathbb{R}^{1+3}} \phi \cdot \tilde{P}_{\kappa_1} u_1 \overline{\tilde{P}_{\kappa_2} u_2} \, dt \, dx = 0. \tag{2.12}$$

*Proof.* We have

$$\int_{\mathbb{R}^{1+3}} \phi \cdot u_1 \overline{u_2} \, dt \, dx = \int_{\mathbb{R}^{1+3}} \widehat{\phi \overline{u_1} u_2} \, d\tau \, d\xi$$

and, with  $\zeta = (\tau, \xi)$ ,

$$\widehat{\overline{u_1} u_2}(\zeta) = \int \widehat{u_1}(\zeta') \widehat{u_2}(\zeta - \zeta') \, d\zeta' = \int \widehat{u_1}(-\zeta') \widehat{u_2}(\zeta - \zeta') \, d\zeta';$$

hence, with  $\zeta_j = (\tau_j, \xi_j)$ ,

$$\int_{\mathbb{R}^{1+3}} \phi \cdot u_1 \overline{u_2} \, dt \, dx = \int \int \widehat{\phi}(\zeta_2 - \zeta_1) \widehat{u_1}(\zeta_1) \widehat{\overline{u_2}}(\zeta_2) \, d\zeta_1 \, d\zeta_2. \tag{2.13}$$



The assumptions imply that we must have

$$|\tau_2 - \tau_1 + \langle \xi_2 - \xi_1 \rangle| \approx 2^j, \quad |\tau_1 + s_1 \langle \xi_1 \rangle| \approx 2^{j_1}, \quad |\tau_2 + s_2 \langle \xi_2 \rangle| \approx 2^{j_2}$$

in order to obtain a nontrivial contribution. This implies

$$|\langle \xi_2 - \xi_1 \rangle + s_1 \langle \xi_1 \rangle - s_2 \langle \xi_2 \rangle| \lesssim 2^{\max(j, j_1, j_2)}. \tag{2.14}$$

(i) By assumption we have  $2^{\max(j, j_1, j_2)} \ll 2^{-\min(k, k_1, k_2)}$ , so that (2.14) contradicts (2.9).

(ii) By assumption we have  $2^{\max(j, j_1, j_2)} \ll 2^{\max(k, k_1, k_2)}$  in Case 1, hence (2.14) contradicts (2.6). Similarly, in Case 2 the estimate (2.14) contradicts (2.7).  $\square$

### 3. Function spaces and linear estimates

For  $1 \leq p \leq \infty$  and  $b \in \mathbb{R}$ , we define

$$\|f\|_{\dot{X}^{\pm, b, p}} = \|(2^{bj} \|Q_j^\pm f\|_{L^2})_{j \in \mathbb{Z}}\|_{\ell^p}.$$

The low frequency part will be treated as a whole, that is, we define

$$\|f\|_{S_{\leq 0}^\pm} = \|f\|_{L_t^\infty L_x^2} + \|f\|_{L_t^2 L_x^6} + \|f\|_{\dot{X}^{\pm, 1/2, \infty}}.$$

By interpolation, the space above provides all the Strichartz estimates for the Schrödinger equation on  $\mathbb{R}^3$ . This is natural since the Klein–Gordon equation in low frequency behaves like the Schrödinger equation.

In high frequency, the Klein–Gordon equation is of wave type and the Strichartz estimates should reflect that. Moreover we need some refinement of the standard Strichartz estimates.

For  $d = 3$  and  $k \in \mathbb{Z}_+$  let  $\Xi_k = 2^k \cdot \mathbb{Z}^d$ . Let  $\gamma^{(1)} : \mathbb{R} \rightarrow [0, 1]$  denote an even smooth function supported in the interval  $[-2/3, 2/3]$  with the property that

$$\sum_{n \in \mathbb{Z}} \gamma^{(1)}(\xi - n) = 1 \quad \text{for } \xi \in \mathbb{R}.$$

Let  $\gamma : \mathbb{R}^d \rightarrow [0, 1]$ ,  $\gamma(\xi) = \gamma^{(1)}(\xi_1) \cdot \dots \cdot \gamma^{(1)}(\xi_d)$ . For  $k \in \mathbb{Z}_+$  and  $n \in \Xi_k$  let

$$\gamma_{k, n}(\xi) = \gamma((\xi - n)/2^k).$$

Clearly,  $\sum_{n \in \Xi_k} \gamma_{k, n} \equiv 1$  on  $\mathbb{R}^d$ . Now, we define  $\Gamma_{k, n}$  to be the Fourier multiplication operator with symbol  $\gamma_{k, n}$ .

There is the following refinement of the classical Strichartz estimate. In the context of Strichartz–Pecher inequalities for the wave equation, the underlying decay estimate after localization to cubes has been proved in [14, (A.59)]; see also [18, Theorem 4.1] for the case  $p = q = 4$ .

**Lemma 3.1.** *Let  $d = 3$  and  $1/p + 1/q = 1/2$  with  $p > 2$ . Then*

$$\sup_{0 \leq k' \leq k} 2^{-(k'+k)/p} \left( \sum_{n \in \mathfrak{E}_{k'}} \|\Gamma_{k',n} P_k e^{\pm it \langle D \rangle} f\|_{L_t^p L_x^q}^2 \right)^{1/2} \lesssim \|f\|_{L^2(\mathbb{R}^3)}. \tag{3.1}$$

*Proof.* By orthogonality, it suffices to prove

$$\|\Gamma_{k',n} P_k e^{\pm it \langle D \rangle} f\|_{L_t^p L_x^q} \lesssim 2^{(k'+k)/p} \|f\|_{L^2(\mathbb{R}^3)},$$

uniformly in  $n \in \mathfrak{E}_{k'}$ . Let  $T = \Gamma_{k',n} P_k e^{\pm it \langle D \rangle}$ . The operator  $TT^*$  is a space-time convolution operator with kernel

$$K_{k',k;n}(t, x) = \int_{\mathbb{R}^3} e^{\pm it \langle \xi \rangle + ix \cdot \xi} \rho_k^2(\xi) \gamma_{k',n}^2(\xi) d\xi.$$

By the  $TT^*$ -argument, it suffices to prove

$$\|TT^*\|_{L_t^{p'} L_x^{q'} \rightarrow L_t^p L_x^q} \lesssim 2^{2(k+k')/p},$$

which reduces to proving the kernel bound

$$|K_{k',k;n}(t, x)| \lesssim 2^{3k'} (1 + 2^{2k'-k} |t|)^{-1}. \tag{3.2}$$

Indeed, by interpolation and Young’s inequality, we obtain

$$\|K_{k',k;n}(t, \cdot) * \phi\|_{L_x^q(\mathbb{R}^3)} \lesssim 2^{3k'(1-2/q)} (1 + 2^{2k'-k} |t|)^{-(1-2/q)} \|\phi\|_{L_x^{q'}(\mathbb{R}^3)},$$

and Hardy–Littlewood–Sobolev with  $1/r = 2/p = 1 - 2/q$  implies

$$\|TT^*\|_{L_t^{p'} L_x^{q'} \rightarrow L_t^p L_x^q} \lesssim 2^{3k'(1-2/q)} \|(1 + 2^{2k'-k} |t|)^{-(1-2/q)}\|_{L_t^{r,w}} \lesssim 2^{2(k+k')/p}.$$

Finally, we give a proof of (3.2): Rescaling yields

$$K_{k',k;n}(t, x) = 2^{3k} K_{k'-k,1,2^{-k}n}(2^k t, 2^k x),$$

where, for  $\langle \xi \rangle_k := (|\xi|^2 + 2^{-2k})^{1/2}$ ,

$$K_{j,1,a}(s, y) = \int_{\mathbb{R}^3} e^{\pm is \langle \xi \rangle_k + iy \cdot \xi} \rho_1^2(\xi) \gamma_{j,a}^2(\xi) d\xi.$$

For  $|a| \approx 1$ , we claim

$$|K_{j,1,a}(s, y)| \lesssim 2^{3j} (1 + 2^{2j} |s|)^{-1}. \tag{3.3}$$

For  $|s| \leq 2^{-2j}$  this is immediate because the domain of integration has volume  $2^{3j}$ , and in the remaining case it can be proved as for the wave equation in [14, (A.70)]. We provide an explicit proof: By a simple covering argument we may replace  $\rho_1^2 \gamma_{j,a}^2$  by a smooth cutoff  $\zeta$  with respect to a thickened spherical cap of size  $2^j$  and denote the

corresponding kernel by  $\tilde{K}_{j,a}$ . By rotation, we may assume that  $y = (0, 0, |y|)$ . We use spherical coordinates:

$$\tilde{K}_{j,a}(s, y) = \int_0^\infty \int_0^{2\pi} \int_0^\pi e^{i(|y|\rho \cos \theta + s(\rho)_k)} \zeta(\theta, \varphi, \rho) \sin(\theta) \rho^2 d\theta d\varphi d\rho.$$

We may choose  $\zeta(\varphi, \theta, \rho) = \zeta_1(\theta)\zeta_2(\varphi)\zeta_3(\rho)$ . The phase of the oscillatory integral is stationary only if  $|y| \approx |s|$ , and the cap is centered near the north pole or south pole, otherwise we get arbitrarily fast decay. We discuss only the first case, where we may further assume that  $|\zeta'_1| \lesssim 2^{-j}$ ,  $\zeta_1$  is supported in an interval of length  $\lesssim 2^j$  in  $[0, \pi)$ , and  $\zeta_3$  is supported in an interval of length  $\lesssim 2^j$  in  $(1/4, 4)$ , with  $|\zeta'_3| \lesssim 2^{-j}$ . We integrate by parts with respect to  $\theta$ :

$$\begin{aligned} \tilde{K}_{j,a}(s, y) &= \frac{i\zeta_1(0)}{|y|} \int_0^\infty \int_0^{2\pi} e^{i(|y|\rho + s(\rho)_k)} \zeta_2(\varphi)\zeta_3(\rho) \rho d\varphi d\rho \\ &\quad - \frac{i}{|y|} \int_0^\infty \int_0^{2\pi} \int_0^\pi e^{i(|y|\rho \cos \theta + s(\rho)_k)} \zeta'_1(\theta) d\theta \zeta_2(\varphi)\zeta_3(\rho) \rho d\varphi d\rho, \end{aligned}$$

and the properties of  $\zeta_1$  and  $\zeta_3$  imply

$$|\tilde{K}_{j,a}(s, y)| \lesssim 2^j |y|^{-1},$$

which completes the proof of (3.3), which implies (3.2). □

**Remark 3.2.** The generalization of Lemma 3.1 to general dimension and nonsharp admissible pairs is obvious, but we do not need it here.

Now, we consider functions  $f \in L_t^\infty(\mathbb{R}; L^2(\mathbb{R}^3; \mathbb{C}^d))$ . We will use  $d = 1$  for the Klein–Gordon part and  $d = 4$  for the Dirac part. For  $k \in \mathbb{Z}, k \geq 0$ , and  $l, k' \in \mathbb{Z}, 0 \leq k', l \leq k$ , we define

$$\|f\|_{L_t^p L_x^q[k;l,k']} := \left( \sum_{\kappa \in \mathcal{K}_l} \sum_{n \in \mathbb{Z}_{k'}} \|\Gamma_{k',n} P_\kappa f\|_{L_t^p L_x^q}^2 \right)^{1/2}.$$

Note that the above norm for  $l = 0$  is similar to the one in (3.1). The general case  $0 \leq l \leq k$  is needed for technical reasons.

For  $k \geq 0$ , we define

$$\begin{aligned} \|f\|_{S_k^\pm} &= \|f\|_{L_t^\infty L_x^2} + \|f\|_{\dot{X}^{\pm, 1/2, \infty}} \\ &\quad + \sup_{0 \leq k', l \leq k} (2^{-(k'+k)/3} \|f\|_{L_t^3 L_x^6[k;l,k']} + 2^{-(k'+k)/6} \|f\|_{L_t^6 L_x^3[k;l,k']}). \end{aligned} \tag{3.4}$$

Note that if  $k' = k$  and  $l = 0$ , that is, no additional localization is provided, the last two norms are simply the standard Strichartz estimates  $L_t^3 L_x^6$  and  $L_t^6 L_x^3$  available for the wave equation in  $\mathbb{R}^3$ .

In the nonlinear estimates we will use the fact that  $\|P_{\leq 0} f\|_{S_{\leq 0}^\pm}$  also dominates (by interpolation and the Sobolev embedding) the localized Strichartz norms (with  $k = 0$ ) available in the high frequency structure.

Next, we consider boundedness properties of certain multipliers.

**Lemma 3.3.** (i) Let  $s_1, s_2 \in \{+, -\}$ . For any  $k_1, k_2 \in \mathbb{N}_0$  with  $1 \leq l \leq \min(k_1, k_2) + 10$ ,  $\kappa_1, \kappa_2 \in \mathcal{K}_l$  with  $d(s_1\kappa_1, s_2\kappa_2) \lesssim 2^{-l}$ , and  $v_1, v_2 \in \mathbb{C}^4$ , we have

$$|\langle \Pi_{s_1}(2^{k_1}\omega(\kappa_1))v_1, \beta \Pi_{s_2}(2^{k_2}\omega(\kappa_2))v_2 \rangle| \lesssim 2^{-l} |v_1| |v_2|. \tag{3.5}$$

Fix  $k \in \mathbb{N}_0$ . All the statements below are made for functions localized at frequency  $2^k$ , i.e. they satisfy  $f = \tilde{P}_k f$ .

(ii) For any  $1 \leq l \leq k + 10$ ,  $\kappa \in \mathcal{K}_l$ , and  $f \in S_k^\pm$ , we have

$$\|[\Pi_\pm(D) - \Pi_\pm(2^k\omega(\kappa))]P_\kappa f\|_{S_k^\pm} \lesssim 2^{-l} \|P_\kappa f\|_{S_k^\pm}, \tag{3.6}$$

and similarly in  $L_t^p L_x^q$ -norms.

(iii) For any  $j \in \mathbb{Z}$ , the operators  $Q_j^\pm$  are uniformly bounded on  $S_k^\pm$ .

(iv) For any  $l \in \mathbb{N}_0$ ,  $\kappa \in \mathcal{K}_l$ , and  $j \in \mathbb{Z}$  with  $j \geq k - 2l - 100$ , the operators  $Q_{>j}^\pm \tilde{P}_\kappa$  and  $Q_{\leq j}^\pm \tilde{P}_\kappa$  are uniformly bounded on  $S_k^\pm$ .

(v) For any  $k' \in \mathbb{N}_0$  and  $j \in \mathbb{Z}$  satisfying  $k' \leq k$  and  $j \geq 2k' - k$ , the operators  $Q_{>j}^\pm$  and  $Q_{\leq j}^\pm$  are uniformly disposable in the sense that

$$\sup_{0 \leq l \leq k} (2^{-(k'+k)/3} \|Q_{>j}^\pm f\|_{L_t^3 L_x^6[k;l,k']} + 2^{-(k'+k)/6} \|Q_{\leq j}^\pm f\|_{L_t^6 L_x^3[k;l,k']}) \lesssim \|f\|_{S_k^\pm}.$$

Further, similar estimates for  $Q_{>j}^\pm$  and  $Q_{\leq j}^\pm$  hold with a bound  $\langle k' \rangle$  as long as  $j \geq -k'$ .

*Proof.* The identity (2.3) implies

$$\begin{aligned} & \langle \Pi_{s_1}(2^{k_1}\omega(\kappa_1))v_1, \beta \Pi_{s_2}(2^{k_2}\omega(\kappa_2))v_2 \rangle \\ &= \langle \beta \Pi_{-s_2}(2^{k_2}\omega(\kappa_2)) \Pi_{s_1}(2^{k_1}\omega(\kappa_1))v_1, v_2 \rangle + s_2 \langle 2^{k_2} \rangle^{-1} \langle \beta \Pi_{s_1}(2^{k_1}\omega(\kappa_1))v_1, v_2 \rangle, \end{aligned}$$

hence (3.5) follows from estimates (2.4) and Cauchy–Schwarz.

In order to prove (3.6), it suffices to consider the case of the + sign. We write the matrix-valued symbol  $p$  of  $2[\Pi_+(D) - \Pi_+(2^k\omega(\kappa))]P_k \tilde{P}_\kappa$  as

$$\begin{aligned} p(\xi) &= 2[\Pi_+(\xi) - \Pi_+(2^k\omega(\kappa))] \tilde{\rho}_k(\xi) \tilde{\eta}_\kappa(\xi) \\ &= \tilde{\rho}_k(\xi) \tilde{\eta}_\kappa(\xi) \left[ \frac{\xi}{\langle \xi \rangle} - \frac{2^k\omega(\kappa)}{\langle 2^k \rangle} \right] \cdot \alpha + \tilde{\rho}_k(\xi) \tilde{\eta}_\kappa(\xi) \left[ \frac{1}{\langle \xi \rangle} - \frac{1}{\langle 2^k \rangle} \right] \beta \\ &=: p_1(\xi) + p_2(\xi). \end{aligned}$$

We further decompose

$$\begin{aligned} p_1(\xi) &= \tilde{\rho}_k(\xi) \tilde{\eta}_\kappa(\xi) \left[ \frac{|\xi|}{\langle \xi \rangle} - \frac{2^k}{\langle 2^k \rangle} \right] \omega(\kappa) \cdot \alpha + \tilde{\rho}_k(\xi) \tilde{\eta}_\kappa(\xi) \frac{|\xi|}{\langle \xi \rangle} \left[ \frac{\xi}{|\xi|} - \omega(\kappa) \right] \cdot \alpha \\ &=: p_{11}(\xi) + p_{12}(\xi). \end{aligned}$$

We denote the Fourier multiplication operators defined by the symbols above by  $P_2(D)$ ,  $P_{11}(D)$ ,  $P_{12}(D)$ . Obviously, the properties of  $\tilde{\rho}_k$  imply that

$$\|P_2(D)\|_{L_x^p \rightarrow L_x^p} \lesssim 2^{-k}, \quad \|P_{11}(D)\|_{L_x^p \rightarrow L_x^p} \lesssim 2^{-k}, \quad \text{for any } 1 < p < \infty,$$

and the properties of  $\tilde{\eta}_\kappa$  imply that

$$\|P_{12}(D)\|_{L_x^p \rightarrow L_x^p} \lesssim 2^{-l} \quad \text{for any } 1 < p < \infty.$$

The claim follows from the definition of the space  $S_k^+$ .

Part (iii) needs to be proved for the Strichartz norms only. For the operator  $Q_j^\pm$  this is an easy consequence of the well-known transference principle. Indeed,

$$Q_j^\pm f(t) = \int e^{it\tau} e^{\mp it\langle D \rangle} \mathcal{F}_t(e^{\pm it\langle D \rangle} f)(\tau) \tilde{\rho}_j(\tau) d\tau,$$

hence by Lemma 3.1 we obtain

$$2^{-(k'+k)/p} \|Q_j^\pm f\|_{L_t^p L_x^q[k;l,k']} \lesssim \|\mathcal{F}_t(e^{\pm it\langle D \rangle} f) \tilde{\rho}_j\|_{L_t^1 L_x^2} \lesssim 2^{j/2} \|\tilde{Q}_j^\pm f\|_{L^2} \approx \|f\|_{\dot{X}^{\pm,1/2,\infty}}.$$

In order to prove (iv), we apply Sobolev inequalities to obtain, for any  $\kappa' \in \mathcal{K}_{l'}$  and  $n \in \mathfrak{E}_{\kappa'}$ ,

$$\begin{aligned} &2^{-(k'+k)/p} \|\Gamma_{\kappa',n} P_{\kappa'} Q_j^\pm P_\kappa f\|_{L_t^p L_x^q} \\ &\lesssim 2^{-k'+k/p} 2^{j(1/2-1/p)} 2^{(k'+\min(2k-2l,2k'))(1/2-1/q)} \|\Gamma_{\kappa',n} P_{\kappa'} Q_j^\pm P_\kappa f\|_{L^2}. \end{aligned}$$

Summing the squares with respect to  $\kappa', n$  yields

$$2^{-(k'+k)/p} \|Q_j^\pm P_\kappa f\|_{L_t^p L_x^q[k;l',k']} \lesssim 2^{\frac{\min(k-2l,2k'-k)-j}{p}} 2^{j/2} \|Q_j^\pm f\|_{L^2}, \quad (3.7)$$

which we finally sum with respect to  $j \geq j_0 \geq k - 2l - 100$  to obtain

$$2^{-(k'+k)/p} \|Q_{>j_0}^\pm P_\kappa f\|_{L_t^p L_x^q[k;l',k']} \lesssim \|f\|_{\dot{X}^{\pm,1/2,\infty}}.$$

The remaining claim in (iv) follows from  $Q_{\leq j}^\pm = I - Q_{>j}^\pm$ .

Part (v) follows similarly from (3.7). The last claim for  $Q_{>j}^\pm$  follows by applying (iii) and (v) to

$$Q_{>j}^\pm = Q_{>2k'-k}^\pm + \sum_{j < j' \leq 2k'-k} Q_{j'}^\pm,$$

because the number of terms in the second sum is bounded by  $\langle k' \rangle$ . The claim for  $Q_{\leq j}^\pm = I - Q_{>j}^\pm$  follows, too.  $\square$

The next lemma shows why the  $S_k^\pm$ -seminorms are useful in the context of the evolution equation.

**Lemma 3.4.** For any  $k \in \mathbb{N}_0$ ,  $u_0 = \tilde{P}_k u_0 \in L^2(\mathbb{R}^3; \mathbb{C}^d)$  and  $f = \tilde{P}_k f \in L^1_t(\mathbb{R}; L^2(\mathbb{R}^3; \mathbb{C}^d))$ , let

$$u(t) = e^{\mp it \langle D \rangle} u_0 + i \int_0^t e^{\mp i(t-s) \langle D \rangle} f(s) ds.$$

Then  $u = \tilde{P}_k u$  is the unique solution of

$$-i \partial_t u \pm \langle D \rangle u = f,$$

and  $u \in C(\mathbb{R}, L^2(\mathbb{R}^3; \mathbb{C}^d))$  and

$$\|u\|_{S_k^\pm} \lesssim \|u_0\|_{L^2(\mathbb{R}^3)} + \sup_{g \in G} \left| \int_{\mathbb{R}^{1+3}} \langle f, g \rangle_{\mathbb{C}^d} dx dt \right| \tag{3.8}$$

provided that the right hand side of (3.8) is finite, where  $G$  is defined as the set of all  $g = \tilde{P}_k g \in L^\infty_t(\mathbb{R}; L^2(\mathbb{R}^3; \mathbb{C}^d))$  such that  $\|g\|_{S_k^\pm} = 1$ .

*Proof.* Without the localization in  $L^3_t L^6_x, L^6_t L^3_x$  the linear theory above is standard using  $X^{s,b}$  theory and the Christ–Kiselev Lemma [7]. It is likely that one can adapt the Christ–Kiselev Lemma to cover the localized versions of  $L^3_t L^6_x, L^6_t L^3_x$  and their dual structures as well, but we do not pursue this strategy here. Instead, we will give a rather short proof using the theory of  $U^2$  and  $V^2$  spaces; see e.g. [15, 11, 16] for details. We recall that for  $1 < p < \infty$  the atomic space  $U_{\pm \langle D \rangle}^p$  is defined via its atoms

$$a(t) = \sum_{k=1}^K \mathbb{1}_{[t_{k-1}, t_k)}(t) e^{\mp it \langle D \rangle} \phi_k, \quad \sum_{k=1}^K \|\phi_k\|_{L^2}^p = 1,$$

where  $\{t_k\}$  is a partition,  $t_K = \infty$ .

As a companion space we use the space  $V_{\pm \langle D \rangle}^p$  of right-continuous functions  $v$  such that  $t \mapsto e^{\pm it \langle D \rangle} v(t)$  is of bounded  $p$ -variation. We have  $V_{\pm \langle D \rangle}^2 \hookrightarrow U_{\pm \langle D \rangle}^p$  for  $p > 2$ .

For  $0 \leq l, k' \leq k$  we define

$$\|u\|_{U_{k;l,k'}^\pm} := \left( \sum_{\kappa \in \mathcal{K}_l} \sum_{n \in \mathbb{Z}_{k'}} \|\Gamma_{k',n} P_\kappa u\|_{U_{\pm \langle D \rangle}^2}^2 \right)^{1/2}. \tag{3.9}$$

Then we have

$$\|u\|_{V_{k;l,k'}^\pm} := \left( \sum_{\kappa \in \mathcal{K}_l} \sum_{n \in \mathbb{Z}_{k'}} \|\Gamma_{k',n} P_\kappa u\|_{V_{\pm \langle D \rangle}^2}^2 \right)^{1/2} \lesssim \|u\|_{U_{k;l,k'}^\pm}. \tag{3.10}$$

It is easy to show that the  $U_{k;l,k'}^\pm$ -norms are decreasing if we localize to smaller scales, i.e.

$$\|u\|_{U_{k;l,k'}^\pm} \lesssim \|u\|_{U_{k;\tilde{l},\tilde{k}'}^\pm} \quad \text{if } \tilde{l} \leq l \text{ and } \tilde{k}' \geq k',$$

and the  $V_{k;l,k'}^\pm$ -norms are increasing if we localize to smaller scales, i.e.

$$\|u\|_{V_{k;l,k'}^\pm} \lesssim \|u\|_{V_{k;\tilde{l},\tilde{k}'}^\pm} \quad \text{if } \tilde{l} \geq l \text{ and } \tilde{k}' \leq k'.$$

Set  $U_k^\pm = U_{k;k,0}^\pm$  and  $V_k^\pm = V_{k;k,0}^\pm$ .

Strichartz estimates for admissible pairs  $(p, q)$  hold for  $U_{\pm(D)}^p$ -functions (which is easily verified for atoms), hence also for  $V_{\pm(D)}^2$ -functions. For any  $0 \leq k', l \leq k$  we have

$$2^{-(k'+k)/p} \left( \sum_{\kappa \in \mathcal{K}_l} \sum_{n \in \Xi_{k'}} \|\Gamma_{k',n} P_\kappa u\|_{L_t^p L_x^q}^2 \right)^{1/2} \lesssim \|u\|_{V_{k,l,k'}^\pm} \lesssim \|u\|_{V_k^\pm}.$$

We also have  $V_k^\pm \hookrightarrow V_{\pm(D)}^2$  and the  $V_{\pm(D)}^2$ -norm dominates both the  $L_t^\infty L_x^2$ -norm and the  $\dot{X}^{\pm,1/2,\infty}$ -seminorm. Hence,

$$\|u\|_{S_k^\pm} \lesssim \|u\|_{V_k^\pm} \lesssim \|u\|_{U_k^\pm}.$$

Now, we can use the  $U^2$  duality theory (see e.g. [11, Prop. 2.10], and [12, Prop. 2.11] for a frequency-localized version) to conclude that

$$\|u\|_{U_k^\pm} \lesssim \|u_0\|_{L^2(\mathbb{R}^3)} + \sup_{h \in H} \left| \int_{\mathbb{R}^{1+3}} \langle f, h \rangle_{\mathbb{C}^d} dx dt \right|,$$

where  $H$  is defined as the set of all  $h = \tilde{P}_k h$  such that  $\|h\|_{V_k^\pm} = 1$ . The claim now follows by using again  $\|g\|_{S_k^\pm} \lesssim \|g\|_{V_k^\pm}$ . □

**Remark 3.5.** In fact, we have proved a stronger result: In the setting of Lemma 3.4, provided that the right hand side of (3.8) is finite, we can upgrade this estimate to

$$\|u\|_{U_k^\pm} \lesssim \|u_0\|_{L^2(\mathbb{R}^3)} + \sup_{g \in G} \left| \int_{\mathbb{R}^{1+3}} \langle f, g \rangle_{\mathbb{C}^d} dx dt \right|.$$

Our resolution space  $S^{\pm,\sigma}$  corresponding to Sobolev regularity  $\sigma$ —used in Subsection 4.2—will be the space of functions in  $C(\mathbb{R}, H^\sigma(\mathbb{R}^3; \mathbb{C}^d))$  such that

$$\|f\|_{S^{\pm,\sigma}} = \|P_{\leq 0} f\|_{S_{\leq 0}^\pm} + \left( \sum_{k \geq 1} 2^{2\sigma k} \|P_k f\|_{S_k^\pm}^2 \right)^{1/2} < \infty,$$

which is obviously a Banach space.

#### 4. Nonlinear estimates and the proof of the main result

Recall (2.2) with the convention  $M = m = 1$  and use the decomposition  $\psi = \Pi_+(D)\psi + \Pi_-(D)\psi$  in the nonlinearity (for all three terms). It then suffices to prove

$$\begin{aligned} \left| \int \langle \Pi_{S_2}(D)[\mathfrak{R}\phi \beta \Pi_{S_1}(D)\psi_1], \psi_2 \rangle dx dt \right| &\lesssim \|\phi\|_{S^{+,1/2+\epsilon}} \|\psi_1\|_{S^{s_1,\epsilon}} \|\psi_2\|_{S^{s_2,-\epsilon}}, \\ \left| \int \langle D \rangle^{-1} \langle \Pi_{S_1}(D)\psi_1, \beta \Pi_{S_2}(D)\psi_2 \rangle \bar{\phi} dx dt \right| &\lesssim \|\phi\|_{S^{+,-1/2-\epsilon}} \|\psi_1\|_{S^{s_1,\epsilon}} \|\psi_2\|_{S^{s_2,\epsilon}}, \end{aligned}$$

for any choice of signs  $s_1, s_2 \in \{+, -\}$ . By symmetry, this follows from

$$\left| \int \phi \langle \Pi_{s_1}(D)\psi_1, \beta \Pi_{s_2}(D)\psi_2 \rangle dx dt \right| \lesssim \|\phi\|_{S^{+,1/2+\epsilon_0}} \|\psi_1\|_{S^{s_1,\epsilon_1}} \|\psi_2\|_{S^{s_2,\epsilon_2}} \quad (4.1)$$

where  $\epsilon_0, \epsilon_1, \epsilon_2 \in \{\pm\epsilon\}$  are such that  $\epsilon_0 + \epsilon_1 + \epsilon_2 = \epsilon$ . More precisely, we will prove this first on the dyadic level, where all integrals are clearly finite (see Lemma 3.4).

4.1. Estimates for dyadic pieces

Our aim will be to identify a function  $G : \mathbb{N}_0^3 \rightarrow (0, \infty)$  such that

$$\sum_{\substack{k, k_1, k_2 \in \mathbb{N}_0 \\ \max(k, k_1, k_2) \sim \text{med}(k, k_1, k_2)}} \frac{G(k, k_1, k_2) a_k b_{k_1} c_{k_2}}{2^{k/2} (\min(k, k_1, k_2) + 1)^{10}} \lesssim \|a\|_{l^2} \|b\|_{l^2} \|c\|_{l^2} \quad (4.2)$$

for all sequences  $a = (a_j)_{j \in \mathbb{N}_0}$  etc. in  $l^2(\mathbb{N}_0)$ . We write  $\mathbf{k} = (k, k_1, k_2)$ .

Clearly, (4.1) is implied by the following key result of this section:

**Proposition 4.1.** *Let  $s_1, s_2 \in \{+, -\}$ . There exists a function  $G$  satisfying (4.2) such that for all  $\phi = P_k \phi$  and  $\psi_i = P_{k_i} \Pi_{s_i}(D)\psi_i, i = 1, 2,$*

$$\left| \int \phi \langle \psi_1, \beta \psi_2 \rangle dx dt \right| \lesssim G(\mathbf{k}) \|\phi\|_{S_k^+} \|\psi_1\|_{S_{k_1}^{s_1}} \|\psi_2\|_{S_{k_2}^{s_2}}. \quad (4.3)$$

*Proof.* We denote the integral on the left hand side of (4.3) by  $I(\mathbf{k})$ . Without restricting the generality of the argument we can assume that  $k_1 \leq k_2$ . We decompose

$$I(\mathbf{k}) = I_0(\mathbf{k}) + I_1(\mathbf{k}) + I_2(\mathbf{k})$$

where

$$\begin{aligned} I_0(\mathbf{k}) &:= \sum_{j \in \mathbb{Z}} \int Q_j^+ \phi \langle Q_{\leq j}^{s_1} \psi_1, \beta Q_{\leq j}^{s_2} \psi_2 \rangle dx dt, \\ I_1(\mathbf{k}) &:= \sum_{j_1 \in \mathbb{Z}} \int Q_{< j_1}^+ \phi \langle Q_{j_1}^{s_1} \psi_1, \beta Q_{\leq j_1}^{s_2} \psi_2 \rangle dx dt, \\ I_2(\mathbf{k}) &:= \sum_{j_2 \in \mathbb{Z}} \int Q_{< j_2}^+ \phi \langle Q_{< j_2}^{s_1} \psi_1, \beta Q_{j_2}^{s_2} \psi_2 \rangle dx dt. \end{aligned}$$

Given the symmetry of the estimate in  $k_1$  and  $k_2$ , we split the argument into two cases.

**Case 1:**  $|k - k_2| \leq 10$ .

*Contribution of  $I_0(\mathbf{k})$ :* We split  $I_0(\mathbf{k}) = I_{01}(\mathbf{k}) + I_{02}(\mathbf{k})$  according to  $j < k_1$  and  $j \geq k_1$ . Then, due to Lemma 2.4, there is no contribution if  $j < k_1$  in the case  $s_1 = +, s_2 = -$ .



For all other choices of signs, we estimate

$$I_{01}(\mathbf{k}) \lesssim \sum_{-k_1 \leq j < k_1} \sum_{\substack{n, n' \in \mathbb{E}_{k_1} \\ |n-n'| \leq k_1}} \|\Gamma_{k_1, n} Q_j^+ \phi\|_{L^2} \|\langle Q_{\leq j}^{s_1} \psi_1, \beta Q_{\leq j}^{s_2} \Gamma_{k_1, n'} \psi_2 \rangle\|_{L^2},$$

where we have used orthogonality, and the nonresonance bound (2.9) to restrict the sum to the range  $j \geq -k_1$ . We conclude from Lemma 2.4 with  $2l = k_1 + k_2 - k - j$  and Lemma 3.3 that

$$\begin{aligned} & \|\langle Q_{\leq j}^{s_1} \psi_1, \beta Q_{\leq j}^{s_2} \Gamma_{k, n'} \psi_2 \rangle\|_{L^2} \\ & \lesssim 2^{-l} \sum_{\substack{\kappa_1, \kappa_2 \in \mathcal{K}_l \\ d(s_1 \kappa_1, s_2 \kappa_2) \lesssim 2^{-l}}} \|Q_{\leq j}^{s_1} P_{\kappa_1} \psi_1\|_{L_t^3 L_x^6} \|Q_{\leq j}^{s_2} P_{\kappa_2} \Gamma_{k_1, n'} \psi_2\|_{L_t^6 L_x^3}. \end{aligned}$$

By Lemma 3.3(v), the operators  $Q_{\leq j}^\pm$  are disposable up to a factor  $\langle k_1 \rangle$ . Then, we apply Cauchy–Schwarz and perform the cube and cap summation to obtain

$$\begin{aligned} I_{01}(\mathbf{k}) & \lesssim \sum_{-k_1 \leq j < k_1} 2^{-j/2} \|\phi\|_{S_k} 2^{-(k_1+k_2-k-j)/2} 2^{2k_1/3} \langle k_1 \rangle \|\psi_1\|_{S_{k_1}^{s_1}} 2^{(k_1+k_2)/6} \langle k_1 \rangle \|\psi_2\|_{S_{k_2}^{s_2}} \\ & \lesssim \langle k_1 \rangle^3 2^{(k_1-k_2)/3} 2^{k/2} \|\phi\|_{S_k^+} \|\psi_1\|_{S_{k_1}^{s_1}} \|\psi_2\|_{S_{k_2}^{s_2}}. \end{aligned}$$

In the range  $j \geq k_1$ , the operators  $Q_{\leq j}^\pm$  are disposable and a similar argument to the one above with  $l = 0$ , i.e. no cap decomposition and no gain from the null structure, gives the bound

$$I_{02}(\mathbf{k}) \lesssim 2^{(k_1-k_2)/3} 2^{k/2} \|\phi\|_{S_k^+} \|\psi_1\|_{S_{k_1}^{s_1}} \|\psi_2\|_{S_{k_2}^{s_2}}.$$

*Contribution of  $I_1(\mathbf{k})$ :* We split  $I_1(\mathbf{k}) = I_{11}(\mathbf{k}) + I_{12}(\mathbf{k})$  according to  $j_1 < k_1$  and  $j_1 \geq k_1$ . Again, by Lemma 2.4 there is no contribution if  $j_1 < k_1$  in the case  $s_1 = +, s_2 = -$ . For all other choices of signs, we can restrict the sum in  $I_{11}$  to  $j_1 \geq -k_1$ , so that by Lemma 2.4 with  $2l = k_1 + k_2 - k - j_1 \sim k_1 - j_1$  we have

$$I_{11}(\mathbf{k}) = \sum_{-k_1 \leq j_1 < k_1} \sum_{\substack{n, n' \in \mathbb{E}_{k_1} \\ |n-n'| \leq k_1}} \sum_{\substack{\kappa_1, \kappa_2 \in \mathcal{K}_l \\ d(s_1 \kappa_1, s_2 \kappa_2) \lesssim 2^{-l}}} \int \Gamma_{k_1, n} Q_{\leq j_1}^+ \phi \cdot \langle P_{\kappa_1} Q_{j_1}^{s_1} \psi_1, \beta P_{\kappa_2} \Gamma_{k_1, n'} Q_{\leq j_1}^{s_2} \psi_2 \rangle dx dt.$$

In view of Lemma 3.3, we decompose

$$\Pi_{s_i}(D) P_{\kappa_i} = [\Pi_{s_i}(D) - \Pi_{s_i}(2^{k_i} \omega(\kappa_i))] P_{\kappa_i} + \Pi_{s_i}(2^{k_i} \omega(\kappa_i)) P_{\kappa_i},$$

and obtain

$$\begin{aligned} & \|\langle P_{\kappa_1} Q_{j_1}^{s_1} \psi_1, \beta P_{\kappa_2} \Gamma_{k_1, n'} Q_{\leq j_1}^{s_2} \psi_2 \rangle\|_{L_t^{3/2} L_x^{6/5}} \\ & \lesssim 2^{-l} \|P_{\kappa_1} Q_{j_1}^{s_1} \psi_1\|_{L^2} \|P_{\kappa_2} \Gamma_{k_1, n'} Q_{\leq j_1}^{s_2} \psi_2\|_{L_t^6 L_x^3}. \end{aligned}$$

By Hölder’s inequality and Cauchy–Schwarz we obtain

$$\begin{aligned}
 I_{11}(\mathbf{k}) &\lesssim \sum_{-k_1 \leq j_1 < k_1} \left\{ 2^{-(k_1-j_1)/2} \|Q_{j_1}^{s_1} \psi_1\|_{L^2} \left( \sum_{n \in \mathfrak{B}_{k_1}} \|\Gamma_{k_1, n} Q_{< j_1}^+ \phi\|_{L_t^3 L_x^6}^2 \right)^{1/2} \right. \\
 &\quad \cdot \left. \left( \sum_{n' \in \mathfrak{B}_{k_1}} \sum_{\kappa_2 \in \mathcal{K}_l} \|P_{\kappa_2} \Gamma_{k_1, n'} Q_{\leq j_1}^{s_2} \psi_2\|_{L_t^6 L_x^3}^2 \right)^{1/2} \right\} \\
 &\lesssim \sum_{-k_1 \leq j_1 < k_1} 2^{-(k_1-j_1)/2} 2^{-j_1/2} \|\psi_1\|_{S_{k_1}^{s_1}} 2^{(k_1+k)/3} \langle k_1 \rangle \|\phi\|_{S_k^+} 2^{(k_1+k_2)/6} \langle k_1 \rangle \|\psi_2\|_{S_{k_2}^{s_2}} \\
 &\lesssim 2^{k/2} \langle k_1 \rangle^3 \|\phi\|_{S_k^+} \|\psi_1\|_{S_{k_1}^{s_1}} \|\psi_2\|_{S_{k_2}^{s_2}},
 \end{aligned}$$

where we have used Lemma 3.3(v).

In the range  $j_1 \geq k_1$ , we forego the gain from the null structure in the above argument and obtain

$$\begin{aligned}
 I_{12}(\mathbf{k}) &\lesssim \sum_{j_1 \geq k_1} 2^{-j_1/2} \|\psi_1\|_{S_{k_1}^{s_1}} 2^{(k_1+k)/3} \|\phi\|_{S_k^+} 2^{(k_1+k_2)/6} \|\psi_2\|_{S_{k_2}^{s_2}} \\
 &\lesssim 2^{k/2} \|\phi\|_{S_k^+} \|\psi_1\|_{S_{k_1}^{s_1}} \|\psi_2\|_{S_{k_2}^{s_2}}
 \end{aligned}$$

since the operators  $Q_{\leq j}^\pm$  are disposable.

*Contribution of  $I_2(\mathbf{k})$ :* As above, we split  $I_2(\mathbf{k}) = I_{21}(\mathbf{k}) + I_{22}(\mathbf{k})$  according to  $j_2 < k_1$  and  $j_2 \geq k_1$ . Again, by Lemma 2.4 there is no contribution if  $j_2 < k_1$  in the case  $s_1 = +$ ,  $s_2 = -$ , whereas for all other choices of signs, we can restrict the sum in  $I_{21}(\mathbf{k})$  to  $j_2 \geq -k_1$ , so that by Lemma 2.4 with  $2l = k_1 + k_2 - k - j_2 \sim k_1 - j_2$  we repeat the argument for  $I_{11}(\mathbf{k})$  to obtain

$$\begin{aligned}
 I_{21}(\mathbf{k}) &\lesssim \sum_{-k_1 \leq j_2 < k_1} 2^{(k_1+k)/3} \langle k_1 \rangle \|\phi\|_{S_k^+} 2^{-(k_1-j_2)/2} 2^{k_1/3} \langle k_1 \rangle \|\psi_1\|_{S_{k_1}^{s_1}} 2^{-j_2/2} \|\psi_2\|_{S_{k_2}^{s_2}} \\
 &\lesssim 2^{k/2} 2^{(k_1-k)/6} \langle k_1 \rangle^3 \|\phi\|_{S_k^+} \|\psi_1\|_{S_{k_1}^{s_1}} \|\psi_2\|_{S_{k_2}^{s_2}}.
 \end{aligned}$$

For the range  $j_2 \geq k_1$ , the same argument as above, but with no gain from the null structure, gives the bound

$$\begin{aligned}
 I_{22}(\mathbf{k}) &\lesssim \sum_{j_2 \geq k_1} 2^{(k_1+k)/3} \|\phi\|_{S_k^+} 2^{k_1/3} \|\psi_1\|_{S_{k_1}^{s_1}} 2^{-j_2/2} \|\psi_2\|_{S_{k_2}^{s_2}} \\
 &\lesssim 2^{k/2} 2^{(k_1-k)/6} \|\phi\|_{S_k^+} \|\psi_1\|_{S_{k_1}^{s_1}} \|\psi_2\|_{S_{k_2}^{s_2}}.
 \end{aligned}$$

**Case 2:**  $|k_1 - k_2| \leq 10$ .

*Contribution of  $I_0(\mathbf{k})$ :* We split  $I_0(\mathbf{k}) = I_{01}(\mathbf{k}) + I_{02}(\mathbf{k})$  according to  $j < k$  and  $j \geq k$ . Then, due to Lemma 2.4, there is no contribution if  $j < k$  in the case  $s_1 = +$ ,  $s_2 = -$  or

$s_1 = -, s_2 = +$  and  $k < \min(k_1, k_2)$ . In all remaining cases, we can restrict the sum in  $I_{01}$  to  $j \geq -k$ , so that

$$I_{01}(\mathbf{k}) \lesssim \sum_{-k \leq j < k} \sum_{\substack{n, n' \in \Xi_k \\ |n - n'| \leq k}} \|Q_j^+ \phi\|_{L^2} \|\langle Q_{\leq j}^{s_1} \Gamma_{k,n} \psi_1, \beta Q_{\leq j}^{s_2} \Gamma_{k,n'} \psi_2 \rangle\|_{L^2}.$$

We conclude from Lemma 2.4 with  $2l = k_1 + k_2 - k - j$  and Lemma 3.3 that

$$\begin{aligned} & \|\langle Q_{\leq j}^{s_1} \Gamma_{k,n} \psi_1, \beta Q_{\leq j}^{s_2} \Gamma_{k,n'} \psi_2 \rangle\|_{L^2} \\ & \lesssim 2^{-l} \sum_{\substack{\kappa_1, \kappa_2 \in \mathcal{K}_l \\ d(s_1 \kappa_1, s_2 \kappa_2) \lesssim 2^{-l}}} \|Q_{\leq j}^{s_1} P_{\kappa_1} \Gamma_{k,n} \psi_1\|_{L_t^3 L_x^6} \|Q_{\leq j}^{s_2} P_{\kappa_2} \Gamma_{k,n'} \psi_2\|_{L_t^6 L_x^3}. \end{aligned}$$

By Lemma 3.3(v), the operators  $Q_{\leq j}^{\pm}$  are disposable up to a factor  $\langle k \rangle$ . Then, we apply Cauchy–Schwarz and perform the cube and cap summation to obtain

$$\begin{aligned} I_{01}(\mathbf{k}) & \lesssim \sum_{-k \leq j < k} 2^{-j/2} \|\phi\|_{S_k} 2^{-(k_1+k_2-k-j)/2} 2^{(k+k_1)/3} \langle k \rangle \|\psi_1\|_{S_{k_1}^{s_1}} 2^{(k+k_2)/6} \langle k \rangle \|\psi_2\|_{S_{k_2}^{s_2}} \\ & \lesssim \langle k \rangle^3 2^{(k-k_1)/2} 2^{k/2} \|\phi\|_{S_k^+} \|\psi_1\|_{S_{k_1}^{s_1}} \|\psi_2\|_{S_{k_2}^{s_2}}. \end{aligned}$$

Let us now consider the range  $j \geq k$ . Now, by Lemma 3.3(v), the operators  $Q_{\leq j}^{\pm}$  are disposable. In the case  $s_1 = +, s_2 = -$  and in the case  $s_1 = -, s_2 = +$  and  $k < \min(k_1, k_2)$ , Lemma 2.4 implies that there is only a contribution if  $j \geq k_1$ . Then, the above argument with  $l = 0$  yields

$$\begin{aligned} I_{02}(\mathbf{k}) & \lesssim \sum_{j \geq k_1} 2^{-j/2} \|\phi\|_{S_k^+} 2^{(k+k_1)/3} \|\psi_1\|_{S_{k_1}^{s_1}} 2^{(k+k_2)/6} \|\psi_2\|_{S_{k_2}^{s_2}} \\ & \lesssim 2^{k/2} \|\phi\|_{S_k^+} \|\psi_1\|_{S_{k_1}^{s_1}} \|\psi_2\|_{S_{k_2}^{s_2}}. \end{aligned}$$

In the case  $s_1 = s_2$ , (2.13) implies that the integral is nonzero only if the frequencies in the supports of  $\widehat{\psi}_1$  and  $\widehat{\psi}_2$  make an angle of at most  $2^{k-k_1}$ , hence, we choose  $l = k_1 - k$ . In the remaining case where  $s_1 = -, s_2 = +$  and  $k \geq \min(k_1, k_2)$  we choose  $l = 0$ . Again, arguing as for  $I_{01}(\mathbf{k})$  we obtain

$$\begin{aligned} I_{02}(\mathbf{k}) & \lesssim \sum_{j \geq k} 2^{-j/2} \|\phi\|_{S_k} 2^{-l} 2^{(k+k_1)/3} \|\psi_1\|_{S_{k_1}^{s_1}} 2^{(k+k_2)/6} \|\psi_2\|_{S_{k_2}^{s_2}} \\ & \lesssim 2^{k/2} \|\phi\|_{S_k^+} \|\psi_1\|_{S_{k_1}^{s_1}} \|\psi_2\|_{S_{k_2}^{s_2}}. \end{aligned}$$

*Contribution of  $I_1(\mathbf{k})$ :* Again, we split  $I_1(\mathbf{k}) = I_{11}(\mathbf{k}) + I_{12}(\mathbf{k})$  according to  $j_1 < k$  and  $j_1 \geq k$ . Then, due to Lemma 2.4, there is no contribution if  $j_1 < k$  in the case  $s_1 = +$ ,

$s_2 = -$ , or  $s_1 = -$ ,  $s_2 = +$  and  $k < \min(k_1, k_2)$ . In all remaining cases, we can restrict the sum in  $I_{11}$  to  $j_1 \geq -k$ , so that by Lemma 2.4 with  $2l = k_1 + k_2 - k - j_1$  we have

$$I_{11}(\mathbf{k}) = \sum_{-k \leq j_1 < k} \sum_{\substack{n, n' \in \Xi_k \\ |n-n'| \leq k}} \sum_{\substack{\kappa_1, \kappa_2 \in \mathcal{K}_l \\ d(s_1 \kappa_1, s_2 \kappa_2) \lesssim 2^{-l}}} \int Q_{< j_1}^+ \phi \cdot \langle P_{\kappa_1} Q_{j_1}^{s_1} \Gamma_{k,n} \psi_1, \beta P_{\kappa_2} \Gamma_{k,n'} Q_{\leq j_1}^{s_2} \psi_2 \rangle dx dt.$$

Using Lemma 3.3, we obtain

$$\begin{aligned} & \| \langle P_{\kappa_1} Q_{j_1}^{s_1} \Gamma_{k,n} \psi_1, \beta P_{\kappa_2} \Gamma_{k,n'} Q_{\leq j_1}^{s_2} \psi_2 \rangle \|_{L_t^{3/2} L_x^{6/5}} \\ & \lesssim 2^{-l} \| P_{\kappa_1} Q_{j_1}^{s_1} \Gamma_{k,n} \psi_1 \|_{L^2} \| P_{\kappa_2} \Gamma_{k,n'} Q_{\leq j_1}^{s_2} \psi_2 \|_{L_t^6 L_x^3}. \end{aligned}$$

By Hölder’s inequality and Cauchy–Schwarz we obtain

$$\begin{aligned} I_{11}(\mathbf{k}) & \lesssim \sum_{-k \leq j_1 < k} \| Q_{< j_1}^+ \phi \|_{L_t^3 L_x^6} 2^{-(k_1+k_2-k-j_1)/2} \| Q_{j_1}^{s_1} \psi_1 \|_{L^2} \| Q_{\leq j_1}^{s_2} \psi_2 \|_{L_t^6 L_x^3[k_2; l, k]} \\ & \lesssim \sum_{-k \leq j_1 < k} 2^{2k/3} \langle k \rangle \| \phi \|_{S_k^+} 2^{-(k_1+k_2-k-j_1)/2} 2^{-j_1/2} \| \psi_1 \|_{S_{k_1}^{s_1}} 2^{(k+k_2)/6} \langle k \rangle \| \psi_2 \|_{S_{k_2}^{s_2}} \\ & \lesssim 2^{k/2} 2^{5(k-k_1)/6} \langle k \rangle^3 \| \phi \|_{S_k^+} \| \psi_1 \|_{S_{k_1}^{s_1}} \| \psi_2 \|_{S_{k_2}^{s_2}}, \end{aligned}$$

where we have also used Lemma 3.3(v).

Let us now consider the case  $j_1 \geq k$ . We use a dichotomy as for  $I_{02}(\mathbf{k})$ . In the case  $s_1 = +$ ,  $s_2 = -$ , or  $s_1 = -$ ,  $s_2 = +$  and  $k < \min(k_1, k_2)$ , Lemma 2.4 implies that there is only a contribution if  $j_1 \geq k_2$ . In that case, from the above argument with  $l = 0$  we obtain

$$\begin{aligned} I_{12}(\mathbf{k}) & \lesssim \sum_{j_1 \geq k_2} 2^{2k/3} \| \phi \|_{S_k^+} 2^{-j_1/2} \| \psi_1 \|_{S_{k_1}^{s_1}} 2^{(k+k_2)/6} \| \psi_2 \|_{S_{k_2}^{s_2}} \\ & \lesssim 2^{k/2} 2^{(k-k_2)/3} \| \phi \|_{S_k^+} \| \psi_1 \|_{S_{k_1}^{s_1}} \| \psi_2 \|_{S_{k_2}^{s_2}}. \end{aligned}$$

In the case  $s_1 = s_2$ , (2.13) implies that the integral is nonzero only if the frequencies in the supports of  $\widehat{\psi}_1$  and  $\widehat{\psi}_2$  make an angle of at most  $2^{k-k_1}$ , hence, we choose  $l = k_1 - k$ . In the remaining case where  $s_1 = -$ ,  $s_2 = +$  and  $k \geq \min(k_1, k_2)$ , we choose  $l = 0$ . By the argument above we obtain

$$\begin{aligned} I_{12}(\mathbf{k}) & \lesssim \sum_{j_1 \geq k} 2^{2k/3} \| \phi \|_{S_k^+} 2^{-l} 2^{-j_1/2} \| \psi_1 \|_{S_{k_1}^{s_1}} 2^{(k+k_2)/6} \| \psi_2 \|_{S_{k_2}^{s_2}} \\ & \lesssim 2^{k/2} 2^{5(k-k_1)/6} \| \phi \|_{S_k^+} \| \psi_1 \|_{S_{k_1}^{s_1}} \| \psi_2 \|_{S_{k_2}^{s_2}}. \end{aligned}$$

Contribution of  $I_2(\mathbf{k})$ : This is treated in the same way as  $I_1(\mathbf{k})$ . □

**Remark 4.2.** Using  $V^2$ -based spaces one can avoid the logarithmic divergencies in Lemma 3.3(v). We expect that one could obtain a result in the critical Besov space  $\dot{B}_{2,1}^{0,\epsilon} \times \dot{B}_{2,1}^{1/2,\epsilon} \times \dot{B}_{2,1}^{-1/2,\epsilon}$ , where  $\epsilon > 0$  accounts for a bit of angular regularity (somewhat strengthening the null structure and this way eliminating any logarithmic factors). This would improve the result in [20] (which corresponds to  $\epsilon = 1$ ) in the massive case; however, we will not pursue these matters here.

4.2. Proof of Theorem 1.1

Again, for notational convenience, let  $m = M = 1$ . Fix  $\epsilon > 0$ . We will construct a solution

$$(\psi_+, \psi_-, \phi_+) \in \mathbf{S}^\epsilon := S^{+,\epsilon} \times S^{-,\epsilon} \times S^{+,1/2+\epsilon}$$

of the system (2.2) in integral form, i.e.

$$\psi_+(t) = e^{-it\langle D \rangle} \Pi_+(D) \psi_0 + i \int_0^t e^{-i(t-s)\langle D \rangle} \Pi_+(D) [\mathfrak{R}\phi_+ \beta(\psi_+ + \psi_-)] ds,$$

$$\psi_-(t) = e^{it\langle D \rangle} \Pi_-(D) \psi_0 + i \int_0^t e^{i(t-s)\langle D \rangle} \Pi_-(D) [\mathfrak{R}\phi_+ \beta(\psi_+ + \psi_-)] ds,$$

$$\phi_+(t) = e^{-it\langle D \rangle} \phi_{+,0} + i \int_0^t e^{-i(t-s)\langle D \rangle} \langle D \rangle^{-1} \langle (\psi_+ + \psi_-), \beta(\psi_+ + \psi_-) \rangle ds,$$

provided that the initial data satisfy

$$\|\psi_0\|_{H^\epsilon(\mathbb{R}^3)} \leq \delta, \quad \|\phi_{+,0}\|_{H^{1/2+\epsilon}(\mathbb{R}^3)} \leq \delta,$$

for sufficiently small  $\delta > 0$ . Let  $T(\psi_+, \psi_-, \phi_+)$  denote the operator defined by the right hand sides of the above formulas.

By the results of the previous subsection and Lemma 3.4 we conclude that

$$\begin{aligned} \|T(\psi_+, \psi_-, \phi_+)\|_{\mathbf{S}^\epsilon} &\lesssim \delta + \|\phi_+\|_{S^{+,1/2+\epsilon}} (\|\psi_+\|_{S^{+,\epsilon}} + \|\psi_-\|_{S^{-,\epsilon}}) + (\|\psi_+\|_{S^{+,\epsilon}} + \|\psi_-\|_{S^{-,\epsilon}})^2 \\ &\lesssim \delta + \|(\psi_+, \psi_-, \phi_+)\|_{\mathbf{S}^\epsilon}^2, \end{aligned}$$

and similar estimates for differences. Hence, in a small closed ball in the complete space  $\mathbf{S}^\epsilon$  we can invoke the contraction mapping principle to obtain a unique solution. Further, continuous dependence on the initial data is an easy consequence.

It remains to prove that these solutions scatter, which we will only do for  $t \rightarrow \infty$ , the other case being similar. It suffices to show that for a solution  $(\psi_+, \psi_-, \phi_+) \in \mathbf{S}^\epsilon$  we have convergence of the integrals, i.e.

$$\begin{aligned} \lim_{t \rightarrow \infty} \int_0^t e^{-i(-s)\langle D \rangle} \Pi_+(D) [\mathfrak{R}\phi_+ \beta(\psi_+ + \psi_-)] ds &\in H^\epsilon(\mathbb{R}^3), \\ \lim_{t \rightarrow \infty} \int_0^t e^{i(-s)\langle D \rangle} \Pi_-(D) [\mathfrak{R}\phi_+ \beta(\psi_+ + \psi_-)] ds &\in H^\epsilon(\mathbb{R}^3), \\ \lim_{t \rightarrow \infty} \int_0^t e^{-i(-s)\langle D \rangle} \langle D \rangle^{-1} \langle (\psi_+ + \psi_-), \beta(\psi_+ + \psi_-) \rangle ds &\in H^{1/2+\epsilon}(\mathbb{R}^3). \end{aligned}$$

We simply observe that this is a by-product of the linear theory provided by Lemma 3.4. Indeed, by Remark 3.5 it follows that at the dyadic level these integrals are in fact in  $U_k^\pm$  and this is square-summable. From this it follows that they are in the space

$$V^2(\mathbb{R}; H^\epsilon(\mathbb{R}^3)) \times V^2(\mathbb{R}; H^\epsilon(\mathbb{R}^3)) \times V^2(\mathbb{R}; H^{1/2+\epsilon}(\mathbb{R}^3)).$$

Functions of bounded 2-variation have limits at infinity [11, Prop. 2.2], which proves the scattering claim.

*Acknowledgments.* The authors thank Sigmund Selberg and Achenef Tesfahun for spotting a couple of typos and flaws in a previous version of the paper.

I.B. was supported by the NSF grant DMS-1600444. S.H. acknowledges support from the German Research Foundation, Collaborative Research Center 701. Part of this research has been carried out while both authors participated in the Trimester Program *Harmonic Analysis and Partial Differential Equations* at the Hausdorff Research Institute for Mathematics in Bonn.

## References

- [1] Bachelot, A.: Problème de Cauchy global pour des systèmes de Dirac–Klein–Gordon. *Ann. Inst. H. Poincaré Phys. Théor.* **48**, 387–422 (1988) [Zbl 0672.35071](#) [MR 0969173](#)
- [2] Beals, M., Bézard, M.: Low regularity local solutions for field equations. *Comm. Partial Differential Equations* **21**, 79–124 (1996) [Zbl 0852.35098](#) [MR 1373766](#)
- [3] Bejenaru, I., Herr, S.: The cubic Dirac equation: Small initial data in  $H^1(\mathbb{R}^3)$ . *Comm. Math. Phys.* **335**, 43–82 (2015) [Zbl 1321.35180](#) [MR 3314499](#)
- [4] Bjorken, J. D., Drell, S. D.: *Relativistic Quantum Mechanics*. McGraw-Hill, New York (1964) [Zbl 0184.54201](#) [MR 0187641](#)
- [5] Bournaveas, N.: Local existence of energy class solutions for the Dirac–Klein–Gordon equations. *Comm. Partial Differential Equations* **24**, 1167–1193 (1999) [Zbl 0931.35134](#) [MR 1697486](#)
- [6] Chadam, J. M., Glassey, R. T.: On certain global solutions of the Cauchy problem for the (classical) coupled Klein–Gordon–Dirac equations in one and three space dimensions. *Arch. Ration. Mech. Anal.* **54**, 223–237 (1974) [Zbl 0285.35042](#) [MR 0369952](#)
- [7] Christ, M., Kiselev, A.: Maximal functions associated to filtrations. *J. Funct. Anal.* **179**, 409–425 (2001) [Zbl 0974.47025](#) [MR 1809116](#)
- [8] D’Ancona, P., Foschi, D., Selberg, S.: Null structure and almost optimal local regularity for the Dirac–Klein–Gordon system. *J. Eur. Math. Soc.* **9**, 877–899 (2007) [Zbl 1187.35191](#) [MR 2341835](#)
- [9] Delort, J.-M., Fang, D.: Almost global existence for solutions of semilinear Klein–Gordon equations with small weakly decaying Cauchy data. *Comm. Partial Differential Equations* **25**, 2119–2169 (2000) [Zbl 0979.35101](#) [MR 1789923](#)
- [10] Germain, P.: Global existence for coupled Klein–Gordon equations with different speeds. *Ann. Inst. Fourier (Grenoble)* **61**, 2463–2506 (2012) (2011) [Zbl 1255.35162](#) [MR 2976318](#)
- [11] Hadac, M., Herr, S., Koch, H.: Well-posedness and scattering for the KP-II equation in a critical space. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **26**, 917–941 (2009) [Zbl 1169.35372](#) [MR 2526409](#)
- [12] Herr, S., Tataru, D., Tzvetkov, N.: Global well-posedness of the energy-critical nonlinear Schrödinger equation with small initial data in  $H^1(\mathbb{T}^3)$ . *Duke Math. J.* **159**, 329–349 (2011) [Zbl 1230.35130](#) [MR 2824485](#)

- [13] Klainerman, S.: Global existence of small amplitude solutions to nonlinear Klein–Gordon equations in four space-time dimensions. *Comm. Pure Appl. Math.* **38**, 631–641 (1985) [Zbl 0597.35100](#) [MR 0803252](#)
- [14] Klainerman, S., Tataru, D.: On the optimal local regularity for Yang–Mills equations in  $\mathbb{R}^{4+1}$ . *J. Amer. Math. Soc.* **12**, 93–116 (1999) [Zbl 0924.58010](#) [MR 1626261](#)
- [15] Koch, H., Tataru, D.: Dispersive estimates for principally normal pseudodifferential operators. *Comm. Pure Appl. Math.* **58**, 217–284 (2005) [Zbl 1078.35143](#) [MR 2094851](#)
- [16] Koch, H., Tataru, D., Vişan, M.: *Dispersive Equations and Nonlinear Waves*. Oberwolfach Seminars 45, Springer Basel (2014) [Zbl 1304.35003](#)
- [17] Schottdorf, T.: Global existence without decay for quadratic Klein–Gordon equations. [arXiv:1209.1518 \[math.AP\]](#) (2012)
- [18] Selberg, S.: Anisotropic bilinear  $L^2$  estimates related to the 3D wave equation. *Int. Math. Res. Notices* **2008**, art. ID rnn 107, 63 pp. [Zbl 1160.35041](#) [MR 2439535](#)
- [19] Sterbenz, J.: Angular regularity and Strichartz estimates for the wave equation (with an appendix by I. Rodnianski). *Int. Math. Res. Notices* **2005**, 187–231 [Zbl 1072.35048](#) [MR 2128434](#)
- [20] Wang, X.: On global existence of 3D charge critical Dirac–Klein–Gordon system. *Int. Math. Res. Notices* **2015**, 10801–10846 [Zbl 1333.35224](#) [MR 3456028](#)