

Supersymmetry and Schrödinger-type operators with distributional matrix-valued potentials

Jonathan Eckhardt, Fritz Gesztesy, Roger Nichols, and Gerald Teschl¹

Abstract. Building on work on Miura’s transformation by Kappeeler, Perry, Shubin, and Topalov, we develop a detailed spectral theoretic treatment of Schrödinger operators with matrix-valued potentials, with special emphasis on distributional potential coefficients.

Our principal method relies on a supersymmetric (factorization) formalism underlying Miura’s transformation, which intimately connects the triple of operators (D, H_1, H_2) of the form

$$D = \begin{pmatrix} 0 & A^* \\ A & 0 \end{pmatrix} \quad \text{in } L^2(\mathbb{R})^{2m}$$

and

$$H_1 = A^* A, \quad H_2 = AA^* \quad \text{in } L^2(\mathbb{R})^m.$$

Here

$$A = I_m(d/dx) + \phi \quad \text{in } L^2(\mathbb{R})^m,$$

with a matrix-valued coefficient $\phi = \phi^* \in L_{\text{loc}}^1(\mathbb{R})^{m \times m}$, $m \in \mathbb{N}$, thus explicitly permitting distributional potential coefficients V_j in H_j , $j = 1, 2$, where

$$H_j = -I_m \frac{d^2}{dx^2} + V_j(x), \quad V_j(x) = \phi(x)^2 + (-1)^j \phi'(x), \quad j = 1, 2.$$

Upon developing Weyl–Titchmarsh theory for these generalized Schrödinger operators H_j , with (possibly, distributional) matrix-valued potentials V_j , we provide some spectral theoretic applications, including a derivation of the corresponding spectral representations for H_j , $j = 1, 2$. Finally, we derive a local Borg–Marchenko uniqueness theorem for H_j , $j = 1, 2$, by employing the underlying supersymmetric structure and reducing it to the known local Borg–Marchenko uniqueness theorem for D .

Mathematics Subject Classification (2010). Primary 34B20, 34B24, 34L05; Secondary 34B27, 34L10, 34L40.

Keywords. Sturm–Liouville operators, distributional coefficients, Weyl–Titchmarsh theory, supersymmetry.

¹ Research supported by the Austrian Science Fund (FWF) under Grant No. Y330.

Contents

1	Introduction	716
2	Weyl–Titchmarsh matrices for supersymmetric Dirac operators	721
3	Supersymmetry and the connection between the Weyl–Titchmarsh matrices for Dirac and generalized Schrödinger-type operators	727
4	Basic spectral theory for H_j , $j = 1, 2$, and some applications	735
A	Supersymmetric Dirac-type operators in a nutshell	753
	References	757

1. Introduction

This paper was inspired by an investigation concerning “the Miura map on the line” by Kappeler, Perry, Shubin, and Topalov [77] in 2005. In it, the authors consider the well-known Miura map,

$$\phi \longmapsto \phi^2 + \phi', \quad (1.1)$$

which relates appropriate classes of solutions of the Korteweg–de Vries (KdV) and modified Korteweg–de Vries (mKdV) equation (cf., e.g., [39], [40], [43], [44], and the literature cited therein). The Miura map is closely related with factorizations of the KdV Lax operator H , the one-dimensional Schrödinger operator in $L^2(\mathbb{R})$, into a product of two first-order operators of the form

$$H = AA^* = -\frac{d^2}{dx^2} + V(x), \quad V(x) = \phi(x)^2 + \phi'(x), \quad (1.2)$$

where

$$A = \frac{d}{dx} + \phi(x), \quad A^* = -\frac{d}{dx} + \phi(x). \quad (1.3)$$

(While these factorizations are formal at this point, their precise mathematical content is discussed in Sections 2 and 3.) In particular, under the assumption $\phi \in L^2_{\text{loc}}(\mathbb{R})$, ϕ real-valued a.e. on \mathbb{R} , this permits the authors in [77] to discuss real-valued distributional potentials $V \in H^{-1}_{\text{loc}}(\mathbb{R})$, and hence accomplish a remarkable extension of the standard theory of self-adjoint one-dimensional Schrödinger operators in $L^2(\mathbb{R})$ which typically deals with the case of real-valued potentials $V \in L^1_{\text{loc}}(\mathbb{R})$ (resp., $V \in L^2_{\text{loc}}(\mathbb{R})$). This program is carried out in [77] by relying on oscillation theoretic techniques and Hartman’s concept of principal and nonprincipal solutions. In particular, the principal focus of [77] is a detailed investigation of the Miura map (1.1), its range, and its geometry on the real line, with special emphasis on function spaces with low regularity.

As it happens, the Miura map (1.1) is intimately connected with an underlying supersymmetric structure which relates a triple of operators (D, H_1, H_2) of the form,

$$\begin{aligned} D &= \begin{pmatrix} 0 & A^* \\ A & 0 \end{pmatrix} \text{ in } L^2(\mathbb{R})^2, \\ H_1 &= A^*A \text{ and } H_2 = AA^* \text{ in } L^2(\mathbb{R}). \end{aligned} \quad (1.4)$$

Most notably in this context, spectral properties of one of D, H_1, H_2 essentially determine the corresponding spectral properties of the remaining two operators in the triple (D, H_1, H_2) (as described in Appendix A). In particular, since, in accordance with (1.3), $A = (d/dx) + \phi(x)$, $A^* = -(d/dx) + \phi(x)$ deal with the (non-distributional) coefficient $\phi \in L^2_{\text{loc}}(\mathbb{R})$ only, so does the Dirac-type operator D . Consequently, spectral theory (including Weyl–Titchmarsh theory) for the standard Dirac operator D should lead in an effective and streamlined manner to spectral and Weyl–Titchmarsh theory for the generalized Schrödinger operators H_1 and H_2 which may harbor distributional potential coefficients V_j , as

$$H_j = -\frac{d^2}{dx^2} + V_j(x), \quad V_j(x) = \phi(x)^2 + (-1)^j \phi'(x), \quad j = 1, 2. \quad (1.5)$$

Realizing this circle of ideas is precisely what is offered in this paper. Moreover, the fact that exploiting the underlying supersymmetric structure is most natural in this context will become clear as we can effortless incorporate two important generalizations as follows:

- we permit more general coefficients ϕ and hence (distributional) coefficients V_j , $j = 1, 2$, as we only need to assume $\phi \in L^1_{\text{loc}}(\mathbb{R})$;
- we actually consider the matrix-valued case in which ϕ and V_j , $j = 1, 2$, are $m \times m$ self-adjoint matrices a.e. on \mathbb{R} .

Before describing the content of this paper, it is appropriate to comment on the history of singular Sturm–Liouville operator with special emphasis on the papers devoted to distributional potentials.

The particular case of point interactions as special distributional coefficients in Schrödinger operators received enormous attention, too numerous to be mentioned here in detail. Hence, we only refer to the standard monographs by Albeverio, Gesztesy, Høegh-Krohn, and Holden [1] and Albeverio and Kurasov [4], and some of the more recent developments in Albeverio, Kostenko, and Malamud [3], Kostenko and Malamud [86] and [87]. We also mention the case of discontinuous Schrödinger operators originally considered by Hald [54], motivated by the

inverse problem for the torsional modes of the earth. For recent development in this direction we refer to Shahriari, Jodayree Akbarfam, and Teschl [128].

The case of Schrödinger operators with strongly singular and oscillating potentials that should be mentioned in this context goes back to studies by Baeteman and Chadan [7], [8], Combescure [21], Combescure and Ginibre [20], Pearson [112], Rofe-Beketov and Hristov [113] and [114], and a more recent contribution treating distributional potentials by Herczyński [56]. The case of very general (i.e., three-coefficient) singular Sturm–Liouville operators including distributional potentials has been studied by Bennewitz and Everitt [12] in 1983 (see also [34, Section I.2]). They restrict their considerations to compact intervals and focus on the special case of a left-definite setting. An extremely thorough and systematic investigation, including even and odd higher-order operators defined in terms of appropriate quasi-derivatives, and in the general case of matrix-valued coefficients (including distributional potential coefficients in the context of Schrödinger-type operators) was presented by Weidmann [135] in 1987. In fact, the general approach in [12] and [135] draws on earlier discussions of quasi-derivatives in Shin [129]–[131], Naimark [111, Chapter V], and Zettl [136]. Still, it appears that the distributional coefficients treated in [12] did not catch on and subsequent authors referring to this paper mostly focused on the various left and right-definite aspects developed therein. Similarly, it seems likely that the extraordinary generality exerted by Weidmann [135] in his treatment of higher-order differential operators obscured the fact that he already dealt with distributional potential coefficients back in 1987.

However, it was not until 1999 that Savchuk and Shkalikov [121] started a new development for Sturm–Liouville (resp., Schrödinger) operators with distributional potential coefficients in connection with areas such as, self-adjointness proofs, spectral and inverse spectral theory, oscillation properties, spectral properties in the non-self-adjoint context, etc. In addition to the important series of papers by Savchuk and Shkalikov [121]–[127], we mention other groups such as Albeverio, Hrynniv, and Mykytyuk [2], Bak and Shkalikov [9], Ben Amara and Shkalikov [10], Ben Amor and Remling [11], Davies [22], Djakov and Mityagin [24]–[27], Eckhardt and Teschl [32], Frayer, Hrynniv, Mykytyuk, and Perry [35], Gesztesy and Weikard [48], Goriunov and Mikhailets [51], [52], Hrynniv [64], Kappeler and Möhr [76], Kappeler, Perry, Shubin, and Topalov [77], Kappeler and Topalov [78], Hrynniv and Mykytyuk [65]–[72], Hrynniv, Mykytyuk, and Perry [73]–[74], Kato [81], Korotyaev [84], [85], Maz'ya and Shaposhnikova [98, Chapter II], Maz'ya and Verbitsky [99]–[102], Mikhailets and Moliboga [103]–[107], Mirzoev and Safanova [109], Mykytyuk and Trush [110], Sadovnichaya [116], [117]. In particular, the paper by Mirzoev and Safanova [109] is closely related to the

present one as it also employs the use of a quasi-derivative of the type $f^{[1]} = f' + \phi f$ to define a Schrödinger-type operator via a Miura-type transformation and appears to be the only paper known to us since Weidmann's 1987 monograph and the very recent [108] that deals with the matrix-valued case, that is, f is \mathbb{C}^m -valued, ϕ is $\mathbb{C}^{m \times m}$ -valued, $m \in \mathbb{N}$. The prime focus of [109] is the computation of deficiency indices of the underlying minimal operator.

It should be mentioned that some of the attraction in connection with distributional potential coefficients in the Schrödinger operator clearly stems from the low-regularity investigations of solutions of the Korteweg–de Vries (KdV) equation. We mention, for instance, Buckmaster and Koch [14], Grudsky and Rybkin [53], Kappeler and Möhr [76], Kappeler and Topalov [79] and [80], and Rybkin [115].

The case of strongly singular potentials at an endpoint and the associated Weyl–Titchmarsh–Kodaira theory for Schrödinger operators can already be found in the seminal paper by Kodaira [83]. A gap in Kodaira's approach was later circumvented by Kac [75]. The theory did not receive much further attention until it was independently rediscovered and further developed by Gesztesy and Zinchenko [50]. This soon lead to a systematic development of Weyl–Titchmarsh theory for strongly singular potentials and we mention, for instance, Eckhardt [28], Eckhardt and Teschl [31], Fulton [36], Fulton and Langer [37], Fulton, Langer, and Luger [38], Kostenko, Sakhnovich, and Teschl [88], [89], [90], [91], [92], and Kurasov and Luger [94].

We also mention that a different approach to general (i.e., three-coefficient) singular Sturm–Liouville operators (which are not necessarily assumed to be bounded from below) on an arbitrary interval $(a, b) \subseteq \mathbb{R}$, has been developed simultaneously in [29] in the special scalar case $m = 1$. This paper systematically develops Weyl–Titchmarsh theory for differential expressions of the type

$$\tau f = \frac{1}{r}(-(p[f' + \phi f])' + \phi p[f' + \phi f] + qf) \quad (1.6)$$

and hence is very close in spirit to the general discussion provided by Weidmann [135]. Here the coefficients p, q, r, ϕ are real-valued and Lebesgue measurable on (a, b) , with $p \neq 0, r > 0$ a.e. on (a, b) , and $p^{-1}, q, r, \phi \in L^1_{\text{loc}}((a, b); dx)$, and f is supposed to satisfy

$$f \in AC_{\text{loc}}((a, b)), \quad p[f' + \phi f] \in AC_{\text{loc}}((a, b)), \quad (1.7)$$

with $AC_{\text{loc}}((a, b))$ denoting the set of locally absolutely continuous functions on (a, b) . In particular, this study includes distributional coefficients. (The paper [29] does not employ the supersymmetric formalism.) Inverse spectral theory for these operators is treated in [30].

It remains to briefly describe the content of this paper: Section 2 recalls the basics of Weyl–Titchmarsh theory for supersymmetric Dirac-type operators $D = \begin{pmatrix} 0 & A^* \\ A & 0 \end{pmatrix}$ in $L^2(\mathbb{R})^{2m}$, where $A = I_m(d/dx) + \phi$ in $L^2(\mathbb{R})^m$ (cf. (2)–(2.5)), with a matrix-valued coefficient $\phi = \phi^* \in L_{\text{loc}}^1(\mathbb{R})^{m \times m}$, $m \in \mathbb{N}$, following the treatment in [16]. In particular, we review Weyl–Titchmarsh theory for D on the half-line and the full real line, including the $2m \times 2m$ matrix-valued Green’s function of D . In Section 3 we exploit the supersymmetric structure of D and analyze the underlying generalized Schrödinger operators $H_1 = A^*A$ and $H_2 = AA^*$ in $L^2(\mathbb{R})^m$. We derive the Weyl–Titchmarsh solutions for H_j , $j = 1, 2$, given those of D described in Section 2, and describe the precise connection between the half-line Weyl–Titchmarsh matrices of H_j , $j = 1, 2$, and D . In addition, we construct the $m \times m$ matrix-valued Green’s functions of H_j , $j = 1, 2$, and the corresponding analogs belonging to the half-lines $[x_0, \infty)$ and $(-\infty, x_0]$ with a Dirichlet boundary condition at x_0 . In our final Section 4 we provide some spectral theoretic applications of the supersymmetric approach outlined in Section 3 and after deriving the fundamental aspects of Weyl–Titchmarsh theory for the generalized Schrödinger operators H_j , $j = 1, 2$, and a discussion of the corresponding spectral representations, we derive a local Borg–Marchenko uniqueness theorem by utilizing the known analog for the Dirac operator D . Supersymmetric Dirac-type operators and associated commutation methods are briefly summarized in Appendix A.

Next, we briefly summarize some of the notation used in this paper: All $m \times m$ matrices $M \in \mathbb{C}^{m \times m}$ will be considered over the field of complex numbers \mathbb{C} . Moreover, I_m denotes the identity matrix in $\mathbb{C}^{m \times m}$ for $m \in \mathbb{N}$, M^* the adjoint (i.e., complex conjugate transpose), M^\top the transpose of the matrix M , $[M_1, M_2] = M_1 M_2 - M_2 M_1$ denotes the standard commutator of two matrices $M_j \in \mathbb{C}^{m \times m}$, $j = 1, 2$.

$AC([a, b])$ (resp., $AC_{\text{loc}}(c, d))$ denotes the set of (locally) absolutely continuous functions on $[a, b]$ (resp., (c, d)). We also agree that $L^2((a, b))^m$, $m \in \mathbb{N}$, without explicit depiction of a measure always denotes $L^2((a, b); dx)^m$, with dx representing the Lebesgue measure restricted to (a, b) , $-\infty \leq a < b \leq \infty$, in particular,

$$L^2((a, b))^m = \left\{ U : (a, b) \longrightarrow \mathbb{C}^m \mid \int_a^b dx \|U(x)\|_{\mathbb{C}^m}^2 < \infty \right\}, \quad m \in \mathbb{N}. \quad (1.8)$$

The identity operator in $L^2((a, b))^m$ will simply be denoted by I .

For ease of notation we will typically use the short cut $[x_0, \pm\infty)$ to denote the half-lines $[x_0, \infty)$ or $(-\infty, x_0]$ for some $x_0 \in \mathbb{R}$.

Finally, let T be a linear operator mapping (a subspace of) a Hilbert space into another, with $\text{dom}(T)$, $\text{ran}(T)$, and $\ker(T)$ denoting the domain, range, and kernel (i.e., null space) of T . The closure of a closable operator S is denoted by \bar{S} . The spectrum and resolvent set of a closed linear operator in a Hilbert space will be denoted by $\sigma(\cdot)$ and $\rho(\cdot)$, respectively.

Acknowledgments. We are indebted to R. Hrynniv and A. Sakhnovich for very helpful discussions.

2. Weyl–Titchmarsh matrices for supersymmetric Dirac operators

In this section we briefly review the Weyl–Titchmarsh theory for Dirac-type operators D in the special supersymmetric case. In particular, D is constructed as a special case of the theory of singular Hamiltonian systems as pioneered by Hinton and Shaw [59]–[63] (see also [57] and [58]) and applied to Dirac-type operators in [16].

Throughout this section we closely follow the treatment in [16] (simplified to the present supersymmetric Dirac-type operator) and hence are making the following assumptions.

Hypothesis 2.1. Suppose $\phi \in L^1_{\text{loc}}(\mathbb{R})^{m \times m}$, $m \in \mathbb{N}$, and $\phi(\cdot) = \phi(\cdot)^*$ a.e. on \mathbb{R} .

Given Hypothesis 2.1 we introduce the maximally defined operators A and A^+ in $L^2(\mathbb{R})^m$ by

$$(Au)(x) = u'(x) + \phi(x)u(x) \text{ for a.e. } x \in \mathbb{R}, \quad (2.1a)$$

$$u \in \text{dom}(A) = \{v \in L^2(\mathbb{R})^m \mid v \in AC_{\text{loc}}(\mathbb{R}); (v' + \phi v) \in L^2(\mathbb{R})^m\}, \quad (2.1b)$$

and

$$(A^+u)(x) = -u'(x) + \phi(x)u(x) \text{ for a.e. } x \in \mathbb{R}, \quad (2.2a)$$

$$u \in \text{dom}(A^+) = \{v \in L^2(\mathbb{R})^m \mid v \in AC_{\text{loc}}(\mathbb{R}); (v' - \phi v) \in L^2(\mathbb{R})^m\}. \quad (2.2b)$$

In addition, we consider the maximally defined Dirac-type operator D in $L^2(\mathbb{R})^{2m}$ by

$$(DU)(x) = \begin{pmatrix} 0 & A^+ \\ A & 0 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} (x) = \begin{pmatrix} (A^+u_2)(x) \\ (Au_1)(x) \end{pmatrix} \text{ for a.e. } x \in \mathbb{R},$$

$$\begin{aligned} U = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} &\in \text{dom}(D) = \text{dom}(A) \oplus \text{dom}(A^+) \\ &= \left\{ V = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \in L^2(\mathbb{R})^{2m} \mid V \in AC_{\text{loc}}(\mathbb{R})^{2m}; \right. \\ &\quad \left. DV \in L^2(\mathbb{R})^{2m} \right\}. \end{aligned}$$

The basic known result on A , A^+ , and D then reads as follows:

Theorem 2.2 ([16], [59], [61], [62]). *Assume Hypothesis 2.1. Then A and A^+ are closed in $L^2(\mathbb{R})^m$ and*

$$A^* = A^+, \quad (A^+)^* = A. \quad (2.3)$$

In addition, D is self-adjoint in $L^2(\mathbb{R})^{2m}$, that is, D is of the form,

$$D = \begin{pmatrix} 0 & A^* \\ A & 0 \end{pmatrix}. \quad (2.4)$$

Proof. By [16, Lemma 2.15], the differential expression

$$\mathcal{D} = \begin{pmatrix} 0 & -I_m(d/dx) + \phi(x) \\ I_m(d/dx) + \phi(x) & 0 \end{pmatrix} \quad \text{for a.e. } x \in \mathbb{R}, \quad (2.5)$$

is in the limit point case at $\pm\infty$. (For a subsequent and more general result we refer to [96], see also [95] and [97] for such proofs under stronger hypotheses on ϕ). Combining this result with the Weyl–Titchmarsh theory developed for singular Hamiltonian systems by Hinton and Shaw in a series of papers [59], [61], [62], yields self-adjointness of the maximal operator associated to the differential expression \mathcal{D} . By (A.2), A and A^+ are hence necessarily closed, and consequently, adjoint to each other, proving (2.3) and (2.4). \square

Because of the special structure (2.4), D is called a *supersymmetric* Dirac-type operator. For a discussion of its general properties we refer to Appendix A.

Because of (2.3), we identify A^+ and A^* from this point on.

In order to discuss $m \times m$ Weyl–Titchmarsh matrices corresponding to D on the half-lines $(-\infty, x_0]$ and $[x_0, \infty)$, we introduce boundary condition parameters $\alpha = (\alpha_1 \ \alpha_2) \in \mathbb{C}^{m \times 2m}$ satisfying the conditions

$$\alpha\alpha^* = I_m, \quad \alpha J \alpha^* = 0, \quad \text{where } J = \begin{pmatrix} 0 & -I_m \\ I_m & 0 \end{pmatrix}. \quad (2.6)$$

Explicitly, this reads

$$\alpha_1\alpha_1^* + \alpha_2\alpha_2^* = I_m, \quad \alpha_2\alpha_1^* - \alpha_1\alpha_2^* = 0. \quad (2.7)$$

In fact, one also has

$$\alpha_1^* \alpha_1 + \alpha_2^* \alpha_2 = I_m, \quad \alpha_2^* \alpha_1 - \alpha_1^* \alpha_2 = 0, \quad (2.8)$$

as is clear from

$$\begin{pmatrix} \alpha_1 & \alpha_2 \\ -\alpha_2 & \alpha_1 \end{pmatrix} \begin{pmatrix} \alpha_1^* & -\alpha_2^* \\ \alpha_2^* & \alpha_1^* \end{pmatrix} = I_{2m} = \begin{pmatrix} \alpha_1^* & -\alpha_2^* \\ \alpha_2^* & \alpha_1^* \end{pmatrix} \begin{pmatrix} \alpha_1 & \alpha_2 \\ -\alpha_2 & \alpha_1 \end{pmatrix}, \quad (2.9)$$

since any left inverse matrix is also a right inverse, and vice versa. Moreover, from (2.8) one obtains

$$\alpha^* \alpha J + J \alpha^* \alpha = J. \quad (2.10)$$

The particular choice where α equals

$$\alpha_0 = (I_m \ 0) \quad (2.11)$$

will play a fundamental role later on.

Next, denote by $U_{\pm}(\zeta, \cdot, x_0, \alpha)$ the $2m \times m$ matrix-valued Weyl–Titchmarsh solutions associated with $\mathcal{D}U = \zeta U$, $\zeta \in \mathbb{C} \setminus \mathbb{R}$, defined by the property that the m columns of U_{\pm} span the deficiency spaces $N(\zeta, \pm\infty)$, $\zeta \in \mathbb{C} \setminus \mathbb{R}$, given by

$$\begin{aligned} N(\zeta, \pm\infty) &= \{V \in L^2((x_0, \pm\infty))^{2m} \mid V \in AC([x_0, x_0 \pm R])^{2m} \text{ for all } R > 0; \\ &\quad \mathcal{D}V = \zeta V \text{ a.e. on } (x_0, \pm\infty)\}, \end{aligned} \quad (2.12)$$

and normalized such that

$$\begin{aligned} U_{\pm}(\zeta, x, x_0, \alpha) &= \begin{pmatrix} u_{\pm,1}(\zeta, x, x_0, \alpha) \\ u_{\pm,2}(\zeta, x, x_0, \alpha) \end{pmatrix} \\ &= \Psi(\zeta, x, x_0, \alpha) \begin{pmatrix} I_m \\ M_{\pm}^D(\zeta, x_0, \alpha) \end{pmatrix} \\ &= \begin{pmatrix} \vartheta_1(\zeta, x, x_0, \alpha) & \varphi_1(\zeta, x, x_0, \alpha) \\ \vartheta_2(\zeta, x, x_0, \alpha) & \varphi_2(\zeta, x, x_0, \alpha) \end{pmatrix} \begin{pmatrix} I_m \\ M_{\pm}^D(\zeta, x_0, \alpha) \end{pmatrix}. \end{aligned} \quad (2.13)$$

Here $M_{\pm}^D(\zeta, x_0, \alpha)$ represents an $m \times m$ matrix, and $\Psi(\zeta, x, x_0, \alpha)$, $\vartheta_j(\zeta, x, x_0, \alpha)$, and $\varphi_j(\zeta, x, x_0, \alpha)$, $j = 1, 2$, are defined as follows: $\Psi(\zeta, x, x_0, \alpha)$ satisfies $\mathcal{D}\Psi = \zeta\Psi$ a.e. on \mathbb{R} , normalized such that

$$\Psi(\zeta, x_0, x_0, \alpha) = (\alpha^* \ J \alpha^*) = \begin{pmatrix} \alpha_1^* & -\alpha_2^* \\ \alpha_2^* & \alpha_1^* \end{pmatrix}. \quad (2.14)$$

Partitioning $\Psi(\zeta, x, x_0, \alpha)$ as follows,

$$\Psi(\zeta, x, x_0, \alpha) = \begin{pmatrix} \vartheta_1(\zeta, x, x_0, \alpha) & \varphi_1(\zeta, x, x_0, \alpha) \\ \vartheta_2(\zeta, x, x_0, \alpha) & \varphi_2(\zeta, x, x_0, \alpha) \end{pmatrix}, \quad (2.15)$$

defines $\vartheta_j(\zeta, x, x_0, \alpha)$ and $\varphi_j(\zeta, x, x_0, \alpha)$, $j = 1, 2$, as $m \times m$ matrices, entire with respect to $\zeta \in \mathbb{C}$, and normalized according to (2.14).

The matrices $M_{\pm}^D(\zeta, x_0, \alpha)$ represent the half-line Weyl–Titchmarsh matrices associated with the Dirac-type operator D , whose basic properties can be summarized as follows:

Theorem 2.3 ([5], [6], [15], [16], [47], [59], [60], [63], [93]). *Suppose Hypothesis 2.1, let $\zeta \in \mathbb{C} \setminus \mathbb{R}$, $x_0 \in \mathbb{R}$, and denote by $\alpha, \gamma \in \mathbb{C}^{m \times 2m}$ matrices satisfying (2.6). Then the following hold.*

- (i) $\pm M_{\pm}^D(\cdot, x_0, \alpha)$ is an $m \times m$ matrix-valued Nevanlinna–Herglotz function of maximal rank m . In particular,

$$\operatorname{Im}(\pm M_{\pm}^D(\zeta, x_0, \alpha)) \geq 0, \quad \zeta \in \mathbb{C}_+, \quad (2.16)$$

$$M_{\pm}^D(\bar{\zeta}, x_0, \alpha) = M_{\pm}^D(\zeta, x_0, \alpha)^*, \quad (2.17)$$

$$\operatorname{rank}(M_{\pm}^D(\zeta, x_0, \alpha)) = m, \quad (2.18)$$

$$\lim_{\varepsilon \downarrow 0} M_{\pm}^D(v + i\varepsilon, x_0, \alpha) \text{ exists for a.e. } v \in \mathbb{R}, \quad (2.19)$$

$$\begin{aligned} M_{\pm}^D(\zeta, x_0, \alpha) \\ = [-\alpha J \gamma^* + \alpha \gamma^* M_{\pm}^D(\zeta, x_0, \gamma)] [\alpha \gamma^* + \alpha J \gamma^* M_{\pm}^D(\zeta, x_0, \gamma)]^{-1}. \end{aligned} \quad (2.20)$$

Local singularities of $\pm M_{\pm}^D(\cdot, x_0, \alpha)$ and $\mp M_{\pm}^D(\cdot, x_0, \alpha)^{-1}$ are necessarily real and at most of first order in the sense that

$$\mp \lim_{\epsilon \downarrow 0} (i\epsilon M_{\pm}^D(v + i\epsilon, x_0, \alpha)) \geq 0, \quad (2.21a)$$

and

$$\pm \lim_{\epsilon \downarrow 0} (i\epsilon M_{\pm}^D(v + i\epsilon, x_0, \alpha)^{-1}) \geq 0, \quad (2.21b)$$

for $v \in \mathbb{R}$.

- (ii) $\pm M_{\pm}^D(\cdot, x_0, \alpha)$ admits the representation

$$\begin{aligned} \pm M_{\pm}^D(\zeta, x_0, \alpha) \\ = F_{\pm}(x_0, \alpha) + \int_{\mathbb{R}} d\Omega_{\pm}^D(v, x_0, \alpha) [(v - \zeta)^{-1} - v(1 + v^2)^{-1}], \end{aligned} \quad (2.22)$$

where

$$F_{\pm}(x_0, \alpha) = F_{\pm}(x_0, \alpha)^*, \quad \int_{\mathbb{R}} \|d\Omega_{\pm}^D(v, x_0, \alpha)\|_{\mathbb{C}^{m \times m}} (1 + v^2)^{-1} < \infty. \quad (2.23)$$

Moreover,

$$\Omega_{\pm}^D((\mu, v], x_0, \alpha) = \lim_{\delta \downarrow 0} \lim_{\varepsilon \downarrow 0} \frac{1}{\pi} \int_{\mu+\delta}^{v+\delta} dv' \operatorname{Im}(\pm M_{\pm}^D(v'+i\varepsilon, x_0, \alpha)). \quad (2.24)$$

(iii) $\operatorname{Im}(M_{\pm}^D(\cdot, x_0, \alpha))$ satisfies

$$\begin{aligned} & \operatorname{Im}(M_{\pm}^D(\zeta, x_0, \alpha)) \\ &= \operatorname{Im}(\zeta) \int_{x_0}^{\pm\infty} dx U_{\pm}(\zeta, x, x_0, \alpha)^* U_{\pm}(\zeta, x, x_0, \alpha) \\ &= \operatorname{Im}(\zeta) \int_{x_0}^{\pm\infty} dx [u_{\pm,1}(\zeta, x, x_0, \alpha)^* u_{\pm,1}(\zeta, x, x_0, \alpha) \\ & \quad + u_{\pm,2}(\zeta, x, x_0, \alpha)^* u_{\pm,2}(\zeta, x, x_0, \alpha)]. \end{aligned} \quad (2.25)$$

For completeness we also recall that the $2m \times 2m$ Green's matrix (i.e., the integral kernel of the resolvent) of D is given in terms of U_{\pm} and M_{\pm} by

$$\begin{aligned} G^D(\zeta, x, x') &= (D - \zeta I)^{-1}(x, x') \\ &= U_{\mp}(\zeta, x, x_0, \alpha) [M_{-}^D(\zeta, x_0, \alpha) - M_{+}^D(\zeta, x_0, \alpha)]^{-1} U_{\pm}(\bar{\zeta}, x', x_0, \alpha)^*, \end{aligned} \quad (2.26)$$

with $x \leq x'$, $x, x' \in \mathbb{R}$, $\zeta \in \mathbb{C} \setminus \mathbb{R}$. Of course, $G^D(\zeta, x, x')$ is independent of the choice of reference point $x_0 \in \mathbb{R}$, and independent of the boundary condition parameter α satisfying (2.6) used in $M_{\pm}^D(\zeta, x_0, \alpha)$ and $U_{\pm}(\zeta, \cdot, x_0, \alpha)$. One also notes that (2.26) extends as usual to all $\zeta \in \rho(D)$. In the particular case $\alpha_0 = (I_m \ 0)$ one obtains

$$U_{\pm}(\zeta, x_0, x_0, \alpha_0) = \begin{pmatrix} u_{\pm,1}(\zeta, x_0, x_0, \alpha_0) \\ u_{\pm,2}(\zeta, x_0, x_0, \alpha_0) \end{pmatrix} = \begin{pmatrix} I_m \\ M_{\pm}^D(\zeta, x_0, \alpha_0) \end{pmatrix}. \quad (2.27)$$

The self-adjoint half-line Dirac operators $D_{\pm}(\alpha)$ in $L^2([x_0, \pm\infty))^{2m}$ associated with a self-adjoint boundary condition at x_0 indexed by $\alpha \in \mathbb{C}^{m \times 2m}$ satisfying (2.6), are of the form

$$(D_{\pm}(\alpha)U)(x) = (\mathcal{D}U)(x) \quad \text{for a.e. } x \in [x_0, \pm\infty), \quad (2.28a)$$

$$\begin{aligned} U &\in \operatorname{dom}(D_{\pm}(\alpha)) \\ &= \{V \in L^2([x_0, \pm\infty))^{2m} \mid V \in AC([x_0, x_0 \pm R])^{2m} \\ & \quad \text{for all } R > 0; \alpha\phi(x_0) = 0; \\ & \quad \mathcal{D}V \in L^2([x_0, \pm\infty))^{2m}\}. \end{aligned} \quad (2.28b)$$

The $m \times m$ matrix-valued spectral function of $D_{\pm}(\alpha)$ then generates the measure $\Omega_{\pm}^D(\cdot, x_0, \alpha)$ in (2.22).

We conclude this section with a brief description of the full-line $2m \times 2m$ Weyl–Titchmarsh matrix $\mathbf{M}^D(\zeta, x_0, \alpha)$ associated with D as described in [59]–[63]:

$$\mathbf{M}^D(\zeta, x_0, \alpha) = (\mathbf{M}_{j,j'}^D(\zeta, x_0, \alpha))_{j,j'=1,2}, \quad \zeta \in \mathbb{C} \setminus \mathbb{R}, \quad (2.29a)$$

$$\mathbf{M}_{0,0}^D(\zeta, x_0, \alpha) = [M_-^D(\zeta, x_0, \alpha) - M_+^D(\zeta, x_0, \alpha)]^{-1}, \quad (2.29b)$$

$$\begin{aligned} \mathbf{M}_{0,1}^D(\zeta, x_0, \alpha) &= 2^{-1}[M_-^D(\zeta, x_0, \alpha) - M_+^D(\zeta, x_0, \alpha)]^{-1} \\ &\quad \times [M_-^D(\zeta, x_0, \alpha) + M_+^D(\zeta, x_0, \alpha)], \end{aligned} \quad (2.29c)$$

$$\begin{aligned} \mathbf{M}_{1,0}^D(\zeta, x_0, \alpha) &= 2^{-1}[M_-^D(\zeta, x_0, \alpha) + M_+^D(\zeta, x_0, \alpha)] \\ &\quad \times [M_-^D(\zeta, x_0, \alpha) - M_+^D(\zeta, x_0, \alpha)]^{-1}, \end{aligned} \quad (2.29d)$$

$$\mathbf{M}_{1,1}^D(\zeta, x_0, \alpha) = M_{\pm}^D(\zeta, x_0, \alpha)[M_-^D(\zeta, x_0, \alpha) - M_+^D(\zeta, x_0, \alpha)]^{-1} M_{\mp}^D(\zeta, x_0, \alpha). \quad (2.29e)$$

The basic results on $\mathbf{M}^D(\cdot, x_0, \alpha)$ then read as follows.

Theorem 2.4 ([47], [59], [60], [63], [93]). *Assume Hypothesis 2.1 and suppose that $\zeta \in \mathbb{C} \setminus \mathbb{R}$, $x_0 \in \mathbb{R}$, and that $\alpha \in \mathbb{C}^{m \times 2m}$ satisfies (2.6). Then the following hold:*

- (i) $\mathbf{M}^D(\cdot, x_0, \alpha)$ is a matrix-valued Nevanlinna–Herglotz function of maximal rank $2m$ with representation

$$\mathbf{M}^D(\zeta, x_0, \alpha) = \mathbf{F}(x_0, \alpha) + \int_{\mathbb{R}} d\Omega^D(v, x_0, \alpha) [(v - \zeta)^{-1} - v(1 + v^2)^{-1}], \quad (2.30)$$

where

$$\mathbf{F}(x_0, \alpha) = \mathbf{F}(x_0, \alpha)^*, \quad \int_{\mathbb{R}} \|d\Omega^D(v, x_0, \alpha)\|_{\mathbb{C}^{2m \times 2m}} (1 + v^2)^{-1} < \infty. \quad (2.31)$$

Moreover,

$$\Omega^D((\mu, \nu], x_0, \alpha) = \lim_{\delta \downarrow 0} \lim_{\varepsilon \downarrow 0} \frac{1}{\pi} \int_{\mu+\delta}^{\nu+\delta} dv' \operatorname{Im}(\mathbf{M}^D(v' + i\varepsilon, x_0, \alpha)). \quad (2.32)$$

- (ii) $\zeta \in \rho(D)$ if and only if $\mathbf{M}^D(\zeta, x_0, \alpha)$ is holomorphic near ζ .

Finally, observe that supersymmetry implies various symmetries for the associated quantities.

Lemma 2.5. *Assume Hypothesis 2.1. The operators $D_{\pm}(\alpha_0)$ are supersymmetric and satisfy $\mathfrak{S}_3 D_{\pm}(\alpha_0) \mathfrak{S}_3 = -D_{\pm}(\alpha_0)$, where*

$$\mathfrak{S}_3 = \begin{pmatrix} I_m & 0 \\ 0 & -I_m \end{pmatrix}. \quad (2.33)$$

Moreover,

$$\mathfrak{S}_3 (\varphi_1(\zeta, x, x_0, \alpha_0) \varphi_2(\zeta, x, x_0, \alpha_0))^{\top} = -(\varphi_1(-\zeta, x, x_0, \alpha_0) \varphi_2(-\zeta, x, x_0, \alpha_0))^{\top},$$

$$\mathfrak{S}_3 (\vartheta_1(\zeta, x, x_0, \alpha_0) \vartheta_2(\zeta, x, x_0, \alpha_0))^{\top} = (\vartheta_1(-\zeta, x, x_0, \alpha_0) \vartheta_2(-\zeta, x, x_0, \alpha_0))^{\top},$$

$$\mathfrak{S}_3 U_{\pm}(\zeta, x, x_0, \alpha_0) = U_{\pm}(-\zeta, x, x_0, \alpha_0),$$

$$M_{\pm}^D(\zeta, x_0, \alpha_0) = -M_{\pm}^D(-\zeta, x_0, \alpha_0),$$

$$d\Omega_{\pm}^D(\nu, x_0, \alpha_0) = d\Omega_{\pm}^D(-\nu, x_0, \alpha_0).$$

Similarly, D is supersymmetric, $\mathfrak{S}_3 D \mathfrak{S}_3 = -D$, and

$$M^D(\zeta, x_0, \alpha_0) = -\mathfrak{S}_3 M^D(-\zeta, x_0, \alpha_0) \mathfrak{S}_3,$$

$$d\Omega^D(\nu, x_0, \alpha_0) = \mathfrak{S}_3 d\Omega^D(-\nu, x_0, \alpha_0) \mathfrak{S}_3.$$

Proof. Clearly $D_{\pm}(\alpha_0)$ are of the form (A.2) in this case and hence they are supersymmetric. Moreover, the symmetries for the solutions follow since both sides satisfy the same differential equation and the same initial conditions, respectively normalizations. The claims for D are immediate from the ones for $D_{\pm}(\alpha_0)$. \square

3. Supersymmetry and the connection between the Weyl–Titchmarsh matrices for Dirac and generalized Schrödinger-type operators

In our principal section we provide the connection with the matrix-valued Weyl–Titchmarsh functions of the supersymmetric Dirac-type operator D described in Section 2 and two naturally associated generalized Schrödinger-type operators H_j , $j = 1, 2$, given by

$$D^2 = \begin{pmatrix} A^* A & 0 \\ 0 & AA^* \end{pmatrix} = H_1 \oplus H_2 \text{ in } L^2(\mathbb{R})^{2m} \simeq L^2(\mathbb{R})^m \oplus L^2(\mathbb{R})^m, \quad (3.1)$$

in particular, we denote

$$H_1 = A^* A, \quad H_2 = AA^*, \quad (3.2)$$

with A and A^* given by (2.1) and (2.2), respectively (cf. Theorem 2.2).

While D contains the locally integrable $m \times m$ matrix-valued coefficient ϕ , the associated generalized Schrödinger operators H_j , $j = 1, 2$, will exhibit distributional potentials and hence are outside the standard Weyl–Titchmarsh theory for Sturm–Liouville operators with locally integrable $m \times m$ matrix-valued potentials. Our supersymmetric approach will enable us to make the transition from the usual L^1_{loc} -potentials in Schrödinger operators to distributional H^{-1}_{loc} -potentials (and more general situations) in an effortless manner, thereby underscoring the power of these supersymmetric arguments.

To describe H_j , $j = 1, 2$, in $L^2(\mathbb{R})$ in detail, we first introduce the following two kinds of quasi-derivatives,

$$\begin{aligned} u^{[1,1]}(x) &= (Au)(x) \\ &= u'(x) + \phi(x)u(x) \quad \text{for a.e. } x \in \mathbb{R}, u \in \text{dom}(A), \end{aligned} \quad (3.3)$$

$$\begin{aligned} v^{[1,2]}(x) &= -(A^*v)(x) \\ &= v'(x) - \phi(x)v(x) \quad \text{for a.e. } x \in \mathbb{R}, v \in \text{dom}(A^*). \end{aligned} \quad (3.4)$$

Thus, one infers,

$$\begin{aligned} (H_1 u)(x) &= (A^* A u)(x) \\ &= (\tau_1 u)(x) \\ &= -(u^{[1,1]})'(x) + \phi(x)u^{[1,1]}(x) \quad \text{for a.e. } x \in \mathbb{R}, \end{aligned} \quad (3.5a)$$

$$\begin{aligned} u \in \text{dom}(H_1) &= \{v \in L^2(\mathbb{R})^m \mid v, v^{[1,1]} \in AC_{\text{loc}}(\mathbb{R})^m; \\ &\quad [(v^{[1,1]})' + \phi v^{[1,1]}] \in L^2(\mathbb{R})^m\}, \end{aligned} \quad (3.5b)$$

and

$$\begin{aligned} (H_2 u)(x) &= (AA^* u)(x) \\ &= (\tau_2 u)(x) \\ &= -(u^{[1,2]})'(x) - \phi(x)u^{[1,2]}(x) \quad \text{for a.e. } x \in \mathbb{R}, \end{aligned} \quad (3.6a)$$

$$\begin{aligned} u \in \text{dom}(H_2) &= \{v \in L^2(\mathbb{R})^m \mid v, v^{[1,2]} \in AC_{\text{loc}}(\mathbb{R})^m; \\ &\quad [(v^{[1,2]})' + \phi v^{[1,2]}] \in L^2(\mathbb{R})^m\}. \end{aligned} \quad (3.6b)$$

Formally, τ_j , $j = 1, 2$, are of the form

$$\tau_j = -I_m \frac{d^2}{dx^2} + V_j(x), \quad V_j(x) = \phi(x)^2 + (-1)^j \phi'(x), \quad j = 1, 2, \quad (3.7)$$

but one notices that, in general, neither ϕ^2 is locally integrable (unless one makes the stronger assumption $\phi \in L^2_{\text{loc}}(\mathbb{R})^{m \times m}$), nor is ϕ' a function (unless one assumes in addition that $\phi \in AC_{\text{loc}}(\mathbb{R})^{m \times m}$).

By inspection, the second-order initial value problems,

$$\begin{aligned} ((\tau_j - z)f)(x) &= g(x) \quad \text{for a.e. } x \in \mathbb{R}, \\ f, f^{[1,j]} &\in AC_{\text{loc}}(\mathbb{R})^m, \\ g &\in L^1_{\text{loc}}(\mathbb{R})^m, \end{aligned} \quad (3.8a)$$

$$f(x_0) = c_0, \quad f^{[1,j]}(x_0) = d_0, \quad j = 1, 2, \quad (3.8b)$$

for some $x_0 \in \mathbb{R}$, $c_0, d_0 \in \mathbb{C}$, are equivalent to the first-order initial value problems

$$\begin{aligned} &\left(\begin{matrix} f(x) \\ f^{[1,j]}(x) \end{matrix} \right)' \\ &= \begin{pmatrix} (-1)^j \phi(x) & 1 \\ -z & (-1)^{j+1} \phi(x) \end{pmatrix} \left(\begin{matrix} f(x) \\ f^{[1,j]}(x) \end{matrix} \right) - \left(\begin{matrix} 0 \\ g(x) \end{matrix} \right) \quad \text{for a.e. } x \in \mathbb{R}, \end{aligned} \quad (3.9a)$$

$$\left(\begin{matrix} f(x_0) \\ f^{[1,j]}(x_0) \end{matrix} \right) = \left(\begin{matrix} c_0 \\ d_0 \end{matrix} \right), \quad j = 1, 2, \quad (3.9b)$$

respectively. Since by Hypothesis 2.1, $\phi \in L^1_{\text{loc}}(\mathbb{R})^{m \times m}$, the initial value problems in (3.9), and hence those in (3.8), are uniquely solvable by [111, Theorem 16.1] (see also [33, Theorem 10.1] and [111, Theorem 16.2]).

Next, suppose that for some $1 \leq p \leq m$, $U = (u_1 \ u_2)^\top$ is a distributional $2m \times p$ solution of $DU = \xi U$, that is,

$$u_j \in AC_{\text{loc}}(\mathbb{R})^{m \times p}, \quad j = 1, 2, \quad (3.10a)$$

$$u_1^{[1,1]} = Au_1 \in L^1_{\text{loc}}(\mathbb{R})^{m \times p}, \quad (3.10b)$$

$$u_2^{[1,2]} = -A^* u_2 \in L^1_{\text{loc}}(\mathbb{R})^{m \times p}. \quad (3.10c)$$

Then, if $\zeta \neq 0$, the supersymmetric structure of D in (2.4) actually implies that also

$$u_1^{[1,1]} = Au_1 = \zeta u_2 \in AC_{\text{loc}}(\mathbb{R})^{m \times p}, \quad (3.11)$$

$$u_2^{[1,2]} = -A^* u_2 = -\zeta u_1 \in AC_{\text{loc}}(\mathbb{R})^{m \times p}, \quad (3.12)$$

and hence that u_j are actually distributional $m \times p$ solutions of $H_j u = \zeta^2 u$, $j = 1, 2$, that is,

$$u_j, u_j^{[1,j]} \in AC_{\text{loc}}(\mathbb{R})^{m \times p}, \quad (u_j^{[1,j]})' \in L_{\text{loc}}^1(\mathbb{R})^{m \times p}, \quad (3.13a)$$

$$\tau_j u_j = -(u_j^{[1,j]})' + (-1)^{j+1} \phi u_j^{[1,j]} = \zeta^2 u_j, \quad j = 1, 2. \quad (3.13b)$$

Thus, applying the L^2 -property (2.25) and equations (3.10)–(3.13) to the Weyl–Titchmarsh solutions $U_{\pm}(\zeta, \cdot, x_0, \alpha)$ associated with the Dirac-type operator D , then shows that $u_{\pm,j}(\zeta, \cdot, x_0, \alpha)$ are Weyl–Titchmarsh solutions associated with H_j , $j = 1, 2$, replacing the complex energy parameter ζ by $z = \zeta^2$. Moreover, introducing the following fundamental system $s_j(z, \cdot, x_0)$, $c_j(z, \cdot, x_0)$, $j = 1, 2$, of $m \times m$ matrix solutions of $\tau_j u = zu$, $z \in \mathbb{C}$, $j = 1, 2$, normalized for arbitrary $z \in \mathbb{C}$ by

$$s_j(z, x_0, x_0) = 0, \quad s_j^{[1,j]}(z, x_0, x_0) = I_m, \quad (3.14)$$

$$c_j(z, x_0, x_0) = I_m, \quad c_j^{[1,j]}(z, x_0, x_0) = 0, \quad j = 1, 2, \quad (3.15)$$

one observes as usual that for fixed $x, x_0 \in \mathbb{R}$, $s_j(\cdot, x, x_0)$, $c_j(\cdot, x, x_0)$ are entire. The connection with the solutions φ_j and ϑ_j , $j = 1, 2$, of D is given by

$$s_1(z, x, x_0) = \zeta^{-1} \varphi_1(\zeta, x, x_0, \alpha_0), \quad c_1(z, x, x_0) = \vartheta_1(\zeta, x, x_0, \alpha_0), \quad (3.16)$$

$$s_2(z, x, x_0) = \zeta^{-1} \vartheta_2(\zeta, x, x_0, \alpha_0), \quad c_2(z, x, x_0) = \varphi_2(\zeta, x, x_0, \alpha_0), \quad z = \zeta^2. \quad (3.17)$$

In addition, introducing the Weyl–Titchmarsh solutions $\psi_{\pm,j}(z, \cdot, x_0)$ for H_j , $j = 1, 2$, via

$$\psi_{\pm,1}(z, \cdot, x_0) = u_{\pm,1}(\zeta, \cdot, x_0, \alpha_0), \quad (3.18)$$

$$\psi_{\pm,2}(z, \cdot, x_0) = u_{\pm,2}(\zeta, \cdot, x_0, \alpha_0) M_{\pm}^D(\zeta, x_0, \alpha_0)^{-1}, \quad (3.19)$$

where $z = \xi^2$ and $\xi \in \mathbb{C} \setminus \mathbb{R}$ (the right-hand sides being independent of the choice of branch for ξ), and the generalized Dirichlet-type $m \times m$ matrix-valued Weyl–Titchmarsh functions $\widehat{M}_{\pm,0,j}(\cdot, x_0)$ of H_j ,

$$\widehat{M}_{\pm,0,1}(z, x_0) = \xi M_{\pm}^D(\xi, x_0, \alpha_0), \quad (3.20)$$

$$\widehat{M}_{\pm,0,2}(z, x_0) = -\xi M_{\pm}^D(\xi, x_0, \alpha_0)^{-1}, \quad (3.21)$$

where $z = \xi^2$ and $\xi \in \mathbb{C} \setminus \mathbb{R}$, one infers from (2.27) that

$$\psi_{\pm,j}(z, \cdot, x_0) = c_j(z, \cdot, x_0) + s_j(z, \cdot, x_0) \widehat{M}_{\pm,0,j}(z, x_0), \quad j = 1, 2. \quad (3.22)$$

where $z \in \mathbb{C} \setminus [0, \infty)$. Indeed, (3.22) follows from combining equations (2.27), (3.11), and (3.12) (for $p = m$), which in turn imply

$$\psi_{\pm,j}(z, x_0, x_0) = I_m, \quad (3.23)$$

$$\psi_{\pm,j}^{[1,j]}(z, x_0, x_0) = \widehat{M}_{\pm,0,j}(z, x_0), \quad j = 1, 2 \quad (3.24)$$

and the unique solvability of the initial value problems in (3.8). We summarize this discussion in the following result.

Theorem 3.1. *Assume Hypothesis 2.1 and let $\alpha_0 = (I_m \ 0)$. Suppose that the corresponding Weyl–Titchmarsh solutions of D are denoted by*

$$U_{\pm}(\xi, \cdot, x_0, \alpha_0) = (u_{\pm,1}(\xi, \cdot, x_0, \alpha_0) \ u_{\pm,2}(\xi, \cdot, x_0, \alpha_0))^{\top}$$

and the corresponding $m \times m$ matrix-valued half-line Weyl–Titchmarsh function of D is given by $M_{\pm}^D(\cdot, x_0, \alpha_0)$. Then the $m \times m$ matrix-valued Weyl–Titchmarsh solutions of H_j , denoted by $\psi_{\pm,j}(z, \cdot, x_0)$, $j = 1, 2$, are given by (3.18) and (3.19), and the $m \times m$ matrix-valued generalized Dirichlet-type Weyl–Titchmarsh functions $\widehat{M}_{\pm,0,j}(\cdot, x_0)$ of H_j , $j = 1, 2$, are given by (3.20) and (3.21). In particular,

$$\widehat{M}_{\pm,0,1}(z, x_0) = \xi M_{\pm}^D(\xi, x_0, \alpha_0) = -z \widehat{M}_{\pm,0,2}(z, x_0)^{-1}, \quad (3.25)$$

with $z = \xi^2$, and $\xi \in \mathbb{C} \setminus \mathbb{R}$.

A version of the equality $\widehat{M}_{\pm,0,1}(z, x_0) = -z \widehat{M}_{\pm,0,2}(z, x_0)^{-1}$, in the special scalar case $m = 1$, and under the stronger hypothesis $\phi \in AC_{loc}(\mathbb{R})$, first appeared in [42, eq. (5.71)], and was quoted again in [46, eq. (A.25)].

The subscript “0” in $\widehat{M}_{\pm,0,j}(z, x_0)$, $j = 1, 2$, indicates that these generalized Weyl–Titchmarsh matrices correspond to a Dirichlet boundary condition at the reference point x_0 in the corresponding generalized half-line Schrödinger operators $H_{\pm,0,j}$, $j = 1, 2$, in $L^2([x_0, \pm\infty))^m$ defined by

$$\begin{aligned} (H_{\pm,0,j}u)(x) &= (\tau_j u)(x) \\ &= -(u^{[1,j]})'(x) + (-1)^{j+1}\phi(x)u^{[1,j]}(x) \quad \text{for a.e. } x \in [x_0, \pm\infty), \end{aligned} \tag{3.26a}$$

$$\begin{aligned} u \in \text{dom}(H_{\pm,0,j}) &= \{v \in L^2([x_0, \pm\infty))^m \mid v, v^{[1,j]} \in AC([x_0, x_0 \pm R])^m \text{ for all } R > 0; \\ &\quad v(x_0) = 0; \\ &\quad [(v^{[1,j]})' + (-1)^j \phi v^{[1,j]}] \in L^2([x_0, \pm\infty))^m\}, \\ &\quad j = 1, 2. \end{aligned} \tag{3.26b}$$

The corresponding Green’s function of $H_{\pm,0,j}$ is then of the familiar form

$$\begin{aligned} G_{+,0,j}(z, x, x') &= (H_{+,0,j} - zI)^{-1}(x, x') \\ &= \begin{cases} s_j(z, x, x_0)\psi_{+,j}(\bar{z}, x', x_0)^*, & x \leqslant x', \\ \psi_{+,j}(z, x, x_0)s_j(\bar{z}, x', x_0)^*, & x' \leqslant x, \end{cases} \\ &\quad x, x' \in [x_0, \infty), z \in \mathbb{C} \setminus [0, \infty), j = 1, 2, \end{aligned} \tag{3.27}$$

and

$$\begin{aligned} G_{-,0,j}(z, x, x') &= (H_{-,0,j} - zI)^{-1}(x, x') \\ &= - \begin{cases} s_j(z, x, x_0)\psi_{-,j}(\bar{z}, x', x_0)^*, & x' \leqslant x, \\ \psi_{-,j}(z, x, x_0)s_j(\bar{z}, x', x_0)^*, & x \leqslant x', \end{cases} \\ &\quad x, x' \in (-\infty, x_0], z \in \mathbb{C} \setminus [0, \infty), j = 1, 2. \end{aligned} \tag{3.28}$$

Similarly, the diagonal terms in (A.33) together with equations (2.17), (2.26) and (3.18)–(3.22) yield the Green’s function for H_j ,

$$\begin{aligned} G_j(z, x, x') &= (H_j - zI)^{-1}(x, x') \\ &= \psi_{\mp, j}(z, x, x_0)[\widehat{M}_{-, 0, j}(z, x_0) - \widehat{M}_{+, 0, j}(z, x_0)]^{-1}\psi_{\pm, j}(\bar{z}, x', x_0)^*, \\ &\quad x \leqslant x', \quad x, x' \in \mathbb{R}, \quad z \in \mathbb{C} \setminus [0, \infty), \quad j = 1, 2. \end{aligned} \tag{3.29}$$

One can show that

$$\begin{aligned} &\widehat{M}_{-, 0, j}(z, x_0) - \widehat{M}_{+, 0, j}(z, x_0) \\ &= W(\psi_{+, j}(\bar{z}, \cdot, x_0)^*, \psi_{-, j}(z, \cdot, x_0)) \\ &= -W(\psi_{-, j}(\bar{z}, \cdot, x_0)^*, \psi_{+, j}(z, \cdot, x_0)), \quad j = 1, 2, \end{aligned} \tag{3.30}$$

where $W(\cdot, \cdot)$, $j = 1, 2$, denote the Wronskians of matrix-valued functions $F, G \in AC_{loc}(\mathbb{R})^{m \times m}$ defined by

$$\begin{aligned} W(F, G)(x) &= F(x)G'(x) - F'(x)G(x) \\ &= F(x)G^{[1, j]}(x) - F^{[1, j]}(x)G(x) + (-1)^j[F(x), \phi(x)]G(x), \\ &\quad \text{for a.e. } x \in \mathbb{R}, \quad j = 1, 2. \end{aligned} \tag{3.31}$$

Of course, (3.18)–(3.25), (3.27)–(3.29), extend as usual to all $z \in \rho(H_{\pm, 0, j})$, respectively, all $z \in \rho(H_j)$, $j = 1, 2$.

We note in passing that

$W(v_j(\bar{z}, \cdot)^*, u_j(z, \cdot))(x)$ is x -independent whenever

$$\begin{aligned} &-[u'_j(z, x) + (-1)^{j+1}\phi(x)u_j(z, x)]' + (-1)^{j+1}\phi(x)[u'_j(z, x) \\ &+ (-1)^{j+1}\phi(x)u_j(z, x)] = zu_j(z, x), \end{aligned} \tag{3.32a}$$

$$\begin{aligned} &-[v'_j(\bar{z}, x)^* + (-1)^{j+1}v_j(\bar{z}, x)^*\phi(x)]' + (-1)^{j+1}[v'_j(\bar{z}, x)^* \\ &+ (-1)^{j+1}v_j(\bar{z}, x)^*\phi(x)]\phi(x) = zv_j(\bar{z}, x)^*, \end{aligned} \tag{3.32b}$$

$$j = 1, 2.$$

This circle of ideas will be further explored in Section 4.

Remark 3.2. In the particular case where $\phi \in AC_{loc}(\mathbb{R})^{m \times m}$ and hence,

$$V_j = [\phi^2 + (-1)^j \phi'] \in L^1_{loc}(\mathbb{R})^{m \times m}, \quad j = 1, 2, \quad (3.33)$$

the relation between the generalized Weyl–Titchmarsh matrices $\hat{M}_{\pm,0,j}(\cdot, x_0)$, defined in (3.20), (3.21), and the standard (Dirichlet-type) Weyl–Titchmarsh matrices $M_{\pm,0,j}(\cdot, x_0)$ is especially simple and reads

$$\hat{M}_{\pm,0,j}(\cdot, x_0) = M_{\pm,0,j}(\cdot, x_0) + (-1)^{j+1} \phi(x_0), \quad j = 1, 2. \quad (3.34)$$

In particular, since $\phi(x_0)$ is z -independent and self-adjoint, the function theoretic (and hence spectral theoretic) content of $\hat{M}_{\pm,0,j}(\cdot, x_0)$ and $M_{\pm,0,j}(\cdot, x_0)$ coincides in this special case as they possess identical matrix measures in their respective Nevanlinna–Herglotz representations. (These matrix measures being generated by the half-line $m \times m$ matrix-valued spectral functions of $H_{\pm,0,j}$.)

Remark 3.3. In the particular scalar case $m = 1$, and under the stronger assumption $\phi \in L^2_{loc}(\mathbb{R})$, the operators H_j , $j = 1, 2$, and especially, the associated Miura transformation,

$$\phi \longmapsto \phi^2 - \phi' \quad (3.35)$$

(i.e., the relation between ϕ and V_1), was studied in great detail in [77]. However, the authors did not directly rely on D and its supersymmetric structure, but instead based their investigations on an oscillation theoretic approach using Hartman’s notion of (non)principal solutions. Subsequently, the authors of [35], [73], and [74] used a Zakharov–Shabat (ZS), or Ablowitz–Kaup–Newell–Segur (AKNS) Dirac-type expression \tilde{D} in connection with their investigation of Miura transformations and inverse scattering theory. Explicitly, \tilde{D} represents a self-adjoint $L^2(\mathbb{R})^2$ -realization associated with the differential expression $\tilde{\mathcal{D}}$ of the form

$$\tilde{\mathcal{D}} = i \begin{pmatrix} -(d/dx) & -\phi(x) \\ \phi(x) & (d/dx) \end{pmatrix} \quad \text{for a.e. } x \in \mathbb{R}. \quad (3.36)$$

To make the connection with the supersymmetric formalism presented in this paper, we introduce the unitary 2×2 matrices

$$\Upsilon = \frac{1}{2} \begin{pmatrix} 1-i & 1-i \\ 1+i & -1-i \end{pmatrix}, \quad \Upsilon^* = \frac{1}{2} \begin{pmatrix} 1+i & 1-i \\ 1+i & -1+i \end{pmatrix} = \Upsilon^{-1}, \quad (3.37)$$

and observe that

$$\Upsilon \tilde{\mathcal{D}} \Upsilon^{-1} = \mathcal{D}, \quad (3.38)$$

with \mathcal{D} in (2.5) the differential expression underlying the supersymmetric Dirac-type operator D .

It is the use of supersymmetry of D in connection with the (standard) assumption of local integrability of the coefficient ϕ that instantly leads to Weyl–Titchmarsh solutions $U_{\pm}(\zeta, x, x_0, \alpha)$ and Weyl–Titchmarsh matrices $M_{\pm}^D(\zeta, x_0, \alpha)$ of D and hence effortlessly via equations (3.18)–(3.22) to those of the generalized Schrödinger-type operators $H_{\pm,0,j}$, H_j , $j = 1, 2$. In particular, it immediately leads to the Green’s functions (3.27)–(3.29) of $H_{\pm,0,j}$ and of H_j , $j = 1, 2$, respectively, and permits the more general hypothesis $\phi \in L_{\text{loc}}^1(\mathbb{R})$ rather than $\phi \in L_{\text{loc}}^2(\mathbb{R})$. In addition, it permits an effortless discussion of the matrix-valued case (the latter cannot easily be obtained via oscillation theoretic methods, cf. the comments preceding [17, Hypothesis 3.6] in this context).

Remark 3.4. The supersymmetric formalism employed in this section relies on nonnegativity of $H_1 = A^*A$ and $H_2 = AA^*$ and so imposes a restriction on the distributional potential coefficients V_j (formally, of the Miura-type $V_j(x) = \phi(x)^2 + (-1)^j \phi'(x)$), $j = 1, 2$. In practice, however, this restriction amounts to dealing with Schrödinger operators bounded from below as adding a sufficiently large positive constant to a given potential (if necessary) will render H_1 and H_2 nonnegative. (Subsequently, this additional constant can be removed.)

In this context we also notice that it is possible to generalize the operator D in (2.4) to the form

$$D_m = \begin{pmatrix} m & A^* \\ A & -m \end{pmatrix}, \quad m \in \mathbb{R}, \quad (3.39)$$

to the effect that then

$$D_m^2 = \begin{pmatrix} A^*A + m^2 I & 0 \\ 0 & AA^* + m^2 I \end{pmatrix} = (H_1 + m^2 I) \oplus (H_2 + m^2 I). \quad (3.40)$$

Finally, we note that these results on generalized Schrödinger operators with distributional potentials extend to general (three-coefficient) Sturm–Liouville operators, but we refrain from further details at this point.

4. Basic spectral theory for H_j , $j = 1, 2$, and some applications

In our final section we provide some spectral theoretic applications of the supersymmetric approach outlined in Section 3. In particular, upon deriving the basic aspects of Weyl–Titchmarsh theory for the generalized Schrödinger operators H_j , $j = 1, 2$, in $L^2(\mathbb{R})^m$ (cf. (3.5), (3.6)) and a discussion of the corresponding spectral representations, we derive a local Borg–Marchenko uniqueness theorem by utilizing the known analog for the Dirac operator D .

Since the Schrödinger operator H_2 as defined in (3.6) with coefficient $\phi(\cdot)$ is realized as a Schrödinger operator of the form H_1 as defined in (3.5) with coefficient $\phi(\cdot)$ replaced by $-\phi(\cdot)$, it suffices to exclusively study spectral theory for Schrödinger operators of the form H_1 . In fact, the forms of many of the subsequent formulas relevant to the spectral theory of H_j , $j = 1, 2$, are independent of the choice $j \in \{1, 2\}$, the exception being (4.33), (4.34) below.

To set the stage for the main results of this section, we begin with a discussion of solutions to the equation $\tau_j u = zu$, $z \in \mathbb{C} \setminus \mathbb{R}$, $j = 1, 2$, with τ_j as defined in (3.13), and their corresponding Wronskian relations.

To this end, suppose $z_k \in \mathbb{C} \setminus \mathbb{R}$, $k = 1, 2$, and that $u_j(z_1, \cdot)$ and $u_j(z_2, \cdot)$ satisfy

$$u_j(z_k, \cdot), u_j^{[1,j]}(z_k, \cdot) \in AC_{loc}(\mathbb{R})^{m \times m}, \quad \tau_j u_j(z_k, \cdot) = z_k u_j(z_k, \cdot), \quad j, k = 1, 2. \quad (4.1)$$

Then one observes that

$$\begin{aligned} & W(u_j(z_1, \cdot)^*, u_j(z_2, \cdot))(x) \\ &= u_j(z_1, x)^* u_j^{[1,j]}(z_2, x) - (u_j^{[1,j]}(z_1, x))^* u_j(z_2, x), \quad j = 1, 2, \end{aligned} \quad (4.2)$$

so that the Wronskian in (4.2) is differentiable almost everywhere. Moreover, (4.1) implies

$$\begin{aligned} \frac{d}{dx}(u_j^{[1,j]}(z_k, x)) &= (-1)^{j+1} \phi(x) u_j^{[1,j]}(z_k, x) - z_k u_j(z_k, x) \\ &\quad \text{for a.e. } x \in \mathbb{R}, \quad j, k = 1, 2. \end{aligned} \quad (4.3)$$

As a result, one computes

$$\begin{aligned} & \frac{d}{dx} W(u_j(z_1, \cdot)^*, u_j(z_2, \cdot))(x) \\ &= \frac{d}{dx} [u_j(z_1, x)^* u_j^{[1,j]}(z_2, x) - (u_j^{[1,j]}(z_1, x))^* u_j(z_2, x)] \\ &= (\overline{z_1} - z_2) u_j(z_1, x)^* u_j(z_2, x) \quad \text{for a.e. } x \in \mathbb{R}. \end{aligned} \quad (4.4)$$

We summarize the above considerations as follows:

Lemma 4.1. *Assume Hypothesis 2.1 and suppose that $z_k \in \mathbb{C} \setminus \mathbb{R}$, $k = 1, 2$. If $u_j(z_k, \cdot)$, $j, k = 1, 2$, satisfy (4.1), then*

$$\begin{aligned} \frac{d}{dx} W(u_j(z_1, \cdot)^*, u_j(z_2, \cdot))(x) &= (\overline{z_1} - z_2) u_j(z_1, x)^* u_j(z_2, x) \\ &\quad \text{for a.e. } x \in \mathbb{R}, \quad j = 1, 2. \end{aligned} \quad (4.5)$$

As an immediate consequence of Lemma 4.1, one has the following result.

Corollary 4.2. *Assume Hypothesis 2.1. Then for $z \in \mathbb{C} \setminus \mathbb{R}$, the following identity holds:*

$$\begin{aligned} \text{Im}(z) \int_{x_0}^x dx' \psi_{\pm,j}(z, x', x_0)^* \psi_{\pm,j}(z, x', x_0) \\ = \text{Im}(\widehat{M}_{\pm,0,j}(z, x_0)) - (2i)^{-1} W(\psi_{\pm,j}(z, \cdot, x_0)^*, \psi_{\pm,j}(z, \cdot, x_0))(x), \\ x \in \mathbb{R}, j = 1, 2. \end{aligned} \quad (4.6)$$

Proof. Let $z \in \mathbb{C} \setminus \mathbb{R}$ be fixed and choose $u_j(z_1, \cdot) = u_j(z_2, \cdot) = \psi_{\pm,j}(z, \cdot, x_0)$, $j = 1, 2$, in (4.5), one obtains

$$\begin{aligned} \frac{d}{dx'} W(\psi_{\pm,j}(z, \cdot, x_0)^*, \psi_{\pm,j}(z, \cdot, x_0))(x') \\ = -2i \text{Im}(z) \psi_{\pm,j}(z, x', x_0)^* \psi_{\pm,j}(z, x', x_0) \quad \text{for a.e. } x' \in \mathbb{R}, j = 1, 2. \end{aligned} \quad (4.7)$$

Integration of both sides of (4.7) from x_0 to x , using the normalizations in (3.23) and (3.24), yields (4.6). \square

Lemma 4.3. *Assume Hypothesis 2.1. Then for any $z \in \mathbb{C} \setminus \mathbb{R}$,*

$$\lim_{x \rightarrow \pm\infty} W(\psi_{\pm,j}(z, \cdot, x_0)^*, \psi_{\pm,j}(z, \cdot, x_0))(x) = 0, \quad j = 1, 2. \quad (4.8)$$

Proof. We provide a proof of (4.8) for the case $j = 1$; an analogous argument is used to settle the case $j = 2$. In order to prove (4.8) for $j = 1$, fix $z \in \mathbb{C} \setminus \mathbb{R}$ and observe that by (3.11) and (3.18),

$$\begin{aligned} W_1(\psi_{\pm,1}(z, \cdot, x_0)^*, \psi_{\pm,1}(z, \cdot, x_0))(x) \\ = \psi_{\pm,1}(z, x, x_0)^* \psi_{\pm,1}^{[1,1]}(z, x, x_0) - \psi_{\pm,1}^{[1,1]}(z, x, x_0)^* \psi_{\pm,1}(z, x, x_0) \\ = \xi u_{\pm,1}(\xi, x, x_0, \alpha_0)^* u_{\pm,2}(\xi, x, x_0, \alpha_0) \\ - \xi u_{\pm,2}(\xi, x, x_0, \alpha_0)^* u_{\pm,1}(\xi, x, x_0, \alpha_0) \\ = (\xi - \bar{\xi}) u_{\pm,2}(\xi, x, x_0, \alpha_0)^* u_{\pm,1}(\xi, x, x_0, \alpha_0) \\ - \bar{\xi} U_{\pm}(\xi, x, x_0, \alpha_0)^* J U_{\pm}(\xi, x, x_0, \alpha_0) \quad \text{for a.e. } x \in \mathbb{R}, z = \xi^2. \end{aligned} \quad (4.9)$$

Since \mathcal{D} is in the limit point case one has the following limit relation (cf., e.g., [62, Corollary 2.3])

$$\lim_{x \rightarrow \pm\infty} U_{\pm}(\xi, x, x_0, \alpha_0)^* J U_{\pm}(\xi, x, x_0, \alpha_0) = 0. \quad (4.10)$$

Thus, in order to prove (4.8), it suffices to show

$$\lim_{x \rightarrow \pm\infty} u_{\pm,2}(\zeta, x, x_0, \alpha_0)^* u_{\pm,1}(\zeta, x, x_0, \alpha_0) = 0. \quad (4.11)$$

To this end, one observes that the function under the limit in (4.11) is differentiable and that, in fact,

$$\begin{aligned} & [u_{\pm,2}(\zeta, \cdot, x_0, \alpha_0)^* u_{\pm,1}(\zeta, \cdot, x_0, \alpha_0)]'(x) \\ &= \zeta u_{\pm,2}(\zeta, x, x_0, \alpha_0)^* u_{\pm,2}(\zeta, x, x_0, \alpha_0) \\ &\quad - \bar{\zeta} u_{\pm,1}(\zeta, x, x_0, \alpha_0)^* u_{\pm,1}(\zeta, x, x_0, \alpha_0) \quad \text{for a.e. } x \in \mathbb{R}. \end{aligned} \quad (4.12)$$

We recall that $u_{\pm,2}(\zeta, \cdot, x_0, \alpha_0), u_{\pm,1}(\zeta, \cdot, x_0, \alpha_0) \in L^2((0, \pm\infty))^{m \times m}$. As a result, one infers that

$$u_{\pm,2}(\zeta, \cdot, x_0, \alpha_0)^* u_{\pm,1}(\zeta, \cdot, x_0, \alpha_0) \in L^1((0, \pm\infty))^{m \times m}, \quad (4.13)$$

the same containment is true for the derivative by (4.12). Moreover,

$$\begin{aligned} & u_{\pm,2}(\zeta, x, x_0, \alpha_0)^* u_{\pm,1}(\zeta, x, x_0, \alpha_0) \\ &= u_{\pm,2}(\zeta, 0, x_0, \alpha_0)^* u_{\pm,1}(\zeta, 0, x_0, \alpha_0) \\ &\quad + \int_0^x dx' [u_{\pm,2}(\zeta, x', x_0, \alpha_0)^* u_{\pm,1}(\zeta, x', x_0, \alpha_0)]'(x'), \quad x \in \mathbb{R}, \end{aligned} \quad (4.14)$$

coupled with the fact that the function appearing under the integral in (4.14) belongs to $L^1((0, \pm\infty))$, affirms the existence of the limits appearing in (4.11). In light of (4.13), both limits must equal zero. \square

Taking limits $x \rightarrow \pm\infty$ throughout (4.6) and using (4.8), one obtains the fundamental identities:

$$\begin{aligned} & \operatorname{Im}(\hat{M}_{\pm,0,j}(z, x_0)) \\ &= \operatorname{Im}(z) \int_{x_0}^{\pm\infty} dx' \psi_{\pm,j}(z, x', x_0)^* \psi_{\pm,j}(z, x', x_0), \quad z \in \mathbb{C} \setminus \mathbb{R}, \quad j = 1, 2. \end{aligned} \quad (4.15)$$

The identities in (4.15) show that $\pm \hat{M}_{\pm,0,j}(z, x_0)$, $j = 1, 2$, are Nevanlinna–Herglotz functions. We summarize this together with some other relevant properties of the generalized Dirichlet-type $m \times m$ matrix-valued Weyl–Titchmarsh functions $\hat{M}_{\pm,0,j}(z, x_0)$, $j = 1, 2$, (associated to H_j) in the following result.

Lemma 4.4. *Assume Hypothesis 2.1 and let $\widehat{M}_{\pm,0,j}(z, x_0)$, $j = 1, 2$, $z \in \mathbb{C} \setminus \mathbb{R}$, denote the generalized Dirichlet-type $m \times m$ matrix-valued Weyl–Titchmarsh functions associated to H_j as defined by (3.20) and (3.21). Then $\pm \widehat{M}_{\pm,0,j}(\cdot, x_0)$, $j = 1, 2$, is an $m \times m$ matrix-valued Nevanlinna–Herglotz function of maximal rank m . In particular,*

$$\operatorname{Im}(\pm \widehat{M}_{\pm,0,j}(z, x_0)) \geq 0, \quad z \in \mathbb{C}_+, \quad (4.16)$$

$$\widehat{M}_{\pm,0,j}(\bar{z}, x_0) = \widehat{M}_{\pm,0,j}(z, x_0)^*, \quad z \in \mathbb{C} \setminus \mathbb{R}, \quad (4.17)$$

$$\operatorname{rank}(\widehat{M}_{\pm,0,j}(z, x_0)) = m, \quad z \in \mathbb{C} \setminus \mathbb{R}, \quad (4.18)$$

$$\lim_{\varepsilon \downarrow 0} \widehat{M}_{\pm,0,j}(\lambda + i\varepsilon, x_0) \text{ exists for a.e. } \lambda \in \mathbb{R}, \quad j = 1, 2. \quad (4.19)$$

Proof. The inequalities in (4.16) follow immediately from equation (4.15). By equations (3.20) and (3.21), $\widehat{M}_{\pm,0,j}(\cdot, x_0)$, $j = 1, 2$, are analytic on $\mathbb{C} \setminus \mathbb{R}$, thus they are Nevanlinna–Herglotz by (4.16). Relation (4.17) (resp., (4.18)) follows from (3.20) and (3.21) via (2.17) (resp., (2.18)). Finally, (4.19) follows from the fact that $\pm \widehat{M}_{\pm,0,j}(\cdot, x_0)$, $j = 1, 2$, are Nevanlinna–Herglotz functions (cf., e.g., [47]). Alternatively, (4.19) can be immediately inferred from (3.20) and (3.21) together with (2.19). \square

Remark 4.5. We used Corollary 4.2 and Lemma 4.3 to prove $\pm \widehat{M}_{\pm,0,j}(\cdot, x_0)$, $j = 1, 2$, are Nevanlinna–Herglotz functions. Alternatively, one can use the following approach based on computing the imaginary parts for $\pm \widehat{M}_{\pm,0,j}(\cdot, x_0)$, $j = 1, 2$, directly using the known representations for $\operatorname{Im}(M_{\pm}^D(\xi, c_0, \alpha_0))$ (cf. (2.25)). We briefly sketch how this approach is carried out.

Note that the system

$$\mathcal{D}\Psi = \xi\Psi, \quad \xi \in \mathbb{C} \setminus \mathbb{R}, \quad (4.20)$$

can be recast as

$$J\Psi' = [\xi I_{2m} + B]\Psi, \quad \xi \in \mathbb{C} \setminus \mathbb{R}, \quad (4.21)$$

where J is the $2m \times 2m$ matrix defined in (2.6), and

$$B = B(x) = \begin{pmatrix} 0 & \phi(x) \\ \phi(x) & 0 \end{pmatrix} \quad \text{for a.e. } x \in \mathbb{R}. \quad (4.22)$$

One can then verify by direct computation that if $\Psi_j = \Psi_j(\xi_j, \cdot)$, $j = 1, 2$, denote solutions of (4.21) with $\xi_j \in \mathbb{C} \setminus \mathbb{R}$, $j = 1, 2$, then

$$(\Psi_1^* \mathfrak{S}_1 \Psi_2)' = -(\xi_2 + \overline{\xi_1}) \Psi_1^* \mathfrak{S}_3 \Psi_2, \quad (4.23)$$

where

$$\mathfrak{S}_1 = \begin{pmatrix} 0 & I_m \\ I_m & 0 \end{pmatrix}, \quad \mathfrak{S}_3 = \begin{pmatrix} I_m & 0 \\ 0 & -I_m \end{pmatrix}. \quad (4.24)$$

Let $z \in \mathbb{C} \setminus \mathbb{R}$ and fix $\xi \in \mathbb{C}$ with $\xi^2 = z$ and $\text{Im}(\xi) > 0$. Upon decomposing ξ and $\pm \widehat{M}_{\pm,0,1}(\cdot, x_0)$ into its real and imaginary parts and using (3.20), one computes

$$\text{Im}(\pm \widehat{M}_{\pm,0,1}(z, x_0)) = \text{Im}(\xi) \text{Re}(M_{\pm}^D(\xi, x_0, \alpha_0)) + \text{Re}(\xi) \text{Im}(M_{\pm}^D(\xi, x_0, \alpha_0)). \quad (4.25)$$

Choosing $\Psi_1 = \Psi_2 = U_{\pm}(\xi, \cdot, x_0, \alpha_0)$ and $\xi_1 = \xi_2 = \xi$ in (4.23) and integrating over $[x_0, \infty)$ yields

$$\begin{aligned} & \text{Re}(M_{\pm}^D(\xi, x_0, \alpha_0)) \\ &= \frac{1}{2} \lim_{x' \rightarrow \pm\infty} U_{\pm}(\xi, x', x_0, \alpha_0^*) \mathfrak{S}_1 U_{\pm}(\xi, x', x_0, \alpha_0) \\ & \quad + \text{Re}(\xi) \int_{x_0}^{\pm\infty} dx [u_{\pm,1}(\xi, x, x_0, \alpha_0)^* u_{\pm,1}(\xi, x, x_0, \alpha_0) \\ & \quad \quad \quad - u_{\pm,2}(\xi, x, x_0, \alpha_0)^* u_{\pm,2}(\xi, x, x_0, \alpha_0)] \quad (4.26) \\ &= \text{Re}(\xi) \int_{x_0}^{\pm\infty} dx [u_{\pm,1}(\xi, x, x_0, \alpha_0)^* u_{\pm,1}(\xi, x, x_0, \alpha_0) \\ & \quad \quad \quad - u_{\pm,2}(\xi, x, x_0, \alpha_0)^* u_{\pm,2}(\xi, x, x_0, \alpha_0)], \end{aligned}$$

applying (4.11). The fact that the supersymmetric nature of D permits the representation (4.26) for the real part of $M_{\pm}^D(\xi, x_0, \alpha_0)$ appears to have gone unnoticed in the literature. By (2.25), the representation in (4.25) can be recast as

$$\begin{aligned} & \text{Im}(\widehat{M}_{\pm,0,1}(z, x_0)) \\ &= 2 \text{Im}(\xi) \text{Re}(\xi) \int_{x_0}^{\pm\infty} dx u_{\pm,1}(\xi, x, x_0, \alpha_0)^* u_{\pm,1}(\xi, x, x_0, \alpha_0) \\ &= \text{Im}(z) \int_{x_0}^{\pm\infty} dx \psi_{\pm,1}(z, x, x_0, \alpha_0)^* \psi_{\pm,1}(z, x, x_0, \alpha_0), \end{aligned}$$

(making use of (3.18)), implying (4.15). The result for $\widehat{M}_{\pm,0,2}(z, x_0)$ follows similarly.

In order to establish spectral theory for H_j , we introduce the $2m \times 2m$ matrix-valued Weyl–Titchmarsh matrix, $\widehat{\mathbf{M}}_j(z, x_0) \in \mathbb{C}^{2m \times 2m}$, $z \in \mathbb{C} \setminus \mathbb{R}$, associated to H_j , $j = 1, 2$, as follows:

$$\widehat{\mathbf{M}}_j(z, x_0) = (\widehat{\mathbf{M}}_{j,k,k'}(z, x_0))_{k,k'=0,1}, \quad z \in \mathbb{C} \setminus \mathbb{R}, \quad j = 1, 2, \quad (4.27)$$

$$\widehat{\mathbf{M}}_{j,0,0}(z, x_0) = W(z)^{-1}, \quad (4.28)$$

$$\widehat{\mathbf{M}}_{j,0,1}(z, x_0) = 2^{-1} W(z)^{-1} [\widehat{\mathbf{M}}_{-,0,j}(z, x_0) + \widehat{\mathbf{M}}_{+,0,j}(z, x_0)], \quad (4.29)$$

$$\widehat{\mathbf{M}}_{j,1,0}(z, x_0) = 2^{-1} [\widehat{\mathbf{M}}_{-,0,j}(z, x_0) + \widehat{\mathbf{M}}_{+,0,j}(z, x_0)] W(z)^{-1}, \quad (4.30)$$

$$\widehat{\mathbf{M}}_{j,1,1}(z, x_0) = \widehat{\mathbf{M}}_{\pm,0,j}(z, x_0) W(z)^{-1} \widehat{\mathbf{M}}_{\mp,0,j}(z, x_0), \quad (4.31)$$

where we have used the abbreviation (cf. (3.31))

$$\begin{aligned} W(z) &= W(\psi_{+,j}(\bar{z}, \cdot, x_0)^*, \psi_{-,j}(z, \cdot, x_0)) \\ &= \widehat{\mathbf{M}}_{-,0,j}(z, x_0) - \widehat{\mathbf{M}}_{+,0,j}(z, x_0), \quad z \in \mathbb{C} \setminus \mathbb{R}, \quad j = 1, 2. \end{aligned} \quad (4.32)$$

With (3.25) and the definitions in (4.27)–(4.31), one readily verifies that

$$\widehat{\mathbf{M}}_1(z, x_0) = \begin{pmatrix} \zeta^{-1} I_m & 0 \\ 0 & I_m \end{pmatrix} \mathbf{M}^D(\zeta, x_0, \alpha_0) \begin{pmatrix} I_m & 0 \\ 0 & \zeta I_m \end{pmatrix}, \quad (4.33)$$

and

$$\begin{aligned} \widehat{\mathbf{M}}_2(z, x_0) &= \begin{pmatrix} -\zeta^{-1} M_+^D(\zeta, x_0, \alpha_0) & 0 \\ 0 & M_-^D(\zeta, x_0, \alpha_0)^{-1} \end{pmatrix} \mathbf{M}^D(\zeta, x_0, \alpha_0) \\ &\quad \times \begin{pmatrix} -M_-^D(\zeta, x_0, \alpha_0) & 0 \\ 0 & \zeta M_+^D(\zeta, x_0, \alpha_0)^{-1} \end{pmatrix}, \quad \zeta^2 = z, \zeta \in \mathbb{C} \setminus \mathbb{R}. \end{aligned} \quad (4.34)$$

In addition, one notes that $\widehat{\mathbf{M}}_j(z, x_0)$ is a $\mathbb{C}^{2m \times 2m}$ -valued Nevanlinna–Herglotz matrix with representation

$$\begin{aligned} \widehat{\mathbf{M}}_j(z, x_0) &= \mathbf{C}_j(x_0) + \int_{\mathbb{R}} d\widehat{\Omega}_j(\lambda, x_0) \left[\frac{1}{\lambda - z} - \frac{\lambda}{1 + \lambda^2} \right], \quad z \in \mathbb{C} \setminus \mathbb{R}, \\ \mathbf{F}_j(x_0) &= \mathbf{F}_j(x_0)^*, \quad \int_{\mathbb{R}} \|d\widehat{\Omega}_j(\lambda, x_0)\|_{\mathbb{C}^{2m}} (1 + \lambda^2)^{-1} < \infty, \quad j = 1, 2. \end{aligned} \quad (4.35)$$

The Stieltjes inversion formula for the nonnegative $2m \times 2m$ matrix-valued measure $d\widehat{\Omega}_j(\cdot, x_0)$ then reads

$$\widehat{\Omega}_j((\lambda_1, \lambda_2], x_0) = \frac{1}{\pi} \lim_{\delta \downarrow 0} \lim_{\varepsilon \downarrow 0} \int_{\lambda_1 + \delta}^{\lambda_2 + \delta} d\lambda \operatorname{Im}(\widehat{M}_j(\lambda + i\varepsilon, x_0)), \quad (4.36)$$

$$\lambda_1, \lambda_2 \in \mathbb{R}, \lambda_1 < \lambda_2, j = 1, 2.$$

In particular, $d\widehat{\Omega}_j(\cdot, x_0)$, $j = 1, 2$, is a 2×2 block matrix-valued measure with $\mathbb{C}^{m \times m}$ -valued entries $d\widehat{\Omega}_{j,\ell,\ell'}(\cdot, x_0)$, $\ell, \ell' = 0, 1$. Since the diagonal entries of $\widehat{M}_j(\cdot, x_0)$ are Nevanlinna–Herglotz functions, the diagonal entries of the measure $d\widehat{\Omega}_j(\cdot, x_0)$ are nonnegative $\mathbb{C}^{m \times m}$ -valued measures. The off-diagonal entries of the measure $d\widehat{\Omega}_j(\cdot, x_0)$ naturally admit decompositions into a linear combination of four nonnegative matrix-valued measures.

Next, we relate the family of spectral projections $\{E_{H_j}(\lambda)\}_{\lambda \in \mathbb{R}}$ of the self-adjoint operator H_j and the $2m \times 2m$ matrix-valued increasing spectral function $\widehat{\Omega}_j(\lambda, x_0)$, $\lambda \in \mathbb{R}$, which generates the matrix-valued measure in the Nevanlinna–Herglotz representation (4.35) of $\widehat{M}_j(\cdot, x_0)$, $j = 1, 2$.

We note that for $F \in C(\mathbb{R})$,

$$(f, F(H_j)g)_{L^2(\mathbb{R})^m} = \int_{\mathbb{R}} d(f, E_{H_j}(\lambda)g)_{L^2(\mathbb{R})^m} F(\lambda), \quad j = 1, 2. \quad (4.37a)$$

with

$$f, g \in \operatorname{dom}(F(H_j))$$

$$= \left\{ h \in L^2(\mathbb{R})^m \mid \int_{\mathbb{R}} d\|E_{H_j}(\lambda)h\|_{L^2(\mathbb{R})^m}^2 |F(\lambda)|^2 < \infty \right\}, \quad j = 1, 2. \quad (4.37b)$$

Theorem 4.6. *Assume Hypothesis 2.1 and let $f, g \in C_0^\infty(\mathbb{R})^m$, $F \in C(\mathbb{R})$, $x_0 \in \mathbb{R}$, and $\lambda_1, \lambda_2 \in \mathbb{R}$, $\lambda_1 < \lambda_2$. Then,*

$$(f, F(H_j)E_{H_j}((\lambda_1, \lambda_2])g)_{L^2(\mathbb{R})^m}$$

$$= (\hat{f}_j(\cdot, x_0), M_F M_{\chi_{(\lambda_1, \lambda_2]}} \hat{g}_j(\cdot, x_0))_{L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0))}, \quad j = 1, 2, \quad (4.38)$$

where we introduced the notation

$$\hat{h}_{j,0}(\lambda, x_0) = \int_{\mathbb{R}} dx c_j(\lambda, x, x_0)^* h(x), \quad (4.39a)$$

$$\hat{h}_{j,1}(\lambda, x_0) = \int_{\mathbb{R}} dx s_j(\lambda, x, x_0)^* h(x) \quad (4.39b)$$

$$\hat{h}_j(\lambda, x_0) = (\hat{h}_{j,0}(\lambda, x_0), \hat{h}_{j,1}(\lambda, x_0))^\top, \quad (4.39c)$$

for $\lambda \in \mathbb{R}$, $h \in C_0^\infty(\mathbb{R})^m$, $j = 1, 2$, and M_G denotes the maximally defined operator of multiplication by $G \in C(\mathbb{R})$ in the Hilbert space $L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0))$,

$$(M_G \hat{h})(\lambda) = G(\lambda) \hat{h}(\lambda) = (G(\lambda) \hat{h}_0(\lambda), G(\lambda) \hat{h}_1(\lambda))^\top \quad \text{for } \widehat{\Omega}_j(\cdot, x_0)\text{-a.e. } \lambda \in \mathbb{R}, \quad (4.40a)$$

$$\begin{aligned} \hat{h} &\in \text{dom}(M_G) \\ &= \{\hat{k} \in L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0)) \mid G\hat{k} \in L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0))\}, \quad j = 1, 2. \end{aligned} \quad (4.40b)$$

Proof. We fix $j \in \{1, 2\}$. Using the weak version of Stone's formula, one obtains

$$\begin{aligned} (f, F(H_j) E_{H_j}((\lambda_1, \lambda_2]) g)_{L^2(\mathbb{R})^m} \\ = \lim_{\delta \downarrow 0} \lim_{\varepsilon \downarrow 0} \frac{1}{2\pi i} \int_{\lambda_1 + \delta}^{\lambda_2 + \delta} d\lambda F(\lambda) [(f, (H_j - (\lambda + i\varepsilon) I_{L^2(\mathbb{R})^m})^{-1} g)_{L^2(\mathbb{R})^m} \\ - (f, (H_j - (\lambda - i\varepsilon) I_{L^2(\mathbb{R})^m})^{-1} g)_{L^2(\mathbb{R})^m}]. \end{aligned} \quad (4.41)$$

Using that the resolvent of H_j is an integral operator with kernel (3.29) in (4.41), and freely interchanging the dx and dx' integrals with the limits and the $d\lambda$ integral (since all integration domains are finite and all integrands are continuous), and employing the expressions (3.22) for $\psi_{\pm,j}(z, x, x_0)$, one obtains

$$\begin{aligned} (f, F(H_j) E_{H_j}((\lambda_1, \lambda_2]) g)_{L^2(\mathbb{R})^m} &= \int_{\mathbb{R}} dx \left(f(x), \left\{ \int_{-\infty}^x dx' \right. \right. \\ &\times \lim_{\delta \downarrow 0} \lim_{\varepsilon \downarrow 0} \frac{1}{2\pi i} \int_{\lambda_1 + \delta}^{\lambda_2 + \delta} d\lambda F(\lambda) \left[[c_j(\lambda, x, x_0) + s_j(\lambda, x, x_0) \widehat{M}_{+,0,j}(\lambda + i\varepsilon, x_0)] \right. \\ &\times W(\lambda + i\varepsilon)^{-1} [c_j(\lambda, x', x_0)^* + \widehat{M}_{-,0,j}(\lambda + i\varepsilon, x_0) s_j(\lambda, x', x_0)^*] g(x') \\ &- [c_j(\lambda, x, x_0) + s_j(\lambda, x, x_0) \widehat{M}_{+,0,j}(\lambda - i\varepsilon, x_0)] \\ &\times W(\lambda - i\varepsilon)^{-1} [c_j(\lambda, x', x_0)^* + \widehat{M}_{-,0,j}(\lambda - i\varepsilon, x_0) s_j(\lambda, x', x_0)^*] g(x')] \Big] \\ &+ \int_x^\infty dx' \lim_{\delta \downarrow 0} \lim_{\varepsilon \downarrow 0} \frac{1}{2\pi i} \int_{\lambda_1 + \delta}^{\lambda_2 + \delta} d\lambda F(\lambda) \\ &\times \left[[c_j(\lambda, x, x_0) + s_j(\lambda, x, x_0) \widehat{M}_{-,0,j}(\lambda + i\varepsilon, x_0)] \right. \\ &\times W(\lambda + i\varepsilon)^{-1} [c_j(\lambda, x', x_0)^* + \widehat{M}_{+,0,j}(\lambda + i\varepsilon, x_0) s_j(\lambda, x', x_0)^*] g(x') \\ &- [c_j(\lambda, x, x_0) + s_j(\lambda, x, x_0) \widehat{M}_{-,0,j}(\lambda - i\varepsilon, x_0)] \\ &\times W(\lambda - i\varepsilon)^{-1} [c_j(\lambda, x', x_0)^* \\ &\left. \left. + \widehat{M}_{+,0,j}(\lambda - i\varepsilon, x_0) s_j(\lambda, x', x_0)^*] g(x') \right] \right\}_{\mathbb{C}^m}. \end{aligned} \quad (4.42)$$

Here we employed (4.17), the fact that for fixed $x \in \mathbb{R}$, $c_j(z, x, x_0)$ and $s_j(z, x, x_0)$ are entire with respect to z , that $c_j(z, \cdot, x_0), s_j(z, \cdot, x_0) \in AC_{\text{loc}}(\mathbb{R}; \mathcal{H})$, and hence that

$$c_j(\lambda \pm i\varepsilon, x, x_0) \underset{\varepsilon \downarrow 0}{=} c_j(\lambda, x, x_0) \pm i\varepsilon(d/dz)c_j(z, x, x_0)|_{z=\lambda} + O(\varepsilon^2), \quad (4.43a)$$

$$s_j(\lambda \pm i\varepsilon, x, x_0) \underset{\varepsilon \downarrow 0}{=} s_j(\lambda, x, x_0) \pm i\varepsilon(d/dz)s_j(z, x, x_0)|_{z=\lambda} + O(\varepsilon^2), \quad (4.43b)$$

with $O(\varepsilon^2)$ being uniform with respect to (λ, x) as long as λ and x vary in compact subsets of \mathbb{R} . Moreover, we used that

$$\varepsilon \|\widehat{M}_j(\lambda + i\varepsilon, x_0)\|_{\mathbb{C}^{2m \times 2m}} \leq C(\lambda_1, \lambda_2, \varepsilon_0, x_0), \quad \lambda \in [\lambda_1, \lambda_2], \quad 0 < \varepsilon \leq \varepsilon_0, \quad (4.44a)$$

$$\varepsilon \|\operatorname{Re}(\widehat{M}_j(\lambda + i\varepsilon, x_0))\|_{\mathbb{C}^{2m \times 2m}} \underset{\varepsilon \downarrow 0}{=} o(1), \quad \lambda \in \mathbb{R}, \quad (4.44b)$$

since $\widehat{M}_j(\cdot, x_0)$, are $\mathbb{C}^{2m \times 2m}$ -valued Nevanlinna–Herglotz functions. Moreover, we utilized (4.17), (4.43), (4.44), and the elementary facts

$$\begin{aligned} & \operatorname{Im}[\widehat{M}_{\pm,0,j}(\lambda + i\varepsilon, x_0)W(\lambda + i\varepsilon)^{-1}] \\ &= \frac{1}{2} \operatorname{Im}[[\widehat{M}_{-,0,j}(\lambda + i\varepsilon, x_0) + \widehat{M}_{+,0,j}(\lambda + i\varepsilon, x_0)]W(\lambda + i\varepsilon)^{-1}], \end{aligned} \quad (4.45a)$$

and

$$\begin{aligned} & \operatorname{Im}[W(\lambda + i\varepsilon)^{-1}\widehat{M}_{\pm,0,j}(\lambda + i\varepsilon, x_0)] \\ &= \frac{1}{2} \operatorname{Im}[W(\lambda + i\varepsilon)^{-1}[\widehat{M}_{-,0,j}(\lambda + i\varepsilon, x_0) + \widehat{M}_{+,0,j}(\lambda + i\varepsilon, x_0)]], \end{aligned} \quad (4.45b)$$

with $\lambda \in \mathbb{R}$, $\varepsilon > 0$. Collecting appropriate terms in (4.42) then yields

$$\begin{aligned} & (f, F(H_j)E_H((\lambda_1, \lambda_2])g)_{L^2(\mathbb{R})^m} \\ &= \int_{\mathbb{R}} dx \left(f(x), \int_{\mathbb{R}} dx' \lim_{\delta \downarrow 0} \lim_{\varepsilon \downarrow 0} \frac{1}{\pi} \int_{\lambda_1 + \delta}^{\lambda_2 + \delta} d\lambda F(\lambda) \right. \\ & \quad \times \left\{ c_j(\lambda, x, x_0) \operatorname{Im}[W(\lambda + i\varepsilon)^{-1}]c_j(\lambda, x', x_0)^* + 2^{-1}c_j(\lambda, x, x_0) \right. \\ & \quad \times \operatorname{Im}[W(\lambda + i\varepsilon)^{-1}[\widehat{M}_{-,0,j}(\lambda + i\varepsilon, x_0) + \widehat{M}_{+,0,j}(\lambda + i\varepsilon, x_0)]]s_j(\lambda, x', x_0)^* \\ & \quad + 2^{-1}s_j(\lambda, x, x_0) \\ & \quad \times \operatorname{Im}[[\widehat{M}_{-,0,j}(\lambda + i\varepsilon, x_0) + \widehat{M}_{+,0,j}(\lambda + i\varepsilon, x_0)]W(\lambda + i\varepsilon)^{-1}]c_j(\lambda, x', x_0)^* \\ & \quad + s_j(\lambda, x, x_0) \operatorname{Im}[\widehat{M}_{-,0,j}(\lambda + i\varepsilon, x_0)W(\lambda + i\varepsilon)^{-1}] \\ & \quad \times \widehat{M}_{+,0,j}(\lambda + i\varepsilon, x_0)]s_j(\lambda, x', x_0)^* \Big\} g(x') \Big)_{\mathbb{C}^m}. \end{aligned} \quad (4.46)$$

Since by (4.36) (for $\ell, \ell' = 0, 1$)

$$\begin{aligned} \int_{(\lambda_1, \lambda_2]} d\widehat{\Omega}_{j, \ell, \ell'}(\lambda, x_0) &= \widehat{\Omega}_{j, \ell, \ell'}((\lambda_1, \lambda_2], x_0) \\ &= \lim_{\delta \downarrow 0} \lim_{\varepsilon \downarrow 0} \frac{1}{\pi} \int_{\lambda_1 + \delta}^{\lambda_2 + \delta} d\lambda \operatorname{Im}(\widehat{M}_{j, \ell, \ell'}(\lambda + i\varepsilon, x_0)), \end{aligned} \quad (4.47)$$

one also has (again for $\ell, \ell' \in \{0, 1\}$)

$$\begin{aligned} \int_{\mathbb{R}} d\widehat{\Omega}_{j, \ell, \ell'}(\lambda, x_0) h(\lambda) & \\ &= \lim_{\varepsilon \downarrow 0} \frac{1}{\pi} \int_{\mathbb{R}} d\lambda \operatorname{Im}(\widehat{M}_{j, \ell, \ell'}(\lambda + i\varepsilon, x_0)) h(\lambda), \quad h \in C_0(\mathbb{R})^m, \end{aligned} \quad (4.48)$$

$$\begin{aligned} \int_{(\lambda_1, \lambda_2]} d\widehat{\Omega}_{j, \ell, \ell'}(\lambda, x_0) k(\lambda) & \\ &= \lim_{\delta \downarrow 0} \lim_{\varepsilon \downarrow 0} \frac{1}{\pi} \int_{\lambda_1 + \delta}^{\lambda_2 + \delta} d\lambda \operatorname{Im}(\widehat{M}_{j, \ell, \ell'}(\lambda + i\varepsilon, x_0)) k(\lambda), \quad k \in C(\mathbb{R})^m. \end{aligned} \quad (4.49)$$

Then using equations (4.27)–(4.31) and (4.39), and interchanging the dx, dx' and $d\widehat{\Omega}_{j, \ell, \ell'}(\cdot, x_0), \ell, \ell' = 0, 1$, integrals once more, one concludes from (4.46) that

$$\begin{aligned} (f, F(H_j)E_{H_j}((\lambda_1, \lambda_2])g)_{L^2(\mathbb{R})^m} & \\ &= \int_{(\lambda_1, \lambda_2]} F(\lambda) (\hat{f}_j(\lambda, x_0), d\widehat{\Omega}_j(\lambda, x_0) \hat{g}_j(\lambda, x_0))_{\mathbb{C}^{2m}}, \end{aligned} \quad (4.50)$$

implying (4.38). \square

Next, we improve on Theorem 4.6 and remove the compact support restrictions on f and g in the usual way, closely following and appropriately adapting the argument of [50, (2.46)–(2.67)]. This leads to a variant of the spectral theorem for (functions of) H_j , $j = 1, 2$. We consider the map

$$\begin{aligned} \tilde{U}_j(x_0): C_0^\infty(\mathbb{R})^m &\longrightarrow L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0)), \\ h &\longmapsto \hat{h}_j(\cdot, x_0) = (\hat{h}_{j,0}(\lambda, x_0), \hat{h}_{j,1}(\lambda, x_0))^\top, \end{aligned} \quad (4.51a)$$

$$\hat{h}_{j,0}(\lambda, x_0) = \int_{\mathbb{R}} dx c_j(\lambda, x, x_0)^* h(x), \quad (4.51b)$$

$$\hat{h}_{j,1}(\lambda, x_0) = \int_{\mathbb{R}} dx s_j(\lambda, x, x_0)^* h(x). \quad (4.51c)$$

Taking $f = g$, $F = 1$, $\lambda_1 \downarrow -\infty$, and $\lambda_2 \uparrow \infty$ in (4.38) then shows that $\tilde{U}_j(x_0)$, $j = 1, 2$, are densely defined isometries in $L^2(\mathbb{R})^m$, which extend by continuity to isometries on $L^2(\mathbb{R})^m$. The latter are denoted by $U_j(x_0)$ and are defined by

$$\begin{aligned} U_j(x_0) : L^2(\mathbb{R})^m &\longrightarrow L^2(\mathbb{R}; d\hat{\Omega}_j(\cdot, x_0)), \\ h &\longmapsto \hat{h}_j(\cdot, x_0) = (\hat{h}_{j,0}(\cdot, x_0), \hat{h}_{j,1}(\cdot, x_0))^\top, \end{aligned} \quad (4.52a)$$

$$\begin{aligned} \hat{h}_j(\cdot, x_0) &= \begin{pmatrix} \hat{h}_{j,0}(\cdot, x_0) \\ \hat{h}_{j,1}(\cdot, x_0) \end{pmatrix} \\ &= \underset{a \downarrow -\infty, b \uparrow \infty}{\text{l.i.m.}} \begin{pmatrix} \int_a^b dx c_j(\cdot, x, x_0)^* h(x) \\ \int_a^b dx s_j(\cdot, x, x_0)^* h(x) \end{pmatrix}, \end{aligned} \quad (4.52b)$$

where l.i.m. refers to the $L^2(\mathbb{R}; d\hat{\Omega}_j(\cdot, x_0))$ -limit.

Theorem 4.7. *Assume Hypothesis 2.1 and let $F \in C(\mathbb{R})$ and $x_0 \in \mathbb{R}$. Then,*

$$U_j(x_0)F(H_j)U_j(x_0)^{-1} = M_F, \quad j = 1, 2, \quad (4.53)$$

in $L^2(\mathbb{R}; d\hat{\Omega}_j(\cdot, x_0))$ (cf. (4.40)). Moreover,

$$\sigma(H_j) = \text{supp}(d\hat{\Omega}_j(\cdot, x_0)), \quad j = 1, 2, \quad (4.54)$$

and the multiplicity of the spectrum of H_j , $j = 1, 2$, is at most equal to $2m$.

Proof. Again, we fix $j \in \{1, 2\}$. One observes that the calculation in (4.50) yields

$$\begin{aligned} (E_{H_j}((\lambda_1, \lambda_2])g)(x) \\ = \int_{(\lambda_1, \lambda_2]} (c_j(\lambda, x, x_0), s_j(\lambda, x, x_0)) d\hat{\Omega}_j(\lambda, x_0) \hat{g}_j(\lambda, x_0), \quad g \in C_0^\infty(\mathbb{R})^m, \end{aligned} \quad (4.55)$$

and, as a result, extends to all $g \in L^2(\mathbb{R})^m$ by continuity. Taking $\lambda_1 \downarrow -\infty$ and $\lambda_2 \uparrow \infty$ in (4.55) and using the spectral family properties

$$\underset{\lambda \downarrow -\infty}{\text{s-lim}} E_{H_j}(\lambda) = 0, \quad \underset{\lambda \uparrow \infty}{\text{s-lim}} E_{H_j}(\lambda) = I_{L^2(\mathbb{R})^m},$$

where $E_{H_j}(\lambda) = E_{H_j}((-\infty, \lambda])$, $\lambda \in \mathbb{R}$, then yields

$$\begin{aligned} g(\cdot) &= \underset{\mu_1 \downarrow -\infty, \mu_2 \uparrow \infty}{\text{l.i.m.}} \int_{(\mu_1, \mu_2]} (c_j(\lambda, \cdot, x_0), s_j(\lambda, \cdot, x_0)) d\hat{\Omega}_j(\lambda, x_0), \\ &\quad \hat{g}_j(\lambda, x_0), \quad g \in L^2(\mathbb{R})^m, \end{aligned} \quad (4.56)$$

where l.i.m. here refers to the limit in $L^2(\mathbb{R})^m$. Next, we show that the maps $U_j(x_0)$ in (4.52) are onto and hence that $U_j(x_0)$ are unitary maps with

$$\begin{aligned} U_j(x_0)^{-1}: L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0)) &\longrightarrow L^2(\mathbb{R})^m, \\ \hat{h} &\longmapsto h_j, \end{aligned} \quad (4.57)$$

$$h_j(\cdot) = \underset{\mu_1 \downarrow -\infty, \mu_2 \uparrow \infty}{\text{l.i.m.}} \int_{(\mu_1, \mu_2]} (c_j(\lambda, \cdot, x_0), s_j(\lambda, \cdot, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \hat{h}(\lambda). \quad (4.58)$$

Letting $W_j(x_0)$ temporarily denote the operators defined by (4.57), one infers that $W_j(x_0)$ are bounded. Indeed, for any $\hat{f} \in C_0^\infty(\mathbb{R})^{2m}$, $g \in C_0^\infty(\mathbb{R})^m$, one computes

$$\begin{aligned} &(g, W_j(x_0) \hat{f})_{L^2(\mathbb{R})^m} \\ &= \int_{\mathbb{R}} dx \left(g(x), \int_{\mathbb{R}} (c_j(\lambda, x, x_0), s_j(\lambda, x, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \hat{f}(\lambda) \right)_{\mathbb{C}^m} \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}} dx ((c_j(\lambda, x, x_0), s_j(\lambda, x, x_0))^* g(x), d\widehat{\Omega}_j(\lambda, x_0) \hat{f}(\lambda))_{\mathbb{C}^{2m}} \\ &= (U_j(x_0)g, \hat{f})_{L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0))}. \end{aligned} \quad (4.59)$$

Since $U_j(x_0)$ are isometries, (4.59) extends by continuity to all $g \in L^2(\mathbb{R})^m$. Thus,

$$\begin{aligned} &\|W_j(x_0) \hat{f}\|_{L^2(\mathbb{R})^m} \\ &= \sup_{g \in L^2(\mathbb{R})^m, g \neq 0} \left| \frac{(g, W_j(x_0) \hat{f})_{L^2(\mathbb{R})^m}}{\|g\|_{L^2(\mathbb{R})^m}} \right| \\ &\leq \sup_{g \in L^2(\mathbb{R})^m, g \neq 0} \frac{\|U_j(x_0)g\|_{L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0))}}{\|g\|_{L^2(\mathbb{R})^m}} \|\hat{f}\|_{L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0))} \\ &= \|\hat{f}\|_{L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0))}, \quad \hat{f} \in C_0^\infty(\mathbb{R})^m. \end{aligned} \quad (4.60)$$

From the limiting relation in (4.57), one also infers that

$$W_j(x_0)U_j(x_0) = I_{L^2(\mathbb{R})^m}. \quad (4.61)$$

To verify that $U_j(x_0)$, $j = 1, 2$, are onto, and hence unitary, it suffices to prove that $W_j(x_0)$ are injective. Suppose that $\hat{f} = (f_0, f_1)^\top \in \ker(W_j(x_0))$. Let

$$\hat{f} \in L^2(\mathbb{R}; \widehat{\Omega}_j(\cdot, x_0)), \quad \lambda_1, \lambda_2 \in \mathbb{R}, \quad \lambda_1 < \lambda_2,$$

and consider

$$\begin{aligned} & (H_j - z I_{L^2(\mathbb{R})^m}) \left(\int_{(\lambda_1, \lambda_2]} (c_j(\lambda, \cdot, x_0), s_j(\lambda, \cdot, x_0)) (\lambda - z)^{-1} d\widehat{\Omega}_j(\lambda, x_0) \widehat{f}(\lambda) \right) \\ &= \int_{(\lambda_1, \lambda_2]} (c_j(\lambda, \cdot, x_0), s_j(\lambda, \cdot, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \widehat{f}(\lambda), \quad z \in \mathbb{C}_+. \end{aligned} \quad (4.62)$$

Then, for $z \in \mathbb{C}_+$,

$$\begin{aligned} & \int_{(\lambda_1, \lambda_2]} (c_j(\lambda, \cdot, x_0), s_j(\lambda, \cdot, x_0)) (\lambda - z)^{-1} d\widehat{\Omega}_j(\lambda, x_0) \widehat{f}(\lambda) \\ &= (H_j - z I_{L^2(\mathbb{R})^m})^{-1} \left(\int_{(\lambda_1, \lambda_2]} (c_j(\lambda, \cdot, x_0), s_j(\lambda, \cdot, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \widehat{f}(\lambda) \right). \end{aligned} \quad (4.63)$$

Taking $s\text{-lim}_{\lambda_1 \downarrow -\infty, \lambda_2 \uparrow \infty}$ in (4.63) implies

$$W_j(x_0)((\cdot - z)^{-1} \widehat{f}) = (H_j - z I_{L^2(\mathbb{R})^m})^{-1} W_j(x_0) \widehat{f}, \quad z \in \mathbb{C}_+. \quad (4.64)$$

Next, suppose that $\widehat{f}_0 = (f_0, f_1)^\top \in \ker(W_j(x_0))$, and take a sequence

$$\{\widehat{f}_n\}_{n \in \mathbb{N}} \subset L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0))$$

such that $\text{supp}(\widehat{f}_n)$ is compact for each $n \in \mathbb{N}$ and

$$\lim_{n \uparrow \infty} \|\widehat{f}_0 - \widehat{f}_n\|_{L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0))} = 0.$$

Then, since each \widehat{f}_n is compactly supported,

$$(W_j(x_0)((\cdot - z)^{-1} \widehat{f}_n))(x) = ((H_j - z I_{L^2(\mathbb{R})^m})^{-1} W_j(x_0) \widehat{f}_n)(x), \quad (4.65)$$

for $x \in \mathbb{R}$, $z \in \mathbb{C}_+$, and $n \in \mathbb{N}$. Consequently, for each $y \in \mathbb{R}$ and all $e \in \mathbb{C}^m$,

$$\begin{aligned} & \int_{x_0}^y dx \int_{\mathbb{R}} (e, (c_j(\lambda, x, x_0), s_j(\lambda, x, x_0)) (\lambda - z)^{-1} d\widehat{\Omega}_j(\lambda, x_0) \widehat{f}_n(\lambda))_{\mathbb{C}^m} \\ &= \int_{\mathbb{R}} \left(\int_{x_0}^y dx (c_j(\lambda, x, x_0), s_j(\lambda, x, x_0))^* e, d\widehat{\Omega}_j(\lambda, x_0) (\lambda - z)^{-1} \widehat{f}_n(\lambda) \right)_{\mathbb{C}^m} \\ &= \int_{x_0}^y dx (e, ((H_j - z I_{L^2(\mathbb{R})^m})^{-1} W_j(x_0) \widehat{f}_n)(x))_{\mathbb{C}^m}. \end{aligned} \quad (4.66)$$

One observes that

$$\begin{aligned} & \int_{x_0}^{\infty} dx (c_j(\lambda, x, x_0), s_j(\lambda, x, x_0))^* \chi_{[x_0, y]}(x) e \\ &= (U_j(x_0) \chi_{[x_0, y]} e)(\cdot) \in L^2(\mathbb{R}; d\widehat{\Omega}_j(\cdot, x_0)). \end{aligned} \quad (4.67)$$

Thus, taking the limit $n \uparrow \infty$ in (4.65) yields

$$\begin{aligned} & \lim_{n \uparrow \infty} \int_{x_0}^y dx \int_{\mathbb{R}} (e, (c_j(\lambda, x, x_0), s_j(\lambda, x, x_0)) d\widehat{\Omega}_j(\lambda, x_0) (\lambda - z)^{-1} \hat{f}_n(\lambda))_{\mathbb{C}^m} \\ &= \int_{\mathbb{R}} (\lambda - z)^{-1} \int_{x_0}^y dx (e, (c_j(\lambda, x, x_0), s_j(\lambda, x, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \hat{f}_0(\lambda))_{\mathbb{C}^m} \\ &= \lim_{n \uparrow \infty} \int_{x_0}^y dx (e, ((H_j - z I_{L^2(\mathbb{R})^m})^{-1} W_j(x_0) \hat{f}_n)(x))_{\mathbb{C}^m} \\ &= \int_{x_0}^y dx (e, ((H_j - z I_{L^2(\mathbb{R})^m})^{-1} W_j(x_0) \hat{f}_0)(x))_{\mathbb{C}^m} = 0, \\ & \quad y \in \mathbb{R}, z \in \mathbb{C}_+, e \in \mathbb{C}^m. \end{aligned} \quad (4.68)$$

Applying the Stieltjes inversion formula to the (finite) complex-valued measure in the 3rd line of (4.68), given by

$$\int_{x_0}^y dx (e, (c_j(\lambda, x, x_0), s_j(\lambda, x, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \hat{f}_0(\lambda))_{\mathbb{C}^m}, \quad (4.69)$$

implies for all $\lambda_1, \lambda_2 \in \mathbb{R}$, $\lambda_1 < \lambda_2$, and $e \in \mathbb{C}^m$, $y \in \mathbb{R}$,

$$\int_{(\lambda_1, \lambda_2]} \int_{x_0}^y dx (e, (c_j(\lambda, x, x_0), s_j(\lambda, x, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \hat{f}_0(\lambda))_{\mathbb{C}^m} = 0. \quad (4.70)$$

Differentiating (4.70) with respect to y , noting that $c_j(\lambda, y, x_0)$ and $s_j(\lambda, y, x_0)$ are continuous in $(\lambda, y) \in \mathbb{R}^2$, and using the dominated convergence theorem, one obtains

$$\int_{(\lambda_1, \lambda_2]} (e, (c_j(\lambda, y, x_0), s_j(\lambda, y, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \hat{f}_0(\lambda))_{\mathbb{C}^m} = 0, \quad (4.71)$$

for $y \in \mathbb{R}$, and $e \in \mathbb{C}^m$. In particular, taking $y = x_0$ in (4.71) and using (3.14) and (3.15), one obtains

$$\begin{aligned} & \int_{(\lambda_1, \lambda_2]} (e_1, (c_j(\lambda, x_0, x_0), s_j(\lambda, x_0, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \hat{f}_0(\lambda))_{\mathbb{C}^m} \\ &= \int_{(\lambda_1, \lambda_2]} ((e_1, 0_m)^\top, d\widehat{\Omega}_j(\lambda, x_0) \hat{f}_0(\lambda))_{\mathbb{C}^{2m}} = 0, \quad e_1 \in \mathbb{C}^m, \end{aligned} \quad (4.72)$$

where 0_m in (4.72) denotes the zero vector in \mathbb{C}^m . Next, applying the quasi-derivative [1, j] with respect to $y \in \mathbb{R}$ to (4.71), yields, for $y \in \mathbb{R}$, and $e \in \mathbb{C}^m$,

$$\int_{(\lambda_1, \lambda_2]} (e, (c_j^{[1,j]}(\lambda, y, x_0), s_j^{[1,j]}(\lambda, y, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \widehat{f}_0(\lambda))_{\mathbb{C}^m} = 0, \quad (4.73)$$

using the fact that, for $y \in \mathbb{R}$ and $e \in \mathbb{C}^m$,

$$\int_{(\lambda_1, \lambda_2]} (e, (\phi(y)c_j(\lambda, y, x_0), \phi(y)s_j(\lambda, y, x_0)) d\widehat{\Omega}_j(\lambda, x_0) \widehat{f}_0(\lambda))_{\mathbb{C}^m} = 0, \quad (4.74)$$

Subsequently, taking $y = x_0$ in (4.73), once more using (3.14) and (3.15) yields

$$\int_{(\lambda_1, \lambda_2]} ((0_m, e_2)^\top, d\widehat{\Omega}_j(\lambda, x_0) \widehat{f}_0(\lambda))_{\mathbb{C}^{2m}} = 0, \quad e_2 \in \mathbb{C}^m. \quad (4.75)$$

Taking $e = (e_1, e_2)^\top$ with $e_1, e_2 \in \mathbb{C}^m$ and adding (4.72) and (4.75), one obtains

$$\int_{(\lambda_1, \lambda_2]} (e, d\widehat{\Omega}_j(\lambda, x_0) \widehat{f}_0(\lambda))_{\mathbb{C}^{2m}} = 0, \quad e \in \mathbb{C}^{2m}. \quad (4.76)$$

Since λ_1 and λ_2 are arbitrary (apart from $\lambda_1 < \lambda_2$), (4.76) implies

$$\widehat{f}_0(\lambda) = 0 \quad \text{for } d\widehat{\Omega}_j(\cdot, x_0)\text{-a.e. } \lambda \in \mathbb{R}. \quad \square$$

The proofs of Theorems 4.6 and 4.7 are adaptation of the proofs of Theorems 2.12 and 2.14 in [50]. This strategy of proof immediately extends to all continuous or discrete second-order problems (such as Sturm–Liouville, Jacobi, and CMV operators) and first-order 2×2 systems (i.e., Dirac-type operators) with matrix-valued coefficients, see, for instance, [18] and [19]. In fact, it also extends to the infinite-dimensional case of bounded operator-valued coefficients (for the case of Schrödinger operators with bounded operator-valued potentials, see [49]).

In our final result, we show that the known local Borg–Marchenko results for Dirac-type operators worked out in [16] immediately imply local Borg–Marchenko results for generalized Schrödinger operators of the form H_j , $j = 1, 2$. For simplicity, we focus on H_1 only.

Theorem 4.8. *Suppose Hypothesis 2.1 holds with ϕ_j , $j = 1, 2$, in place of ϕ . Let $H_{1,1}$ (resp., $H_{1,2}$) denote the operator defined in (3.5) taking $\phi = \phi_1$ (resp., $\phi = \phi_2$) and denote by $\widehat{M}_{1,1}(z, x_0)$ (resp., $\widehat{M}_{1,2}(z, x_0)$) the corresponding $2m \times 2m$ block Weyl–Titchmarsh matrix as defined in (4.27)–(4.31). Then,*

$$\text{if for some } a > 0, \phi_1(x) = \phi_2(x) \text{ for a.e. } x \in (x_0 - a, x_0 + a), \quad (4.77)$$

one obtains

$$\|\widehat{\mathbf{M}}_{1,1}(z, x_0) - \widehat{\mathbf{M}}_{1,2}(z, x_0)\|_{\mathbb{C}^{2m \times 2m}} \underset{\substack{|z| \rightarrow \infty \\ z \in \rho_\theta}}{=} O(z^{1/2} e^{-2 \operatorname{Im}(z^{1/2})a}), \quad (4.78)$$

along any ray $\rho_\theta \subset \mathbb{C}$ with $\arg(z) = \theta \in (0, \pi) \cup (\pi, 2\pi)$, always choosing the branch of the square root with $\operatorname{Im}(z^{1/2}) > 0$ for $z \in \mathbb{C} \setminus [0, \infty)$. On the other hand, suppose that for all $\varepsilon > 0$,

$$\begin{aligned} & \|\widehat{\mathbf{M}}_{1,1}(z, x_0) - \widehat{\mathbf{M}}_{1,2}(z, x_0)\|_{\mathbb{C}^{2m \times 2m}} \\ &= \underset{\substack{|z| \rightarrow \infty \\ z \in \rho_{\theta_\ell}}}{O}(z^{1/2} e^{-2 \operatorname{Im}(z^{1/2})(a-\varepsilon)}), \quad \ell = 1, 2, \end{aligned} \quad (4.79)$$

along a ray $\rho_{\theta_1} \subset \mathbb{C}$ with $\arg(z) = \theta_1$ and $0 < \theta_1 < \pi$ and along a ray $\rho_{\theta_2} \subset \mathbb{C}$ with $\arg(z) = \theta_2$ and $\pi < \theta_2 < 2\pi$. If $m > 1$, assume in addition that $\phi_j \in L^\infty([x_0 - a, x_0 + a])^{m \times m}$, $j = 1, 2$. Then

$$\phi_1(x) = \phi_2(x) \text{ for a.e. } x \in [x_0 - a, x_0 + a]. \quad (4.80)$$

Proof. We begin by fixing some notation. Let $\mathbf{M}_j^D(\zeta, x_0, \alpha_0)$, $j = 1, 2$, denote the $2m \times 2m$ Weyl–Titchmarsh matrix defined by (2.29) corresponding to the Dirac-type operator D_j , $j = 1, 2$ defined by (2.4), that is, the operator (formally) defined as

$$D_j = J \frac{d}{dx} - B_j(x), \quad B_j(x) = \begin{pmatrix} 0 & -\phi_j(x) \\ -\phi_j(x) & 0 \end{pmatrix}, \quad j = 1, 2. \quad (4.81)$$

As a result of (4.33), one estimates

$$\begin{aligned} & \|\widehat{\mathbf{M}}_{1,1}(z, x_0) - \widehat{\mathbf{M}}_{1,2}(z, x_0)\|_{\mathbb{C}^{2m \times 2m}} \\ & \leq C |\zeta| \|\mathbf{M}_1^D(\zeta, x_0, \alpha_0) - \mathbf{M}_2^D(\zeta, x_0, \alpha_0)\|_{\mathbb{C}^{2m \times 2m}}, \quad \zeta^2 = z \in \mathbb{C} \setminus \mathbb{R}, |z| \geq 1, \end{aligned} \quad (4.82)$$

for a z -independent constant $C > 0$.

Evidently, taking $|z| \rightarrow \infty$ along the fixed ray $\rho_\theta \subset \mathbb{C}$ with $\arg(z) = \theta$ and $\theta \in (0, \pi) \cup (\pi, 2\pi)$ implies

$$|\zeta| \rightarrow \infty, \text{ along the fixed ray } \rho_{\theta/2} \subset \mathbb{C}_+ \text{ with } 0 < \arg(\zeta) = \theta/2 < \pi, \quad (4.83)$$

if $\zeta^2 = z$. The assumption in (4.77) implies

$$B_1(x) = B_2(x) \quad \text{for a.e. } x \in (x_0 - a, x_0 + a), \quad (4.84)$$

with $B_j(\cdot)$, $j = 1, 2$, as in (4.81). Consequently, [16, Theorem 5.5] implies

$$\|\mathbf{M}_1^D(\zeta, x_0, \alpha_0) - \mathbf{M}_2^D(\zeta, x_0, \alpha_0)\|_{\mathbb{C}^{2m \times 2m}} \underset{\substack{|\zeta| \rightarrow \infty \\ \zeta \in \rho_{\theta/2}}}{=} O(e^{-2\operatorname{Im}(\zeta)a}). \quad (4.85)$$

As $\operatorname{Re}(\zeta) = 2\theta^{-1} \operatorname{Im}(\zeta)$ for $\zeta \in \rho_{\theta/2}$, (4.85) implies

$$|\zeta| \|\mathbf{M}_1^D(\zeta, x_0, \alpha_0) - \mathbf{M}_2^D(\zeta, x_0, \alpha_0)\|_{\mathbb{C}^{2m \times 2m}} \underset{\substack{|\zeta| \rightarrow \infty \\ \zeta \in \rho_{\theta/2}}}{=} O(\operatorname{Im}(\zeta)e^{-2\operatorname{Im}(\zeta)a}). \quad (4.86)$$

Together, (4.82) and (4.86) yield (4.78).

Assuming (4.79), in order to prove (4.80), it suffices to prove

$$\begin{aligned} & \|\mathbf{M}_1^D(\zeta, x_0, \alpha_0) - \mathbf{M}_2^D(\zeta, x_0, \alpha_0)\|_{\mathbb{C}^{2m \times 2m}} \\ &= \underset{\substack{|\zeta| \rightarrow \infty \\ \zeta \in \rho_{\theta_\ell/2}}}{=} O(e^{-2\operatorname{Im}(\zeta)(a-\epsilon)}), \quad \epsilon > 0, \ell = 1, 2, \end{aligned} \quad (4.87)$$

since $\rho_{\theta_1/2}$ (resp., $\rho_{\theta_2/2}$) is a ray in \mathbb{C} with $0 < \arg(\zeta) = \theta_1/2 < \pi/2$ (resp., $\pi/2 < \arg(\zeta) = \theta_2/2 < \pi$). Indeed, by [16, Theorem 5.5], (4.87) implies

$$B_1(x) = B_2(x) \text{ for a.e. } x \in [x_0 - a, x_0 + a], \quad (4.88)$$

which clearly yields (4.80). In order to prove (4.87), let $\epsilon > 0$. Making use of (4.33), one writes,

$$\begin{aligned} & \|\mathbf{M}_1^D(\zeta, x_0, \alpha_0) - \mathbf{M}_2^D(\zeta, x_0, \alpha_0)\|_{\mathbb{C}^{2m \times 2m}} \\ &= \left\| \begin{pmatrix} \zeta I_m & 0 \\ 0 & I_m \end{pmatrix} [\widehat{\mathbf{M}}_{1,1}(z, x_0) - \widehat{\mathbf{M}}_{1,2}(z, x_0)] \begin{pmatrix} I_m & 0 \\ 0 & \zeta^{-1} I_m \end{pmatrix} \right\|_{\mathbb{C}^{2m \times 2m}} \\ &\leq C |\zeta| \|\widehat{\mathbf{M}}_{1,1}(z, x_0) - \widehat{\mathbf{M}}_{1,2}(z, x_0)\|_{\mathbb{C}^{2m \times 2m}}, \\ & \quad z = \zeta^2, \zeta \in \rho_{\theta_\ell/2}, |\zeta| \geq 1, \ell = 1, 2, \end{aligned} \quad (4.89)$$

for some ζ -independent constant $C > 0$. Choosing $\varepsilon = 2\epsilon$, one computes

$$\begin{aligned} & e^{2\operatorname{Im}(\zeta)(a-\epsilon)} \|\mathbf{M}_1^D(\zeta, x_0, \alpha_0) - \mathbf{M}_2^D(\zeta, x_0, \alpha_0)\|_{\mathbb{C}^{2m \times 2m}} \\ &\leq C |\zeta| e^{2\operatorname{Im}(\zeta)(a-\epsilon)} \|\widehat{\mathbf{M}}_{1,1}(\zeta^2, x_0) - \widehat{\mathbf{M}}_{1,2}(\zeta^2, x_0)\|_{\mathbb{C}^{2m \times 2m}} \end{aligned} \quad (4.90)$$

$$\leq \tilde{C} \operatorname{Im}(\zeta) e^{2\operatorname{Im}(\zeta)(a-\epsilon)} e^{-2\operatorname{Im}(\zeta)(a-\epsilon)} \quad (4.91)$$

$$= \tilde{C} \operatorname{Im}(\zeta) e^{-2\operatorname{Im}(\zeta)\epsilon}, \quad \zeta \in \rho_{\theta_\ell/2}, |\zeta| \gg 1, \ell = 1, 2, \quad (4.92)$$

where $\tilde{C} > 0$ is an appropriate ζ -independent constant. The estimate in (4.90) makes use of (4.89) and the estimate in (4.91) uses the assumption in (4.79) which is applicable since $\zeta \in \rho_{\theta_\ell/2}$, $|\zeta| \gg 1$ implies $z = \zeta^2 \in \rho_{\theta_\ell}$, $|z| \gg 1$. The function of ζ in (4.92) is bounded as $|\zeta| \rightarrow \infty$, $\zeta \in \rho_{\theta_\ell/2}$, $\ell = 1, 2$; hence, (4.92) implies

$$\begin{aligned} & \|M_1^D(\zeta, x_0, \alpha_0) - M_2^D(\zeta, x_0, \alpha_0)\|_{\mathbb{C}^{2m \times 2m}} \\ &= \underset{\substack{|\zeta| \rightarrow \infty \\ \zeta \in \rho_{\theta_\ell/2}}}{O}(e^{-2\operatorname{Im}(\zeta)(a-\epsilon)}), \quad \ell = 1, 2. \end{aligned} \quad (4.93)$$

Since $\epsilon > 0$ was arbitrary, (4.87) is established. \square

The reason for the additional assumption $\phi_j \in L^\infty([x_0 - a, x_0 + a])^{m \times m}$, $j = 1, 2$, in the case $m > 1$, is due to a technical issue as explained in detail in [16, Remark 5.4]. It should be noted in this connection that for matrix-valued Schrödinger operators $H_{1,j}$, $j = 1, 2$, with the standard hypothesis on potentials, that is, $V_j = V_j^* \in L^1_{\text{loc}}(\mathbb{R})^{m \times m}$, the local Borg–Marchenko-type results in Theorem 4.8 were derived in [41] (see also [45]). In this case no analog of the local boundedness assumptions on ϕ_j were necessary.

Finally, in the context of Theorem 4.8, we would also like to point out that an alternative approach to local Borg–Marchenko-type uniqueness results for Dirac-type operators, including a procedure for recovering the potential coefficient, was developed by Sakhnovich [118] (see also [119] and [120, Sects. 2.2.3, 4.2.2]).

A. Supersymmetric Dirac-type operators in a nutshell

In this appendix we briefly summarize some results on supersymmetric Dirac-type operators and commutation methods due to [23], [43], [133], and [134, Chapter 5] (see also [55]).

The standing assumption in this appendix will be the following.

Hypothesis A.1. *Let \mathcal{H}_j , $j = 1, 2$, be separable complex Hilbert spaces and*

$$T : \mathcal{H}_1 \supseteq \operatorname{dom}(T) \rightarrow \mathcal{H}_2 \quad (\text{A.1})$$

be a densely defined, closed, linear operator.

We define the self-adjoint Dirac-type operator in $\mathcal{H}_1 \oplus \mathcal{H}_2$ by

$$Q = \begin{pmatrix} 0 & T^* \\ T & 0 \end{pmatrix}, \quad \operatorname{dom}(Q) = \operatorname{dom}(T) \oplus \operatorname{dom}(T^*). \quad (\text{A.2})$$

Operators of the type Q play a role in supersymmetric quantum mechanics (see, e.g., the extensive list of references in [13]). Then,

$$Q^2 = \begin{pmatrix} T^*T & 0 \\ 0 & TT^* \end{pmatrix} \quad (\text{A.3})$$

and for notational purposes we also introduce

$$H_1 = T^*T \text{ in } \mathcal{H}_1, \quad H_2 = TT^* \text{ in } \mathcal{H}_2. \quad (\text{A.4})$$

In the following, we also need the polar decomposition of T and T^* , that is, the representations

$$T = V_T|T| = |T^*|V_T = V_T T^* V_T \quad \text{on } \text{dom}(T) = \text{dom}(|T|), \quad (\text{A.5})$$

$$T^* = V_{T^*}|T^*| = |T|V_{T^*} = V_{T^*}T V_{T^*} \quad \text{on } \text{dom}(T^*) = \text{dom}(|T^*|), \quad (\text{A.6})$$

$$|T| = V_{T^*}T = T^*V_T = V_{T^*}|T^*|V_T \quad \text{on } \text{dom}(|T|), \quad (\text{A.7})$$

$$|T^*| = V_T T^* = T V_{T^*} = V_T |T| V_{T^*} \quad \text{on } \text{dom}(|T^*|), \quad (\text{A.8})$$

where

$$|T| = (T^*T)^{1/2}, \quad |T^*| = (TT^*)^{1/2}, \quad V_{T^*} = (V_T)^*, \quad (\text{A.9})$$

$$V_{T^*}V_T = P_{\overline{\text{ran}(|T|)}} = P_{\overline{\text{ran}(T^*)}}, \quad V_TV_{T^*} = P_{\overline{\text{ran}(|T^*|)}} = P_{\overline{\text{ran}(T)}}. \quad (\text{A.10})$$

In particular, V_T is a partial isometry with initial set $\overline{\text{ran}(|T|)}$ and final set $\overline{\text{ran}(T)}$ and hence V_{T^*} is a partial isometry with initial set $\overline{\text{ran}(|T^*|)}$ and final set $\overline{\text{ran}(T^*)}$. In addition,

$$V_T = \begin{cases} \overline{T(T^*T)^{-1/2}} = \overline{(TT^*)^{-1/2}T} & \text{on } (\ker(T))^\perp, \\ 0 & \text{on } \ker(T). \end{cases} \quad (\text{A.11})$$

Next, we collect some properties relating H_1 and H_2 .

Theorem A.2 ([23]). *Assume Hypothesis A.1 and let ϕ be a bounded Borel measurable function on \mathbb{R} .*

(i) *One has*

$$\ker(T) = \ker(H_1) = (\text{ran}(T^*))^\perp, \quad \ker(T^*) = \ker(H_2) = (\text{ran}(T))^\perp, \quad (\text{A.12})$$

and

$$V_T H_1^{n/2} = H_2^{n/2} V_T, \quad n \in \mathbb{N}, \quad V_T \phi(H_1) = \phi(H_2) V_T. \quad (\text{A.13})$$

(ii) H_1 and H_2 are essentially isospectral, that is,

$$\sigma(H_1) \setminus \{0\} = \sigma(H_2) \setminus \{0\}, \quad (\text{A.14})$$

in fact,

$T^*T[I_{\mathcal{H}_1} - P_{\ker(T)}]$ is unitarily equivalent to $TT^*[I_{\mathcal{H}_2} - P_{\ker(T^*)}]$. (A.15)

In addition,

$$\begin{aligned} f \in \text{dom}(H_1) \text{ and } H_1 f = \lambda^2 f, \lambda \neq 0, \\ \implies Tf \in \text{dom}(H_2) \text{ and } H_2(Tf) = \lambda^2(Tf), \end{aligned} \quad (\text{A.16})$$

$$\begin{aligned} g \in \text{dom}(H_2) \text{ and } H_2 g = \mu^2 g, \mu \neq 0, \\ \implies T^*g \in \text{dom}(H_1) \text{ and } H_1(T^*g) = \mu^2(T^*g), \end{aligned} \quad (\text{A.17})$$

with multiplicities of eigenvalues preserved.

(iii) One has for $z \in \rho(H_1) \cap \rho(H_2)$,

$$I_{\mathcal{H}_2} + z(H_2 - zI_{\mathcal{H}_2})^{-1} \supseteq T(H_1 - zI_{\mathcal{H}_1})^{-1}T^*, \quad (\text{A.18})$$

$$I_{\mathcal{H}_1} + z(H_1 - zI_{\mathcal{H}_1})^{-1} \supseteq T^*(H_2 - zI_{\mathcal{H}_2})^{-1}T, \quad (\text{A.19})$$

and

$$T^*\phi(H_2) \supseteq \phi(H_1)T^*, \quad T\phi(H_1) \supseteq \phi(H_2)T, \quad (\text{A.20})$$

$$V_{T^*}\phi(H_2) \supseteq \phi(H_1)V_{T^*}, \quad V_T\phi(H_1) \supseteq \phi(H_2)V_T. \quad (\text{A.21})$$

As noted by E. Nelson (unpublished), Theorem A.2 follows from the spectral theorem and the elementary identities,

$$Q = V_Q|Q| = |Q|V_Q, \quad (\text{A.22})$$

$$\ker(Q) = \ker(|Q|) = \ker(Q^2) = (\text{ran}(Q))^\perp = \ker(T) \oplus \ker(T^*), \quad (\text{A.23})$$

$$\begin{aligned} I_{\mathcal{H}_1 \oplus \mathcal{H}_2} + z(Q^2 - zI_{\mathcal{H}_1 \oplus \mathcal{H}_2})^{-1} &= Q^2(Q^2 - zI_{\mathcal{H}_1 \oplus \mathcal{H}_2})^{-1} \\ &\supseteq Q(Q^2 - zI_{\mathcal{H}_1 \oplus \mathcal{H}_2})^{-1}Q, \quad z \in \rho(Q^2), \end{aligned} \quad (\text{A.24})$$

and

$$Q\phi(Q^2) \supseteq \phi(Q^2)Q, \quad (\text{A.25})$$

where

$$V_Q = \begin{pmatrix} 0 & (V_T)^* \\ V_T & 0 \end{pmatrix} = \begin{pmatrix} 0 & V_{T^*} \\ V_T & 0 \end{pmatrix}. \quad (\text{A.26})$$

In particular,

$$\ker(Q) = \ker(T) \oplus \ker(T^*), \quad P_{\ker(Q)} = \begin{pmatrix} P_{\ker(T)} & 0 \\ 0 & P_{\ker(T^*)} \end{pmatrix}, \quad (\text{A.27})$$

and we also recall that

$$\mathfrak{S}_3 Q \mathfrak{S}_3 = -Q, \quad \mathfrak{S}_3 = \begin{pmatrix} I_{\mathcal{H}_1} & 0 \\ 0 & -I_{\mathcal{H}_2} \end{pmatrix}, \quad (\text{A.28})$$

that is, Q and $-Q$ are unitarily equivalent. (For more details on Nelson's trick see also [132, §8.4] and [134, §5.2.3].) We also note that

$$\psi(|Q|) = \begin{pmatrix} \psi(|T|) & 0 \\ 0 & \psi(|T^*|) \end{pmatrix} \quad (\text{A.29})$$

for Borel measurable functions ψ on \mathbb{R} , and

$$\overline{[Q|Q|^{-1}]} = \begin{pmatrix} 0 & (V_T)^* \\ V_T & 0 \end{pmatrix} = V_Q \text{ if } \ker(Q) = \{0\}. \quad (\text{A.30})$$

Finally, we recall the following relationships between Q and H_j , $j = 1, 2$.

Theorem A.3 ([13], [133]). *Assume Hypothesis A.1.*

(i) *Introducing the unitary operator U on $(\ker(Q))^\perp$ by*

$$U = 2^{-1/2} \begin{pmatrix} I_{\mathcal{H}_1} & (V_T)^* \\ -V_T & I_{\mathcal{H}_2} \end{pmatrix} \text{ on } (\ker(Q))^\perp, \quad (\text{A.31})$$

one infers that

$$U Q U^{-1} = \begin{pmatrix} |A| & 0 \\ 0 & -|A^*| \end{pmatrix} \text{ on } (\ker(Q))^\perp. \quad (\text{A.32})$$

(ii) *For $\xi^2 \in \rho(H_1) \cap \rho(H_2)$, one has*

$$(Q - \xi I_{\mathcal{H}_1 \oplus \mathcal{H}_2})^{-1} = \begin{pmatrix} \xi(H_1 - \xi^2 I_{\mathcal{H}_1})^{-1} & T^*(H_2 - \xi^2 I_{\mathcal{H}_2})^{-1} \\ T(H_1 - \xi^2 I_{\mathcal{H}_1})^{-1} & \xi(H_2 - \xi^2 I_{\mathcal{H}_2})^{-1} \end{pmatrix}. \quad (\text{A.33})$$

(iii) *In addition,*

$$\begin{pmatrix} f_1 \\ f_2 \end{pmatrix} \in \text{dom}(Q) \text{ and } Q \begin{pmatrix} f_1 \\ f_2 \end{pmatrix} = \eta \begin{pmatrix} f_1 \\ f_2 \end{pmatrix}, \quad \eta \neq 0, \quad (\text{A.34})$$

$$\implies f_j \in \text{dom}(H_j) \text{ and } H_j f_j = \eta^2 f_j, \quad j = 1, 2.$$

Conversely,

$$\begin{aligned} f \in \text{dom}(H_1) \text{ and } H_1 f = \lambda^2 f, \lambda \neq 0, \\ \Rightarrow \left(\begin{matrix} f \\ \lambda^{-1} T f \end{matrix} \right) \in \text{dom}(Q) \text{ and } Q \left(\begin{matrix} f \\ \lambda^{-1} T f \end{matrix} \right) = \lambda \left(\begin{matrix} f \\ \lambda^{-1} T f \end{matrix} \right). \end{aligned} \quad (\text{A.35})$$

Similarly,

$$\begin{aligned} g \in \text{dom}(H_2) \text{ and } H_2 g = \mu^2 g, \mu \neq 0, \\ \Rightarrow \left(\begin{matrix} \mu^{-1} T^* g \\ g \end{matrix} \right) \in \text{dom}(Q) \text{ and } Q \left(\begin{matrix} \mu^{-1} T^* g \\ g \end{matrix} \right) = \mu \left(\begin{matrix} \mu^{-1} T^* g \\ g \end{matrix} \right). \end{aligned} \quad (\text{A.36})$$

References

- [1] S. Albeverio, F. Gesztesy, R. Høegh-Krohn, and H. Holden, *Solvable models in quantum mechanics*. 2nd revised ed. With an appendix by Pavel Exner. AMS Chelsea Publishing, Providence, R.I., 2005. [MR 2105735](#) [Zbl 1078.81003](#)
- [2] S. Albeverio, R. Hryniv, and Ya. Mykytyuk, On spectra of non-self-adjoint Sturm–Liouville operators. *Selecta Math. (N.S.)* **13** (2008), 571–599. [MR 2403304](#) [Zbl 1149.34003](#)
- [3] S. Albeverio, A. Kostenko, and M. Malamud, Spectral theory of semibounded Sturm–Liouville operators with local interactions on a discrete set. *J. Math. Phys.* **51** (2010), Article Id. 102102, 24 pp. [MR 2761288](#)
- [4] S. Albeverio and P. Kurasov, *Singular perturbations of differential operators*. Solvable Schrödinger type operators. London Mathematical Society Lecture Note Series, 271. Cambridge University Press, Cambridge, 2000. [MR 1752110](#) [Zbl 0945.47015](#)
- [5] G. D. Allen and F. J. Narcowich, *R-operators I. Representation Theory and Applications*, Indiana Univ. Math. J. **25**, 945–963 (1976).
- [6] N. Aronszajn and W. F. Donoghue, On exponential representations of analytic functions in the upper half-plane with positive imaginary part. *J. Analyse Math.* **12** (1964), 113–127. [MR 0168769](#) [Zbl 0138.29601](#)
- [7] M.-L. Baeteman and K. Chadan, The inverse scattering problem for singular oscillating potentials. *Nuclear Phys. A* **255** (1975), 35–44. [MR 0395544](#)
- [8] M.-L. Baeteman and K. Chadan, Scattering theory with highly singular oscillating potentials. *Ann. Inst. H. Poincaré Sect. A (N.S.)* **24** (1976), 1–16. [MR 0400975](#)
- [9] J.G. Bak and A. A. Shkalikov, Multipliers in dual Sobolev spaces and Schrödinger operators with distribution potentials. *Mat. Zametki* **71** (2002), 643–651. In Russian. English transl., *Math. Notes* **71** (2002), 587–594. [MR 1936189](#) [Zbl 1029.46029](#)

- [10] J. Ben Amara and A. A. Shkalikov, Oscillation theorems for Sturm–Liouville problems with distribution potentials. *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* **2009** (2009), 43–49. In Russian. English transl., *Moscow Univ. Math. Bull.* **64** (2009), 132–137. [MR 2664498](#)
- [11] A. Ben Amor and C. Remling, Direct and inverse spectral theory of one-dimensional Schrödinger operators with measures. *Integral Equations Operator Theory* **52** (2005), 395–417. [MR 2184572](#) [Zbl 1107.34064](#)
- [12] C. Bennewitz and W. N. Everitt, On second-order left-definite boundary value problems. In W. N. Everitt and R. T. Lewis (eds.), *Ordinary differential equations and operators*. Proceedings of a symposium held at the University of Dundee, Dundee, March–July 1982. Lecture Notes in Mathematics, 1032. Springer, Berlin etc., 1983, 31–67. [MR 0742634](#) [MR 0742630](#) (collection) [Zbl 0544.34015](#) [Zbl 0514.00013](#) (collection)
- [13] D. Bollé, F. Gesztesy, H. Grosse, W. Schweiger, and B. Simon, Witten index, axial anomaly, and Krein’s spectral shift function in supersymmetric quantum mechanics. *J. Math. Phys.* **28** (1987), 1512–1525. [MR 0894842](#) [Zbl 0643.47005](#)
- [14] T. Buckmaster and H. Koch, The Korteweg–de Vries equation at H^{-1} regularity. Preprint 2012. [arXiv:1112.4657](#) [math.AP]
- [15] R. W. Carey, A unitary invariant for pairs of self-adjoint operators. *J. Reine Angew. Math.* **283/284** (1976), 294–312. [MR 0415366](#) [Zbl 0348.47013](#)
- [16] S. Clark and F. Gesztesy, Weyl–Titchmarsh M -function asymptotics and Borg-type theorems for Dirac operators. *Trans. Amer. Math. Soc.* **354** (2002), 3475–3534. [MR 1911509](#) [Zbl 1017.34019](#)
- [17] S. Clark and F. Gesztesy, On Povzner–Wienholtz-type self-adjointness results for matrix-valued Sturm–Liouville operators. *Proc. Roy. Soc. Edinburgh Sect. A* **133** (2003), 747–758. [MR 2006200](#) [Zbl 1069.47052](#)
- [18] S. Clark, F. Gesztesy, and M. Zinchenko, Weyl–Titchmarsh theory and Borg–Marchenko-type uniqueness results for CMV operators with matrix-valued Verblunsky coefficients. *Oper. Matrices* **1** (2007), 535–592. [MR 2363977](#) [Zbl 1142.34014](#)
- [19] S. Clark, F. Gesztesy, and M. Zinchenko, Borg–Marchenko-type uniqueness results for CMV operators. *Skr. K. Nor. Vidensk. Selsk.* **2008** (2008), 1–18. [MR 2517327](#) [Zbl 1165.47021](#)
- [20] M. Combescure and J. Ginibre, Spectral and scattering theory for the Schrödinger operator with strongly oscillating potentials. *Ann. Inst. H. Poincaré Sect. A (N.S.)* **24** (1976), 17–30. [MR 0400976](#) [Zbl 0336.47007](#)
- [21] M. Combescure, Spectral and scattering theory for a class of strongly oscillating potentials. *Comm. Math. Phys.* **73** (1980), 43–62. [MR 0573612](#) [Zbl 0434.35072](#)
- [22] E. B. Davies, Singular Schrödinger operators in one dimension. *Mathematika* **59** (2013), 141–159. [MR 3028176](#) [Zbl 1282.34087](#)
- [23] P. A. Deift, Applications of a commutation formula. *Duke Math. J.* **45** (1978), 267–310. [MR 0495676](#) [Zbl 0392.47013](#)

- [24] P. Djakov and B. Mityagin, Spectral gap asymptotics of one-dimensional Schrödinger operators with singular periodic potentials. *Integral Transforms Spec. Funct.* **20** (2009), 265–273. [MR 2501800](#) [Zbl 1176.34108](#)
- [25] P. Djakov and B. Mityagin, Spectral gaps of Schrödinger operators with periodic singular potentials. *Dyn. PDE* **6**, no. 2, 95–165 (2009). *Dyn. Partial Differ. Equ.* **6** (2009), 95–165. [MR 2542499](#) [Zbl 1238.34151](#)
- [26] P. Djakov and B. Mityagin, Fourier method for one-dimensional Schrödinger operators with singular periodic potentials. In J. A. Ball, V. Bolotnikov, J. W. Helton, L. Rodman, and I. M. Spitkovsky (eds.), *Topics in operator theory*. Vol. 2. Systems and mathematical physics. A tribute to I. Gohberg on the occasion of his 80th birthday. Proceedings of the 19th International Workshop (IWOTA 2008) held at the College of William and Mary, Williamsburg, VA, July 22–26, 2008. Operator Theory: Advances and Applications, 203. Birkhäuser Verlag, Basel, 2010, 195–236. [MR 2683242](#) [Zbl 1200.34107](#) [Zbl 1181.47003](#) (collection)
- [27] P. Djakov and B. Mityagin, Criteria for existence of Riesz bases consisting of root functions of Hill and 1D Dirac operators. *J. Funct. Anal.* **263** (2012), 2300–2332. [MR 2964684](#) [Zbl 1263.34121](#)
- [28] J. Eckhardt, Inverse uniqueness results for Schrödinger operators using de Branges theory. *Complex Anal. Oper. Theory* **8** (2014), 37–50. [MR 3147711](#) [Zbl 06336119](#)
- [29] J. Eckhardt, F. Gesztesy, R. Nichols, and G. Teschl, Weyl–Titchmarsh theory for Sturm–Liouville operators with distributional potentials. *Opuscula Math.* **33** (2013), 467–563. [MR 3046408](#) [Zbl 1283.34022](#)
- [30] J. Eckhardt, F. Gesztesy, R. Nichols, and G. Teschl, Inverse spectral theory for Sturm–Liouville operators with distributional potentials. *J. Lond. Math. Soc.* (2) **88** (2013), 801–828. [MR 3145132](#) [Zbl 06243087](#)
- [31] J. Eckhardt and G. Teschl, Uniqueness results for one-dimensional Schrödinger operators with purely discrete spectra. *Trans. Amer. Math. Soc.* **365** (2013), 3923–3942. [MR 3042609](#) [Zbl 1286.34022](#)
- [32] J. Eckhardt and G. Teschl, Sturm–Liouville operators with measure-valued coefficients. *J. Anal. Math.* **120** (2013), 151–224. [MR 3095152](#) [Zbl 06247067](#)
- [33] D. E. Edmunds and W. D. Evans, *Spectral theory and differential operators*. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1987. [MR 0929030](#) [Zbl 0628.47017](#)
- [34] W. N. Everitt and L. Markus, *Boundary value problems and symplectic algebra for ordinary differential and quasi-differential operators*. Mathematical Surveys and Monographs, 61. American Mathematical Society, Providence, RI, 1999. [MR 1647856](#) [Zbl 0909.34001](#)
- [35] C. Frayer, R. O. Hryniv, Ya. V. Mykytyuk, and P. A. Perry, Inverse scattering for Schrödinger operators with Miura potentials: I. Unique Riccati representatives and ZS-AKNS systems. *Inverse Problems* **25** (2009), Article Id. 115007, 25 pp. [MR 2558667](#) [Zbl 1181.35330](#)

- [36] C. Fulton, Titchmarsh–Weyl m -functions for second order Sturm–Liouville problems. *Math. Nachr.* **281** (2008), 1418–1475. [MR 2454944](#) [Zbl 1165.34011](#)
- [37] C. Fulton and H. Langer, Sturm–Liouville operators with singularities and generalized Nevanlinna functions. *Complex Anal. Oper. Theory* **4** (2010), 179–243. [MR 2652523](#) [Zbl 1214.34022](#)
- [38] C. Fulton, H. Langer, and A. Luger, Mark Krein’s method of directing functionals and singular potentials. *Math. Nachr.* **285** (2012), 1791–1798. [MR 2988006](#) [Zbl 1259.34017](#)
- [39] F. Gesztesy, On the modified Korteweg–de Vries equation. In J. A. Goldstein, F. Kappel and W. Schappacher (eds.), *Differential equations with applications in biology, physics, and engineering*. Papers from the International Conference on Differential Equations and Applications held in Leibnitz, 1989. Lecture Notes in Pure and Applied Mathematics, 133. Marcel Dekker, New York, 1991, 139–183. [MR 1171468](#) [Zbl 0753.35078](#) [Zbl 0741.00056](#) (collection)
- [40] F. Gesztesy, Quasi-periodic, finite-gap solutions of the modified Korteweg–de Vries equation. In S. Albeverio, J. E. Fenstad, H. Holden, and T. Lindstrøm (eds.), *Ideas and methods in mathematical analysis, stochastics, and applications*. Vol. 1. In memory of R. Høegh-Krohn (1938–1988). Papers from the Symposium on Ideas and Methods in Mathematics and Physics held at the University of Oslo, Oslo, September 1988. Cambridge University Press, Cambridge, 1992, 428–471. [MR 1190516](#) [Zbl 0818.35103](#) [Zbl 0764.00003](#) (collection)
- [41] F. Gesztesy, A. Kiselev, and K. A. Makarov, Uniqueness results for matrix-valued Schrödinger, Jacobi, and Dirac-type operators. *Math. Nachr.* **239/240** (2002), 103–145. [MR 1905666](#) [Zbl 1017.34020](#)
- [42] F. Gesztesy, R. Nowell, and W. Pötz, One-dimensional scattering theory for quantum systems with nontrivial spatial asymptotics. *Differential Integral Equations* **10** (1997), 521–546. [MR 1744860](#) [Zbl 0894.34077](#)
- [43] F. Gesztesy, W. Schweiger, and B. Simon, Commutation methods applied to the mKdV-equation. *Trans. Amer. Math. Soc.* **324** (1991), 465–525. [MR 1029000](#) [Zbl 0728.35106](#)
- [44] F. Gesztesy and B. Simon, Constructing solutions of the mKdV-equation. *J. Funct. Anal.* **89** (1990), 53–60. [MR 1040953](#) [Zbl 0711.35121](#)
- [45] F. Gesztesy and B. Simon, On local Borg–Marchenko uniqueness results. *Comm. Math. Phys.* **211** (2000), 273–287. [MR 1754515](#) [Zbl 0985.34077](#)
- [46] F. Gesztesy, B. Simon, and G. Teschl, Spectral deformations of one-dimensional Schrödinger operators. *J. Anal. Math.* **70** (1996), 267–324. [MR 1444263](#) [Zbl 0951.34061](#)
- [47] F. Gesztesy and E. Tsekanovskii, On matrix-valued Herglotz functions. *Math. Nachr.* **218** (2000), 61–138. [MR 1784638](#) [Zbl 0961.30027](#)
- [48] F. Gesztesy and R. Weikard, Some remarks on the spectral problem underlying the Camassa–Holm hierarchy. In J. A. Ball, M. A. Dritschel, A. F. M. ter Elst, P. Portal, and D. Potapov (eds.), *Operator Theory in Harmonic and Non-commutative*

- Analysis, IWOTA 12.* Operator Theory: Advances and Applications, 240, Birkhäuser, Basel, 2014, 137–188.
- [49] F. Gesztesy, R. Weikard, and M. Zinchenko, On spectral theory for Schrödinger operators with operator-valued potentials. *J. Differential Equations* **255** (2013), 1784–1827. [MR 3072671](#) [Zbl 1291.34139](#)
 - [50] F. Gesztesy and M. Zinchenko, On spectral theory for Schrödinger operators with strongly singular potentials. *Math. Nachr.* **279** (2006), 1041–1082. [MR 2242965](#) [Zbl 1108.34063](#)
 - [51] A. S. Goriunov and V. A. Mikhailets, Resolvent convergence of Sturm–Liouville operators with singular potentials. *Mat. Zametki* **87** (2010), 311–315. In Russian. English transl., *Math. Notes* **87** (2010), 287–292. [MR 2731483](#) [Zbl 1205.34120](#)
 - [52] A. Goriunov and V. Mikhailets, Regularization of singular Sturm–Liouville equations. *Methods Funct. Anal. Topology* **16** (2010), 120–130. [MR 2667807](#) [Zbl 1221.47083](#)
 - [53] S. Grudsky and A. Rybkin, On positive type initial profiles for the KdV equation. *Proc. Amer. Math. Soc.* **142** (2014), 2079–2086. [MR 3182026](#) [Zbl 1294.35122](#)
 - [54] O. Hald, Discontinuous inverse eigenvalue problems. *Comm. Pure Appl. Math.* **37** (1984), 539–577. [MR 0752591](#) [Zbl 0541.34012](#)
 - [55] V. Hardt, A. Konstantinov, and R. Mennicken, On the spectrum of the product of closed operators. *Math. Nachr.* **215** (2000), 91–102. [MR 1768195](#) [Zbl 0965.47002](#)
 - [56] J. Herczyński, On Schrödinger operators with distributional potentials. *J. Operator Theory* **21** (1989), 273–295. [MR 1023316](#) [Zbl 0728.35026](#)
 - [57] D. B. Hinton and A. Schneider, On the Titchmarsh–Weyl coefficients for singular S -Hermitian Systems. I. *Math. Nachr.* **163** (1993), 323–342. [MR 1235076](#) [Zbl 0806.34023](#)
 - [58] D. B. Hinton and A. Schneider, On the Titchmarsh–Weyl coefficients for singular S -Hermitian Systems. II. *Math. Nachr.* **185** (1997), 67–84. [MR 1452477](#) [Zbl 0869.34019](#)
 - [59] D. B. Hinton and J. K. Shaw, On Titchmarsh–Weyl $M(\lambda)$ -functions for linear Hamiltonian systems. *J. Differential Equations* **40** (1981), 316–342. [MR 0621248](#) [Zbl 0472.34014](#)
 - [60] D. B. Hinton and J. K. Shaw, On the spectrum of a singular Hamiltonian system. *Quaestiones Math.* **5** (1982/83), 29–81. [MR 0644777](#) [Zbl 0509.34018](#)
 - [61] D. B. Hinton and J. K. Shaw, Hamiltonian systems of limit point or limit circle type with both endpoints singular. *J. Differential Equations* **50** (1983), 444–464. [MR 0723581](#) [Zbl 0476.34027](#)
 - [62] D. B. Hinton and J. K. Shaw, On boundary value problems for Hamiltonian systems with two singular points. *SIAM J. Math. Anal.* **15** (1984), 272–286. [MR 0731867](#) [Zbl 0539.34016](#)
 - [63] D. B. Hinton and J. K. Shaw, On the spectrum of a singular Hamiltonian system. II. *Quaestiones Math.* **10** (1986), 1–48. [MR 0857767](#) [Zbl 0615.34016](#)

- [64] R. O. Hryniv, Analyticity and uniform stability in the inverse singular Sturm–Liouville spectral problem. *Inverse Problems* **27** (2011), Article Id. 065011, 25 pp. [MR 2805768](#) [Zbl 1226.34012](#)
- [65] R. O. Hryniv and Ya. V. Mykytyuk, 1-D Schrödinger operators with periodic singular potentials. *Methods Funct. Anal. Topology* **7** (2001), 31–42. [MR 1879483](#) [Zbl 0992.47021](#)
- [66] R. O. Hryniv and Ya. V. Mykytyuk, 1-D Schrödinger operators with singular Gordon potentials. *Methods Funct. Anal. Topology* **8** (2002), 36–48. [MR 1903128](#) [Zbl 1013.34081](#)
- [67] R. O. Hryniv and Ya. V. Mykytyuk, Inverse spectral problems for Sturm–Liouville operators with singular potentials. *Inverse Problems* **19** (2003), 665–684. [MR 1984883](#) [Zbl 1074.34007](#)
- [68] R. O. Hryniv and Ya. V. Mykytyuk, Half-inverse spectral problems for Sturm–Liouville operators with singular potentials. *Inverse Problems* **20** (2004), 1423–1444. [MR 2109127](#) [Zbl 1074.34007](#)
- [69] R. O. Hryniv and Ya. V. Mykytyuk, Transformation operators for Sturm–Liouville operators with singular potentials. *Int. J. Pure Appl. Math.* **12** (2004), 311–334. [MR 2049448](#) [Zbl 1070.34041](#)
- [70] R. O. Hryniv and Ya. V. Mykytyuk, Inverse spectral problems for Sturm–Liouville operators with singular potentials. IV. Potentials in the Sobolev space scale. *Proc. Edinb. Math. Soc. (2)* **49** (2006), 309–329. [MR 2243789](#) [Zbl 1105.34004](#)
- [71] R. O. Hryniv and Ya. V. Mykytyuk, Eigenvalue asymptotics for Sturm–Liouville operators with singular potentials. *J. Funct. Anal.* **238** (2006), 27–57. [MR 2234122](#) [Zbl 1108.34021](#)
- [72] R. O. Hryniv and Ya. V. Mykytyuk, Self-adjointness of Schrödinger operators with singular potentials. *Methods Funct. Anal. Topology* **18** (2012), 152–159. [MR 2978191](#) [Zbl 1262.47066](#)
- [73] R. O. Hryniv, Ya. V. Mykytyuk, and P. A. Perry, Inverse scattering for Schrödinger operators with Miura potentials. II. Different Riccati representatives. *Comm. Partial Differential Equations* **36** (2011), 1587–1623. [MR 2825604](#) [Zbl 1232.34113](#)
- [74] R. O. Hryniv, Ya. V. Mykytyuk, and P. A. Perry, Sobolev mapping properties of the scattering transform for the Schrödinger equation. In M. Braverman, L. Friedlander, Th. Kappeler, P. Kuchment, P. Topalov, and J. Weitsman (eds.), *Spectral theory and geometric analysis*. Papers from the International Conference in Honor of M. Shubin’s 65th Birthday held at Northeastern University, Boston, MA, July 29–August 2, 2009. Contemporary Mathematics, 535. American Mathematical Society, Providence, R.I., 2011, 79–93. [MR 2560752](#) [MR 2848603](#) (collection) [Zbl 1221.34230](#) [Zbl 1207.58001](#) (collection)
- [75] I. S. Kac, The existence of spectral functions of generalized second order differential systems with boundary conditions at the singular end. *Mat. Sb. (N.S.)* **68** (110) (1965), 174–227. In Russian. English transl., *AMS Translations* (2) **62**, 204–262. [MR 0209551](#) [Zbl 0144.10003](#)

- [76] T. Kappeler and C. Möhr, Estimates for periodic and Dirichlet eigenvalues of the Schrödinger operator with singular potentials. *J. Funct. Anal.* **186** (2001), 62–91. [MR 1863292](#) [Zbl 1004.34077](#)
- [77] T. Kappeler, P. Perry, M. Shubin, and P. Topalov, The Miura map on the line. *Int. Math. Res. Not.* **2005** (2005), 3091–3133. [MR 2189502](#) [Zbl 1089.35058](#)
- [78] T. Kappeler and P. Topalov, Global fold structure of the Miura map on $L^2(\mathbb{T})$. *Int. Math. Res. Not.* **2004** (2004), 2039–2068. [MR 2062735](#) [Zbl 1076.35111](#)
- [79] T. Kappeler and P. Topalov, Global well-posedness of mKdV in $L^2(\mathbb{T}, \mathbb{R})$. *Comm. Partial Differential Equations* **30** (2005), 435–449. [MR 2131061](#) [Zbl 1080.35119](#)
- [80] T. Kappeler and P. Topalov, Global wellposedness of KdV in $H^{-1}(\mathbb{T}, \mathbb{R})$. *Duke Math. J.* **135** (2006), 327–360. [MR 2267286](#) [Zbl 1106.35081](#)
- [81] M. Kato, Estimates of the eigenvalues of Hill’s operator with distributional coefficients. *Tokyo J. Math.* **33** (2010), 361–364. [MR 2779262](#) [Zbl 1214.47037](#)
- [82] T. Kato, *Perturbation theory for linear operators.*, Corrected printing of the 2nd ed. Grundlehren der mathematischen Wissenschaften, 132. Springer, Berlin etc., 1980. [MR 0407617](#) (2nd ed.) [Zbl Zbl 0435.47001](#)
- [83] K. Kodaira, The eigenvalue problem for ordinary differential equations of the second order and Heisenberg’s theory of S -matrices. *Amer. J. Math.* **71** (1949), 921–945. [MR 0033421](#) [Zbl 0035.27101](#)
- [84] E. Korotyaev, Characterization of the spectrum of Schrödinger operators with periodic distributions. *Int. Math. Res. Not.* **2003** (2003), 2019–2031. [MR 1995145](#) [Zbl 1104.34059](#)
- [85] E. Korotyaev, Sharp asymptotics of the quasimomentum. *Asymptot. Anal.* **80** (2012), 269–287. [MR 3025046](#) [Zbl 1267.34148](#)
- [86] A. S. Kostenko and M. M. Malamud, One-dimensional Schrödinger operator with δ -interactions. *Funktional. Anal. i Prilozhen.* **44** (2010), 87–91. In Russian. English transl., *Funct. Anal. Appl.* **44** (2010), 151–155. [MR 2681961](#) [Zbl 1210.47068](#)
- [87] A. S. Kostenko and M. M. Malamud, 1-D Schrödinger operators with local point interactions on a discrete set. *J. Differential Equations* **249** (2010), 253–304. [MR 2644117](#) [Zbl 1195.47031](#)
- [88] A. Kostenko, A. Sakhnovich, and G. Teschl, Inverse eigenvalue problems for perturbed spherical Schrödinger operators. *Inverse Problems* **26** (2010), Article Id. 105013, 14 pp. [MR 2719774](#) [Zbl 1227.34023](#)
- [89] A. Kostenko, A. Sakhnovich, and G. Teschl, Weyl–Titchmarsh theory for Schrödinger operators with strongly singular potentials. *Int. Math. Res. Not. IMRN* **2012** (2012), 1699–1747. [MR 2920828](#) [Zbl 1248.34027](#)
- [90] A. Kostenko, A. Sakhnovich, and G. Teschl, Commutation methods for Schrödinger operators with strongly singular potentials. *Math. Nachr.* **285** (2012), 392–410. [MR 2899633](#) [Zbl 1250.34069](#)

- [91] A. Kostenko and G. Teschl, On the singular Weyl–Titchmarsh function of perturbed spherical Schrödinger operators. *J. Differential Equations* **250** (2011), 3701–3739. [MR 2773183](#) [Zbl 1219.34037](#)
- [92] A. Kostenko and G. Teschl, Spectral asymptotics for perturbed spherical Schrödinger operators and applications to quantum scattering. *Comm. Math. Phys.* **322** (2013), 255–275. [MR 3073165](#) [Zbl 1284.34130](#)
- [93] S. Kotani and B. Simon, Stochastic Schrödinger operators and Jacobi matrices on the strip. *Comm. Math. Phys.* **119** (1988), 403–429. [MR 0969209](#) [Zbl 0969209](#)
- [94] P. Kurasov and A. Luger, An operator theoretic interpretation of the generalized Titchmarsh–Weyl coefficient for a singular Sturm–Liouville problem. *Math. Phys. Anal. Geom.* **14** (2011), 115–151. [MR 2802457](#) [Zbl 1267.34052](#)
- [95] M. Lesch and M. Malamud, The inverse spectral problem for first order systems on the half line. In V. M. Adamyan, I. Gohberg, M. Gorbachuk, V. Gorbachuk, M. A. Kaashoek, H. Langer, and G. Popov (eds.), *Differential operators and related topics*. Proceedings of the International Conference dedicated to the 90th anniversary of the birth of Mark Krein held in Odessa, August 18–22, 1997. Operator Theory: Advances and Applications, 117. Birkhäuser, Basel, 199–238. [MR 1764962](#)
[MR 1764947](#) (collection) [Zbl 0956.34072](#) [Zbl 1051.47001](#) (collection)
- [96] M. Lesch and M. Malamud, On the deficiency indices and self-adjointness of symmetric Hamiltonian systems. *J. Differential Equations* **189** (2003), 556–615. [MR 1964480](#) [Zbl 1016.37026](#)
- [97] B. M. Levitan and M. Otelbaev, Conditions for selfadjointness of the Schrödinger and Dirac operators. *Trudy Moskov. Mat. Obshch.* **42** (1981), 142–159. In Russian. English transl., *Trans. Moscow. Math. Soc.* **42** (1982), 139–156. [MR 0621997](#)
[Zbl 0525.35025](#)
- [98] V. G. Maz'ya and T. O. Shaposhnikova, *Theory of Sobolev multipliers*. With applications to differential and integral operators. Grundlehren der Mathematischen Wissenschaften, 337. Springer, Berlin etc., 2009. [MR 2457601](#) [Zbl 1157.46001](#)
- [99] V. G. Maz'ya and I. E. Verbitsky, Boundedness and compactness criteria for the one-dimensional Schrödinger operator. In M. Cwikel, M. Engliš, A. Kufner, L.-E. Persson, and G. Sparr, *Function spaces, interpolation theory and related topics*. Proceedings of the International Conference in honour of J. Peetre on his 65th birthday, held at Lund University, Lund, August 17–22, 2000. De Gruyter, Berlin, 2002, 369–382. [MR 1943294](#) [MR 1943279](#) (collection) [Zbl 1034.34101](#) [Zbl 0996.00036](#) (collection)
- [100] V. G. Maz'ya and I. E. Verbitsky, The Schrödinger operator on the energy space: boundedness and compactness criteria. *Acta Math.* **188** (2002), 263–302. [MR 1947894](#) [Zbl 1013.35021](#)
- [101] V. G. Maz'ya and I. E. Verbitsky, Infinitesimal form boundedness and Trudinger's subordination for the Schrödinger operator. *Invent. Math.* **162** (2005), 81–136. [MR 2198326](#) [Zbl 1132.35353](#)

- [102] V. G. Maz'ya and I. E. Verbitsky, Form boundedness of the general second-order differential operator. *Comm. Pure Appl. Math.* **59** (2006), 1286–1329. [MR 2237288](#) [Zbl 1169.35014](#)
- [103] V. A. Mikhailets and V. M. Molyboga, Singular eigenvalue problems on the circle. *Methods Funct. Anal. Topology* **10** (2004), 44–53. [MR 2092532](#) [Zbl 1077.47049](#)
- [104] V. A. Mikhailets and V. M. Molyboga, Uniform estimates for the semi-periodic eigenvalues of the singular differential operators. *Methods Funct. Anal. Topology* **10** (2004), 30–57. [MR 2109216](#) [Zbl 1072.34097](#)
- [105] V. A. Mikhailets and V. M. Molyboga, Singularly perturbed periodic and semiperiodic differential operators. *Ukrain. Mat. Zh.* **59** (2007), 785–797, and *Ukrainian Math. J.* **59** (2007), 858–873. [MR 2367920](#) [Zbl 1150.47032](#)
- [106] V. A. Mikhailets and V. M. Molyboga, One-dimensional Schrödinger operators with singular periodic potentials. *Methods Funct. Anal. Topology* **14** (2008), 184–200. [MR 2432767](#) [Zbl 1164.47047](#)
- [107] V. A. Mikhailets and V. M. Molyboga, Spectral gaps of the one-dimensional Schrödinger operators with singular periodic potentials. *Methods Funct. Anal. Topology* **15**, (2009), 31–40. [MR 2502636](#) [Zbl 1199.34453](#)
- [108] V. A. Mikhailets and V. M. Molyboga, Remarks on Schrödinger operators with singular matrix potentials. *Methods Funct. Anal. Topology* **19** (2013), 161–167. [MR 3098494](#) [Zbl 1289.34239](#)
- [109] K. A. Mirzoev and T. A. Safanova, Singular Sturm–Liouville operators with distribution potential on spaces of vector functions. *Dokl. Akad. Nauk* **441** (2011), 165–168. In Russian. English transl., *Dokl. Math.* **84** (2011), 791–794. [MR 2953790](#) [Zbl 1244.47039](#)
- [110] Ya. V. Mykytyuk and N. S. Trush, Inverse spectral problems for Sturm–Liouville operators with matrix-valued potentials. *Inverse Problems* **26**, Article. Id. 015009 (2010), 36 pp. [MR 575354](#) [Zbl 1195.34022](#)
- [111] M. A. Naimark, *Linear differential operators*. Part II: Linear differential operators in Hilbert space. With additional material by the author, and a supplement by V. È. Ljance. Translated from the Russian by E. R. Dawson. English transl. edited by W. N. Everitt. Frederick Ungar, New York, 1968 [MR 0262880](#) [Zbl 0227.34020](#)
- [112] D. B. Pearson, Scattering theory for a class of oscillating potentials. *Helv. Phys. Acta* **52** (1979), 541–5554. [MR 0566255](#)
- [113] F. S. Rofe-Beketov and E. H. Khristov, Transformation operators and scattering functions for a highly singular potential. *Dokl. Akad. Nauk SSSR* **168** (1966), 1265–1268. In Russian. English transl., *Sov. Math. Dokl.* **7** (1966), 834–837. [MR 0194652](#) [Zbl 0237.34039](#)
- [114] F. S. Rofe-Beketov and E. H. Khristov, Some analytical questions and the inverse Sturm–Liouville problem for an equation with highly singular potential. *Dokl. Akad. Nauk SSSR* **185** (1969), 768–771. In Russian. English transl., *Sov. Math. Dokl.* **10** (1969), 432–435. [MR 0245891](#) [Zbl 0191.10103](#)

- [115] A. Rybkin, Regularized perturbation determinants and KdV conservation laws for irregular initial profiles. In J. A. Ball, V. Bolotnikov, J. W. Helton, L. Rodman, and I. M. Spitkovsky (eds.), *Topics in operator theory*. Vol. 2. Systems and mathematical physics. A tribute to I. Gohberg on the occasion of his 80th birthday. Proceedings of the 19th International Workshop (IWOTA 2008) held at the College of William and Mary, Williamsburg, VA, July 22–26, 2008. Operator Theory: Advances and Applications, 203. Birkhäuser Verlag, Basel, 2010, 427–444. [MR 2683250](#) [Zbl 1191.35235](#) [Zbl 1181.47003](#) (collection)
- [116] I. V. Sadovnichaya, Equiconvergence of expansions in series in eigenfunctions of Sturm–Liouville operators with distribution potentials. *Mat. Sb.* 201 (2010), 61–76. In Russian. English transl., *Sborn. Math.* **201** (2010), 1307–1322. [MR 2760460](#) [Zbl 1143.34324](#)
- [117] I. V. Sadovnichaya, Equiconvergence in Sobolev and Hölder spaces of expansions in eigenfunctions of Sturm–Liouville operators with distribution potentials. *Dokl. Akad. Nauk* **437** (2011), 162–163. In Russian. English transl., *Dokl. Math.* **83** (2011), 169–170. [MR 2849348](#) [Zbl 1252.34096](#)
- [118] A. Sakhnovich, Dirac type and canonical systems: spectral and Weyl–Titchmarsh matrix functions, direct and inverse problems. *Inverse Probl.* **18** (2002), 331–348. [MR 1910241](#) [Zbl 1009.34079](#)
- [119] A. Sakhnovich, Skew-self-adjoint discrete and continuous Dirac-type systems: inverse problems and Borg–Marchenko theorems. *Inverse Problems* **22** (2006), 2083–2101. [MR 2277531](#) [Zbl 1105.47035](#)
- [120] A. L. Sakhnovich, L. A. Sakhnovich, and I. Ya. Roitberg, *Inverse problems and nonlinear evolution equations*. Solutions, Darboux matrices and Weyl–Titchmarsh functions. De Gruyter Studies in Mathematics, 47. De Gruyter, Berlin, 2013. [MR 3098432](#) [Zbl 1283.47003](#)
- [121] A. M. Savchuk and A. A. Shkalikov, Sturm–Liouville operators with singular potentials. *Mat. Zametki* **66** (1999), 897–912. In Russian. English transl., *Math. Notes* **66** (1999), 741–753. [MR 1756602](#) [Zbl 0968.34072](#)
- [122] A. M. Savchuk and A. A. Shkalikov, The trace formula for Sturm–Liouville operators with singular potentials. *Mat. Zametki* **69** (2001), 427–442. In Russian. English transl., *Math. Notes* **69** (2001), 387–400. [MR 1846840](#) [Zbl 0996.34023](#)
- [123] A. M. Savchuk and A. A. Shkalikov, Sturm–Liouville operators with distribution potentials. *Mosk. Mat. Obs.* **64** (2003), 159–212. In Russian. English transl., *Trans. Moscow Math. Soc.* **2003**, 143–192. [MR 2030189](#) [Zbl 1066.34085](#)
- [124] A. M. Savchuk and A. A. Shkalikov, Inverse problem for Sturm–Liouville operators with distribution potentials: reconstruction from two spectra. *Russ. J. Math. Phys.* **12** (2005), 507–514. [MR 2201315](#) [Zbl 05072519](#)
- [125] A. M. Savchuk and A. A. Shkalikov, On the eigenvalues of the Sturm–Liouville operator with potentials from Sobolev spaces. *Mat. Zametki* **80** (2006), 864–884. In Russian. English transl., *Math. Notes* **80** (2006), 814–832. [MR 2311614](#) [Zbl 1129.34055](#)

- [126] A. M. Savchuk and A. A. Shkalikov, On the properties of maps connected with inverse Sturm–Liouville problems. *Tr. Mat. Inst. Steklova* **260** (2008), *Teor. Funkts. i Nelinein. Uravn. v Chastn. Proizvodn.*, 227–247. In Russian. English transl., *Proc. Steklov Inst. Math.* **260** (2008), 218–237. [MR 2489515](#) [Zbl 1233.34010](#)
- [127] A. M. Savchuk and A. A. Shkalikov, Inverse problems for Sturm–Liouville operators with potentials in Sobolev spaces: uniform stability. *Funktional. Anal. i Prilozhen.* **44** (2010), 34–53. In Russian. English transl., *Funct. Anal. Appl.* **44** (2010), 270–285. [MR 2768563](#) [Zbl 1271.34017](#)
- [128] M. Shahriari, A. J. Akbarfam, and G. Teschl, Uniqueness for inverse Sturm–Liouville problems with a finite number of transmission conditions. *J. Math. Anal. Appl.* **395** (2012), 19–29. [MR 2943599](#) [Zbl 1251.34025](#)
- [129] D. Shin, On quasi-differential operators in Hilbert space. C. R. (Doklady) Acad. Sci. URSS (N.S.) **18** (1938), 523–526. In Russian. [Zbl 0019.21404](#)
- [130] D. Shin, Über die Lösungen einer quasi-Differentialgleichung der n -ten Ordnung. *Rec. Math. [Mat. Sbornik] N. S.* **7** (**49**) (1940). 479–532. In Russian. [MR 0002428](#) [JFM 66.0398.01](#) [Zbl 0023.31701](#)
- [131] D. Shin, Quasi-differential operators in Hilbert space. *Rec. Math. [Mat. Sbornik] N. S.* **13** (**55**) (1943), 39–70. In Russian. [MR 0011534](#) [Zbl 0061.26108](#)
- [132] G. Teschl, *Mathematical methods in quantum mechanics*. With applications to Schrödinger operators. 2nd ed. Graduate Studies in Mathematics, 157. American Mathematical Society, Providence, RI, 2014.
- [133] B. Thaller, Normal forms of an abstract Dirac operator and applications to scattering theory. *J. Math. Phys.* **29** (1988), 249–257. [MR 0921790](#) [Zbl 0639.47043](#)
- [134] B. Thaller, *The Dirac equation*. Texts and Monographs in Physics. Springer, Berlin etc., 1992. [MR 1219537](#) [Zbl 0765.47023](#)
- [135] J. Weidmann, *Spectral theory of ordinary differential operators*. Lecture Notes in Mathematics, 1258. Springer, Berlin etc., 1987. [MR 0923320](#) [Zbl 0647.47052](#)
- [136] A. Zettl, Formally self-adjoint quasi-differential operators. *Rocky Mountain J. Math.* **5** (1975), 453–474. [MR 0379976](#) [Zbl 1293.47043](#)

Received 2013 June, 18; revised 2013 October, 14

Jonathan Eckhardt, Faculty of Mathematics, University of Vienna,
Oskar-Morgenstern-Platz 1, 1090 Wien, Austria

home page: <http://homepage.univie.ac.at/jonathan.eckhardt/>

e-mail: jonathan.eckhardt@univie.ac.at

Fritz Gesztesy, Department of Mathematics, University of Missouri, Columbia,
MO 65211, U.S.A.

home page: <http://www.math.missouri.edu/personnel/faculty/gesztesyf.html>

e-mail: gesztesyf@missouri.edu

Roger Nichols, Mathematics Department,
The University of Tennessee at Chattanooga, 415 EMCS Building, Dept. 6956,
615 McCallie Ave, Chattanooga, TN 37403, U.S.A.

home page: <http://www.utc.edu/faculty/roger-nichols/>

e-mail: Roger-Nichols@utc.edu

Gerald Teschl, Faculty of Mathematics, University of Vienna,
Oskar-Morgenstern-Platz 1, 1090 Wien, Austria

Erwin Schrödinger Institute for Mathematical Physics, Boltzmanngasse 9, 1090 Wien,
Austria

home page: <http://www.mat.univie.ac.at/~gerald/>

e-mail: Gerald.Teschl@univie.ac.at