

Asymptotics of the number of the interior transmission eigenvalues

Vesselin Petkov¹ and Georgi Vodev

Abstract. We prove Weyl asymptotics $N(r) = cr^d + \mathcal{O}_\epsilon(r^{d-\kappa+\epsilon})$, for all $0 < \epsilon \ll 1$, for the counting function $N(r) = \#\{\lambda_j \in \mathbb{C} \setminus \{0\}: |\lambda_j| \leq r^2, r > 1\}$, of the interior transmission eigenvalues (ITE), λ_j . Here $d \geq 2$ denotes the space dimension and $0 < \kappa \leq 1$ is such that there are no (ITE) in the region $\{\lambda \in \mathbb{C}: |\operatorname{Im} \lambda| \geq C(|\operatorname{Re} \lambda| + 1)^{1-\frac{\kappa}{2}}\}$ for some $C > 0$.

Mathematics Subject Classification (2010). Primary: 35P20; Secondary: 35P15, 35P25.

Keywords. Interior transmission eigenvalues, Weyl formula with remainder, eigenvalue-free regions.

Contents

1	Introduction and statement of results	2
2	Parametrix of the Dirichlet–to–Neumann map in the elliptic zone	7
3	Analysis of the transmission eigenvalues	13
	References	29

¹ The first author was partially supported by the ANR project Nosevol BS01019 01.

1. Introduction and statement of results

Let $\Omega \subset \mathbb{R}^d$, $d \geq 2$, be a bounded, connected domain with a C^∞ smooth boundary $\Gamma = \partial\Omega$. A complex number $\lambda \in \mathbb{C}$, $\lambda \neq 0$, will be called an interior transmission eigenvalue (ITE) if the following problem has a non-trivial solution:

$$\begin{cases} (\nabla c_1(x)\nabla + \lambda n_1(x)) u_1 = 0 & \text{in } \Omega, \\ (\nabla c_2(x)\nabla + \lambda n_2(x)) u_2 = 0 & \text{in } \Omega, \\ u_1 = u_2, \quad c_1 \partial_\nu u_1 = c_2 \partial_\nu u_2 & \text{on } \Gamma, \end{cases} \quad (1.1)$$

where ν denotes the exterior Euclidean unit normal to Γ , $c_j, n_j \in C^\infty(\bar{\Omega})$, $j = 1, 2$ are strictly positive real-valued functions. The spectral problem for (ITE) is related to a non self-adjoint operator \mathcal{A} (see Section 3) and in the isotropic case $c_1(x) = c_2(x) = 1$ the boundary problem (1.1) is not parameter-elliptic. For these reasons many well-known techniques developed for self-adjoint operators or for parameter-elliptic boundary problems are not applicable. The positive (ITE) are related to the inverse scattering problems. More precisely, if $\lambda = k^2$ is a real (ITE), then the far-field operator $F(\lambda): L^2(\mathbb{S}^{d-1}) \rightarrow L^2(\mathbb{S}^{d-1})$ with kernel the scattering amplitude $s(k, \theta, \omega)$ is not injective and its range is not dense. This is crucial for the so-called linear sampling method (see [5], [1]) which works if we avoid the real (ITE). For this reason the problem of the existence and the discreteness of (ITE) draw the attention of many authors (see the survey [3] for a comprehensive review and a more complete list of references). Secondly, it was proved that we can determine the (ITE) from the far-field operator. Finally, it was established that in some cases the knowledge of all complex (ITE) determines the index of refraction of the scattering obstacle (see [3], [6]). This explains the increasing interest toward (ITE) and the fact that a lot of papers concerning the existence and the spectral properties of (ITE) in relation with the inverse scattering problems of reconstruction have been recently published.

On the other hand, the analysis of the (ITE) leads to some interesting and difficult mathematical spectral problems for non self-adjoint operators. These problems are connected with two *major questions*:

- (A) describe the eigenvalue-free regions in the complex plane;
- (B) find a Weyl asymptotic of the counting function of the eigenvalues.

In contrast to the case of self-adjoint operators these questions are much more difficult and there are no general results. As far as the Weyl asymptotics are concerned, one may study the leading term of the counting function and one can search an optimal remainder. On the other hand, even in the case of boundary

problems for non self-adjoint operators which are parameter-elliptic, the Weyl asymptotics in the literature concern mainly the leading term (see [2] for some results in this direction for non self-adjoint operators).

Question (A) has been investigated by the second author in [27] (see also [10] for a weaker result) and the result in [27] plays an important role in our analysis. In the present paper our purpose is to study question (B). Under some conditions the (ITE) form a discrete set in $\mathbb{C} \setminus \{0\}$ and they have as an accumulation point only infinity (see for instance [11], [24]). Introduce the counting function

$$N(r) := \#\{\lambda_j \in \mathbb{C} \setminus \{0\} : \lambda_j \text{ is (ITE), } |\lambda_j| \leq r^2\}, \quad r > 1,$$

where the eigenvalues are counted with their multiplicity (see Section 3 for the precise definition of the multiplicity). Recently, many works concerning the Weyl asymptotics of $N(r)$ have been published both in the isotropic ($c_1 \equiv c_2 \equiv 1$) and anisotropic cases (see [19], [8], [20], [13], [15], [16], [18], and [9]). In [18] the case when $\Omega = \{x \in \mathbb{R}^d : |x| \leq 1\}$ and $c_1 \equiv c_2 \equiv 1$, $n_1 \equiv 1$, $n_2 = \text{const} \neq 1$ has been investigated and for $d = 1$ an asymptotics of $N(r)$ with remainder has been established (see also [25]). In all other works only the leading term of $N(r)$ was obtained. We should mention that in [13] the anisotropic case has been studied and the asymptotics of $N(r)$ with a remainder is stated. However, the proof has a gap and only the asymptotics with leading term seems to be correct. The isotropic case is more difficult since the boundary problem is not parameter-elliptic and the tools for elliptic boundary problems cannot be applied. In the isotropic case when $n_1(x) \equiv 1$, $n_2(x) > 1$, for all $x \in \bar{\Omega}$, it has been recently established in [9] and [20] the asymptotics

$$N(r) \sim (\tau_1 + \tau_2)r^d, \quad r \rightarrow +\infty, \quad (1.2)$$

where τ_1 and τ_2 are defined below. It is important to remark that in [9] and [20] the analysis is based on the study of some trace class operators leading to an asymptotics

$$\sum_j \frac{1}{|\lambda_j|^p + t} = \alpha t^{-1+\frac{d}{2p}} + o(t^{-1+\frac{d}{2p}}), \quad t \rightarrow +\infty, \quad (1.3)$$

where $p \in \mathbb{N}$ is sufficiently large. Combining this asymptotics with the Tauberian theorem of Hardy and Littlewood, one obtains (1.2) and the remainder is given by the principal part divided by a logarithmic factor. To obtain a sharper remainder one could apply a finer Tauberian theorem (see [17]), but for this purpose it is necessary to establish asymptotics like (1.3) with sharper remainder for t lying on certain parabola in \mathbb{C} . This, however, seems to be a very difficult problem.

In the present work we follow another approach inspired by the paper [4], where asymptotics have been established for the number of the resonances associated to an exterior transmission boundary problem. The purpose is to study the asymptotic behavior of $N(r)$ under the condition

$$c_1(x)n_1(x) \neq c_2(x)n_2(x), \quad \text{for all } x \in \Gamma. \quad (1.4)$$

Our main result is the following

Theorem 1.1. *Assume (1.4) fulfilled. Assume also either the condition*

$$c_1(x) = c_2(x), \quad \partial_\nu c_1(x) = \partial_\nu c_2(x), \quad \text{for all } x \in \Gamma, \quad (1.5)$$

or the condition

$$c_1(x) \neq c_2(x), \quad \text{for all } x \in \Gamma. \quad (1.6)$$

Then, the (ITE) form a discrete set in \mathbb{C} and we have the asymptotics

$$N(r) = (\tau_1 + \tau_2)r^d + \mathcal{O}_\varepsilon(r^{d-\kappa+\varepsilon}), \quad r \rightarrow +\infty, \quad (1.7)$$

for every $0 < \varepsilon \ll 1$, where

$$\tau_j = \frac{\omega_d}{(2\pi)^d} \int_\Omega \left(\frac{n_j(x)}{c_j(x)} \right)^{d/2} dx,$$

ω_d being the volume of the unit ball in \mathbb{R}^d , and $\kappa = \frac{1}{2}$ if (1.5) holds, $\kappa = \frac{2}{5}$ if (1.6) holds. Moreover, if in addition to (1.6) we assume either the condition

$$\frac{n_1(x)}{c_1(x)} \neq \frac{n_2(x)}{c_2(x)}, \quad \text{for all } x \in \Gamma, \quad (1.8)$$

or the condition

$$\frac{n_1(x)}{c_1(x)} = \frac{n_2(x)}{c_2(x)}, \quad \text{for all } x \in \Gamma, \quad (1.9)$$

then (1.7) holds with $\kappa = \frac{1}{2}$.

To prove this theorem we use in an essential way the eigenvalue-free regions obtained in [27]. In fact, we prove in the present paper a *more general result* saying that if there are no interior transmission eigenvalues in a region of the form

$$\{\lambda \in \mathbb{C}: |\operatorname{Im} \lambda| \geq C(|\operatorname{Re} \lambda| + 1)^{1-\frac{\kappa}{2}}\}, \quad C > 0, \quad 0 < \kappa \leq 1, \quad (1.10)$$

then the asymptotics (1.7) with remainder $\mathcal{O}_\varepsilon(r^{d-\kappa+\varepsilon})$ is true. On the other hand, it is proved in [27] that under the assumptions of Theorem 1.1, we have indeed

an eigenvalue-free region (1.10) with κ replaced by $\kappa - \epsilon$, where κ is given by Theorem 1.1. Note that the parametrix construction and the results concerning the Dirichlet-to-Neumann map in [22], Section 11, suggest that for strictly concave domains there are reasons to expect that (1.10) is true with $\kappa = \frac{2}{3}$. It is also worth noticing that if we have an eigenvalue-free region of the form (1.10) with $\kappa = 1 - \epsilon$, for all ϵ , we get asymptotics with an almost optimal remainder term $\mathcal{O}_\epsilon(r^{d-1+\epsilon})$ in (1.7). The existence of such an eigenvalue-free region with $\kappa = 1 - \epsilon$ has been established recently by the second author [28] for strictly concave domains assuming (1.5) fulfilled. According to our result, the problem of bounding the remainder in the Weyl formula for the (ITE) is reduced to that of getting an eigenvalue-free region in \mathbb{C} , and a larger eigenvalue-free region yields a sharper bound for the remainder. To our best knowledge, it seems that our paper is the first one where such a relationship is established.

For reader's convenience, in what follows in this section we will discuss the main steps in the proof of Theorem 1.1. The starting point of our argument is a trace formula (see Section 3 and (3.5)) which allows us to relate the number of the (ITE) with the number of the eigenvalues, ν_j , of two self-adjoint operators for which the Weyl asymptotics are known to hold, together with a trace of an operator given by an integral involving a meromorphic operator-valued function, $T(\lambda)$, and its inverse $T^{-1}(\lambda)$ (see formula (3.6)). The main problem to deal with is to estimate the trace of this integral and it yields the bound $\mathcal{O}_\epsilon(r^{d-\kappa+\epsilon})$ of the remainder. We apply this formula to obtain an asymptotics for the difference $N(r) - N(r/\sqrt{2})$, $r \rightarrow +\infty$, and by a standard argument it is easy to see that this is sufficient to prove (1.7).

Since it is more convenient to work in the semi-classical setting, we reduce our problem to a semi-classical one by introducing a small parameter $h = \frac{\sqrt{2}}{r}$, $r \gg 1$. Thus we are going to count the number of points $\{z_k\}$, $\frac{z_k}{h^2}$ being an (ITE), in a region of the form

$$\{z \in \mathbb{C} : 1 - Ah^{k-\epsilon} \leq |\operatorname{Re} z| \leq 2 + Ah^{k-\epsilon}, |\operatorname{Im} z| \leq h^{k-\epsilon}\}, \quad A > 0,$$

provided we have an eigenvalue-free region (1.10) with $\kappa - \epsilon$ in place of κ (see Proposition 3.7). This requires to make a change of variables $\lambda = z/h^2$ in the trace formula (3.6) and to study the behavior of the integral term when $0 < h \leq h_0(\epsilon)$ and

$$z \in Z = \{z \in \mathbb{C} : 1/2 < |\operatorname{Re} z| < 3, |\operatorname{Im} z| < 1\}.$$

Next we construct a meromorphic function $g_h(z)$ with poles among the points $\{h^2\nu_j\}$ and such that if an (ITE), λ_k , does not belong to the set $\{\nu_j\}$, then $h^2\lambda_k$ is a

zero of $g_h(z)$ and the multiplicities of the corresponding zeros of $g_h(z)$ and (ITE) agree. It should be mentioned that the construction of the function $g_h(z)$ is not trivial and it requires to build a semi-classical parametrix for the corresponding Dirichlet–to–Neumann map $\mathcal{N}(z, h)$ in the elliptic zone. This is carried out in Section 2 by using the parametrix construction in [27].

The estimate of the remainder is reduced to that of the integral

$$\frac{1}{2\pi i} \int_{\gamma_0} \frac{d}{dz} \log g_h(z) dz, \quad (1.11)$$

where $\gamma_0 \subset Z$ is a suitable closed contour chosen so that on γ_0 we have neither zeros nor poles of $g_h(z)$. The main property of the function $g_h(z)$ is the estimate

$$\log |g_h(z)| \leq C_\epsilon h^{1-d-\epsilon}, \quad \text{for all } 0 < \epsilon \ll 1,$$

provided the distance between z and the set $\{h^2\nu_j\}$ is greater than h^M , $M > 0$ being arbitrary (see Lemma 3.4). This estimate plays a crucial role in the estimate of (1.11). Next in Lemma 3.5 we show that for $z \in Z$, $|\operatorname{Im} z| \geq h^{\kappa-\epsilon}$, we also have

$$\log \frac{1}{|g_h(z)|} \leq C_\epsilon h^{1-d-\epsilon}, \quad \text{for all } 0 < \epsilon \ll 1.$$

Moreover, the function $\log g_h(z)$ is holomorphic in $z \in Z$, $|\operatorname{Im} z| \geq h^{\kappa-\epsilon}$ and satisfies the bound

$$\left| \frac{d}{dz} \log g_h(z) \right| \leq \frac{C_\epsilon h^{1-d-2\epsilon}}{|\operatorname{Im} z|} \quad (1.12)$$

in the domain

$$W := \left\{ z \in \mathbb{C} : \frac{2}{3} \leq |\operatorname{Re} z| \leq \frac{5}{2}, \quad 2h^{\kappa-\epsilon} \leq |\operatorname{Im} z| \leq \frac{1}{2} \right\}.$$

The next step consists of choosing a closed contour $\gamma_0 = \gamma_1 \cup \gamma_3 \cup \gamma_2 \cup \gamma_4$, where $\gamma_3 \subset W$, $\gamma_4 \subset W$ are linear segments parallel to the real axis. For the integrals over γ_j , $j = 3, 4$, we apply (1.12) and one gets

$$\left| \int_{\gamma_j} \frac{d}{dz} \log g_h(z) dz \right| \leq C_\epsilon h^{1-d-3\epsilon}, \quad j = 3, 4. \quad (1.13)$$

We take $\gamma_j = [w_j^-, \tilde{w}_j^-] \cup \tilde{\gamma}_j \cup [\tilde{w}_j^+, w_j^+]$, $j = 1, 2$ with suitable contours $\tilde{\gamma}_j$ (see Section 3 for the notation). The estimates of the imaginary parts of the integrals over $\tilde{\gamma}_j$, $j = 1, 2$, are more delicate since these contours cross the positive real axis and we must avoid the points $\{h^2\nu_k\}$. Our argument is similar to

the choice of the contour in [4] and the details are given in Section 3. The main point is Lemma 3.8, where the contours $\tilde{\gamma}_j$ are constructed so that

$$\left| \operatorname{Im} \int_{\tilde{\gamma}_j} \frac{d}{dz} \log g_h(z) dz \right| \leq C_\epsilon h^{-d+\kappa-2\epsilon}, \quad j = 1, 2. \quad (1.14)$$

Combining this with (1.13), we obtain the statement of Proposition 3.7 and by scaling we get the asymptotics of $N(r) - N(r/\sqrt{2})$.

Acknowledgment. Thanks are due to the referee for the comments and remarks concerning the initial version of the paper.

2. Parametrix of the Dirichlet–to–Neumann map in the elliptic zone

Let $f \in H^1(\Gamma)$ and consider the problem

$$\begin{cases} (P(h) - z)u = 0 & \text{in } \Omega, \\ u = f & \text{on } \Gamma, \end{cases} \quad (2.1)$$

where

$$P(h) = -\frac{h^2}{n(x)} \nabla c(x) \nabla,$$

$0 < h \ll 1$, $z \in Z = \{z \in \mathbb{C} : \frac{1}{2} < |\operatorname{Re} z| < 3, |\operatorname{Im} z| < 1\}$, $c, n \in C^\infty(\bar{\Omega})$ being strictly positive functions. The Dirichlet–to–Neumann map is defined by

$$N(z, h)f := \gamma \mathcal{D}_v u : H^{m+1}(\Gamma) \longrightarrow H^m(\Gamma),$$

where $m \geq 0$, $\mathcal{D}_v = -ih\partial_v$ and γ denotes the restriction on Γ . Denote by G_D the Dirichlet self-adjoint realization of the operator $-n^{-1}\nabla c \nabla$ on the Hilbert space $H = L^2(\Omega, n(x)dx)$. It is well-known that the spectrum of G_D consists of a discrete set of positive eigenvalues which are also poles of the resolvent $(\lambda - G_D)^{-1}$. Moreover, if $\nu_k \in \operatorname{spec} G_D$, we have

$$(\lambda - G_D)^{-1} = \frac{\Pi_k}{\lambda - \nu_k}$$

modulo an operator-valued function holomorphic at ν_k , where Π_k is a finite rank projection. The multiplicity of ν_k is defined as being the rank of Π_k . Let $\mathcal{V}(h) := \{\nu_k \in \operatorname{spec} G_D : h^2 \nu_k \in Z\}$. The following properties of the Dirichlet–to–Neumann map are more or less well-known but we will give a proof for the sake of completeness.

Lemma 2.1. *The Dirichlet-to-Neumann map $\mathcal{N}(z, h)$ is a meromorphic operator-valued function in $z \in Z$ with poles at $h^2\nu_k$, $\nu_k \in \mathcal{V}(h)$. Moreover,*

$$\mathcal{N}(z, h) = \frac{\tilde{\Pi}_k(h)}{z - h^2\nu_k} \quad (2.2)$$

modulo an operator-valued function holomorphic at $h^2\nu_k$, where $\tilde{\Pi}_k(h)$ is of rank $\leq \text{mult}(\nu_k)$. If $\delta(z, h) := \min\{1, \text{dist}\{z, \text{spec } h^2G_D\}\} > 0$, then we have the bound

$$\|\mathcal{N}(z, h)\|_{H^{m+1}(\Gamma) \rightarrow H^m(\Gamma)} \leq \frac{Ch}{\delta(z, h)}, \quad (2.3)$$

where $C > 0$ is a constant which may depend on m .

Proof. Clearly, there exists an extension operator $E_m: H^{m+1}(\Gamma) \rightarrow H^{m+3/2}(\Omega)$ such that $\gamma E_m f = f$ and $E_m f$ is supported near Γ . If $f \in H^{m+1}(\Gamma)$ and z/h^2 does not belong to $\text{spec } G_D$, it is easy to see that the solution u of (2.1) can be expressed by the formula

$$u = E_m f - (h^2G_D - z)^{-1}(P(h) - z)E_m f.$$

Hence

$$\mathcal{N}(z, h)f = \gamma \mathcal{D}_\nu E_m f - \gamma \mathcal{D}_\nu (h^2G_D - z)^{-1}(P(h) - z)E_m f. \quad (2.4)$$

It follows from (2.4) that $\mathcal{N}(z, h)$ is a meromorphic operator-valued function in $z \in Z$ with poles among the poles of $(h^2G_D - z)^{-1}$ and that (2.2) holds with

$$\tilde{\Pi}_k(h) = \gamma \mathcal{D}_\nu \Pi_k (P(h) - h^2\nu_k) E_m.$$

This implies $\text{rank } \tilde{\Pi}_k(h) \leq \text{rank } \Pi_k$ as desired. By (2.4) we also have

$$\begin{aligned} & \|\mathcal{N}(z, h)f\|_{H^m(\Gamma)} \\ & \leq Ch \|f\|_{H^{m+1}(\Gamma)} \\ & \quad + Ch \|(h^2G_D - z)^{-1}\|_{H^{m+3/2}(\Omega) \rightarrow H^{m+3/2}(\Omega)} \|E_m f\|_{H^{m+3/2}(\Omega)} \\ & \quad + Ch \|(h^2G_D - z)^{-1}\|_{H^{m-1/2}(\Omega) \rightarrow H^{m+3/2}(\Omega)} \|P(h)E_m f\|_{H^{m-1/2}(\Omega)}. \end{aligned}$$

Clearly, we have

$$\begin{aligned} & \|E_m f\|_{H^{m+3/2}(\Omega)} \leq C \|f\|_{H^{m+1}(\Gamma)}, \\ & \|P(h)E_m f\|_{H^{m-1/2}(\Omega)} \leq Ch^2 \|E_m f\|_{H^{m+3/2}(\Omega)} \leq Ch^2 \|f\|_{H^{m+1}(\Gamma)}. \end{aligned}$$

On the other hand, the coercive estimate

$$\|v\|_{H^{s+2}(\Omega)} \leq C \|G_D v\|_{H^s(\Omega)} + C \|v\|_{H^s(\Omega)}, \quad \text{for all } v \in D(G_D) \cap H^s(\Omega)$$

implies the bounds

$$\begin{aligned} & \| (h^2 G_D - z)^{-1} \|_{H^{m+3/2}(\Omega) \rightarrow H^{m+3/2}(\Omega)} \\ & \leq C \| (h^2 G_D - z)^{-1} \|_{L^2(\Omega) \rightarrow L^2(\Omega)} \\ & \leq \frac{C}{\delta(z, h)}, \\ & \| (h^2 G_D - z)^{-1} \|_{H^{m-1/2}(\Omega) \rightarrow H^{m+3/2}(\Omega)} \\ & \leq C \| (G_D - i)(h^2 G_D - z)^{-1} \|_{H^{m+3/2}(\Omega) \rightarrow H^{m+3/2}(\Omega)} \\ & \quad \times \| (G_D - i)^{-1} \|_{H^{m-1/2}(\Omega) \rightarrow H^{m+3/2}(\Omega)} \\ & \leq \frac{\tilde{C}}{h^2 \delta(z, h)}. \end{aligned}$$

Therefore, (2.3) follows from the above estimates and the proof is complete. \square

Let (x', ξ') be coordinates on $T^*\Gamma$ and denote by $r_0(x', \xi')$ the principal symbol of the Laplace-Beltrami operator, $-\Delta_\Gamma$, on Γ equipped with the Riemannian metric induced by the Euclidean one in \mathbb{R}^d . It is well-known that r_0 is a polynomial function in ξ' , homogeneous of order 2, and $C_2 |\xi'|^2 \geq r_0(x', \xi') \geq C_1 |\xi'|^2$ with constants $C_2 > C_1 > 0$. Set $m(x) = \frac{n(x)}{c(x)}$. Let $\phi \in C^\infty(\mathbb{R})$, $\phi(\sigma) = 1$ for $|\sigma| \leq 1$, $\phi(\sigma) = 0$ for $|\sigma| \geq 2$, and set

$$\chi(x', \xi') = \phi(\delta_0 r_0(x', \xi')),$$

where $0 < \delta_0 \ll 1$. For $(x', \xi') \in \text{supp}(1 - \chi)$, introduce the function

$$\rho(x', \xi', z) = i \sqrt{r_0(x', \xi') - \gamma m(x') z} = i \sqrt{r_0} \left(1 - z \frac{\gamma m}{r_0}\right)^{1/2}.$$

Since

$$\left|z\right| \frac{\gamma m}{r_0} \leq \frac{1}{2}, \quad \text{for all } z \in Z, (x', \xi') \in \text{supp}(1 - \chi),$$

the functions ρ and ρ^{-1} are holomorphic in $z \in Z$ and

$$\text{Im } \rho(x', \xi', z) \geq C \sqrt{r_0(x', \xi')}$$

with some constant $C > 0$. In what follows in this section we will construct a parametrix for the operator $\mathcal{N}(z, h) \text{Op}_h(1 - \chi)$, where $\text{Op}_h(1 - \chi)$ denotes the

$h - \Psi$ DO with symbol $1 - \chi$. In fact, this construction is carried out in [27] and here we will only recall the main points. First, notice that it suffices to make the construction locally and then to glue up all pieces by using a partition of the unity on Γ . Given an arbitrary point $x^0 \in \Gamma$, there exists a small neighborhood $\mathcal{O}(x^0) \subset \bar{\Omega}$ of x^0 and local normal coordinates $(x_1, x') \in \mathcal{O}(x^0)$ such that $x^0 = (0, 0)$, $\Gamma \cap \mathcal{O}(x^0)$ is defined by $x_1 = 0$, x' being coordinates in $\Gamma \cap \mathcal{O}(x^0)$, $x_1 > 0$ in $\Omega \cap \mathcal{O}(x^0)$, and in these coordinates the operator

$$\mathcal{P}(z, h) = -\frac{h^2}{c(x)} \nabla c(x) \nabla - z \frac{n(x)}{c(x)}$$

can be written in the form

$$\mathcal{P}(z, h) = \mathcal{D}_{x_1}^2 + r(x, \mathcal{D}_{x'}) - zm(x) + hq(x, \mathcal{D}_x) + h^2 \tilde{q}(x).$$

Here we have set $\mathcal{D}_{x_1} = -ih\partial_{x_1}$, $\mathcal{D}_{x'} = -ih\partial_{x'}$, $r(x, \xi') = \langle R(x)\xi', \xi' \rangle$, $R = (R_{ij})$ being a symmetric $(d-1) \times (d-1)$ matrix-valued function with smooth real-valued entries, $q(x, \xi) = \langle q(x), \xi \rangle$, $q(x)$ and $\tilde{q}(x)$ being smooth functions. Moreover, we have $r(0, x', \xi') = r_0(x', \xi')$, $r_0(x', \xi')$ being the principal symbol of $-\Delta_\Gamma$ written in the coordinates (x', ξ') . Let $\psi(x') \in C_0^\infty(\Gamma \cap \mathcal{O}(x^0))$, $\psi = 1$ in a neighborhood of x^0 . In [27], it was constructed a parametrix, \tilde{u}_ψ , of (2.1) satisfying the condition $\tilde{u}_\psi|_{x_1=0} = \text{Op}_h(1 - \chi)\psi f$ and having the form

$$\tilde{u}_\psi(x) = (2\pi h)^{-d+1} \int \int e^{\frac{i}{h}\varphi(x, y', \xi', z)} \phi\left(\frac{x_1}{\delta_1}\right) a(x, \xi', z, h) f(y') dy' d\xi',$$

where ϕ is as above and $\delta_1 > 0$ is a small constant independent of x, ξ', h, z . The phase φ is a complex-valued function such that

$$\varphi|_{x_1=0} = -\langle x' - y', \xi' \rangle, \quad \partial_{x_1} \varphi|_{x_1=0} = \rho, \quad \text{Im } \varphi \geq x_1 \text{Im } \rho/2,$$

and the amplitude a satisfies $a|_{x_1=0} = \psi(x')(1 - \chi(x', \xi'))$. More generally, the functions φ and a are of the form

$$\varphi = -\langle x' - y', \xi' \rangle + \sum_{k=1}^{N-1} x_1^k \varphi_k(x', \xi', z) = -\langle x' - y', \xi' \rangle + \tilde{\varphi},$$

$$a = \sum_{k=0}^{N-1} \sum_{j=0}^{N-1} x_1^k h^j a_{k,j}(x', \xi', z),$$

$N \gg 1$ being an arbitrary integer. The phase φ satisfies the eikonal equation mod $\mathcal{O}(x_1^N)$:

$$(\partial_{x_1} \varphi)^2 + r(x, \nabla_{x'} \varphi) - m(x)z = x_1^N \Psi_N(x, \xi', z) \quad (2.5)$$

and a satisfies the equation

$$e^{-\frac{i}{h}\varphi}\mathcal{P}(z, h)e^{\frac{i}{h}\varphi}a = x_1^N A_N(x, \xi', z, h) + h^N B_N(x, \xi', z, h), \quad (2.6)$$

where Ψ_N , A_N and B_N are smooth functions. It was shown in Section 4 of [27] that $a_{k,j} \in S^{-j}$, $j \geq 0$, $k \geq 1$, $\partial_{x_1}^k A_N \in S^2$, $\partial_{x_1}^k B_N \in S^{1-N}$, $k \geq 0$, uniformly in $z \in Z$ and $0 < x_1 \leq \delta_1$. Recall that S^k are the spaces of all functions $a \in C^\infty(T^*\Gamma)$ satisfying the estimates

$$|\partial_{x'}^\alpha \partial_{\xi'}^\beta a(x', \xi')| \leq C_{\alpha,\beta} \langle \xi' \rangle^{k-|\beta|}, \quad \langle \xi' \rangle = (1 + |\xi'|^2)^{1/2}$$

for all multi-indices α and β . Moreover, the functions $a_{k,j}$, A_N , B_N are polynomials in ρ , ρ^{-1} and z , and therefore they are holomorphic in $z \in Z$. As in [27], it is easy to see that

$$\mathcal{P}(z, h)\tilde{u}_\psi = \text{Op}_h(p_\psi)f,$$

where the function

$$p_\psi = e^{\frac{i}{h}\langle x', \xi' \rangle} \left[\mathcal{P}(z, h), \phi\left(\frac{x_1}{\delta_1}\right) \right] e^{-\frac{i}{h}\langle x', \xi' \rangle} e^{\frac{i}{h}\tilde{\varphi}} a + e^{\frac{i}{h}\tilde{\varphi}} \phi\left(\frac{x_1}{\delta_1}\right) (x_1^N A_N + h^N B_N)$$

is holomorphic in z and satisfies the bounds

$$|\partial_x^\alpha p_\psi| \leq C_{\alpha,N} \left(\frac{h}{\langle \xi' \rangle}\right)^{N-\ell-|\alpha|} \quad \text{for } |\alpha| \leq N - \ell \quad (2.7)$$

with some ℓ independent of N and α . The parametrix, $\tilde{\mathcal{N}}_\psi(z, h)$, of the operator $\mathcal{N}(z, h)\text{Op}_h(1 - \chi)\psi$ is defined by

$$\mathcal{D}_{x_1}\tilde{u}_\psi|_{x_1=0} = \tilde{\mathcal{N}}_\psi(z, h)f = \text{Op}_h(\eta_\psi)f,$$

where

$$\eta_\psi = a \frac{\partial \varphi}{\partial x_1} \Big|_{x_1=0} - ih \frac{\partial a}{\partial x_1} \Big|_{x_1=0} = \psi(1 - \chi)\rho - ih \sum_{j=0}^{N-1} h^j a_{1,j},$$

$$a_{1,0} = -\frac{i}{2}q(0, x', 1, \xi'/\rho)\psi - \frac{1}{2\rho} \langle R(0, x')\xi', \nabla_{x'}\psi(x') \rangle.$$

Since

$$\frac{1}{\rho} = \frac{1}{i\sqrt{r_0}} \left(1 - z \frac{\gamma m}{r_0}\right)^{-1/2} = \frac{1}{i\sqrt{r_0}} + \mathcal{O}(\langle \xi' \rangle^{-3}),$$

we deduce that mod S^{-2} the function $a_{1,0}$ is given by the expression

$$\begin{aligned} a_{1,0} &= -\frac{1}{2}q(0, x', 1, \xi'/\sqrt{r_0})\psi + \frac{i}{2\sqrt{r_0}}\langle R(0, x')\xi', \nabla_{x'}\psi(x') \rangle \\ &= \left\langle \frac{i\nabla_{x'}c(0, x')}{2c(0, x')}, \frac{\xi'}{\sqrt{r_0}} \right\rangle \psi + \frac{i\partial_{x_1}c(0, x')}{2c(0, x')} \psi + q_0(x', \xi') \end{aligned} \quad (2.8)$$

with some function $q_0 \in S^0$ independent of the functions c and n .

Let $\{\psi_j\}_{j=1}^J$ be a partition of the unity on Γ . Set

$$p = \sum_{j=1}^J p\psi_j, \quad \eta = \sum_{j=1}^J \eta\psi_j, \quad \tilde{u} = \sum_{j=1}^J \tilde{u}\psi_j.$$

The operator

$$\tilde{N}(z, h) = \sum_{j=1}^J \tilde{N}_{\psi_j}(z, h) = \text{Op}_h(\eta)$$

is an h - Ψ DO on Γ with a principal symbol $\rho(1 - \chi)$, holomorphic in $z \in Z$. Let u_{ψ_j} be the solution of (2.1) with $u_{\psi}|_{\Gamma} = \text{Op}_h(1 - \chi)\psi f$. Then $u = \sum_{j=1}^J u_{\psi_j}$ is the solution of (2.1) with $u|_{\Gamma} = \text{Op}_h(1 - \chi)f$. Moreover, it is easy to see that, if z/h^2 does not belong to $\text{spec } G_D$, we have

$$u = \tilde{u} - (h^2 G_D - z)^{-1} \frac{c}{n} \mathcal{P}(z, h) \tilde{u}$$

which yields the identity

$$N(z, h) \text{Op}_h(1 - \chi) f = \tilde{N}(z, h) f - \gamma \mathcal{D}_v (h^2 G_D - z)^{-1} \frac{c}{n} \text{Op}_h(p) f. \quad (2.9)$$

It follows from (2.7) that if N is taken large enough, the operator

$$F(z, h) := N(z, h) - \tilde{N}(z, h) = N(z, h) \text{Op}_h(\chi) - \gamma \mathcal{D}_v (h^2 G_D - z)^{-1} \frac{c}{n} \text{Op}_h(p)$$

is meromorphic with values in the space of trace class operators on $L^2(\Gamma)$. Let $\mu_j(F)$ be the characteristic values of F . Recall that $\mu_j(F)$ are defined as being the eigenvalues of the self-adjoint operator $(F^* F)^{1/2}$.

Lemma 2.2. *If z/h^2 does not belong to $\text{spec } G_D$, then for every integer $0 \leq m \leq N/4$ we have the bound*

$$\mu_j(F(z, h)) \leq \frac{C}{\delta(z, h)} (hj^{1/(d-1)})^{-2m}, \quad \text{for all } j, \quad (2.10)$$

where the constant $C > 0$ depends on m and N but is independent of z, h, j , and $\delta(z, h)$ is defined in Lemma 2.1

Proof. We will use the well-known fact that the characteristic values of the Laplace-Beltrami operator on a compact Riemannian manifold without boundary (in our case Γ , $\dim \Gamma = d - 1$) satisfy

$$\mu_j((1 - \Delta_\Gamma)^{-m}) \leq C_m j^{-2m/(d-1)} \quad \text{for all } j, \quad (2.11)$$

for every integer $m \geq 0$. On the other hand, by using the trace theorem and Lemma 2.1, we obtain

$$\begin{aligned} & \|F(z, h)\|_{L^2(\Gamma) \rightarrow H^{2m}(\Gamma)} \\ & \leq \|\mathcal{N}(z, h)\|_{H^{2m+1}(\Gamma) \rightarrow H^{2m}(\Gamma)} \|\text{Op}_h(\chi)\|_{L^2(\Gamma) \rightarrow H^{2m+1}(\Gamma)} \\ & \quad + Ch \|(h^2 G_D - z)^{-1}\|_{H^{2m+3/2}(\Omega) \rightarrow H^{2m+3/2}(\Omega)} \|\text{Op}_h(p)\|_{L^2(\Gamma) \rightarrow H^{2m+3/2}(\Omega)} \\ & \leq \frac{Ch}{\delta(z, h)} \|\text{Op}_h(\chi)\|_{L^2(\Gamma) \rightarrow H^{2m+1}(\Gamma)} + \frac{Ch}{\delta(z, h)} \|\text{Op}_h(p)\|_{L^2(\Gamma) \rightarrow H^{2m+3/2}(\Omega)}. \end{aligned}$$

Since the function χ is compactly supported, we have the bound

$$\|\text{Op}_h(\chi)\|_{L^2(\Gamma) \rightarrow H^{2m+1}(\Gamma)} \leq C_m h^{-2m-1}. \quad (2.12)$$

In view of (2.7) we also have

$$\|\text{Op}_h(p)\|_{L^2(\Gamma) \rightarrow H^{2m+3/2}(\Omega)} \leq C_{m,N} h^{N-2m-\ell_1} \quad (2.13)$$

with some ℓ_1 independent of m and N , provided $0 \leq m \leq N/4$ and N being large enough. By (2.12) and (2.13) we conclude

$$\|F(z, h)\|_{L^2(\Gamma) \rightarrow H^{2m}(\Gamma)} \leq \frac{C_m h^{-2m}}{\delta(z, h)}. \quad (2.14)$$

Clearly, (2.10) follows from (2.11) and (2.14) and the proof is complete. \square

3. Analysis of the transmission eigenvalues

For $\lambda \in \mathbb{C} \setminus \{0\}$ define the operator $R(\lambda)v = u$, where $u = (u_1, u_2)$ and $v = (v_1, v_2)$ solve the problem

$$\begin{cases} \left(-\frac{1}{n_1(x)} \nabla c_1(x) \nabla - \lambda \right) u_1 = v_1 & \text{in } \Omega, \\ \left(-\frac{1}{n_2(x)} \nabla c_2(x) \nabla - \lambda \right) u_2 = v_2 & \text{in } \Omega, \\ u_1 = u_2, \quad c_1 \partial_\nu u_1 = c_2 \partial_\nu u_2 & \text{on } \Gamma. \end{cases} \quad (3.1)$$

Denote by $G_D^{(j)}$, $j = 1, 2$, the Dirichlet self-adjoint realization of the operator $-n_j^{-1}\nabla c_j \nabla$ on the Hilbert space $H_j = L^2(\Omega, n_j(x)dx)$. Set $\mathcal{H} = H_1 \oplus H_2$ and define also the operators $K_j(\lambda)f = u$, where u is the solution of the problem

$$\begin{cases} \left(-\frac{1}{n_j(x)}\nabla c_j(x)\nabla - \lambda\right)u = 0 & \text{in } \Omega, \\ u = f & \text{on } \Gamma. \end{cases} \quad (3.2)$$

Differentiating this equation with respect to λ , one obtains easily the identity

$$\frac{dK_j(\lambda)}{d\lambda} = (G_D^{(j)} - \lambda)^{-1}K_j(\lambda). \quad (3.3)$$

Introduce the operator

$$T(\lambda) := c_1\gamma\partial_\nu K_1(\lambda) - c_2\gamma\partial_\nu K_2(\lambda).$$

Proposition 3.1. *If $T(\lambda)^{-1}$ is a meromorphic operator-valued function with residue of finite rank, the same is true for $R(\lambda)$ and we have the formula*

$$R(\lambda) = \begin{pmatrix} R_{11}(\lambda), & R_{12}(\lambda) \\ R_{21}(\lambda), & R_{22}(\lambda) \end{pmatrix}: \mathcal{H} \longrightarrow \mathcal{H}, \quad (3.4)$$

where

$$R_{11}(\lambda) = (G_D^{(1)} - \lambda)^{-1} - K_1(\lambda)T(\lambda)^{-1}c_1\gamma\partial_\nu(G_D^{(1)} - \lambda)^{-1},$$

$$R_{22}(\lambda) = (G_D^{(2)} - \lambda)^{-1} + K_2(\lambda)T(\lambda)^{-1}c_1\gamma\partial_\nu(G_D^{(2)} - \lambda)^{-1},$$

$$R_{12}(\lambda) = K_1(\lambda)T(\lambda)^{-1}c_1\gamma\partial_\nu(G_D^{(2)} - \lambda)^{-1},$$

$$R_{21}(\lambda) = -K_2(\lambda)T(\lambda)^{-1}c_2\gamma\partial_\nu(G_D^{(1)} - \lambda)^{-1}.$$

Moreover, if $\gamma_0 \subset \mathbb{C}$ is a simple closed positively oriented curve which avoids the eigenvalues of $G_D^{(j)}$, $j = 1, 2$, as well as the poles of $T(\lambda)^{-1}$, then we have the identity

$$\begin{aligned} & -\operatorname{tr}_{\mathcal{H}}(2\pi i)^{-1} \int_{\gamma_0} R(\lambda)d\lambda + \sum_{j=1}^2 \operatorname{tr}_{H_j}(2\pi i)^{-1} \int_{\gamma_0} (G_D^{(j)} - \lambda)^{-1}d\lambda \\ & = \operatorname{tr}_{L^2(\Gamma)}(2\pi i)^{-1} \int_{\gamma_0} T(\lambda)^{-1} \frac{dT(\lambda)}{d\lambda} d\lambda. \end{aligned} \quad (3.5)$$

Proof. Clearly, if (u_j, v_j) satisfies (3.1) and λ does not belong to

$$\text{spec } G_D^{(1)} \cup \text{spec } G_D^{(2)},$$

we have

$$u_j = (G_D^{(j)} - \lambda)^{-1} v_j + K_j(\lambda) f,$$

where $f = \gamma u_1 = \gamma u_2$. The boundary condition in (3.1) implies the identity

$$\begin{aligned} 0 &= c_1 \partial_\nu u_1 - c_2 \partial_\nu u_2 \\ &= T(\lambda) f + c_1 \gamma \partial_\nu (G_D^{(1)} - \lambda)^{-1} v_1 - c_2 \gamma \partial_\nu (G_D^{(2)} - \lambda)^{-1} v_2. \end{aligned}$$

Hence

$$\begin{aligned} u_j &= (G_D^{(j)} - \lambda)^{-1} v_j \\ &\quad - K_j(\lambda) T(\lambda)^{-1} (c_1 \gamma \partial_\nu (G_D^{(1)} - \lambda)^{-1} v_1 - c_2 \gamma \partial_\nu (G_D^{(2)} - \lambda)^{-1} v_2) \end{aligned}$$

which clearly implies (3.4). Moreover, if $T(\lambda)^{-1}$ is meromorphic, so are the operators $R_{ij}(\lambda)$, and by (3.4) the operator $R(\lambda)$ is meromorphic, too. Using (3.3) and the cyclicity of the trace (see Lemma 2.2 of [23]), we get

$$\begin{aligned} &\text{tr}_{\mathcal{H}} (2\pi i)^{-1} \int_{\gamma_0} R(\lambda) d\lambda \\ &= \text{tr}_{H_1} (2\pi i)^{-1} \int_{\gamma_0} R_{11}(\lambda) d\lambda + \text{tr}_{H_2} (2\pi i)^{-1} \int_{\gamma_0} R_{22}(\lambda) d\lambda \\ &= \text{tr}_{H_1} (2\pi i)^{-1} \int_{\gamma_0} (G_D^{(1)} - \lambda)^{-1} d\lambda + \text{tr}_{H_2} (2\pi i)^{-1} \int_{\gamma_0} (G_D^{(2)} - \lambda)^{-1} d\lambda \\ &\quad - \text{tr}_{H_1} (2\pi i)^{-1} \int_{\gamma_0} K_1(\lambda) T(\lambda)^{-1} c_1 \gamma \partial_\nu (G_D^{(1)} - \lambda)^{-1} d\lambda \\ &\quad + \text{tr}_{H_2} (2\pi i)^{-1} \int_{\gamma_0} K_2(\lambda) T(\lambda)^{-1} c_2 \gamma \partial_\nu (G_D^{(2)} - \lambda)^{-1} d\lambda \\ &= \text{tr}_{H_1} (2\pi i)^{-1} \int_{\gamma_0} (G_D^{(1)} - \lambda)^{-1} d\lambda + \text{tr}_{H_2} (2\pi i)^{-1} \int_{\gamma_0} (G_D^{(2)} - \lambda)^{-1} d\lambda \\ &\quad - \text{tr}_{L^2(\Gamma)} (2\pi i)^{-1} \int_{\gamma_0} T(\lambda)^{-1} c_1 \gamma \partial_\nu (G_D^{(1)} - \lambda)^{-1} K_1(\lambda) d\lambda \\ &\quad + \text{tr}_{L^2(\Gamma)} (2\pi i)^{-1} \int_{\gamma_0} T(\lambda)^{-1} c_2 \gamma \partial_\nu (G_D^{(2)} - \lambda)^{-1} K_2(\lambda) d\lambda \end{aligned}$$

$$\begin{aligned}
&= \operatorname{tr}_{H_1} (2\pi i)^{-1} \int_{\gamma_0} (G_D^{(1)} - \lambda)^{-1} d\lambda + \operatorname{tr}_{H_2} (2\pi i)^{-1} \int_{\gamma_0} (G_D^{(2)} - \lambda)^{-1} d\lambda \\
&\quad - \operatorname{tr}_{L^2(\Gamma)} (2\pi i)^{-1} \int_{\gamma_0} T(\lambda)^{-1} c_1 \frac{d\gamma \partial_\nu K_1(\lambda)}{d\lambda} d\lambda \\
&\quad + \operatorname{tr}_{L^2(\Gamma)} (2\pi i)^{-1} \int_{\gamma_0} T(\lambda)^{-1} c_2 \frac{d\gamma \partial_\nu K_2(\lambda)}{d\lambda} d\lambda
\end{aligned}$$

which implies (3.5). \square

If $R(\lambda)$ is a meromorphic operator-valued function with residue of finite rank, we define the multiplicity of a pole $\lambda_k \in \mathbb{C}$ of $R(\lambda)$ by

$$\operatorname{mult}(\lambda_k) = \operatorname{rank} (2\pi i)^{-1} \int_{|\lambda - \lambda_k| = \varepsilon} R(\lambda) d\lambda, \quad 0 < \varepsilon \ll 1.$$

Let the curve γ_0 be as in Proposition 3.1 and denote by M_{γ_0} and $M_{\gamma_0}^{(j)}$, $j = 1, 2$, the number (counted with the multiplicity) of the poles of $R(\lambda)$ and the eigenvalues of $G_D^{(j)}$, respectively, in the interior of γ_0 . Proposition 3.1 implies the following

Corollary 3.2. *We have the identity*

$$M_{\gamma_0} = M_{\gamma_0}^{(1)} + M_{\gamma_0}^{(2)} + \operatorname{tr}_{L^2(\Gamma)} (2\pi i)^{-1} \int_{\gamma_0} T(\lambda)^{-1} \frac{dT(\lambda)}{d\lambda} d\lambda. \quad (3.6)$$

Proof. It is easy to see that $R(\lambda) = (\mathcal{A} - \lambda)^{-1}$, where the operator \mathcal{A} is defined by

$$\mathcal{A} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} -\frac{1}{n_1(x)} \nabla c_1(x) \nabla u_1 \\ -\frac{1}{n_2(x)} \nabla c_2(x) \nabla u_2 \end{pmatrix}$$

with domain

$$\begin{aligned}
D(\mathcal{A}) = \{ &(u_1, u_2) \in \mathcal{H}: \nabla c_1(x) \nabla u_1 \in L^2(\Omega), \nabla c_2(x) \nabla u_2 \in L^2(\Omega), \\
&\gamma u_1 = \gamma u_2, c_1 \gamma \partial_\nu u_1 = c_2 \gamma \partial_\nu u_2 \}.
\end{aligned}$$

Hence the finite-rank operator

$$-(2\pi i)^{-1} \int_{|\lambda - \lambda_k| = \varepsilon} R(\lambda) d\lambda = (2\pi i)^{-1} \int_{|\lambda - \lambda_k| = \varepsilon} (\lambda - \mathcal{A})^{-1} d\lambda$$

is in fact a projection (e.g. see [12]), and therefore the rank coincides with the trace. Thus, (3.6) follows from (3.5). \square

Let z and h be as in the previous section and denote by $\mathcal{N}_j, \tilde{\mathcal{N}}_j, F_j, j = 1, 2$, the operators defined by replacing in the definition of $\mathcal{N}, \tilde{\mathcal{N}}, F$ introduced in Section 2 the pair (c, n) by (c_j, n_j) . Clearly, we have the relationship

$$\begin{aligned} hT(z/h^2) &= c_1\mathcal{N}_1(z, h) - c_2\mathcal{N}_2(z, h) \\ &= c_1F_1(z, h) - c_2F_2(z, h) + c_1\tilde{\mathcal{N}}_1(z, h) - c_2\tilde{\mathcal{N}}_2(z, h). \end{aligned} \quad (3.7)$$

In what follows H_h^s will denote the Sobolev space $H^s(\Gamma)$ equipped with the semi-classical norm.

Lemma 3.3. *There exist an invertible, bounded operator*

$$E(z, h): H_h^s \longrightarrow H_h^{s+k} = O(1),$$

with an inverse

$$E(z, h)^{-1}: H_h^s \longrightarrow H_h^{s-k} = O(1),$$

for all $s \in \mathbb{R}$, and trace class operators $L_l(z, h)$ and $L_r(z, h)$ such that

$$E(z, h)(c_1\tilde{\mathcal{N}}_1(z, h) - c_2\tilde{\mathcal{N}}_2(z, h)) = I + L_l(z, h), \quad (3.8)$$

$$(c_1\tilde{\mathcal{N}}_1(z, h) - c_2\tilde{\mathcal{N}}_2(z, h))E(z, h) = I + L_r(z, h), \quad (3.9)$$

where $k = -1$ if (1.5) holds, $k = 1$ if (1.6) holds. Moreover, the operators E, E^{-1}, L_l, L_r are holomorphic with respect to z for $z \in Z$.

Proof. Set $m_j = \frac{n_j}{c_j}, \rho_j = i\sqrt{r_0 - z\gamma m_j}, j = 1, 2$, and let the real-valued function $\chi, 0 \leq \chi \leq 1$ be as in Section 2, with a sufficiently large support. It follows from the parametrix construction in Section 2 that $c_1\tilde{\mathcal{N}}_1 - c_2\tilde{\mathcal{N}}_2 = \text{Op}_h(b)$ with a symbol $b = \sum_{j=0}^N h^j b_j$, where $b_j \in S^{1-j}$ are holomorphic in $z \in Z$, and

$$b_0 = (c_1\rho_1 - c_2\rho_2)(1 - \chi).$$

Let $\chi_0 \in C^\infty(T^*\Gamma)$ be a real-valued compactly supported function such that $0 \leq \chi_0 \leq 1$ and $\chi_0 = 1$ on $\text{supp } \chi$. It suffices to show that the operator $\text{Op}_h(\chi_0 + b)$ is invertible. Indeed, this would imply (3.8) and (3.9) with $E = (\text{Op}_h(\chi_0 + b))^{-1}$ and $L_l = E\text{Op}_h(\chi_0), L_r = \text{Op}_h(\chi_0)E$.

An easy computation shows that

$$b_0 = \frac{\tilde{c}(x')(c_0(x')r_0(x', \xi') - z)}{c_1\rho_1 + c_2\rho_2}(1 - \chi(x', \xi')),$$

where \tilde{c} and c_0 are the restrictions on Γ of the functions

$$c_1n_1 - c_2n_2 \quad \text{and} \quad \frac{c_1^2 - c_2^2}{c_1n_1 - c_2n_2},$$

respectively. Let us see that

$$C_1 \langle \xi' \rangle^k \leq |\chi_0 + b_0| \leq C_2 \langle \xi' \rangle^k \quad (3.10)$$

with some constants $C_1, C_2 > 0$, where $k = -1$ if $c_0(x') \equiv 0$ and $k = 1$ if $c_0(x') \neq 0$, for all $x' \in \Gamma$. Since

$$b_0 = \frac{\tilde{c}(c_0 r_0 - z)}{i(c_1 + c_2)\sqrt{r_0}}(1 - \chi)(1 + \mathcal{O}(\langle \xi' \rangle^{-1})),$$

we have with some positive constants $\tilde{C}, \tilde{C}_1, \tilde{C}_2$,

$$\begin{aligned} 2|\chi_0 + b_0| &\geq |\chi_0 + \operatorname{Re} b_0| + |\operatorname{Im} b_0| \\ &\geq \chi_0 - |\operatorname{Re} b_0| + |\operatorname{Im} b_0| \\ &\geq \chi_0 + \frac{\tilde{C}_1}{\langle \xi' \rangle} |c_0 r_0 - z| (1 - \chi)(1 - \mathcal{O}(\langle \xi' \rangle^{-1})) \\ &\geq \chi_0 + \tilde{C}_2 \langle \xi' \rangle^k (1 - \chi) \\ &\geq \tilde{C} \langle \xi' \rangle^k \end{aligned}$$

which yields the lower bound in (3.10). The upper bound is obvious.

It follows from (3.10) that $(\chi_0 + b_0)^{-1} \in S^{-k}$. Moreover, when $c_1 \equiv c_2$, $\partial_\nu c_1 \equiv \partial_\nu c_2$ on Γ , by (2.8) one concludes that $b_1 \in S^{-2}$. Hence the operator $\operatorname{Op}_h(\chi_0 + b)$ is invertible with an inverse which is an h - Ψ DO with a symbol belonging to the class S^{-k} . In particular, we have

$$\begin{aligned} (\operatorname{Op}_h(\chi_0 + b))^{-1}: H_h^s &\longrightarrow H_h^{s+k} = \mathcal{O}(1), \\ \operatorname{Op}_h(\chi_0 + b): H_h^s &\longrightarrow H_h^{s-k} = \mathcal{O}(1), \end{aligned}$$

for all $s \in \mathbb{R}$. □

Set

$$\mathcal{V}_j(h) := \{v_k \in \operatorname{spec} G_D^{(j)} : h^2 v_k \in Z\}, \quad j = 1, 2.$$

Define the operator \mathcal{K} as follows:

$$\begin{aligned} \mathcal{K}(z, h) &= E(z, h)(c_1 F_1(z, h) - c_2 F_2(z, h)) + L_l(z, h) \quad \text{if } k = -1, \\ \mathcal{K}(z, h) &= (c_1 F_1(z, h) - c_2 F_2(z, h))E(z, h) + L_r(z, h) \quad \text{if } k = 1. \end{aligned}$$

We obtain easily that

$$E(c_1\mathcal{N}_1 - c_2\mathcal{N}_2) = I + \mathcal{K} \quad \text{if } k = -1,$$

$$(c_1\mathcal{N}_1 - c_2\mathcal{N}_2)E = I + \mathcal{K} \quad \text{if } k = 1.$$

Clearly, the operator \mathcal{K} is trace class and meromorphic in $z \in Z$ with poles $\{w_k\}$, $w_k/h^2 \in \mathcal{V}_1(h) \cup \mathcal{V}_2(h)$, and residue of finite rank, so we can define the meromorphic function

$$g_h(z) := \det(I + \mathcal{K}(z, h)).$$

Lemma 3.4. *For all $z \in Z$ such that*

$$\delta^\sharp(z, h) := \min\{1, \text{dist}\{z, \text{spec } h^2 G_D^{(1)} \cup \text{spec } h^2 G_D^{(2)}\}\} > 0$$

we have the bound

$$\log |g_h(z)| \leq C_\varepsilon h^{1-d} \delta^\sharp(z, h)^{-\varepsilon}, \quad \text{for all } 0 < \varepsilon \ll 1. \quad (3.11)$$

Proof. It follows from Lemma 2.2 and the properties of the characteristic values that $\mu_j(\mathcal{K})$ satisfy the bound (2.10) with a new constant $C > 0$ and δ replaced by δ^\sharp . In fact, for $k = -1$ we have

$$\mu_{j_1+j_2-1}(\mathcal{K}) \leq \mu_{j_1}((E(z, h)(c_1 F_1(z, h) - c_2 F_2(z, h))) + \mu_{j_2}(L_l(z, h)).$$

Since the operator $E(x, h)$ is bounded, for the first term on the right hand side we apply Lemma 2.2. On the other hand, $\mu_{j_2}(E\text{Op}_h(\chi_0)) \leq C\mu_{j_2}(\text{Op}_h(\chi_0))$ and for $\mu_{j_2}(\text{Op}_h(\chi_0))$ we obtain easily (2.10) with $\delta(z, h) = 1$ since χ_0 has compact support. Next, if $j = j_1 + j_2 - 1$, then we have $j_1 \geq (j + 1)/2$ or $j_2 \geq (j + 1)/2$. The case $k = 1$ is similar.

Therefore, we have

$$\begin{aligned} \log |g_h(z)| &\leq \sum_{j=1}^{\infty} \log(1 + \mu_j(\mathcal{K})) \\ &\leq \sum_{j=1}^{\infty} \log(1 + C\delta^\sharp(z, h)^{-1} h^{-2m} j^{-2m/(d-1)}) \\ &\leq \int_0^{\infty} \log(1 + C\delta^\sharp(z, h)^{-1} h^{-2m} t^{-2m/(d-1)}) dt \\ &= C_m h^{-d+1} (\delta^\sharp(z, h))^{-\frac{d-1}{2m}} \int_0^{\infty} \log(1 + t^{-2m/(d-1)}) dt \\ &\leq \tilde{C}_m h^{-d+1} \delta^\sharp(z, h)^{-\frac{d-1}{2m}}. \end{aligned} \quad (3.12)$$

Now, given any $0 < \varepsilon \ll 1$, we can take $m \sim \frac{d-1}{2\varepsilon}$ and $N \geq 4m$, and (3.11) follows from (3.12). \square

The next lemma is an almost direct consequence of the results of [27].

Lemma 3.5. *Let κ be as in Theorem 1.1. Then, given any $0 < \epsilon \ll 1$, the operator $I + \mathcal{K}(z, h)$ is invertible on $L^2(\Gamma)$ for $z \in Z$, $|\operatorname{Im} z| \geq h^{\kappa-\epsilon}$, and its inverse satisfies in this region the bound*

$$\|(I + \mathcal{K}(z, h))^{-1}\|_{L^2 \rightarrow L^2} \leq Ch^{-\ell} \quad (3.13)$$

with some constants $C, \ell > 0$. For these values of z we also have

$$\log \frac{1}{|g_h(z)|} \leq C_\epsilon h^{1-d-\epsilon}, \quad \text{for all } 0 < \epsilon \ll 1. \quad (3.14)$$

Moreover, the function $\log g_h(z)$ is holomorphic in $z \in Z$, $|\operatorname{Im} z| \geq h^{\kappa-\epsilon}$ and satisfies the bound

$$\left| \frac{d}{dz} \log g_h(z) \right| \leq \frac{C_\epsilon h^{1-d-2\epsilon}}{|\operatorname{Im} z|} \quad (3.15)$$

in $W := \{z \in \mathbb{C}: \frac{2}{3} \leq |\operatorname{Re} z| \leq \frac{5}{2}, 2h^{\kappa-\epsilon} \leq |\operatorname{Im} z| \leq \frac{1}{2}\}$.

Proof. It follows from the analysis in Section 5 of [27] that, under the assumptions of Theorem 1.1, the operator $c_1 \mathcal{N}_1(z, h) - c_2 \mathcal{N}_2(z, h)$ is invertible for $z \in Z$, $|\operatorname{Im} z| \geq h^{\kappa-\epsilon}$ and

$$\|(c_1 \mathcal{N}_1(z, h) - c_2 \mathcal{N}_2(z, h))^{-1}\|_{H_h^1 \rightarrow L^2} \leq Ch^{-\ell} \quad \text{if } k = -1,$$

$$\|(c_1 \mathcal{N}_1(z, h) - c_2 \mathcal{N}_2(z, h))^{-1}\|_{L^2 \rightarrow H_h^1} \leq Ch^{-\ell} \quad \text{if } k = 1.$$

Now (3.13) follows from these bounds and Lemma 3.3 because

$$(I + \mathcal{K})^{-1} = (c_1 \mathcal{N}_1 - c_2 \mathcal{N}_2)^{-1} E^{-1} \quad \text{if } k = -1,$$

$$(I + \mathcal{K})^{-1} = E^{-1} (c_1 \mathcal{N}_1 - c_2 \mathcal{N}_2)^{-1} \quad \text{if } k = 1.$$

The bound (3.14) can be obtained in precisely the same way as (3.11) by using (3.13) and the formula

$$\frac{1}{g_h(z)} = \det(I - (I + \mathcal{K}(z, h))^{-1} \mathcal{K}(z, h)).$$

Note that the norm $\|(I + \mathcal{K}(z, h))^{-1}\|$ will add a factor $h^{-\frac{l(d-1)}{2m}}$ which for sufficiently large m yields a factor $\mathcal{O}(h^{-\epsilon})$.

Clearly, it follows from the Fredholm theorem that, under the assumptions of Theorem 1.1, the operator-valued function $(I + \mathcal{K}(z, h))^{-1} : L^2(\Gamma) \rightarrow L^2(\Gamma)$ is meromorphic in Z with finite rank residue and holomorphic with respect to $z \in Z$ for $|\operatorname{Im} z| \geq h^{\kappa-\epsilon}$. Therefore the functions $g_h(z)$ and $\frac{1}{g_h(z)}$ are holomorphic in $z \in Z$, $|\operatorname{Im} z| \geq h^{\kappa-\epsilon}$, and hence so is $\log g_h(z)$. Fix an arbitrary $w \in W$. Then the function $f(z) = \log \frac{g_h(z)}{g_h(w)}$ is holomorphic in $z \in Z$, $|\operatorname{Im} z| \geq h^{\kappa-\epsilon}$ and $f(w) = 0$. It follows from the bounds (3.11) and (3.14) that $\operatorname{Re} f(z) \leq \mathcal{O}_\epsilon(h^{1-d-2\epsilon})$ for $z \in Z$, $|\operatorname{Im} z| \geq h^{\kappa-\epsilon}$. In particular, the later estimate holds on the circle $C_w = \{z \in \mathbb{C} : |z - w| = \frac{|\operatorname{Im} w|}{2}\}$ since for every $z \in C_w$ we have $|\operatorname{Im} z| \geq \frac{|\operatorname{Im} w|}{2}$. Applying the Caratheodory theorem (e.g. see 5.5 in [26]), we get

$$|f'(z)| = \mathcal{O}_\epsilon(h^{1-d-2\epsilon})|\operatorname{Im} w|^{-1} \quad \text{for } |z - w| \leq \frac{|\operatorname{Im} w|}{3}.$$

This implies (3.15) because $f'(z) = \frac{d}{dz} \log g_h(z)$. \square

Let $\gamma_0 \subset Z$ be a simple closed positively oriented curve which avoids the eigenvalues of $h^2 G_D^{(j)}$, $j = 1, 2$, as well as the poles of $T(z/h^2)^{-1}$. Denote by $M_{\gamma_0}(h)$ the number of the poles, $\{\lambda_k\}$, of $R(\lambda)$ such that $h^2 \lambda_k$ are in the interior of the domain ω_0 with boundary γ_0 . Similarly, we denote by $M_{\gamma_0}^{(j)}(h)$ the number of the eigenvalues, $\{\nu_k\}$, of $G_D^{(j)}$ such that $h^2 \nu_k$ are in ω_0 . Corollary 3.2 implies the following

Lemma 3.6. *We have the identity*

$$M_{\gamma_0}(h) = M_{\gamma_0}^{(1)}(h) + M_{\gamma_0}^{(2)}(h) + \frac{1}{2\pi i} \int_{\gamma_0} \frac{d}{dz} \log g_h(z) dz. \quad (3.16)$$

Proof. We apply (3.6) and use the identities

$$\begin{aligned} hT(z/h^2) &= E^{-1}(z, h)(I + \mathcal{K}(z, h)), \\ (hT(z/h^2))^{-1} &= (I + \mathcal{K}(z, h))^{-1} E(z, h) \end{aligned}$$

combined with the analyticity of $E(z, h)$ in z and the following well-known formula

$$\operatorname{tr} (I + \mathcal{K}(z, h))^{-1} \frac{d\mathcal{K}(z, h)}{dz} = \frac{d}{dz} \log \det(I + \mathcal{K}(z, h)).$$

The above formula for $\log \det(I + \mathcal{K}(z, h))$ is classical for finite rank perturbations of the identity. For trace class ones this formula follows by an approximation with finite rank operators (see for example, Section 5 in [21]). \square

It follows from (3.16) that $z_0 \in Z \setminus \text{spec}(h^2 G_D^{(1)}) \cup \text{spec}(h^2 G_D^{(2)})$ is a zero of $g_h(z)$ if and only if z_0 is a pole of $R(z/h^2)$ (and hence z_0/h^2 is an interior transmission eigenvalue) and the multiplicities coincide. Similarly, one can see that if \tilde{z}_0 is a pole of $g_h(z)$ with multiplicity \tilde{m}_0 , then $\tilde{z}_0 \in \text{spec}(h^2 G_D^{(1)}) \cup \text{spec}(h^2 G_D^{(2)})$ and $\tilde{m}_0 \leq \tilde{m}_1 + \tilde{m}_2$, where \tilde{m}_j is the multiplicity of \tilde{z}_0/h^2 as an eigenvalue of $G_D^{(j)}$. In what follows we will use the formula (3.16) to prove the following

Proposition 3.7. *For every $0 < \epsilon \ll 1$ and $A > 0$, independent of h , we have the asymptotics*

$$\begin{aligned} I(h) &:= \#\{z_k, z_k/h^2 \text{ is (ITE)}: 1 - Ah^{\kappa-\epsilon} \leq |\text{Re } z_k| \leq 2 + Ah^{\kappa-\epsilon}, |\text{Im } z_k| \leq h^{\kappa-\epsilon}\} \\ &= (2^{d/2} - 1)(\tau_1 + \tau_2)h^{-d} + \mathcal{O}_{\epsilon, A}(h^{-d+\kappa-3\epsilon}), \quad 0 < h \leq h_0(\epsilon, A). \end{aligned} \quad (3.17)$$

Proof. We will consider only the case $\text{Re } z_k > 0$, since the case $\text{Re } z_k < 0$ is similar (and even simpler since the function $g_h(z)$ does not have poles in $\text{Re } z < 0$). Consider the points $w_1^\pm = 1 - Ah^{\kappa-\epsilon} \pm \frac{i}{3}$, $w_2^\pm = 2 + Ah^{\kappa-\epsilon} \pm \frac{i}{3}$, $\tilde{w}_1^\pm = 1 - Ah^{\kappa-\epsilon} \pm i3h^{\kappa-\epsilon}$, $\tilde{w}_2^\pm = 2 + Ah^{\kappa-\epsilon} \pm i3h^{\kappa-\epsilon}$ and set

$$\Theta_1 = \{z \in \mathbb{C} : 1 - 2(A+1)h^{\kappa-\epsilon} \leq \text{Re } z \leq 1 + h^{\kappa-\epsilon}, |\text{Im } z| \leq 4h^{\kappa-\epsilon}\},$$


$$\Theta_2 = \{z \in \mathbb{C} : 2 - h^{\kappa-\epsilon} \leq \text{Re } z \leq 2 + 2(A+1)h^{\kappa-\epsilon}, |\text{Im } z| \leq 4h^{\kappa-\epsilon}\}.$$

The following lemma will be proved later on.

Lemma 3.8. *There exist positively oriented piecewise smooth curves $\tilde{\gamma}_1 \subset \Theta_1$ and $\tilde{\gamma}_2 \subset \Theta_2$, where $\tilde{\gamma}_1$ connects the point \tilde{w}_1^- with \tilde{w}_1^+ , while $\tilde{\gamma}_2$ connects the point \tilde{w}_2^+ with \tilde{w}_2^- , such that*

$$\left| \text{Im} \int_{\tilde{\gamma}_j} \frac{d}{dz} \log g_h(z) dz \right| \leq C_\epsilon h^{-d+\kappa-2\epsilon}, \quad j = 1, 2. \quad (3.18)$$

Now we apply Lemma 3.6 with a contour $\gamma_0 = \gamma_1 \cup \gamma_3 \cup \gamma_2 \cup \gamma_4$, where $\gamma_3 \subset W$ is the segment $[w_1^+, w_2^+]$ on the line passing through the points w_1^+ and w_2^+ , and $\gamma_4 \subset W$ is the segment $[w_2^-, w_1^-]$ on the line passing through the points w_2^- and w_1^- . Next, $\gamma_1 = [w_1^-, \tilde{w}_1^-] \cup \tilde{\gamma}_1 \cup [\tilde{w}_1^+, w_1^+]$, $\gamma_2 = [w_2^+, \tilde{w}_2^+] \cup \tilde{\gamma}_2 \cup [\tilde{w}_2^-, w_2^-]$ (see Figure 1).

Figure 1. Contour γ_0 .

Since $\gamma_j \subset W$, $|\gamma_j| = \mathcal{O}(1)$, $j = 3, 4$, by (3.15) we have

$$\begin{aligned}
 \left| \int_{\gamma_j} \frac{d}{dz} \log g_h(z) dz \right| &\leq \int_{\gamma_j} \left| \frac{d}{dz} \log g_h(z) \right| |dz| \\
 &\leq C_\epsilon h^{-d+1-2\epsilon} \int_{\gamma_j} |dz| \\
 &\leq C_\epsilon h^{-d+1-2\epsilon}, \quad j = 3, 4.
 \end{aligned} \tag{3.19}$$

Applying (3.15) once more, we have

$$\begin{aligned}
 \left| \int_{[w_j^\pm, \tilde{w}_j^\pm]} \frac{d}{dz} \log g_h(z) dz \right| &\leq C_\epsilon h^{-d+1-2\epsilon} \int_{3h^{\kappa-\epsilon}}^{1/2} \frac{d\sigma}{\sigma} \\
 &\leq C_\epsilon h^{-d+1-3\epsilon}, \quad j = 1, 2.
 \end{aligned} \tag{3.20}$$

On the other hand, since the counting function of the eigenvalues of $G_D^{(j)}$ satisfies the Weyl law, we deduce

$$\begin{aligned} M_{\gamma_0}^{(j)}(h) &\leq \#\{v_k \in \text{spec } G_D^{(j)} : 1 - 2(A+1)h^{\kappa-\epsilon} \leq h^2 v_k \leq 2 + 2(A+1)h^{\kappa-\epsilon}\} \\ &= \tau_j \left(\frac{2}{h^2} + \frac{2(A+1)h^{\kappa-\epsilon}}{h^2} \right)^{d/2} - \tau_j \left(\frac{1}{h^2} - \frac{2(A+1)h^{\kappa-\epsilon}}{h^2} \right)^{d/2} + \mathcal{O}_\epsilon(h^{-d+1}) \\ &= (2^{d/2} - 1)\tau_j h^{-d} + \mathcal{O}_{\epsilon,A}(h^{-d+\kappa-\epsilon}) \end{aligned}$$

and similarly

$$\begin{aligned} M_{\gamma_0}^{(j)}(h) &\geq \#\{v_k \in \text{spec } G_D^{(j)} : 1 + h^{\kappa-\epsilon} \leq h^2 v_k \leq 2 - h^{\kappa-\epsilon}\} \\ &= (2^{d/2} - 1)\tau_j h^{-d} - \mathcal{O}_\epsilon(h^{-d+\kappa-\epsilon}). \end{aligned}$$

Consequently,

$$M_{\gamma_0}^{(j)}(h) = (2^{d/2} - 1)\tau_j h^{-d} + \mathcal{O}_{\epsilon,A}(h^{-d+\kappa-\epsilon}). \quad (3.21)$$

Taking together (3.16), (3.18), (3.19), (3.20), and (3.21), we obtain

$$M_{\gamma_0}(h) = (2^{d/2} - 1)(\tau_1 + \tau_2)h^{-d} + \mathcal{O}_{\epsilon,A}(h^{-d+\kappa-3\epsilon}). \quad (3.22)$$

Thus, to establish (3.17), it remains to show that the counting function $I(h)$ satisfies

$$|I(h) - M_{\gamma_0}(h)| \leq C_{\epsilon,A} h^{-d+\kappa-3\epsilon}. \quad (3.23)$$

Given a parameter $\theta > 0$, independent of h , introduce

$$B_j^\pm(\theta) = \{z \in \mathbb{C} : |z - \tilde{w}_j^\pm| \leq \theta h^{\kappa-\epsilon}\}.$$

Clearly, there exists $\theta_0 > 0$ such that $\Theta_j \subset B_j^+(\theta) \cup B_j^-(\theta)$, for all $\theta \geq \theta_0$, $j = 1, 2$. Let $\{z_k^{\pm,j}\}$ be the zeros (repeated with their multiplicities) of $g_h(z)$ in $B_j^\pm(2\theta_0)$ and let $\{y_k^{\pm,j}\}$ be the poles (repeated with their multiplicities) of $g_h(z)$ in $B_j^\pm(4\theta_0)$. Therefore the function

$$f_h^{\pm,j}(z) = g_h(z) \prod_k (z - y_k^{\pm,j})$$

is holomorphic in the interior of $B_j^\pm(4\theta_0)$. Obviously, $\{y_k^{\pm,j}\}$ are among the eigenvalues of the operators $G_D^{(1)}$ and $G_D^{(2)}$ in an interval of the form

$$[1 - \mathcal{O}(h^{\kappa-\epsilon}), 1 + \mathcal{O}(h^{\kappa-\epsilon})] \cup [2 - \mathcal{O}(h^{\kappa-\epsilon}), 2 + \mathcal{O}(h^{\kappa-\epsilon})].$$

Hence, by the Weyl law for the counting function of these eigenvalues, as in the proof of (3.21), we get

$$\#\{y_k^{\pm,j}\} \leq C_{\epsilon,A} h^{-d+\kappa-\epsilon}, \quad j = 1, 2. \quad (3.24)$$

By (3.14) and (3.24), we have

$$\begin{aligned} \log |f_h^{\pm,j}(\tilde{w}_j^{\pm})| &= \log |g_h(\tilde{w}_j^{\pm})| + \sum_k \log |\tilde{w}_j^{\pm} - y_k^{\pm,j}| \\ &\geq -C_{\epsilon} h^{-d+1-\epsilon} - \#\{y_k^{\pm,j}\} C \log \frac{1}{h} \\ &\geq -2C_{\epsilon,A} h^{-d+\kappa-2\epsilon}. \end{aligned} \quad (3.25)$$

On the other hand, applying (3.11) and (3.24), for $z \in B_j^{\pm}(\theta)$, $\theta_0 < \theta < 4\theta_0$, $|z - y_k^{\pm,j}| \geq h^M$, $M \gg 1$, we obtain

$$\begin{aligned} \log |f_h^{\pm,j}(z)| &= \log |g_h(z)| + \sum_k \log |z - y_k^{\pm,j}| \\ &\leq C_{\epsilon} h^{-d+1-\epsilon} + \#\{y_k^{\pm,j}\} M \log \frac{1}{h} \\ &\leq 2C_{\epsilon,A} h^{-d+\kappa-2\epsilon}. \end{aligned} \quad (3.26)$$

We claim that there exists $3\theta_0 < \mu_1 < 4\theta_0$ such that the distance between $\{y_k^{\pm,j}\}$ and the circle $\partial B_j^{\pm}(\mu_1)$ is greater than h^M , provided $M \gg d$. Indeed, if we suppose the contrary, this would imply that the length of the interval

$$J_j^{\pm} := \mathbb{R} \cap (B_j^{\pm}(4\theta_0) \setminus B_j^{\pm}(3\theta_0))$$

is upper bounded by $\#\{y_k^{\pm,j} \in J_j^{\pm}\} h^M = \mathcal{O}(h^{M-d})$, which is impossible if M is taken large enough. This proves the claim. Thus, by (3.26) we have the estimate $\log |f_h^{\pm,j}(z)| = \mathcal{O}_{\epsilon}(h^{-d+\kappa-2\epsilon})$ on $\partial B_j^{\pm}(\mu_1)$, which in turn implies $\log |f_h^{\pm,j}(z)| = \mathcal{O}_{\epsilon}(h^{-d+\kappa-2\epsilon})$ on $B_j^{\pm}(3\theta_0)$. Combining this with (3.25) and the Jensen theorem (see for example 3.6 in [26]), yields for the zeros $z_k^{\pm,j}$ in $B_j^{\pm}(2\theta_0)$ the following bound

$$\#\{z_k^{\pm,j} : z_k^{\pm,j} \in B_j^{\pm}(2\theta_0)\} \leq C_{\epsilon,A} h^{-d+\kappa-2\epsilon}. \quad (3.27)$$

Since the left-hand side of (3.23) is upper bounded by the number of the zeros and the poles of the function $g_h(z)$ in $B_1^+(\theta_0) \cup B_1^-(\theta_0) \cup B_2^+(\theta_0) \cup B_2^-(\theta_0)$, the estimate (3.23) follows from (3.24) and (3.27). \square

Remark 3.9. The bound (3.27) of the number of the zeros $z_k^{\pm, j}$ of $g_h(z)$ in $B_j^{\pm}(2\theta_0)$ does not depend on the statement of Lemma 3.8 but only on the application of the Jensen theorem based on (3.25) and (3.26). We will use (3.27) in the proof of Lemma 3.8 below.

Proof of Lemma 3.8. We will consider only the case $j = 1$, since the case $j = 2$ is similar. Introduce the function

$$\zeta_h(z) := g_h(z) \prod_{w \in \mathcal{M}_1} (z - w)^{-1} \prod_{w \in \mathcal{M}_2} (z - w),$$

where $\mathcal{M}_1 = \{z_k^{+, 1}\} \cup \{z_k^{-, 1}\}$ is the set of all zeros of $g_h(z)$ in $B_1^-(2\theta_0) \cup B_1^+(2\theta_0)$ and $\mathcal{M}_2 = \{y_k^{+, 1}\} \cup \{y_k^{-, 1}\}$ is the set of all poles of $g_h(z)$ in $B_1^-(4\theta_0) \cup B_1^+(4\theta_0)$. Since $\zeta_h(z)$ does not have zeros and poles in $B_1^-(2\theta_0) \cup B_1^+(2\theta_0)$, the function $\log \zeta_h(z)$ is holomorphic in $B_1^-(2\theta_0) \cup B_1^+(2\theta_0)$. We need the following

Lemma 3.10. *The function $\zeta_h(z)$ satisfies the bound*

$$\log |\zeta_h(z)| \leq C_\epsilon h^{-d+1-2\epsilon} \quad \text{for all } z \in B_1^-(\theta) \cup B_1^+(\theta), \quad (3.28)$$

for every $0 < \theta < 2\theta_0$ independent of h .

Proof. Set $U = \bigcup_{w \in \mathcal{M}} \{z \in \mathbb{C} : |z - w| \leq h^M\}$, where $M \gg d$ and $\mathcal{M} = \mathcal{M}_1 \cup \mathcal{M}_2$. Clearly, $U = \bigcup_\nu U_\nu$, where every U_ν is a domain with a piecewise smooth boundary and $U_\nu \cap U_\mu = \emptyset$ if $\nu \neq \mu$. Moreover, we have

$$\sum_\nu \text{measure}(\partial U_\nu) \leq 2\pi h^M \#\{w \in \mathcal{M}\} \leq Ch^{M-d}.$$

Let $\theta < \theta_1 < 2\theta_0$ be independent of h . Let $\{U_{\nu_i}^\pm\}$ be the set of all U_ν such that $U_\nu \cap \partial B_1^\pm(\theta_1) \neq \emptyset$. We now construct a closed curve, $\beta_1^\pm(\theta_1)$ as follows: we keep all arcs on $\partial B_1^\pm(\theta_1)$ having no common points with $\{U_{\nu_i}^\pm\}$ and replace the arc $\partial B_1^\pm(\theta_1) \cap U_{\nu_i}^\pm$ with arcs on $\partial U_{\nu_i}^\pm$ connecting the corresponding end points. Thus we can guarantee that $\beta_1^\pm(\theta_1)$ belongs to an $\mathcal{O}(h^{M-d})$ neighborhood of $\partial B_1^\pm(\theta_1)$ and, moreover, the distance between $\beta_1^\pm(\theta_1)$ and the set \mathcal{M} is greater than h^M . In the same way, as in the proof of (3.25) and (3.26) above, by using (3.11), (3.14), (3.24), and (3.27), we get

$$\log |\zeta_h(z)| \leq C_\epsilon h^{-d+1-2\epsilon} \quad \text{for all } z \in \beta_1^\pm(\theta_1). \quad (3.29)$$

Since $B_1^\pm(\theta)$ is in the interior of the domain bounded by $\beta_1^\pm(\theta_1)$, the estimate (3.29) implies (3.28). \triangle

We will now construct the curve $\tilde{\gamma}_1$. Let $\{U_{v_i}\}$ be the set of all U_v such that $U_v \cap [\tilde{w}_1^-, \tilde{w}_1^+] \neq \emptyset$. We keep all segments on $[\tilde{w}_1^-, \tilde{w}_1^+]$ having no common points with $\{U_{v_i}\}$ and replace the segments on $[\tilde{w}_1^-, \tilde{w}_1^+] \cap U_{v_i}$ with arcs on ∂U_{v_i} connecting the corresponding end points. Thus we get a piecewise smooth curve $\tilde{\gamma}_1$ belonging to an $\mathcal{O}(h^{M-d})$ neighborhood of $[\tilde{w}_1^-, \tilde{w}_1^+]$ and the distance between $\tilde{\gamma}_1$ and the set \mathcal{M} is greater than h^M . Hence $\tilde{\gamma}_1 \subset \Theta_1$. Now we can write

$$\begin{aligned} \int_{\tilde{\gamma}_1} \frac{d}{dz} \log g_h(z) dz &= \int_{[\tilde{w}_1^-, \tilde{w}_1^+]} \frac{d}{dz} \log \zeta_h(z) dz + \sum_{w \in \mathcal{M}_1} \int_{\tilde{\gamma}_1} (z-w)^{-1} dz \\ &\quad - \sum_{w \in \mathcal{M}_2} \int_{\tilde{\gamma}_1} (z-w)^{-1} dz. \end{aligned} \quad (3.30)$$

We will show that

$$\left| \frac{d}{dz} \log \zeta_h(z) \right| \leq C_\epsilon h^{-d+1-\kappa-\epsilon} \quad \text{for all } z \in \Theta_1, \quad (3.31)$$

$$\left| \operatorname{Im} \int_{\tilde{\gamma}_1} (z-w)^{-1} dz \right| \leq 3\pi \quad \text{for all } w \in \mathcal{M}. \quad (3.32)$$

Since the length of the interval $[\tilde{w}_1^-, \tilde{w}_1^+]$ is $6h^{\kappa-\epsilon}$, the estimate (3.31) implies that the absolute value of the first integral on the right-hand side of (3.30) is $\mathcal{O}_\epsilon(h^{-d+1-2\epsilon})$. Thus, (3.18) would follow from (3.30), (3.31), (3.32), and the bounds (3.24) and (3.27).

To prove (3.31), we apply the Caratheodory theorem (see 5.5 in [26]) for the derivative of the function $f_\pm(z) = \log \frac{\zeta_h(z)}{\zeta_h(\tilde{w}_1^\pm)}$. Note that $\log |\zeta_h(\tilde{w}_1^\pm)|$ can be bounded from below in the same way as in (3.25) above. Therefore, applying (3.28), we get for the real part of $f_\pm(z)$ the estimate

$$\operatorname{Re} f_\pm(z) = \log |\zeta_h(z)| - \log |\zeta_h(\tilde{w}_1^\pm)| \leq Ch^{-d+1-2\epsilon} \quad \text{for all } z \in \partial B_1^\pm\left(\frac{3}{2}\theta_0\right).$$

Since $f_\pm(\tilde{w}_1^\pm) = 0$, we conclude by the Caratheodory theorem that $|f'_\pm(z)| = \mathcal{O}_\epsilon(h^{-d+1-\kappa-\epsilon})$ in the disc $B_1^\pm(\theta_0)$, which clearly implies (3.31).

To establish (3.32), observe that if w does not lie on the line connecting the points \tilde{w}_1^- and \tilde{w}_1^+ and if $\sigma_0 > 0$ denotes the distance from w to this line, after a suitable change of variables, we have

$$\left| \operatorname{Im} \int_{\tilde{w}_1^-}^{\tilde{w}_1^+} (z-w)^{-1} dz \right| = \int_a^b \frac{\sigma_0 d\sigma}{\sigma_0^2 + \sigma^2} \leq \int_{-\infty}^{\infty} \frac{d\sigma}{1 + \sigma^2} = \pi. \quad (3.33)$$

Since the integral in the left-hand side of (3.32) differs from the integral in the left-hand side of (3.33) either by 0 or $2\pi i$, the estimate (3.33) implies (3.32) in this case. If w lies on the line connecting the points \tilde{w}_1^- and \tilde{w}_1^+ , then the integral on the left-hand side of (3.32) is a limit of integrals of the first kind, and hence (3.32) will be true in this case, too. This completes the proof of Lemma 3.8. \square

Proof of Theorem 1.1. Let κ be as described in Theorem 1.1. Let A_1 and A_2 be arbitrary real numbers, independent of h , and let $A > \max\{|A_1|, |A_2|\}$ be independent of h . It follows from the proof of Proposition 3.7 (see (3.27)) that

$$\begin{aligned} \#\{z_k, z_k/h^2 \text{ is (ITE)}: 1 - Ah^{\kappa-\epsilon} \leq |\operatorname{Re} z_k| \leq 1 + Ah^{\kappa-\epsilon}, |\operatorname{Im} z_k| \leq \mathcal{O}(h^{\kappa-\epsilon})\} \\ = \mathcal{O}_{\epsilon, A}(h^{-d+\kappa-2\epsilon}), \end{aligned}$$

$$\begin{aligned} \#\{z_k, z_k/h^2 \text{ is (ITE)}: 2 - Ah^{\kappa-\epsilon} \leq |\operatorname{Re} z_k| \leq 2 + Ah^{\kappa-\epsilon}, |\operatorname{Im} z_k| \leq \mathcal{O}(h^{\kappa-\epsilon})\} \\ = \mathcal{O}_{\epsilon, A}(h^{-d+\kappa-2\epsilon}). \end{aligned}$$

Therefore, by (3.17) we get for every $0 < \epsilon \ll 1$

$$\begin{aligned} \#\{z_k, z_k/h^2 \text{ is (ITE)}: 1 - A_1 h^{\kappa-\epsilon} \leq |\operatorname{Re} z_k| \leq 2 + A_2 h^{\kappa-\epsilon}, |\operatorname{Im} z_k| \leq \mathcal{O}(h^{\kappa-\epsilon})\} \\ = (2^{d/2} - 1)(\tau_1 + \tau_2)h^{-d} + \mathcal{O}_{\epsilon, A_1, A_2}(h^{-d+\kappa-3\epsilon}), \quad 0 < h \leq h_1(A_1, A_2, \epsilon). \end{aligned}$$

Choose $h = \frac{\sqrt{2}}{r}$, $r \gg 1$. The above asymptotics yields

$$\begin{aligned} \left\{ \lambda \in \mathbb{C}: \lambda \text{ is (ITE)}, \frac{r^2}{2} - A_1 r^{2-\kappa+\epsilon} \leq |\operatorname{Re} \lambda| \leq r^2 + A_2 r^{2-\kappa+\epsilon}, |\operatorname{Im} \lambda| \leq r^{2-\kappa+\epsilon} \right\} \\ = (1 - 2^{-d/2})(\tau_1 + \tau_2)r^d + \mathcal{O}_{\epsilon, A_1, A_2}(r^{d-\kappa+3\epsilon}), \quad r \geq r_1(A_1, A_2, \epsilon). \end{aligned}$$

Recall that according to the results in [27], there are no (ITE) in the region

$$\left\{ \lambda \in \mathbb{C}: \frac{r^2}{2} \leq |\lambda| \leq r^2, |\operatorname{Im} \lambda| \geq r^{2-\kappa+\epsilon} \right\}$$

for every $0 < \epsilon \ll 1$, provided $r \geq r_0(\epsilon) \gg 1$. On the other hand, it is clear that the region

$$\left\{ \lambda \in \mathbb{C}: \frac{r^2}{2} \leq |\lambda| \leq r^2, |\operatorname{Im} \lambda| \leq r^{2-\kappa+\epsilon} \right\}$$

is contained in the region

$$\left\{ \lambda \in \mathbb{C}: \frac{r^2}{2} - r^{2-\kappa+\epsilon} \leq |\operatorname{Re} \lambda| \leq r^2, |\operatorname{Im} \lambda| \leq r^{2-\kappa+\epsilon} \right\}$$

and contains the region

$$\left\{ \lambda \in \mathbb{C}: \frac{r^2}{2} \leq |\operatorname{Re} \lambda| \leq r^2 - r^{2-\kappa+\epsilon}, |\operatorname{Im} \lambda| \leq r^{2-\kappa+\epsilon} \right\}.$$

Thus we get the asymptotics

$$N(r) - N(r/\sqrt{2}) = (1 - 2^{-d/2})(\tau_1 + \tau_2)r^d + \mathcal{O}_\varepsilon(r^{d-\kappa+\varepsilon}) \quad r \geq r_0(\varepsilon), \quad (3.34)$$

for every $0 < \varepsilon \ll 1$. Here we replace 3ε by ε , which is not important since our argument works for every $0 < \varepsilon \ll 1$. The asymptotics (3.34) yields

$$\begin{aligned} N(r/2^{k/2}) - N(r/2^{(k+1)/2}) \\ = (2^{-kd/2} - 2^{-(k+1)d/2})(\tau_1 + \tau_2)r^d + 2^{-kd/2}\mathcal{O}_\varepsilon(r^{d-\kappa+\varepsilon}) \end{aligned} \quad (3.35)$$

for every integer $k \geq 0$ such that $r2^{-k/2} \geq r_0(\varepsilon)$. Let $k_0(r) \in \mathbb{N}$ be the smallest integer such that $r2^{-k_0(r)/2} < r_0(\varepsilon)$. It is clear that we have

$$N(r/2^{(k_0(r)+1)/2}) \leq N(r_0(\varepsilon)) = R_0(\varepsilon) \quad (3.36)$$

with a constant $R_0(\varepsilon) > 0$ independent of r . Moreover,

$$(2^{-(k_0(r)+1)/2}r)^d \leq (r_0(\varepsilon))^d = R_1(\varepsilon)$$

with $R_1(\varepsilon) > 0$ independent of r . Summing up the asymptotics (3.35) and using (3.36), we get (1.7). Thus the proof of Theorem 1.1 is complete. \square

References

- [1] L. Audibert and H. Haddar, A generalized formulation of the linear sampling method with exact characterization of targets in terms of farfield measurements. *Inverse Problems* **30** (2014), no. 3, article id. 035011. [MR 3168273](#) [Zbl 1291.35377](#)
- [2] K. Kh. Boimatov and A. G. Kostyuchenko, Spectral asymptotics of nonselfadjoint elliptic systems of differential operators in bounded domains. *Mat. Sb.* **181** (1990), no. 12, 1678–1693. In Russian. English translation, *Math. USSR-Sb.* **71** (1992), no. 2, 517–531. [MR 1099521](#) [Zbl 0776.35041](#)
- [3] F. Caconi and H. Haddar, Transmission eigenvalues in inverse scattering theory. In G. Uhlmann (ed.) *Inverse problems and applications: inside out. II*. Mathematical Sciences Research Institute Publications, 60. Cambridge University Press, Cambridge, 2013, 529–580. [MR 3135766](#) [Zbl 1316.35297](#)
- [4] F. Cardoso, G. Popov, and G. Vodev, Asymptotics of the number of resonances in the transmission problem. *Comm. Partial Differential Equations* **26** (2001), no. 9-10, 1811–1859. [MR 1865946](#) [Zbl 1086.35012](#)

- [5] D. Colton and A. Kirsch, A simple method for solving inverse scattering problems in the resonance region. *Inverse Problems* **12** (1996), no. 4, 383–393. [MR 1402098](#) [Zbl 0859.35133](#)
- [6] D. Colton and Y. J. Leung, Complex eigenvalues and inverse spectral problem for transmission eigenvalues. *Inverse Problems* **29** (2013), no. 10, article id. 104008. [MR 3116203](#) [Zbl 1305.34027](#)
- [7] M. Dimassi and J. Sjöstrand, *Spectral asymptotics in semi-classical limit*. London Mathematical Society Lecture Note Series, 268. Cambridge University Press, Cambridge, 1999. [MR 1735654](#) [Zbl 0926.35002](#)
- [8] M. Dimassi and V. Petkov, Upper bound for the counting function of interior transmission eigenvalues. Preprint 2013 [arXiv:1308.2594](#) [math.SP]
- [9] M. Fairman, The interior transmission problem: spectral theory. *SIAM J. Math. Anal.* **46** (2014), no. 1, 803–819. [MR 3165910](#) [Zbl 1294.35049](#)
- [10] M. Hitrik, K. Krupchyk, P. Ola, and L. Päivärinta, The interior transmission problem and bounds on transmission eigenvalues. *Math. Res. Lett.* **18** (2011), no. 2, 279–293. [MR 2784672](#) [Zbl 1241.47057](#)
- [11] M. Hitrik, K. Krupchyk, P. Ola and L. Päivärinta, Transmission eigenvalues for elliptic operators. *SIAM J. Math. Anal.* **43** (2011), no. 6, 2630–2639. [MR 2873234](#) [Zbl 1233.35148](#)
- [12] T. Kato, *Perturbation theory for linear operators*. Die Grundlehren der mathematischen Wissenschaften, 132. Springer-Verlag New York, New York, 1966. [MR 0203473](#) [Zbl 0148.12601](#)
- [13] E. Lakshtanov and B. Vainberg, Remarks on interior transmission eigenvalues, Weyl formula and branching billiards. *J. Phys. A* **45** (2012), no. 12, article id. 125202. [MR 2902328](#) [Zbl 1245.81290](#)
- [14] E. Lakshtanov and B. Vainberg, Bounds on positive interior transmission eigenvalues. *Inverse Problems* **28** (2012), no. 10, article id. 105005. [MR 2987908](#) [Zbl 1256.35036](#)
- [15] E. Lakshtanov and B. Vainberg, Applications of elliptic operator theory to the isotropic interior transmission eigenvalue problem. *Inverse Problems* **29** (2013), no. 10, article id. 104003. [MR 3116198](#) [Zbl 1285.35059](#)
- [16] E. Lakshtanov and B. Vainberg, Weyl type bound on positive interior transmission eigenvalues. *Comm. Partial Differential Equations* **39** (2014), no. 9, 1729–1740. [MR 3246041](#) [Zbl 1301.35078](#)
- [17] P. Malliavin, Un théorème taubérien relié aux estimations des valeurs propres. *Séminaire Jean Leray* (1962–1963), p. 234–231. http://www.numdam.org/item?id=SJL_1962-1963___224_0
- [18] H. Pham and P. Stefanov, Weyl asymptotics of the transmission eigenvalues for a constant index of refraction. *Inverse Probl. Imaging* **8** (2014), no. 3, 795–810. [MR 3295945](#) [Zbl 06377647](#)

- [19] L. Robbiano, Spectral analysis of the interior transmission eigenvalue problem. *Inverse Problems* **29** (2013), no. 10, article id. 104001. [MR 3116196](#) [Zbl 1296.35105](#)
- [20] L. Robbiano, Counting function for interior transmission eigenvalues. *Math. Control Relat. Fields* **6** (2016), no. 1, 167–183. [MR 3448675](#) [Zbl 1332.35242](#)
- [21] J. Sjöstrand, *Lectures on resonances*.
<http://sjostrand.perso.math.cnrs.fr/Coursgbg.pdf>
- [22] J. Sjöstrand, Weyl law for semi-classical resonances with randomly perturbed potentials. *Mém. Soc. Math. Fr. (N.S.)* **136** (2014). [MR 3288114](#) [Zbl 1304.35010](#)
- [23] J. Sjöstrand and G. Vodev, Asymptotics of the number of Rayleigh resonances. *Math. Ann.* **309** (1997), no. 2, 287–306. [MR 1474193](#) [Zbl 0890.35098](#)
- [24] J. Sylvester, Discreteness of transmission eigenvalues via upper triangular compact operators. *SIAM J. Math. Anal.* **44** (2012), no. 1, 341–354. [MR 2888291](#) [Zbl 1238.81172](#)
- [25] J. Sylvester, Transmission eigenvalues in one dimension. *Inverse Problems* **29** (2013), no. 10, article id. 104009. [MR 3116204](#) [Zbl 1294.34079](#)
- [26] E. C. Titchmarsh, *The theory of functions*. Second Edition, Oxford University Press, Oxford, 1939. [JFM 65.0302.01](#)
- [27] G. Vodev, Transmission eigenvalue-free regions. *Comm. Math. Phys.* **336** (2015), no. 3, 1141–1166. [MR 3324140](#) [Zbl 1323.35110](#)
- [28] G. Vodev, Transmission eigenvalues for strictly concave domains. To appear in *Math. Ann.* [doi:10.1007/s00208-015-1329-2](https://doi.org/10.1007/s00208-015-1329-2)

Received September 20, 2014; revised December 14, 2014

Vesselin Petkov, Université de Bordeaux, Institut de Mathématiques de Bordeaux,
351, Cours de la Libération, 33405 Talence, France

e-mail: petkov@math.u-bordeaux1.fr

Georgi Vodev, Université de Nantes, Département de Mathématiques,
2, rue de la Houssinière, 44322 Nantes Cedex, France

e-mail: vodev@math.univ-nantes.fr