

**Green’s function asymptotics
near the internal edges
of spectra of periodic elliptic operators.
Spectral gap interior**

Minh Kha,¹ Peter Kuchment,^{1,2} and Andrew Raich³

Abstract. Precise asymptotics known for the Green function of the Laplacian have found their analogs for bounded below periodic elliptic operators of the second-order below and at the bottom of the spectrum. Due to the band-gap structure of the spectra of such operators, the question arises whether similar results can be obtained near or at the edges of spectral gaps. In a previous work, two of the authors considered the case of a spectral edge. The main result of this article is finding such asymptotics near a gap edge, for “generic” periodic elliptic operators of second-order with real coefficients in dimension $d \geq 2$, when the gap edge occurs at a symmetry point of the Brillouin zone.

Mathematics Subject Classification (2010). 35P99, 35J10, 35J15, 81Q10.

Keywords. Periodic operator, spectrum, Green’s function, dispersion relation.

¹ The author was partially supported by the NSF grant DMS-1517938.

² The author would like to thank the Isaac Newton Institute for Mathematical Sciences for its hospitality during the programme Periodic and Ergodic Problems supported by EPSRC Grant Number EP/K032208/1, where work on this paper was undertaken.

³ The author as partially funded by NSF grant DMS-1405100.

Contents

1	Introduction	1172
2	Assumptions, notation and the main result	1173
3	Proof of the main theorem 2.11 and some remarks	1181
4	On local geometry of the resolvent set	1183
5	A Floquet reduction of the problem	1183
6	Asymptotics of the Green's function	1190
7	The full Green's function asymptotics	1206
8	Some results on parameter-dependent toroidal Ψ DOs	1215
9	Some auxiliary statements	1219
10	Concluding remarks	1230
	References	1230

1. Introduction

The behavior at infinity of the Green function of the Laplacian in \mathbb{R}^n outside and at the boundary of its spectrum is well known. Analogous results below and at the lower boundary of the spectrum have been established for bounded below periodic elliptic operators of the second order in [3, 25] (see also [35] for the discrete version). Due to the band-gap structure of the spectra of such periodic operators, the question arises whether similar results can be obtained at or near the edges of spectral gaps. The corresponding result at the internal edges of the spectrum was established in [23]. The main result of this article, Theorem 2.11, is the description of such asymptotics near the spectral edge for generic periodic elliptic operators of second-order with real coefficients in dimension $d \geq 2$, if the spectral edge is attained at a symmetry point of the Brillouin zone.

It is well known that outside of the spectrum the Green function decays exponentially at infinity, with the rate of decay controlled by the distance to the spectrum. See, e.g., well known Combes-Thomas estimates [7, 4]. However, comparison with the formulas for the case of the Laplacian shows that an additional algebraically decaying factor (depending on the dimension) is lost in this approach. Moreover, the exponential decay in general is expected to be anisotropic, while the operator theory approach can provide only isotropic estimates. The result of this paper provides the exact principal term of asymptotics, thus resolving these issues.

2. Assumptions, notation and the main result

Consider a linear second order elliptic operator in \mathbb{R}^d with periodic coefficients

$$L(x, D) = \sum_{k,l=1}^d D_k(a_{kl}(x)D_l) + V(x) = D^*A(x)D + V(x). \quad (1)$$

Here $A = (a_{kl})_{1 \leq k,l \leq d}$, $D = (D_1, \dots, D_d)$, and

$$D_k := -i \partial_k = -i \frac{\partial}{\partial x_k}.$$

All coefficients a_{kl} , V are smooth real-valued functions on \mathbb{R}^d , periodic with respect to the integer lattice \mathbb{Z}^d in \mathbb{R}^d , i.e., $a_{kl}(x+n) = a_{kl}(x)$ and $V(x+n) = V(x)$, for all $x \in \mathbb{R}^d$, $n \in \mathbb{Z}^d$. The operator L is assumed to be elliptic, i.e., the matrix A is symmetric and

$$\sum_{k,l=1}^d a_{kl}(x)\xi_k\xi_l \geq \theta|\xi|^2, \quad (2)$$

for some $\theta > 0$ and any $x \in \mathbb{R}^d$, $\xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d$. The operator L , with the Sobolev space $H^2(\mathbb{R}^d)$ as the domain, is an unbounded, self-adjoint operator in $L^2(\mathbb{R}^d)$ (see e.g., [30]).

The spectrum of the above operator L in $L^2(\mathbb{R}^d)$ has a *band-gap structure* [8, 19, 20, 27], i.e., it is the union of a sequence of closed bounded intervals (*bands* or *stability zones* of the operator L) $[\alpha_j, \beta_j] \subset \mathbb{R}$ ($j = 1, 2, \dots$):

$$\sigma(L) = \bigcup_{j=1}^{\infty} [\alpha_j, \beta_j], \quad (3)$$

such that $\alpha_j \leq \alpha_{j+1}$, $\beta_j \leq \beta_{j+1}$ and $\lim_{j \rightarrow \infty} \alpha_j = \infty$. The bands can (and do) overlap when $d > 1$, but they may leave open intervals in between, called *spectral gaps*. Thus, a spectral gap is an interval of the form (β_j, α_{j+1}) for some $j \in \mathbb{N}$ for which $\alpha_{j+1} > \beta_j$. We make a convention that the open interval $(-\infty, \alpha_1)$, which contains all real numbers below the bottom of the spectrum of L , is also a spectral gap. However, we will be mostly interested in finite spectral gaps.

In this text, we study Green's function asymptotics for the operator L in a spectral gap, near to a spectral gap edge. More precisely, consider a finite spectral gap (β_j, α_{j+1}) for some $j \in \mathbb{N}$ and a value $\lambda \in (\beta_j, \alpha_{j+1})$ which is close either to the spectral edge β_j or to the spectral edge α_{j+1} . We would like to study the asymptotic behavior when $|x - y| \rightarrow \infty$ of the Green's kernel

$G_\lambda(x, y)$ of the resolvent operator $R_{\lambda,L} := (L - \lambda)^{-1}$. The case of the spectral edges (i.e., $\lambda = \alpha_{j+1}$ or $\lambda = \beta_j$) was studied for the similar purpose in [23]. All asymptotics here and also in [23] are deduced from an assumed “generic” spectral edge behavior of the *dispersion relation* of the operator L , which we will briefly review below.

Let $W = [0, 1]^d \subset \mathbb{R}^d$ be the unit cube, which is a *fundamental domain* of \mathbb{R}^d with respect to the lattice \mathbb{Z}^d (*Wigner–Seitz cell*). The *dual* (or *reciprocal*) *lattice* is $2\pi\mathbb{Z}^d$ and its fundamental domain is $[-\pi, \pi]^d$ (*Brillouin zone*).

The d -dimensional tori with respect to the lattices \mathbb{Z}^d and $2\pi\mathbb{Z}^d$ are denoted by $\mathbb{T} := \mathbb{R}^d/\mathbb{Z}^d$ and $\mathbb{T}^* := \mathbb{R}^d/2\pi\mathbb{Z}^d$, respectively.

Definition 2.1. For any $k \in \mathbb{C}^d$, the subspace $H_k^s(W) \subset H^s(W)$ consists of restrictions to W of functions $f \in H_{\text{loc}}^s(\mathbb{R}^d)$ that satisfy for any $\gamma \in \mathbb{Z}^d$ the *Floquet–Bloch condition* (also known as automorphicity condition or cyclic condition)

$$f(x + \gamma) = e^{ik \cdot \gamma} f(x) \quad \text{for a.e } x \in W. \tag{4}$$

Here H^s denotes the standard Sobolev space of order s . Note that when $s = 0$, the above space coincides with $L^2(W)$ for any k .

Due to periodicity, the operator $L(x, D)$ preserves condition (4) and thus, it defines an operator $L(k)$ in $L^2(W)$ with the domain $H_k^2(W)$. In this model, $L(k)$ is realized as a k -independent differential expression $L(x, D)$ acting on functions in W with boundary conditions depending on k (which can be identified with sections of a linear bundle over the torus \mathbb{T}). An alternative definition of $L(k)$ is as the operator $L(x, D + k)$ in $L^2(\mathbb{T})$ with the domain $H^2(\mathbb{T})$. In the latter model, $L(k)$ acts on the k -independent domain of periodic functions on W as follows:

$$e^{-ik \cdot x} L(x, D)e^{ik \cdot x} = (D + \bar{k})^* A(x)(D + k) + V(x). \tag{5}$$

We use the latter model of $L(k)$ throughout this paper, unless specified differently.

Note that the condition (4) is invariant under translations of k by elements of the dual lattice $2\pi\mathbb{Z}^d$. Moreover, the operator $L(k)$ is unitarily equivalent to $L(k + 2\pi\gamma)$, for any $\gamma \in \mathbb{Z}^d$. In particular, when dealing with real values of k , it suffices to restrict k to the Brillouin zone $[-\pi, \pi]^d$ (or any its shifted copy). It is well-known (see [8, 19, 20, 27]) that the spectrum of L is the union of all the spectra of $L(k)$ when k runs over the Brillouin zone, i.e.

$$\sigma(L) = \bigcup_{k \in [-\pi, \pi]^d} \sigma(L(k)). \tag{6}$$

Hence, the spectrum of L is the range of the multivalued function

$$k \mapsto \lambda(k) := \sigma(L(k)), \quad k \in [-\pi, \pi]^d. \quad (7)$$

By (5), $L(k)$ is self-adjoint in $L^2(\mathbb{T})$ and has domain $H^2(\mathbb{T})$ for each $k \in \mathbb{R}^d$. By the ellipticity of L , each $L(k)$ is bounded from below and has compact resolvent. This forces each of the operators $L(k)$, $k \in \mathbb{R}^d$, to have discrete spectrum in \mathbb{R} . Therefore, we can label its eigenvalues in non-decreasing order:

$$\lambda_1(k) \leq \lambda_2(k) \leq \dots. \quad (8)$$

Hence, we can single out continuous and piecewise-analytic *band functions* $\lambda_j(k)$ for each $j \in \mathbb{N}$, see [34]. The range of the band function λ_j constitutes exactly the band $[\alpha_j, \beta_j]$ of the spectrum of L shown in (3).

Definition 2.2. A *Bloch solution* of the equation $L(x, D)u = 0$ is a solution of the form

$$u(x) = e^{ik \cdot x} \phi(x),$$

where the function ϕ is 1-periodic in each variable x_j for $j = 1, \dots, d$. The vector k is the *quasimomentum* and $z = e^{ik} = (e^{ik_1}, \dots, e^{ik_d})$ is the *Floquet exponent* (or *Floquet multiplier*) of the solution. In our formulation, allowing quasimomenta k to be complex is essential.

Definition 2.3. The (complex) *Bloch variety* B_L of the operator L consists of all pairs $(k, \lambda) \in \mathbb{C}^{d+1}$ such that the equation $Lu = \lambda u$ in \mathbb{R}^d has a non-zero Bloch solution u with a quasimomentum k . Similarly, the *real Bloch variety* $B_{L, \mathbb{R}}$ is $B_L \cap \mathbb{R}^{d+1}$.

The Bloch variety B_L can be treated as the *dispersion relation/curve*, i.e., the graph of the multivalued function $\lambda(k)$:

$$B_L = \{(k, \lambda) : \lambda \in \sigma(L(k))\}.$$

Note that $L(k)$ is non-self-adjoint if $k \notin \mathbb{R}^d$. However, $L(k) - L(0)$ is an operator of lower order for each $k \in \mathbb{C}^d$. Therefore, the spectra of all operators $L(k)$ on the torus \mathbb{T} are discrete (see pp.188-190 in [1]).

Remark 2.4. In fact, the main techniques of Floquet theory (e.g., (6)) apply to non-self-adjoint operators. It is required only that the operators $L(k) = L(x, D + k) : H^2(\mathbb{T}) \rightarrow L^2(\mathbb{T})$ are Fredholm for $k \in \mathbb{C}^d$. The latter condition is always satisfied due to ellipticity and embedding theorems (see Theorem 2.1 in [19]). Unlike the self-adjoint case though, we do not have the band-gap structure as in (3).

Definition 2.5. The (complex) *Fermi surface* $F_{L,\lambda}$ of the operator L at the energy level $\lambda \in \mathbb{C}$ consists of all quasimomenta $k \in \mathbb{C}^d$ such that the equation $Lu = \lambda u$ in \mathbb{R}^d has a non-zero Bloch solution u with a quasimomentum k . For $\lambda = 0$, we simply write F_L instead of $F_{L,0}$. The *real Fermi surface* $F_{L,\mathbb{R}}$ is $F_L \cap \mathbb{R}^d$.

Equivalently, $k \in F_{L,\lambda}$ means the existence of a nonzero periodic solution u of the equation $L(k)u = \lambda u$. In other words, Fermi surfaces are level sets of the dispersion relation.

The following result can be found in Theorem 3.1.7 in [19].

Lemma 2.6. *There exist entire ($2\pi\mathbb{Z}^d$ -periodic in k) functions of finite orders on \mathbb{C}^d and on \mathbb{C}^{d+1} such that the Fermi and Bloch varieties are the sets of all zeros of these functions respectively.*

From this lemma and the proof of Lemma 4.5.1 in [19] (see also [34]), the band functions $\lambda(k)$ are piecewise analytic on \mathbb{C}^d .

From now on, we fix L as a self-adjoint elliptic operator of the form (1), whose band-gap structure is as (3). By adding a constant to the operator L if necessary, we can assume that the spectral edge of interest is 0. It is also enough to suppose that the adjacent spectral band is of the form $[0, a]$ for some $a > 0$ since the case when the spectral edge 0 is the maximum of its adjacent spectral band is treated similarly.

Suppose there is no spectrum for small negative values of λ and hence there is a spectral gap below 0. Thus, there exists at least one band function $\lambda_j(k)$ for some $j \in \mathbb{N}$ such that 0 is the minimal value of this function on the Brillouin zone.

To establish our main result, we need to impose the following analytic assumption on the dispersion curve λ_j as in [23].

Assumption A. There exists $k_0 \in [-\pi, \pi]^d$ and a band function $\lambda_j(k)$ such that

- A1. $\lambda_j(k_0) = 0$,
- A2. $\min_{k \in \mathbb{R}^d, i \neq j} |\lambda_i(k)| > 0$,
- A3. k_0 is the only¹ (modulo $2\pi\mathbb{Z}^d$) minimum of λ_j ,
- A4. $\lambda_j(k)$ is a Morse function near k_0 , i.e., its Hessian matrix $H := \text{Hess}(\lambda_j)(k_0)$ at k_0 is positive definite. In particular, the Taylor expansion of λ_j at k_0 is

$$\lambda_j(k) = \frac{1}{2}(k - k_0)^T H(k - k_0) + O(|k - k_0|^3).$$

¹ Finitely many such points can be also easily handled.

It is known [17] that conditions A1 and A2 ‘generically’ hold (i.e., they can be achieved by small perturbation of coefficients of the operator) for Schrödinger operators. Although this has not been proven, conditions A3 and A4 are widely believed (both in the mathematics and physics literature) to hold ‘generically’. In other words, it is conjectured that for a ‘generic’ selfadjoint second-order elliptic operator with periodic coefficients on \mathbb{R}^d each of the spectral gap’s endpoints is a unique (modulo the dual lattice $2\pi\mathbb{Z}^d$), nondegenerate extremum of a single band function $\lambda_j(k)$ (see e.g., Conjecture 5.1 in [22]). It is known that for a non-magnetic periodic Schrödinger operator, the bottom of the spectrum always corresponds to a non-degenerate minimum of λ_1 , see [16]. A similar statement is correct for a wider class of ‘factorable’ operators [5, 6]. The following condition on k_0 will also be needed.

A5. The quasimomentum k_0 is a *high symmetry point of the Brillouin zone*, i.e., all components of k_0 must be either equal to 0 or to π .

We denote by X the set of such high symmetry points in the Brillouin zone.

It is known [14] that condition A5 is not always satisfied and spectral edges could occur deeply inside the Brillouin zone. However, as it is discussed in [14], in many practical cases (e.g., in the media close to homogeneous) this condition holds.

We would like to introduce a suitable fundamental domain with respect to the dual lattice $2\pi\mathbb{Z}^d$ to work with.

Definition 2.7. Consider the quasimomentum k_0 in our assumptions. By A5, $k_0 = (\delta_1\pi, \delta_2\pi, \dots, \delta_d\pi)$, where $\delta_j \in \{0, 1\}$ for $j \in \{1, \dots, d\}$. We denote by \mathcal{O} the fundamental domain so that k_0 is its center of symmetry, i.e.,

$$\mathcal{O} = \prod_{j=1}^d [(\delta_j - 1)\pi, (\delta_j + 1)\pi].$$

When $k_0 = 0$, \mathcal{O} is just the Brillouin zone.

We now introduce notation that will be used throughout the paper.

Notation 2.8. (a) Let $z_1 \in \mathbb{C}$, $z_2 \in \mathbb{C}^{d-1}$, $z_3 \in \mathbb{C}^d$ and r_i be positive numbers for $i = 1, 2, 3$. Then we denote by $B(z_1, r_1)$, $D'(z_2, r_2)$ and $D(z_3, r_3)$ the open balls (or discs) centered at z_1 , z_2 and z_3 whose radii are r_1 , r_2 and r_3 in \mathbb{C} , \mathbb{C}^{d-1} and \mathbb{C}^d respectively.

(b) The real parts of a complex vector z , or of a complex matrix A are denoted by $\Re(z)$ and $\Re(A)$ respectively.

(c) The standard notation $O(|x - y|^{-n})$ for a function f defined on \mathbb{R}^{2d} means there exist constants $C > 0$ and $R > 0$ such that $|f(x, y)| \leq C|x - y|^{-n}$ whenever $|x - y| > R$. Also, $f(x, y) = o(|x - y|^{-n})$ means that

$$\lim_{|x-y| \rightarrow \infty} \frac{|f(x, y)|}{|x - y|^n} = 0.$$

(d) We often use the notation $A \lesssim B$ to mean that the quantity A is less or equal than the quantity B up to some multiplicative constant factor, which does not affect the arguments.

As we discussed, for each $z \in \mathbb{C}^d$, the operator $L(z)$ has discrete spectrum and is therefore a closed operator with non-empty resolvent set. These operators have the same domain $H^2(\mathbb{T})$ and for each $\phi \in H^2(\mathbb{T})$, $L(z)\phi$ is a $L^2(\mathbb{T})$ -valued analytic function of z , due to (5). Consequently, $\{L(z)\}_{z \in \mathbb{C}^d}$ is an analytic family of type \mathcal{A} in the sense of Kato [18].² Due to A1 and A2, $\lambda_j(k_0)$ is a simple eigenvalue of $L(k_0)$. By using analytic perturbation theory for the family $\{L(z)\}_{z \in \mathbb{C}^d}$ (see e.g., Theorem XII.8 in [27]), there is an open neighborhood V of k_0 in \mathbb{C}^d and some $\epsilon_0 > 0$ such that the following conditions are satisfied.

- P1. λ_j is analytic in a neighborhood of the closure of V .
- P2. $\lambda_j(z)$ has algebraic multiplicity one, i.e., it is a simple eigenvalue of $L(z)$ for any $z \in \bar{V}$.
- P3. The only eigenvalue of $L(z)$ contained in the closed disc $\bar{B}(0, \epsilon_0)$ is $\lambda_j(z)$. Moreover, we may also assume that $|\lambda_j(z)| < \epsilon_0$ for each $z \in V$.
- P4. For each $z \in \bar{V}$, let $\phi(z, x)$ be a nonzero \mathbb{Z}^d -periodic function of x such that it is the unique (up to a constant factor) eigenfunction of $L(z)$ with the eigenvalue $\lambda_j(z)$, i.e., $L(z)\phi(z, \cdot) = \lambda_j(z)\phi(z, \cdot)$. We will also use sometimes the notation ϕ_z for the eigenfunction $\phi(z, \cdot)$.

By elliptic regularity, $\phi(z, x)$ is smooth in x . On a neighborhood of \bar{V} , $\phi(z, \cdot)$ is a $H^2(\mathbb{T})$ -valued holomorphic function.

- P5. By condition A4 and the continuity of³ $\text{Hess}(\lambda_j)$, we can assume that for all $z \in V$,

$$2\Re(\text{Hess}(\lambda_j)(z)) > \min \sigma(\text{Hess}(\lambda_j)(k_0))I_{d \times d}.$$

² It is also an analytic family in the Banach space of bounded linear operators acting from $H^2(\mathbb{T})$ to $L^2(\mathbb{T})$.

³ The Hessian matrix of λ_j .

P6. V is invariant under complex conjugation. Furthermore, the smooth function

$$F(z) := (\phi(z, \cdot), \phi(\bar{z}, \cdot))_{L^2(\mathbb{T})} \quad (9)$$

is non-zero on V , due to analyticity of the mapping $z \mapsto \phi(z, \cdot)$ and the inequality $F(k_0) = \|\phi(k_0)\|_{L^2(\mathbb{T})}^2 > 0$.

The following lemma will be useful when dealing with operators having real and smooth coefficients.

Lemma 2.9. (i) For k in \mathbb{R}^d and $i \in \mathbb{N}$,

$$\lambda_i(k) = \lambda_i(-k). \quad (10)$$

In other words, each band λ_i of L is an even function on \mathbb{R}^d .

(ii) If $k_0 \in X$, we have $\lambda_i(k + k_0) = \lambda_i(-k + k_0)$ for all k in \mathbb{R}^d and $i \in \mathbb{N}$.

Proof. Let ϕ_k be an eigenfunction of $L(k)$ corresponding to $\lambda_j(k)$. This means that ϕ_k is a periodic solution to the equation

$$L(x, \partial + ik)\phi_k(x) = \lambda_j(k)\phi_k(x). \quad (11)$$

Taking the complex conjugate of (11), we get

$$L(x, \partial - ik)\overline{\phi_k}(x) = \lambda_j(k)\overline{\phi_k}(x).$$

Therefore, $\overline{\phi_k}$ is an eigenfunction of $L(-k)$ with eigenvalue $\lambda_j(k)$. This implies the identity (10).

(ii) By (i), $\lambda_i(k + k_0) = \lambda_i(-k - k_0) = \lambda_i(-k + k_0)$ since $2k_0 \in 2\pi\mathbb{Z}^d$. \square

Corollary 2.10. If $\beta \in \mathbb{R}^d$ such that $k_0 + i\beta \in \bar{V}$ then $\lambda_j(k_0 + i\beta) \in \mathbb{R}$.

Proof. Indeed, the statement (ii) of Lemma 2.9 implies that the Taylor series of $\lambda(k)$ at k_0 has only even degree terms and real coefficients. \square

Corollary 2.10 allows us to define near $\beta = 0$ the real analytic function $E(\beta) := \lambda_j(k_0 + i\beta)$ near 0. Since its Hessian at 0 is negative-definite (by A4), there exists a connected and bounded neighborhood V_0 of 0 in \mathbb{R}^d such that $k_0 + iV_0 \subseteq V$ and $\text{Hess}(E)(\beta)$ is negative-definite whenever β belongs to V_0 . Thus, E is strictly concave on V_0 and $\sup_{\beta \in V_0} E(\beta) = E(0) = 0$, $\nabla E(\beta) = 0$ iff $\beta = 0$. Note that at the bottom of the spectrum (i.e., $j = 1$), we could take V_0 as the whole Euclidean space \mathbb{R}^d .

By the Morse lemma and the fact that 0 is a nondegenerate critical point of E , there is a smooth change of coordinates $\Phi: U_0 \rightarrow \mathbb{R}^d$ so that $0 \in U_0 \subset\subset V_0$, U_0 is connected, $\Phi(0) = 0$ and $E(\Phi^{-1}(a)) = -|a|^2$, for all $a \in \Phi(U_0)$. Set

$$K_\lambda := \{\beta \in U_0: E(\beta) \geq \lambda\}$$

and

$$\Gamma_\lambda := \{\beta \in U_0: E(\beta) = \lambda\}$$

for each $\lambda \in \mathbb{R}$. Now, we consider λ to be in the set $\{-|a|^2: a \in \Phi(U_0), a \neq 0\}$. Then K_λ is a strictly convex d -dimensional compact body in \mathbb{R}^d , and $\Gamma_\lambda = \partial K_\lambda$ is a compact hypersurface in \mathbb{R}^d . The compactness of K_λ follows from the equation $-|\Phi(\beta)|^2 = E(\beta) \geq \lambda$ which yields that $|\beta| = |\Phi^{-1}(\Phi(\beta))| \leq \max\{|\Phi^{-1}(a)|: a \in \Phi(U_0), |a|^2 \leq -\lambda\}$. Additionally, $\lim_{\lambda \rightarrow 0^-} \max_{\beta \in K_\lambda} |\beta| = 0$.

Let \mathcal{K}_λ be the Gauss-Kronecker curvature of Γ_λ . Since the Hessian of E is negative-definite on Γ_λ , \mathcal{K}_λ is nowhere-zero. For the value of λ described in the previous paragraph and each $s \in \mathbb{S}^{d-1}$, there is a unique vector $\beta_s \in \Gamma_\lambda$ such that the value of the Gauss map of the hypersurface Γ_λ at this point coincides with s , i.e.

$$\nabla E(\beta_s) = -|\nabla E(\beta_s)|s. \quad (12)$$

This is due to the fact that the Gauss map of a compact, connected oriented hypersurface in \mathbb{R}^d , whose Gauss-Kronecker curvature is nowhere zero, is a diffeomorphism onto the sphere \mathbb{S}^{d-1} (see e.g., Theorem 5, p.104 in [33] or Corollary 3.1 in [11]). Thus, β_s depends smoothly on s . We also see that

$$\lim_{|\lambda| \rightarrow 0} \max_{s \in \mathbb{S}^{d-1}} |\beta_s| = 0.$$

Note that β_s could be defined equivalently by using the support functional h of the strictly convex set K_λ . Recall that for each direction $s \in \mathbb{S}^{d-1}$,

$$h(s) = \max_{\xi \in K_\lambda} \langle s, \xi \rangle.$$

Then β_s is the unique point in Γ_λ such that $\langle s, \beta_s \rangle = h(s)$.

By letting $|\lambda|$ close enough to 0, we can make sure that $(-\lambda)^{1/2} = |a|$ for some $a \in \Phi(U_0)$. Then

$$\{k_0 + it\beta_s, (t, s) \in [0, 1] \times \mathbb{S}^{d-1}\} \subset V. \quad (13)$$

We can now state the main result of the paper.

Theorem 2.11. *Suppose conditions A1–A5 are satisfied. For $\lambda < 0$ sufficiently close to 0 (depending on the dispersion branch λ_j and the operator L), the Green's function G_λ of L at λ admits the following asymptotics as $|x - y| \rightarrow \infty$:*

$$G_\lambda(x, y) = \frac{e^{(x-y)(ik_0-\beta_s)}}{(2\pi|x-y|)^{(d-1)/2}} \frac{|\nabla E(\beta_s)|^{(d-3)/2}}{\det(-\mathcal{P}_s \text{Hess}(E)(\beta_s)\mathcal{P}_s)^{1/2}} \frac{\overline{\phi_{k_0+i\beta_s}(x)\phi_{k_0-i\beta_s}(y)}}{(\phi_{k_0+i\beta_s}, \phi_{k_0-i\beta_s})_{L^2(\mathbb{T})}} + e^{(y-x)\cdot\beta_s} r(x, y). \quad (14)$$

Here $s = (x - y)/|x - y|$, \mathcal{P}_s is the projection from \mathbb{R}^d onto the tangent space of the unit sphere \mathbb{S}^{d-1} at the point s , and when $|x - y|$ is large enough, the remainder term r satisfies $|r(x, y)| \leq C|x - y|^{-d/2}$ for some constant $C > 0$ (independent of s).

This result achieves our stated goal of showing the precise (anisotropic) rates of the exponential decay of the Green's function and capturing the additional algebraic decay factor.

Remark 2.12. In the original version of this paper, the error estimate was ϵ -worse: $|r(x, y)| \leq C_\epsilon|x - y|^{-d/2+\epsilon}$, for arbitrary $\epsilon > 0$. However, N. Filonov has suggested an improvement of the estimates in our proof, which erases this unnecessary ϵ . We are extremely grateful to him for this.

3. Proof of the main theorem 2.11 and some remarks

Theorem 2.11 is a direct consequence of its local (with respect to the direction of $(x - y)$) version.

Theorem 3.1. *Under the hypotheses of Theorem 2.11 and when $\lambda \approx 0$, for each $\omega \in \mathbb{S}^{d-1}$, there are a neighborhood \mathcal{V}_ω in \mathbb{S}^{d-1} containing ω and a smooth function $e(s) = (e_{s,2}, \dots, e_{s,d}): \mathcal{V}_\omega \rightarrow (T_s\mathbb{S}^{d-1})^{d-1}$, which $e(s)$ is an orthonormal basis of the tangent space $T_s\mathbb{S}^{d-1}$ for each unit vector $s \in \mathcal{V}_\omega$, such that following asymptotics*

$$G_\lambda(x, y) = \frac{e^{(x-y)(ik_0-\beta_s)}}{(2\pi|x-y|)^{(d-1)/2}} \left(\frac{|\nabla E(\beta_s)|^{(d-3)}}{\det(-e_{s,p} \cdot \text{Hess}(E)(\beta_s)e_{s,q})_{2 \leq p, q \leq d}} \right)^{1/2} \frac{\overline{\phi_{k_0+i\beta_s}(x)\phi_{k_0-i\beta_s}(y)}}{(\phi_{k_0+i\beta_s}, \phi_{k_0-i\beta_s})_{L^2(\mathbb{T})}} + e^{(y-x)\cdot\beta_s} r(x, y), \quad (15)$$

hold for all (x, y) such that $s = (x - y)/|x - y| \in \mathcal{V}_\omega$. Furthermore, there is a positive constant $C(\omega)$ depending on ω such that $|r(x, y)| \leq C(\omega)|x - y|^{-d/2}$.

Proof of Theorem 2.11. Observe that for any orthonormal basis $\{e_{s,l}\}_{2 \leq l \leq d}$ of the tangent space $T_s \mathbb{S}^{d-1}$,

$$\det(-\mathcal{P}_s \text{Hess}(E)(\beta_s) \mathcal{P}_s) = \det(-e_{s,p} \cdot \text{Hess}(E)(\beta_s) e_{s,q})_{2 \leq p,q \leq d}.$$

Now, using of a finite cover of the unit sphere by neighborhoods \mathcal{V}_{ω_j} in Theorem 3.1, one obtains Theorem 2.11. □

Remark 3.2. (1) The asymptotics (14) (or (15)) resemble formula (1.1) in Theorem 1.1 in [25] when λ is below the bottom of the spectrum of the operator. Moreover, as in Theorem 1.1 in [26], by using the Gauss-Kronecker curvature \mathcal{K}_λ , the main result (14) could be restated as follows:

$$G_\lambda(x, y) = \frac{e^{(x-y)(ik_0-\beta_s)}}{(2\pi|x-y|)^{(d-1)/2}} \frac{1}{|\nabla E(\beta_s)| \mathcal{K}_\lambda(\beta_s)^{1/2}} \frac{\phi_{k_0+i\beta_s}(x) \overline{\phi_{k_0-i\beta_s}(y)}}{(\phi_{k_0+i\beta_s}, \phi_{k_0-i\beta_s})_{L^2(\mathbb{T})}} + e^{(y-x)\cdot\beta_s} O(|x-y|^{-d/2}).$$

(2) Although (14) is an anisotropic formula, it is not hard to obtain from (14) an isotropic upper estimate for the Green’s function G_λ based on the *distance from λ to the spectrum of the operator L* ,⁴ e.g., there are some positive constants C_1, C_2 (depending only on L and λ_j) and C_3 (which may depend on λ) such that whenever $|x-y| > C_3$, the following inequality holds:

$$|G_\lambda(x, y)| \leq C_1 |\lambda|^{(d-3)/4} \frac{e^{-C_2 |\lambda|^{1/2} |x-y|}}{|x-y|^{(d-1)/2}}.$$

(2) If the band edge occurs at finitely many points, rather than a single k_0 , one just needs to combine the asymptotics coming from all these isolated minima.

Now we outline the proof of Theorem 3.1. In Section 5, we introduce the tools of Floquet–Bloch theory to reduce the problem to that of finding the asymptotics of a scalar integral. The purpose of Section 4 is to prepare for Section 5, by shifting an integral from the fundamental domain \mathcal{O} along some purely imaginary directions in \mathbb{C}^d . This reduces finding the asymptotics of the Green’s function G_λ to an auxiliary Green’s function $G_{s,\lambda}$ via the formula (21). Next, we single out a principal term G_0 of the Green’s function $G_{s,\lambda}$ and then represent this kernel G_0 as a scalar integral in (26). We also prove that the error kernel $G_{s,\lambda} - G_0$ decays rapidly (see Theorem 5.4). Then in (29), our reduced Green’s function G_0 can be expressed in terms of the two integrals I and J . Here the integral I

⁴ Recall that the spectral edge is assumed to be zero.

is mainly responsible for the asymptotics of G_0 and the integral J decays fast enough to be included in the remainder term $r(x, y)$ in the asymptotics (14). The first part of Section 6 is devoted to achieving the asymptotics of the main integral I (see Theorem 6.2) by adapting the method similar to the one used in the discrete case [35], while the second part of Section 6 provides an estimate of J (see Proposition 6.6). In order to not overload the main text with technicalities, the proofs of some auxiliary statements are postponed till Sections 7–9.

4. On local geometry of the resolvent set

The following proposition shows that for any $s \in \mathbb{S}^{d-1}$, $k_0 + i\beta_s$ is the only complex quasimomentum having the form of $k + it\beta_s$ where $k \in \mathcal{O}$, $t \in [0, 1]$ such that λ is in the spectrum of the corresponding fiber operator $L(k + it\beta_s)$. In other words, by moving from $k \in \mathcal{O}$ in the direction $i\beta_s$, the first time we *hit the Fermi surface* $F_{L,\lambda}$ (i.e., the spectrum of $L(k)$ meets λ) is at the value of the quasimomentum $k = k_0 + i\beta_s$. This step is crucial for setting up the scalar integral in the next section, which is solely responsible for the main term asymptotics of our Green's function.

Proposition 4.1. *If $|\lambda|$ is small enough (depending on the dispersion branch λ_j and L), then $\lambda \in \rho(L(k + it\beta_s))$ if and only if $(k, t) \neq (k_0, 1)$.*

The proof of this proposition is presented in Subsection 9.3.

5. A Floquet reduction of the problem

We recall here some basic properties of the Floquet transform and then apply this transform to reduce our problem to finding asymptotics of a scalar integral expression, which is close to the one arising when dealing with the Green's function of the Laplacian at a small negative level λ . As in [23], the idea is to show that only the branch of the dispersion relation λ_j appearing in Assumption A dominates the asymptotics.

5.1. The Floquet transform. Let us consider a sufficiently fast decaying function $f(x)$ (to begin with, compactly supported function) on \mathbb{R}^d . We need the following transform that plays the role of the Fourier transform for the periodic case [19, 20, 27]. In fact, it is a version of the Fourier transform on the group \mathbb{Z}^d of periods. We use the following version, which is slightly different from the one used in [23].

Definition 5.1. The Floquet transform \mathcal{F}

$$f(x) \longrightarrow \hat{f}(k, x)$$

maps a function f on \mathbb{R}^d into a function \hat{f} defined on $\mathbb{R}^d \times \mathbb{R}^d$ in the following way:

$$\hat{f}(k, x) := \sum_{\gamma \in \mathbb{Z}^d} f(x + \gamma) e^{-ik \cdot (x + \gamma)}.$$

From the above definition, one can see that \hat{f} is \mathbb{Z}^d -periodic in the x -variable and satisfies a cyclic condition with respect to k :

$$\begin{cases} \hat{f}(k, x + \gamma) = \hat{f}(k, x) & \text{for all } \gamma \in \mathbb{Z}^d, \\ \hat{f}(k + 2\pi\gamma, x) = e^{-2\pi i \gamma \cdot x} \hat{f}(k, x) & \text{for all } \gamma \in \mathbb{Z}^d. \end{cases}$$

Thus, it suffices to consider the Floquet transform \hat{f} as a function defined on $\mathcal{O} \times \mathbb{T}$. Usually, we will regard \hat{f} as a function $\hat{f}(k, \cdot)$ in k -variable in \mathcal{O} with values in the function space $L^2(\mathbb{T})$.

For our purpose, we need to list some well-known results of the Floquet transform (see e.g., [19, 20]).

Lemma 5.2. 1. The transform \mathcal{F} is an isometry of $L^2(\mathbb{R}^d)$ onto

$$\int_{\mathcal{O}}^{\oplus} L^2(\mathbb{T}) = L^2(\mathcal{O}, L^2(\mathbb{T}))$$

and of $H^2(\mathbb{R}^d)$ into

$$\int_{\mathcal{O}}^{\oplus} H^2(\mathbb{T}) = L^2(\mathcal{O}, H^2(\mathbb{T})).$$

2. The inversion \mathcal{F}^{-1} is given by the formula

$$f(x) = (2\pi)^{-d} \int_{\mathcal{O}} e^{ik \cdot x} \hat{f}(k, x) dk, \quad x \in \mathbb{R}^d. \tag{16}$$

By using cyclic conditions of \hat{f} , we obtain an alternative inversion formula

$$f(x) = (2\pi)^{-d} \int_{\mathcal{O}} e^{ik \cdot x} \hat{f}(k, x - \gamma) dk, \quad x \in W + \gamma. \tag{17}$$

3. The action of any \mathbb{Z}^d -periodic elliptic operator L (not necessarily self-adjoint) in $L^2(\mathbb{R}^d)$ under the Floquet transform \mathcal{F} is given by

$$\mathcal{F}L(x, D)\mathcal{F}^{-1} = \int_{\mathcal{O}}^{\oplus} L(x, D + k) dk = \int_{\mathcal{O}}^{\oplus} L(k) dk,$$

where $L(k)$ is defined in (5).

Equivalently,

$$\widehat{L}f(k) = L(k)\hat{f}(k), \quad \text{for all } f \in H^2(\mathbb{R}^d).$$

4. (**A Paley–Wiener theorem for \mathcal{F}**) Let $\phi(k, x)$ be a function defined on $\mathbb{R}^d \times \mathbb{R}^d$ such that for each k , it belongs to the Sobolev space $H^s(\mathbb{T})$ for $s \in \mathbb{R}^+$ and satisfies the cyclic condition in k -variable. Then

- (1) Suppose the mapping $k \rightarrow \phi(k, \cdot)$ is a C^∞ -map from \mathbb{R}^d into the Hilbert space $H^s(\mathbb{T})$. Then $\phi(k, x)$ is the Floquet transform of a function $f \in H^s(\mathbb{R}^d)$ such that for any compact set K in \mathbb{R}^d and any $N > 0$, the norm $\|f\|_{H^s(K+\gamma)} \leq C_N |\gamma|^{-N}$. In particular, by Sobolev's embedding theorem, if $s > d/2$, then the pointwise estimation holds:

$$|f(x)| \leq C_N (1 + |x|)^{-N}, \quad \text{for all } N > 0.$$

- (2) Suppose the mapping $k \rightarrow \phi(k, \cdot)$ is an analytic map from \mathbb{R}^d into the Hilbert space $H^s(\mathbb{T})$. Then $\phi(k, x)$ is the Floquet transform of a function $f \in H^s(\mathbb{R}^d)$ such that for any compact set K in \mathbb{R}^d , one has $\|f\|_{H^s(K+\gamma)} \leq C e^{-C|\gamma|}$. In particular, by Sobolev's embedding theorem, if $s > d/2$, then the pointwise estimation holds:

$$|f(x)| \leq C e^{-C|x|}.$$

5.2. The Floquet reduction. The Green's function G_λ of L at λ is the Schwartz kernel of the resolvent operator $R_\lambda = (L - \lambda)^{-1}$. Fix a $\lambda < 0$ such that the statement of Proposition 4.1 holds. For any $s \in \mathbb{S}^{d-1}$ and $t \in [0, 1]$, we consider the following operator with real coefficients on \mathbb{R}^d :

$$L_{t,s} := e^{t\beta_s \cdot x} L e^{-t\beta_s \cdot x}. \quad (18)$$

For simplicity, we write $L_s := L_{1,s}$ and note that $L_{0,s} = L$. Due to self-adjointness of L , the adjoint of $L_{t,s}$ is

$$L_{t,s}^* = L_{-t,s}. \quad (19)$$

By definition, $L_{t,s}(k) = L(k + it\beta_s)$ for any k in \mathbb{C}^d and therefore, (6) yields

$$\sigma(L_{t,s}) = \bigcup_{k \in \mathcal{O}} \sigma(L(k + it\beta_s)) \supseteq \{\lambda_j(k + it\beta_s)\}_{k \in \mathcal{O}}. \quad (20)$$

The Schwartz kernel $G_{s,\lambda}$ of the resolvent operator $R_{s,\lambda} := (L_s - \lambda)^{-1}$ is

$$G_{s,\lambda}(x, y) = e^{\beta_s \cdot x} G_\lambda(x, y) e^{-\beta_s \cdot y} = e^{\beta_s \cdot (x-y)} G_\lambda(x, y). \quad (21)$$

Thus, instead of finding asymptotics of G_λ , we can focus on the asymptotics of $G_{s,\lambda}$.

By (20) and Proposition 4.1, λ is not in the spectrum of $L_{t,s}$ for any $s \in \mathbb{S}^{d-1}$ and $t \in [0, 1)$. Let us consider

$$R_{t,s,\lambda} f := (L_{t,s} - \lambda)^{-1} f, \quad f \in L^2_{\text{comp}}(\mathbb{R}^d),$$

where L^2_{comp} stands for compactly supported functions in L^2 .

Applying Lemma 5.2, we have

$$\widehat{R_{t,s,\lambda} f}(k) = (L_{t,s}(k) - \lambda)^{-1} \hat{f}(k), \quad (t, k) \in [0, 1) \times \mathcal{O}.$$

We consider the sesquilinear form

$$(R_{t,s,\lambda} f, \varphi) = (2\pi)^{-d} \int_{\mathcal{O}} ((L_{t,s}(k) - \lambda)^{-1} \hat{f}(k), \hat{\varphi}(k)) dk,$$

where $\varphi \in L^2_{\text{comp}}(\mathbb{R}^d)$.

In the next lemma (see Subsection 9.3), we show the weak convergence of $R_{t,s,\lambda}$ in L^2_{comp} as $t \nearrow 1$ and introduce the limit operator $R_{s,\lambda} = \lim_{t \rightarrow 1^-} R_{t,s,\lambda}$. The limit operator $R_{s,\lambda}$ is central in our study of the asymptotics of the Green's function.

Lemma 5.3. *Let $d \geq 2$. Under Assumption A, the following equality holds:*

$$\lim_{t \rightarrow 1^-} (R_{t,s,\lambda} f, \varphi) = (2\pi)^{-d} \int_{\mathcal{O}} ((L_s(k) - \lambda)^{-1} \hat{f}(k), \hat{\varphi}(k)) dk. \quad (22)$$

The integral in the right hand side of (22) is absolutely convergent for f, φ in $L^2_{\text{comp}}(\mathbb{R}^d)$. Thus, the Green's function $G_{s,\lambda}$ is the integral kernel of the operator $R_{s,\lambda}$ defined as follows

$$\widehat{R_{s,\lambda} f}(k) = (L_s(k) - \lambda)^{-1} \hat{f}(k). \quad (23)$$

5.3. Singling out the principal term in $R_{s,\lambda}$. By (23), the Green's function $G_{s,\lambda}$ is the integral kernel of the operator $R_{s,\lambda}$ with the domain $L^2_{\text{comp}}(\mathbb{R}^d)$. The inversion formula (16) gives

$$R_{s,\lambda} f(x) = (2\pi)^{-d} \int_{\mathcal{O}} e^{ik \cdot x} (L_s(k) - \lambda)^{-1} \hat{f}(k, x) dk, \quad x \in \mathbb{R}^d.$$

The purpose of this part is to single out the part of the above integral that is responsible for the leading term of the Green's function asymptotics.

To find the Schwartz kernel of $R_{s,\lambda}$, it suffices to consider functions $f \in C_c^\infty(\mathbb{R}^d)$. Our first step is to localize the integral around the point k_0 . Let us consider a connected neighborhood V of k_0 on which there exist nonzero \mathbb{Z}^d -periodic (in x) functions $\phi_z(x), z \in V$ satisfying 1) $L(z)\phi_z = \lambda_j(z)\phi_z$ and 2) each ϕ_z spans the eigenspace corresponding to the eigenvalue $\lambda_j(z)$ of the operator $L(z)$. According to P3, $\lambda_j(V) \subseteq B(0, \epsilon_0)$ and $\partial B(0, \epsilon_0) \subseteq \rho(L(z))$ when $z \in V$. For such z , let $P(z)$ be the Riesz projection of $L(z)$ that projects $L^2(\mathbb{T})$ onto the eigenspace spanned by ϕ_z , i.e.,

$$P(z) = -\frac{1}{2\pi i} \oint_{|\alpha|=\epsilon_0} (L(z) - \alpha)^{-1} d\alpha.$$

Taking the adjoint, we get

$$P(z)^* = -\frac{1}{2\pi i} \oint_{|\alpha|=\epsilon_0} (L(\bar{z}) - \alpha)^{-1} d\alpha = P(\bar{z}),$$

which is the Riesz projection from $L^2(\mathbb{T})$ onto the eigenspace spanned by $\phi_{\bar{z}}$. Recall that due to (13), by choosing $|\lambda|$ small enough, there exists $r_0 > 0$ (independent of s) such that $k \pm i\beta_s \in V$ for $k \in \bar{D}(k_0, r_0) \cap \mathbb{R}^d$. We denote $P_s(k) := P(k + i\beta_s)$ for such real k . Then $P_s(k)$ is the projector onto the eigenspace spanned by $\phi(k + i\beta_s)$ and $P_s(k)^* = P(k - i\beta_s)$. Additionally, due to P6,

$$P_s(k)g = \frac{(g, \phi(k - i\beta_s))_{L^2(\mathbb{T})}}{(\phi(k + i\beta_s), \phi(k - i\beta_s))_{L^2(\mathbb{T})}} \phi(k + i\beta_s), \quad \text{for all } g \in L^2(\mathbb{T}). \quad (24)$$

Let η be a cut-off smooth function on \mathcal{O} such that $\text{supp}(\eta) \Subset D(k_0, r_0)$ and $\eta = 1$ around k_0 .

We decompose $\hat{f} = \eta\hat{f} + (1 - \eta)\hat{f}$. When $k \neq k_0$, the operator $L_s(k) - \lambda$ is invertible by Proposition 4.1. Hence, the following function is well-defined and smooth with respect to (k, x) on $\mathbb{R}^d \times \mathbb{R}^d$:

$$\hat{u}_g(k, x) = (L_s(k) - \lambda)^{-1}(1 - \eta(k))\hat{f}(k, x).$$

Using Lemma 5.2, smoothness of \hat{u}_g implies that u_g has rapid decay in x . Now we want to solve

$$(L_s(k) - \lambda)\hat{u}(k) = \eta(k)\hat{f}(k). \quad (25)$$

Let $Q_s(k) = I - P_s(k)$ and we denote the ranges of projectors $P_s(k), Q_s(k)$ by $R(P_s(k)), R(Q_s(k))$ respectively. We are interested in decomposing the solution \hat{u} into a sum of the form $\hat{u}_1 + \hat{u}_2$ where $\hat{u}_1 = P_s(k)\hat{u}_1$ and $\hat{u}_2 = Q_s(k)\hat{u}_2$. Let $\hat{f}_1 = P_s(k)\eta(k)\hat{f}$ and $\hat{f}_2 = Q_s(k)\eta(k)\hat{f}$. Observe that since the Riesz

projection $P_s(k)$ commutes with the operator $L_s(k)$ and $R(P_s(k))$ is invariant under the action of $L_s(k)$, we have $Q_s(k)L_s(k)P_s(k) = P_s(k)L_s(k)Q_s(k) = 0$ and $Q_s(k)L_s(k)Q_s(k) = L_s(k)Q_s(k)$. Thus, the problem of solving (25) can be reduced to the following block-matrix structure form

$$\left(\begin{array}{c|c} (L_s(k) - \lambda)P_s(k) & 0 \\ \hline 0 & (L_s(k) - \lambda)Q_s(k) \end{array} \right) \begin{pmatrix} \hat{u}_1 \\ \hat{u}_2 \end{pmatrix} = \begin{pmatrix} \hat{f}_1 \\ \hat{f}_2 \end{pmatrix}.$$

When k is close to k_0 ,

$$B(0, \epsilon_0) \cap \sigma(L_s(k)|_{R(Q_s(k))}) = B(0, \epsilon_0) \cap \sigma(L(k + i\beta_s)) \setminus \{\lambda_j(k + i\beta_s)\} = \emptyset.$$

Since $\lambda = \lambda_j(k_0 + i\beta_s) \in B(0, \epsilon_0)$, λ must belong to $\rho(L_s(k)|_{R(Q_s(k))})$. Hence, the operator function $\hat{u}_2(k) = (L_s(k) - \lambda)^{-1}Q_s(k)\hat{f}_2(k)$ is well-defined and smooth in k and hence by Lemma 5.2 again, u_2 has rapid decay when $|x| \rightarrow \infty$. More precisely, we have the following claim:

Theorem 5.4. *For each $s \in \mathbb{S}^{d-1}$, let $K_s(x, y)$ be the Schwartz kernel of the operator T_s acting on $L^2(\mathbb{R}^d)$ as follows:*

$$T_s = \mathcal{F}^{-1} \left(\int_0^\oplus T_s(k) dk \right) \mathcal{F},$$

where \mathcal{F} is the Floquet transform (see Definition 5.1) and

$$T_s(k) = (1 - \eta(k))(L_s(k) - \lambda)^{-1} + \eta(k)((L_s(k) - \lambda)|_{R(Q_s(k))})^{-1}Q_s(k).$$

Then the kernel $K_s(x, y)$ is continuous away from the diagonal and furthermore, as $|x - y| \rightarrow \infty$, we have

$$\sup_{s \in \mathbb{S}^{d-1}} |K_s(x, y)| = O(|x - y|^{-N}), \quad \text{for all } N > 0.$$

The proof of this claim shall be provided in Section 7.

The u_1 term contributes the leading asymptotics for the Schwartz kernel $G_{s,\lambda}$. Therefore, we only need to solve the equation $(L_s(k) - \lambda)P_s(k)\hat{u}_1 = \hat{f}_1$ on the one-dimensional range of $P_s(k)$.

Applying (24), we can rewrite

$$\hat{f}_1(k) = \frac{\eta(k)(\hat{f}, \phi(k - i\beta_s))_{L^2(\mathbb{T})}}{(\phi(k + i\beta_s), \phi(k - i\beta_s))_{L^2(\mathbb{T})}} \phi(k + i\beta_s),$$

so that equation becomes

$$\begin{aligned} & (L_s(k) - \lambda) \frac{(\hat{u}_1, \phi(k - i\beta_s))_{L^2(\mathbb{T})}}{(\phi(k + i\beta_s), \phi(k - i\beta_s))_{L^2(\mathbb{T})}} \phi(k + i\beta_s) \\ &= \frac{\eta(k)(\hat{f}, \phi(k - i\beta_s))_{L^2(\mathbb{T})}}{(\phi(k + i\beta_s), \phi(k - i\beta_s))_{L^2(\mathbb{T})}} \phi(k + i\beta_s). \end{aligned}$$

So,

$$\begin{aligned} & \frac{(\lambda_j(k + i\beta_s) - \lambda)(\hat{u}_1, \phi(k - i\beta_s))_{L^2(\mathbb{T})}}{(\phi(k + i\beta_s), \phi(k - i\beta_s))_{L^2(\mathbb{T})}} \phi(k + i\beta_s) \\ &= \frac{\eta(k)(\hat{f}, \phi(k - i\beta_s))_{L^2(\mathbb{T})}}{(\phi(k + i\beta_s), \phi(k - i\beta_s))_{L^2(\mathbb{T})}} \phi(k + i\beta_s). \end{aligned}$$

In addition to the equation $\hat{u}_1 = P_s(k)\hat{u}_1$, \hat{u}_1 must also satisfy

$$(\lambda_j(k + i\beta_s) - \lambda)(\hat{u}_1, \phi(k - i\beta_s))_{L^2(\mathbb{T})} = \eta(k)(\hat{f}, \phi(k - i\beta_s))_{L^2(\mathbb{T})}.$$

Thus, we define

$$\hat{u}_1(k, \cdot) := \frac{\eta(k)\phi(k + i\beta_s, \cdot)(\hat{f}, \phi(k - i\beta_s))_{L^2(\mathbb{T})}}{(\phi(k + i\beta_s), \phi(k - i\beta_s))_{L^2(\mathbb{T})}(\lambda_j(k + i\beta_s) - \lambda)}.$$

By the inverse Floquet transform (16),

$$u_1(x) = (2\pi)^{-d} \int_{\mathbb{O}} e^{ik \cdot x} \frac{\eta(k)\phi(k + i\beta_s, x)(\hat{f}, \phi(k - i\beta_s))_{L^2(\mathbb{T})}}{(\phi(k + i\beta_s), \phi(k - i\beta_s))_{L^2(\mathbb{T})}(\lambda_j(k + i\beta_s) - \lambda)} dk,$$

for any $x \in \mathbb{R}^d$.

5.4. A reduced Green's function. We are now ready for setting up a *reduced Green's function* G_0 , whose asymptotic behavior reflects exactly the leading term of the asymptotics of the Green's function $G_{s,\lambda}$. We introduce $G_0(x, y)$ (roughly speaking) as the Schwartz kernel of the restriction of the operator $R_{s,\lambda}$ onto the one-dimensional range of P_s (which is the direct integral of idempotents $P_s(k)$) as follows:

$$u_1(x) = \int_{\mathbb{R}^d} G_0(x, y) f(y) dy, \quad x \in \mathbb{R}^d,$$

where f is in $L^2_{\text{comp}}(\mathbb{R}^d)$.

We recall from (9) that $F(k + i\beta_s)$ is the inner product $(\phi(k + i\beta_s), \phi(k - i\beta_s))_{L^2(\mathbb{T})}$. As in [23], we notice that

$$\begin{aligned}
 u_1(x) &= (2\pi)^{-d} \int_{\mathcal{O}} \int_{\mathbb{T}} e^{ik \cdot x} \eta(k) \hat{f}(k, y) \\
 &\quad \frac{\overline{\phi(k - i\beta_s, y)} \phi(k + i\beta_s, x)}{F(k + i\beta_s)(\lambda_j(k + i\beta_s) - \lambda)} dy dk \\
 &= (2\pi)^{-d} \int_{\mathcal{O}} \eta(k) \int_{[0,1]^d} \sum_{\gamma \in \mathbb{Z}^d} f(y - \gamma) e^{ik \cdot (x + \gamma - y)} \\
 &\quad \frac{\overline{\phi(k - i\beta_s, y)} \phi(k + i\beta_s, x)}{F(k + i\beta_s)(\lambda_j(k + i\beta_s) - \lambda)} dy dk \\
 &= (2\pi)^{-d} \int_{\mathcal{O}} \eta(k) \sum_{\gamma \in \mathbb{Z}^d} \int_{[0,1]^d - \gamma} f(y) e^{ik \cdot (x - y)} \\
 &\quad \frac{\overline{\phi(k - i\beta_s, y + \gamma)} \phi(k + i\beta_s, x)}{F(k + i\beta_s)(\lambda_j(k + i\beta_s) - \lambda)} dy dk \\
 &= (2\pi)^{-d} \int_{\mathcal{O}} \eta(k) \sum_{\gamma \in \mathbb{Z}^d} \int_{[0,1]^d - \gamma} f(y) e^{ik \cdot (x - y)} \\
 &\quad \frac{\overline{\phi(k - i\beta_s, y)} \phi(k + i\beta_s, x)}{F(k + i\beta_s)(\lambda_j(k + i\beta_s) - \lambda)} dy dk \\
 &= (2\pi)^{-d} \int_{\mathbb{R}^d} f(y) \left(\int_{\mathcal{O}} \eta(k) e^{ik \cdot (x - y)} \right. \\
 &\quad \left. \frac{\overline{\phi(k - i\beta_s, y)} \phi(k + i\beta_s, x)}{F(k + i\beta_s)(\lambda_j(k + i\beta_s) - \lambda)} dk \right) dy.
 \end{aligned}$$

Therefore, our *reduced Green's function* is

$$G_0(x, y) = (2\pi)^{-d} \int_{\mathcal{O}} \eta(k) e^{ik \cdot (x - y)} \frac{\phi(k + i\beta_s, x) \overline{\phi(k - i\beta_s, y)}}{F(k + i\beta_s)(\lambda_j(k + i\beta_s) - \lambda)} dk. \quad (26)$$

6. Asymptotics of the Green's function

Let (e_1, \dots, e_d) be the standard orthonormal basis in \mathbb{R}^d . Fixing $\omega \in \mathbb{S}^{d-1}$, we would like to show that the asymptotics (15) will hold for all (x, y) such that $x - y$ belongs to a conic neighborhood containing ω . Without loss of generality, suppose that $\omega \neq e_1$.

Now let \mathcal{R}_s be the rotation in \mathbb{R}^d such that $\mathcal{R}_s(s) = e_1$ and \mathcal{R}_s leaves the orthogonal complement of the subspace spanned by $\{s, e_1\}$ invariant. We define $e_{s,j} := \mathcal{R}_s^{-1}(e_j)$, for all $j = 2, \dots, d$. Then, $\langle s, e_{s,p} \rangle = \langle e_1, e_p \rangle = 0$ and $\langle e_{s,p}, e_{s,q} \rangle = \langle e_p, e_q \rangle = \delta_{p,q}$ for $p, q > 1$. In other words,

$$\{s, e_{s,2}, \dots, e_{s,d}\} \text{ is an orthonormal basis of } \mathbb{R}^d.$$

Then around ω , we pick a compact coordinate patch \mathcal{V}_ω , so that the $\mathbb{R}^{d(d-1)}$ -valued function $e(s) = (e_{s,l})_{2 \leq l \leq d}$ is smooth in a neighborhood of \mathcal{V}_ω .

We use the same notation for \mathcal{R}_s and its \mathbb{C} -linear extension to \mathbb{C}^d .

6.1. The asymptotics of the leading term of the Green’s function. We introduce the function $\rho(k, x, y)$ on $D(k_0, r_0) \times \mathbb{R}^d \times \mathbb{R}^d$ as follows:

$$\rho(k, x, y) = \frac{\phi(k + i\beta_s, x) \overline{\phi(k - i\beta_s, y)}}{F(k + i\beta_s)},$$

where F is defined in (9) and $D(k_0, r_0)$ is described in Subection 5.3.

Due to Proposition 9.6, the function ρ is in $C^\infty(\overline{D(k_0, r_0)} \times \mathbb{R}^d \times \mathbb{R}^d)$. For each (x, y) , the Taylor expansion around k_0 of $\rho(k)$ gives

$$\rho(k, x, y) = \rho(k_0, x, y) + \rho'(k, x, y)(k - k_0), \tag{27}$$

where $\rho' \in C^\infty(\overline{D(k_0, r_0)} \times \mathbb{R}^d \times \mathbb{R}^d, \mathbb{C}^d)$. Note that for $z \in V$, $\phi(z, x)$ is \mathbb{Z}^d -periodic in x and thus, ρ and ρ' are $\mathbb{Z}^d \times \mathbb{Z}^d$ -periodic in (x, y) . Since our integrals are taken with respect to k , it is safe to write $\rho(k_0)$ instead of $\rho(k_0, x, y)$. We often omit the variables x, y in ρ if no confusion can arise.

Let $\mu(k) := \eta(k + k_0)$ be a cut-off function supported near 0, where η is introduced in Subsection 5.3. We define

$$I := (2\pi)^{-d} \int_{\mathfrak{O}} e^{i(k-k_0) \cdot (x-y)} \frac{\mu(k - k_0)}{\lambda_j(k + i\beta_s) - \lambda} dk, \tag{28a}$$

$$J := (2\pi)^{-d} \int_{\mathfrak{O}} e^{i(k-k_0) \cdot (x-y)} \frac{\mu(k - k_0)(k - k_0)\rho'(k, x, y)}{\lambda_j(k + i\beta_s) - \lambda} dk. \tag{28b}$$

Hence, we can represent the reduced Green’s function as

$$G_0(x, y) = e^{ik_0 \cdot (x-y)} (\rho(k_0)I + J). \tag{29}$$

The rest of this subsection is devoted to computing the asymptotics of the main integral I , which gives the leading term in asymptotic expansion of the reduced Green’s function $G_0(x, y)$ as $|x - y| \rightarrow \infty$.

By making the change of variables $\xi = (\xi_1, \xi') = \mathcal{R}_s(k - k_0)$, we have

$$I = (2\pi)^{-d} \int_{\mathbb{R}^d} e^{i|x-y|\xi_1} \frac{\mu(\xi_1, \xi')}{(\lambda_j \circ \mathcal{R}_s^{-1})(\xi + \mathcal{R}_s(k_0 + i\beta_s)) - \lambda} d\xi. \tag{30}$$

We introduce the following function defined on some neighborhood of 0 in \mathbb{C}^d :

$$W_s(z) := (\lambda_j \circ \mathcal{R}_s^{-1})(-iz + \mathcal{R}_s(k_0 + i\beta_s)) - \lambda.$$

It is holomorphic near 0 (on $i\mathcal{R}_s(V)$) and $W_s(0) = 0$. Then $W_s(iz)$ is the analytic continuation to the domain $\mathcal{R}_s(V)$ of the denominator of the integrand in (30). For a complex vector $z = (z_1, \dots, z_d) \in \mathbb{C}^d$, we write $z = (z_1, z')$, where $z' = (z_2, \dots, z_d)$.

The following proposition provides a factorization of W_s that is crucial for computing the asymptotics of the integral I .

Proposition 6.1. *There exist $r > 0$ and $\epsilon > 0$ (independent of $s \in \mathcal{V}_\omega$), such that W_s has the decomposition⁵*

$$W_s(z) = (z_1 - A_s(z'))B_s(z), \quad \text{for all } z = (z_1, z') \in B(0, r) \times D'(0, \epsilon). \tag{31}$$

Here the functions A_s, B_s are holomorphic in $D'(0, \epsilon)$ and $B(0, r) \times D'(0, \epsilon)$ respectively such that $A_s(0) = 0$ and B_s is non-vanishing on $B(0, r) \times D'(0, \epsilon)$. Also, these functions and their derivatives depend continuously on s . Moreover for $z' \in D'(0, \epsilon)$,

$$A_s(z') = \frac{1}{2}z' \cdot Q_s z' + O(|z'|^3), \tag{32}$$

where $O(|z'|^3)$ is uniform in s when $z' \rightarrow 0$ and Q_s is the positive definite $(d - 1) \times (d - 1)$ matrix

$$Q_s = -\frac{1}{|\nabla E(\beta_s)|} (e_{s,p} \cdot \text{Hess}(E)(\beta_s) e_{s,q})_{2 \leq p, q \leq d}. \tag{33}$$

Proof. By Cauchy–Riemann equations for W_s and (12),

$$\frac{\partial W_s}{\partial z_1}(0) = \frac{\partial W_s}{\partial \xi_1}(0) = -i \nabla \lambda_j(k_0 + i\beta_s) \cdot \mathcal{R}_s^{-1} e_1 = -\nabla E(\beta_s) \cdot s = |\nabla E(\beta_s)| > 0. \tag{34}$$

⁵ See Notation 2.8 (a) in Section 2 for the definitions of $B(0, r)$ and $D'(0, \epsilon)$.

Thus 0 is a simple zero of W_s . Due to smoothness in s of W_s and β_s , we have

$$c := \min_{s \in \mathcal{V}_\omega} \frac{\partial W_s}{\partial z_1}(0) \geq \min_{s \in \mathbb{S}^{d-1}} |\nabla E(\beta_s)| > 0. \quad (35)$$

Applying the Weierstrass preparation theorem (see Theorem 9.2), we obtain the decomposition (31) on a neighborhood of 0.

To show that this neighborhood can be chosen such that it does not depend on s , we have to chase down how the neighborhood is constructed in the proof of Theorem 7.5.1 in [15] (only the first three lines of the proof there matter) and then show that all steps in this construction can be done independently of s .

In the first step of the construction, we need $r > 0$ such that $W_s(z_1, 0') \neq 0$ when $0 < |z_1| < 2r$. The mapping

$$(s, z) \mapsto \frac{\partial W_s}{\partial z_1}(z) = -i \nabla \lambda_j(-i \mathcal{R}_s^{-1} z + k_0 + i \beta_s) \cdot s$$

is jointly continuous on $\mathcal{V}_\omega \times \mathcal{R}_s(V)$ and the value of this mapping at $z = 0$ is greater or equal than c due to (34) and (35). Therefore, $\left| \frac{\partial W_s}{\partial z_1}(z) \right| > c/2$ in some open neighborhood $X_s \times Y_s$ of $(s, 0)$ in $\mathcal{V}_\omega \times \mathbb{C}^d$. By compactness, $\mathcal{V}_\omega \subseteq \bigcup_{k=1}^N X_{s_k}$ for a finite collection of points s_1, \dots, s_N on the sphere. Let Y be the intersection of all Y_{s_k} and let $r > 0$ be such that $D(0, 2r) \subseteq Y$. Note that r is independent of s . We claim r has the desired property. Observe that for $|z| < 2r$, we have $\left| \frac{\partial W_s}{\partial z_1}(z) \right| > \frac{c}{2}$ for any s in \mathcal{V}_ω . For a proof by contradiction, suppose that there is some z_1 such that $0 < |z_1| < 2r$ and $W_s(z_1, 0') = 0 = W_s(0, 0')$ for some s . Applying Rolle's theorem to the function $t \in [0, 1] \mapsto W_s(tz_1, 0')$ yields $\frac{\partial W_s}{\partial z_1}(tz_1, 0') = 0$ for some $t \in (0, 1)$. Consequently, $(tz_1, 0') \notin D(0, 2r)$ while $|tz_1| < |z_1| < 2r$ (contradiction!).

For the second step of the construction, we want some $\delta > 0$ (independent of s) such that $W_s(z) \neq 0$ when $|z_1| = r, |z'| < \delta$. This can be done in a similar manner. Let $S(0, r) \subset \mathbb{C}$ be the circle with radius r . Now we consider the smooth mapping $W: (s, z_1, z') \mapsto W_s(z_1, z')$ where $z_1 \in S(0, r)$. Its value at each point $(s, z_1, 0')$ is equal to $W_s(z_1, 0')$, which is non-zero due to the choice of r in the first step of the construction. Thus, it is also non-zero in some open neighborhood $\tilde{X}_{s, z_1} \times \tilde{Y}_{s, z_1} \times \tilde{Z}_{s, z_1}$ of $(s, z_1, 0')$ in $\mathcal{V}_\omega \times S(0, r) \times \mathbb{C}^{d-1}$. We select points $s_1, \dots, s_M \in \mathcal{V}_\omega$ and $\gamma_1, \dots, \gamma_M \in S(0, r)$ so that the union of all $\tilde{X}_{s_k, \gamma_k} \times \tilde{Y}_{s_k, \gamma_k}, 1 \leq k \leq M$ covers the compact set $\mathcal{V}_\omega \times S(0, r)$. Next we choose $\delta > 0$ so that $D'(0, \delta)$ is contained in the intersection of these \tilde{Z}_{s_k, z_k} . Note that δ is independent of s and also z_1 . Of course $W_s(z_1, z') \neq 0$ for all s and $z \in \{|z_1| = r, |z'| < \delta\}$. According to [15], the decomposition (31) holds in the polydisc $\{|z_1| < r, |z'| < \delta\}$.

Also, from the proof of Theorem 7.5.1 in [15], the function A_s is defined via the following formula

$$z_1 - A_s(z') = \exp\left(\frac{1}{2\pi i} \int_{|\omega|=r} \left(\frac{\partial W_s(\omega, z')}{\partial \omega} / W_s(\omega, z')\right) \log(z_1 - \omega) d\omega\right). \quad (36)$$

The mappings $(s, z') \mapsto A_s(z')$ and $(s, z) \mapsto B_s(z)$ are jointly continuous due to (31) and (36). There exists $0 < \epsilon \leq \delta$ such that $\max_{s \in \mathcal{V}_\omega} |A_s(z')| < r$ whenever $|z'| < \epsilon$. We have the identity (31) on $B(0, r) \times D'(0, \epsilon)$. Now, we show that this is indeed the neighborhood that has the desired properties. Since $|z'| < \epsilon$ implies that the points $z = (A_s(z'), z') \in B(0, r) \times D'(0, \epsilon)$, we can evaluate (31) at these points to obtain

$$W_s(A_s(z'), z') = 0, \quad z' \in D'(0, \epsilon). \quad (37)$$

By differentiating (37), we have

$$\frac{\partial W_s}{\partial z_p}(A_s(z'), z') + \frac{\partial W_s}{\partial z_1}(A_s(z'), z') \frac{\partial A_s}{\partial z_p}(z') = 0, \quad \text{for } p = 2, \dots, d. \quad (38)$$

Observe that from the above construction, the term $\frac{\partial W_s}{\partial z_1}(A_s(z'), z')$ is always non-zero whenever $|z'| < \epsilon$. Consequently, all first-order derivatives of A_s are jointly continuous in (s, z) . Similarly, we deduce by induction on $n \in \mathbb{N}^d$ that all derivatives of the function A_s depend continuously on s since after taking differentiation of the equation (37) up to order n , the n -order derivative term always goes with the nonzero term $\frac{\partial W_s}{\partial z_1}(A_s(z'), z')$ and the remaining terms in the sum are just lower order derivatives. Hence the same conclusion holds for all derivatives of B_s by differentiating (31).

In particular, set $z' = 0$ in (38) to obtain

$$\frac{\partial W_s}{\partial z_p}(0) + \frac{\partial W_s}{\partial z_1}(0) \frac{\partial A_s}{\partial z_p}(0) = 0, \quad \text{for } p = 2, \dots, d. \quad (39)$$

Note that for $p > 1$,

$$\frac{\partial W_s}{\partial z_p}(z) = -i \nabla \lambda_j (-i \mathcal{R}_s^{-1} z + k_0 + i \beta_s) \cdot \mathcal{R}_s^{-1} e_p. \quad (40)$$

By substituting $z = 0$,

$$\begin{aligned} \frac{\partial W_s}{\partial z_p}(0) &= -i \nabla \lambda_j(k_0 + i\beta_s) \cdot \mathcal{R}_s^{-1} e_p \\ &= -\nabla E(\beta_s) \cdot e_{s,p} \\ &= -|\nabla E(\beta_s)| s \cdot e_{s,p} \\ &= 0. \end{aligned} \quad (41)$$

Equations (34), (39), and (41) imply

$$\frac{\partial A_s}{\partial z_p}(0) = 0, \quad \text{for } p = 2, \dots, d. \quad (42)$$

Taking a partial derivative with respect to z_q ($q > 1$) of (40) at $z = 0$, we see that

$$\begin{aligned} \frac{\partial^2 W_s}{\partial z_p \partial z_q}(0) &= \sum_{m=1}^d -\nabla \left(\frac{\partial \lambda_j}{\partial z_m}(k_0 + i\beta_s) \right) \cdot \mathcal{R}_s^{-1} e_p (\mathcal{R}_s^{-1} e_q)_m \\ &= - \sum_{m,n=1}^d \frac{\partial^2 \lambda_j}{\partial z_m \partial z_n}(k_0 + i\beta_s) (e_{s,p})_m (e_{s,q})_n \\ &= e_{s,q} \cdot \text{Hess}(E)(\beta_s) e_{s,p}. \end{aligned} \quad (43)$$

A second differentiation of (38) at $z = (A_s(z'), z')$ gives

$$\begin{aligned} 0 &= \left(\frac{\partial^2 W_s}{\partial z_p \partial z_q}(z) + \frac{\partial W_s}{\partial z_1}(z) \frac{\partial^2 A_s}{\partial z_p \partial z_q}(z') \right) \\ &\quad + \left(\frac{\partial^2 W_s}{\partial z_1 \partial z_q}(z) \frac{\partial A_s}{\partial z_p}(z') + \frac{\partial^2 W_s}{\partial z_p \partial z_1}(z) \frac{\partial A_s}{\partial z_q}(z') + \frac{\partial^2 W_s}{\partial z_1^2}(z) \frac{\partial A_s}{\partial z_p}(z') \frac{\partial A_s}{\partial z_q}(z') \right). \end{aligned} \quad (44)$$

At $z = 0$, the sum in the second bracket of (44) is zero due to (42). Thus,

$$\frac{\partial^2 A_s}{\partial z_p \partial z_q}(0) = - \left(\frac{\partial W_s}{\partial z_1}(0) \right)^{-1} \frac{\partial^2 W_s}{\partial z_p \partial z_q}(0) \quad (2 \leq p, q \leq d). \quad (45)$$

Together with (34) and (43), the above equality becomes

$$\frac{\partial^2 A_s}{\partial z_p \partial z_q}(0) = - \frac{1}{|\nabla E(\beta_s)|} (e_{s,p} \cdot \text{Hess}(E)(\beta_s) e_{s,q})_{2 \leq p, q \leq d} = Q_s. \quad (46)$$

Consequently, by (42) and (46), the Taylor expansion of A_s at 0 implies (32).

Finally, the remainder term $O(|z'|^3)$ in the Taylor expansion (32), denoted by $R_{s,3}(z')$, can be estimated as follows:

$$\begin{aligned} |R_{s,3}(z')| &\lesssim |z'|^3 \max_{|\alpha|=3, 0 \leq t \leq 1} \left| \frac{\partial^\alpha A_s}{\partial z^\alpha}(tz') \right| \\ &\lesssim |z'|^3 \max_{|\alpha|=3, |y| \leq |z'|} \left| \frac{\partial^\alpha A_s}{\partial z^\alpha}(y) \right|. \end{aligned}$$

Due to the continuity of third-order derivatives of A_s on $\mathcal{V}_\omega \times D'(0, \epsilon)$,

$$\lim_{|z'| \rightarrow 0} \max_{s \in \mathcal{V}_\omega} \frac{|R_{s,3}(z')|}{|z'|^3} < \infty. \tag{47}$$

This proves the last claim of this proposition. □

We can now let the size of the support of η ($\Subset \mathcal{O}$) be small enough such that the decomposition (31) in Proposition 6.1 holds on the support of μ , i.e., $\text{supp}(\mu) \Subset B(0, r) \times D'(0, \epsilon)$. Therefore, from (30), we can represent the integral I as follows:

$$\begin{aligned} I &= (2\pi)^{-d} \int_{\mathbb{R}^d} e^{i|x-y|\xi_1} \frac{\mu(\xi_1, \xi')}{W_s(i\xi)} d\xi_1 d\xi' \\ &= (2\pi)^{-d} \int_{|\xi'| < \epsilon} \int_{\mathbb{R}} \frac{e^{i|x-y|\xi_1} \tilde{\mu}_s(\xi_1, \xi')}{i\xi_1 - A_s(i\xi')} d\xi_1 d\xi', \end{aligned} \tag{48}$$

where $\tilde{\mu}_s(\xi) = \mu(\xi)(B_s(i\xi))^{-1}$. We extend $\tilde{\mu}_s$ to a smooth compactly supported function on \mathbb{R}^d by setting $\tilde{\mu}_s = 0$ outside its support. Since all derivatives of $\tilde{\mu}_s$ depend continuously on s , they are uniformly bounded in s . Let $\nu_s(t, \xi')$ be the Fourier transform in the variable ξ_1 of the function $\tilde{\mu}_s(-\xi_1, \xi')$ for each $\xi' \in \mathbb{R}^{d-1}$, i.e.,

$$\nu_s(t, \xi') = \int_{-\infty}^{+\infty} e^{it\xi_1} \tilde{\mu}_s(\xi_1, \xi') d\xi_1.$$

By applying the Lebesgue Dominated Convergence Theorem, the function ν_s is continuous in (s, t, ξ') on $\mathcal{V}_\omega \times \mathbb{R}^d$. For such ξ' , $\nu_s(\cdot, \xi')$ is a Schwartz function in t on \mathbb{R} . Due to Lemma 9.1, for any $N > 0$, $\nu_s(t, \xi') = O(|t|^{-N})$ uniformly in s and ξ' as $t \rightarrow \infty$. We also choose ϵ small enough such that whenever $|\xi'| < \epsilon$, the absolute value of the remainder term $O(|\xi'|^3)$ in (32) is bounded from above by $\frac{1}{4}\xi' \cdot Q_s \xi'$. Note that ϵ is still independent of s , because the term $O(|\xi'|^3)/|\xi'|^3$ is uniformly bounded by the quantity in (47). Meanwhile, each positive definite matrix Q_s dominates the positive matrix $\gamma_\omega I_{(d-1) \times (d-1)}$, where $\gamma_\omega > 0$ is the

smallest among all the eigenvalues of all matrices Q_s ($s \in \mathcal{V}_\omega$). This implies that if $0 < |\xi'| < \epsilon$, then

$$\begin{aligned} \Re(i\xi_1 - A_s(i\xi')) &= -\Re(A_s(i\xi')) \\ &= -\Re\left(-\frac{1}{2}\xi' \cdot Q_s \xi' + O(|\xi'|^3)\right) \\ &= \frac{1}{2}\xi' \cdot Q_s \xi' - \Re(O(|\xi'|^3)) \\ &> \frac{1}{4}\gamma_\omega |\xi'|^2 \\ &> 0. \end{aligned}$$

We thus can obtain the following integral representation for a factor in the integrand of I (see (48)):

$$\frac{1}{i\xi_1 - A_s(i\xi')} = \int_{-\infty}^0 e^{(i\xi_1 - A_s(i\xi'))w} dw, \quad (\xi_1, \xi') \in \mathbb{R} \times (D'(0, \epsilon) \setminus \{0\}). \quad (49)$$

Therefore,

$$\begin{aligned} I &= \frac{1}{(2\pi)^d} \int_{|\xi'| < \epsilon} \int_{-\infty}^0 e^{-wA_s(i\xi')} \int_{-r}^r e^{i(w+|x-y|\xi_1)\xi_1} \tilde{\mu}_s(\xi_1, \xi') d\xi_1 dw d\xi' \\ &= \frac{1}{(2\pi)^d} \int_{|\xi'| < \epsilon} \int_{-\infty}^{|x-y|} e^{(-t+|x-y|)A_s(i\xi')} \nu_s(t, \xi') dt d\xi'. \end{aligned} \quad (50)$$

Now our remaining task is to prove the following asymptotics of the integral I :

Theorem 6.2. *We have*

$$I = \frac{|\nabla E(\beta_s)|^{(d-3)/2} |x-y|^{-(d-1)/2}}{(2\pi)^{(d-1)/2} \det(-e_{s,p} \cdot \text{Hess}(E)(\beta_s)e_{s,q})_{2 \leq p,q \leq d}^{1/2}} + O(|x-y|^{-d/2}). \quad (51)$$

Here the term $O(|x-y|^{-d/2})$ is uniform in $s \in \mathcal{V}_\omega$ as $|x-y| \rightarrow \infty$.

The next lemma reduces the leading term of the right hand side of (51) to a scalar integral as follows.

Lemma 6.3. *We have*

$$\begin{aligned} &\int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}} \exp\left(-\frac{1}{2}x' \cdot Q_s x'\right) \nu_s(t, 0) dt dx' \\ &= \frac{(2\pi)^{(d+1)/2} |\nabla E(\beta_s)|^{(d-3)/2}}{\det(-e_{s,p} \cdot \text{Hess}(E)(\beta_s)e_{s,q})_{2 \leq p,q \leq d}^{1/2}}. \end{aligned}$$

Proof. By applying the Fourier inversion formula to v_s , we get

$$\frac{1}{2\pi} \int_{\mathbb{R}} v_s(t, 0) dt = \tilde{\mu}_s(0) = (B_s(0))^{-1} = \left(\frac{\partial W_s}{\partial z_1}(0)\right)^{-1} = \frac{1}{|\nabla E(\beta_s)|}. \tag{52}$$

Here (34) is used in the last equality. Thus,

$$\begin{aligned} & \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}} \exp\left(-\frac{1}{2}x' \cdot Q_s x'\right) v_s(t, 0) dt dx' \\ &= \frac{1}{(\det Q_s)^{1/2}} \int_{\mathbb{R}} \int_{\mathbb{R}^{d-1}} \exp\left(-\frac{1}{2}|u'|^2\right) v_s(t, 0) du' dt \\ &= \frac{(2\pi)^{(d-1)/2}}{(\det Q_s)^{1/2}} \int_{\mathbb{R}} v_s(t, 0) dt \\ &= \frac{(2\pi)^{(d+1)/2}}{(\det Q_s)^{1/2} |\nabla E(\beta_s)|} \\ &= \frac{(2\pi)^{(d+1)/2} |\nabla E(\beta_s)|^{(d-3)/2}}{\det(-e_{s,p} \cdot \text{Hess}(E)(\beta_s) e_{s,q})_{2 \leq p, q \leq d}^{1/2}}. \end{aligned}$$

Note that we use the change of variables $u' := Q_s^{1/2} x'$, (52), and (33) in the first, the third, and the last equality respectively. □

For clarity, we introduce the notation $x_0 := |x - y|$. The purpose of the following two lemmas is to truncate some unnecessary (rapidly decreasing) parts of the main integrals we are interested in.

Lemma 6.4. *i) For any $\alpha \in (0, 1)$ and $n > 0$, one has*

$$\sup_{s \in \mathcal{V}_\omega} \int_{|\xi'| < \epsilon} \int_{(-\infty, -x_0^\alpha) \cup (x_0^\alpha, x_0)} \exp((x_0 - t)A_s(i\xi')) v_s(t, \xi') dt d\xi' = O(x_0^{-n})$$

and

$$\sup_{s \in \mathcal{V}_\omega} \int_{\mathbb{R}^{d-1}} \int_{|t| > x_0^\alpha} \exp\left(-\frac{1}{2}x' \cdot Q_s x'\right) v_s(t, 0) dt dx' = O(x_0^{-n}).$$

ii) *For any $\beta < 1/2$, $n > 0$ and each fixed $t \in [-x_0/2, x_0/2]$, one obtains*

$$\sup_{s \in \mathcal{V}_\omega} \int_{\epsilon \sqrt{x_0 - t} > |x'| \geq x_0^\beta} \left| \exp\left((x_0 - t)A_s\left(\frac{ix'}{\sqrt{x_0 - t}}\right)\right) \right| dx' = O(x_0^{-n})$$

and

$$\sup_{s \in \mathcal{V}_\omega} \int_{|x'| \geq x_0^\beta} \exp\left(-\frac{1}{2}x' \cdot Q_s x'\right) dx' = O(x_0^{-n}).$$

Proof. i) We recall that $\sup_{s, \xi'} |v_s(t, \xi')| = O(|t|^{-n})$ for any $n > 0$. Observe that when $t \leq x_0$, $|e^{(x_0-t)A_s(i\xi')}| \leq 1$. Thus, we have

$$\begin{aligned} & \sup_{s \in \mathcal{V}_\omega} \left| \int_{|\xi'| < \epsilon} \int_{(-\infty, -x_0^\alpha) \cup (x_0^\alpha, x_0)} \exp((x_0 - t)A_s(i\xi')) v_s(t, \xi') dt d\xi' \right| \\ & \lesssim \int_{|\xi'| < \epsilon} \int_{(-\infty, -x_0^\alpha) \cup (x_0^\alpha, x_0)} |t|^{-n/\alpha-1} dt d\xi' \\ & \lesssim \int_{|t| > x_0^\alpha} |t|^{-n/\alpha-1} dt \\ & = O(x_0^{-n}). \end{aligned} \tag{53}$$

Since $|\exp(-\frac{1}{2}x' \cdot Q_s x')| \leq 1$, the second integral in this part also decays rapidly by the same argument.

ii) When $t < x_0$, we can substitute $\xi' = x'(x_0 - t)^{-1/2}$ into (32) to obtain

$$(x_0 - t) \cdot A_s\left(\frac{ix'}{\sqrt{x_0 - t}}\right) = -\frac{1}{2}x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0 - t}}\right). \tag{54}$$

Due to our choice of ϵ and the definition of γ_ω , we get the following estimate when $|x'| < \epsilon\sqrt{x_0 - t}$:

$$\begin{aligned} & \sup_{s \in \mathcal{V}_\omega} \left| \exp\left((x_0 - t)A_s\left(\frac{ix'}{\sqrt{x_0 - t}}\right)\right) \right| \\ & = \sup_{s \in \mathcal{V}_\omega} \exp\left(-\frac{1}{2}x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0 - t}}\right)\right) \\ & \leq \exp\left(-\frac{1}{4}\gamma_\omega |x'|^2\right). \end{aligned} \tag{55}$$

Hence, the two integrals in the statement can be estimated from above by

$$\begin{aligned} \int_{|x'| \geq x_0^\beta} \exp\left(-\frac{1}{4}\gamma_\omega |x'|^2\right) dx' & \lesssim \int_{x_0^\beta}^\infty \exp\left(-\frac{1}{4}\gamma_\omega r^2\right) r^{d-2} dr \\ & \lesssim \int_{x_0^\beta}^\infty r^{-n/\beta-1-(d-2)} r^{d-2} dr \\ & = O(x_0^{-n}). \end{aligned} \tag{56}$$

□

Lemma 6.5. *If $\alpha \in (0, 1)$, we have*

$$\begin{aligned} & \sup_{s \in \mathcal{V}_\omega} \int_{|t| \leq x_0^\alpha} \int_{|x'| < \epsilon \sqrt{x_0 - t}} \left(\left(1 - \frac{t}{x_0}\right)^{-(d-1)/2} - 1 \right) \\ & \quad \exp\left((x_0 - t)A_s\left(\frac{ix'}{\sqrt{x_0 - t}}\right)\right) \\ & \quad v_s\left(t, \frac{x'}{\sqrt{x_0 - t}}\right) dx' dt \\ & = O(x_0^{2\alpha-1}). \end{aligned}$$

Proof. As we argued in the proof of Lemma 6.4 (ii), this integral is majorized by

$$\begin{aligned} & \int_{-x_0^\alpha}^{x_0^\alpha} \int_{|x'| < \epsilon \sqrt{x_0 - t}} \exp\left(-\frac{1}{4}x' \cdot Q_s x'\right) \cdot \left|v_s\left(t, \frac{x'}{\sqrt{x_0 - t}}\right)\right| \\ & \quad \cdot \left|\left(1 - \frac{t}{x_0}\right)^{-(d-1)/2} - 1\right| dx' dt. \end{aligned}$$

Since v_s is uniformly bounded on $\mathbb{R} \times D'(0, \epsilon)$, it suffices to estimate the factor $\left(1 - \frac{t}{x_0}\right)^{-(d-1)/2} - 1$. But this is straightforward, since

$$\begin{aligned} \int_{-x_0^\alpha}^{x_0^\alpha} \left|\left(1 - \frac{t}{x_0}\right)^{-(d-1)/2} - 1\right| dt & \leq 2x_0^\alpha \left(\left(1 - x_0^{\alpha-1}\right)^{-(d-1)/2} - 1\right) \\ & = O(x_0^{2\alpha-1}). \end{aligned} \quad \square$$

Proof of Theorem 6.2. Thanks to Lemma 6.3, it is enough to prove the relation

$$I = (2\pi)^{-d} x_0^{-(d-1)/2} \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}} \exp\left(-\frac{1}{2}x' \cdot Q_s x'\right) v_s(t, 0) dt dx' + O(x_0^{-d/2}).$$

Due to Lemma 6.4 (i) with $\alpha = 1/4$, we only need to show that

$$\tilde{I} = x_0^{-(d-1)/2} \int_{\mathbb{R}^{d-1}} \int_{|t| \leq x_0^{1/4}} \exp\left(-\frac{1}{2}x' \cdot Q_s x'\right) v_s(t, 0) dt dx' + O(x_0^{-d/2})$$

where

$$\tilde{I} = \int_{|\xi'| < \epsilon} \int_{|t| \leq x_0^{1/4}} \exp((x_0 - t)A_s(ix')) v_s(t, \xi') dt d\xi'.$$

Then we substitute $x' = \xi' \sqrt{x_0 - t}$ to the integral \tilde{I} to get

$$\begin{aligned} \tilde{I} &= x_0^{-(d-1)/2} \int_{|t| \leq x_0^{1/4}} \int_{|x'| < \epsilon \sqrt{x_0 - t}} \left(1 - \frac{t}{x_0}\right)^{-(d-1)/2} \\ &\quad \exp\left((x_0 - t)A_s\left(\frac{ix'}{\sqrt{x_0 - t}}\right)\right) v_s\left(t, \frac{x'}{\sqrt{x_0 - t}}\right) dx' dt. \end{aligned}$$

By Lemma 6.5 with $\alpha = 1/4$, we have

$$\begin{aligned} &\int_{|t| \leq x_0^{1/4}} \int_{|x'| < \epsilon \sqrt{x_0 - t}} \exp\left((x_0 - t)A_s\left(\frac{ix'}{\sqrt{x_0 - t}}\right)\right) v_s\left(t, \frac{x'}{\sqrt{x_0 - t}}\right) dx' dt \\ &= x_0^{(d-1)/2} \tilde{I} + O(x_0^{-1/2}). \end{aligned} \quad (56)$$

Next, it is clear that for $|t| \leq x_0^{1/4}$, one has

$$\begin{aligned} \left|v_s\left(t, \frac{x'}{\sqrt{x_0 - t}}\right) - v_s(t, 0)\right| &\leq \frac{|x'|}{\sqrt{x_0 - t}} \sup_{s, \xi'} |\nabla_{\xi'} v_s(t, \xi')| \\ &\lesssim \frac{|x'|}{\sqrt{x_0}} \sup_{s, \xi'} |\nabla_{\xi'} v_s(t, \xi')|. \end{aligned}$$

Also, from the definition of the function v_s , it follows that

$$\sup_{s, \xi'} |\nabla_{\xi'} v_s(t, \xi')| = O(|t|)^{-n} \quad \text{for any } n > 0.$$

Consequently,

$$\begin{aligned} &\int_{|t| \leq x_0^{1/4}} \int_{|x'| < \epsilon \sqrt{x_0 - t}} \left| \exp\left((x_0 - t)A_s\left(\frac{ix'}{\sqrt{x_0 - t}}\right)\right) \right. \\ &\quad \left. \left(v_s\left(t, \frac{x'}{\sqrt{x_0 - t}}\right) - v_s(t, 0)\right) \right| dx' dt \\ &\lesssim \frac{1}{\sqrt{x_0}} \int_{\mathbb{R}^{d-1}} \exp\left(-\frac{1}{4}\gamma_\omega |x'|^2\right) |x'| dx' \cdot \int_{\mathbb{R}} \sup_{s, \xi'} |\nabla_{\xi'} v_s(t, \xi')| dt \\ &= O(x_0^{-1/2}). \end{aligned} \quad (57)$$

Using (56), (57), and Lemma 6.4 (i), it remains to derive the relation

$$\begin{aligned} &\int_{|t| \leq x_0^{1/4}} \int_{|x'| < \epsilon \sqrt{x_0 - t}} \exp\left((x_0 - t)A_s\left(\frac{ix'}{\sqrt{x_0 - t}}\right)\right) v_s(t, 0) dx' dt \\ &= \int_{|t| \leq x_0^{1/4}} \int_{\mathbb{R}^{d-1}} \exp\left(-\frac{1}{2}x' \cdot Q_s x'\right) v_s(t, 0) dx' dt + O(x_0^{-1/2}). \end{aligned} \quad (58)$$

Due to Lemma 6.4 (ii) with $\beta = 1/6$, we obtain

$$\begin{aligned} \sup_{s \in \mathcal{V}_\omega} \int_{\epsilon \sqrt{x_0-t} > |x'| \geq x_0^{1/6}} \left| \exp \left((x_0 - t) A_s \left(\frac{ix'}{\sqrt{x_0 - t}} \right) \right) \right| dx' &= O(x_0^{-n}), \\ \sup_{s \in \mathcal{V}_\omega} \int_{|x'| \geq x_0^{1/6}} \exp \left(-\frac{1}{2} x' \cdot Q_s x' \right) dx' &= O(x_0^{-n}). \end{aligned}$$

On the other hand,

$$\begin{aligned} &\sup_{s \in \mathcal{V}_\omega} \int_{|x'| < x_0^{1/6}} \left| \exp \left((x_0 - t) A_s \left(\frac{ix'}{\sqrt{x_0 - t}} \right) \right) - \exp \left(-\frac{1}{2} x' \cdot Q_s x' \right) \right| dx' \\ &= \sup_{s \in \mathcal{V}_\omega} \int_{|x'| < x_0^{1/6}} \exp \left(-\frac{1}{2} x' \cdot Q_s x' \right) \left| \exp \left(O \left(\frac{|x'|^3}{\sqrt{x_0}} \right) \right) - 1 \right| dx' \\ &\lesssim \int_{|x'| < x_0^{1/6}} \exp \left(-\frac{1}{2} \gamma_\omega |x'|^2 \right) \frac{|x'|^3}{\sqrt{x_0}} dx' \\ &= O(x_0^{-1/2}). \end{aligned}$$

Hence, we deduce

$$\begin{aligned} &\int_{|x'| < \epsilon \sqrt{x_0-t}} \exp \left((x_0 - t) A_s \left(\frac{ix'}{\sqrt{x_0 - t}} \right) \right) dx' - \int_{\mathbb{R}^{d-1}} \exp \left(-\frac{1}{2} x' \cdot Q_s x' \right) dx' \\ &= O(x_0^{-1/2}) \end{aligned}$$

for each $t \in [-x_0^{1/4}, x_0^{1/4}]$. Finally, we multiply the above relation with $v_s(t, 0)$ and then integrate over the interval $[-x_0^{1/4}, x_0^{1/4}]$. Since $\sup_s |v_s(t, 0)|$ is integrable over \mathbb{R} , the right hand side is still $O(x_0^{-1/2})$. Thus, we derive (58) as we wish. \square

6.2. Estimates of the integral J . In this part, we want to show that the expression J decays as $O(|x - y|^{-d/2})$. Thus, taking into account (51), we conclude that J does not contribute to the leading term of the reduced Green’s function.

In (27), we set the coordinate functions of ρ' as (ρ_1, \dots, ρ_d) . Let us introduce the smooth function $\mu^{(l)}(k, x, y) = \rho_l(k + k_0, x, y) \mu(k)$ for any $k \in \mathbb{R}^d$. The support of $\mu^{(l)}$ (as a function of k for each pair (x, y)) is contained in the support of μ and $\mu^{(l)}(k, \cdot, \cdot)$ is $\mathbb{Z}^d \times \mathbb{Z}^d$ -periodic. We denote the components of a vector k in \mathbb{R}^d as (k_1, \dots, k_d) . Observe that J is the sum of integrals J_l ($1 \leq l \leq d$) if we define

$$J_l := (2\pi)^{-d} \int_{\mathcal{O}} e^{i(k-k_0) \cdot (x-y)} \frac{\mu^{(l)}(k - k_0, x, y)(k - k_0)_l}{\lambda_j(k + i\beta_s) - \lambda} dk. \quad (59)$$

Proposition 6.6. *As $|x - y| \rightarrow \infty$, we have $J_1 = O(|x - y|^{-(d+1)/2})$ and $J_l = O(|x - y|^{-d/2})$ if $l > 1$. In particular, $J = O(|x - y|^{-d/2})$.*

Proof. Indeed, to treat these integrals, we need to re-examine the calculation in the previous subsection done for the integral I . After applying the orthogonal transformation \mathcal{R}_s on each integral J_l , we rewrite them under the form of (48) as

$$J_l = (2\pi)^{-d} \int_{|\xi'| < \epsilon} \int_{\mathbb{R}} e^{i|x-y|\xi_1} \frac{\tilde{\mu}_s^{(l)}(\xi_1, \xi', x, y)\xi_l}{i\xi_1 - A_s(i\xi')} d\xi_1 d\xi', \tag{60}$$

where $\tilde{\mu}_s^{(l)}(\xi, x, y)$ is $\mu^{(l)}(\xi, x, y)(B_s(i\xi))^{-1}$ on the support of $\mu^{(l)}$ and vanishes elsewhere. Let $v_s^{(l)}(t, \xi', x, y)$ be the Fourier transform in ξ_1 of $\tilde{\mu}_s^{(l)}(\xi_1, \xi', x, y)$. If the parameter s is viewed as another argument of our functions here, then $v_s^{(l)}(\cdot, \xi', x, y)$ is a Schwartz function for each quadruple (s, ξ', x, y) . It is elementary to check that the Fourier transform $v_s^{(l)}(t, \xi', x, y)$ is jointly continuous on $\mathcal{V}_\omega \times \mathbb{R} \times \mathbb{R}^{d-1} \times \mathbb{R}^d \times \mathbb{R}^d$ due to the corresponding property of $\tilde{\mu}_s^{(l)}(\xi, x, y)$. Periodicity in (x, y) of $v_s^{(l)}$ and Lemma 9.1 imply the following decay:

$$\lim_{t \rightarrow \infty} |t|^N \sup_{(s, \xi', x, y) \in \mathcal{V}_\omega \times \overline{D'(0, \epsilon)} \times \mathbb{R}^d \times \mathbb{R}^d} |v_s^{(l)}(t, \xi', x, y)| = 0, \quad N \geq 0. \tag{61}$$

In particular,

$$\max_{1 \leq l \leq d} \sup_{(s, t, \xi', x, y) \in \mathcal{V}_\omega \times \mathbb{R} \times \overline{D'(0, \epsilon)} \times \mathbb{R}^d \times \mathbb{R}^d} |v_s^{(l)}(t, \xi', x, y)| < \infty \tag{62}$$

and

$$S := \max_{1 \leq l \leq d} \int_{\mathbb{R}} \sup_{(s, \xi', x, y) \in \mathcal{V}_\omega \times \overline{D'(0, \epsilon)} \times \mathbb{R}^d \times \mathbb{R}^d} |v_s^{(l)}(t, \xi', x, y)| dt < \infty. \tag{63}$$

Recall that when $0 < |\xi'| < \epsilon$, $\Re(A_s(i\xi')) < 0$ and thus from (62),

$$\lim_{t \rightarrow -\infty} e^{(-t+|x-y|)A_s(i\xi')} v_s^{(1)}(t, \xi', x, y) = 0. \tag{64}$$

Case 1: $l = 1$. Using (49), (64), and integration by parts, we obtain

$$\begin{aligned} J_1 &= \frac{1}{(2\pi)^d} \int_{|\xi'| < \epsilon} \int_{-\infty}^0 e^{-wA_s(i\xi')} \int_{-r}^r \xi_1 e^{i(w+|x-y|\xi_1)} \tilde{\mu}_s^{(1)}(\xi_1, \xi', x, y) d\xi_1 dw d\xi \\ &= -\frac{i}{(2\pi)^d} \int_{|\xi'| < \epsilon} \int_{-\infty}^{|x-y|} e^{(-t+|x-y|)A_s(i\xi')} \frac{d}{dt} v_s^{(1)}(t, \xi', x, y) dt d\xi' \\ &= -\frac{i}{(2\pi)^d} \int_{|\xi'| < \epsilon} \left(v_s^{(1)}(|x-y|, \xi', x, y) \right. \\ &\quad \left. + \int_{-\infty}^{|x-y|} A_s(i\xi') e^{(-t+|x-y|)A_s(i\xi')} v_s^{(1)}(t, \xi', x, y) dt \right) d\xi'. \end{aligned} \tag{65}$$

Recall the notation $x_0 = |x - y|$. The term

$$\int_{|\xi'| < \epsilon} v_s^{(1)}(x_0, \xi', x, y) d\xi'$$

decays rapidly in x_0 , due to (61). We decompose the other term

$$\int_{-\infty}^{x_0} A_s(i\xi') e^{(x_0-t)A_s(i\xi')} v_s^{(1)}(t, \xi', x, y) dt$$

into two parts, where the first integral is taking over $(x_0/2, x_0]$ and the second one over $(-\infty, x_0/2]$. The first part decays rapidly, as in Lemma 6.4 (i). Now we need to prove that the second part decays as $O(x_0^{(d+1)/2})$. To do this, we use the change of variables $x' = \xi' \sqrt{x_0 - t}$ to rewrite the remaining integral as

$$\begin{aligned} & x_0^{(d+1)/2} \int_{|\xi'| < \epsilon} \int_{-\infty}^{x_0/2} A_s(i\xi') e^{(-t+x_0)A_s(i\xi')} v_s^{(1)}(t, \xi', x, y) dt d\xi' \\ &= \int_{-\infty}^{x_0/2} \left(1 - \frac{t}{x_0}\right)^{-(d+1)/2} \int_{|x'| < \epsilon \sqrt{x_0-t}} \left(-\frac{1}{2}x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0-t}}\right)\right) \\ & \quad \exp\left(-\frac{1}{2}x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0-t}}\right)\right) v_s^{(1)}\left(t, \left(\frac{x'}{\sqrt{x_0-t}}\right), x, y\right) dx' dt. \end{aligned} \tag{66}$$

From (63), we derive

$$\int_{-\infty}^{x_0/2} \left(1 - \frac{t}{x_0}\right)^{-(d+1)/2} \sup_{(s, \xi', x, y) \in \mathcal{V}_\omega \times \overline{D'(0, \epsilon)} \times \mathbb{R}^d \times \mathbb{R}^d} |v_s^{(1)}(t, \xi', x, y)| dt \leq 2^{\frac{d+1}{2}} S. \tag{67}$$

On the other hand, we recall that

$$\Re\left(-\frac{1}{2}x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0-t}}\right)\right) \leq -\frac{1}{4}\gamma_\omega |x'|^2.$$

The exponential term is majorized as follows:

$$\begin{aligned} & \left| \left(-\frac{1}{2}x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0-t}}\right)\right) \exp\left(-\frac{1}{2}x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0-t}}\right)\right) \right| \\ & \leq \left(\frac{1}{2}x' \cdot Q_s x' + O(\epsilon|x'|^2)\right) \exp\left(-\frac{1}{4}\gamma_\omega |x'|^2\right). \end{aligned}$$

Consequently,

$$\begin{aligned} & \int_{|x'| < \epsilon \sqrt{x_0 - t}} \left| \left(-\frac{1}{2} x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0 - t}}\right) \right) \right. \\ & \quad \left. \exp\left(-\frac{1}{2} x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0 - t}}\right)\right) \right| dx' \\ & \lesssim \int_{\mathbb{R}^{d-1}} |x'|^2 \exp\left(-\frac{1}{4} \gamma_\omega |x'|^2\right) dx' \\ & < \infty. \end{aligned} \tag{68}$$

Combining (65) through (68), we deduce $J_1 = O(x_0^{-(d+1)/2})$.

Case 2: $l > 1$. Using (49) and decomposing J_l into two parts as in Case 1, we get

$$\begin{aligned} J_l &= \frac{1}{(2\pi)^d} \int_{|\xi'| < \epsilon} \int_{-\infty}^0 \xi_l e^{-w A_s(i\xi')} \int_{-r}^r e^{i(w+|x-y)\xi_1} \tilde{\mu}_s^{(l)}(\xi_1, \xi', x, y) d\xi_1 dw d\xi \\ &= \frac{1}{(2\pi)^d} \int_{|\xi'| < \epsilon} \int_{-\infty}^{|x-y|} \xi_l e^{(-t+|x-y)A_s(i\xi')} v_s^{(l)}(t, \xi', x, y) dt d\xi' \\ &= \frac{1}{(2\pi)^d} \int_{|\xi'| < \epsilon} \int_{-\infty}^{|x-y|/2} \xi_l e^{(-t+|x-y)A_s(i\xi')} v_s^{(l)}(t, \xi', x, y) dt d\xi' \\ & \quad + o(|x-y|^{-d/2}). \end{aligned} \tag{69}$$

By changing the variables as before,

$$\begin{aligned} & x_0^{d/2} \int_{|\xi'| < \epsilon} \int_{-\infty}^{x_0/2} \xi_l e^{(-t+x_0)A_s(i\xi')} v_s^{(l)}(t, \xi', x, y) dt d\xi' \\ &= \int_{-\infty}^{x_0/2} \left(1 - \frac{t}{x_0}\right)^{-d/2} \\ & \quad \int_{|x'| < \epsilon \sqrt{x_0 - t}} x'_l \exp\left(-\frac{1}{2} x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0 - t}}\right)\right) \\ & \quad v_s^{(l)}\left(t, \left(\frac{x'}{\sqrt{x_0 - t}}\right), x, y\right) dx' dt. \end{aligned} \tag{70}$$

In a similar manner, we obtain

$$\begin{aligned} & \int_{-\infty}^{x_0/2} \left(1 - \frac{t}{x_0}\right)^{-d/2} \\ & \int_{|x'| < \epsilon \sqrt{x_0-t}} \left| x'_l \exp\left(-\frac{1}{2}x' \cdot Q_s x' + O\left(\frac{|x'|^3}{\sqrt{x_0-t}}\right)\right) \right| \\ & \left| v_s^{(l)}\left(t, \left(\frac{x'}{\sqrt{x_0-t}}\right), x, y\right) \right| dx' dt \\ & \leq 2^{d/2} S \int_{\mathbb{R}^{d-1}} |x'| \exp\left(-\frac{1}{4}\gamma_\omega |x'|^2\right) dx' < \infty. \end{aligned}$$

This final estimate and (69) and (70) imply $J_l = O(x_0^{-d/2})$. □

7. The full Green’s function asymptotics

The main purpose of this section is to give a detailed proof of Theorem 5.4. Essentially, this theorem is needed for showing that full Green’s function $G_{s,\lambda}$ has the same asymptotics as the reduced Green’s function G_0 as $|x - y| \rightarrow \infty$.

First, we recall that for each unit vector s , $T_s(k) = (1 - \eta(k))(L_s(k) - \lambda)^{-1} + \eta(k)((L_s(k) - \lambda)|_{R(Q_s(k))})^{-1} Q_s(k)$ and the operator T_s is unitarily equivalent (via the Floquet transform) to the direct integral of the operators $T_s(k)$ over \mathcal{O} . Now we observe that the kernel of each projector $P_s(k)$ (see Subsection 5.3) is the smooth function:

$$\frac{\phi(k + i\beta_s, x)\overline{\phi(k - i\beta_s, y)}}{F(k + i\beta_s)},$$

for each k in the support of η . Thus, $(1 - \eta(k))P_s(k)$ is a finite rank smoothing operator on \mathbb{T} . Moreover, we also have $(L_s(k) - \lambda)T_s(k) = T_s(k)(L_s(k) - \lambda) = I - \eta(k)P_s(k)$. Each $T_s(k)$ is a parametrix (i.e., an inverse modulo a smoothing operator) of the elliptic operator $L_s(k) - \lambda$ when $(s, k) \in \mathcal{S}^{d-1} \times \mathcal{O}$. This suggests to study parametrices of the family of elliptic operators $L_s(k) - \lambda$ simultaneously.

7.1. Parameter-dependent periodic pseudodifferential operators. First, we briefly recall some basic definitions of *periodic* (or *toroidal*) pseudodifferential operators (i.e., Ψ DO on the torus \mathbb{T}). We also introduce some useful classes of symbols with parameters and describe some of their properties that we will use.

There are several approaches to defining pseudodifferential operators on the torus. The standard approach based on Hörmander’s symbol classes (see e.g., [31]) uses local smooth structure on the torus \mathbb{T} and thus ignores the group structure on

\mathbb{T} . An alternative approach uses Fourier series with the difference calculus and avoids using local coordinate charts on \mathbb{T} (the details in Chapter 4 in [28]).⁶ To make a distinction, Ruzhansky and Turunen in [28] refer to the symbols in the first approach as *Euclidean symbols* and the symbols in the latter one as *toroidal symbols* (see Section 4.5 in [28]). We recall their definitions for only the Kohn-Nirenberg symbol classes, which we need here.

Definition 7.1. Let m be a real number.

(a) The class $S^m(\mathbb{T} \times \mathbb{R}^d)$ consists of all smooth functions $\sigma(x, \xi)$ on $\mathbb{T} \times \mathbb{R}^d$ such that for any multi-indices α, β ,

$$|D_\xi^\alpha D_x^\beta \sigma(x, \xi)| \leq C_{\alpha\beta} (1 + |\xi|)^{m-|\alpha|},$$

for some constant $C_{\alpha,\beta}$ that depends only on α, β . Symbols in $S^m(\mathbb{T} \times \mathbb{R}^d)$ are called *Euclidean symbols* of order m on \mathbb{T} .

(b) The class $S^m(\mathbb{T} \times \mathbb{Z}^d)$ consists of all functions $\sigma(x, \xi)$ on $\mathbb{T} \times \mathbb{Z}^d$ such that for each $\xi \in \mathbb{Z}^d$, $\sigma(\cdot, \xi) \in C^\infty(\mathbb{T})$ and for any multi-indices α, β ,

$$|\Delta_\xi^\alpha D_x^\beta \sigma(x, \xi)| \leq C_{\alpha\beta} (1 + |\xi|)^{m-|\alpha|},$$

for some constant $C_{\alpha,\beta}$ that depends only on α, β . Here we recall the definition of the forward difference operator Δ_ξ^α with respect to the variable ξ , see [28]. Let f be a complex-valued function defined on \mathbb{Z}^d and $1 \leq j \leq d$. Then we define

$$\Delta_j f(\xi) := f(\xi_1, \dots, \xi_{j-1}, \xi_j + 1, \xi_{j+1}, \dots, \xi_d) - f(\xi),$$

and for any multi-index α ,

$$\Delta_\xi^\alpha := \Delta_1^{\alpha_1} \dots \Delta_d^{\alpha_d}.$$

Symbols in $S^m(\mathbb{T} \times \mathbb{Z}^d)$ are called *toroidal symbols* of order m on \mathbb{T} .

(c) The intersection of all the classes $S^m(\mathbb{T} \times \mathbb{R}^d)$ ($S^m(\mathbb{T} \times \mathbb{Z}^d)$) is denoted by $S^{-\infty}(\mathbb{T} \times \mathbb{R}^d)$ ($S^{-\infty}(\mathbb{T} \times \mathbb{Z}^d)$), which are also called *smoothing symbols*.

Due to Theorem 4.5.3 in [28], a symbol is toroidal of order m if and only if it could be extended in ξ to an Euclidean symbol of the same order m . Such an extension is unique modulo a smoothing symbol. Consequently, we will use the notation $S^m(\mathbb{T})$ for both classes $S^m(\mathbb{T} \times \mathbb{R}^d)$ and $S^m(\mathbb{T} \times \mathbb{Z}^d)$. The two approaches are essentially equivalent in defining pseudodifferential operators on \mathbb{T} whenever the symbol is in the class $S^m(\mathbb{T})$. According to [13], this motivates us to define periodic pseudodifferential operators as follows.

⁶ A different approach to periodic Ψ DOs is introduced by A. Sobolev in [32].

Definition 7.2. Given a symbol $\sigma(x, \xi) \in S^m(\mathbb{T})$, we denote by $\text{Op}(\sigma)$ the corresponding periodic pseudodifferential operator defined by

$$(\text{Op}(\sigma)f)(x) := \sum_{\xi \in \mathbb{Z}^d} \sigma(x, \xi) \tilde{f}(\xi) e^{2\pi i \xi \cdot x}, \tag{71}$$

where $\tilde{f}(\xi)$ is the Fourier coefficient of f at ξ . The right hand side of (71) converges absolutely if, for instance, $f \in C^\infty(\mathbb{T})$.

We also use the notation $\text{Op}(S^m(\mathbb{T}))$ for the set of all periodic pseudodifferential operators $\text{Op}(\sigma)$ with $\sigma \in S^m(\mathbb{T})$.

Since we must deal with parameters s and k , we introduce a suitable class of symbols depending on parameters $(s, k) \in \mathbb{S}^{d-1} \times \mathcal{O}$.

Definition 7.3. The parameter-dependent class $\tilde{S}^m(\mathbb{T})$ consists of symbols $\sigma(s, k; x, \xi)$ satisfying the following conditions.

- For each $(s, k) \in \mathbb{S}^{d-1} \times \mathcal{O}$, the function $\sigma(s, k; \cdot, \cdot)$ is a symbol in the class $S^m(\mathbb{T})$.
- Consider any multi-indices α, β, γ . Then for each $s \in \mathbb{S}^{d-1}$, the function $\sigma(s, \cdot; \cdot, \cdot)$ is smooth on $\mathcal{O} \times \mathbb{T} \times \mathbb{R}^d$, and furthermore,

$$\sup_{s \in \mathbb{S}^{d-1}} |D_k^\alpha D_\xi^\beta D_x^\gamma \sigma(s, k; x, \xi)| \leq C_{\alpha\beta\gamma} (1 + |\xi|)^{m - |\alpha| - |\beta|},$$

for some constant $C_{\alpha\beta\gamma} > 0$ that is independent of s, k, x , and ξ .

Thus, taking derivatives of a symbol in $\tilde{S}^m(\mathbb{T})$ with respect to k improves decay in ξ . We also denote

$$\tilde{S}^{-\infty}(\mathbb{T}) := \bigcap_{m \in \mathbb{R}} \tilde{S}^m(\mathbb{T}).$$

Definition 7.4. For each $m \in \mathbb{R} \cup \{-\infty\}$, we denote by $\text{Op}(\tilde{S}^m(\mathbb{T}))$ the set of all families of periodic pseudodifferential operators $\{\text{Op}(\sigma(s, k; \cdot, \cdot))\}_{(s,k) \in \mathbb{S}^{d-1} \times \mathcal{O}}$, where σ runs over the class $\tilde{S}^m(\mathbb{T})$.

Example 7.5. (1) Suppose that $|\lambda|$ is small enough so that $\max_{s \in \mathbb{S}^{d-1}} |\beta_s| < 1$. Then the family of symbols $\{(1 + (\xi + k + i\beta_s)^2)^{m/2}\}_{(s,k)}$ belongs to the class $\tilde{S}^m(\mathbb{T})$ for any $m \in \mathbb{R}$.

(2) If $a_\alpha(x) \in C^\infty(\mathbb{T})$ and $m \geq 0$, then

$$\left\{ \sum_{|\alpha| \leq m} a_\alpha(x) (\xi + k + i\beta_s)^\alpha \right\}_{(s,k)} \in \tilde{S}^m(\mathbb{T}).$$

- (3) The family of elliptic operators $\{(L_s(k) - \lambda)\}_{(s,k)}$ is in $\text{Op}(\tilde{S}^2(\mathbb{T}))$.
- (4) If $a = \{a(s, k; x, \xi)\}_{(s,k)} \in \tilde{S}^l(\mathbb{T})$ and $b = \{b(s, k; x, \xi)\}_{(s,k)} \in \tilde{S}^m(\mathbb{T})$ then $ab = \{ab(s, k; x, \xi)\}_{(s,k)} \in \tilde{S}^{l+m}(\mathbb{T})$.
- (5) $a(s, k; x, \xi) \in \tilde{S}^l(\mathbb{T})$ implies $D_k^\alpha D_\xi^\beta D_x^\gamma a(s, k; x, \xi) \in \tilde{S}^{l-|\alpha|-|\beta|}(\mathbb{T})$.

The following result will be needed in the next subsection.

Theorem 7.6. *There exists a family of parametrices $\{A_s(k)\}_{(s,k)}$ in the class $\text{Op}(\tilde{S}^{-2}(\mathbb{T}))$ for the family of elliptic operators $\{(L_s(k) - \lambda)\}_{(s,k)}$.*

The reader can refer to Section 8 for the proof of this result as well as some other basic properties of parameter-dependent toroidal Ψ DOs.

7.2. Decay of the Schwartz kernel of T_s

Lemma 7.7. *For all k on a sufficiently small neighborhood of the support of η , λ (< 0) is in the resolvent of the operator $L_s(k)Q_s(k)$ acting on $L^2(\mathbb{T})$. Furthermore, for such k , we have the following identity:*

$$((L_s(k) - \lambda)|_{R(Q_s(k))})^{-1} Q_s(k) = \lambda^{-1} P_s(k) + (L_s(k)Q_s(k) - \lambda)^{-1}. \tag{72}$$

Proof. In the block-matrix form, $(L_s(k)Q_s(k) - \lambda)$ is

$$\left(\begin{array}{c|c} -\lambda P_s(k) & 0 \\ \hline 0 & (L_s(k) - \lambda)|_{R(Q_s(k))} \end{array} \right). \tag{73}$$

This gives the first claim of this lemma. The inverse of (73) is

$$\left(\begin{array}{c|c} -\lambda^{-1} P_s(k) & 0 \\ \hline 0 & ((L_s(k) - \lambda)|_{R(Q_s(k))})^{-1} \end{array} \right),$$

which proves the identity (72). □

The identity (72) implies that for each (s, k) , the operator

$$\eta(k)((L_s(k) - \lambda)|_{R(Q_s(k))})^{-1} Q_s(k)$$

is a periodic pseudodifferential operator in $S^{-2}(\mathbb{T})$. Thus, each of the operators $T_s(k)$ is also in $S^{-2}(\mathbb{T})$ and its symbol is smooth in (s, k) since $P_s(k)$ and $Q_s(k)$ are smooth in (s, k) . Actually, more information about the family of operators $\{T_s(k)\}_{(s,k)}$ and their Schwartz kernels can be obtained.

At first, we want to introduce a class of families of operators whose kernels behave nicely.

Definition 7.8. We denote by \mathcal{S} the set consisting of families of smoothing operators $\{U_s(k)\}_{(s,k)}$ acting on \mathbb{T} so that the following properties hold.

- For any $m_1, m_2 \in \mathbb{R}$, the operator $U_s(k)$ is smooth in k as a $B(H^{m_1}(\mathbb{T}), H^{m_2}(\mathbb{T}))$ -valued function.⁷
- The following uniform condition holds for any multi-index α :

$$\sup_{s,k} \|D_k^\alpha U_s(k)\|_{B(H^{m_1}(\mathbb{T}), H^{m_2}(\mathbb{T}))} < \infty.$$

If the family of smoothing operators $\{U_s(k)\}_{(s,k)}$ is in $\text{Op}(\tilde{\mathcal{S}}^{-\infty}(\mathbb{T}))$, then this family also belongs to \mathcal{S} .

In order to obtain information on Schwartz kernels of a family of operators in \mathcal{S} , we need to use the following standard lemma on Schwartz kernels of integral operators acting on \mathbb{T} .

Lemma 7.9. *Let A be a bounded operator in $L^2(\mathbb{T})$. Suppose that the range of A is contained in $H^m(\mathbb{T})$, where $m > d/2$ and in addition,*

$$\|Af\|_{H^m(\mathbb{T})} \leq C \|f\|_{H^{-m}(\mathbb{T})}$$

for all $f \in L^2(\mathbb{T})$.

Then A is an integral operator whose kernel $K_A(x, y)$ is bounded and uniformly continuous on $\mathbb{T} \times \mathbb{T}$ and the following estimate holds:

$$|K_A(x, y)| \leq \gamma_0 C, \tag{74}$$

where γ_0 is a constant depending only on d and m .

The fact can be found in Lemma 2.2 in [2].

Now we can state a useful property of Schwartz kernels of a family of operators in \mathcal{S} .

Corollary 7.10. *If $\{U_s(k)\}_{(s,k)}$ is a family of smoothing operators in \mathcal{S} , then the Schwartz kernel $K_{U_s}(k, x, y)$ of the operator $U_s(k)$ satisfies*

$$\sup_{s,k,x,y} |D_k^\alpha K_{U_s}(k, x, y)| < \infty,$$

for any multi-index α .

⁷ We remind the reader that $B(E, F)$ denotes the space of all bounded linear operators from the Banach space E to F .

Proof. We pick any $m > d/2$. Then by Definition 7.8, we have

$$\sup_{s,k} \|D_k^\alpha U_s(k) f\|_{H^m(\mathbb{T})} \leq C_\alpha \|f\|_{H^{-m}(\mathbb{T})}.$$

Applying Lemma 7.9, the estimates (74) hold for kernels $D_k^\alpha K_{U_s}(k, x, y)$ of the operators $D_k^\alpha U_s(k)$ uniformly in (s, k) . \square

We now go back to the family of operators $T_s(k)$.

Proposition 7.11. *There is a family of periodic pseudodifferential operators $\{A_s(k)\}_{(s,k)}$ in $\text{Op}(\tilde{S}^{-2}(\mathbb{T}))$ such that the family of operators $\{T_s(k) - A_s(k)\}_{(s,k)}$ belongs to \mathcal{S} .*

Proof. Due to Theorem 7.6, there is a family of operators $\{A_s(k)\}_{(s,k)}$ in $\text{Op}(\tilde{S}^{-2}(\mathbb{T}))$ and a family of operators $\{R_s(k)\}_{(s,k)}$ in $\text{Op}(\tilde{S}^{-\infty}(\mathbb{T}))$ such that

$$(L_s(k) - \lambda)A_s(k) = I - R_s(k).$$

Since $T_s(k)(L_s(k) - \lambda) = I - \eta(k)P_s(k)$, we deduce that

$$T_s(k) = A_s(k) - \eta(k)P_s(k)A_s(k) + T_s(k)R_s(k). \quad (75)$$

Now it remains to show that the two families of smoothing operators $\{T_s(k)R_s(k)\}_{(s,k)}$ and $\{\eta(k)P_s(k)A_s(k)\}_{(s,k)}$ are in \mathcal{S} . Let us fix any two real numbers m_1, m_2 and a multi-index α . Notice that $(L_s(k) - \lambda)$ is analytic in k as a $B(H^{m_2}(\mathbb{T}), H^{m_2-2}(\mathbb{T}))$ -valued function and also,

$$\sup_{s,k} \|D_k^\alpha (L_s(k) - \lambda)\|_{B(H^{m_2}(\mathbb{T}), H^{m_2-2}(\mathbb{T}))} < \infty.$$

Due to Lemma 7.7,

$$T_s(k) = (1 - \eta(k))(L_s(k) - \lambda)^{-1} + \eta(k)\lambda^{-1}P_s(k) + \eta(k)(L_s(k)Q_s(k) - \lambda)^{-1}.$$

Thus, $T_s(k)$ is smooth in k as a $B(H^{m_2-2}(\mathbb{T}), H^{m_2}(\mathbb{T}))$ -valued function and moreover,

$$\sup_{s,k} \|D_k^\alpha T_s(k)\|_{B(H^{m_2-2}(\mathbb{T}), H^{m_2}(\mathbb{T}))} < \infty. \quad (76)$$

Since $\{R_s(k)\}$ is in $\text{Op}(\tilde{S}^{-\infty}(\mathbb{T}))$, $R_s(k)$ is smooth in k as a $B(H^{m_1}(\mathbb{T}), H^{m_2-2}(\mathbb{T}))$ -valued function and furthermore,

$$\sup_{s,k} \|D_k^\alpha R_s(k)\|_{B(H^{m_1}(\mathbb{T}), H^{m_2-2}(\mathbb{T}))} < \infty. \quad (77)$$

By (76), (77), and Leibnitz’s rule, we deduce that $T_s(k)R_s(k)$ is smooth in k as a $B(H^{m_1}(\mathbb{T}), H^{m_2}(\mathbb{T}))$ -valued function and the corresponding uniform estimate also holds. Hence, we conclude that the family $\{T_s(k)R_s(k)\}_{(s,k)}$ belongs to \mathcal{S} . Meanwhile, since $\{\eta(k)P_s(k)\}_{(s,k)}$ is in \mathcal{S} and $\{D_k^\alpha A_s(k)\}_{(s,k)}$ is a toroidal pseudodifferential operator of order $2 - |\alpha|$ for any multi-index α , we could repeat the above argument to show that the family $\{\eta(k)P_s(k)A_s(k)\}_{(s,k)}$ is also in \mathcal{S} . \square

We need the following important estimate of Schwartz kernels of operators $T_s(k)$.

Corollary 7.12. *Let $K_s(k, x, y)$ be the Schwartz kernel of the operator $T_s(k)$. Let $N > d - 2$. If α is a multi-index such that $|\alpha| = N$, then each $D_k^\alpha K_s(k, x, y)$ is a continuous function on $\mathbb{T} \times \mathbb{T}$ and the following estimate also holds uniformly with respect to (x, y) :*

$$\sup_{(s,k) \in \mathbb{S}^{d-1} \times \mathcal{O}} |D_k^\alpha K_s(k, x, y)| < \infty.$$

Proof. Due to Proposition 7.11, the operator $T_s(k)$ is a sum of operators $A_s(k)$ and $U_s(k)$ such that $\{A_s(k)\}_{(s,k)} \in \text{Op}(\tilde{\mathcal{S}}^{-2}(\mathbb{T}))$ and $\{U_s(k)\}_{(s,k)} \in \mathcal{S}$. In particular,

$$K_s(k, x, y) = K_{A_s}(k, x, y) + K_{U_s}(k, x, y).$$

Recall that in the distributional sense, the Schwartz kernel $K_{A_s}(k, x, y)$ of the periodic pseudodifferential operator $A_s(k)$ is given by

$$\sum_{\xi \in \mathbb{Z}^d} \sigma(s, k; x, \xi) e^{2\pi i \xi \cdot (x-y)},$$

where $\sigma(s, k; x, \xi)$ is the symbol of the operator $A_s(k)$.

Since $\{\sigma(s, k; x, \xi)\}_{(s,k)}$ is in $\tilde{\mathcal{S}}^{-2}(\mathbb{T})$,

$$|e^{2\pi i \xi \cdot (x-y)} D_k^\alpha \sigma(s, k; x, \xi)| \lesssim (1 + |\xi|)^{-2-N}.$$

Since $-(2 + N) < -d$, the sum

$$\sum_{\xi \in \mathbb{Z}^d} D_k^\alpha \sigma(s, k; x, \xi) e^{2\pi i \xi \cdot (x-y)}$$

converges absolutely and moreover,

$$\sup_{(s,k,x,y) \in \mathbb{S}^{d-1} \times \mathcal{O} \times \mathbb{T} \times \mathbb{T}} |D_k^\alpha K_{A_s}(k, x, y)| \lesssim \sum_{\xi \in \mathbb{Z}^d} (1 + |\xi|)^{-(d+1)} < \infty.$$

Combining this with Corollary 7.10, we complete the proof. \square

Notation 7.13. Let ψ be a function on \mathbb{R}^d and γ be a vector in \mathbb{R}^d , then $\tau_\gamma\psi$ is the γ -shifted version of ψ . Namely, it is defined as follows:

$$\tau_\gamma\psi(\cdot) = \psi(\cdot + \gamma).$$

We denote by \mathcal{P} the subset of $C_0^\infty(\mathbb{R}^d)$ consisting of all functions ψ such that its support is connected, and if γ is a *non-zero vector in \mathbb{Z}^d* , then the support of $\tau_\gamma\psi$ does not intersect with the support of ψ .

Definition 7.14. Since \mathbb{R}^d is the universal covering space of \mathbb{T} , we can consider the covering map

$$\pi: \mathbb{R}^d \longrightarrow \mathbb{R}^d / \mathbb{Z}^d = \mathbb{T}.$$

In particular, $\pi(x + \gamma) = \pi(x)$ for any $x \in \mathbb{R}^d$ and $\gamma \in \mathbb{Z}^d$.

A *standard fundamental domain* (with respect to the covering map π) is of the form $[0, 1]^d + \gamma$ for some vector γ in \mathbb{R}^d . Thus, a standard fundamental domain is a fundamental domain of \mathbb{R}^d with respect to the lattice \mathbb{Z}^d .

Using Definition 5.1 of the Floquet transform \mathcal{F} , we obtain the following formula.

Lemma 7.15. *Let ϕ and θ be any two smooth functions in \mathcal{P} . Then the Schwartz kernel $K_{s,\phi,\theta}$ of the operator $\phi T_s \theta$ satisfies the following identity for any $(x, y) \in \mathbb{R}^d \times \mathbb{R}^d$:*

$$K_{s,\phi,\theta}(x, y) = \frac{1}{(2\pi)^d} \int_{\mathcal{O}} e^{ik \cdot (x-y)} \phi(x) K_s(k, \pi(x), \pi(y)) \theta(y) dk.$$

Proof. Since both $\phi, \theta \in \mathcal{P}$, there are standard fundamental domains W_ϕ and $W_\theta \subset \mathbb{R}^d$ so that

$$\text{supp}(\phi) \subset \overset{\circ}{W}_\phi, \quad \text{supp}(\theta) \subset \overset{\circ}{W}_\theta.$$

Then, it suffices to show that $\langle \phi T_s \theta f, g \rangle$ equals

$$\frac{1}{(2\pi)^d} \int_{W_\phi} \int_{W_\theta} \int_{\mathcal{O}} e^{ik \cdot (x-y)} (\phi \bar{g})(x) K_s(k, \pi(x), \pi(y)) (\theta f)(y) dk dy dx,$$

for any f, g in $C^\infty(\mathbb{R}^d)$.

We observe that

$$\begin{aligned} \langle \phi T_s \theta f, g \rangle &= \langle \mathcal{F} \phi T_s \theta f, \mathcal{F} g \rangle \\ &= \frac{1}{(2\pi)^d} \left\langle (\mathcal{F} \phi \mathcal{F}^{-1}) \left(\int_{\mathcal{O}}^\oplus T_s(k) dk \right) \mathcal{F}(\theta f), \mathcal{F} g \right\rangle \\ &= \frac{1}{(2\pi)^d} \left\langle \left(\int_{\mathcal{O}}^\oplus T_s(k) dk \right) \mathcal{F}(\theta f), \mathcal{F}(\bar{\phi} g) \right\rangle. \end{aligned}$$

Since $\theta \in \mathcal{P}$, for any y in W_θ , we have

$$\mathcal{F}(\theta f)(k, \pi(y)) = (\theta f)(y)e^{-ik \cdot y}.$$

Similarly,

$$\mathcal{F}(\bar{\phi}g)(k, \pi(x)) = (\bar{\phi}g)(x)e^{-ik \cdot x}, \quad \text{for all } x \in W_\phi.$$

We also have

$$\left(\int_{\mathcal{O}}^{\oplus} T_s(k) dk \right) (\mathcal{F}(\theta f))(k, \pi(x)) = T_s(k)(\mathcal{F}(\theta f)(k, \cdot))(\pi(x)).$$

Consequently,

$$\begin{aligned} & \left\langle \left(\int_{\mathcal{O}}^{\oplus} T_s(k) dk \right) \mathcal{F}(\theta f), \mathcal{F}(\bar{\phi}g) \right\rangle \\ &= \int_{\mathcal{O}} \int_{W_\phi} T_s(k) (\mathcal{F}(\theta f)(k, \cdot))(\pi(x)) \overline{(\bar{\phi}g)(x)e^{-ik \cdot x}} dx dk \\ &= \int_{\mathcal{O}} \int_{W_\phi} \int_{W_\theta} K_s(k, \pi(x), \pi(y)) \mathcal{F}(\theta f)(k, \pi(y)) (\phi \bar{g})(x) e^{ik \cdot x} dy dx dk \\ &= \int_{\mathcal{O}} \int_{W_\phi} \int_{W_\theta} e^{ik \cdot (x-y)} K_s(k, \pi(x), \pi(y)) (\theta f)(y) (\phi \bar{g})(x) dy dx dk. \end{aligned}$$

Using Fubini's theorem to rewrite the above integral, we have the desired identity. \square

Proposition 7.16. *Consider any two smooth compactly supported functions ϕ and θ on \mathbb{R}^d such that their supports are disjoint. Then the kernel $K_{s,\phi,\theta}(x, y)$ is continuous on $\mathbb{R}^d \times \mathbb{R}^d$ and moreover, it satisfies the following decay:*

$$\sup_s |K_{s,\phi,\theta}(x, y)| \leq C_N |\phi(x)\theta(y)| \cdot |x - y|^{-N},$$

for any $N > d - 2$. Here, the constant C_N is independent of ϕ and θ .

Proof. By using partitions of unity, any smooth compactly supported function can be written as a finite sum of smooth functions in the set \mathcal{P} . Thus, we can assume without loss of generality that both ϕ and θ belong to \mathcal{P} .

First, observe that for any $(k, n) \in \mathcal{O} \times \mathbb{Z}^d$,

$$T_s(k + 2\pi n) = \mathcal{M}_n^{-1} T_s(k) \mathcal{M}_n,$$

where \mathcal{M}_n is the multiplication operator on $L^2(\mathbb{T})$ by the exponential function $e^{2\pi i n \cdot x}$.

Hence,

$$\nabla_k^\alpha K_s(k + 2\pi n, \pi(x), \pi(y)) = e^{-2\pi i n \cdot \pi(x)} \nabla_k^\alpha K_s(k, \pi(x), \pi(y)) e^{2\pi i n \cdot \pi(y)},$$

for any multi-index α . Since $e^{2\pi i n \cdot x} = e^{2\pi i n \cdot \pi(x)}$ for any $x \in \mathbb{R}^d$, we obtain

$$e^{i(k+2\pi n) \cdot (x-y)} \nabla_k^\alpha K_s(k + 2\pi n, \pi(x), \pi(y)) = e^{ik \cdot (x-y)} \nabla_k^\alpha K_s(k, \pi(x), \pi(y)). \tag{78}$$

Applying Lemma 7.15, we then use integration by parts (all boundary terms vanish when applying integration by parts due to (78)) to derive that for any $|\alpha| = N$,

$$(2\pi)^d (i(x - y))^\alpha K_{s,\phi,\theta}(x, y) = \phi(x)\theta(y) \int_{\mathfrak{O}} e^{ik \cdot (x-y)} \nabla_k^\alpha K_s(k, \pi(x), \pi(y)) dk.$$

Suppose $N > d - 2$. Then by applying Corollary 7.12, the above integral is absolutely convergent and it is also uniformly bounded in (s, x, y) . Consequently, the kernel $K_{s,\phi,\theta}(x, y)$ is continuous. Furthermore,

$$\begin{aligned} \sup_s |K_{s,\phi,\theta}(x, y)| &\lesssim |\phi(x)\theta(y)| \cdot \min_{|\alpha|=N} |(x - y)^\alpha|^{-1} \\ &\lesssim |\phi(x)\theta(y)| \cdot |x - y|^{-N}. \end{aligned} \quad \square$$

We now have enough tools to approach our goal.

Proof of Theorem 5.4. Let us fix a point (s, x) in $\mathbb{S}^{d-1} \times \mathbb{R}^d$. Now we consider a point $y = x + st$, where t is a real number. When $|t| > 0$, we can choose two cut-off functions ϕ and θ such that ϕ and θ equal 1 on some neighborhoods of x and y , respectively, and also, the supports of these two functions are disjoint. Then, Proposition 7.16 implies that the kernel $K_s(x, y)$ is continuous at (x, y) since it coincides with $K_{s,\phi,\theta}$ on a neighborhood of (x, y) . This yields the first statement about the continuity off diagonal of K_s . Again, by Proposition 7.16, we obtain

$$\sup_s |K_s(x, y)| = \sup_s |K_{s,\phi,\theta}(x, y)| \leq C_N |x - y|^{-N},$$

which proves the last statement. □

8. Some results on parameter-dependent toroidal Ψ DOs

The aim in this section is to provide some results needed to complete the proof of Theorem 7.6. We adopt the approach of [13] to periodic elliptic differential operators.

The next two theorems are straightforward modifications of the proofs for non-parameter toroidal Ψ DOs.

Theorem 8.1 (asymptotic summation theorem). *Given families of symbols $b_l \in \tilde{S}^{m-l}(\mathbb{T})$, where each family $b_l = \{b_l(s, k)\}_{(s,k)}$ for $l = 0, 1, \dots$, there exists a family of symbols b in $\tilde{S}^m(\mathbb{T})$ such that*

$$\{b(s, k) - \sum_{i < l} b_i(s, k)\}_{(s,k)} \in \tilde{S}^{m-l}(\mathbb{T}). \tag{79}$$

We will write $b \sim \sum_l b_l$ if b satisfies (79).

Proof. Step 1. Let $n = m + \epsilon$ for some $\epsilon > 0$. Then

$$|b_l(s, k; x, \xi)| \leq C_l(1 + |\xi|)^{m-l} = \frac{C_l(1 + |\xi|)^{n-l}}{(1 + |\xi|)^\epsilon}.$$

Thus, there is a sequence $\{\eta_l\}_{l \geq 1}$ such that $\eta_l \rightarrow +\infty$ and

$$|b_l(s, k; x, \xi)| < \frac{1}{2^l}(1 + |\xi|)^{n-l}$$

for $|\xi| > \eta_l$. Let $\rho \in C^\infty(\mathbb{R})$ satisfy that $0 \leq \rho \leq 1$, $\rho(t) = 0$ whenever $|t| < 1$ and $\rho(t) = 1$ whenever $|t| > 2$. We define

$$b(s, k; x, \xi) = \sum_l \rho\left(\frac{|\xi|}{\eta_l}\right) b_l(s, k; x, \xi).$$

Since only a finite number of summands are non-zero on any compact subset of $\mathbb{T} \times \mathbb{R}^d$, $b(s, \cdot; \cdot, \cdot) \in C^\infty(\mathcal{O} \times \mathbb{T} \times \mathbb{R}^d)$. Moreover, $b(s, k) - \sum_{r < l} b_r(s, k)$ is equal to

$$\sum_{r < l} \left(\rho\left(\frac{|\xi|}{\eta_r}\right) - 1\right) b_r(s, k) + \rho\left(\frac{|\xi|}{\eta_l}\right) b_l(s, k) + \sum_{r > l} \rho\left(\frac{|\xi|}{\eta_r}\right) b_r(s, k).$$

The first summand is compactly supported while the second summand is in $S^{m-l}(\mathbb{T})$. Now let $\epsilon < 1$. Then, the third summand is bounded from above by

$$\sum_{r > l} \frac{1}{2^r} (1 + |\xi|)^{n-r} \leq (1 + |\xi|)^{n-l-1} \leq (1 + |\xi|)^{m-l}.$$

Consequently,

$$\sup_{s \in \mathbb{S}^{d-1}} \left| b(s, k) - \sum_{r < l} b_r(s, k) \right| \leq C(1 + |\xi|)^{m-l}.$$

Step 2. For $|\alpha| + |\beta| + |\gamma| \leq N$, one can choose η_l such that

$$\sup_{s \in \mathbb{S}^{d-1}} \left| D_k^\alpha D_\xi^\beta D_x^\gamma b_l(s, k; x, \xi) \right| \leq \frac{1}{2^l} (1 + |\xi|)^{n-l-|\alpha|-|\beta|}$$

for $\eta_l < |\xi|$. The same argument as in Step 1 implies that

$$\sup_{s \in \mathbb{S}^{d-1}} \left| D_k^\alpha D_\xi^\beta D_x^\gamma (b(s, k) - \sum_{r < l} b_r(s, k)) \right| \leq C_N (1 + |\xi|)^{m-l-|\alpha|-|\beta|}. \tag{80}$$

Step 3. The sequence of η_l 's in Step 2 depends on N . We denote this sequence by $\eta_{l,N}$ to indicate this dependence on N . By induction, we can assume that for all l , $\eta_{l,N} \leq \eta_{l,N+1}$. Applying the Cantor diagonal process to this family of sequences, i.e., let $\eta_l = \eta_{l,l}$ then b has the property (80) for every N . \square

Theorem 8.2 (composition formula). *Let $a = \{a(s, k)\}$ be a family of symbols in $\tilde{S}^l(\mathbb{T})$ and $Q(x, D) = \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha$ be a differential operators of order $m \geq 0$ with smooth periodic coefficients $a_\alpha(x)$. Then the family of periodic pseudodifferential operators $\{Q(x, D + k + i\beta_s) \text{Op}(a(s, k))\}_{(s,k)} \in \text{Op}(\tilde{S}^{l+m}(\mathbb{T}))$. Indeed, we have*

$$Q(x, D + k + i\beta_s) \text{Op}(a(s, k)) = \text{Op}((Q \circ a)(s, k)),$$

where

$$(Q \circ a)(s, k; x, \xi) = \sum_{|\alpha| \leq m} \frac{1}{\alpha!} D_\xi^\alpha Q(x, \xi + k + i\beta_s) D_x^\alpha a(s, k; x, \xi) \tag{81}$$

Proof. The composition formula (81) is obtained for each (s, k) is standard in pseudodifferential operator theory (see e.g., [13, 28, 31]). We only need to check that the family of symbols $\{(Q \circ a)(s, k; x, \xi)\}_{(s,k)}$ is in $\tilde{S}^{l+m}(\mathbb{T})$. But this fact follows easily from (81) and Leibnitz's formula. \square

We now finish the proof of Theorem 7.6.

Theorem 8.3 (inversion formula). *There exists a family $a = \{a(s, k)\}_{(s,k)}$ in $\tilde{S}^{-2}(\mathbb{T})$ and a family $r = \{r(s, k)\}_{(s,k)}$ in $\tilde{S}^{-\infty}(\mathbb{T})$ such that*

$$(L_s(k) - \lambda) \text{Op}(a(s, k)) = I - \text{Op}(r(s, k)).$$

Proof. Let

$$L_0(s, k; x, \xi) := \sum_{|\alpha|=2} a_\alpha(x) (\xi + k + i\beta_s)^\alpha,$$

$$\|a\|_\infty := \sum_{|\alpha|=2} \|a_\alpha(\cdot)\|_{L^\infty(\mathbb{T})},$$

and

$$M := \max_{(s,k) \in \mathbb{S}^{d-1} \times \mathbb{O}} (|k|^2 + \theta^{-1} \|a\|_\infty |\beta_s|^2 + \theta^{-1}),$$

where θ is the ellipticity constant in (2). Whenever $|\xi| > (2M)^{1/2}$,

$$\begin{aligned} |L_0(s, k; x, \xi)| &\geq \Re(L_0(s, k; x, \xi)) \\ &\geq \theta|\xi + k|^2 - \sum_{|\alpha|=2} a_\alpha(x)(\beta_s)^\alpha \\ &\geq \theta\left(\frac{|\xi|^2}{2} - |k|^2\right) - \|a\|_\infty |\beta_s|^2 \\ &> 1. \end{aligned}$$

Let $\rho \in C^\infty(\mathbb{R})$ be a function satisfying $\rho(t) = 0$ when $|t| < (2M)^{1/2}$ and $\rho(t) = 1$ when $|t| > 2M^{1/2}$. We define the function

$$a_0(s, k)(x, \xi) = \rho(|\xi|) \frac{1}{L_0(s, k; x, \xi)}. \tag{82}$$

Then $a_0 := \{a_0(s, k)\}_{(s,k)}$ is well-defined and belongs to $\tilde{S}^{-2}(\mathbb{T})$. The next lemma is the final piece we need to complete the proof of the theorem.

Lemma 8.4. (i) *If $b = \{b(s, k)\}_{(s,k)} \in \tilde{S}^l(\mathbb{T})$ then $b - (L - \lambda) \circ (a_0 b) \in \tilde{S}^{l-1}(\mathbb{T})$.*

(ii) *There exists a sequence of families of symbols $a_l = \{a_l(s, k)\}_{(s,k)}$ in $\tilde{S}^{-2-l}(\mathbb{T})$, $l = 0, 1, \dots$ and a sequence of families of symbols $r_l = \{r_l(s, k)\}_{(s,k)}$ in $\tilde{S}^{-l}(\mathbb{T})$, $l = 0, 1, \dots$ such that a_0 is the family of symbols in (82), $r_0(s, k) = 1$ for every (s, k) and for all l ,*

$$(L - \lambda) \circ a_l = r_l - r_{l+1}.$$

Proof. (i) Let

$$p(s, k) = (L(s, k) - \lambda)(x, \xi) - L_0(s, k; x, \xi)$$

so that $p = \{p(s, k)\}_{(s,k)} \in \tilde{S}^1(\mathbb{T})$ and hence, $p \circ (a_0 b)$ is in $\tilde{S}^{l-1}(\mathbb{T})$ due to Theorem 8.2. Moreover, $b - L_0 a_0 b = (1 - \rho(|\xi|))b$ is a family of symbols whose ξ -supports are compact and thus it is in $\tilde{S}^{-\infty}(\mathbb{T})$. We can now derive again from the composition formula (81) when $P := L_0$ that

$$(L - \lambda) \circ (a_0 b) = L_0 \circ (a_0 b) + p \circ (a_0 b) = L_0 a_0 b + \dots = b + \dots,$$

where the dots are the terms in $\tilde{S}^{l-1}(\mathbb{T})$.

(ii) Recursively, let $a_l = a_0 r_l$ and $r_{l+1} = r_l - (L - \lambda) \circ a_l$. By part (i), $r_{l+1} \in \tilde{S}^{-(l+1)}(\mathbb{T})$. △

Now let a be the asymptotic sum of the families of symbols a_l , i.e., $a \sim \sum_l a_l$. Then

$$(L - \lambda) \circ a \sim \sum_l (L - \lambda) \circ a_l = \sum_l r_l - r_{l+1} = r_0 = 1,$$

which implies that $1 - (L - \lambda) \circ a \sim 0$. In other words, this means that $r := 1 - (L - \lambda) \circ a \in \tilde{\mathcal{S}}^{-\infty}(\mathbb{T})$. Hence, there exists a family of symbols a in $\tilde{\mathcal{S}}^{-2}(\mathbb{T})$ and a family of symbols r in $\tilde{\mathcal{S}}^{-\infty}(\mathbb{T})$ satisfying $(L - \lambda) \circ a = 1 - r$. Finally, an application of Theorem 8.2 completes the proof of Theorem 7.6. \square

9. Some auxiliary statements

9.1. A lemma on the principle of non-stationary phase

Lemma 9.1. *Let M be a compact manifold (with or without boundary) and $a: \mathbb{R} \times M \rightarrow \mathbb{C}$ be a smooth function with compact support. Then for any $N > 0$, there exists a constant $C_N > 0$ so that the following estimate holds for any non-zero $t \in \mathbb{R}$:*

$$\sup_{x \in M} \left| \int_{-\infty}^{\infty} e^{ity} a(y, x) dy \right| \leq C_N |t|^{-N}. \tag{83}$$

Here C_N depends only on N , the diameter R of the y -support of a and $\sup_{x,y} |\partial_y^N a|$.

Proof. Let $t \neq 0$. Applying integration by parts repeatedly (N -times), it follows that

$$\begin{aligned} \left| \int_{-\infty}^{\infty} e^{ity} a(y, x) dy \right| &= |t|^{-N} \left| \int_{-\infty}^{\infty} e^{ity} \partial_y^N a(y, x) dy \right| \\ &\leq R \sup_{x,y} |\partial_y^N a| \cdot |t|^{-N}. \end{aligned} \tag{84}$$

9.2. The Weierstrass preparation theorem

Theorem 9.2. *Let $f(t, z)$ be an analytic function of $(t, z) \in \mathbb{C}^{1+n}$ in a neighborhood of $(0, 0)$ such that $(0, 0)$ is a simple zero of f , i.e.,*

$$f(0, 0) = 0, \quad \frac{\partial f}{\partial t}(0, 0) \neq 0.$$

Then there is a unique factorization

$$f(t, z) = (t - A(z))B(t, z),$$

where A, B are analytic in a neighborhood of 0 and $(0, 0)$ respectively. Moreover, $B(0, 0) \neq 0$ and $A(0) = 0$.

The proof of a more general version of this theorem could be found in Theorem 7.5.1 in [15].

9.3. Proofs of Proposition 4.1 and Lemma 5.3

Remark 9.3. Consider a domain \mathcal{D} of \mathbb{C}^d and let $f: \mathcal{D} \rightarrow \mathbb{C}$ be a holomorphic function. For $z \in \mathbb{C}^d$, write $z = x + iy$ where $x, y \in \mathbb{R}^d$. Now we fix a vector β in \mathbb{R}^d and denote $\mathcal{D}_\beta = (\mathcal{D} - i\beta) \cap \mathbb{R}^d$. If this intersection is non-empty, we may consider the restriction $k \rightarrow f(k + i\beta)$ as a real analytic function defined on a subdomain \mathcal{D}_β of \mathbb{R}^d . Thanks to Cauchy–Riemann equations of f , we do not need to make any distinction between derivatives of f with respect to x (when f is viewed as a real analytic one) or z (when f is considered as a complex analytic one) at every point in \mathcal{D}_β since

$$\frac{\partial f}{\partial x_l}(k + i\beta) = \frac{\partial f}{\partial z_l}(k + i\beta) = -i \frac{\partial f}{\partial y_l}(k + i\beta), \quad 1 \leq l \leq d.$$

For higher order derivatives, we use induction and the above identity to obtain

$$\partial_x^\alpha f(k + i\beta) = \partial_z^\alpha f(k + i\beta) = (-i)^{|\alpha|} \partial_y^\alpha f(k + i\beta),$$

for any multi-index α . We use these facts implicitly for the function λ_j . When dealing with the analytic function $f = \lambda_j$ in this part, denote $\partial^\alpha \lambda_j$ to indicate either its x or z -derivatives.

We also want to mention this simple relation between derivatives of λ_j and E :

$$\partial^\alpha E(\beta) = \partial_y^\alpha \lambda_j(k_0 + i\beta) = i^{|\alpha|} \partial^\alpha \lambda_j(k_0 + i\beta).$$

Proof of Proposition 4.1. We recall from Section 2 that V is an open neighborhood of k_0 in \mathbb{C}^d such that properties P1–P6 are satisfied. Note that V depends only on the local structure at k_0 of the dispersion branch λ_j of L . Denote $\mathcal{O}_s = \{k + it\beta_s: k \in \mathcal{O}, t \in [0, 1]\}$ for each $s \in \mathbb{S}^{d-1}$. For $C > 0$ (which is defined later), set $M_{s,C} = \mathcal{O}_s \cap \{z \in \mathbb{C}^d: |\Re(z) - k_0| < C\}$ and $N_{s,C} = \mathcal{O}_s \setminus M_{s,C}$. For C and $|\lambda|$ small enough, we can suppose $M_{s,C} \Subset V$ since β_s is small too. We also assume that $|\lambda| \leq \epsilon_0$.⁸

⁸ Recall the definition of ϵ_0 from P3.




Figure 1. An illustration of the regions $M_{s,C}$ and $N_{s,C}$ when $k_0 \equiv 0$.

For any point $z = k + it\beta_s \in M_{s,C}$, we want to show if $\lambda \in \sigma(L(z))$, it forces $z = k_0 + i\beta_s$. By P3, this is the same as showing the equation $\lambda_j(z) = \lambda$ has no solution z in $M_{s,C}$ except for the trivial solution $z = k_0 + i\beta_s$. Suppose for contradiction $\lambda_j(k + it\beta_s) = \lambda = \lambda(\beta_s)$ for some $t \in [0, 1]$ and k in $\{k \in \mathcal{O} \mid 0 < |k - k_0| < C\}$. By Taylor expanding around $k_0 + it\beta_s$, there is some $\gamma \in (0, 1)$ such that

$$\begin{aligned} & \lambda - \lambda_j(k_0 + it\beta_s) \\ &= \left((k - k_0) \cdot \nabla \lambda_j(k_0 + it\beta_s) + \sum_{|\alpha|=3} \frac{(k - k_0)^\alpha}{\alpha!} \partial^\alpha \lambda_j(k_0 + it\beta_s) \right) \\ &+ \left(\frac{1}{2} (k - k_0) \cdot \text{Hess}(\lambda_j)(k_0 + it\beta_s)(k - k_0) \right. \\ &\quad \left. + \sum_{|\alpha|=4} \frac{(k - k_0)^\alpha}{\alpha!} \partial^\alpha \lambda_j(\gamma(k - k_0) + k_0 + it\beta_s) \right). \end{aligned} \tag{84}$$

If $|\alpha|$ is odd, then by Remark 9.3 and the fact that E is real, we have

$$\partial^\alpha \lambda_j(k_0 + it\beta_s) = \frac{1}{i^{|\alpha|}} \partial^\alpha E(t\beta_s) \in i\mathbb{R}.$$

Taking the real part of equation (84) to get

$$\begin{aligned} E(\beta_s) - E(t\beta_s) &= -\frac{1}{2} (k - k_0) \cdot \text{Hess}(E)(t\beta_s)(k - k_0) \\ &+ \sum_{|\alpha|=4} \frac{(k - k_0)^\alpha}{\alpha!} \Re(\partial^\alpha \lambda_j(\gamma(k - k_0) + k_0 + it\beta_s)). \end{aligned}$$

The left-hand side is bounded above by $(1 - t)\lambda \leq 0$ because of the concavity of E ($E(t\beta_s) \geq tE(\beta_s) = t\lambda$). On the other hand, by P5,

$$-\frac{1}{2}(k - k_0) \cdot \text{Hess}(E)(t\beta_s)(k - k_0) \geq \frac{1}{4}|k - k_0|^2 \min \sigma(\text{Hess}(\lambda_j)(k_0))$$

and

$$\left| \sum_{|\alpha|=4} \frac{(k - k_0)^\alpha}{\alpha!} \Re(\partial^\alpha \lambda_j(\gamma(k - k_0) + k_0 + it\beta_s)) \right| \leq C(d)|k - k_0|^4 \max_{z \in \bar{V}, |\alpha|=4} |\partial^\alpha \lambda_j(z)|.$$

We simply choose

$$C^2 < \frac{\min \sigma(\text{Hess}(\lambda_j)(k_0))}{C(d) \max_{z \in \bar{V}, |\alpha|=4} |\partial^\alpha \lambda_j(z)|}$$

to get a contradiction if $k \neq k_0$.

For the remaining part, we just need to treat points $k + it\beta_s$ in $N_{s,C}$. We have $\lambda \in \rho(L(k))$, for all $k \in \mathbb{R}^d$. The idea is to adapt the upper-semicontinuity of the spectrum of an analytic family of type \mathcal{A} on \mathbb{C}^d , following [18]. For any $k \in \mathcal{O}$ and $z \in \mathbb{C}^d$, the composed operators $(L(k + z) - L(k))(L(k) - \lambda)^{-1}$ are closed and defined on $L^2(\mathbb{T})$ and by closed graph theorem, these are bounded operators. Clearly,

$$L(k + z) - \lambda = (1 + (L(k + z) - L(k))(L(k) - \lambda)^{-1})(L(k) - \lambda).$$

Thus, λ is in the resolvent of $L(k + z)$ if the operator

$$1 + (L(k + z) - L(k))(L(k) - \lambda)^{-1}$$

is invertible. Hence, it is enough to show that there is some positive constant τ such that for any $k \in \mathcal{O}$ and $|z| < \tau$,

$$\|(L(k + z) - L(k))(L(k) - \lambda)^{-1}\|_{\text{op}} < 1/2, \quad |k - k_0| \geq C, \quad (85)$$

where the operator norm on $L^2(\mathbb{T})$ is denoted by $\|\cdot\|_{\text{op}}$. If $|\lambda|$ is small enough so that we have $\max_{s \in \mathbb{S}^{d-1}} |\beta_s| < \tau$ and then (85) implies that $\lambda \in \rho(L(k + it\beta_s))$ for any $t \in [0, 1]$.

Finally, we will use some energy estimates of linear elliptic equations and spectral theory to obtain (85). Observe that,

$$L(k+z) - L(k) = z \cdot A(x)(D+k) + (D+k) \cdot A(x)z + z \cdot A(x)z.$$

For $v \in H^1(\mathbb{T})$ and $|z| < 1$, there is some constant $C_1 > 0$ (independent of z) such that

$$\|(z \cdot A(x)(D+k) + (D+k) \cdot A(x)z + z \cdot A(x)z)v\|_{L^2(\mathbb{T})} \leq C_1|z| \cdot \|v\|_{H^1(\mathbb{T})}. \quad (86)$$

Set $v := (L(k) - \lambda)^{-1}u$ for $u \in L^2(\mathbb{T})$. Ellipticity of $L(k)$ yields $v \in H^2(\mathbb{T})$ and in particular, we obtain (86) for such v . Testing the equation $(L(k) - \lambda)v = u$ with the function v , we derive the standard energy estimate

$$\|Dv\|_{L^2(\mathbb{T})} \leq C_2(\|v\|_{L^2(\mathbb{T})} + \|u\|_{L^2(\mathbb{T})}). \quad (87)$$

Note that both C_1 and C_2 in (86) and (87) are independent of k and λ since we take k in the bounded set \mathcal{O} and consider $|\lambda|$ to be small enough.

Suppose that $|\lambda|$ is less than one-half of the length of the gap between the dispersion branches λ_j and λ_{j-1} . Due to functional calculus of the self-adjoint operator $L(k)$, we get

$$\|(L(k) - \lambda)^{-1}\|_{\text{op}} = \text{dist}(\lambda, \sigma(L(k)))^{-1} = \min\{(\lambda_j(k) - \lambda), (\lambda - \lambda_{j-1}(k))\}^{-1}.$$

Now let $\delta_1 = -\frac{1}{2} \max \lambda_{j-1}(k) > 0$ and $\delta_2 = \min_{k \in \mathcal{O}, |k-k_0| \geq C} \lambda_j(k)$. Then due to A3, $\delta_2 > 0$. Moreover,

$$\lambda - \lambda_{j-1}(k) \geq \lambda - \max_{k \in \mathcal{O}} \lambda_{j-1}(k) > \delta_1,$$

and

$$\lambda_j(k) - \lambda \geq \min_{k \in \mathcal{O}, |k-k_0| \geq C} \lambda_j(k) - \lambda > \delta_2.$$

Hence,

$$\|(L(k) - \lambda)^{-1}\|_{\text{op}} < \delta := \min\{\delta_1, \delta_2\}^{-1}. \quad (88)$$

In other words, $\|v\|_{L^2(\mathbb{T})} \leq \delta\|u\|_{L^2(\mathbb{T})}$. Applying this fact together with (86) and (87), we have

$$\begin{aligned} \|(L(k+z) - L(k))(L(k) - \lambda)^{-1}u\|_{L^2(\mathbb{T})} &\leq |z|C_1\|v\|_{H^1(\mathbb{T})} \\ &\leq |z|C_1C_2(\|v\|_{L^2(\mathbb{T})} + \|u\|_{L^2(\mathbb{T})}) \\ &\leq |z|C_1C_2(1 + \delta)\|u\|_{L^2(\mathbb{T})}. \end{aligned}$$

Now (85) is a consequence of the above estimate if we let

$$\tau \leq \min\left(\frac{1}{2C_1C_2(1 + \delta)}, 1\right). \quad \square$$

Proof of Lemma 5.3. From Lemma 2.6, the complex Bloch variety $\Sigma := B_L$ of the operator L is an analytic subset of codimension one in \mathbb{C}^{d+1} . By [19, 34], there exist an entire scalar function $h(k, \mu)$ and an entire operator-valued function $I(k, \mu)$ on \mathbb{C}^{d+1} such that

- (1) h vanishes only on Σ and has simple zeros on Σ , i.e., its normal derivative is not zero at all smooth parts of Σ ;
- (2) $n \mathbb{C}^{d+1} \setminus \Sigma, (L(k) - \mu)^{-1} = h(k, \mu)^{-1} I(k, \mu)$.

In particular, $(L_{t,s}(k) - \lambda)^{-1} = h(k + it\beta_s, \lambda)^{-1} I(k + it\beta_s, \lambda)$ for $k \in \mathbb{R}^d$ and $t \in [0, 1)$ by Proposition 4.1. Due to Assumption A and P2, if $k_0 + it\beta_s \in V$, the k -variable function $h(k, \lambda)^{-1}$ is equal (up to a non-vanishing analytic factor) to $(\lambda_j(k + it\beta_s) - \lambda)^{-1}$ on an open disc $D(k_0, 2\varepsilon) \subseteq V$ in \mathbb{C}^d for some $\varepsilon > 0$. Hence, we can write the sesquilinear form for such values of k as

$$(R_{t,s,\lambda} f, \varphi) = R_1 + R_2,$$

where

$$R_1 = (2\pi)^{-d} \int_{\mathbb{O} \cap D(k_0, \varepsilon)} \frac{(M(k, \lambda) \hat{f}(k), \hat{\varphi}(k))}{\lambda_j(k + it\beta_s) - \lambda} dk$$

and

$$R_2 = (2\pi)^{-d} \int_{\mathbb{O} \setminus D(k_0, \varepsilon)} (L(k + it\beta_s) - \lambda)^{-1} \hat{f}(k), \hat{\varphi}(k) dk.$$

Here $M(k, \lambda)$ is a $L^2(\mathbb{T})$ -valued analytic function on $D(k_0, \varepsilon)$ when $|\lambda|$ is small. Since f and φ have compact supports, their Floquet transforms $\hat{f}(k), \hat{\varphi}(k)$ are analytic with respect to k . To prove (22), we apply the Lebesgue Dominated Convergence Theorem. For R_1 , it suffices to show that the denominator in the integrand when $t \rightarrow 1^-$ is integrable over $D(k_0, \varepsilon)$ for $d \geq 2$. Indeed,

$$|\lambda_j(k + i\beta_s) - \lambda| \geq \delta \left| i \nabla E(\beta_s) \cdot (k - k_0) - \frac{1}{2} (k - k_0) \cdot \text{Hess}(E)(\beta_s)(k - k_0) \right|,$$

for some $\delta > 0$ if ε is chosen small enough so that in the Taylor expansion of λ_j at $k_0 + i\beta_s$, the remainder term $O(|k - k_0|^3)$ is dominated by the quadratic term $|k - k_0|^2$. Furthermore,

$$\begin{aligned} & \left| i \nabla E(\beta_s) \cdot (k - k_0) - \frac{1}{2} (k - k_0) \cdot \text{Hess}(E)(\beta_s)(k - k_0) \right|^2 \\ & \geq C(|(k - k_0, s)|^2 + |k - k_0|^4), \end{aligned}$$

for some constant $C > 0$ (independent of k). Now let $v := (k - k_0)$ and so the right hand side of the above estimate is just $|\langle v, s \rangle|^2 + |v|^4$ (up to a constant factor). One can apply Hölder's inequality to obtain

$$|\langle v, s \rangle|^2 + |v|^4 \geq |\langle v, s \rangle|^2 + |v'|^4 \geq C |\langle v, s \rangle|^{3/2} |v'|,$$

where $v = (v_1, v') \in \mathbb{R} \times \mathbb{R}^{d-1}$. Thus, we deduce

$$|\lambda_j(k + i\beta_s) - \lambda|^{-1} \leq C |\langle v, s \rangle|^{-3/4} |v'|^{-1/2}. \quad (89)$$

Since the function $|x|^{-n}$ is integrable near 0 in \mathbb{R}^d if and only if $n < d$, $|v'|^{-1/2}$ and $|\langle v, s \rangle|^{-3/4}$ are integrable near 0 in \mathbb{R}^{d-1} and \mathbb{R} respectively. Therefore, the function in the right hand side of (89) is integrable near 0.

The integrability of R_2 as $t \rightarrow 1^-$ follows from (85) in the proof of Proposition 4.1. Indeed,

$$\begin{aligned} & \| (L(k + it\beta_s) - \lambda)^{-1} \|_{\text{op}} \\ &= \| (1 - (L(k + it\beta_s) - L(k))(\lambda - L(k))^{-1})^{-1} (\lambda - L(k))^{-1} \|_{\text{op}} \\ &\leq \frac{\| (L(k) - \lambda)^{-1} \|_{\text{op}}}{1 - \| (L(k + it\beta_s) - L(k))(\lambda - L(k))^{-1} \|_{\text{op}}}. \end{aligned} \quad (90)$$

By decreasing $|\lambda|$, if necessary, and repeating the arguments when showing (85) and (88) we derive

$$1 - \| (L(k + it\beta_s) - L(k))(\lambda - L(k))^{-1} \|_{\text{op}} \geq 1/2, \quad \text{for all } k \in \mathcal{O} \setminus D(k_0, \varepsilon) \quad (91)$$

and

$$\sup_{k \in \mathcal{O} \setminus D(k_0, \varepsilon)} \| (L(k) - \lambda)^{-1} \|_{\text{op}} < \infty. \quad (92)$$

Thanks to (90), (91), (92), the Cauchy–Schwarz inequality, and Lemma 5.2, we have

$$\begin{aligned} & \sup_{t \in [0, 1]} |(L(k + it\beta_s) - \lambda)^{-1} \hat{f}(k), \hat{\varphi}(k)| \\ & \leq 2 \| (L(k) - \lambda)^{-1} \|_{\text{op}} \cdot \| \hat{f}(k) \|_{L^2(\mathbb{T})} \| \hat{\varphi}(k) \|_{L^2(\mathbb{T})} \\ & \lesssim \| \hat{f}(k) \|_{L^2(\mathbb{T})} \| \hat{\varphi}(k) \|_{L^2(\mathbb{T})}, \quad \text{for all } k \in \mathcal{O} \setminus D(k_0, \varepsilon), \end{aligned}$$

and

$$\int_{\mathcal{O} \setminus D(k_0, \varepsilon)} \| \hat{f}(k) \|_{L^2(\mathbb{T})} \| \hat{\varphi}(k) \|_{L^2(\mathbb{T})} dk \leq \| f \|_{L^2(\mathbb{R}^d)} \| \varphi \|_{L^2(\mathbb{R}^d)} < \infty.$$

This completes the proof of our lemma. \square

9.4. Regularity of eigenfunctions $\phi(z, x)$. In this subsection, we study the regularity of the eigenfunctions $\phi(z, x)$ of the operator $L(z)$ with corresponding eigenvalue $\lambda_j(z)$ (see P4). It is known that for each $z \in V$, the eigenfunction $\phi(z, x)$ is smooth in x . We will claim that these eigenfunctions are smooth in (z, x) when z is near to k_0 . The idea is that initially, $\phi(z, \cdot)$ is an analytic section of the Hilbert bundle $V \times H^2(\mathbb{T})$ and then by ellipticity, it is also an analytic section of the bundle $V \times H^m(\mathbb{T})$ for any $m > 0$ (for statements related to Fredholm morphisms between analytic Banach bundles, see e.g., [36]) and hence smoothness will follow.

For the sake of completeness, we provide the proof of the above claim by applying standard bootstrap arguments in the theory of elliptic differential equations.

Lemma 9.4. *The function $\partial_x^\alpha \phi(z, x)$ is jointly continuous on $V \times \mathbb{R}^d$ for any multi-index α .*

Proof. By periodicity, it suffices to restrict x to \mathbb{T} . Let $K := \bar{V}$. Due to P4, the function $z \mapsto \phi(z, \cdot)$ is a $H^2(\mathbb{T})$ -valued analytic on some neighborhood of K . Thus,

$$\sup_{z \in K} \|\phi(z, \cdot)\|_{H^2(\mathbb{T})} < \infty.$$

Then, we can apply bootstrap arguments for the equation

$$L(z)\phi(z, \cdot) = \lambda_j(z)\phi(z, \cdot)$$

to see that $M_m := \sup_{z \in K} \|\phi(z, \cdot)\|_{H^m(\mathbb{T})}$ is finite for any nonnegative integer m .

Now we consider z and z' in K . Let $\phi_{z,z'}(x) := \phi(z, x) - \phi(z', x)$. Then, $\phi_{z,z'}$ is a (classical) solution of the equation

$$L(z)\phi_{z,z'} = f_{z,z'},$$

where $f_{z,z'} := (\lambda_j(z)\phi(z, \cdot) - \lambda_j(z')\phi(z', \cdot)) + (L(z') - L(z))\phi(z', \cdot)$.

By induction, we will show that for any $m \geq 0$,

$$\|\phi_{z,z'}\|_{H^m(\mathbb{T})} \lesssim |z - z'|. \tag{93}$$

The case $m = 0$ is clear because P4 implies that $z \mapsto \|\phi(z, \cdot)\|_{L^2(\mathbb{T})}$ is Lipschitz continuous.

Next, we assume that (93) holds for m . As in (86),

$$\|(L(z) - L(z'))\phi(z', \cdot)\|_{H^m(\mathbb{T})} \lesssim |z - z'| \cdot \|\phi(z', \cdot)\|_{H^{m+1}(\mathbb{T})} \lesssim M_{m+1}|z - z'|. \quad (94)$$

Using triangle inequalities, (93), (94), and the analyticity of λ_j , we get

$$\begin{aligned} \|f_{z,z'}\|_{H^m(\mathbb{T})} &\lesssim \|\lambda_j(z)\phi(z, \cdot) - \lambda_j(z')\phi(z', \cdot)\|_{H^m(\mathbb{T})} \\ &\quad + \|(L(z) - L(z'))\phi(z', \cdot)\|_{H^m(\mathbb{T})} \\ &\lesssim |\lambda_j(z)| \cdot \|\phi_{z,z'}\|_{H^m(\mathbb{T})} + M_m |\lambda_j(z) - \lambda_j(z')| + M_{m+1}|z - z'| \\ &\lesssim |z - z'|. \end{aligned} \quad (95)$$

Notice that for any $m \geq 0$, the following standard energy estimate holds (see e.g., [9, 12, 24]):

$$\|\phi_{z,z'}\|_{H^{m+2}(\mathbb{T})} \lesssim \|f_{z,z'}\|_{H^m(\mathbb{T})} + \|\phi_{z,z'}\|_{L^2(\mathbb{T})}. \quad (96)$$

Combining (95) and (96), we deduce that $\|\phi_{z,z'}\|_{H^{m+2}(\mathbb{T})} \lesssim |z - z'|$. Hence, (93) holds for $m + 2$. This finishes our induction.

Applying the Sobolev embedding theorem, we get $\|\phi_{z,z'}\|_{C^m(\mathbb{T})} \lesssim |z - z'|$ for any $m \geq 0$. In other words, $\phi \in C(K, C^m(\mathbb{T}))$ for any m . Since $C(K \times \mathbb{T}) = C(K, C(\mathbb{T}))$, this completes the proof. \square

Notation 9.5. Consider a z -parameter family of linear partial differential operators $\{L(z)\}$ where $z \in \mathbb{R}^d$. Suppose $L(x, \xi, z)$ is the symbol of $L(z)$. Whenever it makes sense, the differential operator $\frac{\partial L(z)}{\partial z_l}$ is the one whose symbol is $\frac{\partial L}{\partial z_l}(x, \xi, z)$ for any $l \in \{1, 2, \dots, d\}$.

Proposition 9.6. Assume D is an open disc centered at k_0 in \mathbb{R}^d such that $D \pm i\beta_s \in V$ for any $s \in \mathbb{S}^{d-1}$. Then all eigenfunctions $\phi(k \pm i\beta_s, x)$ are smooth on a neighborhood of $\bar{D} \times \mathbb{R}^d$. Furthermore, all derivatives of $\phi(k \pm i\beta_s, x)$ are bounded on $\bar{D} \times \mathbb{R}^d$ uniformly in s , i.e., for any multi-indices α, β :

$$\sup_{(s,k,x) \in \mathbb{S}^{d-1} \times \bar{D} \times \mathbb{R}^d} |\partial_k^\alpha \partial_x^\beta \phi(k \pm i\beta_s, x)| < \infty.$$

Proof. Pick any open disc D' in \mathbb{R}^d so that $\bar{D} \pm i\beta_s \subset D' \pm i\beta_s \subseteq V$. We will prove that all eigenfunctions are smooth on the domain $D' \times \mathbb{R}^d$. Also, it is enough to consider the function $\phi(k + i\beta_s)$ since the other one is treated similarly.

First, we show that $\frac{\partial \phi}{\partial k_l}(k + i\beta_s, x)$ is continuous for any $1 \leq l \leq d$. By Lemma 9.4, the function $(k, x) \mapsto \phi(k + i\beta_s, x)$ is continuous on $D' \times \mathbb{T}$. We consider any two complex-valued test functions $\varphi \in C_c^\infty(D')$ and $\psi \in C^\infty(\mathbb{T})$. Testing the equation of the eigenfunction $\phi(k + i\beta_s, x)$ with ψ and $\frac{\partial \varphi}{\partial k_l}$, we derive

$$\int_{D'} \int_{\mathbb{T}} (L(k + i\beta_s) - \lambda_j(k + i\beta_s)) \phi(k + i\beta_s, x) \overline{\psi(x)} \frac{\partial \varphi}{\partial k_l}(k) dx dk = 0.$$

Observe that

$$L(k + i\beta_s)^* = L(k - i\beta_s)$$

and

$$\left(\frac{\partial L(k - i\beta_s)}{\partial k_l} \right)^* = \frac{\partial L(k + i\beta_s)}{\partial k_l}.$$

We integrate by parts to derive

$$\begin{aligned} 0 &= \int_{D'} ((L(k + i\beta_s) - \lambda_j(k + i\beta_s)) \phi(k + i\beta_s, x), \psi(x))_{L^2(\mathbb{T})} \frac{\partial \varphi}{\partial k_l}(k) dk \\ &= \int_{D'} (\phi(k + i\beta_s, x), (L(k - i\beta_s) - \overline{\lambda_j}(k + i\beta_s)) \psi(x))_{L^2(\mathbb{T})} \frac{\partial \varphi}{\partial k_l}(k) dk \\ &= \int_{D'} \left(-\frac{\partial \phi}{\partial k_l}(k + i\beta_s, x), (L(k - i\beta_s) - \overline{\lambda_j}(k + i\beta_s)) \psi(x) \right)_{L^2(\mathbb{T})} \varphi(k) dk \\ &\quad - \int_{D'} \left(\phi(k + i\beta_s, x), \frac{\partial L(k - i\beta_s)}{\partial k_l} \psi(x) - \frac{\partial \overline{\lambda_j}}{\partial k_l}(k + i\beta_s) \psi(x) \right)_{L^2(\mathbb{T})} \varphi(k) dk \\ &= \int_{D'} \left((-L(k + i\beta_s) + \lambda_j(k + i\beta_s)) \frac{\partial \phi}{\partial k_l}(k + i\beta_s, x), \psi(x) \right)_{L^2(\mathbb{T})} \varphi(k) dk \\ &\quad - \int_{D'} \left(\left(\frac{\partial L(k + i\beta_s)}{\partial k_l} - \frac{\partial \lambda_j}{\partial k_l}(k + i\beta_s) \right) \phi(k + i\beta_s, x), \psi(x) \right)_{L^2(\mathbb{T})} \varphi(k) dk. \end{aligned} \tag{97}$$

We introduce

$$\phi_l(k, x) := \frac{\partial \phi}{\partial k_l}(k + i\beta_s, x),$$

$$G(k) := -L(k + i\beta_s) + \lambda_j(k + i\beta_s),$$

$$H(k, x) := \left(\frac{\partial L(k + i\beta_s)}{\partial k_l} - \frac{\partial \lambda_j}{\partial k_l}(k + i\beta_s) \right) \phi(k + i\beta_s, x).$$

By invoking the previous lemma, the Lipschitz continuity of the $C^{m+2}(\mathbb{T})$ -valued function $\phi(k + i\beta_s, \cdot)$ implies that the mapping $k \mapsto H(k, \cdot)$ must be Lipschitz as a $C^m(\mathbb{T})$ -valued function on $\overline{D'}$ for any $m \geq 0$. On the other hand, the $H^2(\mathbb{T})$ -valued function $\phi_I(k, \cdot)$ is also Lipschitz on D' due to P4. Hence, both $(G(k)\phi_I(k, \cdot), \psi)_{L^2(\mathbb{T})}$ and $(H(k, \cdot), \psi)_{L^2(\mathbb{T})}$ are continuous on D' for any test function ψ . The continuity let us conclude from (97) that for every $k \in D'$, $\phi_I(k, \cdot)$ is a weak solution of the equation

$$G(k)\phi_I(k, x) = H(k, x). \quad (98)$$

We interpret (98) in the classical sense since all the coefficients of this equation are smooth. Consider any k_1, k_2 in D' and subtract the equation corresponding to k_1 from the one corresponding to k_2 to obtain the equation for the oscillation function $\phi_I(k_1, \cdot) - \phi_I(k_2, \cdot)$:

$$G(k_1)(\phi_I(k_1, x) - \phi_I(k_2, x)) = (G(k_2) - G(k_1))\phi_I(k_2, x) + (H(k_1, x) - H(k_2, x)).$$

Note that due to regularities of λ_j , H and the fact that the differential operator $G(k)$ depends analytically on k , we get

$$\|H(k_1, \cdot) - H(k_2, \cdot)\|_{H^m(\mathbb{T})} + \|(G(k_1) - G(k_2))\phi_I(k_2, \cdot)\|_{H^m(\mathbb{T})} = O(|k_1 - k_2|)$$

for all $m \in \mathbb{N}$. Combining this with the uniform boundedness in k of the supremum norms of all coefficients of the differential operator $G(k_1)$, we obtain

$$\|\phi_I(k_1, \cdot) - \phi_I(k_2, \cdot)\|_{H^m(\mathbb{T})} = O(|k_1 - k_2|),$$

by using energy estimates as in the proof of Lemma 9.4. An application of the Sobolev embedding theorem shows that $\partial_x^\beta \phi_I(k, x)$ is continuous on $D' \times \mathbb{T}$ for any multi-index β .

To deduce continuity of higher derivatives $\partial_x^\beta \partial_k^\alpha \phi(k + i\beta_s)$ ($|\alpha| > 1$, $|\beta| \geq 0$), we induct on $|\alpha|$ and repeat the arguments of the $|\alpha| = 1$ case.

Finally, the last statement of this proposition also follows since all of our estimates hold uniformly in s . \square

Observation 9.7. 1. Property P4 is crucial in order to bootstrap regularities of eigenfunctions $\phi(k \pm i\beta_s)$.

2. If one just requires $\phi(k \pm i\beta_s) \in C^m(\overline{D} \times \mathbb{R}^d)$ for certain $m > 0$ then the smoothness on coefficients of L could be relaxed significantly (see [12, 24]).

10. Concluding remarks

The condition that the potentials A , V are infinitely differentiable is an overkill. The Fredholm property of the corresponding Floquet operators is essential, which can be obtained under much weaker assumptions.

The main result of this article assumes the central symmetry (evenness) of the relevant branch of the dispersion curve $\lambda_j(k)$, which does not hold for instance for operators with periodic magnetic potentials [29, 10]. Note that the result of [23] at the spectral edge does not require such a symmetry. It seems that in the inside-the-gap situation one also should not need such a symmetry. However, the authors have not been able to do so, and thus were limited to the case of high symmetry points of the Brillouin zone.

In the case when λ is below the whole spectrum, the result of this paper implies the Theorem 1.1 in [25] for self-adjoint operators.

Acknowledgements. The work of the first two authors were partially supported by the NSF grant DMS-1517938. The third author is partially funded by NSF grant DMS-1405100. The authors express their gratitude to the NSF for the support. P. Kuchment would like to thank the Isaac Newton Institute for Mathematical Sciences for its hospitality during the programme Periodic and Ergodic Problems supported by EPSRC Grant Number EP/K032208/1, where work on this paper was undertaken.

As it was mentioned in Remark 2.12, N. Filonov has suggested a modification to our proof that leads to the better error estimate, currently formulated in the main theorem. The authors are extremely grateful to him for this contribution, as well as for other many comments that were very important for improvement of the text.

References

- [1] Sh. Agmon, *Lectures on elliptic boundary value problems*. Prepared for publication by B. F. Jones, Jr., with the assistance of G. W. Batten, Jr. Revised edition of the 1965 original. AMS Chelsea Publishing, Providence, R.I., 2010. [MR 2589244](#) [Zbl 1221.35002](#)
- [2] Sh. Agmon, On kernels, eigenvalues, and eigenfunctions of operators related to elliptic problems. *Comm. Pure Appl. Math.* **18** (1965), 627–663. [MR 0198287](#) [Zbl 0151.20203](#)
- [3] M. Babillot, Théorie du renouvellement pour des chaînes semi-markoviennes transientes. *Ann. Inst. H. Poincaré Probab. Statist.* **24** (1988), no. 4, 507–569. [MR 0978023](#) [Zbl 0681.60095](#)

- [4] J. M. Barbaroux, J. M. Combes, and P. D. Hislop, Localization near band edges for random Schrödinger operators. *Helv. Phys. Acta* **70** (1997), no. 1-2, 16–43. Papers honouring the 60th birthday of Klaus Hepp and of Walter Hunziker, Part II (Zürich, 1995). [MR 1441595](#) [Zbl 0866.35077](#)
- [5] M. Birman and T. Suslina, Threshold effects near the lower edge of the spectrum for periodic differential operators of mathematical physics. In A. A. Borichev and N. K. Nikolski (eds.), *Systems, approximation, singular integral operators, and related topics*. Papers from the 11th International Workshop on Operator Theory and Applications (IWOTA 2000) held at the Université Bordeaux I, Bordeaux, June 13–16, 2000, 71–107. [MR 1882692](#) [Zbl 1032.35145](#)
- [6] M. Birman and T. Suslina, Periodic second-order differential operators. Threshold properties and averaging. *Algebra i Analiz* **15** (2003), no. 5, 1–108. In Russian. English translation, *St. Petersburg Math. J.* **15** (2003), no. 5, 639–714. [MR 2068790](#)
- [7] J. M. Combes and L. Thomas, Asymptotic behaviour of eigenfunctions for multiparticle Schrödinger operators. *Comm. Math. Phys.* **34** (1973), 251–270. [MR 0391792](#) [Zbl 0271.35062](#)
- [8] M. S. P. Eastham, *The spectral theory of periodic differential equations*. Texts in Mathematics (Edinburgh). Scottish Academic Press, Edinburgh, and Hafner Press, New York, 1973. [MR 3075381](#) [Zbl 0287.34016](#)
- [9] L. C. Evans, *Partial differential equations*. Graduate Studies in Mathematics, 19. American Mathematical Society, Providence, R.I., 1998. [MR 1625845](#) [Zbl 0902.35002](#)
- [10] J. Feldman, H. Knörrer and E. Trubowitz, Asymmetric Fermi surfaces for magnetic Schrödinger operators. *Comm. Partial Differential Equations* **25** (2000), no. 1-2, 319–336. [MR 1737550](#) [Zbl 0943.35015](#)
- [11] M. Ghomi, Gauss map, topology, and convexity of hypersurfaces with nonvanishing curvature. *Topology* **41** (2002), no. 1, 107–117. [MR 1871243](#) [Zbl 0993.57009](#)
- [12] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*. Second edition. Grundlehren der Mathematischen Wissenschaften, 224. Springer-Verlag, Berlin etc., 1983. [MR 0737190](#) [Zbl 0562.35001](#)
- [13] V. Guillemin, Notes on elliptic operators. Lecture notes. MIT, Cambridge, MA, 2005. <http://math.mit.edu/~vwg/classnotes-spring05.pdf>
- [14] J. M. Harrison, P. Kuchment, A. Sobolev, and B. Winn, On occurrence of spectral edges for periodic operators inside the Brillouin zone. *J. Phys. A* **40** (2007), no. 27, 7597–7618. [MR 2369966](#) [Zbl 1141.82340](#)
- [15] L. Hörmander, *The analysis of linear partial differential operators*. I. Distribution theory and Fourier analysis. Grundlehren der Mathematischen Wissenschaften, 256. Springer-Verlag, Berlin etc., 1983. [MR 0717035](#) [Zbl 0521.35001](#)
- [16] W. Kirsch and B. Simon, Comparison theorems for the gap of Schrödinger operators. *J. Funct. Anal.* **75** (1987), no. 2, 396–410. [MR 0916759](#) [Zbl 0661.35062](#)

- [17] F. Klopp and J. Ralston, Endpoints of the spectrum of periodic operators are generically simple. *Methods Appl. Anal.* **7** (2000), no. 3, 459–463. [MR 1869296](#) [Zbl 1040.35050](#)
- [18] T. Kato, *Perturbation theory for linear operators*. Second edition. Grundlehren der Mathematischen Wissenschaften, 132. Springer-Verlag, Berlin etc., 1976. [MR 0407617](#) [Zbl 0342.47009](#)
- [19] P. Kuchment, *Floquet theory for partial differential equations*. Operator Theory: Advances and Applications, 60. Birkhäuser Verlag, Basel, 1993. [MR 1232660](#) [Zbl 0789.35002](#)
- [20] P. Kuchment, An overview of periodic elliptic operators. *Bull. Amer. Math. Soc. (N.S.)* **53** (2016), no. 3, 343–414. [MR 3501794](#) [Zbl 1346.35170](#)
- [21] P. Kuchment and Y. Pinchover, Integral representations and Liouville theorems for solutions of periodic elliptic equations. *J. Funct. Anal.* **181** (2001), no. 2, 402–446. [MR 1821702](#) [Zbl 0986.35028](#)
- [22] P. Kuchment and Y. Pinchover, Liouville theorems and spectral edge behavior on abelian coverings of compact manifolds. *Trans. Amer. Math. Soc.* **359** (2007), no. 12, 5777–5815. [MR 2336306](#) [Zbl 1127.58015](#)
- [23] P. Kuchment and A. Raich, Green’s function asymptotics near the internal edges of spectra of periodic elliptic operators. *Math. Nachr.* **285** (2012), no. 14–15, 1880–1894. [MR 2988010](#) [Zbl 1257.35080](#)
- [24] O. A. Ladyzhenskaya and N. N. Ural’tseva, *Linear and quasilinear elliptic equations*. Translated from the Russian by Scripta Technica. Translation editor, L. Ehrenpreis. Academic Press, New York and London, 1968. [MR 0244627](#) [Zbl 0164.13002](#)
- [25] M. Murata and T. Tsuchida, Asymptotics of Green functions and Martin boundaries for elliptic operators with periodic coefficients. *J. Differential Equations* **195** (2003), no. 1, 82–118. [MR 2019244](#) [Zbl 1036.31005](#)
- [26] M. Murata and T. Tsuchida, Asymptotics of Green functions and the limiting absorption principle for elliptic operators with periodic coefficients. *J. Math. Kyoto Univ.* **46** (2006), no. 4, 713–754. [MR 2320348](#) [Zbl 1142.35013](#)
- [27] M. Reed and B. Simon, *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press, New York and London, 1978. [MR 0493421](#) [Zbl 0401.47001](#)
- [28] M. Ruzhansky and V. Turunen, Pseudo-differential operators and symmetries. Background analysis and advanced topics. Pseudo-Differential Operators. Theory and Applications, 2. Birkhäuser Verlag, Basel, 2010. [MR 2567604](#) [Zbl 1193.35261](#)
- [29] R. G. Shterenberg, An example of a periodic magnetic Schrödinger operator with a degenerate lower edge of the spectrum. *Algebra i Analiz* **16** (2004), no. 2, 177–185. In Russian. English translation, *St. Petersburg Math. J.* **16** (2004), no. 2, 417–422. [MR 2068347](#) [Zbl 1075.35022](#)

- [30] M. A. Šubin, Spectral theory and the index of elliptic operators with almost-periodic coefficients. *Uspekhi Mat. Nauk* **34** (1979), no. 2(206), 95–135. In Russian. English translation, *Russian Math. Surveys* **34** (1979), no. 2, 109–158. [MR 0535710](#) [Zbl 0448.47032](#)
- [31] M. A. Šubin, *Pseudodifferential operators and spectral theory*. Translated from the 1978 Russian original by S. I. Andersson. Second edition. Springer-Verlag, Berlin, 2001. [MR 1852334](#) [Zbl 0980.35180](#)
- [32] A. Sobolev, Periodic operators: the method of gauge transform. Lectures at the I. Newton Institute, Cambridge, 2015. <https://www.newton.ac.uk/event/pepw01/timetable>
- [33] J. A. Thorpe, Elementary topics in differential geometry. Corrected reprint of the 1979 original. Undergraduate Texts in Mathematics. Springer-Verlag, Berlin, 1994. [MR 1329997](#) [Zbl 0404.53001](#)
- [34] C. H. Wilcox, Theory of Bloch waves. *J. Analyse Math.* **33** (1978), 146–167. [MR 0516045](#) [Zbl 0408.35067](#)
- [35] W. Woess, *Random walks on infinite graphs and groups*. Cambridge Tracts in Mathematics, 138. Cambridge University Press, Cambridge, 2000. [MR 1743100](#) [Zbl 0951.60002](#)
- [36] M. G. Zaïdenberg, S. G. Kreĭn, P. A. Kučment, and A. A. Pankov, Banach bundles and linear operators. *Uspehi Mat. Nauk* **30** (1975), no. 5(185), 101–157. In Russian. English translation, *Russian Math. Surveys* **30** (1975), no. 5, 115–175]. [MR 0415661](#) [Zbl 0317.47018](#) [Zbl 0335.47017](#) (transl.)

Received September 1, 2015; revised April 13, 2016

Minh Kha, Department of Mathematics, Texas A&M University, College Station, TX 77843-3368, USA

e-mail: kha@math.tamu.edu

Current address: Department of Mathematics, University of Arizona, Tucson, AZ 85721-0089, USA

e-mail: minhkha@math.arizona.edu

Peter Kuchment, Department of Mathematics, Texas A&M University, College Station, TX 77843-3368, USA

e-mail: kuchment@math.tamu.edu

Andrew Raich, Department of Mathematical Sciences, University of Arkansas, Fayetteville, AR 72701, USA

e-mail: araich@uark.edu