

A multidimensional Borg–Levinson theorem for magnetic Schrödinger operators with partial spectral data

Yavar Kian

Abstract. We consider the multidimensional Borg–Levinson theorem of determining both the magnetic field dA and the electric potential V , appearing in the Dirichlet realization of the magnetic Schrödinger operator $H = (-i\nabla + A)^2 + V$ on a bounded domain $\Omega \subset \mathbb{R}^n$, $n \geq 2$, from partial knowledge of the boundary spectral data of H . The full boundary spectral data are given by the set $\{(\lambda_k, \partial_\nu \varphi_k|_{\partial\Omega}) : k \geq 1\}$, where $\{\lambda_k : k \geq 1\}$ is the non-decreasing sequence of eigenvalues of H , $\{\varphi_k : k \geq 1\}$ an associated Hilbertian basis of eigenfunctions and ν is the unit outward normal vector to $\partial\Omega$. We prove that some asymptotic knowledge of $(\lambda_k, \partial_\nu \varphi_k|_{\partial\Omega})$ with respect to $k \geq 1$ determines uniquely the magnetic field dA and the electric potential V .

Keywords. Inverse spectral problem, Borg–Levinson theorem, magnetic Schrödinger operators.

Mathematics Subject Classification (2010). Primary: 35R30, 35J10; Secondary: 35P99.

Contents

1	Introduction	236
2	Notations and preliminary results	240
3	Representation formulas	245
4	Proof of the main result	261
	References	267

1. Introduction

1.1. Statement of the problem. We consider $\Omega \subset \mathbb{R}^n$, $n \geq 2$, a $\mathcal{C}^{1,1}$ bounded and connected domain such that $\mathbb{R}^n \setminus \Omega$ is also connected. We set $\Gamma = \partial\Omega$. Let $A \in W^{1,\infty}(\Omega, \mathbb{R}^n)$, $V \in L^\infty(\Omega, \mathbb{R})$ and consider the magnetic Schrödinger operator $H = (-i\nabla + A)^2 + V$ acting on $L^2(\Omega)$ with domain $D(H) = \{v \in H_0^1(\Omega): (-i\nabla + A)^2 v \in L^2(\Omega)\}$.

Let $A_j \in W^{1,\infty}(\Omega, \mathbb{R}^n)$, $V_j \in L^\infty(\Omega, \mathbb{R})$, $j = 1, 2$, and consider the magnetic Schrödinger operators $H_j = H$ for $A = A_j$ and $V = V_j$, $j = 1, 2$. We say that H_1 and H_2 are gauge equivalent if there exists $p \in W^{2,\infty}(\Omega, \mathbb{R}) \cap H_0^1(\Omega)$ such that $H_2 = e^{-ip} H_1 e^{ip}$.

It is well known that H is a selfadjoint operator. By the compactness of the embedding $H_0^1(\Omega) \hookrightarrow L^2(\Omega)$, the spectrum of H is purely discrete. From now on, we denote by \mathbb{N}^* the set of integer $\mathbb{N}^* = \{1, 2, \dots\}$. We note $\{\lambda_k: k \in \mathbb{N}^*\}$ the non-decreasing sequence of eigenvalues of H and $\{\varphi_k: k \in \mathbb{N}^*\}$ an associated Hilbertian basis of eigenfunctions. In the present paper we consider the Borg–Levinson inverse spectral problem of determining uniquely H , modulo gauge equivalence, from partial knowledge of the boundary spectral data $\{(\lambda_k, \partial_\nu \varphi_k|_\Gamma): k \in \mathbb{N}^*\}$ with ν the outward unit normal vector to Γ . Namely, we prove that some asymptotic knowledge of $(\lambda_k, \partial_\nu \varphi_k|_\Gamma)$ with respect to $k \in \mathbb{N}^*$ determines uniquely the operator H modulo gauge transformation.

1.2. Borg–Levinson inverse spectral problems . It is Ambartsumian who first investigated in 1929 the inverse spectral problem of determining the real potential V appearing in the Schrödinger operator $H = -\Delta + V$, acting in $L^2(\Omega)$, from partial spectral data of H . For $\Omega = (0, 1)$, he proved in [1] that $V = 0$ if the spectrum of the Neumann realization of H equals $\{k^2: k \in \mathbb{N}\}$. For the same operator, but endowed with homogeneous Dirichlet boundary conditions, Borg [6] and Levinson [19] established that the Dirichlet spectrum $\{\lambda_k: k \in \mathbb{N}^*\}$ does not uniquely determine V . They showed that additional spectral data, namely $\{\|\varphi_k\|_{L^2(0,1)}: k \in \mathbb{N}^*\}$, where $\{\varphi_k: k \in \mathbb{N}^*\}$ is an $L^2(0, 1)$ -orthogonal basis of eigenfunctions of H obeying the condition $\varphi_k'(0) = 1$, is needed. Gel'fand and Levitan proved in [12] that uniqueness is still valid upon substituting the terminal velocity $\varphi_k'(1)$ for $\|\varphi_k\|_{L^2(0,1)}$ in the one-dimensional Borg and Levinson theorem.

In 1988, Nachman, Sylvester, and Uhlmann [22] and Novikov [25] proposed a multidimensional formulation of the result of Borg and Levinson. Namely, they proved that the boundary spectral data $\{(\lambda_k, \partial_\nu \varphi_k|_{\partial\Omega}): k \in \mathbb{N}^*\}$, where ν denotes the outward unit normal vector to $\partial\Omega$ and (λ_k, φ_k) is the k^{th} eigenpair of $-\Delta + V$, determines uniquely the Dirichlet realization of the operator $-\Delta + V$. The initial

formulation of the multidimensional Borg–Levinson theorem by [22] and [25] has been improved in several ways by various authors. Isozaki [14] (see also [9]) extended the result of [22] when finitely many eigenpairs remain unknown, and, recently, Choulli and Stefanov [10] claimed uniqueness in the determination of V from the asymptotic behavior of $(\lambda_k, \partial_\nu \varphi_k|_\Gamma)$ with respect to k . Moreover, Canuto and Kavian [7, 8] considered the determination of the conductivity c , the electric potential V and the weight ρ from the boundary spectral data of the operator $\rho^{-1}(-\operatorname{div}(c\nabla\cdot) + V)$ acting on the weighted space $L^2_\rho(\Omega)$ endowed with either Dirichlet or Neumann boundary conditions. Namely, [7, 8] proved that the boundary spectral data of $\rho^{-1}(-\operatorname{div}(c\nabla\cdot) + V)$ determines uniquely two of the three coefficients c , V and ρ . The case of magnetic Schrödinger operator has been treated by [27] who determined both the magnetic field dA and the electric potential V of the operator $H = (-i\nabla + A)^2 + V$. Here the 2-form dA of a vector valued function $A = (a_1, \dots, a_n)$ is defined by

$$dA = \sum_{i < j} (\partial_{x_j} a_i - \partial_{x_i} a_j) dx_j \wedge dx_i.$$

All the above mentioned results were obtained with Ω bounded and operators of purely discrete spectral type. In some recent work [16] examined a Borg–Levinson inverse problem stated in an infinite cylindrical waveguide for Schrödinger operators with purely absolutely continuous spectral type. More precisely, [16] proved that a real potential V which is 2π -periodic along the axis of the waveguide is uniquely determined by some asymptotic knowledge of the boundary Floquet spectral data of the Schrödinger operator $-\Delta + V$ with Dirichlet boundary conditions.

Finally, let us mention for the sake of completeness that the stability issue in the context of Borg–Levinson inverse problems was examined in [4, 5, 9, 10, 16] and that [2, 5, 15] established related results on Riemannian manifolds. We also precise that [21, 28, 29] have proved stability estimates in the recovery of coefficients from the hyperbolic Dirichlet–to–Neumann map which is equivalent to the determination of general Schrödinger operators from boundary spectral data.

1.3. Main result. Let $A_j \in W^{1,\infty}(\Omega, \mathbb{R}^n)$, $V_j \in L^\infty(\Omega, \mathbb{R})$ and consider the magnetic Schrödinger operators $H_j = H$ for $A = A_j$ and $V = V_j$, $j = 1, 2$. Further we note $(\lambda_{j,k}, \varphi_{j,k})$, $k \geq 1$, the k^{th} eigenpair of H_j , for $j = 1, 2$. Our main result can be stated as follows.

Theorem 1.1. *We fix Ω_1 an arbitrary open neighborhood of Γ in Ω ($\Gamma \subset \overline{\Omega_1}$ and $\Omega_1 \subsetneq \Omega$). For $j = 1, 2$, let $V_j \in L^\infty(\Omega, \mathbb{R})$ and let $A_j \in \mathcal{C}^1(\overline{\Omega}, \mathbb{R}^n)$ fulfill*

$$A_1(x) = A_2(x), \quad x \in \Omega_1. \tag{1.1}$$

Assume that the conditions

$$\lim_{k \rightarrow +\infty} |\lambda_{1,k} - \lambda_{2,k}| = 0, \quad \sum_{k=1}^{+\infty} \|\partial_\nu \varphi_{1,k} - \partial_\nu \varphi_{2,k}\|_{L^2(\Gamma)}^2 < \infty \tag{1.2}$$

hold simultaneously. Then, we have $dA_1 = dA_2$ and $V_1 = V_2$.

Note that condition (1.1) corresponds to the knowledge of the magnetic potential on a neighborhood of the boundary.

Let us observe that, as mentioned by [10, 16], Theorem 1.1 can be considered as a uniqueness theorem under the assumption that the spectral data are asymptotically “very close.” Conditions (1.2) are similar to the one considered by [16] and they are weaker than the requirement that

$$|\lambda_{1,k} - \lambda_{2,k}| \leq Ck^{-\alpha}, \quad \|\partial_\nu \varphi_{1,k} - \partial_\nu \varphi_{2,k}\|_{L^2(\Gamma)} \leq Ck^{-\beta}$$

for some $\alpha > 1$ and $\beta > 1 - \frac{1}{2n}$, considered in [10, Theorem 2.1]. Note also that conditions (1.2) are weaker than the knowledge of the boundary spectral data with a finite number of data missing considered by [14].

Let us remark that there is an obstruction to uniqueness given by the gauge invariance of boundary spectral data for magnetic Schrödinger operators. More precisely, let $p \in \mathcal{C}_0^\infty(\Omega \setminus \overline{\Omega_1}) \setminus \{0\}$ and assume that $A_1 = \nabla p + A_2 \neq A_2$, $V_1 = V_2$. Then, we have $H_1 = e^{-ip} H_2 e^{ip}$ and one can check that we can choose the spectral data of H_1 and H_2 in such a way that the conditions

$$\partial_\nu \varphi_{1,k}|_\Gamma = \partial_\nu \varphi_{2,k}|_\Gamma, \quad \lambda_{1,k} = \lambda_{2,k}, \quad k \in \mathbb{N}^*$$

are fulfilled. Therefore, conditions (1.1) and (1.2) are fulfilled but $H_1 \neq H_2$. Nevertheless, assuming (1.1) fulfilled, the conditions $dA_1 = dA_2$ and $V_1 = V_2$ imply that H_1 and H_2 are gauge equivalent. Therefore, Theorem 1.1 is equivalent to the unique determination of magnetic Schrödinger operators modulo gauge transformation from the asymptotic knowledge of the boundary spectral data given by conditions (1.2).

The problem under examination in this text is a Borg–Levinson inverse problem for the magnetic Schrödinger operator $H = (-i\nabla + A)^2 + V$. To our best knowledge, there are only two multi-dimensional Borg–Levinson uniqueness

result for magnetic Schrödinger operators available in the mathematical literature, [15, Theorem B] and [27, Theorem 3.2] (we refer also to [24] for related inverse scattering results). In [15], the authors considered general magnetic Schrödinger operators with smooth coefficients on a smooth connected Riemannian manifold and they proved unique determination of these operators modulo gauge invariance from the knowledge of the boundary spectral data with a missing finite number of data. In [27], Serov treated this problem on a bounded domain of \mathbb{R}^n , and he proved that, for $A \in W^{1,\infty}(\Omega, \mathbb{R}^n)$ and $V \in L^\infty(\Omega, \mathbb{R})$, the full boundary spectral data $\{(\lambda_k, \partial_\nu \varphi_k|_\Gamma) : k \in \mathbb{N}^*\}$ determines uniquely dA and V . In contrast to [15, 27], in the present paper we prove that the asymptotic knowledge of the boundary spectral data, given by the conditions (1.2), is sufficient for the unique determination of dA and V . To our best knowledge, conditions (1.2) are the weakest conditions on boundary spectral data that guaranty uniqueness of magnetic Schrödinger operators modulo gauge transformation. Moreover, our uniqueness result is stated with conditions similar to [16, Theorem 1.4], which seems to be the most precise Borg–Levinson uniqueness result so far for Schrödinger operators without magnetic potential ($A = 0$).

An important ingredient in our analysis is a suitable representation that allows to express the magnetic potential A and the electric potential V in terms of Dirichlet–to–Neumann map associated to the equations $(-i\nabla + A)^2 u + Vu - \lambda u = 0$ for some $\lambda \in \mathbb{C}$. In [14] Isozaki applied a similar approach to the Schrödinger operator $-\Delta + V$ with Dirichlet boundary condition¹ and [10, 16] applied the representation formulas of [14]. Inspired by the construction of complex geometric optics solutions of [3, 11, 17, 18, 23, 26, 30] we prove that the approach of [10, 14, 16] can be extended to magnetic Schrödinger operators. More precisely, we derive two representation formulas that allow to recover both the magnetic field and the electric potential of magnetic Schrödinger operators which means recovery of both coefficients of order one and zero in contrast to [10, 14, 16] where only determination of coefficients of order zero is considered. This paper is the first where the extension of the approach developed by [14] to more general coefficients than coefficients of order zero is considered. Note also that our approach make it possible to prove this extension without imposing important assumptions of regularity of the admissible coefficients.

We believe that the approach developed in the present paper can be used for results of stability in the determination of the magnetic field dA and the electric potential V similar to [16, Theorem 1.3]. Indeed, following the strategy set in this paper we expect a stability estimate associated to the the determination of the

¹ This argument was inspired by the Born approximation method of the scattering theory.

magnetic field dA . The main issue comes from the stability in the determination of the electric potential V . Nevertheless, we believe that this problem can be solved by adapting suitably the argument developed in [32] related to the inversion of the d operator on differential forms restricted to the right subspaces.

1.4. Outline. This paper is organized as follows. In Section 2 we consider some useful preliminary results concerning solutions of equations of the form $(-i\nabla + A)^2u + Vu - \lambda u = 0$ for some $\lambda \in \mathbb{C} \setminus \sigma(H)$. In Section 3 we introduce two representation formulas making the connection between the Dirichlet-to-Neumann map associated with the previous equations and the couple (A, V) of magnetic and electric potential. Finally, in Section 4 we combine all these results and we prove Theorem 1.1.

2. Notations and preliminary results

In this section we introduce some notations and we give some properties of solution of the equation $(-i\nabla + A)^2u + Vu - \lambda u = 0$. We denote by $\langle f, \psi \rangle$ the duality between $\psi \in H^{1/2}(\Gamma)$ and f belonging to the dual $H^{-1/2}(\Gamma)$ of $H^{1/2}(\Gamma)$. However, when in $\langle f, \psi \rangle$ both f and ψ belong to $L^2(\Gamma)$, to make things simpler $\langle \cdot, \cdot \rangle$ can be interpreted as the scalar product of $L^2(\Gamma)$, namely

$$\langle f, \psi \rangle = \int_{\Gamma} f(x) \overline{\psi(x)} d\sigma(x).$$

We introduce the operator H defined as

$$Hu := (-i\nabla + A)^2u + Vu, \quad (2.1)$$

for $u \in D(H) := \{\psi \in H_0^1(\Omega) : (-i\nabla + A)^2\psi \in L^2(\Omega)\}$. Recall that H is associated to the quadratic form b given by

$$b(u, v) = \int_{\Omega} (-i\nabla + A)u(x) \cdot \overline{(-i\nabla + A)v(x)} dx + \int_{\Omega} V(x)u(x)\overline{v(x)} dx,$$

for $u, v \in H_0^1(\Omega)$. Moreover, the spectrum of H is discrete and composed of the non-decreasing sequence of eigenvalues denoted by $\sigma(H) = \{\lambda_k : k \geq 1\}$. If we write $V = V^+ - V^-$, with $V^\pm := \max(0, \pm V)$, we have that the spectrum $\sigma(H)$ of H is contained into $[-\|V^-\|_{L^\infty(\Omega)}, +\infty)$. According to [13, Theorem 2.2.2.3], we can show that $D(H)$ embedded continuously into $H^2(\Omega)$. Therefore the eigenfunctions $(\varphi_k)_{k \geq 1}$ of H , that form an Hilbertian basis, are lying in $H^2(\Omega)$ and we have $\partial_\nu \varphi_k|_{\Gamma} \in H^{1/2}(\Gamma)$.

From now on, we fix $f \in H^{1/2}(\Gamma)$ and $\lambda \in \mathbb{C} \setminus \sigma(H)$ and we consider the problem

$$\begin{cases} (-i\nabla + A)^2 u + Vu - \lambda u = 0, & \text{in } \Omega, \\ u(x) = f(x), & x \in \Gamma. \end{cases} \tag{2.2}$$

We start with two results related to the asymptotic behavior of solutions of (2.2) as $\lambda \rightarrow -\infty$.

Lemma 2.1. *For any $f \in H^{1/2}(\Gamma)$ and $\lambda \in \mathbb{C} \setminus \sigma(H)$, there exists a unique solution $u \in H^1(\Omega)$ to (2.2) which can be written as*

$$u_\lambda := u = \sum_{k \geq 1} \frac{\alpha_k}{\lambda - \lambda_k} \varphi_k, \tag{2.3}$$

where for convenience we set

$$h_k := \partial_\nu \varphi_k|_\Gamma, \quad \text{and} \quad \alpha_k := \langle f, h_k \rangle. \tag{2.4}$$

Moreover, we have

$$\|u_\lambda\|_{L^2(\Omega)}^2 = \sum_{k \geq 1} \frac{|\alpha_k|^2}{|\lambda - \lambda_k|^2} \rightarrow 0 \quad \text{as } \lambda \rightarrow -\infty.$$

Proof. Since $\lambda \notin \sigma(H)$, one can easily check that (2.2) admits a unique solution $u_\lambda \in H^1(\Omega)$. Moreover, u_λ can be written in terms of the eigenvalues and eigenfunctions λ_k, φ_k . Indeed, $u_\lambda \in L^2(\Omega)$ can be expressed in the Hilbert basis $(\varphi_k)_{k \geq 1}$ as

$$u_\lambda = \sum_{k \geq 1} (u_\lambda, \varphi_k) \varphi_k$$

with (\cdot, \cdot) the scalar product with respect to $L^2(\Omega)$. Since $u_\lambda \in H^1(\Omega)$ and $\Delta u_\lambda = -2iA \cdot \nabla u_\lambda + (-i \operatorname{div}(A) + |A|^2 + V)u_\lambda \in L^2(\Omega)$, we have $\nabla u_\lambda \in H_{\operatorname{div}}(\Omega) = \{v \in L^2(\Omega; \mathbb{C}^n) : \operatorname{div}(v) \in L^2(\Omega)\}$. Thus, taking the scalar product of the first equation in (2.2) with φ_k and applying the Green formula we obtain

$$\langle f, h_k \rangle = (\lambda - \lambda_k) (u, \varphi_k),$$

which yields the expression given by (2.3). The fact that $\|u_\lambda\| \rightarrow 0$ as $\lambda \rightarrow -\infty$ is a consequence of the fact that we may fix $c_0 > \|V\|_{L^\infty(\Omega)}$ large enough so that if λ is real and such that $\lambda \leq -c_0$, we have $|\lambda - \lambda_k|^2 \geq |c_0 - \lambda_k|^2$ for all $k \geq 1$, and thus

$$\frac{|\alpha_k|^2}{|\lambda - \lambda_k|^2} \leq \frac{|\alpha_k|^2}{|c_0 - \lambda_k|^2},$$

so that we may apply Lebesgue’s dominated convergence as $\lambda \rightarrow -\infty$. □

Lemma 2.2. For all $\lambda < -\|V\|_{L^\infty(\Omega)} - 6\|A\|_{L^\infty(\Omega, \mathbb{R}^n)}^2$, the solution u_λ of (2.2) satisfies

$$\|\nabla u_\lambda\|_{L^2(\Omega \setminus \Omega_1)} \leq C \|u_\lambda\|_{L^2(\Omega_1)} \quad (2.5)$$

with C depending only on Ω and Ω_1 .

Proof. Let us denote by $\chi \in \mathcal{C}_0^\infty(\Omega, \mathbb{R})$ a function satisfying $\chi = 1$ on $\Omega \setminus \Omega_1$. Then, since $\nabla u_\lambda \in H_{\text{div}}(\Omega)$, multiplying (2.2) by $\chi^2 \overline{u_\lambda}$ and applying the Green formula we obtain

$$\begin{aligned} 0 &= \int_{\Omega} (-i\nabla + A)^2 u_\lambda \chi^2 \overline{u_\lambda} dx + \int_{\Omega} (V - \lambda) \chi^2 |u_\lambda|^2 dx \\ &= \int_{\Omega} |\chi \nabla u_\lambda|^2 dx \\ &\quad + 2 \int_{\Omega} (\chi \nabla u_\lambda) \cdot \overline{\nabla \chi u_\lambda} dx \\ &\quad + \int_{\Omega} [2i(u_\lambda \chi) A \cdot \overline{u_\lambda \nabla \chi} + i \chi u_\lambda A \cdot \overline{\chi \nabla u_\lambda} + \chi \nabla u_\lambda \cdot \overline{i A \chi u_\lambda}] dx \\ &\quad + \int_{\Omega} (|A|^2 + V - \lambda) \chi^2 |u_\lambda|^2 dx. \end{aligned} \quad (2.6)$$

Applying the Cauchy–Schwarz inequality we find

$$\begin{aligned} &\|\chi \nabla u_\lambda\|_{L^2(\Omega)}^2 + (-\|A\|_{L^\infty(\Omega, \mathbb{R}^n)}^2 - \|V\|_{L^\infty(\Omega)} - \lambda) \|u_\lambda\|_{L^2(\Omega)}^2 \\ &\leq 2\|u_\lambda \nabla \chi\|_{L^2(\Omega)} \|\chi \nabla u_\lambda\|_{L^2(\Omega)} \\ &\quad + 2\|A\|_{L^\infty(\Omega)} \|\chi u_\lambda\|_{L^2(\Omega)} \|u_\lambda \nabla \chi\|_{L^2(\Omega)} \\ &\quad + 2\|A\|_{L^\infty(\Omega)} \|\chi u_\lambda\|_{L^2(\Omega)} \|\chi \nabla u_\lambda\|_{L^2(\Omega)} \\ &\leq 4\|u_\lambda \nabla \chi\|_{L^2(\Omega)}^2 \\ &\quad + \frac{\|\chi \nabla u_\lambda\|_{L^2(\Omega)}^2}{4} \\ &\quad + \|A\|_{L^\infty(\Omega, \mathbb{R}^n)}^2 \|\chi u_\lambda\|_{L^2(\Omega)}^2 \\ &\quad + \|u_\lambda \nabla \chi\|_{L^2(\Omega)}^2 \\ &\quad + 4\|A\|_{L^\infty(\Omega, \mathbb{R}^n)}^2 \|\chi u_\lambda\|_{L^2(\Omega)}^2 \\ &\quad + \frac{\|\chi \nabla u_\lambda\|_{L^2(\Omega)}^2}{4}. \end{aligned}$$

From this estimate, we deduce

$$\frac{\|\chi \nabla u_\lambda\|_{L^2(\Omega)}^2}{2} + (-\|V\|_{L^\infty(\Omega)} - 6\|A\|_{L^\infty(\Omega, \mathbb{R}^n)}^2 - \lambda) \|\chi u_\lambda\|_{L^2(\Omega)}^2 \leq 5\|u_\lambda \nabla \chi\|_{L^2(\Omega)}^2.$$

Using the fact that $\lambda < -\|V\|_{L^\infty(\Omega)} - 6\|A\|_{L^\infty(\Omega, \mathbb{R}^n)}^2$, we obtain

$$\begin{aligned} \|\nabla u_\lambda\|_{L^2(\Omega \setminus \Omega_1)}^2 &\leq \|\chi \nabla u_\lambda\|_{L^2(\Omega)}^2 \\ &\leq 10\|u_\lambda \nabla \chi\|_{L^2(\Omega)}^2 \\ &\leq 10\|\nabla \chi\|_{L^\infty(\Omega)}^2 \|u_\lambda\|_{L^2(\Omega_1)}^2 \\ &\leq C \|u_\lambda\|_{L^2(\Omega_1)}^2. \end{aligned}$$

From this estimate we deduce (2.5). □

It is clear that the series (2.3) giving u_λ in terms of α_k , λ_k and φ_k , converges only in $L^2(\Omega)$ and thus we cannot deduce an expression of the normal derivative $\partial_\nu u_\lambda$ in terms of α_k , λ_k and h_k . To avoid this difficulty, in a similar way to [16], we have the following lemma:

Lemma 2.3. *Let $f \in H^{1/2}(\Gamma)$ be fixed and for $\lambda, \mu \in \mathbb{C} \setminus \sigma(H)$ let u_λ and u_μ be the solutions given by Lemma 2.1. If we set $v := v_{\lambda, \mu} := u_\lambda - u_\mu$, then*

$$\partial_\nu v = \sum_{k \geq 1} \frac{(\mu - \lambda)\alpha_k}{(\lambda - \lambda_k)(\mu - \lambda_k)} h_k, \tag{2.7}$$

the convergence taking place in $H^{1/2}(\Gamma)$.

Proof. Let $v_{\lambda, \mu} := u_\lambda - u_\mu$; One verifies that $v_{\lambda, \mu}$ solves

$$\begin{cases} (-i\nabla + A)^2 v_{\lambda, \mu} + V v_{\lambda, \mu} - \lambda v_{\lambda, \mu} = (\lambda - \mu)u_\mu, & \text{in } \Omega, \\ v_{\lambda, \mu}(x) = 0, & x \in \Gamma. \end{cases} \tag{2.8}$$

Since $(u_\mu|_{\varphi_k}) = \alpha_k/(\mu - \lambda_k)$, it follows that

$$v_{\lambda, \mu} = \sum_{k \geq 1} \frac{(\lambda - \mu)\alpha_k}{(\lambda_k - \lambda)(\mu - \lambda_k)} \varphi_k,$$

the convergence taking place in $D(H)$. Since the operator $v \mapsto \partial_\nu v$ is continuous from $D(H)$ into $H^{1/2}(\Gamma)$, the result of the lemma follows. □

The next lemma states essentially that if for $j = 1$ or $j = 2$ we have two magnetic potentials A_j , two electric potentials V_j and $u_j := u_{j,\mu}$ solutions of

$$\begin{cases} (-i\nabla + A_j)^2 u_j + V_j u_j - \mu u_j = 0, & \text{in } \Omega, \\ u_j(x) = f(x), & x \in \Gamma, \end{cases} \quad (2.9)$$

then $u_{1,\mu}$ and $u_{2,\mu}$ are close as $\mu \rightarrow -\infty$: in some sense the influence of the potentials A_j and V_j are dimmed when $\mu \rightarrow -\infty$. More precisely we have:

Lemma 2.4. *Let $V_j \in L^\infty(\Omega, \mathbb{R})$ and $A_j \in W^{1,\infty}(\Omega, \mathbb{R}^n)$ be given for $j = 1$ or $j = 2$, and denote by H_j the corresponding operator defined by (2.1). We assume that condition (1.1) is fulfilled. For $f \in H^{1/2}(\Gamma)$ and $\mu \in (-\infty, \mu_*) \subset \mathbb{C} \setminus \sigma(H)$, let $u_{j,\mu} := u_j$ be the solution of (2.9). Then $z_\mu := u_{1,\mu} - u_{2,\mu} \in H^2(\Omega)$ converge to 0 in $H^2(\Omega)$ as $\mu \rightarrow -\infty$. In particular $\partial_\nu z_\mu \rightarrow 0$ in $L^2(\Gamma)$ as $\mu \rightarrow -\infty$.*

Proof. Since the trace map $v \mapsto \partial_\nu v$ is continuous from $H^2(\Omega)$ to $L^2(\Gamma)$, it is enough to show that $z_\mu \in H^2(\Omega)$ and $\|z_\mu\|_{H^2(\Omega)} \rightarrow 0$ when $\mu \rightarrow -\infty$. We fix $\mu < \mu_*$ with $\mu_* < -\|V\|_{L^\infty(\Omega)} - 6\|A\|_{L^\infty(\Omega, \mathbb{R}^n)}^2$ less than the constants given by Lemma 2.1 for $A = A_j$, $V = V_j$, $j = 1, 2$. Without lost of generality we assume that $H_j - \mu_*$ is positive, $j = 1, 2$. One verifies that z_μ solves the equation

$$\begin{cases} (-i\nabla + A_1)^2 z_\mu + V_1 z_\mu - \mu z_\mu \\ = -2i(A_2 - A_1) \cdot \nabla u_{2,\mu} + (p_2 - p_1)u_{2,\mu}, & \text{in } \Omega, \\ z_\mu(x) = 0, & x \in \Gamma \end{cases} \quad (2.10)$$

with

$$p_j = -i \operatorname{div}(A_j) + |A_j|^2 + V_j, \quad j = 1, 2.$$

That is, denoting by $R_{1,\mu} = (H_1 - \mu I)^{-1}$ the resolvent of the operator $H_1 := (-i\nabla + A_1)^2 + V_1$, we have

$$z_\mu = R_{1,\mu}(-2i(A_2 - A_1) \cdot \nabla u_{2,\mu} + (p_2 - p_1)u_{2,\mu}) = \sum_{k=1}^{+\infty} \frac{(w_\mu, \varphi_{1,k})}{(\lambda_{1,k} - \mu)} \varphi_{1,k}$$

with $w_\mu = -2i(A_2 - A_1) \cdot \nabla u_{2,\mu} + (p_2 - p_1)u_{2,\mu}$ and $(\lambda_{1,k})_{k \geq 1}$, $(\varphi_{1,k})_{k \geq 1}$ respectively the eigenvalues of H_1 and an Hilbertian basis of eigenfunctions associated to these eigenvalues. Since $w_\mu \in L^2(\Omega)$, z_μ is lying in $D(H_1)$ and by the same way in $H^2(\Omega)$. It remains to show that $\|z_\mu\|_{H^2(\Omega)} \rightarrow 0$ when $\mu \rightarrow -\infty$.

Since $D(H_1)$ embedded continuously into $H^2(\Omega)$ there exists a generic constant C depending on A_1, V_1 and Ω such that

$$\|z_\mu\|_{H^2(\Omega)}^2 \leq C \sum_{k=1}^{\infty} |\lambda_{1,k} - \mu_*|^2 |(z_\mu, \varphi_{1,k})|^2 \leq C \|w_\mu\|_{L^2(\Omega)}^2.$$

On the other hand, condition (1.1) implies

$$\|w_\mu\|_{L^2(\Omega)} \leq C (\|\nabla u_{2,\mu}\|_{L^2(\Omega \setminus \Omega_1)} + \|u_{2,\mu}\|_{L^2(\Omega)})$$

with C independent of μ . Then, according to Lemma 2.1 and (2.5), we obtain

$$\limsup_{\mu \rightarrow -\infty} \|w_\mu\|_{L^2(\Omega)} \leq C \limsup_{\mu \rightarrow -\infty} \|u_{2,\mu}\|_{L^2(\Omega)} = 0.$$

Thus, we have

$$\limsup_{\mu \rightarrow -\infty} \|\partial_\nu z_\mu\|_{L^2(\Gamma)} \leq C \limsup_{\mu \rightarrow -\infty} \|z_\mu\|_{H^2(\Omega)} \leq C \limsup_{\mu \rightarrow -\infty} \|w_\mu\|_{L^2(\Omega)} = 0.$$

This completes the proof. □

Armed with these results, we will prove Theorem 1.1 by using some asymptotic properties of solutions of (2.2) with respect to λ . For this purpose, like in [10, 14, 16] we use representation formulas that will allow us to make a connection between the boundary spectral data and the potentials A and V .

3. Representation formulas

From now on, for all $x = (x_1, \dots, x_n) \in \mathbb{C}^n$ and $y = (y_1, \dots, y_n) \in \mathbb{C}^n$, we denote by $x \cdot y$ the quantity

$$x \cdot y = \sum_{k=1}^n x_k y_k$$

and for all $x \in \mathbb{R}^n$ we denote by x^\perp the subspace of \mathbb{R}^n defined by $\{y \in \mathbb{R}^n : y \cdot x = 0\}$. Moreover, we set $A_j \in C^1(\bar{\Omega}, \mathbb{R}^n)$, $V_j \in L^\infty(\Omega, \mathbb{R})$, $j = 1, 2$, and we assume that condition (1.1) is fulfilled. For $j = 1, 2$ and $\lambda \in \mathbb{C} \setminus \mathbb{R}$, we associate to the problem

$$\begin{cases} (-i\nabla + A_j)^2 u_j + V_j u_j - \lambda u_j = 0, & \text{in } \Omega, \\ u_j(x) = f(x), & x \in \Gamma, \end{cases} \tag{3.1}$$

the Dirichlet–to–Neumann map

$$\Lambda_{j,\lambda}: H^{\frac{1}{2}}(\partial\Omega) \ni f \longmapsto (\partial_\nu + iA_j \cdot \nu)u_{j,\lambda}|_\Gamma,$$

where $u_{j,\lambda}$ solves (3.1). The goal of this section is to apply the Dirichlet–to–Neumann maps $\Lambda_{j,\lambda}$ to some suitable ansatz associated with (3.1) in order to get two representation formulas involving the magnetic potentials A_j and the electric potentials V_j , $j = 1, 2$. A similar approach was developed by [14] and [10, 16] used the representation of [14]. The idea is to establish the link between the electric and magnetic potentials and the boundary spectral data by mean of an expression involving the Dirichlet–to–Neumann maps $\Lambda_{1,\lambda}$, $\Lambda_{2,\lambda}$. We start with two general representation formulas, stated in the next subsection, where some properties of the ansatz will not be completely specified. This will allow us to clarify the main goal of these formulas. Then, in Section 3.2 we will introduce the remaining properties of our ansatz and establish some asymptotic properties from our representations which will be one of the main points of our analysis.

3.1. General representation formulas. In this subsection we introduce the first formulation of two representation formulas involving respectively the Dirichlet–to–Neumann maps $\Lambda_{1,\lambda}$, $\Lambda_{2,\lambda}$ and some ansatz associated with problem (3.1). In [14], Isozaki considered such formulas for Schrödinger operators $-\Delta + V$ with an electric potential V , in other words for Schrödinger operators with a variable coefficient of order zero. In our case we need to extend this strategy to Schrödinger operators with both magnetic and electric potentials, which means an extension to Schrödinger operators with variable coefficients of order zero and one. In addition, we need to consider ansatz that allow to recover both the magnetic field and the electric potential. Therefore, we fix $\eta_1, \eta_2 \in \mathbb{S}^{n-1} = \{y \in \mathbb{R}^n, |y| = 1\}$ and we consider some ansatz, associated with (3.1), of the form

$$\Phi_{1,\lambda}(x) = e^{i\sqrt{\lambda}\eta_1 \cdot x} g_1(x), \quad \Phi_{2,\lambda}(x) = e^{-i\sqrt{\lambda}\eta_2 \cdot x} g_2(x), \quad x \in \Omega, \quad j = 1, 2. \quad (3.2)$$

For $\lambda > 0$, these ansatzs are the principal term of the standard geometric optics solutions. In contrast to the complex geometric optics solutions taking the form $e^{\xi_j \cdot x} h_j(x)$, $j = 1, 2$, where the entire vector $\xi_j \in \mathbb{C}^n$ is complex valued, here we consider these ansatzs only with complex frequency. The expression g_1 and g_2 appearing in (3.2), are respectively a solution of the transport equations

$$i\eta_1 \cdot \nabla g_1 - (\eta_1 \cdot A_{1,\#})g_1 = 0, \quad i\eta_2 \cdot \nabla g_2 + (\eta_2 \cdot A_{2,\#})g_2 = 0, \quad (3.3)$$

with $A_{j,\#}$ some smooth function close to the magnetic potential A_j , $j = 1, 2$. More precisely, we define $A_{j,\#} \in \mathcal{C}_0^\infty(\mathbb{R}^n, \mathbb{R}^n)$, $j = 1, 2$, some smooth approximations

on $\bar{\Omega}$ of A_j . Then, we consider solutions of the transport equations (3.3) given by

$$\begin{aligned} g_1(x) &:= e^{i\psi_1(x)}, \\ g_2(x) &:= b_2(x)e^{-i\psi_2(x)}, \\ \psi_j(x) &:= -\int_{-\infty}^0 \eta_j \cdot A_{j,\#}(x + s\eta_j) ds, \quad \eta_2 \cdot \nabla b_2(x) = 0, \end{aligned}$$

for $x \in \mathbb{R}^n$. Therefore, we consider ansatz associated with (3.1) taking the form

$$\Phi_{1,\lambda}(x) := e^{i\sqrt{\lambda}\eta_1 \cdot x} e^{i\psi_1(x)}, \quad \Phi_{2,\lambda}(x) := e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2(x) e^{-i\psi_2(x)}, \quad x \in \Omega. \tag{3.4}$$

We assume in addition that $b_2 \in W^{2,\infty}(\mathbb{R}^n)$ and we recall that ψ_j solves the equation

$$\eta_j \cdot \nabla \psi_j(x) = -\eta_j \cdot A_{j,\#}, \quad j = 1, 2, \quad x \in \mathbb{R}^n.$$

For the time being, we consider general ansatz of the form (3.4) with the properties describe above. Additional information about the parameter λ , the function $A_{j,\#}$, the vector η_j , $j = 1, 2$, and the function b_2 will be given in Section 3.2. In a similar way to [11, 17, 18, 23, 26, 30], in the construction of our ansatz we consider some smooth approximations of the magnetic potentials instead of the magnetic potentials to obtain sufficiently smooth functions $\Phi_{j,\lambda}$, $j = 1, 2$. Using this approach, we can weaken the regularity assumption imposed on admissible magnetic potential from $W^{3,\infty}(\Omega)$ to $\mathcal{C}^1(\bar{\Omega})$. Further, for $j = 1, 2$, we put

$$S_j(\lambda, \eta_1, \eta_2) = \langle \Lambda_{j,\lambda} \Phi_{1,\lambda}, \overline{\Phi_{2,\lambda}} \rangle = \int_{\Gamma} (\Lambda_{j,\lambda} \Phi_{1,\lambda}) \Phi_{2,\lambda}(x) d\sigma(x). \tag{3.5}$$

In other words, we apply $\Lambda_{j,\lambda}$, $j = 1, 2$, to ansatz of the form (3.2) with $g_1 = e^{i\psi_1(x)}$ and $g_2 = b_2(x)e^{-i\psi_2(x)}$. We recall that quantities similar to S_1 and S_2 have also been used by [11, 14, 16, 17, 18, 23, 26, 30]. Let us also mention that, like in [14, 16], the ansatz (3.4) do not depend on the potential V_1 and V_2 which are coefficients of order zero of the equation (3.1). On the other hand, the ansatz (3.4) depend on the magnetic potentials A_1 and A_2 which are coefficients of order one of the equation (3.1). By modifying the construction of [14, 16] with the new expression g_j , $j = 1, 2$, we will extend the approach of [14, 16] to Schrödinger operators with magnetic potentials. From now on, for the sake of simplicity we will systematically omit the subscripts λ in $\Phi_{j,\lambda}$, $j = 1, 2$, in the remaining of this text. In view of determining the behavior of $S_1 - S_2$, as $\mathfrak{I}\lambda \rightarrow +\infty$, we introduce the following representations associated with S_1 and S_2 .

Proposition 3.1. For all $\lambda \in \mathbb{C} \setminus \mathbb{R}$ and $\eta_j \in \mathbb{S}^{n-1}$, $j = 1, 2$, the scalar products $S_j(\lambda, \eta_1, \eta_2)$ have the following expression

$$\begin{aligned}
 S_1(\lambda, \eta_1, \eta_2) &= 2\sqrt{\lambda} \int_{\Omega} \eta_2 \cdot (A_1 - A_{2,\#}) e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} b_2 e^{i(\psi_1(x) - \psi_2(x))} dx \\
 &\quad + \int_{\Omega} (V_1 - q_{12}) e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} b_2 e^{i(\psi_1(x) - \psi_2(x))} dx \\
 &\quad - i \int_{\Gamma} e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} e^{i(\psi_1(x) - \psi_2(x))} \\
 &\quad \quad (b_2 \sqrt{\lambda} \eta_2 + b_2 \nabla \psi_2 + i \nabla b_2 + b_2 A_1) \cdot \nu d\sigma(x) \\
 &\quad - \int_{\Omega} [(H_1 - \lambda)^{-1} (2\sqrt{\lambda} \eta_1 \cdot (A_1 - A_{1,\#}) + q_{11}) \Phi_1] \\
 &\quad \quad (2\sqrt{\lambda} \eta_2 \cdot (A_1 - A_{2,\#}) b_2 + V_1 b_2 - q_{12}) e^{-i\sqrt{\lambda} \eta_2 \cdot x} e^{-i\psi_2} dx,
 \end{aligned} \tag{3.6}$$

and

$$\begin{aligned}
 S_2(\lambda, \eta_1, \eta_2) &= \int_{\Omega} [2\sqrt{\lambda} \eta_2 \cdot (A_2 - A_{2,\#}) + V_2 - q_{22}] e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} b_2 e^{i(\psi_1(x) - \psi_2(x))} dx \\
 &\quad - i \int_{\partial\Omega} e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} e^{i(\psi_1(x) - \psi_2(x))} \\
 &\quad \quad (b_2 \sqrt{\lambda} \eta_2 + b_2 \nabla \psi_2 + i \nabla b_2 + b_2 A_2) \cdot \nu d\sigma(x) \\
 &\quad - \int_{\Omega} [(H_2 - \lambda)^{-1} (2\sqrt{\lambda} \eta_1 \cdot (A_2 - A_{1,\#}) + q_{21}) \Phi_1] \\
 &\quad \quad (2\sqrt{\lambda} \eta_2 \cdot (A_2 - A_{2,\#}) b_2 + V_2 b_2 - q_{22}) e^{-i\sqrt{\lambda} \eta_2 \cdot x} e^{-i\psi_2} dx.
 \end{aligned} \tag{3.7}$$

Here we denote by q_{11} , q_{12} , q_{21} , q_{22} the expressions

$$\begin{aligned}
 q_{11} &= -i \operatorname{div}(A_1) + |A_1|^2 + V_1(x) + 2A_1 \cdot \nabla \psi_1 - i \Delta \psi_1 + |\nabla \psi_1|^2, \\
 q_{12} &= \Delta b_2 - 2i \nabla \psi_2 \cdot \nabla b_2 - 2i \nabla b_2 \cdot A_1 \\
 &\quad + (-i \Delta \psi_2 - |\nabla \psi_2|^2 - 2 \nabla \psi_2 \cdot A_1 - i \operatorname{div}(A_1) - |A_1|^2) b_2, \\
 q_{21} &= -i \operatorname{div}(A_2) + |A_2|^2 + V_2(x) + 2A_2 \cdot \nabla \psi_1 - i \Delta \psi_1 + |\nabla \psi_1|^2, \\
 q_{22} &= \Delta b_2 - 2i \nabla \psi_2 \cdot \nabla b_2 - 2i \nabla b_2 \cdot A_2 \\
 &\quad + (-i \Delta \psi_2 - |\nabla \psi_2|^2 - 2 \nabla \psi_2 \cdot A_2 - i \operatorname{div}(A_2) - |A_2|^2) b_2.
 \end{aligned}$$

Moreover, H_j , $j = 1, 2$, denotes the selfadjoint operator $(-i \nabla + A_j) + V_j$ acting on $L^2(\Omega)$ with domain

$$D(H_j) = \{v \in H_0^1(\Omega) : (-i \nabla + A_j)v \in L^2(\Omega)\}.$$

Note that formulas (3.6) and (3.7) contain expressions involving the magnetic potentials A_1, A_2 and the electric potentials V_1, V_2 , expressions on the boundary $\partial\Omega$ and expressions described by the resolvent $(H_j - \lambda)^{-1}, j = 1, 2$. Using condition (1.1) one can check that the expressions on $\partial\Omega$ of S_1 and S_2 coincide and applying the decay of the resolvent $(H_j - \lambda)^{-1}, j = 1, 2$, as $\Im\lambda \rightarrow +\infty$ we will show in the next subsection that, for some suitable choice of our ansatz, the expressions

$$\begin{aligned}
 & - \int_{\Omega} [(H_1 - \lambda)^{-1}(2\sqrt{\lambda}\eta_1 \cdot (A_1 - A_{1,\#}) + q_{11})\Phi_1] \\
 & \quad (2\sqrt{\lambda}\eta_2 \cdot (A_1 - A_{2,\#})b_2 + V_1b_2 - q_{12})e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} dx, \\
 & - \int_{\Omega} [(H_2 - \lambda)^{-1}(2\sqrt{\lambda}\eta_1 \cdot (A_2 - A_{1,\#}) + q_{21})\Phi_1] \\
 & \quad (2\sqrt{\lambda}\eta_2 \cdot (A_2 - A_{2,\#})b_2 + V_2b_2 - q_{22})e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} dx,
 \end{aligned}$$

vanish as $\Im\lambda \rightarrow +\infty$. Thus, what will remain in the asymptotic expansion of $S_1 - S_2$, as $\Im\lambda \rightarrow +\infty$, will be two expressions involving $A_1 - A_2$ and $V_1 - V_2$. These two expressions, that will be given in the next subsection, are one of the main ingredients in our proof. The remaining of this subsection will be devoted to the proof of Proposition 3.1.

Proof of Proposition 3.1. Let us first remark that the expressions (3.6) and (3.7) correspond to some asymptotic expansion of the expression $S_j, j = 1, 2$, with respect to $\sqrt{\lambda}$ ². We will prove (3.6) and (3.7) by combining properties of the ansatz (3.2), with properties of solutions of (3.1) when $f = \Phi_1$. This proof will be divided into two steps, first for S_1 then for S_2 . We start by showing that for $j = 1, 2$ and $f = \Phi_1$ problem (3.1) admits a unique solution $u_j \in H^2(\Omega)$ taking the form

$$u_1 = \Phi_1 - (H_1 - \lambda)^{-1}[2\sqrt{\lambda}\eta_1 \cdot (A_1 - A_{1,\#}) + q_{11}]\Phi_1, \tag{3.8}$$

$$u_2 = \Phi_1 - (H_2 - \lambda)^{-1}[2\sqrt{\lambda}\eta_1 \cdot (A_2 - A_{1,\#}) + q_{21}]\Phi_1. \tag{3.9}$$

Then, combining these formulas with the properties of the ansatz (3.2) and applying the Green formula, we derive (3.6) and (3.7).

We start with the expression of $S_1(\lambda, \eta_1, \eta_2)$. Let us first prove (3.8). Recall that

$$(-i\nabla + A_1)^2u + V_1u - \lambda u = -\Delta u - 2iA_1 \cdot \nabla u + qu - \lambda u$$

²This statement will be clarified in the next subsection where we will give additional information about the parameter λ and the vectors η_1, η_2 .

with $q(x) = -i \operatorname{div}(A_1)(x) + |A_1(x)|^2 + V_1(x)$. Therefore, in light of (3.4) we have

$$\begin{aligned} & (-i\nabla + A_1)^2 \Phi_1 + V_1 \Phi_1 - \lambda \Phi_1 \\ &= (\lambda + 2\sqrt{\lambda} \eta_1 \cdot \nabla \psi_1 - i\Delta \psi_1 + |\nabla \psi_1|^2) \Phi_1 \\ &\quad + (2\sqrt{\lambda} \eta_1 \cdot A_1 + 2A_1 \cdot \nabla \psi_1) \Phi_1 \\ &\quad + q \Phi_1 - \lambda \Phi_1 \\ &= 2\sqrt{\lambda} (\eta_1 \cdot \nabla \psi_1 + \eta_1 \cdot A_1) \Phi_1 + q_{11} \Phi_1 \end{aligned}$$

with $q_{11} = q + 2A_1 \cdot \nabla \psi_1 - i\Delta \psi_1 + |\nabla \psi_1|^2$. On the other hand, since ψ_1 satisfies $\eta_1 \cdot \nabla \psi_1 + \eta_1 \cdot A_{1,\#} = 0$, we deduce that

$$(-i\nabla + A_1)^2 \Phi_1 + V_1 \Phi_1 - \lambda \Phi_1 = [2\sqrt{\lambda} \eta_1 \cdot (A_1 - A_{1,\#}) + q_{11}] \Phi_1. \quad (3.10)$$

Now consider u_1 the solution of

$$\begin{cases} (-i\nabla + A_1)^2 u_1 + V_1 u_1 - \lambda u_1 = 0, & \text{in } \Omega, \\ u_1(x) = \Phi_1(x), & x \in \partial\Omega. \end{cases}$$

Note that, with our assumptions one can check that $D(H_1) = H_0^1(\Omega) \cap H^2(\Omega)$. In view of (3.10), we can split u_1 into two terms $u_1 = \Phi_1 + v_1$ with v_1 the solution of

$$\begin{cases} (-i\nabla + A_1)^2 v_1 + V_1 v_1 - \lambda v_1 = -[2\sqrt{\lambda} \eta_1 \cdot (A_1 - A_{1,\#}) + q_{11}] \Phi_1, & \text{in } \Omega, \\ v_1(x) = 0, & x \in \partial\Omega. \end{cases}$$

Then, $u_1 \in H^2(\Omega)$ take the form (3.8). Using this formula we will complete the proof of (3.6). Since

$$S_1 = \int_{\partial\Omega} (\partial_\nu + iA_1 \cdot \nu) u_1(x) e^{-i\sqrt{\lambda} \eta_2 \cdot x} b_2 e^{-i\psi_2(x)} d\sigma(x), \quad (3.11)$$

from (3.5), applying Green formula, we get

$$\begin{aligned} S_1 &= \int_{\Omega} \operatorname{div}((\nabla + iA_1(x)) u_1(x) e^{-i\sqrt{\lambda} \eta_2 \cdot x} b_2 e^{-i\psi_2(x)}) dx \\ &= \int_{\Omega} (\nabla + iA_1)^2 u_1 e^{-i\sqrt{\lambda} \eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\ &\quad + \int_{\Omega} (\nabla + iA_1) u_1 \cdot (\nabla - iA_1) e^{-i\sqrt{\lambda} \eta_2 \cdot x} b_2 e^{-i\psi_2} dx. \end{aligned} \quad (3.12)$$

Doing the same with the second term on the right hand side of this formula, we find out that

$$\begin{aligned} & \int_{\Omega} (\nabla + iA_1)u_1 \cdot (\nabla - iA_1)e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\ &= -i \int_{\Gamma} u_1(x) e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} (\sqrt{\lambda}b_2\eta_2 + b_2\nabla\psi_2 + i\nabla b_2 + b_2A_1) \cdot \nu d\sigma(x) \\ & \quad - \int_{\Omega} u_1(x) (\nabla - iA_1)^2 e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx. \end{aligned}$$

In light of (3.4) and the identity $u_1|_{\Gamma} = \Phi_1$, this entails

$$\begin{aligned} & \int_{\Omega} (\nabla + iA_1)u_1 \cdot (\nabla - iA_1)e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\ &= -i \int_{\Gamma} e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} e^{i(\psi_1(x) - \psi_2(x))} \\ & \quad (\sqrt{\lambda}b_2\eta_2 + b_2\nabla\psi_2 + i\nabla b_2 + b_2A_1) \cdot \nu d\sigma(x) \\ & \quad - \int_{\Omega} u_1(x) (\nabla - iA_1)^2 e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx. \end{aligned}$$

Moreover, one can check that

$$\begin{aligned} & (\nabla - iA_1)^2 e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} \\ &= (-\lambda b_2 - 2\sqrt{\lambda}(\eta_2 \cdot \nabla\psi_2 + A_1 \cdot \eta_2)b_2 \\ & \quad - 2i\sqrt{\lambda}\eta_2 \cdot \nabla b_2 + q_{12})e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} \end{aligned}$$

with

$$\begin{aligned} q_{12} &= \Delta b_2 - 2i\nabla\psi_2 \cdot \nabla b_2 - 2i\nabla b_2 \cdot A_1 \\ & \quad + (-i\Delta\psi_2 - |\nabla\psi_2|^2 - 2\nabla\psi_2 \cdot A_1 - i\operatorname{div}(A_1) - |A_1|^2)b_2. \end{aligned}$$

Combining this with the fact that ψ_2 satisfies

$$\eta_2 \cdot \nabla\psi_2 + \eta_2 \cdot A_{2,\#} = 0$$

and b_2 solves

$$\eta_2 \cdot \nabla b_2 = 0,$$

we deduce that

$$\begin{aligned} & (\nabla - iA_1)^2 e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} \\ &= ([-\lambda - 2\sqrt{\lambda}\eta_2 \cdot (A_1 - A_{2,\#})]b_2 + q_{12})e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2}. \end{aligned}$$

Therefore, we find

$$\begin{aligned}
& \int_{\Omega} (\nabla + iA_1)u_1 \cdot (\nabla - iA_1)e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\
&= -i \int_{\Gamma} e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} e^{i(\psi_1(x) - \psi_2(x))} \\
&\quad (\sqrt{\lambda}b_2\eta_2 + b_2\nabla\psi_2 + i\nabla b_2 + b_2A_1) \cdot \nu d\sigma(x) \\
&\quad - \int_{\Omega} u_1(x)(-\lambda b_2 - 2\sqrt{\lambda}\eta_2 \cdot (A_1 - A_{2,\#})b_2 + q_{12})e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} dx.
\end{aligned}$$

Then, from (3.8) we get

$$\begin{aligned}
& \int_{\Omega} (\nabla + iA_1)u_1 \cdot (\nabla - iA_1)e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\
&= -i \int_{\Gamma} e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} e^{i(\psi_1(x) - \psi_2(x))} \\
&\quad (\sqrt{\lambda}b_2\eta_2 + b_2\nabla\psi_2 + i\nabla b_2 + b_2A_1) \cdot \nu d\sigma(x) \\
&\quad + \lambda \int_{\Omega} u_1 e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\
&\quad + 2\sqrt{\lambda} \int_{\Omega} \eta_2 \cdot (A_1 - A_{2,\#})e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} b_2 e^{i(\psi_1(x) - \psi_2(x))} dx \\
&\quad - \int_{\Omega} q_{12} e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} b_2 e^{i(\psi_1(x) - \psi_2(x))} dx \\
&\quad - \int_{\Omega} [(H_1 - \lambda)^{-1}(2\sqrt{\lambda}\eta_1 \cdot (A_1 - A_{1,\#}) + q_{11})\Phi_1] \\
&\quad (2\sqrt{\lambda}\eta_2 \cdot (A_1 - A_{2,\#})b_2 - q_{12})e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} dx.
\end{aligned} \tag{3.13}$$

Next, taking into account the fact that $(\nabla + iA_1)^2 u_1 = (V_1 - \lambda)u_1$ in Ω , we obtain

$$\begin{aligned}
& \int_{\Omega} (\nabla + iA_1)^2 u_1 e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\
&= \int_{\Omega} (V_1 - \lambda)u_1 e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\
&= -\lambda \int_{\Omega} u_1 e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\
&\quad + \int_{\Omega} V_1 e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} b_2 e^{i(\psi_1(x) - \psi_2(x))} dx \\
&\quad - \int_{\Omega} V_1 [(H_1 - \lambda)^{-1}(2\sqrt{\lambda}\eta_1 \cdot (A_1 - A_{1,\#}) + q_{11})\Phi_1] e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx.
\end{aligned}$$

Finally, we deduce (3.6) from (3.12) and (3.13).

Now let us consider (3.7). For this purpose, we start by proving formula (3.9). In a similar way to (3.8), we have

$$(-i\nabla + A_2)^2 \Phi_1 + V_2 \Phi_1 - \lambda \Phi_1 = 2\sqrt{\lambda}(\eta_1 \cdot \nabla \psi_1 + A_2 \cdot \eta_1) \Phi_1 + q_{21} \Phi_1$$

with

$$q_{21} = -i \operatorname{div}(A_2) + |A_2|^2 + V_2(x) + 2A_2 \cdot \nabla \psi_1 - i\Delta \psi_1 + |\nabla \psi_1|^2.$$

Then, since ψ_1 is a solution of $\eta_1 \cdot \nabla \psi_1 + \eta_1 \cdot A_{1,\#} = 0$, we deduce that

$$(-i\nabla + A_2)^2 \Phi_1 + V_2 \Phi_1 - \lambda \Phi_1 = (2\sqrt{\lambda} \eta_1 \cdot (A_2 - A_{1,\#}) + q_{21}) \Phi_1.$$

Moreover, one can check that the solution u_2 of

$$\begin{cases} (-i\nabla + A_2)^2 u_2 + V_2 u_2 - \lambda u_2 = 0, & \text{in } \Omega, \\ u_2(x) = \Phi_1(x), & x \in \partial\Omega \end{cases}$$

is given by (3.9). Repeating our previous arguments, we deduce

$$\begin{aligned} S_2 &= \int_{\Omega} (\nabla + iA_2)^2 u_2 e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\ &\quad + \int_{\Omega} (\nabla + iA_2) u_2 \cdot (\nabla - iA_2) e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx. \end{aligned} \quad (3.14)$$

On the other hand, using the fact that ψ_2 is a solution of the equation $\eta_2 \cdot \nabla \psi_2 + \eta_2 \cdot A_{2,\#} = 0$, we get

$$\begin{aligned} &\int_{\Omega} (\nabla + iA_2) u_2 \cdot (\nabla - iA_2) e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2} dx \\ &= -i \int_{\Gamma} e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} e^{i(\psi_1(x) - \psi_2(x))} \\ &\quad (\sqrt{\lambda} b_2 \eta_2 + b_2 \nabla \psi_2 + i\nabla b_2 + b_2 A_2) \cdot \nu d\sigma(x) \\ &\quad - \int_{\Omega} u_2(x) (-\lambda b_2 - 2\sqrt{\lambda} \eta_2 \cdot (A_2 - A_{2,\#}) b_2 + q_{22}) e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} dx \end{aligned} \quad (3.15)$$

with

$$\begin{aligned} q_{22} &= \Delta b_2 - 2i\nabla \psi_2 \cdot \nabla b_2 - 2i\nabla b_2 \cdot A_2 \\ &\quad + (-i\Delta \psi_2 - |\nabla \psi_2|^2 - 2\nabla \psi_2 \cdot A_2 - i \operatorname{div}(A_2) - |A_2|^2) b_2. \end{aligned}$$

Combining this with (3.9)–(3.14) and repeating our previous arguments we obtain (3.7). \square

3.2. Asymptotic properties of $S_1 - S_2$ and representation formulas for $A_1 - A_2$ and $V_1 - V_2$. In this subsection we will apply formulas (3.6) and (3.7) in order to derive two expressions involving $A_1 - A_2$ and $V_1 - V_2$ from the asymptotic expansion of $S_1 - S_2$, as $\Im\lambda \rightarrow +\infty$. For this purpose, we start by specifying our choice for the parameter λ , the function $A_{j,\#}$, the vector η_j , $j = 1, 2$, and the function b_2 appearing in (3.2). Let us first define the parameter λ and the vectors η_1, η_2 . We consider an arbitrary $\xi \in \mathbb{R}^n \setminus \{0\}$ and pick $\eta \in \mathbb{S}^{n-1}$ such that $\eta \cdot \xi = 0$. Then, for $\tau > |\xi|$ we put

$$B_\tau = \sqrt{1 - \frac{|\xi|^2}{4\tau^2}}, \tag{3.16a}$$

$$\eta_1(\tau) = B_\tau \eta - \frac{\xi}{2\tau}, \tag{3.16b}$$

$$\eta_2(\tau) = B_\tau \eta + \frac{\xi}{2\tau}, \tag{3.16c}$$

$$\lambda(\tau) = (\tau + i)^2, \tag{3.16d}$$

in such a way that

$$\begin{cases} \eta_1, \eta_2 \in \mathbb{S}^{n-1}, \\ \sqrt{\lambda}(\eta_1 - \eta_2) \rightarrow -\xi, & \text{as } \tau \rightarrow +\infty, \\ \Im\lambda \rightarrow +\infty, & \text{as } \tau \rightarrow +\infty, \\ \Im\sqrt{\lambda}\eta_1, \Im\sqrt{\lambda}\eta_2 \text{ are bounded with respect to } \tau > |\xi|. \end{cases} \tag{3.17}$$

In order to get a suitable expression of the functions $A_{j,\#}$, we first need to extend identically the magnetic potentials A_j , $j = 1, 2$. For this purpose we set $\tilde{\Omega}$ an arbitrary open bounded set of \mathbb{R}^n such that $\bar{\Omega} \subset \tilde{\Omega}$ and we define $\tilde{A}_1 \in \mathcal{C}_0^1(\tilde{\Omega}, \mathbb{R}^n)$ such that $\tilde{A}_1|_\Omega = A_1$. Then, we define \tilde{A}_2 by

$$\tilde{A}_2(x) = \begin{cases} A_2(x) & \text{for } x \in \Omega, \\ \tilde{A}_1(x) & \text{for } x \in \tilde{\Omega} \setminus \Omega. \end{cases}$$

In view of (1.1), it is clear that $\tilde{A}_2 \in \mathcal{C}_0^1(\tilde{\Omega}, \mathbb{R}^n)$. We define the functions $A_{j,\#} \in \mathcal{C}_0^\infty(\mathbb{R}^n; \mathbb{R}^n)$, $j = 1, 2$, by

$$A_{j,\#}(x) := \chi_\delta * \tilde{A}_j(x) = \int_{\mathbb{R}^n} \chi_\delta(x - y) \tilde{A}_j(y) dy,$$

where $\chi_\delta(x) = \delta^{-n} \chi(\delta^{-1}x)$, with $\delta > 0$, is the usual mollifier with $\chi \in \mathcal{C}_0^\infty(\mathbb{R}^n)$,

$\text{supp}(\chi) \subset \{x \in \mathbb{R}^n : |x| \leq 1\}$, $\chi \geq 0$ and $\int_{\mathbb{R}^n} \chi dx = 1$. From now on we set $\delta = \tau^{-\frac{1}{3}}$ and we recall that

$$\psi_j(x) = - \int_{-\infty}^0 \eta_j \cdot A_{j,\#}(x + s\eta_j) ds.$$

We set also

$$b_2(x) = e^{i\omega \cdot x} y \cdot \nabla \left[\exp \left(-i \int_{\mathbb{R}} \eta_2 \cdot A_{\#}(x + s\eta_2) ds \right) e^{-i\omega \cdot x} \right], \tag{3.18}$$

where

$$A_{\#} = A_{2,\#} - A_{1,\#}, \quad \omega = B_{\tau} \xi - \frac{|\xi|^2 \eta}{2\tau} \in \eta_2^{\perp}, \quad B_{\tau} = \sqrt{1 - \frac{|\xi|^2}{4\tau^2}},$$

and

$$b(x) = e^{ix \cdot \xi} y \cdot \nabla \left[\exp \left(-i \int_{\mathbb{R}} \eta \cdot A(x + s\eta) ds \right) e^{-ix \cdot \xi} \right],$$

$$\psi(x) = \int_{-\infty}^0 \eta \cdot A(x + s\eta) ds.$$

Here $y \in \mathbb{S}^{n-1} \cap \eta^{\perp}$, $y \cdot \nabla$ denotes the derivative in the $y = (y_1, \dots, y_n)$ direction given by

$$y \cdot \nabla = \sum_{j=1}^n y_j \partial_{x_j}$$

and A is the function defined by $A_2 - A_1$ on Ω extended by 0 outside Ω . Note that, in view of condition (1.1) we have $A \in \mathcal{C}_0^1(\Omega)$. Since $\tilde{A}_j \in \mathcal{C}_0^1(\mathbb{R}^n, \mathbb{R}^n)$, we find

$$\|A_{j,\#} - A_j\|_{L^\infty(\Omega)} \leq \|A_{j,\#} - \tilde{A}_j\|_{L^\infty(\mathbb{R}^n)} \leq C\delta = C\tau^{-\frac{1}{3}} \tag{3.19}$$

with C depending on Ω and any $M \geq \max_{j=1,2} \|\tilde{A}_j\|_{W^{1,\infty}(\mathbb{R}^n)}$. On the other hand, one can check that

$$\|\partial_x^\alpha A_{j,\#}\|_{L^\infty(\mathbb{R}^m)} \leq C\delta^{-|\alpha|+1} = C\tau^{\frac{|\alpha|-1}{3}}, \quad \alpha \in \mathbb{N}^n \setminus \{0\}, \tag{3.20}$$

where C depends on Ω and any $M \geq \max_{j=1,2} \|\tilde{A}_j\|_{W^{1,\infty}(\mathbb{R}^n)}$.

Remark 3.2. Let us observe that, our anstazs are related to the principal part of the complex geometric optics solutions of [31] and the extension of this construction to magnetic Schrödinger operators by [11, 17, 18, 23, 26, 30]. Nevertheless, in contrast to the complex geometric optics solutions of [31], the large parameter

of the ansatz (3.2), that will be sent to $+\infty$ for the uniqueness result, is given by $\Im\lambda$ where the parameter λ appears explicitly in (3.1). This makes it possible to construct ansatzs bounded with respect to the large parameter and to use the resolvent $(H_j - \lambda)^{-1}$, $j = 1, 2$, for the construction of a remainder term that admits a decay with respect to the large parameter $\Im\lambda$. Moreover, in contrast to the geometric optics solutions of [31], whose principal parts take the form $e^{\xi_j \cdot x}$ with $\xi_j \cdot \xi_j = 0$, our construction is not restricted to dimension $n \geq 3$. Indeed, our construction is subjected only to the condition (3.17), already considered by [14], which requires only the two orthogonal vectors η and ξ appearing in (3.16). For this reason, in contrast to the construction of [31], that requires three orthogonal vectors, our construction works also for $n = 2$.

From now on, our goal is to derive from (3.6) and (3.7) two formulas from some asymptotic properties of $S_1 - S_2$ as $\tau \rightarrow +\infty$. For this purpose we need the following intermediate result which follows from (3.19) and (3.20).

Lemma 3.3. *Let the condition introduced above be fulfilled. Then, we have*

$$\sup_{\tau > |\xi| + 1} \|b_2\|_{L^\infty(\mathbb{R}^n)} < \infty \tag{3.21}$$

and

$$\lim_{\tau \rightarrow +\infty} b_2(x) = b(x), \quad \lim_{\tau \rightarrow +\infty} \psi_1(x) - \psi_2(x) = \psi(x), \quad x \in \mathbb{R}^n. \tag{3.22}$$

Proof. Note first that

$$b_2(x) = \left(-i\omega \cdot y - i \int_{\mathbb{R}} \eta_2 \cdot y \cdot \nabla A_{\#}(x + s\eta_2) ds \right) \exp \left(-i \int_{\mathbb{R}} \eta_2 \cdot A_{\#}(x + s\eta_2) ds \right). \tag{3.23}$$

On the other hand, we have $|\omega| \leq 1 + |\xi|$ and, since $\tilde{A}_2 - \tilde{A}_1$ is compactly supported and $\tilde{A}_2 - \tilde{A}_1 \in \mathcal{C}_0^1(\mathbb{R}^n, \mathbb{R}^n)$, we find $y \cdot \nabla A_{\#} = \chi_\delta * (y \cdot \nabla(\tilde{A}_2 - \tilde{A}_1))$. Therefore, we obtain

$$\begin{aligned} \|b_2\|_{L^\infty(\mathbb{R}^n)} &\leq 1 + |\xi| + C \|\chi_\delta\|_{L^1(\mathbb{R}^n)} \|y \cdot \nabla(\tilde{A}_2 - \tilde{A}_1)\|_{L^\infty(\mathbb{R}^n, \mathbb{R}^n)} \\ &\leq 1 + |\xi| + CM \end{aligned}$$

with C a generic constant depending only on Ω and $M \geq \max_{j=1,2} \|\tilde{A}_j\|_{W^{1,\infty}(\mathbb{R}^n)}$.

From this last estimate we deduce (3.21). Now let us prove (3.22). Since \tilde{A}_1 and \tilde{A}_2 coincide outside of Ω , we have $\tilde{A}_2 - \tilde{A}_1 = A$. Therefore, we deduce that $A_{\#} = \chi_{\delta} * A$ and

$$|y \cdot \nabla A_{\#}(x + s\eta_2) - y \cdot \nabla A(x + s\eta)| \leq |y \cdot \nabla A_{\#}(x + s\eta_2) - y \cdot \nabla A_{\#}(x + s\eta)| + |y \cdot \nabla A_{\#}(x + s\eta) - y \cdot \nabla A(x + s\eta)|. \tag{3.24}$$

The second term on the right hand side of this estimate can be rewritten as

$$y \cdot \nabla A_{\#}(x + s\eta) - y \cdot \nabla A(x + s\eta) = \chi_{\delta} * [y \cdot \nabla A](x + s\eta) - y \cdot \nabla A(x + s\eta)$$

and since $A \in C_0^1(\mathbb{R}^n)$, we get

$$\lim_{\tau \rightarrow +\infty} y \cdot \nabla A_{\#}(x + s\eta) - y \cdot \nabla A(x + s\eta) = 0, \quad x \in \mathbb{R}^n, s \in \mathbb{R}. \tag{3.25}$$

For the first term on the right hand side of (3.24), using the fact that for τ sufficiently large we have

$$\eta_2 = \eta + \frac{\xi}{2\tau} + o_{\tau \rightarrow +\infty} \left(\frac{1}{\tau} \right)$$

and applying (3.20), we get

$$|y \cdot \nabla A_{\#}(x + s\eta_2) - y \cdot \nabla A_{\#}(x + s\eta)| \leq \|A_{\#}\|_{W^{2,\infty}(\mathbb{R}^n)} |s(\eta - \eta_1)| \leq C |s| \tau^{-\frac{2}{3}}$$

with C depending on ξ, Ω, \tilde{A}_1 and \tilde{A}_2 . In view of this estimate we have

$$\lim_{\tau \rightarrow +\infty} y \cdot \nabla A_{\#}(x + s\eta_2) - y \cdot \nabla A_{\#}(x + s\eta) = 0, \quad x \in \mathbb{R}^n, s \in \mathbb{R}.$$

Combining this last result with (3.24) and (3.25), we get

$$\lim_{\tau \rightarrow +\infty} y \cdot \nabla A_{\#}(x + s\eta) = y \cdot \nabla A(x + s\eta), \quad x \in \mathbb{R}^n, s \in \mathbb{R}.$$

Then, using the fact that $\text{supp}(A_{\#}) \subset \Omega + \{x \in \mathbb{R}^n : |x| \leq \delta\}$ and (3.20), by the dominate convergence theorem we get that

$$\lim_{\tau \rightarrow +\infty} \int_{\mathbb{R}} y \cdot \nabla A_{\#}(x + s\eta_2) ds = \int_{\mathbb{R}} y \cdot \nabla A(x + s\eta) ds, \quad x \in \mathbb{R}^n.$$

Putting this together with (3.23) and the fact that $\omega \rightarrow \xi$, $\eta_2 \rightarrow \eta$ as $\tau \rightarrow +\infty$, we obtain

$$\begin{aligned} \lim_{\tau \rightarrow +\infty} b_2(x) &= \left(-i\xi \cdot y + -i \int_{\mathbb{R}} \eta \cdot y \cdot \nabla A(x + s\eta) ds \right) \\ &\quad \exp \left(-i \int_{\mathbb{R}} \eta \cdot A(x + s\eta) ds \right) \\ &= b(x), \quad x \in \mathbb{R}^n. \end{aligned}$$

Using similar arguments we deduce that

$$\lim_{\tau \rightarrow +\infty} \psi_1(x) - \psi_2(x) = \psi(x) = \int_{-\infty}^0 \eta \cdot A(x + s\eta) ds, \quad x \in \mathbb{R}^n.$$

This completes the proof of the lemma. □

Applying (3.6), (3.7), and (3.19)–(3.22), and sending $\tau \rightarrow +\infty$, we obtain our first formula involving the magnetic potentials A_1, A_2 .

Proposition 3.4. *Fix $\xi \in \mathbb{R}^n \setminus \{0\}$ and $\eta \in \mathbb{S}^{n-1}$ such that $\eta \cdot \xi = 0$. Let λ, η_1 and η_2 be defined by (3.16) and let b_2 be defined by (3.18). Then, we have*

$$\lim_{\tau \rightarrow +\infty} \frac{S_1 - S_2}{\sqrt{\lambda}} = 2 \int_{\Omega} \eta \cdot (A_1 - A_2) e^{-i\xi \cdot x} b e^{i\psi(x)} dx. \tag{3.26}$$

Proof. With reference to (3.4) and (3.16) we have

$$|\Phi_1(x)| = e^{-\eta_1 \cdot x} \quad \text{and} \quad |e^{-i\sqrt{\lambda}\eta_2 \cdot x}| = e^{\eta_2 \cdot x}$$

for all $x \in \Omega$, hence

$$\|\Phi_1\|_{L^2(\Omega)}^2 = \int_{\Omega} e^{-2\eta_1 \cdot x} dx \leq C$$

and

$$\|e^{-i\sqrt{\lambda}\eta_2 \cdot x}\|_{L^2(\Omega)}^2 \leq C$$

since $|\eta_1| = |\eta_2| = 1$. Moreover, in view of (3.16), we have the estimate

$$\|(H_j - \lambda)^{-1}\|_{\mathcal{B}(L^2(\Omega))} = \frac{1}{\text{dist}(\lambda, \sigma(H_j))} \leq \frac{1}{|\Im \lambda|} = \frac{1}{2\tau}, \quad j = 1, 2.$$

In addition, in light of (3.20), we get

$$\|\psi_j\|_{W^{2,\infty}(\Omega)} \leq C\delta = C\tau^{\frac{1}{3}}, \quad \|b_j\|_{W^{2,\infty}(\Omega)} \leq C\delta^2 = C\tau^{\frac{2}{3}}$$

with C a generic constant depending on ξ , Ω and \tilde{A}_j , $j = 1, 2$. Putting these estimates together with (1.1), (3.6), (3.7), and (3.19), we deduce that

$$\frac{S_1 - S_2}{\sqrt{\lambda}} = 2 \int_{\Omega} \eta_2 \cdot (A_1 - A_2) e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} b_2 e^{i(\psi_1(x) - \psi_2(x))} dx + o_{\tau \rightarrow +\infty}(\tau^{-\frac{1}{3}}).$$

Combining this with (3.17), (3.21), and (3.22), and applying the dominate convergence theorem we deduce (3.26). □

Using similar arguments and assuming that the magnetic potentials are known ($A_1 = A_2$), we obtain our second formula involving the electric potentials V_1, V_2 .

Proposition 3.5. *Assume that $A_1 = A_2$. Fix $\xi \in \mathbb{R}^n \setminus \{0\}$ and $\eta \in \mathbb{S}^{n-1}$ such that $\eta \cdot \xi = 0$. Let λ , η_1 and η_2 be defined by (3.16) and $b_2 = 1$. Then, we have*

$$\lim_{\tau \rightarrow +\infty} S_1 - S_2 = \int_{\Omega} (V_1 - V_2) e^{-i\xi \cdot x} dx. \tag{3.27}$$

Proof. Note that for $A_1 = A_2$ we have $q_{11} - V_1 = q_{21} - V_2$, $q_{12} = q_{22}$, $A_{1,\#} = A_{2,\#}$. Therefore, we deduce that (3.6) and (3.7) imply

$$\begin{aligned} S_1 - S_2 &= \int_{\Omega} (V_1 - V_2) e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} e^{i(\psi_1(x) - \psi_2(x))} dx \\ &\quad - \int_{\Omega} [\lambda((H_1 - \lambda)^{-1} - (H_2 - \lambda)^{-1}) Q_1] Q_2 dx \\ &\quad - \int_{\Omega} [\sqrt{\lambda}(H_1 - \lambda)^{-1} Q_1] V_1 e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} dx \\ &\quad - \int_{\Omega} [\sqrt{\lambda}(H_1 - \lambda)^{-1} V_1 \Phi_1] Q_2 dx \\ &\quad - \int_{\Omega} [(H_1 - \lambda)^{-1} V_1 \Phi_1] V_1 e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} dx \\ &\quad + \int_{\Omega} [\sqrt{\lambda}(H_2 - \lambda)^{-1} Q_1] V_2 e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} dx \\ &\quad + \int_{\Omega} [\sqrt{\lambda}(H_2 - \lambda)^{-1} V_2 \Phi_1] Q_2 dx \\ &\quad + \int_{\Omega} [(H_2 - \lambda)^{-1} V_2 \Phi_1] V_2 e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2} dx, \end{aligned} \tag{3.28}$$

where

$$Q_1 = 2\eta_1 \cdot (A_1 - A_{1,\#})\Phi_1 + \frac{(q_{11} - V_1)\Phi_1}{\sqrt{\lambda}},$$

$$Q_2 = \left(2\eta_2 \cdot (A_1 - A_{1,\#}) - \frac{q_{12}}{\sqrt{\lambda}}\right) e^{-i\sqrt{\lambda}\eta_2 \cdot x} e^{-i\psi_2}.$$

On the other hand, since $H_2 - \lambda = H_1 - \lambda - (V_1 - V_2)$, for τ sufficiently large we have

$$\begin{aligned} (H_1 - \lambda)^{-1} - (H_2 - \lambda)^{-1} &= (H_1 - \lambda)^{-1} (\text{Id} - (\text{Id} - (V_1 - V_2)(H_1 - \lambda)^{-1})^{-1}) \\ &= -(H_1 - \lambda)^{-1} \sum_{k=1}^{\infty} ((V_1 - V_2)(H_1 - \lambda)^{-1})^k. \end{aligned}$$

Combining this with the fact that $\Im\lambda = 2\tau$, $|\lambda| \leq |\tau^2 - 1| + 2\tau$, and the fact that

$$\|(H_1 - \lambda)^{-1}\|_{\mathcal{B}(L^2(\Omega))} + \|(V_1 - V_2)(H_1 - \lambda)^{-1}\|_{\mathcal{B}(L^2(\Omega))} \leq \frac{C}{|\Im\lambda|} = \frac{C}{2\tau}$$

with C depending only on V_1 , V_2 and Ω , we deduce that

$$\sup_{\tau > |\xi| + 1} \|\lambda ((H_1 - \lambda)^{-1} - (H_2 - \lambda)^{-1})\|_{\mathcal{B}(L^2(\Omega))} < \infty. \quad (3.29)$$

In addition, (3.19) and (3.20) imply

$$\lim_{\tau \rightarrow +\infty} \|Q_1\|_{L^\infty(\Omega)} = \lim_{\tau \rightarrow +\infty} \|Q_2\|_{L^\infty(\Omega)} = 0.$$

Putting this result together with (3.17), (3.28), and (3.29), we obtain

$$\limsup_{\tau \rightarrow +\infty} |(S_1 - S_2) - \int_{\Omega} (V_1 - V_2) e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} e^{i(\psi_1(x) - \psi_2(x))} dx| = 0.$$

On the other hand, repeating the arguments of Lemma 3.3, we find

$$\lim_{\tau \rightarrow +\infty} \psi_1(x) - \psi_2(x) = \psi(x) = \int_{-\infty}^0 \eta \cdot A(x + s\eta) ds = 0$$

since $A_1 = A_2$. Thus, applying the dominated convergence theorem we obtain

$$\lim_{\tau \rightarrow +\infty} \int_{\Omega} (V_1 - V_2) e^{i\sqrt{\lambda}(\eta_1 - \eta_2) \cdot x} e^{i(\psi_1(x) - \psi_2(x))} dx = \int_{\Omega} (V_1 - V_2) e^{-ix \cdot \xi} dx$$

and we deduce (3.27). \square

Armed with formulas (3.26) and (3.27), in the next section we will complete the proof of Theorem 1.1.

4. Proof of the main result

This section is devoted to the proof of our main result. In all this section, for $j = 1$ and $j = 2$, we consider two magnetic potentials A_j and electric potentials V_j satisfying the assumptions of Theorem 1.1 and we denote by H_j the associated operators defined by (2.1) for $A = A_j$ and $V = V_j$. Let $(\lambda_{j,k}, \varphi_{j,k})_{k \geq 1}$ be a sequence of eigenvalues and eigenfunctions of H_j . In order to prove Theorem 1.1, in light of (3.26)–(3.27), we prove first that the condition

$$\lim_{\tau \rightarrow +\infty} \frac{S_1(\lambda(\tau), \eta_1(\tau), \eta_2(\tau)) - S_2(\lambda(\tau), \eta_1(\tau), \eta_2(\tau))}{\sqrt{\lambda(\tau)}} = 0 \tag{4.1}$$

implies $dA_1 = dA_2$. Then, we show that for $A_1 = A_2$ the condition

$$\lim_{\tau \rightarrow +\infty} S_1(\lambda(\tau), \eta_1(\tau), \eta_2(\tau)) - S_2(\lambda(\tau), \eta_1(\tau), \eta_2(\tau)) = 0 \tag{4.2}$$

implies $V_1 = V_2$. Finally, we complete the proof by proving that conditions (1.1) and (1.2) imply (4.1) and (4.2).

We start by proving that (4.1) implies $dA_1 = dA_2$.

Lemma 4.1. *Let $\eta_1(\tau)$, $\eta_2(\tau)$ and $\lambda(\tau)$ be fixed by (3.16) and b_2 be defined by (3.18). Assume that (4.1) is fulfilled. Then, we have $dA_1 = dA_2$.*

Proof. Combining (4.1) with (3.26) we deduce that for all $\xi \in \mathbb{R}^n \setminus \{0\}$, $\eta \in \mathbb{S}^{n-1}$, satisfying $\eta \cdot \xi = 0$, we get

$$\int_{\Omega} \eta \cdot (A_2 - A_1) e^{-i\xi \cdot x} b(x) e^{i\psi(x)} dx = 0.$$

Here b takes the form

$$b(x) = e^{ix \cdot \xi} y \cdot \nabla \left[\exp \left(-i \int_{\mathbb{R}} \eta \cdot A(x + s\eta) ds \right) e^{-ix \cdot \xi} \right]$$

with $y \in \mathbb{S}^{n-1} \cap \eta^\perp$. Then, applying Fubini’s theorem, we obtain

$$\begin{aligned} 0 &= \int_{\mathbb{R}^n} \eta \cdot A(x) e^{-i\xi \cdot x} b(x) e^{i\psi(x)} dx \\ &= \int_{\eta^\perp} \int_{\mathbb{R}} \eta \cdot A(x' + t\eta) e^{i\psi(x' + t\eta)} b(x') e^{-i\xi \cdot x'} dt dx'. \end{aligned}$$

Here we use the fact that $b(x) = b(x - (x \cdot \eta)\eta)$ and $\xi \cdot \eta = 0$. On the other hand, for all $x' \in \eta^\perp$ and $t \in \mathbb{R}$, we have

$$\begin{aligned} \eta \cdot A(x' + t\eta)e^{i\psi(x'+t\eta)} &= \eta \cdot A(x' + t\eta) \exp\left(i \int_{-\infty}^t \eta \cdot A(x' + s\eta) ds\right) \\ &= -i\partial_t \exp\left(i \int_{-\infty}^t \eta \cdot A(x' + s\eta) ds\right). \end{aligned}$$

Therefore, we find

$$\begin{aligned} &\int_{\mathbb{R}^n} \eta \cdot A(x)e^{-i\xi \cdot x} b(x)e^{i\psi(x)} dx \\ &= -i \int_{\eta^\perp} \left[\int_{\mathbb{R}} \partial_t \exp\left(i \int_{-\infty}^t \eta \cdot A(x' + s\eta) ds\right) dt \right] b(x')e^{-i\xi \cdot x'} dx' \\ &= -i \int_{\eta^\perp} \left[\exp\left(i \int_{\mathbb{R}} \eta \cdot A(x' + s\eta) ds\right) - 1 \right] b(x')e^{-i\xi \cdot x'} dx'. \end{aligned}$$

It follows

$$\int_{\eta^\perp} \left[\exp\left(i \int_{\mathbb{R}} \eta \cdot A(x' + s\eta) ds\right) - 1 \right] b(x')e^{-i\xi \cdot x'} dx' = 0. \quad (4.3)$$

We fix $i, j \in \{1, \dots, n\}$ such that $i < j$ and we assume that $\xi \in \{\xi = (\xi_1, \dots, \xi_n) : \xi_i \neq 0\}$. We can choose

$$\eta = \frac{\xi_j e_i - \xi_i e_j}{\sqrt{\xi_i^2 + \xi_j^2}} \quad \text{and} \quad y = \frac{\xi_i e_i + \xi_j e_j}{\sqrt{\xi_i^2 + \xi_j^2}} \in \eta^\perp.$$

Here (e_1, \dots, e_n) is the canonical basis of \mathbb{R}^n defined by $e_1 = (1, 0, \dots, 0), \dots, e_n = (0, \dots, 0, 1)$. Then, (4.3) implies

$$\begin{aligned} &\int_{\eta^\perp} \left[\exp\left(i \int_{\mathbb{R}} \eta \cdot A(x' + s\eta) ds\right) - 1 \right] y \\ &\quad \cdot \nabla \left[\exp\left(-i \int_{\mathbb{R}} \eta \cdot A(x' + s\eta) ds\right) e^{-ix' \cdot \xi} \right] dx' = 0. \end{aligned}$$

Integrating by parts we get

$$\begin{aligned} &\frac{-i}{\sqrt{\xi_i^2 + \xi_j^2}} \cdot \int_{\mathbb{R}^n} (\xi_j y \cdot \nabla a_i(x) - \xi_i y \cdot \nabla a_j(x)) e^{-ix \cdot \xi} dx \\ &= -i \int_{\eta^\perp} \left(\int_{\mathbb{R}} \eta \cdot [y \cdot \nabla A](x' + s\eta) ds \right) e^{-ix' \cdot \xi} dx' \\ &= 0 \end{aligned}$$

with $A = (a_1, \dots, a_n)$. Integrating again by parts, we find

$$\begin{aligned} & \int_{\mathbb{R}^n} (\xi_j a_i - \xi_i a_j) e^{-ix \cdot \xi} dx \\ &= \frac{y \cdot \xi}{\sqrt{\xi_i^2 + \xi_j^2}} \int_{\mathbb{R}^n} (\xi_j a_i - \xi_i a_j) e^{-ix \cdot \xi} dx \\ &= \frac{-i}{\sqrt{\xi_i^2 + \xi_j^2}} \cdot \int_{\mathbb{R}^n} (\xi_j y \cdot \nabla a_i(x) - \xi_i y \cdot \nabla a_j(x)) e^{-ix \cdot \xi} dx \\ &= 0 \end{aligned}$$

and it follows that $\mathcal{F}[\partial_{x_j} a_i - \partial_{x_i} a_j](\xi) = 0$ for all $\xi \in \{\xi = (\xi_1, \dots, \xi_n) : \xi_i \neq 0\}$. On the other hand, since $\partial_{x_j} a_i - \partial_{x_i} a_j$ is compactly supported, $\mathcal{F}(\partial_{x_j} a_i - \partial_{x_i} a_j)(\xi)$ is continuous in $\xi \in \mathbb{R}^n$ and it follows $\mathcal{F}(\partial_{x_j} a_i - \partial_{x_i} a_j) = 0$ on \mathbb{R}^n . From this last result, we deduce that $\partial_{x_j} a_i - \partial_{x_i} a_j = 0$ which implies that $dA_1 = dA_2$. \square

Now assuming that $A_1 = A_2$, we show in the next lemma that (4.2) implies $V_1 = V_2$.

Lemma 4.2. *Let $\eta_1(\tau)$, $\eta_2(\tau)$ and $\lambda(\tau)$ be fixed by (3.16) and $b_2 = 1$. Assume that $A_1 = A_2$ and (4.2) is fulfilled. Then, we have $V_1 = V_2$.*

Proof. Fix $\xi \in \mathbb{R}^n \setminus \{0\}$ and choose $\eta \in \mathbb{S}^{n-1} \cap \xi^\perp$. Fix also $b = 1$. Thus, combining (3.27) and (4.2), we find

$$\int_{\mathbb{R}^n} V(x) e^{-ix \cdot \xi} dx = 0$$

with $V = V_1 - V_2$ extended by 0 outside of Ω . It follows that $V_1 = V_2$. \square

According to Lemma 4.1, 4.2, the proof of Theorem 1.1 will be completed if we show that conditions (1.2) imply conditions (4.1), (4.2). For this purpose, we adapt the approach of [16] to magnetic Schrödinger operators. Let $f \in H^{\frac{1}{2}}(\Gamma)$ being fixed, with the notations of Lemmas 2.1 and 2.3, we denote by $v_{j,\lambda,\mu} := u_{j,\lambda} - u_{j,\mu}$ the solution of (2.8) where V is replaced by V_j and A by A_j . We fix also $h_{j,k} := \partial_\nu \varphi_{j,k}|_\Gamma$ $\alpha_{j,k} := \langle f, h_{j,k} \rangle$. Recalling that in Lemma 2.4 we have set $z_\mu = u_{1,\mu} - u_{2,\mu}$, in a similar way to [16], writing the above identity for $j = 1$ and $j = 2$, applying (1.1) and then subtracting the resulting equations, we end up with a new relation, namely

$$\begin{aligned} (\partial_\nu + iA_1 \cdot \nu)u_{1,\lambda}|_\Gamma - (\partial_\nu + iA_2 \cdot \nu)u_{2,\lambda}|_\Gamma &= i(A_1 - A_2) \cdot \nu f + \partial_\nu u_{1,\lambda} - \partial_\nu u_{2,\lambda} \\ &= \partial_\nu z_\mu + \partial_\nu v_{1,\lambda,\mu} - \partial_\nu v_{2,\lambda,\mu}. \end{aligned} \tag{4.4}$$

Now let us set

$$F_j(\lambda, \mu, f) := \partial_\nu v_{j,\lambda,\mu|_\Gamma}, \quad j = 1, 2.$$

According to (2.7), we have

$$\begin{aligned} F(\lambda, \mu, f) &:= F_1(\lambda, \mu, f) - F_2(\lambda, \mu, f) \\ &= \sum_{k=1}^{+\infty} \left[\frac{(\mu - \lambda)\alpha_{1,k}}{(\lambda - \lambda_{1,k})(\mu - \lambda_{1,k})} h_{1,k} - \frac{(\mu - \lambda)\alpha_{2,k}}{(\lambda - \lambda_{2,k})(\mu - \lambda_{2,k})} h_{2,k} \right]. \end{aligned} \tag{4.5}$$

Consider the following intermediate results.

Lemma 4.3. *Let η_1, η_2, λ be given by (3.16). Consider $\Phi_j, j = 1, 2$, with Φ_1 introduced in the previous section and $\Phi_2 = e^{-i\sqrt{\lambda}\eta_2 \cdot x} b_2 e^{-i\psi_2}$, where b_2 is defined by (3.18) or $b_2 = 1$. Then, we have*

$$\sup_{\tau > 1} \sum_{k=1}^{\infty} \left| \frac{\langle \Phi_1, h_{j,k} \rangle}{\lambda_{j,k} - \lambda} \right|^2 < \infty, \quad \sup_{\tau > 1} \sum_{k=1}^{\infty} \left| \frac{\langle \overline{\Phi_2}, h_{2,k} \rangle}{\lambda_{2,k} - \lambda} \right|^2 < \infty, \quad j = 1, 2. \tag{4.6}$$

Proof. We start with the first estimate of (4.6) for $j = 1$. According to Lemma 2.1 the solution $u_{1,\lambda}$ of (2.2) for $f = \Phi_1, A = A_1$ and $V = V_1$, is given by

$$u_{1,\lambda} = \sum_{k=1}^{\infty} \frac{\langle \Phi_1, h_{1,k} \rangle}{\lambda - \lambda_{1,k}} \varphi_{1,k}.$$

Therefore,

$$\|u_{1,\lambda}\|_{L^2(\Omega)}^2 = \sum_{k=1}^{\infty} \left| \frac{\langle \Phi_1, h_{1,k} \rangle}{\lambda_{1,k} - \lambda} \right|^2. \tag{4.7}$$

On the other hand, in view of (3.8), we have

$$\|u_{1,\lambda}\|_{L^2(\Omega)} \leq \|\Phi_1\|_{L^2(\Omega)} + \left\| \sqrt{\lambda}(H_1 - \lambda)^{-1} \left[2\eta_1 \cdot (A_1 - A_{1,\#}) + \frac{q_{11}}{\sqrt{\lambda}} \right] \right\|_{L^2(\Omega)}.$$

Here q_{11} is the expression introduced in Lemma 3.1. Combining this with the fact that

$$\|\sqrt{\lambda}(H_1 - \lambda)^{-1}\|_{\mathcal{B}(L^2(\Omega))} \leq \frac{|\tau + i|}{|\Im \lambda|} = \frac{|\tau + i|}{2\tau} \leq 1$$

and the fact that, according to (3.19) and (3.20),

$$\lim_{\tau \rightarrow +\infty} \|\eta_1 \cdot (A_1 - A_{1,\#})\|_{L^\infty(\Omega)} = \lim_{\tau \rightarrow +\infty} \left\| \frac{q_{11}}{\sqrt{\lambda}} \right\|_{L^\infty(\Omega)} = 0$$

we deduce the first estimate of (4.6) for $j = 1$. In a same way, for $j = 2$ using the fact that according to (3.20) we have

$$(-i\nabla + A_2)^2\Phi_1 + V_2\Phi_1 - \lambda\Phi_1 = \underset{\tau \rightarrow +\infty}{\mathcal{O}}(\tau)$$

and repeating our previous arguments we deduce the first estimate (4.6) for $j = 2$. For the second estimate of (4.6), repeating the previous arguments we find

$$(-i\nabla + A_2)^2\overline{\Phi_2} + V_2\overline{\Phi_2} - \bar{\lambda}\overline{\Phi_2} = \overline{(-i\nabla + A_2)^2\Phi_2 + V_2\Phi_2 - \lambda\Phi_2} = \underset{\tau \rightarrow +\infty}{\mathcal{O}}(\tau).$$

Combining this estimate with the fact that

$$\left| \frac{\langle \overline{\Phi_2}, h_{2,k} \rangle}{\lambda_{2,k} - \lambda} \right| = \left| \frac{\langle \overline{\Phi_2}, h_{2,k} \rangle}{\lambda_{2,k} - \bar{\lambda}} \right|$$

since $\lambda_{2,k} \in \mathbb{R}$, we deduce the second estimate of (4.6) by repeating the above arguments. □

From now on we set

$$\begin{aligned} G(\lambda, \mu, \Phi_1, \Phi_2) &:= \langle F(\lambda, \mu, \Phi_1), \overline{\Phi_2} \rangle \\ &::= \sum_{k=1}^{+\infty} (\mu - \lambda) \left[\frac{\langle \Phi_1, h_{1,k} \rangle \langle h_{1,k}, \overline{\Phi_2} \rangle}{(\lambda - \lambda_{1,k})(\mu - \lambda_{1,k})} - \frac{\langle \Phi_1, h_{2,k} \rangle \langle h_{2,k}, \overline{\Phi_2} \rangle}{(\lambda - \lambda_{2,k})(\mu - \lambda_{2,k})} \right]. \end{aligned}$$

Combining estimates (4.6) with Lemma 4.3, 4.4, 4.5 of [16], we obtain the following.

Lemma 4.4. *Let the conditions of Theorem 1.1 be fulfilled and let η_1, η_2, λ be given by (3.16). Then, $G(\lambda, \mu, \Phi_1, \Phi_2)$ converge to $G_*(\lambda, \Phi_1, \Phi_2)$ as $\mu \rightarrow -\infty$ and $G_*(\lambda, \Phi_1, \Phi_2)$ converge to 0 as $\tau \rightarrow +\infty$. Here we consider both the case b_2 given by (3.18) and the case $b_2 = 1$.*

Armed with Lemma 4.4, we are now in position to complete the proof of Theorem 1.1.

Proof of Theorem 1.1. Note first that according (4.4), for

$$M = \|V_1\|_{L^\infty(\Omega)} + \|V_2\|_{L^\infty(\Omega)},$$

we have

$$S_1(\lambda, \eta_1, \eta_2) - S_2(\lambda, \eta_1, \eta_2) = \langle \partial_v z_\mu, e^{i\sqrt{\lambda}\eta_2 \cdot x} \overline{b_2} e^{i\psi_2} \rangle + G(\lambda, \mu, \Phi_1, \Phi_2),$$

for $\mu \in (-\infty, -M)$, where λ, η_1, η_2 are fixed by (3.16), b_2 is given by (3.18) or $b_2 = 1$ and $z_\mu = u_{1,\mu} - u_{2,\mu}$ with $u_{j,\mu}, j = 1, 2$, the solution of (2.8) where λ is replaced by μ, V by V_j, A by A_j and f by Φ_1 . In view of Lemma 2.4 and Lemma 4.4, sending $\mu \rightarrow -\infty$ we get

$$S_1(\lambda, \eta_1, \eta_2) - S_2(\lambda, \eta_1, \eta_2) = G_*(\lambda, \Phi_1, \Phi_2).$$

Then, in view of Lemma 4.4, conditions (4.1) and (4.2) are fulfilled and according to Lemma 4.1 we have $dA_1 = dA_2$. Therefore, condition (1.1) implies that for $A = A_2 - A_1$ extended by 0 outside of Ω we have $dA = 0$ on \mathbb{R}^n . Thus, there exists $p \in W^{2,\infty}(\mathbb{R}^n)$ given by

$$p(x) = \int_0^1 x \cdot A(tx) dt$$

such that $A = \nabla p$ on \mathbb{R}^n . Since $\mathbb{R}^n \setminus \Omega$ is connected, applying the fact that $A = 0$ on $\mathbb{R}^n \setminus \Omega$, upon eventually subtracting a constant we may assume that $p|_{\mathbb{R}^n \setminus \Omega} = 0$ which implies that $p|_\Gamma = 0$. Now let us consider the operator $H_3 = (-i\nabla + A_1) + V_2$ acting on $L^2(\Omega)$ with Dirichlet boundary condition and let $(\lambda_{3,k}, \varphi_{3,k})_{k \geq 1}$ be a sequence of eigenvalues and eigenfunctions of H_3 . Since $A_1 = A_2 - \nabla p$ one can check that $H_3 = e^{ip} H_2 e^{-ip}$. From this identity we deduce that

$$\lambda_{3,k} = \lambda_{2,k}, \quad k \geq 1.$$

Moreover, for all $k \geq 1$ we can choose $\varphi_{3,k} = e^{ip} \varphi_{2,k}$ and deduce that the condition

$$\partial_\nu \varphi_{3,k} = \partial_\nu \varphi_{2,k}, \quad k \geq 1$$

is also fulfilled. Thus, conditions (1.2) imply that

$$\lim_{k \rightarrow +\infty} |\lambda_{1,k} - \lambda_{3,k}| = 0 \quad \text{and} \quad \sum_{k=1}^{+\infty} \|\partial_\nu \varphi_{1,k} - \partial_\nu \varphi_{3,k}\|_{L^2(\Gamma)}^2 < \infty.$$

Then, repeating the arguments of Lemma 4.4 we obtain

$$\lim_{\tau \rightarrow +\infty} \tilde{S}_1(\lambda(\tau), \eta_1(\tau), \eta_2(\tau)) - \tilde{S}_3(\lambda(\tau), \eta_1(\tau), \eta_2(\tau)) = 0,$$

where

$$\tilde{S}_j(\lambda, \eta_1, \eta_2) = \langle \Lambda_{j,\lambda} \Phi_1, e^{i\sqrt{\lambda} \eta_2 \cdot x} e^{i\tilde{\psi}_2} \rangle, \quad j = 1, 3$$

with

$$\tilde{\psi}_2(x) = \int_{-x}^{x \cdot \eta_2} \eta_2 \cdot A_{1,\#}(x + (s - x \cdot \eta_2)\eta_2) ds, \quad b_2 = 1$$

and $\Lambda_{3,\lambda}$ the Dirichlet-to-Neumann map associated to problem (2.2) for $A = A_1$ and $V = V_2$. Then, in view of Lemma 4.2 we have $V_1 = V_2$. This completes the proof of Theorem 1.1. □

Acknowledgements. The author would like to thank the anonymous referee for valuable comments and helpful remarks. The author is grateful to Otared Kavian and Eric Soccorsi for their remarks and suggestions.

References

- [1] V. A. Ambartsumian, Über eine Frage der Eigenwerttheorie. *Z. Phys.* **53** (1929), 690–695. [JFM 55.0868.01](#)
- [2] M. Belishev and Y. Kurylev, To the reconstruction of a Riemannian manifold via its spectral data (BC-method). *Comm. Partial Differential Equations* **17** (1992), no. 5–6, 767–804. [MR 1177292 Zbl 0812.58094](#)
- [3] M. Bellassoued and M. Choulli, Stability estimate for an inverse problem for the magnetic Schrödinger equation from the Dirichlet–to–Neumann map. *J. Funct. Anal.* **258** (2010), no. 1, 161–195. [MR 2557958 Zbl 1180.35436](#)
- [4] M. Bellassoued, M. Choulli, and M. Yamamoto, Stability estimate for an inverse wave equation and a multidimensional Borg–Levinson theorem. *J. Differential Equations* **247** (2009), no. 2, 465–494. [MR 2523687 Zbl 1171.35123](#)
- [5] M. Bellassoued and D. Dos Santos Ferreira, Stability estimates for the anisotropic wave equation from the Dirichlet–to–Neumann map. *Inverse Probl. Imaging* **5** (2011), no. 4, 745–773. [MR 2852371 Zbl 1250.58012](#)
- [6] G. Borg, Eine Umkehrung der Sturm–Liouvilleschen Eigenwertaufgabe. *Acta Math.* **78** (1946), 1–96. [MR 0015185 Zbl 0063.00523](#)
- [7] B. Canuto and O. Kavian, Determining coefficients in a class of heat equations via boundary measurements. *SIAM J. Math. Anal.* **32** (2001), no. 5, 963–986. [MR 1828313 Zbl 0981.35096](#)
- [8] B. Canuto and O. Kavian, Determining two coefficients in elliptic operators via boundary spectral data: a uniqueness result. *Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8)* **7** (2004), no. 1, 207–230. [MR 2044267 Zbl 1178.35152](#)
- [9] M. Choulli, *Une introduction aux problèmes inverses elliptiques et paraboliques*. Mathématiques & Applications (Berlin), 65. Springer-Verlag, Berlin, 2009. [MR 2554831 Zbl 1192.35187](#)
- [10] M. Choulli and P. Stefanov, Stability for the multi-dimensional Borg–Levinson theorem with partial spectral data. *Comm. Partial Differential Equations* **38** (2013), no. 3, 455–476. [MR 3019446 Zbl 1302.35433](#)
- [11] D. Dos Santos Ferreira, C. Kenig, J. Sjöstrand, and G. Uhlmann, Determining a magnetic Schrödinger operator from partial Cauchy data. *Comm. Math. Phys.* **271** (2007), no. 2, 467–488. [MR 2287913 Zbl 1148.35096](#)

- [12] I. M. Gel'fand and B. M. Levitan, On the determination of a differential equation from its spectral function. *Izvestiya Akad. Nauk SSSR. Ser. Mat.* **15** (1951). 309–360. In Russian. English translation, *Amer. Math. Soc. Transl. (2)* **1** (1955), 253–304. [MR 0045281](#) [MR 0073805](#) (transl.) [Zbl 0044.09301](#) [Zbl 0066.33603](#) (transl.)
- [13] P. Grisvard, *Elliptic problems in nonsmooth domains*. Monographs and Studies in Mathematics, 24. Pitman (Advanced Publishing Program), Boston, MA, 1985. [MR 0775683](#) [Zbl 0695.35060](#)
- [14] H. Isozaki, Some remarks on the multi-dimensional Borg–Levinson theorem. *J. Math. Kyoto Univ.* **31** (1991), no. 3, 743–753. [MR 1127097](#) [Zbl 0753.35121](#)
- [15] A. Katchalov and Y. Kurylev, Multidimensional inverse problem with incomplete boundary spectral data. *Comm. Partial Differential Equations* **23** (1998), no. 1-2, 55–95. [MR 1608496](#) [Zbl 0904.65114](#)
- [16] O. Kavian, Y. Kian, and E. Soccorsi, Uniqueness and stability results for an inverse spectral problem in a periodic waveguide. *J. Math. Pures Appl. (9)* **104** (2015), no. 6, 1160–1189. [MR 3420415](#) [Zbl 1335.35300](#)
- [17] K. Krupchyk, M. Lassas, and G. Uhlmann, Inverse problems with partial data for a magnetic Schrödinger operator in an infinite slab and on a bounded domain. *Comm. Math. Phys.* **312** (2012), no. 1, 87–126. [MR 2914058](#) [Zbl 1238.35188](#)
- [18] K. Krupchyk and G. Uhlmann, Uniqueness in an inverse boundary problem for a magnetic Schrodinger operator with a bounded magnetic potential. *Comm. Math. Phys.* **327** (2014), no. 3, 993–1009. [MR 3192056](#) [Zbl 1295.35366](#)
- [19] N. Levinson, The inverse Sturm–Liouville problem. *Mat. Tidsskr. B.* **1949** (1949). 25–30. [MR 0032067](#) [Zbl 0041.42310](#) [Zbl 0045.36402](#)
- [20] J.-L. Lions and E. Magenes, *Problèmes aux limites non homogènes et applications*. Vol. 1. Travaux et Recherches Mathématiques, 17. Dunod, Paris, 1968. [MR 0247243](#) [Zbl 0165.10801](#)
- [21] C. Montalto, Stable determination of a simple metric, a co-vector field and a potential from the hyperbolic Dirichlet–to–Neumann map. *Comm. Partial Differential Equations* **39** (2014), no. 1, 120–145. [MR 3169781](#) [Zbl 1302.35234](#)
- [22] A. Nachman, J. Sylvester, and G. Uhlmann, An n -dimensional Borg–Levinson theorem. *Comm. Math. Phys.* **115** (1988), no. 4, 595–605. [MR 0933457](#) [Zbl 0644.35095](#)
- [23] G. Nakamura, Z. Sun, and G. Uhlmann, Global identifiability for an inverse problem for the Schrödinger equation in a magnetic field. *Math. Ann.* **303** (1995), no. 3, 377–388. [MR 1354996](#) [Zbl 0843.35134](#)
- [24] F. Nicoleau, A stationary approach to inverse scattering for Schrödinger operators with first order perturbation. *Comm. Partial Differential Equations* **22** (1997), no. 3-4, 527–553. [MR 1443048](#) [Zbl 0882.35086](#)
- [25] N. G. Novikov, Multidimensional inverse spectral problems for the equation $-\Delta\psi + (v(x) - Eu(x))\psi = 0$. *Funktsional. Anal. i Prilozhen.* **22** (1988), no. 4, 11–22, 96. In Russian. English translation, *Funct. Anal. Appl.* **22** (1988), no. 4, 263–272. [MR 0976992](#) [Zbl 0689.35098](#)

- [26] M. Salo, Inverse problems for nonsmooth first order perturbations of the Laplacian. *Ann. Acad. Sci. Fenn. Math. Diss.* **139** (2004), 67 pp. [MR 2105191](#) [Zbl 1059.35175](#)
- [27] V. S. Serov, Borg–Levinson theorem for magnetic Schrödinger operator. *Bull. Greek Math. Soc.* **57** (2010), 321–332. [MR 3617885](#)
- [28] P. Stefanov and G. Uhlmann, Stability estimates for the hyperbolic Dirichlet to Neumann map in anisotropic media. *J. Funct. Anal.* **154** (1998), no. 2, 330–358. [MR 1612709](#) [Zbl 0915.35066](#)
- [29] P. Stefanov and G. Uhlmann, Stable determination of generic simple metrics from the hyperbolic Dirichlet–to–Neumann map. *Int. Math. Res. Not.* **2005**, no. 17, 1047–1061. [MR 2145709](#) [Zbl 1088.53027](#)
- [30] Z. Sun, An inverse boundary value problem for the Schrödinger operator with vector potentials. *Comm. Partial Differential Equations* **18** (1993), no. 1-2, 83–124. [MR 1211726](#) [Zbl 0781.35073](#)
- [31] J. Sylvester and G. Uhlmann, A global uniqueness theorem for an inverse boundary value problem. *Ann. of Math. (2)* **125** (1987), no. 1, 153–169.
- [32] L. Tzou, Stability estimates for coefficients of magnetic Schrödinger equation from full and partial boundary measurements. *Comm. Partial Differential Equations* **33** (2008), no. 10-12, 1911–1952. [MR 2475324](#) [Zbl 1157.35108](#)

Received October 13, 2015

Yavar Kian, Aix Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France
e-mail: yavar.kian@univ-amu.fr