

Anderson localisation for infinitely many interacting particles in Hartree–Fock theory

Raphael Ducatez

Abstract. We prove the occurrence of Anderson localisation for a system of infinitely many particles interacting with a short range potential, within the ground state Hartree–Fock approximation. We assume that the particles hop on a discrete lattice and that they are submitted to an external periodic potential which creates a gap in the non-interacting one particle Hamiltonian. We also assume that the interaction is weak enough to preserve a gap. We prove that the mean-field operator has exponentially localised eigenvectors, either on its whole spectrum or at the edges of its bands, depending on the strength of the disorder.

Mathematics Subject Classification (2010). 47A10, 81Q10, 81V70.

Keywords. Anderson localisation, Hartree–Fock theory, multiscale analysis.

Contents

1	Introduction	1020
2	Model and results	1021
3	Construction of the mean-field Hamiltonian: proof of Theorem 2.1	1026
4	Local influence: proof of Theorem 2.3	1027
5	Wegner estimate: proof of Theorem 2.4	1031
6	Multiscale analysis	1035
7	Numerical simulations	1044
	References	1047

1. Introduction

In 1958, the physicist P.W. Anderson predicted that, in a random medium, diffusion could disappear and waves stay localised [3]. Since then, *Anderson localisation* has played an important role to explain several properties of physical systems.

There are many works on the mathematical side, starting with the simpler one dimension case in the beginning of the 80s. In dimension 3, the first proof of Anderson localisation was provided by Fröhlich and Spencer [18] who developed a method now called *multiscale analysis*. There are now other approaches which include for instance the *fractional moment method* proposed by Aizenman and Molchanov [1] or the techniques proposed by Imbrie [20].

All these works are restricted to the one wave problem, which is well adapted to optics and acoustics. In condensed matter, the interaction between the particles is expected to play an important role and in this case one speaks of *many-body localisation*. This question is not fully understood and is very actively studied in physics [8, 9, 2, 19]. A typical system of interest is a crystal composed of quantum electrons and classical nuclei placed on a random perturbation of a perfect lattice [11, 24]. This is a very complicated system since the Coulomb interaction is long range and there are of the order of 10^{23} particles, usually mathematically treated as an infinite number. Due to screening effects, the Coulomb potential is often replaced by an effective short range interaction and this is what we are going to do in this work.

Finite interacting systems have been recently considered in several works [15, 2, 21, 17] but infinite systems have not been studied thoroughly. One should however mention a series of works on superfluidity and Bose-Einstein condensation in the Lieb–Liniger 1D Bose gas in a strong random potential [31, 22], and the very recent article of Seiringer and Warzel [30] on the 1D Tonks-Girardeau gas.

It is very hard to deal with the exact interacting Schrödinger problem for an infinite system. A useful and widespread approximation is the Hartree–Fock model, where the particles are treated as independent objects but see a field which depends on their own states and is then self-consistently optimised. For random systems this model has been recently introduced by E. Cancès, S. Lahbabi and M. Lewin in [13]. The authors were able to construct a solution of the random nonlinear Hartree–Fock equation for an infinite system of fermions with short range interaction, but the phenomenon of Anderson localisation was only investigated numerically in [23]. We should also mention the paper of Chen and Rodnianski [14], where the Boltzmann limit of a similar model is studied with a dynamical approach.

The Hartree Fock model and its generalisation including the pairing term are widely used in condensed matter physics [32, 34, 10, 7]. Understanding the interplay between disorder and interactions is indeed a great challenge [29]. For 1D disordered photonic lattices this was for instance studied in [25, 28].

In this work, we complete the program initiated in [13] in the case of a discrete system with a periodic background. In short, we show that the unique solution of the Hartree–Fock equation

$$\begin{cases} H_{\min} = -\Delta + V_0 + V_\omega + W * \gamma(x, x) - W(x - y)\gamma(x, y), \\ \gamma = \mathbb{1}_{<\mu}(H_{\min}), \end{cases} \quad (1)$$

provides a mean-field operator H_{\min} which has exponentially localised eigenvectors, either on its whole spectrum or at the edges of its bands. Our main assumption is that the periodic potential V_0 is sufficiently strong to create a gap for $-\Delta + V_0$ and that the random part V_ω as well as the interaction W do not alter this gap. We take the chemical potential μ in this gap. Our argument is to adapt the well known multiscale analysis in order to include the nonlinear terms.

The paper is organised as follows. In the next section, we properly define the discrete Hartree–Fock model and state our main localisation results. Sections 3 to 6 are devoted to the proof of our results. Then in section 7 we present some simple one dimensional numerical simulations in order to illustrate our findings. We also look at some cases which are not covered by our theorems and find, for instance, an interesting delocalisation phenomenon at the Fermi level in the absence of a gap, which deserves further investigations.

2. Model and results

We consider a system of fermions on a subset Λ of the lattice \mathbb{Z}^d and which are submitted to an external potential V . Without interactions the system is described by the one-particle Hamiltonian

$$H^\Lambda = -\Delta + V. \quad (2)$$

In the canonical basis $(\delta_x)_{x \in \Lambda}$ of $\ell^2(\Lambda)$, the operator H^Λ is defined by

$$(\delta_x, H^\Lambda \delta_y) = \begin{cases} V(x) & \text{if } x = y, \\ 1 & \text{if } |x - y| = 1, \\ 0 & \text{otherwise,} \end{cases} \quad \text{for all } x, y \in \Lambda. \quad (3)$$

When Λ is the whole lattice \mathbb{Z}^d we will use the simpler notation $H = H^{\mathbb{Z}^d}$. Our theorems will actually hold for finite as well as for infinite domains. Since H will later be perturbed by a non linear term describing the interactions between the particles, we will often call H the linear part.

The potential V describes a crystal lattice which is randomly perturbed. It is therefore assumed to be of the form

$$V = V_0 + V_\omega, \quad (4)$$

where V_0 is a periodic potential (of an arbitrary period) and V_ω is a random potential. We use the Anderson tight binding model where the value of V_ω is chosen independently at each site of Λ with the same random probability law \mathbb{P} :

$$V_\omega = \sum_{i \in \Lambda} v_i(\omega) \delta_i. \quad (5)$$

Here the v_i are iid random variables.

Under suitable regularity assumptions on \mathbb{P} , it is well known that H displays Anderson localisation. This means that there exist small intervals at the edges of the spectral bands where the spectrum is pure point with exponentially decaying eigenvectors (“Lifshitz tail”). Moreover, if the random potential V_ω is strong enough (compared to $-\Delta$) then the whole spectrum is pure point with localised eigenvectors.

We will show that if the interactions between the particles are small enough then the same holds for the interacting model. Before introducing interactions, we first discuss the precise assumptions that we will use for \mathbb{P} and V_0 .

(A1) Regularity of \mathbb{P} . We assume that \mathbb{P} has a bounded support and that it has a density ρ with respect to the Lebesgue measure. Moreover we assume that there exist $d_1 < d_2 < \dots < d_k$ such that $\text{Supp}(\mathbb{P}) \subset [d_1, d_k]$ and ρ is Lipschitz on the intervals $]d_i, d_i + 1[$, $i = 1, \dots, (k - 1)$.

We think that this assumption can be weakened in several possible ways but we will not discuss this for the sake of simplicity. With no loss of generality, we can assume that $0 \in \text{Supp}(\mathbb{P})$.

(A2) Gap. We assume that H has a gap $[a, b]$ in its (deterministic) spectrum.

This assumption implies that the periodic potential V_0 is not constant and strong enough. Indeed the spectrum of H is almost surely equal to

$$\sigma(H) = \sigma(-\Delta + V_0) + \text{supp}(\mathbb{P}),$$

when the support of \mathbb{P} is an interval. In particular it is enough to assume that $-\Delta + V_0$ has a gap of size $G_0 > 2|\text{supp}(\mathbb{P})|$. When the domain Λ is large enough then H^Λ will have a gap as well. We call the size of the gap $G = b - a$, and choose a chemical potential μ in the middle $\mu = (a + b)/2$.

Now we turn to the definition of the interacting model. We assume that the interaction is translation-invariant and decays fast enough.

(A3) Short range interaction. We assume that there exists $\nu > 0$ such that

$$|W(x - y)| \leq C e^{-\nu|x-y|}. \quad (6)$$

In the Hartree–Fock model (see for example [4, 6, 26, 27, 33, 5]) the system in the domain Λ is completely described by its one-particle density matrix which is an orthogonal projection γ on $\ell^2(\Lambda)$. More precisely, for a finite system, the rank- N projection $\gamma = \sum_{i=1}^N |\phi_i\rangle\langle\phi_i|$ corresponds to the N -particle wave function Ψ ,

$$\Psi(x_1, x_2, \dots, x_N) = \frac{1}{\sqrt{N!}} \det(\phi_i(x_j)), \quad (7)$$

called a Slater determinant. In a finite domain Λ , the many-body energy of this state is

$$\begin{aligned} & \left(\Psi, \left[\sum_i^N -\Delta_i + V(x_i) + \frac{1}{2} \sum_{i \neq j} W(x_i - x_j) \right] \Psi \right) \\ &= \text{Tr}(H^\Lambda \gamma) + \frac{1}{2} \sum_{x, y \in \Lambda} W(x - y) \gamma(x, x) \gamma(y, y) \\ & \quad - \frac{1}{2} \sum_{x, y \in \Lambda} W(x - y) |\gamma(x, y)|^2. \end{aligned} \quad (8)$$

We define the effective interaction A_{eff} by

$$A_{\text{eff}}(\gamma)(x, y) = \left[\sum_n W(n - y) \gamma(n, n) \delta_{x=y} \right] - W(x - y) \gamma(x, y) \quad (9)$$

for any γ positive bounded operator. It is the derivative of the interaction part of the energy.

Any minimiser γ of the energy, with fixed rank N , is a solution of the non-linear equation which involve an effective Hamiltonian:

$$\begin{cases} H_{\min} = -\Delta + V + A_{\text{eff}}(\gamma), \\ \gamma = \mathbb{1}_{<\mu'}(H_{\min}), \end{cases} \quad (10)$$

In other words γ is the projection on the N first eigenfunctions of H_{\min} which itself depends of γ . To be more precise the equation (10) hold under the condition that $\lambda_{N+1}(H_{\min}) > \lambda_N(H_{\min})$ which is known to be automatically satisfied when $W > 0$, see [5, 6].

We will first show that our problem is well defined for a bounded domain Λ as well as for the infinite domain $\Lambda = \mathbb{Z}^d$.

Theorem 2.1 (HF ground states for infinitely many particles). *Let V' be a bounded function such that $-\Delta + V'$, defined on a subset Λ of \mathbb{Z}^d , has a gap $[a', b']$ in its spectrum, with $G' = b' - a'$ and $\mu' = (b' + a')/2$. If $\|W\|_{\ell^1} < G'/6$, then there exists a unique solution of the system $\gamma = \mathbb{1}_{\leq \mu'}(-\Delta + V' + A_{\text{eff}}(\gamma))$. If Λ is finite, the trace is preserved:*

$$\text{Tr}(\mathbb{1}_{\leq \mu'}(-\Delta + V' + A_{\text{eff}}(\gamma))) = \text{Tr}(\mathbb{1}_{\leq \mu'}(-\Delta + V'))$$

and, because of uniqueness, our solution is as well the unique minimiser of the energy among all Hartree–Fock states with fixed particles number $N = \text{Tr}(\mathbb{1}_{\leq \mu'}(-\Delta + V'))$.

This result will be shown in Section 3. The continuous case is more complicated and has been studied in [24]. When Λ is a large cube and $V' = V = V_0 + V_\omega$ then the number of particles $\text{Tr}(\mathbb{1}_{\leq \mu'}(-\Delta + V'))$ is proportional to the volume, which is the interesting physical case. For $\Lambda = \mathbb{Z}^d$, $H_{\min}(\omega)$ exists and its spectrum, as in the linear model, does not depend on ω almost surely. The reason is that H_{\min} is stationary with respect to space translations, which follows from the uniqueness in the theorem.

The main result of our paper is the following:

Theorem 2.2 (Anderson localisation in Hartree–Fock theory). *Under the assumptions (A1), (A2), and (A3), then the following holds:*

- (1) *there is ϵ such that if $\|W\|_{\ell^1} < \epsilon$ then there are small intervals at the edges of the bands of the spectrum of H_{\min} , where the spectrum is pure point with exponentially decaying eigenvectors;*
- (2) *there exists ϵ which only depends on C and ν in (6) such that if $\|\rho\|_\infty + \|\rho'\|_\infty < \epsilon$ and $\|W\|_{\ell^1} < G/6$, then the whole spectrum of H_{\min} is pure point and its eigenvectors are exponentially decaying in space.*

We remark that because of (A2), the second part of the statement can only be applied in cases where V_0 is also very large. This theorem will be shown by using the multiscale analysis on H_{\min} . We will proceed in three steps.

Step 1. We show that because of the gap, γ_{\min} only depends locally on the potential. From a practical point of view, in order to know how γ_{\min} looks like in a box of size L after solving the minimising problem for \mathbb{Z}^d , solving the minimising problem for a box of size $2L$ will be enough to have a very good approximation. In a more mathematical formulation:

Theorem 2.3 (locality). *We assume that (A2) and (A3) hold. There exist $a > 0$, $\nu > 0$ and $C > 0$ such that, if $\|W\|_{\ell^1} < aG$, then for any modification of the potential $V_\omega \rightarrow V_\omega + \delta V$ so that $V_\omega + \delta V \in \text{supp}(\mathbb{P})$, the change induced to the minimising projector given by Theorem 2.1 satisfies*

$$\sup_{y_2 \in \Lambda} |\gamma_{\min}(V + \delta V)(y_1, y_2) - \gamma_{\min}(V)(y_1, y_2)| \leq \|\delta V\| C e^{-\nu d(y_1, \text{supp}(\delta V))}. \quad (11)$$

Here $d(y_1, \text{supp}(\delta V))$ is the distance between y_1 and the support of the perturbation δV . The constants do not depend on the choice of the domain Λ .

This theorem will be proved in Section 4.

Step 2. We then show in section 5 a kind of Wegner estimate. We denote by $(H_{\min})|_\Lambda$ the submatrices $\mathbf{1}_\Lambda H_{\min} \mathbf{1}_\Lambda$, restricted to $\ell^2(\Lambda)$ where Λ is a finite cube in \mathbb{Z}^d .

Theorem 2.4 (Wegner estimate). *Assuming (A1), (A2), (A3), there exists $a > 0$ such that if $\|W\|_{\ell^1} < aG$ then there exists a constant C so that*

$$\begin{aligned} \mathbb{P}[d(\sigma[(H_{\min})|_\Lambda], \lambda) < \epsilon] \\ \leq C |\Lambda| \sqrt{\epsilon} (|\text{Supp}(\rho)|^{-1/2} + \|\rho\|_\infty |\text{Supp}(\rho)|^{1/2} + \|\rho'\|_\infty |\text{Supp}(\rho)|^{3/2}), \end{aligned} \quad (12)$$

for any $\lambda \in \mathbb{C}$.

This result says that there is no arbitrary small interval where we can find for sure an eigenvalue of $(H_{\min})|_\Lambda$.

Step 3. In the last step we perform the multiscale analysis. This is explained in Section 6.

3. Construction of the mean-field Hamiltonian: proof of Theorem 2.1

The aim of this section is to prove that our operator H_{\min} is well defined. We will show that finding the unique solution γ_{\min} in presence of a gap can be done using a fixed point lemma.

In this subsection we solve our system

$$\begin{cases} H_{\min} = -\Delta + V' + A_{\text{eff}}(\gamma), \\ \gamma = \mathbb{1}_{<\mu'}(H_{\min}), \end{cases} \quad (13)$$

under the assumption that μ' is inside a gap $[a', b']$ in the spectrum, where $\mu' = (a' + b')/2$. We introduce $G' = b' - a'$.

Let \mathcal{C} be a loop in the complex plane surrounding a part I of the spectrum of an operator H . We make the assumption that the loop does not cross the spectrum (which implies that there exist gaps above and below I). Then

$$\mathbb{1}_I(H) = \frac{1}{2i\pi} \oint_{\mathcal{C}} (H - z)^{-1} dz \quad (14)$$

is the projector on the spectral subspace associated with I .

Let us define an application that gives this projector.

Definition 3.1 (Fixed point map). Let \mathcal{C} be a fixed loop in the complex plane. For all γ orthogonal projector and $H_{\text{eff}}(\gamma) = -\Delta + V' + A_{\text{eff}}(\gamma)$, we define

$$F(\gamma) = \frac{1}{2i\pi} \oint_{\mathcal{C}} (H_{\text{eff}}(\gamma) - z)^{-1} dz. \quad (15)$$

This application enable us to reformulate our system (13) as

$$F(\gamma) = \gamma, \quad (16)$$

where the loop \mathcal{C} crosses the real axis at μ' . Recall that H_{\min} is bounded so that we can always enclose all of its spectrum below μ' . Because $\|A_{\text{eff}}\|$ is bounded by $2\|W\|_{\ell^1}$, we always have

$$d(\sigma(H_{\text{eff}}(\gamma)), \mu') \geq \frac{b' - a'}{2} - 2\|W\|_{\ell^1}. \quad (17)$$

So $(b' - a')/2 > 2\|W\|_{\ell^1}$ is enough to ensure that \mathcal{C} never crosses the spectrum of H_{eff} and F is always well defined. In order to solve (16) we will show that if $\|W\|_{\ell^1}$ is small enough then F is a contraction.

Proof of Theorem 2.1. Let γ_1 and γ_2 be two orthogonal projectors. Then we have

$$\begin{aligned} & F(\gamma_1) - F(\gamma_2) \\ &= -\frac{1}{2i\pi} \oint_{\mathcal{C}} (H_{\text{eff}}(\gamma_2) - z)^{-1} (H_{\text{eff}}(\gamma_1) - H_{\text{eff}}(\gamma_2)) (H_{\text{eff}}(\gamma_1) - z)^{-1} dz \\ &= -\frac{1}{2i\pi} \oint_{\mathcal{C}} (H_{\text{eff}}(\gamma_2) - z)^{-1} (A_{\text{eff}}(\gamma_1) - A_{\text{eff}}(\gamma_2)) (H_{\text{eff}}(\gamma_1) - z)^{-1} dz. \end{aligned}$$

The map F does not depend on the choice of the surrounding loop provided it encloses the appropriate part of the spectrum. Expanding it continuously to infinity, we can replace it in this formula by $\mathcal{C} = \mu' + i\mathbb{R}$ and write $z = \mu' + is$. We estimate $A_{\text{eff}}(\gamma_1) - A_{\text{eff}}(\gamma_2) = A_{\text{eff}}(\gamma_1 - \gamma_2)$ with $2\|W\|_{\ell^1}\|\gamma_1 - \gamma_2\|$ and $(H_{\text{eff}}(\gamma_2) - \mu' + is)^{-1} \leq ((G'/2 - 2\|W\|_{\ell^1})^2 + s^2)^{-1/2}$. Therefore

$$\|F(\gamma_1) - F(\gamma_2)\| \leq \frac{\|W\|_{\ell^1}\|\gamma_1 - \gamma_2\|}{\pi} \int_{\mathbb{R}} ((G'/2 - 2\|W\|_{\ell^1})^2 + s^2)^{-1} ds, \quad (18)$$

so that

$$\|F(\gamma_1) - F(\gamma_2)\| \leq \frac{\|W\|_{\ell^1}\|\gamma_1 - \gamma_2\|}{(G'/2 - 2\|W\|_{\ell^1})}. \quad (19)$$

Now, if $\|W\|_{L^1}$ is smaller than $G'/6$, then F is contracting, so it has a unique fixed point. \square

This concludes the first section, we have shown that in the presence of a gap, (13) has always a unique solution.

4. Local influence: proof of Theorem 2.3

The aim of this section is to show that under hypothesis (A2) and (A3) the random potential in a domain Λ_2 will only have a very small influence on A_{eff} in Λ_1 if Λ_2 is far enough from Λ_1 . This implies a weak form of independence between the submatrices $(H_{\min})_{|\Lambda_1}$ and $(H_{\min})_{|\Lambda_2}$ which is necessary for the multiscale analysis. The key tool is a Combes–Thomas estimate.

4.1. Combes–Thomas estimate. Because we want to use it for more general operators than just the Laplacian, we have written again the details of the proof.

Definition 4.1 (exponential off-diagonal decay operator). We will say that an operator K on $L^2(\Lambda)$ has exponential off-diagonal decay if there exist a rate $M > 0$, and a constant C so that $(\delta_x, K\delta_y) \leq C \exp(-M|x - y|)$ for all $x, y \in \Lambda$.

In our case, we note that $-\Delta + V + A_{\text{eff}}$ is an exponential decay operator if $W(x - y)$ decays exponentially.

Lemma 4.1 (Combes–Thomas estimate). *Let K be an exponential off-diagonal decay operator and Σ be its spectrum. Let $\lambda \in \mathbb{C}$ so that $d(\lambda, \Sigma) > 0$. Then there exist $\nu > 0$ and $C > 0$ such that*

$$(\delta_x, (K - \lambda)^{-1} \delta_y) \leq C e^{-\nu|x-y|}. \tag{20}$$

Proof. Let $f(z) := e^{-\nu|x-z|}$ we have

$$\begin{aligned} (\delta_x, (K - \lambda)^{-1} \delta_y) &= (\delta_x, f(z)^{-1} f(z) (K - \lambda)^{-1} f(z)^{-1} f(z) \delta_y) \\ &= f(x)^{-1} f(y) (\delta_x, f(z) (K - \lambda)^{-1} f(z)^{-1} \delta_y) \\ &= e^{-\nu|x-y|} (\delta_x, (f(z) K f(z)^{-1} - \lambda)^{-1} \delta_y). \end{aligned} \tag{21}$$

Then we have

$$((f(z) K f(z)^{-1} - K)u)(n) = \sum_{m \in \Lambda} (e^{-\nu(|x-n|-|x-m|)} - 1) K(m, n) u(m), \tag{22}$$

so

$$\|(f(z) K f(z)^{-1} - K)u\| \leq C \sum_{m \in \Lambda} (e^{\nu(|n-m|)} - 1) e^{-M(n-m)} |u(m)| < \infty,$$

and therefore

$$\|(f(z) K f(z)^{-1} - H)\| \leq C \|(e^{\nu|x|} - 1) e^{-M|x|}\|_{\ell^1}.$$

Because of the dominated convergence theorem, this converges to 0 with ν going to 0. So there exist $\nu > 0$ so that $\|(f(z) K f(z)^{-1} - H)\| < d(\lambda, \Sigma)$ and

$$d(\lambda, \Sigma(f(z) K f(z)^{-1})) \geq d(\lambda, \Sigma) - \|f(z) K f(z)^{-1} - H\| > 0.$$

So there exists C' such that

$$\|(f(z) K f(z)^{-1} - \lambda)^{-1}\| \leq \frac{1}{d(\lambda, \Sigma(f(z) K f(z)^{-1}))} = C'$$

and using (21) we find

$$(\delta_x, (K - \lambda)^{-1} \delta_y) \leq C' e^{-\nu|x-y|}. \quad \square$$

4.2. Local influence. We prove here Theorem 2.3. We use again the map defined in Section 3.

$$F_V(\gamma) = \frac{1}{2i\pi} \oint_{\mathcal{C}} (-\Delta + V + A_{\text{eff}}(\gamma) - z)^{-1} dz, \tag{23}$$

where \mathcal{C} is the loop enclosing the whole part of the spectrum below the middle of the gap μ . We will denote by $\gamma_{\min}(V)$ the solution of the system given by Theorem 2.1 and recall that $F_V(\gamma_{\min}(V)) = \gamma_{\min}(V)$.

Proof of Theorem 2.3. Because we can write $\delta V = \sum_{k=1}^K \delta V / K$ and apply our theorem K times we can suppose δV arbitrary small. We are looking for the new fixed point for $F_{V+\delta V}$ which is the limit of $(F_{V+\delta V})^n(\gamma)$. We start from $\gamma = \gamma_{\min}(V)$ and remark that

$$\begin{aligned} (\gamma_{\min}(V + \delta V) - \gamma) &= \lim_{n \rightarrow \infty} ((F_{V+\delta V})^n(\gamma) - \gamma) \\ &= \sum_{n=0}^{\infty} ((F_{V+\delta V})^{n+1}(\gamma) - (F_{V+\delta V})^n(\gamma)). \end{aligned} \tag{24}$$

Step 1. We evaluate the first term of the sum $F_{V+\delta V}(\gamma) - \gamma = F_{V+\delta V}(\gamma) - F_V(\gamma)$, as follows

$$\begin{aligned} &F_{V+\delta V}(\gamma) - F_V(\gamma) \\ &= \frac{1}{2i\pi} \oint_{\mathcal{C}} \frac{1}{H_{\text{eff}}(\gamma) + \delta V - z} - \frac{1}{H_{\text{eff}}(\gamma) - z} dz \\ &= \frac{1}{2i\pi} \oint_{\mathcal{C}} \frac{1}{H_{\text{eff}}(\gamma) - z} \left(\delta V \sum_{k \leq 0} ((H_{\text{eff}}(\gamma) - z)^{-1} \delta V)^k \right) \frac{1}{H_{\text{eff}}(\gamma) - z} dz \\ &= \frac{1}{2i\pi} \oint_{\mathcal{C}} \frac{1}{H_{\text{eff}}(\gamma) - z} B(\delta V) \frac{1}{H_{\text{eff}}(\gamma) - z} dz, \end{aligned}$$

where

$$B(\delta V) = \delta V \sum_{k \leq 0} ((H_{\text{eff}}(\gamma) - z)^{-1} \delta V)^k. \tag{25}$$

This sum converges as soon as $\delta V < \frac{G}{6}$. Note that if the support of δV has a bounded support Ω so has $B(\delta V)$. We will use the Combes–Thomas estimate (20) for y_1, y_2 outside Ω . Remark that if we choose the loop \mathcal{C} correctly, v does not

depend on z but only on the size on the gap. We find

$$\begin{aligned}
 & |(F_{V+\delta V}(\gamma))(y_1, y_2) - (F_V(\gamma))(y_1, y_2)| \\
 &= \oint_e \left(y_1, \frac{1}{H_{\text{eff}}(\gamma) - z} B(\delta V) \frac{1}{H_{\text{eff}}(\gamma) - z} y_2 \right) dz \\
 &\leq \frac{1}{2i\pi} \oint_e C^2 \sum_{x, x' \in \Omega} e^{-\nu|y_1-x|} (x, B(\delta V)x') e^{-\nu|x'-y_2|} dz \\
 &\leq C^2 |\mathcal{C}| e^{-\nu d(y_1, \Omega)} \sum_{x, x' \in \Omega} |(x, B(\delta V)x') e^{-\nu|x'-y_2|}| \\
 &\leq C' \|\delta V\| e^{-\nu d(y_1, \Omega)},
 \end{aligned}$$

where C' is just a constant. With W small enough, there exists $\tau < 1$ so that

$$[A_{\text{eff}}(F_{V+\delta V}(\gamma)) - A_{\text{eff}}(\gamma)](y_1, y_2) \leq \tau \|\delta V\| e^{-\nu d(y_1, \Omega)}. \tag{26}$$

Step 2. We evaluate the remainder of the sum. We repeat the previous argument with

$$\delta A_{\text{eff}} = A_{\text{eff}}(F_{V+\delta V}^n(\gamma)) - A_{\text{eff}}(F_{V+\delta V}^{n-1}(\gamma)) \tag{27}$$

instead of δV . There is only one little difference: B does not have a bounded support any more but still has an off-diagonal exponential decay (proved by iteration with constant $\nu > \nu' > 0$). We just check this does not bring more difficulties:

$$\begin{aligned}
 & |(F_{V+\delta V}^n(\gamma))(y_1, y_2) - (F_{V+\delta V}^{n-1}(\gamma))(y_1, y_2)| \\
 &\leq C^2 \sum_{x, x' \in \Omega} e^{-\nu|y_1-x|} (x, B(\delta A_{\text{eff}})x') e^{-\nu|x'-y_2|} dy \\
 &\leq C^2 \|\delta A_{\text{eff}}\| \sum_{x, x' \in \mathbb{Z}^2} e^{-\nu|y_1-x|} e^{-\nu' d(x, \Omega)} e^{-\nu' d(x', \Omega)} e^{-\nu|x'-y_2|} \\
 &\leq C_2 \|\delta A_{\text{eff}}\| e^{-\nu' d(y_1, \Omega)} \sum_{x, x' \in \mathbb{Z}^2} e^{\nu' d(y_1, \Omega) - \nu|y_1-x| - \nu' d(x, \Omega)} e^{-\nu d(x', \Omega)} e^{-\nu|x'-y_2|} \\
 &\leq C'_2 \|(F_{V+\delta V}^{n-1}(\gamma)) - F_{V+\delta V}^{n-2}(\gamma)\| e^{-\nu' d(y_1, \Omega)},
 \end{aligned}$$

where C'_2 is just another constant. We can conclude by iteration that

$$|A_{\text{eff}}(F_{V+\delta V}^n(\gamma))(y_1, y_2) - A_{\text{eff}}(F_V^{n-1}(\gamma))(y_1, y_2)| \leq \|\delta V\| \tau^n e^{-\nu' d(y_1, \Omega)}, \tag{28}$$

and so

$$\begin{aligned} & |(A_{\text{eff}}(\gamma_{\min}(V + \delta V)) - A_{\text{eff}}(\gamma_{\min}(V)))(y_1, y_2)| \\ &= \lim_{n \rightarrow \infty} |A_{\text{eff}}(F_{V+\delta V}^n(\gamma))(y_1, y_2) - A_{\text{eff}}(\gamma)(y_1, y_2)| \\ &\leq \|\delta V\| \frac{1}{1-\tau} e^{-\nu d(y_1, \text{supp}(\delta V))}, \end{aligned}$$

as we wanted. \square

5. Wegner estimate: proof of Theorem 2.4

In this section, we prove the Wegner-type estimate in Theorem 2.4.

The idea of the proof is the following. At first sight we do not have any idea of how A_{eff} looks like. But because of the gap, if

$$\gamma = \mathbb{1}_{\leq \mu}(-\Delta + V + A_{\text{eff}}(\gamma)) \quad (29)$$

then

$$\gamma = \mathbb{1}_{\leq (\mu+\alpha)}(-\Delta + V + A_{\text{eff}}(\gamma)) = \mathbb{1}_{\leq (\mu)}(-\Delta + V - \alpha \mathbb{1}_{\mathbb{Z}^d} + A_{\text{eff}}(\gamma)) \quad (30)$$

for any $\alpha \in \mathbb{R}$ with $2|\alpha|$ smaller than the gap. So if we could add α to the random potential with α a smooth random variable, in this case every eigenvalue of H_{\min} would just be offset by α no matter what the non linear part is and we are done. In our case, we will make the change of variable $(V_\omega(x)) \rightarrow (\alpha = \frac{1}{|\Lambda|} \sum_{x \in \Lambda} V_\omega(x), V_\omega(x) - \alpha)$ for $x \in \Lambda$ and we expect that the conditional density of α is smooth enough and that the change induced to γ is small.

Proof of Theorem 2.4. Let $\Lambda = \Lambda_L(n)$ be the cube in \mathbb{Z}^d of size L with its center in n and $\Lambda_{2L}(n)$ the cube twice bigger. Because of (30) and (11)

$$\begin{aligned} & \left\| \frac{d}{d\alpha} ((\gamma_{\min})(V + \alpha \mathbb{1}_{\Lambda_{2L}(n)})|_{\Lambda_L(n)}) \right\| \\ &= \left\| \frac{d}{d\alpha} ((\gamma_{\min})|_{\Lambda_L(n)}(V + \alpha \mathbb{1}_{\mathbb{Z}^d} - \alpha \mathbb{1}_{\Lambda_{2L}(n)^c})) \right\| \\ &= \left\| \frac{d}{d\alpha} ((\gamma_{\min})|_{\Lambda_L(n)}(V - \alpha \mathbb{1}_{\Lambda_{2L}(n)^c})) \right\| \\ &\leq C e^{-\nu d(\Lambda_L(n), \Lambda_{2L}(n)^c)} \\ &\leq C e^{-\nu L}. \end{aligned}$$

We suppose that L is large enough so that $2\|W\|_{\ell^1} C e^{-\nu L} \leq 1/2$ and we obtain that

$$\alpha \rightarrow \|\mathbb{1}_{\Lambda_L(n)} A_{\text{eff}}(V + \alpha \mathbb{1}_{\Lambda_{2L}(n)}) \mathbb{1}_{\Lambda_L(n)}\| \tag{31}$$

is $1/2$ Lipschitz. Under this hypothesis, for any $\lambda_i(\alpha)$ eigenvalue of

$$(-\Delta + V + \alpha \mathbb{1}_{\Lambda_{2L}(n)} + A_{\text{eff}}(\gamma_{\min}(V + \alpha \mathbb{1}_{\Lambda_{2L}(n)})))|_{\Lambda_L(n)},$$

we have

$$\frac{d}{d\alpha} \lambda_i(\alpha) \geq 1 - \left\| \frac{d}{d\alpha} (\mathbb{1}_{\Lambda_L(n)} A_{\text{eff}}(V + \alpha \mathbb{1}_{\Lambda_{2L}(n)}) \mathbb{1}_{\Lambda_L(n)}) \right\| \geq \frac{1}{2}. \tag{32}$$

Let $\lambda \in \mathbb{R}$ and $\epsilon > 0$. Let $D_0 = \{d_1 < d_2 < \dots < d_k\}$ be so that $\rho(s)$ is Lipschitz on $]d_n, d_{n+1}[$. Let f and δ be two positive functions that will be chosen later. We define the following events

$$O_x := \{\omega: \text{for all } y \text{ such that } |y - V_\omega(x)| < \delta(\epsilon), \rho(y) > f(\epsilon), \text{ and } d(V_\omega(x), D_0) > \delta(\epsilon)\} \tag{33}$$

for any $x \in \Lambda_{2L}(n)$. We now estimate

$$\begin{aligned} & \mathbb{P}(d(\sigma[(H_{\min})|_{\Lambda_L(n)}], \lambda) < \epsilon) \\ & \leq \mathbb{P}\left(\bigcup_{x \in \Lambda_{2L}(n)} O_x^c\right) + \mathbb{P}(\cap_{x \in \Lambda_{2L}(n)} O_x \cap d(\sigma[(H_{\min})|_{\Lambda_L(n)}], \lambda) < \epsilon). \end{aligned} \tag{34}$$

We will deal with each term separately. Starting with the left term, we erase the indices because the probability does not depend of the position and argue as follows:

$$\begin{aligned} \mathbb{P}\left(\bigcup_{x \in \Lambda_{2L}(n)} O_x^c\right) & \leq |\Lambda_{2L}(n)| \mathbb{P}(O_x^c) \\ & = 2^d |\Lambda| \mathbb{P}(O^c) \\ & \leq 2^d |\Lambda| (\mathbb{P}(d(V_\omega, D_0) < \delta(\epsilon)) \\ & \quad + \mathbb{P}(V_\omega: \text{there exists } y: |y - V_\omega| < \delta(\epsilon), \rho(y) < f(\epsilon))) \\ & \leq 2^d |\Lambda| (\mathbb{P}(d(V_\omega, D_0) < \delta(\epsilon)) \\ & \quad + \mathbb{P}(V_\omega: \rho(V_\omega) < f(\epsilon) + \delta(\epsilon) \|\rho'\|_\infty)) \\ & \leq 2^d |\Lambda| (\|\rho\|_\infty 2\delta(\epsilon) \#D_0 + (\delta(\epsilon) \|\rho'\|_\infty + f(\epsilon)) |\text{supp}(\phi)|). \end{aligned}$$

The right term in (34) can be estimated by introducing the mean and the resolvent, using

$$\mathbb{1}_{[\lambda - \epsilon, \lambda + \epsilon]}(\alpha) \leq \frac{2\epsilon^2}{((\lambda - \alpha)^2 + \epsilon^2)} = 2\epsilon \Im\left(\frac{1}{\lambda - \alpha + i\epsilon}\right). \tag{35}$$

We simplify a bit the notation using $\cap_x O_x$ instead of $\cap_{x \in \Lambda_{2L}(n)} O_x$. We get

$$\begin{aligned} & \mathbb{P}(\cap_x O_x \cap d(\sigma[(H_{\min})|_{\Lambda_L(n)}], \lambda) < \epsilon) \\ &= \mathbb{E}[\mathbb{1}_{d(\sigma(H_{\min}^\wedge), \lambda) < \epsilon} \mathbb{1}_{\cap_x O_x}] \\ &\leq 2\epsilon \mathbb{E}[\mathfrak{S}(\text{Tr}[(H_{\min})|_{\Lambda_L(n)} - \lambda + i\epsilon]^{-1}) \mathbb{1}_{\cap_x O_x}]. \end{aligned}$$

We now make a change of variable for $V_\omega(x) \in \Lambda_{2L}(n)$:

$$(V_\omega(x)) \longrightarrow \left(\alpha = \frac{1}{2^d |\Lambda|} \sum_{x \in \Lambda_{2L}(n)} V_\omega(x), V_\omega(x) - \alpha \right). \tag{36}$$

We write $\tilde{V}_\omega(x) = V_\omega(x) - \alpha$ and $\xi_{\tilde{V}}(\alpha)$ for the conditional random density of the mean knowing \tilde{V} . We first integrate over α , then over \tilde{V} (we denote the expectation by $\mathbb{E}_{\tilde{V}}$):

$$\begin{aligned} & \mathbb{P}(\cap_x O_x \cap d(\sigma[(H_{\min})|_{\Lambda_L(n)}], \lambda) < \epsilon) \\ &\leq \mathbb{E}_{\tilde{V}} \left[2\epsilon \int \mathfrak{S}(\text{Tr}[(H_{\min})|_{\Lambda_L} - \lambda + i\epsilon]^{-1}) \mathbb{1}_{\cap_x O_x} \xi_{\tilde{V}}(\alpha) d\alpha \right] \\ &\leq 2|\epsilon| \mathbb{E}_{\tilde{V}} \left[\sum_{\lambda_i \in \sigma((H_{\min})|_{\Lambda_L})} \int \mathfrak{S}\left(\frac{1}{(\lambda_i(\alpha) - \lambda + i\epsilon)}\right) \mathbb{1}_{\cap_x O_x} \xi_{\tilde{V}}(\alpha) d\alpha \right]. \end{aligned}$$

We estimate the integral with a change of variable $\alpha' = (\lambda_i(\alpha) - \lambda)$, $d\alpha' = (\frac{d}{d\alpha} \lambda_i) d\alpha$. Recall that

$$\int \frac{\epsilon}{(\alpha')^2 + \epsilon^2} \frac{\xi_{\tilde{V}}(\alpha) \mathbb{1}_{\cap_x O_x}}{\frac{d}{d\alpha} \lambda_i} d\alpha' \leq \pi \sup \left[\frac{|\xi_{\tilde{V}}(\alpha) \mathbb{1}_{\cap_x O_x}|}{\frac{d}{d\alpha} \lambda_i} \right]. \tag{37}$$

So we have

$$\mathbb{P}(\cap_x O_x \cap d(\sigma[(H_{\min})|_{\Lambda_L(n)}], \lambda) < \epsilon) \leq 2\pi |\epsilon| |\Lambda_L| \mathbb{E}_{\tilde{V}} \left[\sup \left[\frac{|\xi_{\tilde{V}}(\alpha) \mathbb{1}_{\cap_x O_x}|}{\frac{d}{d\alpha} \lambda_i} \right] \right].$$

Finally, because of (32), $\frac{d}{d\alpha} \lambda_i > 1/2$ and we get

$$\mathbb{P}(\cap_x O_x \cap d(\sigma[(H_{\min})|_{\Lambda_L(n)}], \lambda) < \epsilon) \leq 4\pi |\epsilon| |\Lambda_L| \sup[\xi_{\tilde{V}} \mathbb{1}_{\cap_x O_x}]. \tag{38}$$

From now on, it is enough to have an estimate on $\xi_{\tilde{V}} \mathbb{1}_{\cap_x O_x}$. A computation gives

$$\xi_{\tilde{V}}(\alpha) d\alpha = \mathbb{P}\left(\sum_x V_\omega(x) \in [\alpha, \alpha + d\alpha] | \tilde{V}\right) = \frac{\prod_{i \in \Lambda} \rho(\tilde{V}_i + \alpha)}{\int \prod_{i \in \Lambda} \rho(\tilde{V}_i + \alpha') d\alpha'}. \tag{39}$$

Let $\alpha_0 \in \mathbb{R}$. If we do not have $\tilde{V}_\omega(x) + \alpha_0 \in O_x$ for all x , then $\xi_{\tilde{V}}(\alpha_0)\mathbb{1}_{\cap_x O_x} = 0$ and we have nothing else to do. So we can assume that for all α , $|\alpha - \alpha_0| < \delta(\epsilon) \implies \rho(\tilde{V}(x) + \alpha) > f(\epsilon)$ for all $x \in \Lambda_{2L}(n)$ and all $|\alpha - \alpha_0| < \delta(\epsilon)$. So we have

$$\frac{\frac{d}{d\alpha} \prod_{x \in \Lambda_{2L}} \rho(\tilde{V} + \alpha)}{\prod_{x \in \Lambda_{2L}} \rho(V + \alpha)} = \sum_{x \in \Lambda_{2L}} \frac{\rho'(\tilde{V}_\omega(x) + \alpha)}{\rho(\tilde{V}_\omega(x) + \alpha)} \leq 2^d |\Lambda_L| \frac{\|\rho'\|_\infty}{f(\epsilon)}. \tag{40}$$

From this differential equation we get

$$\prod_{x \in \Lambda_{2L}} \rho(\tilde{V}_\omega(x) + \alpha) \geq \exp\left(-(|\alpha - \alpha_0|)2^d |\Lambda_L| \frac{\|\rho'\|_\infty}{f(\epsilon)}\right) \prod_{x \in \Lambda_{2L}} \rho(\tilde{V}_\omega(x) + \alpha_0)$$

and, after integrating,

$$\begin{aligned} & \int_{\alpha_0 - \delta(\epsilon)}^{\alpha_0 + \delta(\epsilon)} \prod \rho(\tilde{V}_\omega(x) + \alpha) d\alpha \\ & \geq \left[1 - \exp(-|\delta(\epsilon)|2^d |\Lambda_L| \frac{\|\rho'\|_\infty}{f(\epsilon)})\right] \frac{f(\epsilon)}{2^d |\Lambda_L| \|\rho'\|_\infty} \prod \rho(\tilde{V}_\omega(x) + \alpha_0). \end{aligned}$$

Therefore

$$\frac{\prod \rho(\tilde{V}_\omega(x) + \alpha_0)}{\int \prod \rho(\tilde{V}_\omega(x) + \alpha) d\alpha} \leq \frac{2^d |\Lambda_L| \|\rho'\|_\infty}{f(\epsilon) \left(1 - \exp\left(-|\delta(\epsilon)|2^d |\Lambda_L| \frac{\|\rho'\|_\infty}{f(\epsilon)}\right)\right)}$$

and hence we have

$$\frac{\prod \rho(\tilde{V}_\omega(x) + \alpha_0)}{\int \prod \rho(\tilde{V}_\omega(x) + \alpha) d\alpha} \leq 2 \max\left(\frac{2^d |\Lambda_L| \|\rho'\|_\infty}{f(\epsilon)}, \frac{1}{\delta(\epsilon)}\right). \tag{41}$$

We finally obtain

$$\xi_{\tilde{V}_i} \mathbb{1}_{\cap_x O_x} \leq 2 \max\left(\frac{2^d |\Lambda_L| \|\rho'\|_\infty}{f(\epsilon)}, \frac{1}{\delta(\epsilon)}\right), \tag{42}$$

for all \hat{V} and all α_0 .

To conclude, putting (34), (35), (38), and (42) together, we have

$$\begin{aligned} & \mathbb{P}(d(\sigma[(H_{\min})|_{\Lambda_L(n)}], \lambda) < \epsilon) \\ & \leq 2^d |\Lambda| \left[\|\rho\|_\infty \delta(\epsilon) \# |D_0| + (\delta(\epsilon) \|\rho'\|_\infty \right. \\ & \quad \left. + f(\epsilon)) * |\text{supp}(\rho)| + 8\pi\epsilon \max\left(\frac{|\Delta| \|\rho'\|_\infty}{f(\epsilon)}, \frac{1}{\delta(\epsilon)}\right) \right], \end{aligned}$$

from which we can conclude (12) choosing $f(\epsilon) = \sqrt{\epsilon}/|\text{supp}(\rho)|^{-3/2}$ and $\delta(\epsilon) = \sqrt{\epsilon} * |\text{supp}(\rho)|^{1/2}$. Hence

$$\begin{aligned} & \mathbb{P}(d(\sigma[(H_{\min})|_{\Lambda_L(n)}], \lambda) < \epsilon) \\ & \leq C |\Lambda| [|\text{supp}(\rho)|^{-1/2} + \|\rho\|_{\infty} |\text{supp}(\rho)|^{1/2} + \|\rho'\|_{\infty} |\text{supp}(\rho)|^{3/2}], \end{aligned}$$

where the constant C only depends on the cardinal of D_0 . □

In particular, with the change $V \rightarrow \ell V$, we deduce that

$$\mathbb{P}(d(\sigma[(H_{\min})|_{\Lambda_L(n)}], \lambda) < \epsilon) \longrightarrow 0 \tag{43}$$

when $\ell \rightarrow \infty$.

6. Multiscale analysis

We will now start the proof of the multiscale analysis. There will be very little differences with the proof we can find in [16, Part 10] and we will follow the method exposed there step by step. But because it is a more general case, we have written the proof again.

6.1. The setting. For any operator K with off-diagonal exponential decay and $\Lambda \subset \mathbb{Z}^d$, we define a border operator Γ by

$$\Gamma_{K,\Lambda}(x, y) = \begin{cases} K(x, y) & \text{if } (x \in \Lambda \text{ and } y \notin \Lambda) \text{ or } (y \in \Lambda \text{ and } x \notin \Lambda), \\ 0 & \text{otherwise.} \end{cases}$$

The following proposition is a form of the Schur complement formula.

Proposition 6.1. *Let K with off-diagonal exponential decay, Λ a box of size L , and $\lambda \in \mathbb{C} \setminus \mathbb{R}$. Then*

$$(K - \lambda)^{-1}(x, y) = - \sum_{u \in \Lambda, v \notin \Lambda} (K^\Lambda - \lambda)^{-1}(x, u) \Gamma_{K,\Lambda}(u, v) (K - \lambda)^{-1}(v, y) \tag{44}$$

for any $x \in \Lambda$ and any $y \notin \Lambda$, where

$$K^\Lambda(x, y) = \begin{cases} K(x, y) & \text{if } x \in \Lambda \text{ and } y \in \Lambda, \\ 0 & \text{otherwise,} \end{cases}$$

is the restriction of K to Λ .

Proof. We can divide K into the following three parts

$$K = K^\Lambda + \Gamma_{K,\Lambda} + K^{\Lambda^c} \tag{45}$$

where Λ^c is the complement of Λ . We here use the resolvent formula

$$\begin{aligned} (K - \lambda)^{-1} &= (K^\Lambda - \lambda)^{-1} + (K^{\Lambda^c} - \lambda)^{-1} \\ &\quad - ((K^\Lambda - \lambda)^{-1} + (K^{\Lambda^c} - \lambda)^{-1})\Gamma_{K,\Lambda}(K - \lambda)^{-1}. \end{aligned} \tag{46}$$

Just remark now that $(K^\Lambda - \lambda)^{-1}(x, y) = 0$ and $(K^{\Lambda^c} - \lambda)^{-1}(x, y) = 0$ if $x \in \Lambda$ and $y \notin \Lambda$. □

We now apply the multiscale method. Let $\Lambda_L(n)$ be the box of side length $2L + 1$ centered at $n \in \mathbb{Z}^d$. We replace the random potential V_ω by an arbitrary constant outside the box $\Lambda_{2L}(n)$ in order to make the mean field Hamiltonian inside $\Lambda_L(n)$ independent of what is happening outside $\Lambda_{2L}(n)$:

$$\widehat{V}_\omega^{\Lambda_L(n)}(x) = \begin{cases} V_\omega(x) & \text{if } x \in \Lambda_{2L}(n), \\ 0 & \text{otherwise.} \end{cases}$$

Recall that $0 \in \text{supp}(\mathbb{P})$. From this potential we can obtain with Theorem 2.1 the minimiser $\gamma_{\min}(\widehat{V}_\omega^{\Lambda_L(n)})$ and the mean-field Hamiltonian $H_{\min}(\widehat{V}_\omega^{\Lambda_L(n)})$. We denote its restriction to $\Lambda_L(n)$ by

$$\widehat{H}(n, L) := (H_{\min}(\widehat{V}_\omega^{\Lambda_L(n)}))|_{\Lambda_L(n)}.$$

We introduce this Hamiltonian because of two properties. First it is independent of what is happening outside $\Lambda_{2L}(n)$. Second, it is a good approximation of $(H_{\min})|_{\Lambda_L(n)}$. Indeed, from Theorem 2.3 we have

$$\|(H_{\min})|_{\Lambda_L(n)} - \widehat{H}(n, L)\| < De^{-\nu L} \tag{47}$$

where D does not depend on n and L .

Definition 6.1 (L-resonance). A number $\lambda \in \mathbb{R}$ is called L-resonant for the box $\Lambda_L(n)$ if there exists A_c with $\|A_c\| \leq 2D \exp(-\nu L)$ and

$$d(\lambda, \sigma[\widehat{H}(n, L) + A_c]) \leq \exp(-\sqrt{L}), \tag{48}$$

where D is the constant defined in (47)

Remark that our definition of non-resonance is equivalent to

$$d(\lambda, \sigma[\widehat{H}(n, L)]) > \exp(-\sqrt{L}) - 2D \exp(-\nu L). \tag{49}$$

We have added the operator A_c in the above definition to handle the difference between $(H_{\min})|_{\Lambda_L(n)}$ and $\widehat{H}(n, L)$. This corresponds to Definition 9.1 in [16].

Definition 6.2 ((L, ζ, λ) -good box). The box $\Lambda_L(n)$ is called an (L, ζ, λ) -good box if

- (1) it is not L -resonant;
- (2) for any $x \in \Lambda_{\sqrt{L}}(n)$, $y \notin \Lambda_L(n)$ and A_c with $\|A_c\| \leq 2D \exp(-\nu L)$,

$$\sum_v |\hat{H}(n, L) + A_c - \lambda)^{-1}(x, v)| |\Gamma_{\hat{H}(n, L) + A_c, \Lambda_L(n)}(v, y)| \leq \exp(-\zeta|y - x|). \tag{50}$$

6.2. From a scale to another. Let L_0 be not too small and set $L_k = L_0^{\alpha^k}$ with $1 < \alpha < 2$. In this subsection, we prove the following proposition.

Proposition 6.2. *If the following conditions are satisfied*

- (1) *for any 4 boxes of side length L_k in $\Lambda_{L_{k+1}}(n)$, separated from each other by a distance of at least $2L_k$, there is at least one which is (L_k, ζ, λ) -good with $\zeta > 20/\sqrt{L_k}$,*
- (2) *no box in $\Lambda_{L_{k+1}}(n)$ of side length $4L_k, 12L_k, 20L_k$ is L_k -resonant,*
- (3) *the domain $\Lambda_{L_{k+1}}(n)$ is not L_{k+1} -resonant,*

then the cube $\Lambda_{L_{k+1}}(n)$ is $(L_{k+1}, \zeta_{k+1}, \lambda)$ -good with a decay satisfying $\zeta_{k+1} > 20/\sqrt{L}$.

This proposition corresponds to Theorem 10.20 in [16].

Proof of Proposition 6.2. Let A_c be so that

$$\|A_c\| \leq 2De^{-\nu L_{k+1}}.$$

Let $(\hat{H}(n, L_{k+1}))|_{\Lambda_{L_k}(m)}$ be the restriction to the box $\Lambda_{L_k}(m)$ of $\hat{H}(n, L_{k+1})$, with $\Lambda_k(m) \subset \Lambda_{k+1}(n)$. Because of Theorem 2.3, we have

$$\|(\hat{H}(n, L_{k+1}))|_{\Lambda_{L_k}(m)} - \hat{H}(m, L_k)\| \leq De^{-\nu L_k}$$

and so

$$\|(\hat{H}(n, L_{k+1}) + A_c)|_{\Lambda_{L_k}(m)} - \hat{H}(m, L)\| \leq 2De^{-\nu L_k}$$

for L_k big enough.

Let $K = \hat{H}(n, L_{k+1}) + A_c$ and for simplicity we will just write Γ_Λ instead of $\Gamma_{K,\Lambda}$. Because of what we have just said, if $\Lambda_{L_k}(m)$ is (L_k, ζ, E) good then

$$\sum_v (K^{\Lambda_{L_k}(m)} - \lambda)^{-1}(x, v) |\Gamma(v, y)| \leq \exp(-\zeta|y - x|) \tag{51}$$

and

$$\|(K^{\Lambda_{jL_k}(m)} - \lambda)^{-1}\| \leq \exp(\sqrt{jL_k}) \tag{52}$$

if $\Lambda_{jL_k}(m)$ is not jL_k -resonant for $j = 4, 12, 20$.

The idea is to use equation (44) as many times as we want. For any v appearing in the equation (44), we can define another box $\Lambda(v)$ with $y \notin \Lambda(v)$ and repeat the formula with v instead of x . Proceeding this way again and again, we get after iteration

$$(K - \lambda)^{-1}(x, y) = \sum_{(u_i, v_i)_{i=1..n}} (K^{\Lambda_1} - \lambda)^{-1}(x, u_1) \Gamma_{\Lambda_1}(u_1, v_1) (K^{\Lambda_2} - \lambda)^{-1}(v_1, u_2) \Gamma_{\Lambda_2}(u_2, v_2) \dots (K - \lambda)^{-1}(v_n, y). \tag{53}$$

We will write the indices of the sum as a tree \mathcal{T} of chains $\mathcal{X} = (u_i, v_i, \Lambda_i)_{i \leq n}$ with $v_i \in \Lambda_i$, $u_{i+1} \in \Lambda_i$, $v_{i+1} \notin \Lambda_i$ and $y \notin \Lambda_i$. We first sum over the u_i 's so as to reduce our chains to $\mathcal{X} = (v_i, \Lambda_i)_{i \leq n}$ and we introduce an upper bound $R_{\mathcal{X}}$ such that

$$R_{\mathcal{X}} \geq \sum_{(u_i)_{i=1..n}} |(K^{\Lambda_1} - \lambda)^{-1}(x, u_1) \Gamma_{\Lambda_1}(u_1, v_1) (K^{\Lambda_2} - \lambda)^{-1}(v_1, u_2) \Gamma_{\Lambda_2}(u_2, v_2) \dots \Gamma(u_n, v_n)|. \tag{54}$$

Then, equation (53) gives

$$|(K - \lambda)^{-1}(x, y)| \leq \|(K - \lambda)^{-1}\| \sum_{\mathcal{X} \text{ leaves of } \mathcal{T}} R_{\mathcal{X}}. \tag{55}$$

This formula is very general and is valid for any expansion. Different choices for the construction of the tree exist in the literature and we will follow that of [16]. The goal is to get at least one good box at each step. The choice of the Λ_i and the construction of the tree \mathcal{T} of \mathcal{X} and $R_{\mathcal{X}}$ are made according to the following algorithm.

We start from x so we define $v_0 = x$ and $R = 1$. The choice of Λ_{i+1} will depend on v_i .

- If we get close to the boundary or to y , ($d(v_i, \partial\Lambda_{L_{k+1}}(n)) < L_k$ or $d(v_i, y) < L_k$) then we stop. The construction of this chain is over and we carry on with the other branches of the tree.

- Otherwise

- if $\Lambda_{L_k}(v_i)$ is an (L_k, ζ, E) -good box then

$$\begin{aligned} &R_{\mathcal{X}}(K - \lambda)^{-1}(v_i, y) \\ &\leq R_{\mathcal{X}} \sum_{u_{i+1}, v_{i+1}} (K^{\Lambda_i} - \lambda)^{-1}(v_i, u_{i+1}) \\ &\quad \Gamma_{\Lambda_i}(u_{i+1}, v_{i+1})(K - \lambda)^{-1}(v_{i+1}, y) \\ &\leq \sum_{v_{i+1} \notin \Lambda_{L'}(v_i)} R_{\mathcal{X}} \exp(-\zeta|v_{i+1} - v_i|)(K - \lambda)^{-1}(v_{i+1}, y) \end{aligned}$$

so for each v_{i+1} outside $\Lambda_L(v_i)$, we set

$$R_{\mathcal{X}+v_{i+1}} = R_{\mathcal{X}} \exp(-\zeta|v_{i+1} - v_i|) \tag{56}$$

and carry on the algorithm with the new chain $\mathcal{X} + v_{i+1}$;

- else if $\Lambda_{L_k}(u_i)$ is not a good box, choose $j = 4$ or 12 or L_k such that for every v in $\Lambda_{2jL_k} \setminus \Lambda_{jL_k}$, $\Lambda_{L_k}(v)$ is a good box and Λ_{jL_k} is not resonant. It is always possible to do this because of the following remark: either 3 boxes are far away from each other then there are 3 boxes M_1, M_2, M_3 of size $4L_k$ separated by at least $2L_k$ so that every cube $\Lambda_{L_k}(m) \subset \Lambda_{L_{k+1}}(m)$ whose center is not included in $\bigcup_{i=1,2,3} M_i$ are (L_k, ζ, λ) -good. Or two of them are close and the other is far away then there are two boxes M_1 of size $12L_k$ and M_2 of size $4L_k$ separated by at least $2L_k$ so that every cube in $\Lambda_{L_k}(m) \subset \Lambda_{L_{k+1}}(m)$ whose center is not included in M_i are (L_k, ζ, λ) -good. Or the three of them are together then there exist one box M_1 of size $20L_k$ so that every cube $\Lambda_{L_k}(m) \subset \Lambda_{L_{k+1}}(m)$ whose center is not included in M_1 are (L_k, ζ, λ) -good. We can assume that the good box decay is smaller than the off diagonal decay parameter ν , $\zeta < \nu$. We then have

$$\begin{aligned} &R_{\mathcal{X}}(K - \lambda)^{-1}(v_i, y) \\ &\leq R_{\mathcal{X}} \sum_{u_{i+1}, v_{i+1}} (K^{\Lambda_{i+1}} - \lambda)^{-1}(v_i, u_{i+1}) \\ &\quad \Gamma_{\Lambda_{i+1}}(u_{i+1}, v_{i+1})(K - \lambda)^{-1}(v_{i+1}, y) \\ &\leq R_{\mathcal{X}} \sum_{u_{i+1}, v_{i+1} \in \Lambda_{2jL_k}(v_i)} (K^{\Lambda_{i+1}} - \lambda)^{-1}(v_i, u_{i+1}) \\ &\quad \Gamma_{\Lambda_{i+1}}(u_{i+1}, v_{i+1})(K - \lambda)^{-1}(v_{i+1}, y) \\ &\quad + \sum_{u_{i+1}, v_{i+1} \notin \Lambda_{2jL_k}(v_i)} (K^{\Lambda_i} - \lambda)^{-1}(v_i, u_{i+1}) \\ &\quad \Gamma_{\Lambda_{i+1}}(u_{i+1}, v_{i+1})(K - \lambda)^{-1}(v_{i+1}, y) \end{aligned}$$

$$\begin{aligned}
 &\leq R_{\mathcal{X}} \sum_{\substack{u_{i+1}, v_{i+1} \in \Lambda_{2jL_k}(v_i) \\ u'_{i+1}, v'_{i+1} \notin \Lambda_{L_k}(v_{i+1})}} (K^{\Lambda_{i+1}} - \lambda)^{-1}(v_i, u_{i+1}) \Gamma(u_{i+1}, v_{i+1}) (K^{\Lambda_{L_k}(v_{i+1})} - \lambda)^{-1}(v_{i+1}, u'_{i+1}) \\
 &\quad \Gamma_{\Lambda_{L_k}(v_{i+1})}(u'_{i+1}, v'_{i+1}) (K - \lambda)^{-1}(v'_{i+1}, y) \\
 &\quad + \sum_{v_{i+1} \notin \Lambda_{2jL_k}(u_i)} (jL_k)^d C e^{\sqrt{jL_k}} \exp(-\nu(|v_{i+1} - v_i| - jL_k)) (K - \lambda)^{-1}(v_{i+1}, y) \\
 &\leq R_{\mathcal{X}} \sum_{v_{i+1}} (2jL_k)^d e^{\sqrt{jL_k}} \exp(-\zeta \max(|v_{i+1} - v_i| - 2jL_k, L_k)) (K - \lambda)^{-1}(v_{i+1}, y) \\
 &\quad + \sum_{v_{i+1} \notin \Lambda_{2jL_k}(u_i)} (jL_k)^d C e^{\sqrt{jL_k}} \exp(-\nu(|v_{i+1} - v_i| - jL_k)) (K - \lambda)^{-1}(v_{i+1}, y).
 \end{aligned}$$

Therefore we can set

$$\begin{aligned}
 R_{\mathcal{X}+v_{i+1}} &= R_{\mathcal{X}} 2C (2jL_k)^d \exp(\sqrt{jL_k}) \\
 &\quad \exp(-\zeta \max(L_k, |v_{i+1} - v_i| - 2jL_k)) \tag{57}
 \end{aligned}$$

and as previously, we carry on with the new chain $\mathcal{X} + v_{i+1}$.




Figure 1. Schematic representation a typical chain \mathcal{X} used in the proof of Proposition 6.2.

We have finished the description of the algorithm. We define the "good path length" $P_g(x, v)$ as the minimum length between x and v when any cube $M = \Lambda_{2jL_k}(v)$ containing bad boxes defined in the procedure can be crossed for free. We easily check that (56) and (57) imply that

$$R_{\mathcal{X}} \leq \exp(-\zeta \max(\text{length}(\mathcal{X})L_k, P_g(x, v_i))). \tag{58}$$

for every chain \mathcal{X} . From this, our algorithm gives us the following estimate:

Proposition 6.3. *We have*

$$\sum_x R_{\mathcal{X}} \leq \frac{1}{1 - (L_{k+1})^d \exp(-\zeta L_k)} (L_{k+1})^{d \frac{L_{k+1}}{L_k}} \exp(-\zeta(P_g(x, y))). \tag{59}$$

Proof. We have

$$\begin{aligned} \sum_{(\mathcal{X})} R_{\mathcal{X}} &= \sum_{N=1}^{\infty} \sum_{\mathcal{X}, N=\text{length}(\mathcal{X})} R_{\mathcal{X}} \\ &\leq \sum_{N=1}^{\infty} \sum_{\mathcal{X}, N=\text{length}(\mathcal{X})} \exp(-\zeta \max(NL_k, P_g(x, y))) \\ &\leq \sum_{N=1}^{\infty} \sum_{\mathcal{X}, N=\text{length}(\mathcal{X})} \exp(-\zeta \left(P_g(x, y) + \max\left(0, N - \frac{L_{k+1}}{L_k}\right) L_k \right)) \\ &\leq (L_{k+1})^{d \frac{L_{k+1}}{L_k}} \exp(-\zeta(P_g(x, y))) \sum_{N=0}^{\infty} L_{k+1}^{Nd} \exp(-\zeta NL_k) \\ &\leq \frac{1}{1 - (L_{k+1})^d \exp(-\zeta L_k)} (L_{k+1})^{d \frac{L_{k+1}}{L_k}} \exp(-\zeta(P_g(x, y))), \end{aligned}$$

because at each step i there are only L_{k+1}^d possible v_i 's. △

The fact that there are only 3 bad boxes implies that the good path length is close to the usual distance $P_g(x, y) \geq |x - y| - 20L_k$. We can now conclude. Let y be outside $\Lambda_{L_{k+1}}$. Then

$$\begin{aligned} &\sum_v |(\widehat{H}(n, L + 1) + A_c - \lambda)^{-1}(x, v)| |\Gamma(v, y)| \\ &\leq \sum_v \|((\widehat{H}(n, L + 1) + A_c - \lambda)^{-1})\| \\ &\quad \frac{1}{1 - 1 - (L_{k+1})^d \exp(-\zeta L_k)} (L_{k+1})^{L_{k+1}/L_k} \\ &\quad \exp(-\zeta(|x - v| - 20)) C \exp(-\nu|v - y|) \\ &\leq C'(L_{k+1})^d \exp(\log(L_k)L_k^{\alpha-1}) \exp(\sqrt{L_{k+1}}) \exp(-\zeta(|x - y| - 20)) \end{aligned}$$

In order to conclude, we just remark that

$$\sqrt{L_k} + \log(L_k)L_k^{\alpha-1} + d \log(L_k) = o(|x - y|).$$

So we can choose

$$\zeta_{k+1} = \zeta_k - \frac{\sqrt{L_k} + \log(L_k)L_k^{\alpha-1} + d \log(L_k)}{L_k}, \tag{60}$$

which finishes the argument. □

6.3. The multiscale

Theorem 6.1. *Assume that there exists a gap big enough in the spectrum, that the law of potential has a density Lipschitz by part and that $\|W\|_{L^1}$ is small enough compared to the gap. Then, let L_0 be large enough, $\lambda \in \mathbb{R}$ $\zeta > 1/\sqrt{L_0}$, $p > 2d$ and $1 < \alpha < 2p/(p + 2d)$. If for any cubes $\Lambda_{L_0}(n_0)$, $\Lambda_{L_0}(m_0)$ separated by at least $2L_0$,*

$$\begin{aligned} &\mathbb{P}(\text{there exists } \lambda \in I: \Lambda_{L_0}(n) \text{ and } \Lambda_{L_0}(m) \text{ are not } (L_0, \zeta, \lambda)\text{-good}) \\ &\leq \frac{1}{L_0^{2p}}, \end{aligned} \tag{61}$$

then for any k , and any cubes $\Lambda_{L_k}(n_k)$, $\Lambda_{L_k}(m_k)$ separated by at least $2L_k$, we have

$$\begin{aligned} &\mathbb{P}(\text{there exists } \lambda \in I: \Lambda_{L_k}(n_k) \text{ and } \Lambda_{L_k}(m_k) \text{ are not } (L_k, \zeta, \lambda) \text{ good}) \\ &\leq \frac{1}{L_k^{2p}}, \end{aligned} \tag{62}$$

where $L_{k+1} = L_k^\alpha$

This theorem is similar to Theorem 10.22 of [16].

Proof. The demonstration is done by iteration using Proposition 6.2. Suppose there exists $\lambda \in I$ such that $\Lambda_{L_{k+1}}(n)$ and $\Lambda_{L_{k+1}}(m)$ are not $(L_{k+1}, \zeta, \lambda)$ -good. Then for each box one of the hypothesis of Proposition 6.2 fails. So either one of the boxes admits 4 separated bad sub-boxes, or there exists λ such that for the two boxes, one of their sub-boxes shows a resonance at λ . The probability that the hypothesis over the existence of 4 bad boxes is not true can be estimated by iteration. Indeed, 4 cubes means 2 pairs. Because of independence, the 4-cubes

probability will be the 2-cubes probability squared and because there are only L^d cubes, hence L^{4d} 4 cubes combinations we can estimate this probability by

$$\begin{aligned} &\leq CL^{4d} \left(\frac{1}{(L_k)^{2p}} \right)^2 \\ &\leq C \frac{1}{(L_{k+1})^{4\frac{p}{\alpha} - 4d}} \\ &\leq \frac{1}{4} \frac{1}{L_{k+1}^{2p}}. \end{aligned}$$

The last inequality is true for L_k big enough because

$$2p < 4\frac{p}{\alpha} - 4d$$

so

$$\alpha < \frac{2p}{p + 2d}.$$

The probability of the non resonance hypothesis is controlled by Wegner estimate (12) as it is done Theorem 10.22 of [16] with $\mathcal{O}(L^d \sqrt{e^{-\sqrt{L_k}}}) = o(L^{-2p})$. □

From this and (12) we can deduce the following corollaries. The proof can be found again in [16, Part 9 and Part 11].

Corollary 6.1. *In presence of strong disorder, meaning*

$$(|\text{supp}(\rho)|^{-1/2} + \|\rho\|_\infty |\text{supp}(\rho)|^{1/2} + \|\rho'\|_\infty |\text{supp}(\rho)|^{3/2})$$

small enough, then H_{\min} has pure point spectrum and its eigenvectors are localised in space.

Furthermore the Lifshitz tail is not modified too much for $\|W\|$ very small because

$$\mathbb{P}[d(\sigma[(H_{\min})|_\Delta], \lambda) < \epsilon] \leq \mathbb{P}[d(\sigma[H^\Lambda], \lambda) < \epsilon + 2\|W\|_{\ell^1}], \tag{63}$$

so we also get the following

Corollary 6.2. *There is ϵ such that if $\|W\|_{\ell^1} < \epsilon$ there are small intervals at the edges of the bands of the spectrum of H_{\min} , where the spectrum is pure point with exponentially decaying eigenvectors.*

The two results conclude the proof of Theorem 2.2.

7. Numerical simulations

In this section, we present some simple numerical simulations in order to illustrate our theorems. Due to the computational cost we restrict ourselves to the one dimensional case, which however is known to present stronger localisation effects than in higher dimensions. It would be interesting to generalise our simulation to dimension 2 and 3.

We take $\Lambda = [0, L]$ with L ranging from two hundred to a few thousands lattice sites, $200 \leq L \leq 2000$. The discrete one-dimensional Laplacian is defined in (3). The deterministic potential V_0 is 2-periodic:

$$V_0(n) = \begin{cases} \xi & \text{if } n \text{ is even,} \\ -\xi & \text{if } n \text{ is odd,} \end{cases} \quad (64)$$

where $\xi > 0$ is a parameter. The probability of the random potential V_ω is the uniform law over the interval $[0, \zeta]$ where ζ is another parameter. For $W(x - y)$, we use a simple next-to-nearest neighbour interaction of the form

$$W(x - y) = \begin{cases} q & \text{if } x - y = 0, \\ q/2 & \text{if } |x - y| = 1, \\ q/4 & \text{if } |x - y| = 2 \text{ or } 3, \\ 0 & \text{otherwise,} \end{cases} \quad (65)$$

where q is another parameter. Our model depends therefore on three parameters ξ , ζ and q . When $\xi > 2 + \zeta$ the spectrum of the linear Hamiltonian $-\Delta + V$ is composed of two distinct intervals. We then choose q such as to keep a gap in the spectrum and ensure that the map F is contracting (Theorem 2.1). In this model the particles fill half of the energy states, that is, there are $N = L/2$ particles.

7.1. Illustration of Theorems 2.1, 2.3, 2.4 and 2.2 . In order to construct the solution γ_{\min} and the associated mean-field Hamiltonian H_{\min} , we use the fixed point algorithm employed in the proof of Theorem 2.1.

we have tested that adding a dirac at the site 250 to the potential V induces a perturbation in the non linear minimiser, which decays exponentially fast (Theorem 2.3). In Figure 2 we plot the relative density $\gamma_{\min}(V + \delta_{250})(x, x) - \gamma_{\min}(V)(x, x)$.

We conclude the illustration of our results with the case of Theorem 2.2. Since we deal with a one-dimensional system all the eigenvectors are localised, even in a regime of parameters which is not covered by the second part of Theorem 2.2. This is shown in Figure 3. It is an interesting open problem to prove a stronger localisation result in the one-dimensional Hartree–Fock model.

7.2. Closing the gap: insulators and metals. We have increased the intensity q of the interaction up to the point where the gap closes. For a large interaction the fixed point algorithm used to construct the solution in Theorem 2.1 does not work. Instead we have used the optimal damping algorithm of [12] which works perfectly. In general, we have observed that the eigenvectors are less localised except at the edges of the spectrum. There is no sign of a phase transition. The situation is more interesting if we erase the gap by choosing $\xi = 0$. A small delocalisation phenomenon seems to appear at the Fermi energy μ , see Figure 4. We hope to be able to understand these phenomena rigorously in the future.




Figure 2. The relative density $\gamma_{\min}(V + \delta_{250})(x, x) - \gamma_{\min}(V)(x, x)$ with $\xi = 1$, $\zeta = 1$, $q = 2$, $L = 500$.




Figure 3. Left: Standard deviation of all the eigenvectors of the mean field operator H_{\min} in terms of their eigenvalue. Here the size of the domain is $L = 1000$, hence a value of 10 shows localisation. Right: an eigenvector chosen at random. The values of the parameters are $\xi = 1$, $\zeta = 1$ and $q = 2$.




Figure 4. Standard deviation of the nonlinear eigenvectors in terms of their eigenvalue, with the periodic potential dropped out ($\xi = 0$). Here there are only $L/4$ particles. The Fermi level is $\mu = 3.5$ and there is no gap. A small delocalisation seems to appear at the Fermi level. The other parameters are $\zeta = 4$ and $q = 4$.

Acknowledgement. I would like to thank Mathieu Lewin for his help and all the time he gave me during this work.

References

- [1] M. Aizenman and S. Molchanov, Localization at large disorder and at extreme energies: an elementary derivation. *Comm. Math. Phys.* **157** (1993), no. 2, 245–278. [MR 1244867](#) [Zbl 0782.60044](#)
- [2] M. Aizenman and S. Warzel, Localization bounds for multiparticle systems. *Comm. Math. Phys.* **290** (2009), no. 3, 903–934. [MR 2525644](#) [Zbl 1186.82055](#)
- [3] P. W. Anderson, Absence of diffusion in certain random lattices. *Phys. Rev.* **109** (1958), 1492–1505.
- [4] V. Bach, Error bound for the Hartree–Fock energy of atoms and molecules. *Comm. Math. Phys.* **147** (1992), no. 3, 527–548. [MR 1175492](#) [Zbl 0771.46038](#)
- [5] V. Bach, E. H. Lieb, M. Loss, and J. P. Solovej, There are no unfilled shells in unrestricted Hartree–Fock theory. *Phys. Rev. Lett.* **72** (1994), no. 19, 2981–2983.

- [6] V. Bach, E. H. Lieb, and J. P. Solovej, Generalized Hartree–Fock theory and the Hubbard model. *J. Statist. Phys.* **76** (1994), no. 1-2, 3–89. [MR 1297873](#) [Zbl 0839.60095](#)
- [7] D. Basko, I. Aleiner, and B. Altshuler, Metal-insulator transition in a weakly interacting many-electron system with localized single-particle states. *Ann. of Phys.* **321** (2006), no. 5, 1126–1205. [Zbl 1091.82014](#)
- [8] D. Basko, I. Aleiner, and B. Altshuler, Metal-insulator transition in a weakly interacting many-electron system with localized single-particle states. *Ann. of Phys.* **321** (2006), no. 5, 1126–1205. [Zbl 1091.82014](#)
- [9] D. M. Basko, I. L. Aleiner, and B. L. Altshuler, *Problems of condensed matter physics*. Chapter “On the problem of many-body localization.” International Series of Monographs on Physics. Oxford University Press, Oxford, 2007.
- [10] D. Belitz and T. R. Kirkpatrick, The Anderson–Mott transition. *Rev. Mod. Phys.* **66** (1994), 261–380.
- [11] X. Blanc and M. Lewin, Existence of the thermodynamic limit for disordered quantum Coulomb systems. *J. Math. Phys.* **53** (2012), 095209. Special issue in honor of E. H. Lieb’s 80th birthday. [MR 2905791](#) [Zbl 1278.82031](#)
- [12] É. Cancès and C. Le Bris, On the convergence of SCF algorithms for the Hartree–Fock equations. *M2AN Math. Model. Numer. Anal.* **34** (2000), no. 4, 749–774. [MR 1784484](#) [Zbl 1090.65548](#)
- [13] E. Cancès, S. Lahbabi, and M. Lewin, Mean-field models for disordered crystals. *J. Math. Pures Appl.* (9) **100** (2013), no. 2, 241–274. [MR 3073215](#) [Zbl 1290.82015](#)
- [14] T. Chen and I. Rodnianski, Boltzmann limit for a homogeneous Fermi gas with Dynamical Hartree–Fock interactions in a random medium. *J. Stat. Phys.* **142** (2011), no. 5, 1000–1051. [MR 2781719](#) [Zbl 1220.82075](#)
- [15] V. Chulaevsky and Y. Suhov, Multi-particle Anderson localisation: induction on the number of particles. *Math. Phys. Anal. Geom.* **12** (2009), no. 2, 117–139. [MR 2497319](#) [Zbl 2497319](#)
- [16] M. Disertori, W. Kirsch, and A. Klein, *Random Schrödinger operators*. Panoramas et Synthèses, 25. Société Mathématique de France, Paris, 2008. [MR 2516524](#) [Zbl 1151.35002](#)
- [17] M. Fauser and S. Warzel, Multiparticle localization for disordered systems on continuous space via the fractional moment method. *Rev. Math. Phys.* **27** (2015), no. 4, 1550010, 42 pp. [MR 3350554](#) [Zbl 1335.47046](#)

- [18] J. Fröhlich and T. Spencer, Absence of diffusion in the Anderson tight binding model for large disorder or low energy. *Comm. Math. Phys.* **88** (1983), no. 2, 151–184. [MR 0696803](#) [Zbl 0519.60066](#)
- [19] I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, Interacting electrons in disordered wires: Anderson localization and low- T transport. *Phys. Rev. Lett.* **95** (2005), 206603.
- [20] J. Z. Imbrie, Multi-scale Jacobi method for Anderson localization. *Comm. Math. Phys.* **341** (2016), no. 2, 491–521. [MR 3440194](#) [Zbl 1337.82009](#)
- [21] A. Klein and S. Nguyen, The bootstrap multiscale analysis for the multi-particle Anderson model. *J. Stat. Phys.* **151** (2013), no. 5, 938–973. [JFM 1272.82021](#)
- [22] M. Könenberg, T. Moser, R. Seiringer, and J. Yngvason, Superfluid behavior of a Bose–Einstein condensate in a random potential. *New J. Phys.* **17** (2015), no. 1, 013022.
- [23] S. Lahbabi, *Mathematical study of quantum and classical models for random materials in the atomic scale*. Ph.D. thesis. Université de Cergy-Pontoise, Cergy-Pontoise, 2013.
- [24] S. Lahbabi, The reduced Hartree–Fock model for short-range quantum crystals with nonlocal defects. *Ann. Henri Poincaré* **15** (2014), no. 7, 1403–1452. [MR 3225735](#) [Zbl 1328.82052](#)
- [25] Y. Lahini, A. Avidan, F. Pozzi, M. Sorel, R. Morandotti, D. N. Christodoulides, and Y. Silberberg, Anderson localization and nonlinearity in one-dimensional disordered photonic lattices. *Phys. Rev. Lett.* **100** (2008), 013906.
- [26] E. H. Lieb and B. Simon, The Hartree–Fock theory for Coulomb systems. *Comm. Math. Phys.* **53** (1977), no. 3, 185–194. [MR 0452286](#)
- [27] P.-L. Lions, Solutions of Hartree–Fock equations for Coulomb systems. *Comm. Math. Phys.* **109** (1987), no. 1, 33–97. [MR 879032](#) [Zbl 0618.35111](#)
- [28] A. S. Pikovsky and D. L. Shepelyansky, Destruction of Anderson localization by a weak nonlinearity. *Phys. Rev. Lett.* **100** (2008), 094101.
- [29] M. Segev, Y. Silberberg, and D. N. Christodoulides, Anderson localization of light. *Nature Photonics* **7** (2013), no. 3, 197–204.
- [30] R. Seiringer and S. Warzel, Decay of correlations and absence of superfluidity in the disordered Tonks–Girardeau gas. *New J. Phys.* **18** (2016), March, 035002, 14 pp. [MR 3484875](#)
- [31] R. Seiringer, J. Yngvason, and V. A. Zagrebnov, Disordered Bose–Einstein condensates with interaction in one dimension. *J. Stat. Mech. Theory Exp.* **2012** (2012), no. 11, P11007.

- [32] H. Shiba, A Hartree–Fock theory of transition-metal impurities in a superconductor. *Progr. Theoret. Phys.* **50** (1973), no. 1, 50–73.
- [33] J. P. Solovej, The ionization conjecture in Hartree–Fock theory. *Ann. of Math. (2)* **158** (2003), no. 2, 509–576. [MR 2018928 Zbl 1106.81081](#)
- [34] K. Yamada, Perturbation expansion for the Anderson Hamiltonian. II. *Progr. Theoret. Phys.* **53** (1975), no. 4, 970–986.

Received March 31, 2016; revised August 7, 2016

Raphael Ducatez, CEREMADE, UMR CNRS 7534, Université Paris-Dauphine, Place de Lattre de Tassigny, 75775 Paris cedex 16, France

e-mail: ducatez@ceremade.dauphine.fr