

Conformal invariants from nodal sets.

II. Manifolds with boundary

Graham Cox,¹ Dmitry Jakobson,² Mikhail Karpukhin,³ and Yannick Sire

Abstract. In this paper, we study conformal invariants that arise from nodal sets and negative eigenvalues of conformally covariant operators on manifolds with boundary. We also consider applications to curvature prescription problems on manifolds with boundary. We relate Dirichlet and Neumann eigenvalues and put the results developed here for the Escobar problem into the more general framework of boundary operators of arbitrary order.

Mathematics Subject Classification (2010). 58J50, 53A30, 53A55, 53C21.

Keywords. Spectral geometry, conformal geometry, nodal sets, manifolds with boundary.

1. Introduction

The Yamabe problem on a closed Riemannian manifold is to find a metric of constant scalar curvature in a given conformal class; it generalizes the uniformization theorem for Riemann surfaces. It was solved by Yamabe, Trudinger, Aubin and Schoen. A related problem is to *prescribe* scalar curvature in a given conformal class; the two-dimensional version (prescribing Gauss curvature on a surface) is the *Nirenberg problem*. The problem of prescribing scalar curvature *without* fixing a conformal class was solved by Kazdan and Warner [26] (the solution is known as the *trichotomy theorem*). Fixing a conformal class makes the problem more difficult.

The conformal Laplacian appears naturally in the study of the Yamabe and curvature prescription problems. Nonzero eigenfunctions in its kernel provide obstructions to prescribing curvature; see [3, 4] for precise statements. In addition

¹ Graham Cox acknowledges the support of NSERC grant RGPIN-2017-04259.

² Dmitry Jakobson was supported by NSERC and FQRNT grants and Peter Redpath Fellowship of McGill University (Canada).

³ Mikhail Karpukhin was supported by Schulich Fellowship of McGill University (Canada) at initial stages of this project.

to the conformal Laplacian, in the papers [3, 4, 15] the authors considered general conformally covariant operators (including the Paneitz operator), and studied negative and zero eigenvalues of those operators on manifolds without boundary. They also considered applications to the associated curvature prescription problems (including the Q -curvature), and constructed conformal invariants determined by eigenfunctions in the kernel of the corresponding operator. In the current paper, we aim to generalize those results to manifolds with boundary.

The Yamabe problem on manifolds with boundary was solved in a series of works by Escobar [10, 11], Marques [29], Chen [8] and Mayer and Ndiaye [30]. They proved that on any compact manifold with boundary of dimension $n \geq 3$, there exists in every conformal class a scalar flat metric in the interior with constant mean curvature of the boundary.

The problem of *prescribing* scalar curvature R_g on the manifold M and mean curvature h_g on the boundary ∂M was considered in [13] and other papers. It was shown in [13, Corollary 2] that the solution attaining (R_g, h_g) is unique in the conformal class of $[g]$ if $R_g \leq 0$ and $h_g \leq 0$.

In the current paper we consider general conformally covariant operators on manifolds with boundary, and generalize the conformal invariants constructed in [3, 4] to this setting. We extend the results of [15], showing that 0 is generically *not* an eigenvalue in certain natural boundary value problems. We also answer some natural questions about the number of negative eigenvalues for conformally covariant boundary value problems.

Main results. We summarize below several important results of the present paper. The first theorem extends the main result of [15] to manifolds with boundary.

Theorem 1.1. *Generically, 0 is not an eigenvalue of the conformal Laplacian on a manifold with boundary.*

The proof follows easily from Proposition 3.1 of Section 3 by methods of [15].

The next result shows that the number of negative eigenvalues of the conformal Laplacian on manifolds with boundary can be made arbitrarily large.

Theorem 1.2. *For any compact manifold with boundary and any natural number m , there exists a conformal class \mathcal{C} such that for any metric $g \in \mathcal{C}$ one has $\lambda_m(P_{1,g}^R) < 0$, where the operator $P_{1,g}^R$ is defined below in (6).*

The proof in Section 4 uses the ideas of El Sayed [9].

The next result provides an obstruction to prescribing curvature in a conformal class $[g]$ with non-trivial $\ker(P_{1,g}^R)$, formulated as an inequality for a *single nodal domain* of a function $u \in \ker(P_{1,g}^R)$.

Theorem 1.3. *Let $(Q, f) \in \mathcal{R}([g])$ be the scalar curvature and the mean curvature of some metric in the conformal class $[g]$. Then, there is a pair of positive functions $\omega_i \in C^\infty(M)$, $\omega_b \in C^\infty(\partial M)$ such that for any nonzero $u \in \ker(P_{1,g}^R)$ and any nodal domain Ω of u ,*

$$\int_{\Omega} Q|u|\omega_i dv_g + \int_{\partial\Omega \cap \partial M} f|u|\omega_b d\sigma_g < 0.$$

Finally, Theorem 6.1 (the main result in Section 6) provides a generalization of a result in [14] to conformally covariant operators, relating the number of negative eigenvalues for the Dirichlet and Robin problems to the number of negative eigenvalues for a conformally covariant version of the Dirichlet-to-Neumann map.

Outline of the paper. In Section 2 we give examples of conformally covariant boundary value problems and describe natural extensions of the results in [3, 4] to manifolds with boundary. In Section 3, we show that generically the conformal Laplacian on manifolds with boundary does not have the eigenvalue 0. In Section 4, we show that the number of negative eigenvalues of the conformal Laplacian on manifolds with boundary can be arbitrarily large. In Section 5 we explore some natural curvature prescription problems (in a given conformal class) on manifolds with boundary; we also study the relationship between nodal sets of eigenfunctions in the kernel of the conformal Laplacian in the interior of the manifold, and the corresponding nodal sets on the boundary. In Section 6, we study inequalities between Dirichlet and Robin eigenvalues, and establish Friedlander-type results in the conformal setting. In Section 7 we study a continuous family of GJMS operators and their conformal extensions to the boundary of Poincaré–Einstein manifolds. In Section 8 we outline some natural questions that we have not considered in this paper.

2. Conformally covariant boundary value problems

A key observation used in [3, 4] (and many other papers) is that eigenfunctions that belong to the kernel of a conformally covariant operator (on a manifold without boundary) are multiplied by a positive function under a conformal change of the

metric. As a consequence, their nodal sets and nodal domains are preserved; other consequences of this fact were explored in [3, 4].

Here we consider a manifold M (of dimension $n \geq 3$) with smooth boundary ∂M . Given a conformally covariant operator P on M , we consider eigenfunctions $P\phi = \lambda\phi$ on M . We are interested in boundary value problems for P where eigenfunctions corresponding to $\lambda = 0$ transform in the same way as when $\partial M = \emptyset$.

We start with a discussion of the conformal Laplacian,

$$P_{1,g} = \Delta_g + \frac{n-2}{4(n-1)}R_g, \quad (1)$$

where Δ_g is the positive-definite Laplacian for g , and R_g is the scalar curvature. If $\hat{g} = e^{2\omega}g$ is a metric in the conformal class $[g]$, then $P_{1,g}$ transforms according to the law

$$P_{1,\hat{g}} = e^{-(n/2+1)\omega} P_{1,g} e^{(n/2-1)\omega}.$$

Accordingly,

$$\ker P_{1,\hat{g}} = e^{-(n/2-1)\omega} \ker P_{1,g}. \quad (2)$$

Consider now a boundary value problem for $P_{1,g}$ on $(M, \partial M)$. In the interior we have $P_{1,g}\phi = \lambda\phi$. Next, we need to specify boundary conditions on ∂M . We would like the transformation formula (2) to hold for eigenfunctions with $\lambda = 0$.

Standard boundary conditions include *Dirichlet BC*

$$\begin{cases} P_{1,g}\phi(x) = \lambda\phi(x), & x \in M, \\ \phi(x) = 0, & x \in \partial M, \end{cases} \quad (3)$$

and *Neumann BC*

$$\begin{cases} P_{1,g}\phi(x) = \lambda\phi(x), & x \in M, \\ \partial_\nu\phi(x) = 0, & x \in \partial M. \end{cases} \quad (4)$$

Here $\nu = \nu_g$ is the unit (with respect to the metric g) outer normal vector field on ∂M . We remark that the direction of ν is preserved under conformal transformations.

Example 2.1. If 0 is an eigenvalue of $P_{1,g}$ on a closed manifold, then it is an eigenvalue of the Dirichlet problem on any nodal domain of the corresponding eigenfunction.

We next state the first result about boundary value problems:

Proposition 2.2. *Let $\hat{g} = e^{2\omega} g$ be two metrics on M lying in the same conformal class.*

- (a) *Solutions of the Dirichlet BVP (3) with $\lambda = 0$ transform under the rule (2).*
- (b) *Solutions of the Neumann BVP (4) with $\lambda = 0$ transform under the rule (2), provided ω satisfies $\partial_\nu \omega(x) = 0$ for all $x \in \partial M$.*

Proof. The condition (3) is clearly invariant under the transformation (2). For the condition (4), we remark that

$$\partial_\nu(e^{-(n/2-1)\omega} \phi) = e^{-(n/2-1)\omega} ((1 - n/2)\partial_\nu \omega + \partial_\nu \phi)$$

and both expressions in the bracket vanish if $\partial_\nu \omega = 0$. □

It follows that choosing boundary conditions as in Proposition 2.2 allows one to develop the theory analogous to that in [3, 4]. In particular, the following Proposition can be proved easily by the methods of [3].

Proposition 2.3. *Let P be a conformally covariant operator on $(M, \partial M)$ whose kernel transforms according to (2). Let g be a Riemannian metric such that $\dim \ker P_g \geq 1$. Then the following quantities give rise to invariants of the conformal class $[g]$.*

- (1) *The dimension k of $\ker P_g$*
- (2) *The number of negative eigenvalues of P_g*
- (3) *Nodal sets $\mathcal{N}(u)$ and nodal domains of any eigenfunction $u \in \ker P_g$*
- (4) *Non-empty intersections of nodal sets of eigenfunctions in $\ker P_g$ and their complements, provided $k \geq 2$*
- (5) *Let $k \geq 2$, and let (u_1, \dots, u_k) be a basis of $\ker P_g$. Let*

$$\tilde{M} = M \setminus \left(\bigcap_{i=1}^k \mathcal{N}(u_i) \right).$$

Define $\Phi_g: \tilde{M} \rightarrow \mathbb{R}P^{k-1}$ by

$$\Phi_g(x) = (u_1(x) : \dots : u_k(x)).$$

Then the orbit of $\Phi_g(\tilde{M})$ in $\mathbb{R}P^{k-1}$ under the action of $GL_k(\mathbb{R})$ is conformally invariant.

- (6) *The same results hold for ∂M and the restrictions of eigenfunctions to ∂M .*
- (7) *Let $p = 2n/(n - 2)$, and let $u \in \ker P_g$. Then the density $|u(x)|^p d\text{vol}(x)$ is conformally invariant. In particular, the following quantities are conformally invariant: $\|u\|_p$, as well as $\int_{\Omega} |u(x)|^p d\text{vol}(x)$, where Ω is any domain in M . Similar results hold for restrictions of u to submanifolds N of M , with suitably adjusted exponents $p = p(N)$.*

Next, consider the boundary operator B_g given by

$$B_g u(x) = \partial_{\nu_g} u(x) + \frac{n - 2}{2(n - 1)} h_g u(x),$$

where $h_g = \text{tr}_g II$ is the mean curvature of the boundary. We have

Proposition 2.4. *The operator B_g is conformally covariant, i.e.*

$$B_{e^{2\omega} g}(e^{(-\frac{n}{2}+1)\omega} u) = e^{-\frac{n}{2}\omega} B_g u.$$

Proof. By definition, the second fundamental form is given by $II_{ij} = \frac{1}{2} \partial_{\nu} g_{ij}$. Thus, one has the following formula for the mean curvature

$$h_g = \frac{1}{2} (\partial_{\nu_g} g_{ij}) g^{ij}. \tag{5}$$

Note that $\nu_{e^{2\omega} g} = e^{-\omega} \nu_g$. Therefore, formula (5) implies

$$\begin{aligned} h_{e^{2\omega} g} &= \frac{1}{2} e^{-3\omega} (\partial_{\nu_g} (e^{2\omega} g_{ij})) g^{ij} \\ &= e^{-\omega} (\partial_{\nu_g} \omega) g^{ij} g_{ij} + \frac{1}{2} e^{-\omega} (\partial_{\nu_g} g_{ij}) g^{ij} \\ &= e^{-\omega} ((n - 1) \partial_{\nu_g} \omega + h_g). \end{aligned}$$

Applying this transformation law to B_g , we obtain

$$\begin{aligned} &B_{e^{2\omega} g}(e^{(-\frac{n}{2}+1)\omega} u) \\ &= e^{-\omega} \partial_{\nu_g}(e^{(-\frac{n}{2}+1)\omega} u) + e^{-\omega} \frac{n - 2}{2(n - 1)} ((n - 1) \partial_{\nu_g} \omega + h_g) e^{(-\frac{n}{2}+1)\omega} u \\ &= e^{-\frac{n}{2}\omega} B_g u. \end{aligned} \quad \square$$

Proposition 2.4 allows us to define two types of conformally covariant eigenvalue problems on M .

Conformal Robin BVP. We set

$$\begin{cases} P_{1,g}\phi(x) = \lambda\phi(x), & x \in M, \\ B_g\phi(x) = 0, & x \in \partial M. \end{cases} \tag{6}$$

Proposition 2.4 implies that solutions of the Conformal Robin BVP (6) with $\lambda = 0$ transform under the rule (2). In the following we denote the operator $P_{1,g}$ with conformal Robin BC as $P_{1,g}^R$.

Conformal Steklov BVP (conformal Dirichlet-to-Robin map). Suppose that $\lambda = 0$ is *not* an eigenvalue of the Dirichlet BVP (3). Then for any $u \in C^\infty(\partial M)$ there exists a unique solution $\mathcal{H}_g(u)$ (see e.g. [27, §1.3]) to

$$\begin{cases} P_{1,g}(\mathcal{H}_g(u))(x) = 0, & x \in M, \\ \mathcal{H}_g(u) = u, & x \in \partial M. \end{cases} \tag{7}$$

We define the *Conformal Dirichlet-to-Robin map* $\mathcal{D}_g: C^\infty(\partial M) \rightarrow C^\infty(\partial M)$ as

$$\mathcal{D}_g(u) = B_g(\mathcal{H}_g(u)). \tag{8}$$

We have then

Proposition 2.5. *The operator \mathcal{D}_g is conformally covariant, i.e.*

$$\mathcal{D}_{e^{2\omega}g}(e^{(-\frac{n}{2}+1)\omega}u) = e^{-\frac{n}{2}\omega}\mathcal{D}_g(u).$$

Proof. Conformal covariance of $P_{1,g}$ implies that

$$\mathcal{H}_{e^{2\omega}g}(e^{(-\frac{n}{2}+1)\omega}u) = e^{(-\frac{n}{2}+1)\omega}\mathcal{H}_g(u).$$

Proposition 2.4 and the definition of \mathcal{D}_g then imply

$$\begin{aligned} \mathcal{D}_{e^{2\omega}g}(e^{(-\frac{n}{2}+1)\omega}u) &= B_{e^{2\omega}g}(e^{(-\frac{n}{2}+1)\omega}\mathcal{H}_g(u)) \\ &= e^{-\frac{n}{2}\omega}B_g(\mathcal{H}_g(u)) \\ &= e^{-\frac{n}{2}\omega}\mathcal{D}_g(u). \end{aligned} \quad \square$$

Remark 2.6. It is easy to see that $(\ker P_{1,g}^R)|_{\partial M} = \ker \mathcal{D}_g$. Also, the number of negative eigenvalues of \mathcal{D}_g is equal to the number of negative eigenvalues of the Robin boundary value problem (6) minus the number of negative eigenvalues for the Dirichlet problem (3). This is proved in Theorem 6.1. Both of these claims remain true when 0 is a Dirichlet eigenvalue, provided \mathcal{D}_g is replaced by the operator $\hat{\mathcal{D}}_g$ defined below.

If 0 is an eigenvalue of the Dirichlet problem (3), then $\ker P_{1,g}^D$ is nontrivial. Define

$$S = \left\{ u \in C^\infty(\partial M) : \int_{\partial M} u \partial_\nu w \, d\sigma_g = 0 \text{ for all } w \in \ker P_{1,g}^D \right\}.$$

The boundary value problem (7) has a solution for any $u \in S$. Moreover, the solution is unique if we require it to be orthogonal to $\ker P_{1,g}^D$ in $L^2(M)$. Denoting this unique solution by $\hat{\mathcal{H}}_g(u)$, we define the operator $\hat{\mathcal{D}}_g : S \rightarrow S$ by

$$\hat{\mathcal{D}}_g u = \Pi_S B_g(\hat{\mathcal{H}}_g(u)), \tag{9}$$

where Π_S is the $L^2(\partial M)$ -orthogonal projection onto \bar{S} . If $w \in \ker P_{1,g}^D$, then $B_g w = \partial_\nu w \in S^\perp$, so $\Pi_S(B_g w) = 0$ and we in fact have

$$\hat{\mathcal{D}}_g u = \Pi_S B_g(\psi)$$

where ψ is any solution to the boundary value problem (7). Alternatively, one can easily show that there exists a unique solution $\tilde{\mathcal{H}}_g(u)$ to (7) with the property that $B_g(\tilde{\mathcal{H}}_g(u)) \in S$, and so

$$\hat{\mathcal{D}}_g u = B_g(\tilde{\mathcal{H}}_g(u)).$$

3. Generically there is no zero eigenvalue: proof of Theorem 1.1

Let g be a metric such that $\dim \ker P_{1,g}^R = k > 0$. For simplicity we assume $R_g \equiv -1$ and $h_g \equiv 0$. The goal of this section is the following proposition.

Proposition 3.1. *There exists $g_t = g + t\tilde{g}$ such that for all small enough $t \ll 1$ one has $\dim \ker P_{1,g_t}^R < k$.*

Repeating the deformation inductively one gets that for a generic metric, 0 is not an eigenvalue of $P_{1,g}^R$. Our proof closely follows [15], where the same statement is proved for closed manifolds.

Our main observation is that if $\tilde{g} \equiv 0$ in a neighbourhood of ∂M , then the computations of [15, §4] follow through verbatim. Let $E_0 = \ker P_{1,g}^R$ be the 0-eigenspace and let $c_n = \frac{n-2}{4(n-1)}$. Consider the operator

$$Q_{g,\tilde{g}} = \Pi_{E_0} \frac{d}{dt} \Big|_{t=0} (\Delta_{g_t} + c_n R_{g_t}) \Big|_{E_0}$$

on E_0 . Then Proposition 3.1 is equivalent to the statement that $Q_{g,\tilde{g}} \neq 0$.

In [15] one obtains that if $\text{tr}_g \tilde{g} = 0$, then

$$(Q_{g, \tilde{g}} \psi, \psi) = (\tilde{g}, c_n \psi (2\text{Hess}^\circ(\psi) - \psi \text{Ric}^\circ) + (2c_n - 1)(d\psi \otimes d\psi)^\circ),$$

where \circ denotes the traceless part of a bilinear form.

Assume that $q = c_n \psi (2\text{Hess}^\circ(\psi) - \psi \text{Ric}^\circ) + (2c_n - 1)(d\psi \otimes d\psi)^\circ \neq 0$. Then there exists $p \in \text{int}M$ such that $q(p) \neq 0$. Let $p \in U \subset \text{int}M$ be a neighbourhood of p and let $\rho_\varepsilon \in C_0^\infty(U)$ be a cut-off function such that $\rho_\varepsilon(p) = 1$. Set $\tilde{g} = \rho_\varepsilon q$, then $(Q_{g, \tilde{g}} \psi, \psi) > 0$ for $\varepsilon \ll 1$, i.e. $Q_{g, \tilde{g}} \neq 0$.

Assume now that $q \equiv 0$. Then, since $R_g \equiv -1$ and $h_g \equiv 0$, ψ is an eigenfunction of Δ_g with eigenvalue c_n , i.e it is orthogonal to constant function. As a result, the nodal set $N(\psi)$ intersects $\text{int} M$ nontrivially. Let us restrict $q = 0$ to $N(\psi) \cap \text{int} M$. Then, by definition of q one obtains $(d\psi \otimes d\psi)^\circ = 0$ there. The latter is equivalent to

$$d\psi \otimes d\psi = \frac{1}{n} |\psi|_g^2 g,$$

which implies $d\psi \equiv 0$. This is impossible by, e.g. [20, 21, 22], as in [15].

4. Number of negative eigenvalues: proof of Theorem 1.2

The proof is very similar to the proof of Proposition 3.2 in the paper of El Sayed [9]. Let g_0 be some metric on M . The Rayleigh quotient of $P_{1,g}^R$ is given by

$$R_g[u] = \frac{\int_M |\nabla u|^2 dv_g + \frac{n-2}{4(n-1)} \int_M R_g u^2 dV_g + \frac{n-2}{2(n-1)} \int_{\partial M} h_g u^2 d\sigma_g}{\int_M u^2 dV_g}.$$

The plan is to locally change the metric g_0 to a new metric g_p around a point p to create a function u_p supported in the neighbourhood of p such that $R_{g_p}[u_p] < 0$. Doing so at m different points creates an m -dimensional negative subspace for the Rayleigh quotient.

Let $p_i, i = 1, \dots, m$ be a collection of points in the interior of M and let $p_i \in U_i$ be sufficiently small pairwise disjoint geodesic balls lying in the interior of M . We identify each U_i with a copy of $\mathbb{S}^n \setminus \{p\}$ for some fixed point $p \in \mathbb{S}^n$. By a result of Aubin [2] there exists metric g on \mathbb{S}^n such that the Yamabe invariant $Y_g(\mathbb{S}^n) < 0$. Therefore there exists a function $u \in C^\infty(\mathbb{S}^n)$ such that $R_g[u] < -2\varepsilon$ for some $\varepsilon > 0$. Let $V_\delta \subset V_{2\delta} \subset \mathbb{S}^n$ be two small balls around p small enough

so that there exists a cut-off function φ_δ with $\varphi_\delta \equiv 1$ on $S^n \setminus V_{2\delta}$, $\varphi_\delta = 0$ on V_δ , $0 \leq \varphi_\delta \leq 1$ and $\|\nabla\varphi_\delta\|_{L^2_g(S^n)}^2 \leq \delta$. As $\dim M \geq 3$, such balls exist for all $\delta > 0$. Then one has $R_g[u\varphi_\delta] \rightarrow R_g[u]$ as $\delta \rightarrow 0$.

Thus, for small enough $\delta > 0$ one has $R_g[u\varphi_\delta] < -\varepsilon$. We let u_i be the functions defined as $u\varphi_\delta$ in the neighbourhood U_i of p_i . We define the metric g_k on M to be any metric equal to g on $U_i \setminus V_\delta$, where V_δ lies in the corresponding copy of S^n . Then there exist pairwise disjoint functions u_1, \dots, u_m such that $R_{g_n}[u_i] < 0$ for all i . Thus, the same is true for any linear combination of these functions, i.e. $\lambda_m(P_{1,g_n}^R) < 0$.

5. Curvature prescription

5.1. Background. A natural extension of the Yamabe problem on a manifold M with boundary ∂M is to prescribe the scalar curvature R_g in the interior of M and the mean curvature h_g on ∂M . Existence and uniqueness for this problem was considered e.g. in the papers [12, 13]. In the paper [4, §5], the authors obtained several results that give obstructions to prescribing curvature in terms of eigenfunctions lying in the kernel of the conformal Laplacian. In this section, we generalize those results to manifolds with boundary.

We start with the following result:

Theorem 5.1. *For any $u \in \ker P_{1,g}^R$ one has the following identity*

$$\frac{n-2}{4(n-1)} \int_M R_g u \, dv_g + \frac{n-2}{2(n-1)} \int_{\partial M} h_g u \, d\sigma_g = 0.$$

Proof. By definition one has $\frac{n-2}{4(n-1)} R_g = P_{1,g}(1)$. Integrating by parts we obtain

$$\int_M P_{1,g}(1)u \, dv_g = \int_M P_{1,g}(u) \, dv_g + \int_{\partial M} \partial_n u \, d\sigma_g = -\frac{n-2}{2(n-1)} \int_{\partial M} h_g u \, d\sigma_g. \quad \square$$

This theorem has consequences for the curvature prescription problem which we describe below. For a fixed conformal class of metrics $\mathcal{C} = [g]$ one defines the following curvature map

$$\mathcal{R}: \mathcal{C} \rightarrow C^\infty(M) \times C^\infty(\partial M), \quad \mathcal{R}(g) = (R_g, h_g).$$

Our goal is to determine the image $\mathcal{R}(\mathcal{C})$. It turns out that non-trivial elements $u \in \ker(P_{1,g}^R)$ constitute obstructions to this curvature prescription problem. An easy example is the following corollary.

Corollary 5.2. *For any $u \in \ker P_{1,g}^R \setminus \{0\}$, the pair $(u, u|_{\partial M})$ does not belong to $\mathcal{R}(\mathcal{C})$.*

Proof. By Theorem 5.1 and formula (2), for any metric $\hat{g} = e^{2\omega}g$ one has

$$\frac{n-2}{4(n-1)} \int_M R_{\hat{g}} e^{-(n/2-1)\omega} u \, dv_{\hat{g}} + \frac{n-2}{2(n-1)} \int_{\partial M} h_{\hat{g}} e^{-(n/2-1)\omega} u \, d\sigma_{\hat{g}} = 0.$$

Assume the contrary, i.e. $\mathcal{R}(\hat{g}) = (u, u|_{\partial M})$. Then the previous formula implies

$$\frac{n-2}{4(n-1)} \int_M e^{-(n/2-1)\omega} u^2 \, dv_{\hat{g}} + \frac{n-2}{2(n-1)} \int_{\partial M} e^{-(n/2-1)\omega} u^2 \, d\sigma_{\hat{g}} = 0,$$

which is impossible. □

Similarly, any pair that has the same or the opposite strict signs as $(u, u|_{\partial M})$ can not belong to $\mathcal{R}(\mathcal{C})$.

Corollary 5.3. *Let M be a manifold with boundary equipped with a conformal class \mathcal{C} , let $g \in \mathcal{C}$ and $u \in \ker P_{1,g}^R \setminus \{0\}$. Then any pair (Q, f) such that (Q, f) has the same or the opposite strict sign as $(u, u|_{\partial M})$ can not belong to $\mathcal{R}(\mathcal{C})$.*

Examples of such pairs include $(e^{\omega_1}u^{p_1}, e^{\omega_2}u^{p_2}|_{\partial M})$ for any pair of odd natural numbers (p_1, p_2) and $(\omega_1, \omega_2) \in C^\infty(M) \times C^\infty(\partial M)$.

5.2. Obstructions from nodal domains. Next, we would like to generalize [4, Thm. 5.5, Cor. 5.6] to manifolds with boundary. There is an extensive literature studying the relationship between the spectral theory of conformally covariant operators on a manifold M , and the behaviour of a related conformally covariant operators on ∂M , see e.g. [18, 5, 7, 16]. The following discussion deals with the relationship between nodal sets of eigenfunctions of a conformally covariant operator P on M , and the nodal sets of eigenfunctions of a related operator Q on ∂M . Clearly, if the nodal set $\mathcal{N}(u)$ of an eigenfunction u of P intersects the boundary, the limiting value of u on ∂M is equal to 0. Conversely, let us suppose that $u(x)$ is an eigenfunction of a conformally covariant BVP, and that $u(y) = 0$ for some $y \in \partial M$. A natural question is whether there exists a sequence $x_i \in \text{int } M$ such that $u(x_i) = 0$ and $x_i \rightarrow y$. In other words: is a nodal set on the boundary always a limit of an interior nodal set? This is the case by a simple application of the strong maximum principle. Indeed, assume the contrary, i.e. that $u(y) = 0$ but $u(x) > 0$ for x in the pointed neighbourhood of y . Then by Hopf's lemma (see, e.g. [24, Theorem 2.8(b)]) one has $\partial_\nu u(y) < 0$. At the same time, the

boundary condition $B_g(u)(y) = \partial_\nu u(y) + \frac{n-2}{2(n-1)}h_g u(y) = 0$ yields $\partial_\nu u(y) = 0$, a contradiction.

We now describe the obstruction in question. We prove

Theorem 5.4. *Let $u \in \ker(P_{1,g}^R)$ and let Ω be a nodal domain of u . Then, for all $v \in C^\infty(M)$*

$$\int_{\Omega} |u| P_{1,g}(v) dv_g = - \int_{\partial\Omega \setminus \partial M} v |\nabla u|_g d\sigma_g - \int_{\partial\Omega \cap \partial M} |u| B_g(v) d\sigma_g. \tag{10}$$

Proof. Up to a change of sign, we can assume that $u > 0$ on Ω . Integrating by parts we obtain

$$\int_{\Omega} u P_{1,g}(v) dv_g = \int_{\Omega} v P_{1,g}(u) dv_g - \int_{\partial\Omega} (u \partial_\nu v - v \partial_\nu u) d\sigma_g.$$

Decomposing the second summand on the right hand side into the sum of integrals over $\partial\Omega \cap \partial M$ and $\partial\Omega \setminus \partial M$ we obtain

$$\int_{\partial\Omega \cap \partial M} (u \partial_\nu v - v \partial_\nu u) d\sigma_g = \int_{\partial\Omega \cap \partial M} u \left(\partial_\nu v + \frac{n-2}{2(n-1)} h_g v \right) d\sigma_g,$$

$$\int_{\partial\Omega \setminus \partial M} (u \partial_\nu v - v \partial_\nu u) d\sigma_g = - \int_{\partial\Omega \setminus \partial M} v \partial_\nu u d\sigma_g.$$

Since on $\partial\Omega \setminus \partial M$ one has $\partial_\nu u = -|\nabla u|_g$ the proof is complete. □

Theorem 5.5. *Let $(Q, f) \in \mathcal{R}([g])$ be the scalar curvature and mean curvature of some metric in the conformal class $[g]$. Then, there is a pair of positive functions $\omega_i \in C^\infty(M)$, $\omega_b \in C^\infty(\partial M)$ such that for any $u \in \ker(P_{1,g}^R)$ and any nodal domain Ω of u ,*

$$\int_{\Omega} Q |u| \omega_i dv_g + \int_{\partial\Omega \cap \partial M} f |u| \omega_b d\sigma_g < 0.$$

Proof. By Theorem 5.4 one has

$$\int_{\Omega} |u| P_{1,g}(v) dv_g + \int_{\partial\Omega \cap \partial M} |u| B_g(v) d\sigma_g < 0$$

for any positive function $v \in C^\infty(M)$. Let $\hat{g} = e^{2\omega}g$ be the metric such that $\mathcal{R}(\hat{g}) = (Q, f)$. Let us apply the previous inequality for $v = 1$ and the metric \hat{g} , taking into account the transformation law (2)

$$\frac{n-2}{4(n-1)} \int_{\Omega} e^{-(n/2-1)\omega} |u| Q \, dv_{\hat{g}} + \frac{n-2}{2(n-1)} \int_{\partial\Omega \cap \partial M} e^{-(n/2-1)\omega} |u| f \, d\sigma_{\hat{g}} < 0.$$

Equivalently,

$$\frac{n-2}{4(n-1)} \int_{\Omega} e^{(n/2+1)\omega} |u| Q \, dv_g + \frac{n-2}{2(n-1)} \int_{\partial\Omega \cap \partial M} e^{\frac{n+1}{2}\omega} |u| f \, d\sigma_g < 0.$$

Setting $\omega_i = \frac{n-2}{4(n-1)}e^{(n/2+1)\omega}$ and $\omega_b = \frac{n-2}{2(n-1)}e^{\frac{n+1}{2}\omega}$ completes the proof. \square

Corollary 5.6. For any $(Q, f) \in \mathcal{R}([g])$ and any nodal domain Ω of $u \in \ker(P_{1,g}^R)$, either Q or f must be negative for at least one point of $\bar{\Omega}$.

Theorem 5.7. Let $u \in \ker P_{1,g}^R$ and let Ω be a nodal domain of u . Then for any $\hat{g} = e^{2\omega}g \in [g]$ one has

$$-\int_{\partial\Omega \setminus \partial M} e^{(1-\frac{n}{2})\omega} |\nabla \hat{u}|_{\hat{g}} \, d\sigma_{\hat{g}} = \frac{n-2}{4(n-1)} \int_{\Omega} |u| R_g \, dv_g + \frac{n-2}{2(n-1)} \int_{\partial\Omega \cap \partial M} h_g |u| \, d\sigma_g,$$

i.e. the expression on the left hand side is a conformal invariant.

Proof. Applying Theorem 5.4 for the metric \hat{g} and $v = e^{(1-\frac{n}{2})\omega}$ yields

$$-\int_{\partial\Omega \setminus \partial M} e^{(1-\frac{n}{2})\omega} |\nabla \hat{u}|_{\hat{g}} \, d\sigma_{\hat{g}} = \int_{\Omega} \hat{u} |P_{1,\hat{g}}(v)| \, dv_{\hat{g}} + \int_{\partial\Omega \cap \partial M} \hat{u} |B_{\hat{g}}(v)| \, d\sigma_{\hat{g}}.$$

By the transformation laws for $P_{1,g}$ and B_g one has $P_{1,\hat{g}}(v) = e^{-(n/2+1)\omega} P_{1,g}(1)$ and $B_{\hat{g}} = e^{-\frac{n}{2}\omega} B_g(1)$. Therefore one has

$$\begin{aligned} -\int_{\partial\Omega \setminus \partial M} e^{(1-\frac{n}{2})\omega} |\nabla \hat{u}|_{\hat{g}} \, d\sigma_{\hat{g}} &= \int_{\Omega} |e^{-(n/2-1)\omega} u| e^{-(n/2+1)\omega} P_{1,g}(1) e^{n\omega} \, dv_g \\ &\quad + \int_{\partial\Omega \cap \partial M} |e^{-(n/2-1)\omega} u| e^{-\frac{n}{2}\omega} B_g(1) e^{(n-1)\omega} \, d\sigma_g. \quad \square \end{aligned}$$

Remark 5.8. Suppose there is a function $u \in \ker P_{1,g}$ (not necessarily $\ker P_{1,g}^R$!) such that it has a nodal domain $\Omega \Subset M$. Then by the maximum principle for Schrödinger operators, for any $(Q, f) \in \mathcal{R}([g])$ the function Q must change sign inside Ω .

6. Inequalities between Dirichlet and Neumann eigenvalues

In the paper [14], Friedlander obtained inequalities between Dirichlet and Neumann eigenvalues of the Laplacian using the equality

$$N_N(\lambda) - N_D(\lambda) = N_-(\mathcal{D}(\lambda)), \tag{11}$$

where $N_N(\lambda)$ (resp. $N_D(\lambda)$) denotes the Neumann (resp. Dirichlet) eigenvalue counting function, where $\mathcal{D}(\lambda)$ denotes the Dirichlet-to-Neumann operator, and where $N_-(\mathcal{D}(\lambda))$ denotes the number of its negative eigenvalues. This equality was given a geometric interpretation by Mazzeo in [31].

We prove a Friedlander-type result for the conformal Laplacian. Let N_R and N_D denote the number of negative eigenvalues for the conformal Laplacian with Robin and Dirichlet boundary conditions, respectively, and recall the definition (9) of the conformal Dirichlet-to-Neumann operator $\widehat{\mathcal{D}}_g$.

Theorem 6.1. $N_R - N_D = N_-(\widehat{\mathcal{D}}_g) + \dim \ker P_{1,g}^D$.

Proof. For each $s \in [0, \infty)$ consider the operator $P_{1,g}(s)$ defined to be the conformal Laplacian $P_{1,g}$ with boundary conditions

$$B_g u + s u = 0.$$

This is the unbounded operator on $L^2(M)$ generated by the bilinear form

$$a_s(u, v) = \int_M \left(\nabla u \cdot \nabla v + \frac{n-2}{4(n-1)} R_g u v \right) dv_g + \int_{\partial M} \left(\frac{n-2}{2(n-1)} h_g + s \right) u v d\sigma_g$$

with form domain $H^1(M)$. From this one sees that each $P_{1,g}(s)$ is self-adjoint and the family is analytic with respect to s .

We show that the eigenvalues of $P_{1,g}(s)$ are strictly increasing. Suppose $\lambda = \lambda(s)$ is an analytic curve of eigenvalues, with normalized eigenfunctions $u = u(s)$, so that $a_s(u, v) = \lambda \langle u, v \rangle$ for all $v \in H^1(M)$. Substituting $v = u'$ yields

$$a_s(u, u') = \lambda \langle u, u' \rangle.$$

On the other hand, differentiating and then letting $v = u$ yields

$$a'_s(u, u) + a_s(u', u) = \lambda \langle u', u \rangle + \lambda' \|u\|^2$$

and hence

$$\lambda' = a'_s(u, u) = \int_{\partial M} u^2,$$

which is strictly positive. (If u vanishes on ∂M , the boundary condition would imply that $\partial_\nu u$ also vanishes, which contradicts the unique continuation principle.)

Note that $P_{1,g}(0) = P_{1,g}^R$ has N_R negative eigenvalues. On the other hand, the proof of [1, Theorem 2.4] implies that the k th ordered eigenvalue $\lambda_k(s)$ tends to the k th Dirichlet eigenvalue λ_k^D as $s \rightarrow \infty$. This implies $P_{1,g}(s)$ has $N_D + \dim \ker P_{1,g}^D$ negative eigenvalues for all sufficiently large s .

Therefore, the difference $N_R - N_D - \dim \ker P_{1,g}^D$ equals the number of eigenvalue curves that have $\lambda(s) = 0$ for some finite $s > 0$. This occurs when there is a nontrivial solution to the Robin boundary value problem

$$\begin{aligned} P_{1,g}\psi &= 0 && \text{in } M, \\ B_g\psi + s\psi &= 0 && \text{on } \partial M. \end{aligned} \tag{12}$$

If 0 is not a Dirichlet eigenvalue, this is equivalent to $\mathcal{D}_g u + su = 0$, where $u = \psi|_{\partial M}$. Moreover, the multiplicity of $\lambda = 0$ as an eigenvalue of the above Robin problem equals the multiplicity of $-s$ as an eigenvalue of \mathcal{D}_g .

When 0 is a Dirichlet eigenvalue we have to consider $\widehat{\mathcal{D}}_g$ instead of \mathcal{D}_g . If ψ solves (12), it must be the case that $u = \psi|_{\partial M} \in S$, hence

$$\widehat{\mathcal{D}}_g u = \Pi_S B_g \psi = -s \Pi_S u = -su.$$

Therefore $-s$ is an eigenvalue of $\widehat{\mathcal{D}}_g$. If ψ_1 and ψ_2 are linearly independent solutions to (12), the corresponding functions u_1 and u_2 on the boundary must be linearly independent as well. Otherwise, there would exist a linear combination $\psi = c_1\psi_1 + c_2\psi_2$ with $P_{1,g}\psi = 0$, $\psi|_{\partial M} = 0$, and $B_g\psi = 0$, hence $\partial_\nu\psi = 0$, which is only possible if $\psi \equiv 0$, a contradiction. Therefore, the multiplicity of $-s$ is at least as large as the multiplicity of $\lambda = 0$ for (12).

Conversely, if $-s$ is an eigenvalue of $\widehat{\mathcal{D}}_g$, with eigenfunction u , there exists a function ψ on M such that $P_{1,g}\psi = 0$ and $\Pi_S B_g\psi = -su$. Given ψ , there exists a unique $v \in \ker P_{1,g}^D$ so that $B_g(\psi + v) \in S$. Then $P_{1,g}(\psi + v) = 0$ and

$$B_g(\psi + v) = \Pi_S B_g(\psi + v) = \Pi_S B_g(\psi) = -s\psi|_{\partial M} = -s(\psi + v)|_{\partial M},$$

hence 0 is an eigenvalue of (12), with eigenfunction $\psi + v$.

Suppose u_1 and u_2 are linearly independent eigenfunctions of $\widehat{\mathcal{D}}_g$, with the same eigenvalue $-s$. If the corresponding eigenfunctions $\psi_i + v_i$ of (12) are not linearly independent, there will exist constants c_1 and c_2 so that we have $c_1(\psi_1 + v_1) + c_2(\psi_2 + v_2) = 0$. Restricting to the boundary yields $c_1u_1 + c_2u_2 = 0$, a contradiction. This proves that the multiplicity of $\lambda = 0$ for (12) equals the multiplicity of $-s$ for $\widehat{\mathcal{D}}_g$, and thus completes the proof. \square

Remark 6.2. Repeating the above argument with $P_{1,g}$ replaced by $P_{1,g} - \lambda$, we obtain a similar equality in terms of the counting functions $N_R(\lambda)$ and $N_D(\lambda)$ for the conformal Laplacian. However, these quantities are only conformally invariant when $\lambda = 0$.

7. A continuous family of GJMS operators and their conformal extensions to Poincaré–Einstein manifolds

In this section, we put the previous results in a much more general framework which has been developed in recent years. First we consider higher order extensions of the conformal Laplacian, i.e. the so-called GJMS operators [17]. The original construction of these operators did not provide useful information on their analytic properties, due to the nature itself of the construction. The important result on their analytic properties came with the paper by Graham and Zworski [18]. Roughly speaking, they prove that if $(M^n, [g])$ is a smooth compact manifold endowed with a conformal structure, then the GJMS operators P_k can be realized as residues at the values $\gamma = k$ of the meromorphic family $S(n/2 + \gamma)$ of scattering operators associated to the Laplacian on any Poincaré–Einstein manifold (X, G) for which $(M^n, [g])$ is the conformal infinity. These are the ‘trivial’ poles of the scattering operator $S(s)$; typically $S(s)$ has infinitely many other poles, which are called resonances. In this geometric framework, multiplying this scattering family by some regularizing factor, one obtains a holomorphic family of elliptic pseudo-differential operators denoted P_γ^g (which patently depends on the filling (X, G)) for $\gamma \in (0, n/2)$. This realization of the GJMS operators (and their continuous in γ counterpart) has led to important new understanding of them, including, for example, the basic fact that P_γ^g is symmetric with respect to dv_g . Hence even though the family P_γ^g is not entirely canonically associated to $(M, [g])$, its study can still illuminate the truly canonical operators which occur as special values, i.e. the GJMS operators.

Let us sum up the main properties of these operators: $P_0^g = \text{Id}$, and more generally, P_k^g is the k^{th} GJMS operator; P_γ^g is a classical elliptic pseudodifferential operator of order 2γ with principal symbol $\sigma_{2\gamma}(P_\gamma^g) = |\xi|_g^{2\gamma}$; P_γ^g is Fredholm on L^2 when $\gamma > 0$; if P_γ^g is invertible, then $P_{-\gamma}^g = (P_\gamma^g)^{-1}$; and, most importantly,

$$\hat{g} = u^{\frac{4}{n-2\gamma}} g \implies P_\gamma^g(uf) = u^{\frac{n+2\gamma}{n-2\gamma}} P_\gamma^{\hat{g}}(f) \quad (13)$$

for any smooth function f . Generalizing the formulæ for scalar curvature ($\gamma = 1$) and the Paneitz–Branson Q -curvature ($\gamma = 2$), for any $0 < \gamma < n/2$ we define

the Q -curvature of order γ associated to a metric g by

$$Q_\gamma^g = P_\gamma^g(1). \tag{14}$$

It is then natural to define a “generalized Yamabe problem”: given a metric g on a compact manifold M , find $u > 0$ so that if $\hat{g} = u^{4/(n-2\gamma)}g$, then $Q_\gamma^{\hat{g}}$ is constant. This amounts to solving

$$P_\gamma^g u = Q_\gamma^{\hat{g}} u^{\frac{n+2\gamma}{n-2\gamma}}, \quad u > 0, \tag{15}$$

for $Q_\gamma^{\hat{g}} = \text{const}$. From the analytic point of view, this problem exhibits the same features as the standard Yamabe problem (lack of compactness, for instance) with the additional major difficulty that the equation involves a pseudodifferential operator.

In the last decade or so, several works have studied this generalized Yamabe problem. The first major tool is a result of Chang and Gonzalez [7] realizing the operators P_γ^g for $\gamma \in (0, 1)$ as boundary operators, i.e. Dirichlet-to-Neumann type operators. It relies heavily on the original construction based on scattering of Graham and Zworski, reformulating it in a suitable form. However, a drawback of the result by Chang and Gonzalez is that the operator defined on the Poincaré-Einstein manifold (X, G) is not conformally covariant except in the case $\gamma = 1/2$, which corresponds precisely to the Escobar problem we considered in the previous sections. This was later resolved by Case and Chang [6], where they reinterpret the scattering theory of Graham and Zworski in terms of smooth metric measure spaces. We now introduce this framework.

Smooth metric measure spaces and conformally covariant operators of order 2γ with $\gamma \in (0, 1)$. We will refrain from going deeply in the theory of Case and Chang since it is quite technical and would obscure our goal of applying their construction to the spectral geometry of their operators. Notice first that the most popular range of powers is $\gamma \in (0, 1)$ since in this case the problem enjoys a standard maximum principle. However, as already noted, the case $\gamma = 2$ is also of major importance since it corresponds to the Paneitz–Branson operator. We will thus focus on the range $\gamma \in (0, 1)$, but several results mentioned in this section can be obtained for a wider range (under appropriate assumptions). We now describe their construction. Notice that we change slightly the notations below to be consistent with the previous discussion starting this section.

A smooth metric measure space is a four-tuple $(\bar{X}^{n+1}, G, v^m \text{dvol}, \mu)$ determined by a Riemannian manifold (\bar{X}, G) with boundary $M = \partial X$, the Riemannian volume element dvol associated to G , a nonnegative function $v \in C^\infty(\bar{X})$

with $v^{-1}(0) = M$ and constants $m \in \mathbb{R} \setminus \{1 - n\}$ and $\mu \in \mathbb{R}$. In the interior of \bar{X} we define a smooth function ϕ by

$$v^m = e^{-\phi}.$$

We say that $(\bar{X}^{n+1}, G, v^m \text{dvol}_G, \mu)$ and $(\bar{X}^{n+1}, \hat{G}, \hat{v}^m \text{dvol}_{\hat{G}}, \mu)$ are pointwise conformally equivalent if there exists a smooth function u such that

$$\hat{g} = u^{-2}g$$

and

$$\hat{v} = u^{-1}v.$$

We then define the weighted Laplacian (in relation to the Bakry-Emery Ricci tensor) and the weighted GJMS operator of order 2 by

$$\Delta_\phi = \Delta - \nabla\phi$$

and

$$L_{2,\phi}^m = -\Delta_\phi + \frac{m+n-1}{2}J_\phi^m,$$

where

$$J_\phi^m = \frac{1}{2(m+n)}(R - 2mv^{-1}\Delta v - m - (m-1)v^{-2}|\nabla v|^2 + m\mu v^{-2}).$$

The important property for us is conformality in the following sense: if $\hat{g} = u^{-2}g$ and $\hat{v} = u^{-1}v$, then

$$\widehat{L_{2,\phi}^m} w = u^{(m+n+3)/2} L_{2,\phi}^m (u^{-(m+n-1)/2} w)$$

for every smooth w . We can now state the main theorem (see [6], Theorem 4.1 and Lemma 6.1) in the case $\gamma \in (0, 1)$.

Theorem 7.1. *Let (X^{n+1}, M^n, G_+) be a Poincaré–Einstein manifold and $\gamma \in (0, 1)$. Suppose that $\lambda_1(-\Delta_{G_+}) > \frac{n^2}{4} - \gamma^2$ and set $m_0 = 1 - 2\gamma$. Fix a representative h of the conformal boundary and let r be the geodesic defining function associated to h . Then there exists a unique defining function ρ such that*

$$\rho = r + c_\gamma Q_\gamma r^{1+2\gamma} + O(r^3)$$

where c_γ is just a normalizing constant and

$$J_{\phi_0}^{m_0} = 0.$$

Consider now the smooth metric measure space

$$(\bar{X}^{n+1}, G := \rho^2 G_+, \rho^{m_0} \text{dvol}, m_0 - 1).$$

Given f smooth, the function U is the solution of

$$\begin{cases} L_{2,\phi_0}^{m_0} U = 0, & x \in X, \\ U = f, & x \in M, \end{cases} \tag{16}$$

if and only if $u = \rho^{n-s} U$ is the solution of

$$\begin{cases} -\Delta_{G_+} u - s(n-s)u = 0, & x \in X, \\ u = Fr^{n-s} + Gr^s, & \\ F|_{r=0} = f, & x \in M, \end{cases} \tag{17}$$

for $s = \frac{n}{2} + \gamma$. Moreover U is such that

$$P_\gamma f - \frac{n-2\gamma}{2} c_\gamma Q_\gamma f = \frac{c_\gamma}{2\gamma} \lim_{\rho \rightarrow 0} \rho^{m_0} \partial_\rho U.$$

Applying the previous theorem, the generalized Yamabe problem (15) for $\gamma \in (0, 1)$ is equivalent to solving the BVP

$$\begin{cases} -\Delta_{\phi_0} U = 0, & x \in X, \\ \lim_{\rho \rightarrow 0} \rho^{1-2\gamma} \partial_\rho U - \frac{n-2\gamma}{2} c_\gamma Q_\gamma U = d_\gamma U^{\frac{n+2\gamma}{n-2\gamma}}, & x \in M, \end{cases} \tag{18}$$

for some harmless constants c_γ and d_γ (with signs as defined in [6]). For $\gamma = 1/2$ this is exactly the Escobar problem [11]. Notice that by Hopf’s lemma (see [6]), one can prove that the problem (18) admits a strong maximum principle. The point of the BVP (18) is that, by integration by parts on the weighted smooth space and the conformality of the operators, one can deduce obstructions by nodal domains for curvature prescription problems in exactly the same way as in Section 5. Similarly, one can define a version of the conformal Robin BVP and obtain analogues of the whole set of previous results, adapting straightforwardly the same computations. In particular, having in mind the program for the conformal Laplacian on manifolds with boundary, one could study the analogue of the Kazdan–Warner trichotomy.

The previous construction is developed here in the case $\gamma \in (0, 1)$. Case and Chang used the setting of smooth metric measure spaces to give the analogue of the previous theorem for the whole family of operators P_γ^g . However, in that case the formulas are much more involved and it starts being cumbersome to report the results in this section.

8. Further questions

In this section, we formulate several open problems that seem interesting to the authors, continuing the list in [4, §7].

- (A) We showed in Section 3 that generically, 0 is not an eigenvalue of the conformal Laplacian on manifolds with boundary, for the natural boundary value problems considered in this paper. The following is a natural analogue of Conjecture C in [4, §7.3] for manifolds with boundary: among metrics where 0 is an eigenvalue of the conformal Laplacian, generically its multiplicity is equal to 1.
- (B) Relative conformal invariants: can the results in [34] be generalized from dimension 2 to higher dimensions?

It is known that on some closed manifolds ([19, 33]) the space of Yamabe-positive metrics can have infinitely many connected components. On the other hand, the space of Yamabe-negative metrics is connected and has trivial homotopy groups ([28, 23]). The number of negative eigenvalues of the conformal Laplacian for one of the conformally invariant boundary value problems considered in Section 2 provides a natural grading of the space of Yamabe-negative metrics; this leads naturally to Problem K raised in [4, §7]. Below, we formulate its natural counterpart for manifolds with boundary.

- (C) Let $[g_0]$ and $[g_1]$ be two conformal classes of metrics on $(M, \partial M)$ such that P_{g_0} and P_{g_1} have at least k negative eigenvalues for one of the conformally invariant boundary value problems considered in Section 2. Is it possible to connect g_0 and g_1 by a curve of metrics g_t such that P_{g_t} has at least k negative eigenvalues for the corresponding boundary value problems?

Acknowledgements. The authors would like to thank Asma Hassannezhad and A. Rod Gover for stimulating discussions and very interesting remarks about preliminary versions of this paper. In addition, the authors would like to thank Ailana Fraser, Pengfei Guan and Richard Schoen for useful discussions. The authors would like to thank BIRS, CRM, Oberwolfach, McGill and Johns Hopkins University for their hospitality.

References

- [1] W. Arendt and R. Mazzeo, Spectral properties of the Dirichlet-to-Neumann operator on Lipschitz domains. *Ulmer Seminaire* **12** (2007), 28–38.

- [2] T. Aubin, *Some nonlinear problems in Riemannian geometry*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1998 [MR 1636569](#) [Zbl 0896.53003](#)
- [3] Y. Canzani, R. Gover, D. Jakobson and R. Ponge, Nullspaces of conformally invariant operators. Applications to Q_k -curvature. *Electron. Res. Announc. Math. Sci.* **20** (2013), 43–50. [MR 3044498](#) [Zbl 1282.53027](#)
- [4] Y. Canzani, R. Gover, D. Jakobson and R. Ponge, Conformal invariants from nodal sets. I. Negative eigenvalues and curvature prescription. *Int. Math. Res. Not. IMRN* **2014**, no. 9, 2356–2400. [MR 3207371](#) [Zbl 1321.58022](#)
- [5] J. Case, Boundary operators associated with the Paneitz operator. *Indiana Univ. Math. J.* **67** (2018), no. 1, 293–327. [MR 3776023](#) [Zbl 1409.58017](#)
- [6] J. Case and S.-Y. Alice Chang, On fractional GJMS operators. *Comm. Pure Appl. Math.* **69** (2016), no. 6, 1017–1061. [MR 3493624](#) [Zbl 1348.35292](#)
- [7] S.-Y. A. Chang and M. del Mar Gonzalez, Fractional Laplacian in conformal geometry. *Adv. Math.* **226** (2011), no. 2, 1410–1432. [MR 2737789](#) [Zbl 1214.26005](#)
- [8] S. S. Chen, Conformal deformation to scalar flat metrics with constant mean curvature on the boundary in higher dimensions. Preprint, 2009. [arXiv:0912.1302](#) [math.DG]
- [9] S. El Sayed, Second eigenvalue of the Yamabe operator and applications. *Calc. Var. Partial Differential Equations* **50** (2014), no. 3–4, 665–692. [MR 3216829](#) [Zbl 1304.35467](#)
- [10] J. Escobar, The Yamabe problem on manifolds with boundary. *J. Differential Geom.* **35** (1992), no. 1, 21–84. [MR 1152225](#) [Zbl 0771.53017](#)
- [11] J. Escobar, Conformal deformation of a Riemannian metric to a scalar flat metric with constant mean curvature on the boundary. *Ann. of Math. (2)* **136** (1992), no. 1, 1–50. [MR 1173925](#) [Zbl 0766.53033](#)
- [12] J. Escobar, Conformal metrics with prescribed mean curvature on the boundary. *Calc. Var. Partial Differential Equations* **4** (1996), no. 6, 559–592. [MR 1416000](#) [Zbl 0867.53034](#)
- [13] J. Escobar, Uniqueness and non-uniqueness of metrics with prescribed scalar and mean curvature on compact manifolds with boundary. *J. Funct. Anal.* **202** (2003), no. 2, 424–442. [MR 1990532](#) [Zbl 1041.53026](#)
- [14] L. Friedlander, Some inequalities between Dirichlet and Neumann eigenvalues. *Arch. Rational Mech. Anal.* **116** (1991), no. 2, 153–160. [MR 1143438](#) [Zbl 0789.35124](#)
- [15] A. R. Gover, A. Hassannezhad, D. Jakobson, and M. Levitin, Zero and negative eigenvalues of the conformal Laplacian. *J. Spectr. Theory* **6** (2016), no. 4, 793–806. [MR 3584184](#) [Zbl 1362.58017](#)
- [16] A. R. Gover and L. J. Peterson. Conformal boundary operators, T -curvatures, and conformal fractional Laplacians of odd order. Preprint, 2018. [arXiv:1802.08366](#) [math.DG]

- [17] C. R. Graham, R. Jenne, L. J. Mason, and G. A. Sparling, Conformally invariant powers of the Laplacian. I. Existence. *J. London Math. Soc.* (2) **46** (1992), no. 3, 557–565. [MR 1190438](#) [Zbl 0726.53010](#)
- [18] C. Graham and M. Zworski, Scattering matrix in conformal geometry. *Invent. Math.* **152** (2003), no. 1, 89–118. [MR 1965361](#) [Zbl 1030.58022](#)
- [19] M. Gromov and H. B. Lawson, The classification of simply connected manifolds of positive scalar curvature. *Ann. of Math.* (2) **111** (1980), no. 3, 423–434. [MR 0577131](#) [Zbl 0463.53025](#)
- [20] Q. Han, Singular sets of solutions to elliptic equations. *Indiana Univ. Math. J.* **43** (1994), no. 3, 983–1002. [MR 1305956](#) [Zbl 0817.35020](#)
- [21] Q. Han, R. Hardt, and F. H. Lin, Geometric measure of singular sets of elliptic equations. *Comm. Pure Appl. Math.* **51** (1998), no. 11–12, 1425–1443. [MR 1639155](#) [Zbl 0940.35065](#)
- [22] R. Hardt, M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, and N. Nadirashvili, Critical sets of solutions to elliptic equations. *J. Differential Geom.* **51** (1999), no. 2, 359–373. [MR 1728303](#) [Zbl 1144.35370](#)
- [23] M. Katagiri, On the topology of the moduli space of negative constant scalar curvature metrics on a Haken manifold. *Proc. Japan Acad. Ser. A Math. Sci.* **75** (1999), no. 7, 126–128. [MR 1729860](#) [Zbl 0978.53077](#)
- [24] J. L. Kazdan, *Applications of partial differential equations to problems in geometry*. Lecture notes. Preprint, 1983–1993. <https://www2.math.upenn.edu/~kazdan/japan/japan.pdf>
- [25] J. L. Kazdan and F. W. Warner, Curvature functions for compact 2-manifolds. *Ann. of Math.* (2) **99** (1974), 14–47. [MR 0343205](#) [Zbl 0273.53034](#)
- [26] J. L. Kazdan and F. W. Warner, Scalar curvature and conformal deformation of Riemannian structure. *J. Differential Geometry* **10** (1975), 113–134. [MR 0365409](#) [Zbl 0296.53037](#)
- [27] M. Lassas, T. Liimatainen, and M. Salo, The Calderón problem for the conformal Laplacian. Preprint, 2016. [arXiv:1612.07939](#) [math.AP]
- [28] J. Lohkamp, The space of negative scalar curvature metrics. *Invent. Math.* **110** (1992), no. 2, 403–407. [MR 1185590](#) [Zbl 0771.58006](#)
- [29] F. Marques, Conformal deformations to scalar-flat metrics with constant mean curvature on the boundary. *Comm. Anal. Geom.* **15** (2007), no. 2, 381–405. [MR 2344328](#) [Zbl 1132.53021](#)
- [30] M. Mayer and C.-B. Ndiaye, Barycenter technique and the Riemann mapping problem of Escobar. *J. Differential Geom.* **107** (2017), no. 3, 519–560. [MR 3715348](#) [Zbl 1410.53043](#)
- [31] R. Mazzeo, Remarks on a paper of Friedlander concerning inequalities between Neumann and Dirichlet eigenvalues. *Internat. Math. Res. Notices* **1991**, no. 4, 41–48. [MR 1121164](#) [Zbl 0752.58035](#)

- [32] T. Parker and S. Rosenberg, Invariants of conformal Laplacians. *J. Differential Geom.* **25** (1987), no. 2, 199–222. [MR 0880183](#) [Zbl 0644.53038](#)
- [33] J. Rosenberg, Manifolds of positive scalar curvature: a progress report. In J. Cheeger and K. Grove (eds.), *Surveys in differential geometry*. Vol. XI. International Press, Somerville, MA, 2007, 259–294. [MR 2408269](#) [Zbl 1171.53028](#)
- [34] E. Schippers, Quadratic differentials and conformal invariants. *J. Anal.* **24** (2016), no. 2, 209–228. [MR 3755820](#) [Zbl 1362.30035](#)

Received May 13, 2019

Graham Cox, Department of Mathematics and Statistics,
Memorial University of Newfoundland, St. John's, NL A1C 5S7, Canada

e-mail: gcox@mun.ca

Dmitry Jakobson, Department of Mathematics and Statistics, 805 Sherbrooke Street West,
Montreal, Quebec, Canada H3A0B9

e-mail: dmitry.jakobson@mcgill.ca

Mikhail Karpukhin, Mathematics 253-37, Caltech, Pasadena, CA 91125, USA

e-mail: mikhailk@caltech.edu

Yannick Sire, Department of Mathematics, Johns Hopkins University,
404 Krieger Hall, 3400 N. Charles Street, Baltimore, MD 21218, USA

e-mail: ysire1@jhu.edu