

# Existence of metrics maximizing the first eigenvalue on non-orientable surfaces

Henrik Matthiesen and Anna Siffert

**Abstract.** We prove the existence of metrics maximizing the first eigenvalue normalized by area on closed, non-orientable surfaces assuming two spectral gap conditions. These spectral gap conditions are proved by the authors in [21].

**Mathematics Subject Classification (2020).** 35P15, 49Q05, 49Q10, 58E11, 58E20.

**Keywords.** Laplace operator, topological spectrum, harmonic map, minimal surface, shape optimization.

## 1. Introduction

For a closed Riemannian surface  $(\Sigma, g)$  the spectrum of the Laplace operator acting on smooth functions, is purely discrete and can be written as

$$0 = \lambda_0 < \lambda_1(\Sigma, g) \leq \lambda_2(\Sigma, g) \leq \lambda_3(\Sigma, g) \leq \dots \longrightarrow \infty,$$

where we repeat an eigenvalue as often as its multiplicity requires. Via the variational characterization of eigenvalues this extends to a much larger class of metrics with lower regularity properties. For details we refer to [18], which in particular covers the metrics considered here.

The pioneering work of Hersch [13] and Yang and Yau [36] raised the natural question, whether there are metrics  $g$  that maximize the scale-invariant quantities

$$\bar{\lambda}_1(\Sigma) := \lambda_1(\Sigma, g) \operatorname{area}(\Sigma, g)$$

if  $\Sigma$  is a closed surface of fixed topological type (see also [15, 19] for the case of non-orientable surfaces). Such maximizers have remarkable properties. In fact, they always arise as immersed minimal surfaces (of possibly high codimension) in a sphere [9] and are unique in their conformal class unless they are branched

immersions into the two sphere [6, 23, 31]. By a slight abuse of notation, we also call  $\Sigma$ , endowed with a maximizing metric, a ‘maximizer’.

For the statement of our results and related work, we need to introduce some notation. We write  $\Sigma_\gamma$  for a closed orientable surface of genus  $\gamma$ . Similarly,  $\Sigma_\delta^K$  denotes a closed non-orientable surface of non-orientable genus  $\delta$ . Here,  $K$  stands for Klein. We briefly elaborate on these notions in Appendix A. Furthermore, we use the common notation

$$\Lambda_1(\gamma) = \sup_g \lambda_1(\Sigma_\gamma, g) \text{ area}(\Sigma_\gamma, g),$$

and similarly,

$$\Lambda_1^K(\delta) = \sup_g \lambda_1(\Sigma_\delta^K, g) \text{ area}(\Sigma_\delta^K, g),$$

with the supremum taken over all smooth metrics on  $\Sigma_\gamma$ , respectively  $\Sigma_\delta^K$ .

Explicit values for  $\Lambda_1(\gamma)$  or  $\Lambda_1^K(\delta)$  are only known in very few cases. However, in all of these cases not only the values but also explicit maximizing metrics are known.

The case of the sphere is due to Hersch. We have  $\Lambda_1(\mathbb{S}^2) = 8\pi$  with unique maximizer the round metric [13]. His arguments are very elegant and a cornerstone in the development of the subject. For the real projective plane, we have  $\Lambda_1(\mathbb{RP}^2) = 12\pi$  with unique maximizer the round metric [19]. The proof extends the ideas from [13] in a conceptually very nice way.

The first result for higher genus surfaces is due to Nadirashvili, namely  $\Lambda_1(T^2) = 8\pi^2/\sqrt{3}$  with unique maximizer the flat equilateral torus [25]. Nadirashvili’s arguments are very different from the previously employed methods. The crucial step in his proof is to obtain the existence of a maximizer. Using [23] (see also [6]) it follows that such a maximizer necessarily has to be flat. The sharp bound follows then from earlier work of Berger [2].

For the Klein bottle,  $\Lambda_1(K) = 12\pi E(2\sqrt{2}/3)$  with unique maximizer a metric of revolution [6, 8, 25]. Here  $E$  is the complete elliptic integral of the second kind.

There is also a conjecture concerning the sharp bound on genus 2 surfaces [14], a proof of which has very recently been given by Nayatani and Shoda in [31].

Let us also mention that there are quite some results concerning similar questions for higher order eigenvalues, see e.g. [29, 33] and [26, 30, 17] for the case of  $\mathbb{S}^2$  and [27, 16] for the case of  $\mathbb{RP}^2$ .

The growing interest in finding maximizers for eigenvalue functionals on surfaces starting from Nadirashvili’s paper [25] is certainly connected to the

connection of the problem to minimal surfaces in spheres. Similarly, for the Steklov eigenvalue problem, there is a connection to free boundary minimal surfaces in Euclidean balls. In pioneering work Fraser and Schoen showed the existence of maximizers for the first Steklov eigenvalue on surfaces with boundary of genus 0 [11].

Recently, Petrides used many of the ideas in [11] to prove the following beautiful result concerning metrics realizing  $\Lambda_1(\gamma)$ .

**Theorem 1.1** (Theorem 2 in [32]). *If  $\Lambda_1(\gamma - 1) < \Lambda_1(\gamma)$ , there is a metric  $g$  on  $\Sigma = \Sigma_\gamma$ , which is smooth away from finitely many conical singularities, such that*

$$\lambda_1(\Sigma, g) \text{ area}(\Sigma, g) = \Lambda_1(\gamma).$$

We extend this to non-orientable surfaces. Since non-orientable surfaces can degenerate to non-orientable surfaces as well as orientable ones, we need to make two instead of only a single spectral assumption.

**Theorem 1.2.** *If  $\Lambda_1^K(\delta - 1) < \Lambda_1^K(\delta)$  and  $\Lambda_1(\lfloor(\delta - 1)/2\rfloor) < \Lambda_1^K(\delta)$ , there is a metric  $g$  on  $\Sigma = \Sigma_\delta^K$ , which is smooth away from at most finitely many conical singularities, such that*

$$\lambda_1(\Sigma, g) \text{ area}(\Sigma, g) = \Lambda_1^K(\delta).$$

Our methods are very similar to those in [32]. In addition to the cases already handled by Petrides, we also need to take care of degenerating one-sided geodesics.

The non-strict inequality  $\Lambda_1(\gamma - 1) \leq \Lambda_1(\gamma)$  was proved by Colbois and El Soufi in [7] using a result of Anné [1]. It is easy to obtain the non-strict versions of the inequalities assumed in Theorem 1.2 along the same lines, see also [20]. In [20] we prove the monotonicity  $\Lambda_1(\gamma - 1) < \Lambda_1(\gamma)$  under some extra assumptions on the maximizing metric using a relatively simple gluing construction. In [21], by means of a much more complicated gluing construction and using Theorem 1.1 and Theorem 1.2, we prove all of the spectral gap conditions assumed in Theorem 1.1 and Theorem 1.2 above. In particular, this implies the existence of maximizing metrics on closed surfaces of any topological type.

**Theorem 1.3** (Theorem 1.3 in [21] using Theorem 1.1 and Theorem 1.2). *Let  $\Sigma$  be a closed surface. Then there is a metric  $g$  on  $\Sigma$ , which is smooth away from at most finitely many conical singularities, such that*

$$\lambda_1(\Sigma, h) \text{ area}(\Sigma, h) \leq \lambda_1(\Sigma, g) \text{ area}(\Sigma, g)$$

for any smooth metric  $h$  on  $\Sigma$ .

**Outline.** In Section 2 we prove that the set of orientable, hyperbolic surfaces with injectivity radius bounded below is a compact subset of the moduli space. This is a version of the Mumford compactness criterion for non-orientable surfaces. We use this result in the main section, namely Section 3, in which we prove Theorem 1.2.

**Acknowledgements.** The authors would like to thank the Max Planck Institute for Mathematics in Bonn for financial support and excellent working conditions.

## 2. Compactness for non-orientable surfaces

The Mumford compactness criterion [24] states that the set of orientable, hyperbolic surfaces with injectivity radius bounded below is a compact subset of the moduli space. In this section we show that the Mumford argument for orientable surfaces holds for the non-orientable case as well.

Given any Riemannian metric  $g_0$  on  $\Sigma = \Sigma_\delta^K$ , the Poincaré uniformization theorem asserts that we can find a new metric on  $\Sigma$  which is conformal to  $g_0$  and has constant curvature  $+1, 0$ , or  $-1$ , depending on the sign of  $\chi(\Sigma)$ . Assuming  $\delta \geq 3$ , these metrics have curvature  $-1$ . Let  $h_k$  be a sequence of such metrics on  $\Sigma$  with injectivity radius bounded uniformly from below,  $\text{inj}(\Sigma, h_k) \geq c > 0$ . The goal is to prove that there exist diffeomorphisms  $\sigma_k$  of  $\Sigma$  and a hyperbolic metric  $h$  of  $\Sigma$ , such that  $\sigma_k^* h_k$  converges smoothly to  $h$  as  $k \rightarrow \infty$ . Our strategy is to apply the Mumford compactness criterion to the orientation double covers of the surfaces  $(\Sigma, h_k)$ .

Consider the orientation double cover  $\widehat{\Sigma} = \Sigma_{\delta-1}$  of  $\Sigma$  endowed with the pullback metrics of  $h_k$ , denoted by  $\widehat{h}_k$ . Since  $\delta \geq 3$ , these are orientable hyperbolic surface of genus  $\delta - 1$  and may thus be regarded (if we also fix a marking) as elements in Teichmüller space  $\mathcal{T}_{\delta-1}$ , which in addition admit fixed point free, isometric, orientation reversing involutions  $\iota_k$ .

We have the following lemma.

**Lemma 2.1.** *Assume that  $\inf_k \text{inj}(\Sigma_\delta^K, h_k) > 0$ . Then there exists a sequence of diffeomorphisms  $\tau_k: \Sigma_{\delta-1} \rightarrow \Sigma_{\delta-1}$ , such that, up to taking a subsequence,  $\tau_k^* \widehat{h}_k \rightarrow \widehat{h}$  in  $C^\infty$ . Moreover,  $(\Sigma_{\delta-1}, \widehat{h})$  admits a fixed point free, isometric, orientation reversing involution  $\iota$ , which is obtained as a  $C^1$ -limit of the involutions  $\tau_k^{-1} \circ \iota_k \circ \tau_k$ .*

*Proof.* As above, we simply write  $\Sigma$  instead of  $\Sigma_\delta^K$ , and  $\widehat{\Sigma}$  instead of  $\Sigma_{\delta-1}$ . It is elementary to see that  $\text{inj}(\widehat{\Sigma}, \widehat{h}_k) \geq \text{inj}(\Sigma, h_k)$ . Therefore, we can apply the Mumford compactness criterion [24] and find diffeomorphisms  $\tau_k$  and a limit metric  $\widehat{h}$  as asserted.

It remains to show that we can find the involution  $\iota$ . Since  $\tau_k^* \widehat{h}_k \rightarrow \widehat{h}$  in  $C^\infty$ , we have the uniform Lipschitz bound

$$\begin{aligned} d_{\widehat{h}}((\tau_k^{-1} \circ \iota_k \circ \tau_k)(p), (\tau_k^{-1} \circ \iota_k \circ \tau_k)(q)) &\leq C d_{\tau_k^* \widehat{h}_k}((\tau_k^{-1} \circ \iota_k \circ \tau_k)(p), (\tau_k^{-1} \circ \iota_k \circ \tau_k)(q)) \\ &= C d_{\tau_k^* \widehat{h}_k}(p, q) \\ &\leq C d_{\widehat{h}}(p, q). \end{aligned}$$

Similarly, we obtain a uniform bound on  $\|D\iota\|_{C^{0,1}(\widehat{\Sigma}, \widehat{h})}$ . Since  $\widehat{\Sigma}$  is compact, it follows from Arzelà–Ascoli, that, up to taking a subsequence,  $\tau_k^{-1} \circ \iota_k \circ \tau_k \rightarrow \iota$  in  $C^1(\widehat{\Sigma}, \widehat{h})$ . We have

$$\begin{aligned} d_{\widehat{h}}(\iota(p), \iota(q)) &\leq \lim_{k \rightarrow \infty} d_{\tau_k^* \widehat{h}_k}(\iota(p), (\tau_k^{-1} \circ \iota_k \circ \tau_k)(p)) \\ &\quad + \lim_{k \rightarrow \infty} d_{\tau_k^* \widehat{h}_k}((\tau_k^{-1} \circ \iota_k \circ \tau_k)(p), (\tau_k^{-1} \circ \iota_k \circ \tau_k)(q)) \\ &\quad + \lim_{k \rightarrow \infty} d_{\tau_k^* \widehat{h}_k}((\tau_k^{-1} \circ \iota_k \circ \tau_k)(q), \iota(q)) \\ &\leq C \lim_{k \rightarrow \infty} d_{C^0(\widehat{\Sigma}, \widehat{h})}(\tau_k^{-1} \circ \iota_k \circ \tau_k, \iota) \\ &\quad + \lim_{k \rightarrow \infty} d_{\tau_k^* \widehat{h}_k}((\tau_k^{-1} \circ \iota_k \circ \tau_k)(p), (\tau_k^{-1} \circ \iota_k \circ \tau_k)(q)) \\ &= d_{\widehat{h}}(p, q), \end{aligned} \tag{2.2}$$

using that  $\tau_k^* \widehat{h}_k \rightarrow \widehat{h}$  in  $C^\infty$ , and  $\tau_k^{-1} \circ \iota_k \circ \tau_k \rightarrow \iota$  in  $C^0(\widehat{\Sigma}, \widehat{h})$ . Observe that  $\iota$  is an involution again, hence (2.2) implies that actually

$$d_{\widehat{h}}(\iota(p), \iota(q)) = d_{\widehat{h}}(p, q).$$

By the Myers–Steenrod theorem it thus follows that  $\iota$  is a smooth, isometric involution.

We need to show that  $\iota$  does not have any fixed points. But this is a consequence of the general bound  $d_{\tau_k^* \widehat{h}_k}((\tau_k^{-1} \circ \iota_k \circ \tau_k)(p), p) \geq c > 0$  for some uniform  $c$ . To prove this let  $c > 0$  be such that  $B_{\widehat{h}}(x, 2c) \subset \widehat{\Sigma}$  is strictly geodesically convex for any  $x \in \widehat{\Sigma}$ . Then  $B_{\tau_k^* \widehat{h}_k}(x, c)$  is strictly geodesically convex for  $k \geq K$  sufficiently large. Assume now that there is  $k \geq K$ , such that  $d_{\widehat{h}_k}((\tau_k^{-1} \circ \iota_k \circ \tau_k)(p), p) < c$ .

Let  $\gamma$  be the unique minimizing geodesic connecting  $p$  to  $(\tau_k^{-1} \circ \iota_k \circ \tau_k)(p)$ . Since  $\tau_k^{-1} \circ \iota_k \circ \tau_k$  is an isometry, we need to have  $\text{im}(\tau_k^{-1} \circ \iota_k \circ \tau_k \circ \gamma) = \text{im } \gamma$ . Since  $\iota_k$  is fixed point free,  $\gamma$  is non-constant. Therefore,  $\tau_k^{-1} \circ \iota_k \circ \tau_k$  restricted to  $\text{im } \gamma$  induces an involution of the interval  $[0, 1]$  mapping 0 to 1 and vice versa. But such an involution needs to have a fixed point. It follows that  $\iota_k$  has a fixed point for large  $k$ , which is a contradiction.

Finally, note that  $\iota$  is orientation reversing by  $C^0$ -convergence. □

Since  $\tau_k^{-1} \circ \iota_k \circ \tau_k \rightarrow \iota$  in  $C^1$  it follows that the metric  $\hat{h}$  on  $\hat{\Sigma}$  is  $\iota$ -invariant. Therefore, it induces a smooth hyperbolic metric  $h$  on  $\Sigma$ . Moreover, the hyperbolic metrics on  $\Sigma$  induced from  $\tau_k^* h_k$  and  $\tau_k^{-1} \circ \iota_k \circ \tau_k$  converge smoothly to  $h$  on  $\Sigma$ . Finally, observe that the diffeomorphisms  $\tau_k$  induce diffeomorphisms  $\sigma_k$  of  $\Sigma$ , such that  $\sigma_k^* h_k$  are the metrics described above and converge smoothly to  $h$ .

Thus we have proved the following proposition.

**Proposition 2.3.** *Let  $(h_k)$  be a sequence of hyperbolic metrics on  $\Sigma_\delta^K$  such that their injectivity radius is uniformly bounded from below  $\text{inj}(\Sigma_\delta^K, h_k) \geq c > 0$ . Then there are diffeomorphisms  $\sigma_k$  of  $\Sigma_\delta^K$  and a hyperbolic metric  $h$ , such that  $\sigma_k^* h_k \rightarrow h$  smoothly.*

### 3. Maximizing the first eigenvalue

In this section we extend [32, Theorem 2] to the non-orientable case. The strategy is the same as in [32]. That is, we first use that we can maximize the first eigenvalue in each conformal class. We then pick a maximizing sequence, consisting of maximizers in their own conformal class. This has the advantage, that these metrics can be studied in terms of sphere-valued harmonic maps. Using these harmonic maps it is possible to estimate the first eigenvalue along the maximizing sequence in case that the conformal class degenerates. To do so, we extend the results from [37] to non-orientable surfaces.

For fixed non-orientable genus  $\delta \geq 3$ , let  $c_k$  be a sequence of conformal classes on  $\Sigma = \Sigma_\delta^K$  represented by hyperbolic metrics  $h_k$ , such that

$$\lim_{k \rightarrow \infty} \sup_{g \in c_k} \lambda_1(\Sigma, g) \text{ area}(\Sigma, g) = \Lambda_1^K(\delta).$$

We will now use the following result due to Nadirashvili–Sire (with an extra assumption not relevant for our purposes) and, independently, Petrides.

**Theorem 3.1** ([28, Theorem 2.1] or [32, Theorem 1]). *For each conformal class  $c_k$  as above, there is a metric  $g_k$ , which is smooth away from finitely many conical singularities, such that*

$$\lambda_1(\Sigma, g_k) \operatorname{area}(\Sigma, g_k) = \sup_{g \in c_k} \lambda_1(\Sigma, g) \operatorname{area}(\Sigma, g).$$

From now on we assume that  $g_k \in c_k$  is picked as in the preceding theorem. Moreover, we assume that they are normalized to have

$$\operatorname{area}(\Sigma, g_k) = 1.$$

Since these metrics are maximizers, there is a family of first eigenfunctions  $u_1^k, \dots, u_{\ell(k)+1}^k$ , such that  $\Phi_k = (u_1^k, \dots, u_{\ell(k)+1}^k): (\Sigma, h_k) \rightarrow \mathbb{S}^{\ell(k)}$  is a harmonic map [10]. Since the multiplicity of  $\lambda_1$  is uniformly bounded in terms of the topology of  $\Sigma$  [3, 5], we may pass to a subsequence, such that  $\ell(k)$  is some constant number  $l$ . Moreover, in this situation the maximizing metrics can be recovered by

$$g_k = \frac{|\nabla \Phi_k|_{h_k}^2}{\lambda_1(\Sigma, g_k)} h_k.$$

In view of Proposition 2.3, we want to show the following proposition.

**Proposition 3.2.** *The injectivity radius of  $h_k$  is uniformly bounded from below, provided that  $\Lambda_1^K(\delta) > \Lambda_1(\delta - 1)$ , and  $\Lambda_1^K(\delta) > \Lambda_1^K(\delta - 1)$ .*

We will argue by contradiction and assume  $\operatorname{inj}(\Sigma, h_k) \rightarrow 0$ . The Margulis lemma implies that we can find closed geodesics  $\gamma_k^1, \dots, \gamma_k^s$  in  $(\Sigma, h_k)$ , such that their lengths go to zero, i.e.  $l_{h_k}(\gamma_i^k) \rightarrow 0$ , as  $k \rightarrow \infty$ . We assume that  $s$  is chosen maximal with this property.

Each of these geodesics is either one-sided or two-sided. If a such a geodesic is two-sided, tubular neighborhoods are described by the classical collar lemma for hyperbolic surfaces [4]. In the second case we may apply the collar lemma to the orientation double cover as follows.

Let  $c$  be a one-sided closed geodesic in  $\Sigma$ . We write  $\widehat{\Sigma}$  for the orientation double cover and  $\tau$  for the non-trivial deck transformation. The lifts of  $c$  to  $\widehat{\Sigma}$  can not be closed, since in this case they would be disjoint and it would follow that  $c$  is two-sided. Thus the lifts  $c_1$  and  $c_2$  are geodesic segments with  $\tau \circ c_1 = c_2$ . Let  $\mathcal{C}$  be a collar around the closed geodesic  $c_2 * c_1$ . It is not very difficult to see that the action of  $\tau$  near  $c_2 * c_1$  is just given by rotation about  $\pi$  and reflection at  $c_2 * c_1$ . Therefore,  $\tau$  maps  $\mathcal{C}$  to itself (by the explicit construction of  $\mathcal{C}$ ), so that we can use  $\mathcal{C}/\tau$  as a tubular neighborhood of  $c$ .

Our first goal is to prove that for the situation at hand the volume, measured with respect to  $g_k$ , either concentrates in the neighborhood of a pinching geodesic, or in one connected component of the complement of these neighborhoods. Before stating and proving this result we need to introduce some notation, which we borrow from Section 4 in [32].

We write  $s_1$  for the number of one-sided closed geodesics with length going to 0. Moreover, we denote by  $s_2$  the number of such geodesics that are two-sided. Clearly,  $s = s_1 + s_2$  and  $0 \leq s_1, s_2 \leq s$ . From now on we assume that the closed geodesics  $\gamma_k^i$  are ordered such that the first  $s_1$  geodesics are one-sided. Moreover, we write  $l_k^i = l_{h_k}(\gamma_k^i)$  for the hyperbolic length of the short geodesics.

For all  $s_1 + 1 \leq i \leq s$  the collar theorem [4] asserts the existence of an open neighborhood  $P_k^i$  of  $\gamma_k^i$  isometric to the following truncated hyperbolic cylinder

$$\mathcal{C}_k^i = \{(t, \theta) \mid -w_k^i < t < w_k^i, 0 \leq \theta < 2\pi\}$$

with

$$w_k^i = \frac{\pi}{l_k^i} \left( \pi - 2 \arctan \left( \sinh \frac{l_k^i}{2} \right) \right)$$

endowed with the metric

$$\left( \frac{l_k^i}{2\pi \cos \left( \frac{l_k^i}{2\pi} t \right)} \right)^2 (dt^2 + d\theta^2).$$

Below we identify  $(\theta, t) = (0, t)$  with  $(\theta, t) = (2\pi, t)$ . Thus the closed geodesic  $\gamma_k^i$  corresponds to  $\{t = 0\}$ .

By the discussion above and the collar theorem again, we get that for all  $1 \leq i \leq s_1$ , there exists an open neighborhood  $P_k^i$  of  $\gamma_k^i$  isometric to the following truncated Möbius strip

$$\mathcal{M}_k^i = \{(t, \theta) \mid -w_k^i < t < w_k^i, 0 \leq \theta < 2\pi\} / \sim$$

with

$$w_k^i = \frac{\pi}{2l_k^i} (\pi - 2 \arctan(\sinh l_k^i))$$

endowed with the metric

$$\left( \frac{2l_k^i}{2\pi \cos \left( \frac{2l_k^i}{2\pi} t \right)} \right)^2 (dt^2 + d\theta^2).$$

Moreover, the equivalence relation  $\sim$  is given by identifying  $(t, \theta, ) \sim (-t, \theta + \pi)$ , where  $\theta + \pi \in \mathbb{R}/2\pi\mathbb{R}$ . Hence, the closed geodesic  $\gamma_k^i$  corresponds to  $\{t = 0\}$ .

We denote by  $\Sigma_k^1, \dots, \Sigma_k^r$  the connected components of  $\Sigma \setminus \bigcup_{i=1}^s P_k^i$ . Consequently,  $\Sigma$  can be written as the disjoint union

$$\Sigma = \left( \bigcup_{i=1}^s P_k^i \right) \cup \left( \bigcup_{j=1}^r \Sigma_k^j \right).$$

For  $s_1 + 1 \leq i \leq s$  and  $0 < b < w_k^i$  we denote by  $P_k^i(b)$  the truncated hyperbolic cylinder whose length, compared to  $P_k^i$ , is reduced by  $b$ , i.e.,

$$P_k^i(b) = \{(t, \theta), -w_k^i + b < t < w_k^i - b\}.$$

Analogously, for  $1 \leq i \leq s_1$  and  $0 < b < w_k^i$ , we introduce

$$P_k^i(b) = \{(t, \theta), -w_k^i + b < t < w_k^i - b\} / \sim.$$

Finally, we denote by  $\Sigma_k^j(b)$  the connected components of  $\Sigma \setminus \bigcup_{i=1}^s P_k^i(b)$  which contains  $\Sigma_k^j$ .

We are now ready to prove the above mentioned result, namely, that the volume either concentrates in the neighborhood of a pinching geodesic  $P_k^i$ , or in one connected component  $\Sigma_k^j$  of the complement of these neighborhoods.

**Lemma 3.3.** *There exists  $D > 0$  such that one of the two following assertions is true:*

(1) *there exists an  $i \in \{1, \dots, s\}$  such that*

$$\text{area}_{g_k}(P_k^i(a_k)) \geq 1 - \frac{D}{a_k}$$

*for all sequences  $a_k \rightarrow +\infty$  with  $\frac{a_k}{w_k^i} \rightarrow 0$  as  $k \rightarrow +\infty$  for all  $1 \leq i \leq s$ ;*

(2) *there exists a  $j \in \{1, \dots, r\}$  such that*

$$\text{area}_{g_k}(\Sigma_k^j(9a_k)) \geq 1 - \frac{D}{a_k}$$

*for all sequences  $a_k \rightarrow +\infty$  with  $\frac{a_k}{w_k^i} \rightarrow 0$  as  $k \rightarrow +\infty$  for all  $1 \leq i \leq s$ .*

*Proof.* The proof of Claim 11 in [32] can easily be adapted to the present situation. First recall the rough strategy of the proof: construct suitable test functions for  $\lambda_1(\Sigma, g_k)$  in the  $P_k^i$  and the  $\Sigma_k^j$ 's, apply the min-max formula for the first eigenvalue and prove the claim by contradiction. More precisely, on  $\hat{\Sigma}$ , the test functions are constructed with linear decay in the  $t$  variable in neck regions of

the type  $\widehat{P}_k^i(2a_k) \setminus \widehat{P}_k^i(3a_k)$  and  $\widehat{P}_k^i(1a_k) \setminus \widehat{P}_k^i(2a_k)$ , respectively, where the hat indicates that we consider the preimages under the covering map  $\widehat{\Sigma} \rightarrow \Sigma$ . By conformal invariance, the Dirichlet energy of these can be estimated using the hyperbolic metric and decays like  $a_k^{-1}$ . From the construction it is clear that these functions are invariant under the relevant involutions. From this point on, one can just follow the arguments in [32].  $\square$

Below we consider the two possible cases of the preceding lemma separately. The following lemma deals with the first case, i.e. when the volume concentrates in one of the  $P_k^i$ . We show that in this case we would have  $\Lambda_1^K(\delta) \leq 8\pi$  if  $\gamma_k^i$  is 2-sided; and  $\Lambda_1^K(\delta) \leq 12\pi$  if  $\gamma_k^i$  is 1-sided.

**Lemma 3.4.** *Suppose that there exists an  $i \in \{1, \dots, s\}$  such that*

$$\text{area}_{g_k}(P_k^i(a_k)) \geq 1 - \frac{D}{a_k}$$

for all sequences  $a_k \rightarrow +\infty$  with  $\frac{a_k}{w_k^i} \rightarrow 0$  as  $k \rightarrow +\infty$  for all  $1 \leq i \leq s$ .

- (1) *If  $\gamma_k^i$  is 2-sided, then  $\Lambda_1^K(\delta) \leq 8\pi$ .*
- (2) *If  $\gamma_k^i$  is 1-sided, then  $\Lambda_1^K(\delta) \leq 12\pi$ .*

*Proof.* In [32], Petrides proved the first statement by following ideas of Girouard (see [12]). The proof of the second statement is carried out analogously.

By assumption, there exists an  $i \in \{1, \dots, s\}$ , such that the volume concentrates on  $P_k := P_k^i$ . On  $P_k$  we have coordinates  $(t, \theta)$  as above (on  $\mathcal{M}_k$ ). By the assumptions on the volume and  $a_k$ , we can find cut-off functions  $\eta_k$  which are 1 on  $P_k(a_k)$  and 0 outside  $P_k$ , and satisfy

$$\int_{\Sigma} |\nabla \eta_k|^2 dv_{g_k} \rightarrow 0. \tag{3.5}$$

We denote by  $\mathcal{C} = (-\infty, \infty) \times \mathbb{S}^1$  the infinite cylinder with its canonical coordinates  $(t, \theta) \in (-\infty, \infty) \times [0, 2\pi)$ . Let  $\phi: \mathcal{C} \rightarrow \mathbb{S}^2 \subset \mathbb{R}^3$  be given by

$$\phi(t, \theta) = \frac{1}{e^{2t} + 1} (2e^t \cos(\theta), 2e^t \sin(\theta), e^{2t} - 1).$$

Observe that this induces a map  $\psi: \mathcal{M} \rightarrow \mathbb{R}P^2(\sqrt{3})$  if we divide by the  $\mathbb{Z}/2$ -actions that we have on both sides. More precisely,  $\mathcal{M} = \mathcal{C} / \sim$ , where  $(t, \theta) \sim (-t, \theta + \pi)$  as above, and on  $\mathbb{S}^2$  we simply take the antipodal map. If we denote by  $v: \mathbb{R}P^2(\sqrt{3}) \rightarrow \mathbb{S}^4$  the Veronese map, the concatenation  $v \circ \phi: \mathcal{M} \rightarrow \mathbb{S}^4$

is a conformal map [12]. We may regard  $\mathcal{M}_k \subset \mathcal{M}$  using Fermi coordinates as introduced above.

By a theorem of Hersch [13], there exists a conformal diffeomorphism  $\tau_k$  of  $S^4$ , such that

$$\int_{P_k} (\pi \circ \tau_k \circ v \circ \phi) \eta_k dv_{g_k} = 0,$$

where  $\pi: S^4 \hookrightarrow \mathbb{R}^5$  is the standard embedding. Set  $u_k^i = (\pi_i \circ \tau_k \circ v \circ \phi) \eta_k$ . By construction, we have

$$\sum_{i=1}^5 \int_{\mathcal{M}_k} (u_k^i)^2 dv_{g_k} \geq 1 - \frac{D}{a_k},$$

since  $\text{area}_{g_k}(P_\alpha^i(a_k)) \geq 1 - \frac{D}{a_k}$ . Using conformal invariance and (3.5), one easily finds that

$$\int_{\Sigma} |\nabla u_k|_{g_k}^2 dv_{g_k} \leq 12\pi + o(1).$$

For details we refer to [12]. Consequently, there is  $i = i(k) \in \{1, \dots, 5\}$ , such that

$$\lambda_1(\Sigma, g_k) \leq \frac{\int_{\mathcal{M}} |\nabla u_k^i|_{g_k}^2 dv_{g_k}}{\int_{\mathcal{M}} (u_k^i)^2 dv_{g_k}} \leq 12\pi + o(1).$$

This finally implies

$$\Lambda_1^K(\delta) \leq \limsup_{k \rightarrow \infty} \lambda_1(\Sigma, g_k) \leq 12\pi,$$

which establishes the claim. □

We are thus left with the case second case from Lemma 3.3. In this case, we have the following lemma, which concludes the proof of Proposition 3.2.

**Lemma 3.6.** *Suppose that the second alternative from 3.3 holds, then either*

(i)  $\Lambda_1^K(\delta) \leq \Lambda_1^K(\delta - 1)$ , or

(ii)  $\Lambda_1^K(\delta) \leq \Lambda_1(\gamma)$ ,

where  $\gamma = \lfloor (\delta - 1)/2 \rfloor$ .

*Proof.* Again, we apply the machinery from [32] to the orientation cover. The essential point is to keep track of the geometry of the corresponding involutions. Denote by  $(\widehat{\Sigma}, \widehat{h}_k)$  the orientation covers of  $(\Sigma, h_k)$ , and by  $\iota_k$  the corresponding deck transformations.

We can then identify the spectrum of the Laplacian for any metric  $g$  in  $[h_k]$  with the spectrum of the Laplacian acting only on the even functions on  $(\widehat{\Sigma}, \widehat{g})$ . We consider the associated harmonic maps  $\Phi_k: (\Sigma, g_k) \rightarrow \mathbb{S}^l$ . By conformal invariance, we can also view these as harmonic maps from  $(\Sigma, h_k)$  to  $\mathbb{S}^l$ . In this situation, the metric can be recovered by

$$g_k = \frac{|\nabla \Phi_k|_{h_k}^2}{\lambda_1(\Sigma, g_k)} h_k,$$

see [32, proof of Theorem 1]. By pulling back the  $\Phi_k$ 's to  $\widehat{\Sigma}$ , we obtain even harmonic maps  $\widehat{\Phi}_k: (\widehat{\Sigma}, \widehat{h}_k) \rightarrow \mathbb{S}^l$ , such that

$$\widehat{g}_k = \frac{|\nabla \widehat{\Phi}_k|_{\widehat{h}_k}^2}{\lambda_1(\Sigma, g_k)} \widehat{h}_k.$$

With out loss of generality, we may assume that the volume (with respect to  $g_k$ ) concentrates in  $\Sigma_k^1(9a_k)$ . Denote by  $\widehat{\Sigma}_k^1(9a_k)$  its preimage under the covering projection. Note that this preimage might be disconnected. As in [32, Section 4], there are a compact Riemann surface  $\overline{\Sigma}$  and diffeomorphisms  $\tau_k: \overline{\Sigma} \setminus \{p_1, \dots, p_r\} \rightarrow \widehat{\Sigma}_k^1(9a_k)$ . Moreover, the hyperbolic metrics  $\widehat{h}_k = \tau_k^* h_k$  converge in  $C_{loc}^\infty(\overline{\Sigma} \setminus \{p_1, \dots, p_r\})$  to a hyperbolic metric  $\bar{h}$ .

Observe, that we can restrict and pullback the involutions  $\iota_k$  to get involutions  $\bar{\iota}_k$  of  $\overline{\Sigma} \setminus \{p_1, \dots, p_r\}$ . Clearly, these involutions are isometric with respect to the hyperbolic metrics  $\bar{h}_k$ .

In a next step, we construct a fixed point free limit involution on  $\overline{\Sigma}$ . For the compact subsets  $\overline{\Sigma}_c := \{x \in \overline{\Sigma} \mid \text{inj}_x(\overline{\Sigma}, \bar{h}) \geq c\}$ , we can argue exactly as in the proof of Lemma 2.1 to get limit involutions  $\bar{\sigma}_n$  on  $\overline{\Sigma}_{1/n}$ . Since any isometric involution must map  $\overline{\Sigma}_c$  to itself, we may take subsequences, such that for  $m \geq n$ , we have  $\bar{\sigma}_m|_{\overline{\Sigma}_{1/n}} = \bar{\sigma}_n$ . Using a standard diagonal argument, we find a limit involution on  $\overline{\Sigma} \setminus \{p_1, \dots, p_r\}$ . Clearly, this involution extends to an involution  $\bar{\iota}$  on all of  $\overline{\Sigma}$ . Moreover,  $\bar{\iota}$  is fixed point free: Arguing again as in Lemma 2.1, we can not have fixed points different from the  $p_i$ 's. If say  $p_1$  is fixed under  $\bar{\iota}$ , the involution is just rotation by  $\pi$  in a disc centered at  $p_1$ . By  $C^0$ -convergence away from  $p_1$ , we see that the involutions  $\widehat{\iota}_k$  act just via rotation on the collars around the degenerating geodesic. But this is impossible, since this implies that  $\widehat{\iota}_k$  is orientation preserving.

By [37], the pullbacks  $\bar{\Phi}_k$  of the harmonic maps  $\hat{\Phi}_k$  along the diffeomorphisms  $\tau_k$  are then harmonic maps that converge in  $C_{\text{loc}}^\infty(\bar{\Sigma} \setminus \{p_1, \dots, p_r, x_1, \dots, x_s\})$  to a limit harmonic map  $\bar{\Phi}$ . Clearly,  $\bar{\Phi}$  is invariant under  $\bar{\iota}$ . Note, that no energy can be lost at the points  $x_i$  or  $p_i$ . By construction, no volume concentrates near the closed geodesics bounding  $\hat{\Sigma}_k^1(9a_k) \subset \hat{\Sigma}$ , which implies that no energy is lost at the points  $p_i$ . Observe next, that the points  $x_i$  always come in pairs by the invariance of the harmonic maps. Moreover, from the construction of the limit involution, it is clear, that two such points are bounded away from each other. Therefore, energy concentration of the harmonic maps in a point  $x_i$  implies that the volume with respect to the metric  $g_k$  concentrates at a point in  $\Sigma$ . But by [18, Lemma 2.1 and 3.1] this implies

$$\Lambda_1^K(\delta) = \lim_{k \rightarrow \infty} (\Sigma, g_k) \leq 8\pi.$$

Finally, the energy identity from [37] implies that there is also no energy lost in the necks. Let  $\bar{h}_0$  be the hyperbolic metric in the conformal class of the cusp compactification of  $(\bar{\Sigma} \setminus \{p_1, \dots, p_r\}, [\bar{h}])$ . Since  $\bar{\Phi}: (\bar{\Sigma} \setminus \{p_1, \dots, p_s\}, [\bar{h}_0]) \rightarrow S^l$  has finite energy,  $\bar{\Phi}$  extends to a harmonic map  $(\bar{\Sigma}, [\bar{h}_0]) \rightarrow S^l$  [34, Theorem 3.6]. Moreover, this extension is certainly invariant under  $\bar{\iota}$ . In conclusion,  $\bar{\Phi}: (\bar{\Sigma}, \bar{h}_0) \rightarrow S^l$  is an  $\bar{\iota}$  invariant harmonic map with energy

$$\int_{\bar{\Sigma}} |\nabla \bar{\Phi}|^2 dv_{\bar{h}_0} = \lim_{k \rightarrow \infty} \int_{\hat{\Sigma}} |\nabla \hat{\Phi}_k| dv_{\hat{h}_k}.$$

We consider the metric

$$\bar{g} = \frac{|\nabla \bar{\Phi}|_{\bar{h}_0}^2}{\Lambda_1^K(\delta)} \bar{h}_0$$

and observe that it is invariant under the involution  $\bar{\iota}$ , so that it descends to a metric  $g$  on  $\bar{\Sigma}/\bar{\iota}$ . Since there is no energy lost along the sequence  $\bar{\Phi}_k$  of harmonic maps, we have

$$\text{area}(\bar{\Sigma}/\bar{\iota}, g) = 1.$$

Using that the capacity of a point relative to any ball is 0 [22, Chapter 2.2.4], it is easy to construct  $\bar{\iota}_k$ -invariant cut-offs  $\eta_{\varepsilon,k}$  on  $\bar{\Sigma}$  with the following two properties. For  $\varepsilon$  small, there are neighborhoods  $U_\varepsilon \subset V_\varepsilon$  of  $\{p_1, \dots, p_r\}$  such that

- $\bigcap_{\varepsilon > 0} V_\varepsilon = \{p_1, \dots, p_r\}$ ,
- $\eta_{\varepsilon,k} = 0$  in  $U_\varepsilon$ ,
- $\eta_{\varepsilon,k} = 1$  outside  $V_\varepsilon$ , and
- $\int_{\bar{\Sigma}} |\nabla \eta_{\varepsilon,k}|^2 dv_{\bar{g}} \leq \varepsilon^2$ .

We write  $\bar{g}_k = \tau_k^*(\hat{g}_k)$ . Let  $u$  be the lift of a first eigenfunction of  $(\bar{\Sigma}/\bar{i}, g)$  to  $\bar{\Sigma}$ . Using  $\eta_{\varepsilon,k}u$  as a test function on  $\hat{\Sigma}_k(9a_k)$  for  $k$  large enough, we find with the help of the dominated convergence theorem that

$$\begin{aligned} \Lambda_1^K(\delta) &= \lim_{k \rightarrow \infty} \lambda_1(\Sigma, g_k) \\ &\leq \limsup_{\varepsilon \rightarrow 0} \lim_{k \rightarrow \infty} \frac{\int_{\bar{\Sigma}} |\nabla(\eta_{\varepsilon,k}u)|^2 dv_{\hat{g}_k}}{\int_{\bar{\Sigma}} |\eta_{\varepsilon,k}u|^2 dv_{\bar{g}_k} - \left(\int_{\bar{\Sigma}} \eta_{\varepsilon,k}u dv_{\bar{g}_k}\right)^2} \\ &\leq \limsup_{\varepsilon \rightarrow 0} \frac{\int_{\bar{\Sigma}} |\nabla u|^2 dv_{\bar{g}} + C\varepsilon}{\int_{\bar{\Sigma} \setminus V_\varepsilon} |u|^2 dv_{\bar{g}} - \left(\int_{\bar{\Sigma}} u dv_{\bar{g}}\right)^2} \\ &\leq \frac{\int_{\hat{\Sigma}} |\nabla u|^2 dv_{\bar{g}}}{\int_{\bar{\Sigma}} |u|^2 dv_{\bar{g}}} \\ &\leq \lambda_1(\bar{\Sigma}/\bar{i}, g). \end{aligned}$$

If  $\bar{\Sigma}$  is disconnected, it has two connected components and the genus of each component is at most  $\lfloor(\delta - 1)/2\rfloor$ . Therefore, the quotient  $\bar{\Sigma}/\bar{i}$  is an orientable surface of genus at most  $\lfloor(\delta - 1)/2\rfloor$  in this case and thus

$$\Lambda_1^K(\delta) \leq \Lambda_1(\lfloor(\delta - 1)/2\rfloor)$$

thanks to [7]. In case  $\bar{\Sigma}$  is connected, the quotient is non-orientable of non-orientable genus at most  $\delta - 1$  and we have that

$$\Lambda_1^K(\delta) \leq \Lambda_1^K(\delta - 1)$$

again thanks to [7].<sup>1</sup> □

Thanks to the weak inequality  $\Lambda_1^K(\delta + 1) \geq \Lambda_1^K(\delta - 1)$  and the fact that  $\Lambda_1^K(2) > 12\pi$ , we can always rule out the first scenario from Lemma 3.3. Thanks to [19] and [6, 8, 25] our main result Theorem 1.2 follows from the theorem below.

**Theorem 3.7.** *Let  $\delta \geq 3$ . If  $\Lambda_1^K(\delta) > \max\{\Lambda_1^K(\delta - 1), \Lambda_1(\lfloor(\delta - 1)/2\rfloor)\}$ , there is a metric smooth away from finitely many singularities on  $\Sigma_\delta^K$  that achieves  $\Lambda_1^K(\delta)$ .*

*Proof.* By the assumptions, Proposition 2.3, and Proposition 3.2, we can take hyperbolic metrics  $h_k \rightarrow h$  in  $C^\infty$ , such that

$$\lim_{k \rightarrow \infty} \sup_{g \in [h_k]} \lambda_1(\Sigma, g) \text{ area}(\Sigma, g) = \Lambda_1^K(\delta).$$

---

<sup>1</sup> More precisely, the non-orientable version of the result.

As above, we take unit volume metrics  $g_k \in [h_k]$ , such that

$$\lambda_1(\Sigma, g_k) = \sup_{g \in [h_k]} \lambda_1(\Sigma, g) \text{ area}(\Sigma, g).$$

For the corresponding sequence of harmonic maps  $\Phi_k: (\Sigma, h_k) \rightarrow S^l$  no bubbling can occur since this would imply  $\Lambda_1^K(\delta) \leq 8\pi$ , by the same argument as above. Therefore, we can take a subsequence such that  $\Phi_k \rightarrow \Phi$  in  $C^\infty$ , which implies that  $g_k \rightarrow g = \frac{|\nabla\Phi|_h^2}{\Lambda_1^K(\delta)}h$  in  $C^\infty$ . In particular,

$$\lambda_1(\Sigma, g) \text{ area}(\Sigma, g) = \Lambda_1^K(\delta)$$

and  $g$  is smooth away from the branch points of  $\Phi$ . The number of branch points is finite and the branch points correspond to conical singularities of  $g$  [35].  $\square$

### A. Topology of surfaces

For convenience of the reader and the authors, we review here the notion of non-orientable genus.

Recall the classification of closed surfaces. The classes of closed orientable and non-orientable surfaces are both uniquely described up to diffeomorphism by the Euler characteristic. More precisely, any closed orientable surface is diffeomorphic to a surface of the form

$$\Sigma_\gamma = S^2 \# \underbrace{T^2 \# \dots \# T^2}_\gamma \text{ times},$$

and any closed non-orientable surface is diffeomorphic to a surface of the form

$$\Sigma_\delta^K = S^2 \# \underbrace{\mathbb{R}P^2 \# \dots \# \mathbb{R}P^2}_\delta \text{ times}.$$

These two families provide – up to diffeomorphism – a complete list of all orientable respectively non-orientable closed surfaces. We call  $\gamma$  the *genus* of  $\Sigma_\gamma$  and  $\delta$  the *non-orientable genus* of  $\Sigma_\delta^K$ . Note that with this convention, the real projective plane has non-orientable genus 1. We have  $\chi(\Sigma_\gamma) = 2 - 2\gamma$  and  $\chi(\Sigma_\delta^K) = 2 - \delta$ , so that the orientation cover of  $\Sigma_\delta^K$  is given by  $\Sigma_{\delta-1}$ . Some authors prefer to refer to the genus of the orientation cover as the non-orientable genus. As explained above these two definitions differ. Moreover, recall that we have the relation

$$S^2 \# \underbrace{\mathbb{R}P^2 \# \dots \# \mathbb{R}P^2}_\delta \text{ times} \cong S^2 \# \underbrace{T^2 \# \dots \# T^2}_k \text{ times} \# \underbrace{\mathbb{R}P^2 \# \dots \# \mathbb{R}P^2}_{(\delta-2k) \text{ times}},$$

if  $2k < \delta$ .

**Acknowledgements.** Anna Siffert gratefully acknowledges the following funding: the project was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 427320536 – SFB 1442, as well as under Germany’s Excellence Strategy EXC 2044 390685587, Mathematics Münster: Dynamics-Geometry-Structure.

## References

- [1] C. Anné, Spectre du laplacien et écrasement d’anses. *Ann. Sci. École Norm. Sup.* (4) **20** (1987), no. 2, 271–280. [MR 0911759](#) [Zbl 0634.58035](#)
- [2] M. Berger, Sur les premières valeurs propres des variétés riemanniennes. *Compositio Math.* **26** (1973), 129–149. [MR 0316913](#) [Zbl 0257.53048](#)
- [3] G. Besson, Sur la multiplicité de la première valeur propre des surfaces riemanniennes. *Ann. Inst. Fourier (Grenoble)* **30** (1980), no. 1, 109–128. [MR 0576075](#) [Zbl 0417.30033](#)
- [4] P. Buser, *Geometry and spectra of compact Riemann surfaces*. Progress in Mathematics, 106. Birkhäuser Boston, Boston, MA, 1992. [MR 1183224](#) [Zbl 0770.53001](#)
- [5] S. Y. Cheng, Eigenfunctions and nodal sets. *Comment. Math. Helv.* **51** (1976), no. 1, 43–55. [MR 0397805](#) [Zbl 0334.35022](#)
- [6] D. Cianci, M. A. Karpukhin, and V. Medvedev, On branched minimal immersions of surfaces by first eigenfunctions. *Ann. Global Anal. Geom.* **56** (2019), no. 4, 667–690. [MR 4029852](#) [Zbl 1427.53080](#)
- [7] B. Colbois and A. El Soufi, Extremal eigenvalues of the Laplacian in a conformal class of metrics: the “conformal spectrum”. *Ann. Global Anal. Geom.* **24** (2003), no. 4, 337–349. [MR 2015867](#) [Zbl 1036.58026](#)
- [8] A. El Soufi, H. Giacomini, and J. Mustapha, A unique extremal metric for the least eigenvalue of the Laplacian on the Klein bottle. *Duke Math. J.* **135** (2006), no. 1, 181–202. [MR 2259925](#) [Zbl 1109.58029](#)
- [9] A. El Soufi and S. Ilias, Riemannian manifolds admitting isometric immersions by their first eigenfunctions. *Pacific J. Math.* **195** (2000), no. 1, 91–99. [MR 1781616](#) [Zbl 1030.53043](#)
- [10] A. El Soufi and S. Ilias, Extremal metrics for the first eigenvalue of the Laplacian in a conformal class. *Proc. Amer. Math. Soc.* **131** (2003), no. 5, 1611–1618. [MR 1950293](#) [Zbl 1027.58010](#)
- [11] A. Fraser and R. Schoen, Sharp eigenvalue bounds and minimal surfaces in the ball. *Invent. Math.* **203** (2016), no. 3, 823–890. [MR 3461367](#) [Zbl 1337.35099](#)
- [12] A. Girouard, Fundamental tone, concentration of density to points and conformal degeneration on surfaces. *Canad. J. Math.* **61** (2009), no. 3, 548–565. [MR 2514484](#) [Zbl 1198.53039](#)

- [13] J. Hersch, Quatre propriétés isopérimétriques de membranes sphériques homogènes. *C. R. Acad. Sci. Paris Sér. A-B* **270** (1970), A1645–A1648. [MR 0292357](#)  
[Zbl 0224.73083](#)
- [14] D. Jakobson, M. Levitin, N. Nadirashvili, N. Nigam, and I. Polterovich, How large can the first eigenvalue be on a surface of genus two? *Int. Math. Res. Not.* **2005**, no. 63, 3967–3985. [MR 2202582](#) [Zbl 1114.58026](#)
- [15] M. A. Karpukhin, Upper bounds for the first eigenvalue of the Laplacian on non-orientable surfaces. *Int. Math. Res. Not. IMRN* **2016**, no. 20, 6200–6209. [MR 3579963](#)  
[Zbl 1404.58045](#)
- [16] M. A. Karpukhin, Index of minimal spheres and isoperimetric eigenvalue inequalities. *Invent. Math.* **223** (2021), no. 1, 335–377. [MR 4199444](#) [Zbl 07306974](#)
- [17] M. A. Karpukhin, N. Nadirashvili, A. V. Penskoi, I. Polterovich, An isoperimetric inequality for Laplace eigenvalues on the sphere. Preprint, 2017. [arXiv:1706.05713](#)
- [18] G. Kokarev, Variational aspects of Laplace eigenvalues on Riemannian surfaces. *Adv. Math.* **258** (2014), 191–239. [MR 3190427](#) [Zbl 1296.58020](#)
- [19] P. Li and S. T. Yau, A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces. *Invent. Math.* **69** (1982), no. 2, 269–291. [MR 0674407](#) [Zbl 0503.53042](#)
- [20] H. Matthiesen and A. Siffert, Sharp bounds for the first eigenvalue on some degenerating surfaces. Preprint, 2019. [arXiv:1909.02974](#)
- [21] H. Matthiesen and A. Siffert, Handle attachment and the normalized first eigenvalue. Preprint, 2019. [arXiv:1909.03105](#)
- [22] V. Maz'ya, *Sobolev spaces with applications to elliptic partial differential equations*. Second, revised and augmented edition. Grundlehren der Mathematischen Wissenschaften, 342. Springer, Berlin etc., 2011. [MR 2777530](#) [Zbl 1217.46002](#)
- [23] S. Montiel and A. Ros, Minimal immersions of surfaces by the first eigenfunctions and conformal area. *Invent. Math.* **83** (1986), no. 1, 153–166. [MR 0813585](#)  
[Zbl 0584.53026](#)
- [24] D. Mumford, A remark on Mahler's compactness theorem. *Proc. Amer. Math. Soc.* **28** (1971), 289–294. [MR 0276410](#) [Zbl 0215.23202](#)
- [25] N. Nadirashvili, Berger's isoperimetric problem and minimal immersions of surfaces. *Geom. Funct. Anal.* **6** (1996), no. 5, 877–897. [MR 1415764](#) [Zbl 0868.58079](#)
- [26] N. Nadirashvili, Isoperimetric inequality for the second eigenvalue of a sphere. *J. Differential Geom.* **61** (2002), no. 2, 335–340. [MR 1972149](#) [Zbl 1071.58024](#)
- [27] N. Nadirashvili and A. V. Penskoi, An isoperimetric inequality for the second non-zero eigenvalue of the Laplacian on the projective plane. *Geom. Funct. Anal.* **28** (2018), no. 5, 1368–1393. [MR 3856795](#) [Zbl 1400.58010](#)
- [28] N. Nadirashvili and Y. Sire, Conformal spectrum and harmonic maps. *Mosc. Math. J.* **15** (2015), no. 1, 123–140, 182. [MR 3427416](#) [Zbl 1354.58027](#)

- [29] N. Nadirashvili and Y. Sire, Maximization of higher order eigenvalues and applications. *Mosc. Math. J.* **15** (2015), no. 4, 767–775. [MR 3438833](#) [Zbl 1335.35167](#)
- [30] N. Nadirashvili and Y. Sire, Isoperimetric inequality for the third eigenvalue of the Laplace–Beltrami operator on  $S^2$ . *J. Differential Geom.* **107** (2017), no. 3, 561–571. [MR 3715349](#) [Zbl 1385.53027](#)
- [31] S. Nayatani and T. Shoda, Metrics on a closed surface of genus two which maximize the first eigenvalue of the Laplacian. *C. R. Math. Acad. Sci. Paris* **357** (2019), no. 1, 84–98. [MR 3907601](#) [Zbl 1408.58023](#)
- [32] R. Petrides, Existence and regularity of maximal metrics for the first Laplace eigenvalue on surfaces. *Geom. Funct. Anal.* **24** (2014), no. 4, 1336–1376. [MR 3248488](#) [Zbl 1310.58022](#)
- [33] R. Petrides, On the existence of metrics which maximize Laplace eigenvalues on surfaces. *Int. Math. Res. Not. IMRN* **2018**, no. 14, 4261–4355. [MR 3830572](#) [Zbl 1410.58015](#)
- [34] J. Sacks and K. Uhlenbeck, The existence of minimal immersions of 2-spheres. *Ann. of Math. (2)* **113** (1981), no. 1, 1–24. [MR 0604040](#) [Zbl 0462.58014](#)
- [35] S. Salamon, Harmonic and holomorphic maps. In E. Vesentini (ed.), *Geometry seminar “Luigi Bianchi” II – 1984*. Lectures given at the second seminar held at the Scuola Normale Superiore, Pisa, 1984. Lecture Notes in Mathematics, 1164. Springer-Verlag, Berlin, 1985, 161–224. [MR 0829230](#) [Zbl 0591.53031](#)
- [36] P. C. Yang and S. T. Yau, Eigenvalues of the Laplacian of compact Riemann surfaces and minimal submanifolds. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **7** (1980), no. 1, 55–63. [MR 0577325](#) [Zbl 0446.58017](#)
- [37] M. Zhu, Harmonic maps from degenerating Riemann surfaces. *Math. Z.* **264** (2010), no. 1, 63–85. [MR 2564932](#) [Zbl 1213.53086](#)

Received September 8, 2019

Henrik Matthiesen, Max Planck Institute for Mathematics, Vivatsgasse 7, 53111 Bonn, Germany

Current address: Department of Mathematics, University of Chicago, 5734 S. University Ave, Chicago, IL 60637, USA

e-mail: [hmatthiesen@math.uchicago.edu](mailto:hmatthiesen@math.uchicago.edu)

Anna Siffert, Mathematisches Institut, Westfälische Wilhelms-Universität Münster, Einsteinstr. 62, 48149 Münster, Germany

e-mail: [ASiffert@uni-muenster.de](mailto:ASiffert@uni-muenster.de)