

Distinguished self-adjoint extension and eigenvalues of operators with gaps. Application to Dirac–Coulomb operators

Jean Dolbeault, Maria J. Esteban, and Eric Séré

Abstract. We consider a linear symmetric operator in a Hilbert space that is neither bounded from above nor from below, admits a block decomposition corresponding to an orthogonal splitting of the Hilbert space and has a variational gap property associated with the block decomposition. A typical example is the Dirac–Coulomb operator defined on $C_c^\infty(\mathbb{R}^3 \setminus \{0\}, \mathbb{C}^4)$. In this paper we define a distinguished self-adjoint extension with a spectral gap and characterize its eigenvalues in that gap by a min-max principle. This has been done in the past under technical conditions. Here we use a different, geometric strategy, to achieve that goal by making only minimal assumptions. Our result applied to the Dirac–Coulomb-like Hamiltonians covers sign-changing potentials as well as molecules with an arbitrary number of nuclei having atomic numbers less than or equal to 137.

1. Introduction and main result

In three space dimensions, the *free Dirac operator* is of the form $D = -i\alpha \cdot \nabla + \beta$ with

$$\beta = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix}, \quad \alpha_k = \begin{pmatrix} 0 & \sigma_k \\ \sigma_k & 0 \end{pmatrix} \quad (k = 1, 2, 3),$$

$\sigma_1, \sigma_2, \sigma_3$ being the Pauli matrices (see [30]). The *Dirac–Coulomb operator* is $D_V = D + V$ where V is the Coulomb potential $-\frac{\nu}{|x|}$ ($\nu > 0$) or, more generally, the convolution of $-\frac{1}{|x|}$ with an extended charge density. Usually, one first defines $D_{-\nu/|x|}$ on the so-called *minimal domain* $C_c^\infty(\mathbb{R}^3 \setminus \{0\}, \mathbb{C}^4)$. The resulting minimal operator is symmetric but not closed in the Hilbert space $L^2(\mathbb{R}^3, \mathbb{C}^4)$. It is essentially self-adjoint when ν lies in the interval $(0, \sqrt{3}/2]$. In other words, its closure is self-adjoint and there is no other self-adjoint extension. For larger constants ν one must define a distinguished, physically relevant, self-adjoint extension and this can be done when $\nu \leq 1$.

2020 Mathematics Subject Classification. Primary 47B25; Secondary 47A75, 49R50, 81Q10.

Keywords. Variational methods, self-adjoint operators, symmetric operators, quadratic forms, spectral gaps, eigenvalues, min-max principle, Rayleigh-Ritz quotients, Dirac operators.

The essential spectrum of this extension is $\mathbb{R} \setminus (-1, 1)$, which is neither bounded from above nor from below. In atomic physics, its eigenvalues in the gap $(-1, 1)$ are interpreted as discrete electronic energy levels.

Important contributions to the construction of distinguished self-adjoint realisations of Dirac–Coulomb operators were made in the 1970s, see, e.g., [17, 18, 23, 24, 27, 34–36]. In these papers, general classes of potentials V are considered, but in the case $V = -\nu/|x|$ one always assumes that ν is smaller than 1.

Reliable computations of the discrete electronic energy levels in the spectral gap $(-1, 1)$ are a central issue in Relativistic Quantum Chemistry. For this purpose, Talman [29] and Datta and Devaiah [2] proposed a min-max principle involving Rayleigh quotients and the decomposition of four-spinors into their so-called large and small two-components. A related min-max principle based on another decomposition using the free-energy projectors $\mathbb{1}_{\mathbb{R}_{\pm}}(D)$ was proposed in [14] and justified rigorously in [5] for $\nu \in (0, \frac{2}{\pi/2+2/\pi})$. An abstract version of these min-max principles deals with a self-adjoint operator A defined in a Hilbert space \mathcal{H} and satisfying a *variational gap* condition, to be specified later, related to a block decomposition under an orthogonal splitting

$$\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-. \quad (1.1)$$

Such an abstract principle was proved for the first time in [16], but its hypotheses were rather restrictive and the application to the distinguished self-adjoint realization of D_V only gave Talman’s principle for bounded electric potentials (see also [19, 33] for related abstract principles). In [15], an improved abstract min-max principle was applied to D_V with the splitting given by the free-energy projectors, for the unbounded potential $-\nu/|x|$ with $\nu \in (0, 0.305]$. In [4], thanks to a different abstract approach, the range of essential self-adjointness $\nu \in (0, \sqrt{3}/2]$ was dealt with, both for Talman’s splitting and the free projectors. The articles [10, 11, 21, 22, 26] followed and the full range $\nu \in (0, 1]$ is now covered.

Using some of the tools of [4], Esteban and Loss [12, 13] proposed a new strategy to build a distinguished, Friedrichs-like, self-adjoint extension of an abstract *symmetric* operator with variational gap and applied it to the minimal Dirac–Coulomb operator, with $\nu \in (0, 1]$. In [10, 11], connections were established between this new approach and the earlier constructions for Dirac–Coulomb operators.

Important closability issues had been overlooked in some arguments of [4] and some domain invariance questions had not been addressed properly in [12, 13] (see the beginning of Section 3.2). In [26] these issues are clarified and the self-adjoint extension problem considered in [12, 13] is connected to the min-max principle for the eigenvalues of self-adjoint operators studied in [4]. The abstract results in [26] have many important applications, but some examples are not covered yet, due to an essential self-adjointness assumption made on one of the blocks. In the corrigendum [8]

of [4], we present another way of correcting the arguments of [4] thanks to a new geometric viewpoint. In the present work, by adopting this viewpoint, we are able to completely relax the essential self-adjointness assumption of [26]. Additionally, our variational gap assumption is more general, as it covers a class of multi-center Dirac–Coulomb Hamiltonians in which the lower min-max levels fall below the threshold of the continuous spectrum (see, e.g., [6] for a study of such operators): we shall use the image that some eigenvalues *dive* into the negative continuum.

Before going into the detail of our assumptions and results, we fix some general notations that will be used in the whole paper. We consider a Hilbert space \mathcal{H} with scalar product $\langle \cdot, \cdot \rangle$ and associated norm $\| \cdot \|$. When the sum $V + W$ of two subspaces V, W of \mathcal{H} is direct in the algebraic sense, we use the notation $V \dot{+} W$. We reserve the notation $V \oplus W$ to topological sums. We adopt the convention of using the same letter to denote a quadratic form $q(\cdot)$ and its polar form $q(\cdot, \cdot)$. We use the notations $\mathcal{D}(q)$ for the domain of a quadratic form q , $\mathcal{D}(L)$ for the domain of a linear operator L and $\mathcal{R}(L)$ for its range. The space $\mathcal{D}(L)$ is endowed with the norm

$$\|x\|_{\mathcal{D}(L)} := \sqrt{\|x\|^2 + \|Lx\|^2}, \quad \text{for all } x \in \mathcal{D}(L).$$

We denote the resolvent set, spectrum, essential spectrum and discrete spectrum of a self-adjoint operator T by $\rho(T)$, $\sigma(T)$, $\sigma_{\text{ess}}(T)$ and $\sigma_{\text{disc}}(T)$ respectively.

Let us briefly recall the standard Friedrichs extension method. Let $S: \mathcal{D}(S) \rightarrow \mathcal{H}$ be a densely defined operator. Assume that S is symmetric, which means that $\langle Sx, y \rangle = \langle x, Sy \rangle$ for all $x, y \in \mathcal{D}(S)$. If the quadratic form $s(x) := \langle x, Sx \rangle$ associated to S is bounded from below, i.e., if

$$\ell_1 := \inf_{x \in \mathcal{D}(S) \setminus \{0\}} \frac{s(x)}{\|x\|^2} > -\infty,$$

then S has a natural self-adjoint extension T , which is called the *Friedrichs extension* of S and can be constructed as follows (see, e.g. [25] for more details). First of all, since the quadratic form s is bounded from below and associated to a densely defined symmetric operator, it is closable in \mathcal{H} . Denote its closure by \bar{s} . Take $\ell < \ell_1$, so that $\bar{s}(\cdot, \cdot) - \ell \langle \cdot, \cdot \rangle$ is a scalar product on $\mathcal{D}(\bar{s})$ giving it a Hilbert space structure. By the Riesz isomorphism theorem, for each $f \in \mathcal{H}$, there is a unique $u_f \in \mathcal{D}(\bar{s})$ such that $\bar{s}(v, u_f) - \ell \langle v, u_f \rangle = \langle v, f \rangle$ for all $v \in \mathcal{D}(\bar{s})$. Note that u_f is also the unique minimizer of the functional $\mathcal{I}_f(u) := \frac{1}{2}(\bar{s}(u) - \ell \|u\|^2) - \langle u, f \rangle$ in $\mathcal{D}(\bar{s})$. The map $f \mapsto u_f$ is linear, bounded and self-adjoint for $\langle \cdot, \cdot \rangle$. Its inverse is $T - \ell \text{id}_{\mathcal{H}}$ and one easily checks that T does not depend on ℓ : this operator is just the restriction of S^* to $\mathcal{D}(\bar{s}) \cap \mathcal{D}(S^*)$. An important property of the Friedrichs extension is that the eigenvalues of T below its essential spectrum, if they exist, can be characterized by

the classical Courant–Fisher min-max principle: for every positive integer k , the level

$$\ell_k := \inf_{\substack{V \text{ subspace of } \mathcal{D}(S) \\ \dim V = k}} \sup_{x \in V \setminus \{0\}} \frac{s(x)}{\|x\|^2}$$

is either the bottom of $\sigma_{\text{ess}}(T)$ (in the case $\ell_j = \ell_k$ for all $j \geq k$) or the k -th eigenvalue of T (counted with multiplicity) below $\sigma_{\text{ess}}(T)$.

In the special case of the Laplacian in a bounded domain Ω of \mathbb{R}^d with smooth boundary, $S = -\Delta: C_c^\infty(\Omega) \rightarrow L^2(\Omega)$, one has $\mathcal{D}(\bar{s}) = H_0^1(\Omega)$ and the construction of the Friedrichs extension T corresponds to the weak formulation in $H_0^1(\Omega)$ of the Dirichlet problem: $-\Delta u = f$ in Ω , $u = 0$ on $\partial\Omega$. In other words, u_f is the unique function in $H_0^1(\Omega)$ such that for all $v \in H_0^1(\Omega)$, $\int_\Omega \nabla u_f \cdot \nabla v dx = \int_\Omega f v dx$. So, T is the self-adjoint realization of the Dirichlet Laplacian. By regularity theory, we learn that $\mathcal{D}(T) = H^2(\Omega) \cap H_0^1(\Omega)$.

From now on, in this paper, we consider a dense subspace F of \mathcal{H} and a symmetric operator $A: F \rightarrow \mathcal{H}$. We do *not* assume that the quadratic form $a(x) := \langle x, Ax \rangle$ is bounded from below, so we cannot apply the standard Friedrichs extension theorem to A . We introduce an orthogonal splitting $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$ of \mathcal{H} as in (1.1). We denote by

$$\Lambda_\pm: \mathcal{H} \rightarrow \mathcal{H}_\pm$$

the orthogonal projectors associated to this splitting. We make the following assumptions:

$$F_+ := \Lambda_+ F \text{ and } F_- := \Lambda_- F \text{ are subspaces of } F \tag{H1}$$

and

$$\lambda_0 := \sup_{x_- \in F_- \setminus \{0\}} \frac{a(x_-)}{\|x_-\|^2} < +\infty. \tag{H2}$$

We also make the *variational gap assumption* that

$$\text{for some } k_0 \geq 1, \text{ we have } \lambda_{k_0} > \lambda_{k_0-1} = \lambda_0 \tag{H3}$$

where the min-max levels λ_k ($k \geq 1$) are defined by

$$\lambda_k := \inf_{\substack{V \text{ subspace of } F_+ \\ \dim V = k}} \sup_{x \in (V \oplus F_-) \setminus \{0\}} \frac{a(x)}{\|x\|^2}. \tag{1.2}$$

In order to construct a distinguished self-adjoint extension of A , for $E > \lambda_0$ we are going to decompose the quadratic form $a - E \|\cdot\|^2$ as the difference of two quadratic forms q_E and \bar{b}_E with q_E bounded from below and closable, while \bar{b}_E is positive and closed. Before stating our main result, let us define these quadratic forms.

We first introduce a quadratic form b on F_- :

$$b(x_-) = -a(x_-) = \langle x_-, (-\Lambda_- A \upharpoonright_{F_-})x_- \rangle \quad \text{for all } x_- \in F_-. \tag{1.3}$$

For $E > \lambda_0$, it is convenient to define the associated form

$$b_E(x_-) = b(x_-) + E\|x_-\|^2 \quad \text{for all } x_- \in F_-. \tag{1.4}$$

As a consequence of assumption (H2) and of the symmetry of $-\Lambda_- A \upharpoonright_{F_-}$, we have that

$$b_E \text{ is positive definite for all } E > \lambda_0 \text{ and } b \text{ is closable in } \mathcal{H}_-. \tag{b}$$

We denote by \bar{b} the closure of b and by $\bar{b}_E = \bar{b} + E\|\cdot\|^2$ the closure of b_E , their domain being $\mathcal{D}(\bar{b})$. We can consider the Friedrichs extension B of $-\Lambda_- A \upharpoonright_{F_-}$. For every parameter $E > \lambda_0$, the operator $B + E: \mathcal{D}(B) \rightarrow \mathcal{H}_-$ is invertible with bounded inverse. This allows us to define the operator $L_E: F_+ \rightarrow \mathcal{D}(B)$ such that

$$L_E x_+ := (B + E)^{-1} \Lambda_- A x_+, \quad \text{for all } x_+ \in F_+. \tag{1.5}$$

We then introduce the subspace

$$\Gamma_E := \{x_+ + L_E x_+ : x_+ \in F_+\} \subset F_+ \oplus \mathcal{D}(B). \tag{1.6}$$

Making an abuse of terminology justified by the isomorphism

$$F_+ \oplus \mathcal{D}(B) \approx F_+ \times \mathcal{D}(B),$$

we call Γ_E the graph of L_E . On this space, we define a quadratic form q_E by

$$q_E(x_+ + L_E x_+) := \langle x_+, (A - E)x_+ \rangle + \langle L_E x_+, (B + E)L_E x_+ \rangle. \tag{1.7}$$

Denoting by $\bar{\Gamma}_E$ the closure of Γ_E in \mathcal{H} and by Π_E the orthogonal projection on $\bar{\Gamma}_E$, we may write

$$q_E(x) = \langle x, S_E x \rangle, \quad \text{for all } x \in \Gamma_E,$$

where

$$S_E := \Pi_E(\Lambda_+(A - E)\Lambda_+ + \Lambda_-(B + E)\Lambda_-) \upharpoonright_{\Gamma_E}. \tag{1.8}$$

The operator S_E is symmetric and densely defined in the Hilbert space $(\bar{\Gamma}_E, \langle \cdot, \cdot \rangle \upharpoonright_{\bar{\Gamma}_E \times \bar{\Gamma}_E})$. It is one of the two Schur complements associated with the block decomposition of the operator $A - E \text{id}_{\mathcal{H}}$ under the orthogonal splitting $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$. Further details on q_E, S_E are given in Section 2. In particular, in Section 2.1 the decomposition of $a - E\|\cdot\|^2$ in terms of q_E, \bar{b}_E is given. Note that in [4] (before its corrigendum [8]) as well as in [12, 13, 26], the form \bar{b} was already present and a form analogous to q_E was defined, but its domain was F_+ instead of Γ_E .

The main result of this paper is as follows.

Theorem 1. *Let A be a densely defined symmetric operator on the Hilbert space \mathcal{H} with domain F . Assume (H1), (H2), and (H3) and take $E > \lambda_0$. With the above notations, the quadratic forms b and q_E are bounded from below, b is closable in \mathcal{H}_- , q_E is closable in $\bar{\Gamma}_E$ and they satisfy*

$$\mathcal{D}(\bar{q}_E) \cap \mathcal{D}(\bar{b}) = \{0\}.$$

The operator A admits a unique self-adjoint extension \tilde{A} such that

$$\mathcal{D}(\tilde{A}) \subset \mathcal{D}(\bar{q}_E) \dot{+} \mathcal{D}(\bar{b}).$$

The domain of this extension is

$$\mathcal{D}(\tilde{A}) = \mathcal{D}(A^*) \cap (\mathcal{D}(\bar{q}_E) \dot{+} \mathcal{D}(\bar{b}))$$

and it does not depend on E .

Writing

$$\lambda_\infty := \lim \lambda_k \in (\lambda_0, \infty]$$

one has

$$\lambda_\infty = \inf(\sigma_{\text{ess}}(\tilde{A}) \cap (\lambda_0, +\infty)).$$

In addition, the numbers λ_k ($k \geq 1$) satisfying $\lambda_0 < \lambda_k < \lambda_\infty$ are all the eigenvalues – counted with multiplicity – of \tilde{A} in the spectral gap $(\lambda_0, \lambda_\infty)$.

Theorem 1 deserves some comments.

- In some situations, one encounters a symmetric operator A that does not satisfy (H1) but has the weaker property $\Lambda_\pm \mathcal{D}(A) \subset \mathcal{D}(\bar{A})$, where \bar{A} denotes the closure of A . This happens for instance if one defines a Dirac–Coulomb operator on a “minimal” domain consisting of compactly supported smooth functions, and one considers the splitting associated with the free energy projectors $\Lambda_\pm = \mathbb{1}_{\mathbb{R}_\pm}(D)$; see the example of Section 6.2. In such a case one can replace A by its symmetric extension $\bar{A} \upharpoonright_{\Lambda_+ \mathcal{D}(A) \oplus \Lambda_- \mathcal{D}(A)}$. Then (H1) is automatically satisfied by the new domain and it remains to check that the new operator satisfies (H2)–(H3) before applying Theorem 1.

- In the earlier works [4–6, 10, 15, 16, 21, 22, 26] on the min-max principle in gaps, one assumes that $\lambda_1 > \lambda_0$, which amounts to consider assumption (H3) with $k_0 = 1$. Allowing $k_0 \geq 2$ can be important in some applications; see Section 6. The abstract min-max principle for eigenvalues in the case $k_0 \geq 2$ was first considered in [7], but in that paper (H2) was replaced by a much more restrictive assumption. Moreover, the proof in [7] was based on the arguments of [4], so it suffered from the same closability issue solved in [26] and the corrigendum [8] of [4]: the closure of L_E was used but its existence was not proved.

• Compared with [13, 26], another important novelty is that for constructing \tilde{A} we do not need the operator $-\Lambda_-A \upharpoonright_{F_-}$ to be self-adjoint or essentially self-adjoint in \mathcal{H}_- . This assumption was used in [26] to prove that L_E is closable, while in the present work this closability is not needed thanks to a new geometric viewpoint: instead of trying to close L_E we consider the subspace $\bar{\Gamma}_E$, which of course always exists, but is not necessarily a graph. As pointed out in [26], essential self-adjointness of $-\Lambda_-A \upharpoonright_{F_-}$ holds true in many important situations. However, there are also interesting examples for which it does not hold true. An application to Dirac–Coulomb operators in which the essential self-adjointness of $-\Lambda_-A \upharpoonright_{F_-}$ does not hold true is described in Section 6.2. Let us give a simpler example: on the domain $F := (C_c^\infty(\Omega, \mathbb{R}))^2$ consider the operator

$$A \begin{pmatrix} u \\ v \end{pmatrix} := \begin{pmatrix} -\Delta u \\ \Delta v \end{pmatrix} \tag{1.9}$$

taking values in $\mathcal{H} = (L^2(\Omega, \mathbb{R}))^2$, where Ω is a bounded open subset of \mathbb{R}^d with smooth boundary. In this case one takes

$$\Lambda_+ \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} u \\ 0 \end{pmatrix}, \quad \Lambda_- \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 \\ v \end{pmatrix}$$

and (H1) holds true. If $\lambda(\Omega) > 0$ is the smallest eigenvalue of the Dirichlet Laplacian on Ω , we have $\lambda_0 = -\lambda(\Omega)$ in (H2) and $\lambda_1 = \lambda(\Omega) > \lambda_0$, so (H3) with $k_0 = 1$ holds true. But $-\Lambda_-A \upharpoonright_{F_-}$ is the Laplacian defined on the minimal domain $\{0\} \times C_c^\infty(\Omega, \mathbb{R})$, and it is well known that this operator is not essentially self-adjoint in $\{0\} \times L^2(\Omega, \mathbb{R})$, so one cannot apply the abstract results of [13, 26]. In this example, the distinguished extension given by Theorem 1 is easily obtained as follows. One checks that

$$\begin{aligned} \mathcal{D}(\bar{b}) &= \{0\} \times H_0^1(\Omega, \mathbb{R}), \\ \mathcal{D}(\bar{q}_E) &= H_0^1(\Omega, \mathbb{R}) \times \{0\}, \\ \mathcal{D}(A^*) &= (H^2(\Omega, \mathbb{R}))^2. \end{aligned}$$

So, denoting by $\Delta^{(D)}$ the Dirichlet Laplacian with domain $H^2(\Omega, \mathbb{R}) \cap H_0^1(\Omega, \mathbb{R})$, one finds that

$$\tilde{A} = \begin{pmatrix} -\Delta^{(D)} & 0 \\ 0 & \Delta^{(D)} \end{pmatrix}.$$

• In [26], it is proved that the extension \tilde{A} is unique among the self-adjoint extensions whose domain is included in $\Lambda_+ \mathcal{D}(\bar{q}_E) \oplus \mathcal{H}_-$, assuming that the operator $-\Lambda_-A \upharpoonright_{F_-}$ is essentially self-adjoint. But the above example shows that without this assumption, such a uniqueness result does not hold true in general. Indeed,

since $\Delta: C_c^\infty(\Omega, \mathbb{R}) \rightarrow L^2(\Omega, \mathbb{R})$ is not essentially self-adjoint, the operator A given by (1.9) has infinitely many self-adjoint extensions with domains included in $\Lambda_+ \mathcal{D}(\bar{q}_E) \oplus \mathcal{H}_-$. For instance, one can take

$$\hat{A} = \begin{pmatrix} -\Delta^{(D)} & 0 \\ 0 & \Delta^{(N)} \end{pmatrix}$$

with $\Delta^{(N)}$ the self-adjoint extension of Δ associated with the Neumann boundary condition $\nabla v \cdot n = 0$, where n denotes the outward normal unit vector on $\partial\Omega$. Obviously, $\hat{A} \neq \tilde{A}$.

- As we will see in Section 6.1, when dealing with the Dirac–Coulomb operator $D_{-v/|x|}$ with Talman’s splitting, it is natural to choose $F = C_c^\infty(\mathbb{R}^3 \setminus \{0\}, \mathbb{C}^4)$. Then the large and small two-components appearing in Talman’s min-max principle are taken in $C_c^\infty(\mathbb{R}^3 \setminus \{0\}, \mathbb{C}^2)$. But other functional spaces can be used for these components. In [21, 22] an abstract min-max principle is stated in the setting of quadratic forms and applied to Talman’s min-max principle with $H^{1/2}(\mathbb{R}^3, \mathbb{C}^2)$ as space of large and small two-components, under the condition $\nu \in (0, 1)$. However, it seems that some closability issues are present in the proof of the abstract principle, as in [4]. We do not know whether the geometric approach of the present paper could be adapted to the framework of [21] in order to avoid these closability issues without additional assumptions. Note that, by a completely different approach, Talman’s min-max principle is proved in [10] for all $\nu \in (0, 1]$, with arbitrary spaces of large and small two-components lying between $C_c^\infty(\mathbb{R}^3 \setminus \{0\}, \mathbb{C}^2)$ and $H^{1/2}(\mathbb{R}^3, \mathbb{C}^2)$.

Concerning the proof of Theorem 1, we emphasize three main facts.

(1) *The quadratic form $q_E(x) = \langle x, S_E x \rangle$ is bounded from below for all $E > \lambda_0$, so that it has a closure \bar{q}_E in $\bar{\Gamma}_E$ and S_E has a Friedrichs extension T_E .* This fact will allow us to define the distinguished extension \tilde{A} as the restriction of A^* to $\mathcal{D}(A^*) \cap (\mathcal{D}(\bar{q}_E) + \mathcal{D}(\bar{b}))$. We will prove its symmetry thanks to a formula expressing the product $\langle (\tilde{A} - E)X, U \rangle$ in terms of \bar{q}_E and \bar{b}_E , for $X \in \mathcal{D}(\tilde{A})$ and $U \in \mathcal{D}(\bar{q}_E) + \mathcal{D}(\bar{b})$. This formula will be deduced by density arguments from a decomposition of $\|a - E\| \cdot \|^2$ as the difference of q_E and \bar{b}_E .

(2) *The self-adjoint operator T_E is invertible for all $\lambda_0 < E < \lambda_{k_0}$.* Combined with the (obvious) invertibility of $B + E$, this fact will allow us to construct the inverse of the distinguished extension $\tilde{A} - E$, by using once again the formula relating $\langle (\tilde{A} - E)X, U \rangle$ to \bar{q}_E and \bar{b}_E . Then, by a classical argument, we will conclude that \tilde{A} is self-adjoint.

(3) Although we are not able to prove that $\bar{\Gamma}_E$ is a graph above \mathcal{H}_+ , we will see that the sum $\bar{\Gamma}_E + \mathcal{D}(\bar{b})$ is direct in the algebraic sense. More importantly, if we denote by $\pi_E: \bar{\Gamma}_E \dot{+} \mathcal{D}(\bar{b}) \rightarrow \bar{\Gamma}_E$ and $\pi'_E: \bar{\Gamma}_E \dot{+} \mathcal{D}(\bar{b}) \rightarrow \mathcal{D}(\bar{b})$ the associated projec-

tors, the linear map

$$X \in \mathcal{D}(\tilde{A}) \mapsto (\pi_E X, \pi'_E X) \in \bar{\Gamma}_E \times \mathcal{D}(\bar{b})$$

is continuous for the norms $\|X\|_{\mathcal{D}(\tilde{A})}$ and $\|\pi_E X\| + \|\pi'_E X\|$. Thanks to this fact, we will be able to give a relation between the spectra of \tilde{A} and T_E which will allow us to prove the min-max principle for the eigenvalues of \tilde{A} above λ_0 .

For $k_0 = 1$ the facts (1) and (2) are a consequence of the positivity of q_E for $E \in (\lambda_0, \lambda_1)$ and of the Riesz isomorphism theorem. When $k_0 \geq 2$ the positivity is lost, but these two key facts still hold true for other reasons to be given in the proofs of Proposition 6 and Lemma 7 and in Remark 8.

The paper is organized as follows. In Section 2, we study the quadratic form q_E under assumptions (H1) and (H2). In Section 3, under the additional condition (H3) we prove that q_E is bounded from below, then we study its closure \bar{q}_E and the Friedrichs extension T_E . The self-adjoint extension \tilde{A} is constructed in Section 4 and the abstract version of Talman’s principle for its eigenvalues is proved in Section 5, which ends the proof of Theorem 1. Section 6 is devoted to Dirac–Coulomb operators with charge configurations that are not covered by earlier abstract results.

2. The quadratic form q_E

The results of this section are essentially contained in the earlier works [4, 13, 26], we recall them here for the reader’s convenience. We first give a more intuitive interpretation of the objects L_E, q_E that have been defined in the introduction. Then we define a sequence of min-max levels for q_E that will be related to the min-max levels λ_k of A in Section 5.

2.1. A family of maximization problems

In this section we motivate the definition of L_E and q_E given in the introduction. Consider the eigenvalue equation $(A - E)x = 0$ with unknowns $x \in F$ and $E \in \mathbb{R}$. Writing $x_+ := \Lambda_+ x, y_- := \Lambda_- x$ and projecting both sides of the equation on \mathcal{H}_- , one gets

$$\Lambda_- A x_+ + \Lambda_- (A - E) y_- = 0$$

which is also the Euler–Lagrange equation for the problem

$$\sup_{y_- \in F_-} \langle x_+ + y_-, (A - E)(x_+ + y_-) \rangle.$$

Given x_+ in F_+ , one can try to look for a solution y_- . In general, the problem is not solvable in F_- but one can consider a larger space in which a solution exists.

We denote by $L_E x_+$ the generalized solution. In order to make these explanations more precise, we need to express the quadratic form $a - E \| \cdot \|^2$ in terms of q_E, \bar{b}_E .

Given $x_+ \in F_+$ and $E > \lambda_0$, let $\varphi_{E,x_+} : F_- \rightarrow \mathbb{R}$ be defined by

$$\varphi_{E,x_+}(y_-) := \langle x_+ + y_-, (A - E)(x_+ + y_-) \rangle, \quad \text{for all } y_- \in F_-,$$

One easily sees that φ_{E,x_+} has a unique continuous extension to $\mathcal{D}(\bar{b})$ which is the strictly concave function

$$\bar{\varphi}_{E,x_+} : y_- \in \mathcal{D}(\bar{b}) \mapsto \langle x_+, (A - E)x_+ \rangle + 2 \operatorname{Re} \langle Ax_+, y_- \rangle - \bar{b}_E(y_-).$$

The main result of this section is the following.

Proposition 2. *Let A be a symmetric operator on the Hilbert space \mathcal{H} . Assume hypotheses (H1) and (H2), take $E > \lambda_0$, and remember the definition (1.7) of q_E . Then*

- *one has the decomposition*

$$\begin{aligned} \langle X, (A - E)X \rangle &= q_E(\Lambda_+ X + L_E \Lambda_+ X) \\ &\quad - \bar{b}_E(\Lambda_- X - L_E \Lambda_+ X), \quad \text{for all } X \in F; \end{aligned} \quad (2.1)$$

- *for each $x_+ \in F_+$, $L_E x_+$ is the unique maximizer of $\bar{\varphi}_{E,x_+}$ and one has*

$$\begin{aligned} q_E(x_+ + L_E x_+) &= \bar{\varphi}_{E,x_+}(L_E x_+) \\ &= \max_{y_- \in F_-} \bar{\varphi}_{E,x_+}(y_-) \\ &= \sup_{y_- \in F_-} \varphi_{E,x_+}(y_-). \end{aligned} \quad (2.2)$$

Proof. If $X \in F$, taking $x_+ := \Lambda_+ X \in F_+$, $y_- := \Lambda_- X \in F_-$ and $z_- := y_- - L_E x_+ \in \mathcal{D}(B)$, we obtain

$$\begin{aligned} \langle X, (A - E)X \rangle &= \langle x_+, (A - E)x_+ \rangle + 2 \operatorname{Re} \langle Ax_+, y_- \rangle - \langle y_-, (B + E)y_- \rangle \\ &= \langle x_+, (A - E)x_+ \rangle + 2 \operatorname{Re} \langle L_E x_+, (B + E)y_- \rangle \\ &\quad - \langle y_-, (B + E)y_- \rangle \\ &= \langle x_+, (A - E)x_+ \rangle + \langle L_E x_+, (B + E)L_E x_+ \rangle \\ &\quad + \operatorname{Re} \langle L_E x_+, (B + E)z_- \rangle - \operatorname{Re} \langle z_-, (B + E)y_- \rangle \\ &= \langle x_+, (A - E)x_+ \rangle + \langle L_E x_+, (B + E)L_E x_+ \rangle \\ &\quad - \langle z_-, (B + E)z_- \rangle, \end{aligned}$$

which proves (2.1). Now, given $x_+ \in F_+$ this identity can be rewritten in the form

$$\varphi_{E,x_+}(y_-) = q_E(x_+ + L_E x_+) - \bar{b}_E(y_- - L_E x_+), \quad \text{for all } y_- \in F_-.$$

By density of F_- in the Hilbert space $(\mathcal{D}(\bar{b}), \bar{b}_E(\cdot, \cdot))$ one thus has

$$\bar{\varphi}_{E,x_+}(y_-) = q_E(x_+ + L_E x_+) - \bar{b}_E(y_- - L_E x_+), \quad \text{for all } y_- \in \mathcal{D}(\bar{b})$$

and by the positivity of \bar{b}_E one concludes that (2.2) holds true, which completes the proof. ■

2.2. The min-max levels for q_E

If assumptions (H1) and (H2) hold true, to each $E > \lambda_0$ we may associate the nondecreasing sequence of min-max levels $(\ell_k(E))_{k \geq 1}$ defined by

$$\ell_k(E) := \inf_{\substack{V \text{ subspace of } \Gamma_E \\ \dim V = k}} \sup_{x \in V \setminus \{0\}} \frac{q_E(x)}{\|x\|^2} \in [-\infty, +\infty). \tag{2.3}$$

We may also define the (possibly infinite) multiplicity numbers

$$m_k(E) := \text{card}\{k' \geq 1 : \ell_{k'}(E) = \ell_k(E)\} \geq 1. \tag{2.4}$$

In this section we analyze the dependence on E of the quadratic form q_E and its associated min-max levels. The results are summarized in the following proposition.

Proposition 3. *Assume that (H1) and (H2) of Theorem 1 are satisfied. Then*

- *for all $\lambda_0 < E < E'$ and for all $x_+ \in F_+$, we have*

$$\|x_+ + L_{E'}x_+\| \leq \|x_+ + L_E x_+\| \leq \frac{E' - \lambda_0}{E - \lambda_0} \|x_+ + L_{E'}x_+\| \tag{2.5}$$

and

$$\begin{aligned} (E' - E)\|x_+ + L_{E'}x_+\|^2 &\leq q_E(x_+ + L_E x_+) - q_{E'}(x_+ + L_{E'}x_+) \\ &\leq (E' - E)\|x_+ + L_E x_+\|^2; \end{aligned} \tag{2.6}$$

- *for every positive integer k and all $\lambda > \lambda_0$, one has*

$$\ell_k(\lambda) \leq \lambda_k - \lambda; \tag{2.7}$$

- *for every positive integer k , if $\lambda_k > \lambda_0$ then for all $\lambda > \lambda_0$, one has*

$$\ell_k(\lambda) \geq (\lambda_k - \lambda) \left(\frac{\lambda - \lambda_0}{\lambda_k - \lambda_0} \right)^2. \tag{2.8}$$

As a consequence, when $\lambda_k > \lambda_0$ the min-max level $\ell_k(\lambda)$ is finite for every $\lambda > \lambda_0$. It is positive when $\lambda_0 < \lambda < \lambda_k$, negative when $\lambda > \lambda_k$ and one has $\ell_k(\lambda) = 0$ if and only if $\lambda = \lambda_k$. Therefore, the formula $m_k(\lambda_k) = \text{card}\{k' \geq 1 : \lambda_{k'} = \lambda_k\}$ holds true.

Proof. Both formula (2.5) and (2.6) as well as their detailed proof can be found in [4, Lemma 2.1] and [26, Lemma 2.4], so here we just give the main arguments. In order to prove (2.5) one can start from the fact that for all $t \geq -\lambda_0$,

$$(t + E')^{-1} \leq (t + E)^{-1} \leq \frac{E' - \lambda_0}{E - \lambda_0} (t + E')^{-1}.$$

Then one can use the inclusion $\sigma(B) \subset [-\lambda_0, \infty)$ and the definition of L_E . In order to prove (2.6), one notices that this formula is equivalent to the two inequalities $\bar{\varphi}_E(L_{E'}x_+) \leq q_E(x_+ + L_E x_+)$ and $\bar{\varphi}_{E'}(L_E x_+) \leq q_{E'}(x_+ + L_{E'} x_+)$, which both hold true thanks to (2.2).

We now prove (2.7) and (2.8). By definition of λ_k , for each $\varepsilon > 0$ there is a k -dimensional subspace V_ε of F_+ such that for all $x_+ \in V_\varepsilon$ and $y_- \in F_-$, $a(X) \leq (\lambda_k + \varepsilon)\|X\|^2$ with $X = x_+ + y_-$. If $E \in (\lambda_0, \infty)$, then this inequality can be rewritten as

$$\varphi_{E,x_+}(y_-) \leq (\lambda_k - E + \varepsilon)\|x_+ + y_-\|^2.$$

By a density argument, one infers that the inequality

$$\bar{\varphi}_{E,x_+}(y_-) \leq (\lambda_k - E + \varepsilon)\|x_+ + y_-\|^2$$

holds true for all $y_- \in \mathcal{D}(\bar{b})$. Choosing $y_- = L_E x_+$ and using (2.2), one gets the estimate $q_E(x) \leq (\lambda_k - E + \varepsilon)\|x\|^2$ with $x = x_+ + L_E x_+$, hence

$$\sup_{x \in W_\varepsilon \setminus \{0\}} \frac{q_E(x)}{\|x\|^2} \leq \lambda_k - E + \varepsilon$$

with $W_\varepsilon := \{x \in \Gamma_E : \Lambda_+ x \in V_\varepsilon\}$. Since ε is arbitrary and $\dim(W_\varepsilon) = k$, we conclude that (2.7) holds true.

On the other hand, using once again the definition of λ_k , we find that for each $\varepsilon > 0$ and each k -dimensional subspace W of Γ_{λ_k} , there is a nonzero vector x_ε in the k -dimensional space $V := \Lambda_+ W \subset F_+$ and a vector $y_\varepsilon \in F_-$ such that $a(X_\varepsilon) \geq (\lambda_k - \varepsilon)\|X_\varepsilon\|^2$ with $X_\varepsilon = x_\varepsilon + y_\varepsilon$. If $\lambda_k > \lambda_0$, after imposing $\varepsilon < \lambda_k - \lambda_0$ we get $\varphi_{\lambda_k - \varepsilon, x_\varepsilon}(y_\varepsilon) \geq 0$, hence, invoking (2.2), $q_{\lambda_k - \varepsilon}(x_\varepsilon + L_{\lambda_k - \varepsilon} x_\varepsilon) \geq 0$. Then, using (2.5) and (2.6) with the choices $E = \lambda_k - \varepsilon$, $E' = \lambda_k$, we get

$$\begin{aligned} q_{\lambda_k}(x_\varepsilon + L_{\lambda_k} x_\varepsilon) &\geq q_{\lambda_k - \varepsilon}(x_\varepsilon + L_{\lambda_k - \varepsilon} x_\varepsilon) - \varepsilon\|x_\varepsilon + L_{\lambda_k - \varepsilon} x_\varepsilon\|^2 \\ &\geq -\varepsilon \left(\frac{\lambda_k - \lambda_0}{\lambda_k - \varepsilon - \lambda_0} \right)^2 \|x_\varepsilon + L_{\lambda_k} x_\varepsilon\|^2. \end{aligned}$$

Since W and ε are arbitrary, we thus have $\ell_k(\lambda_k) \geq 0$. Combining this with (2.7), we see that $\ell_k(\lambda_k) = 0$.

It remains to study the case $\lambda_k > \lambda_0$ and $\lambda \in (\lambda_0, \infty) \setminus \{\lambda_k\}$. We take an arbitrary k -dimensional subspace \widehat{W} of Γ_λ . We define

$$V := \Lambda_+ \widehat{W} \subset F_+$$

and

$$W := \{x = x_+ + L_{\lambda_k} x_+ : x_+ \in V\} \subset \Gamma_{\lambda_k}.$$

Then W is also k -dimensional, so one has $\sup_{x \in W \setminus \{0\}} \frac{q_{\lambda_k}(x)}{\|x\|^2} \geq 0$, from what we have just seen. So, by compactness of the unit sphere for $\|\cdot\|$ of the k -dimensional space W and the continuity of q_{λ_k} on this space, there is $x_0 \in V$ such that $\|x_0 + L_{\lambda_k} x_0\| = 1$ and $q_{\lambda_k}(x_0 + L_{\lambda_k} x_0) \geq 0$. In order to bound $q_\lambda(x_0 + L_\lambda x_0)$ from below, we use (2.5) and (2.6) with $E = \min(\lambda, \lambda_k)$ and $E' = \max(\lambda, \lambda_k)$. We get

$$\begin{aligned} q_\lambda(x_0 + L_\lambda x_0) &\geq (\lambda_k - \lambda) \|x_0 + L_{\lambda_k} x_0\|^2 \\ &\geq (\lambda_k - \lambda) \left(\frac{\lambda - \lambda_0}{\lambda_k - \lambda_0} \right)^2 \|x_0 + L_\lambda x_0\|^2. \end{aligned}$$

Since \widehat{W} is arbitrary, we conclude that (2.8) holds true.

The last statements of Proposition 3 – finiteness and sign of $\ell_k(\lambda)$, the fact that λ_k is the unique zero of ℓ_k – are an immediate consequence of (2.7) and (2.8). Note that this characterization of λ_k as unique solution of a nonlinear equation was already stated and proved in [4, Lemma 2.2 (c)] and [26, Lemma 2.8 (iii)]. ■

Remark 4. Assumptions (H1) and (H2) are rather easy to check in practice, but checking (H3) is more delicate. The second point in Proposition 3 provides a way to do this: one just has to prove that for some $k_0 \geq 1$ and $E_0 > \lambda_0$ the level $\ell_{k_0}(E_0)$ is nonnegative, which implies that $\lambda_{k_0} \geq E_0$. In Section 6 we will apply this method to one-center and multi-center Dirac–Coulomb operators.

Remark 5. The numerical calculation of eigenvalues in a spectral gap is a delicate issue, due to a well-known phenomenon called *spectral pollution*: as the discretization is refined, one sometimes observes more and more *spurious* eigenvalues that do not approximate any eigenvalue of the exact operator (see [20]). It is possible to eliminate these spurious eigenvalues thanks to Talman’s min-max principle. A method inspired of Talman’s work was proposed in [6, 9]. The idea was to calculate each eigenvalue λ_k as the unique solution of the problem $\ell_k(\lambda) = 0$. This method is free of spectral pollution, but solving nonlinear equations has a computational cost. The estimates (2.7) and (2.8) proved in the present work suggest a fast and stable iterative algorithm that could reduce this cost. Starting from a value $E^{(0)}$ comprised between λ_0 and λ_k , one can compute a sequence of approximations by the formula $E^{(j+1)} = E^{(j)} + \ell_k(E^{(j)})$. From (2.7), one proves by induction that for all $j \geq 0$,

one has $E^{(j)} \in (\lambda_0, \lambda_k)$, $E^{(j+1)} - E^{(j)} = \ell_k(E^{(j)}) > 0$ and $E^{(j)}$ converges monotonically to λ_k . Moreover, combining the inequalities (2.7) and (2.8) one finds that for $|h|$ small, $h + \ell_k(\lambda_k + h) = \mathcal{O}(h^2)$. So, $E^{(j)}$ converges quadratically to λ_k . It would be interesting to perform numerical tests of this algorithm in practical situations.

3. The closure \bar{q}_E and the Friedrichs extension T_E

In this section, under assumptions (H1), (H2), and (H3) we prove that the form q_E is bounded from below and closable, so that the Schur complement S_E has a Friedrichs extension T_E . We then relate the spectrum of T_E to the min-max levels λ_k . Finally, we construct a natural isomorphism between the domains of \bar{q}_E and $\bar{q}_{E'}$ for all $E, E' > \lambda_0$.

3.1. Construction of \bar{q}_E and T_E

In this section we are going to prove the following result.

Proposition 6. *Let A be a symmetric operator on the Hilbert space \mathcal{H} . Assuming (H1), (H2), and (H3) and with the above notations,*

- *for each $E > \lambda_0$, the quadratic form $q_E(x) = \langle x, S_E x \rangle$ is bounded from below hence closable in $\bar{\Gamma}_E$ and S_E has a Friedrichs extension T_E ;*
- *if $E \in (\lambda_0, \lambda_\infty) \setminus \{\lambda_k : k \geq k_0\}$, then $T_E : \mathcal{D}(T_E) \rightarrow \bar{\Gamma}_E$ is invertible with bounded inverse; if $\lambda_0 < \lambda_k < \lambda_\infty$, then 0 is the k -th eigenvalue of T_{λ_k} counted with multiplicity; moreover, its multiplicity is finite and equal to $\text{card}\{k' \geq 1 : \lambda_{k'} = \lambda_k\}$. If $\lambda_k = \lambda_\infty$ for some positive integer k , then $0 = \min \sigma_{\text{ess}}(T_{\lambda_k})$.*

The main tool in the proof of Proposition 6 is the following result.

Lemma 7. *Under assumptions (H1), (H2), and (H3), for every $E > \lambda_0$, there is $\kappa_E > 0$ such that $q_E + \kappa_E \|\cdot\|^2 \geq \|\cdot\|^2$ on Γ_E .*

Proof. We distinguish two cases depending on the value of $k_0 = \min\{k \geq 1 : \lambda_k > \lambda_0\}$.

When $k_0 = 1$, one has $\lambda_1 > \lambda_0$ and $q_{\lambda_1}(x) \geq 0$ for all $x \in \Gamma_{\lambda_1}$. So, using the inequalities (2.5) and (2.6), one finds that for all $E > \lambda_0$ and $x \in \Gamma_E$, $q_E(x) + \kappa_E \|x\|^2 \geq \|x\|^2$, with

$$\kappa_E := 1 + \max\{0, (E - \lambda_1)\} \left(\frac{E - \lambda_0}{\lambda_1 - \lambda_0} \right)^2.$$

When $k_0 \geq 2$, we need a different argument and the formula for κ_E is less explicit. As in the case $k_0 = 1$, we just have to find a constant κ_E for some $E > \lambda_0$; then

the inequalities (2.5) and (2.6) will immediately imply its existence for all $E > \lambda_0$. We take $E \in (\lambda_0, \lambda_{k_0})$. Since $\lambda_{k_0-1} = \lambda_0 < E$, by the second point of Proposition 3 we have $\ell_{k_0-1}(E) \in [-\infty, 0)$. So, there is a $(k_0 - 1)$ -dimensional subspace W of Γ_E such that

$$\ell' := \sup_{w \in W \setminus \{0\}} \frac{q_E(w)}{\|w\|^2} \in (-\infty, 0).$$

Let $C := \sup\{\|S_E w\| : w \in W, \|w\| \leq 1\}$. This constant is finite, since the space W is finite-dimensional. We now consider an arbitrary vector x in Γ_E and we look for a lower bound on $q_E(x)$. We distinguish two cases.

First case: $x \in W$. Then $q_E(x) = \langle x, S_E x \rangle \geq -C\|x\|^2$.

Second case: $x \notin W$. Then the vector space $\text{span}\{x\} \oplus W$ has dimension k_0 . Since $\lambda_{k_0} > E > \lambda_0$, by the third point of Proposition 3 we obtain $\ell_{k_0}(E) > 0$, so there is a vector $w_0 \in W$ such that $q_E(x + w_0) \geq 0$. Then we have

$$\begin{aligned} q_E(x) &= q_E(x + w_0) - 2 \operatorname{Re}\langle x, S_E w_0 \rangle - q_E(w_0) \\ &\geq -2C\|x\|\|w_0\| + |\ell'|\|w_0\|^2 \geq -\frac{C^2}{|\ell'|}\|x\|^2. \end{aligned}$$

In all cases, if we choose $\kappa_E = 1 + \max\{C, C^2/|\ell'|\}$, we get $q_E(x) + \kappa_E\|x\|^2 \geq \|x\|^2$. This completes the proof of the lemma. ■

Proof of Proposition 6. As mentioned in the introduction, we have $q_E(x) = \langle x, S_E x \rangle$, where $S_E: \Gamma_E \rightarrow \bar{\Gamma}_E$ is the Schur complement of the block decomposition of $A - E$ under the splitting $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$ given by formula (1.8). The operator S_E is densely defined in the Hilbert space $\bar{\Gamma}_E$ and it is clearly symmetric, moreover we have just seen that q_E is bounded from below, so q_E is closable in $\bar{\Gamma}_E$. We denote its closure by \bar{q}_E . We call T_E the Friedrichs extension of S_E in $\bar{\Gamma}_E$. With

$$\ell_\infty(E) := \lim_{k \rightarrow \infty} \ell_k(E),$$

the classical min-max principle implies that the levels $\ell_k(E)$ such that $\ell_k(E) < \ell_\infty(E)$ are all the eigenvalues of T_E below $\ell_\infty(E)$ counted with multiplicity, and one has $\ell_\infty(E) = \inf \sigma_{\text{ess}}(T_E)$. So we have the following cases.

If $E \in (\lambda_0, \lambda_\infty) \setminus \{\lambda_k : k \geq 1\}$, then by Proposition 3, one has $0 < \ell_\infty(E)$ and 0 is not in the set $\{\ell_k(E) : k \geq 1\}$. As a consequence, it is not in the spectrum of T_E , so T_E is invertible with bounded inverse.

If $E = \lambda_k$ with $\lambda_0 < \lambda_k < \lambda_\infty$, then, using once again Proposition 3, we find that $\ell_k(E) = 0$ and $\ell_\infty(E) > 0$. So, 0 is an eigenvalue of T_E of finite multiplicity equal to $m_k(\lambda_k)$ where m_k has been defined in (2.4). From Proposition 3, $m_k(\lambda_k)$ equals $\text{card}\{k' \geq 1 : \lambda_{k'} = \lambda_k\}$.

If $\lambda_k = \lambda_\infty$, then for all $k' \geq k$ one has $\lambda_{k'} = \lambda_k$, so $0 = \ell_{k'}(\lambda_k)$ by Proposition 3. In other words, $0 = \ell_\infty(\lambda_k)$. Then the classical min-max theorem implies that $0 = \min \sigma_{\text{ess}}(T_{\lambda_k})$.

Proposition 6 is thus proved. ■

Remark 8. When $k_0 = 1$ and $\lambda_0 < E < \lambda_1$, the closed quadratic form \bar{q}_E is positive definite and the invertibility of T_E is just a consequence of the Riesz isomorphism theorem.

3.2. A family of isomorphisms

In the earlier works [4] (before its corrigendum [8]) and [12, 13, 26], q_E was seen as a quadratic form on F_+ and the domain of its closure was independent of E . Note, however, that the existence of the closure was claimed without proof in [4] and this was a serious gap. Moreover, the proof of the domain invariance was based on an incorrect estimate in [12, Proposition 2] and was incomplete in [13], but this domain invariance problem is easily fixed using [4, inequalities (10)–(11)] which are called (2.5)–(2.6) in the present work.

In our situation, since we do not know whether L_E is closable or not, it is essential to define q_E on Γ_E and then to close it in $\bar{\Gamma}_E$. So, the domain of q_E cannot be independent of E : indeed, it is a subspace of $\bar{\Gamma}_E$ which itself depends on E . However, if we endow each space $\mathcal{D}(\bar{q}_E)$ with the norm $\|x\|_{\mathcal{D}(\bar{q}_E)} := \sqrt{\bar{q}_E(x) + \kappa_E \|x\|^2}$, there is a natural isomorphism $\hat{i}_{E,E'}$ between any two Banach spaces $\mathcal{D}(\bar{q}_E)$ and $\mathcal{D}(\bar{q}_{E'})$, as explained in the next proposition.

Proposition 9. *Under conditions (H1), (H2), and (H3), for $E, E' \in (\lambda_0, \infty)$, the linear map $i_{E,E'}: x_+ + L_E x_+ \mapsto x_+ + L_{E'} x_+$ can be uniquely extended to an isomorphism $\hat{i}_{E,E'}$ between the Banach spaces $\mathcal{D}(\bar{q}_E)$ and $\mathcal{D}(\bar{q}_{E'})$ which can itself be uniquely extended to an isomorphism $\bar{i}_{E,E'}$ between $\bar{\Gamma}_E$ and $\bar{\Gamma}_{E'}$ for the norm $\|\cdot\|$. Moreover, one has the formula*

$$\bar{i}_{E,E'}(x) = x + (E - E')(B + E')^{-1} \Lambda_- x, \quad \text{for all } x \in \bar{\Gamma}_E. \tag{3.1}$$

Proof. The linear map $i_{E,E'}$ is obviously a bijection between Γ_E and $\Gamma_{E'}$, of inverse $i_{E',E}$. The estimates (2.5) and (2.6) of Proposition 3 imply that $i_{E,E'}$ is an isomorphism for the norm $\|\cdot\|$ as well as for the norms $\|\cdot\|_{\mathcal{D}(\bar{q}_E)}$ and $\|\cdot\|_{\mathcal{D}(\bar{q}_{E'})}$, hence the existence and uniqueness of the successive continuous extensions $\hat{i}_{E,E'}$ and $\bar{i}_{E,E'}$, that are isomorphisms of inverses $\hat{i}_{E',E}$ and $\bar{i}_{E',E}$.

From the formula $L_E x_+ = (B + E)^{-1} \Lambda_- A x_+$ one easily gets the resolvent identity

$$L_{E'} x_+ = L_E x_+ + (E - E')(B + E')^{-1} L_E x_+, \quad \text{for all } x_+ \in F_+$$

which is the same as

$$\bar{i}_{E,E'}(x) = x + (E - E')(B + E')^{-1}\Lambda_-x, \quad \text{for all } x \in \Gamma_E.$$

By continuity of $\bar{i}_{E,E'}$, $(B + E')^{-1}$ and Λ_- for the norm $\|\cdot\|$, one can extend the above formula to $\bar{\Gamma}_E$ and this ends the proof of the proposition. ■

Proposition 9 has the following consequence which will be useful in the next section.

Corollary 10. *Assume that conditions (H1), (H2), and (H3) hold true. Let $E, E' > \lambda_0$. Then*

$$\mathcal{D}(\bar{q}_{E'}) + \mathcal{D}(\bar{b}) = \mathcal{D}(\bar{q}_E) + \mathcal{D}(\bar{b}). \tag{3.2}$$

Proof. From formula (3.1), for each $x \in \mathcal{D}(\bar{q}_E)$ one has $\hat{i}_{E,E'}(x) - x \in \mathcal{D}(B)$, hence

$$\mathcal{D}(\bar{q}_{E'}) \subset \mathcal{D}(\bar{q}_E) + \mathcal{D}(B) \subset \mathcal{D}(\bar{q}_E) + \mathcal{D}(\bar{b}),$$

which of course implies the inclusion

$$\mathcal{D}(\bar{q}_{E'}) + \mathcal{D}(\bar{b}) \subset \mathcal{D}(\bar{q}_E) + \mathcal{D}(\bar{b}).$$

Exchanging E and E' , one gets the reverse inclusion, so (3.2) is proved. ■

4. The distinguished self-adjoint extension

In this section, we continue with the proof of Theorem 1 by constructing the distinguished self-adjoint extension \tilde{A} and studying some properties of its domain that will be useful in the sequel. But before doing this, we need to establish a decomposition of the product $\langle (A^* - E)X, U \rangle$ under weak assumptions on the vectors X, U .

4.1. A useful identity

In this section we state and prove an identity that plays a crucial role in the construction and study of \tilde{A} .

Proposition 11. *Assume that conditions (H1), (H2), and (H3) hold true. Let $x, u \in \mathcal{D}(\bar{q}_E)$ and $z_-, v_- \in \mathcal{D}(\bar{b})$ be such that $X = x + z_- \in \mathcal{D}(A^*)$. Then, with $U = u + v_-$, we have*

$$\langle (A^* - E)X, U \rangle = \bar{q}_E(x, u) - \bar{b}_E(z_-, v_-) \tag{4.1}$$

for every $E > \lambda_0$.

Proof. Formula (2.1) of Proposition 2 exactly says that for all $X = x + z_- \in F$ with $x = \Lambda_+X + L_E\Lambda_+X$ and $z_- = \Lambda_-X - L_E\Lambda_+X$, one has

$$\langle X, (A - E)X \rangle = \langle x, S_E x \rangle - \langle z_-, (B + E)z_- \rangle.$$

This relation between quadratic forms directly implies a formula involving their polar forms: for all $X = x + z_- \in F$ and $U = u + v_- \in F$ with $x = \Lambda_+X + L_E\Lambda_+X$, $u = \Lambda_+U + L_E\Lambda_+U$, $z_- = \Lambda_-X - L_E\Lambda_+X$ and $v_- = \Lambda_-U - L_E\Lambda_+U$, one has

$$\langle x + z_-, (A - E)U \rangle = \langle x, S_E u \rangle - \langle z_-, (B + E)v_- \rangle. \tag{4.2}$$

In order to prove Proposition 11, we have to generalize (4.2) to larger classes of vectors X, U . We proceed in two steps.

First step: we fix $U = u + v_-$ in F , with $u = \Lambda_+U + L_E\Lambda_+U$ and $v_- = \Lambda_-U - L_E\Lambda_+U$. If $x = 0$ and $X = z_- \in F_-$, the identity (4.2) holds true and reduces to

$$\langle z_-, (A - E)U \rangle = -\langle z_-, (B + E)v_- \rangle. \tag{4.3}$$

Both sides of (4.3) are continuous in z_- for the norm $\| \cdot \|$, and we recall that F_- is dense in \mathcal{H}_- . So (4.3) remains true for all $z_- \in \mathcal{H}_-$.

Now, in the special case $X = x_+ \in F_+$, $x = x_+ + L_E x_+$ and $z_- = -L_E x_+$, the identity (4.2) becomes

$$\langle x_+, (A - E)U \rangle = \langle x, S_E u \rangle + \langle L_E x_+, (B + E)v_- \rangle.$$

We may also apply (4.3) to $z_- = -L_E x_+$ and we get

$$-\langle L_E x_+, (A - E)U \rangle = \langle L_E x_+, (B + E)v_- \rangle.$$

Subtracting these two identities, we find

$$\langle x, (A - E)U \rangle = \langle x, S_E u \rangle, \quad \text{for all } x \in \Gamma_E. \tag{4.4}$$

Both sides of (4.4) are continuous in x for the norm $\| \cdot \|$, so (4.4) remains true for all $x \in \bar{\Gamma}_E$.

Then, for $x \in \bar{\Gamma}_E$ and $z_- \in \mathcal{H}_-$, we may add (4.3) and (4.4). We conclude that (4.2) remains valid for all $x \in \bar{\Gamma}_E$ and $z_- \in \mathcal{H}_-$. When $x \in \mathcal{D}(\bar{q}_E)$ and $z_- \in \mathcal{D}(\bar{b})$, this identity may be rewritten in the form

$$\langle x + z_-, (A - E)U \rangle = \bar{q}_E(x, u) - \bar{b}_E(z_-, v_-). \tag{4.5}$$

In particular, when $X = x + z_- \in \mathcal{D}(A^*)$ with $x \in \mathcal{D}(\bar{q}_E)$ and $z_- \in \mathcal{D}(\bar{b})$, one obtains the equality

$$\langle (A^* - E)X, U \rangle = \bar{q}_E(x, u) - \bar{b}_E(z_-, v_-). \tag{4.6}$$

This formula is the same as (4.1) under the additional assumptions $U \in F$, $u \in \Gamma_E$ and $v_- = U - u \in \mathcal{D}(B)$.

Second step: we fix $X = x + z_- \in \mathcal{D}(A^*)$ with $x \in \mathcal{D}(\bar{q}_E)$, $z_- \in \mathcal{D}(\bar{b})$. If $u = 0$ and $v_- = U \in F_-$, the identity (4.6) holds true and reduces to

$$\langle (A^* - E)X, v_- \rangle = -\bar{b}_E(z_-, v_-). \tag{4.7}$$

Both sides of (4.7) are continuous in v_- for the norm $\|\cdot\|_{\mathcal{D}(\bar{b})}$ and we recall that F_- is dense in $\mathcal{D}(\bar{b})$ for this norm. So (4.7) remains true for all $v_- \in \mathcal{D}(\bar{b})$.

Now, in the special case $U = u_+ \in F_+$, $u = u_+ + L_E u_+$ and $v_- = -L_E u_+$, the identity (4.6) becomes

$$\langle (A^* - E)X, u_+ \rangle = \bar{q}_E(x, u) + \bar{b}_E(z_-, L_E u_+).$$

We may also apply (4.7) to $v_- = -L_E u_+ \in \mathcal{D}(B) \subset \mathcal{D}(\bar{b})$ and we get

$$-\langle (A^* - E)X, L_E u_+ \rangle = \bar{b}_E(z_-, L_E u_+).$$

Subtracting these two identities, we find

$$\langle (A^* - E)X, u \rangle = \bar{q}_E(x, u), \quad \text{for all } u \in \Gamma_E. \tag{4.8}$$

Both sides of (4.8) are continuous in u for the norm $\|\cdot\|_{\mathcal{D}(\bar{q}_E)}$, and we recall that Γ_E is dense in $\mathcal{D}(\bar{q}_E)$ for this norm. So (4.8) remains true for all $u \in \mathcal{D}(\bar{q}_E)$.

Then, for $u \in \mathcal{D}(\bar{q}_E)$ and $v_- \in \mathcal{D}(\bar{b})$, we may add (4.7) and (4.8) and we finally get (4.1) in the general case. ■

4.2. Construction of the self-adjoint extension

In this section we prove

Proposition 12. *Under conditions (H1), (H2), and (H3), given $E > \lambda_0$ the operator A admits a unique self-adjoint extension \tilde{A} such that $\mathcal{D}(\tilde{A}) \subset \mathcal{D}(\bar{q}_E) + \mathcal{D}(\bar{b})$. This extension is independent of E and defined by*

$$\tilde{A}x := A^*x, \quad \text{for all } x \in \mathcal{D}(\tilde{A}) \tag{4.9}$$

where

$$\mathcal{D}(\tilde{A}) := (\mathcal{D}(\bar{q}_E) + \mathcal{D}(\bar{b})) \cap \mathcal{D}(A^*). \tag{4.10}$$

Moreover, for each E in $(\lambda_0, \lambda_\infty) \setminus \{\lambda_k : k_0 \leq k < \infty\}$, the operator $\tilde{A} - E$ is invertible with bounded inverse given by the formula

$$(\tilde{A} - E)^{-1} = T_E^{-1} \circ \Pi_E - (B + E)^{-1} \circ \Lambda_-. \tag{4.11}$$

Proof. For $E > \lambda_0$, the operator \tilde{A} defined by (4.9)–(4.10) is indeed an extension of A , since

$$\mathcal{D}(A) = F \subset (\Gamma_E + \mathcal{D}(B)) \cap \mathcal{D}(A^*) \subset \mathcal{D}(\tilde{A}) \quad \text{and} \quad A^* \upharpoonright_{\mathcal{D}(A)} = A.$$

By Corollary 10, $\mathcal{D}(\tilde{A})$ is independent of E , as well as $\tilde{A} = A^* \upharpoonright_{\mathcal{D}(\tilde{A})}$. Moreover, the extension \tilde{A} is symmetric: this immediately follows from Proposition 11.

Now, given $f \in \mathcal{H}$ and $E > \lambda_0$, we want to study the equation $(\tilde{A} - E)X = f$. For this purpose, we introduce the following problem written in weak form.

Problem. Find $(x, z_-) \in \mathcal{D}(\bar{q}_E) \times \mathcal{D}(\bar{b})$ such that

$$\begin{cases} \bar{q}_E(x, u) = \langle f, u \rangle, & \text{for all } u \in \mathcal{D}(\bar{q}_E), \\ \bar{b}_E(z_-, v_-) = -\langle f, v_- \rangle, & \text{for all } v_- \in \mathcal{D}(\bar{b}). \end{cases} \quad (\mathcal{P}_f)$$

We recall the identity (4.5), which is a special case of formula (4.1) stated in Proposition 11: if $(x, z_-) \in \mathcal{D}(\bar{q}_E) \times \mathcal{D}(\bar{b})$ then, for all $U \in F$, one has

$$\langle X, (A - E)U \rangle = \bar{q}_E(x, u) - \bar{b}_E(z_-, v_-)$$

with $X = x + z_-$, $u = \Lambda_+ U + L_E \Lambda_+ U$ and $v_- = U - u$. Thanks to this identity, we see that for any solution (x, z_-) of (\mathcal{P}_f) , the sum $X = x + z_-$ satisfies

$$\langle X, (A - E)U \rangle = \langle f, U \rangle, \quad \text{for all } U \in F.$$

As a consequence, X is in $\mathcal{D}(A^*)$ and solves $(A^* - E)X = f$. But this vector is also in $\mathcal{D}(\bar{q}_E) + \mathcal{D}(\bar{b})$, so it solves $(\tilde{A} - E)X = f$.

On the other hand, we can rewrite (\mathcal{P}_f) in terms of the Friedrichs extensions T_E and B .

Problem. Find $(x, z_-) \in \mathcal{D}(T_E) \times \mathcal{D}(B)$ such that

$$\begin{cases} T_E x = \Pi_E(f), \\ (B + E)z_- = -\Lambda_-(f). \end{cases} \quad (4.12)$$

Since $E > \lambda_0$, the operator $B + E$ is invertible with bounded inverse, and by Proposition 6 the same is true with T_E if E is in $(\lambda_0, \lambda_\infty) \setminus \{\lambda_k, k \geq k_0\}$. Then (4.12) has a unique solution given by

$$\begin{cases} x = T_E^{-1} \circ \Pi_E(f), \\ z_- = -(B + E)^{-1} \circ \Lambda_-(f) \end{cases} \quad (4.13)$$

and the vector $X = (T_E^{-1} \circ \Pi_E - (B + E)^{-1} \circ \Lambda_-)(f)$ solves $(\tilde{A} - E)X = f$.

The above discussion shows that for E in $(\lambda_0, \lambda_\infty) \setminus \{\lambda_k, k \geq k_0\}$ the symmetric operator $\tilde{A} - E$ is surjective and admits the bounded operator

$$T_E^{-1} \circ \Pi_E - (B + E)^{-1} \circ \Lambda_-$$

as a right inverse. But it is well known that the surjectivity of a symmetric operator implies its injectivity, since its kernel is orthogonal to its range. So, $\tilde{A} - E$ is invertible and (4.11) holds true. Another classical result is that a densely defined surjective symmetric operator is always self-adjoint: see, e.g., [28, Corollary 3.12]. Applying this to $\tilde{A} - E$, we conclude that \tilde{A} is self-adjoint.

The self-adjoint extension \tilde{A} is thus built. Its uniqueness among those whose domain is contained in $\mathcal{D}(\tilde{q}_E) + \mathcal{D}(\tilde{b})$ is almost trivial. Indeed, if \hat{A} is a self-adjoint extension of A , we must have $\mathcal{D}(\hat{A}) \subset \mathcal{D}(A^*)$, hence, if in addition $\mathcal{D}(\hat{A}) \subset \mathcal{D}(\tilde{q}_E) + \mathcal{D}(\tilde{b})$ then $\mathcal{D}(\hat{A}) \subset \mathcal{D}(\tilde{A})$, which automatically implies $\hat{A} = \tilde{A}$ since both operators are self-adjoint. This completes the proof of Proposition 12. ■

4.3. Direct sums

Recall that in (1.6) we defined the graph Γ_E of L_E as

$$\Gamma_E := \{x_+ + L_E x_+ : x_+ \in F_+\} \subset F_+ \oplus \mathcal{D}(B).$$

A natural question is whether its closure $\bar{\Gamma}_E$ in \mathcal{H} has the graph property $\bar{\Gamma}_E \cap \mathcal{H}_- = \{0\}$. A partial answer to this question is given in the next lemma.

Lemma 13. *Under conditions (H1) and (H2), and with the above notations,*

$$\bar{\Gamma}_E \cap \mathcal{H}_- \subset ((B + E)(F_-))^\perp. \tag{4.14}$$

Proof. The arguments below are essentially contained in the proof of [26, Lemma 2.2], but we repeat them here for the reader’s convenience. If $y \in \bar{\Gamma}_E \cap \mathcal{H}_-$, then there is a sequence (x_n) in F_+ such that $\|x_n\| \rightarrow 0$ and $\|L_E x_n - y\| \rightarrow 0$. Then, for $z \in (B + E)(F_-)$, we may write $\langle y, z \rangle = \lim \langle L_E x_n, z \rangle$. On the other hand,

$$|\langle L_E x_n, z \rangle| = |\langle x_n, A(B + E)^{-1} z \rangle| \leq \|x_n\| \|A(B + E)^{-1} z\| \rightarrow 0,$$

so $\langle y, z \rangle = 0$. ■

If one assumes as in [26] that $\Lambda_- A \upharpoonright_{F_-}$ is essentially self-adjoint, then the subspace $(B + E)(F_-)$ of \mathcal{H}_- is dense in \mathcal{H}_- and one concludes that $\bar{\Gamma}_E$ has the graph property. But we do not make this assumption, and for this reason we cannot infer from (4.14) that $\bar{\Gamma}_E \cap \mathcal{H}_- = \{0\}$. In other words, we do not know whether the operator L_E is closable or not. This is why we have to resort to a geometric strategy in

which the linear subspace $\bar{\Gamma}_E$ replaces the possibly nonexistent closure of L_E . Here is the main difference between the present work and [26].

While we may have $\bar{\Gamma}_E \cap \mathcal{H}_- \neq \{0\}$, the following property holds true, as a consequence of Lemma 13.

Proposition 14. *Under conditions (H1) and (H2) and with the above notations,*

$$\bar{\Gamma}_E \cap \mathcal{D}(\bar{b}) = \{0\}.$$

Proof. From (4.14), we have

$$\begin{aligned} \bar{\Gamma}_E \cap \mathcal{D}(\bar{b}) &= (\bar{\Gamma}_E \cap \mathcal{H}_-) \cap \mathcal{D}((B + E)^{1/2}) \\ &\subset ((B + E)(F_-))^\perp \cap \mathcal{D}((B + E)^{1/2}) \\ &= (B + E)^{-1/2}((B + E)^{1/2}F_-)^\perp = \{0\}, \end{aligned}$$

since $(B + E)^{1/2}F_-$ is dense in \mathcal{H}_- . ■

Proposition 14 tells us that the sum of $\bar{\Gamma}_E$ and $\mathcal{D}(\bar{b})$ is algebraically direct. Let us denote by $\pi_E: \bar{\Gamma}_E \dot{+} \mathcal{D}(\bar{b}) \rightarrow \bar{\Gamma}_E$ and $\pi'_E: \bar{\Gamma}_E \dot{+} \mathcal{D}(\bar{b}) \rightarrow \mathcal{D}(\bar{b})$ the associated projectors. In Section 5 we will need some information on the continuity of the restrictions $\pi_E \upharpoonright_{\mathcal{D}(\tilde{A})}$ and $\pi'_E \upharpoonright_{\mathcal{D}(\tilde{A})}$. These operators are not necessarily continuous for the $\|\cdot\|$ norm, but we have the following result.

Proposition 15. *Under assumptions (H1), (H2), and (H3), for all $E > \lambda_0$, one has*

$$\pi_E(\mathcal{D}(\tilde{A})) \subset \mathcal{D}(T_E) \quad \text{and} \quad \pi'_E(\mathcal{D}(\tilde{A})) \subset \mathcal{D}(B).$$

As a consequence, the domain of \tilde{A} may also be written as

$$\mathcal{D}(\tilde{A}) = (\mathcal{D}(T_E) \dot{+} \mathcal{D}(B)) \cap \mathcal{D}(A^*). \tag{4.15}$$

Moreover, the operator $\pi_E \upharpoonright_{\mathcal{D}(\tilde{A})}$ is continuous for the norms $\|\cdot\|_{\mathcal{D}(\tilde{A})}$, $\|\cdot\|_{\mathcal{D}(T_E)}$ and the operator $\pi'_E \upharpoonright_{\mathcal{D}(\tilde{A})}$ is continuous for the norms $\|\cdot\|_{\mathcal{D}(\tilde{A})}$, $\|\cdot\|_{\mathcal{D}(B)}$. More precisely, there is a positive constant C_E such that for all $X \in \mathcal{D}(\tilde{A})$,

$$\|\pi'_E(X)\|_{\mathcal{D}(B)} \leq C_E \|\Lambda_-(\tilde{A} - E)X\| \quad \text{and} \quad \|\pi_E(X)\|_{\mathcal{D}(T_E)} \leq C_E \|X\|_{\mathcal{D}(\tilde{A})}.$$

The constant C_E remains uniformly bounded when E stays away from λ_0 and ∞ .

Proof. Note that formula (4.11) for the inverse of $\tilde{A} - E$ already proves the two inclusions $\pi_E(\mathcal{D}(\tilde{A})) \subset \mathcal{D}(T_E)$ and $\pi'_E(\mathcal{D}(\tilde{A})) \subset \mathcal{D}(B)$ when E is in $(\lambda_0, \lambda_\infty) \setminus \{\lambda_k: k_0 \leq k < \infty\}$. But we want to prove a statement for all values of E in (λ_0, ∞) and this requires some additional work.

In the arguments below, the constant C_E changes value from line to line but we keep the same notation for the sake of simplicity. We shall use the weak form (\mathcal{P}_f) of the equation $(\tilde{A} - E)X = f$ and the equivalent system of strong equations (4.12), introduced in the proof of Proposition 12. In that proof, f was given, $X = x + z_-$ was unknown and it was shown that for each $E > \lambda_0$ the solvability of (\mathcal{P}_f) is a sufficient condition for the solvability of $(\tilde{A} - E)X = f$. But it turns out that this condition is also necessary. Indeed, taking $E > \lambda_0$, $X \in \mathcal{D}(\tilde{A})$ and defining

$$x := \pi_E(X), \quad z_- := \pi'_E(X), \quad f := (\tilde{A} - E)X,$$

we can apply formula (4.1) of Proposition 11 with the successive choices $U = u \in \mathcal{D}(\bar{q}_E)$, $U = v_- \in \mathcal{D}(\bar{b})$ and this tells us that (x, z_-) satisfies (\mathcal{P}_f) . Then, the second equation of the equivalent system (4.12) implies that

$$z_- = -(B + E)^{-1} \Lambda_- (\tilde{A} - E)X,$$

so z_- is in $\mathcal{D}(B)$ with an estimate of the form

$$\|z_-\|_{\mathcal{D}(B)} \leq C_E \|\Lambda_- (\tilde{A} - E)X\|.$$

This estimate on z_- implies in turn the estimate $\|x\| \leq C_E \|X\|_{\mathcal{D}(\tilde{A})}$, since $x = X - z_-$, $\|X\| \leq \|X\|_{\mathcal{D}(\tilde{A})}$ and $\|z_-\| \leq \|z_-\|_{\mathcal{D}(B)}$. Moreover, the first equation in (4.12) exactly means that x is in $\mathcal{D}(T_E)$ and $T_E x = \Pi_E (\tilde{A} - E)X$, so we finally get the estimate

$$\|x\|_{\mathcal{D}(T_E)} \leq C_E \|X\|_{\mathcal{D}(\tilde{A})}.$$

We thus have the desired inclusions $\pi_E(\mathcal{D}(\tilde{A})) \subset \mathcal{D}(T_E)$ and $\pi'_E(\mathcal{D}(\tilde{A})) \subset \mathcal{D}(B)$, hence $\mathcal{D}(\tilde{A}) \subset \mathcal{D}(T_E) \dot{+} \mathcal{D}(B)$. Then, remembering the definition

$$\mathcal{D}(\tilde{A}) = (\mathcal{D}(\bar{q}_E) \dot{+} \mathcal{D}(\bar{b})) \cap \mathcal{D}(A^*)$$

and the inclusions $\mathcal{D}(T_E) \subset \mathcal{D}(\bar{q}_E)$, $\mathcal{D}(B) \subset \mathcal{D}(\bar{b})$, one easily gets (4.15). This ends the proof of Proposition 15. ■

Remark 16. In Section 5, we do not use all the information contained in Proposition 15: we only need the weaker estimates

$$\|\pi'_E(X)\| \leq C_E \|\Lambda_- (\tilde{A} - E)X\| \quad \text{and} \quad \|\pi_E(X)\| \leq C_E \|X\|_{\mathcal{D}(\tilde{A})}. \quad (4.16)$$

4.4. Variational interpretation when $k_0 = 1$

In the special case $k_0 = 1$, for $\lambda_0 < E < \lambda_1$ the quadratic form \bar{q}_E is positive definite as well as \bar{b}_E and the existence and uniqueness of a solution to the weak problem (\mathcal{P}_f)

directly follows from the Riesz isomorphism theorem. One can even give an interpretation of (\mathcal{P}_f) that generalizes the minimization principle for the Friedrichs extension of semibounded operators mentioned in the introduction. We describe it in this short section, as a side remark.

Assuming that $E \in (\lambda_0, \lambda_1)$ and given $f \in \mathcal{H}$, let us consider the inf-sup problem

$$I_{E,f} = \inf_{x_+ \in F_+} \sup_{y_- \in F_-} \left\{ \frac{1}{2} \langle x_+ + y_-, (A - E)(x_+ + y_-) \rangle - \langle f, x_+ + y_- \rangle \right\}.$$

Of course, in general, $I_{E,f}$ is not attained, but using the decomposition (2.1) and replacing $F = F_+ \oplus F_-$ by the larger space $\mathcal{D}(\bar{q}_E) \dot{+} \mathcal{D}(\bar{b})$, one can transform it into a min-max:

$$\begin{aligned} I_{E,f} &= \inf_{x_+ \in F_+} \sup_{z_- \in \mathcal{D}(B)} \left\{ \frac{1}{2} q_E(x_+ + L_E x_+) - \langle f, x_+ + L_E x_+ \rangle - \frac{1}{2} \bar{b}_E(z_-) - \langle f, z_- \rangle \right\} \\ &= \inf_{x_+ \in F_+} \left\{ \frac{1}{2} q_E(x_+ + L_E x_+) - \langle f, x_+ + L_E x_+ \rangle \right\} \\ &\quad - \inf_{z_- \in \mathcal{D}(B)} \left\{ \frac{1}{2} \bar{b}_E(z_-) + \langle f, z_- \rangle \right\} \\ &= \min_{x \in \mathcal{D}(\bar{q}_E)} \left\{ \frac{1}{2} \bar{q}_E(x) - \langle f, x \rangle \right\} - \min_{z_- \in \mathcal{D}(\bar{b})} \left\{ \frac{1}{2} \bar{b}_E(z_-) + \langle f, z_- \rangle \right\}. \end{aligned}$$

Each of these last two convex minimization problems has a unique solution, and the system of Euler–Lagrange equations solved by the two minimizers is just (\mathcal{P}_f) , so their sum is $X = (\tilde{A} - E)^{-1} f$.

5. The min-max principle

In this section, we establish the min-max principle for the eigenvalues of \tilde{A} that constitutes the last part of Theorem 1.

Proposition 17. *Under assumptions (H1), (H2), and (H3), for $k \geq k_0$, the numbers λ_k satisfying $\lambda_k < \lambda_\infty$ are all the eigenvalues of \tilde{A} in the spectral gap $(\lambda_0, \lambda_\infty)$ counted with multiplicity. Moreover, one has*

$$\lambda_\infty = \inf(\sigma_{\text{ess}}(\tilde{A}) \cap (\lambda_0, \infty)).$$

Even if our assumptions are weaker and our formalism slightly different, the arguments in the proof of Proposition 17 are essentially the same as in [4, § 2] (but some details are missing in that reference) and [26, Section 2.6]. This proof is based on two facts:

- a relation between the min-max levels λ_k and the spectrum of T_E which is provided by the second part of Proposition 6;
- a relation between the spectra of T_E and \tilde{A} which is provided by the next lemma, and whose proof relies on Proposition 11 and on the estimates (4.16) of Remark 16.

Lemma 18. *Under assumptions (H1), (H2), and (H3), let $E > \lambda_0$ and let r be a positive integer. The two following properties are equivalent:*

- (i) for all $\delta > 0$, $\text{Rank}(\mathbb{1}_{(-\delta, \delta)}(T_E)) \geq r$;
- (ii) for all $\varepsilon > 0$, $\text{Rank}(\mathbb{1}_{(E-\varepsilon, E+\varepsilon)}(\tilde{A})) \geq r$.

In other words, $0 \in \sigma_{\text{ess}}(T_E)$ if and only if $E \in \sigma_{\text{ess}}(\tilde{A})$; $0 \in \sigma_{\text{disc}}(T_E)$ if and only if $E \in \sigma_{\text{disc}}(\tilde{A})$ and when this happens they have the same multiplicity; $0 \in \rho(T_E)$ if and only if $E \in \rho(\tilde{A})$.

Proof. If (i) holds true, for each $\delta > 0$ there is a subspace \mathcal{X}_δ of $\mathcal{R}(\mathbb{1}_{(-\delta, \delta)}(T_E))$ of dimension r (we recall the notation $\mathcal{R}(L)$ for the range of an operator L). Then we have $\mathcal{X}_\delta \subset \mathcal{D}(T_E) \subset \mathcal{D}(\tilde{q}_E)$. Using Proposition 11 and the second estimate of (4.16), we find that, for all $x \in \mathcal{X}_\delta$ and $Y \in \mathcal{D}(\tilde{A})$,

$$\begin{aligned} |\langle x, (\tilde{A} - E)Y \rangle| &= |\tilde{q}_E(x, \pi_E(Y))| = |\langle T_E x, \pi_E(Y) \rangle| \\ &\leq \delta \|x\| \|\pi_E(Y)\| \leq C_E \delta \|x\| \|Y\|_{\mathcal{D}(\tilde{A})}. \end{aligned}$$

Assume, in addition, that the property (ii) does not hold true. This means that, for some $\varepsilon_0 > 0$, $\text{Rank}(\mathbb{1}_{(E-\varepsilon_0, E+\varepsilon_0)}(\tilde{A})) \leq r - 1$. Then for each $\delta > 0$ there is $x_\delta \in \mathcal{X}_\delta$ such that $\|x_\delta\| = 1$ and $\mathbb{1}_{(E-\varepsilon_0, E+\varepsilon_0)}(\tilde{A})x_\delta = 0$. So, there is $Y_\delta \in \mathcal{D}(\tilde{A})$ such that $(\tilde{A} - E)Y_\delta = x_\delta$ and $\|Y_\delta\| \leq \varepsilon_0^{-1}$. We thus get $\langle x_\delta, (\tilde{A} - E)Y_\delta \rangle = \|x_\delta\|^2 = 1$ and $C_E \|x_\delta\| \|Y_\delta\|_{\mathcal{D}(\tilde{A})}$ is bounded independently of δ . So, taking δ small enough we obtain $|\langle x_\delta, (\tilde{A} - E)Y_\delta \rangle| > C_E \delta \|x_\delta\| \|Y_\delta\|_{\mathcal{D}(\tilde{A})}$ and this is absurd. We have thus proved by contradiction that (i) implies (ii).

It remains to show that (ii) implies (i). If (ii) holds true, then for each $\varepsilon > 0$ there is a subspace \mathcal{Y}_ε of $\mathcal{R}(\mathbb{1}_{(E-\varepsilon, E+\varepsilon)}(\tilde{A}))$ of dimension r and we have $\mathcal{Y}_\varepsilon \subset \mathcal{D}(\tilde{A}) \subset \mathcal{D}(T_E) \dot{+} \mathcal{D}(B)$. Using Proposition 11 we find that for all $x \in \mathcal{D}(T_E)$ and $Y \in \mathcal{Y}_\varepsilon$,

$$|\langle T_E x, \pi_E(Y) \rangle| = |\tilde{q}_E(x, \pi_E(Y))| = |\langle x, (\tilde{A} - E)Y \rangle| \leq \varepsilon \|x\| \|Y\|.$$

Moreover, for all $Y \in \mathcal{Y}_\varepsilon$, from the first estimate of (4.16) one has

$$\|\pi'_E(Y)\| \leq C_E \|\Lambda_-(\tilde{A} - E)Y\| \leq C_E \varepsilon \|Y\|.$$

So, imposing $\varepsilon \leq \frac{1}{2C_E}$ and using the triangular inequality, we get the estimate $\|Y\| \leq 2\|\pi_E(Y)\|$ for all $Y \in \mathcal{Y}_\varepsilon$. As a consequence, the subspace $V_\varepsilon := \pi_E(\mathcal{Y}_\varepsilon) \subset \mathcal{D}(T_E)$ is r -dimensional and for all $x \in \mathcal{D}(T_E)$ and $y \in V_\varepsilon$, one has

$$|\langle T_E x, y \rangle| \leq 2\varepsilon \|x\| \|y\|.$$

Assume, in addition, that (i) does not hold true. This means that there exists $\delta_0 > 0$ such that $\text{Rank}(\mathbb{1}_{(-\delta_0, \delta_0)}(T_E)) \leq r - 1$. Then for each small ε there is y_ε in V_ε such that $\|y_\varepsilon\| = 1$ and $\mathbb{1}_{(-\delta_0, \delta_0)}(T_E)y_\varepsilon = 0$. So, there is $x_\varepsilon \in \mathcal{D}(T_E)$ such that $T_E x_\varepsilon = y_\varepsilon$ and $\|x_\varepsilon\| \leq \delta_0^{-1}$. We thus get $\langle T_E x_\varepsilon, y_\varepsilon \rangle = \|y_\varepsilon\|^2 = 1$ and $\|x_\varepsilon\| \|y_\varepsilon\| \leq \delta_0^{-1}$. So, taking ε small enough we get $|\langle T_E x_\varepsilon, y_\varepsilon \rangle| > 2\varepsilon \|x_\varepsilon\| \|y_\varepsilon\|$ and this is absurd. We have thus proved by contradiction that (ii) implies (i), so the two properties are equivalent.

Now, given $E > \lambda_0$, 0 is in $\sigma_{\text{ess}}(T_E)$ if and only if (i) holds true for every r , and this is equivalent to saying that (ii) holds true for every r , which exactly means that $E \in \sigma_{\text{ess}}(\tilde{A})$. Similarly, we can say that 0 is in $\sigma_{\text{disc}}(T_E)$ and has multiplicity μ_E as an eigenvalue if and only if (i) holds true for μ_E but not for $\mu_E - 1$, and this is equivalent to saying that (ii) holds true for μ_E but not for $\mu_E - 1$, which exactly means that $E \in \sigma_{\text{disc}}(\tilde{A})$ with multiplicity μ_E . The last statement on $\rho(\tilde{A})$ follows immediately, since for any operator L , $\sigma_{\text{ess}}(L)$, $\sigma_{\text{disc}}(L)$ and $\rho(L)$ form a partition of \mathbb{C} . This ends the proof of the lemma. ■

Proof of Proposition 17. Let us define

$$\underline{\lambda} := \inf(\sigma_{\text{ess}}(\tilde{A}) \cap (\lambda_0, \infty)) \in [\lambda_0, \infty].$$

By Proposition 6, if $E \in (\lambda_0, \lambda_\infty)$, then 0 is either an element of $\rho(T_E)$ or an eigenvalue of T_E of finite multiplicity μ_E . The second case occurs when $E = \lambda_k$ for some positive integer k . Then $\mu_E = \text{card}\{k' : \lambda_{k'} = \lambda_k\}$. So, by Lemma 18, $(\lambda_0, \lambda_\infty) \cap \sigma_{\text{ess}}(\tilde{A})$ is empty hence $\lambda_\infty \leq \underline{\lambda}$, and the levels λ_k in $(\lambda_0, \lambda_\infty)$ are all the eigenvalues of \tilde{A} in this open interval, counted with multiplicity.

It remains to prove that $\underline{\lambda} \leq \lambda_\infty$. The nontrivial case is when the sequence (λ_k) is bounded, so that $\lambda_\infty \in (\lambda_0, \infty)$. If the sequence is nonstationary and bounded, then $\{\lambda_k : k \geq k_0\}$ is an infinite subset of $\sigma_{\text{disc}}(\tilde{A})$, so its limit point λ_∞ is in $\sigma_{\text{ess}}(\tilde{A})$. If the sequence is stationary, let k be such that $\lambda_k = \lambda_\infty$. Then, by Proposition 6, $0 \in \sigma_{\text{ess}}(T_{\lambda_\infty})$ so, by Lemma 18, we find once again that $\lambda_\infty \in \sigma_{\text{ess}}(\tilde{A})$. In conclusion, one always has $\underline{\lambda} \leq \lambda_\infty$ and this ends the proof of Proposition 17. ■

Proof of Theorem 1. Propositions 12, 14, and 17 together imply Theorem 1. ■

6. Applications to Dirac–Coulomb operators

In this section, we consider the three-dimensional *Dirac–Coulomb operator* $D_V = D + V$ mentioned in the introduction. We assume that V is a linear combination of Coulomb potentials $|x - x_j|^{-1}$ due to J distinct point-like charges located at x_1, \dots, x_J . If we define D_V on the minimal domain $F = C_c^\infty(\mathbb{R}^3 \setminus \{x_1, \dots, x_J\}, \mathbb{C}^4)$, it is obviously symmetric in the Hilbert space $\mathcal{H} = L^2(\mathbb{R}^3, \mathbb{C}^4)$. Thanks to Theorem 1

we are going to construct a distinguished self-adjoint realization of D_V and give a min-max principle for its eigenvalues, under some conditions on the coefficients of the linear combination. In each case, assumption (H3) will be checked by the method of Remark 4.

6.1. The attractive case

In this section we assume that $V(x) = -\sum_{j=1}^J \frac{v_j}{|x-x_j|}$ is an attractive potential generated by J distinct point-like nuclei, each having Z_j protons with $0 < Z_j \leq Z_* \approx 137.04$ so that $0 < v_j = Z_j/Z_* \leq 1$ (we allow non-integer values of Z_j). We are going to use Talman’s splitting $\Lambda_+\psi = \begin{pmatrix} \phi \\ 0 \end{pmatrix}$, $\Lambda_-\psi = \begin{pmatrix} 0 \\ \chi \end{pmatrix}$ of four-spinors $\psi = \begin{pmatrix} \phi \\ \chi \end{pmatrix}$ into upper and lower two-spinors, also called *large* and *small two-components*. Then $\Lambda_+F = \mathfrak{F} \times \{0\}$ and $\Lambda_-F = \{0\} \times \mathfrak{F}$ with $\mathfrak{F} := C_c^\infty(\mathbb{R}^3 \setminus \{x_1, \dots, x_J\}, \mathbb{C}^2)$. With the standard notation $\sigma = (\sigma_1, \sigma_2, \sigma_3)$ for the collection of Pauli matrices, we recall (see [30]) that

$$D_V \begin{pmatrix} \phi \\ \chi \end{pmatrix} = \begin{pmatrix} -i\sigma \cdot \nabla \chi + (1 + V)\phi \\ -i\sigma \cdot \nabla \phi - (1 - V)\chi \end{pmatrix}.$$

Assumptions (H1) and (H2) are easily checked with $\lambda_0 = -1$. It remains to check assumption (H3). By Remark 4, it suffices to show that for some $k_0 \geq 1$, $\ell_{k_0}(0) \geq 0$. Indeed, this inequality implies that $\lambda_{k_0} \geq 0 > \lambda_0$. So, we are led to study the quadratic form q_0 . For $\phi \in \mathfrak{F}$ and $\psi_+ = \begin{pmatrix} \phi \\ 0 \end{pmatrix}$, the quantity $q_0(\psi_+ + L_0\psi_+)$ is a function of ϕ , V and in the rest of the section it is more convenient to denote it by $\alpha^V(\phi)$. With this notation, we have

$$\alpha^V(\phi) = \int_{\mathbb{R}^3} \left\{ \frac{|\sigma \cdot \nabla \phi|^2}{1 - V} + (1 + V)|\phi|^2 \right\}, \quad \text{for all } \phi \in \mathfrak{F}. \tag{6.1}$$

We start by the potential $V(x) = -\nu|x|^{-1}$ with $0 < \nu \leq 1$, corresponding to a unique point-like nucleus. We recall the Hardy–Dirac inequality

$$\alpha^{-|\cdot|^{-1}}(\phi) = \int_{\mathbb{R}^3} \left\{ \frac{|\sigma \cdot \nabla \phi|^2}{1 + |x|^{-1}} + (1 - |x|^{-1})|\phi|^2 \right\} \geq 0, \quad \text{for all } \phi \in C_c^\infty(\mathbb{R}^3 \setminus \{0\}, \mathbb{C}^2) \tag{6.2}$$

proved in [3, 4]. Since $\alpha^{-\nu|\cdot|^{-1}} \geq \alpha^{-|\cdot|^{-1}}$, (6.2) implies that $\ell_1(0) \geq 0$ and assumption (H3) is satisfied with $k_0 = 1$. Then, using Theorem 1 we can define a distinguished self-adjoint extension of D_V for $0 < \nu \leq 1$ and we can also characterize all the eigenvalues of this extension in the spectral gap $(-1, 1)$ by the min-max principle (1.2). This is not a new result; see [4, 10, 12, 26], and it is known that V can be replaced by more general attractive potentials that are bounded from below by $-|x|^{-1}$.

We now assume that $J \geq 2$. In such a case, the distinguished self-adjoint extension was constructed in [17, 24] in the subcritical case $\nu_i < 1$ for all i by a method completely different from the one considered in the present work. Talman’s min-max principle for the eigenvalues of the extension was studied in [11], also in the subcritical case. But that paper appealed to the abstract result of [7] and as mentioned in the introduction, the arguments in [7] suffered from the same closability issue as [4]. Theorem 1 solves this issue, moreover it provides a unified treatment: construction of the extension and justification of the min-max principle even in the critical case, i.e., when some of the coefficients ν_i are equal to 1. But of course, in order to apply this theorem we have to check (H3) and this is more delicate than in the one-center case. Indeed, when the total number of protons $\sum_j Z_j$ is larger than 137.04, if the nuclei are close to each other one expects some eigenvalues of the distinguished extension to *dive* into the negative continuum. If this happens, the corresponding min-max levels λ_k should become equal to λ_0 . To check assumption (H3) in such a situation, let us prove by contradiction that for *some* $k_0 \geq 1$, the inequality $\ell_{k_0}(0) \geq 0$ holds true.

Otherwise, there exists a sequence $(G_k)_{k \geq 1}$ of k -dimensional subspaces of \mathfrak{F} such that $\alpha^V(\phi) < 0$ for all $\phi \in G_k \setminus \{0\}$. So, one can construct by induction a sequence (ϕ_k) of wave functions such that $\phi_k \in G_k$ and $\langle \phi_k, \phi_l \rangle_{L^2(\mathbb{R}^3, \mathbb{C}^2)} = \delta_{kl}$. Then ϕ_k converges weakly to 0 in $L^2(\mathbb{R}^3, \mathbb{C}^2)$. In order to derive a contradiction, one can try to prove that for k large enough, $\alpha^V(\phi_k) \geq 0$. In the subcritical case $\nu_i < 1$ for all i this has been done in [11, Section 6, Step 4, p. 1448f]. We give below a proof that is also valid in the critical case. In what follows, the constant C changes from line to line but we keep the same notation for the sake of simplicity.

With $\delta := \frac{1}{2} \min_{1 \leq j < j' \leq K} |x_j - x_{j'}|$ one takes $R > \delta + \max_{1 \leq j \leq J} |x_j|$ (to be chosen later) and a partition of unity $(\theta_j)_{0 \leq j \leq J+1}$ consisting of smooth functions with values in $[0, 1]$ such that $\sum_{j=0}^{J+1} \theta_j^2 = 1$, $\text{supp}(\theta_0) \subset B(0, 2R) \setminus \bigcup_{j=1}^J B(x_j, \delta/2)$, $\text{supp}(\theta_j) \subset B(x_j, \delta)$ for $1 \leq j \leq J$ and $\text{supp}(\theta_{J+1}) \cap B(0, R) = \emptyset$. The pointwise IMS formula [1, Lemma 4.1] for the Pauli operator gives

$$|\sigma \cdot \nabla \phi_k|^2 = \sum_{j=0}^{J+1} |\sigma \cdot \nabla(\theta_j \phi_k)|^2 - \left(\sum_{j=0}^{J+1} |\nabla \theta_j|^2 \right) |\phi_k|^2,$$

so, remembering that $\|\phi_k\|_{L^2(\mathbb{R}^3)}^2 = 1$, one gets

$$\begin{aligned} \alpha^V(\phi_k) &= \sum_{j=0}^{J+1} \alpha^V(\theta_j \phi_k) - \int_{\mathbb{R}^3} \left(\sum_{j=0}^{J+1} |\nabla \theta_j|^2 \right) \frac{|\phi_k|^2}{1 - V} \\ &= \frac{1}{2} + \sum_{j=0}^J \alpha^V(\theta_j \phi_k) + \left(\alpha^V(\theta_{J+1} \phi_k) - \frac{1}{2} \|\theta_{J+1} \phi_k\|_{L^2(\mathbb{R}^3)}^2 \right) \end{aligned}$$

$$\begin{aligned}
 & - \int_{\mathbb{R}^3} \left(\frac{1 - \theta_{J+1}^2}{2} + \frac{1}{1 - V} \sum_{j=0}^{J+1} |\nabla \theta_j|^2 \right) |\phi_k|^2 \\
 & \geq \frac{1}{2} + \sum_{j=0}^J \mathfrak{q}^V(\theta_j \phi_k) + \left(\mathfrak{q}^V(\theta_{J+1} \phi_k) - \frac{1}{2} \|\theta_{J+1} \phi_k\|_{L^2(\mathbb{R}^3)}^2 \right) \\
 & \quad - C \int_{B(0, 2R)} |\phi_k|^2.
 \end{aligned}$$

From now on, we fix R such that $-V \leq 1/4$ on $\mathbb{R}^3 \setminus B(0, R)$. Then one has

$$\mathfrak{q}^V(\theta_{J+1} \phi_k) - \frac{1}{2} \|\theta_{J+1} \phi_k\|_{L^2(\mathbb{R}^3)}^2 \geq \frac{1}{4} \|\theta_{J+1} \phi_k\|_{H^1(\mathbb{R}^3)}^2.$$

Let

$$\begin{aligned}
 M := 1 + \max \{ & \sup_{x \in \text{supp}(\theta_0)} -V(x); \quad \sup_{x \in \text{supp}(\theta_1)} (-V(x) - |x - x_1|^{-1}); \dots; \\
 & \sup_{x \in \text{supp}(\theta_J)} (-V(x) - |x - x_J|^{-1}) \}.
 \end{aligned}$$

Then

$$\mathfrak{q}^V(\theta_0 \phi_k) \geq \frac{1}{M} \|\theta_0 \phi_k\|_{H^1(\mathbb{R}^3)}^2 - C \int_{\mathbb{R}^3} |\theta_0 \phi_k|^2$$

and, introducing the rescaled functions $\hat{\phi}_{j,k}(y) := (\theta_j \phi_k)(x_j + M^{-1}y)$ for $1 \leq j \leq J$, one finds

$$\mathfrak{q}^V(\theta_j \phi_k) \geq \frac{1}{M^2} \mathfrak{q}^{-|\cdot|^{-1}}(\hat{\phi}_{j,k}) - C \int_{\mathbb{R}^3} |\theta_j \phi_k|^2.$$

Gathering these estimates, one gets the lower bound

$$\begin{aligned}
 \mathfrak{q}^V(\phi_k) & \geq \frac{1}{2} + \frac{1}{M^2} \sum_{j=1}^J \mathfrak{q}^{-|\cdot|^{-1}}(\hat{\phi}_{j,k}) + \frac{1}{M} \|\theta_0 \phi_k\|_{H^1(\mathbb{R}^3)}^2 \\
 & \quad + \frac{1}{4} \|\theta_{J+1} \phi_k\|_{H^1(\mathbb{R}^3)}^2 - C \int_{B(0, 2R)} |\phi_k|^2. \tag{6.3}
 \end{aligned}$$

From the Hardy–Dirac inequality (6.2), each of the terms $\mathfrak{q}^{-|\cdot|^{-1}}(\hat{\phi}_{j,k})$ is nonnegative, so the assumptions that $\|\phi_k\|_{L^2(\mathbb{R}^3)} = 1$ and $\mathfrak{q}^V(\phi_k) < 0$ imply that the quantities $\mathfrak{q}^{-|\cdot|^{-1}}(\hat{\phi}_{j,k})$, $\|\theta_0 \phi_k\|_{H^1}$ and $\|\theta_{J+1} \phi_k\|_{H^1}$ are uniformly bounded. But from [10, Theorem 1.9], for $0 \leq s < 1/2$ there is a positive constant κ_s such that

$$\mathfrak{q}^{-|\cdot|^{-1}}(\phi) + \|\phi\|_{L^2(\mathbb{R}^3)}^2 \geq \kappa_s \|\phi\|_{H^s(\mathbb{R}^3)}^2, \quad \text{for all } \phi \in \mathfrak{F}.$$

Applying this inequality to the functions $\hat{\phi}_{j,k}$ ($1 \leq j \leq J$), one easily finds that the sequence $(\phi_k)_{k \geq 1}$ is bounded in $H^s(\mathbb{R}^3)$, hence precompact in $L^2_{loc}(\mathbb{R}^3)$. Since this sequence converges weakly to zero in $L^2(\mathbb{R}^3)$, one concludes that

$$\lim_{k \rightarrow \infty} \int_{B(0,2R)} |\phi_k|^2 = 0.$$

Combining this information with (6.3) one finds that for k large enough, $q^V(\phi_k) \geq 0$ and this is a contradiction.

In conclusion, the assumptions of Theorem 1 are satisfied in our multi-center example, with k_0 possibly larger than 1.

6.2. The sign-changing case

We now consider a potential of the form

$$V(x) = -\frac{\nu_1}{|x|} + \frac{\nu_2}{|x - x_0|} \quad \text{with } x_0 \neq 0, 0 < \nu_1 \leq 1, \text{ and } 0 < \nu_2 \leq \frac{2}{\frac{\pi}{2} + \frac{2}{\pi}}.$$

The corresponding Dirac–Coulomb operator D_V is obviously symmetric if we define it on the “minimal” domain $C_c^\infty(\mathbb{R}^3 \setminus \{0, x_0\}, \mathbb{C}^4)$. But Talman’s decomposition in upper and lower spinors cannot be used: due to the unbounded repulsive term $\frac{\nu_2}{|x - x_0|}$, (H2) would not be satisfied. Instead, for the splitting we choose the free-energy projectors

$$\Lambda_\pm = \mathbb{1}_{\mathbb{R}_\pm}(D).$$

We recall (see [30]) that

$$D\Lambda_\pm = \Lambda_\pm D = \pm\sqrt{1 - \Delta}\Lambda_\pm = \pm\Lambda_\pm\sqrt{1 - \Delta}.$$

In momentum space (i.e., after Fourier transform), Λ_\pm becomes the multiplication operator by the matrix

$$M_\pm(p) = \frac{1}{2} \left(I_4 \pm \frac{\alpha \cdot p + \beta}{\sqrt{|p|^2 + 1}} \right).$$

This matrix depends smoothly on p and is bounded on \mathbb{R}^3 as well as its derivatives. As a consequence, the multiplication by M_\pm preserves the Schwartz class $\mathcal{S}(\mathbb{R}^3, \mathbb{C}^4)$. So, the same is true for Λ_\pm in position space. But this nonlocal operator does not preserve the compact support property, so (H1) does not hold for the domain $C_c^\infty(\mathbb{R}^3 \setminus \{0, x_0\}, \mathbb{C}^4)$. Since $\Lambda_+ C_c^\infty(\mathbb{R}^3 \setminus \{0, x_0\}, \mathbb{C}^4) \subset \mathcal{S}(\mathbb{R}^3, \mathbb{C}^4) \subset \mathcal{D}(\overline{D_V})$, one can either replace the minimal domain by

$$F = \Lambda_+ C_c^\infty(\mathbb{R}^3 \setminus \{0, x_0\}, \mathbb{C}^4) \oplus \Lambda_- C_c^\infty(\mathbb{R}^3 \setminus \{0, x_0\}, \mathbb{C}^4)$$

as mentioned in the first comment after Theorem 1, or by

$$F = \mathcal{S}(\mathbb{R}^3, \mathbb{C}^4).$$

In what follows, A is the restriction of $\overline{D_V}$ to one of these two domains. We do not need to specify which one: the arguments proving (H2)–(H3) are the same in both cases.

By the upper bound on ν_2 , it follows from an inequality of Tix [32] that the Brown–Ravenhall operator $-\Lambda_-(A + 1 - \nu_2) \upharpoonright_{F_-} = \Lambda_-(\sqrt{1 - \Delta} - V - 1 + \nu_2) \upharpoonright_{F_-}$ is nonnegative, so (H2) holds true with $\lambda_0 \leq -1 + \nu_2$. In order to bound λ_1 from below, we can use [4, inequality (38)]. This inequality involves a parameter $\nu \in (0, 1)$ and is stated for all functions $\psi_+ \in F_+$. One easily passes to the limit $\nu \rightarrow 1$ with ψ_+ fixed and this gives us the inequality

$$\begin{aligned} \langle \psi_+, \sqrt{1 - \Delta} \psi_+ \rangle_{L^2(\mathbb{R}^3)} - \int_{\mathbb{R}^3} \frac{|\psi_+|^2}{|x|} \\ + \left\langle \Lambda_- \frac{1}{|x|} \psi_+, (B_{-|x|^{-1}})^{-1} \Lambda_- \frac{1}{|x|} \psi_+ \right\rangle_{L^2(\mathbb{R}^3)} \geq 0 \end{aligned} \quad (6.4)$$

for all $\psi_+ \in F_+$. Here, we denote by $B_{\mathcal{V}}$ the Friedrichs extension of the Brown–Ravenhall operator $\Lambda_-(\sqrt{1 - \Delta} - \mathcal{V}) \upharpoonright_{F_-}$, for any electric potential \mathcal{V} such that this operator is bounded from below. Inequality (6.4) exactly says that if one chooses $(\nu_1, \nu_2) = (1, 0)$, then there holds $q_0(\psi_+ + L_0 \psi_+) \geq 0$ for all $\psi_+ \in F_+$, so $\ell_1(0) \geq 0$, hence $\lambda_1 \geq 0 > \lambda_0$. This remains true for $0 < \nu_1 \leq 1$ and $0 < \nu_2 \leq \frac{2}{\pi/2 + 2/\pi}$, since the min-max level λ_1 is a non-decreasing function of V . Thus, Theorem 1 can be applied with $k_0 = 1$ in order to find a distinguished self-adjoint extension of D_V and to characterize its eigenvalues by a min-max principle.

Note that by [31, Corollary 3], the operator $-\Lambda_- A \upharpoonright_{F_-}$ is not essentially self-adjoint for $\nu_2 > 3/4$. So, the abstract result [26, Theorem 1.1] cannot be applied in this case.

Acknowledgments. The authors wish to thank the referees for their very careful reading of the manuscript, and for many comments and suggestions that greatly improved the quality of this paper.

Funding. Jean Dolbeault has been partially supported by the Project EFI (ANR-17-CE40-0030) and Maria J. Esteban and Eric Séré by the project molQED (ANR-17-CE29-0004) of the French National Research Agency (ANR).

References

- [1] R. Bosi, J. Dolbeault, and M. J. Esteban, [Estimates for the optimal constants in multipolar Hardy inequalities for Schrödinger and Dirac operators](#). *Commun. Pure Appl. Anal.* **7** (2008), no. 3, 533–562 Zbl 1162.35329 MR 2379440
- [2] S. N. Datta and G. Devaiah, [The minimax technique in relativistic Hartree–Fock calculations](#). *Pramana* **30** (1988), no. 5, 387–405
- [3] J. Dolbeault, M. J. Esteban, M. Loss, and L. Vega, [An analytical proof of Hardy-like inequalities related to the Dirac operator](#). *J. Funct. Anal.* **216** (2004), no. 1, 1–21 Zbl 1060.35120 MR 2091354
- [4] J. Dolbeault, M. J. Esteban, and E. Séré, [On the eigenvalues of operators with gaps. Application to Dirac operators](#). *J. Funct. Anal.* **174** (2000), no. 1, 208–226 Zbl 0982.47006 MR 1761368
- [5] J. Dolbeault, M. J. Esteban, and E. Séré, [Variational characterization for eigenvalues of Dirac operators](#). *Calc. Var. Partial Differential Equations* **10** (2000), no. 4, 321–347 Zbl 0968.49025 MR 1767717
- [6] J. Dolbeault, M. J. Esteban, and E. Séré, [A variational method for relativistic computations in atomic and molecular physics](#). *Int. J. Quantum Chem.* **93** (2003), no. 3, 149–155
- [7] J. Dolbeault, M. J. Esteban, and E. Séré, [General results on the eigenvalues of operators with gaps, arising from both ends of the gaps. Application to Dirac operators](#). *J. Eur. Math. Soc. (JEMS)* **8** (2006), no. 2, 243–251 Zbl 1157.47306 MR 2239275
- [8] J. Dolbeault, M. J. Esteban, and E. Séré, [Corrigendum to: “On the eigenvalues of operators with gaps. Application to Dirac operators”](#). *J. Funct. Anal.* **184** (2023), no. 1, article no. 109651 Zbl 0982.47006 MR 4500240
- [9] J. Dolbeault, M. J. Esteban, E. Séré, and M. Vanbreugel, [Minimization methods for the one-particle Dirac equation](#). *Phys. Rev. Lett.* **85** (2000), 4020–4023
- [10] M. J. Esteban, M. Lewin, and É. Séré, [Domains for Dirac–Coulomb min-max levels](#). *Rev. Mat. Iberoam.* **35** (2019), no. 3, 877–924 Zbl 1450.81039 MR 3960263
- [11] M. J. Esteban, M. Lewin, and É. Séré, [Dirac–Coulomb operators with general charge distribution I. Distinguished extension and min-max formulas](#). *Ann. H. Lebesgue* **4** (2021), 1421–1456 Zbl 1486.81094 MR 4353967
- [12] M. J. Esteban and M. Loss, [Self-adjointness for Dirac operators via Hardy–Dirac inequalities](#). *J. Math. Phys.* **48** (2007), no. 11, article no. 112107 Zbl 1152.81423 MR 2370231
- [13] M. J. Esteban and M. Loss, [Self-adjointness via partial Hardy-like inequalities](#). In *Mathematical results in quantum mechanics*, pp. 41–47, World Scientific, Hackensack, NJ, 2008 Zbl 1156.81369 MR 2466677
- [14] M. J. Esteban and E. Séré, [Existence and multiplicity of solutions for linear and non-linear Dirac problems](#). In *Partial differential equations and their applications (Toronto, ON, 1995)*, pp. 107–118, CRM Proc. Lecture Notes 12, American Mathematical Society, Providence, RI, 1997 Zbl 0889.35084 MR 1479240
- [15] M. Griesemer, R. T. Lewis, and H. Siedentop, [A minimax principle for eigenvalues in spectral gaps: Dirac operators with Coulomb potentials](#). *Doc. Math.* **4** (1999), 275–283 Zbl 1115.47300 MR 1701887

- [16] M. Griesemer and H. Siedentop, [A minimax principle for the eigenvalues in spectral gaps](#). *J. London Math. Soc. (2)* **60** (1999), no. 2, 490–500 Zbl [0952.47022](#) MR [1724845](#)
- [17] M. Klaus, [Dirac operators with several Coulomb singularities](#). *Helv. Phys. Acta* **53** (1980), no. 3, 463–482 (1981) MR [611770](#)
- [18] M. Klaus and R. Wüst, [Characterization and uniqueness of distinguished selfadjoint extensions of Dirac operators](#). *Comm. Math. Phys.* **64** (1978/79), no. 2, 171–176 Zbl [0408.47022](#) MR [519923](#)
- [19] M. Kraus, M. Langer, and C. Tretter, [Variational principles and eigenvalue estimates for unbounded block operator matrices and applications](#). *J. Comput. Appl. Math.* **171** (2004), no. 1-2, 311–334 Zbl [1066.49030](#) MR [2077211](#)
- [20] M. Lewin and É. Séré, [Spectral pollution and how to avoid it \(with applications to Dirac and periodic Schrödinger operators\)](#). *Proc. Lond. Math. Soc. (3)* **100** (2010), no. 3, 864–900 Zbl [1192.47016](#) MR [2640293](#)
- [21] S. Morozov and D. Müller, [On the minimax principle for Coulomb–Dirac operators](#). *Math. Z.* **280** (2015), no. 3-4, 733–747 Zbl [1320.49032](#) MR [3369348](#)
- [22] D. Müller, [Minimax principles, Hardy–Dirac inequalities, and operator cores for two and three dimensional Coulomb–Dirac operators](#). *Doc. Math.* **21** (2016), 1151–1169 Zbl [1350.49074](#) MR [3578211](#)
- [23] G. Nenciu, [Self-adjointness and invariance of the essential spectrum for Dirac operators defined as quadratic forms](#). *Comm. Math. Phys.* **48** (1976), no. 3, 235–247 Zbl [0349.47014](#) MR [421456](#)
- [24] G. Nenciu, [Distinguished self-adjoint extension for Dirac operator with potential dominated by multicenter Coulomb potentials](#). *Helv. Phys. Acta* **50** (1977), no. 1, 1–3 MR [462346](#)
- [25] M. Reed and B. Simon, *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness*. Academic Press, New York and London, 1975 Zbl [0308.47002](#) MR [0493420](#)
- [26] L. Schimmer, J. P. Solovej, and S. Tokus, [Friedrichs extension and min-max principle for operators with a gap](#). *Ann. Henri Poincaré* **21** (2020), no. 2, 327–357 Zbl [1432.49065](#) MR [4056270](#)
- [27] U.-W. Schmincke, [Distinguished selfadjoint extensions of Dirac operators](#). *Math. Z.* **129** (1972), 335–349 Zbl [0252.35062](#) MR [326448](#)
- [28] K. Schmüdgen, *Unbounded self-adjoint operators on Hilbert space*. Grad. Texts Math. 265, Springer, Dordrecht, 2012 MR [2953553](#) Zbl [1257.47001](#)
- [29] J. D. Talman, [Minimax principle for the Dirac equation](#). *Phys. Rev. Lett.* **57** (1986), no. 9, 1091–1094 MR [854208](#)
- [30] B. Thaller, *The Dirac equation*. Texts Monogr. Phys., Springer, Berlin, 1992 Zbl [0765.47023](#) MR [1219537](#)
- [31] C. Tix, [Self-adjointness and spectral properties of a pseudo-relativistic Hamiltonian due to Brown and Ravenhall](#). 1997, mp-arc [97-441](#)
- [32] C. Tix, [Strict positivity of a relativistic Hamiltonian due to Brown and Ravenhall](#). *Bull. London Math. Soc.* **30** (1998), no. 3, 283–290 Zbl [0939.35134](#) MR [1608118](#)

- [33] C. Tretter, *Spectral theory of block operator matrices and applications*. Imperial College Press, London, 2008 Zbl 1173.47003 MR 2463978
- [34] R. Wüst, A convergence theorem for selfadjoint operators applicable to Dirac operators with cutoff potentials. *Math. Z.* **131** (1973), 339–349 Zbl 0274.47008 MR 355655
- [35] R. Wüst, Distinguished self-adjoint extensions of Dirac operators constructed by means of cut-off potentials. *Math. Z.* **141** (1975), 93–98 Zbl 0311.47020 MR 365233
- [36] R. Wüst, Dirac operations with strongly singular potentials. Distinguished self-adjoint extensions constructed with a spectral gap theorem and cut-off potentials. *Math. Z.* **152** (1977), no. 3, 259–271 Zbl 0361.35051 MR 437948

Received 23 June 2022; revised 21 March 2023.

Jean Dolbeault

CEREMADE (CNRS UMR 7534), PSL University, Université Paris-Dauphine,
Place du Maréchal de Lattre de Tassigny, 75775 Paris Cedex 16, France;
dolbeaul@ceremade.dauphine.fr

Maria J. Esteban

CEREMADE (CNRS UMR 7534), PSL University, Université Paris-Dauphine,
Place du Maréchal de Lattre de Tassigny, 75775 Paris Cedex 16, France;
esteban@ceremade.dauphine.fr

Eric Séré

CEREMADE (CNRS UMR 7534), PSL University, Université Paris-Dauphine,
Place du Maréchal de Lattre de Tassigny, 75775 Paris Cedex 16, France;
sere@ceremade.dauphine.fr