© 2024 European Mathematical Society Published by EMS Press



Gal Binyamini · Christopher Daw

Effective computations for weakly optimal subvarieties

Received August 24, 2021; revised May 3, 2022

Abstract. Ren and the second author established that the weakly optimal subvarieties (e.g. maximal weakly special subvarieties) of a subvariety V of a Shimura variety arise in finitely many families. In this article, we refine this theorem by (1) constructing a finite collection of algebraic families whose fibres are precisely the weakly optimal subvarieties of V; (2) obtaining effective degree bounds on the weakly optimal locus and its individual members; (3) describing an effective procedure to determine the weakly optimal locus.

Keywords. Unlikely intersections, Zilber–Pink, Shimura varieties, effectivity, differential-algebraic geometry, multiplicity estimates

1. Introduction

This article is concerned with effective results on the (geometric side of the) Zilber–Pink conjecture for (pure) Shimura varieties. The conjecture itself is as follows.

Conjecture 1 (Zilber–Pink). Let *S* be a Shimura variety and let *V* be a Hodge generic, irreducible, algebraic subvariety of *S*. Then the intersection of *V* with the union of the special subvarieties of *S* of codimension at least dim V + 1 is not Zariski dense in *V*.

By [2, Theorem 12.4], Conjecture 1 is equivalent to the seemingly stronger variations involving atypical intersections and optimal subvarieties, the latter of which states that V contains only finitely many optimal subvarieties. We state this version and the necessary definitions in Section 3.3.

In [9], Ren and the second author outline a Pila–Zannier strategy for proving Conjecture 1, applying the familiar combination of o-minimality, functional transcendence, and arithmetic (see, in particular, [15]). The unconditional aspect of that strategy is what might be considered the geometric Zilber–Pink conjecture for Shimura varieties. Gao [12] refers to its generalization to mixed Shimura varieties as *a finiteness result à la Bogomolov*. The

Mathematics Subject Classification (2020): Primary 14G35; Secondary 11G18, 14Q20

Gal Binyamini: Department of Mathematics, Weizmann Institute of Science, 7610001 Rehovot, Israel; gal.binyamini@weizmann.ac.il

Christopher Daw: Department of Mathematics and Statistics, University of Reading, RG6 6AH Reading, UK; chris.daw@reading.ac.uk

statement is as follows. We refer to Section 3.4 for the relevant facts and definitions, recalling here only that an optimal subvariety is, in particular, a weakly optimal subvariety.

Theorem 2 ([9, Proposition 6.3]). Let $S = \Gamma \setminus X$ be a Shimura variety and let V be an irreducible algebraic subvariety of S. There exists a finite set Σ of pairs $(X_{\mathbf{H}}, X_1 \times X_2)$ with $X_{\mathbf{H}}$ a pre-special subvariety of X and $X_{\mathbf{H}} = X_1 \times X_2$ a \mathbb{Q} -splitting of $X_{\mathbf{H}}$ such that, for any weakly optimal subvariety W of V, the weakly special closure of W in S is equal to the image in S of $X_1 \times \{x_2\}$ for some $(X_{\mathbf{H}}, X_1 \times X_2) \in \Sigma$ and some $x_2 \in X_2$.

In this article, we give a refined version of this theorem (see Theorem 19). We summarize this refinement as follows. We denote by P the so-called *standard principal bundle* associated with S and, by Ω , a Chow variety parametrizing certain subvarieties of the compact dual \check{X} of X (see Sections 4.1, 5.1, and 6 for the details). Similarly, for a triple $T = (X_{\rm H}, X_1 \times X_2)$ as above, we denote by P_T the standard principal bundle associated with a Shimura variety S_T corresponding to $X_{\rm H}$, and by Ω_T a Chow subvariety contained in Ω parametrizing certain subvarieties of $\check{X}_{\rm H}$. As such, we obtain algebraic morphisms

$$\begin{array}{cccc} P_T \times \Omega_T & \stackrel{\iota_{P_T}}{\longrightarrow} & P \times \Omega \\ & & & \downarrow^{\pi_T} & & \downarrow \\ S_2 & \xleftarrow{\phi_T} & S_T & \longrightarrow & S \end{array}$$

where S_2 is a Shimura variety corresponding to X_2 . For a subset Π of $P \times \Omega$, we denote by Π_T the union of the Zariski closures of the fibres of the map ϕ_T restricted to $\pi_T(\iota_{P_T}^{-1}(\Pi))$.

Theorem 3. Let *S* be a Shimura variety and let *V* be an irreducible algebraic subvariety of *S*. For every $d \in \mathbb{N}$, there exists a locally closed algebraic subset $\Pi(d)$ of $P \times \Omega$ and, for each of the irreducible components $\Pi(d)^{\circ}$ of $\Pi(d)$, an associated triple $T = (X_{\mathbf{H}}, X_1 \times X_2)$ such that the following holds:

- (i) Let Π(d)° be an irreducible component of Π(d) and let T be its associated triple. Then Π(d)[°]_T is constructible.
- (ii) Let W be a weakly optimal subvariety of V of weakly special defect d. Then there exists an irreducible component $\Pi(d)^{\circ}$ of $\Pi(d)$ (with its associated triple T) such that W is the image in S of some irreducible component of a fibre of $\phi_T|_{\Pi(d)_T^{\circ}}$.
- (iii) Let $\Pi(d)^{\circ}$ denote an irreducible component of $\Pi(d)$ (with its associated triple T). If W is an irreducible component of a fibre of $\phi_T|_{\Pi(d)_T^{\circ}}$, then the image in S of W is a weakly optimal subvariety of V of weakly special defect d.

In short, we show that the weakly optimal subvarieties of V are precisely the fibres of finitely many constructible families.

However, our construction allows us to go further, applying estimates from differential algebraic geometry to obtain effective upper bounds for the degrees of the families $\Pi(d)$, as well as the individual weakly optimal subvarieties. Moreover, our construction also

produces an effective description of this *unlikely intersection* locus, in the form of an explicit system of algebraic equations.

To formulate such a result explicitly, we fix a coordinate system on the set $P \times \Omega$ as follows. Let $U \subset S$ denote an open dense subset such that P trivializes over U (that is, if **G** is the algebraic group over \mathbb{Q} associated with S, then $P|_U \simeq U \times \mathbf{G}(\mathbb{C})$). Fixing a faithful representation of **G** we may realize it as a subvariety of some affine space. We use (1) a very-ample power of the Baily–Borel line bundle for a set of projective coordinates on U; (2) the matrix entries as coordinates on $\mathbf{G}(\mathbb{C})$; (3) the standard Chow coordinates on the Chow variety Ω . All degrees on $U \times \mathbf{G}(\mathbb{C}) \times \Omega$ below are taken with respect to the Segre product of these three coordinate systems.

If X is a locally closed subset of $U \times \mathbf{G}(\mathbb{C}) \times \Omega$, then it is of the form $Y \setminus Z$, where Y and Z are Zariski closed. We define the *complexity* of X to be deg(Y) + deg(Z). We give a bound for the complexity of the sets $\Pi(d)$, as well as a procedure for computing the equations and inequations defining $\Pi(d)$, in terms of the equations defining V. The construction depends on certain data associated with S, namely, (1) the equations describing the projective embedding of $U \times \mathbf{G}(\mathbb{C}) \times \Omega$ and (2) the equations describing the canonical connection ∇ on P (written in the chosen coordinate system on $U \times \mathbf{G}(\mathbb{C})$). Throughout the paper, with the exception of Section 9, our effective bounds are assumed to depend on this data (we simply say *depending only on S*). In the final Section 9, we address the issue of obtaining fully effective computations in the case $S = \mathcal{A}_g$ using Gauss–Manin connections.

Theorem 4. The complexity of $\Pi(d)$ is bounded by $f_S(\deg(V))$ for some polynomial f_S depending only on S. Moreover, we produce an explicit system of equations and inequations for $\Pi(d)$ of degree bounded by $f_S(\deg(V))$.

According to Theorem 3, the sets $\Pi(d)$ provide a complete parametrization for all families of weakly optimal subvarieties of V of weakly special defect d. Theorem 4 therefore provides a method for explicitly computing these families and controlling their degrees. From this, we obtain degree bounds on individual weakly optimal subvarieties.

Theorem 5. Let $d \in \mathbb{N}$ and let W be a weakly optimal subvariety of V of weakly special defect d. Then

$$\deg(W) \le f_S(\deg(V))$$

for some polynomial f_S depending only on S.

1.1. Fully effective computation

Even though our main construction is effective, to carry out this procedure in practice one would need to obtain

- (1) an explicit system of equations for the projective embedding of *S* with respect to (some power of the) Baily–Borel line bundle;
- (2) an explicit description of the canonical connection ∇ with respect to a prescribed trivialization of the corresponding bundle *P*;

(3) an explicit system of equations for the subvariety V with respect to the projective embedding.

Computing the first two of these items for a given Shimura variety S appears to be a non-trivial task. Moreover, computing equations for some subvariety V of interest is by itself a non-trivial problem. For example, consider the case $S = \mathcal{A}_g$, the space of principally polarized abelian varieties of dimension g, and V the closure \mathcal{T}_g of the open Torelli locus $\mathcal{T}_g^{\circ} \subset \mathcal{A}_g$. Computing the weakly special locus is a problem of significant interest due to its relation to the Coleman–Oort conjecture on the finiteness of the set of isomorphism classes of genus-g Jacobians with complex multiplication. We refer to [24] for an excellent survey and simply recall here that it is conjectured that for $g \gg 1$ the set of positive-dimensional special subvarieties of \mathcal{T}_g intersecting \mathcal{T}_g° is empty. In combination with the André–Oort conjecture for \mathcal{A}_g (established by Tsimerman [27]), this would imply the Coleman–Oort conjecture. Note, however, that computing a set of equations for V in this case is the famous Schottky problem, a subject of substantial independent study, still not fully resolved.

Fortunately, taking advantage of the functoriality of the canonical connections on Shimura varieties, we are able to alternatively carry out the computation for $V \subset S$ described via a moduli interpretation instead of an explicit projective embedding. We focus our attention on the case $S = A_g$. In light of the moduli interpretation of A_g , it is natural to describe a subvariety $V \subset A_g$ as a family of genus-g principally polarized abelian varieties, or the Jacobians of a family of genus-g curves. In this case, the computation of the canonical connection translates into the computation of the Gauss–Manin connection for the corresponding family. This is a classical problem and it is well known, going back to the work of Manin [19], that it can be carried out explicitly.

We will focus here on the case of families of Jacobians, which lends itself more readily to effective computation, as curves are relatively simple to describe using explicit equations. This is also the case required in principle to treat the Coleman–Oort conjecture and related problems on the Torelli locus. Our methods could in principle also be used to carry out explicit computations with more general constructions, such as Prym varieties, or even with general families of polarized abelian varieties presented by explicit equations. However, since it is far less common to present general abelian varieties in this way, we do not pursue this matter explicitly.

Let V denote an algebraic variety, which we assume for simplicity to be smooth. Let $T \rightarrow V$ denote a smooth curve over V, by which we mean that T is smooth and the map $T \rightarrow V$ is submersive.

To coincide with our general formalism, we should work with a neat subgroup $\Gamma \subset \mathbf{GSp}_{2g}(\mathbb{Z})$. We therefore denote by $f: \tilde{V} \to V$ an étale cover, $\tilde{T} \to \tilde{V}$ the base change of T by f, and choose f such that \tilde{T} is compatible with an N-level structure (say for N = 3). We denote by $\iota: \tilde{V} \to \mathcal{A}_{g,N}$ the corresponding moduli map.

Theorem 6. Given an explicit system of equations for (a projective embedding of) the family $T \to V$, one can explicitly construct an affine cover $\{V_{\alpha} \subset V\}$ such that for each $V' = V_{\alpha}$:

(1) $\iota^* P|_{f^{-1}(V')} \simeq V' \times \mathbf{GSp}(\mathbb{C}),$

(2) for each d, one can explicitly construct a system of algebraic equations and inequations for sets $\Pi'(d) \subset V' \times \mathbf{GSp}(\mathbb{C})$ such that $f^*\Pi'(d) = \iota^*\Pi(d)|_{f^{-1}(V')}$.

Theorem 6 in principle provides an effective method for deciding whether weakly special subvarieties exist within the open Torelli locus \mathcal{T}_g° for any given g. One first has to explicitly write down equations for a family $T \to V$ parametrizing all genus-g curves. One then computes explicitly the equations for the defect-zero set $\iota^*\Pi(0)$ corresponding to families of weakly special subvarieties contained in V. Finally, one may use effective commutative algebra methods (for instance, Gröbner basis algorithms) to determine whether the family is empty.

Remark 7. In the particular case of checking for the existence of weakly special subvarieties, the smoothness assumption on V can be dropped. One can first effectively construct a smooth open dense subset $V' \subset V$ and apply the preceding process to V', and then proceed by induction on dimension with $V \setminus V'$.

One may apply a similar process to look for weakly special or weakly optimal subvarieties of other families of interest, for instance, the family of (Jacobians of) hyperelliptic curves of a given genus g. We stress that, to perform this computation using Theorem 4, one would need to obtain an explicit description of the hyperelliptic locus $V \subset A_g$, which is a non-trivial problem.

1.2. Literature review

Effective results on the André–Oort and the Zilber–Pink conjectures are still relatively sparse. This work improves upon a previous work of the second author with Javanpeykar and Kühne [7] which, by entirely different methods, gave effective degree bounds for so-called *non-facteur* maximal special subvarieties. We refer to the introduction of the latter for references to several earlier works. More recently, the first author has obtained effective results on the André–Oort conjecture for products of modular curves [3] and, with Masser, for Hilbert modular varieties [4]. In May 2021, Pila and Scanlon announced effective results for function field versions of the Zilber–Pink conjectures for varieties supporting a variation of Hodge structures, also using differential-algebraic methods [25]. In September 2021, Urbanik released an algorithm capable of computing the set of all weakly special subvarieties of degree at most d of any smooth quasi-projective algebraic variety equipped with a polarizable variation of Hodge structures [30].

2. Generalities

2.1. Analytic and algebraic sets

For a complex analytic set X and $x \in X$, we write $\dim_x X$ for the dimension of X at x, as defined in [13]. We write dim X for the supremum of $\dim_x X$ for all $x \in X$. We recall

that, if X is irreducible, $\dim_X X$ is constant on X. If X is (explicitly) an algebraic variety, then dim X will refer to its dimension as an algebraic variety. All algebraic subvarieties are assumed to be (Zariski) closed, unless stated otherwise.

2.2. Degrees

If X is a complex algebraic variety and $k \in \mathbb{N}$, we denote by $A_k X$ the group of k-cycles modulo rational equivalence on X (see [10, Section 1.3]). We define the degree deg(α) of $\alpha \in A_k X$ as in [10, Definition 1.4]. In particular, deg(α) = 0 if k > 0. If $\alpha \in A_k X$ and L is a line bundle on X, we obtain, for any positive integer $d \leq k$, a class

$$c_1(L)^d \cap \alpha \in A_{k-d}X$$

(see [10, Definition 2.5]). If V is an irreducible subvariety of X, we define the degree $\deg_L(V)$ of V with respect to L to be the degree of

$$c_1(L)^{\dim V} \cap [V] \in A_0 X$$

where $[V] \in A_{\dim V} X$ denotes the class of the dim V-cycle given by V. If V is a (not necessarily irreducible) subvariety of X, we define the deg_L(V) to be the sum of the deg_L(V_i) as V_i varies over the irreducible components of V.

2.3. Algebraic groups

For an algebraic group G, we denote by G° the connected component (in the Zariski topology) of G containing the identity, and we denote by the corresponding mathfrak letter g its Lie algebra.

We include "connected" in our definitions of reductive and semisimple algebraic groups. For a reductive algebraic group **G**, we denote by \mathbf{G}^{ad} the quotient of **G** by its centre $\mathbf{Z}(\mathbf{G})$, and we denote by \mathbf{G}^{der} the derived subgroup of **G**. If **G** is defined over (a field containing) \mathbb{R} , we denote the connected component (in the archimedean topology) $\mathbf{G}(\mathbb{R})^+$ of $\mathbf{G}(\mathbb{R})$ containing the identity by the italic letter *G*, retaining any superscripts or subscripts, and we write $\mathbf{G}(\mathbb{R})_+$ for the inverse image of G^{ad} under the natural map $\mathbf{G}(\mathbb{R}) \to \mathbf{G}^{ad}(\mathbb{R})$. We write $\mathbf{G}(\mathbb{Q})_+$ for $\mathbf{G}(\mathbb{R})_+ \cap \mathbf{G}(\mathbb{Q})$.

If **G** is a reductive algebraic group over a field of characteristic zero, and **H** is a reductive algebraic subgroup of **G**, then we write $N_G(H)$ (resp. $Z_G(H)$) for the normalizer (resp. the centralizer) of **H** in **G**. We recall that $N_G(H)^\circ$ and $Z_G(H)^\circ$ are both reductive. We have an almost direct product decomposition $N_G(H)^\circ = H \cdot Z_G(H)^\circ$.

2.4. Arithmetic groups

Let **G** denote a reductive algebraic group over \mathbb{Q} and, via a faithful representation, consider **G** as a subgroup of **GL**_n for some $n \in \mathbb{N}$. The definitions that follow are independent of this representation, and hence we can and do make use of them without reference to such a representation.

An *arithmetic subgroup* of $G(\mathbb{Q})$ is any subgroup commensurable with $G(\mathbb{Q}) \cap$ $GL_n(\mathbb{Z})$. An element of $G(\mathbb{Q})$ is *neat* if the subgroup of $\overline{\mathbb{Q}}$ generated by its eigenvalues (considering it as an element of $GL_n(\overline{\mathbb{Q}})$) is torsion free. A subgroup of $G(\mathbb{Q})$ is neat if all of its elements are neat. In particular, a neat subgroup is torsion free.

3. Shimura varieties and the Zilber-Pink conjecture

3.1. Shimura data

Let S denote the Deligne torus (that is, the Weil restriction from \mathbb{C} to \mathbb{R} of \mathbb{G}_m). By a *Shimura datum*, we refer to a pair (**G**, **X**), where **G** is a reductive algebraic group defined over \mathbb{Q} and **X** is a **G**(\mathbb{R})-conjugacy class of morphisms $\mathbb{S} \to \mathbf{G}_{\mathbb{R}}$ such that the conditions SV1, SV2, and SV3 of [21, p. 50] hold. Furthermore, we impose the condition

(SV0) **G** is the generic Mumford–Tate group on **X**.

Condition (SV0) means that **G** is the smallest algebraic subgroup **H** of **G** defined over \mathbb{Q} such that $x(\mathbb{S})$ is contained in $\mathbf{H}_{\mathbb{R}}$ for all $x \in \mathbf{X}$. We recall that **X** is naturally a disjoint union of hermitian symmetric domains. We refer the reader to [21] for more details regarding the theory of Shimura varieties.

In this article, in order to simplify technical issues, we will assume that our ambient Shimura datum (G, X) satisfies the condition that $Z(G)(\mathbb{R})$ is compact.

3.2. Shimura varieties

Let (\mathbf{G}, \mathbf{X}) be a Shimura datum such that $\mathbf{Z}(\mathbf{G})(\mathbb{R})$ is compact and let X be a connected component of \mathbf{X} . As in [8], we refer to the pair (\mathbf{G}, X) as a *Shimura datum component*. Let Γ be an arithmetic subgroup of $\mathbf{G}(\mathbb{Q})$ contained in $\mathbf{G}(\mathbb{Q})_+$. Then Γ acts on X and, by the theorem of Baily–Borel [1], the quotient $S = \Gamma \setminus X$ naturally possesses the structure of an irreducible quasi-projective complex algebraic variety. Indeed, by [1, Lemma 10.8], the line bundle of holomorphic forms of maximal degree on X descends to an ample line bundle L_{Γ} on S. Note that, if Γ is neat, then S is non-singular (see [26, Facts 2.3]). We let k_{Γ} denote the smallest integer such that $L_{\Gamma}^{\otimes k_{\Gamma}}$ is very ample.

We will refer to the irreducible variety *S* as the Shimura variety *associated with* (**G**, *X*) and Γ . We will denote by π the natural complex analytic map $X \to S$.

3.3. Special subvarieties and the Zilber–Pink conjecture

Recall the situation described in Section 3.2. Let $(\mathbf{H}, \mathbf{X}_{\mathbf{H}})$ denote a Shimura subdatum of (\mathbf{G}, \mathbf{X}) (in particular, \mathbf{H} is the generic Mumford–Tate group on $\mathbf{X}_{\mathbf{H}}$), and let $X_{\mathbf{H}}$ denote a connected component of $\mathbf{X}_{\mathbf{H}}$ contained in X. For any arithmetic subgroup $\Gamma_{\mathbf{H}}$ of $\mathbf{H}(\mathbb{Q})$ contained in $\mathbf{H}(\mathbb{Q})_+$, we obtain a Shimura variety $S_{\mathbf{H}} = \Gamma_{\mathbf{H}} \setminus X_{\mathbf{H}}$ and, when $\Gamma_{\mathbf{H}}$ is contained in Γ , the natural complex analytic map $\Gamma_{\mathbf{H}} \setminus X_{\mathbf{H}} \to \Gamma \setminus X$ is a finite (hence closed) morphism of algebraic varieties (see [26, Facts 2.6]). We refer to the image of any such map as a *special subvariety* of S.

It is straightforward to show that the intersection of two special subvarieties is a finite union of special subvarieties. In particular, for any subvariety W of S, there exists a smallest special subvariety $\langle W \rangle$ of S containing W. In light of this, we define the *defect* $\delta(W)$ of W by

$$\delta(W) = \dim \langle W \rangle - \dim W.$$

Now fix a subvariety V of S. We define an irreducible subvariety W of V to be *optimal* in V if, whenever $W \subsetneq Y$ for some other irreducible subvariety Y of V, we have $\delta(W) < \delta(Y)$. (In particular, V is an optimal subvariety of V itself.)

Note that an optimal subvariety W of V such that $\delta(W) = 0$ is a maximal special subvariety of V. Observe also that an optimal subvariety W of V is necessarily an irreducible component of $\langle W \rangle \cap V$.

We denote by Opt(V) the set of optimal subvarieties of V. The central problem in the area of *unlikely intersections* in Shimura varieties is (equivalent to) the following (see [9]).

Conjecture 8 (Zilber–Pink). Let V be a subvariety of a Shimura variety S. Then the set Opt(V) is finite.

3.4. Weakly special and weakly optimal subvarieties

Recall the situation described in Section 3.2. Let $(\mathbf{H}, \mathbf{X}_{\mathbf{H}})$ denote a Shimura subdatum of (\mathbf{G}, \mathbf{X}) and let $X_{\mathbf{H}}$ denote a connected component of $\mathbf{X}_{\mathbf{H}}$ contained in X. Then the image $S_{\mathbf{H}}$ of $X_{\mathbf{H}}$ in S is a special subvariety. Decompose \mathbf{H}^{ad} as a product $\mathbf{H}_1 \times \mathbf{H}_2$ of two (permissibly trivial) normal Q-subgroups. In this way, we obtain a decomposition $X_{\mathbf{H}} = X_1 \times X_2$, and we will refer to a decomposition of this form as a Q-*splitting*. For any $x_2 \in X_2$, the image $S_{\mathbf{H},x_2}$ of $X_1 \times \{x_2\}$ in S is a closed irreducible algebraic subvariety and we refer to any subvariety of this form as a *weakly special* subvariety of S. We remark that any special subvariety is weakly special, whereas $S_{\mathbf{H},x_2}$ is special if and only if the Mumford–Tate group of x_2 is a torus (or, equivalently, if x_2 is a pre-special point of X_2 , to use another terminology).

It is straightforward to show that the intersection of two weakly special subvarieties is a finite union of weakly special subvarieties. In particular, for any subvariety W of S, there exists a smallest weakly special subvariety $\langle W \rangle_{ws}$ of S containing W. In light of this, we define the *weakly special defect* $\delta_{ws}(W)$ of W by

$$\delta_{ws}(W) = \dim \langle W \rangle_{ws} - \dim W.$$

Now fix a subvariety V of S. We define an irreducible subvariety W of V to be *weakly* optimal in V if, whenever $W \subsetneq Y$ for some other irreducible subvariety Y of V, we have $\delta_{ws}(W) < \delta_{ws}(Y)$. (In particular, V is a weakly optimal subvariety of V itself.)

Note that a weakly optimal subvariety W of V such that $\delta_{ws}(W) = 0$ is a maximal weakly special subvariety of V. Observe also that a weakly optimal subvariety W of V is necessarily an irreducible component of $\langle W \rangle_{ws} \cap V$. By [9, Corollary 4.5], any optimal subvariety of V is weakly optimal.

Remark 9. Note that any point $z \in S$ is a weakly special subvariety (according to our convention, at least). In particular, $\delta_{ws}(z) = 0$. Hence, $z \in V$ is a weakly optimal subvariety of V if and only if it is not contained in a positive-dimensional weakly special subvariety contained in V.

4. The (weak) hyperbolic Ax–Schanuel conjecture

4.1. Subvarieties of X

Recall the situation described in Section 3.1 and let *X* be an irreducible component of **X**. Recall that *X* is a *G*-conjugacy class of homomorphisms $\mathbb{S} \to \mathbf{G}_{\mathbb{R}}$. By extending scalars to \mathbb{C} and pre-composing with $\mathbb{G}_m \to \mathbb{G}_m^2 \cong \mathbb{S}_{\mathbb{C}}$, where the first map is given by $z \mapsto (z, 1)$, we obtain from each point $x \in X$ a cocharacter $\mu_x : \mathbb{G}_m \to \mathbf{G}_{\mathbb{C}}$ such that

(*) in the action of \mathbb{G}_m on Lie($\mathbf{G}_{\mathbb{C}}$), obtained via restriction of the adjoint representation, only the characters z, 1, and z^{-1} occur.

In this way, we obtain an embedding of X into a $\mathbf{G}(\mathbb{C})$ -conjugacy class X^{co} of cocharacters of $\mathbf{G}_{\mathbb{C}}$ satisfying (*). For each $\mu \in X^{co}$ and $r \in \{1, 0, -1\}$, we define V_{μ}^{r} to be the character subspace of Lie($\mathbf{G}_{\mathbb{C}}$) on which \mathbb{G}_{m} acts (according to the action obtained from μ) via the character z^{r} . Then Lie($\mathbf{G}_{\mathbb{C}}$) = $\bigoplus_{r} V_{\mu}^{r}$, and we obtain a filtration F_{μ} of Lie($\mathbf{G}_{\mathbb{C}}$) by setting $F_{\mu}^{p} = \bigoplus_{r \geq p} V_{\mu}^{r}$. In this way, we obtain a $\mathbf{G}(\mathbb{C})$ -invariant surjective map $\mu \mapsto F_{\mu}$ from X^{co} to a $\mathbf{G}(\mathbb{C})$ -orbit of filtrations \check{X} . Note that \check{X} is a complex projective flag variety known as the *compact dual* of X. The composite map $X \to \check{X}$ is a complex analytic $\mathbf{G}(\mathbb{R})_{+}$ -invariant embedding, known as the *Borel embeddding* of X, and we identify X with its image, which is an open subset of \check{X} . In particular, dim $\check{X} = \dim X$. We define a subvariety of X to be any irreducible analytic component of $X \cap Y$ for any algebraic subvariety Y of \check{X} . As noted in [9, paragraph following Theorem 5.4], this definition agrees with the definition therein.

4.2. Pre-special and pre-weakly special subvarieties

Recall the situation described in Section 3.1 and let X be an irreducible component of X. If $(\mathbf{H}, \mathbf{X}_{\mathbf{H}})$ is a Shimura subdatum of (\mathbf{G}, \mathbf{X}) and $X_{\mathbf{H}}$ is a connected component of $\mathbf{X}_{\mathbf{H}}$ contained in X, we obtain a commutative diagram of complex analytic $\mathbf{H}(\mathbb{R})_+$ -invariant embeddings



and we identify all objects with their images in X.

Lemma 10. We have $X \cap \check{X}_{\mathbf{H}} = X_{\mathbf{H}}$.

Proof. By [14, Proposition VI.B.11], the intersection is contained in $X \cap \mathbf{X_H}$. Therefore, let $x_1, x_2 \in X \cap \mathbf{X_H}$ and let X_i denote the *H*-orbit of x_i in *X* (in other words, the connected component of $\mathbf{X_H}$ containing x_i). Let K_i denote the maximal compact subgroup of *G* stabilizing x_i and let $G = P_i K_i$ denote the corresponding Cartan decomposition. We also have Cartan decompositions $H = (P_i \cap H)(K_i \cap H)$.

Writing $x_2 = \alpha x_1$ for some $\alpha \in G$, we have $K_2 = \alpha K_1 \alpha^{-1}$ and $P_2 = \alpha P_1 \alpha^{-1}$. Since Cartan decompositions are conjugate, we also have $K_2 \cap H = h(K_1 \cap H)h^{-1}$ and $P_2 \cap H = h(P_1 \cap H)h^{-1}$ for some $h \in H$.

We set $\gamma = h^{-1}\alpha$ and write $\gamma = pk$ for some $p \in P_1$ and some $k \in K_1$. We deduce that $K_1 \cap H = pK_1p^{-1} \cap H$ and (trivially) $P_1 \cap H = pP_1p^{-1} \cap H$. Using [28, Lemme 3.11], as in the proof of [29, Lemma 3.7] we deduce that p^2 centralizes H. Since H is Zariski dense in $\mathbf{H}_{\mathbb{R}}$, it follows that $p^2 \in \mathbf{Z}_{\mathbf{G}}(\mathbf{H})(\mathbb{R})$. Therefore, since $p^2 \in G$, we conclude that $p^2 \in K_1$ and so p^2 is trivial. It follows that p is fixed by the Cartan involution associated with K_1 and so p = 1. We conclude that $x_2 = hx_1$ and so $x_2 \in X_1$, which finishes the proof.

In particular, $X_{\mathbf{H}}$ is a subvariety of X, and we call such a subvariety a *pre-special* subvariety. A similar discussion shows that, for a \mathbb{Q} -splitting $X_{\mathbf{H}} = X_1 \times X_2$ as above and a point $x_2 \in X_2$, the set $X_{\mathbf{H},x_2} = X_1 \times \{x_2\}$ is again a subvariety of X (we define $\check{X}_{\mathbf{H},x_2}$ analogously), and we refer to such a subvariety as a *pre-weakly special* subvariety of X.

Remark 11. It is an easy consequence of [29, Lemma 3.7] and the fact that $G_{\mathbb{R}}$ possesses only finitely many $G(\mathbb{R})$ -conjugacy classes of semisimple subgroups that the pre-weakly special subvarieties of X belong to finitely many $G(\mathbb{R})$ -orbits. It follows that, for a given embedding of \check{X} into projective space, there exists a $D \in \mathbb{N}$ such that, for any pre-weakly special subvariety $X_{\mathbf{H},x_2}$ of X, the degree of $\check{X}_{\mathbf{H},x_2}$ is at most D.

4.3. Intersection components

Recall the situation described in Section 3.2 and let *V* be an irreducible subvariety of *S*. We define an intersection component of $\pi^{-1}(V)$ to be an irreducible analytic component of the intersection of $\pi^{-1}(V)$ with a subvariety of *X*. For any intersection component *A* of $\pi^{-1}(V)$, there exists a smallest subvariety $\langle A \rangle_{Zar}$ of *X* containing *A* (from which it follows that *A* is automatically an irreducible analytic component of $\langle A \rangle_{Zar} \cap \pi^{-1}(V)$). In light of this, we define the *Zariski defect* $\delta_{Zar}(A)$ of *A* by

$$\delta_{\operatorname{Zar}}(A) = \dim \langle A \rangle_{\operatorname{Zar}} - \dim A.$$

We say that *A* is *Zariski optimal* in $\pi^{-1}(V)$ if, whenever $A \subsetneq B$ for some other intersection component *B* of $\pi^{-1}(V)$, we have $\delta_{Zar}(A) < \delta_{Zar}(B)$. The weak hyperbolic Ax–Schanuel conjecture, which follows (see [9, Lemma 5.16]) from the hyperbolic Ax–Schanuel conjecture, proven by Mok–Pila–Tsimerman [23], is the following.

Theorem 12 (weak hyperbolic Ax–Schanuel). Let A be a Zariski optimal intersection component of $\pi^{-1}(V)$. Then $\langle A \rangle_{Zar}$ is pre-weakly special.

5. Standard principal bundles and canonical foliations

5.1. Standard principal bundles

Recall the situation described in Section 3.2 and suppose that Γ is neat. Since $\mathbf{Z}(\mathbf{G})(\mathbb{R})$ is compact, the stabilizer in $\mathbf{G}(\mathbb{R})$ of any point in X is compact. Therefore, since Γ is torsion free, it acts without fixed points on X. It follows that Γ is (isomorphic to) the fundamental group $\pi_1(S)$ of S and that

$$P = \Gamma \setminus (X \times \mathbf{G}(\mathbb{C}))$$

is a principal complex analytic $G(\mathbb{C})$ -bundle over *S*. (The action of Γ on $X \times G(\mathbb{C})$ is diagonal and on the left, and the action of $G(\mathbb{C})$ is given by $h \cdot [x, g] = [x, gh^{-1}]$, where we use [x, g] to denote the class of $(x, g) \in X \times G(\mathbb{C})$ in *P*.) Furthermore, there is a canonical flat connection ∇ on *P*.

Following convention, we refer to $P = (P, \nabla)$ as the *standard principal bundle* associated with (**G**, *X*) and Γ . By [20, Proposition 3.2], *P* is complex algebraic as a bundle over the algebraic variety *S*. We let $\pi_P : P \to S$ denote the natural (complex algebraic) morphism $[x, g] \mapsto \pi(x)$.

Note that there is also a natural $\mathbf{G}(\mathbb{C})$ -equivariant algebraic map $\beta: P \to \check{X}$ defined by

$$\beta([x,g]) = g^{-1}F_{\mu_x}.$$

We observe that the composite of β with the natural embedding $X \to P$ given by $x \mapsto [x, 1]$ yields the Borel embedding of X into \check{X} .

5.2. Trivializations

Recall the situation described in Section 5.1. Let $p = [x, g] \in P$ and let $U \subset X$ be an open neighbourhood of x such that

$$\gamma \in \Gamma$$
 and $U \cap \gamma U \neq \emptyset \implies \gamma = 1$.

Such a U exists by [22, Proposition 2.5] and the fact that Γ is torsion free. It follows immediately that $\pi_U : U \to \pi(U)$ is biholomorphic and we obtain a (complex analytic) trivialization

$$\varphi_U : \pi(U) \times \mathbf{G}(\mathbb{C}) \to \pi_P^{-1}(\pi(U))$$

of *P* over $\pi(U)$ defined by $(s, g) \mapsto [\pi_U^{-1}(s), g]$.

5.3. Flat structures

Recall again the situation described in Section 5.1. Choose an open covering \mathcal{C} of X, stable under translation by Γ and such that each point $x \in X$ is contained in an arbitrarily small $U \in \mathcal{C}$. We claim that we can choose \mathcal{C} such that

$$U_1, U_2 \in \mathcal{C} \text{ and } U_1 \cap U_2 \neq \emptyset \implies U_1 \cap \gamma U_2 = \emptyset \text{ for all } \gamma \in \Gamma \setminus \{1\}.$$
 (1)

To see this, equip X with its usual metric $d : X \times X \to \mathbb{R}$, and consider $f : X \times \mathbf{G}(\mathbb{R})_+ \to X \times X$ defined by $(x, g) \mapsto (x, gx)$. Recall that both d and f are proper maps. In particular, for a compact interval $I \subset \mathbb{R}$, the preimage $(d \circ f)^{-1}(I)$ is a compact subset of $X \times \mathbf{G}(\mathbb{R})_+$ and, therefore, so too is its projection Θ to $\mathbf{G}(\mathbb{R})_+$. Since Γ is a discrete subgroup of $\mathbf{G}(\mathbb{R})_+$, we conclude that $\Gamma \cap \Theta$ is finite. Therefore, since Γ acts freely on X, there exists C > 0 such that

$$\gamma \neq 1 \implies d(x, \gamma x) > C$$
 for all $x \in X$.

Therefore, in order to define \mathcal{C} , choose around each point $x \in X$ a system of arbitraily small relatively compact open neighbourhoods U such that

$$x, y \in U \implies d(x, y) < C/2,$$

and such that \mathcal{C} is stable under translation by Γ . Now let $U_1, U_2 \in \mathcal{C}$ and suppose that $x_1 \in U_1 \cap U_2$. Furthermore, suppose that $x_2 \in U_1 \cap \gamma U_2$ for some $\gamma \in \Gamma \setminus \{1\}$. We conclude that

$$C < d(x_1, \gamma x_1) \le d(x_1, x_2) + d(x_2, \gamma x_1) < C/2 + C/2 = C,$$

which is a contradiction, yielding the claim.

Note that condition (1) with $U_2 = U_1 = U$ implies that we have trivializations

$$\varphi_U : \pi(U) \times \mathbf{G}(\mathbb{C}) \to \pi_P^{-1}(\pi(U))$$

as before, for any $U \in \mathcal{C}$. Now suppose that $\pi(U_1) \cap \pi(U_2) \neq \emptyset$ for $U_1, U_2 \in \mathcal{C}$. We obtain a transition map

$$\varphi_{U_2}^{-1} \circ \varphi_{U_1} : \pi(U_1) \cap \pi(U_2) \times \mathbf{G}(\mathbb{C}) \to \pi(U_1) \cap \pi(U_2) \times \mathbf{G}(\mathbb{C}),$$

which sends (s, g) to $(s, \gamma_s^{-1}g)$, where $\gamma_s \in \Gamma$ is the unique element such that

$$\pi_{U_1}^{-1}(s) = \gamma_s \pi_{U_2}^{-1}(s) \in U_1 \cap \gamma_s U_2.$$

However, by (1), $\gamma = \gamma_s$ is constant on $\pi(U_1) \cap \pi(U_2)$.

We refer to a covering \mathcal{C} of X satisfying the properties above as a *flat structure* for P. In particular, a flat structure \mathcal{C} comes with an associated set $\{\varphi_U\}_{U \in \mathcal{C}}$ of trivializations. Note that, if \mathcal{C} and \mathcal{C}' are both flat structures for P, then $\mathcal{C} \cap \mathcal{C}'$ (whose members are precisely those of the form $U \cap U'$ for $U \in \mathcal{C}$ and $U' \in \mathcal{C}'$) is also a flat structure for P.

5.4. The canonical foliation

Recall again the situation described in Section 5.1. By [18, Section 1B], we obtain a canonical foliation \mathcal{F} of P. For any $p \in P$, we can obtain the leaf \mathcal{L}_p of \mathcal{F} through p as follows. Let \mathcal{C} be a flat structure for P, write $p = [x, g_p]$, and let $U \in \mathcal{C}$ be such that $x \in U$. Then

$$\varphi_U^{-1}(\mathcal{L}_p \cap \pi_P^{-1}(\pi(U))) = \pi(U) \times \{g_p\}.$$

In other words, \mathcal{L}_p is given locally by the images of the horizontal sections.

5.5. Intersection dimensions

Recall the situation described in Section 5.4 and, for a point $p \in P$, let \mathcal{L}_p denote the leaf of \mathcal{F} through p. Let V denote an irreducible subvariety of S. We will make repeated use of the following lemma.

Lemma 13. Let $p \in P$ and let Y denote a subvariety of \check{X} such that $p \in \pi_P^{-1}(V) \cap \beta^{-1}(Y)$. For any choice of representation $p = [x, g_p]$, we have

$$\dim_p(\mathcal{L}_p \cap \beta^{-1}(Y) \cap \pi_P^{-1}(V)) = \dim_x(g_pY \cap \pi^{-1}(V)).$$

Proof. Write $p = [x, g_p]$. By definition, $x \in g_p Y \cap \pi^{-1}(V)$. Fix a flat structure \mathcal{C} for P and consider $U \in \mathcal{C}$ such that $x \in U$. From Section 5.4, we have

$$\varphi_U^{-1}(\mathcal{L}_p \cap \pi_P^{-1}(\pi(U))) = \pi(U) \times \{g_p\}$$
(2)

and, from the definitions, we have

$$\varphi_U^{-1}(\pi_P^{-1}(V) \cap \pi_P^{-1}(\pi(U))) = (\pi(U) \cap V) \times \mathbf{G}(\mathbb{C}).$$
(3)

Finally, we claim that

$$\varphi_U^{-1}(\beta^{-1}(Y) \cap \pi_P^{-1}(\pi(U))) = \{ (\pi(gy), g) : y \in \check{Y}, g \in \mathbf{G}(\mathbb{C}), gy \in U \}.$$
(4)

To see (4), first note that

$$\beta^{-1}(Y) = \{ [gy, g] : y \in Y, g \in \mathbf{G}(\mathbb{C}) \}.$$

In particular,

$$\beta^{-1}(Y) \cap \pi_P^{-1}(\pi(U)) = \{ [gy, g] : y \in Y, g \in \mathbf{G}(\mathbb{C}), \pi(gy) \in \pi(U) \}.$$

Now, if $\pi(gy) \in \pi(U)$, then $gy \in \gamma U$ for some $\gamma \in \Gamma$. That is, $\gamma^{-1}gy = \pi_U^{-1}(\pi(gy))$ and so

$$\varphi_U^{-1}([gy,g]) = (\pi(gy), \gamma^{-1}g) = (\pi(\gamma^{-1}gy), \gamma^{-1}g),$$

which is an element belonging to the set on the right hand side of (4). On the other hand, if $y \in Y, g \in \mathbf{G}(\mathbb{C})$, and $gy \in U$, then $[gy,g] \in \beta^{-1}(Y) \cap \pi_P^{-1}(\pi(U))$ maps to $(\pi(gy),g)$. This establishes (4).

Combining (2)–(4), we conclude that $\varphi_U^{-1}(\mathscr{L}_p \cap \beta^{-1}(gY) \cap \pi_P^{-1}(V) \cap \pi_P^{-1}(\pi(U)))$ is equal to the set of tuples $(\pi(g_p y), g_p)$, where $y \in Y$ is such that $g_p y \in U \cap \pi^{-1}(V)$. Therefore, applying π_U^{-1} to the first factor, we obtain

$$\varphi_U^{-1}(\mathcal{L}_p \cap \beta^{-1}(Y) \cap \pi_P^{-1}(V) \cap \pi_P^{-1}(\pi(U))) \cong g_p Y \cap U \cap \pi^{-1}(V),$$
(5)

which proves the result.

6. The main construction

Recall the situation described in Section 5.5:

• (G, X) is a Shimura datum component such that $Z(G)(\mathbb{R})$ is compact;

- Γ is a neat arithmetic subgroup of G(Q) contained in G(Q)₊ equal to π₁(S), where S is the Shimura variety associated with (G, X) and Γ;
- V is an irreducible subvariety of S;
- *P* is the standard principal bundle associated with (\mathbf{G}, X) and Γ ;
- $\pi: X \to S$ and $\pi_P: P \to S$ denote the natural maps;
- $\beta: P \to \check{X}$ denotes the map defined in Section 5.1;
- \mathcal{F} denotes the canonical foliation of *P* (see Section 5.4);
- \mathcal{L}_p denotes the leaf of \mathcal{F} through $p \in P$.

Fix an embedding of \check{X} into projective space and let $D \in \mathbb{N}$ be as in Remark 11. Let $\Omega(k) = \Omega(\check{X}, k, D^{k+1})$ denote the (quasi-projective) Chow variety of closed irreducible complex subvarieties of \check{X} of codimension at most k and degree at most D^{k+1} . Let

$$\Omega = \bigcup_{k=0}^{\dim \check{X}} \Omega(k).$$

Consider the algebraic subvariety $\Theta = \Theta(V)$ of $P \times \Omega$ consisting of the tuples (p, Y) such that $p \in \pi_P^{-1}(V) \cap \beta^{-1}(Y)$. Here we slightly abuse notation by using *Y* to denote the Chow coordinate representing an irreducible variety as well as the variety itself. However, the correspondence between the Chow coordinate of *Y* and the points of *Y* is of course algebraic, and Θ is indeed Zariski closed. Note that there is a natural morphism $\Theta \to S$ given by the projection to *P* composed with π_P .

We define a function

$$d = d(V) : \Theta \to \mathbb{N} \cup \{0\}$$

by setting $d(p, Y) = \dim_p(\mathcal{L}_p \cap \beta^{-1}(Y) \cap \pi_p^{-1}(V)).$

For any $(p, Y_1) \in \Theta$, we let $\delta_1(p, Y_1)$ be the statement that

$$\dim Y_1 - d(p, Y_1) < \dim Y_2 - d(p, Y_2)$$

for all $Y_2 \in \Omega$ such that $Y_1 \subsetneq Y_2$. Similarly, we let $\delta_2(p, Y_1)$ be the statement that

$$d(p, Y_2) < d(p, Y_1)$$

for all $Y_2 \in \Omega$ such that $\beta(p) \in Y_2 \subsetneq Y_1$.

We define $\Pi = \Pi(V)$ to be the set of tuples $(p, Y) \in \Theta$ for which $\delta_1(p, Y)$ and $\delta_2(p, Y)$ hold. For any $d \in \mathbb{N}$, we let $\Pi(d)$ denote the set of tuples $(p, Y) \in \Pi$ such that

$$\dim Y - d(p, Y) = d. \tag{6}$$

Recall the notion of complexity of locally closed sets introduced before Theorem 4. The proof of the following results relies on differential-algebraic tools developed in Section 7, and is presented below to avoid breaking the logical flow of the paper. However, the reader may easily verify that the contents of Section 7 are self-contained and do not rely on Proposition 14.

Proposition 14. The sets $\Pi(d)$ for $d \in \mathbb{N}$ are locally closed subsets of Θ . The complexity of $\Pi(d)$ is bounded by $f_S(\deg(V))$ for some polynomial f_S depending only on S. Moreover, one can derive an explicit system of equations and inequations for $\Pi(d)$, as described in Theorem 4.

The (algebraic, left) action of $\mathbf{G}(\mathbb{C})$ on $P \times \Omega$ defined by g(p, Y) = (gp, gY) preserves the $\Pi(d)$ and their irreducible components.

Proof. Consider the foliation \mathcal{F}_0 on $P \times \Omega$ given by the direct product of (1) the canonical foliation of P and (2) the trivial foliation by zero-dimensional leafs on Ω .

Applying Proposition 20 below to the sets $\Sigma(\Theta, \mathcal{F}_0, k)$ we conclude that the sets

$$A(k) := \{ (p, Y) \in \Theta : d(p, Y) \ge k \}$$

$$\tag{7}$$

are Zariski closed with degrees bounded by a polynomial as claimed, and that it is possible to effectively compute equations for these sets. (This is our principal ingredient from differential algebra, and we will apply this below to deduce the same result for the sets $\Pi(d)$.)

Let $d \in \mathbb{N}$. Since Ω is a disjoint union of Chow varieties corresponding to different dimensions and degrees, it will be enough to consider each of these components separately. We now restrict, therefore, to one of these Chow varieties and assume that dim *Y* and deg(*Y*) are fixed.

The set $\Delta(d) \subset P \times \Omega$ defined by (6) is given by $A(\dim Y - d) \setminus A(\dim Y - d + 1)$. It is therefore locally closed. We claim further that the condition $\delta_1(p, Y)$ is open in $\Delta(d)$. To see this, let $\overline{\Omega}$ denote the projective closure of Ω and consider

$$\Delta_1 := \{ (p, Y_1, Y_2) \in P \times \Omega' \times \Omega : Y_1 \subsetneq^* Y_2, d \ge \dim Y_2 - d(p, Y_2) \}.$$

Here we write $Y_1 \subsetneq^* Y_2$ to mean that Y_1 is strictly contained in *each* component of the support of Y_2 . This is a Zariski closed condition, and Δ_1 is therefore closed for the same reason that A(k) is closed (with similar degree bounds, etc.). Since $\overline{\Omega}$ is projective, the projection $\pi_{\Theta}(\Delta_1)$ is closed as well and the standard resultant constructions from elimination theory produce effective systems of equations for this set as well.

By definition, in $\Delta(d)$ the condition $\delta_1(p, Y)$ essentially agrees with the complement of $\pi_{\Theta}(\Delta_1)$, except for a minor technicality: in $\delta_1(p, Y)$ we quantify over Y_2 in the open Chow variety Ω , whereas in the definition of Δ_1 we used the closed $\overline{\Omega}$. It is, however, easy to see that quantifying over $\overline{\Omega}$ gives an equivalent condition. Indeed, the points of the closed Chow variety $\overline{\Omega}$ represent effective cycles. If there exists $Y_2 \in \overline{\Omega}$ with $Y_1 \subsetneq^* Y_2$ such that

$$\dim Y_1 - d(p, Y_1) < \dim Y_2 - d(p, Y_2),$$

then the same must be true for one of the irreducible components of the support of Y_2 (note that here it is crucial that we used the refined relation \subsetneq^*). To conclude, in $\Delta(d)$ the condition $\delta_1(p, Y)$ is given by the complement of $\pi_{\Theta}(\Delta_1)$, and is therefore locally closed with the stated degree bounds and explicit equations. In an entirely analogous way, one checks that $\delta_2(p, Y)$ is open in Δ , and this concludes the proof of the local closedness, as well as the degree bounds for $\Pi(d)$.

The fact that $\mathbf{G}(\mathbb{C})$ preserves the $\Pi(d)$ is immediate from the definitions. Considering the action as a morphism $\mathbf{G}(\mathbb{C}) \times P \times \Omega \rightarrow P \times \Omega$ yields the remaining claims.

Lemma 15. Let $d \in \mathbb{N}$, let $(p, Y) \in \Pi(d)$, and write $p = [x, g_p]$. Then

- (i) $x \in g_p Y \cap \pi^{-1}(V)$;
- (ii) if A denotes an irreducible analytic component of $g_p Y \cap \pi^{-1}(V)$ passing through x such that

$$\dim A = \dim_X(g_p Y \cap \pi^{-1}(V)),$$

A is a Zariski optimal intersection component of $\pi^{-1}(V)$;

- (iii) writing X_A for the pre-weakly special subvariety $\langle A \rangle_{Zar}$ of X (see Theorem 12), we have $g_p Y = \check{X}_A$;
- (iv) $\delta_{\operatorname{Zar}}(A) = d$.

Proof. We will imitate the proof of [9, Lemma 6.14]. The fact that $x \in g_p Y \cap \pi^{-1}(V)$ follows from the definition of Π . Note also that

$$\dim_x(g_p Y \cap \pi^{-1}(V)) = d(p, Y)$$

by (5). Therefore, let A be as in (ii). By definition, A is an intersection component and $\langle A \rangle_{Zar}$ is contained in $g_p Y$. Therefore, since $(p, Y) \in \Pi(d)$, we have

$$\delta_{\operatorname{Zar}}(A) \leq \dim g_p Y - \dim A = \dim Y - d(p, Y) = d.$$

Let *B* be an intersection component of $\pi^{-1}(V)$ containing *A* such that $\delta_{\text{Zar}}(B) \leq \delta_{\text{Zar}}(A)$. We can and do assume that *B* is Zariski optimal, and therefore, by Theorem 12, $\langle B \rangle_{\text{Zar}}$ is equal to a pre-weakly special subvariety X_B of *X*.

Let Z be an irreducible component of $g_p Y \cap \check{X}_B$ containing A. Observe that either $Z = g_p Y$, $Z = \check{X}_B$, or codim Z > codim $g_p Y$. In all cases, the degree of Z in \check{X} is at most $D^{\operatorname{codim} Z+1}$ and so $Z \in \Omega$. However, $\beta(p) \in g_p^{-1}Z \subseteq Y$, and

$$d(p, g_p^{-1}Z) = \dim_x(Z \cap \pi^{-1}(V)) = \dim A = d(p, Y).$$

Therefore, since $\delta_2(p, Y)$ holds, we conclude that $Z = g_p Y$. In particular, $g_p Y$ is contained in \check{X}_B .

On the other hand,

$$\dim \check{X}_B - d(p, g_p^{-1}\check{X}_B) = \dim \check{X}_B - \dim_x (\check{X}_B \cap \pi^{-1}(V))$$
$$\leq \delta_{\operatorname{Zar}}(B) \leq \delta_{\operatorname{Zar}}(A) \leq \dim Y - d(p, Y),$$

and since $\delta_1(p, Y)$ holds, it follows that $g_p Y = \check{X}_B$. Therefore, B = A and

$$\delta_{\operatorname{Zar}}(B) = \dim Y - d(p, Y) = d.$$

Lemma 16. Let A be a Zariski optimal intersection component of $\pi^{-1}(V)$ and let $d = \delta_{\text{Zar}}(A)$. Let X_A denote the pre-weakly special subvariety $\langle A \rangle_{\text{Zar}}$ of X (see Theorem 12). Then, for any p = [x, 1] with $x \in A$ satisfying

$$\dim_x(\dot{X}_A \cap \pi^{-1}(V)) = \dim A,$$

we have $(p, \check{X}_A) \in \Pi(d)$.

The proof of Lemma 16 is very similar to the proof of [9, Lemma 6.13]. However, note that there is an error in the statement of [9, Lemma 6.13]: the term "pre-weakly special" should be replaced by "Zariski optimal". This also occurs in [9, Proposition 6.10] and its proof.

Proof of Lemma 16. Let p = [x, 1] with $x \in A$ satisfying

$$\dim_x(\check{X}_A \cap \pi^{-1}(V)) = \dim A.$$

Since, X_A is pre-weakly special, $(p, \check{X}_A) \in \Theta$ and we will now show that $(p, \check{X}_A) \in \Pi$.

To that end, suppose that $\delta_1(p, \check{X}_A)$ does not hold. Therefore, there exists $Y \in \Omega$ such that $\check{X}_A \subsetneq Y$ and

$$\dim \dot{X}_A - d(p, \dot{X}_A) \ge \dim Y - d(p, Y).$$
(8)

Recall that $d(p, \check{X}_A) = \dim_x(\check{X}_A \cap \pi^{-1}(V)) = \dim A$. Let *B* be an irreducible analytic component of $Y \cap \pi^{-1}(V)$ passing through *x* such that dim $B = \dim_x(Y \cap \pi^{-1}(V))$. In other words,

$$d(p,Y) = \dim_p(\mathcal{L}_p \cap \beta^{-1}(Y) \cap \pi_P^{-1}(V)) = \dim_x(Y \cap \pi^{-1}(V)) = \dim B,$$

and we obtain $\delta_{\text{Zar}}(B) \leq \delta_{\text{Zar}}(A)$. It follows, as in the proof of [9, Lemma 6.13], that B = A. However, this contradicts (8), and so $\delta_1(p, \check{X}_A)$ holds.

Now suppose that $\delta_2(p, \check{X}_A)$ does not hold. Therefore, there exists $Y \in \Omega$ such that $Y \subsetneq \check{X}_A$ and $d(p, Y) = d(p, \check{X}_A)$. However, this implies that

$$\dim_x(Y \cap \pi^{-1}(V)) = \dim_x(\check{X}_A \cap \pi^{-1}(V)).$$

from which it follows that A is contained in $Y \subsetneq \check{X}_A$. However, this contradicts the fact that $X_A = \langle A \rangle_{\text{Zar}}$, and so $\delta_2(p, \check{X}_A)$ holds.

Finally, since $d(p, X_A) = \dim A$, we have

$$\dim \check{X}_A - d(p, \check{X}_A) = \dim X_A - \dim A = d,$$

and so $(p, \check{X}_A) \in \Pi(d)$.

Lemma 17. Let Π° denote an irreducible component of Π . There exists a pre-special subvariety $X_{\mathbf{H}}$ of X and a \mathbb{Q} -splitting $X_{\mathbf{H}} = X_1 \times X_2$ such that, for any $(p, Y) \in \Pi^{\circ}$, if we write p = [x, g], then $gY = \gamma \check{X}_{\mathbf{H}, x_2}$ for some $\gamma \in \Gamma$ and some $x_2 \in X_2$.

Proof. Fix a flat structure \mathcal{C} for P and let $(p_0, Y_0) \in \Pi^\circ$. Write $p_0 = [x_0, g_0]$ and let $U_0 \in \mathcal{C}$ be such that $x_0 \in U_0$. We have a biholomorphic map

$$\tilde{U}_0 = \pi_P^{-1}(\pi(U_0)) \times \Omega \xrightarrow{f_{U_0}} \pi(U_0) \times \mathbf{G}(\mathbb{C}) \times \Omega$$

given by $(p, Y) \mapsto (\varphi_{U_0}^{-1}(p), Y)$. We write $\prod_{U_0}^{\circ} = \prod^{\circ} \cap \tilde{U}_0$.

Observe that, for any pre-special subvariety $X_{\mathbf{H}}$ of X and any \mathbb{Q} -splitting $X_1 \times X_2$ of $X_{\mathbf{H}}$, the set $\mathcal{R}(X_{\mathbf{H}}, X_1, X_2)$ of points $(s, g, Y) \in S \times \mathbf{G}(\mathbb{C}) \times \Omega$ such that $gY = \check{X}_{\mathbf{H}, x_2}$ for some $x_2 \in \check{X}_2$ is a closed algebraic subvariety. To see this, let $\mathcal{S}(X_{\mathbf{H}}, X_1, X_2)$ denote the closed algebraic subvariety of tuples

$$(y, g, Y, x_2) \in \check{X} \times \mathbf{G}(\mathbb{C}) \times \Omega \times \check{X}_2$$

such that $y \in gY \cap \check{X}_{\mathbf{H},x_2}$ and let f denote the natural projection from $\mathscr{S}(X_{\mathbf{H}}, X_1, X_2)$ to $\mathbf{G}(\mathbb{C}) \times \Omega \times \check{X}_2$. Observe that the set of points $(g, Y, x_2) \in \mathbf{G}(\mathbb{C}) \times \Omega \times \check{X}_2$ satisfying

$$\dim f^{-1}((g, Y, x_2)) \ge \max \{\dim Y, \dim X_1\}$$

constitutes a closed algebraic subvariety (to simplify the exposition, one may assume that dim *Y* is constant on Ω). Now the observation follows from the fact that, because \check{X}_2 is projective, the natural projection from $\mathbf{G}(\mathbb{C}) \times \Omega \times \check{X}_2$ to $\mathbf{G}(\mathbb{C}) \times \Omega$ is closed.

By Lemma 15, $f_{U_0}(\Pi_{U_0}^{\circ})$ is contained in the union of the $\mathcal{R}(X_{\mathbf{H}}, X_1, X_2)$ as $X_{\mathbf{H}}$ varies over the pre-special subvarieties of X and $X_1 \times X_2$ varies over the Q-splittings of $X_{\mathbf{H}}$. Furthermore, after possibly replacing U_0 with a subset (also belonging to \mathcal{C}), we may assume that $f_{U_0}(\Pi_{U_0}^{\circ})$ is connected. Therefore, since their union is countable, $f_{U_0}(\Pi_{U_0}^{\circ})$ is contained in one of the $\mathcal{R}(X_{\mathbf{H}}, X_1, X_2)$, which we denote \mathcal{R} .

By definition, $P \times \Omega$ is covered by the union of the \tilde{U} as U varies over the elements of \mathcal{C} . Therefore, since Π° is path-connected and the transition functions associated with the trivializations of P are given by elements of Γ , we conclude that there exists a prespecial subvariety $X_{\mathbf{H}}$ of X and a \mathbb{Q} -splitting $X_{\mathbf{H}} = X_1 \times X_2$ such that, for any $(p, Y) \in$ Π , if we write p = [x, g], then $gY = \gamma \check{X}_{\mathbf{H}, x_2}$ for some $\gamma \in \Gamma$ and some $x_2 \in \check{X}_2$. To conclude the proof, we recall that $x \in gY$. Hence, $x \in X \cap \gamma \check{X}_{\mathbf{H}, x_2}$ and so $x_2 \in X_2$, by Lemma 10.

With each $d \in \mathbb{N}$ and each irreducible component $\Pi(d)^{\circ}$ of $\Pi(d)$, we associate a triple $T = (X_{\mathbf{H}}, X_1, X_2)$, where $X_{\mathbf{H}}$ is a pre-special subvariety $X_{\mathbf{H}}$ of X and $X_{\mathbf{H}} = X_1 \times X_2$ is a Q-splitting, such that Lemma 17 holds for all $(p, Y) \in \Pi(d)^{\circ}$. With the triple T we associate the standard principal bundle $P_T = \Gamma_{\mathbf{H}} \setminus (X_{\mathbf{H}} \times \mathbf{H}(\mathbb{C}))$ associated with $(\mathbf{H}, X_{\mathbf{H}})$ and $\Gamma_{\mathbf{H}} = \Gamma \cap \mathbf{H}(\mathbb{Q})_+$. (This is indeed a principal bundle since $\Gamma_{\mathbf{H}}$ is neat and, by [29, Remark 2.3], $\mathbf{Z}(\mathbf{H})(\mathbb{R})$ is compact.) We let Ω_T denote the subvariety of Ω comprising the subvarieties of \check{X} of the form $\check{X}_{\mathbf{H},x_2}$ for some $x_2 \in \check{X}_2$. By [20, Section 3], the natural map

$$P_T \times \Omega_T \xrightarrow{\iota_{P_T}} P \times \Omega$$

is algebraic, and it is easy to check that it is injective. We let $\Pi(d)_T$ denote the locally closed subset

$$\iota_{P_T}^{-1}(\Pi(d)^\circ) \subset P_T \times \Omega_T,$$

and we let $\Pi(d)_{S_T}$ denote its image in $S_T = \Gamma_{\mathbf{H}} \setminus X_{\mathbf{H}}$ under the natural map.

If we let $\mathbf{H}_1 \times \mathbf{H}_2$ denote the decomposition of \mathbf{H}^{ad} giving rise to the \mathbb{Q} -splitting $X_{\mathbf{H}} = X_1 \times X_2$, and we let Γ_2 denote the image of $\Gamma_{\mathbf{H}}$ in $\mathbf{H}_2(\mathbb{Q})$, we obtain a diagram

$$X_{2} \longleftrightarrow X_{H} \longleftrightarrow X$$
$$\downarrow^{\pi_{T,2}} \qquad \downarrow^{\pi_{T}} \qquad \downarrow^{\pi}$$
$$\Gamma_{2} \backslash X_{2} \xleftarrow{\phi_{T}} S_{T} \xrightarrow{\iota_{T}} S$$

We let $\Pi(d)_{S_T}$ denote the union of the Zariski closures of the fibres of the map ϕ_T restricted to $\Pi(d)_{S_T}$ and we let $\phi_{T,d}$ denote the map

$$\overline{\Pi}(d)_{S_T} \to \phi_T(\Pi(d)_{S_T}).$$

Lemma 18. The sets $\overline{\Pi}(d)_{S_T}$ are constructible.

Proof. It is a general fact that for any constructible map $f : X \to Y$, the union of the Zariski closures of the fibres is constructible. Since we did not find a suitable reference we give the details below.

Up to taking affine covers one may assume that X and Y are affine. The union of the Zariski closures can be defined by

$$\{x \in X : \forall_{P \in \mathbb{C}[X]} [(\forall_{x' \in f^{-1}(f(x))} P(x') = 0) \Rightarrow P(x) = 0]\}.$$
(9)

This would be constructible if one could restrict to quantifying over $P \in \mathbb{C}[X]$ of degree bounded by some $N \in \mathbb{N}$, that is, if one could show that the Zariski closures of $f^{-1}(y)$ are set-theoretically cut out by some polynomials of uniformly bounded degree. Equivalently, it suffices to show that the Zariski closures of these sets have uniformly bounded degrees, which is standard.

We have the following structure theorem for weakly optimal subvarieties.

Theorem 19. *Let* $d \in \mathbb{N}$ *.*

- (i) Let W be a weakly optimal subvariety of V such that $\delta_{ws}(W) = d$. Then there exists an irreducible component $\Pi(d)^\circ$ of $\Pi(d)$ such that, if T denotes the triple associated with $\Pi(d)^\circ$, then $W = \iota_T(W_T)$ for some irreducible component W_T of a fibre of $\phi_{T,d}$.
- (ii) Let $\Pi(d)^{\circ}$ denote an irreducible component of $\Pi(d)$ and let $T = (X_{\mathbf{H}}, X_1, X_2)$ denote the triple associated with $\Pi(d)^{\circ}$. If W is an irreducible component of a fibre of $\phi_{T,d}$, then $\iota_T(W)$ is a weakly optimal subvariety of V such that $\delta_{ws}(W) = d$ and $\langle W \rangle_{ws} = \pi(X_{\mathbf{H},x_2})$ for some $x_2 \in X_2$.

Proof of (i). Let A be an irreducible analytic component of $\pi^{-1}(W)$. By [9, Proposition 6.9], A is a Zariski optimal intersection component of $\pi^{-1}(V)$ and so, by Theorem 12, $\langle A \rangle_{\text{Zar}}$ is a pre-weakly special subvariety X_A of X. A simple calculation shows that $\pi(X_A) = \langle W \rangle_{\text{ws}}$ and so $\delta_{\text{Zar}}(A) = \delta_{\text{ws}}(W) = d$.

By Lemma 16, for any p = [x, 1] with $x \in A$ satisfying

$$\dim_x(X_A \cap \pi^{-1}(V)) = \dim A, \tag{10}$$

we have $(p, \check{X}_A) \in \Pi(d)$. Since (10) defines an open subset of A, there exists an open subset U of A and an irreducible component $\Pi(d)^\circ$ of $\Pi(d)$ such that $(p, \check{X}_A) \in \Pi(d)^\circ$ for all p = [x, 1] with $x \in U$.

Let $T = (X_{\mathbf{H}}, X_1, X_2)$ be the triple associated with $\Pi(d)^\circ$. Then $\check{X}_A = \gamma \check{X}_{\mathbf{H}, x_2}$ for some $\gamma \in \Gamma$ and some $x_2 \in X_2$. Therefore, $(p, \gamma^{-1}\check{X}_A) \in P_T \times \Omega_T$ for all $p = [\gamma^{-1}x, 1]$ with $x \in U$. In fact, by Proposition 14, these points also belong to $\Pi(d)^\circ$ as they belong to the $\mathbf{G}(\mathbb{C})$ -orbits of the points above.

Let W_T denote the irreducible component $\pi_T(\gamma^{-1}A)$ of $\iota_T^{-1}(W)$. Then W_T is a weakly optimal subvariety of $V_T = \iota_T^{-1}(V)$ and $\langle W_T \rangle_{ws}$ is equal to $S_{\mathbf{H},x_2} = \pi_T(X_{\mathbf{H},x_2})$. In particular, W_T is an irreducible component of $S_{\mathbf{H},x_2} \cap V_T$, which is the fibre of $V_T \to \phi_T(V_T)$ over the point $z_2 = \pi_{T,2}(x_2)$. However, since $\Pi(d)_{S_T}$ contains $\pi_T(\gamma^{-1}U)$, we deduce that W_T is contained in $\overline{\Pi}(d)_{S_T}$, and therefore is an irreducible component of the fibre of $\phi_{T,d}$ over z_2 .

Proof of (ii). Let $z \in \Pi(d)_{S_T} \cap W$ (observe that $\Pi(d)_{S_T} \cap W$ is Zariski dense in W, and in particular is non-empty). Write $z = \pi_T(x)$ for some $x = (x_1, x_2) \in X_{\mathbf{H}} = X_1 \times X_2$ (in particular, W is an irreducible component of the fibre of $\phi_{T,d}$ over $\pi_{T,2}(x_2)$) and choose $(p, Y) \in \Pi(d)_T$ lying above z. Clearly, we can choose p = [x, h] for some $h \in \mathbf{H}(\mathbb{C})$, and so $hY = \check{X}_{\mathbf{H}, y_2}$ for some $y_2 \in X_2$. In fact, since $x \in hY$, we conclude that $y_2 = x_2$.

Let *B* denote an irreducible analytic component of $hY \cap \pi^{-1}(V)$ containing *x* such that

$$\dim B = \dim_x(hY \cap \pi^{-1}(V)).$$

By Lemma 15, *B* is a Zariski optimal intersection component of $\pi^{-1}(V)$ such that $\delta_{\text{Zar}}(B) = d$ and, writing X_B for the pre-weakly special subvariety $\langle B \rangle_{\text{Zar}}$ of *X*, we have $hY = \check{X}_{\mathbf{H},x_2} = \check{X}_B$. By [9, Proposition 6.9], $W_B = \pi(B)$ is a weakly optimal subvariety of *V*, and we see that $\delta_{ws}(W_B) = d$ and $\langle W_B \rangle_{ws} = \pi(X_{\mathbf{H},x_2})$.

Since $\Pi(d)_{S_T} \cap W$ is Zariski dense in W, we conclude that $\iota_T(W)$ is contained in the union of the W_B as obtained above. This is a finite union since each W_B is a weakly optimal subvariety of V with weakly special closure $\pi(X_{\mathbf{H},x_2})$, and $\pi(X_{\mathbf{H},x_2})$ does not depend on z. Since $\iota_T(W)$ is irreducible, it is contained in one such subvariety, and we conclude that

$$\dim W \le \dim B = \dim X_1 - d. \tag{11}$$

On the other hand, since $\Pi(d)$ is locally closed, $\Pi(d)^{\circ}$ contains a Zariski open subset U that is disjoint from the other irreducible components of $\Pi(d)$. Then $U_T = \iota_{P_T}^{-1}(U)$ is Zariski open in $\Pi(d)_T$ and so, since it is dense and constructible, the image U_{S_T} of U_T in $\Pi(d)_{S_T}$ contains an open subset of $\Pi(d)_{S_T}$.

Suppose then that $z \in U_{S_T}$ and $(p, Y) \in U_T$. By Lemma 16, for any $y \in B$ such that

$$\dim B = \dim_{\mathcal{V}}(hY \cap \pi^{-1}(V))$$

and p = [y, 1], we have $(p, hY) \in \Pi(d)$. Since this condition defines an open subset of *B* containing *x*, we conclude that there exists an open subset U_B of *B* containing *x* such that, for any $y \in U_B$ and p = [y, 1], we have $(p, hY) \in \Pi(d)_T$. It follows that $\pi_T(B)$ is contained in the fibre of $\phi_{T,d}$ over $\pi_{X_2}(x_2)$.

Therefore, since U_{S_T} contains an open subset of $\Pi(d)_{S_T}$, the irreducible components of the fibres of $\phi_{T,d}$ are of dimension at least dim $B = \dim X_1 - d$. Hence, we conclude from (11) that they are pure of dimension dim $X_1 - d$ and this concludes the proof.

Observe that Theorem 19 implies [9, Proposition 6.3], which establishes that the weakly optimal subvarieties of V come from finitely many triples $(X_{\rm H}, X_1, X_2)$. The novelty in Theorem 19 is that the fibres of the $\phi_{T,d}$ are precisely the weakly optimal subvarieties of V of weakly special defect d.

We recall that [9, Theorem 7.2] established that the union V^{an} of the positive-dimensional weakly optimal subvarieties of V of weakly special defect at most d is a Zariski closed subset of V. If V^{an} is strictly contained in V, the Zilber–Pink conjecture can be reduced to arithmetic (see [9, Theorem 14.3]).

7. Ingredients from differential-algebraic geometry

7.1. The main statement

Let \bar{X} denote a proper smooth complex algebraic variety of dimension d, and let $X \subset \bar{X}$ denote an open dense subset. Let L denote a very ample line bundle on \bar{X} . In this section, the *degree* deg(Y) of a subvariety $Y \subset X$ is taken to mean the degree of the Zariski closure $\bar{Y} \subset \bar{X}$ with respect to L.

Let \mathcal{F} denote a non-singular *n*-dimensional foliation of *X*. We further assume for simplicity that *X* is affine and that \mathcal{F} is generated by *n* commuting regular vector fields $\boldsymbol{\xi} = (\xi_1, \ldots, \xi_n)$. We may thus think of regular/rational functions *P* on *X* as restrictions of polynomials/rational functions on a suitable \mathbb{A}^N and we use deg(*P*) to denote the (minimal) degree of such a representative. Similarly, we denote by deg($\boldsymbol{\xi}$) the maximum among the degrees of the coefficients of ξ_1, \ldots, ξ_n thought of as fields $\xi_i : X \to T \mathbb{A}^N$.

The assumption above can always be achieved by passing to an affine cover of X. Identifying X as an affine subvariety of \mathbb{A}^N and choosing generic linear coordinates x_1, \ldots, x_N on \mathbb{A}^N , there are unique rational vector fields ξ_1, \ldots, ξ_n tangent to \mathcal{F} of the form

$$\xi_i = \frac{\partial}{\partial x_i} + \sum_{j=n+1}^N c_{ij}(x) \frac{\partial}{\partial x_j}$$
(12)

where c_{ij} are rational functions. Moreover, $[\xi_i, \xi_j] = 0$ by the Frobenius theorem. The results below can be applied, after this affine covering process, to any foliation \mathcal{F} as above.

For any algebraic subvariety $V \subset X$ and $k \in \mathbb{N}$, let

$$\Sigma(V, \mathcal{F}, k) = \{ p \in V : \dim(\mathcal{L}_p \cap V) \ge k \}.$$

Our main tool is the following.

Proposition 20. Let $V \subset X$ be an algebraic subvariety. Then

$$\deg(\Sigma(V,\mathcal{F},k)) \le P_N(\deg(X),\deg(V),\deg(\xi))$$

for some explicit polynomial P_N depending on N. Moreover, equations for $\Sigma(V, \mathcal{F}, k)$ can be effectively computed from the equations defining X, V and $\boldsymbol{\xi}$.

Remark 21. The explicit choice of affine coordinates is not strictly necessary to state the degree bound in Proposition 20. However, it is convenient for establishing the effective nature of our construction (i.e. to clarify the sense in which *equations* for $\Sigma(V, \mathcal{F}, k)$ are to be effectively computed).

We prove Proposition 20 below after describing how it relates to our concrete context involving flat connections on a Shimura variety.

7.2. Flat connections and foliations

Recall the situation described in Section 5.1. Let U be an affine Zariski open subset of S such that $\pi_P^{-1}(U) \cong U \times \mathbf{G}(\mathbb{C})$ and suppose that x_1, \ldots, x_n is a system of étale coordinates on U. The connection $\nabla \in \Omega^1(P, \mathfrak{g}_{\mathbb{C}})$ can be written, with respect to these coordinates, as

$$\nabla = \sum_{i=1}^{n} \Omega_i dx_i$$

where Ω_i is an algebraic morphism $U \to \mathfrak{g}_{\mathbb{C}}$. Choosing a faithful representation $\mathfrak{g} \to \mathfrak{gl}_M$, we may write the Ω_i as matrices with entries given by polynomials in the x_1, \ldots, x_n .

The vector fields $\frac{\partial}{\partial x_i}$ on U lift to vector fields ξ_i on $\pi_P^{-1}(U)$, which in our choice of coordinates can be written

$$\xi_i = \frac{\partial}{\partial x_i} + \Omega_i \cdot g.$$

The vector fields ξ_i commute by the flatness of ∇ . By definition, their integral manifolds \mathcal{L}_p at a point $p \in P$ are given by (germs of) horizontal sections of ∇ .

7.3. Multiplicity estimates

In order to prove Proposition 20, we will require the following multiplicity estimate due to Gabrielov–Khovanskii [11].

Theorem 22 ([11, Theorem 1]). Let $P = (P_1, ..., P_n) \in \mathcal{O}(X)^n$ and let $p \in X$. Suppose that the restriction of P to the leaf \mathcal{L}_p of \mathcal{F} through p has an isolated zero at p. Then

 $\operatorname{mult}_{p} P|_{\mathcal{L}_{p}} < f_{N}(\operatorname{deg}(\boldsymbol{\xi}), \operatorname{deg}(P))$

for some explicit polynomial f_N .

We will use this result to characterize the locus of points where the intersection of \mathcal{L}_p with the vanishing locus of *P* is positive-dimensional. For this purpose, we are interested in expressing the condition that a tuple of functions admits a common zero of multiplicity at least *k* by means of differential-algebraic conditions.

Let $F = (F_1, \ldots, F_n)$ denote an *n*-tuple of holomorphic functions in some domain $\Omega \subset \mathbb{C}^n$. The problem above is addressed in [5] by means of a collection $\{M^{\alpha}\}$ of "multiplicity operators" of order k. These are polynomial partial differential operators of order k, i.e. polynomial combinations of F_1, \ldots, F_n and their first k derivatives. We will usually denote a multiplicity operator of order k by $M^{(k)}$ and write $M_p^{(k)}F$ for $[M^{(k)}(F)](p)$.

Proposition 23 ([5, Proposition 5]). We have $\operatorname{mult}_p F > k$ if and only if $M_p^{(k)}F = 0$ for all multiplicity operators of order k.

For every $p \in X$, we let $\phi_p : B \to \mathcal{L}_p$ denote the germ of a holomorphic map, for some open ball $B \subset \mathbb{C}^n$ centred at the origin, satisfying $\partial \phi_p / \partial x_i = \xi_i$ for i = 1, ..., n. We refer to this map as the ξ -chart on \mathcal{L}_p . When $P = (P_1, ..., P_n) \in \mathcal{O}(X)^n$ we may apply the multiplicity operator $M^{(k)}$ to P by evaluating the derivatives along $\xi_1, ..., \xi_n$, which amounts to computing, for each point $p \in X$, the multiplicity operator of $P|_{\mathcal{L}_p}$ in the ξ -chart.

Lemma 24. For any multiplicity operator $M^{(k)}$ of order k we have

 $\deg(M^{(k)}P) \le f_N(\deg(P), \deg(\boldsymbol{\xi}), k)$

for some explicit polynomial f_N .

Proof. This follows easily since $M^{(k)}$ is defined by expanding a determinant, of size polynomial in k, with entries defined in terms of P and its ξ_i -derivatives up to order k.

7.4. Proof of Proposition 20

Recall the situation described in Section 7.1. It is classical that $V \subset X$ is then cut out by polynomials of degree at most deg(V). Denote the set of these polynomials by \mathcal{P} . Then we have

$$\Sigma(V,\mathcal{F},k) = \bigcap_{\substack{\mathcal{P}' \subset \mathcal{P} \\ \#\mathcal{P}' = n-k+1}} \Sigma(V(\mathcal{P}'),\mathcal{F},k).$$

Indeed, the inclusion \subset is obvious. For the other inclusion, suppose $p \notin \Sigma(V, \mathcal{F}, k)$ so that $\dim(V \cap \mathcal{L}_p) < k$. In particular, $\mathcal{L}_p \not\subset V$, so there exists an equation $P_1 \in \mathcal{P}$ not identically zero on \mathcal{L}_p . If n - k > 1 then, similarly, no component of the intersection of \mathcal{L}_p with

the vanishing locus of P_1 is contained in V, so there exists P_2 not vanishing on any of these components. Reiterating n - k steps of this form, we obtain $\mathcal{P}' = \{P_1, \ldots, P_{n-k+1}\}$ with dim $(V(\mathcal{P}') \cap \mathcal{L}_p) = k - 1$, so $p \notin \Sigma(V(\mathcal{P}'), \mathcal{F}, k)$.

By Bézout's Theorem, $\deg(\Sigma(V, \mathcal{F}, k))$ is bounded by a polynomial in the maximum of the $\deg(\Sigma(V(\mathcal{P}'), \mathcal{F}, k))$ for \mathcal{P}' as above. Hence, it is enough to prove Proposition 20 assuming that V is a complete intersection defined by $P = (P_1, \ldots, P_{n-k+1})$.

We now make a similar reduction involving the foliation \mathcal{F} . Namely,

$$\Sigma(V, \mathcal{F}, k) = \bigcap_{\substack{\mathcal{F}' \subset \mathcal{F} \\ \dim \mathcal{F}' = n-k+1}} \Sigma(V, \mathcal{F}', 1),$$
(13)

where the intersection is taken over foliations \mathcal{F}' generated by linear combinations of n-k+1 of the vector fields comprising $\boldsymbol{\xi}$. Again the inclusion \subset is obvious. For the other inclusion, suppose $p \notin \Sigma(V, \mathcal{F}, k)$, so that $\dim(V \cap \mathcal{L}_p) < k$. Intersecting with linear hyperplanes passing through the origin in the $\boldsymbol{\xi}$ -chart on \mathcal{L}_p , we find, similarly to the previous step, k-1 such hyperplanes defining a subleaf \mathcal{L}'_p with $\dim(V \cap \mathcal{L}'_p) = 0$. Noting that \mathcal{L}'_p is a leaf of a subfoliation \mathcal{F}' as above finishes the proof.

By Bézout's Theorem, as above, it suffices, replacing \mathcal{F} by the \mathcal{F}' , to prove Proposition 20 in the case k = 1. In this case we have

$$\Sigma(V, \mathcal{F}, 1) = \{ p \in V : \dim(\mathcal{L}_p \cap V) \ge 1 \}$$
$$= \{ p \in V : \operatorname{mult}_p P | \mathcal{L}_p = \infty \}$$
$$= \{ p \in V : \operatorname{mult}_p P | \mathcal{L}_p \ge \mu \},$$

where μ is the multiplicity bound of Theorem 22. Finally, according to Proposition 23, the right hand side is the zero locus of all multiplicity operators $M_p^{(\mu)}P$ taken with respect to the foliation \mathcal{F} . The degrees of all of these polynomials are bounded by Lemma 24. Applying Bézout's Theorem concludes the proof of the degree bound.

Finally, we indicate how to effectively obtain a system of equations for $\Sigma(V, \mathcal{F}, k)$. The only step above which is not effective a priori is (13), where one intersects an infinite collection of sets. To deal with this, we first note that it would suffice to consider \mathcal{F}' in some open-dense subset of the Grassmannian of n - k + 1-dimensional subspaces of the span of $\boldsymbol{\xi}$. We can generate such $\mathcal{F}'(c)$ as the span of $\boldsymbol{\xi}'_1(c), \ldots, \boldsymbol{\xi}'_{n-k+1}(c)$ with

$$\xi'_{i} = \xi_{i} + \sum_{j=n-k+2}^{n} c_{ij}\xi_{j}$$
(14)

and $(c_{ij}) \in \mathbb{A}^{(n-k+1)(k-1)}$. Repeating the construction above with $\mathcal{F}'(c)$ and treating c_{ij} as independent variables, we obtain a system of equations $E_1(x,c) = \cdots = E_Q(x,c) = 0$ such that $x \in \Sigma(V, \mathcal{F}, k)$ if and only if the equations vanish at (x, c) for every c. It is then clear that $\Sigma(V, \mathcal{F}, k)$ is cut out by the coefficients of E_1, \ldots, E_Q viewed as polynomials in the c-variables.

8. Effective bounds for degrees of weakly optimal subvarieties

Recall the situation described in Section 6. For i = 1, ..., n, let U_i denote a Zariski open subvariety of S such that $\bigcup_{i=1}^{n} U_i = S$ and $\pi_P^{-1}(U_i) \cong U_i \times \mathbf{G}(\mathbb{C})$ is trivial. Furthermore, we can and do assume that, under the embedding $S \to \mathbb{P}^N$ given by $L_{\Gamma}^{\otimes k_{\Gamma}}$, each U_i is contained in one of the standard affine charts. Similarly, let Ω_i denote Zariski open subvarieties of Ω (already considered as a subvariety of some \mathbb{P}^M), each contained in a standard affine chart, such that $\Omega = \bigcup_{i=1}^{n} \Omega_i$. Fix an embedding of **G** into some affine space \mathbb{A}^L . For i, j = 1, ..., n, let U_{ij} denote the Zariski open subset $U_i \times \mathbf{G}(\mathbb{C}) \times \Omega_j$ of $P \times \Omega$. The following is an immediate corollary of Proposition 14.

Lemma 25. Let $d \in \mathbb{N}$ and let $\Pi(d)^{\circ}$ denote an irreducible component of $\Pi(d)$. For any $i, j \in \{1, ..., n\}$, the degree of the Zariski closure of the locally closed subset $U_{ij} \cap \Pi(d)^{\circ}$ of $P \times \Omega$, considered as a subvariety of \mathbb{A}^{N+L+M} , is bounded by $f_S(\deg_{L_{\Gamma}^{\otimes k_{\Gamma}}}(V))$ for some polynomial f_S depending only on S.

The main result of this section is the following.

Theorem 26. Let $d \in \mathbb{N}$ and let W be a weakly optimal subvariety of V such that $\delta_{ws}(W) = d$. Then

$$\deg_{L_{\Gamma}^{\otimes k_{\Gamma}}}(W) \le f_{S}(\deg_{L_{\Gamma}^{\otimes k_{\Gamma}}}(V))$$

for some polynomial f_S depending only on S.

Proof. Since k_{Γ} depends only on S, we may replace $L_{\Gamma}^{\otimes k_{\Gamma}}$ with L_{Γ} .

Recall the situation reached at the end of the proof of Theorem 19 (i). Let $\Pi(d)_S$ denote the image of $\Pi(d)^\circ$ in S and let V_0 denote its Zariski closure. Let $V_{T,0}$ denote the Zariski closure of $\Pi(d)_{S_T}$ in S_T . Observe that $\phi_T(V_{T,0})$ is closed as ϕ_T extends to a (projective) morphism on the Baily–Borel compactifications.

We will prove the theorem in a succession of lemmas. The following lemma will be used in the proofs of Lemma 28 (ii, iii).

Lemma 27. The subvariety $V_{T,0}$ is irreducible and $\iota_T(V_{T,0}) = V_0$.

Proof. We appeal to the commutative diagram of natural morphisms



where $\iota_T^*(P \times \Omega)$ is the pullback of $P \times \Omega$ to S_T . Recall that ι_{P_T} is injective.

First observe that the $G(\mathbb{C})$ -orbit of (the image of) $\Pi(d)_T$ in $P \times \Omega$ is $\Pi(d)^\circ$. Indeed, the left-right inclusion is clear from Proposition 14. Therefore, let $(p, Y) \in \Pi(d)^\circ$ and

write p = [x, g]. Then $gY = \gamma \check{X}_{\mathbf{H}, x_2}$ for some $\gamma \in \Gamma$ and some $x_2 \in X_2$. We rewrite $p = [\gamma^{-1}x, \gamma^{-1}g]$ and we see that $\gamma^{-1}g(p, Y) \in P_T \times \Omega_T$, which proves the claim.

Let Z_T denote the Zariski closure of (the image of) $\Pi(d)_T$ in $\iota_T^*(P \times \Omega)$. Then, by the preceding paragraph, (the image of) $\mathbf{G}(\mathbb{C})Z_T$ is a Zariski dense subset of $\Pi(d)^\circ$. Let Y_T denote an irreducible component of Z_T such that $\mathbf{G}(\mathbb{C})Y_T$ is also dense in $\Pi(d)^\circ$. Then, since the Zariski closure of $f(\Pi(d)^\circ)$ is V_0 , we see that $f(\mathbf{G}(\mathbb{C})Y_T) = f(Y_T)$ is dense in V_0 and, by the commutativity of the diagram, equal to $\iota_T(Y_S_T)$, where $Y_{S_T} = g_T(Y_T)$.

Now let Y'_T denote another irreducible component of Z_T . We claim that $g_T(Y'_T)$ is contained in the Zariski closure of Y_{S_T} . To see this, observe that Y'_T is contained in the Zariski closure of $\mathbf{G}(\mathbb{C})Y_T$ in $\iota_T^*(P \times \Omega)$. Therefore, $g_T(Y'_T)$ is contained in the Zariski closure of $g_T(\mathbf{G}(\mathbb{C})Y_T) = g_T(Y_T)$.

It follows that the Zariski closure of Y_{S_T} is equal to $V_{T,0}$. Hence, $V_{T,0}$ is irreducible. The fact that $\iota_T(V_{T,0}) = V_0$ now follows easily from the facts that ι_T is closed and V_0 is equal to the Zariski closure of $f(\Pi(d)^\circ)$.

The following lemma will reduce the proof to Lemma 30 below.

- **Lemma 28.** (i) $\deg_{L_{\Gamma}}(V_0) \leq f_S(\deg_{L_{\Gamma}}(V))$ for some polynomial f_S depending only on S;
- (ii) $\deg_{L_{\Gamma_{u}}}(V_{T,0}) \leq r_{\mathbf{G}}^{3} \deg_{L_{\Gamma}}(V_{0})$, where $r_{\mathbf{G}}$ denotes the rank of \mathbf{G} ;
- (iii) $\deg_{L_{\Gamma}}(W) \leq r_{\mathbf{G}}^3 \deg_{L_{\Gamma_{\mathbf{u}}}}(W_T).$

Proof. There exist $i, j \in \{1, ..., n\}$ such that the Zariski closure of $U_{ij} \cap \Pi(d)^{\circ}$ dominates V_0 . Therefore, (i) follows easily from Lemma 25.

(ii) follows immediately from Lemma 27, [7, Lemma 4.1] (a corollary of [17, Proposition 5.3.10]) and [7, Lemma 4.2]; and (iii) follows from (ii).

The following lemma will be used in the proof of Lemma 30.

Lemma 29. (i) W_T is an irreducible component of the fibre of $V_{T,0} \rightarrow \phi_T(V_{T,0})$ over z_2 ; (ii) dim W_T = dim $X_1 - d$ is the generic dimension of the fibres of $V_{T,0} \rightarrow \phi_T(V_{T,0})$.

Proof. (i) Observe that $V_{T,0}$ is contained in V_T and W_T is contained in $V_{T,0}$. Therefore, since W_T is an irreducible component of the fibre of $V_T \rightarrow \phi_T(V_T)$ over z_2 , the claim follows.

(ii) First observe that the Zariski closures of $\Pi(d)_{S_T}$ and $\overline{\Pi}(d)_{S_T}$ coincide, that is, they are both $V_{T,0}$. Also, $\phi_T(V_{T,0})$ is contained in the Zariski closure of $\phi_T(\Pi(d)_{S_T})$. However, $\phi_T(V_{T,0})$ is closed and contains $\phi_T(\Pi(d)_{S_T})$. Therefore, $\phi_T(V_{T,0})$ is equal to the Zariski closure of $\phi_T(\Pi(d)_{S_T})$. Since, by Theorem 19, the fibres of $\phi_{T,d}$ are pure of dimension dim $X_1 - d$, the claim follows.

It remains to prove the following lemma.

Lemma 30. We have

$$\deg_{L_{\Gamma_{\mathbf{H}}}}(W_T) \leq \deg_{L_{\Gamma_{\mathbf{H}}}}(V_{T,0}).$$

Proof. In order to simplify notation, we will, for the remainder of the proof, replace $\Gamma_{\mathbf{H}}$ with Γ , S_T with S, $V_{T,0}$ with V, W_T with W, and dim W with d (as opposed to dim $X_1 - d$). We will also reassign ι to be the (proper) closed embedding $V \to S$. Finally, for i = 1, 2, we let Γ_i denote the image of Γ in $\mathbf{H}_i(\mathbb{Q})$, we let f denote the natural morphism $S \to S_1 \times S_2$, where $S_i = \Gamma_i \setminus X_i$ is the Shimura variety associated with (\mathbf{H}_i, X_i) and Γ_i , and we let ϕ_i denote the projection $S \to S_i$ (which factors as f composed with the natural projection f_i from $S_1 \times S_2$ to S_i).

By the projection formula, the degree of $c_1(L_{\Gamma})^d \cap [\iota(V)]$ is equal to the degree of $c_1(\iota^*L_{\Gamma})^d \cap [V] \in A_0(V)$. Let $V_2 = \phi_2(\iota(V))$, which, as explained above, is closed. There exists a Zariski open subset $U_2 \subset V_2$ such that, if $U = (\phi_2 \circ \iota)^{-1}(U_2)$, then $(\phi_2 \circ \iota)|_U$ is flat. In particular, its fibres are equidimensional, of dimension d (by Lemma 29 (ii)).

Now consider the excision exact sequence

$$A_0(V \setminus U) \xrightarrow{i_*} A_0(V) \xrightarrow{j^*} A_0(U)$$

where i_* is the pushforward associated with the (proper) closed embedding $i: V \setminus U \to V$, and j^* is the pullback associated with the (flat) inclusion $j: U \to V$. We see that the degree of $c_1(\iota^* L_{\Gamma})^{\dim V} \cap [V]$ is at least the degree of

$$j^*(c_1(\iota^*L_{\Gamma})^{\dim V} \cap [V]) = c_1(j^*\iota^*L_{\Gamma})^{\dim V} \cap [U] \in A_0(U).$$

Next, as in [17, Proposition 5.3.2(1)], we have

$$L_{\Gamma} = f^{*}(L_{\Gamma_{1}} \boxtimes L_{\Gamma_{1}}) = f^{*}(f_{1}^{*}L_{\Gamma_{1}} \otimes f_{2}^{*}L_{\Gamma_{2}}) = \phi_{1}^{*}L_{\Gamma_{1}} \otimes \phi_{2}^{*}L_{\Gamma_{2}}.$$

Therefore,

$$j^*\iota^*L_{\Gamma} = j^*\iota^*\phi_1^*L_{\Gamma_1} \otimes j^*\iota^*\phi_2^*L_{\Gamma_2}$$

and so

$$c_1(j^*\iota^*L_{\Gamma})^{\dim V} \cap [U]$$

= $\sum_{r=0}^{\dim V} {\dim V \choose r} \left[c_1(j^*\iota^*\phi_1^*L_{\Gamma_1})^r \cap \left(c_1(j^*\iota^*\phi_2^*L_{\Gamma_2})^{\dim V-r} \cap [U] \right) \right].$

Since $\phi_2 \circ \iota \circ j$ is flat,

$$c_1(j^*\iota^*\phi_2^*L_{\Gamma_2})^{\dim V-r} \cap [U] = j^*\iota^*\phi_2^*(c_1(L_{\Gamma_2})^{\dim V-r} \cap [U_2]) \in A_r(U),$$

where $j^* \iota^* \phi_2^*$ is the flat pullback $A_0(U_2) \to A_r(U)$. Therefore, since L_{Γ_2} is ample, these classes can be represented by non-negative sums of *r*-cycles (which are zero if r > d). Hence, since L_{Γ_1} is ample, each summand of the above sum can be represented by a non-negative sum of 0-cycles. In particular, $c_1(j^*\iota^*\phi_2^*L_{\Gamma_2})^{\dim V-d} \cap [U]$ can be represented by the cycle associated to finitely many fibres of $(\phi_2 \circ \iota \circ j)|_U$. For such a fibre *F*, we have

$$c_1(j^*\iota^*\phi_1^*L_{\Gamma_1})^d \cap [F] = c_1(j^*\iota^*L_{\Gamma})^d \cap [F] = \deg_{j^*\iota^*L_{\Gamma}}(F) = \deg_{L_{\Gamma}}(\iota(j(F)))$$

where the first equality can be deduced from a binomial expression as above. Therefore, since $(\phi_2 \circ \iota \circ j)|_U$ is flat, and all fibres of a flat family of subschemes of a projective space have the same Hilbert polynomial, we conclude that, if *W* is an irreducible component of $(\phi_2 \circ \iota \circ j)^{-1}(z)$ for some $z \in U_2$, then

$$\deg_{L_{\Gamma}}(W) \le \deg_{L_{\Gamma}}(V),$$

as claimed.

Therefore, by Lemma 29 (i), it remains to deal with the case when W is an irreducible component of $(\phi_2 \circ \iota)^{-1}(z)$ for some $z \in V_2 \setminus U_2$. To that end, let C be an irreducible algebraic curve in V_2 passing through z such that $C \cap U_2 \neq \emptyset$ (just choose a point in U_2 and use the fact that, for any two points x and y in an irreducible algebraic variety Z, there exists an irreducible algebraic curve $C \subset Z$ such that $x, y \in C$). Let Y denote an irreducible component of $(\phi_2 \circ \iota)^{-1}(C)$ containing W. Note that $W \subsetneq Y$ as

$$\dim Y \ge \dim V - \dim V_2 + 1 = d + 1 > \dim W$$

Therefore, the morphism $Y \to C$ is dominant. Let \tilde{C} denote the normalization of C, and let \tilde{Y} denote the fibre product $Y \times_C \tilde{C}$. Note that \tilde{Y} is irreducible and $\tilde{Y} \to \tilde{C}$ is dominant. Since \tilde{C} is regular, it follows from [16, Proposition 9.7] that $\tilde{Y} \to \tilde{C}$ is flat. As such, its fibres all have the same degree with respect to $\eta^* L_{\Gamma}$.

Let $\eta : \tilde{Y} \to Y$ denote the natural map. It is finite and surjective. Therefore, if we let \tilde{W} denote an irreducible component of $\eta^{-1}(W)$, we have

$$\deg_{L_{\Gamma}}(W) \le \deg_{\eta^* L_{\Gamma}}(W)$$

and, since \tilde{W} is a fibre of $\tilde{Y} \to \tilde{C}$ and C passes through U_2 , the claim follows.

The theorem now follows immediately, combining Lemmas 28 (i)–(iii) and 30. ■

9. Effective determination of the weakly optimal locus

Recall the setup described in Section 1.1. This section is devoted to the proof of Theorem 6.

Throughout the proof, we think of the family T as embedded in projective space with respect to the prescribed embedding. We begin by computing an affine cover $\{V_{\alpha}\}$ of V, such that over each V_{α} once can select an explicit set of sections $\omega_1, \ldots, \omega_g$ for the sheaf of relative differentials $\Omega^1_{T/V}$ and an additional set of meromorphic differentials of the second kind (i.e. with vanishing residues) $\omega_{g+1}, \ldots, \omega_{2g}$ for the sheaf $\Omega^1_{T/V}(N \cdot D)$ where D is a hyperplane divisor and $N \gg 1$, such that $\omega_1, \ldots, \omega_{2g}$ are pointwise linearly independent everywhere. Such a choice of differentials can be computed explicitly for (families of) algebraic curves by classical methods. For the holomorphic differentials, see e.g. [6, Theorem 9.3.1] where the explicit computation is attributed to Abel and Riemann. For meromorphic differentials of the second type see e.g. [6, Propositions 9.3.8, 9.3.9]. Below we continue with V replaced by one of the V_{α} , and assume that the sections above are pointwise linearly independent over V.

Having computed a base for the de Rham cohomology $H^1(T/V)$ we may now compute the Gauss–Manin connection $\nabla : H^1(T/V) \to H^1(T/V) \otimes \Omega_V^1$ explicitly. The fact that the Gauss–Manin connection admits a purely algebraic construction is essentially due to Manin [19]. The approach of Manin is fully explicit and there is no difficulty in principle carrying it out computationally.

The sections $\omega_1, \ldots, \omega_{2g}$ provide a trivialization of $H^1(T/V)$. Thinking of $V \times \mathbf{GL}_{2g}(\mathbb{C})$ as a principal $\mathbf{GL}_{2g}(\mathbb{C})$ -bundle with respect to the action $g(v, \Pi) = (v, \Pi g^{-1})$, we may express ∇ as a flat connection on this trivial bundle as follows:

$$d\Pi = \Omega \cdot \Pi, \quad \Omega \in \mathfrak{gl}_n(\Lambda^1_V). \tag{15}$$

In fact, the construction that follows could be expressed in terms of this $\mathbf{GL}_{2g}(\mathbb{C})$ connection. However, to stress the relationship with the general formalism of Shimura
varieties considered in the first part of the paper, we show that one can explicitly compute
a $\mathbf{GSp}_{2g}(\mathbb{C})$ -bundle $P \subset V \times \mathbf{GL}_{2g}(\mathbb{C})$ compatible with ∇ .

The existence of such a bundle *P* follows from the fact that ∇ preserves the symplectic form on $H^1(T/V)$ induced by duality from the intersection form on $H_1(T/V, \mathbb{Z})$. More explicitly, let $\delta_1(v), \ldots, \delta_{2g}(v)$ denote a (possibly multivalued) choice of symplectic basis of $H_1(T/V, \mathbb{Z})$, i.e. with the intersection form (δ_i, δ_j) given by

$$J = \begin{pmatrix} 0 & I_g \\ -I_g & 0 \end{pmatrix}.$$
 (16)

Then

$$\Pi(v) = \begin{pmatrix} \oint_{\delta_1(v)} \omega_1 & \cdots & \oint_{\delta_{2g}(v)} \omega_1 \\ \vdots & \ddots & \vdots \\ \oint_{\delta_1(v)} \omega_{2g} & \cdots & \oint_{\delta_{2g}(v)} \omega_{2g} \end{pmatrix} = \begin{pmatrix} A(v) & B(v) \\ C(v) & D(v) \end{pmatrix}$$
(17)

is a section of ∇ and $\Pi J \Pi^T = \Lambda$ defines a regular mapping from V to $\mathbf{GL}_{2g}(\mathbb{C})$ (singlevaluedness follows from the fact that the monodromy of $\Pi(v)$ respects the intersection form, and regularity then follows from GAGA). Thus the subset of $V \times \mathbf{GL}_{2g}(\mathbb{C})$ defined by $\Pi J \Pi^T = \Lambda$ is a ∇ -invariant principal $\mathbf{Sp}_{2g}(\mathbb{C})$ -bundle with respect to the action

$$g \cdot (v, \Pi) = (v, \Pi \cdot g^{-1}), \quad g \in \mathbf{Sp}_{2g}(\mathbb{C}).$$
(18)

However, as we will see below, this bundle is not defined over a number field, and we will show instead how to construct the corresponding $\mathbf{GSp}_{2g}(\mathbb{C})$ -bundle explicitly over a number field.

By definition, $\Lambda(v)_{ij} = (\omega_i, \omega_j)_v$ is the matrix representing the symplectic form on $H^1(X_v)$. The explicit computation of this form reduces to the following bilinear relation for meromorphic differentials of the second kind. **Lemma 31.** Let ω , η be two meromorphic differentials of the second kind. Then

$$\frac{1}{2\pi i} \left(\sum_{j=1}^{g} \oint_{\delta_{j}} \omega \oint_{\delta_{j+g}} \eta - \oint_{\delta_{j+g}} \omega \oint_{\delta_{j}} \eta \right) = \sum_{P} \operatorname{res}_{P}(f\eta)$$
(19)

where *P* ranges over the poles of ω and η , and *f* is any primitive of ω . Note that since η has no residues, res_{*P*}($f\eta$) is independent of the choice of the primitive.

The left hand side of (19) is, by definition, the symplectic pairing (ω, η) up to the constant $2\pi i$. The right hand side can be explicitly computed in local coordinates around $P \in V$, i.e. it depends only on finitely many Laurent coefficients of ω, η in local coordinates around the poles. Using this, one may explicitly compute each entry of $2\pi\Lambda(v)$ as a regular function on V.

Having computed $2\pi\Lambda(v)$, we further simplify the computation as follows. The Riemann bilinear relations imply that $\omega_1, \ldots, \omega_g$ span an isotropic space, and $\Lambda(v)$ defines a non-degenerate pairing between this space and the span of $\omega_{g+1}, \ldots, \omega_{2g}$. By elementary linear algebra, one may now replace each section ω_{g+j} by a linear combination ω'_{g+j} such that $\omega_1, \ldots, \omega_g$ and $\omega'_{g+1}, \ldots, \omega'_{2g}$ form a standard symplectic basis. Assume without loss of generality that we have made such a choice, so that $\Lambda(v) \equiv 2\pi J$. With this choice, ∇ restricts to a connection on the trivial bundle $P = V \times \mathbf{GSp}_{2g}(\mathbb{C})$ with the connection equation

$$d\Pi = \Omega \cdot \Pi, \quad \Omega \in \mathfrak{sp}_{2g}(\Lambda_V^1), \tag{20}$$

and the left $\mathbf{GSp}_{2g}(\mathbb{C})$ -action given as before by $g(s, \Pi) = (s, \Pi g^{-1})$.

Denote by $X = \mathcal{H}_g$ the Siegel upper half-space, by \check{X} the compact dual, and by $X' \subset \check{X}$ the set of symmetric $g \times g$ matrices. We have a $\mathbf{GSp}_{2g}(\mathbb{C})$ -equivariant rational map $\beta : P \to X'$ given by $\beta(v, \Pi) = B^{-1}A$ where A and B are the blocks given in (17). One can verify that, since $\Pi \in \mathbf{GSp}_{2g}(\mathbb{C})$, the image of β , when defined, is an element of X'. The map β extends to a regular map $\beta : P \to \check{X}$.

Recall that we denote by $f: \tilde{V} \to V$ an étale cover, $\tilde{T} \to \tilde{V}$ the base change of T by f, and choose f such that \tilde{T} is compatible with an N-level structure (say for N = 3). We denote by $\iota: \tilde{V} \to \mathcal{A}_{g,N}$ the corresponding moduli map.

Proposition 32. We have $f^*(P, \nabla) \simeq \iota^*(\mathcal{P}, \nabla_0)$ where \mathcal{P} denotes the canonical bundle on $\mathcal{A}_{g,N}$ and ∇_0 its canonical connection.

Proof. By functioniality of the Gauss–Manin connection, if $\tilde{\nabla}$ denotes the connection on $\tilde{P} \to \tilde{V}$ then $\tilde{\nabla} = f^* \nabla$. It will therefore suffice to prove that $(\tilde{P}, \tilde{\nabla}) \simeq \iota^*(\mathcal{P}, \nabla_0)$. Thus, we may assume, without loss of generality, that the family $T \to V$ already respects the N-level structure and that we have a map $\iota : V \to S = \mathcal{A}_{g,N}$. Denote by $\Gamma \subset \mathbf{GSp}_{2g}(\mathbb{Z})$ the neat subgroup corresponding to the N-level structure.

Choose a generic $v_0 \in V$ and consider the period map $\Pi(v)$ defined by (17) around v_0 . Since $\delta_1, \ldots, \delta_{2g}$ form a symplectic basis, the Riemann bilinear relations imply that $\Pi(v) \in X$ globally (i.e. after arbitrary analytic continuation). By definition of the moduli interpretation, ι is given by

$$\iota(v) = \pi(\beta(v, \Pi(v))), \quad \pi : X \to \Gamma \backslash X.$$
(21)

Indeed, $\Pi(v)$ is the period matrix of the fibre T_v . Hence, the Jacobian of T_v is given by the lattice spanned by the columns of $(A \ B)$, and $\beta(v, \Pi(v)) = B^{-1}(v)A(v) \in X$ is the point representing this Jacobian in X. In particular, write $x_0 = \beta(v_0, \Pi(v_0))$ so that $\pi(x_0) = \iota(v_0)$.

We will show that $(P, \nabla) \simeq \iota^*(\mathcal{P}, \nabla_0)$ by showing that they define the same Γ representation of $\pi_0(V, v_0)$. Let $\gamma \in \pi_0(V, v_0)$ be a closed loop, and let $\gamma_X \subset X$ be the
curve obtained by lifting γ to X. Then the endpoint of γ_X is a point $g \cdot x_0$ for some $g \in \Gamma$.
According to (21) and the equivariance of β , the monodromy of ∇ along γ is g, as it is
the unique element of Γ mapping x_0 to $g \cdot x_0$. On the other hand, the monodromy of $\iota^* \nabla_0$ along γ is the monodromy of ∇_0 along $\iota(\gamma)$, which equals g for the same reason. This
shows that the representations are indeed the same.

One can repeat the proof of Proposition 14 with the bundle (P, ∇) in place of (\mathcal{P}, ∇_0) to define sets $\Pi'(d) \subset P$. By Proposition 32 it follows that $f^*\Pi'(d) = \iota^*\Pi(d)$ over $f^{-1}(V)$ as claimed.

Acknowledgments. C. Daw would like to thank the University of Oxford for having him as a Visiting Research Fellow. Both authors would like to thank the anonymous referee for his/her comments.

Funding. G. Binyamini was supported by the ISRAEL SCIENCE FOUNDATION (grant No. 1167/17) and by funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 802107). This research was generously supported by Shimon and Golde Picker - Weizmann Annual Grant. C. Daw would like to thank the EPSRC for its support via a New Investigator Award (EP/S029613/1).

References

- Baily, W. L., Jr., Borel, A.: Compactification of arithmetic quotients of bounded symmetric domains. Ann. of Math. (2) 84, 442–528 (1966) Zbl 0154.08602 MR 216035
- [2] Barroero, F., Dill, G. A.: Distinguished categories and the Zilber–Pink cojecture. arXiv:2103.07422 (2021)
- [3] Binyamini, G.: Some effective estimates for André–Oort in Y(1)ⁿ. J. Reine Angew. Math. 767, 17–35 (2020) Zbl 1476.11091 MR 4160301
- [4] Binyamini, G., Masser, D.: Effective André–Oort for non-compact curves in Hilbert modular varieties. C. R. Math. Acad. Sci. Paris 359, 313–321 (2021) Zbl 1470.11159 MR 4245899
- [5] Binyamini, G., Novikov, D.: Multiplicity operators. Israel J. Math. 210, 101–124 (2015) Zbl 1336.32010 MR 3430270
- [6] Brieskorn, E., Knörrer, H.: Plane algebraic curves. Birkhäuser, Basel (1986) Zbl 0588.14019 MR 886476
- [7] Daw, C., Javanpeykar, A., Kühne, L.: Effective estimates for the degrees of maximal special subvarieties. Selecta Math. (N.S.) 26, art. 2, 31 pp. (2020) Zbl 1430.14057 MR 4041504
- [8] Daw, C., Orr, M.: Unlikely intersections with *E* × CM curves in A₂. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5) 22, 1705–1745 (2021) Zbl 1483.11122 MR 4360602

- [9] Daw, C., Ren, J.: Applications of the hyperbolic Ax–Schanuel conjecture. Compos. Math. 154, 1843–1888 (2018) Zbl 1427.11057 MR 3867286
- [10] Fulton, W.: Intersection theory. 2nd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete 2, Springer, Berlin (1998) Zbl 0085.14002 MR 1644323
- [11] Gabrielov, A., Khovanskii, A.: Multiplicity of a Noetherian intersection. In: Geometry of differential equations, American Mathematical Society Translations (2) 186, American Mathematical Society, Providence, RI, 119–130 (1998) Zbl 0916.32008 MR 1732408
- [12] Gao, Z.: Mixed Ax–Schanuel for the universal abelian varieties and some applications. Compos. Math. 156, 2263–2297 (2020) Zbl 1477.11118 MR 4190045
- [13] Grauert, H., Remmert, R.: Coherent analytic sheaves. Grundlehren der mathematischen Wissenschaften 265, Springer, Berlin (1984) Zbl 0537.32001 MR 755331
- [14] Green, M., Griffiths, P., Kerr, M.: Mumford–Tate groups and domains. Annals of Mathematics Studies 183, Princeton University Press, Princeton, NJ (2012) Zbl 1248.14001 MR 2918237
- [15] Habegger, P., Pila, J.: O-minimality and certain atypical intersections. Ann. Sci. École Norm. Sup. (4) 49, 813–858 (2016) Zbl 1364.11110 MR 3552014
- [16] Hartshorne, R.: Algebraic geometry. Graduate Texts in Mathematics 52, Springer, New York (1977) Zbl 0367.14001 MR 463157
- [17] Klingler, B., Yafaev, A.: The André–Oort conjecture. Ann. of Math. (2) 180, 867–925 (2014) Zbl 1377.11073 MR 3245009
- [18] Lawson, H. B., Jr.: Foliations. Bull. Amer. Math. Soc. 80, 369–418 (1974) Zbl 0293.57014 MR 343289
- [19] Manin, Yu. I.: Rational points on algebraic curves over function fields. Izv. Akad. Nauk SSSR Ser. Mat. 27, 1395–1440 (1963) (in Russian) Zbl 0178.55102 MR 157971
- [20] Milne, J. S.: Automorphic vector bundles on connected Shimura varieties. Invent. Math. 92, 91–128 (1988) Zbl 0684.14006 MR 931206
- [21] Milne, J. S.: Introduction to Shimura varieties. In: Harmonic analysis, the trace formula, and Shimura varieties, Clay Mathematics Proceedings 4, American Mathematical Society, Providence, RI, 265–378 (2005) Zbl 1148.14011 MR 2192012
- [22] Milne, J. S.: Modular functions and modular forms. https://www.jmilne.org/math/ CourseNotes/mf.html
- [23] Mok, N., Pila, J., Tsimerman, J.: Ax–Schanuel for Shimura varieties. Ann. of Math. (2) 189, 945–978 (2019) Zbl 1481.14048 MR 3961087
- [24] Moonen, B., Oort, F.: The Torelli locus and special subvarieties. In: Handbook of moduli. Vol. II, Advanced Lectures in Mathematics 25, International Press, Somerville, MA, 549–594 (2013) Zbl 1322.14065 MR 3184184
- [25] Pila, J., Scanlon, T.: Effective transcendental Zilber–Pink for variations of Hodge structures. arXiv:2105.05845 (2021)
- [26] Pink, R.: A combination of the conjectures of Mordell–Lang and André–Oort. In: Geometric methods in algebra and number theory, Progress in Mathematics 235, Birkhäuser Boston, Boston, MA, 251–282 (2005) Zbl 1200.11041 MR 2166087
- [27] Tsimerman, J.: The André–Oort conjecture for Ag. Ann. of Math. (2) 187, 379–390 (2018)
 Zbl 1415.11086 MR 3744855
- [28] Ullmo, E.: Equidistribution de sous-variétés spéciales. II. J. Reine Angew. Math. 606, 193– 216 (2007) Zbl 1137.11043 MR 2337648
- [29] Ullmo, E., Yafaev, A.: Galois orbits and equidistribution of special subvarieties: towards the André–Oort conjecture. Ann. of Math. (2) 180, 823–865 (2014) Zbl 1328.11070 MR 3245008
- [30] Urbanik D.: Sets of special subvarieties of bounded degree. Compos. Math. 159, 616–657 (2023) Zbl 1510.14012 MR 4560541