

# Existence of solutions for a class of fractional Kirchhoff variational inequality

Shenbing Deng, Wenshan Luo, César E. Torres Ledesma, and  
George W. Alama Quiroz

**Abstract.** We are concerned with the following fractional Kirchhoff variational inequality:

$$\begin{aligned} & (a + b[u]^2) \int_{\mathbb{R}^3} (-\Delta)^{\frac{s}{2}} u (-\Delta)^{\frac{s}{2}} (v - u) dx + \int_{\mathbb{R}^3} (1 + \lambda V(x)) u (v - u) dx \\ & \geq \int_{\mathbb{R}^3} f(u) (v - u) dx \quad \forall v \in \mathbb{K}, \end{aligned}$$

where  $s \in (\frac{3}{4}, 1)$ ,  $\lambda > 0$ . In this paper, by applying penalization techniques from Bensoussan and Lions (1978) combined with mountain pass theorem, we show the existence and concentration behavior of positive solution to the cited variational inequality. This result extend some results established by Alves, Barros and Torres [J. Math. Anal. Appl. 494 (2021)] to the fractional case.

## 1. Introduction

In this paper, we focus our attention on the following fractional Kirchhoff inequality, for  $u \in \mathbb{K}$ :

$$\begin{aligned} & (a + b[u]^2) \int_{\mathbb{R}^3} (-\Delta)^{\frac{s}{2}} u (-\Delta)^{\frac{s}{2}} (v - u) dx + \int_{\mathbb{R}^3} (1 + \lambda V(x)) u (v - u) dx \\ & \geq \int_{\mathbb{R}^3} f(u) (v - u) dx \quad \forall v \in \mathbb{K}, \end{aligned} \tag{1.1}$$

where  $s \in (\frac{3}{4}, 1)$ ,  $\lambda > 0$ ,

$$[u]^2 = \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(u(x) - u(y))^2}{|x - y|^{3+2s}} dx dy,$$

and the function  $V: \mathbb{R}^3 \rightarrow \mathbb{R}$  is a continuous potential verifying the following assumptions:

(V<sub>1</sub>)  $V(x) \geq 0 \forall x \in \mathbb{R}^3$ ;

(V<sub>2</sub>) there exists an open, connected and bounded domain  $\Omega \subset \mathbb{R}^3$  with smooth boundary such that  $\Omega := \text{int}(V^{-1}(\{0\})) \neq \emptyset$ ;

---

2020 Mathematics Subject Classification. Primary 35A15; Secondary 35J86, 49J40.

Keywords. Fractional Kirchhoff variational inequality, variational methods, critical nonlinearity.

(V<sub>3</sub>) there exists  $M_0 > 0$  such that the set  $\mathcal{L} = \{x \in \mathbb{R}^3 : V(x) \leq M_0\}$  is nonempty and  $|\mathcal{L}| < \infty$ , where  $|A|$  denotes the Lebesgue measure of  $A$  on  $\mathbb{R}^3$ .

Letting

$$E := \left\{ u \in H^s(\mathbb{R}^3); \int_{\mathbb{R}^3} V(x)|u|^2 dx < \infty \right\},$$

we define

$$\mathbb{K} := \{v \in E; v \geq \varphi \text{ a.e. in } \Omega\},$$

with  $\varphi \in H^s(\mathbb{R}^3)$ ,  $\varphi^+ \neq 0$  and  $\text{supp}(\varphi^+) \subset \Omega$ . Moreover, on the nonlinearity  $f$ , we require that

$$f(t) = \mu t^{q-1} + t^{2_s^*-1}$$

for any  $t > 0$ , with  $4 < q < 2_s^*$ ,  $\mu > 0$  and  $f$  vanishes in  $(-\infty, 0)$ .

Variational inequalities are well known in the literature of applied mathematics and lead to many applications. According to Rodrigues [32], the theory of variational inequalities born in Italy in the sixties with the work of Fichera in 1963 on the elasticity problem and the work of Stampacchia in 1964 in the frame of potential theory in connection with capacity. The classical example of a variational problem consists of an elastic membrane, with vertical displacement  $u$  on a domain  $\Omega$  with  $u = u_0$  along  $\partial\Omega$  and it is forced to lie below some obstacle, that is,  $u \leq \psi$ . Then, at the equilibrium, whenever the membrane does not touch the obstacle, the elasticity provides a balance of the tension of the membrane that, geometrically, reflects into a balance of the principal curvatures of the surface described by  $u$ . On the other hand, when the membrane sticks to the obstacle, its principal curvatures are expected to adapt to those of  $\psi$ . Moreover, if an external force  $-f$  is switched on, the rest configuration of the membrane will be such that the elastic tension of the membrane equilibrates the force. These physical considerations lead to the classical variational inequality

$$\int_{\Omega} \nabla u(\nabla v - \nabla u) dx \geq \int_{\Omega} f(x)(v(x) - u(x)) dx \quad (1.2)$$

for any test function  $v$ , with  $v \leq \psi$  and  $v = u_0$  along  $\partial\Omega$ .

Many extensions of this problem have been considered in the literature, particularly for taking into account nonlinear elastic reactions of the membrane, non-commutative effects and nonlocal interactions [3, 4, 9, 10, 16, 25, 26, 28–30]. When replacing the local elastic reaction with a nonlocal one, with the purpose of taking into account the long-range interactions of particles, for instance, the standard Laplacian  $-\Delta$  might be replaced with the fractional Laplacian  $(-\Delta)^s$ , equation (1.2) becomes

$$\begin{aligned} & \iint_{\mathbb{R}^{2N} \setminus \mathcal{Q}} \frac{[u(x) - u(y)][v(x) - v(y) - u(x) + u(y)]}{|x - y|^{N+2s}} dx dy \\ & \geq \int_{\Omega} f(x)(v(x) - u(x)) dx, \end{aligned}$$

where  $\mathcal{Q} = (\mathbb{R}^N \setminus \Omega) \times (\mathbb{R}^N \setminus \Omega)$ . This kind of variational problems has been extensively studied in [9, 19, 33, 34, 37] and the references therein.

Recently, Alves, Barros and Torres Ledesma [3] pointed out there are three methods for researching variational inequalities: the nonsmooth critical point theory [12, 13, 23, 27], the minimax principles [17, 38, 39] and the penalization method [8, 28, 29]. Moreover, they established the existence of positive solutions for the variational inequality

$$\begin{cases} u \in \mathbb{K}, \\ \int_{\mathbb{R}^3} \nabla u \nabla (v - u) dx + \int_{\mathbb{R}^3} (1 + \lambda V(x)) u (v - u) dx \\ \geq \int_{\mathbb{R}^3} f(u) (v - u) dx \quad \forall v \in \mathbb{K}, \end{cases} \quad (1.3)$$

where the definition of these letters in this inequality is similar to our paper when  $s = 1$ .

Elliptic problems with critical growth like

$$-\Delta u + \lambda V(x)u = \mu u^{q-1} + u^{2^*-1} \quad \text{in } \mathbb{R}^3 \quad (1.4)$$

have been extensively studied by many authors. In fact, Alves and Barros [2] considered the existence and multiplicity for (1.4), under  $V(x)$  satisfies  $(V_1)$ – $(V_3)$ . Based on the important research in [2], Alves et al. [3, 4] solved (1.3) by using the penalization method under different conditions on  $V$ . Moreover, as for a class of problems where  $\lambda$  is large enough to get the existence result in (1.4), we refer to [7, 11, 14] and references therein.

The equation related to the variational inequality (1.1) is given by

$$(a + b[u]^2)(-\Delta)^s u + (1 + \lambda V(x))u = \mu |u|^{q-2}u + |u|^{2^*_s-2}u. \quad (1.5)$$

The Kirchhoff part of problem (1.5) is due to the work of Kirchhoff [24], in which, in 1883, he studied the hyperbolic equation

$$\rho \frac{\partial^2 u}{\partial t^2} - \left( \frac{p_0}{h} + \frac{E}{2L} \int_0^L |u_x|^2 dx \right) u_{xx} = 0$$

that extends the classical D'Alembert wave equation, by considering the effects of the changes in the length of the strings during the vibrations. As  $s \in (\frac{3}{4}, 1)$ , the fractional Laplacian  $(-\Delta)^s$  is defined by

$$(-\Delta)^s \Psi(x) = C(3, s) \text{P.V.} \int_{\mathbb{R}^3} \frac{\Psi(x) - \Psi(y)}{|x - y|^{3+2s}} dy, \quad \Psi \in \mathcal{S}(\mathbb{R}^3),$$

where P.V. stands for the Cauchy principle value and  $\mathcal{S}$  is the Schwartz space of rapidly decaying functions. The operator  $(-\Delta)^s$  can be also defined via Fourier transform and viewed as a pseudo-differential operator of symbol  $|\xi|^{2s}$ , see [5], that is,

$$\mathcal{F}((-\Delta)^s u)(\xi) = |\xi|^{2s} \mathcal{F}(u)(\xi) \quad \text{for } \xi \in \mathbb{R}^3,$$

where  $\mathcal{F}$  denotes the Fourier transform, i.e., for function  $w$  in the Schwartz class,

$$\mathcal{F}(w)(\xi) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int_{\mathbb{R}^N} e^{-i\xi x} w(x) dx.$$

When  $s = 1$ , Alimohammady et al. [1] studied the multiplicity of positive solutions for a Kirchhoff type problem, under conditions  $(V_1)$ – $(V_3)$  and  $a = 1$  in (1.5). Besides, Fan [15] proved multiple positive solutions of a Kirchhoff type problem on a bounded domain, when there are competing potentials in (1.5) and  $s = 1$ . For more Kirchhoff problems, we refer to [21, 22] and references therein.

When  $s \in (0, 1)$ , in [18], Fiscella and Valdinoci researched the existence of solutions in (1.5), when there is no potential and replace the nonlinearity with  $\lambda f(u)$  in (1.5). Moreover, Chen considered existence of a fractional  $p$ -Kirchhoff type problem, when there is  $\lambda$  in the generalized Choquard nonlinearity. For more fractional Kirchhoff equation, we refer to [6, 31, 36] and so on. Moreover, Frites and Moussaoui in [20] and Zuo et al. in [40] have explored the variational Kirchhoff inequality based on the nonsmooth critical point theory due to Szulkin.

Since we did not find in the literature any paper dealing with the existence of non-negative solutions for problem (1.1) in  $\mathbb{R}^N$ , motivated by the previous exposition and by the ideas of [1, 4], in the present paper, we intend to prove that (1.1) has a nontrivial weak solution. To the best of our knowledge, it is the first time to study fractional Kirchhoff variational inequality by using the penalization method.

The main result of this paper can be stated as follows.

**Theorem 1.1.** *Suppose that  $V$  satisfies  $(V_1)$ – $(V_3)$ . Then there exist  $\lambda_* > 0$ ,  $\mu_* > 0$  and  $b_* > 0$ , problem (1.1) has at least one nontrivial weak solution  $u_\lambda$  for  $\lambda \geq \lambda_*$ ,  $\mu \geq \mu_*$  and  $b < b_*$ . Furthermore, for any sequence  $\lambda_n \rightarrow +\infty$ , there exists a subsequence, still denoted by  $\{\lambda_n\}$ , such that  $\{u_{\lambda_n}\}$  converges strongly in  $H^s(\mathbb{R}^3)$  to a function  $u$  with  $u = 0$  a.e. in  $\Omega^c$ , where  $u$  is a solution of the following variational inequality:*

$$\begin{aligned} & \left( a + b \int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{3+2s}} dx dy \right) \\ & \quad \times \int_Q \frac{(u(x) - u(y))((v - u)(x) - (v - u)(y))}{|x - y|^{3+2s}} dx dy \\ & \quad + \int_\Omega u(v - u) dx \\ & \geq \int_\Omega (\mu|u|^{q-2} + |u|^{2_s^*-2})u(v - u) dx, \end{aligned}$$

where  $Q := \mathbb{R}^6 \setminus (\Omega^c \times \Omega^c)$  and, for every  $v \in \tilde{\mathbb{K}}$ , where

$$\tilde{\mathbb{K}} = \{v \in H_0^s(\Omega); v \geq \varphi \text{ a.e. in } \Omega\}.$$

In order to prove Theorem 1.1, we use penalization techniques due to Bensoussan and Lions [8], that is, considering problem (1.1), we introduce the penalized problem defined as

$$\begin{aligned} & (a + b[u^+]^2)(-\Delta)^s u + (1 + \lambda V(x))u - \frac{1}{\varepsilon}(\varphi - u)^+ \chi_\Omega \\ & = \mu(u^+)^{q-1} + (u^+)^{2_s^*-1} \quad \text{in } \mathbb{R}^3, \end{aligned} \tag{1.6}$$

where  $\epsilon > 0$ . Associated to (1.6), we have the energy functional

$$J_{\lambda,\epsilon}(u) = \frac{1}{2}\|u\|_{\lambda}^2 + \frac{b}{4}[u]^4 + \frac{1}{2\epsilon} \int_{\Omega} [(\varphi - u)^+]^2 dx - \int_{\mathbb{R}^3} F_+(u) dx,$$

where

$$\begin{aligned} P: E_{\lambda} &\rightarrow E'_{\lambda}, \\ u &\rightarrow \langle P(u), v \rangle = - \int_{\Omega} (\varphi - u)^+ v dx \end{aligned}$$

is the penalization operator. Then, by applying mountain pass arguments, we study the existence of at least one critical point of the functional  $J_{\lambda,\epsilon}$  which is a weak solution of penalized problem (1.6). By taking  $\epsilon = \frac{1}{n}$  with  $n$  large enough, we denote this mountain pass solution as  $u_n$  and we show that  $(u_n)$  is bounded in  $E_{\lambda}$ ; hence, up to a subsequence, there is  $u \in E_{\lambda}$  such that

$$u_n \rightharpoonup u \quad \text{in } E_{\lambda}.$$

This limit is such that  $P(u) = 0$  and hence a weak solution of problem (1.1).

**Remark 1.2.** Compared with the previous results, Theorem 1.1 can be regarded as an extension of [3, Theorem 1.1] under the fractional Kirchhoff situation. Due to the Kirchhoff term and the critical term, we cannot ensure in a standard way that weak limits of a bounded Palais–Smale sequence of the energy functional are critical points of it. Besides, if we change this inequality into an equation, a penalization term will be generated naturally. Therefore, we need to get some restrictions on  $b$  for our compactness.

The paper is organized as follows. In Section 2, we give some preliminaries about definitions and properties of the function space. In Section 3, we study the penalized problem (1.6). In Section 4, we get Theorem 1.1.

**Notation.** • The letter  $C$  changes from line to line.

- $B_r(x)$  denotes the ball in  $\mathbb{R}^3$  centered at  $x \in \mathbb{R}^3$  with radius  $r$ .
- $\Gamma^c$  means  $\mathbb{R}^3 \setminus \Gamma$ , where  $\Gamma \subset \mathbb{R}^3$ .
- $|\cdot|_r$  means the norm in  $L^r(\mathbb{R}^3)$  and  $|u|_{r(M)} := (\int_M |u|^r dx)^{\frac{1}{r}}$ , where  $M \subset \mathbb{R}^3$  and  $u \in L^r(M)$ .

## 2. Abstract setting and preliminary results

In this preliminary section, we fix the notation and we recall some technical results. For  $s \in (\frac{3}{4}, 1)$ , we define the fractional Sobolev space  $H^s(\mathbb{R}^3)$  as

$$H^s(\mathbb{R}^3) := \{u \in L^2(\mathbb{R}^3) : [u] < \infty\},$$

where  $[u]$  is the so-called Gagliardo semi-norm of  $u$ . It is well known that  $H^s(\mathbb{R}^3)$  is a Hilbert space, under the norm

$$\|u\|^2 := \iint_{\mathbb{R}^6} \frac{|u(x) - u(y)|^2}{|x - y|^{3+2s}} dx dy + \int_{\mathbb{R}^3} |u|^2 dx.$$

We denote by  $D^{s,2}(\mathbb{R}^3)$  the completion of  $C_c^\infty(\mathbb{R}^3)$  as

$$D^{s,2}(\mathbb{R}^3) = \{u \in L^{2_s^*}(\mathbb{R}^3) : [u] < \infty\}.$$

When  $\lambda > 0$ , we denote by  $E_\lambda$  the Hilbert space

$$E_\lambda := \left\{ u \in H^s(\mathbb{R}^3) : \int_{\mathbb{R}^3} V(x)u^2(x) dx < \infty \right\},$$

endowed with the norm

$$\|u\|_\lambda^2 := a[u]^2 + \int_{\mathbb{R}^3} (1 + \lambda V(x))u^2 dx.$$

The scalar product in  $E_\lambda$  is, for  $v \in E_\lambda$ ,

$$\langle u, v \rangle_\lambda := a \iint_{\mathbb{R}^6} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{3+2s}} dx dy + \int_{\mathbb{R}^3} (1 + \lambda V(x))uv dx.$$

By [5], since  $E_\lambda \subset H^s(\mathbb{R}^3)$ , we can obtain the embedding  $E_\lambda \rightarrow L^r(\mathbb{R}^3)$  is continuous for all  $r \in [2, 2_s^*]$  and locally compact for all  $r \in [1, 2_s^*)$ . The following Sobolev inequality can be found in [5, Theorem 1.1.8]: for all  $u \in D^{s,2}(\mathbb{R}^3)$ , there exists  $S > 0$  such that

$$S|u|_{2_s^*}^2 \leq [u]^2.$$

As in [4, 8], we get the penalized problem of (1.1),

$$\begin{aligned} & (a + b[u]^2)(-\Delta)^s u + (1 + \lambda V(x))u - \frac{1}{\varepsilon}(\varphi - u)^+ \chi_\Omega \\ & = \mu(u^+)^{q-1} + (u^+)^{2_s^*-1} \quad \text{in } \mathbb{R}^3, \end{aligned} \quad (2.1)$$

where  $u^+ = \max\{u, 0\}$  and  $\varepsilon > 0$  is the penalization parameter. Let

$$\langle P(u), v \rangle = - \int_{\Omega} (\varphi - u)^+ v dx.$$

We know  $P$  is the penalty operator and  $\frac{1}{\varepsilon} \int_{\Omega} (\varphi - u)^+ v dx$  is the penalization term. The associated energy functional associated to problem (2.1) is

$$J_{\lambda,\varepsilon}(u) = \frac{1}{2}\|u\|_\lambda^2 + \frac{b}{4}[u]^4 + \frac{1}{2\varepsilon} \int_{\Omega} [(\varphi - u)^+]^2 dx - \int_{\mathbb{R}^3} F_+(u) dx,$$

where

$$F_+(t) = \int_0^t f_+(s) ds \quad \text{and} \quad f_+(t) = \mu(t^+)^{q-1} + (t^+)^{2_s^*-1} \quad \forall t \in \mathbb{R}.$$

It is easy to check that  $J_{\lambda,\varepsilon} \in C^1(E_\lambda, \mathbb{R})$  and its differential is defined as, for any  $v \in E_\lambda$ ,

$$\begin{aligned} J'_{\lambda,\varepsilon}(u)v & = \langle u, v \rangle_\lambda + b[u]^2 \iint_{\mathbb{R}^6} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{3+2s}} dx dy \\ & \quad - \frac{1}{\varepsilon} \int_{\Omega} (\varphi - u)^+ v dx - \int_{\mathbb{R}^3} f_+(u)v dx. \end{aligned}$$

### 3. The penalized problem

We stress that, by the definition of  $f$ ,  $J_{\lambda,\epsilon}$  possesses a mountain pass geometry [35].

**Lemma 3.1.** (i) *There exist constants  $r, \rho > 0$ , independent of  $\lambda$  and  $\epsilon$ , such that*

$$J_{\lambda,\epsilon}(u) \geq \rho \quad \text{for } \|u\|_\lambda = r.$$

(ii) *There is  $e \in E_\lambda$  with  $\|e\|_\lambda > r$  and  $J_{\lambda,\epsilon}(e) < 0$ .*

*Proof.* (i) By Sobolev embeddings, we derive that

$$J_{\lambda,\epsilon}(u) \geq \frac{1}{2}\|u\|_\lambda^2 - \frac{C_1\mu}{q}\|u\|_\lambda^q - \frac{C_2}{2_s^*}\|u\|_\lambda^{2_s^*}.$$

As  $4 < q < 2_s^*$ , if we choose  $r > 0$  satisfying

$$r < \min\left\{\left(\frac{3 \cdot 2_s^*}{8C_2}\right)^{\frac{1}{2_s^*-2}}, \left(\frac{3q}{8C_1\mu}\right)^{\frac{1}{q-2}}\right\},$$

we obtain

$$J_{\lambda,\epsilon}(u) \geq \frac{1}{8}r^2 := \rho \quad \text{for } \|u\|_\lambda = r.$$

(ii) Since  $\text{supp}(\varphi^+) \subset \Omega$ , then we have

$$J_{\lambda,\epsilon}(\varphi^+) \leq \frac{1}{2}\|\varphi^+\|^2 + \frac{C}{4}\|\varphi^+\|^4 \leq \frac{1}{2}\|\varphi\|^2 + \frac{C}{4}\|\varphi\|^4,$$

where  $C := \frac{b}{a}$ . Choose  $\|\varphi\|$  small enough such that

$$\frac{1}{2}\|\varphi\|^2 + \frac{C}{4}\|\varphi\|^4 < \rho,$$

which implies that  $J_{\lambda,\epsilon}(\varphi^+) < \rho$  and

$$\begin{aligned} J_{\lambda,\epsilon}(t\varphi^+) &= \frac{t^2}{2}\|\varphi^+\|_\lambda^2 + \frac{bt^4}{4}[\varphi^+]^4 \\ &\quad - \frac{t^q\mu}{q} \int_\Omega |\varphi^+|^q dx - \frac{t^{2_s^*}}{2_s^*} \int_\Omega |\varphi^+|^{2_s^*} dx \quad \text{for } t > 1. \end{aligned}$$

As  $4 < q < 2_s^*$ , it follows that  $\lim_{t \rightarrow \infty} J_{\lambda,\epsilon}(t\varphi^+) = -\infty$ . Hence, taking  $e := (1 + t_0)\varphi^+$  for  $t_0$  large enough, we get  $\|e\|_\lambda > r$  and  $J_{\lambda,\epsilon}(e) < 0$ .  $\blacksquare$

From [35, Theorem 1.15], there is a (PS) $_c$  sequence  $\{u_n\} \subset E_\lambda$  such that  $J_{\lambda,\epsilon}(u_n) \rightarrow c_{\lambda,\epsilon}$  and  $J'_{\lambda,\epsilon}(u_n) \rightarrow 0$ , where  $c_{\lambda,\epsilon}$  is the mountain pass level characterized by

$$c_{\lambda,\epsilon} = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} J_{\lambda,\epsilon}(\gamma(t)),$$

and

$$\Gamma = \{\gamma \in C([0, 1], E_\lambda) : \gamma(0) = \varphi^+ \text{ and } \gamma(1) = e\}.$$

We can see the following results.

**Lemma 3.2.** *The  $(\text{PS})_{c_{\lambda,\epsilon}}$  sequence  $\{u_n\}$  of  $J_{\lambda,\epsilon}$  is bounded in  $E_\lambda$ .*

*Proof.* Let  $\{u_n\} \subset E_\lambda$  be a  $(\text{PS})_{c_{\lambda,\epsilon}}$  sequence for  $J_{\lambda,\epsilon}$ , that is,

$$J_{\lambda,\epsilon}(u_n) \rightarrow c \quad \text{and} \quad J'_{\lambda,\epsilon}(u_n) \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

Since  $[(\varphi - u_n)^+]^2 + (\varphi - u_n)^+ u_n \geq (\varphi - u_n)^+ \varphi$ , as  $q > 4$ , then

$$\frac{1}{2\epsilon} \int_{\Omega} [(\varphi - u_n)^+]^2 dx + \frac{1}{q\epsilon} \int_{\Omega} (\varphi - u_n)^+ u_n dx \geq \frac{1}{q\epsilon} \int_{\Omega} (\varphi - u_n)^+ \varphi dx.$$

Therefore, from Hölder inequality and Sobolev embedding,

$$\begin{aligned} J_{\lambda,\epsilon}(u_n) - \frac{1}{q} J'_{\lambda,\epsilon}(u_n) u_n &\geq \left(\frac{1}{2} - \frac{1}{q}\right) \|u_n\|_\lambda^2 + \left(\frac{1}{4} - \frac{1}{q}\right) b[u_n]^4 \\ &\quad + \frac{1}{q\epsilon} \int_{\Omega} (\varphi - u_n)^+ \varphi dx + \left(\frac{1}{q} - \frac{1}{2_s^*}\right) \int_{\mathbb{R}^3} |u_n^+|^{2_s^*} dx \\ &\geq \left(\frac{1}{2} - \frac{1}{q}\right) \|u_n\|_\lambda^2 - \frac{1}{q\epsilon} \int_{\Omega} (|\varphi| + |u_n|) |\varphi| dx \\ &\geq \frac{q-2}{2q} \|u_n\|_\lambda^2 - \frac{1}{q\epsilon} |\varphi|_2^2 - \frac{1}{q\epsilon} |\varphi|_2 \|u_n\|_\lambda, \end{aligned}$$

which yields

$$\frac{q-2}{2q} \|u_n\|_\lambda^2 \leq c_{\lambda,\epsilon} + \frac{1}{q\epsilon} |\varphi|_2^2 + o_n(1) + (o_n(1) + \frac{1}{q\epsilon} |\varphi|_2) \|u_n\|_\lambda.$$

Thus,  $\{u_n\}$  is bounded in  $E_\lambda$ . ■

**Lemma 3.3.** *Given  $\tau > 0$ , there is  $\mu_* = \mu_*(\tau) > 0$  such that*

$$c_{\lambda,\epsilon} < \frac{q-2}{2q} S^{\frac{3}{2_s^*}} - \tau \quad \text{for all } \lambda > 0, \epsilon > 0 \quad \text{and} \quad \mu \geq \mu_*.$$

*Proof.* Firstly, we fix the path  $\gamma \in \Gamma$ ,

$$\begin{aligned} \gamma: [0, 1] &\rightarrow E_\lambda, \\ t &\mapsto \gamma(t) = (1 + tt_0)\varphi^+, \end{aligned}$$

where  $t_0$  is defined as in Lemma 3.1. Since  $(1 + tt_0)\varphi^+ \geq \varphi$ , we get

$$\int_{\Omega} [(\varphi - (1 + tt_0)\varphi^+)^+]^2 dx = 0,$$

and then

$$\begin{aligned} J_{\lambda,\epsilon}(\gamma(t)) &= \frac{(1 + tt_0)^2}{2} \|\varphi^+\|_\lambda^2 + \frac{b(1 + tt_0)^4}{4} [\varphi^+]^4 \\ &\quad - \frac{\mu(1 + tt_0)^q}{q} \int_{\Omega} |\varphi^+|^q dx - \frac{(1 + tt_0)^{2_s^*}}{2_s^*} \int_{\Omega} |\varphi^+|^{2_s^*} dx \\ &\leq \frac{(1 + tt_0)^2}{2} \|\varphi^+\|_\lambda^2 + \frac{b(1 + tt_0)^4}{4} [\varphi^+]^4 - \frac{\mu(1 + tt_0)^q}{q} \int_{\Omega} |\varphi^+|^q dx. \end{aligned}$$



Taking  $A := \frac{1}{2}\|\varphi^+\|^2$ ,  $B := \frac{b[\varphi^+]^4}{4}$  and  $C := \frac{1}{q} \int_{\mathbb{R}^3} |\varphi^+|^q dx$ , then we consider the following function:

$$g_\mu(t) := At^2 + Bt^4 - \mu Ct^q \quad \text{for all } t \in [0, \infty).$$

From the second derivative of  $g_\mu(t)$ , for any  $\mu > 0$ ,  $g_\mu(t)$  is bounded up. Let the maximum point of  $g_\mu$  be  $t_{\max} > 0$ ; thus

$$\begin{aligned} c_{\lambda, \epsilon} &\leq \max_{t \geq 0} J_{\lambda, \epsilon}(\gamma(t)) \\ &\leq \max_{t \geq 0} \left\{ \frac{(1 + tt_0)^2}{2} \|\varphi^+\|^2 + \frac{b(1 + tt_0)^4}{4} [\varphi^+]^4 - \frac{\mu(1 + tt_0)^q}{q} \int_{\Omega} |\varphi^+|^q dx \right\} \\ &= \max_{t \geq 0} g_\mu(t) = At_{\max}^2 + Bt_{\max}^4 - \mu Ct_{\max}^q < \frac{q-2}{2q} S^{\frac{3}{2s}} - \tau \end{aligned}$$

for all  $\lambda > 0$ ,  $\epsilon > 0$  and  $\mu \geq \mu_*$ , where  $\mu_*$  is large enough. ■

**Lemma 3.4.**  $\{u_n^+\}$  is also a  $(PS)_{c_{\lambda, \epsilon}}$  sequence of  $J_{\lambda, \epsilon}$ .

*Proof.* Let  $u_n^- = \min\{u_n, 0\}$ , then  $u_n = u_n^+ + u_n^-$  and

$$-\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{u_n^+(x)u_n^-(y) + u_n^+(y)u_n^-(x)}{|x-y|^{3+2s}} dx dy \geq 0. \quad (3.1)$$

By Lemma 3.2,  $\{u_n^-\}$  is bounded, then from (3.1),

$$\begin{aligned} o_n(1) &= J'_{\lambda, \epsilon}(u_n)u_n^- \\ &= \langle u_n, u_n^- \rangle_\lambda + b[u_n]_b^2 \iint_{\mathbb{R}^6} \frac{(u_n(x) - u_n(y))(u_n^-(x) - u_n^-(y))}{|x-y|^{3+2s}} dx dy \\ &\quad - \frac{1}{\epsilon} \int_{\Omega} (\varphi - u_n)^+ u_n^- dx \\ &\geq \|u_n^-\|_\lambda^2, \end{aligned}$$

which implies that  $\|u_n^-\|_\lambda^2 = o_n(1)$ . Up to a subsequence, we get

$$\begin{aligned} u_n^- &\rightharpoonup 0 \quad \text{in } E_\lambda, \\ u_n^- &\rightarrow 0 \quad \text{in } L_{\text{loc}}^r(\mathbb{R}^3) \quad \text{for } r \in [1, 2^*), \\ u_n^-(x) &\rightarrow 0 \quad \text{a.e. in } \mathbb{R}^3. \end{aligned}$$

Then we note that  $\|u_n\|_\lambda = \|u_n^+\|_\lambda + o_n(1)$ . Besides, by Lebesgue dominated convergence theorem,

$$\int_{\Omega} [(\varphi - u_n)^+]^2 dx = \int_{\Omega} [(\varphi - u_n^+)]^2 dx + o_n(1).$$

Then

$$\begin{aligned}
 J_{\lambda,\varepsilon}(u_n) &= \frac{1}{2}\|u_n\|_\lambda^2 + \frac{b}{4}[u_n]^4 + \frac{1}{2\varepsilon} \int_\Omega [(\varphi - u_n)^+]^2 dx \\
 &\quad - \frac{\mu}{q} \int_{\mathbb{R}^3} |u_n^+|^q dx - \frac{1}{2_s^*} \int_{\mathbb{R}^3} |u_n^+|^{2_s^*} dx \\
 &= \frac{1}{2}\|u_n^+\|_\lambda^2 + \frac{b}{4}[u_n^+]^4 + \frac{1}{2\varepsilon} \int_\Omega [(\varphi - u_n^+)^+]^2 dx \\
 &\quad - \frac{\mu}{q} \int_{\mathbb{R}^3} |u_n^+|^q dx - \frac{1}{2_s^*} \int_{\mathbb{R}^3} |u_n^+|^{2_s^*} dx + o_n(1) \\
 &= J_{\lambda,\varepsilon}(u_n^+) + o_n(1).
 \end{aligned}$$

Hence

$$J_{\lambda,\varepsilon}(u_n^+) = c_{\lambda,\varepsilon} + o_n(1).$$

Following, we are going to show  $J'_{\lambda,\varepsilon}(u_n^+) = o_n(1)$ . Now let  $v \in E_\lambda$  with  $\|v\|_\lambda \leq C$  for some  $C > 0$ . By Lebesgue dominated convergence theorem, we have

$$\int_\Omega (\varphi - u_n)^+ v dx = \int_\Omega (\varphi - u_n^+)^+ v dx + o_n(1),$$

and

$$\int_{\mathbb{R}^3} (1 + \lambda V(x)) u_n^- v dx = o_n(1).$$

Thus,

$$\begin{aligned}
 &\iint_{\mathbb{R}^6} \frac{(u_n^-(x) - u_n^-(y))(v(x) - v(y))}{|x - y|^{3+2s}} dx dy \\
 &= \langle u_n^-, v \rangle_\lambda - \int_{\mathbb{R}^3} (1 + \lambda V(x)) u_n^- v dx = o_n(1).
 \end{aligned}$$

Therefore, from Hölder inequality and Lemma 3.2, we obtain

$$\begin{aligned}
 &J'_{\lambda,\varepsilon}(u_n)v \\
 &= \langle u_n^+, v \rangle_\lambda + \langle u_n^-, v \rangle_\lambda + b[u_n]^2 \iint_{\mathbb{R}^6} \frac{(u_n(x) - u_n(y))(v(x) - v(y))}{|x - y|^{3+2s}} dx dy \\
 &\quad - \frac{1}{\varepsilon} \int_\Omega (\varphi - u_n)^+ v dx - \mu \int_{\mathbb{R}^3} (u_n^+)^{q-1} v dx - \int_{\mathbb{R}^3} (u_n^+)^{2_s^*-1} v dx \\
 &= \langle u_n^+, v \rangle_\lambda + \langle u_n^-, v \rangle_\lambda \\
 &\quad + b([u_n^+]^2 + o_n(1)) \left( \iint_{\mathbb{R}^6} \frac{(u_n^+(x) - u_n^+(y))(v(x) - v(y))}{|x - y|^{3+2s}} dx dy + o_n(1) \right) \\
 &\quad - \frac{1}{\varepsilon} \int_\Omega (\varphi - u_n^+)^+ v dx - \mu \int_{\mathbb{R}^3} (u_n^+)^{q-1} v dx - \int_{\mathbb{R}^3} (u_n^+)^{2_s^*-1} v dx + o_n(1) \\
 &= J'_{\lambda,\varepsilon}(u_n^+)v + \langle u_n^-, v \rangle_\lambda + o_n(1),
 \end{aligned}$$

from where we get

$$\begin{aligned} |J'_{\lambda,\varepsilon}(u_n^+)v| &\leq |J'_{\lambda,\varepsilon}(u_n)v| + |\langle u_n^-, v \rangle_\lambda| + o_n(1) \\ &\leq \|J'_{\lambda,\varepsilon}(u_n)\|_{E'_\lambda} \|v\|_\lambda + \|u_n^-\|_\lambda \|v\|_\lambda + o_n(1). \end{aligned}$$

Therefore, since  $J'_{\lambda,\varepsilon}(u_n) = o_n(1)$ , we conclude that

$$\|J'_{\lambda,\varepsilon}(u_n^+)\|_{E'_\lambda} = o_n(1). \quad \blacksquare$$

As a byproduct of the above lemma, we obtain the clearly upper boundedness of  $\{u_n\}$ .

**Lemma 3.5** ([3, Lemma 3.6]). *If  $\{u_n\}$  is given by Lemma 3.2, then for any  $c_{\lambda,\varepsilon} > 0$ ,*

$$\limsup_{n \rightarrow \infty} \|u_n\|_\lambda^2 \leq \frac{2q}{q-2} c_{\lambda,\varepsilon}.$$

**Lemma 3.6.** *Let  $K > 0$  be independent on  $\lambda$  and let  $\{u_n\} \subset E_\lambda$  be a  $(PS)_c$  sequence for  $J_{\lambda,\varepsilon}$  with  $0 < c < K$ . Then, as  $\lambda_n \rightarrow \infty$ , there exists  $u \in H_0^s(\Omega)$  such that  $u_n \rightharpoonup u$  in  $E_{\lambda_n}$  and  $u_n \rightarrow u$  in  $L^r(\mathbb{R}^3)$  for all  $r \in [2, 2_s^*)$ .*

*Proof.* Lemma 3.5 yields that

$$\|u_n\|^2 \leq \|u_n\|_{\lambda_n}^2 \leq \frac{2q}{q-2} K.$$

We assume that  $u_n \rightharpoonup u$  in  $H^s(\mathbb{R}^3)$ . By the Fatou lemma, we have

$$\int_{\mathbb{R}^3} V(x)|u|^2 dx \leq \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^3} V(x)|u_n|^2 dx \leq \liminf_{n \rightarrow \infty} \frac{\|u_n\|_{\lambda_n}^2}{\lambda_n} = 0,$$

which implies that  $u(x) = 0$  almost everywhere in  $\mathbb{R}^3 \setminus V^{-1}(0)$ . Thus,  $u \in H_0^s(\Omega)$ .

Let  $F := \{x \in \mathbb{R}^3 : V(x) \leq M_0\}$ . Then

$$\int_{F^c} |u_n|^2 dx \leq \frac{1}{\lambda_n M_0} \int_{F^c} \lambda_n V(x) |u_n|^2 dx \leq \frac{2q}{(q-2)\lambda_n M_0} K \rightarrow 0.$$

For any  $r \in (1, \frac{2_s^*}{2})$ , and fixing  $r' = \frac{r}{r-1}$ , as  $R$  is large enough, from  $(V_3)$ , we have

$$\begin{aligned} \int_{F \cap B_R^c} |u_n - u|^2 dx &\leq |u_n - u|_{2r}^2 |\mathcal{L}(F \cap B_R^c)|^{\frac{1}{r'}} \\ &\leq C \|u_n - u\|^2 |\mathcal{L}(F \cap B_R^c)|^{\frac{1}{r'}} \rightarrow 0. \end{aligned}$$

On the other hand,  $u_n \rightarrow u$  in  $L^2(B_R)$ . This shows that  $u_n \rightarrow u$  in  $L^2(\mathbb{R}^3)$  as  $\lambda_n \rightarrow \infty$ . Following, by using interpolation, we obtain  $u_n \rightarrow u$  in  $L^r(\mathbb{R}^3)$  for all  $r \in [2, 2_s^*)$ .  $\blacksquare$

**Proposition 3.7.** *There exist  $\lambda^* = \lambda^*(\tau) > 0$  and  $b^*(\tau) > 0$  such that, for all  $\lambda \geq \lambda^*$  and  $b \in (0, b^*)$ ,  $J_{\lambda,\varepsilon}$  satisfies the  $(PS)_c$  condition at the level  $c \in (0, \frac{q-2}{2q} S^{\frac{3}{2s}} - \tau)$ , where  $\tau$  is from Lemma 3.3.*

*Proof.* Let  $\{u_n\}$  be a  $(PS)_c$  sequence of  $J_{\lambda,\varepsilon}$ . From Lemmas 3.2 and 3.4,  $\{u_n\}$  is a non-negative bounded sequence; hence, up to a subsequence, there are  $u \in E_\lambda$  and  $A \geq 0$  such that

$$\begin{aligned} u_n &\rightharpoonup u && \text{in } E_\lambda, \\ u_n &\rightarrow u && \text{in } L^r_{\text{loc}}(\mathbb{R}^3) \quad \text{for any } r \in [1, 2_s^*), \\ u_n(x) &\rightarrow u(x) && \text{a.e. in } \mathbb{R}^3, \\ [u_n]^2 &\rightarrow A^2. \end{aligned}$$

Setting the functional

$$\begin{aligned} P_{\lambda,\varepsilon}(u) &= \frac{1}{2} \|u\|_\lambda^2 + \frac{bA^2}{2} [u]^2 + \frac{1}{2\varepsilon} \int_\Omega [(\varphi - u)^+]^2 dx \\ &\quad - \frac{\mu}{q} \int_{\mathbb{R}^3} u^q dx - \frac{1}{2_s^*} \int_{\mathbb{R}^3} u^{2_s^*} dx, \end{aligned}$$

we have

$$P_{\lambda,\varepsilon}(u_n) = J_{\lambda,\varepsilon}(u_n) + \frac{bA^4}{4} + o_n(1),$$

and

$$J'_{\lambda,\varepsilon}(u_n)u_n - P'_{\lambda,\varepsilon}(u_n)u_n = o_n(1).$$

Thus,

$$P_{\lambda,\varepsilon}(u_n) = c + \frac{bA^4}{4} + o_n(1) \quad \text{and} \quad P'_{\lambda,\varepsilon}(u_n)u_n = o_n(1).$$

For any  $h \in C_0^\infty(\mathbb{R}^3)$ , taking  $h$  as a test function in (2.1), we obtain  $P'_{\lambda,\varepsilon}(u) = 0$ . Therefore,  $P_{\lambda,\varepsilon}(u) \geq 0$ .

Now, setting  $v_n := u_n - u$ , firstly by Lebesgue dominated convergence theorem, we get

$$\int_\Omega [(\varphi - u_n)^+]^2 dx = \int_\Omega [(\varphi - u)^+]^2 dx + o_n(1).$$

Therefore, by the Brezis–Lieb Lemma in [35] and from Lemma 3.3, it follows that

$$\begin{aligned} &P_{\lambda,\varepsilon}(u_n) - P_{\lambda,\varepsilon}(u) + o_n(1) \\ &= \frac{1}{2} \|v_n\|_\lambda^2 + \frac{bA^2}{2} [v_n]^2 - \frac{\mu}{q} \int_{\mathbb{R}^3} |v_n|^q dx - \frac{1}{2_s^*} \int_{\mathbb{R}^3} |v_n|^{2_s^*} dx \\ &= c + \frac{bA^4}{4} - P_{\lambda,\varepsilon}(u) < \frac{q-2}{2q} S^{\frac{3}{2s}} - \tau + \frac{bA^4}{4}. \end{aligned} \tag{3.2}$$

Then, taking  $b^*(\tau)$  small enough, for any  $b \in (0, b^*)$ , we can assume that

$$-\tau + \frac{bA^4}{4} \leq 0.$$

Thus,

$$P_{\lambda,\varepsilon}(u_n) - P_{\lambda,\varepsilon}(u) + o_n(1) < \frac{q-2}{2q} S^{\frac{3}{2s}}.$$

Besides, also by Lebesgue dominated convergence theorem, we get

$$\int_{\Omega} (\varphi - u_n)^+ u_n dx = \int_{\Omega} (\varphi - u)^+ u dx + o_n(1);$$

then we have

$$\begin{aligned} o_n(1) &= P'_{\lambda,\varepsilon}(u_n)u_n = P'_{\lambda,\varepsilon}(u_n)u_n - P'_{\lambda,\varepsilon}(u)u \\ &= \|v_n\|_{\lambda}^2 + bA^2[v_n]^2 - \mu \int_{\mathbb{R}^3} |v_n|^q dx - \int_{\mathbb{R}^3} |v_n|^{2_s^*} dx. \end{aligned} \quad (3.3)$$

Fixed  $\lambda > 0$ , assume  $\|v_n\|_{\lambda}^2 + bA^2[v_n]^2 \rightarrow l_1(\lambda)$  and  $\int_{\mathbb{R}^3} |v_n|^{2_s^*} dx \rightarrow l_2$ . If  $l_1(\lambda) = 0$ , then  $u_n \rightarrow u$  in  $E_{\lambda}$ . Thus, we may assume  $l_1(\lambda) > 0$ . Since Lemma 3.6, we have

$$\lim_{n \rightarrow \infty} \mu \int_{\mathbb{R}^3} |v_n|^q dx = o_{\lambda}(1).$$

Therefore, from (3.3), as  $n \rightarrow \infty$ , we may get  $l_2 = l_2(\lambda) = l_1(\lambda) + o_{\lambda}(1)$  and

$$\|v_n\|^2 \leq C(\|v_n\|_{\lambda}^q + \|v_n\|_{\lambda}^{2_s^*}).$$

As  $q \in (4, 2_s^*)$ , we get, for any  $t \in \mathbb{R}$ ,

$$|t|^q \leq \frac{1}{2C}|t|^2 + C|t|^{2_s^*}.$$

Then we find

$$\begin{aligned} l_1(\lambda) &= \lim_{n \rightarrow \infty} (\|v_n\|_{\lambda}^2 + bA^2[v_n]^2) \geq \lim_{n \rightarrow \infty} \|v_n\|_{\lambda}^2 \\ &\geq \left( \frac{1}{2C(C+1)} \right)^{\frac{2}{2_s^*-2}} := C_1 > 0, \end{aligned} \quad (3.4)$$

where  $C_1 > 0$  does not depend on  $\lambda$ . Since  $n \rightarrow \infty$ , by Sobolev inequality,

$$S \leq \frac{\|v_n\|_{\lambda}^2}{|v_n|_{2_s^*}^2} \leq \frac{l_1(\lambda)}{l_2(\lambda)^{\frac{2}{2_s^*}}} = \frac{l_1(\lambda)}{(l_1(\lambda) + o_{\lambda}(1))^{\frac{2}{2_s^*}}}.$$

From (3.4), we have

$$\liminf_{\lambda \rightarrow \infty} l_1(\lambda) \geq S^{\frac{3}{2_s^*}}.$$

Thus, from (3.2), we get

$$\frac{q-2}{2q} S^{\frac{3}{2_s^*}} > \liminf_{\lambda \rightarrow \infty} \left( \frac{1}{2} - \frac{1}{2_s^*} \right) l_1(\lambda) \geq \frac{q-2}{2q} S^{\frac{3}{2_s^*}},$$

which is absurd. Then we complete this proof.  $\blacksquare$

**Remark 3.8.** From Lemma 3.1, Lemma 3.3 and Proposition 3.7, there exist  $\mu_*$ ,  $\lambda^*$ ,  $b^* > 0$ , for every  $\mu \geq \mu_*$ ,  $\lambda \geq \lambda^*$  and  $b \in (0, b^*)$ , problem (2.1) has at least one weak solution  $u_{\varepsilon}$ . Besides, from Lemma 3.4,  $u_{\varepsilon}$  is non-negative and nontrivial.

#### 4. Proof of Theorem 1.1

From now on, making the change of notations

$$\epsilon = \frac{1}{n}, \quad u_n = u_{\frac{1}{n}}, \quad J_n = J_{\lambda, \epsilon} \quad \text{and} \quad J_n(u_n) = c_n = c_{\lambda, \epsilon},$$

for any  $v \in E_\lambda$ , we get

$$\begin{aligned} \langle u_n, v \rangle_\lambda + b[u_n]^2 \iint_{\mathbb{R}^6} \frac{(u_n(x) - u_n(y))(v(x) - v(y))}{|x - y|^{3+2s}} dx dy \\ - \frac{1}{\epsilon} \int_{\Omega} (\varphi - u_n)^+ v dx \\ = \int_{\mathbb{R}^3} f(u_n) v dx. \end{aligned} \quad (4.1)$$

From Lemma 3.2, there is  $u \in E_\lambda$  such that, up to a subsequence,  $u_n \rightharpoonup u$  in  $E_\lambda$ .

**Lemma 4.1** ([3, Lemma 3.11]). *If  $u$  is given above, then  $u \in \mathbb{K}$ .*

**Remark 4.2.** Take  $v := u_n - u$  in (4.1). Since

$$\langle P(u_n), u_n - u \rangle = \langle P(u_n) - P(u), u_n - u \rangle \geq 0,$$

we may get

$$\begin{aligned} \langle u_n, u_n - u \rangle_\lambda + b[u_n]^2 \iint_{\mathbb{R}^6} \frac{(u_n(x) - u_n(y))((u_n - u)(x) - (u_n - u)(y))}{|x - y|^{3+2s}} dx dy \\ \leq \int_{\mathbb{R}^3} f(u_n)(u_n - u) dx. \end{aligned} \quad (4.2)$$

**Lemma 4.3.** *We have  $u_n \rightarrow u$  in  $E_\lambda$  as  $\lambda \rightarrow \infty$ .*

*Proof.* Firstly, we show  $u_n \rightarrow u$  in  $H^s(\mathbb{R}^3)$ . Assume by contradiction that

$$u_n \not\rightarrow u \quad \text{in } H^s(\mathbb{R}^3),$$

that is,

$$\|u_n - u\|^2 \rightarrow B > 0.$$

Setting  $v_n = u_n - u$ , from [3, Lemma 3.12], we may have

$$\int_{\mathbb{R}^3} f(u_n) v_n dx = \int_{\mathbb{R}^3} f(v_n) v_n + \int_{\mathbb{R}^3} f(u) v_n + o_n(1).$$

Moreover, as

$$\int_{\mathbb{R}^3} f(u) v_n = o_n(1),$$

we have

$$\int_{\mathbb{R}^3} f(u_n)v_n dx = \int_{\mathbb{R}^3} f(v_n)v_n + o_n(1). \quad (4.3)$$

Since

$$\begin{aligned} & \iint_{\mathbb{R}^6} \frac{(u_n(x) - u_n(y))(u(x) - u(y))}{|x - y|^{3+2s}} dx dy \\ & \leq \iint_{\mathbb{R}^6} \frac{|(u_n(x) - u_n(y))(u(x) - u(y))|}{|x - y|^{3+2s}} dx dy \leq [u_n][u], \end{aligned}$$

thus from (4.2), we obtain

$$\begin{aligned} & \int_{\mathbb{R}^3} f(u_n)(u_n - u) dx \\ & \geq \langle u_n, u_n - u \rangle_\lambda \\ & \quad + b[u_n]^2 \iint_{\mathbb{R}^6} \frac{(u_n(x) - u_n(y))((u_n - u)(x) - (u_n - u)(y))}{|x - y|^{3+2s}} dx dy \\ & = \langle u_n, u_n - u \rangle_\lambda \\ & \quad + b[u_n]^2 \left( [u_n]^2 - \iint_{\mathbb{R}^6} \frac{(u_n(x) - u_n(y))(u(x) - u(y))}{|x - y|^{3+2s}} dx dy \right) \\ & \geq \langle u_n, u_n - u \rangle_\lambda + b[u_n]^2([u_n]^2 - [u_n][u]) \\ & = \langle u_n, u_n - u \rangle_\lambda + b[u_n]^3([u_n] - [u]). \end{aligned} \quad (4.4)$$

Moreover, from the Brezis–Lieb Lemma in [35], we get

$$[v_n]^2 = [u_n]^2 - [u]^2 + o_n(1) \geq 0;$$

thus,  $[u_n] = [u] + o_n(1)$  or  $[u_n] \geq [u]$ . Then by (4.3) and (4.4), we have

$$\int_{\mathbb{R}^3} f(v_n)v_n + o_n(1) \geq \langle u_n, u_n - u \rangle_\lambda = \|v_n\|_\lambda^2.$$

This is equivalent to

$$\|v_n\|^2 \leq \|v_n\|_\lambda^2 \leq \mu|v_n|_q^q + |v_n|_{2_s^*}^{2_s^*} + o_n(1). \quad (4.5)$$

On the other hand, since

$$\|v_n\|^2 \leq \|v_n\|_\lambda^2 \leq \|u_n\|_\lambda^2 + o_n(1),$$

from Lemma 3.5, we have

$$\limsup_{n \rightarrow \infty} \|v_n\|^2 < S^{\frac{3}{2s}},$$

that is,

$$B < S^{\frac{3}{2s}}. \quad (4.6)$$

Moreover, by (4.5),

$$S \leq \frac{\|v_n\|^2}{|v_n|_{2_s^*}^2} \leq \frac{\|v_n\|^2}{(\|v_n\|^2 - \mu|v_n|_q^q + o_n(1))^{\frac{2}{2_s^*}}}.$$

Taking  $\lambda \rightarrow \infty$  and  $n \rightarrow \infty$ , from Lemma 3.6, we will see

$$S \leq \frac{\|v_n\|^2}{(\|v_n\|^2 + o_\lambda(1) + o_n(1))^{\frac{2}{2_s^*}}} = \frac{B}{B^{\frac{2}{2_s^*}}};$$

then  $B \geq S^{\frac{3}{2_s}}$ . Because of (4.6),  $S^{\frac{3}{2_s}} \leq B < S^{\frac{3}{2_s}}$ , which shows a contradiction. Thus,  $\|v_n\| \rightarrow 0$ , and from (4.5), we have  $\|v_n\|_\lambda \rightarrow 0$  when  $n \rightarrow \infty$  and  $\lambda$  is large enough. ■

**Remark 4.4.** For any  $v \in \mathbb{K}$ , taking  $v - u_n$  in (4.1), from Lemmas 4.1 and 4.3, we can obtain  $u$  is a non-negative and nontrivial solution for (1.1).

To complete our concentration, we verify the following theorem.

**Theorem 4.5.** *Let  $\{u_n\} \subset E_{\lambda_n}$  be a sequence satisfying*

$$\begin{aligned} & (a + b[u_n]^2) \int_{\mathbb{R}^3} (-\Delta)^{\frac{s}{2}} u_n (-\Delta)^{\frac{s}{2}} (v - u_n) dx + \int_{\mathbb{R}^3} (1 + \lambda_n V(x)) u_n (v - u_n) dx \\ & \geq \int_{\mathbb{R}^3} f(u_n) (v - u_n) dx \end{aligned}$$

as  $\lambda_n \rightarrow \infty$  and for all  $v \in \mathbb{K}$ . Then there are a subsequence of  $\{u_n\}$ , still denoted by itself, and  $u \in H^s(\mathbb{R}^3)$  such that  $u_n \rightharpoonup u$  in  $H^s(\mathbb{R}^3)$ . Moreover,

- (i)  $u = 0$  a.e. in  $\Omega^c$ ;
- (ii)  $\|u_n - u\|_{\lambda_n}^2 \rightarrow 0$ ;
- (iii) we may obtain

$$\begin{aligned} & u_n \rightarrow u \quad \text{in } H^s(\mathbb{R}^3), \\ & \lambda_n \int_{\mathbb{R}^3} V(x) u_n^2 dx \rightarrow 0, \\ & \|u_n\|_{\lambda_n}^2 \rightarrow \int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{3+2s}} dx dy + \int_\Omega |u|^2 dx, \end{aligned}$$

where  $Q := \mathbb{R}^3 \times \mathbb{R}^3 \setminus (\Omega^c \times \Omega^c)$ .

- (iv)  $u$  is a solution of the variational inequality

$$\begin{aligned} & (a + b[u]_Q^2) \iint_Q \frac{(u(x) - u(y))((w - u)(x) - (w - u)(y))}{|x - y|^{3+2s}} dx dy + \int_\Omega u(w - u) dx \\ & \geq \int_\Omega (\mu|u|^{q-2} + |u|^{2^*-2}) u(w - u) dx \end{aligned}$$

for all  $w \in \tilde{\mathbb{K}}$ , where

$$\tilde{\mathbb{K}} := \{w \in H_0^s(\Omega), w \geq \varphi \text{ a.e. in } \Omega\}.$$



*Proof.* From Lemma 3.5,  $\{u_n\}$  is bounded in  $H^s(\mathbb{R}^3)$ , so up to a subsequence, there exists  $u \in H^s(\mathbb{R}^3)$  such that

$$u_n \rightharpoonup u \quad \text{in } H^s(\mathbb{R}^3).$$

(i) The same process as in Lemma 3.6 works to prove  $u = 0$  a.e. in  $\Omega^c$ .

(ii) Since  $u_n \rightharpoonup u$  in  $H^s(\mathbb{R}^3)$ , if we set  $v_n = u_n - u$ , by the same arguments explored in the proof of Lemma 4.3, we can show that  $v_n \rightarrow 0$  in  $H^s(\mathbb{R}^3)$ , and

$$\|v_n\|_{\lambda_n}^2 \leq \int_{\mathbb{R}^3} f(v_n)v_n \, dx + o_n(1).$$

Thus, we obtain, as  $n \rightarrow \infty$ ,

$$v_n \rightarrow 0 \quad \text{in } E_{\lambda_n}.$$

(iii) From (i)–(ii), we get

$$\lambda_n \int_{\mathbb{R}^3} V(x)|u_n|^2 \, dx = \lambda_n \int_{\mathbb{R}^3} V(x)|u_n - u|^2 \, dx \leq \|u_n - u\|_{\lambda_n}^2 \rightarrow 0,$$

and

$$\|u_n\|_{\lambda_n}^2 \rightarrow \int_{\mathcal{Q}} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} \, dx \, dy + \int_{\Omega} |u|^2 \, dx.$$

(iv) For any  $w \in \tilde{\mathbb{K}}$ , we may get

$$\begin{aligned} \langle u_n, w - u_n \rangle_{\lambda_n} + b[u_n]^2 & \iint_{\mathbb{R}^6} \frac{(u_n(x) - u_n(y))((w - u_n)(x) - (w - u_n)(y))}{|x - y|^{3+2s}} \, dx \, dy \\ & \geq \int_{\mathbb{R}^3} f(u_n)(w - u_n) \, dx. \end{aligned}$$

Thus, from (i)–(iii), taking  $n \rightarrow \infty$ , we obtain

$$\begin{aligned} & \left( a + b \int_{\mathcal{Q}} \frac{|u(x) - u(y)|^2}{|x - y|^{3+2s}} \, dx \, dy \right) \\ & \quad \times \int_{\mathcal{Q}} \frac{(u_n(x) - u_n(y))((w - u_n)(x) - (w - u_n)(y))}{|x - y|^{3+2s}} \, dx \, dy \\ & \quad + \int_{\Omega} u(w - u) \, dx \geq \int_{\Omega} (\mu u^{q-2} + u^{2^*_s-2})u(w - u) \, dx. \end{aligned}$$

■

**Acknowledgments.** The authors warmly thank the anonymous referees for their useful and nice comments on the paper.

**Funding.** S. Deng and W. Luo have been supported by National Natural Science Foundation of China, No. 11971392, and Natural Science Foundation of Chongqing, China, cstc2019jcyjX0022.

## References

- [1] M. Alimohammady, C. O. Alves, and H. Kaffash Amiri, [Existence and multiplicity of positive solutions for a class of Kirchhoff Laplacian type problems](#). *J. Math. Phys.* **60** (2019), no. 10, article no. 101503 Zbl 1427.35012 MR 4015627
- [2] C. O. Alves and L. M. Barros, [Existence and multiplicity of solutions for a class of elliptic problem with critical growth](#). *Monatsh. Math.* **187** (2018), no. 2, 195–215 Zbl 1421.35142 MR 3850308
- [3] C. O. Alves, L. M. Barros, and C. E. Torres Ledesma, [Existence of solution for a class of variational inequality in whole  \$\mathbb{R}^N\$  with critical growth](#). *J. Math. Anal. Appl.* **494** (2021), no. 2, article no. 124672 Zbl 1459.35204 MR 4164961
- [4] C. O. Alves, L. M. Barros, and C. E. Torres Ledesma, [Existence of solution for a class of variational inequality in whole  \$\mathbb{R}^N\$  with critical growth: the local mountain pass case](#). *Mediterr. J. Math.* **20** (2023), no. 5, article no. 239 Zbl 1518.35378 MR 4603356
- [5] V. Ambrosio, [Nonlinear fractional Schrödinger equations in  \$\mathbb{R}^N\$](#) . Front. Ellip. Parab. Probl., Birkhäuser/Springer, Cham, 2021 MR 4264520
- [6] V. Ambrosio, [A Kirchhoff type equation in  \$\mathbb{R}^N\$  involving the fractional  \$\(p, q\)\$ -Laplacian](#). *J. Geom. Anal.* **32** (2022), no. 4, article no. 135 Zbl 1485.35003 MR 4376646
- [7] T. Bartsch and Z.-Q. Wang, [Multiple positive solutions for a nonlinear Schrödinger equation](#). *Z. Angew. Math. Phys.* **51** (2000), no. 3, 366–384 Zbl 0972.35145 MR 1762697
- [8] A. Bensoussan and J.-L. Lions, [Applications des inéquations variationnelles en contrôle stochastique](#). Méth. Math. Infor. 6, Dunod, Paris, 1978 MR 513618
- [9] L. A. Caffarelli, S. Salsa, and L. Silvestre, [Regularity estimates for the solution and the free boundary of the obstacle problem for the fractional Laplacian](#). *Invent. Math.* **171** (2008), no. 2, 425–461 Zbl 1148.35097 MR 2367025
- [10] S. Challal, A. Lyaghfour, and J. F. Rodrigues, [On the  \$A\$ -obstacle problem and the Hausdorff measure of its free boundary](#). *Ann. Mat. Pura Appl. (4)* **191** (2012), no. 1, 113–165 Zbl 1235.35285 MR 2886164
- [11] M. Clapp and Y. Ding, [Positive solutions of a Schrödinger equation with critical nonlinearity](#). *Z. Angew. Math. Phys.* **55** (2004), no. 4, 592–605 Zbl 1060.35130 MR 2107669
- [12] J.-N. Corvellec, M. Degiovanni, and M. Marzocchi, [Deformation properties for continuous functionals and critical point theory](#). *Topol. Methods Nonlinear Anal.* **1** (1993), no. 1, 151–171 Zbl 0789.58021 MR 1215263
- [13] M. Degiovanni and M. Marzocchi, [A critical point theory for nonsmooth functionals](#). *Ann. Mat. Pura Appl. (4)* **167** (1994), 73–100 Zbl 0828.58006 MR 1313551
- [14] Y. Ding and K. Tanaka, [Multiplicity of positive solutions of a nonlinear Schrödinger equation](#). *Manuscripta Math.* **112** (2003), no. 1, 109–135 Zbl 1038.35114 MR 2005933
- [15] H. Fan, [Multiple positive solutions for a class of Kirchhoff type problems involving critical Sobolev exponents](#). *J. Math. Anal. Appl.* **431** (2015), no. 1, 150–168 Zbl 1319.35050 MR 3357580
- [16] M. Fang, [Degenerate elliptic inequalities with critical growth](#). *J. Differential Equations* **232** (2007), no. 2, 441–467 Zbl 1166.35334 MR 2286387
- [17] G. M. Figueiredo, M. Furtado, and M. Montenegro, [An obstacle problem in a plane domain with two solutions](#). *Adv. Nonlinear Stud.* **14** (2014), no. 2, 327–337 Zbl 1301.49023 MR 3194357
- [18] A. Fiscella and E. Valdinoci, [A critical Kirchhoff type problem involving a nonlocal operator](#). *Nonlinear Anal.* **94** (2014), 156–170 Zbl 1283.35156 MR 3120682

- [19] S. Frassu, E. M. Rocha, and V. Staicu, [The obstacle problem at zero for the fractional  \$p\$ -Laplacian](#). *Set-Valued Var. Anal.* **30** (2022), no. 1, 207–231 Zbl 07490854 MR 4389882
- [20] O. Frites and T. Moussaoui, [Existence of positive solutions for a variational inequality of Kirchhoff type](#). *Arab J. Math. Sci.* **21** (2015), no. 2, 127–135 Zbl 1325.47110 MR 3376105
- [21] X. He and W. Zou, [Existence and concentration behavior of positive solutions for a Kirchhoff equation in  \$\mathbb{R}^3\$](#) . *J. Differential Equations* **252** (2012), no. 2, 1813–1834 Zbl 1235.35093 MR 2853562
- [22] X. He and W. Zou, [Ground states for nonlinear Kirchhoff equations with critical growth](#). *Ann. Mat. Pura Appl. (4)* **193** (2014), no. 2, 473–500 Zbl 1300.35016 MR 3180929
- [23] A. Ioffe and E. Schwartzman, [Metric critical point theory. I. Morse regularity and homotopic stability of a minimum](#). *J. Math. Pures Appl. (9)* **75** (1996), no. 2, 125–153 Zbl 0852.58018 MR 1380672
- [24] G. Kirchhoff, *Mechanik*, Teubner, Leipzig, 1883
- [25] K. Lan, [A variational inequality theory in reflexive smooth Banach spaces and applications to  \$p\$ -Laplacian elliptic inequalities](#). *Nonlinear Anal.* **113** (2015), 71–86 Zbl 1304.49020 MR 3281846
- [26] K. Q. Lan, [Positive weak solutions of semilinear second order elliptic inequalities via variational inequalities](#). *J. Math. Anal. Appl.* **380** (2011), no. 2, 520–530 Zbl 1216.35184 MR 2794411
- [27] P. Magrone, D. Mugnai, and R. Servadei, [Multiplicity of solutions for semilinear variational inequalities via linking and  \$\nabla\$ -theorems](#). *J. Differential Equations* **228** (2006), no. 1, 191–225 Zbl 1130.35072 MR 2254429
- [28] M. Matzeu and R. Servadei, [Semilinear elliptic variational inequalities with dependence on the gradient via mountain pass techniques](#). *Nonlinear Anal.* **72** (2010), no. 11, 4347–4359 Zbl 1273.35144 MR 2606792
- [29] M. Matzeu and R. Servadei, [Stability for semilinear elliptic variational inequalities depending on the gradient](#). *Nonlinear Anal.* **74** (2011), no. 15, 5161–5170 Zbl 1220.35078 MR 2810697
- [30] P. D. Panagiotopoulos, *Hemivariational inequalities*. Springer, Berlin, 1993 Zbl 0826.73002 MR 1385670
- [31] P. Pucci, M. Xiang, and B. Zhang, [Multiple solutions for nonhomogeneous Schrödinger–Kirchhoff type equations involving the fractional  \$p\$ -Laplacian in  \$\mathbb{R}^N\$](#) . *Calc. Var. Partial Differential Equations* **54** (2015), no. 3, 2785–2806 Zbl 1329.35338 MR 3412392
- [32] J.-F. Rodrigues, *Obstacle problems in mathematical physics*. North-Holland Math. Stud. 134, North-Holland Publishing Co., Amsterdam, 1987 Zbl 0606.73017 MR 880369
- [33] R. Servadei and E. Valdinoci, [Lewy-Stampacchia type estimates for variational inequalities driven by \(non\)local operators](#). *Rev. Mat. Iberoam.* **29** (2013), no. 3, 1091–1126 Zbl 1275.49016 MR 3090147
- [34] K. Teng, [Two nontrivial solutions for hemivariational inequalities driven by nonlocal elliptic operators](#). *Nonlinear Anal. Real World Appl.* **14** (2013), no. 1, 867–874 Zbl 1259.35229 MR 2969880
- [35] M. Willem, *Minimax theorems*. Progr. Nonlinear Differential Equations Appl. 24, Birkhäuser Boston, Inc., Boston, MA, 1996 MR 1400007
- [36] K. Wu and G. Gu, [Existence of positive solutions for fractional Kirchhoff equation](#). *Z. Angew. Math. Phys.* **73** (2022), no. 2, article no. 45 Zbl 1492.74039 MR 4379085
- [37] M. Xiang, [A variational inequality involving nonlocal elliptic operators](#). *Fixed Point Theory Appl.* (2015), 2015:148, 9 MR 3385687

- [38] J. F. Yang, [Positive solutions of an obstacle problem](#). *Ann. Fac. Sci. Toulouse Math.* (6) **4** (1995), no. 2, 339–366 Zbl [0866.49017](#) MR [1344725](#)
- [39] J. F. Yang, [Positive solutions of quasilinear elliptic obstacle problems with critical exponents](#). *Nonlinear Anal.* **25** (1995), no. 12, 1283–1306 Zbl [0838.49008](#) MR [1355723](#)
- [40] J. Zuo, T. An, and W. Liu, [A variational inequality of Kirchoff-type in  \$\mathbb{R}^N\$](#) . *J. Inequal. Appl.* (2018), article no. 329 Zbl [1498.49023](#) MR [3881333](#)

Received 24 July 2023; revised 10 October 2023.

**Shenbing Deng**

School of Mathematics and Statistics, Southwest University, 400715 Chongqing, P. R. of China; [shbdeng@swu.edu.cn](mailto:shbdeng@swu.edu.cn)

**Wenshan Luo**

School of Mathematics and Statistics, Southwest University, 400715 Chongqing, P. R. of China; [swluoswumaths@163.com](mailto:swluoswumaths@163.com)

**César E. Torres Ledesma**

Departamento de Matemáticas, Instituto de Investigación en Matemáticas, Universidad Nacional de Trujillo, Av. Juan Pablo II s/n, 13006 Trujillo, Peru; [ctl\\_576@yahoo.es](mailto:ctl_576@yahoo.es)

**George W. Alama Quiroz**

Facultad de Estudios Generales-UPN-Campus Virtual, Universidad Privada del Norte, Urb. San Isidro 2da etapa, 13006 Trujillo, Peru; [george.alama@upn.edu.pe](mailto:george.alama@upn.edu.pe)