

# A New Minimal Point Theorem in Product Spaces

A. Göpfert, Chr. Tammer and C. Zălinescu

**Abstract.** We derive a minimal point theorem for a subset  $A$  in a cone in product spaces under a weak assumption concerning the boundedness of the considered set  $A$ . Using this result we improve two vectorial variants of Ekeland's variational principle. Finally, a new characterization of well-based cones is given.

**Keywords:** *Minimal point theorems, Ekeland's variational principle, well-based cones*

**AMS subject classification:** 49 J 40, 90 C 29, 46 A 40, 90 C 26

Assume that  $(X, d)$  is a complete metric space,  $Y$  is a separated locally convex space,  $Y^*$  is its topological dual,  $K \subset Y$  is a convex cone, i.e.  $K + K \subset K$  and  $[0, \infty) \cdot K \subset K$ ,

$$K^+ = \{y^* \in Y^* : \langle y, y^* \rangle \geq 0 \text{ for all } y \in K\}$$

is the dual cone of  $K$  and

$$K^\# = \{y^* \in Y^* : \langle y, y^* \rangle > 0 \text{ for all } y \in K \setminus \{0\}\}.$$

In this note we suppose that  $K$  is pointed, i.e.  $K \cap (-K) = \{0\}$ . The cone  $K$  determines an order relation on  $Y$ , denoted in the sequel by  $\leq_K$ ; so, for  $y_1, y_2 \in Y$ ,  $y_1 \leq_K y_2$  if  $y_2 - y_1 \in K$ . It is well known that " $\leq_K$ " is reflexive, transitive and antisymmetric. Let  $k^0 \in K \setminus \{0\}$ ; using the element  $k^0$  we introduce an order relation on  $X \times Y$ , denoted by " $\preceq_{k^0}$ ", in the following manner:

$$(x_1, y_1) \preceq_{k^0} (x_2, y_2) \quad \text{iff} \quad y_1 + k^0 d(x_1, x_2) \leq_K y_2.$$

Note that " $\preceq_{k^0}$ " is reflexive, transitive and antisymmetric. That is, our notations are those of [3].

The essential idea for the derivation of a minimal point theorem (cf. [2, 8]) in general product spaces  $X \times Y$ , as well as of the vectorial Ekeland principle, consists in including the ordering cone  $K \subset Y$  in a "larger" cone  $B \subset Y$ :  $K \setminus \{0\} \subset \text{int } B$ . We will use  $B$  to define a suitable functional  $z_B : Y \rightarrow \mathbb{R}$ . Moreover, we will replace the usual boundedness condition of the projection  $P_Y A$  of  $A$  onto  $Y$  by a weaker one.

---

A. Göpfert: Dept. Math. & Inform. Univ., D-06099 Halle  
Chr. Tammer: Dept. Math. & Inform. Univ., D-06099 Halle  
C. Zălinescu: Fac. Math. Univ. "Al. I. Cuza", 6600 Iași, Romania

**Theorem 1.** Assume that there exists a proper convex cone  $B \subset Y$  such that  $K \setminus \{0\} \subset \text{int } B$ . Suppose that the set  $A \subset X \times Y$  satisfies the condition

(H1) for every  $\preceq_{k^0}$ -decreasing sequence  $((x_n, y_n)) \subset A$  with  $x_n \rightarrow x \in X$  there exists  $y \in Y$  such that  $(x, y) \in A$  and  $(x, y) \preceq_{k^0} (x_n, y_n)$  for every  $n \in \mathbb{N}$

and that  $P_Y(A) \cap (\tilde{y} - \text{int } B) = \emptyset$  for some  $\tilde{y} \in Y$ . Then for every  $(x_0, y_0) \in A$  there exists  $(\bar{x}, \bar{y}) \in A$ , minimal with respect to  $\preceq_{k^0}$ , such that  $(\bar{x}, \bar{y}) \preceq_{k^0} (x_0, y_0)$ .

**Proof.** Let

$$z_B : Y \rightarrow \mathbb{R}, \quad z_B(y) = \inf\{t \in \mathbb{R} : y \in tk^0 - \text{cl } B\}.$$

By [3: Lemma 7],  $z_B$  is a continuous sublinear function such that  $z_B(y + tk^0) = z_B(y) + t$  for all  $t \in \mathbb{R}$  and  $y \in Y$ , and for every  $\lambda \in \mathbb{R}$

$$\begin{aligned} \{y \in Y : z_B(y) \leq \lambda\} &= \lambda k^0 - \text{cl } B \\ \{y \in Y : z_B(y) < \lambda\} &= \lambda k^0 - \text{int } B. \end{aligned}$$

Moreover, if  $y_2 - y_1 \in K \setminus \{0\}$ , then  $z_B(y_1) < z_B(y_2)$ . Observe that for  $(x, y) \in A$  we have that  $z_B(y - \tilde{y}) \geq 0$ . Otherwise for some  $(x, y) \in A$  we have  $z_B(y - \tilde{y}) < 0$ . It follows that there exists  $\lambda > 0$  such that  $y - \tilde{y} \in -\lambda k^0 - \text{cl } B$ . Hence

$$y \in \tilde{y} - (\lambda k^0 + \text{cl } B) \subset \tilde{y} - (\text{int } B + \text{cl } B) \subset \tilde{y} - \text{int } B$$

which is a contradiction. Since  $0 \leq z_B(y - \tilde{y}) \leq z_B(y) + z_B(-\tilde{y})$ , it follows that  $z_B$  is bounded from below on  $P_Y(A)$ . Let us construct a sequence  $((x_n, y_n))_{n \geq 0} \subset A$  as follows: having  $(x_n, y_n) \in A$  we take  $(x_{n+1}, y_{n+1}) \in A$ ,  $(x_{n+1}, y_{n+1}) \preceq_{k^0} (x_n, y_n)$ , such that

$$z_B(y_{n+1}) \leq \inf \left\{ z_B(y) : (x, y) \in A \text{ and } (x, y) \preceq_{k^0} (x_n, y_n) \right\} + \frac{1}{n+1}.$$

Of course, the sequence  $((x_n, y_n))$  is  $\preceq_{k^0}$ -decreasing. It follows that

$$y_{n+p} + k^0 d(x_{n+p}, x_n) \leq_K y_n \quad \forall n, p \in \mathbb{N}^*$$

so that

$$d(x_{n+p}, x_n) \leq z_B(y_n) - z_B(y_{n+p}) \leq \frac{1}{n} \quad \forall n, p \in \mathbb{N}^*.$$

It follows that  $(x_n)$  is a Cauchy sequence in the complete metric space  $(X, d)$ , and so  $(x_n)$  is convergent to some  $\bar{x} \in X$ . By condition (H1) there exists  $\bar{y} \in Y$  such that  $(\bar{x}, \bar{y}) \in A$  and  $(\bar{x}, \bar{y}) \preceq_{k^0} (x_n, y_n)$  for every  $n \in \mathbb{N}$ .

Let us show that  $(\bar{x}, \bar{y})$  is the desired element. Indeed,  $(\bar{x}, \bar{y}) \preceq_{k^0} (x_0, y_0)$ . Suppose that  $(x', y') \in A$  is such that  $(x', y') \preceq_{k^0} (\bar{x}, \bar{y})$  ( $\preceq_{k^0} (x_n, y_n)$  for every  $n \in \mathbb{N}$ ). Thus  $z_B(y') + d(x', \bar{x}) \leq z_B(\bar{y})$ , whence

$$d(x', \bar{x}) \leq z_B(\bar{y}) - z_B(y') \leq z_B(y_n) - z_B(y') \leq \frac{1}{n} \quad \forall n \geq 1.$$

It follows that  $d(x', \bar{x}) = z_B(\bar{y}) - z_B(y') = 0$ . Hence  $x' = \bar{x}$ . As  $y' \leq_K \bar{y}$ , if  $y' \neq \bar{y}$ , then  $\bar{y} - y' \in K \setminus \{0\}$ , whence  $z_B(y') < z_B(\bar{y})$ , which is a contradiction. Therefore  $(x', y') = (\bar{x}, \bar{y})$  ■

Comparing with [3: Theorem 4], note that the present condition on  $K$  is stronger (because in this case  $K^\# \neq \emptyset$ ), while the condition on  $A$  is weaker ( $A$  may be not contained in a half-space). Note that when  $K$  and  $k^0$  are as in Theorem 1, Corollaries 2 and 3 from [3] may be improved. In the next result  $Y^* = Y \cup \{\infty\}$  with  $\infty \notin Y$ ; we consider that  $y \leq_K \infty$  for every  $y \in Y$ . We consider also a function  $f : X \rightarrow Y^*$  and  $\text{dom } f = \{x \in X : f(x) \neq \infty\}$ .

In the following corollary we derive a variational principle of Ekeland's type for objective functions which take values in a general space  $Y$  (cf. [2, 3, 5 - 7]) under a weaker assumption with respect to the usual lower semicontinuity. For the case  $Y = \mathbb{R}$ , assumption (H4) in Corollary 2 is fulfilled for decreasingly semicontinuous real-valued functions as in the paper [4].

**Corollary 2.** *Let  $f : X \rightarrow Y^*$ . Assume that there exists a proper convex cone  $B \subset Y$  such that  $K \setminus \{0\} \subset \text{int } B$  and  $f(X) \cap (\tilde{y} - B) = \emptyset$  for some  $\tilde{y} \in Y$ . Also, suppose that*

(H3)  $\{x' \in X : f(x') + k^0 d(x', x) \leq_K f(x)\}$  is closed for every  $x \in X$

or

(H4) for every sequence  $(x_n) \subset \text{dom } f$  with  $x_n \rightarrow x$  and  $(f(x_n)) \leq_K$ -decreasing,  $f(x) \leq_K f(x_n)$  for every  $n \in \mathbb{N}$ , and  $K$  is closed in the direction  $k^0$ .

Then for every  $x_0 \in \text{dom } f$  there exists  $\bar{x} \in X$  such that

$$f(\bar{x}) + k^0 d(\bar{x}, x_0) \leq_K f(x_0)$$

and

$$\forall x \in X : f(x) + k^0 d(\bar{x}, x) \leq_K f(\bar{x}) \implies x = \bar{x}.$$

We say that  $K$  is closed in the direction  $k^0$  if  $K \cap (y - \mathbb{R}_+ k^0)$  is closed for every  $y \in K$ . The proof of Corollary 2 is similar to those of Corollaries 2 and 3 in [3].

As mentioned in [3], condition (H1) is verified if  $K$  is a well based convex cone,  $Y$  is a Banach space and  $A$  is closed. As usually (cf. [1]), a convex set  $S$  is said to be a base for a convex cone  $K \subset Y$  if

$$K = \mathbb{R}_+ S = \{\lambda y : \lambda \geq 0 \text{ and } y \in S\} \quad \text{and} \quad 0 \notin \text{cl } S.$$

The cone  $K$  is called well based if  $K$  has a bounded base  $S$ . Concerning well based convex cones in normed spaces we have the following characterization.

**Proposition 3.** *Let  $Y$  be a normed vector space and  $K \subset Y$  a proper convex cone. Then  $K$  is well based if and only if there exist  $k^0 \in K$  and  $z^* \in K^+$  such that  $\langle k^0, z^* \rangle > 0$  and*

$$K \cap S_1 \subset k^0 + \{y \in Y : \langle y, z^* \rangle > 0\}$$

where  $S_1 = \{y \in Y : \|y\| = 1\}$  is the unit sphere in  $Y$ .

**Proof.** Suppose first that  $K$  is well based with bounded base  $S$ ; therefore  $0 \notin \text{cl } S$  and  $K = [0, \infty) \cdot S$ . Then there exists  $z^* \in Y^*$  such that  $1 \leq \langle y, z^* \rangle$  for all  $y \in S$ . Consider  $\tilde{S} := \{k \in K : \langle k, z^* \rangle = 1\}$ . It follows that  $\tilde{S}$  is a base of  $K$ ; moreover, since

$\tilde{S} \subset [0, 1] \cdot S$ ,  $\tilde{S}$  is also bounded. Taking  $k^1 \in K \setminus \{0\}$  we have  $K \cap S_1 \subset \lambda k^1 + B_+$  for some  $\lambda > 0$ , where  $B_+ = \{y \in Y : \langle y, z^* \rangle > 0\}$ . Otherwise

$$\forall n \in \mathbb{N}^* \exists k_n \in K \cap S_1 : \quad k_n \notin \frac{1}{n} k^1 + B_+.$$

Therefore  $\langle k_n, z^* \rangle \leq \frac{1}{n} \langle k^1, z^* \rangle$  for every  $n \geq 1$ . But, because  $\tilde{S}$  is a base,  $k_n = \lambda_n b_n$  with  $\lambda_n > 0$  and  $b_n \in \tilde{S}$ ; it follows that  $1 = \|k_n\| = \lambda_n \|b_n\| \leq \lambda_n M$  with  $M > 0$  (because  $\tilde{S}$  is bounded). Therefore

$$M^{-1} \leq \lambda_n = \langle \lambda_n b_n, z^* \rangle = \langle k_n, z^* \rangle \leq n^{-1} \langle k^1, z^* \rangle \quad \forall n \in \mathbb{N}^*$$

whence  $M^{-1} \leq 0$ , which is a contradiction. Thus there exists  $\lambda > 0$  such that  $K \cap S_1 \subset \lambda k^1 + B_+$ . Taking  $k^0 := \lambda k^1$  the conclusion follows.

Suppose now that  $K \cap S_1 \subset k^0 + B_+$  for some  $k^0 \in K$  and  $z^* \in K^+$  with  $\langle k^0, z^* \rangle = c > 0$ , where  $B_+$  is defined as above. Consider  $S = \{k \in K : \langle k, z^* \rangle = 1\}$ . Let  $k \in K \setminus \{0\}$ . Then  $\|k\|^{-1} k = k^0 + y$  for some  $y \in B_+$ . It follows that  $\langle k, z^* \rangle > c \|k\| > 0$ ; therefore  $z^* \in K^\#$  and so  $k \in (0, \infty) \cdot S$ . Since  $\text{cl } S \subset \{y \in Y : \langle y, z^* \rangle = 1\}$ , we have that  $S$  is a base of  $K$ . Let now  $y \in S \subset K$ . Then  $\|y\|^{-1} y \in K \cap S_1$ . There exists  $z \in B_+$  such that  $\|y\|^{-1} y = k^0 + z$ . We get

$$1 = \langle y, z^* \rangle = \|y\| \langle k^0 + z, z^* \rangle \geq c \|y\|$$

whence  $\|y\| \leq c^{-1}$ . Therefore  $S$  is bounded, and so  $K$  is well-based ■

## References

- [1] Borwein, J. M.: *Continuity and differentiability properties of convex operators*. Proc. London Math. Soc. 44 (1982), 420 - 444.
- [2] Göpfert, A. and Chr. Tammer: *A new maximal point theorem*. Z. Anal. Anw. 14 (1995), 379 - 390.
- [3] Göpfert, A., Tammer, Chr. and C. Zălinescu: *On the vectorial Ekeland's variational principle and minimal points in product spaces*. Nonlin. Anal.: Theory, Methods and Appl. (to appear).
- [4] Gajek, L. and D. Zagrodny: *Geometric Variational Principle*. Diss. Math. 340 (1995), 55 - 71.
- [5] Isac, G.: *The Ekeland's principle and the Pareto  $\epsilon$ -efficiency*. In: Multi-objective Programming and Goal Programming. Theory and Applications (ed.: M. Tamiz; Lect. Notes Econ. Math. Syst.: Vol. 432). Berlin: Springer-Verlag 1996, pp. 148 - 163.
- [6] Khanh, P. O.: *On Caristi-Kirk's theorem and Ekeland's variational principle for Pareto extrema*. Preprint. Warsaw: Polish Acad. Sci., Inst. Math., Preprint 357 (1986), 1 - 7.
- [7] Nemeth, A. B.: *A nonconvex vector minimization problem*. Nonlin. Anal.: Theory, Methods and Appl. 10 (1986), 669 - 678.
- [8] Phelps, R. R.: *Convex Functions, Monotone Operators and Differentiability*, 2nd Ed. Lect. Notes Math. 1364 (1993).