

A Compactness Criterion of Mixed Krasnoselskiĭ-Riesz Type in Regular Ideal Spaces of Vector Functions

M. Väth

Abstract. We present a compactness criterion in ideal spaces of vector-valued functions. In the case of real functions, the criterion gives a precise formula for the measure of non-compactness. In the Lebesgue-Bochner spaces $L_p(\mathbb{R}^n, U)$ the result can be interpreted as a Riesz compactness criterion and generalizes a theorem of Orlicz and Szulfa.

Keywords: *Riesz compactness criterion, Krasnoselskiĭ compactness criterion, ideal spaces, Köthe spaces, Banach function spaces, spaces of measurable functions, vector-valued functions*

AMS subject classification: Primary 46 E 30, 46 E 40, secondary 45 N 10, 46 G 10

1. The main result

Let S be a measure space and $(U, |\cdot|)$ be a Banach space. We call a function $x : S \rightarrow U$ (*strongly Bochner*) *measurable* if, on each set of finite measure, x may be approximated a.e. (in the sense of the Lebesgue extension of the measure space) by a sequence of simple functions. This is the definition used, e.g., in [9]. For a measurable set $E \subseteq S$ we define the projection P_E by $P_E x(s) = \chi_E(s)x(s)$.

A normed linear space $(X, \|\cdot\|)$ of (classes of) measurable functions $x : S \rightarrow U$ is called a *preideal space*, if the relations $x \in X$ and $|y(s)| \leq |x(s)|$ for a measurable function y imply that $y \in X$ and $\|y\| \leq \|x\|$. If X is complete, it is called an *ideal space*. In some literature, preideal spaces are called (*normed*) *Köthe spaces*, and ideal spaces are called *Banach function spaces*; but sometimes additional requirements are imposed on these spaces. The proofs of properties for such spaces which are not given in this paper can be found in [27, 30, 31].

To each preideal space X one associates a preideal space $X_{\mathbb{R}}$ of real functions $x : S \rightarrow \mathbb{R}$ in the obvious way by the relation

$$x \in X \iff |x| \in X_{\mathbb{R}}.$$

M. Väth: University of Würzburg, Department of Mathematics, Am Hubland, D-97074 Würzburg; e-mail: vaeth@cip.mathematik.uni-wuerzburg.de

The paper was written while the author was visiting Caltech as a DAAD fellow. Financial support by the DAAD is gratefully acknowledged. The author thanks W. A. J. Luxemburg for stimulating discussions and valuable remarks on the subject.

X is an ideal space if and only if $X_{\mathbb{R}}$ is an ideal space. We will often not distinguish between X and $X_{\mathbb{R}}$ and thus write, e.g., $\chi_E \in X$. We say that a family $M \subseteq X$ has *equicontinuous norm* in X , if

$$\inf_{\text{mes } E < \infty} \sup_{x \in M} \|P_{E^c} x\| = 0 \quad \text{and} \quad \lim_{\delta \rightarrow 0} \sup_{\text{mes } D \leq \delta} \sup_{x \in M} \|P_D x\| = 0.$$

The space X is called *regular*, if all singletons $M = \{x\}$ with $x \in X$ have equicontinuous norm. For example, the Lebesgue-Bochner spaces $X = L_p(S, U)$ ($1 \leq p \leq \infty$) are regular if $p < \infty$. It will be convenient to introduce the shortcut

$$M(s) = \{x(s) : x \in M\}$$

for a set $M \subseteq X$.

We are interested in estimating the Hausdorff measure of non-compactness of the set M in relation to the Hausdorff measure of non-compactness of the sets $M(s)$. Such estimates play an important role in the study of integral and functional equations of vector-valued functions (see, e.g., [20, 21, 23, 26]). Recall that the Hausdorff measure of non-compactness of a set A in a metric space Y is defined as the infimum of all $\varepsilon > 0$ such that A admits a finite ε -net in Y .

We denote the Hausdorff measure of non-compactness by $\chi(A)$ (or by $\chi_Y(A)$, to emphasize the dependence on Y). Observe that $\chi_Y(A)$ increases, in general, if we replace Y by a subspace $Y_0 \subseteq Y$ with $A \subseteq Y_0$. Thus, it is not surprising that in order to get "good" estimates one has to impose geometric conditions on the space (in our case, on U). Indeed, for an analogous result in the space of continuous functions [4, 19] one loses the factor 2 if the space U does not have certain geometric properties (the factor 2 can not be decreased by the examples in [12]). The geometric property which turned out useful for that case, is the following

Definition 1. We say that a Banach space U has the *L -retraction property*, if for each separable subspace $U_0 \subseteq U$ there exists a function $R : U \rightarrow U$ with the following properties:

1. The range of R is separable, and $R(u) = u$ on U_0 .
2. R satisfies a Lipschitz condition with constant L .

If $L = 1$, we say that U has the *retraction property*.

We shall also need another geometric property:

Definition 2. We say that a Banach space U has the *(p, q) -exhaustion property*, if each separable subspace $U_0 \subseteq U$ is contained in a separable subspace $V \subseteq U$ with the following properties: There are bounded linear projections $P_k : V \rightarrow V$ and numbers q_k such that:

1. The range $U_k = P_k(V)$ is a finite-dimensional subspace, where $U_1 \subseteq U_2 \dots$ and $V = \overline{\cup U_k}$.
2. We have $\limsup \|P_k\| \leq p$, $\limsup q_k \leq q$, and

$$\|u - P_k u\| \leq q_k \text{ dist}(u, U_k) \quad (u \in U_0). \quad (1)$$

If $p = q = 1$, we say that U has the *exhaustion property*.

For applications, it is usually sufficient to put $q_k = \|I - P_k\|$: For this choice, estimate (1) is always satisfied, even for all $u \in V$. Indeed, for any $u_k \in U_k$ we have $|u - P_k u| = |(I - P_k)(u - u_k)| \leq \|I - P_k\| |u - u_k|$.

Let us give some examples. Recall that a Banach space U is called *weakly compactly generated*, if there is some weakly compact set $K \subseteq U$ whose linear span is dense in U . All separable spaces and all reflexive spaces are weakly compactly generated.

Example 1. Each separable Banach space U has the retraction property. More general, if one assumes the axiom of choice, each weakly compactly generated Banach space has the retraction property. Indeed, if U is weakly compactly generated, the existence of a (linear!) projection R onto a separable subspace containing U_0 is proved in [3] with the axiom of choice (see also [8: Chapter 5, §2/Theorem 3]; for related results see [7]).

Example 2. Each Hilbert space has the retraction and the exhaustion property: One may choose R as the projection onto the element of best approximation in U_0 which is the orthonormal projection onto U_0 . Observe that in this way a countable form of the axiom of choice is sufficient (we shall assume the so-called axiom of dependent choices throughout).

Example 3. Each separable Banach space U with a (Schauder) base has the retraction and the (p, q) -exhaustion property. Indeed, it is well-known that the canonical projections P_k are bounded (uniformly, by the uniform boundedness principle); see, e.g., [24: Chapter I, §3] or [14: Chapter 1]. Hence, we may put $p = \limsup \|P_k\|$ and $q = \limsup \|I - P_k\|$. In particular, each Banach space with a monotone base (i.e. $\|P_k\| \equiv 1$) has the $(1, 2)$ -exhaustion property.

Example 4. Each space $U = L_p(S, \mathbb{R}^n)$ ($1 \leq p < \infty$) with the counting measure on S (e.g. $U = l_p$ or $U = l_p(S)$) has the retraction and the exhaustion property. Indeed, if $U_0 \subseteq U$ is separable, it has countable support E . Put $R = P_E$, and observe that the canonical projections P_k for the canonical base of $V = P_E U$ satisfy $\|P_k\| = \|I - P_k\| = 1$.

Example 5. Each finite-dimensional space has the retraction and the exhaustion property.

To formulate our results as general as possible, we recall two other measures of non-compactness:

The *Kuratowski measure of non-compactness* α of a set A in a metric space is defined as the infimum of all $\varepsilon > 0$ such that A admits a covering of finitely many sets of diameter less than ε .

The *inner Hausdorff measure of non-compactness* χ_i of a set A in a metric space is defined as the infimum of all $\varepsilon > 0$ such that A has a finite ε -net in A , i.e. $\chi_i(A) = \chi_A(A)$.

The Kuratowski and the inner Hausdorff measures of non-compactness have the advantage that they do not depend on the underlying space (only on the metric). The following estimates hold:

$$\chi(A) \leq \chi_i(A) \leq \alpha(A) \leq 2\chi(A).$$

One should be aware that χ_i is (in contrast to χ and α) not monotone, in general.

Our main results are the following sufficient compactness criteria (which, as we shall see, are even necessary if U has finite dimension). For a subset $M \subseteq X$ of a preideal space, we define the expression

$$\omega(M) = \inf_{(E_k)} \sup_{x \in M} \inf_{u_1, \dots, u_n \in U} \left\| x - \sum_{k=1}^n u_k \chi_{E_k} \right\| \tag{2}$$

where the infimum is taken over all systems of finitely many pairwise disjoint sets E_1, \dots, E_n with $\chi_{E_k} \in X$.

Theorem 1. *Let X be a regular preideal space of functions $x : S \rightarrow U$ with a Banach space U , and $M \subseteq X$ be bounded and have equicontinuous norm in X . Assume that there is a function $y \in X$ with*

$$\sup_n \chi_i(M_0(s) \cap K_n) \leq y(s) \text{ a.e. } (M_0 \subseteq M \text{ countable}) \tag{3}$$

for some bounded Borel sets $K_n \supseteq \{u \in U : |u| \leq n\}$. If M is countable, it suffices that (3) holds for $M_0 = M$. If U has the L -retraction property, we may replace (3) by

$$\sup_n L\chi(M_0(s) \cap K_n) \leq y(s) \text{ a.e. } (M_0 \subseteq M \text{ countable}). \tag{4}$$

Then we have the estimate

$$\chi_i(M) \leq 2(\omega(M) + 2\|y\|). \tag{5}$$

Moreover, if M is separable (in X), we even have

$$\alpha(M) \leq 2(\omega(M) + 2\|y\|). \tag{6}$$

If U has the (p, q) -exhaustion property, we have

$$\chi_i(M) \leq 2(p\omega(M) + q\|y\|), \tag{7}$$

and if additionally M is separable, even

$$\chi_X(M) \leq p\omega(M) + q\|y\|. \tag{8}$$

For the scalar case $U = \mathbb{R}$, Theorem 1 implies that a bounded set $M \subseteq X$ is precompact if it has absolutely continuous norm and satisfies $\omega(M) = 0$. This is a special case of Krasnoselskii's compactness criterion in the spaces $L_p(S, \mathbb{R})$ (see, e.g., [13: Lemma 1.1]): A set $M \subseteq L_p(S, \mathbb{R})$ is precompact if and only if it is precompact in measure and has absolutely continuous norm. In this sense, the condition $\omega(M) = 0$ means that M is precompact in measure. We will make this more precise in Corollary 1. On the other hand, the condition $\omega(M) = 0$ may be interpreted as some "equicontinuity" condition (in the norm) for the set M : This condition means that it is possible to approximate the functions in M uniformly by "step functions" with fixed steps. Thus, one might suspect

some connections between the condition $\omega(M) = 0$ and the compactness criterion of Riesz: A set $M \subseteq L_p(\mathbb{R}, \mathbb{R})$ is precompact if and only if it is equicontinuous in the norm in the sense that

$$\lim_{h \rightarrow 0} \sup_{x \in M} \|x(\cdot + h) - x(\cdot)\|_p = 0. \tag{9}$$

This result was generalized for ideal spaces in [11]. The connection of (9) with $\omega(M) = 0$ will become clear in Theorem 3.

Let us remark that the assumption that X be only a preideal space is slightly “too general” in Theorem 1, since each regular preideal space is a dense subspace of a regular ideal space (at least, if the underlying measure space is σ -finite). To see this, observe that it suffices to prove that X is a dense subspace of an ideal space \tilde{X} , because the regular part of \tilde{X} is closed by [27: Theorem 3.3.2]. To verify that X is a subspace of an ideal space, it suffices to consider $U = \mathbb{R}$ in view of [27: Theorem 3.2.1]. But for this case the claim has been proved in [17] (see also [15, 16]) (X has the property $(A, 0)$ by the regularity).

In the space $X = L_1(S, U)$ we may even drop the conditions on the geometry of U and weaken the separability assumption on M for (8), if we modify (2):

Theorem 2. *Let $X = L_1(S, U)$ with a Banach space U , and $M \subseteq X$ be bounded and have equicontinuous norm in X . Assume that there is a function $y \in X$ such that (3) holds. If M is countable, it suffices that (3) holds for $M_0 = M$; if U has the L -retraction property, we may replace (3) by (4). Put*

$$\omega_1(M) = \inf_{(E_k)} \sup_{x \in M} \left\| x - \sum_{k=1}^n \left(\frac{1}{\text{mes } E_k} \int_{E_k} x(s) ds \right) \chi_{E_k} \right\| \tag{10}$$

where the infimum is taken over all systems of finitely many pairwise disjoint sets E_1, \dots, E_n of positive finite measure. Then we have the estimate

$$\chi_X(M) \leq \omega_1(M) + 2\|y\|. \tag{11}$$

If either M is separable (in X) or U is separable, we even have the estimate

$$\chi_X(M) \leq \omega_1(M) + \|y\|. \tag{12}$$

For Theorem 1 it is worth noting that, for a separable Banach space U , the set $M \subseteq X$ is usually separable in X , because X itself is separable: We call a measure space S *separable*, if the system of measurable sets of finite measure with the metric $d(A, B) = \text{mes}(A \Delta B)$ is separable. In particular, $S = \mathbb{R}^n$ is separable by this definition. This is the definition used in [30]. It can be proved that each regular preideal space X of functions $x : S \rightarrow U$ is separable if S and U both are separable. This fact is not evident, but we skip the proof.

From this point of view, it is not so surprising that we already get a better estimate in Theorem 2 if U is separable. However, the question remains open whether one also gets a sharper estimate in Theorem 1, if U is separable but M (and thus S) is not.

Observe that we always have $\omega(M) \leq \omega_1(M)$. Hence, Theorem 1 might still provide a slightly better estimate for $\chi_X(M)$ than Theorem 2 (if U has a nice geometry and M is separable). However, if one is only interested in the question whether M is precompact, the theorems are equivalent:

Lemma 1. Let $X = L_1(S, U)$ with a Banach space U , and $M \subseteq X$. Then we have

$$\omega(M) \leq \omega_1(M) \leq 2\omega(M). \quad (13)$$

Proof. For any finite system of pairwise disjoint sets E_1, \dots, E_n of positive finite measure and any $x, y \in X$ we have

$$\begin{aligned} & \left\| \sum_{k=1}^n \left(\frac{1}{\text{mes} E_k} \int_{E_k} x(s) ds \right) \chi_{E_k} - \sum_{k=1}^n \left(\frac{1}{\text{mes} E_k} \int_{E_k} y(s) ds \right) \chi_{E_k} \right\| \\ &= \sum_{k=1}^n \left| \int_{E_k} x(s) ds - \int_{E_k} y(s) ds \right| \\ &\leq \sum_{k=1}^n \int_{E_k} |x(s) - y(s)| ds \\ &\leq \|x - y\|. \end{aligned}$$

Hence, two applications of the triangle inequality imply

$$\begin{aligned} & \left\| x - \sum_{k=1}^n \left(\frac{1}{\text{mes} E_k} \int_{E_k} x(s) ds \right) \chi_{E_k} \right\| \\ &\leq \left\| y - \sum_{k=1}^n \left(\frac{1}{\text{mes} E_k} \int_{E_k} y(s) ds \right) \chi_{E_k} \right\| + 2\|x - y\|. \end{aligned} \quad (14)$$

For any $C > \omega(M)$ we find pairwise disjoint sets E_1, \dots, E_n of positive finite measure such that

$$\sup_{z \in M} \inf_{u_1, \dots, u_n \in U} \left\| z - \sum_{k=1}^n u_k \chi_{E_k} \right\| < C.$$

In particular, for any $x \in M$ we find a function $y = \sum u_k \chi_{E_k}$ with $\|x - y\| \leq C$. Now (14) implies

$$\left\| x - \sum_{k=1}^n \left(\frac{1}{\text{mes} E_k} \int_{E_k} x(s) ds \right) \chi_{E_k} \right\| \leq 2\|x - y\| \leq 2C$$

and so $\omega_1(M) \leq 2C$ ■

From the proof we can also see the following stability property of ω and ω_1 :

Lemma 2. Let X be a preideal space of functions $x : S \rightarrow U$, and $M, M_0 \subseteq X$. If there is some $\delta > 0$ such that for each function $x \in M$ there exists a function $y \in M_0$ with $\|x - y\| \leq \delta$, then $\omega(M) \leq \omega(M_0) + \delta$. In the case $X = L_1(S, U)$, we also have $\omega_1(M) \leq \omega_1(M_0) + 2\delta$.

Proof. The statement for $\omega(M)$ is a straightforward application of the triangle inequality. Thus, let $X = L_1(S, U)$. For any $C > \omega_1(M_0)$, we find finitely many pairwise disjoint sets E_1, \dots, E_n of positive finite measure such that

$$\left\| y - \sum_{k=1}^n \left(\frac{1}{\text{mes} E_k} \int_{E_k} y(s) ds \right) \chi_{E_k} \right\| \leq C \quad (y \in M_0).$$

For any $x \in M$ we find some $y \in M$ with $\|x - y\| \leq \delta$. By (14) and the triangle inequality, we get

$$\left\| x - \sum_{k=1}^n \left(\frac{1}{\text{mes}E_k} \int_{E_k} x(s) ds \right) \chi_{E_k} \right\| \leq C + 2\delta.$$

Since $x \in M$ was arbitrary, this shows $\omega_1(M) \leq C + 2\delta$ ■

Except for possibly the precompactness of $M(s)$, the conditions of Theorems 1 and 2 are even necessary for the precompactness of M :

Proposition 1. *Let X be a regular preideal space of functions $x : S \rightarrow U$ with a Banach space U . Then we have*

$$\chi_X(M) \geq \omega(M), \tag{15}$$

and, in the case $X = L_1(S, U)$, also

$$\chi_X(M) \geq \frac{1}{2}\omega_1(M). \tag{16}$$

Moreover,

$$\chi_X(M) \geq \inf_{\text{mes}E < \infty} \sup_{x \in M} \|P_{E^c}x\| \tag{17}$$

and

$$\chi_X(M) \geq \limsup_{\delta \rightarrow 0} \sup_{\text{mes}D \leq \delta} \sup_{x \in M} \|P_Dx\|. \tag{18}$$

In particular, if M is precompact, then M has equicontinuous norm and satisfies $\omega(M) = 0$ (and $\omega_1(M) = 0$ in the case $X = L_1(S, U)$).

Proof. Observe that (16) is a consequence of (15) and (13). Let us now prove the statement for the case that $M \subseteq X$ is a finite set.

That the right-hand side of (17) and (18) vanishes in this case follows immediately from the definition of regular spaces and the triangle inequality. To see that $\omega(M) = 0$, let $\varepsilon > 0$ be given and choose a set E of finite measure such that $\|P_Ex\| < \varepsilon$ for each $x \in M$. Since the support of P_Ex has finite measure, there is a sequence y_n of simple functions with $y_n(s) \rightarrow P_Ex(s)$ and $|y_n(s)| \leq |P_Ex(s)|$ for almost all s (see, e.g., [14: Lemma 4.1.1]). Since X is regular, Lebesgue's dominated convergence theorem for regular preideal spaces [27: Theorem 3.3.6] implies $\|y_n - P_Ex\| \rightarrow 0$ ($n \rightarrow \infty$). Thus, for any $\varepsilon > 0$ and any $x \in M$, we find a simple function $y = \sum u_k \chi_{E_k}$ with $\chi_{E_k} \in X$ and $\|y - P_Ex\| \leq \varepsilon$. By considering a common refinement, we may assume that the sets E_k are pairwise disjoint and independent of x (M is finite!). Since $\|x - y\| = \|(P_Ex - y) + P_{E^c}x\| \leq 2\varepsilon$, we have $\omega(M) \leq 2\varepsilon$.

Now we attack the general case. Given $C > \chi_X(M)$, we find a finite C -net $N \subseteq X$ for M . By what we had proved, we find for each $\varepsilon > 0$ some set E of finite measure, some $\delta > 0$, and finitely many pairwise disjoint sets E_1, \dots, E_n with $\chi_{E_k} \in X$ such that for each $y \in N$ we have the estimates $\|P_{E^c}y\| \leq \varepsilon$, $\|P_Dy\| \leq \varepsilon$ ($\text{mes}D \leq \delta$), and

$$\inf_{u_1, \dots, u_n \in U} \left\| y - \sum_{k=1}^n u_k \chi_{E_k} \right\| \leq \varepsilon.$$

For each $x \in M$, we find some $y \in N$ with $\|x - y\| \leq C$. But now the triangle inequality implies $\|P_{E^\varepsilon}x\| \leq C + \varepsilon$, $\|P_Dx\| \leq C + \varepsilon$ ($\text{mes}D \leq \delta$), and

$$\inf_{u_1, \dots, u_n \in U} \left\| x - \sum_{k=1}^n u_k \chi_{E_k} \right\| \leq C + \varepsilon.$$

We may conclude that (17), (18) and (15) are satisfied ■

Even for $U = \mathbb{R}^n$ our compactness criteria are apparently new, and give even a precise formula for the Hausdorff measure of non-compactness for sets with equicontinuous norm (be aware that the apparently simpler formula in L_p given in [1] is false; see also [22]):

Corollary 1. *Let X be a regular preideal space of functions $x : S \rightarrow U$ with a finite-dimensional space U . Then a set $M \subseteq X$ is precompact if and only if the following holds:*

1. M is bounded and has equicontinuous norm.
2. $\omega(M) = 0$ or, if $X = L_1(S, U)$, equivalently $\omega_1(M) = 0$.

Moreover, if M is bounded and has equicontinuous norm and the measure space S is separable, we have the identity

$$\chi_X(M) = \omega(M).$$

Under the assumption that M be precompact in measure, another formula for $\chi_X(M)$ was given in [29] (related results in L_p can also be found in [28]).

Let us remark that the precompactness of $M(s)$ is far from being necessary for the precompactness of M . Let us give a typical class of counterexamples in $L_p([0, 1], U)$:

Example 6. Let $X = L_p([0, 1], U)$ with $1 \leq p < \infty$. Given an arbitrary countable set $U_0 \subseteq U$, we may define a precompact countable set $M \subseteq X$ such that $M(s) \equiv U_0$ for all s .

Indeed, without loss of generality let $0 \in U_0$ (otherwise we consider a translation of U_0 and add a constant function afterwards). Choose a sequence $u_k \in U_0$ such that $U_0 = \{u_1, u_2, \dots\}$ and $|u_k| \leq 2^{\frac{k}{2p}}$ for all sufficiently large k (repeat some elements, if necessary). Now define a sequence x_n in the following way: For $n = 2^k + j$ ($j = 0, \dots, 2^k - 1$) put $x_n = u_k \chi_{[j2^{-k}, (j+1)2^{-k}]}$. For all sufficiently large n we then have $\|x_n\| \leq 2^{-\frac{k}{2p}} \rightarrow 0$, hence the set $M = \{x_1, x_2, \dots\}$ is precompact. Moreover, $M(s) = U_0$ for each s .

However, in the theory of vector-valued integral and functional equations, one usually has an estimate for $\chi(M(s))$, and our results are applicable. However, $\omega(M) = 0$ may be interpreted as some "equicontinuity" condition on the family M of functions. Thus, it is not surprising that this condition may also be formulated similarly to the compactness criterion of Riesz, at least for locally compact groups:

Theorem 3. *Let S be a locally compact Hausdorff group with a left Haar measure and unit element 0 . Let X be a regular ideal space of locally integrable functions $x : S \rightarrow U$ which contains all functions χ_K with compact $K \subseteq S$. Let $M \subseteq X$ satisfy*

$$\lim_{h \rightarrow 0} \sup_{x \in M} \sup_{t \in S} \|x^{h \circ t} - x^t\| = \lim_{h \rightarrow 0} \sup_{x \in M} \|x^{-h} - x\| = 0 \tag{19}$$

where we have put $x^h(s) = x(h \circ s)$ and $x^{-h}(s) = x(s \circ h)$. Assume that either $\text{supp } M$ exists and is contained in a compact set, or that M has equicontinuous norm and even

$$\lim_{\delta \rightarrow 0} \sup_{\text{mes } D \leq \delta} \sup_{t \in S} \sup_{x \in M} \|P_D x^t\| = 0. \tag{20}$$

Then $\omega(M) = 0$.

Proof. Let us first assume that $\text{supp } M \subseteq K$ for some compact set K . Let $\epsilon > 0$ be given. Choose some compact neighborhood H of 0 such that

$$\|x^{-h} - x\| < \epsilon \quad (x \in M, h \in H).$$

Observe that $0 < \text{mes } H < \infty$. Since all functions $x \in X$ are locally integrable, we may define the "inverse" convolutions ("Steklov functions")

$$Fx(t) = \frac{1}{\text{mes } H} \int_H x(t \circ s) ds = \int_S x^t(s)y(s) ds$$

where we have put $y(s) = (\text{mes } H)^{-1} \chi_H$. Observe that the function $w(t, s) = x(t \circ s)y(s)$ vanishes for $s \notin H$ or for $t \notin H^{-1} \circ K =: K_0$. Since K_0 and H are compact and thus have finite measure, we may apply Fubini-Tonelli's theorem on the measure space $K_0 \times H$: The Borel function $v(t, s) = t \circ s$ has a (compact) range of finite measure and the property that preimages of null sets are null sets. Indeed, if $N \subseteq S$ is a null set, it is contained in a Borel null set N_0 . Then $v^{-1}(N_0)$ is a Borel set, in particular measurable, and by Fubini-Tonelli we have

$$\text{mes}(v^{-1}(N)) \leq \int_{K_0} \int_H \chi_{v^{-1}(N_0)}(t, s) ds dt = \int_{K_0} \int_H \chi_{N_0}(t \circ s) ds dt = 0.$$

We may conclude that w is measurable on $K_0 \times H$. Indeed, if x_n are simple Borel functions on the range of v which converge a.e. to x , then $x_n \circ v$ is measurable and converges a.e. to $x \circ v$. Hence, also the function $z(t, s) = (x(t \circ s) - x(t))y(s)$ is measurable on $K_0 \times H$. Since X is regular, this implies by [27: Theorem 4.4.2] that the abstract function $Z : H \rightarrow X$, $Z(s) = (x^{-s} - x)y(s)$ is measurable. By [27: Theorem 4.4.3] the integral over this abstract function (the following formula shows that it exists) may be evaluated as the pointwise integral (a.e.), and thus we have the estimate

$$\begin{aligned} \|Fx - x\| &= \left\| t \mapsto \int_S (x^t(s) - x(t))y(s) ds \right\| \\ &= \left\| t \mapsto \int_H (x^{-s}(t) - x(t))y(s) ds \right\| \\ &= \left\| \int_H Z(s) ds \right\| \leq \int_H \|Z(s)\| ds = \int_S \|x^{-s} - x\| y(s) ds \leq \epsilon. \end{aligned}$$

It follows with Lemma 2 for the set $FM = \{Fx : x \in M\}$ that $\omega(M) \leq \omega(FM) + \varepsilon$.

To estimate $\omega(FM)$, we argue as follows: Since all functions in $P_H X$ are integrable, the function y must even belong to the associate space $(P_H X)'$ (see, e.g., [27: Theorem 3.4.2]), i.e. we may apply Hölder's inequality for ideal spaces: For all $t, h \in S$, and all $x \in X$, we get the estimate

$$|Fx(h \circ t) - Fx(t)| = \left| \int_S (x^{h \circ t}(s) - x^t(s))y(s) ds \right| \leq \|x^{h \circ t}(s) - x^t\| \|y\|_{(P_H X)'}$$

By assumption, we find an open neighborhood O of 0 such that for all $h \in O$ the right-hand side is less than $\frac{\varepsilon}{\|\chi_{K_0}\|}$ for each $x \in M$ and each $t \in S$. From what we have shown above, the support of FM is contained in K_0 . By the compactness there exist finitely many $t_j \in S$ such that K_0 is covered by the sets $O \circ t_j$. By considering a common refinement, we may divide K_0 into finitely many pairwise disjoint sets E_1, \dots, E_m such that each E_k is contained in some set $O \circ t_{j(k)}$. Given $x \in M$, we put $u_k = Fx(t_{j(k)})$. Then for each $s \in E_k$, $s = h \circ t_{j(k)}$ with some $h \in O$, the estimate

$$|Fx(s) - u_k| = |Fx(h \circ t_{j(k)}) - Fx(t_{j(k)})| \leq \frac{\varepsilon}{\|\chi_{K_0}\|}$$

holds. Summing up these inequalities, we find

$$\left| Fx(s) - \sum_{k=1}^n u_k \chi_{E_k}(s) \right| \leq \varepsilon \frac{\chi_{K_0}(s)}{\|\chi_{K_0}\|}$$

Taking the norm in X for the functions on both sides of this inequality, and observing that E_1, \dots, E_m was independent from $x \in M$, we may conclude that $\omega(FM) \leq \varepsilon$. Hence, $\omega(M) \leq 2\varepsilon$.

Let us now assume that M has equicontinuous norm and (20) holds. Given $\varepsilon > 0$, we find a set E of finite measure and $\delta > 0$ such that $\|P_{E^c}x\| \leq \varepsilon$ and $\|P_Dx\| \leq \varepsilon$ ($\text{mes}D \leq \delta$). Since E is of finite measure and inner-regular, it contains a compact set $K \subseteq E$ with $\text{mes}(E \setminus K) \leq \delta$. We may conclude that $\|P_{K^c}x\| = \|P_{E^c}x + P_{E \setminus K}x\| \leq 2\varepsilon$ for all $x \in M$. Hence, Lemma 2 implies $\omega_X(M) \leq \omega_X(P_K M) + 2\varepsilon$. Thus, if we can apply the statement for $P_K M$ in place of M , we are done.

To show that $P_K M$ satisfies (19), we first prove that the measure of the sets

$$K_{t,h} = (t^{-1} \circ K)\Delta(t^{-1} \circ h^{-1} \circ K) \quad \text{and} \quad K_h = K\Delta(K \circ h^{-1})$$

tends to 0 as $h \rightarrow 0$, uniformly for $t \in S$. Indeed, given $\varepsilon > 0$, there is a continuous function z with compact support such that $\|z - \chi_K\|_{L_1} < \varepsilon$. Then we have $\|z^h - \chi_K^h\|_{L_1} < \varepsilon$ and $\|z^{-h} - \chi_K^{-h}\|_{L_1} < \Delta(h)\varepsilon$, where Δ denotes the modular function of the Haar measure. We thus find

$$\begin{aligned} \text{mes}(K_{t,h}) &= \|\chi_K^t - \chi_K^{h \circ t}\|_{L_1} \\ &= \|\chi_K - \chi_K^h\|_{L_1} \\ &\leq \|\chi_K - z\|_{L_1} + \|z - z^h\|_{L_1} + \|z^h - \chi_K^h\|_{L_1} \\ &\leq 2\varepsilon + \|z - z^h\|_{L_1} \end{aligned} \tag{21}$$

and

$$\begin{aligned} \text{mes}(K_h) &= \|\chi_K - \chi_K^{-h}\|_{L_1} \\ &\leq \|\chi_K - z\|_{L_1} + \|z - z^{-h}\|_{L_1} + \|z^h - \chi_K^h\|_{L_1} \\ &\leq (1 + \Delta(h))\varepsilon + \|z - z^{-h}\|_{L_1}. \end{aligned} \tag{22}$$

In view of the equicontinuity of z and $\Delta(h) \rightarrow 1$ as $h \rightarrow 0$, we thus have $\sup_t \text{mes}(K_{t,h}) \rightarrow 0$ and $\text{mes}(K_h) \rightarrow 0$, as claimed.

Hence, by (20), we find for each $\varepsilon > 0$ some neighborhood H of 0 such that for all $h \in H, t \in S, x \in X$ the inequalities $\|P_{K_{t,h}}x^t\| < \varepsilon$ and $\|P_{K_h}x\| < \varepsilon$ hold. In view of

$$\begin{aligned} |(P_Kx)^{hot}(s) - (P_Kx)^t(s)| &= \left| \chi_K^{hot}(s)(x^{hot}(s) - x^t(s)) - (\chi_K^{hot}(s) - \chi_K^t(s))x^t(s) \right| \\ &\leq |x^{hot}(s) - x^t(s)| + |P_{K_{t,h}}x^t(s)| \end{aligned}$$

and

$$\begin{aligned} |(P_Kx)^{-h}(s) - (P_Kx)(s)| &= \left| \chi_K^{-h}(s)(x^{-h}(s) - x(s)) + (\chi_K^{-h}(s) - \chi_K(s))x(s) \right| \\ &\leq |x^{-h}(s) - x(s)| + |P_{K_h}x(s)| \end{aligned}$$

we thus find that

$$\limsup_{h \rightarrow 0} \sup_{z \in M} \sup_{t \in S} \|(P_Kx)^{hot} - (P_Kx)^h\| \leq \varepsilon$$

and

$$\limsup_{h \rightarrow 0} \sup_{z \in M} \|(P_Kx)^{-h} - P_Kx\| \leq \varepsilon.$$

Hence, the set P_KM satisfies (19), as claimed ■

If the ideal space X is even invariant under left-translations of S in the sense that $x \in X$ implies $x^t \in X$ and $\|x\| = \|x^t\|$ (like, e.g., $X = L_p(S, U)$), then condition (19) may of course equivalently be replaced by

$$\lim_{h \rightarrow 0} \sup_{x \in M} \|x^h - x\| = \lim_{h \rightarrow 0} \sup_{x \in M} \|x^{-h} - x\| = 0.$$

Moreover, in this case condition (20) may be dropped, since it already is a consequence of the fact that M has equicontinuous norm.

Corollary 2. *Let $X = L_p(\mathbb{R}^n, U)$ ($1 \leq p < \infty$) with a Banach space U . If U has the retraction property, we put $L = 1$, otherwise $L = 2$. Let $M \subseteq X$ have equicontinuous norm and satisfy*

$$\lim_{h \rightarrow 0} \sup_{x \in M} \int_S |x(s+h) - x(s)|^p ds = 0.$$

Then for each $y \in X$ which satisfies $y(s) \geq \chi(M_0(s))$ a.e. for each countable $M_0 \subseteq M$, we have

$$\chi_X(M) \leq 2L\|y\|.$$

Moreover, the factor 2 may be dropped if either M is separable and U has the exhaustion property, or if $p = 1$ and either M or U are separable. If U has the retraction property,

the family $\mathcal{M} = \{\chi(M_0(\cdot)) : M_0 \subseteq M \text{ countable}\}$ consists of measurable functions, and thus we may choose $y = \sup \mathcal{M}$.

The measurability of the function $\chi(M_0(\cdot))$ has been proved in [4] (see also [12, 19]). Here, $\sup \mathcal{M}$ denotes the smallest upper bound with respect to the order "almost everywhere"; this supremum exists (and is measurable) by a theorem of Kantorovich, if the underlying measure space is σ -finite (see, e.g., [30]).

A special version of Corollary 2 for $L_1([a, b], U)$ with a separable Banach space U and countable M has been proved in [20] (observe that Corollary 2 shows that it is not necessary for that result that M be countable; moreover, our boundedness assumptions are much weaker). An analogous version of the mentioned result from [20] for the Kuratowski measure of non-compactness can be found in [25]. Applications of these special versions to Volterra equations in Banach spaces can be found in the earlier mentioned papers [20, 21, 23, 26].

2. Proofs of Theorems 1 and 2

The heart of the proofs is the following result which has implicitly been shown in [18: Propostion 1.4] (see also [19: Proposition 2]).

Lemma 3. *Let U be a normed space, and $U_1 \subseteq U_2 \subseteq \dots$ be finite-dimensional subspaces. Then for any countable bounded $M \subseteq U$, $M = \{u_1, u_2, \dots\}$, we have the estimate*

$$\chi_U(M) \leq \lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \text{dist}(u_n, U_k). \quad (23)$$

Moreover, if U is separable, then there exists a sequence of finite-dimensional subspaces $U_1 \subseteq U_2 \subseteq \dots$ with $U = \overline{\cup U_k}$; and for any such sequence we have the equality

$$\chi_U(M) = \lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \text{dist}(u_n, U_k). \quad (24)$$

Proof. Let C be larger than the right-hand side of (23). Then we find some k and n_0 with the following property: For any $n \geq n_0$ there is some $v_n \in U_k$ with $\|u_n - v_n\| \leq C$. Since the set of all v_n is a bounded subset of the finite-dimensional space U_k , we find for any $\varepsilon > 0$ a finite ε -net $N \subseteq U_k$ for this set. In particular, for any $n \geq n_0$ we find some $u \in N$ with $\|u_n - u\| \leq C + \varepsilon$. Hence, $N \cup \{u_1, \dots, u_{n_0}\}$ is a finite $(C + \varepsilon)$ -net for M , and so $\chi_U(M) \leq C + \varepsilon$. This proves (23).

If U is separable, there exists a countable dense subset $\{w_1, w_2, \dots\}$, and one may choose $U_k = \text{span}\{w_1, \dots, w_k\}$. To see (24), let $C > \chi_U(M)$ and N be a finite C -net for M . Given $\varepsilon > 0$, we find some k with $\text{dist}(u, U_k) \leq \varepsilon$ for each $u \in N$. This implies $\text{dist}(u_n, U_k) \leq C + \varepsilon$ for each n , and it follows that the right-hand side of (24) is bounded by $C + \varepsilon$, hence bounded by $\chi_U(M)$ ■

The crucial point in Lemma 3 is that it allows to calculate χ from the "outside", i.e. without explicit knowledge of a finite ε -net. To get an estimate if U does not have a nice geometry, we need the following fact.

Lemma 4. *Each separable Banach space U_0 is isometrically and linearly embedded into a separable Banach space V with the (1, 2)-exhaustion property.*

Proof. By [5: Chapter XI, Theorem 9 (§8)], U_0 can be mapped isometrically and linearly onto a closed linear subspace of the space $V = C([0, 1])$. Since V has a monotone base (namely the classical Schauder system; see, e.g., [24: Chapter I, §3] or [14: Chapter 1]), Example 3 implies that V has the (1, 2)-exhaustion property ■

We emphasize that the proof of Lemma 4 does not require the (uncountable) axiom of choice: The Hahn-Banach extension theorem (which is invoked in the cited result) can be proved “constructively” in separable spaces (see [10]).

We do not know whether any separable Banach space may be embedded (isometrically) into a separable Banach space with the (p_0, q_0) -property with $q_0 < 2$. If this is the case, one may strengthen Theorem 1. More precisely, if $C([0, 1])$ may be embedded into a separable Banach space with the (p_0, q_0) -exhaustion property, then our proof shows that we may replace (5) and (6) by the estimates

$$\chi_i(M) \leq 2(p_0\omega(M) + q_0\|y\|) \quad \text{and} \quad \alpha(M) \leq 2(p_0\omega(M) + q_0\|y\|),$$

respectively. However, it is not very reasonable that better constants than in Lemma 4 are possible: As a matter of fact, the “universal” space $U = C([0, 1])$ has the (p, q) -exhaustion property only for $p \geq 1$ and $q \geq 2$. The latter follows from the result in [6] which states that any compact operator K in U satisfies $\|I + K\| = 1 + \|K\|$: Let $U_0 = U$, and $P_k : V = U \rightarrow U_k$ be as in Definition 2. Since P_k is compact, we have $\|I - P_k\| = 1 + \|P_k\| \geq 2$. Thus, for any $\varepsilon > 0$ we find some $u \in U_0 = V$, $u \neq 0$, such that $|u - P_k u| \geq (2 - \varepsilon)|u| \geq (2 - \varepsilon)\text{dist}(u, U_k)$. With the notation of Definition 2, this means $q_k \geq 2$ for each k , and so $q \geq 2$, as claimed.

Proof of Theorem 1. Let us first reduce the statement to the case that the functions in M are uniformly dominated by a function $\rho_{\chi_E} \in X$, where E has finite measure:

Suppose that $M \subseteq X$ has equicontinuous norm. Given $\varepsilon > 0$, let E_0 be a set of finite measure with

$$\sup_{x \in M} \|P_{E_0^c} x\| < \varepsilon,$$

and $\delta > 0$ be such that

$$\sup_{\text{mes } D \leq \delta} \sup_{x \in M} \|P_D x\| < \varepsilon.$$

Since E_0 has finite measure, $S_0 = \text{supp } P_{E_0} X$ exists [27: Theorem 2.2.4], and by [27: Theorem 2.2.5] there is a set $E \subseteq S_0$ with $\text{mes}(S_0 \setminus E) < \delta$ such that $\chi_E \in X$. For all $x \in X$ we have $P_{E^c} x = P_{E_0^c} x + P_{S_0 \setminus E} x$, and so

$$\|P_{E^c} x\| \leq 2\varepsilon.$$

Note that $P_E M$ is bounded in X . By [27: Corollary 3.1.3], the set $P_E M$ thus is bounded in measure. In particular, we find some natural number n_0 such that the measure of the set $\{s : |P_E x(s)| > n_0\}$ is less than δ for each $x \in M$. Then also the measure of the

set $D(x) = \{s \in E : x(s) \notin K_{n_0}\}$ is less than δ for each $x \in M$. Now we consider the set

$$M_\epsilon = \{P_{E \setminus D(x)}x : x \in M\}.$$

For each $x \in M$ the corresponding function $z = P_{E \setminus D(x)}x \in M_\epsilon$ satisfies

$$\|x - z\| = \|P_{E^c}x + P_{D(x)}x\| \leq 3\epsilon.$$

Since each measure $\gamma \in \{\chi, \alpha, \chi_i\}$ is continuous with respect to the Hausdorff distance (see, e.g., [1]), we find for each $\epsilon_0 > 0$ some $\epsilon > 0$ such that the sizes $\gamma(M)$ and $\gamma(M_\epsilon)$ differ by at most ϵ_0 . By Lemma 2, the sizes $\omega(M_\epsilon)$ and $\omega(M)$ differ by at most ϵ . Moreover, the corresponding estimate (3) respectively (4) holds for M_ϵ in place of M by our construction. Thus, it suffices to prove the statement for M_ϵ in place of M (for all sufficiently small $\epsilon > 0$). But the set M_ϵ has by construction the additional required property (observe that K_{n_0} is bounded by some number $\rho > 0$).

Thus, we assume without loss of generality that all functions in M are uniformly dominated by $\rho\chi_E \in X$ with $\text{mes}E < \infty$ (and hence take their values in some K_{n_0}).

First, assume additionally that M is countable. Since the support of each function in M has finite measure, the functions are essentially separably valued (see [9: Section III.6/Theorem 10]). This means that after modifying the (countably many!) functions on a null set, we may assume that they have separable range. For any $C > \omega(M)$ we can find a finite partition E_1, \dots, E_m of E (with measurable sets E_i) with the following property: For each $x \in M$ there is some function $z = z_x$ of the form

$$z = \sum_{i=1}^m u_i \chi_{E_i} \tag{25}$$

with $u_i \in U$ such that $\|x - z\| < C$. Let H denote the closed linear hull of all values of the functions in M and all values of the functions z_x . Then H is separable.

If U has the L -retraction property, we choose R corresponding to Definition 1 (for the subspace H), and denote the closed linear hull of the range of R by U_0 ; otherwise, we put $U_0 = H$ and $R = I$. In both cases, we have

$$\chi_{U_0}(M(s)) \leq y(s) \quad \text{a.e.}$$

Indeed, if U has the L -retraction property, and $N \subseteq U$ is a finite ϵ -net for $M(s)$, then $R(N) \subseteq U_0$ is a finite $L\epsilon$ -net for $R(M(s)) = M(s)$; hence $\chi_{U_0}(M(s)) \leq L\chi(M(s)) \leq y(s)$. In the other case, our assumptions imply $\chi_{U_0}(M(s)) \leq \chi_i(M(s)) \leq y(s)$.

In order to prove (8), choose P_n, V, U_n, p , and q corresponding to Definition 2. If we want to prove (6) instead, we choose V as in Lemma 4, and then P_n and U_n as in Definition 2 (corresponding to $U_0 := V$); in this case we put $p = 1, q = 2$, and tacitly identify U_0 with a subspace of V in the following.

To clarify notation, we denote by X_V the preideal space of functions $x : S \rightarrow V$ which has the same real form than X . Let X_k denote the finite-dimensional subspace of all functions of the form (25) with $u_i \in U_k$. We claim that for each $\epsilon > 0$ there is some K such that

$$\text{dist}_{X_V}(x, X_k) \leq pC + q\|\text{dist}_V(x(\cdot), U_k)\| + \epsilon \quad (x \in M, k \geq K). \tag{26}$$

Indeed, we have

$$\begin{aligned} |P_k z_x(s) - x(s)| &= |P_k(z_x(s) - x(s)) + (P_k x(s) - x(s))| \\ &\leq \|P_k\| |z_x(s) - x(s)| + q_k \text{dist}_V(x(s), U_k) \end{aligned}$$

which implies that

$$\|P_k z_x - x\| \leq \|P_k\| \|z_x - x\| + q_k \|\text{dist}_V(x(\cdot), U_k)\|.$$

Since $P_k z_x \in X_k$, this proves (26).

For definiteness, let $M = \{x_1, x_2, \dots\}$. Putting $x_{n,k}(s) = \sup_{m \geq n} \text{dist}_V(x_m(s), U_k)$, we have by (26) that

$$\sup_{m \geq n} \text{dist}_{X_V}(x_m, X_k) \leq q \|x_{n,k}\| + pC + \varepsilon. \tag{27}$$

Since $|x_{n,k}(s)| \leq \rho_{X_E}(s)$ and

$$x_{n,k}(s) \rightarrow y_k(s) = \limsup_{n \rightarrow \infty} \text{dist}_V(x_n(s), U_k),$$

we may pass to the limit $n \rightarrow \infty$ in (27) by Lebesgue's dominated convergence theorem for preideal spaces (see [27: Theorem 3.3.6]). Thus, we get

$$\limsup_{n \rightarrow \infty} \text{dist}_{X_V}(x_n, X_k) \leq q \|y_k\| + pC + \varepsilon. \tag{28}$$

Observe that (24) implies

$$y_k(s) \rightarrow \lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \text{dist}_V(x_n(s), U_k) = \chi_V(M(s)) \leq \chi_{U_0}(M(s)) \leq y(s).$$

Hence, passing to the limit $k \rightarrow \infty$ in (28), we find

$$\lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \text{dist}_{X_V}(x_n, X_k) \leq q \|y\| + pC + \varepsilon.$$

Now (23) implies $\chi_{X_V}(M) \leq q \|y\| + pC + \varepsilon$. Letting $\varepsilon \rightarrow 0$, we thus find $\chi_{X_V}(M) \leq pC + q \|y\|$. Now the proof of (8) is completed, since $V \subseteq U$ implies $X_V \subseteq X$, and so $\chi_X(M) \leq \chi_{X_V}(M)$. For the proof of (6), we observe that $\alpha(M) \leq 2\chi_{X_V}(M)$ and that $\alpha(M)$ is the same in the space X as in the space X_V .

Now we consider the case that M is not necessarily countable: If M is separable in X , there is a countable subset $M_0 \subseteq M$ with $\overline{M_0} = M$ (see, e.g., [2: Lemma 2.6]). Then we have $\gamma(M) = \gamma(M_0)$ for $\gamma \in \{\alpha, \chi_X, \chi_i\}$. Since evidently $\omega(M_0) \leq \omega(M)$, we get estimate (6) respectively (8) by applying the corresponding estimate for the countable set M_0 .

In general, there exists a countable $M_0 \subseteq M$ such that $\chi_i(M) \leq \alpha(M_0) \leq 2\chi_X(M_0)$ (see, e.g., [4]). Applying estimate (6) respectively (8) for M_0 in place of M , we get the desired estimate (5) respectively (7) ■

The proof of Theorem 2 is based on the following result.

Theorem 4. *Let $X = L_1(S, U)$, and let $M \subseteq X$ be uniformly dominated by an integrable function. Let y be integrable such that*

$$\chi_i(M_0(s)) \leq y(s) \text{ a.e.} \quad (M_0 \subseteq M \text{ countable}). \tag{29}$$

If M is countable, (29) needs to hold only for $M_0 = M$. If U has the L -retraction property, we may replace (29) by

$$L\chi(M_0(s)) \leq y(s) \text{ a.e.} \quad (M_0 \subseteq M \text{ countable}). \tag{30}$$

Then the estimate

$$\chi_i\left(\left\{\int_S x(s) ds : x \in M\right\}\right) \leq 2 \int_S y(s) ds$$

holds. Moreover, if either M is countable or U is separable, we even have

$$\chi\left(\left\{\int_S x(s) ds : x \in M\right\}\right) \leq \int_S y(s) ds. \tag{31}$$

Theorem 4 is proved in [4]. Actually, the result in [4] is only formulated for the case that $L = 1$ and that estimate (29) respectively (30) holds everywhere, but an inspection of the proof shows that the result also holds for $L > 1$ and that the exceptional null set may depend on M_0 . Also, the proof of Theorem 4 is based on Lemma 3. It is worth noting that the idea to use Lemma 3 to prove results like Theorem 4 is apparently due to Mönch [18, 19] (although the proof of [19: Proposition 3] contained a small mistake in the application of Fatou’s lemma which however can be avoided by using Lebesgue’s theorem instead). We remark that in [12] it is shown by means of an example that, if U does not have the retraction property, one may not replace (29) by (30) with $L < 2$ (even for $S = [0, 1]$, $U = l_\infty$ and countable $M \subseteq C(S, U)$).

Let us note that the condition that $M \subseteq X = L_1(S, U)$ be uniformly dominated by an integrable function in Theorem 4 can actually be replaced by the weaker condition that all functions in M have σ -finite support and

$$\lim_{n \rightarrow \infty} \sup_{x \in M} \int_{D_n} |x(s)| ds = 0 \tag{32}$$

for each sequence of measurable sets $D_n \downarrow \emptyset$. Indeed, under the additional assumption that M is a bounded subset of $X = L_1(S, U)$, this has been proved in [4]. The general case may be established by modifying the proof in [4] by isolating certain atoms of the measure space. However, we shall not apply this more general result (although this would allow to prove a slight generalization of Theorem 2).

Proof of Theorem 2. With the same argument as in the proof of Theorem 1, we can reduce the statement to the case that all functions in M are uniformly dominated by a function $\rho \chi_E \in X$. Thus, let us assume this.

For any $C > \omega_1(M)$ we find finitely many pairwise disjoint sets E_1, \dots, E_n of positive finite measure such that

$$\sup_{x \in M} \left\| x - \sum_{k=1}^n \left(\frac{1}{\text{mes} E_k} \int_{E_k} x(s) ds \right) \chi_{E_k} \right\| < C. \tag{33}$$

By Theorem 4, the Hausdorff measure of non-compactness (in U) of the set

$$V_k = \left\{ \int_{E_k} x(s) ds : x \in M \right\} \subseteq U$$

is bounded by

$$c_k := 2 \int_{E_k} y(s) ds$$

(and we may even drop the factor 2, if U is separable or M is countable). This estimate means that, given $\varepsilon > 0$, we find a finite $(c_k + \varepsilon)$ -net $N_k \subseteq U$ for V_k . Now put

$$N = \left\{ \sum_{k=1}^n \frac{1}{\text{mes} E_k} u_k \chi_{E_k} : u_k \in N_k \right\}.$$

For each $x \in M$, there exist $u_k \in N_k$ such that

$$\left| \int_{E_k} x(s) ds - u_k \right| \leq c_k + \varepsilon \quad (k = 1, \dots, n).$$

The function $z = \sum (\text{mes} E_k)^{-1} u_k \chi_{E_k} \in N$ thus satisfies

$$\begin{aligned} & \left\| \sum_{k=1}^n \left(\frac{1}{\text{mes} E_k} \int_{E_k} x(s) ds \chi_{E_k} \right) - z \right\| \\ &= \sum_{k=1}^n \left| \int_{E_k} x(s) ds - u_k \right| \leq \sum_{k=1}^n (c_k + \varepsilon) \leq 2 \int_S y(s) ds + n\varepsilon = 2\|y\| + n\varepsilon. \end{aligned}$$

By (33) and the triangle inequality, this implies $\|x - z\| \leq C + 2\|y\| + n\varepsilon$. Hence, $N \subseteq X$ is a finite $(C + 2\|y\| + n\varepsilon)$ -net for M , i.e. $\chi_X(M) \leq C + 2\|y\| + n\varepsilon$. Now first letting $\varepsilon \rightarrow 0$ (n depends on C !) and then $C \rightarrow \omega_1(M)$, we find (11). If U is separable or M is countable, we may drop the factor 2 in all above formulas.

The case that M is uncountable but separable in X , is exceptional. To get the better estimate (12) in this case, we choose a countable dense $M_0 \subseteq M$ (recall [2: Lemma 2.6]). Then $\chi_X(M) = \chi_X(M_0)$, and by what we have proved so far, $\chi_X(M_0) \leq \omega_1(M_0) + \|y\| \leq \omega_1(M) + \|y\|$ ■

Let us remark that Theorems 1 and 2 hold slightly more general:

1. Instead of requiring that M have equicontinuous norm, it suffices to require that each countable subset of M have equicontinuous norm.

2. Actually, it is not necessary to assume that the sets K_n are bounded.

The generalization of Statement 1 is of interest, e.g., for $X = L_p(S, U)$, if S is not σ -finite. For example, (32) is equivalent to the statement that M has equicontinuous norm in $L_1(S, U)$, provided that M has σ -finite support [27: Lemma 3.3.4]. But the latter is true for countable sets $M \subseteq L_1(S, U)$ while not necessarily for the whole set M .

The generalization of Statement 2 is only interesting in view of the fact that χ_i is not monotone, in general.

Let us briefly sketch, how the proof of Theorem 1 has to be modified to cover these cases. For Statement 1 it suffices to change the order of argumentation: One first has to reduce the statement to countable M (with the same arguments as in the proof). Then the reduction to the case that all functions in M be uniformly dominated by some $\rho \chi_E$ requires only the equicontinuity of the bounded set M .

The changes for Statement 2 are more difficult to describe. Actually, our proof shows for the case that M is countable and all functions are uniformly dominated by some $\rho \chi_E$ slightly more than as is claimed in the statement. Namely, it is not required that $\chi_i(M(s) \cap K_n) \leq y(s)$ but it suffices that $\chi_H(M(s)) \leq y(s)$ for some separable subspace $H \subseteq U$ which contains (essentially) all values of the functions of M . Observing that χ_H is monotone, the other parts of the proof actually reduce the theorem to this special case, if we replace $D(x)$ by $D(x) = \{s : |x(s)| > n_0\}$ (put $\rho = n_0$).

It would lead too far to describe the necessary changes in the proof of Theorem 2 here in detail. Theorem 4 has to be modified appropriately (in particular, one has to introduce the sets K_n already in the statement of Theorem 4).

Let us finally note that Theorem 1 also holds (in principle) for the case that X is only quasinormed, i.e. instead of the triangle inequality of the norm, we only have

$$\|x + y\| \leq c(\|x\| + \|y\|)$$

with some constant $c < \infty$; the most prominent example is $X = L_p(S, U)$ ($0 < p < 1$). In this case, one has to replace inequalities (5) - (8) by the respective estimates

$$\begin{aligned} \chi_i(M) &\leq 2c^4(c\omega(M) + 2\|y\|) \\ \alpha(M) &\leq 2c^5(c\omega(M) + 2\|y\|) \end{aligned} \quad (34)$$

$$\begin{aligned} \chi_i(M) &\leq 2c^4(cp\omega(M) + q\|y\|) \\ \chi_X(M) &\leq c^4(cp\omega(M) + q\|y\|). \end{aligned} \quad (35)$$

Moreover, for countable M one may divide the right-hand sides of (34) and (35) by c .

Of course, one may discuss whether it makes sense at all to consider the Hausdorff and Kuratowski measures of non-compactness in non-metric spaces.

References

- [1] Akhmerov, R. R. et al.: *Measures of Noncompactness and Condensing Operators*. Basel - Boston - Berlin: Birkhäuser Verlag 1992.
- [2] Alt, H. W.: *Lineare Funktionalanalysis*, 2nd ed. Berlin - Heidelberg - New York: Springer-Verlag 1992.
- [3] Amir, D. and J. Lindenstrauss: *The structure of weakly compact sets in Banach spaces*. Ann. Math. (2) 88 (1968), 34 - 46.
- [4] Appell, J., Văth, M., and A. Vignoli: *Compactness and existence results for ordinary differential equations in Banach spaces*. Z. Anal. Anw. 18 (1999)3 (to appear).
- [5] Banach, S.: *Introduction to Linear Operations*. Amsterdam - New York - Oxford: North-Holland 1987; translation of: *Théorie des Opérations Linéaires*. Warsaw: Polish Sci. Publ. 1979.
- [6] Daugavet, I. K.: *On a property of completely continuous operators in the space C* (in Russian). Uspekhi Mat. Nauk 18 (1963)5, 157 - 158.
- [7] Day, M. M.: *Normed linear spaces*, 3rd ed. Berlin - Heidelberg - New York: Springer-Verlag 1973.
- [8] Diestel, J.: *Geometry of Banach Spaces - Selected Topics*. Lect. Notes Math. 485 (1975), 1 - 282.
- [9] Dunford, N. and J. T. Schwartz: *Linear Operators*. Vol. I: *General Theory*, 3rd ed. New York: Int. Publ. 1966.
- [10] Garnir, H. G., de Wilde, M., and J. Schmets: *Analyse fonctionnelle. Théorie constructive des espaces linéaires à semi-normes*. Tome I: *Théorie générale*. Basel - Stuttgart: Birkhäuser Verlag 1968.
- [11] Gribanov, Yu. I.: *Two compactness criteria for sets in Banach function spaces* (in Russian). Kazan. Gos. Univ. Uchen. Zap. 125 (1966)2, 32 - 39.
- [12] Heinz, H.-P.: *On the behaviour of measures of noncompactness with respect to differentiation and integration of vector-valued functions*. Nonlin. Anal. 7 (1983), 1351 - 1371.
- [13] Krasnoselskiĭ, M. A. et al.: *Integral Operators in Spaces of Summable Functions* (in Russian). Moscow: Nauka 1958; Engl. transl.: Leyden: Noordhoff 1976.
- [14] Lindenstrauss, J. and L. Tzafriri: *Classical Banach Spaces*. Vol. I: *Sequence spaces*. Berlin - Heidelberg - New York: Springer-Verlag 1977.
- [15] Luxemburg, W. A. J.: *Notes on Banach function spaces, Part XVI_A*. Proc. Netherl. Acad. Sci. (A) 68 (1965), 646 - 657; Indag. Math. 27 (1965), 646 - 657.
- [16] Luxemburg, W. A. J.: *Notes on Banach function spaces, Part XVI_B*. Proc. Netherl. Acad. Sci. (A) 68 (1965), 658 - 667; Indag. Math. 27 (1965), 658 - 667.
- [17] Luxemburg, W. A. J. and A. C. Zaanen: *Notes on Banach function spaces, Part XII*. Proc. Netherl. Acad. Sci. (A) 67 (1964), 519 - 529; Indag. Math. 26 (1964), 519 - 529.
- [18] Mönch, H.: *Boundary value problems for nonlinear ordinary differential equations in Banach spaces*. Nonlinear Anal. 4 (1980), 985 - 999.
- [19] Mönch, H. and G.-F. von Harten: *On the Cauchy problem for ordinary differential equations in Banach spaces*. Arch. Math. (Basel) 39 (1982), 153 - 160.
- [20] Orlicz, W. and S. Szuffla: *On some classes of nonlinear Volterra integral equations in Banach spaces*. Bull. Acad. Polon. Sci. Sér. Sci. Math. 30 (1982), 239 - 250.
- [21] Orlicz, W.: *On the structure of L_ϕ solution sets of integral equations in Banach spaces*. Studia Math. 77 (1984), 465 - 477.

- [22] Otáhal, A.: *Measure of noncompactness of subsets of Lebesgue spaces*. Cas. Pestovani. Mat. 103 (1978), 67 – 72.
- [23] Pluciennik, R. and S. Szufła: *Nonlinear Volterra integral equations in Orlicz spaces*. Demonstratio Math. 17 (1984), 515 – 532.
- [24] Singer, I.: *Bases in Banach spaces*, Vol. I. Berlin - Heidelberg - New York: Springer-Verlag 1970.
- [25] Szufła, S.: *Appendix to the paper "An existence theorem for the Urysohn integral equation in Banach spaces"*. Comment. Math. Univ. Carolinae 25 (1984), 763 – 764.
- [26] Szufła, S.: *On the Aronszajn property for integral equations in Banach space*. Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8) 83 (1989), 93 – 99.
- [27] Váth, M.: *Ideal Spaces*. Lect. Notes Math. 1664 (1997), 1 – 146.
- [28] Yezakova, N. A.: *On the Hausdorff internal measure of noncompactness* (in Russian). In: *Operator Theory in Functional Spaces* (ed.: E. M. Semenov). Interuniv. Collect. of Sci. Works. Voronezh: Izd. Univ. 1983, pp. 37 – 45.
- [29] Yezakova, N. A.: *On measures of non-compactness in regular spaces*. Z. Anal. Anw. 15 (1996), 299 – 307.
- [30] Zaanen, A. C.: *Integration*. Amsterdam: North-Holland Publ. Comp. 1967.
- [31] Zabreiko, P. P.: *Ideal spaces of functions*, Part I (in Russian). Vestnik Jaroslav. Univ. 8 (1974), 12 – 52.

Received 28.09.1998