

Regularizability of Ill-Posed Problems and the Modulus of Continuity

Radu Ioan Boț, Bernd Hofmann and Peter Mathé

*Dedicated to Ulrich Tautenhahn, a friend and co-author,
who passed away too early at the age of 60.*

Abstract. The regularization of linear ill-posed problems is based on their conditional well-posedness when restricting the problem to certain classes of solutions. Given such class one may consider several related real-valued functions, which measure the well-posedness of the problem on such class. Among those functions the modulus of continuity is best studied. For solution classes which enjoy the additional feature of being star-shaped at zero, the authors develop a series of results with focus on continuity properties of the modulus of continuity. In particular it is highlighted that the problem is conditionally well-posed if and only if the modulus of continuity is right-continuous at zero. Those results are then applied to smoothness classes in Hilbert space. This study concludes with a new perspective on a concavity problem for the modulus of continuity, recently addressed by two of the authors in [Trudy Inst. Mat. i Mekh. UrO RAN 18 (2012)(1), 34–41].

Keywords. Linear ill-posed problems, modulus of continuity, conditional stability

Mathematics Subject Classification (2010). Primary 47A52, secondary 65J20

1. Introduction

We shall consider linear ill-posed problems, given in the form

$$y^\delta = Ax + \delta\xi, \quad (1)$$

where $A: X \rightarrow Y$ denotes an injective and bounded linear operator acting between Banach spaces X and Y . If the range $\mathcal{R}(A) \subset Y$ is non-closed, then the Problems (1) are ill-posed, meaning that the inverse mapping $A^{-1} : \mathcal{R}(A) \subset Y \rightarrow X$ is not bounded and hence not continuous. However,

R. I. Boț, B. Hofmann: Department of Mathematics, Chemnitz University of Technology, 09107 Chemnitz, Germany;

radu.bot@mathematik.tu-chemnitz.de; hofmannb@mathematik.tu-chemnitz.de

P. Mathé: Weierstraß Institute for Applied Analysis and Stochastics, Mohrenstraße 39, 10117 Berlin, Germany; peter.mathe@wias-berlin.de

one may restrict a problem as posed in (1) by introducing the *a priori information* that $x \in M \subseteq X$, for a non-empty subset M . By doing so we may study the following questions.

Question 1.1. *Does the mapping $A: M \rightarrow Y$ have a bounded inverse?*

If this is the case, then the mapping $\Omega_M: [0, +\infty) \rightarrow [0, +\infty]$, given as

$$\Omega_M(\delta) := \sup \{ \|x - x'\| : x, x' \in M, \|Ax - Ax'\| \leq \delta \}, \quad (2)$$

is of interest. The above function Ω_M is analyzed in various studies, and we mention [7, § 2.3] for a monograph.

Question 1.2. *Is the Problem (1) regularizable on M in the sense of Tikhonov?*

This question is related to the reconstruction of x from noisy data, and we introduce the error criterion. If S is any mapping of the form

$$y^\delta \in Y \rightarrow S(y^\delta) \in X,$$

then we let

$$e(S, x, \delta) := \sup_{y^\delta: \|Ax - y^\delta\| \leq \delta} \|S(y^\delta) - x\|, \quad \delta > 0,$$

and the corresponding *uniform error* of any reconstruction S on the set M will be given by

$$e(S, M, \delta) := \sup_{x \in M} e(S, x, \delta), \quad \delta > 0.$$

By *regularizability* on M in the sense of Tikhonov we mean that there is a family R_δ of reconstructions such that for each $x \in M$ we have that

$$e(R_\delta, x, \delta) \rightarrow 0 \quad \text{as } \delta \rightarrow 0.$$

The question whether a problem is regularizable received attention very early in the analysis of ill-posed problems (for an early study we refer to [12], and we also mention the monograph [1, Chapter 1]). If the above is the case then we call the restriction of A^{-1} on

$$A(M) := \{z \in Y : z = Ax, x \in M\}$$

regularizable on $A(M)$. In addition we may ask for *uniform regularizability* of the Problem (1) on the set M , thus asking whether a family R_δ of reconstructions exists for which $e(R_\delta, M, \delta) \rightarrow 0$ as $\delta \rightarrow 0$.

While the first question studies intrinsic continuity features of the problem, the latter question asks for the ability to find reconstruction methods, which are capable to recover x based on noisy data y^δ as δ is getting smaller. In this study we will not address the regularizability problem in the sense of Tikhonov. Instead we analyze uniform regularizability for classes M .

2. Modulus of continuity

For a variety of classes of subsets M of X that contain the zero element, instead of the function Ω_M from (2) one can consider the function $\omega_M : [0, +\infty) \rightarrow [0, +\infty]$, defined as

$$\omega_M(\delta) := \sup \{ \|x\| : x \in M, \|Ax\| \leq \delta \},$$

usually called *modulus of continuity*. Clearly, for $0 \in M$ we have that $\omega_M(\delta) \leq \Omega_M(\delta)$ for all $\delta \geq 0$. The modulus of continuity is obviously a non-decreasing function of δ .

2.1. Elementary properties. The following elementary properties are easily verified.

Lemma 2.1. *Let $M, N \subseteq X$ be two subsets. Then for all $\delta \geq 0$ it holds*

- (i) $\omega_{M \cup N}(\delta) = \max \{ \omega_M(\delta), \omega_N(\delta) \}$.
- (ii) $\omega_M(\delta) = \omega_{-M}(\delta)$.

In particular, we have that $\omega_{M \cup -M}(\delta) = \omega_M(\delta)$.

Below we shall confine ourselves to sets M from the following class of sets.

Definition 2.2. A subset $M \subseteq X$ is said to be *star-shaped at zero* if $0 \in M$ and if $x \in M$ implies $\alpha x \in M$ for all $0 \leq \alpha \leq 1$. Equivalently, for each $C \geq 1$ we have that $\frac{1}{C}M \subseteq M$.

Lemma 2.3. *Suppose that M is star-shaped at zero. Then for all $\delta \geq 0$ and all $C \geq 1$ it holds*

$$\omega_M(C\delta) \leq C\omega_M(\delta) \quad \text{and} \quad \Omega_M(C\delta) \leq C\Omega_M(\delta).$$

In particular, we have that $\frac{\omega_M(t)}{t} \leq \frac{\omega_M(s)}{s}$ as well as $\frac{\Omega_M(t)}{t} \leq \frac{\Omega_M(s)}{s}$ whenever $0 < s \leq t$.

Proof. Let $C \geq 1$ be arbitrary. Then for all $\delta \geq 0$ it holds

$$\begin{aligned} \omega_M(C\delta) &= \sup \{ \|x\| : x \in M, \|Ax\| \leq C\delta \} \\ &= C \sup \left\{ \left\| \frac{1}{C}x \right\| : \frac{1}{C}x \in \frac{1}{C}M, \left\| A \frac{1}{C}x \right\| \leq \delta \right\} \\ &= C \sup \left\{ \|z\| : z \in \frac{1}{C}M, \|Az\| \leq \delta \right\} \\ &= C\omega_{\frac{1}{C}M}(\delta) \leq C\omega_M(\delta). \end{aligned}$$

Taking into account that $\Omega_M = \omega_{M-M}$, where we let

$$M - M := \{ x - y; x, y \in M \},$$

and the fact that the difference of two star-shaped at zero sets is again star-shaped at zero, the assertion for Ω_M follows. □

We summarize the above elementary findings.

Proposition 2.4. *Suppose that the set $M \subseteq X$ is star-shaped at zero.*

- (i) *Then the mapping ω_M is non-decreasing, $\omega_M(0) = 0$, and the mapping $\delta \rightarrow \frac{\omega_M(\delta)}{\delta}$ is non-increasing on $(0, +\infty)$.*
- (ii) *Likewise the mapping Ω_M is non-decreasing, $\Omega_M(0) = 0$, and the mapping $\delta \rightarrow \frac{\Omega_M(\delta)}{\delta}$ is non-increasing on $(0, +\infty)$.*

These elementary properties have immediate consequences for the continuity properties of both ω_M and Ω_M .

2.2. Modulus functions.

Definition 2.5. We agree to call a mapping $f: [0, +\infty) \rightarrow [0, +\infty]$ a *modulus function* if it is non-decreasing, $f(0) = 0$, and $t \rightarrow \frac{f(t)}{t}$ is non-increasing on $(0, +\infty)$. It is called *proper* if for some $t > 0$ it has a finite value $f(t) < +\infty$.

With this notion both the functions ω_M and Ω_M are modulus functions provided that the set M is star-shaped at zero.

Proposition 2.6. *Let f be any proper modulus function. Then*

- (i) *the values $f(t)$ are finite for every $t > 0$.*
- (ii) *If $f(t_0) = 0$ for some $t_0 > 0$ then $f(t) = 0$ for all $t \geq 0$.*
- (iii) *The mapping f is continuous on $(0, +\infty)$, and*
- (iv) *for every pair $t_1, t_2 > 0$ we have that $f(t_1 + t_2) \leq f(t_1) + f(t_2)$.*

Proof. Suppose that f is finite at $t_0 > 0$. Then it is finite for every $0 < t \leq t_0$ by monotonicity. Also, if $t > t_0$ then $\frac{f(t)}{t} \leq \frac{f(t_0)}{t_0} < +\infty$, which proves the assertion (i).

For the second assertion (ii), suppose that $f(t_0) = 0$, and $t_0 > 0$. Then, due to monotonicity we have that $f(t) = 0$, $0 \leq t \leq t_0$. For any $t_1 > t_0$ we see that $0 \leq f(t_1) = f(\frac{t_1}{t_0}t_0) \leq \frac{t_1}{t_0}f(t_0) = 0$.

For proving (iii) let $t > 0$ be any real number. If $f(t) = 0$ then by item (ii), f is identically zero, and the assertion is obvious. Otherwise, we first prove right continuity of f at t . Let $t_n \searrow t$. Then $1 \leq \frac{f(t_n)}{f(t)} \leq \frac{t_n}{t} \rightarrow 1$. Similarly, if $t_n \nearrow t$ then $1 \leq \frac{f(t)}{f(t_n)} \leq \frac{t}{t_n} \rightarrow 1$, which proves the continuity of f at $t > 0$.

The proof of the sub-additivity in item (iv) is well-known, and we recall this here for convenience. Plainly, $t_1, t_2 \leq t_1 + t_2$, and therefore

$$f(t_1 + t_2) = t_1 \frac{f(t_1 + t_2)}{t_1 + t_2} + t_2 \frac{f(t_1 + t_2)}{t_1 + t_2} \leq t_1 \frac{f(t_1)}{t_1} + t_2 \frac{f(t_2)}{t_2} = f(t_1) + f(t_2),$$

and this completes the proof of the proposition. \square

Remark 2.7. Proposition 2.6(iv) yields that $|f(t_1) - f(t_2)| \leq f(|t_1 - t_2|)$, $t_1, t_2 \geq 0$. Therefore, proper modulus functions are sub-additive, and continuous at every $t > 0$.

Within the classical context, the modulus of continuity, say ω , of a real function on a bounded interval is non-decreasing, $\omega(0) = 0$, and sub-additive, see e.g. [8, § 6.1]. Therefore any such function is called *modulus of continuity* if, in addition, it is continuous at zero.

Corollary 2.8. *A proper modulus function is continuous on $[0, +\infty)$ if and only if it is right-continuous at zero.*

We conclude this subsection with gathering more, and important, properties of modulus functions, we refer to [8, Lemma 6.1.4].

Proposition 2.9. *For every proper modulus function f which is right-continuous at zero there is a concave right-continuous proper modulus function f_* with*

$$f(t) \leq f_*(t) \leq 2f(t), \quad t > 0.$$

The constant 2 cannot be improved, in general.

We finally mention the following result.

Corollary 2.10. *If a proper modulus function f does not vanish identically then it tends to zero at most linearly, i.e., $t = \mathcal{O}(f(t))$ as $t \searrow 0$.*

Proof. Suppose that f does not vanish identically and that it is finite for $t_0 > 0$. According to Proposition 2.6(i), f is finite on $[0, +\infty)$. Then the assertion is immediate from the fact that $\frac{f(t)}{t}$ is non-increasing, which implies that for all $0 < t \leq t_0$ we have that $f(t) \geq \frac{f(t_0)}{t_0}t$. □

2.3. Conditional well-posedness. The right-continuity at zero of the functions ω_M and Ω_M is intimately related to the continuity of the inverse A^{-1} , when considered as acting from $A(M) \subset Y$ to X . Indeed, we have the following proposition.

Proposition 2.11. *Let $M \subseteq X$ be star-shaped at zero. Then the modulus ω_M is right-continuous at zero if and only if the mapping A^{-1} restricted to $A(M)$ is continuous at zero.*

Proof. One has that ω_M is right-continuous at zero if and only if

$$\begin{aligned} & \forall \varepsilon > 0 \exists \bar{\delta} > 0 \text{ such that } \forall \delta < \bar{\delta} \text{ it holds } \omega_M(\delta) < \varepsilon \\ \Leftrightarrow & \forall \varepsilon > 0 \exists \bar{\delta} > 0 \text{ such that } \forall \delta < \bar{\delta} \forall y \in A(M) \text{ with } \|y\| \leq \delta \text{ it holds } \|A^{-1}y\| < \varepsilon \\ \Leftrightarrow & \forall \varepsilon > 0 \exists \bar{\delta} > 0 \text{ such that } \forall y \in A(M) \text{ with } \|y\| < \bar{\delta} \text{ it holds } \|A^{-1}y\| < \varepsilon, \end{aligned}$$

which is nothing else than $A^{-1}: A(M) \rightarrow X$ is continuous at zero. □

We summarize the preceding discussion.

Definition 2.12. We call the Problem (1) *conditionally well-posed* on M if the mapping A^{-1} restricted to $A(M)$ is continuous.

A famous theorem by A. N. Tikhonov, see [11], asserts that the problem is conditionally well-posed whenever the set $M \subset X$ is compact. For some further discussion and examples concerning conditional well-posedness we also refer to [4, 5, 14].

Theorem 2.13. *Suppose that the set $M \subseteq X$ is star-shaped at zero. Then the following assertions are equivalent:*

- (i) *The Problem (1) is conditionally well-posed on M .*
- (ii) *The modulus of continuity ω_M is right-continuous at zero.*
- (iii) *The modulus of continuity ω_M is continuous on $[0, \infty)$.*

Proof. The equivalence of the first two assertions is a consequence of Proposition 2.11. The equivalence of the last two assertions follows from Corollary 2.8. \square

Tikhonov's result translates to the following statement.

Proposition 2.14. *If the set $M \subset X$ is compact and star-shaped at zero, then we have $\omega_M(\delta) \rightarrow 0$ as $\delta \rightarrow 0$, and hence the modulus of continuity ω_M is continuous on $[0, \infty)$.*

We present the following examples.

Example 2.15. Suppose that the operator A is injective and has a non-closed range. By denoting with $B_X := \{x \in X : \|x\| \leq 1\}$ the closed unit ball of X , we show that for all $\delta > 0$ it holds

$$\omega_{B_X}(\delta) = \sup\{\|x\| : \|x\| \leq 1, \|Ax\| \leq \delta\} = 1.$$

Indeed, consider a fixed $\delta > 0$. Obviously, $\omega_{B_X}(\delta) \leq 1$. Since the range of A is non-closed, there is no $K > 0$ such that $\|x\| \leq K\|Ax\|$ for all $x \in X$. Thus there exists a sequence $\{\tilde{x}_n\}_{n \geq 1} \subset X \setminus \{0\}$ such that $\frac{\|\tilde{x}_n\|}{\|A\tilde{x}_n\|} \rightarrow +\infty$ as $n \rightarrow +\infty$. By defining for all $n \geq 1$ $x_n := \frac{1}{\|\tilde{x}_n\|}\tilde{x}_n$, one has that $\|x_n\| = 1$ and $\|Ax_n\| \rightarrow 0$ as $n \rightarrow +\infty$. Thus there exists $n(\delta) \geq 1$ such that $\|Ax_n\| \leq \delta$ for all $n \geq n(\delta)$ and this provides the desired assertion. Consequently, the modulus of continuity ω_M is not right-continuous at zero for $M = B_X$.

Example 2.16. If $\{0\} \neq L \subset X$ is a finite-dimensional linear subspace then we have $\omega_M(\delta) \asymp \delta$ as $\delta \rightarrow 0$ for $M = L$. Indeed, the image space $A(L)$ is a closed linear subspace in Y , and therefore $A^{-1}: A(L) \rightarrow L$ is a bounded linear

operator. Thus there is a constant $C < +\infty$ for which $\|x\| \leq C \|Ax\|$, $x \in L$. But this yields that $\omega_L(\delta) \leq C\delta$ for all $\delta > 0$. Next, if ω_L would vanish, say at $\delta_0 > 0$, then $\{x \in L : \|Ax\| \leq \delta_0\} = \{0\}$, which is not true. In the light of Corollary 2.10 the above assertion is proved.

The above extremal decay rate, together with Lemma 2.1 gives rise to the following

Conjecture 2.17. Let $M \subset X$ be any set which is star-shaped at zero, and let L be a finite-dimensional subspace of X . If ω_M does not vanish identically then $\omega(M + L, \delta) \asymp \omega(M, \delta)$ as $\delta \rightarrow 0$.

2.4. Regularizability. We now highlight the problem of the continuity of the modulus of continuity to the regularizability problem. To this end we introduce the local companion to the function Ω_M from (2), and we introduce, given any $x \in M$, the function

$$\Omega_M^{loc}(x, \delta) := \sup \{ \|x - x'\| : x' \in M, \|Ax - Ax'\| \leq \delta \}.$$

Plainly, $\sup_{x \in M} \Omega_M^{loc}(x, \delta) = \Omega_M(\delta)$.

In agreement with the usual nomenclature we call a reconstruction S *interpolatory at level δ* if for each $x \in M$, and data y^δ with $\|Ax - y^\delta\| \leq \delta$, we have that $S(y^\delta) \in M$ and $\|AS(y^\delta) - y^\delta\| \leq \delta$.

Lemma 2.18. *Let S be any reconstruction.*

(i) *If S is interpolatory, then we have for each $x \in M$ that*

$$e(S, x, \delta) \leq \Omega_M^{loc}(x, \frac{2}{\delta}), \quad \delta > 0.$$

(ii) *If the set M is centrally-symmetric, then*

$$\omega_M(\delta) \leq e(S, M, \delta), \quad \delta > 0.$$

(iii) *If the set M is convex and centrally-symmetric, then for each $x \in M$ it holds*

$$\Omega_M^{loc}(x, \delta) \leq \Omega_M(\delta) = 2\omega_M\left(\frac{\delta}{2}\right), \quad \delta > 0$$

Thus, for convex centrally-symmetric sets M we have

$$\omega_M(\delta) \leq e(S, M, \delta) \leq 2\omega_M(\delta), \quad \delta > 0, \tag{3}$$

for any reconstruction S which is interpolatory at the level δ .

Proof. Given data y^δ and interpolatory reconstruction S we let $x' := S(y^\delta) \in M$. Then $\|x - S(y^\delta)\| \leq \Omega_M^{loc}(x, \frac{\delta}{2})$. Since this is true for any data y^δ with $\|Ax - y^\delta\| \leq \delta$ the proof of the first assertion (i) can be completed.

For the second assertion (ii) we argue as follows. For every $x \in M$, and due to symmetry also $-x \in M$ with $\|Ax\| = \|A(-x)\| \leq \delta$ the data $y^\delta := 0$ are possible data, and we bound, using the triangle inequality

$$e(S, M, \delta) \geq \max \{ \|S(0) - x\|, \|S(0) + x\| \} \geq \|x\|.$$

Therefore we conclude that $e(S, M, \delta) \geq \sup \{ \|x\| : x \in M, \|Ax\| \leq \delta \} = \omega_M(\delta)$.

To prove item (iii) we fix any $x \in M$, and let $x' \in M$ be arbitrary. Then $\hat{x} := \frac{x-x'}{2} \in M$, and we have that $\|A\hat{x}\| \leq \frac{\delta}{2}$, provided that $\|Ax - Ax'\| \leq \delta$. Thus, $\|x - x'\| = 2\|\hat{x}\| \leq 2\omega_M(\frac{\delta}{2})$. Next we prove that $2\omega_M(\frac{\delta}{2}) \leq \Omega_M(\delta)$. To this end, given $0 < \varepsilon \leq \omega_M(\frac{\delta}{2})$, let $z \in M$, $\|Az\| \leq \frac{\delta}{2}$, be such that $\|z\| \geq \omega_M(\frac{\delta}{2}) - \varepsilon$. This yields that $z, -z \in M$, $\|A(z - (-z))\| \leq \delta$, and

$$\|z - (-z)\| = 2\|z\| \geq 2\omega_M\left(\frac{\delta}{2}\right) - 2\varepsilon,$$

hence that $\Omega_M(\delta) \geq 2\omega_M(\frac{\delta}{2})$. The bounds (3) are now easy consequences. \square

Remark 2.19. Indeed, interpolatory reconstructions always exist. To check this, fix $x \in M$ and data y^δ with $\|Ax - y^\delta\| \leq \delta$ and consider the set

$$D(y^\delta) := \{ z \in M : \|Az - y^\delta\| \leq \delta \}.$$

Since $x \in D(y^\delta)$ this set is non-empty, and any selection $S(y^\delta) \in D(y^\delta)$ will yield an interpolatory reconstruction. If the set $M \subset X$ is convex and compact, then there is even a continuous selection by *Michael's continuous selection theorem*, we refer to [2, § 7] for details and extensions. For compact sets M the construction of interpolatory reconstructions may be achieved by solving the optimization problem

$$x_{qu} := \arg \min_{z \in M} \|Az - y^\delta\|,$$

which exists due to the compactness of M . It is readily checked that $x_{qu} \in D(y^\delta)$. This construction goes back to Ivanov [6] and it is called *method of quasi-solutions*, there.

Corollary 2.20. *For convex centrally-symmetric sets M the Problem (1) is uniformly regularizable on M if and only if the modulus of continuity ω_M is right-continuous at zero.*

3. Smoothness classes in Hilbert space

Here and in the subsequent section let X and Y be separable Hilbert spaces. We recall that the linear operator $A : X \rightarrow Y$ is assumed to be bounded and injective. For that case one can consider its self-adjoint companion $H := A^*A$, where we set $a := \|H\| = \|A\|^2$.

The typical smoothness classes as considered in inverse problems, and we mention *source sets* expressing general smoothness assumptions, and more recently, *level sets*, are based on the distribution function

$$F_x^2(t) := \|\chi_{(0,t]}(H)x\|^2 = \int_0^t d\|E_\tau x\|^2, \quad t \geq 0,$$

which is well-defined and finite for each $x \in X$. Above, we let $\chi_{(0,t]}$ be the characteristic function of the interval $(0, t]$, and $E_t = E_t(H)$, $0 \leq t \leq a$, be the spectral resolution of the operator H . The following elementary properties are easily seen, for a further discussion and consequences cf. [3].

Lemma 3.1. *Let $x \in X$ be arbitrary.*

- (i) *The function $t \mapsto F_x(t)$ is right-continuous and non-decreasing.*
- (ii) *If the operator H is injective then $F_x(0) = 0$.*
- (iii) *For all $0 < t \leq a$ we have that $F_{\chi_{(0,t]}(H)x}(\tau) = F_{\chi_{(0,t]}(H)x}(t)$, $\tau \geq t$.*

For a class $M \subset X$ we consider the associated function \bar{F}_M , given as

$$\bar{F}_M(t) := \sup_{x \in M} F_x(t), \quad t \geq 0.$$

This function is finite whenever M is bounded, and we thus will assume boundedness of M , throughout. It is also non-decreasing, and we have $\bar{F}_M(0) = 0$. As it will turn out, the right-continuity of the function $\bar{F}_M(t)$ at zero and the right-continuity of the function ω_M at zero are closely related, and we will dwell into this, now. Best results are obtained for smoothness which is expressed through sets M , which are determined in the vicinity of zero of the distribution function $F_x(t)$, $t > 0$, only.

Definition 3.2 (spectral smoothness). We call a smoothness class M *spectral*, if for each $t > 0$ we have that $x \in M$ yields that $\chi_{(0,t]}(H)x \in M$.

Definition 3.3 (index function). We call a function $\varphi : (0, a] \rightarrow (0, \infty)$ *index function* if it is continuous and increasing with $\lim_{t \searrow 0} \varphi(t) = 0$.

Example 3.4. For an index function φ we assign the smoothness class \mathcal{M}_φ as

$$\mathcal{M}_\varphi := \{x = \varphi(H)v : \|v\| \leq 1\},$$

i.e., the image of the unit ball under the mapping $\varphi(H)$. Such classes are spectral since with $x = \varphi(H)v \in \mathcal{M}_\varphi$ we also have that

$$\chi_{(0,t]}(H)x = \chi_{(0,t]}(H)\varphi(H)v = \varphi(H)\chi_{(0,t]}(H)v,$$

and $\|\chi_{(0,t]}(H)v\| \leq \|v\| \leq 1$. We mention that

$$\bar{F}_{\mathcal{M}_\varphi}(t) = \|\chi_{(0,t]}(H)\varphi(H)\| \leq \sup_{0 < s \leq t} \varphi(s) = \varphi(t), \quad 0 < t \leq a,$$

such that $\bar{F}_{\mathcal{M}_\varphi}$ is right-continuous at zero exactly if φ was an index function.

Example 3.5. For an index function φ we assign the level set \mathcal{E}_φ as

$$\mathcal{E}_\varphi := \{x \in X : F_x(t) \leq \varphi(t), \quad 0 < t \leq a\}.$$

In view of Lemma 3.1(iii) such classes also constitute spectral smoothness classes, and $\bar{F}_{\mathcal{E}_\varphi}$ is right-continuous at zero for index functions φ .

Proposition 3.6. *Suppose that the set $M \subset X$ is bounded. If the function \bar{F}_M is right-continuous at zero then ω_M is also right-continuous at zero. Moreover, for spectral smoothness classes M the converse also holds true.*

Proof. To prove the first assertion we observe that for $0 < t < a$ we can estimate the norm square as

$$\|x\|^2 = \int_0^t dF_x^2(\tau) + \int_t^a dF_x^2(\tau) \leq F_x^2(t) + \frac{1}{t} \int_t^a \tau dF_x^2(\tau).$$

From this it follows $\|x\|^2 \leq F_x^2(t) + \frac{1}{t}\|Ax\|^2$ and taking the supremum over the set $\{x \in M : \|Ax\| \leq \delta\}$ we obtain for $\delta > 0$ and all sufficiently small $t > 0$

$$\omega_M^2(\delta) \leq \bar{F}_M^2(t) + \frac{\delta^2}{t}. \tag{4}$$

By setting $t := \delta$ and under the condition $\bar{F}_M(t) \rightarrow 0$ as $t \searrow 0$ the upper bound of $\omega_M^2(\delta)$ in (4) tends to zero as $\delta \searrow 0$. Hence $\lim_{\delta \searrow 0} \omega_M(\delta) = 0$. The second assertion is proved by contraposition. Without loss of generality we assume that M belongs to the unit ball in X . Suppose that there is some $\varepsilon > 0$ such that for all $t > 0$ we have that $\bar{F}_M(t) = \sup_{x \in M} \|\chi_{(0,t]}(H)x\| > \varepsilon$. Thus we can find $\tilde{x}_t \in M$ with $\|\chi_{(0,t]}(H)\tilde{x}_t\| \geq \varepsilon$. We assign $x_t := \chi_{(0,t]}(H)\tilde{x}_t$, $t > 0$, and $x_t \in M$ since the set M was assumed to be spectral. We thus have that for this $\varepsilon > 0$ we can find a family $x_t \in M$, $t > 0$ with $\|x_t\| \geq \varepsilon$. We claim that $\|Ax_t\| \leq t$. Indeed, taking into account item (iii) of Lemma 3.1 we bound

$$\|Ax_t\|^2 = \int_0^a \tau dF_{x_t}^2(\tau) = \int_0^t \tau dF_{x_t}^2(\tau) \leq t \int_0^t dF_{x_t}^2(\tau) \leq t \|x_t\|^2 \leq t.$$

Consequently we see that $\omega_M(\sqrt{t}) \geq \varepsilon$, $t > 0$, which is a contradiction. The proof is complete. \square

Remark 3.7. Tight bounds for the modulus of continuity $\omega_{\mathcal{M}_\varphi}$ can be obtained under additional geometric (convexity) assumptions by means of general interpolation results within the framework of *variable Hilbert scales*, and we mention [13] for an early work on this. More recently this is pursued within the framework of *conditionally stability estimates*, see the recent study [14].

Evidently, Proposition 3.6 provides us with a characterization for the convex and centrally-symmetric smoothness classes \mathcal{M}_φ and \mathcal{E}_φ .

In many cases the set M is an ellipsoid in Hilbert space, i.e., there is an operator $G: Z \rightarrow X$, for a Hilbert space Z such that

$$M(G) := \{Gv : \|v\| \leq 1\}, \tag{5}$$

the image of the unit ball in Z under the mapping G . Such ellipsoids $M(G)$ need not be spectral smoothness classes, in general. An obvious exception occurs for commuting operators G and H .

Example 3.8. In case that $G = \varphi(H): X \rightarrow X$ is a function of H , where φ is any index function we have $M(G) = \mathcal{M}_\varphi$. Hence the concept of ellipsoidal sets generalizes general smoothness classes from Example 3.4.

For ellipsoids $M(G)$ we can rewrite

$$\bar{F}_{M(G)}(t) = \sup_{\|v\| \leq 1} \|\chi_{(0,t]}(H)Gv\| = \|\chi_{(0,t]}(H)G\|, \quad 0 < t \leq a.$$

where the latter is the operator norm of $\chi_{(0,t]}(H)G: Z \rightarrow X$.

The following was proved in [5, Thm. 4.4], and this shows that the bound in Lemma 2.18(iii) can be attained for ellipsoidal sets.

Theorem 3.9. *For each ellipsoid $M(G)$ and each $\delta > 0$ there is a linear reconstruction method S_δ such that $e(S_\delta, M(G), \delta) = \omega_{M(G)}(\delta)$.*

For compact operators A we have the following characterization.

Proposition 3.10. *Suppose that A is a compact operator and $H := A^*A$. The function $\bar{F}_{M(G)}$ is right-continuous at zero exactly if the operator G is compact.*

Proof. Since the operator H is self-adjoint and compact it has a monotone Schmidt representation (svd) in the form $Hx = \sum_{j=1}^\infty s_j \langle x, u_j \rangle u_j$, $x \in X$ with $\|H\| = s_1 \geq s_2 \geq \dots \geq 0$, and orthonormal system u_j , $j = 1, 2, \dots$. Then the operator $\chi_{(0,t]}(H)$ is a finite co-dimensional orthogonal projection, and hence it can be written as $\chi_{(0,t]}(H) = I - P_N$, where P_N is the projection onto the finite dimensional space $X_N := \text{span}(\{u_j, s_j > t\})$. Thus, we see that $\sup_{x \in M(G)} F_x(t) = \|(I - P_N)G\|$. As $t \searrow 0$ the dimension of the spaces X_N will increase to $+\infty$. From this we conclude that $\sup_{x \in M(G)} F_x(t) \rightarrow 0$ if and only if the operator G is approximable by finite rank operators, and thus compact. In Hilbert space, as a consequence of its (metric) approximation property, the notions of compactness and approximability coincide, see e.g. [9, Chapter 10]. \square

Conjecture 3.11. Suppose that A is compact and that the set $M(G)$ is an ellipsoid generated by the operator G , see (5). The modulus of continuity $\omega_{M(G)}$ is right-continuous at zero if and only if G is compact.

Remark 3.12. For commuting operators G and H this holds true, in view of Proposition 3.6.

4. On the concavity of the modulus of continuity

Now, under the setting and notation of the previous section we (re)prove (see [4, Theorem]) the concavity of the function $\omega_M^2(\sqrt{\delta})$, $0 \leq \delta < \infty$, for $M = \mathcal{M}_\varphi$, and $M = \mathcal{E}_\varphi$ (for an index function φ , see Examples 3.4 and 3.5), by using some tools of convex analysis.

For the beginning we notice that, according to the spectral theorem for bounded self-adjoint linear operators in Hilbert spaces (see [10, Chapter VII]), there exist a measurable space $(\Omega, \mathcal{A}, \mu)$, a unitary transformation $U : X \rightarrow L^2(\Omega, \mathcal{A}, \mu)$ and a measurable function

$$f : \Omega \rightarrow \sigma(H) \setminus \{0\} \subseteq (0, \|H\|] \subset \mathbb{R}$$

such that $T_f := UHU^* : L^2(\Omega, \mathcal{A}, \mu) \rightarrow L^2(\Omega, \mathcal{A}, \mu)$ is a multiplication operator defined as

$$[T_f h](\omega) := f(\omega) h(\omega) \quad \text{for all } \omega \in \Omega.$$

By [4, Proposition 3] we get for an arbitrary set $M \subseteq X$ that

$$\omega_M(\delta) := \omega_{M,A}(\delta) = \omega_{M,H^{\frac{1}{2}}}(\delta) = \omega_{UM,T_{\sqrt{f}}}(\delta) \quad \text{for all } \delta \geq 0. \tag{6}$$

In the formula above the second lower index in the modulus of the continuity denotes the bounded linear operator to which this is associated, while $UM := \{Ux : x \in M\}$ stands for the image of the set M through the operator U .

For the images of the sets \mathcal{M}_φ and \mathcal{E}_φ through the unitary transformation U we have, according to [4, Lemma 1], the representations

$$UM_\varphi = \{g \in L^2(\Omega, \mathcal{A}, \mu) : g = \varphi(f)h, \|h\|_{L^2(\Omega, \mathcal{A}, \mu)} \leq 1\},$$

and, respectively,

$$UE_\varphi = \left\{ g \in L^2(\Omega, \mathcal{A}, \mu) : \int_{0 < f(\omega) \leq t} g^2(\omega) d\mu(\omega) \leq \varphi^2(t) \quad \forall t \in (0, a] \right\}.$$

The main result of this section follows.

Theorem 4.1. *Let $A : X \rightarrow Y$ be an injective and bounded linear operator with non-closed range $\mathcal{R}(A)$ mapping between separable Hilbert spaces X and Y and let $\varphi : (0, \|A\|^2] \rightarrow (0, +\infty)$ be an arbitrary index function. Then the functions*

$$\delta \mapsto \omega_{\mathcal{M}_\varphi}^2(\sqrt{\delta}) \quad \text{and} \quad \delta \mapsto \omega_{\mathcal{E}_\varphi}^2(\sqrt{\delta})$$

are concave on the interval $[0, +\infty)$.

Proof. To prove that $\delta \mapsto \omega_{\mathcal{M}_\varphi}^2(\sqrt{\delta})$ is concave on $[0, +\infty)$, we let be $\Theta(t) := \sqrt{t} \varphi(t)$ for all $0 < t \leq a$. According to (6) and by making use of the representation given for $U\mathcal{M}_\varphi$ above, it holds, for all $\delta \geq 0$,

$$\begin{aligned} & -\omega_{\mathcal{M}_\varphi}^2(\sqrt{\delta}) \\ &= \inf \left\{ -\|g\|_{L^2(\Omega, \mathcal{A}, \mu)}^2 : g = \varphi(f)h, \|h\|_{L^2(\Omega, \mathcal{A}, \mu)} \leq 1, \|\sqrt{f}g\|_{L^2(\Omega, \mathcal{A}, \mu)}^2 \leq \delta \right\} \\ &= \inf \left\{ -\|\varphi(f)h\|_{L^2(\Omega, \mathcal{A}, \mu)}^2 : \|h\|_{L^2(\Omega, \mathcal{A}, \mu)} \leq 1, \|\Theta(f)h\|_{L^2(\Omega, \mathcal{A}, \mu)}^2 \leq \delta \right\}. \end{aligned} \quad (7)$$

It will be convenient to consider the duality pairing $\langle \cdot, \cdot \rangle$ between $L^\infty(\Omega, \mathcal{A}, \mu)$ and $L^1(\Omega, \mathcal{A}, \mu)$. Because both the functions $\varphi(f)$ and $\Theta(f)$ are uniformly bounded, and the function $k := h^2 \geq 0$ belongs to $L^1(\Omega, \mathcal{A}, \mu)$, the representation (7) rewrites as

$$-\omega_{\mathcal{M}_\varphi}^2(\sqrt{\delta}) = \inf \left\{ \langle -\varphi^2(f), k \rangle : \langle 1, k \rangle \leq 1, \langle \Theta^2(f), k \rangle \leq \delta, k \geq 0 \right\},$$

which results in $-\omega_{\mathcal{M}_\varphi}^2(\sqrt{\delta}) = \inf_{k \in L^1(\Omega, \mathcal{A}, \mu)} \Psi(k, \delta)$, where the function $\Psi : L^1(\Omega, \mathcal{A}, \mu) \times \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ is given as

$$\Psi(k, \delta) = \begin{cases} \langle -\varphi^2(f), k \rangle, & \langle 1, k \rangle \leq 1, \langle \Theta^2(f), k \rangle \leq \delta, k \geq 0, \delta \geq 0, \\ \infty, & \text{otherwise.} \end{cases}$$

The function Ψ is a convex function in both variables, and this fact implies the convexity of marginal function $\delta \mapsto -\omega_{\mathcal{M}_\varphi}^2(\sqrt{\delta})$ (cf. [15, Theorem 2.1.3(v)]), which further yields the concavity of $\omega_{\mathcal{M}_\varphi}^2(\sqrt{\cdot})$ on $[0, +\infty)$, and which completes the proof. The proof of the the concavity of $\delta \mapsto \omega_{\mathcal{E}_\varphi}^2(\sqrt{\delta})$ is similar and hence omitted. \square

Acknowledgement. The research of Radu Ioan Boț and Bernd Hofmann was partially supported by Deutsche Forschungsgemeinschaft (DFG) under grants BO 2516/4-1 and HO 1454/8-1, respectively.

References

- [1] Bakushinsky, A. and Goncharsky, A., *Ill-Posed Problems: Theory and Applications* (Transl. from Russian by I. V. Kochikov). Math. Appl. 301. Dordrecht: Kluwer 1994.
- [2] Bessaga, C. and Pełczyński, A., *Selected Topics in Infinite-Dimensional Topology*. Monogr. Mat. 58. Warsaw: PWN–Polish Scientific Publishers 1975.
- [3] Flemming, J., Hofmann, B. and Mathé, P., Sharp converse results for the regularization error using distance functions. *Inverse Probl.* 27 (2011)(2), 025006 (18 pp).
- [4] Hofmann, B. and Mathé, P., A note on the modulus of continuity for ill-posed problems in Hilbert space. *Trudy Inst. Mat. i Mekh. UrO RAN* 18 (2012)(1), 34 – 41.
- [5] Hofmann, B., Mathé, P. and Schieck, M., Modulus of continuity for conditionally stable ill-posed problems in Hilbert space. *J. Inverse Ill-Posed Probl.* 16 (2008)(6), 567 – 585.
- [6] Ivanov, V. K., Approximate solution of operator equations of the first kind. *Zh. Vychisl. Mat. Mat. Fiz.* 6 (1966), 1089 – 1094.
- [7] Ivanov, V. K., Vasin, V. V. and Tanana, V. P., *Theory of Linear Ill-Posed Problems and its Applications* (Transl. and rev. from the 1978 Russian original (Moscow)). Inverse Ill-posed Probl. Ser. Utrecht: VSP 2002.
- [8] Korneĭchuk, N., *Exact Constants in Approximation Theory*. (Transl. from Russian by K. Ivanov). Encyclopedia Math. Appl. 38. Cambridge: Cambridge Univ. Press 1991.
- [9] Pietsch, A., *Operator Ideals*. (Transl. from German by the author). Amsterdam: North-Holland 1980.
- [10] Reed, M. and Simon, B., *Methods of Modern Mathematical Physics. I. Functional Analysis*. New York: Academic Press 1980.
- [11] Tikhonov, A. N., On the stability of inverse problems. *C. R. (Doklady) Acad. Sci. URSS (N.S.)* 39 (1943), 176 – 179.
- [12] Vinokurov, V. A., Regularization by continuous mappings. *Zh. Vychisl. Mat. Mat. Fiz.* 11 (1971), 1584 – 1586.
- [13] Tautenhahn, U., Optimality for ill-posed problems under general source conditions. *Numer. Funct. Anal. Optim.* 19 (1998)(3-4), 377 – 398.
- [14] Tautenhahn, U., Hämarik, U., Hofmann, B. and Shao, Y., Conditional stability estimates for ill-posed PDE problems by using interpolation. Preprint Ser. Faculty Math. Preprint 2011-16. Chemnitz Univ. of Technol. 2011. (available at <http://nbn-resolving.de/urn:nbn:de:bsz:ch1-qucosa-72654>).
- [15] Zălinescu, C., *Convex Analysis in General Vector Spaces*. Singapore: World Scientific 2002.

Received January 3, 2012; revised August 27, 2012