

Curvature theory of boundary phases: the two-dimensional case

ANDREA BRAIDES[†]

*Dipartimento di Matematica, Università di Roma 'Tor Vergata', via della Ricerca Scientifica,
00133 Rome, Italy*

AND

ANDREA MALCHIODI[‡]

*Institute for Advanced Study, School of Mathematics, 1 Einstein Drive, Princeton, NJ 08540,
USA*

[Received 7 March 2001 and in revised form 20 June 2002]

We describe the behaviour of minimum problems involving non-convex surface integrals in 2D, singularly perturbed by a curvature term. We show that their limit is described by functionals which take into account energies concentrated on vertices of polygons. Non-locality and non-compactness effects are highlighted.

Keywords: Surface energies; curvature functionals; phase transitions; Γ -convergence; non convex problems

1. Introduction

The starting point of this work is the study of minimum problems related to the equilibrium of elastic crystals (see for example, [15, 16] for the variational formulation, [8, 9] for a derivation of the model from statistical considerations, [3] for its links with Ising systems, and [20, 25] for an analogous derivation as a singular perturbation of the Allen–Cahn model). The model problem we have in mind is that of finding sets minimizing a (possibly highly anisotropic) ‘perimeter functional’ (i.e. a line integral on the boundary, that reduces to the usual perimeter if ∂E is regular and ψ is identically 1), of the form

$$\min \left\{ \int_{\partial E} \psi(\nu_E) d\mathcal{H}^1 : E_0 \subseteq E \right\}, \quad (1)$$

where the minimum is computed among all sets $E \subset \mathbb{R}^2$ with boundary of class C^1 and containing a fixed open set E_0 . Here, ψ is a Borel function, ν_E denotes the (appropriately oriented) tangent to E and \mathcal{H}^1 is the one-dimensional (Hausdorff) surface measure. Another model problem is that of *local minimizers* of the same anisotropic perimeter, related to

$$\min \left\{ \int_{\partial E} \psi(\nu_E) d\mathcal{H}^1 : |E_0 \Delta E| \leq \delta \right\}, \quad (2)$$

where $\delta > 0$ is a fixed constant ($A \Delta B$ stands for the symmetric difference of the sets A and B).

[†] Corresponding author. Email: braides@mat.uniroma2.it

[‡] Email: malchiod@ias.edu

Problems of the above type, and some of their perturbations, can be tackled following the so-called direct method of the calculus of variations. First, problems (1) and (2) can be ‘relaxed’ by admitting as competing sets all sets with finite perimeter (see [5, 18]). Then, if ψ is larger than a fixed constant on S^1 and if its homogeneous positive extension of degree one outside S^1 is a *convex* function, classical results imply that the surface integral in (1) and (2) is lower semicontinuous and coercive in the appropriate topology of the L^1 -convergence of characteristic functions. The application of the direct method of the calculus of variations thus yields the existence of minimizing sets of finite perimeter and, if ψ^2 is smooth and strictly convex (hypotheses are usually made on ψ^2 since ψ is positively homogeneous and hence non-convex on radial directions), regularity results for minimal surfaces assure that such minimizers are regular. On the other hand if ψ^2 is not convex, then the minimum problems (1) or (2) may not possess solutions. It can be seen (see for example, [21]) that the application of the direct method of the calculus of variations gives minimizing sequences with increasingly wiggly boundaries (even though with equi-bounded total area). Their limits can be described (see [4]) as minimizers of a ‘relaxed’ problem of the same type: in the case of (1), for example,

$$\min \left\{ \int_{\partial E} \bar{\psi}(v_E) d\mathcal{H}^1 : E_0 \subseteq E \right\}, \quad (3)$$

where the new surface energy density $\bar{\psi}$ is simply the convex envelope of the one-homogeneous extension of ψ to \mathbb{R}^2 . This process may lead to non-strictly convex integrands, which in turn may yield non-unique and non-smooth solutions. In this case, it may be necessary to consider higher-order terms in the surface energy to explain solutions with sharp corners and facets (see also [30]; a similar phenomenon is studied in [19]). Note that so far the problem can be framed in an n -dimensional framework, upon replacing curves by hypersurfaces.

In this paper we study, in a genuinely two-dimensional setting, the case when we add a singular perturbation by a curvature term in (1) (or analogously in (2)), obtaining a minimum problem of the form

$$\min \left\{ \int_{\partial E} \left(\psi(v_E) + \varepsilon^2 \kappa^2 \right) d\mathcal{H}^1 : E_0 \subseteq E \right\}, \quad (4)$$

where now the minimum is taken among sets with C^2 boundary and $\kappa(x)$ denotes the curvature of ∂E at x . In this way, oscillating boundaries are penalized if they introduce large curvatures.

In a way similar to [22, 24, 25], in order to understand the behaviour of minimizers for (4) we may study the (equivalent) ε -rescaled minimum problem

$$\min \left\{ \int_{\partial E} \left(\frac{\psi(v_E) - \bar{\psi}(v_E)}{\varepsilon} + \varepsilon \kappa^2 \right) d\mathcal{H}^1 : E_0 \subseteq E \right\}. \quad (5)$$

We assume for simplicity that $\psi(v_E) = \bar{\psi}(v_E)$ only on a finite number of directions v_1, \dots, v_N ($N > 2$). One can easily check that under this assumption ψ must satisfy

$$\psi(v) > \frac{\sin(v_{i+1} - v)}{\sin(v_{i+1} - v_i)} \psi(v_i) + \frac{\sin(v - v_i)}{\sin(v_{i+1} - v_i)} \psi(v_{i+1}), \quad \forall v \in (v_i, v_{i+1}), \forall i = 1, \dots, N$$

(we identify v_{N+1} with v_1). Note that this condition rules out a smooth behaviour near v_1, \dots, v_N as in the energies considered in [19]. The problem can then be rewritten as

$$\min \left\{ \int_{\partial E} \left(\frac{\varphi(v_E)}{\varepsilon} + \varepsilon \kappa^2 \right) d\mathcal{H}^1 : E_0 \subseteq E \right\}, \quad (6)$$

where $\varphi : S^1 \rightarrow [0, +\infty)$ vanishes only on those preferred directions.




FIG. 1. Two approximations with simple vertices.

Our main result is to describe the asymptotic behaviour as $\varepsilon \rightarrow 0$ of problem (6), showing that the minimizers E_ε tend, up to translations, to sets E which in turn minimize a limit energy. This limit energy can be computed by using the techniques of Γ -convergence (see [10, 11, 13]). We define the functional F_ε on sets of finite perimeter as

$$F_\varepsilon(E) = \begin{cases} \int_{\partial E} \left(\frac{\varphi(v_E)}{\varepsilon} + \varepsilon \kappa^2 \right) d\mathcal{H}^1 & \text{if } E \text{ is of class } C^2 \\ +\infty & \text{otherwise,} \end{cases} \tag{7}$$

and we compute their Γ -limit G with respect to the L^1 and L^1_{loc} -convergence of characteristic functions. As an example, in the simplest case when φ is *symmetric* with respect to both axes and the preferred directions coincide with the coordinate directions, the domain of the limit G is simply the set of the coordinate polyrectangles and $G(E) = c \#(V(E))$, where $V(E)$ is the set of vertices of the polyrectangle E . The constant c can be computed as

$$c = 2 \int_S \sqrt{\varphi(s)} d\mathcal{H}^1(s), \tag{8}$$

where S is the minimal arc in S^1 connecting $(1, 0)$ and $(0, 1)$. Hence, the limit problem is trivially

$$\min \left\{ c \#(V(E)) : E \text{ a coordinate polyrectangle, } E_0 \subseteq E \right\} \tag{9}$$

and the minimizers of the limit problem are simply all coordinate rectangles containing E_0 . Note that the limits of minimizers E_ε of (4) minimize both (3) and (9), so that they are coordinate rectangles (since they must minimize the number of vertices) containing E_0 of minimal perimeter (since the energy in (3) coincides with the Euclidean perimeter on polyrectangles).

In the general case, we show that the domain of the limit energy consists of those polygons whose tangents belong to the set of the preferred directions $\{v_1, \dots, v_N\}$, and that the limit energy G is much more complicated than (9). If E contains only simple vertices (which can be also phrased: if ∂E is locally Lipschitz) we define

$$F(E) = \sum \left\{ g(v^-(v), v^+(v)) : v \in V(E) \right\}, \tag{10}$$




FIG. 2. Approximation giving a non-local effect.

where g is given by

$$g(v_1, v_2) = 2 \int_{A(v_1, v_2)} \sqrt{\varphi(s)} \, d\mathcal{H}^1(s) \tag{11}$$

($A(v_1, v_2)$ is the minimal arc connecting v_1 and v_2 in S^1) and $v^\pm(v)$ are the two tangents at v . If, loosely speaking, E is such that approximating sequences E_ε may be chosen ‘uniformly close’ to E then we prove that $G(E) = F(E)$. In the general case, the value $G(E)$ is obtained as

$$G(E) = \inf \left\{ \liminf_j F(E_j) : E_j \rightarrow E, E_j \text{ with simple vertices} \right\}. \tag{12}$$

This formula hides two types of degenerate behaviours. First of all, we have to take into account that when two or more vertices coincide the set E may be approximated in many different ways and (12) highlights that the approximation of minimal energy must be chosen. Figure 1 shows two different approximations E_j with simple vertices of a set E with a ‘double’ vertex. In addition, the energy G may be *non-local*: in a sense, a polygon may be completed by adding segments pointing in some of the preferred directions, which must be considered as degenerate parts of E ; the energy $G(E)$ takes into account the ‘minimal’ of such completions. In Fig. 2 the corresponding optimal approximation is pictured. This effect is analogous to that highlighted in [6] for functionals depending on the square of the curvature. As a consequence of formula (12), we get that the study of minimizers of problems involving G corresponds to the analysis of minimizing sequences of corresponding problems involving F . In particular, we deduce that the limit problem of (6) admits as solutions all the convex polygons with tangents in the preferred directions.

Once the form of the Γ -limit is computed, we may apply our results also to other problems for which the solution is less immediate, such as

$$\min \left\{ \int_{\partial E} \left(\frac{\varphi(v_E)}{\varepsilon} + \varepsilon \kappa^2 \right) d\mathcal{H}^1 : |E_0 \Delta E| \leq \delta \right\}, \tag{13}$$

or

$$\min \left\{ \int_{\partial E} \left(\frac{\varphi(v_E)}{\varepsilon} + \varepsilon \kappa^2 \right) d\mathcal{H}^1 + |E_0 \Delta E| \right\}, \tag{14}$$

where E_0 is some fixed set. The latter problem is also of interest in some models in Image Processing where energies depending on curvatures and on (the number of) vertices are considered

(see [12, 23, 27]). Note that the solution to problem (14) may be given by a set E satisfying $G(E) < F(E)$ (see the example in Section 6.2).

Finally, we note that, since the solutions of the limit problem are polygons with fixed orientations, it is very tempting to link this approximation result to the theory of crystalline growth as recently developed (see [7, 17, 28, 29]), where non-strictly convex ψ are considered.

The paper is organized as follows. Section 2 contains the statement of the main results in terms of Γ -convergence and the necessary notation. In Sections 3 and 4 we prove the lower and upper bounds for the limit energy. In Section 5 some cases are dealt with when the limit energy is *local*: i.e., it can be written as a sum of energies concentrated on vertices. Finally, in Section 6 we consider the pathological case when we do not have a boundedness condition on the perimeter, giving a qualitative description of the shape of sequences with equi-bounded energy. We also give an example where Γ -limits computed in the L^1 and L^1_{loc} topology differ.

2. Main results

2.1 Statement of the main results

For every open set $E \subseteq \mathbb{R}^2$ of class C^2 and every $\varepsilon > 0$, we define the energy

$$F_\varepsilon(E) = \int_{\partial E} \left(\frac{1}{\varepsilon} \varphi(\nu) + \varepsilon \kappa^2 \right) d\mathcal{H}^1 \tag{15}$$

where $\nu = \nu(x)$ is the *tangent direction* to ∂E in x , defined in such a way that $(\nu_2, -\nu_1)$ coincides with the outer unit normal to ∂E in x . With \mathcal{H}^1 we denote the one-dimensional Hausdorff measure. The quantity $\kappa = \kappa(x)$ denotes the *curvature* of ∂E in x , and $\varphi : S^1 \rightarrow [0, +\infty)$ (we identify S^1 with $\mathbb{R} \bmod 2\pi$) is a continuous function with the following property:

$$\exists \nu_1, \dots, \nu_N \in S^1, \quad \nu_1 < \nu_2 < \dots < \nu_N < \nu_{N+1} = \nu_1 + 2\pi \quad \text{such that } \varphi^{-1}(0) = \{\nu_1, \dots, \nu_N\}.$$

We will always assume that

$$|\nu_i - \nu_{i+1}| < \pi, \quad i = 1, \dots, N.$$

We will identify sets E with their *characteristic function* χ_E , and then the functional given by formula (15) will be identified with the functional $F_\varepsilon : L^1(\mathbb{R}^2) \rightarrow [0, +\infty]$ given by

$$F_\varepsilon(u) = \begin{cases} \int_{\partial E} \left(\frac{1}{\varepsilon} \varphi(\nu) + \varepsilon \kappa^2 \right) d\mathcal{H}^1 & \text{if } u = \chi_E \text{ and } E \text{ is of class } C^2 \\ +\infty & \text{otherwise.} \end{cases} \tag{16}$$

With an additional slight abuse of notation, we say that a sequence of sets $(E_n) \subseteq \mathbb{R}^2$ converges to $E \subseteq \mathbb{R}^2$ in $L^1(\mathbb{R}^2)$ if $\chi_{E_n} \rightarrow \chi_E$ in $L^1(\mathbb{R}^2)$.

For $\theta_1, \theta_2 \in S^1$, $\theta_1 \neq \theta_2 + \pi$, let $A_{(\theta_1, \theta_2)}$ denote the shortest of the two arcs in S^1 connecting θ_1 and θ_2 . We assume that $A_{(\theta_1, \theta_2)}$ is oriented in the direction going from θ_1 to θ_2 . We define $g : S^1 \times S^1 \rightarrow [0, +\infty)$ in the following way:

$$g(\theta_1, \theta_2) = \begin{cases} 2 \int_{A_{(\theta_1, \theta_2)}} \sqrt{\varphi(\nu)} d\mathcal{H}^1(\nu) & \text{if } \theta_i \in \{\nu_1, \dots, \nu_N\} \quad i = 1, 2 \\ +\infty & \text{otherwise.} \end{cases} \tag{17}$$

Note that $g(\theta_2, \theta_1) = g(\theta_1, \theta_2)$.

An *admissible polygon* is a set $P \subseteq \mathbb{R}^2$ whose boundary is a polygonal composed of segments whose directions lie in the set $\{\nu_1, \dots, \nu_N\}$. We set

$$\mathcal{P} = \{P : P \text{ is an admissible polygon}\}.$$

We also define the class

$$\mathcal{R} = \left\{ P \in \mathcal{P} : \partial P \text{ is piecewise } C^1 \right\},$$

and we call *regular admissible polygons* the elements of \mathcal{R} . The difference between a general admissible polygon and a regular admissible polygon is that each vertex of the second is the endpoint of exactly two sides.

Given a polygon P in \mathbb{R}^2 , we define the set $V(P) \subseteq \mathbb{R}^2$ of the *vertices of P* to be

$$V(P) = \{x \in \partial P : \partial P \text{ is not } C^1 \text{ at } x\}.$$

We also define the functional $F_{\mathcal{R}} : \mathcal{P} \rightarrow \mathbb{R}$ in the following way:

$$F_{\mathcal{R}}(P) = \begin{cases} \sum_{v \in V(P)} g(\nu^-(v), \nu^+(v)), & \text{if } P \in \mathcal{R}; \\ +\infty, & \text{if } P \notin \mathcal{R}. \end{cases}$$

Here, $\nu^-(v), \nu^+(v)$ denote the directions of the two sides intersecting in $v \in V(P)$. This functional will be identified with a functional $F_{\mathcal{R}} : L^1(\mathbb{R}^2) \rightarrow [0, +\infty]$ in the same spirit of (16).

We also set

$$G = sc^-(F_{\mathcal{R}}),$$

where sc^- denotes the sequential lower semi-continuous envelope, understood in the sense of the L^1 -topology with uniform bounds on the perimeters, namely

$$sc^-(F_{\mathcal{R}})(E) := \inf \left\{ \liminf_n F_{\mathcal{R}}(E_n) : E_n \rightarrow E \text{ in } L^1(\mathbb{R}^2), \sup_n \mathcal{H}^1(\partial E_n) < +\infty \right\}.$$

REMARK 2.1 It can be easily checked that G is finite only on (characteristic functions of) admissible polygons. Moreover, given an admissible polygon P , there always exists a sequence (P_n) of regular polygons which converge to P in $L^1(\mathbb{R}^2)$, and for which $\sup_n \mathcal{H}^1(\partial P_n) < +\infty$ and $\sup_n F_{\mathcal{R}}(P_n) < +\infty$. In fact, it is sufficient to take

$$P_n = \left\{ x \in P : \text{dist}(x, \partial P) \leq \frac{1}{n} \right\}. \quad (18)$$

Note that in general the sequence given by formula (18) does not recover the infimum in the definition of $G(E)$.

REMARK 2.2 Given an admissible polygon P , there always exists a sequence (P_n) of regular polygons which converge to P in $L^1(\mathbb{R}^2)$, and for which $G(P) = F_{\mathcal{R}}(P_n)$ for sufficiently large n . In fact, whenever the quantity $F_{\mathcal{R}}(P_n)$ remains bounded, it ranges over a finite set of numbers, and the infimum is always attained.

Our main result is the following Γ -convergence theorem (for a general introduction to the subject we refer to [11, 13]).

THEOREM 2.1 For $\varepsilon > 0$, let $F_\varepsilon : L^1(\mathbb{R}^2) \rightarrow [0, +\infty]$ be the functional given by formula (16). Then

$$\Gamma\text{-}\lim_{\varepsilon \rightarrow 0} F_\varepsilon = G \tag{19}$$

with respect to the convergence in $L^1(\mathbb{R}^2)$ with uniform bound of the perimeter. By (19) we mean:

- (i) (closure) if $\sup_\varepsilon \mathcal{H}^1(E_\varepsilon) < +\infty$, $\sup_\varepsilon F_\varepsilon(E_\varepsilon) < +\infty$ and $E_\varepsilon \rightarrow u$ in $L^1(\mathbb{R}^2)$ then there exists $P \in \mathcal{P}$ such that $u = \chi_P$;
- (ii) (Γ -lim inf inequality) for all $P \in \mathcal{P}$ and for all $E_\varepsilon \rightarrow P$ in $L^1(\mathbb{R}^2)$ with $\sup_\varepsilon \mathcal{H}^1(E_\varepsilon) < +\infty$, we have $G(P) \leq \liminf_\varepsilon F_\varepsilon(E_\varepsilon)$;
- (iii) (Γ -lim sup inequality) for all $P \in \mathcal{P}$ there exists $E_\varepsilon \rightarrow P$ in $L^1(\mathbb{R}^2)$ with $\sup_\varepsilon \mathcal{H}^1(E_\varepsilon) < +\infty$ such that $G(P) = \lim_\varepsilon F_\varepsilon(E_\varepsilon)$.

REMARK 2.3 (Convergence of minimum problems) From Theorem 2.1 we obtain the convergence of the minimum values of problems (13) and (14) to the minimum values

$$\min \left\{ G(P) : P \in \mathcal{P}, |E_0 \Delta P| \leq \delta \right\} = \inf \left\{ \sum_{v \in V(P)} g(v^-(v), v^+(v)) : P \in \mathcal{R}, |E_0 \Delta P| \leq \delta \right\},$$

and

$$\min \left\{ G(P) + |E_0 \Delta P| : P \in \mathcal{P} \right\} = \inf \left\{ \sum_{v \in V(P)} g(v^-(v), v^+(v)) + |E_0 \Delta P| : P \in \mathcal{R} \right\},$$

respectively, provided that there exists a sequence of minimizers with equi-bounded perimeter. This property is a well known result of Γ -convergence, once we notice that the equi-boundedness of the perimeters ensures compactness of the minimizing sequence (upon, possibly, a translation), and that the constraints or the additional terms are ‘compatible’ with Γ -convergence. To check this for problem (13), it is sufficient to notice that a slight modification of the argument in the proof of Theorem 2.1(iii) allows us to suppose $|E_0 \Delta E| \leq \delta$. On the other hand, it is clear that the addition of the perturbation in (14) is compatible, since it is continuous with respect to the L^1 -convergence.

REMARK 2.4 Theorem 2.1 remains valid if F_ε takes the form

$$F_\varepsilon(E) = \int_{\partial E} \left(\frac{1}{\varepsilon} \varphi(v) + \varepsilon \kappa^2 \right) d\mathcal{H}^1 + c \mathcal{H}^1(\partial E), \tag{20}$$

with $c > 0$, i.e. if we add a term proportional to the length of ∂E . In this case, we similarly modify $F_{\mathcal{R}}(E)$ by setting

$$F_{\mathcal{R}}(E) = \sum_{v \in V(E)} g(v^-(v), v^+(v)) + c \mathcal{H}^1(\partial E)$$

on \mathcal{R} . Note that in this case the equi-boundedness condition on the perimeter is redundant.

We refer to Section 6 for the case when we drop the equi-boundedness condition on the perimeters and we consider the L^1_{loc} convergence. We conclude this section by deducing a convergence result for the minimum problems in (1) as an example of application of Theorem 2.1.

COROLLARY 2.1 Let ψ and $\bar{\psi}$ be as in the Introduction. Let E_0 be a bounded connected open set and let E_ε be minimizers for the problems

$$m_\varepsilon = \min \left\{ \int_{\partial E} \left(\psi(v_E) + \varepsilon^2 \kappa^2 \right) d\mathcal{H}^1 : E_0 \subseteq E \right\}.$$

Then, to within a translation and a passage to a subsequence, (E_ε) converges to a polygon P which minimizes both

$$m = \min \left\{ \int_{\partial E} \bar{\psi}(v_E) d\mathcal{H}^1 : E_0 \subseteq E \right\} \tag{21}$$

and

$$m^{(1)} = \min \left\{ \sum_{v \in V(E)} g(v^-(v), v^+(v)) : E_0 \subseteq E, E \in \mathcal{R} \right\}. \tag{22}$$

Proof. We just sketch the proof, including details only for the passages involving Γ -convergence.

By a relaxation argument (see [4]) and the density of sets with regular boundary we may suppose that E_ε converges to a minimizer \bar{E} of (21), which is connected since such is E_0 . On the other hand, E_ε is also a minimizer of

$$m_\varepsilon^{(1)} = \min \left\{ \int_{\partial E} \left(\frac{\psi(v_E)}{\varepsilon} + \varepsilon \kappa^2 \right) d\mathcal{H}^1 - \frac{m}{\varepsilon} : E_0 \subseteq E \right\}.$$

Define $\varphi = \psi - \bar{\psi}$. By using the construction of Section 4, it is easily seen that we have

$$m \leq \int_{\partial E_\varepsilon} \bar{\psi}(v_{E_\varepsilon}) d\mathcal{H}^1 \leq m + o(\varepsilon),$$

and that E_ε is an $o(1)$ -minimizer of

$$\begin{aligned} \tilde{m}_\varepsilon^{(1)} &= \min \left\{ \int_{\partial E} \left(\frac{\psi(v_E) - \bar{\psi}(v_E)}{\varepsilon} + \varepsilon \kappa^2 \right) d\mathcal{H}^1 : E_0 \subseteq E \right\} \\ &= \min \left\{ \int_{\partial E} \left(\frac{\varphi(v_E)}{\varepsilon} + \varepsilon \kappa^2 \right) d\mathcal{H}^1 : E_0 \subseteq E \right\}. \end{aligned}$$

We may apply Theorem 2.1 and Remark 2.4 as the perimeter of E_ε is equi-bounded since $\psi \geq c$. We then obtain that \bar{E} is a (convex) polygon minimizing both (21) and (22). \square

2.2 Notation

We introduce some preliminary notation and definitions.

Given a polygon P in \mathbb{R}^2 , we define a *side* of P to be the closure of a component of $\partial P \setminus V(P)$; we also define

$$\bar{s}(P) = \min \{ \text{length of } s : s \text{ is a side of } P \}.$$

If $\gamma^i : [a_i, b_i] \rightarrow \mathbb{R}^2$ $i = 1, 2$ are two curves with $\gamma^1(b_1) = \gamma^2(a_2)$, we define $\gamma^1 * \gamma^2 : [a_1, b_1 + b_2 - a_2] \rightarrow \mathbb{R}^2$ as

$$(\gamma^1 * \gamma^2)(t) = \begin{cases} \gamma^1(t) & t \in [a_1, b_1] \\ \gamma^2(t - b_1 + a_2) & t \in [b_1, b_1 + b_2 - a_2]. \end{cases}$$

Similarly, we define inductively

$$\gamma^1 * \dots * \gamma^k = \left(\gamma^1 * \dots * \gamma^{k-1} \right) * \gamma^k.$$

Given a curve $c : [a, b] \rightarrow \mathbb{R}^2$, we denote by $\text{im}(c)$ its image. If c is of class C^2 , and $t \in [a, b]$ is such that $c'(t) \neq 0$, we define $\kappa(c(t))$ to be the curvature of c at $c(t)$.

Given two sequences $(A_n), (B_n)$ of subsets of \mathbb{R}^2 such that $A_n \cap B_n = \emptyset$ for all n , and given $\nu \in S^1$, we say that (B_n) aligns with (A_n) in the direction ν (or that (B_n) and (A_n) are in line with the direction ν) if for every $\delta > 0$ it is

$$\left| \frac{x - y}{|x - y|} - \nu \right| < \delta, \quad \forall x \in A_n, \forall y \in B_n, \quad \text{for } n \text{ sufficiently large.}$$

We say that a family of curves $\gamma_n : (a_n, b_n) \rightarrow \mathbb{R}^2$ aligns in the direction ν if for every $\eta > 0$ and for every sequence of pairs (x_n, y_n) , $x_n, y_n \in \text{im}(\gamma_n)$, with $|x_n - y_n| > \eta$, and such that $\gamma_n^{-1}(x_n) > \gamma_n^{-1}(y_n)$, the sequence (x_n) aligns with (y_n) in the direction ν .

Given a piecewise C^1 curve $\gamma : S^1 \rightarrow \mathbb{R}^2$, and given a point x which does not belong to $\text{im}(\gamma)$, we define $\text{ind}(x, \gamma)$ to be the winding number of γ around x , namely (in complex notation)

$$\text{ind}(\gamma, x) = \frac{1}{2\pi i} \int_{S^1} \frac{\dot{\gamma}(t)}{\gamma(t) - x} dt.$$

Finally, we say that two segments $[x_1, x_2], [y_1, y_2] \subseteq \mathbb{R}^2$ do not intersect transversally if

$$(x_1, x_2) \cap (y_1, y_2) = \emptyset. \tag{NT}$$

Given $\theta_1, \theta_2 \in S^1$, the sum $\theta_1 + \theta_2$ will denote, unless it is explicitly remarked, the sum as elements of the group S^1 endowed with its natural structure (i.e the sum of \mathbb{R} modulo 2π).

3. The Γ -liminf inequality

This section is devoted to the proof of the Γ -lim inf inequality in Theorem 2.1.

We consider sequences $(E_n) \subseteq \mathbb{R}^2$, $\varepsilon_n \rightarrow 0^+$ for which

- (H₁) $\chi_{E_n} \rightarrow u$ in $L^1(\mathbb{R}^2)$;
- (H₂) $\sup_n \mathcal{H}^1(\partial E_n) < +\infty$;
- (H₃) $\sup_n F_{\varepsilon_n}(E_n) < +\infty$.

Our first aim is to prove that the sequence (E_n) converges in $L^1(\mathbb{R}^2)$ to some admissible polygon P . In fact we have the following result.

PROPOSITION 3.1 Let $\varepsilon_n \rightarrow 0$ and let (E_n) satisfy hypotheses (H₁)–(H₃). Then there exists an admissible polygon $P \in \mathcal{P}$ such that $u = \chi_P$, and for which there holds

$$G(u) \leq \liminf_n F_{\varepsilon_n}(E_n). \tag{23}$$

Before proving Proposition 3.1 we introduce some preliminary results.

LEMMA 3.1 Let $a, b, \delta \in \mathbb{R}$, $a < b$, $\delta > 0$, and let $v_i \in \varphi^{-1}(0)$. Then for every curve $\eta : [a, b] \rightarrow A_{(v_i, v_{i+1})}$ of class C^1 with

$$\eta(a) = v_i + \delta, \quad \eta(b) = v_{i+1} - \delta,$$

we have

$$\int_a^b \left(\frac{1}{\varepsilon} \varphi(\eta(t)) + \varepsilon \|\dot{\eta}(t)\|^2 \right) dt \geq g(v_i, v_{i+1}) + o_\delta(1), \tag{24}$$

where $o_\delta(1) \rightarrow 0$ as $\delta \rightarrow 0$.

Proof. This is a simple consequence of the Young inequality: in fact we obtain

$$\begin{aligned} \int_a^b \left(\frac{1}{\varepsilon} \varphi(\eta(t)) + \varepsilon \|\dot{\eta}(t)\|^2 \right) dt &\geq 2 \int_a^b \sqrt{\varphi(\eta(t))} |\dot{\eta}(t)| dt \\ &\geq 2 \int_{v_i+\delta}^{v_{i+1}-\delta} \sqrt{\varphi(\eta(t))} dt \geq 2 \int_{v_i}^{v_{i+1}} \sqrt{\varphi(\eta(t))} dt + o_\delta(1), \end{aligned}$$

which is the desired inequality. □

Now we consider a family of curves $\gamma_n : S^1 \rightarrow \mathbb{R}^2$ of class C^2 with the following properties:

$$\sup_n \int_{S^1} \left(\frac{1}{\varepsilon_n} \varphi \left(\frac{\dot{\gamma}_n}{|\dot{\gamma}_n|} \right) + \varepsilon_n \left(\frac{d}{dt} \frac{\dot{\gamma}_n}{|\dot{\gamma}_n|} \right)^2 \right) dt = M < +\infty, \tag{25}$$

$$\sup_n \int_{S^1} |\dot{\gamma}_n| dt < +\infty. \tag{26}$$

We suppose also that the curves γ_n are parametrized proportionally to their arc length, namely that there holds

$$|\dot{\gamma}_n(t)| = \frac{1}{2\pi} \int_{S^1} |\dot{\gamma}_n| ds; \quad \text{for all } t \in S^1 \text{ and for all } n \in \mathbb{N}.$$

We want to describe the limit shape of the curves γ_n when $n \rightarrow +\infty$. In order to do this, we set for $\delta > 0$

$$S_\delta = S^1 \setminus ([v_1 - \delta, v_1 + \delta] \cup \dots \cup [v_N - \delta, v_N + \delta]),$$

and

$$C(\delta) = \inf_{v \in S_\delta} \varphi(v). \tag{27}$$

If $\eta : [a, b] \rightarrow S_\delta$ is a curve of class C^1 , then there holds clearly

$$\int_a^b \left(\frac{1}{\varepsilon} \varphi(\eta(t)) + \varepsilon \|\dot{\eta}(t)\|^2 \right) dt \geq \frac{1}{\varepsilon} (b - a) C(\delta); \tag{28}$$

hence, using (25) and (28) with $\eta = \dot{\gamma}_n$ and $\varepsilon = \varepsilon_n$, we deduce

$$\mathcal{H}^1(\{t \in [0, T_n] : \dot{\gamma}_n(t) \in S_\delta\}) \leq \frac{\varepsilon_n}{C(\delta)} \int_{\dot{\gamma}_n \in S_\delta} \left(\frac{1}{\varepsilon_n} \varphi \left(\frac{\dot{\gamma}_n(t)}{|\dot{\gamma}_n|} \right) + \varepsilon_n \kappa^2(\gamma_n(t)) \right) dt \leq \frac{\varepsilon_n M}{C(\delta)}.$$

From this inequality we deduce the existence of a sequence $\delta_n \rightarrow 0$ such that

$$\mathcal{L}^1(I_n) \rightarrow 0 \quad \text{as } n \rightarrow +\infty, \tag{29}$$

where we have set

$$I_n = \left\{ t \in S^1 : \frac{\dot{\gamma}_n(t)}{|\dot{\gamma}_n(t)|} \in S_{\delta_n} \right\}.$$

Since S_δ is open, the components of I_n are at most countable: denote by $I_n^j = (\sigma_n^j, \theta_n^j)$, $j = 1, \dots, k_n$, those components of I_n for which $\dot{\gamma}_n(a_n^j) \neq \dot{\gamma}_n(b_n^j)$. From assumption (H_3) and from Lemma 3.1 it follows that $\sup_n k_n < +\infty$ and so, passing to a subsequence, we can assume that $k_n = \bar{k}$ for all n . We also set

$$J_n = S^1 \setminus \bigcup_{j=1}^{\bar{k}} I_n^j. \tag{30}$$

LEMMA 3.2 Let $(\theta_n^h, \sigma_n^{h+1})$ be a component of J_n such that

$$\dot{\gamma}_n(\theta_n^h) = |\dot{\gamma}_n| (v_i \pm \delta_n), \quad \text{for some } v_i \in \{v_1, \dots, v_N\}.$$

Then $\gamma_n|_{(\theta_n^h, \sigma_n^{h+1})}$ aligns in the direction v_i .

Proof. Let $\eta > 0$, and let $\alpha_n, \beta_n \in (\theta_n^h, \sigma_n^{h+1})$ be such that $|\gamma_n(\alpha_n) - \gamma_n(\beta_n)| > \eta$. Then, since γ_n is parametrized proportionally to the arc length, there holds

$$\eta < |\gamma_n(\alpha_n) - \gamma_n(\beta_n)| \leq \int_{\alpha_n}^{\beta_n} |\dot{\gamma}_n(t)| \, dt = |\dot{\gamma}_n| \mathcal{L}_1((\alpha_n, \beta_n)), \tag{31}$$

so in particular we have

$$\frac{\eta}{\sup_j |\dot{\gamma}_j|} \leq \frac{\eta}{|\dot{\gamma}_n|} < \mathcal{L}_1((\alpha_n, \beta_n)) < 2\pi.$$

Hence by equation (26) the quantities $\mathcal{L}_1((\alpha_n, \beta_n))$ are uniformly bounded from above and from below. Set

$$\rho_n = \int_{(\alpha_n, \beta_n) \setminus I_n} \dot{\gamma}_n(t) \, dt; \quad \tau_n = \int_{(\alpha_n, \beta_n) \cap I_n} \dot{\gamma}_n(t) \, dt.$$

Equations (26) and (29) imply that $\tau_n \rightarrow 0$ as $n \rightarrow +\infty$. We also have

$$\int_{(\alpha_n, \beta_n) \setminus I_n} \dot{\gamma}_n(t) \, dt = |\dot{\gamma}_n| \mathcal{L}_1((\alpha_n, \beta_n) \setminus I_n) v_i + \int_{(\alpha_n, \beta_n) \setminus I_n} (\dot{\gamma}_n(t) - |\dot{\gamma}_n| v_i) \, dt,$$

so from (31) and the definition of I_n we deduce

$$\rho_n = |\dot{\gamma}_n| \mathcal{L}_1((\alpha_n, \beta_n)) v_i + o(1). \tag{32}$$

From this expression and from the fact that $\rho_n \rightarrow 0$ it follows that

$$\frac{\gamma_n(\beta_n) - \gamma_n(\alpha_n)}{|\gamma_n(\beta_n) - \gamma_n(\alpha_n)|} - v_i = \frac{\rho_n + \tau_n}{|\rho_n + \tau_n|} - v_i = \frac{\rho_n}{|\rho_n|} - v_i + o(1) = o(1).$$

This concludes the proof. □

The next lemma shows that γ_n , restricted to a component of J_n , converges uniformly to a segment in direction v_i parametrized by arc length.

LEMMA 3.3 Let $(\theta_n^h, \sigma_n^{h+1})$ be a component of J_n as in Lemma 3.2. Then, given any $\rho > 0$, there exists $n_\rho \in \mathbb{N}$ such that

$$\|\gamma_n(\beta_n) - \gamma_n(\alpha_n) - |\dot{\gamma}_n|(\beta_n - \alpha_n)v_i\| < \rho, \quad \forall \alpha_n, \beta_n \in (\theta_n^h, \sigma_n^{h+1}), \quad \forall n \geq n_\rho. \quad (33)$$

Proof. This follows easily from $\gamma_n(\beta_n) - \gamma_n(\alpha_n) = \rho_n + \tau_n$, equation (32), and the fact that $\tau_n \rightarrow 0$ as $n \rightarrow +\infty$. \square

Let us now introduce some additional notation. We define the class

$$\mathcal{C} = \left\{ \{\gamma^1, \dots, \gamma^k\} \mid \gamma^i : S^1 \rightarrow \mathbb{R}^2 \text{ is piecewise } C^1, \frac{\dot{\gamma}^i}{|\dot{\gamma}^i|} \in \{v_1, \dots, v_N\} \text{ a.e. in } S^1, i = 1, \dots, k \right\}.$$

Let $\gamma = \{\gamma^1, \dots, \gamma^k\} \in \mathcal{C}$. Then for all i $\text{im}(\gamma^i)$ is composed by a finite number of segments with directions v_{j_1}, \dots, v_{j_i} . We define $\tilde{F} : \mathcal{C} \rightarrow \mathbb{R}$ in the following way:

$$\tilde{F}(\gamma) = \sum_{i=1}^k \sum_{h=1}^{j_i} g(v_{j_h}, v_{j_{h+1}}).$$

PROPOSITION 3.2 Let ε_n and let (E_n) satisfy hypotheses (H_1) – (H_3) above. Let $\gamma_n^j, j = 1, \dots, l$ (passing to a subsequence we can suppose that the number l is independent of n) be parameterizations of the components of ∂E_n . Then there exist a polygon $P \in \mathcal{P}$ such that $u = \chi_P$, there exist integers $h, k, k \leq h \leq l$, and there exists $\gamma = \{\gamma^1, \dots, \gamma^k\} \in \mathcal{C}$ with the following properties:

- (Γ_1) $\gamma_n^j \rightarrow \gamma^j, j = 1, \dots, k$, uniformly on S^1 , and $\gamma_n^j \rightarrow x^j \in \mathbb{R}^2, j = k + 1, \dots, h$, uniformly on S^1 .
- (Γ_2) the segments of $\text{im}(\gamma)$ do not intersect transversally;
- (Γ_3) for a.e. $x \in \mathbb{R}^2$, it is $\sum_{i=1}^k \text{ind}(\gamma^i, x) \in \{0, 1\}$, and $\chi_P(x) = \sum_{i=1}^k \text{ind}(\gamma^i, x)$;
- (Γ_4) $\tilde{F}(\gamma) \leq \liminf_n F_{\varepsilon_n}(E_n)$.

Proof. Let $i \in \{1, \dots, h\}$, and consider the sequence of curves γ_n^i which parameterize the i -th component of ∂E_n . This sequence satisfies conditions (25) and (26), hence we can repeat for them the constructions above. Let J_n^i be the counterpart of the set J_n for the curve γ_n^i . We can also suppose that the number of components of J_n^i is a constant k^i independent of n . From Lemma 3.3 it follows that

$$\text{up to translation, } \gamma_n^i \rightarrow \gamma^i \text{ uniformly on } S^1, \quad \text{for some curve } \gamma^i \in \mathcal{C}, \quad (34)$$

or

$$\text{up to translation, } \gamma_n^i \rightarrow x^i \text{ uniformly on } S^1, \quad \text{for some point } x^i \in \mathbb{R}^2. \quad (35)$$

Up to a permutation of the indices, there exist $h, k \in \mathbb{N}$, $0 \leq k \leq h \leq l$ such that $(\gamma_n^1), \dots, (\gamma_n^k)$ converge uniformly in S^1 to some $\gamma^1, \dots, \gamma^k \in \mathcal{C}$, and that $(\gamma_n^{k+1}), \dots, (\gamma_n^h)$ converge uniformly in S^1 to some points $x^{k+1}, \dots, x^h \in \mathbb{R}^2$. Define γ to be $\gamma = \{\gamma^1, \dots, \gamma^k\}$, so that also $\gamma \in \mathcal{C}$. Condition (I_1) is automatically satisfied. Condition (I_2) follows easily from the fact that the sets E_n are of class C^2 .

From equations (34) and (35) we deduce

$$\mathcal{H}^2(B_\gamma) = 0, \quad \text{where} \quad B_\gamma = \left(\bigcup_{i=1}^k \text{im}(\gamma^i) \right) \cup \left(\bigcup_{i=k+1}^h x^i \right). \tag{36}$$

By the continuity of the winding number with respect to the uniform convergence we have

$$\lim_n \sum_{i=1}^h \text{ind}(\gamma_n^i, x) = \sum_{i=1}^h \text{ind}(\gamma^i, x), \quad \text{for all } x \in \mathbb{R}^2 \setminus B_\gamma,$$

hence, since the index is integer-valued there holds

$$\sum_{i=1}^h \text{ind}(\gamma_n^i, x) = \sum_{i=1}^k \text{ind}(\gamma^i, x), \quad \text{for } n \text{ large and for all } x \in \mathbb{R}^2 \setminus B_\gamma.$$

From this we can deduce that, setting

$$P = \left\{ x \in \mathbb{R}^2 \setminus B_\gamma : \lim_n \sum_{i=1}^h \text{ind}(\gamma_n^i, x) = 1 \right\},$$

we have

$$\begin{cases} x \in P & \Rightarrow & x \in E_n & \text{for } n \text{ large;} \\ x \notin P & \Rightarrow & x \notin E_n & \text{for } n \text{ large.} \end{cases}$$

This implies that

$$\chi_{E_n} \rightarrow \chi_P \quad \text{as } n \rightarrow +\infty, \quad \text{a.e. in } \mathbb{R}^2,$$

and proves condition (I_3) . Property (I_4) follows from Lemma 3.1. □

LEMMA 3.4 Suppose that $\gamma \in \mathcal{C}$ satisfies conditions (I_1) and (I_2) in Proposition 3.2. Then there exists a sequence of regular polygons $(P_n) \subseteq \mathcal{R}$ such that

$$\chi_{P_n} \rightarrow \chi_P \quad \text{in } L^1(\mathbb{R}^2); \quad F_{\mathcal{R}}(P_n) \leq \tilde{F}(\gamma). \tag{37}$$

Proof. For the proof of this Lemma we refer to [14]. □

Finally, we are in position to prove Proposition 3.1.

Proof of Proposition 3.1. Let P be the polygon given by Proposition 3.2, and let $(P_n) \subseteq \mathcal{R}$ be the sequence of regular polygons given by Lemma 3.4. Then, by equation (37) and by property (I_4) there holds

$$F_{\mathcal{R}}(P_n) \leq \tilde{F}(\gamma) \leq \liminf_n F_{\varepsilon_n}(E_n).$$

Finally, by the definition of G we have

$$G(P) \leq \liminf_n F_{\mathcal{R}}(P_n) \leq \liminf_n F_{\varepsilon_n}(E_n).$$

This concludes the proof. □

4. The Γ -limsup inequality

The goal of this section is to prove the Γ -lim sup inequality in Theorem 2.1. Starting with a regular admissible polygon P , we modify it near its vertices and we obtain a sequence of sets E_n of class C^2 which converge to P and such that $F_{\varepsilon_n}(E_n)$ is as small as possible. Then we treat the general case of an admissible polygon by approximating it with regular polygons.

PROPOSITION 4.1 Let $P \in \mathcal{R}$ be an admissible regular polygon. Then, given any sequence $\varepsilon_n \rightarrow 0^+$, there exists a sequence of sets (E_n) of class C^2 such that

$$\chi_{E_n} \rightarrow \chi_P \text{ in } L^1(\mathbb{R}^2); \quad \limsup_n F_{\varepsilon_n}(E_n) \leq F_{\mathcal{R}}(P).$$

Proof. Let v be a vertex of P : since P is regular, there are exactly two sides of P intersecting v . Without loss of generality, we can suppose that the directions of these sides, which we denote by l_1 and l_2 , are v_1 and v_2 respectively. Let $\lambda : \left(-\frac{1}{2} |l_1|, \frac{1}{2} |l_2|\right) \rightarrow \mathbb{R}^2$ be defined by

$$\lambda(t) = \begin{cases} v - t v_1, & t \in \left[-\frac{1}{2} |l_1|, 0\right]; \\ v + t v_2, & t \in \left[0, \frac{1}{2} |l_2|\right]. \end{cases} \tag{38}$$

The curve λ defined in this way parametrizes part of l_1 for $t < 0$ and part of l_2 for $t > 0$. Our aim is to find a sequence of regular curves $\lambda_n : \left[-\frac{1}{2} |l_1|, \frac{1}{2} |l_2|\right] \rightarrow \mathbb{R}^2$ with the following properties:

$$\lambda_n \rightarrow \lambda \quad \text{uniformly on } \left[-\frac{1}{2} |l_1|, \frac{1}{2} |l_2|\right]; \tag{39}$$

$$\lim_n \int_{\left(-\frac{1}{2} |l_1|, \frac{1}{2} |l_2|\right)} \left(\frac{1}{\varepsilon_n} \varphi \left(\frac{\dot{\lambda}_n}{|\dot{\lambda}_n|} \right) + \varepsilon_n \kappa^2(\lambda_n) \right) dt = g(v_1, v_2). \tag{40}$$

Since φ is assumed to be of class C^1 in $S^1 \setminus \{v_1, \dots, v_N\}$, the following Cauchy problem:

$$\begin{cases} y'(t) = \sqrt{\varphi(y(t))} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} y(t) \\ y(0) = \frac{v_1 + v_2}{2}. \end{cases} \tag{41}$$

admits a unique maximal solution $u : (a, b) \rightarrow S^1$, with $-\infty \leq a < 0, 0 < b \leq +\infty$. It is immediate to check that u is a C^1 increasing function which tends to v_1 (respectively, v_2) as $t \rightarrow a$ (respectively, $t \rightarrow b$).

For every $c, d \in (a, b)$, with $c < 0 < d$ (c and d will be taken sufficiently close to a and b), define $e = c - (u(c) - v_1)$ and $f = d + (v_2 - u(d))$; note that $e < c < d < f$. We can find a nondecreasing function $\eta : [e, f] \rightarrow A_{(v_1, v_2)}$ of class C^1 , such that

$$\eta(e) = v_1; \quad \dot{\eta}(e) = 0; \tag{42}$$

$$\eta(f) = v_2; \quad \dot{\eta}(f) = 0; \tag{43}$$

$$\begin{cases} \eta(t) = u(t), & t \in (c, d); \\ |\dot{\eta}(t)| < 2|u(c) - v_1|, & t \in (e, c); \\ |\dot{\eta}(t)| < 2|v_2 - u(d)|, & t \in (d, f). \end{cases} \quad (44)$$

For $\varepsilon > 0$, let η_ε denote the unique continuous extension of η to the interval $\left[-\frac{1}{\varepsilon} \frac{1}{2} |l_1|, \frac{1}{\varepsilon} \frac{1}{2} |l_2|\right]$ for which

$$\eta_\varepsilon(t) = \begin{cases} v_1, & t \in \left[-\frac{1}{\varepsilon} \frac{1}{2} |l_1|, e\right]; \\ v_2, & t \in \left[f, \frac{1}{\varepsilon} \frac{1}{2} |l_2|\right]. \end{cases} \quad (45)$$

Finally, for $\varepsilon_n \rightarrow 0^+$, define $\lambda_n : \left[-\frac{1}{2} |l_1|, \frac{1}{2} |l_2|\right] \rightarrow \mathbb{R}^2$ to be

$$\lambda_n(t) = v + \int_0^t \eta_{\varepsilon_n} \left(\frac{s}{\varepsilon_n} \right) ds, \quad t \in \left[-\frac{1}{2} |l_1|, \frac{1}{2} |l_2|\right].$$

Since η_{ε_n} is an S^1 -valued curve of class C^1 , it follows that λ_n is of class C^2 and is parametrized by arc length. For $t < 0$ it turns out that

$$\begin{aligned} \lambda_n(t) - \lambda(t) &= v + \int_0^t \eta_{\varepsilon_n} \left(\frac{s}{\varepsilon_n} \right) ds - v - t v_1 \\ &= \int_0^{\varepsilon_n e} \eta_{\varepsilon_n} \left(\frac{s}{\varepsilon_n} \right) ds + \int_{\varepsilon_n e}^t \eta_{\varepsilon_n} \left(\frac{s}{\varepsilon_n} \right) ds - t v_1. \end{aligned}$$

Since $|\eta_{\varepsilon_n}| = 1$, and since $\eta_{\varepsilon_n}(t) = v_1$ for $t < e$, it follows that

$$\lambda_n(t) - \lambda(t) \rightarrow 0, \quad \text{uniformly for } t \in \left[-\frac{1}{2} |l_1|, 0\right].$$

In the same way one can show that

$$\lambda_n(t) - \lambda(t) \rightarrow 0, \quad \text{uniformly for } t \in \left[0, \frac{1}{2} |l_1|, \right],$$

so we have proved (39).

Using the definition of λ_n and the change of variable $\frac{s}{\varepsilon_n} = y$, we find

$$\int_{-\frac{1}{2} |l_1|}^{\frac{1}{2} |l_2|} \left(\frac{1}{\varepsilon_n} \varphi \left(\frac{\dot{\lambda}_n}{|\dot{\lambda}_n|} \right) + \varepsilon_n \kappa^2(\lambda_n) \right) ds = \int_{-\frac{1}{2} \frac{1}{\varepsilon_n} |l_1|}^{\frac{1}{2} \frac{1}{\varepsilon_n} |l_2|} \left(\varphi(\eta_{\varepsilon_n}) + (\dot{\eta}_{\varepsilon_n})^2 \right) dy;$$

then, taking into account equation (45), one has

$$\int_{-\frac{1}{2} \frac{1}{\varepsilon_n} |l_1|}^{\frac{1}{2} \frac{1}{\varepsilon_n} |l_2|} \left(\varphi(\eta_{\varepsilon_n}) + (\dot{\eta}_{\varepsilon_n})^2 \right) dt = \int_e^f \left(\varphi(\eta_{\varepsilon_n}) + (\dot{\eta}_{\varepsilon_n})^2 \right) dt.$$

Dividing the interval (e, f) into (e, c) , (c, d) and (d, f) , by equation (41) we get

$$\begin{aligned} \int_e^f \left(\varphi(\eta_{\varepsilon_n}) + (\dot{\eta}_{\varepsilon_n})^2 \right) dt &\leq |c - e| \left(\sup_{(e,c)} \varphi + \sup_{(e,c)} \dot{\eta}_{\varepsilon_n}^2 \right) \\ &\quad + g(v_1, v_2) + |f - d| \left(\sup_{(d,f)} \varphi + \sup_{(d,f)} \dot{\eta}_{\varepsilon_n}^2 \right). \end{aligned}$$

Using the expression of e, f , and taking into account (44), we deduce

$$\int_e^f \left(\varphi(\eta_{\varepsilon_n}) + (\dot{\eta}_{\varepsilon_n})^2 \right) dt \leq g(v_1, v_2) + |u(c) - v_1| \left(\sup_{(e,c)} \varphi + 4|u(c) - v_1|^2 \right) + |v_2 - u(d)| \left(\sup_{(e,c)} \varphi + 4|v_2 - u(d)|^2 \right).$$

Hence, choosing $c = c(n)$ and $d = d(n)$ depending on n and such that

$$|u(c) - v_1| + |v_2 - u(d)| \rightarrow 0 \quad \text{as } n \rightarrow +\infty,$$

also (40) follows.

Now consider a component Θ of ∂P . Let v_1, \dots, v_{i_Θ} denote an ordering of the vertices of Θ along the parametrization of λ , and let λ_j be the curve defined above corresponding to the vertex $v_j, j = 1, \dots, i_\Theta$. Then we can choose as parametrization for Θ the piecewise- C^2 curve λ_Θ given by

$$\lambda_\Theta = \lambda_1 * \dots * \lambda_{i_\Theta}.$$

For $j \in \{1, \dots, i_\Theta\}$, let $\lambda_{j,n}$ be a sequence of curves which satisfy (39) and (40) with $\lambda = \lambda_j$ and $v^-(v_j), v^+(v_j)$ instead of v_1 and v_2 . If we consider the sequence of curves

$$\lambda_{p,n} = \lambda_{1,n} * \dots * \lambda_{i_\Theta,n}, \quad n \in \mathbb{N},$$

they will converge uniformly to λ_Θ on their domain (a_Θ, b_Θ) . In general the curve $\lambda_{\Theta,n}$ is not closed, but since λ_Θ is closed there holds

$$\lambda_{\Theta,n}(a_\Theta) - \lambda_{\Theta,n}(b_\Theta) \rightarrow 0, \quad \text{as } n \rightarrow +\infty.$$

Consider the curve $\lambda_{1,n}$. Since the directions of its two rectilinear parts are linearly independent, it is sufficient to modify slightly the length of these parts in such a way that $\lambda_{\Theta,n}$ transforms into a closed curve $\bar{\lambda}_{\Theta,n}$.

Repeating this procedure for all the components of ∂P we obtain a set E_n whose boundary is parametrized by the union of the curves $(\bar{\lambda}_{\Theta,n})_\Theta$. The sequence E_n will satisfy the required properties in the proposition. \square

REMARK 4.1 From the proof of Proposition 4.1 it follows that we can choose λ_n satisfying (40) and

$$\lambda_n \text{ coincides with } \lambda \text{ in a neighbourhood of } \left\{ -\frac{1}{2}|l_1|, \frac{1}{2}|l_2| \right\}; \quad \|\lambda_n - \lambda\|_\infty \leq 2(|e(n)| + |f(n)|) \varepsilon_n, \tag{46}$$

where $e(n) = c(n) - u(c(n)) + v_1$ and $f(n) = d(n) + u(c(n)) - v_2$.

As an immediate consequence of Proposition 4.1 we have the following corollary.

COROLLARY 4.1 (**Γ -lim sup inequality**) Let $P \in \mathcal{P}$ be an admissible polygon. Then, for every (ε_n) with $\varepsilon_n \rightarrow 0^+$ there exists a sequence of sets E_n of class C^2 such that

$$E_n \rightarrow P \text{ in } L^1(\mathbb{R}^2) \quad \text{and} \quad \limsup_n F_{\varepsilon_n}(E_n) \leq G(P).$$

Proof. By Remark 2.2, there exists a sequence $(P_k)_k \subseteq \mathcal{R}$ of regular polygons such that

$$\chi_{P_k} \rightarrow \chi_P \text{ in } L^1(\mathbb{R}^2); \quad \limsup_k F_{\mathcal{R}}(P_k) = G(P); \quad \sup_k \mathcal{H}^1(\partial P_k) < +\infty.$$

Then, by Proposition 4.1, for every $k \in \mathbb{N}$ there exists a sequence (E_n^k) of sets of class C^2 such that

$$E_n^k \rightarrow P_k; \quad F_{\varepsilon_n}(E_n^k) \rightarrow F_{\mathcal{R}}(P_k), \quad \text{as } n \rightarrow +\infty.$$

Hence we can choose a sequence of natural numbers $n(k)$ with $n(k_2) > n(k_1)$ if $k_2 > k_1$ such that

$$\left\| \chi_{E_n^k} - \chi_{P_k} \right\|_{L^1(\mathbb{R}^2)} \leq \frac{1}{k}, \quad F_{\varepsilon_n}(E_n^k) \leq F_{\mathcal{R}}(P_k) + \frac{1}{k}.$$

So, if we choose

$$E_n = E_n^k, \quad \text{for } n(k) \leq n < n(k+1),$$

the sequence (E_n) satisfies the desired properties. □

5. Some local cases

In this section we study some specific cases for which the Γ -limit G has a local expression, namely it is the sum over the vertices of an energy depending only on each single vertex.

5.1 A non-symmetric case

In this section we treat the following particular case. We assume that the function φ satisfies the conditions

- (i) $\varphi \in C^1(S^1)$;
- (ii) $\varphi^{-1}(0) = \{v_1, \dots, v_N\}$, and for all i we have $-v_i \notin \varphi^{-1}(0)$ if $v_i \in \varphi^{-1}(0)$.

Under these hypotheses we will prove that $\Gamma\text{-lim}_{\varepsilon \rightarrow 0} F_\varepsilon$ has a local expression. Namely, to every vertex of an admissible polygon P is associated a quantity $E(v)$, and $\Gamma\text{-lim}_{\varepsilon \rightarrow 0} F_\varepsilon(P)$ is the sum of $E(v)$ over the vertices v of P , see Proposition 5.4. In order to state this result precisely we introduce some additional notation.

Let P be an admissible polygon, and let v be a vertex of P . Let l_1, \dots, l_{2k} be the sides of P which intersect at v . If condition (i) above is satisfied, then for each of these segments l_j , $j = 1, \dots, 2k$, is uniquely determined a tangent direction $v(l_j) = v_{i_j} \in \varphi^{-1}(0)$.

To each l_j we can associate an orientation $\sigma_v(l_j)$ with respect to v , namely we set

$$\begin{cases} \sigma_v(l_j) = -1, & \text{if } l_j \text{ is oriented toward } v; \\ \sigma_v(l_j) = 1, & \text{if } -l_j \text{ is oriented toward } v, \end{cases} \quad j = 1, \dots, 2k.$$

If the segments l_1, \dots, l_{2k} , are ordered in such a way that $\sigma_v(l_j) v_{i_1} < \sigma_v(l_j) v_{i_2} < \dots < \sigma_v(l_j) v_{i_{2k}}$, then clearly it must be

$$\sigma_v(l_j) \cdot \sigma_v(l_{j+1}) = -1, \quad j = 1, \dots, 2k - 1, \quad \text{and} \quad \sigma_v(l_{2k}) \cdot \sigma_v(l_1) = -1.$$

DEFINITION 5.1 An admissible decomposition ω of v is a partition of l_1, \dots, l_{2k} in pairs (l_i^-, l_i^+) , $i = 1, \dots, k$, such that

$$\sigma_v(l_i^-) = -1, \quad \sigma_v(l_i^+) = 1; \quad i = 1, \dots, k, \tag{AD_1}$$

and

$$[v(l_i^-), v(l_i^+)] \cap [v(l_j^-), v(l_j^+)] = \emptyset, \quad i, j = 1, \dots, k, \quad i \neq j, \tag{AD_2}$$

where $[v(l_h^-), v(l_h^+)]$ in the above formula denotes the segment in \mathbb{R}^2 joining $v(l_h^-)$ and $v(l_h^+)$. We set also

$$\Omega_v = \{\omega \mid \omega \text{ is an admissible decomposition for } v\}.$$

REMARK 5.2 Every vertex $v \in V(P)$ admits an admissible decomposition. In fact, if the vectors $v_{i_1}, \dots, v_{i_{2k}}$, are ordered in such a way that $v_{i_1} < v_{i_2} < \dots < v_{i_{2k}}$, then one can take

$$l_i^- = l_{2i-1}, \quad l_i^+ = l_{2i} \quad i = 1, \dots, k.$$

To each admissible decomposition $\omega = \{(l_i^-, l_i^+)\}_i$ of a vertex v , we associate the energy $\psi(\omega)$ defined by

$$\psi(\omega) = \sum_{i=1}^k g(v(l_i^-), v(l_i^+)), \tag{47}$$

and we define

$$E(v) = \min \{\psi(\omega) \mid \omega \in \Omega_v\}. \tag{48}$$

LEMMA 5.3 Let $\gamma \in \mathcal{C}$ satisfy conditions (Γ_2) and (Γ_3) in Proposition 3.2, and let P be the polygon associated to γ from (Γ_3) . Let $v \in V(P)$ and let l_1, \dots, l_{2k} be the segments of γ which intersect v . Let l_1^-, \dots, l_k^- be the segments of $\{l_1, \dots, l_{2k}\}$ which are oriented toward v , and let l_1^+, \dots, l_k^+ be the elements of $\{l_1, \dots, l_{2k}\}$ which, following the parametrization of γ , are after l_1^-, \dots, l_k^- respectively. Then $\omega_v^\gamma = (l_j^-, l_j^+)$, $j = 1, \dots, k$, is an admissible decomposition of v .

Proof. Property (AD_1) is immediate to verify. Condition (AD_2) is equivalent to the fact that adjacent sides must have opposite orientations. \square

PROPOSITION 5.4 Suppose φ satisfies conditions (i) and (ii) above and let P be an admissible polygon. Then

$$G(\chi_P) = \sum_{v \in V(P)} E(v). \tag{49}$$

Proof. Let us prove first the Γ -lim inf inequality. Let $\varepsilon_n \rightarrow 0$, let (E_n) satisfy hypotheses $(H_1) - (H_3)$, and let $u = \chi_P$. Let $\gamma \in \mathcal{C}$ be given by Proposition 3.2. Then, if ω_v^γ is given by Lemma 5.3, there holds

$$\tilde{F}(\gamma) = \sum_{v \in V(P)} E(\omega_v^\gamma).$$

Finally, using equation (48) and property (Γ_4) in Proposition 3.2 we get

$$\sum_{v \in V(P)} E(v) \leq \sum_{v \in V(P)} E(\omega_\gamma^v) = \tilde{F}(\gamma) \leq \liminf_n F_{\varepsilon_n}(E_n).$$

This proves the Γ -lim inf inequality; let us now turn to the Γ -lim sup inequality.

Let $v \in V(P)$ and let $\bar{\omega}_v$ be an admissible decomposition of v which realizes the minimum energy, namely for which

$$\psi(\bar{\omega}_v) = E(v).$$

The set of the admissible decompositions $\bar{\omega}_v$, when v ranges over $V(P)$, determines an element $\gamma \in \mathcal{C}$ in the following way.

Given a side l^1 of P , are uniquely determined two vertices v_1 and v_2 and two indices i_1 and i_2 for which, if we set $\bar{\omega}_1 = \{(l_{i_1,1}^+, l_{i_1,1}^-)\}_i$ and $\bar{\omega}_2 = \{(l_{i_2,2}^+, l_{i_2,2}^-)\}_i$, we have

$$l^1 = l_{i_1,1}^+ = l_{i_2,2}^-.$$

Let $l^2 = l_{i_2,2}^-$; reasoning as above, there exist an unique vertex v_3 an unique index i_3 for which, if we set $\bar{\omega}_3 = \{(l_{i_3,3}^+, l_{i_3,3}^-)\}_i$, there holds

$$l^2 = l_{i_2,2}^+ = l_{i_3,3}^-.$$

Continuing in this way, we obtain a first segment l^{j_1} for which $l^{j_1} = l_{i_1,1}^-$. Let $c^i : [\alpha^i, \beta^i] \rightarrow \mathbb{R}^2$, $i = 1, \dots, j$ be parameterizations of the sides l^i , and consider the closed curve γ^1 defined by

$$\gamma^1 = c^1 * \dots * c^j.$$

Up to reparameterizations, we can suppose that γ^1 is defined on S^1 . In the same way, we define the curves $\gamma^2, \dots, \gamma^k : S^1 \rightarrow \mathbb{R}^2$ until all the remaining sides of P are considered.

Now we fix a number $M > 0$, a sequence of positive numbers δ_n converging to zero, and we consider the set

$$A_n = \{\cup \bar{B}_{M\delta_n}(v) \mid v \in V(P)\}.$$

Let γ^1 be the curve defined above, and let $\xi_n^1 = \{t \in S^1 : \gamma^1(t) \in A_n\}$. The set ξ_n^1 is a finite union of closed intervals $[\alpha_n^{1,i}, \beta_n^{1,i}]$, $i = 1, \dots, j_1$, and we denote by $(\sigma_n^{1,i}, \tau_n^{1,i})$, $i = 1, \dots, j_1$, the components of $S^1 \setminus \xi_n^1$, where we have taken $\sigma_n^{1,i} = \beta_n^{1,i}$. Setting $\bar{c}_n^{1,i} = \gamma^1|_{[\alpha_n^{1,i}, \beta_n^{1,i}]}$, and $\hat{c}_n^{1,i} = \gamma^1|_{[\sigma_n^{1,i}, \tau_n^{1,i}]}$, it is clear that

$$\gamma^1 = \bar{c}_n^{1,1} * \hat{c}_n^{1,1} * \bar{c}_n^{1,2} * \dots * \bar{c}_n^{1,j_1} * \hat{c}_n^{1,j_1}.$$

Of course, we can write a similar expression for $\gamma^2, \dots, \gamma^k$.

We observe that the maps $\bar{c}_n^{i,l}$, $i = 1, \dots, k$, $l = 1, \dots, j_i$, are union of two rectilinear curves with directions $v_-^{i,l}$ and $v_+^{i,l}$ (following the order of the parametrization), while the curves $\hat{c}_n^{i,l}$ are rectilinear with direction $v_+^{i,l}$.

We define also the curves

$$\tilde{c}_n^{i,l}(t) = \bar{c}_n^{i,l}(t) + \delta_n \left(v_-^{i,l} + v_+^{i,l} \right), \quad t \in [\alpha_n^{i,l}, \beta_n^{i,l}];$$

where the above sum $v_-^{i,l} + v_+^{i,l}$ is now a sum of elements in \mathbb{R}^2 . It follows from property (AD_2) that the images of the curves $\tilde{c}_n^{i,l}$ are all disjoint when i varies from 1 to k , and l varies from 1 to j_i . We have also

$$\frac{(\tilde{c}_n^{i,l})'(\beta_n^{i,l})}{|(\tilde{c}_n^{i,l})'(\beta_n^{i,l})|} = v_+^{i,l} = v_-^{i,l+1} = \frac{(\tilde{c}_n^{i,l+1})'(\alpha_n^{i,l+1})}{|(\tilde{c}_n^{i,l+1})'(\alpha_n^{i,l+1})|}, \quad \text{for all } i = 1, \dots, k, l = 1, \dots, j_i. \quad (50)$$

Now we choose a function $\eta : [0, 1] \rightarrow [0, 1]$ of class C^∞ and which satisfies the following properties:

$$\begin{cases} \eta = 0 \text{ in a neighbourhood of } 0; \\ \eta = 1 \text{ in a neighbourhood of } 1; \\ \eta' \geq 0; \quad |\eta'| \leq 2; \quad |\eta''| \leq 4, \end{cases} \quad (51)$$

and for $a, b > 0$, let $\eta_{a,b} : [0, 1] \rightarrow \mathbb{R}^2$ be defined by

$$\eta_{a,b}(t) = \begin{pmatrix} a t \\ b \eta(t) \end{pmatrix}; \quad t \in [0, 1].$$

Using simple computations, one can check that

$$|\kappa(\eta_{a,b}(t))| \leq 4 \frac{b}{a}, \quad \text{for all } t \in [0, 1]. \quad (52)$$

We recall $\kappa(\eta_{a,b}(t))$ denotes the curvature of $\eta_{a,b}$ at $\eta_{a,b}(t)$.

Fix $i \in \{1, \dots, k\}$, $l \in \{1, \dots, j_i\}$, and consider the points $\tilde{c}_n^{i,l}(\beta_n^{i,l})$ and $\tilde{c}_n^{i,l+1}(\alpha_n^{i,l+1})$; then by equation (50) there exist unique numbers $a, b > 0$, and an unique affine isometry T of \mathbb{R}^2 for which the curve $T \circ \eta_{a,b}$ possesses the following properties (we omit the dependence of a, b, T on the indices i, l and n):

$$\begin{cases} T \circ \eta_{a,b}(0) = \tilde{c}_n^{i,l}(\beta_n^{i,l}); & T \circ \eta_{a,b}(1) = \tilde{c}_n^{i,l+1}(\alpha_n^{i,l+1}); \\ (T \circ \eta_{a,b})'(0) = v_+^{i,l}; & (T \circ \eta_{a,b})'(1) = v_+^{i,l}. \end{cases}$$

One can easily check that

$$|b| \leq 2 \delta_n, \quad a \geq \frac{1}{2} \bar{s}; \quad \text{for } n \text{ large};$$

see Section 2.2 for the definition of \bar{s} . From these equations and from (52), it follows that

$$\left| \frac{(T \circ \eta_{a,b})'}{|(T \circ \eta_{a,b})'|} - v_i \right| \leq 8 \frac{\delta_n}{\bar{s}(P)}; \quad |\kappa(T \circ \eta_{a,b})| \leq 16 \frac{\delta_n}{\bar{s}(P)}. \quad (53)$$

Denote by $\tilde{C}_n^{i,l}$ the curve $\eta_{a,b}$, where a, b are chosen as above depending on i, l, n , and consider

$$\tilde{\gamma}_n^i = \tilde{c}_n^{1,1} * \tilde{C}_n^{1,1} * \tilde{c}_n^{1,2} * \dots * \tilde{c}_n^{1,j_1} * \tilde{C}_n^{1,j_1}.$$

It follows from the first equation in (53) that if M is sufficiently large, then the curves $\tilde{\gamma}_n^i$, $i = 1, \dots, k$ are simple, mutually disjoint, and the union of their images is the boundary of a piecewise C^2 set $\tilde{E}_n \subseteq \mathbb{R}^2$. It is clear that $\tilde{E}_n \rightarrow P$ in $L^1(\mathbb{R}^2)$.

Let $\varepsilon_n \rightarrow 0$: for every $i \in \{1, \dots, k\}$ and every $l \in \{1, \dots, j_i\}$, let $a^{i,l}, b^{i,l}$, etc., be the analogous of a, b, c, d in the proof of Proposition 4.1 when we consider $v^{i,l}, v_-^{i,l}$ and $v_+^{i,l}$. Since φ is assumed to be of class C^1 , we can choose $\delta_n \rightarrow 0$ and $e^{i,l}(n), f^{i,l}(n)$ with the following properties:

$$(i) \lim_n \frac{\delta_n}{\varepsilon_n (|e^{i,l}(n)| + |f^{i,l}(n)|)} = +\infty \quad \text{for all } i \in \{1, \dots, k\} \text{ and every } l \in \{1, \dots, j_i\};$$

$$(ii) \lim_n \frac{1}{\varepsilon_n} C \left(\frac{8}{s} \delta_n \right) = 0;$$

see (27) for the definition of $C(\delta)$.

We have

$$\int_{[0,1]} \frac{1}{\varepsilon_n} \varphi \left(\frac{(\tilde{C}_n^{i,l})'}{|\tilde{C}_n^{i,l}|} \right) dt + \varepsilon_n \int_{[0,1]} \kappa^2(\tilde{C}_n^{i,l}) dt \leq \frac{1}{\varepsilon_n} C \left(\frac{8}{s} \delta_n \right) + \varepsilon_n \left(\frac{16}{s} \right)^2 \delta_n^2.$$

From property (ii) above and from (53), it follows that

$$\lim_n \left(\int_{[0,1]} \frac{1}{\varepsilon_n} \varphi \left(\frac{(\tilde{C}_n^{i,l})'}{|\tilde{C}_n^{i,l}|} \right) dt + \varepsilon_n \int_{[0,1]} \kappa^2(\tilde{C}_n^{i,l}) dt \right) = 0. \tag{54}$$

By Remark 4.1, for every $i \in \{1, \dots, k\}$, every $l \in \{1, \dots, j_i\}$ and every n sufficiently large it is possible to choose a curve $\bar{C}_n^{i,l} : [\alpha_n^{i,l}, \beta_n^{i,l}] \rightarrow \mathbb{R}^2$ such that

$$|\bar{C}_n^{i,l}(t) - \tilde{c}_n^{i,l}(t)| \leq 2\varepsilon_n (|e^{i,l}(n)| + |f^{i,l}(n)|); \tag{55}$$

$$\bar{C}_n^{i,l} \text{ coincides with } \tilde{c}_n^{i,l} \text{ in a neighbourhood of } \{\alpha_n^{i,l}, \beta_n^{i,l}\}; \tag{56}$$

$$\int_{[\alpha_n^{i,l}, \beta_n^{i,l}]} \frac{1}{\varepsilon_n} \varphi \left(\frac{(\bar{C}_n^{i,l})'}{|\bar{C}_n^{i,l}|} \right) dt + \int_{[\alpha_n^{i,l}, \beta_n^{i,l}]} \varepsilon_n \kappa^2(\bar{C}_n^{i,l}) dt \rightarrow g(v_-^{i,l}, v_+^{i,l}). \tag{57}$$

Let $\tilde{\gamma}^i$ be the curve defined by

$$\tilde{\gamma}_n^i = \bar{C}_n^{i,1} * \tilde{c}_n^{i,1} * \bar{C}_n^{i,2} * \dots * \bar{C}_n^{i,j_i} * \tilde{c}_n^{i,j_i}.$$

From (56) it follows that the curve $\tilde{\gamma}_n^i, i = 1, \dots, k$, are curves of class C^2 , while (55) implies that they are simple, mutually disjoint, and the union of their images is the boundary of a C^2 set $\bar{E}_n \subseteq \mathbb{R}^2$. Again, $\bar{E}_n \rightarrow P$ in $L^1(\mathbb{R}^2)$. Moreover from (57) one can deduce that

$$\limsup_n F_{\varepsilon_n}(\bar{E}_n) \leq \sum_{v \in V(P)} E(v).$$

This concludes the proof. □

5.2 A symmetric case

In this section we treat the case in which the admissible polygons are polyrectangles, and the function φ is symmetric with respect to the axes x and y . A direct proof of Theorem 5.1 is also presented in [10], Appendix B.

THEOREM 5.1 Let $\mathbf{e}_1, \mathbf{e}_2$ be the canonical basis of \mathbb{R}^2 , and suppose that φ satisfies the conditions

$$\varphi^{-1}(0) = \{v_1, \dots, v_4\}, \quad \text{where} \quad v_1 = \mathbf{e}_1, \quad v_2 = \mathbf{e}_2, \quad v_3 = -\mathbf{e}_1, \quad v_4 = -\mathbf{e}_2, \quad (58)$$

and

$$g_0 := g(v_i, v_{i+1}) \quad \text{is independent of } i = 1, \dots, 4. \quad (59)$$

Then the admissible polygons are polyrectangles, and for every $P \in \mathcal{P}$

$$G(P) = g_0 \times \#\{\text{vertices of } P\}.$$

Proof. Let us prove first the Γ -lim inf inequality. We note that if $P \in \mathcal{R}$, then one has

$$F_{\mathcal{R}}(P) = \#\{\text{vertices of } P\} = \#\{\text{sides of } P\}. \quad (60)$$

Let $E \in \mathcal{P}$, and let $E_k \in \mathcal{R}$, $E_k \rightarrow E$ in $L^1(\mathbb{R}^2)$. Then, since it must be $\#\{\text{sides of } E_k\} \geq \#\{\text{sides of } E\}$ for k large, it follows from (60) that

$$F_{\mathcal{R}}(E_k) \geq \#\{\text{sides of } E\} \geq \#\{\text{vertices of } E\}, \quad \text{for } k \text{ large.}$$

Hence we have also

$$G(E) = sc^-(F_{\mathcal{R}})(E) \geq \#\{\text{vertices of } E\},$$

which is the Γ -lim inf inequality. Let us prove now the Γ -lim sup inequality. Given a polyrectangle E , and given a number $\sigma > 0$, consider the set E_σ defined by

$$E_\sigma = \{x \in E : \text{dist}(x, \partial E) \leq \sigma\}.$$

Then, if σ is sufficiently small, $E_\sigma \in \mathcal{R}$, and $\#\{\text{sides of } E_\sigma\} \leq \#\{\text{sides of } E\}$. This concludes the proof. \square

6. Pathological cases

In this section we consider the case in which the uniform boundedness of the perimeter is not required in the definition of convergence. In this situation, it is possible to have the convergence in the $L^1_{\text{loc}}(\mathbb{R}^2)$ sense without having convergence in $L^1(\mathbb{R}^2)$, so we are led to defining

$$\overline{G}(E) = \inf \{\liminf_n F_{\varepsilon_n}(E_n) : E_n \rightarrow E \text{ in } L^1_{\text{loc}}(\mathbb{R}^2)\}.$$

We recall that, by Theorem 2.1, $G(E) = \inf \{\liminf_n F_{\varepsilon_n}(E_n) : E_n \rightarrow E \text{ in } L^1(\mathbb{R}^2), \sup_n \mathcal{H}_1(\partial E_n) < +\infty\}$, so it is clearly $\overline{G}(E) \leq G(E)$. In Section 6.1 we describe the asymptotic shape of the subsequences (E_n) for which $\sup_n F_{\varepsilon_n}(E_n) < +\infty$, highlighting similarities with Section 3. However, in general $\overline{G} < G$. In Section 6.2 we are able to exhibit a function φ and a polygon P for which $\overline{G}(P)$ is strictly less than $G(P)$.

6.1 *Asymptotic shape of minimizers*

In this section we describe the limit shape of a sequence of sets (E_n) for which just condition (H_3) holds, while condition (H_2) —the uniform boundedness of the perimeter—is lifted.

We suppose that ∂E_n possesses just one connected component; the general case requires only simple modifications. Let γ_n be a parametrization of ∂E_n proportional to the arc length. First, we note that Lemmas 3.1 and 3.2 remain unchanged, so we can define the quantities $\delta_n \rightarrow 0$, I_n and J_n with $|I_n| \rightarrow 0$, just as we did in Section 3. In general, we do not have uniform convergence on the components of J_n as in Lemma 3.3. However, it can be recovered under a suitable rescaling.

LEMMA 6.1 Let J_n be defined as in (30), and let $(\theta_n^h, \sigma_n^{h+1})$ be a component of J_n such that

$$\dot{\gamma}_n(\theta_n^h) = |\dot{\gamma}_n| (v_i \pm \delta_n) \quad \text{for some } v_i \in \{v_1, \dots, v_N\},$$

and such that $|\gamma_n(\theta_n^h) - \gamma_n(\sigma_n^{h+1})| \rightarrow +\infty$ as $n \rightarrow +\infty$. Let $\tilde{\gamma}_n : (\theta_n^h, \sigma_n^{h+1})$ be defined by

$$\tilde{\gamma}_n(t) = \frac{1}{|\gamma_n(\theta_n^h) - \gamma_n(\sigma_n^{h+1})|} (\gamma_n(t) - \gamma_n(\theta_n^h)).$$

Then we have

$$\sup_{t \in (\theta_n^h, \sigma_n^{h+1})} |\tilde{\gamma}_n(t) - v_i t| \rightarrow 0, \quad \text{as } n \rightarrow +\infty.$$

Proof. We have $|\tilde{\gamma}_n(t)| \leq C$ on $(\theta_n^h, \sigma_n^{h+1})$, and moreover

$$\int_{(\theta_n^h, \sigma_n^{h+1})} \frac{1}{\varepsilon_n} \varphi \left(\frac{\dot{\tilde{\gamma}}_n}{|\dot{\tilde{\gamma}}_n|} \right) dt \leq \int_{(\theta_n^h, \sigma_n^{h+1})} \frac{1}{\varepsilon_n} \varphi \left(\frac{\dot{\gamma}_n}{|\dot{\gamma}_n|} \right) dt.$$

Hence, considering the curve $\tilde{\gamma}_n$, we are in the same situation of Lemma 3.3, so our statement follows. \square

Passing to a subsequence, we find an integer k , and k sequences of points $(x_n^1), \dots, (x_n^k)$ such that

$$\text{dist}(\gamma_n(I_n), \{x_n^1, \dots, x_n^k\}) \rightarrow 0, \quad \text{as } n \rightarrow +\infty.$$

In this case, the mutual distances between the points x_n^i can diverge. However, it turns out that the sequences of points $\{x_n^1, \dots, x_n^k\}$ arrange in ‘clusters’, and the limit shape of some rescaled portion of E_n is still polygonal.

In fact, let

$$d_n^1 = \sup \{|x_n^i - x_n^j| : i, j \in \{1, \dots, k\}, i \neq j\},$$

and consider the sequence of sets

$$E_n^1 = (d_n^1)^{-1} (E_n - x_n^1).$$

Let γ_n^1 be a parametrization of ∂E_n^1 . Then, there exists a number $k^1 \leq k$ and k_1 sequences of points $(x_n^{1,1}), \dots, (x_n^{1,k_1})$ such that

$$\text{dist}(\gamma_n^1(I_n), \{x_n^{1,1}, \dots, x_n^{1,k_1}\}) \rightarrow 0, \quad \text{as } n \rightarrow +\infty.$$

From Lemma 6.1, it is easy to see that the sequence E_n^1 converges in $L^1(\mathbb{R}^2)$ to some admissible polygon $P^1 \in \mathcal{P}$.

If we choose a different rescaling for the set E_n , we can obtain some ‘finer’ structures of these sets. In fact, consider the set of indices $\{i_1, \dots, i_j\} \subseteq \{1, \dots, k\}$, for which

$$\lim_n (d_n^1)^{-1} |x_n^{i_l} - x_n^1| \rightarrow 0, \quad l = 1, \dots, j,$$

and define d_n^2 to be

$$d_n^2 = \sup\{|x_n^{i_l} - x_n^{i_h}| : l, h \in \{i_1, \dots, i_j\}, l \neq h\};$$

it is clear that $(d_n^1)^{-1} d_n^2 \rightarrow 0$. Consider the sequence of sets E_n^2

$$E_n^2 = (d_n^2)^{-1} (E_n - x_n^1).$$

Then, using the arguments above, one can check that $E_n^2 \rightarrow P^2$ in $L^1_{\text{loc}}(\mathbb{R}^2)$, where $P^2 \subseteq \mathbb{R}^2$ is a set whose boundary is composed of segments, half-lines or lines oriented in the directions $\{v_1, \dots, v_N\}$. In some sense, P^2 could be considered as a polygon with some sides of infinite length.

Of course, the same result holds true if one considers suitable rescalings at the points x_n^i for $i \neq 1$.

6.2 An example in which $\overline{\mathbf{G}} \neq \mathbf{G}$

In this section we consider the following particular case, namely $\varphi^{-1}(0) = \{v_1, \dots, v_5\}$ with

$$v_1 = (1, 0); \quad v_2 = (0, 1); \quad v_3 = \frac{\sqrt{2}}{2}(-1, 1); \quad v_4 = \frac{\sqrt{2}}{2}(-1, -1); \quad v_5 = (0, -1), \quad (61)$$

and

$$g(v_1, v_2) = g(v_2, v_3) = g(v_4, v_5) = g(v_1, v_2) = 1; \quad g(v_5, v_1) = 5. \quad (62)$$

Let $p_i, q_i \in \mathbb{R}^2, i = 1, \dots, 3$, be given by

$$p_1 = (0, 0), \quad p_2 = (1, 0), \quad p_3 = (1, 1); \quad q_1 = (2, 0), \quad q_2 = (3, 0), \quad q_3 = (2, 1),$$

and let P be the polygon defined as follows (see Fig. 3(a)):

$$P = \left\{ \sum_{i=1}^3 t_i p_i \mid t_i \geq 0, \sum_{i=1}^3 t_i = 1 \right\} \cup \left\{ \sum_{i=1}^3 t_i q_i \mid t_i \geq 0, \sum_{i=1}^3 t_i = 1 \right\}.$$

It is clear from (61) that $P \in \mathcal{P}$. We show that in this case $\overline{G}(P)$ is strictly less than $G(P)$.

In fact, let $(E_n) \subseteq \mathcal{R}$ be a sequence of sets of class C^2 as in Fig. 3(b). It is clear that the boundary of E_n has just one component and from (62) one can check that $F_{\mathcal{R}}(P_n) = 17 + o(1)$, where $o(1) \rightarrow 0$ as $n \rightarrow +\infty$.

Now, suppose by contradiction that $G(P) = \overline{G}(P) \leq 17$, namely that there exists $(E_n) \subseteq \mathbb{R}^2$ with

$$E_n \rightarrow E \text{ in } L^1(\mathbb{R}^2), \quad \sup_n \mathcal{H}_1(\partial E_n) < +\infty, \quad \lim_n F_{\varepsilon_n}(E_n) \leq 17.$$




FIG. 3. A set with $\overline{G}(P) < G(P)$ and its optimal approximations

Passing to a subsequence, we can assume that the number of the components of ∂E_n is a fixed number k independent of n . By Lemma 3.1, it turns out that $F_{\varepsilon_n}(E_n) \geq 9k + o(1)$, so, since we are assuming that $F_{\varepsilon_n}(E_n) \leq 17 + o(1)$, it follows that $k = 1$.

Let $\gamma_n : S^1 \rightarrow \mathbb{R}^2$ be a parametrization of ∂E_n proportional to the arc length. Then we can apply Proposition 3.2, and we find a curve $\gamma : S^1 \rightarrow \mathbb{R}^2$, $\gamma \in \mathcal{C}$, for which $\gamma_n \rightarrow \gamma$ uniformly on S^1 , and for which $P = \{x \in \mathbb{R}^2 : \text{ind}(\gamma, x) = 1\}$.

Consider the set

$$A = \{t \in S^1 : 1 < (\gamma)_x(t) < 2, \dot{\gamma}(t) \in \{v_3, v_4\}\}.$$

Since γ has just one component, it must be $A \neq \emptyset$, and since $-v_3$ and $-v_4$ do not belong to $\varphi^{-1}(0)$, it should be $\gamma(A) \subseteq \partial P$, which is a contradiction.

REMARK 6.1 It is possible to have $\overline{G}(P) < G(P)$ also when the (strong) L^1 convergence is required in the definition of \overline{G} . In fact, if φ is of class C^1 , one could choose a sequence of approximating sets (E_n) as in Fig. 3(c). Reasoning as in Section 5, one can prove that $F_{\varepsilon_n}(E_n) = 17 + o(1)$.

7. Acknowledgements

AB is part of the MURST project *Calculus of Variations*; AM has been supported by MURST under the project *Variational Methods and Nonlinear Differential Equations* and by a Fulbright Fellowship for the academic year 2000–2001. We thank Antonio Di Carlo for a useful critical reading of the manuscript. A great part of the present work was accomplished at SISSA, Trieste.

REFERENCES

1. ALBERTI, G. & BELLETTINI, G. A non-local anisotropic model for phase transitions: asymptotic behaviour of rescaled energies. *European J. Appl. Math.* **9**, (1998) 261–284.
2. ALBERTI, G. & BELLETTINI, G. A nonlocal anisotropic model for phase transitions. I. The optimal profile problem. *Math. Ann.* **310**, (1998) 527–560.
3. ALBERTI, G., BELLETTINI, G., CASSANDRO, M., & PRESUTTI, E. Surface tension in Ising systems with Kac potentials. *J. Statist. Phys.* **82**, (1996) 743–796.

4. AMBROSIO, L. & BRAIDES, A. Functionals defined on partitions in sets of finite perimeter. II. Semicontinuity, relaxation and homogenization. *J. Math. Pures Appl.* **69**, (1990) 307–333.
5. AMBROSIO, L., FUSCO, N., & PALLARA, D. *Functions of Bounded Variation and Free Discontinuity Problems*. Oxford University Press, Oxford (2000).
6. BELLETTINI, G., DAL MASO, G., & PAOLINI, M. Semicontinuity and relaxation properties of a curvature depending functional in 2D. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **20**, (1993) 247–297.
7. BELLETTINI, G., GUGLIONE, R., & NOVAGA, M. Approximation to driven motion by crystalline curvature in two dimensions. *Adv. Math. Sci. Appl.* **10**, (2000) 467–493.
8. BODINEAU, T. The Wulff construction in three and more dimensions. *Comm. Math. Phys.* **207**, (1999) 197–229.
9. BODINEAU, T., IOFFE, D., & VELENIK, Y. Rigorous probabilistic analysis of equilibrium crystal shapes. Probabilistic techniques in equilibrium and nonequilibrium statistical physics. *J. Math. Phys.* **41**, (2000) 1033–1098.
10. BRAIDES, A. *Approximation of Free-Discontinuity Problems*. Springer, Berlin (1998).
11. BRAIDES, A. *Γ -convergence for Beginners*. Oxford University Press, Oxford (2002).
12. COSCIA, A. On curvature sensitive image segmentation. *Nonlinear Anal.* **39**, (2000) 711–730.
13. DAL MASO, G. *An Introduction to Γ -convergence*. *Progress in Nonlinear Differential Equations and their Applications* **8**. Birkhäuser, Boston (1993).
14. DAL MASO, G. & ROMANELLI, M. Paper in preparation.
15. FONSECA, I. Variational methods for elastic crystals. *Arch. Rational Mech. Anal.* **97**, (1987) 189–220.
16. FONSECA, I. The lower quasiconvex envelope of the stored energy function for an elastic crystal. *J. Math. Pures Appl.* **67**, (1988) 175–195.
17. GIGA, Y., GURTIN, M. E., & MATIAS, J. On the dynamics of crystalline motions. *Japan J. Indust. Appl. Math.* **15**, (1998) 7–50.
18. GIUSTI, E. *Minimal Surfaces and Functions of Bounded Variation*. *Monographs in Mathematics* **80**. Birkhäuser, Boston, MA (1984).
19. GUGLIONE, R. & PAOLINI, M. Numerical simulations of crystalline motion by mean curvature with Allen-Cahn relaxation. *Free Boundary Problems, Theory and Applications (Zakopane, 1995)*, Pitman Res. Notes Math. Ser. vol. 363. Longman, Harlow (1996) pp. 203–216.
20. GURTIN, M. E. Some results and conjectures in the gradient theory of phase transitions, in “Metastability and incompletely posed problems, Proc. Workshop, Minneapolis/Minn. 1984/85”. *IMA Vol. Math. Appl.* **3**, (1987) 135–146.
21. GURTIN, M. E., SONER, H. M., & SOUGANIDIS, P. E. Anisotropic motion of an interface relaxed by the formation of infinitesimal wrinkles. *J. Differential Equations* **119**, (1995) 54–108.
22. LUCKHAUS, S. & MODICA, L. The Gibbs-Thompson relation within the gradient theory of phase transitions. *Arch. Rational Mech. Anal.* **107**, (1989) 71–83.
23. MANTEGAZZA, C. Curvature varifolds with boundary. *J. Differential Geom.* **43**, (1996) 807–843.
24. MODICA, L. The gradient theory of phase transitions and the minimal interface criterion. *Arch. Rational Mech. Anal.* **98**, (1987) 123–142.
25. MODICA, L. & MORTOLA, S. Un esempio di Γ^- -convergenza. *Boll. Un. Mat. Ital. B* **14**, (1977) 285–299.
26. MUMFORD, D. Elastica and computer vision. *Algebraic Geometry and its Applications (West Lafayette, IN, 1990)*. Springer, New York (1994) pp. 491–506.
27. NITZBERG, M., MUMFORD, D., & SHIOTA, T. Filtering, Segmentation and Depth. *Lecture Notes in Computer Science*, **662**. Springer, Berlin (1993).
28. TAYLOR, J. E. On the global structure of crystalline surfaces. *Discrete Comput. Geom.* **6**, (1991) 225–262.
29. TAYLOR, J. E. Constructions and conjectures in crystalline nondifferential geometry. *Differential Geometry*, Pitman Monographs Surveys Pure Appl. Math. vol. 52. Longman, Harlow (1991) pp. 321–336.
30. TAYLOR, J. E. & CAHN, J. W. Diffuse interfaces with sharp corners and facets: phase field models with strongly anisotropic surfaces. With an appendix by Jason Yunger. *Phys. D* **112**, (1998) 381–411.