

## Consistency result for a non monotone scheme for anisotropic mean curvature flow

ERIC BONNETIER

*LJK, Université de Joseph Fourier, B.P. 53, 38041 Grenoble Cedex 9, France*

*E-mail: [eric.bonnetier@imag.fr](mailto:eric.bonnetier@imag.fr)*

ELIE BRETIN

*ICJ, INSA de Lyon, 69621 Villeurbanne Cedex, France*

*E-mail: [bretin@cmap.polytechnique.fr](mailto:bretin@cmap.polytechnique.fr)*

ANTONIN CHAMBOLLE

*CMAP, Ecole Polytechnique, 91128 Palaiseau, France*

*E-mail: [antonin.chambolle@polytechnique.fr](mailto:antonin.chambolle@polytechnique.fr)*

[Received 3 June 2010 and in revised form 1 February 2012]

In this paper, we propose a new scheme for anisotropic motion by mean curvature in  $\mathbb{R}^d$ . The scheme consists of a phase-field approximation of the motion, where the nonlinear diffusive terms in the corresponding anisotropic Allen–Cahn equation are linearized in the Fourier space. In real space, this corresponds to the convolution with a specific kernel of the form

$$K_{\phi,t}(x) = \mathcal{F}^{-1} \left[ e^{-4\pi^2 t \phi^\circ(\xi)} \right] (x).$$

We analyse the resulting scheme, following the work of Ishii–Pires–Souganidis on the convergence of the Bence–Merriman–Osher algorithm for isotropic motion by mean curvature. The main difficulty here, is that the kernel  $K_{\phi,t}$  is not positive and that its moments of order 2 are not in  $L^1(\mathbb{R}^d)$ . Still, we can show that in one sense the scheme is consistent with the anisotropic mean curvature flow.

*2010 Mathematics Subject Classification:* Primary 53A99, 45L05, 35B40.

*Keywords:* Anisotropic mean curvature flow, BMO algorithm, phase field approximation

### 1. Introduction and motivation

In the last decades, a lot of attention has been devoted to the motion of interfaces, and particularly to motion by mean curvature. Applications concern image processing (denoising, segmentation), material sciences (motion of grain boundaries in alloys, crystal growth), biology (modelling of vesicles and blood cells). This paper is interested in numerical schemes for the anisotropic mean curvature flow, that is, the “gradient flow” of an anisotropic perimeter

$$P_\phi(\Omega) = \int_{\partial\Omega} \phi^\circ(n(x)) d\sigma \tag{1}$$

where  $n(x)$  is the outer normal to  $\partial E$  at  $x$  and  $\phi^\circ$  is a convex, one-homogeneous surface tension (the isotropic case corresponds to  $\phi^\circ(n(x)) = |n(x)| = 1$ ).

There is an important literature on numerical methods for the isotropic and anisotropic curvature flows. These can be roughly classified into three categories: Parametric methods [7, 8, 23, 24], Level

set formulations [19, 27, 35–37] or Phase field approaches [10, 17, 34, 38]. See for instance [24] for a complete review and comparison between these three different strategies.

In this work, we will consider a new scheme, proposed in [14], based on a phase field representation. It relies on the introduction of a specific anisotropic Laplacian (pseudo-differential) operator, which can be used both in a standard phase-field approximation (an anisotropic Allen–Cahn equation), or in a convolution/thresholding scheme [11, 32] which can be thought as a limiting case of the Allen–Cahn equation. The basic idea in [11] is to alternate the diffusion (with the heat equation) and the sharpening (by thresholding) of the characteristic functions of a set; [32] study a more general variant where the diffusion is replaced with the convolution by quite general kernels.

In the phase-field approach, anisotropic flows can be tackled either by a modified version of the Bence–Merriman–Osher algorithm [11], where the heat equation is replaced with a nonlinear variant built upon the anisotropy  $\phi^\circ$  [15], or by replacing the heat equation with the convolution with a nonnegative, nonsymmetric kernel  $f$  as in [32]. However, in the latter case, the inverse problem of finding an appropriate kernel  $f$ , given the anisotropy  $\phi^\circ$ , is solved only in 2D [41]. Some progress was done recently in relating the convolution kernel with  $\phi^\circ$  in [22], but the inverse problem is still considered untractable in higher dimension. In fact, it is not even clear that any anisotropy, even smooth, can be obtained in the framework of [32].

The aim of this work is to study a simple construction, proposed in [14] of a kernel  $f$  for all kind of anisotropy  $\phi^\circ$  in all dimension. This kernel can be seen as the fundamental solution of the heat equation with a particular pseudo-differential operator which can be seen as an anisotropic Laplacian. The most interesting feature of this approach is that the diffusion can be solved efficiently using the Fourier Transform, as proposed in [16] for the isotropic Allen–Cahn equation (and as can also be done for [32, 41]). A few numerical experiments with this approach have been already shown in [14] with smooth, but also crystalline or even non-convex anisotropies. Although the approach seems (numerically) to perform well in all these cases, a full justification is still missing. The essential contribution of this paper is to extend the consistency proof (see Theorem (2)) of Ishii–Pires–Souganidis in the case of our specific kernel, for smooth, uniformly elliptic anisotropies. The main issue is that in this case, the kernel does not satisfy the assumptions which are needed in [32]. In particular, it is not even nonnegative, so that our scheme is non-monotone and a complete proof of convergence is still missing.

In the next section, we introduce our notation and a precise framework. We give a short introduction to level set formulations, phase field approximations and the Bence–Merriman–Osher algorithm in the case of the isotropic and anisotropic mean curvature flow. Next, we introduce our anisotropic heat kernel and establish some of its properties. Our main consistency result (Theorem 2) is given in Section 4. We show the consistency of an anisotropic Bence–Merriman–Osher scheme built upon that kernel. The last section shows numerical evidence of the convergence of a slightly modified scheme, which corresponds to a splitting of the anisotropic Allen–Cahn equation (hence the thresholding is replaced with a reaction term which only enhances the slope of the diffuse interface). Computationally, the scheme proves very efficient and very fast, even when the anisotropy is not smooth.

In comparison with other existing methods, our approach can be easily implemented: Contrarily to methods based on a parametric representations, it does not require special care in handling topological changes or in the case of 3D computations. Besides, it avoids direct discretization of the non linear anisotropic Allen Cahn equation (e.g., by finite elements) [10, 38]. Indeed, in our

approach the non linear diffusion operator is replaced by an approximate linear operator, whose resolution can be easily performed by Fourier transform.

## 2. Preliminaries

### 2.1 Motion by isotropic mean curvature

The simplest case of motion by isotropic mean curvature concerns the evolution of a set  $\Omega_t \subset \mathbb{R}^d$  with a boundary  $\Gamma_t$  of codimension 1, whose normal velocity  $V_n$  is proportional to its mean curvature  $\kappa$

$$V_n(x) = \kappa(x), \quad \text{a.e. } x \in \Gamma_t, \quad (2)$$

with the convention that  $\kappa$  is negative if  $\Omega_t$  is a convex set. It at  $t = 0$  the initial set  $\Omega_0$  is smooth, then the evolution is well-defined until some time  $T > 0$  when singularities may develop [2].

Viscosity solutions provide a more general framework, that defines evolution past singularities, or evolution from non-smooth initial sets. If  $g$  is a level set function of  $\Omega_0$ , i.e.,

$$\Omega_0 = \{x \in \mathbb{R}^d ; g(x) \leq 0\}, \quad \Gamma_0 = \{x \in \mathbb{R}^d ; g(x) = 0\},$$

and if  $u$  denotes the viscosity solution to the geometric evolution equation

$$\begin{cases} u_t = \operatorname{div} \left( \frac{\nabla u}{|\nabla u|} \right) |\nabla u|, \\ u(0, x) = g(x), \end{cases}$$

then the generalized mean curvature flow  $\Omega_t$  starting from  $\Omega_0$  is defined by the 0-level set of  $u$  [20, 26, 27, 37]

$$\Omega_t = \{x \in \mathbb{R}^d ; u(t, x) \leq 0\}, \quad \partial\Gamma_t = \{x \in \mathbb{R}^d ; u(t, x) = 0\}.$$

Alternatively, one can define the motion by mean curvature as the limit of diffuse interface approximations obtained by solving the Allen–Cahn equation

$$\frac{\partial u}{\partial t} = \Delta u - \frac{1}{\epsilon^2} (W'(u)), \quad (3)$$

where  $\epsilon$  is a small parameter (that determines the width of the diffuse interface) and where  $W(s) = \frac{s^2(1-s)^2}{2}$  is a double well potential. This equation can be viewed as a gradient flow for the energy

$$J_\epsilon(u) = \int_{\mathbb{R}^d} \left( \frac{\epsilon}{2} |\nabla u|^2 + \frac{1}{\epsilon} W(u) \right) dx.$$

Modica and Mortola [33, 34] have shown that  $J_\epsilon$  approximates (in the sense of  $\Gamma$ -convergence) the surface energy  $c_W J$  where

$$J(\Omega) = \int_{\partial\Omega} 1 \, d\sigma \quad \text{and} \quad c_W = \int_0^1 \sqrt{2W(s)} \, ds.$$

Existence, uniqueness, and a comparison principle have been established for (3) (see, for example, Chapters 14 and 15 in [2] and the references therein).

Let  $u_\epsilon$  solve (3) with the initial condition

$$u_\epsilon(x, 0) = q\left(\frac{d(x, \Omega_0)}{\epsilon}\right),$$

where  $d(x, \Omega)$  denotes the signed distance of a point  $x$  to the set  $\Omega$  and where the profile  $q$  is defined by

$$q = \arg \min \left\{ \int_{\mathbb{R}} \left( \frac{1}{2} \gamma'^2 + W(\gamma) \right); \gamma \in H_{loc}^1(\mathbb{R}), \gamma(-\infty) = +1, \gamma(+\infty) = -1, \gamma(0) = \frac{1}{2} \right\}.$$

The set

$$\Omega_{t,\epsilon} = \left\{ x \in \mathbb{R}^d; u_\epsilon(x, t) \geq \frac{1}{2} \right\},$$

approximates  $\Omega(t)$  at the rate of convergence  $O(\epsilon^2 |\log \epsilon|^2)$  in the case of smooth motion by mean curvature [9, 17].

In the case of generalized motion by mean curvature, convergence has been shown [3, 26] provided that the interior of the set  $\Gamma_t$  remains empty (i.e., no fattening occurs).

About numerical point of view, convergence has been established for a finite element method in [39] and for a finite difference method in [18]. A splitting spectral Fourier method is also been addressed in [14, 16].

The Bence–Merriman–Osher algorithm [11] is yet another approximation to motion by mean curvature. Given a closed set  $E \subset \mathbb{R}^d$ , and denoting  $\chi_E$  its characteristic function, one defines

$$T_h E = \left\{ x \in \mathbb{R}^d; u(x, h) \geq \frac{1}{2} \right\},$$

where  $u$  solves the heat equation

$$\begin{cases} \frac{\partial u}{\partial t}(x, t) = \Delta u(x, t), & t > 0, \quad x \in \mathbb{R}^d, \\ u(x, 0) = \chi_E(x). \end{cases}$$

Setting  $E_h(t) = T^{[t/h]} E$ , where  $[t/h]$  is the integer part of  $t/h$ , Evans [25], and Barles and Georgelin [4] have shown that  $E_h(t)$  converges to  $E_t$ , the evolution by mean curvature from  $E$ . Remark also that this algorithm can be seen as a splitting algorithm for the Allen–Cahn equation in the limited case  $\epsilon \rightarrow 0$ . See also [41] where an efficient numerical resolution is presented.

## 2.2 Motion by anisotropic mean curvature

We use the framework of the Finsler geometry as described in [10]. Let  $\phi : \mathbb{R}^d \rightarrow [0, +\infty[$  denote a strictly convex function in  $C^2(\mathbb{R}^d \setminus \{0\})$ , which is 1-homogeneous and bounded, i.e.,

$$\begin{cases} \phi(t\xi) = |t|\phi(\xi) & \xi \in \mathbb{R}^d, t \in \mathbb{R}, \\ \lambda|\xi| \leq \phi(\xi) \leq \Lambda|\xi| & \xi \in \mathbb{R}^d, \end{cases}$$

for two positive constants  $0 < \lambda \leq \Lambda < +\infty$ . We assume that its dual function  $\phi^\circ : \mathbb{R}^d \rightarrow [0, +\infty[$ , defined by

$$\phi^\circ(\xi^*) = \sup \{ \xi^* \cdot \xi; \phi(\xi) \leq 1 \}$$

is also in  $C^2(\mathbb{R}^N \setminus \{0\})$ ). Given a smooth set  $E$  and a smooth function  $u : \mathbb{R}^d \rightarrow \mathbb{R}$  such that  $\partial E = \{x \in \mathbb{R}^d ; u(x) = 0\}$ , we define

- the Cahn–Hoffman vector field  $n_\phi = \phi_\xi^o(\nabla u)$ ,
- the  $\phi$ -curvature  $\kappa_\phi = \operatorname{div}(n_\phi)$ .

We say that  $E(t)$  is the evolution from  $E$  by  $\phi$ -curvature, if at each time  $t$ , the normal velocity  $V_n$  is given by

$$V_n = -\kappa_\phi n_\phi.$$

As in the case of isotropic flows, one can define motion by  $\phi$ -curvature using a level set formulation, i.e., following the level lines of the solution to the anisotropic evolution equation

$$u_t = \phi^o(\nabla u) \phi_{\xi\xi}^o(\nabla u) : \nabla^2 u. \quad (4)$$

Existence, uniqueness and a comparison principle have been established in [5, 6, 19, 21]. The anisotropic surface energy

$$J(\Omega) = \int_{\partial\Omega} \phi^o(n) d\sigma$$

can be approximated by the Ginzburg–Landau-like energy

$$J_{\epsilon,\phi}(u) = \int_{\mathbb{R}^d} \left( \frac{\epsilon}{2} \phi^o(\nabla u)^2 + \frac{1}{\epsilon} W(u) \right) dx,$$

and its gradient flow leads to the anisotropic Allen–Cahn equation [1]

$$\frac{\partial u}{\partial t} = \Delta_\phi u - \frac{1}{\epsilon^2} W'(u). \quad (5)$$

The operator  $\Delta_\phi := \operatorname{div} \left( \phi_\xi^o(\nabla u) \phi^o(\nabla u) \right)$  is called the anisotropic Laplacian.

This equation can be numerically solved by a semi finite elements method (see [38] for instance) but the complexity of this algorithm is much greater than in the isotropic case because it needs a resolution of a new linear system at each iteration in time.

The Bence–Merriman–Osher algorithm has also been extended to anisotropic motion by mean curvature. One generalization was proposed by Chambolle and Novaga [15] as follows: Given a closed set  $E$ , let  $T_h(E) = \{x \in \mathbb{R}^d ; u(x, h) \geq \frac{1}{2}\}$ , where  $u(x, t)$  is the solution to

$$\begin{cases} \frac{\partial u}{\partial t}(x, t) = \Delta_\phi u(x, t), & t > 0, \quad x \in \mathbb{R}^d, \\ u(x, 0) = \chi_E(x). \end{cases} \quad (6)$$

Define then  $E_h(t) = T_h^{[t/h]} E$ . The convergence of  $E_h(t)$  to the generalized anisotropic mean curvature flow from  $E$  is established in [15]. The result holds for very general anisotropic surface tensions and even in the crystalline case. However, as for the phase field formulation, because of the strongly nonlinear character of  $\Delta_\phi$ , the numerical resolution of (6) is much harder and not also efficient than in the isotropic case.

Another generalization of the Bence–Merriman–Osher algorithm has been studied by Ishii, Pires and Souganidis [32]. The main idea is to represent the solution  $u$  of (6) as the convolution of  $\chi_E$  with a geometric kernel. More precisely, Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function which satisfies the following conditions

(A<sub>1</sub>) Positivity and symmetry:

$$f(x) \geq 0, \quad f(-x) = f(x), \quad \text{and} \quad \int_{\mathbb{R}^d} f(x) dx = 1.$$

(A<sub>2</sub>) Boundedness of the moments:

$$\int_{\mathbb{R}^d} |x|^2 f(x) dx < +\infty,$$

$$0 < \int_{p^\perp} (1 + |x|^2) f(x) d\mathcal{H}^{d-1} < \infty, \quad \text{for all } p \in \mathbb{S}^{d-1}.$$

(A<sub>3</sub>) Smoothness:

$$p \rightarrow \int_{p^\perp} f(x) d\mathcal{H}^{d-1} \quad \text{and} \quad p \rightarrow \int_{p^\perp} x_i x_j f(x) d\mathcal{H}^{d-1} \quad \text{are continuous on } \mathbb{S}^{d-1}.$$

Here  $p^\perp$  denotes the orthogonal complement of the vector  $p$ , i.e.,  
 $p^\perp = \{x \in \mathbb{R}^d ; \langle x, p \rangle = 0\}$ .

Given  $E \subset \mathbb{R}^d$ , define  $T_h E = \{x \in \mathbb{R}^d ; u(x, h) \geq \frac{1}{2}\}$ , where

$$u(x, h) = \int_{\mathbb{R}^d} \tilde{K}_h(y) \chi_E(y - x) dy,$$

with the kernel

$$\tilde{K}_t(x) = \frac{1}{t^{d/2}} f(\sqrt{t}x), \quad x \in \mathbb{R}^d.$$

They showed [32] that  $T_h^{[t/h]} E$  converges to the set  $E(t)$  obtained from  $E$  as the generalized motion by anisotropic mean curvature via the geometric evolution equation

$$u_t = F(D^2u, Du),$$

where

$$F(X, p) = \left( \int_{p^\perp} f(x) d\mathcal{H}^{d-1}(x) \right)^{-1} \left( -\frac{1}{2} \int_{p^\perp} \langle Xx, x \rangle f(x) d\mathcal{H}^{d-1}(x) \right).$$

This algorithm appears to be more efficient than the last one (with the non linear operator  $\Delta_\phi$ ), but raises a natural question: Given an anisotropy  $\phi^\rho$ , can one find a kernel  $f$ , so that the generalized front  $\partial E(t)$  defined by the associated evolution equation evolves by  $\phi$ -mean curvature? This problem has been addressed by Ruuth and Merriman [42] in dimension 2. They propose a class of kernels and study the corresponding numerical schemes, which prove very efficient. However, their approach cannot be generalized to higher dimensions. In contrast, our algorithm is not specific to the dimension 2.

### 2.3 A new algorithm for motion by anisotropic mean curvature

In this work, our objective is to extend Ishii–Pires–Souganidis' analysis to study the following algorithm. Starting from a bounded closed set  $E \subset \mathbb{R}^d$ , we define an operator  $T_h E$  by

$$T_h E = \left\{ x \in \mathbb{R}^d ; u(x, h) \geq \frac{1}{2} \right\}, \quad (7)$$

where  $u$  solves the following parabolic equation:

$$\begin{cases} \frac{\partial u}{\partial t}(x, t) = \tilde{\Delta}_\phi u(x, t), & t > 0, \quad x \in \mathbb{R}^d, \\ u(x, 0) = \chi_E(x). \end{cases} \quad (8)$$

Denoting by  $\mathcal{F}(u)$  the Fourier transform of a function  $u$ ,

$$\mathcal{F}(u)(\xi) = \int_{\mathbb{R}^d} u(x) e^{-2\pi i x \cdot \xi} dx,$$

the operator  $\tilde{\Delta}_\phi$  is defined by

$$\tilde{\Delta}_\phi u = \mathcal{F}^{-1} \left( -4\pi^2 \phi^o(\xi)^2 \mathcal{F}(u)(\xi) \right).$$

and can be seen as a linearization of  $\Delta_\phi$  in the Fourier space. The solution  $u$  of (8) can be expressed as a convolution product of the characteristic function of  $E$  and of the anisotropic kernel

$$K_{\phi, t}(x) = \mathcal{F}^{-1} \left( e^{-4\pi^2 t \phi^o(\xi)^2} \right) (x).$$

However, this kernel (more precisely  $K_{\phi, t=1}$ ) does not satisfy the hypotheses  $(A_1)$  and  $(A_2)$  above:  $K_{\phi, 1}$  is not positive and  $x \rightarrow \int_{\mathbb{R}^d} |x|^2 K_\phi(x)$  is not in  $L^1(\mathbb{R})$ . But we will show that the associated Hamiltonian flow is

$$\begin{aligned} F(X, p) &= \left( \int_{p^\perp} K_\phi d\mathcal{H}^{d-1} \right)^{-1} \left( \frac{1}{2} \int_{p^\perp} \langle Xx, x \rangle K_\phi(x) d\mathcal{H}^{d-1} \right) \\ &= \phi^o(p) \phi_{\xi\xi}^o(p) : X, \end{aligned}$$

which establishes a link between  $K_\phi$  and  $\phi$ -anisotropic mean curvature flow.

### 3. The operator $\tilde{\Delta}_\phi$ and properties of the anisotropic kernel $K_\phi$

Let  $\phi = \phi(\xi)$  denote a strictly convex smooth Finsler metric and let  $\phi^o$  denote its dual (see [10]). We assume that  $\phi^o$  is a 1-homogenous, symmetric function in  $C^\infty(\mathbb{R}^d \setminus \{0\})$  that satisfies

$$\lambda|\xi| \leq \phi^o(\xi) \leq \Lambda|\xi|. \quad (9)$$

In particular, it follows that for any  $\xi \in \mathbb{R}^d$  and  $t \in \mathbb{R}$ ,

$$\begin{cases} \phi^o(t\xi) = |t| \phi^o(\xi), \\ \phi_\xi^o(t\xi) = \frac{t}{|t|} \phi_\xi^o(\xi), \\ \phi_\xi^o(\xi) \cdot \xi = \phi^o(\xi). \end{cases}$$

The associated anisotropic mean curvature is defined as the anisotropic Laplacian operator

$$\Delta_\phi u = \operatorname{div} \left( \phi^o(\nabla u) \phi_\xi^o(\nabla u) \right), \quad \forall u \in H^2(\Omega).$$

A direct computation shows that for any  $\xi \in \mathbb{R}^d$ ,

$$\begin{cases} \Delta_\phi [\cos(2\pi\xi \cdot x)] = -4\pi^2 \phi^o(\xi)^2 \cos(2\pi\xi \cdot x), \\ \Delta_\phi [\sin(2\pi\xi \cdot x)] = -4\pi^2 \phi^o(\xi)^2 \sin(2\pi\xi \cdot x), \end{cases}$$

i.e., that plane waves are eigenfunctions of the anisotropic Laplacian (albeit nonlinear). We define  $\tilde{\Delta}_\phi : H^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$  by

$$\tilde{\Delta}_\phi u = \mathcal{F}^{-1} \left[ -4\pi^2 \phi^o(\xi)^2 \mathcal{F}[u](\xi) \right].$$

Given an initial condition  $u_0 \in L^2(\mathbb{R}^d)$ , we study the solution  $u$  of

$$\begin{cases} u_t(t, x) = \tilde{\Delta}_\phi u(t, x), \\ u(0, x) = u_0. \end{cases}$$

The function  $u$  can also be expressed as the convolution product  $u = K_{\phi,t} * u_0$ , where the anisotropic heat kernel  $K_{\phi,t}$  is defined by

$$K_{\phi,t} = \mathcal{F}^{-1} \left[ e^{-4\pi^2 t \phi^o(\xi)^2} \right].$$

We also set  $K_\phi = K_{\phi,1}$ . In the rest of this section, we establish some properties of this operator.

**PROPOSITION 1 (Regularity of  $\hat{K}_\phi$ )** The function  $\hat{K}_\phi : \xi \rightarrow e^{-4\pi^2 \phi^o(\xi)^2}$  is in  $W^{d+1,1}(\mathbb{R}^d)$ , and  $D^{d+2} \hat{K}_\phi$  is a function.

*Proof.* First, we notice that

$$D \hat{K}_\phi(\xi) = -8\pi^2 \phi_\xi^o(\xi) \phi^o(\xi) e^{-4\pi^2 \phi^o(\xi)^2},$$

and

$$\begin{aligned} D^2 \hat{K}_\phi(\xi) &= 64\pi^4 \phi^o(\xi)^2 \left( \phi_\xi^o(\xi) \otimes \phi_\xi^o(\xi) \right) e^{-4\pi^2 \phi^o(\xi)^2} \\ &\quad - 8\pi^2 \left( \phi^o(\xi) \phi_{\xi\xi}^o(\xi) + \phi_\xi^o(\xi) \otimes \phi_\xi^o(\xi) \right) e^{-4\pi^2 \phi^o(\xi)^2}. \end{aligned}$$

We note that  $\phi_\xi^o$  is discontinuous at  $\xi = 0$ . Nevertheless, we next prove that the  $d - 1^{th}$  derivative of  $D^2 \hat{K}_\phi$  belongs to  $L^1(\mathbb{R}^d)^{d^2}$ . Assume that  $f = D^{n+2} \hat{K}_\phi$  is an integrable function on  $\mathbb{R}^d$  for some integer  $n < d$ . The homogeneity of  $\phi^o$  shows the existence of a constant  $C_n$  such that

$$|D^{n+2} \hat{K}_\phi| \leq C_n \frac{1}{|\xi|^n} e^{-\lambda|\xi|^2}, \quad \text{for all } \xi \in \mathbb{R}^d \setminus \{0\}.$$

Since  $f$  is smooth away from  $\xi = 0$ , the distributional derivative of  $f$  is the sum of a function and of possibly a Dirac mass at  $\xi = 0$ :

$$Df = \{\nabla f\} + c \delta,$$



where  $c$  is a constant and  $\nabla f$  denotes the pointwise derivative of  $f$ . Let  $\varphi \in \mathfrak{D}(\mathbb{R}^d)^{d^{n+2}}$  and let  $\epsilon > 0$ . Then

$$\begin{aligned} \langle Df, \varphi \rangle &= -\langle f, \operatorname{div} \varphi \rangle = -\int_{\mathbb{R}^d} f \cdot \operatorname{div} \varphi \, dx \\ &= -\int_{\mathbb{R}^d \setminus B(0, \epsilon)} f \cdot \operatorname{div} \varphi \, dx - \int_{B(0, \epsilon)} f \cdot \operatorname{div} \varphi \, dx \\ &= \int_{\mathbb{R}^d \setminus B(0, \epsilon)} \nabla f \cdot \varphi \, dx - \int_{\partial B(0, \epsilon)} f \cdot (\varphi \cdot \vec{n}) \, d\sigma - \int_{B(0, \epsilon)} f \cdot \operatorname{div} \varphi \, dx. \end{aligned}$$

Since we assumed that  $f \in L^1(\mathbb{R}^d)^{d^{n+2}}$ , the last integral above tends to 0, as  $\epsilon \rightarrow 0$ . Moreover as  $n < d$ , we have

$$\begin{aligned} \left| \int_{\partial B(0, \epsilon)} f \cdot \varphi \cdot \vec{n} \, d\sigma \right| &\leq \|\varphi\|_{L^\infty} \int_{\partial B(0, \epsilon)} C_n \frac{1}{|\xi|^n} e^{-\lambda|\xi|^2} \, d\sigma \\ &\leq \|\varphi\|_{L^\infty} C_n \int_{\partial B(0, \epsilon)} \epsilon^{-n} \, d\sigma \leq C_n \|\varphi\|_{L^\infty} \epsilon^{d-1-n}, \end{aligned}$$

so that

$$\lim_{\epsilon \rightarrow 0} \left| \int_{\partial B(0, \epsilon)} f \cdot \varphi \cdot \vec{n} \, d\sigma \right| = 0.$$

It follows that  $c = 0$ , which concludes the proof.  $\square$

**PROPOSITION 2** (Decay properties of  $K_\phi$ ) Let  $s \in [0, 1]$ . There exists a constant  $C_{\phi^o, s}$ , which only depends on the anisotropy  $\phi^o$  and on  $s$ , such that

$$|K_\phi(x)| \leq \frac{C_{\phi^o, s}}{1 + |x|^{d+1+s}}, \quad \forall x \in \mathbb{R}^d. \quad (10)$$

**REMARK 1** The case  $s = 0$  is easy: According to Proposition 1, the function  $\Delta^{\frac{d+1}{2}} \hat{K}_\phi(\xi)$  is in  $L^1(\mathbb{R}^d)$ . The continuity of the Fourier transform from  $L^1$  to  $L^\infty$  shows that

$$\|(1 + |x|^{d+1})K_\phi\|_{L^\infty} \leq C \|\hat{K}_\phi(\xi) + \Delta^{\frac{d+1}{2}} \hat{K}_\phi(\xi)\|_{L^1(\mathbb{R}^d)},$$

and since  $\hat{K}_\phi(\xi) = e^{-4\pi^2 \phi^o(\xi)^2}$ ,

$$|K_\phi(x)| \leq \frac{C_{\phi^o, 0}}{1 + |x|^{d+1}}, \quad \forall x \in \mathbb{R}^d.$$

The proof uses properties of interpolation spaces [12]. Consider  $X, Y$  two Banach spaces, and for  $u \in X + Y$  and  $t \in \mathbb{R}^+$ , let

$$k(t, u) = \inf_{u=u_0+u_1} \{\|u_0\|_X + t\|u_1\|_Y\}.$$

For  $s \in [0, 1]$  and  $p \geq 1$ , the interpolation space  $[X, Y]_{s, p}$  between  $X$  and  $Y$  is defined by

$$[X, Y]_{s, p} = \left\{ u \in X + Y ; t^{-s} K(t, u) \in L^p \left( \frac{1}{t} \right) \right\}.$$

In particular, given a strictly positive function  $h : \mathbb{R}^d \rightarrow \mathbb{R}$ , consider the weighted space  $L_h^\infty$  defined by

$$L_h^\infty(\mathbb{R}^d) = \{u \in L^\infty(\mathbb{R}^d); \sup_{x \in \mathbb{R}^d} \{h(x)u(x)\} < \infty\}.$$

One can interpolate between  $L^\infty$  and  $L_h^\infty$  according to the following lemma.

LEMMA 1 Let  $h$  be a strictly positive function  $\mathbb{R}^d \rightarrow \mathbb{R}$ , and let  $s \in ]0, 1[$ . Then

$$[L^\infty(\mathbb{R}^d), L_h^\infty(\mathbb{R}^d)]_{s, \infty} = L_{h^s}^\infty(\mathbb{R}^d).$$

*Proof.* 1) Assume that  $u \in L_{h^s}^\infty(\mathbb{R}^d)$ . There exists a constant  $C$  such that for a.e.  $x \in \mathbb{R}^d$ ,

$$|u(x)| \leq \frac{C}{h(x)^s}. \quad (11)$$

To estimate  $k(t, u) = \inf_{u=u_0+u_1} \{\|u_0\|_{L^\infty} + t\|u_1\|_{L_h^\infty}\}$ , we note that

- If  $t \geq 1$ , the choice  $u_0 = u$  and  $u_1 = 0$ , shows that  $K(t, u) \leq \|u\|_{L^\infty}$ .
- If  $t < 1$ , we consider the set  $A = \{x \in \mathbb{R}^d; |u(x)|h(x) \leq t^{s-1}\}$ , and we choose  $u_0 = \chi_{A^c} u$  and  $u_1 = \chi_A u$ , so that  $\|u_1\|_{L_h^\infty} \leq t^{s-1}$ . Moreover, we remark that for all  $x \in A^c$ ,  $|u(x)|h(x) \geq t^{s-1}$  so that, in view of (11),

$$|u_0(x)| \leq Ch(x)^{-s} \leq C|u_0(x)|^s t^{s(1-s)},$$

and thus  $k(t, u) \leq (C + 1)t^s$ .

In summary, these estimates show that

$$K(t, u) \leq \begin{cases} \|u\|_{L^\infty} & \text{if } t \geq 1, \\ (C + 1)t^s & \text{if } t < 1, \end{cases}$$

which proves that  $u \in [L^\infty, L_h^\infty]_{s, \infty}$ .

2) Conversely, we consider  $u \in [L^\infty, L_h^\infty]_{s, \infty}$ . For all  $t > 0$ , there exists a decomposition  $u = u_{0,t} + u_{1,t}$  such that

$$|u_{0,t}|_{L^\infty} + t|u_{1,t}|_{L_h^\infty} \leq Ct^s.$$

It follows that for all  $t > 0$ , we have

$$h(x)^s |u(x)| \leq |h(x)^s |u_{0,t}(x) + u_{1,t}(x)| \leq C (h(x)^s t^s + h(x)^{s-1} t^{s-1}).$$

Choosing  $t = h(x)^{-1}$  in the above inequality shows that for all  $x \in \mathbb{R}^d$ ,  $h(x)^s |u(x)| \leq 2C$ , which concludes the proof.  $\square$

We use the following properties of interpolation spaces:

- (P<sub>1</sub>) if  $T$  is continuous from  $X \rightarrow \tilde{X}$  and from  $Y \rightarrow \tilde{Y}$ , then  $T$  is continuous from  $[X, Y]_{s, p}$  to  $[\tilde{X}, \tilde{Y}]_{s, p}$ .
- (P<sub>2</sub>) if  $p < p'$ , then  $[X, Y]_{s, p} \subset [X, Y]_{s, p'}$  for any  $0 < s < 1$  and  $p \geq 1$ .
- (P<sub>3</sub>)  $[L^\infty(\mathbb{R}^d), L_{(1+|\cdot|)}^\infty(\mathbb{R}^d)]_{s, \infty} = L_{(1+|\cdot|)^s}^\infty(\mathbb{R}^d)$  for any  $0 < s < 1$ .

In the following, we consider the case where  $T$  is the Fourier transform,  $X = L^1(\mathbb{R}^d)$ ,  $Y = L^\infty(\mathbb{R}^d)$ ,  $\tilde{X} = W^{1,1}(\mathbb{R}^d)$  and  $\tilde{Y} = L^\infty_{(1+|x|)}(\mathbb{R}^d)$ .

*Proof of Proposition 2.* We claim that it suffices to show that for any  $0 < s < 1$

$$u(\xi) := \Delta^{\frac{d+1}{2}} \hat{K}_\phi(\xi) \in [X, Y]_{s,1}. \quad (12)$$

Indeed, the inclusion  $[X, Y]_{s,1} \subset [X, Y]_{s,\infty}$  implies then that  $u \in [X, Y]_{s,\infty}$ , so that in view of  $(P_1)$  and  $(P_3)$  we obtain

$$\hat{u} \in [\tilde{X}, \tilde{Y}]_{s,\infty} = [L^\infty(\mathbb{R}^d), L^\infty_{(1+|x|)}(\mathbb{R}^d)]_{s,\infty} = L^\infty_{(1+|x|)^s}(\mathbb{R}^d),$$

and consequently

$$|(1+|x|^s)\hat{u}(x)| = |(1+|x|^{d+1})K_\phi(x)(1+|x|^s)| \leq C_{\phi^o,s}, \quad \text{for all } x \in \mathbb{R}^d.$$

It follows that for some constant  $C_{\phi^o,s}$

$$|K_\phi(x)| \leq \frac{C_{\phi^o,s}}{1+|x|^{d+1+s}}, \quad \text{for all } x \in \mathbb{R}^d.$$

We now prove (12). The homogeneity of  $\phi^o$  shows that for some  $c_1 > 0$  and  $c_2 > 0$ , and for  $\xi \in \mathbb{R}^d \setminus \{0\}$ ,

$$|u(\xi)| \leq \frac{c_1}{|\xi|^{d-1}} e^{-\lambda|\xi|^2} \quad \text{and} \quad |\nabla u(\xi)| \leq \frac{c_2}{|\xi|^d} e^{-\lambda|\xi|^2},$$

which shows that  $u \in X = L^1(\mathbb{R}^d)$ . However,  $u$  may not belong to  $Y = L^\infty(\mathbb{R}^d)$ . We now estimate  $k(u, t)$ , for  $t \in \mathbb{R}^+$ . If  $t \geq 1$ , we set  $u_0 = u$ ,  $u_1 = 0$ , so that

$$k(t, u) \leq \|u\|_X, \quad \forall t \geq 1. \quad (13)$$

If  $t < 1$ , consider the function  $\rho_t(\xi)$  defined by

$$\rho_t(\xi) = \begin{cases} 0 & \text{if } |x| \leq t, \\ 1 & \text{if } |x| > 2t, \\ \sin\left(\frac{\pi}{2} \frac{|\xi|-t}{t}\right) & \text{otherwise.} \end{cases}$$

We choose  $u_0 = (1 - \rho_t)u$  and  $u_1 = \rho_t u$  and check that

$$\|u_0\|_{L^1(\mathbb{R}^d)} \leq \int_{B(0,2t)} |u(\xi)| d\xi \leq \int_{B(0,2t)} \frac{C}{|\xi|^{d-1}} d\xi \leq 2C |\mathbf{S}^d| t.$$

Moreover,

$$\begin{aligned} \|\nabla u_1\|_{L^1(\mathbb{R}^d)} &\leq \|\nabla \rho_t u + \rho_t \nabla u\|_{L^1(\mathbb{R}^d)} \\ &\leq \int_{\mathbb{R}^d \setminus B(0,t)} |\nabla \rho_t| |u(\xi)| d\xi + \int_{\mathbb{R}^d \setminus B(0,t)} |\nabla u(\xi)| d\xi \\ &\leq \frac{\pi}{2t} \int_{B(0,2t) \setminus B(0,t)} \frac{C}{|\xi|^{d-1}} e^{-\lambda|\xi|^2} d\xi + \int_{\mathbb{R}^d \setminus B(0,t)} \frac{C}{|\xi|^d} e^{-\lambda|\xi|^2} d\xi. \end{aligned}$$

First, we have

$$\frac{\pi}{2t} \int_{B(0,2t) \setminus B(0,t)} \frac{C}{|\xi|^{d-1}} e^{-\lambda|\xi|^2} d\xi \leq \frac{C\pi}{2t} |\mathbf{S}^d| \int_t^{2t} dr \leq \frac{|\mathbf{S}^d| C \pi}{2}.$$

Second,

$$\begin{aligned} \int_{\mathbb{R}^d \setminus B(0,t)} \frac{C}{|\xi|^d} e^{-\lambda|\xi|^2} d\xi &\leq \int_{B(0,1) \setminus B(0,t)} \frac{C}{|\xi|^d} e^{-\lambda|\xi|^2} d\xi + \int_{\mathbb{R}^d \setminus B(0,1)} \frac{C}{|\xi|^d} e^{-\lambda|\xi|^2} d\xi \\ &\leq C |\mathbf{S}^d| \int_t^1 \frac{1}{r} dr + C |\mathbf{S}^d| \int_1^\infty e^{-\lambda r^2} dr \\ &\leq C |\mathbf{S}^d| \left( |\ln(t)| + \frac{1}{\sqrt{\lambda}} \frac{\sqrt{\pi}}{2} \right), \end{aligned}$$

so that

$$\|u_1\|_Y \leq C \left[ |\mathbf{S}^d| \left( \frac{\pi}{2} + \frac{1}{\sqrt{\lambda}} \frac{\sqrt{\pi}}{2} + |\ln(t)| \right) \right].$$

Consequently, this decomposition of  $u$  shows that

$$k(u, t) \leq C(1 + |\ln(t)|)t, \quad \forall t < 1, \quad (14)$$

for some constant  $C > 0$ . In summary,

$$k(u, t) \leq \begin{cases} \|u\|_X & \text{if } t \geq 1, \\ C(1 + |\ln(t)|)t & \text{if } t < 1, \end{cases}$$

and therefore we obtain

$$\begin{aligned} \|t^{-s} k(t, u)\|_{L^1(1/t)}^1 &= \int_{\mathbb{R}^+} |k(t, u) t^{-s}| \frac{1}{t} dt \\ &\leq \int_0^1 \frac{(C_0 + C_1 |\ln(t)|)}{t^s} dt + \int_1^\infty \frac{\|u\|_X^1}{t^{1+s}} dt < +\infty, \end{aligned}$$

which proves that  $u \in [X, Y]_{s,1}$  as claimed.  $\square$

**COROLLARY 1** For any  $s \in [0, 1[$  and  $p \in \mathbf{S}^d$ ,

$$|x|^{1+s} K_\phi \in L^1(\mathbb{R}^d), \quad (K_\phi)|_{p^\perp} \in L^1(\mathbb{R}^{d-1}), \quad (x \otimes x K_\phi)|_{p^\perp} \in L^1(\mathbb{R}^{d-1}).$$

**PROPOSITION 3** (Decay of averages of  $K_\phi$  on spheres) The integral

$$I(R) = \int_{\partial B(0,R)} K_\phi d\mathcal{H}^{d-1}$$

is strictly positive, and decays rapidly as

$$\frac{R^{d-1} |\mathbf{S}^{d-1}|}{(4\pi)^{d/2} \Lambda^d} e^{-\frac{R^2}{4\Lambda^2}} \leq I(R) \leq \frac{R^{d-1} |\mathbf{S}^{d-1}|}{(4\pi)^{d/2} \lambda^d} e^{-\frac{R^2}{4\lambda^2}},$$

where  $\lambda$  and  $\Lambda$  are bounds for  $\phi^\circ$  as in (9).

*Proof.* Since the measure  $\mu := \delta_{\partial B(0,R)}$  has finite mass, its Fourier transform is the continuous and bounded function

$$\hat{\mu}(\xi) = \int_{\mathbb{R}^d} e^{-2\pi i x \cdot \xi} d\mu = \int_{\partial B(0,R)} e^{-2\pi i x \cdot \xi}.$$

As  $\mu$  is radially symmetric,  $\hat{\mu}$  can be expressed in the form

$$\hat{\mu}(\xi) = R^{d-1} J(R|\xi|),$$

where  $J$  is a function  $\mathbb{R}^+ \rightarrow \mathbb{R}$ . It follows that

$$\begin{aligned} I(R) &= \langle \delta_{\partial B(0,R)}, K_\phi \rangle = \left\langle R^{d-1} J(R|\xi|), e^{-4\pi^2 \phi^o(\xi)^2} \right\rangle \\ &= R^{d-1} \int_{\mathbb{S}^{d-1}} \int_0^{+\infty} r^{d-1} J(Rr) e^{-4\pi^2 \phi^o(\theta)^2 r^2} dr d\mathcal{H}^{d-1}. \end{aligned} \quad (15)$$

We use the particular case when  $\phi^o(\xi)$  is isotropic, i.e.,  $\phi^o(\xi) = |\xi|$  to estimate the previous integral. In this case,  $K_\phi = \frac{1}{(4\pi)^{d/2}} e^{-\frac{x^2}{4}}$  is the heat kernel, and by a direct calculation we see that the corresponding integral is  $I(R) = \langle \delta_{B(0,R)}, K_\phi \rangle = \frac{R^{d-1} |\mathbb{S}^{d-1}|}{(4\pi)^{d/2}} e^{-\frac{R^2}{4}}$ . Comparing this expression to (15) and using the radial symmetry of  $K_\phi$  shows that

$$\int_0^{+\infty} r^{d-1} J(Rr) e^{-4\pi^2 r^2} dr = \frac{1}{(4\pi)^{d/2}} e^{-\frac{R^2}{4}},$$

or, after a change of variable, that

$$\int_0^{+\infty} r J(Rr) e^{-4\pi^2 \phi^o(\theta)^2 r^2} dr = \frac{1}{(4\pi)^{d/2} \phi^o(\theta)^d} e^{-\frac{R^2}{4\phi^o(\theta)^2}}. \quad (16)$$

Returning to a general kernel  $K_\phi$ , we deduce from (15) and (16) that

$$I(R) = \frac{R^{d-1}}{(4\pi)^{d/2}} \int_{\mathbb{S}^{d-1}} \frac{1}{\phi^o(\theta)^d} e^{-\frac{R^2}{4\phi^o(\theta)^2}} d\mathcal{H}^{d-1},$$

which in view of (9) concludes the proof.  $\square$

**PROPOSITION 4 (Positivity on hyperplanes)** For all  $p \in \mathbb{S}^{d-1}$ , the integral  $\int_{p^\perp} K_\phi d\mathcal{H}^{d-1}$  is well defined, and satisfies

$$\int_{p^\perp} K_\phi d\mathcal{H}^{d-1} = \frac{1}{2\sqrt{\pi} \phi^o(p)}.$$

In particular, we have

$$\frac{1}{2\sqrt{\pi} \Lambda} \leq \int_{p^\perp} K_\phi d\mathcal{H}^{d-1} \leq \frac{1}{2\sqrt{\pi} \lambda}.$$

*Proof.* Let  $p \in \mathbb{S}^{d-1}$ . We already know from Corollary 1 that  $\int_{p^\perp} K_\phi d\mathcal{H}^{d-1}$  is well defined. Consider for  $\mu > 0$ , the approximating functions  $f_\mu$ , defined by

$$\begin{cases} f_\mu(x) = K_\phi(x) e^{-\pi|x|^2/\mu^2}, \\ \hat{f}_\mu(\xi) = e^{-4\pi^2 \phi^o(\xi)^2} * \frac{1}{\mu^2} e^{-\pi\mu^2|\xi|^2}. \end{cases}$$

The function  $f_\mu$  belongs to the Schwartz space  $S(\mathbb{R}^d)$ . Moreover,  $\hat{f}_\mu \rightarrow \hat{K}_\phi$  in  $W^{d-1,1}(\mathbb{R}^d)$ , and the trace theorem [31] shows that one also has

$$\lim_{\mu \rightarrow \infty} \int_{\mathbb{R}} \hat{f}_\mu(sp) ds = \int_{\mathbb{R}} \hat{K}_\phi(sp) ds. \quad (17)$$

On the other hand, it follows from the Lebesgue dominated convergence theorem and from (10) that

$$\lim_{\mu \rightarrow \infty} \int_{p^\perp} f_\mu d\mathcal{H}^{d-1} = \int_{p^\perp} K_\phi d\mathcal{H}^{d-1}. \quad (18)$$

As  $f_\mu \in S(\mathbb{R}^d)$ , we infer that

$$\int_{p^\perp} f_\mu d\mathcal{H}^{d-1} = \langle \delta_{p^\perp}, f_\mu \rangle = \langle \delta_p, \mathfrak{F}[f_\mu] \rangle = \int_{\mathbb{R}} \hat{f}_\mu(sp) ds,$$

so that (17) and (18) yield

$$\begin{aligned} \int_{p^\perp} K_\phi d\mathcal{H}^{d-1} &= \int_{\mathbb{R}} \hat{K}_\phi(sp) ds = \int_{\mathbb{R}} e^{-4\pi^2 s^2 \phi^o(p)^2} ds \\ &= \int_{\mathbb{R}} e^{-\pi(2\sqrt{\pi}\phi^o(p)s)^2} ds = \frac{1}{2\sqrt{\pi}\phi^o(p)}, \end{aligned}$$

which concludes the proof.  $\square$

**PROPOSITION 5 (Moments of order 2)** Let  $p \in \mathbb{S}^{d-1}$ . Then  $\frac{1}{2} \int_{p^\perp} x \otimes x K_\phi d\mathcal{H}^{d-1}$  is well defined and satisfies

$$\frac{1}{2} \int_{p^\perp} x \otimes x K_\phi d\mathcal{H}^{d-1} = \phi_{\xi\xi}^o(p) \frac{1}{2\sqrt{\pi}}.$$

*Proof.* Corollary 1 states that the integral  $\int_{p^\perp} |x|^2 K_\phi d\mathcal{H}^{d-1}$  is well defined. Recalling the sequence  $f_\mu$  used in the previous proposition, we observe that  $D^2 \hat{f}_\mu \rightarrow D^2 \hat{K}_\phi$  in  $W^{d-1,1}(\mathbb{R}^d)$ , so that the trace theorem implies

$$\lim_{\mu \rightarrow \infty} \int_{\mathbb{R}} D^2 \hat{f}_\mu(sp) ds = \int_{\mathbb{R}} D^2 \hat{K}_\phi(sp) ds. \quad (19)$$

From Proposition 2 and the Lebesgue dominated convergence, we obtain

$$\lim_{\mu \rightarrow \infty} \int_{p^\perp} x \otimes x f_\mu(x) d\mathcal{H}^{d-1} \rightarrow \int_{p^\perp} x \otimes x K_\phi(x) d\mathcal{H}^{d-1}. \quad (20)$$

Moreover, we have

$$\begin{aligned} \int_{p^\perp} x \otimes x f_\mu(x) d\mathcal{H}^{d-1} &= \langle \delta_{p^\perp}, x \otimes x f_\mu \rangle = -\frac{1}{4\pi^2} \langle \delta_p, D^2 \hat{f}_\mu \rangle \\ &= -\frac{1}{4\pi^2} \int_{\mathbb{R}} D^2 \hat{f}_\mu(sp) ds, \end{aligned}$$

so that in view of (19)

$$\int_{p^\perp} x \otimes x K_\phi(x) d\mathcal{H}^{d-1} = -\frac{1}{4\pi^2} \int_{\mathbb{R}} D^2 \hat{K}_\phi(sp) ds.$$

We next estimate the above right-hand side by a direct calculation:

$$\begin{aligned} -\frac{1}{4\pi^2} \int_{\mathbb{R}} D^2 \hat{K}_\phi(sp) ds &= \left[ 2\phi^o(p)\phi_{\xi\xi}^o(p) + 2\phi_\xi^o(p) \otimes \phi_\xi^o(p) \right] \int_{\mathbb{R}} e^{-4\pi^2 s^2 \phi^o(p)^2} ds \\ &\quad - \left[ 2\phi_\xi^o(p) \otimes \phi_\xi^o(p) \right] \int_{\mathbb{R}} 8\pi^2 s^2 \phi^o(p)^2 e^{-4\pi^2 s^2 \phi^o(p)^2} ds. \end{aligned}$$

Further, we see by integration by parts that

$$\begin{aligned} \int_{\mathbb{R}} 8\pi^2 s^2 \phi^o(p)^2 e^{-4\pi^2 s^2 \phi^o(p)^2} ds &= \int_{\mathbb{R}} \left\{ 4\pi^2 2s \phi^o(p)^2 e^{-4\pi^2 s^2 \phi^o(p)^2} \right\} \{s\} ds \\ &= \int_{\mathbb{R}} e^{-4\pi^2 s^2 \phi^o(p)^2} ds = \frac{1}{2\sqrt{\pi} \phi^o(p)}, \end{aligned}$$

and we conclude that

$$\frac{1}{2} \int_{p^\perp} x \otimes x K_\phi(x) d\mathcal{H}^{d-1} = \phi_{\xi\xi}^o(p) \frac{1}{2\sqrt{\pi}}.$$

□

COROLLARY 2 (The operator  $F(X, p)$ ) Given  $X \in \mathbb{R}^{d \times d}$  and  $p \in \mathbb{S}^{d-1}$ , let

$$F(X, p) = \left( \int_{p^\perp} K_\phi(x) d\mathcal{H}^{d-1} \right)^{-1} \left( \frac{1}{2} \int_{p^\perp} \langle Xx, x \rangle K_\phi(x) d\mathcal{H}^{d-1} \right). \quad (21)$$

This operator is elliptic and satisfies

$$F(X, p) = \phi^o(p)\phi_{\xi\xi}^o(p) : X. \quad (22)$$

*Proof.* Equation (22) is a direct consequence of Propositions 4 and 5, while the ellipticity of  $F$  follows from the convexity of  $\phi^o$ . □

REMARK 2 In the next section, we introduce an algorithm for motion by anisotropic mean curvature, and show its consistency with an evolution equation of the form  $u_t = -F(D^2u, \frac{\nabla u}{|\nabla u|})$ , where  $F$  is defined by (21). The expression (22) shows that this operator is precisely the one corresponding to motion by anisotropic mean curvature (see [10]).

PROPOSITION 6 (Positivity of order moment  $s$ ) Let  $V$  be a subspace of  $\mathbb{R}^d$  of dimension  $1 \leq m \leq d$ , and let  $0 < s < 2$ . Then

$$\int_V |x|^s K_\phi d\mathcal{H}^m > 0.$$

*Proof.* We first consider the case  $m = d$  and  $V = \mathbb{R}^d$ . we consider the finite part  $Pf \left( \frac{1}{|x|^{d+s}} \right)$  as a temperate distribution, defined for  $\varphi \in S(\mathbb{R}^d)$  by

$$\left\langle Pf \left( \frac{1}{|x|^{d+s}} \right), \varphi \right\rangle = \lim_{\epsilon \rightarrow 0} \left\{ \int_{\mathbb{R}^d \setminus B(0, \epsilon)} \frac{\varphi(x) - \varphi(0)}{|x|^{d+s}} dx \right\}.$$

This function happens to be the Fourier transform of the distribution  $|x|^s$ . More precisely,

$$\mathcal{F}[|x|^s] = C_{s,d} Pf \left( \frac{1}{|2\pi\xi|^{d+s}} \right), \quad \text{with } C_{s,d} = 2^{s+d} \pi^{d/2} \frac{\Gamma((s+d)/2)}{\Gamma(-s/2)} \quad (23)$$

(see for instance [30],  $\Gamma$  denotes the Gamma function). We can thus write

$$\int_{\mathbb{R}^d} |x|^s K_\phi dx = \langle |x|^s, K_\phi \rangle = \left\langle C_{s,d} Pf \left( \frac{1}{|2\pi\xi|^{d+s}} \right), e^{-4\pi^2\phi^o(\xi)^2} \right\rangle \quad (24)$$

$$= C_{s,d} \lim_{\epsilon \rightarrow 0} \int_{\mathbb{R}^d \setminus B(0,\epsilon)} \frac{e^{-4\pi^2\phi^o(\xi)^2} - 1}{|2\pi\xi|^{d+s}} > 0, \quad (25)$$

a strictly positive quantity, in view of the sign of  $C_{s,d}$ .

Suppose now that  $m < d$  and consider the subspace  $V = \text{Vect}\{e_1, \dots, e_m\}$ . We write  $x = (x', x'')$ ,  $\xi = (\xi', \xi'' \text{prime})$ , with  $x', \xi' \in V$ . A straightforward computation shows that

$$\begin{aligned} \int_V |x'|^s K_\phi d\mathcal{H}^m &= \langle |x'|^s, K_\phi(x', 0) \rangle_{D'(\mathbb{R}^m), D(\mathbb{R}^m)} \\ &= \left\langle \mathcal{H}_{\xi''=0}^{d-m} \otimes |x'|^s, K_\phi(x', x'') \right\rangle_{D'(\mathbb{R}^d), D(\mathbb{R}^d)} \\ &= \left\langle C_{s,m} Pf \left( \frac{1}{|2\pi\xi'|^{m+s}} \right), h(\xi') \right\rangle_{D'(\mathbb{R}^m), D(\mathbb{R}^m)}, \end{aligned}$$

where the function  $h : \mathbb{R}^m \rightarrow \mathbb{R}$  is defined by

$$h(\xi') = \int_{\mathbb{R}^{d-m}} e^{-4\pi^2\phi^o((\xi', \xi'')^2)} d\xi''.$$

The next lemma states that  $h$  is  $C^1$  and maximal at  $\xi' = 0$ , which in view of (23) and of the sign of  $C_{s,m}$  concludes the proof.  $\square$

LEMMA 2 The function  $h : \mathbb{R}^m \rightarrow \mathbb{R}$ , defined by

$$h(\xi') = \int_{\mathbb{R}^{d-m}} e^{-4\pi^2\phi^o((\xi', \xi'')^2)} d\xi''$$

is  $C^1$ , with fast decay as  $|\xi'| \rightarrow \infty$ , and is maximal at  $\xi' = 0$ .

*Proof.* Recalling (9), we first remark that

$$e^{-4\pi^2\phi^o(\xi', \xi'')^2} \leq e^{-4\pi^2\lambda^2|\xi|^2} \leq e^{-4\pi^2\lambda^2|\xi'|^2},$$

so that the functions  $\xi' \rightarrow e^{-4\pi^2\phi^o(\xi', \xi'')^2}$  and their derivatives are uniformly bounded in  $L^1(\mathbb{R}^{d-m})$ . The  $C^1$  regularity of  $h$  is thus a consequence of the Lebesgue theorem. The above estimate also shows that

$$|h(\xi')| \leq \int_{\mathbb{R}^{d-m}} e^{-4\pi^2\lambda^2(\xi_1^2 + \xi_2^2 + \dots + \xi_d^2)} d\xi_{m+1} \dots d\xi_d \leq \frac{1}{2\lambda^m \sqrt{\pi}^m} e^{-4\pi^2\lambda^2\xi'^2}.$$






FIG. 1.

To determine the maximal value of  $h$ , we consider the sets  $A_{\xi', t}$ , defined for all  $\xi' \in \mathbb{R}^m$  and  $t \in ]0, 1[$  by

$$A_{\xi', t} = \left\{ \xi'' \in \mathbb{R}^{d-m} ; e^{-4\pi^2 \phi^o((\xi', \xi''))^2} \geq t \right\}.$$

Fix  $\xi'_0 \in \mathbb{R}^m$ . The set  $A_{\xi'_0, t}$  can be defined as the intersection of the hyperplane  $\{\xi \in \mathbb{R}^d ; \xi' = \xi'_0\}$  with the Frank shape

$$B_{\phi^o, t} = \left\{ \xi \in \mathbb{R}^d ; \phi^o(\xi) \leq \frac{1}{2\pi} \sqrt{-\ln(t)} \right\}.$$

The set  $B_{\phi^o, t}$  is convex since  $\phi^o$  is convex. Moreover, from the symmetry of  $\phi^o$ , ( $\phi^o(\xi) = \phi^o(-\xi)$ ), we have

$$|A_{\xi'_0, t}| = |A_{-\xi'_0, t}|.$$

Next, let

$$\begin{aligned} \tilde{A}_{\xi'_0, t} &= \frac{1}{2} (A_{\xi'_0, t} + A_{-\xi'_0, t}) \\ &= \left\{ \xi'' \in \mathbb{R}^{d-m} ; \exists (\xi''_1, \xi''_2) \in A_{\xi'_0, t} \times A_{-\xi'_0, t}, \quad \xi'' = \frac{1}{2} (\xi''_1 + \xi''_2) \right\}. \end{aligned}$$

We remark that the convexity of  $\phi^o$  implies that  $\tilde{A}_{\xi'_0, t} \subset A_{0, t}$ . Indeed, let  $\xi'' \in \tilde{A}_{\xi'_0, t}$ ,

$$\begin{aligned} \phi^o((0, \xi'')) &= \phi^o\left(\frac{1}{2}((\xi'_0, \xi''_1) + (-\xi'_0, \xi''_2))\right) \\ &\leq \frac{1}{2} (\phi^o((\xi'_0, \xi''_1)) + \phi^o((-\xi'_0, \xi''_2))) \leq \frac{1}{2\pi} \sqrt{-\ln(t)}, \end{aligned}$$

so that  $e^{-4\pi^2 \phi^o((0, \xi''))^2} \geq t$ , i.e.,  $\xi'' \in A_{0, t}$ . Invoking the Brunn–Minkowski inequality, we obtain

$$|\tilde{A}_{\xi'_0, t}|^{1/(d-m)} = \frac{1}{2} |A_{\xi'_0, t} + A_{-\xi'_0, t}|^{1/(d-m)} \quad (26)$$

$$\geq \frac{1}{2} (|A_{\xi'_0, t}|^{1/(d-m)} + |A_{-\xi'_0, t}|^{1/(d-m)}) \geq |A_{\xi'_0, t}|^{1/(d-m)}, \quad (27)$$

and finally that,

$$|A_{0,t}| \geq |\tilde{A}_{\xi'_0,t}| \geq |A_{\xi'_0,t}|.$$

As this equality holds for any  $\xi'_0 \in \mathbb{R}^m$ , it follows that  $h$  is maximal at  $\xi' = 0$ .  $\square$

#### 4. The Bence–Merriman–Osher-like algorithm

Barles and Souganidis [6] have studied the convergence of a general approximation scheme to viscosity solutions of nonlinear second-order parabolic PDE's of the type

$$u_t + F(D^2u, Du) = 0. \quad (28)$$

The main assumption on the function  $F$  is its ellipticity, i.e.,  $F$  satisfies

$$\forall p \in \mathbb{R}^d \setminus \{0\}, \forall X, Y \in \mathbf{M}_s^{d \times d}, \quad X \leq Y \Leftrightarrow F(X, p) \leq F(Y, p). \quad (29)$$

Let  $BUC(\mathbb{R}^d)$  denote the space of bounded uniformly continuous functions on  $\mathbb{R}^d$ . Thus, Barles and Souganidis study a family of operators  $G_h : BUC(\mathbb{R}^d) \rightarrow BUC(\mathbb{R}^d)$  for  $h > 0$ , which satisfy, for all  $u, v \in BUC(\mathbb{R}^d)$ :

- *Continuity*

$$\forall c \in \mathbb{R}, \quad G_h(u + c) = G_h u, \quad (30)$$

- *Monotonicity*

$$u \leq v \Leftrightarrow G_h u \leq G_h v + o(h), \quad (31)$$

(see Remark 2.1 in [6])

- *Consistency*

$$\forall \varphi \in C^\infty(\mathbb{R}^d), \quad \begin{cases} \lim_{h \rightarrow 0} h^{-1}(G_h(\varphi) - \varphi)(x) \leq -F_*(D^2\varphi(x), D\varphi(x)), \\ \lim_{h \rightarrow 0} h^{-1}(G_h(\varphi) - \varphi)(x) \geq -F^*(D^2\varphi(x), D\varphi(x)). \end{cases} \quad (32)$$

For all  $T > 0$  and for all partitions  $P = \{O = t_0 < \dots < t_n = T\}$  of  $[0, T]$ , one can then define a sequence of functions  $u_P : \mathbb{R}^d \times [0, T] \rightarrow \mathbb{R}$  by

$$u_P(., t) = \begin{cases} G_{t-t_i}(u_P(., t_i)) & \text{if } t \in (t_i, t_{i+1}], \\ g & \text{if } t = 0. \end{cases} \quad (33)$$

If additionally the following condition holds,

- *Stability*

$$\left\{ \begin{array}{l} \text{there exists } \omega \in C([0, \infty], [0, \infty]), \text{ independent of } P \text{ and depending} \\ \text{on } g \text{ only through the modulus of continuity of } g, \\ \text{such that } \omega(0) = 0 \text{ and for all } t \in [0, t], \\ \|u_P(., t) - g\|_{L^\infty} \leq \omega(t), \end{array} \right. \quad (34)$$

then the following theorem holds [6]:

**THEOREM 1** Assume that  $G_h : BUC(\mathbb{R}^d) \rightarrow BUC(\mathbb{R}^d)$  satisfies (30), (31), (32), and (34) for all  $T > 0$ ,  $g \in BUC(\mathbb{R}^d)$  and all partitions  $P$  of  $[0, T]$ . Then,  $u_P$  defined in (33) converges uniformly in  $\mathbb{R} \times [0, T]$  to the viscosity solution of (28).

This result was used by H. Ishii, G. Pires and P. E. Souganidis in [32] to study anisotropic mean curvature flow. These authors introduce a kernel  $f$ , which satisfies:

$$(H_1) \quad f(x) \geq 0, \quad f(-x) = f(x) \quad \text{for all } x \in \mathbb{R}^d, \quad \text{and} \quad \int_{\mathbb{R}^d} f(x) dx = 1,$$

$$(H_2) \quad \int_{p^\perp} (1 + |x|^2) |f(x)| d\mathcal{H}^{d-1} < \infty \quad \text{for all } p \in \mathbb{S}^{d-1},$$

$$(H_3) \quad \begin{cases} \text{the functions } p \rightarrow \int_{p^\perp} f(x) d\mathcal{H}^{d-1} & p \rightarrow \int_{p^\perp} x_i x_j f(x) d\mathcal{H}^{d-1}, \\ 1 \leq i, j \leq d, & \text{are continuous on } \mathbb{S}^{d-1}, \end{cases}$$

$$(H_4) \quad \int_{\mathbb{R}^d} |x|^2 |f(x)| dx < \infty,$$

$$(H_5) \quad \text{For all collections } \{R(\rho)\}_{0 < \rho < 1} \subset \mathbb{R} \text{ such that } R(\rho) \rightarrow \infty \text{ and } \rho R(\rho)^2 \rightarrow 0 \text{ as } \rho \rightarrow 0, \text{ and for all functions } g : \mathbb{R}^{d-1} \rightarrow \mathbb{R} \text{ of the form } g(\xi) = a + \langle A\xi, \xi \rangle \text{ where } a \in \mathbb{R} \text{ and where } A \text{ is a symmetric matrix,}$$

$$\lim_{\rho \rightarrow 0} \sup_{U \in \mathbf{O}(d)} \sup_{0 < r < \rho} \left| \int_{B(0, R(\rho))} f_U(\xi, r g(\xi)) g(\xi) d\xi - \int_{\mathbb{R}^{d-1}} f_U(\xi, 0) g(\xi) d\xi \right| = 0,$$

where  $\mathbf{O}(n)$  denotes the group of  $d \times d$  orthogonal matrices, and where  $f_U : \mathbb{R}^d \rightarrow \mathbb{R}$  is defined for all  $U \in \mathbf{O}(d)$  by  $f_U(x) = f(U^*x)$ .

Theorem 1 has been applied to schemes for anisotropic mean curvature motion (see Theorem 3.3 in [32]) with  $G_h$  defined by

$$G_h \Psi(x) = \sup \{ \lambda \in \mathbb{R} ; S_h \mathbb{1}_{\Psi \geq \lambda}(x) \geq \theta_h \} \quad (35)$$

$$= \inf \{ \lambda \in \mathbb{R} ; S_h \mathbb{1}_{\Psi \geq \lambda}(x) < \theta_h \}, \quad (36)$$

where

$$S_h g(x) = h^{-d/2} f(\cdot / \sqrt{h}) * g(x) = h^{-d/2} \int_{\mathbb{R}^d} f(y / \sqrt{h}) g(x - y) dy, \quad \theta_h = \frac{1}{2} + c\sqrt{h},$$

and where  $F(X, p)$  is given by

$$F(X, p) = - \left( \int_{p^\perp} f(x) d\mathcal{H}^{d-1}(x) \right)^{-1} \left( \frac{1}{2} \int_{p^\perp} \langle Xx, x \rangle f(x) d\mathcal{H}^{d-1}(x) + c|p| \right)$$

(the last term in this integral models a forcing term).

In this section, we follow the proof in [32] to show a consistency result in our case when  $f$  is a non positive kernel and does not have moments of order two (i.e.,  $x \rightarrow |x|^2 f(x) \notin L^1(\mathbb{R}^d)$ ). We introduce two operators  $G_h^+$  and  $G_h^-$  defined by

$$G_h^+ \Psi(x) = \sup \{ \lambda \in \mathbb{R} ; S_h \mathbb{1}_{\Psi \geq \lambda}(x) \geq \theta_h \}, \quad (37)$$

$$G_h^- \Psi(x) = \inf \{ \lambda \in \mathbb{R} ; S_h \mathbb{1}_{\Psi \geq \lambda}(x) < \theta_h \}, \quad (38)$$

which are not necessarily equal as our kernel is not being nonnegative. To adapt these results to our context we modify the assumptions (H<sub>1</sub>), (H<sub>4</sub>) and (H<sub>5</sub>) as follows

( $H'_1$ )  $\int_{p^\perp} f(x) d\mathcal{H}^{d-1} > 0$  for all  $p \in \mathbb{S}^{d-1}$ ,  $f(-x) = f(x)$  and  $\int_{\mathbb{R}^d} f(x) dx = 1$ ,

( $H'_4$ )  $\int_{\mathbb{R}^d} |x|^{2-\mu} |f(x)| dx < \infty$  for  $0 < \mu < 2$ ,

( $H'_5$ ) Assume that  $\mu \in ]0, 1/2]$ . Then for all collections  $\{R(\rho)\}_{0 < \rho < 1} \subset \mathbb{R}$  such that  $R(\rho) \rightarrow \infty$  and  $\rho R(\rho)^{2-\mu} \rightarrow 0$  as  $\rho \rightarrow 0$ , and for all functions  $g : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$  of the form  $g(\xi) = a + \langle A\xi, \xi \rangle$  where  $a \in \mathbb{R}$  and where  $A$  is a symmetric matrix,

$$\lim_{\rho \rightarrow 0} \sup_{U \in \mathcal{O}(d)} \sup_{0 < r < \rho} \left| \int_{B(0, R(\rho))} f_U(\xi, rg(\xi)) g(\xi) d\xi - \int_{\mathbb{R}^{d-1}} f_U(\xi, 0) g(\xi) d\xi \right| = 0,$$

$$\lim_{\rho \rightarrow 0} \sup_{U \in \mathcal{O}(d)} \sup_{0 < r < \rho} \left| \int_{B(0, R(\rho))} |f_U(\xi, rg(\xi))| g(\xi) d\xi - \int_{\mathbb{R}^{d-1}} |f_U(\xi, 0)| g(\xi) d\xi \right| = 0.$$

In this last statement,  $B(0, R(\rho))$  denotes the  $(n-1)$ -dimensional ball, centered at 0 and of radius  $R(\rho)$ .

4.1  $K_\phi$  satisfies ( $H_2$ ,  $H_3$ ) and ( $H'_1$ ,  $H'_4$ ,  $H'_5$ )

We remark that  $\hat{K}_\phi(\xi) = \hat{K}_\phi(-\xi)$  and  $\mathfrak{F}(K_\phi)(0) = 1$ , so that

$$K_\phi(-x) = K_\phi(x) \quad \text{for all } x \in \mathbb{R}^d, \quad \text{and} \quad \int_{\mathbb{R}^d} K_\phi(x) dx = 1.$$

Moreover, Proposition 4 shows that

$$\int_{p^\perp} K_\phi(x) d\mathcal{H}^{d-1} \geq \frac{1}{(4\pi)^{d/2} \Lambda^d} > 0 \quad \text{for all } p \in \mathbb{S}^{d-1},$$

so that ( $H'_1$ ) is satisfied. Propositions 4 and 5 also imply that  $K_\phi$  satisfies ( $H_2$ ), i.e.,

$$\int_{p^\perp} (1 + |x|^2) |K_\phi(x)| d\mathcal{H}^{d-1} < \infty \quad \text{for all } p \in \mathbb{S}^{d-1}. \quad (39)$$

Concerning ( $H_3$ ), we note that

$$\frac{1}{2} \int_{p^\perp} x \otimes x K_\phi(x) d\mathcal{H}^{d-1} = \frac{1}{2\sqrt{\pi}} \phi_{\xi\xi}^o(p),$$

and that

$$\int_{p^\perp} K_\phi d\mathcal{H}^{d-1} = \frac{1}{2\sqrt{\pi} \phi^o(p)}.$$

Since  $\phi^o$  is smooth on  $\mathbb{R}^d \setminus \{0\}$  and positive (in particular  $\phi^o \geq \lambda$  on  $\mathbf{S}^d$ ) we see that the functions

$$p \rightarrow \int_{p^\perp} K_\phi(x) d\mathcal{H}^{d-1} \quad p \rightarrow \int_{p^\perp} x_i x_j K_\phi(x) d\mathcal{H}^{d-1}, \quad 1 \leq i, j \leq d,$$

are continuous on  $\mathbf{S}^d$ .

We next prove that if  $0 < \mu < 2$ , then

$$\int_{\mathbb{R}^d} |x|^{2-\mu} |f(x)| dx < \infty.$$

Indeed, Proposition 2 with  $s = 1 - \mu/2$  shows that

$$\begin{aligned} \int_{\mathbb{R}^d} |x|^{2-\mu} |f(x)| dx &\leq \int_{\mathbb{R}^d} \frac{C_{\phi^o,s} |x|^{2-\mu}}{1 + |x|^{d+1+(1-\mu/2)}} dx \leq \int_{\mathbb{R}^d} \frac{C}{1 + |x|^{d+\mu/2}} dx \\ &\leq C |\mathbf{S}^d| \int_0^\infty \frac{1}{(1+r^{1+\mu/2})} dr < \infty, \end{aligned}$$

for some generic constant  $C$ .

It remains to prove  $(H'_5)$ : Let  $0 < \mu < 1/2$  and let  $R : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that, as  $\rho \rightarrow 0$ ,  $R(\rho) \rightarrow \infty$  and  $\rho R(\rho)^{2-\mu} \rightarrow 0$ . Setting  $f_U(x) = K_\phi(U^*x)$ , we consider

$$\begin{aligned} &\int_{\mathbb{R}^{d-1}} |(f_U(\tilde{x}, rg(\tilde{x})) \mathbb{1}_{B(0,R(\rho))}(\tilde{x}) - f_U(\tilde{x}, 0)) g(\tilde{x})| d\tilde{x} \\ &\leq \int_{B(0,R(\rho))^c} |f_U(\tilde{x}, 0)| |g(\tilde{x})| d\tilde{x} + \int_{B(0,R(\rho)^{\frac{2-\mu}{2}})} |(f_U(\tilde{x}, rg(\tilde{x})) - f_U(\tilde{x}, 0)) g(\tilde{x})| d\tilde{x} \\ &\quad + \int_{B(0,R(\rho)) \setminus B(0,R(\rho)^{\frac{2-\mu}{2}})} |(f_U(\tilde{x}, rg(\tilde{x})) - f_U(\tilde{x}, 0)) g(\tilde{x})| d\tilde{x}. \quad (40) \end{aligned}$$

From the decay assumptions on  $K_\phi$  (see Proposition 2) we have

$$|f_U(\tilde{x}, rg(\tilde{x})) g(\tilde{x})| \leq \frac{C}{1 + |\tilde{x}|^{d-1+s}},$$

where  $C$  does not depend on  $U$  and  $r$ . Then, it holds that

$$\begin{aligned} \int_{B(0,R(\rho))^c} |f_U(\tilde{x}, 0) g(\tilde{x})| d\tilde{x} &\leq C \int_{B(0,R(\rho))^c} \frac{1}{1 + |\tilde{x}|^{d-1+s}} d\tilde{x} \\ &\leq C |\mathbf{S}^{d-1}| \int_{R(\rho)}^\infty \frac{1}{1 + |r|^{1+s}} dr \\ &\leq C |\mathbf{S}^{d-1}| R(\rho)^{-s}, \end{aligned}$$

and

$$\begin{aligned}
& \int_{B(0, R(\rho)) \setminus B(0, R(\rho)^{\frac{2-\mu}{2}})} \left| (f_U(\tilde{x}, rg(\tilde{x})) - f_U(\tilde{x}, 0))g(\tilde{x}) \right| d\tilde{x} \\
& \leq 2C \int_{B(0, R(\rho)) \setminus B(0, R(\rho)^{\frac{2-\mu}{2}})} \frac{1}{1 + |\tilde{x}|^{d-1+s}} d\tilde{x} \\
& \leq 2C |\mathbf{S}^{d-1}| \int_{R(\rho)^{\frac{2-\mu}{2}}}^{R(\rho)} \frac{1}{1 + |r|^{1+s}} dr \\
& \leq C |\mathbf{S}^{d-1}| (R(\rho)^{\frac{-(2-\mu)s}{2}} - R(\rho)^{-s}),
\end{aligned}$$

and implies that the first and the third term in (40) converge to 0 uniformly with respect to  $U$  and  $r$  as  $\rho \rightarrow 0$ . Moreover, using smoothness property of  $K_\phi (C^\infty(\mathbb{R}^d))$ , we have

$$|\nabla f_U(\tilde{x}, rg(\tilde{x}))| |g(\tilde{x})| \leq \frac{\tilde{C}}{1 + |\tilde{x}|^{d-1+s}} \in L^1(\mathbb{R}^{d-1}),$$

uniformly on  $U$  and  $r$ , and with  $r < \rho$ ,

$$\begin{aligned}
& \int_{B(0, R(\rho)^{\frac{2-\mu}{2}})} \left| (f_U(\tilde{x}, rg(\tilde{x})) - f_U(\tilde{x}, 0))g(\tilde{x}) \right| d\tilde{x} \\
& \leq rR(\rho)^{2-\mu} \int_{B(0, R(\rho)^{\frac{2-\mu}{2}})} \left\{ \sup_{U \in \mathcal{O}(d)} \sup_{0 \leq s \leq rg(\tilde{x})} \{ |\partial_{x_n} f_U(\tilde{x}, s)g(\tilde{x})| \} \right\} d\tilde{x} \\
& \leq C\rho R(\rho)^{2-\mu},
\end{aligned}$$

for some generic constant  $C$ .

We conclude that

$$\lim_{\rho \rightarrow 0} \sup_{U \in \mathcal{O}(d)} \sup_{0 < r < \rho} \left| \int_{B(0, R(\rho))} f_U(\tilde{x}, rg(\tilde{x}))g(\tilde{x}) d\tilde{x} - \int_{\mathbb{R}^{d-1}} f_U(\tilde{x}, 0)g(\tilde{x}) d\tilde{x} \right| = 0.$$

The second statement in  $(H'_2)$  is established similarly.

#### 4.2 The consistency proof

**THEOREM 2** Let  $\varphi \in C^2(\mathbb{R}^d)$ . For all  $z \in \mathbb{R}^d$  and  $\epsilon > 0$ , there exists  $\delta > 0$  such that for all  $x \in B(z, \delta)$  and  $h \in (0, \delta]$ , if  $\nabla\phi(x) \neq 0$  we have

$$\begin{aligned}
G_h^- \varphi(x) & \leq \varphi(x) + (-F(D^2\varphi(z), D\varphi(z)) + \epsilon)h, \\
G_h^+ \varphi(x) & \geq \varphi(x) + (-F(D^2\varphi, D\varphi(z)) - \epsilon)h.
\end{aligned}$$

*Proof.* We closely follow the argument in [32].

1. We only prove the first inequality. The other one is obtained similarly.

2. Without loss of generality, we can assume that  $z = 0$ . Let us fix  $a \in \mathbb{R}$ , such that

$$a > -F(D^2\varphi(0), D\varphi(0)).$$

The inequality is proved if we can exhibit a  $\delta > 0$  such that, for all  $x \in B(0, \delta)$  and  $h \in (0, \delta]$ ,

$$S_h \mathbb{1}_{\varphi \geq \varphi(x) + ah}(x) < \theta_h.$$

3. Fix  $\delta_1 > 0$ , such that  $D\varphi \neq 0$  on  $B(0, \delta_1)$  and choose a continuous family  $\{U(x)\}_{x \in B(0, \delta_1)} \subset O(d)$ , such that for all  $x \in B(0, \delta_1)$ ,

$$U(x) \left( \frac{D\varphi(x)}{|D\varphi(x)|} \right) = e_d,$$

where  $e_d$  denotes the unit vector with components  $(0, 0, \dots, 0, 1) \in \mathbb{R}^d$ . Note that if  $x \in B(0, \delta_1)$ , then

$$S_h \mathbb{1}_{\varphi \geq \varphi(x) + ah} = \int_{\mathbb{R}^d} f_{U(x)}(y) \mathbb{1}_{\varphi \geq \varphi(x) + ah}(x - \sqrt{h}U(x)^*y) dy.$$

4. Choosing  $\delta$  smaller if necessary,  $(H'_1)$  implies the inequality

$$a > -F(D^2\varphi, D\varphi) \quad \text{in } B(0, \delta_1),$$

or in other words,

$$\begin{aligned} \frac{1}{2} \int_{\mathbb{R}^{d-1}} \langle P^*U(x)D^2\varphi(x)U(x)^*P\xi, \xi \rangle f_{U(x)}(\xi, 0) d\xi - a \int_{\mathbb{R}^{d-1}} f_{U(x)}(\xi, 0) d\xi \\ < -c|D\varphi(x)|, \end{aligned} \quad (41)$$

where  $P$  denotes the  $d \times (d-1)$  matrix with components  $P_{ij} = \delta_{ij}$ .

5. We next fix  $\epsilon > 0$ , and  $\delta_2 \in (0, \delta_1[$ , such that for all  $x \in B(0, \delta_2)$ ,

$$\begin{aligned} \frac{1}{2} \int_{\mathbb{R}^{d-1}} \langle P^*U(0)(D^2\varphi(0) + 3\epsilon^2 I)U(0)^*P\xi, \xi \rangle f_{U(x)}(\xi, 0) d\xi \\ - (a - \epsilon^2) \int_{\mathbb{R}^{d-1}} f_{U(x)}(\xi, 0) d\xi < -(\xi + \epsilon)|D\varphi(0)|. \end{aligned}$$

6. The Taylor theorem yields a  $\gamma > 0$  such that for all  $h > 0$ ,  $y \in \mathbb{R}^d$ , and  $x \in B(0, \delta_2)$ , if  $\sqrt{h}|y| \leq \gamma$ , then

$$\begin{aligned} \varphi(x - \sqrt{h}U(x)^*y) &\leq \varphi(x) - \sqrt{h} \langle D\varphi(x), U(x)^*y \rangle \\ &\quad + \frac{h}{2} \langle U(x)(D^2\varphi(x) + \epsilon^2 I)U(x)^*y, y \rangle \\ &\leq \varphi(x) - \sqrt{h}|D\varphi(x)|y_d + Chy_d^2 \\ &\quad + \frac{h}{2} \langle P^*U(x)(D^2\varphi(x) + 2\epsilon^2 I)U(x)^*Py', y' \rangle, \end{aligned}$$

and

$$\begin{aligned}
\varphi(x - \sqrt{h}U(x)^*y) &\geq \varphi(x) - \sqrt{h} \langle D\varphi(x), U(x)^*y \rangle \\
&\quad + \frac{h}{2} \langle U(x)(D^2\varphi(x) - \epsilon^2 I)U(x)^*y, y \rangle \\
&\geq \varphi(x) - \sqrt{h}|D\varphi(x)|y_d - Chy_d^2 \\
&\quad + \frac{h}{2} \langle P^*U(x)(D^2\varphi(x) - 2\epsilon^2 I)U(x)^*Py', y' \rangle,
\end{aligned}$$

where we write  $y = (y', y_d) \in \mathbb{R}^{d-1} \times \mathbb{R}$ , and where  $C$  is a positive constant.

7. Reducing  $\gamma$  and  $\delta_2$  if necessary, the previous inequalities imply that for  $y \in B(0, \gamma/\sqrt{h})$  and  $x \in B(0, \delta_2)$ ,

- if  $\varphi(x - \sqrt{h}U(x)^*y) \geq \varphi(x) + ah$ , then

$$\begin{aligned}
y_d &\leq \frac{\sqrt{h}}{|D\varphi(x)| - C\sqrt{h}y_d} \left( -a + \frac{1}{2} \langle P^*U(x)(D^2\varphi(x) + 2\epsilon^2 I)U(x)^*Py', y' \rangle \right) \\
&\leq \frac{\sqrt{h}}{|D\varphi(0)|} \left( -a + \epsilon^2 + \frac{1}{2} \langle P^*U(0)(D^2\varphi(0) + 3\epsilon^2 I)U(0)^*Py', y' \rangle \right),
\end{aligned}$$

- if

$$y_d \leq \frac{\sqrt{h}}{|D\varphi(0)|} \left( -a - \epsilon^2 + \frac{1}{2} \langle P^*U(0)(D^2\varphi(0) - 3\epsilon^2 I)U(0)^*Py', y' \rangle \right),$$

then

$$\varphi(x - \sqrt{h}U(x)^*y) \geq \varphi(x) + ah.$$

We define

$$\begin{cases} a^\epsilon = (a - \epsilon^2)|D\varphi(0)|^{-1}, \\ a_\epsilon = (a + \epsilon^2)|D\varphi(0)|^{-1}, \\ A^\epsilon = |D\varphi(0)|^{-1}P^*U(0)(D^2\varphi(0) + 3\epsilon^2 I)U(0)^*P, \\ A_\epsilon = |D\varphi(0)|^{-1}P^*U(0)(D^2\varphi(0) - 3\epsilon^2 I)U(0)^*P, \end{cases}$$

and for  $y' \in \mathbb{R}^{d-1}$

$$g^\epsilon(y') = \left( -a^\epsilon + \frac{1}{2} \langle A^\epsilon y', y' \rangle \right), \quad g_\epsilon(y') = \left( -a_\epsilon + \frac{1}{2} \langle A_\epsilon y', y' \rangle \right).$$

We also set

$$V_{h,x} = \left\{ y \in \mathbb{R}^d ; \varphi(x - \sqrt{h}U(x)^*y) \geq \varphi(x) + ah \right\},$$

and

$$\begin{cases} E_{\epsilon,h,x}^+ = \left\{ y \in \mathbb{R}^d ; y_d \leq \sqrt{h}g_\epsilon(y') \right\}, \\ E_{\epsilon,h,x}^- = \left\{ y \in \mathbb{R}^d ; y_d \leq \sqrt{h}g^\epsilon(y') \right\}. \end{cases}$$



We check that for all  $x \in B(0, \delta_2)$ ,

$$\begin{cases} (V_{h,x} \cap B(0, \gamma/\sqrt{h})) & \subset (E_{\epsilon,h,x}^+ \cap B(0, \gamma/\sqrt{h})), \\ (E_{\epsilon,h,x}^- \cap B(0, \gamma/\sqrt{h})) & \subset (V_{h,x} \cap B(0, \gamma/\sqrt{h})). \end{cases}$$

8. The assumption  $(H_4)$  yields the existence of a decreasing function  $\omega \in C([0, \infty), [0, \infty))$  such that  $\omega(R) \rightarrow 0$  as  $R \rightarrow \infty$ , and

$$\int_{B(0,R)^c} |f(y)||y|^{2-\mu} dy \leq \omega(R)^2, \text{ for all } R \geq 0.$$

For each  $0 < t < 1$ , we define the family of sets  $R(t) \in (0, \infty)$  by

$$\omega(R(t)) = tR(t)^{2-\mu}, \quad (42)$$

which satisfy  $(H_5')$ . We then choose  $\tau \in (0, 1)$  such that

$$R(t) \leq \gamma/t, \text{ for all } t \in (0, \tau]. \quad (43)$$

9. Let

$$\rho = \sqrt{h}, \quad T(\rho) = B_{n-1}(0, R(\rho)) \times \mathbb{R} \subset \mathbb{R}^d.$$

For all  $h \in ]0, \tau^2)$  and for all  $x \in B(0, \delta_2)$ , we estimate

$$\begin{aligned} \int_{V_{h,x}} f_{U(x)}(y) dy &= \int_{\mathbb{R}^d} f_{U(x)}(y) \mathbb{1}_{\varphi \geq \varphi(x) + ah}(x - \sqrt{h}U^*(x)y) dy \\ &\leq \int_{V_{h,x} \cap B(0, R(\rho))} f_{U(x)}(y) dy + \int_{B(0, R(\rho))^c} |f_{U(x)}(y)| dy \\ &\leq \int_{E_{\epsilon,h,x}^+ \cap B(0, R(\rho))} f_{U(x)}(x) dx + \int_{B(0, R(\rho))^c} |f_{U(x)}(y)| dy \\ &\quad + \int_{(E_{\epsilon,h,x}^+ \setminus E_{\epsilon,h,x}^-) \cap B(0, R(\rho))} |f_{U(x)}(y)| dy \\ &\leq \int_{E_{\epsilon,h,x}^+ \cap T(\rho)} f_{U(x)}(y) dy + \int_{(E_{\epsilon,h,x}^+ \setminus E_{\epsilon,h,x}^-) \cap T(\rho)} |f_{U(x)}(y)| dy \\ &\quad + 3 \int_{B(0, R(\rho))^c} |f_{U(x)}(y)| dy. \end{aligned}$$

10. For the last integral above, we have

$$\int_{B(0, R(\rho))^c} |f_{U(x)}(y)| dy \leq \frac{1}{R(\rho)^{2-\mu}} \int_{B(0, R(\rho))^c} |y|^{2-\mu} |f_{U(x)}(y)| dy \leq \omega(R(\rho))\rho,$$

and moreover, since  $K_\phi$  is symmetric,

$$\frac{1}{2} = \int_{y_d \leq 0} f_{U(x)}(y) dy \leq \int_{T(\rho) \cap \{y_d \leq 0\}} |f_{U(x)}(y)| dy + \omega(R(\rho))\rho.$$

We note that

$$\begin{aligned}
& \int_{T(\rho) \cap E_{\epsilon, h, x}^+} f_{U(x)}(y) dy \\
&= \int_{T(\rho) \cap \{y_d \leq \rho g^\epsilon(y')\}} f_{U(x)}(y) dy \\
&= \int_{T(\rho) \cap \{y_d \leq 0\}} f_{U(x)}(y) dy + \int_{B_{n-1}(0, R(\rho))} d\xi \int_0^{\rho g^\epsilon(y')} f_{U(x)}(\xi, r) dr \\
&= \int_{T(\rho) \cap \{y_d \leq 0\}} f_{U(x)}(y) dy + \int_0^\rho dr \int_{B_{n-1}(0, R(\rho))} f_{U(x)}(\xi, r g(\xi)) g^\epsilon(\xi) d\xi.
\end{aligned}$$

It follows from  $(H'_\xi)$  that as  $\rho \rightarrow 0$ ,

$$\frac{1}{\rho} \left\{ \int_{T(\rho) \cap E_{\epsilon, h, x}^+} f_{U(x)}(y) dy - \int_{T(\rho) \cap \{y_d \leq 0\}} f_{U(x)}(y) dy \right\} \rightarrow \int_{\mathbb{R}^{d-1}} f_{U(x)}(\xi, 0) g^\epsilon(\xi) d\xi,$$

uniformly with respect to  $x$ . Possibly reducing  $\tau$  we may assume that for  $x \in B(0, \delta_2)$ ,

$$\frac{1}{\rho} \left\{ \int_{T(\rho) \cap E_{\epsilon, h, x}^+} f_{U(x)}(y) dy - \int_{T(\rho) \cap \{y_d \leq 0\}} f_{U(x)}(y) dy \right\} \leq \int_{\mathbb{R}^{d-1}} f_{U(x)}(\xi, 0) g^\epsilon(\xi) d\xi + \epsilon^2.$$

Using same argument, we also conclude that

$$\begin{aligned}
& \int_{T(\rho) \cap (E_{\epsilon, h, x}^+ \setminus E_{\epsilon, h, x}^-)} |f_{U(x)}(y)| dy = \left\{ \int_{T(\rho) \cap \{0 \leq y_d \leq \rho g^\epsilon(y')\}} |f_{U(x)}(y)| dy \right\} \\
& \quad - \left\{ \int_{T(\rho) \cap \{0 \leq y_d \leq \rho g_\epsilon(y')\}} |f_{U(x)}(y)| dy \right\} \\
& \leq \rho \int_{\mathbb{R}^{d-1}} |f_{U(x)}|(\xi, 0) (g^\epsilon(\xi) - g_\epsilon(\xi)) d\xi + \rho \epsilon^2 \\
& \leq \rho \epsilon^2 \left( 1 + \int_{\mathbb{R}^{d-1}} (2 + 3|\xi|^2) |f_{U(x)}|(\xi, 0) d\xi \right) \\
& \leq C_0 \rho \epsilon^2,
\end{aligned}$$

where

$$C_0 = \sup_{x \in B(0, \delta_2)} \left\{ 1 + \int_{\mathbb{R}^{d-1}} (2 + 3|\xi|^2) |f_{U(x)}|(\xi, 0) d\xi \right\}.$$

11. Finally, noting that from (41),

$$\int_{\mathbb{R}^{d-1}} f_{U(x)}(\xi, 0) g^\epsilon(\xi) d\xi \leq -c - \epsilon,$$

we get

$$\begin{aligned} \int_{\mathbb{R}^d} f(x) \mathbb{1}_{\varphi \geq \varphi(x) + ah}(x - \sqrt{h}z) dz &\leq \frac{1}{2} + \int_{\mathbb{R}^{d-1}} f_{U(x)}(\xi, 0) g(\xi) d\xi \\ &\quad + \rho(\epsilon^2 + 4\omega(R(\rho)) + C_0\epsilon^2) \\ &\leq \frac{1}{2} + \rho(-c - \epsilon + \epsilon^2 + 4\omega(R(\rho)) + C_0\epsilon^2) \\ &< \theta_h, \end{aligned}$$

for  $\epsilon$  sufficiently small. □

Even if the function  $\phi$  is regular,  $G_h^+ \phi$  and  $G_h^- \phi$  need not be equal and continuous. However, it is easy to check that if  $\varphi = \mathbb{1}_\Omega$  is a characteristic function then  $G_h^+ \mathbb{1}_\Omega = G_h^- \mathbb{1}_\Omega$ . The next proposition shows that if  $\varphi$  is smooth,  $G_h^- \varphi(x) = G_h^+ \varphi(x) + o(h)$ , so that one could conceivably build a Bence Merriman Osher type scheme using either  $G_h^+$  or  $G_h^-$ .

**PROPOSITION 7** Let  $\varphi \in C^2(\mathbb{R}^d)$ . Let  $x \in \mathbb{R}^d$  such as  $\nabla\varphi(x) \neq 0$ , then

$$G_h^- \varphi(x) = G_h^+ \varphi(x) + o(h).$$

*Proof.* Let  $x \in \mathbb{R}^d$  such as  $\nabla\varphi(x) \neq 0$  and for all  $h > 0$  let

$$\epsilon(h) = G_h^+ \varphi(x) - G_h^- \varphi(x).$$

Introduce also  $g_h(\lambda) : \mathbb{R} \rightarrow \mathbb{R}$  defined by

$$g_h(\lambda) = S_h \chi_{\varphi \geq \lambda}(x) = \int_{\mathbb{R}^d} K_{\phi, h}(y) \chi_{\varphi \geq \lambda}(x - y) dy.$$

This function may not be continuous. We claim that its jumps are bounded by  $o(\sqrt{h})$ . Indeed, for all  $\lambda \in \mathbb{R}$ , one can express  $g_h(\lambda)$  as

$$\begin{aligned} g_h(\lambda) &= \int_{B(0, \sigma)} K_{\phi, h}(y) \chi_{\{\varphi \geq \lambda\}}(x - y) dy + \int_{\mathbb{R}^d \setminus B(0, \sigma)} K_{\phi, h}(y) \chi_{\{\varphi \geq \lambda\}}(x - y) dy \\ &= \tilde{g}_h(\lambda) + R_h(\lambda), \end{aligned}$$

where  $\sigma$  is chosen sufficiently small so that  $|\nabla\varphi(y)| > 0$  for all  $y \in B(x, \sigma)$ . Let  $0 < \mu < 1$ , let

$$\omega(R) = \int_{B(0, R)^c} |y|^{2-\mu} |K_\phi(y)| dy,$$

and let  $R(t)$  be defined by the equality  $\omega(R(t)) = tR(t)^{2-\mu}$ . Note that  $(H_4')$  implies that  $\sqrt{h}R(h)^{2-\mu} \rightarrow 0$  as  $h \rightarrow 0$ , so that  $\sqrt{h}R(h)^{1-\mu/2} < \sigma$  for  $h$  sufficiently small, it follows that

$$|R_h(\lambda)| \leq \int_{\mathbb{R}^d \setminus B(0, \sqrt{h}R(h)^{1-\mu/2})} |K_{\phi, h}(y)| dy.$$

Moreover, changing variables, we see that

$$\begin{aligned} \int_{\mathbb{R}^d \setminus B(0, \sqrt{h}R(h)^{1-\mu/2})} |K_{\phi,h}(y)| dy &\leq \int_{\mathbb{R}^d \setminus B(0, R(h)^{1-\mu/2})} |K_{\phi}(y)| dy \\ &\leq \frac{1}{R(h)^{(2-\mu)^2/2}} \int_{\mathbb{R}^d \setminus B(0, R(h)^{1-\mu/2})} |y|^{2-\mu} |K_{\phi}(y)| dy \\ &\leq \frac{\omega(R(h))}{R(h)^{(2-\mu)^2/2}}. \end{aligned}$$

Since  $0 < (2 - \mu)/2 < 1$ , it follows that

$$|R_h(\lambda)| \leq \left( \frac{\omega(R(h))}{R(h)^{2-\mu}} \right)^{1-\mu/2} = h^{1-\mu/2} = o(\sqrt{h}).$$

Further, the fact that  $|\nabla\varphi(y)| > 0$  on  $B(x, \sigma)$  show that  $\tilde{g}_h$  is continuous in  $\lambda$ , which proves the claim.

Recall that

$$\begin{cases} G_h^- \varphi(x) = \inf \{s \in \mathbb{R} ; S_h \chi_{\varphi \geq s}(x) < \theta_h\}, \\ G_h^+ \varphi(x) = \sup \{s \in \mathbb{R} ; S_h \chi_{\varphi \geq s}(x) \geq \theta_h\}, \end{cases}$$

it follows from the claim above that

$$S_h \chi_{\varphi \geq G_h^- \varphi(x)}(x) = \theta_h + o(\sqrt{h}), \quad \text{and} \quad S_h \chi_{\varphi \geq G_h^+ \varphi(x)}(x) = \theta_h + o(\sqrt{h}),$$

and consequently

$$\int_{\mathbb{R}^d} K_{\phi,h}(y) \chi_{G_h^- \varphi(x) \leq \varphi \leq G_h^- \varphi(x) + \epsilon(h)}(x-y) dy = o(\sqrt{h}).$$

One can use the same argument as in the consistency proof, (in particular see point 7) to show that asymptotically, the above integral behaves like

$$\int_{\mathbb{R}^d} K_{\phi,h}(y) \chi_{G_h^- \varphi(x) \leq \varphi \leq G_h^- \varphi(x) + \epsilon(h)}(x-y) dy = \frac{\epsilon(h)}{|\nabla\varphi(x)|\sqrt{h}} \int_{p^\perp} K_{\phi}(x) d\mathcal{H}^{d-1}(x) + o(\sqrt{h}),$$

where  $p = \frac{\nabla\varphi(x)}{|\nabla\varphi(x)|}$ . In conclusion, as  $\int_{p^\perp} K_{\phi}(x) d\mathcal{H}^{d-1}(x) > 0$ , we deduce that

$$\epsilon(h) = \frac{|\nabla\varphi(x)|}{\int_{p^\perp} K_{\phi}(x) d\mathcal{H}^{d-1}(x)} o(h),$$

which proves the proposition.  $\square$

**REMARK 3** Our consistency result sheds light on the relationship between the kernel  $K_{\phi}$  and the evolution equation (4). Proving convergence of a Bence Merriman Osher type algorithm in our context seems to be very difficult (if true at all). The argument of [32] does not apply here. The main difficulty is that  $G_h^{\pm} \varphi$  may not be continuous, even if  $\varphi$  is regular. Further, we can only show monotonicity of the operators  $G_h^{\pm}$  up to  $o(h)$  for smooth functions whose gradients do not vanish. The source of these difficulties is really the thresholding in the definition of  $G_h^{\pm}$ .

## 5. Numerical simulations

In the previous section, we proved a consistency result for a Bence Merriman Osher-type algorithm. Here we numerically investigate the convergence properties of a related scheme, based on a phase-field discretization. Both schemes consist in a diffusion step followed by a correction step. In the case of the BMO scheme, the correction is a simple thresholding, while the correction is obtained via a reaction term for the phase field scheme. More precisely, in the second case, given a small parameter  $\epsilon > 0$ , we set

$$G_{h,\epsilon}\varphi(x) = T_{h,\epsilon}(K_{\phi,h} * \varphi),$$

where  $T_{h,\epsilon}$  is defined as follows: Given  $\lambda \in \mathbb{R}$ ,  $T_{h,\epsilon}(\lambda) = \psi(\lambda)$  where  $\psi$  is the solution of the ODE

$$\begin{cases} \psi_t &= -\frac{1}{\epsilon^2}W'(\psi), \\ \psi(0) &= \lambda, \end{cases}$$

and  $W$  a double well potential with wells located at  $\psi = 0$  and  $\psi = 1$ . Note that if  $\varphi = \mathbb{1}_\Omega$  is a characteristic function, then

$$\lim_{\epsilon \rightarrow 0} G_{h,\epsilon}\mathbb{1}_\Omega = G_h^+\mathbb{1}_\Omega = G_h^-\mathbb{1}_\Omega,$$

which shows a formal relationship in the correction step of the BMO and phase field schemes.

The advantage of the phase field scheme, is that it produces smoother interfaces, which avoids numerical errors due to aliasing. Moreover, we wanted to test our method for approximating anisotropic diffusion on computations of Wulff shapes, a problem where one has to impose a volume constraint. Such constraint is easier to handle with a phase field scheme, where one can explicitly compute the associated Lagrange multiplier. The next paragraph, describes the phase-field algorithm for the operator  $\tilde{\Delta}_\phi$ .

### 5.1 The $\tilde{\Delta}_\phi$ -phase field model and its discretization

As an approximation to the anisotropic Allen–Cahn equation (5), we consider the following phase-field model

$$\begin{cases} u_t = \tilde{\Delta}_\phi u - \frac{1}{\epsilon^2}W'(u), \\ u(x, 0) = q\left(\frac{\text{dist}(x, \partial E)}{\epsilon}\right). \end{cases} \quad (44)$$

We also report tests, where we estimate the  $L^1$ -error on anisotropic Wulff sets (the sets which minimize the anisotropic perimeter under a volume constraint). To impose volume conservation, we consider a conserved phase-field model, of the form

$$\begin{cases} u_t(x, t) = \tilde{\Delta}_\phi u(x, t) - \frac{1}{\epsilon^2}W'(u(x, t)) + \frac{1}{\epsilon}\lambda(t)\sqrt{2W(u(x, t))}, \\ u(x, 0) = q\left(\frac{\text{dist}(x, \Omega_0)}{\epsilon}\right). \end{cases} \quad (45)$$

The parameter

$$\lambda(t) = \frac{\int_{\mathbb{R}^d} W'(u(x, t)) dx}{\epsilon \int_{\mathbb{R}^d} \sqrt{2W(u(x, t))} dx},$$

can be seen as a Lagrange multiplier, which preserves the mass of  $u$ . See [13] where schemes of this form have been studied for isotropic mean curvature with a volume constraint.

We now describe the numerical method we use for solving the PDE's (44) and (45). Several studies of classical numerical schemes for the Allen–Cahn equation have already been conducted in the past: see for instance, [16, 18, 24, 28, 29, 39, 40]. Here, the computational domain is the fixed box  $Q = [-1/2, 1/2]^d \subset \mathbb{R}^d$ ,  $d = 2, 3$ . The initial datum is  $u_0 = q(\frac{\text{dist}(x, \partial\Omega)}{\epsilon})$ , where  $\Omega_0$  is a smooth bounded set strictly contained  $Q$ . We assume that during the evolution, the set  $\Omega_{\epsilon, t} := \{u_\epsilon(x, t) \geq 1/2\}$  remains strictly inside  $Q$ , so that we may impose periodic boundary conditions on  $\partial Q$ .

Our strategy consists in representing  $u$  as a Fourier series in  $Q$ , and in using a splitting method. First, one applies the diffusion operator, which given the form of  $\tilde{\Delta}_\phi$ , merely amounts to a multiplication in the Fourier space. The interesting feature of our approach is that this step is fast and very accurate. Next, the reaction term is applied.

More precisely,  $u_\epsilon(x, t_n)$  at time  $t_n = t_0 + n\delta t$  is approximated by

$$u_\epsilon^P(x, t_n) = \sum_{\max_{1 \leq i \leq d} |p_i| \leq P} u_{\epsilon, p}(t_n) e^{2i\pi p \cdot x}.$$

In the diffusion step, we set

$$u_\epsilon^P(x, t_n + 1/2) = \sum_{\max_{1 \leq i \leq d} |p_i| \leq P} u_{\epsilon, p}(t_n) e^{-4\pi^2 \delta t \phi''(p)^2} e^{2i\pi p \cdot x}.$$

We then integrate the reaction terms

$$u_\epsilon^P(x, t_n + 1) = u_\epsilon^P(x, t_n + 1/2) - \delta t \epsilon^2 W'_{i, \epsilon}(u_\epsilon^P(x, t_n + 1/2)).$$

In practice, the first step is performed via a fast Fourier transform, with a computational cost  $O(P^d \ln(P))$ .



The corresponding numerical scheme turns out to be stable when solving (44), under the condition  $\delta t \leq M\epsilon^2$ , where  $M = \left[ \sup_{t \in [0, 1]} \{W''(t)\} \right]^{-1}$ . Numerically, we observed that this condition is also sufficient for the conserved potential in (45). In the simulations, we used  $W(s) = \frac{1}{2}s^2(1-s)^2$ .

The isotropic version of our splitting scheme has been studied in [13]. It is shown there that this scheme converges with the same rate as phase-field approximations based on a spatial discretization by finite differences or by finite elements. Its advantages are greater precision, and unconditional stability.

## 5.2 Test of convergence in dimension 2

We consider following anisotropic densities

$$\begin{aligned} \phi_1^o(\xi) &= \|\xi\|_{\ell^4} = (|\xi_1|^4 + |\xi_2|^4)^{\frac{1}{4}}, \\ \phi_2^o(\xi) &= \|\xi\|_{\ell^{\frac{4}{3}}} = \left( |\xi_1|^{\frac{4}{3}} + |\xi_2|^{\frac{4}{3}} \right)^{\frac{3}{4}}, \\ \phi_3^o(\xi) &= \left( |\xi_1|^{1.001} + \frac{1}{2}\xi_1 + \frac{\sqrt{3}}{2}|\xi_2|^{1.001} + \left| \frac{1}{2}\xi_1 - \frac{\sqrt{3}}{2}\xi_2 \right|^{1.001} \right)^{\frac{1}{1.001}}. \end{aligned}$$

FIG. 2. Wulff Set (black) and Frank diagram (grey) for the anisotropic densities  $(\phi_1, \phi_1^o)$ ,  $(\phi_2, \phi_2^o)$  and  $(\phi_3, \phi_3^o)$ FIG. 3.  $\Omega(t)$  at different times for the anisotropic densities  $\phi_1, \phi_2, \phi_3$ 

See Figure 2 for a representation of their Wulff sets  $B_{\phi_i}$  and Frank diagrams  $B_{\phi_i^o}$ .

### 1. Evolution from a Wulff set.

We consider the equation

$$\begin{cases} \partial_t u &= \tilde{\Delta}_\phi u - \frac{1}{\epsilon^2} W'(u), \\ u(0, x) &= q(\text{dist}(x, \Omega_0)/\epsilon), \end{cases}$$

where the initial set  $\Omega_0$  is a Wulff set of radius  $R_0 = 0.25$

$$\Omega_0 = \{x \in \mathbb{R}^2; \phi(x) \leq R_0\}.$$

It is well known that the set  $\Omega(t)$  obtained from  $\Omega_0$  through evolution by anisotropic mean curvature is a Wulff set with radius  $R(t) = \sqrt{R_0^2 - 2t}$ , which decreases to a point at the extinction time  $t_{ext} = \frac{R_0^2}{2}$ . In these simulations, the number of Fourier modes is  $P = 2^8$ , and the time step and phase-field parameter are chosen to be  $\delta_t = 1/P^2$  and  $\epsilon = 1/P$ . In Figure 3 the interface  $\Omega(t)$  is plotted at different times. We observe a good agreement between the theoretical and computed curves, in spite of the smoothing of the corners of the latter.

### 2. Convergence to the Wulff set

This smoothing of corners actually depends on the thickness  $\epsilon$  of the diffuse interface, as




FIG. 4. From left to right:  $\Omega(t)$  at different times with anisotropy  $\phi_1^o$ ,  $\Omega(t)$  at different times with anisotropy  $\phi_2^o$ , error estimate  $\epsilon \rightarrow \|\mathbb{1}_{B_\phi^\epsilon} - \mathbb{1}_{B_{\phi, R_0}}\|_{L^1(\mathbb{R}^d)}$  in logarithmic scale ( $\phi_1^o$  in red and  $\phi_2^o$  in blue)

evidenced in the next series of tests, of evolution by anisotropic mean curvature under a volume constraint according to (45). The initial set  $\Omega_0$  is a circle centered at 0, of the same volume as  $\Omega^* = \{x \in \mathbb{R}^d ; \phi(x) < R_0\}$ . The evolution  $\Omega_t$  from  $\Omega_0$  is expected to converge to the Wulff set  $\Omega^*$ .

The left and middle of Figure 4 represent the final sets  $\Omega_\epsilon^*$  obtained from the resolution of anisotropic Allen–Cahn equation, with respective anisotropic densities  $\phi_1$  and  $\phi_2$ , and for different value of  $\epsilon$ . We observe that the smaller  $\epsilon$ , the better the approximation of the Wulff set. On the right-hand side of Figure 4, the  $L^1$  error

$$\epsilon \rightarrow \|\mathbb{1}_{\Omega_\epsilon^*} - \mathbb{1}_{\Omega^*}\|_{L^1(\mathbb{R}^d)},$$

is plotted in a logarithmic scale. This graph indicates that this error is of order  $\epsilon$ .

### 5.3 Some 3D simulations

As final illustrations, we consider the anisotropic densities

$$\begin{cases} \phi_4^o(\xi) &= \sqrt{\xi_1^2 + \xi_2^2} + |\xi_3|, \\ \phi_5^o(\xi) &= |\xi_1| + |\xi_2| + |\xi_3|. \end{cases}$$

The corresponding Wulff sets and Frank diagrams are plotted in Figure 5.

We report in Figure 6 (respectively in Figure 7) the evolution by  $\phi_4^o$  (resp.  $\phi_5^o$ ) anisotropic mean curvature from an initial torus. The number of Fourier modes is  $P = 2^7$ , the time step and diffuse interface thickness are  $\delta_t = 1/P^2$  and  $\epsilon = 1/P$ .






FIG. 5. Frank diagram and Wulff set:  $B_{\phi_4^o}$ ,  $B_{\phi_4}$ ,  $B_{\phi_5^o}$ ,  $B_{\phi_5}$



FIG. 6.  $\phi_4^0(\xi)$ -evolution from an initial torus, at different timesFIG. 7.  $\phi_5^0(\xi)$ -evolution from an initial torus, at different times

## REFERENCES

1. ALLEN, S. M., & CAHN, J. W. A microscopic theory for antiphase boundary motion and its application to antiphase domain coarsening. *Acta Metall.* **27** (1979), 1085–1095.
2. AMBROSIO, L. Geometric evolution problems, distance function and viscosity solutions, *Calculus of variations and partial differential equations* (Pisa, 1996), Springer, Berlin, 2000, pp. 5–93. [Zb10956.35002 MR1757696](#)
3. BARLES, G. *Solutions de viscosité des équations de Hamilton-Jacobi*, Mathématiques & Applications (Berlin) [Mathematics & Applications], vol. 17, Springer-Verlag, Paris (1994). [Zb10819.35002 MR1613876](#)
4. BARLES, G., & GEORGELIN, C. A simple proof of convergence for an approximation scheme for

- computing motions by mean curvature. *SIAM J. Numer. Anal.* **32** (1995), 484–500. [Zb10831.65138](#) [MR1324298](#)
5. BARLES, G., SONER, H. M., & SOUGANIDIS, P. E. Front propagation and phase field theory. *SIAM J. Control Optim.* **31** (1993), 439–469. [Zb10785.35049](#) [MR1205984](#)
  6. BARLES, G., & SOUGANIDIS, P. E. Convergence of approximation schemes for fully nonlinear second order equations. *Asymptotic Anal.* **4** (1991), 271–283. [Zb10729.65077](#) [MR1115933](#)
  7. BARRETT, J. W., GARCKE, H., & NÜRNBERG, R. On the parametric finite element approximation of evolving hypersurfaces in  $\mathbb{R}^3$ . *J. Comput. Phys.* **227** (2008), 4281–4307. [Zb11145.65068](#) [MR2406538](#)
  8. BARRETT, J. W., GARCKE, H., & NÜRNBERG, R. A variational formulation of anisotropic geometric evolution equations in higher dimensions. *Numer. Math.* **109** (2008), 1–44. [Zb11149.65082](#) [MR2377611](#)
  9. BELLETTINI, G., & PAOLINI, M. Quasi-optimal error estimates for the mean curvature flow with a forcing term. *Differential Integral Equations* **8** (1995), 735–752. [Zb10820.49019](#) [MR1306590](#)
  10. BELLETTINI, G., & PAOLINI, M. Anisotropic motion by mean curvature in the context of Finsler geometry. *Hokkaido Math. J.* **25** (1996), 537–566. [Zb10873.53011](#) [MR1416006](#)
  11. BENCE, J., MERRIMAN, B., & OSHER, S. Diffusion generated motion by mean curvature. In: *Computational Crystal Growers Workshop, J. Taylor ed. Selected Lectures in Math., Amer. Math. Soc.* (1992), 73–83.
  12. BERGH, J., & LÖFSTRÖM, J. *Interpolation spaces. An introduction.* Springer, Berlin (1976). [Zb10344.46071](#) [MR0482275](#)
  13. BRASSEL, M., & BRETIN, E. A modified phase field approximation for mean curvature flow with conservation of the volume. *Math. Methods Appl. Sci.* **34** (2011), 1157–1180. [Zbl pre05914672](#) [MR2838769](#)
  14. BRETIN, E. Méthode de champ de phase et mouvement par courbure moyenne, Ph.D. thesis, Institut national polytechnique de Grenoble, 2009.
  15. CHAMBOLLE, A., & NOVAGA, M. Convergence of an algorithm for the anisotropic and crystalline mean curvature flow. *SIAM J. Math. Anal.* **37** (2006), 1978–1987. [Zb11116.35074](#) [MR2213402](#)
  16. CHEN, L., & SHEN, J. Applications of semi-implicit Fourier-spectral method to phase field equations. *Comput. Phys. Commun.* **108** (1998), 147–158. [Zb11017.65533](#)
  17. CHEN, X. Generation and propagation of interfaces for reaction-diffusion equations. *J. Differential Equations* **96** (1992), 116–141. [Zb10765.35024](#) [MR1153311](#)
  18. CHEN, X., ELLIOTT, C. M., GARDINER, A., & ZHAO, J. J. Convergence of numerical solutions to the Allen–Cahn equation. *Appl. Anal.* **69** (1998), 47–56. [Zb10992.65096](#) [MR1708186](#)
  19. CHEN, Y. G., GIGA, Y., & GOTO, S. Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. *Proc. Japan Acad. Ser. A Math. Sci.* **65** (1989), 207–210. [Zb10735.35082](#) [MR1030181](#)
  20. CHEN, Y. G., GIGA, Y., & GOTO, S. Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. *J. Differential Geom.* **33** (1991), 749–786. [Zb10696.35087](#) [MR1100211](#)
  21. CRANDALL, M. G., ISHII, H., & LIONS, P. L. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)* **27** (1992), 1–67. [Zbl 0755.35015](#) [MR1118699](#)
  22. DA LIO, F., FORCADEL, N., & MONNEAU, R. Convergence of a non-local eikonal equation to anisotropic mean curvature motion. Application to dislocation dynamics. *J. Eur. Math. Soc. (JEMS)* **10** (2008), 1061–1104. [Zb11158.35324](#) [MR2443929](#)
  23. DECKELNICK, K., & DZIUK, G. Discrete anisotropic curvature flow of graphs. *M2AN Math. Model. Numer. Anal.* **33** (1999), 1203–1222. [Zb10948.65138](#) [MR1736896](#)
  24. DECKELNICK, K., DZIUK, G., & ELLIOTT, C. M. Computation of geometric partial differential equations and mean curvature flow. *Acta Numer.* **14** (2005), 139–232. [Zb11113.65097](#) [MR2168343](#)
  25. EVANS, L. C. Convergence of an algorithm for mean curvature motion, *Indiana University mathematics journal* **42** (1993), 553–557. [Zb10802.65098](#) [MR1237058](#)

26. EVANS, L. C., SONER, H. M., & SOUGANIDIS, P. E. Phase transitions and generalized motion by mean curvature. *Comm. Pure Appl. Math.* **45** (1992), 1097–1123. [Zb10801.35045](#) [MR1177477](#)
27. EVANS, L. C., & SPRUCK, J. Motion of level sets by mean curvature. I, *J. Differential Geom.* **33** (1991), 635–681. [Zb10726.53029](#) [MR1100206](#)
28. GARCKE, H., NESTLER, B., & STOTH, B., Anisotropy in multiphase systems: A phase-field approach. *Interfaces and Free Boundaries* **1** (1999), 175–198. [Zb10959.35169](#) [MR1867130](#)
29. GARCKE, H., NESTLER, B., & STOTH, B. A multi phase field concept: Numerical simulations of moving phase boundaries and multiple junctions, *SIAM J. Appl. Math.* **60** (1999), 295–315. [Zb10942.35095](#) [MR1740846](#)
30. IGEL'FAND, I. M., & SHILOV, G. E. *Generalized Functions*, vol. 2. Academic Press, New York [Harcourt Brace Jovanovich Publishers], Spaces of fundamental and generalized functions, Translated from the Russian by Morris D. Friedman, Amiel Feinstein and Christian P. Peltzer., (1968 [1977]). [Zb10159.18301](#) [MR0230128](#)
31. GRISVARD, P. *Elliptic Problems in Nonsmooth Domains*, vol. 24 of Monographs and Studies in Mathematics. Pitman, Boston, MA (Advanced Publishing Program) (1985). [Zb10695.35060](#) [MR0775683](#)
32. ISHII, H., PIRES, G. E., & SOUGANIDIS, P. E. Threshold dynamics type approximation schemes for propagating fronts. *J. Math. Soc. Japan* **51** (1999), 267–308. [Zb10935.53006](#) [MR1674750](#)
33. MODICA, L., & MORTOLA, S. Il limite nella  $\Gamma$ -convergenza di una famiglia di funzionali ellittici. *Boll. Un. Mat. Ital. A (5)* **14** (1977), 526–529. [Zb10364.49006](#) [MR0473971](#)
34. MODICA, L., & MORTOLA, S. Un esempio di  $\Gamma$ -convergenza, *Boll. Un. Mat. Ital. B (5)* **14** (1977), 285–299. [Zb10356.49008](#) [MR0445362](#)
35. OSHER, S., & FEDKIW, R. *Level Set Methods and Dynamic Implicit Surfaces*, Springer-Verlag, New York, Applied Mathematical Sciences (2003). [Zb11026.76001](#) [MR1939127](#)
36. OSHER, S., & PARAGIOS, N. *Geometric Level Set Methods in Imaging, Vision and Graphics*, Springer-Verlag, New York (2003). [Zb11027.68137](#)
37. OSHER, S., & SETHIAN, J. A. Fronts propagating with curvature-dependent speed: Algorithms based on Hamilton-Jacobi formulations. *J. Comput. Phys.* **79** (1988), 12–49. [Zb10659.65132](#) [MR0965860](#)
38. PAOLINI, M. An efficient algorithm for computing anisotropic evolution by mean curvature, Curvature flows and related topics (Levico, 1994), GAKUTO Internat. Ser. Math. Sci. Appl., vol. 5, Gakkōtoshō, Tokyo, 1995, pp. 199–213. [Zb10838.73079](#) [MR1365309](#)
39. PAOLINI, M. A quasi-optimal error estimate for a discrete singularly perturbed approximation to the prescribed curvature problem. *Math. Comp.* **66** (1997), 45–67. [Zb10854.35008](#) [MR1361810](#)
40. PAOLINI, M., & VERDI, C. Asymptotic and numerical analyses of the mean curvature flow with a space-dependent relaxation parameter. *Asymptotic Anal.* **5** (1992), 553–574. [Zb10757.65078](#) [MR1169358](#)
41. RUUTH, S. J. Efficient algorithms for diffusion-generated motion by mean curvature. *J. Comput. Phys.* **144** (1998), 603–625. [Zb10946.65093](#) [MR1638032](#)
42. RUUTH, S. J., & MERRIMAN, B. Convolution-generated motion and generalized Huygens' principles for interface motion. *SIAM J. Appl. Math.* **60** (2000), 868–890. [Zb10958.65021](#) [MR1740854](#)