

On the regularity of the interface of a thermodynamically consistent two-phase Stefan problem with surface tension

JAN PRÜSS

Institut für Mathematik, Martin-Luther-Universität Halle-Wittenberg, 60120 Halle, Germany

E-mail: jan.pruess@mathematik.uni-halle.de

YUANZHEN SHAO

Department of Mathematics, Vanderbilt University, Nashville, TN 37240, USA

E-mail: yuanzhen.shao@vanderbilt.edu

GIERI SIMONETT

Department of Mathematics, Vanderbilt University, Nashville, TN 37240, USA

E-mail: gieri.simonett@vanderbilt.edu

[Received 16 December 2014 and in revised form 28 September 2015]

We study the regularity of the free boundary arising in a thermodynamically consistent two-phase Stefan problem with surface tension by means of a family of parameter-dependent diffeomorphisms, L_p -maximal regularity theory, and the implicit function theorem.

2010 Mathematics Subject Classification: Primary 35R35, 82C26, 35K55, 35B65

Keywords: Free boundary problems; phase transitions; the Stefan problem; regularity of moving interfaces; real analytic solutions; maximal regularity; the implicit function theorem.

1. Introduction

The main objective of this article is to develop a technique relying on a family of parameter-dependent diffeomorphisms, maximal regularity theory, and the implicit function theorem to prove regularity of moving interfaces occurring in free boundary problems. As an application, we prove that the moving interface in a thermodynamically consistent two-phase Stefan problem is jointly C^k -smooth in time and space, for $k \in \mathbb{N} \cup \{\infty, \omega\}$ with ω being the symbol of real analyticity, as long as several physical quantities, that is, the coefficients of the heat conductivity, kinetic undercooling, and the free energy, enjoy appropriate regularity assumptions.

The idea of establishing regularity of solutions to differential equations by means of the implicit function theorem in conjunction with a translation argument was first introduced by S.B. Angenent in [8] to prove analyticity of the free boundary in one dimensional porous medium equations, and has proved itself a useful tool later in many publications. See for example [9, 27, 28, 50, 58]. More precisely, to study the regularity of the solution to a differential equation, one introduces parameters representing translation in space and time into the solution to the given differential equation. Then one studies the parameter-dependent equation satisfied by this transformed solution. The implicit function theorem yields the smooth dependence upon the parameters of the solution to the parameter-dependent problem. This regularity property is then inherited by the original solution. An advantage of this technique is reflected by its power to prove analyticity of solutions to differential equations, which cannot be attained through the classical method of bootstrapping.

A well-known approach to free boundary problems is to transform the original problem with a moving boundary, or separating interface, which we denote by $\Gamma(t)$, into one with a fixed reference manifold Σ by means of the Hanzawa transformation, see [39]. Then the problem of establishing the regularity of the free boundary $\Gamma(t)$ is transferred into establishing the regularity of the height function parameterizing $\Gamma(t)$ over Σ . However, applying the aforementioned translation technique to the height function on the surface Σ causes an essential challenge, considering for instance the usual translation $(t, x) \mapsto (t + \lambda, x + \mu)$, because of the global nature of these translations. Hence we desire an alternative that only shifts the variables “locally”. The idea of localizing translations was first introduced by J. Escher, J. Prüss and G. Simonett in [26] to study regularity of solutions to elliptic and parabolic equations in Euclidean space. The basic building block of [26] consists of rescaling translations by some cutoff function. This technique was later generalized in Y. Shao [73] to introduce a family of parameter-dependent diffeomorphisms acting on functions or tensor fields on Riemannian manifolds by means of a smooth atlas in order to study the regularity of solutions to geometric evolution equations. But in view of the physical quantities in the bulk phases adjacent to $\Gamma(t)$, e.g., the temperature function in the case of phase transitions, or the velocity and the pressure field in the case of two-phase fluids, we need introduce a localized translation not only on the fixed reference surface Σ , but also in a neighborhood of Σ . This adds one more degree of complexity to the aforementioned technique for geometric flows. In Section 3, we will build up a complete theory of parameter-dependent diffeomorphisms for free boundary problems.

Free boundary problems form an important field of applied analysis. They deal with solving partial differential equations in a given domain, a part of whose boundary is a priori unknown. That portion of the boundary is called the free boundary or the moving boundary. In addition to the standard boundary conditions that are needed in order to solve the prescribed partial differential equations, an additional condition must be imposed at the free boundary. One then seeks to determine both the free boundary and the solution of the differential equation. This field has drawn great attention over decades due to its applications to physics, chemistry, medicine, material science, and so forth.

Great progress has been seen in the studies of regularity of free boundaries during the past half century. Many mathematicians have made contributions to the theory of free boundary problems, among them H.W. Alt, H. Berestycki, L.A. Caffarelli, A. Friedman, D. Kinderlehrer, L. Nirenberg, J. Spruck, G. Stampacchia [2, 3, 12, 14, 16, 43–46], E. DiBenedetto [24]. We also refer the reader to [33, 46, 67–69] for a historical account of the field.

The Stefan problem, arguably, is the most studied free boundary problem, with over 1200 mathematical publications devoted to the topic. It was first introduced in 1889 by J. Stefan. We refer the reader to the books [53, 71, 77] for further information. The Stefan problem describes phase transitions in liquid-solid systems and accounts for heat diffusion and exchange of latent heat in a homogeneous medium, wherein the liquid and solid phases are separated by a closed moving interface $\Gamma(t)$. The basic physical law governing this process is conservation of energy. To be more precise, let $\Omega \subset \mathbb{R}^{m+1}$ be a bounded domain of class C^2 , $m \geq 1$. Ω is occupied by a material that can undergo phase changes: at time t , phase i occupies the subdomain $\Omega_i(t)$ of Ω , respectively, with $i = 1, 2$. We assume that $\partial\Omega_1(t) \cap \partial\Omega = \emptyset$; this means that no *boundary contact* can occur. The closed compact hypersurface $\Gamma(t) := \partial\Omega_1(t) \subset \Omega$ forms the interface between the phases. Then the *Stefan problem with surface tension*, or the *Stefan problem with Gibbs–Thomson correction*, can

be formulated as follows:

$$\left\{ \begin{array}{ll} \partial_t \vartheta - d \Delta \vartheta = 0 & \text{in } \Omega \setminus \Gamma(t), \\ \partial_{\nu_{\partial\Omega}} \vartheta = 0 & \text{on } \partial\Omega, \\ V(t) - \llbracket d \partial_{\nu_\Gamma} \vartheta \rrbracket = 0 & \text{on } \Gamma(t), \\ \vartheta + \sigma \mathcal{H} = 0 & \text{on } \Gamma(t), \\ \vartheta(0) = \vartheta_0, \quad \Gamma(0) = \Gamma_0. \end{array} \right. \quad (1.1)$$

Here $\vartheta(t) = \vartheta_1(t)\chi_{\Omega_1(t)} + \vartheta_2(t)\chi_{\Omega_2(t)}$, where ϑ_i denotes the relative temperature distribution in phase i , $\nu_\Gamma(t)$ the outer normal field of $\partial\Omega_1(t)$, $V(t)$ the normal velocity of $\Gamma(t)$, $\mathcal{H}(t) = \mathcal{H}(\Gamma(t)) = -\text{div}_{\Gamma(t)} \nu_\Gamma(t)/m$ the mean curvature of $\Gamma(t)$, and $\llbracket v \rrbracket = v_2|_{\Gamma(t)} - v_1|_{\Gamma(t)}$ the jump of a quantity v across $\Gamma(t)$. The sign of the mean curvature \mathcal{H} is chosen to be negative at a point $x \in \Gamma$ if $\Omega_1 \cap \mathbb{B}(x, r)$ is convex for some sufficiently small $r > 0$. Thus if Ω_1 is a ball of radius R then $\mathcal{H} = -1/R$ for its boundary Γ . The function d agrees with a positive constant d_i in $\Omega_i(t)$, which stand for the heat conductivities in different phases. The condition that $V(t) = \llbracket d \partial_\nu \vartheta \rrbracket$ is caused by the law of conservation of energy and is usually called the *Stefan condition*.

If a pair (ϑ, Γ) solves the system of equations (1.1), Γ is called the free boundary to the Stefan problem. In the case that the condition $\vartheta + \sigma \mathcal{H} = 0$ is replaced by

$$\vartheta = 0 \quad \text{on } \Gamma(t), \quad (1.2)$$

i.e., if $\sigma = 0$, the resulting problem is usually referred to as the *classical Stefan problem*. Here we mention the monographs by B. Chalmers [20, Chapter 1], P. Hartman [40], A. Visintin [77], and the research papers by G. Caginalp [19], M. E. Gurtin [34–36], J. S. Langer [48], W. W. Mullins and R. F. Sekerka [56, 57], A. Visintin [76], where this law has been motivated and derived based on various mathematical and physical principles. As has been explained in [65, Section 1], see also [64], condition (1.2) can be understood as a first order approximation of the relation

$$\llbracket \psi(\theta) \rrbracket + \sigma \mathcal{H} = 0 \quad \text{on } \Gamma(t) \quad (1.3)$$

around the melting temperature, where θ is the absolute temperature and ψ is the free energy of the system.

In the absence of one of ϑ_i , that is, one of ϑ_i is identically equal to some constant, the problem is called the one-phase Stefan problem, otherwise it is alluded to as the two-phase Stefan problem.

In the one-dimensional case, different aspects of the *classical Stefan problem* were extensively studied before the 1980s. In higher dimensions, global existence and uniqueness of weak solutions was first established by S.L. Kamenomostskaja in [42], see also A. Friedman [30]. Existence of local classical solutions was obtained by E.I. Hanzawa [39] and A.M. Meřmanov [52], provided that the initial data are smooth enough and satisfy some higher order compatibility conditions.

In the case of the one-phase *classical Stefan problem*, regularity of the free boundary for weak solutions is studied in [14, 15, 32, 43, 44]. The formulation of the problem as a parabolic variational inequality was initiated by G. Duvaut in [25]. After that, it has been applied in [32] to get the Lipschitz continuity of the free boundary under some conditions on the initial geometry and the given data. Later in a paper of L.A. Caffarelli [14], the author proved that the free boundary is C^1 in space and time and the temperature is C^2 up to the free boundary near a density point for the states set, namely the solid phase. In particular, if the free boundary is Lipschitz continuous

then it is actually C^1 . In [15], the author identified conditions to guarantee a locally Lipschitz free boundary. Almost at the same time, D. Kinderlehrer and L. Nirenberg [43] established via the partial hodograph and Legendre transformations in conjunction with a variational inequality argument smoothness of the free boundary as well as the solution. The conditions in [43] can be verified by the results in [14]. In a subsequent paper [44], the same authors showed under some integral conditions by the same technique that the free boundary and the solution are analytic in the space variables and in the second Gevrey class for the time variable. Regularization of the free boundary for large time was obtained by H. Matano in [51], i.e., any weak solution eventually becomes smooth. Continuity of the temperature distribution for weak solutions is proved in [18].

Regularity of the free boundary for weak solutions or viscosity solutions has been explored in [10, 11, 47] for the multi-dimensional two-phase Stefan problem based on a non-degeneracy condition. The non-degeneracy condition states, roughly speaking, that the heat fluxes are not vanishing simultaneously on the free boundary, which is automatically satisfied when the free boundary is regular enough, say belongs to the Hölder class of order more than 1. Assuming that the free boundary is locally Lipschitz continuous, it was proven in [10] that the solution is classical, i.e., the free boundary and the viscosity solution are C^1 . Later the authors showed that the non-degeneracy condition can be replaced by some flatness condition. In the presence of the non-degeneracy condition, H. Koch verified that the C^1 free boundary and the temperature are in fact smooth. Through a different approach by imposing a mild regularity condition on the initial data, it was shown by J. Prüss, J. Saal and G. Simonett in [58] that the free boundary and the temperature are analytic under the assumption that the free boundary can be expressed as the graph of some function. Continuity of the temperature distribution for weak solutions was obtained in [17, 23, 24, 78].

Although the Stefan problem with the *Gibbs–Thomson correction* (1.2) has been around for many decades, only few analytical results concerning existence of solutions can be found in the literature. A. Friedman and F. Reitich [31] considered the case with small surface tension $0 < \sigma \ll 1$ and linearized the problem about $\sigma = 0$. Assuming the existence of a smooth solution for the case $\sigma = 0$, that is, for the *classical Stefan problem*, the authors proved existence and uniqueness of a weak solution for the *linearized* problem and then investigated the effect of small surface tension on the shape of $\Gamma(t)$. Existence of long time weak solutions was first established by Luckhaus [49], using a discretized problem and a capacity-type estimate for approximating solutions. The weak solutions obtained have a sharp interface, but are highly non-unique. See also Röger [70], and Almgren and Wang [1]. Existence of classical solutions, but without uniqueness, was proved by E.V. Radkevich in [66]. In A.M. Meirmanov [54], the way in which a spherical ball of ice in a supercooled fluid melts down was investigated. It was presented by J. Escher, J. Prüss and G. Simonett in [27] that there exists a unique local strong solution and the free boundary immediately regularizes to be analytic in space and time provided that the initial data satisfy some mild regularity assumptions. This result is based on the assumption that the free boundary is given by the graph of some function. In J. Prüss and G. Simonett [59], linearized stability and instability of equilibria was investigated. Nonlinear stability results were obtained by M. Hadžić, Y. Guo [37, 38] and J. Prüss, G. Simonett and R. Zacher [65], see also the monograph [64].

In this paper we consider a general model for phase transitions, formulated in [65], that is thermodynamically consistent, see [7] and [41] for related work. It involves the thermodynamic quantities of absolute temperature, free energy, internal energy, and entropy, and is complemented by constitutive equations for the free energies and the heat fluxes in the bulk regions. An important

assumption is that there be no entropy production on the interface. In particular, the interface is assumed to carry no mass and no energy except for surface tension.

To be more precise, we choose Ω , $\Omega_i(t)$ and $\Gamma(t)$ as in (1.1). By the thermodynamically consistent two-phase *Stefan problem with surface tension*, possibly with kinetic undercooling, we mean the following problem: find a family of closed compact hypersurfaces $\{\Gamma(t)\}_{t \geq 0}$ contained in Ω and an appropriately smooth function $\theta : \mathbb{R}_+ \times \bar{\Omega} \rightarrow \mathbb{R}$ such that

$$\left\{ \begin{array}{ll} \kappa(\theta)\partial_t\theta - \operatorname{div}(d(\theta)\nabla\theta) = 0 & \text{in } \Omega \setminus \Gamma(t), \\ \partial_{\nu_\Omega}\theta = 0 & \text{on } \partial\Omega, \\ \llbracket\theta\rrbracket = 0 & \text{on } \Gamma(t), \\ \llbracket\psi(\theta)\rrbracket + \sigma\mathcal{H} = \gamma(\theta)V & \text{on } \Gamma(t), \\ \llbracket d(\theta)\partial_{\nu_\Gamma}\theta\rrbracket = (l(\theta) - \gamma(\theta)V)V & \text{on } \Gamma(t), \\ \theta(0) = \theta_0, \quad \Gamma(0) = \Gamma_0. & \end{array} \right. \quad (1.4)$$

Here θ denotes the (absolute) temperature. Several quantities are derived from the free energies $\psi_i(\theta)$ as follows:

- $\epsilon_i(\theta) = \psi_i(\theta) + \theta\eta_i(\theta)$, the internal energy in phase i ,
- $\eta_i(\theta) = -\psi'_i(\theta)$, the entropy,
- $\kappa_i(\theta) = \epsilon'_i(\theta) = -\theta\psi''_i(\theta) > 0$, the heat capacity,
- $l(\theta) = \theta\llbracket\psi'(\theta)\rrbracket = -\theta\llbracket\eta(\theta)\rrbracket$, the latent heat.

Furthermore, $d_i(\theta) > 0$ denotes the coefficient of heat conduction in Fourier’s law, $\gamma(\theta) \geq 0$ the coefficient of kinetic undercooling. As is commonly done, we assume that there exists a unique (constant) *melting temperature* θ_m , characterized by the equation $\llbracket\psi(\theta_m)\rrbracket = 0$. Finally, system (1.4) is to be completed by constitutive equations for the free energies ψ_i in the bulk phases $\Omega_i(t)$.

In the sequel we drop the index i , as there is no danger of confusion; we just keep in mind that the coefficients depend on the phases. The temperature is assumed to be continuous across the interface, as indicated by the condition $\llbracket\theta\rrbracket = 0$ in (1.4). However, the free energy and the heat conductivities depend on the respective phases, and hence the jumps $\llbracket\psi(\theta)\rrbracket$, $\llbracket\kappa(\theta)\rrbracket$, $\llbracket\eta(\theta)\rrbracket$, $\llbracket d(\theta)\rrbracket$ are in general non-zero at the interface. We assume that the coefficient of surface tension is constant.

In this paper, we will prove the following regularity result for $k \in \mathbb{N} \cup \{\infty, \omega\}$.

THEOREM 1.1 ($\gamma \equiv 0$). Let $p > m + 3$, $\gamma = 0$, $\sigma > 0$. Suppose that $d_i \in C^{k+2}(0, \infty)$, $\psi_i \in C^{k+3}(0, \infty)$ for $i = 1, 2$ such that

$$\kappa_i(u) = -u\psi''_i(u) > 0, \quad d_i(u) > 0, \quad u \in (0, \infty).$$

Assume the *regularity conditions*

$$\theta_0 \in W_p^{2-2/p}(\Omega \setminus \Gamma_0) \cap C(\bar{\Omega}), \quad \theta_0 > 0, \quad \Gamma_0 \in W_p^{4-3/p},$$

the *compatibility conditions*

$$\partial_{\nu_\Omega}\theta_0 = 0, \quad \llbracket\psi(\theta_0)\rrbracket + \sigma\mathcal{H}(\Gamma_0) = 0, \quad \llbracket d(\theta_0)\partial_{\nu_{\Gamma_0}}\theta_0\rrbracket \in W_p^{2-6/p}(\Gamma_0),$$

and the *well-posedness condition*

$$l(\theta_0) \neq 0 \quad \text{on } \Gamma_0.$$

Then there exists a unique L_p -solution (u, Γ) for the Stefan problem with surface tension (1.4) on some possibly small but nontrivial time interval $J = [0, T]$, and

$$\mathfrak{M} := \bigcup_{t \in (0, T)} \{t\} \times \Gamma(t)$$

is a C^k -manifold in \mathbb{R}^{m+2} . In particular, each manifold $\Gamma(t)$ is C^k for $t \in (0, T)$.

The result in the presence of kinetic undercooling reads as

THEOREM 1.2 ($\gamma > 0$). Let $p > m + 3$, $\sigma > 0$. Suppose that $d_i, \gamma \in C^{k+2}(0, \infty)$, $\psi_i \in C^{k+3}(0, \infty)$ for $i = 1, 2$ such that

$$\kappa_i(u) = -u\psi_i''(u) > 0, \quad d_i(u) > 0, \quad \gamma(u) > 0, \quad u \in (0, \infty).$$

Assume the *regularity conditions*

$$\theta_0 \in W_p^{2-2/p}(\Omega \setminus \Gamma_0) \cap C(\bar{\Omega}), \quad \theta_0 > 0, \quad \Gamma_0 \in W_p^{4-3/p},$$

and the *compatibility conditions*

$$\partial_{\nu_\Omega} \theta_0 = 0, \quad (\llbracket \psi(\theta_0) \rrbracket + \sigma \mathcal{H}(\Gamma_0))(l(\theta_0) - \llbracket \psi(\theta_0) \rrbracket - \sigma \mathcal{H}(\Gamma_0)) = \gamma(\theta_0) \llbracket d(\theta_0)_{\Gamma_0} \theta_0 \rrbracket.$$

Then there exists a unique L_p -solution (θ, Γ) for the Stefan problem with surface tension (1.4) on some possibly small but nontrivial time interval $J = [0, T]$, and

$$\mathfrak{M} := \bigcup_{t \in (0, T)} \{t\} \times \Gamma(t)$$

is a C^k -manifold in \mathbb{R}^{m+2} . In particular, each manifold $\Gamma(t)$ is C^k for $t \in (0, T)$.

Remark 1.3. (a) For $k \in \mathbb{N} \cup \{\infty\}$, under the conditions in Theorems 1.1 and 1.2, we can show that the temperature satisfies

$$\theta \in C^k\left(\left((0, T) \times \Omega\right) \setminus \mathfrak{M}\right).$$

See Remark 4.5 for a justification. However, in the case $k = \omega$, additional work is needed to establish the interior analyticity of θ in the bulk phases. In order to keep this already long paper at a reasonable length, we will refrain from establishing this here.

(b) According to [65, Theorem 3.9] we obtain a solution of (1.4) in the state manifold $\mathfrak{S}\mathfrak{M}$ on a maximal interval of existence $[0, t_*)$. The regularity assertions of Theorem 1.1 and Theorem 1.2, respectively, then hold true on $[0, t_*)$. □

Notation. Throughout this paper, we always assume that

- E denotes a finite dimensional Banach space.
- $m + 3 < p < \infty$, and $s \geq 0$, unless stated otherwise.

Given two Banach spaces X, Y , the notation $\mathfrak{L}(X, Y)$ stands for the set of all bounded linear operators from X to Y , and $\mathfrak{L}is(X, Y)$ denotes the set of all bounded linear isomorphisms from X to Y .

For any topological sets U and V , $U \subset\subset V$ means that $\bar{U} \subset \overset{\circ}{V}$ with \bar{U} compact.

2. Parameterization over a fixed interface

Let $\Omega \subset \mathbb{R}^{m+1}$ be a bounded domain with boundary $\partial\Omega$ of class C^2 , and suppose that $\Gamma_0 \subset \Omega$ is a closed embedded hypersurface of class C^2 , that is, a C^2 -manifold which is the boundary of a bounded domain $\Omega_1(0) \subset \Omega$. Set $\Omega_2(0) := \Omega \setminus \bar{\Omega}_1(0)$. Following [63, 64], we may approximate Γ_0 by an m -dimensional real analytic compact closed oriented reference hypersurface (Σ, g) , with g the tangential metric induced by the Euclidean metric g_{m+1} , in the sense that there exists a function $h_0 \in C^2(\Sigma, (-a, a))$ such that the map

$$\Lambda_{h_0} : \Sigma \rightarrow \mathbb{R}^{m+1} : [\mathbf{p} \mapsto \mathbf{p} + h_0(\mathbf{p})\nu_\Sigma(\mathbf{p})]$$

is a diffeomorphism from Σ onto Γ_0 , where ν_Σ is the outer normal of Σ . The positive constant a depends on the inner and outer ball conditions of Σ . It is well-known that Σ admits a a -tubular neighborhood \mathbb{T}_a , which means that the map

$$\Lambda : \Sigma \times (-a, a) \rightarrow \mathbb{R}^{m+1} : (\mathbf{p}, r) \mapsto \mathbf{p} + r\nu_\Sigma(\mathbf{p})$$

is a diffeomorphism from the fiber bundle $\Sigma \times (-a, a)$ onto $\text{im}(\Lambda) := \mathbb{T}_a$. For sufficiently small a , $\bar{\mathbb{T}}_a \subset \Omega$. Σ bounds a domain Ω_1 , and we set $\Omega_2 = \Omega \setminus \bar{\Omega}_1$. The inverse Λ^{-1} can be decomposed as

$$\Lambda^{-1} : \mathbb{T}_a \rightarrow \Sigma \times (-a, a) : z \mapsto (\Pi(z), d_\Sigma(z)),$$

where $\Pi(z)$ is the metric projection of z onto Σ and $d_\Sigma(z)$ denotes the signed distance from z to Σ such that $|d_\Sigma(z)| = \text{dist}(z, \Sigma)$ and $d_\Sigma(z) < 0$ iff $z \in \Omega_1$. It follows from the inverse function theorem that the maps Π and d_Σ are both real analytic.

We may use the map Λ to parameterize the unknown free boundary $\Gamma(t)$ over Σ by a height function $h(t) : \Sigma \rightarrow \mathbb{R}$ via

$$\Gamma(t) := \{\mathbf{p} + h(t, \mathbf{p})\nu_\Sigma(\mathbf{p}), \quad \mathbf{p} \in \Sigma\}, \quad t \geq 0,$$

for small $t \geq 0$, at least. $\Gamma(t)$ bounds a bounded region $\Omega_1(t)$. Put $\Omega_2(t) = \Omega \setminus \bar{\Omega}_1(t)$. Choosing an auxiliary function $\zeta \in \mathcal{D}((-2a/3, 2a/3), [0, 1])$ such that $\zeta|_{(-a/3, a/3)} \equiv 1$, we may extend the above diffeomorphism onto $\bar{\Omega}$ via

$$\mathcal{E}_h(t, z) = z + \zeta(d_\Sigma(z))h(t, \Pi(z))\nu_\Sigma(\Pi(z)) =: z + \Upsilon(h)(t, z).$$

We have transformed the time varying regions $\Omega_i(t)$ to the fixed domains Ω_i . This is the direct mapping method, also known as Hanazawa transformation. By means of this transformation, we obtain the following transformed problem for $\vartheta(t, z) := \theta(t, \mathcal{E}_h(t, z))$:

$$\left\{ \begin{array}{ll} \kappa(\vartheta)\partial_t \vartheta + \mathcal{Q}(\vartheta, h)\vartheta = \kappa(\vartheta)\mathcal{R}(h)\vartheta & \text{in } \Omega \setminus \Sigma, \\ \partial_{\nu_\Omega} \vartheta = 0 & \text{on } \partial\Omega, \\ \llbracket \vartheta \rrbracket = 0 & \text{on } \Sigma, \\ \llbracket \psi(\vartheta) \rrbracket + \sigma \mathcal{H}(h) = \gamma(\vartheta)\beta(h)\partial_t h & \text{on } \Sigma, \\ \{l(\vartheta) - \gamma(\vartheta)\beta(h)\partial_t h\}\beta(h)\partial_t h + \mathcal{B}(\vartheta, h)\vartheta = 0 & \text{on } \Sigma, \\ \vartheta(0) = \vartheta_0, \quad h(0) = h_0. & \end{array} \right. \quad (2.1)$$

Here $\mathfrak{Q}(\vartheta, h)$ and $\mathfrak{B}(\vartheta, h)$ denote the transformations of $-\operatorname{div}(d\nabla)$ and $-[d\partial_{\nu_\Gamma}]$, respectively. Moreover, $\mathfrak{H}(h)$ stands for the mean curvature of the hypersurface

$$\Gamma_h := \Lambda_h(\Sigma).$$

The term $\beta(h)\partial_t h$ represents the normal velocity V with $\beta(h) := (\nu_\Sigma|_{\nu_\Gamma}(h))$, where $\nu_\Gamma(h)$ is the outer normal field of Γ_h , and

$$\mathfrak{R}(h)\vartheta := \partial_t \vartheta - \partial_t \theta \circ \mathfrak{E}_h.$$

It is shown in [65] that $\nu_\Gamma(h) = \beta(h)(\nu_\Sigma - \alpha(h))$, and

$$\beta(h) = (1 + |\alpha(h)|^2)^{-1/2}, \quad \mathfrak{R}(h)\vartheta = (\nabla\vartheta|[I + \nabla\Upsilon(h)^\top]^{-1}\partial_t\Upsilon(h)).$$

Here with the Weingarten map $L_\Sigma = -\nabla_\Sigma\nu_\Sigma$, we have

$$\alpha(h) := M_0(h)\nabla_\Sigma h, \quad \text{where } M_0(h) := (I - hL_\Sigma)^{-1}.$$

With $\partial_\nu := \partial_{\nu_\Sigma}$, the operator $\mathfrak{B}(\vartheta, h)$ becomes

$$\begin{aligned} \mathfrak{B}(\vartheta, h)\vartheta &= -[d(\vartheta)\partial_{\nu_\Gamma}\theta] \circ \mathfrak{E}_h = -([d(\vartheta)(I - M_1(h))\nabla\vartheta]|_{\nu_\Gamma}) \\ &= -\beta(h)([d(\vartheta)(I - M_1(h))\nabla\vartheta]|_{\nu_\Sigma - \alpha(h)}) \\ &= -\beta(h)[d(\vartheta)\partial_\nu\vartheta] + \beta(h)([d(\vartheta)\nabla\vartheta]|(I - M_1(h))^\top\alpha(h)), \end{aligned}$$

where $M_1(h) := [(I + \nabla\Upsilon(h)^\top)^{-1}\nabla\Upsilon(h)^\top]^\top$, and finally

$$\begin{aligned} \mathfrak{Q}(\vartheta, h)\vartheta &= -\operatorname{div}(d(\vartheta)\nabla\vartheta) \circ \mathfrak{E}_h = -\left((I - M_1(h))\nabla|d(\vartheta)(I - M_1(h))\nabla\vartheta\right) \\ &= -d(\vartheta)\Delta\vartheta + d(\vartheta)[M_1(h) + M_1^\top(h) - M_1(h)M_1^\top(h)] : \nabla^2\vartheta \\ &\quad - d'(\vartheta)|\nabla\vartheta|^2 + d(\vartheta)\left((I - M_1(h)) : \nabla M_1(h)|\nabla\vartheta\right). \end{aligned}$$

We recall that, for matrices $A, B \in \mathbb{R}^{n \times n}$, $A : B = \sum_{i,j=1}^n a_{ij}b_{ij} = \operatorname{tr}(AB^\top)$ denotes the inner product.

We set

$$\begin{aligned} \mathbb{E}_1(J) &:= \{\vartheta \in H_p^1(J; L_p(\Omega)) \cap L_p(J; H_p^2(\Omega \setminus \Sigma)) : \llbracket \vartheta \rrbracket = 0, \partial_{\nu_\Omega}\vartheta = 0\}, \\ \mathbb{E}_2(J) &:= \begin{cases} W_p^{3/2-1/2p}(J; L_p(\Sigma)) \cap W_p^{1-1/2p}(J; H_p^2(\Sigma)) \cap L_p(J; W_p^{4-1/p}(\Sigma)), & \gamma \equiv 0, \\ W_p^{2-1/2p}(J; L_p(\Sigma)) \cap L_p(J; W_p^{4-1/p}(\Sigma)), & \gamma > 0, \end{cases} \\ \mathbb{E}(J) &:= \mathbb{E}_1(J) \times \mathbb{E}_2(J), \end{aligned}$$

that is, $\mathbb{E}(J)$ denotes the solution space for (2.1). Similarly, we define

$$\begin{cases} \mathbb{F}_1(J) := L_p(J; L_p(\Omega)), \\ \mathbb{F}_2(J) := W_p^{1-1/2p}(J; L_p(\Sigma)) \cap L_p(J; W_p^{2-1/p}(\Sigma)), \\ \mathbb{F}_3(J) := W_p^{1/2-1/2p}(J; L_p(\Sigma)) \cap L_p(J; W_p^{1-1/p}(\Sigma)), \\ \mathbb{F}_4 := [W_p^{2-2/p}(\Omega \setminus \Sigma) \cap C(\bar{\Omega})] \times W_p^{4-3/p}(\Sigma). \end{cases} \quad (2.2)$$

A left subscript zero means vanishing time trace at $t = 0$, whenever it exists. So for example ${}_0\mathbb{E}_2(J) = \{h \in \mathbb{E}_2(J) : h(0) = \partial_t h(0) = 0\}$ for $p > 3$.

Whenever (J) is replaced by (J, U) , or $(J, U; E)$ for any set U , e.g., in $\mathbb{E}(J)$, it always means that in the corresponding spaces the original spatial domain is replaced by U and the functions become E -valued in the latter case. For example,

$$\mathbb{F}_2(J, \tau_a; E) := W_p^{1-1/2p}(J; L_p(\tau_a, E)) \cap L_p(J; W_p^{2-1/p}(\tau_a, E)).$$

We equip Σ with a normalized real analytic atlas $\mathfrak{A} = (\mathcal{O}_\kappa, \varphi_\kappa)_{\kappa \in \mathfrak{K}}$, where \mathfrak{K} is a finite index set, in the sense that $\varphi_\kappa(\mathcal{O}_\kappa) = \mathbb{Q}^m := (-1, 1)^m$, for every κ . Let $\psi_\kappa := [\varphi_\kappa]^{-1}$. Then we may endow Ω with a real analytic atlas $\mathfrak{A}_\epsilon = (\mathcal{O}_{\epsilon, \kappa}, \varphi_{\epsilon, \kappa})_{\kappa \in \mathfrak{K}_\epsilon}$ with $\mathfrak{K}_\epsilon := \mathfrak{K} \cup \{\kappa_1, \kappa_2\}$ by the following construction. When $\kappa \in \mathfrak{K}$

$$\mathcal{O}_{\epsilon, \kappa} := \tau_{a, \kappa} := \Lambda(\mathcal{O}_\kappa \times (-a, a)), \quad \varphi_{\epsilon, \kappa}(z) := (\varphi_\kappa \circ \Pi(z), d_\Sigma(z)), \quad z \in \mathcal{O}_{\epsilon, \kappa}.$$

Let $\mathcal{Q}_a := \mathbb{Q}^m \times (-a, a)$ and

$$\psi_{\epsilon, \kappa} := [\varphi_{\epsilon, \kappa}]^{-1} : \mathcal{Q}_a \rightarrow \mathcal{O}_{\epsilon, \kappa} : [(x, y) \mapsto \psi_\kappa(x) + y\nu_\Sigma(\psi_\kappa(x))]$$

with $x \in \mathbb{Q}^m$ and $y \in (-a, a)$. For $\kappa \notin \mathfrak{K}$, we set

$$(\mathcal{O}_{\epsilon, \kappa_i}, \varphi_{\epsilon, \kappa_i}) := (\mathcal{O}_{\epsilon, \kappa_i}, \text{id}_{\mathcal{O}_{\epsilon, \kappa_i}}), \quad i = 1, 2,$$

where $\mathcal{O}_{\epsilon, \kappa_i} := \Omega_i \setminus \bar{\tau}_{a/6}$ for $i = 1, 2$. It is immediate from our construction that the atlas \mathfrak{A}_ϵ is real analytically compatible with the Euclidean structure of Ω . Moreover, all the transition maps are C^∞ -continuous and have bounded derivatives, which we refer to as BC^∞ -continuous.

A family $(\pi_\kappa)_{\kappa \in \mathfrak{K}}$ is called a localization system subordinate to \mathfrak{A} if

$$\pi_\kappa \in \mathfrak{D}(\mathcal{O}_\kappa, [0, 1]) \text{ and } (\pi_\kappa^2)_{\kappa \in \mathfrak{K}} \text{ is a partition of unity subordinate to } (\mathcal{O}_\kappa)_{\kappa \in \mathfrak{K}}.$$

Based on [5, Lemma 3.2], Σ admits a localization system $(\pi_\kappa)_{\kappa \in \mathfrak{K}}$ subordinate to \mathfrak{A} . Using the cut-off function ζ introduced above, we may construct a new localization system $(\pi_{\epsilon, \kappa})_{\kappa \in \mathfrak{K}_\epsilon}$ subordinate to \mathfrak{A}_ϵ as follows:

(L1) For any $\kappa \in \mathfrak{K}$, $\pi_{\epsilon, \kappa} \in \mathfrak{D}(\mathcal{O}_{\epsilon, \kappa}, [0, 1])$ is defined by

$$\pi_{\epsilon, \kappa}(z) := \begin{cases} \zeta(d_\Sigma(z))\pi_\kappa(\Pi(z)), & z \in \mathcal{O}_{\epsilon, \kappa}, \\ 0, & z \notin \mathcal{O}_{\epsilon, \kappa}. \end{cases}$$

(L2) For $i = 1, 2$, $\pi_{\epsilon, \kappa_i} \in C^\infty(\bar{\mathcal{O}}_{\epsilon, \kappa_i}, [0, 1])$ is defined by

$$\pi_{\epsilon, \kappa_i}(z) = \begin{cases} \frac{1 - \zeta(d_\Sigma(z))}{\sqrt{(1 - \zeta(d_\Sigma(z)))^2 + \zeta^2(d_\Sigma(z))}}, & z \in \Omega_i \cap \tau_a, \\ 1, & z \in \bar{\Omega}_i \setminus \tau_a. \end{cases}$$

$(\pi_{\epsilon, \kappa}^2)_{\kappa \in \mathfrak{K}_\epsilon}$ forms a localization system subordinate to $(\mathcal{O}_{\epsilon, \kappa})_{\kappa \in \mathfrak{K}_\epsilon}$. We put

$$\mathbb{X}_{\epsilon, \kappa} := \begin{cases} \mathbb{R}^{m+1}, & \kappa \in \mathfrak{K}, \\ \mathcal{O}_{\epsilon, \kappa_i}, & i = 1, 2. \end{cases}$$

For any finite dimensional Banach space E , the maps $\mathcal{R}_{\epsilon,\kappa}^c$ and $\mathcal{R}_{\epsilon,\kappa}$ are defined by

$$\mathcal{R}_{\epsilon,\kappa}^c : L_{1,loc}(\Omega, E) \rightarrow L_{1,loc}(\mathbb{X}_{\epsilon,\kappa}, E) : u \mapsto \psi_{\epsilon,\kappa}^* \pi_{\epsilon,\kappa} u, \quad \kappa \in \mathfrak{K}_\epsilon,$$

and

$$\mathcal{R}_{\epsilon,\kappa} : L_{1,loc}(\mathbb{X}_{\epsilon,\kappa}, E) \rightarrow L_{1,loc}(\Omega, E) : u \mapsto \pi_{\epsilon,\kappa} \varphi_{\epsilon,\kappa}^* u, \quad \kappa \in \mathfrak{K}_\epsilon.$$

Here and in the following it is understood that a partially defined and compactly supported map is automatically extended over the whole base manifold by identifying it to be zero outside its original domain. Moreover, let

$$\mathcal{R}_\epsilon^c : L_{1,loc}(\Omega, E) \rightarrow \prod_{\kappa \in \mathfrak{K}_\epsilon} L_{1,loc}(\mathbb{X}_{\epsilon,\kappa}, E) : u \mapsto (\mathcal{R}_{\epsilon,\kappa}^c u)_{\kappa \in \mathfrak{K}_\epsilon},$$

and

$$\mathcal{R}_\epsilon : \prod_{\kappa \in \mathfrak{K}_\epsilon} L_{1,loc}(\mathbb{X}_{\epsilon,\kappa}, E) \rightarrow L_{1,loc}(\Omega, E) : (u_\kappa)_\kappa \mapsto \sum_{\kappa \in \mathfrak{K}_\epsilon} \pi_{\epsilon,\kappa} \varphi_{\epsilon,\kappa}^* u_\kappa.$$

On the manifold Σ , similar maps \mathcal{R}_κ^c , \mathcal{R}^c , \mathcal{R}_κ , and \mathcal{R} are defined in terms of π_κ , φ_κ , and ψ_κ . See [73].

2.1 Function spaces

For any open subset $U \subset \mathbb{R}^n$, the Banach space $BC^k(U, E)$ is defined by

$$BC^k(U, E) := (\{u \in C^k(U, E) : \|u\|_{k,\infty} < \infty\}, \|\cdot\|_{k,\infty}).$$

The closed linear subspace $BUC^k(U, E)$ of $BC^k(U, E)$ consists of all functions $u \in BC^k(U, E)$ such that $\partial^\alpha u$ is uniformly continuous for all $|\alpha| \leq k$. Moreover,

$$BC^\infty(U, E) := \bigcap_k BC^k(U, E) = \bigcap_k BUC^k(U, E).$$

It is a Fréchet space equipped with the natural projective topology.

For $0 < s < 1$, $0 < \delta \leq \infty$ and $u \in E^U$, the seminorm $[\cdot]_{s,\infty}^\delta$ is defined by

$$[u]_{s,\infty}^\delta := \sup_{h \in (0,\delta)^n} \frac{\|u(\cdot + h) - u(\cdot)\|_\infty}{|h|^s}, \quad [\cdot]_{s,\infty} := [\cdot]_{s,\infty}^\infty.$$

Let $k < s < k + 1$. The Hölder space $BC^s(U, E)$ is defined as

$$BC^s(U, E) := (\{u \in BC^k(U, E) : \|u\|_{s,\infty} < \infty\}, \|\cdot\|_{s,\infty}),$$

where $\|u\|_{s,\infty} := \|u\|_{k,\infty} + \max_{|\alpha|=k} [\partial^\alpha u]_{s-k,\infty}$.

In order to have a general theory that is applicable to situations other than the Stefan problem, we also introduce the *little Hölder space* of order $s \geq 0$, which is defined by

$$bc^s(U, E) := \text{the closure of } BC^\infty(U, E) \text{ in } BC^s(U, E).$$

By [6, formula (11.13), Corollary 11.2, Theorem 11.3], we have

$$bc^k(U, E) = BUC^k(U, E),$$

and for $k < s < k + 1$

$$u \in BC^s(U, E) \text{ belongs to } bc^s(U, E) \text{ iff } \lim_{\delta \rightarrow 0} [\partial^\alpha u]_{s-[s], \infty}^\delta = 0, \quad |\alpha| = [s].$$

The spaces $\mathfrak{F}^s(\Sigma, E)$ with $\mathfrak{F} \in \{bc, BC, W_p, H_p\}$ are defined in terms of the smooth atlas \mathfrak{A} , that is, $u \in \mathfrak{F}^s(\Sigma, E)$ iff $\psi_\kappa^* u := u \circ \psi_\kappa \in \mathfrak{F}^s(\mathbb{Q}^m, E)$ for all $\kappa \in \mathfrak{K}$. See [5] and [73].

Throughout, for any finite index set \mathbb{A} and Banach spaces E_α , $\alpha \in \mathbb{A}$, it is understood that the space $\mathbb{E} := \prod_{\alpha \in \mathbb{A}} E_\alpha$ is equipped with the maximum norm. For $\mathfrak{F} \in \{bc, BC, BUC, W_p, H_p\}$, we put

$$\mathfrak{F}_\epsilon^s := \prod_{\kappa \in \mathfrak{K}_\epsilon} \mathfrak{F}^s(\mathbb{X}_{\epsilon, \kappa}, E), \quad \mathfrak{F}^s := \prod_{\kappa \in \mathfrak{K}} \mathfrak{F}_\kappa^s,$$

where $\mathfrak{F}_\kappa^s = \mathfrak{F}^s(\mathbb{R}^m, E)$.

PROPOSITION 2.1 Suppose that $\mathfrak{B} = \mathfrak{F}$ when $\mathfrak{F} \in \{bc, W_p, H_p\}$, or $\mathfrak{B} \in \{BC, BUC\}$ when $\mathfrak{F} = BC$. Then

- (a) \mathcal{R} is a retraction from \mathfrak{B}^s onto $\mathfrak{F}^s(\Sigma, E)$ with \mathcal{R}^c as a coretraction. Moreover, the case $s < 0$ and $\mathfrak{F} \in \{W_p, H_p\}$ is also admissible.
- (b) \mathcal{R}_ϵ is a retraction from \mathfrak{B}_ϵ^s onto $\mathfrak{F}^s(\Omega, E)$ with \mathcal{R}_ϵ^c as a coretraction.

Proof. (a) The assertions are special cases of [6, Theorem 9.3] and [73, Propositions 2.1, 2.2].

- (b) For $\kappa \in \mathfrak{K}$, the boundedness of the maps $\mathcal{R}_{\epsilon, \kappa}$ and $\mathcal{R}_{\epsilon, \kappa}^c$ follows from the proofs in [5, Section 6, 7] and [6, Section 12]. The continuity of the maps $\mathcal{R}_{\epsilon, \kappa_i}^c$ and $\mathcal{R}_{\epsilon, \kappa_i}$ is straightforward. \square

Let $J := [0, T]$ for some $T > 0$. Due to the temporal independence of the above retraction-coretraction systems, we readily infer that

PROPOSITION 2.2 Let $r \geq 0$. Suppose that $\mathcal{C} \in \{C, W_p\}$, and $\mathfrak{B} = \mathfrak{F}$ when $\mathfrak{F} \in \{bc, W_p, H_p\}$, or $\mathfrak{B} \in \{BC, BUC\}$ when $\mathfrak{F} = BC$. Then

- (a) \mathcal{R} is a retraction from $\mathcal{C}^r(J; \mathfrak{B}^s)$ onto $\mathcal{C}^r(J; \mathfrak{F}^s(\Sigma, E))$ with \mathcal{R}^c as a coretraction. Moreover, the case $s < 0$ and $\mathfrak{F} \in \{W_p, H_p\}$ is also admissible.
- (b) \mathcal{R}_ϵ is a retraction from $\mathcal{C}^r(J; \mathfrak{B}_\epsilon^s)$ onto $\mathcal{C}^r(J; \mathfrak{F}^s(\Omega, E))$ with \mathcal{R}_ϵ^c as a coretraction.

3. Parameter-dependent diffeomorphisms

3.1 Model diffeomorphisms in Euclidean space

Given $z_c \in \mathbb{T}_{a/3, \kappa_c} \setminus \Sigma \subset \Omega$ for some $\kappa_c \in \mathfrak{K}$, let

$$(x_c, y_c) = (\varphi_{\kappa_c} \circ \Pi(z_c), d_\Sigma(z_c)).$$

We may choose a sufficiently small positive constant ε_0 such that

$$\mathbb{B}^m(x_c, 5\varepsilon_0) \subset \mathbb{Q}^m, \quad \mathbb{B}(y_c, 5\varepsilon_0) \subset (-1)^i(0, a/3).$$

Let $\mathbb{B}_{\varepsilon, \alpha} := \mathbb{B}^m(x_c, \varepsilon) \times (-\alpha, \alpha)$. We pick several auxiliary functions in the following manner:

- $\varpi \in \mathfrak{D}(-13a/18, 13a/18), [0, 1]$, and $\varpi|_{(-2a/3, 2a/3)} \equiv 1$.
 - $\chi_m \in \mathfrak{D}(\mathbb{B}^m(x_c, 2\varepsilon_0), [0, 1])$, and $\chi_m|_{\mathbb{B}^m(x_c, \varepsilon_0)} \equiv 1$.
 - $\chi \in \mathfrak{D}(\mathbb{B}(y_c, 2\varepsilon_0), [0, 1])$, and $\chi|_{\mathbb{B}(y_c, \varepsilon_0)} \equiv 1$.
 - $\zeta_\varepsilon \in \mathfrak{D}(\psi_{\kappa_c}(\mathbb{B}_{5\varepsilon_0, 17a/18}), [0, 1])$, and $\zeta_\varepsilon|_{\psi_{\kappa_c}(\mathbb{B}_{4\varepsilon_0, 8a/9})} \equiv 1$. We set $\zeta := \zeta_\varepsilon|_\Sigma$
- Then we can introduce a localized translation on \mathbb{Q}_a as follows

$$\theta_{\mu, \eta} : (x, y) \mapsto (x + \chi_m(x)\varpi(y)\mu, y + \chi_m(x)\chi(y)\eta), \quad (\mu, \eta) \in \mathbb{B}(0, r_0),$$

with $\mu \in \mathbb{R}^m, \eta \in \mathbb{R}$ and sufficiently small $r_0 > 0$. Henceforth, $\mathbb{B}(x_0, r_0)$ always denotes the ball with radius r_0 centered at x_0 in \mathbb{R}^n . The dimension n of the ball is not distinguished as long as it is clear from the context.

The related partial translations in horizontal/vertical directions can be defined separately as

$$\begin{cases} \theta_\mu : (x, y) \mapsto (x + \chi_m(x)\varpi(y)\mu, y), \\ \bar{\theta}_\eta : (x, y) \mapsto (x, y + \chi_m(x)\chi(y)\eta). \end{cases} \tag{3.1}$$

For r_0 small and every $(\mu, \eta) \in \mathbb{B}(0, r_0)$

$$\theta_{\mu, \eta} = \theta_\mu \circ \bar{\theta}_\eta. \tag{3.2}$$

It is not hard to show that for sufficiently small r_0 , we have $\theta_{\mu, \eta}(\mathbb{B}_{3\varepsilon_0, 7a/9}) \subset \mathbb{B}_{3\varepsilon_0, 7a/9}$ for any $(\mu, \eta) \in \mathbb{B}(0, r_0)$. (3.1) implies that

$$\theta_{\mu, \eta}^* = \bar{\theta}_\eta^* \circ \theta_\mu^*, \quad \text{and} \quad \theta_{\mu, \eta}^{\mu, \eta} := [\theta_{\mu, \eta}]_* = \theta_\mu^* \circ \bar{\theta}_\eta^*. \tag{3.3}$$

Since θ_μ and $\bar{\theta}_\eta$ are the restrictions of truncated translations in the sense of those defined in [26] in horizontal and vertical direction, respectively, the readers should have no difficulty convincing themselves that all the results in [26] are at our disposal for functions defined on an open subset $U \subset \mathbb{R}^{m+1}$ and the transformations θ_μ^* and $\bar{\theta}_\eta^*$. Taking advantage of these considerations, we will state some properties of $\theta_{\mu, \eta}^*$ for later use in the following with brief proofs.

Hereafter, we assume $O \subset U \subset \mathbb{R}^{m+1}$ are open, and O contains $\bar{\mathbb{B}}_{3\varepsilon_0, 7a/9}$. We conclude from $\theta_{\mu, \eta}(U) \subset U$ that $\theta_{\mu, \eta}^* : \mathfrak{F}^s(U, E) \rightarrow E^U$. In the rest of this section, we always assume the following, unless mentioned otherwise,

$$1 < p < \infty, \quad n, l \in \mathbb{N}_0, \quad k \in \mathbb{N}_0 \cup \{\infty, \omega\}.$$

PROPOSITION 3.1 Suppose that $\mathfrak{F} \in \{bc, BUC, W_p, H_p\}$ and $u \in C^{n+k}(O, E) \cap \mathfrak{F}^s(U, E)$ with $s \in [0, n]$ when $\mathfrak{F} \in \{bc, W_p, H_p\}$, or with $s = n$ when $\mathfrak{F} = BUC$. Then for sufficiently small r_0 ,

- (a) $\theta_{\mu, \eta} \in \text{Diff}^\infty(U), (\mu, \eta) \in \mathbb{B}(0, r_0)$.
- (b) $\theta_{\mu, \eta}^* \in \mathfrak{L}(\mathfrak{F}^s(U, E)), [\theta_{\mu, \eta}^*]^{-1} = \theta_{\mu, \eta}^{\mu, \eta}, (\mu, \eta) \in \mathbb{B}(0, r_0)$. Moreover, there exists a constant $M > 0$ such that

$$\|\theta_{\mu, \eta}^*\|_{\mathfrak{L}(\mathfrak{F}^s(U, E))} \leq M, \quad (\mu, \eta) \in \mathbb{B}(0, r_0).$$

- (c) $[(\mu, \eta) \mapsto \theta_{\mu, \eta}^* u] \in C^k(\mathbb{B}(0, r_0), \mathfrak{F}^s(U, E))$.

Proof. (a) The statement is immediate from (3.3) and [26, Proposition 2.2].

(b) We infer from (a) that

$$\theta_{\mu,\eta}^* \circ \theta_{\mu,\eta}^{\mu,\eta} = \theta_{\mu,\eta}^{\mu,\eta} \circ \theta_{\mu,\eta}^* = \text{id}_{\mathfrak{F}^s(U,E)}.$$

[26, Proposition 2.4] implies that for r small enough

$$\theta_{\mu}^*, \bar{\theta}_{\eta}^* \in \mathfrak{Lis}(\mathfrak{F}^s(U, E)), \quad (\mu, \eta) \in \mathbb{B}(0, r_0).$$

Moreover, there exists a constant $M > 0$ such that

$$\|\theta_{\mu}^*\|_{\mathfrak{L}(\mathfrak{F}^s(U,E))} + \|\bar{\theta}_{\eta}^*\|_{\mathfrak{L}(\mathfrak{F}^s(U,E))} \leq M, \quad (\mu, \eta) \in \mathbb{B}(0, r_0).$$

The assertion now follows from (3.3).

(c) It is a simple matter to see that the assertion holds for the transformation θ_{μ}^* and $\bar{\theta}_{\eta}^*$, respectively, namely that we have

$$[\mu \mapsto \theta_{\mu}^* u] \in C^k(\mathbb{B}(0, r_0), \mathfrak{F}^s(U, E)), \quad [\eta \mapsto \bar{\theta}_{\eta}^* u] \in C^k(\mathbb{B}(0, r_0), \mathfrak{F}^s(U, E)) \quad (3.4)$$

with $\partial_{\mu}^{\alpha}[\theta_{\mu}^* u] = \chi_m^{|\alpha|} \varpi^{|\alpha|}[\theta_{\mu}^* \partial_x^{\alpha} u]$, and $\partial_{\eta}^{\beta}[\bar{\theta}_{\eta}^* u] = \chi_m^{\beta} \chi^{\beta}[\bar{\theta}_{\eta}^* \partial_y^{\beta} u]$. Then we can obtain from (3.3), (3.4), the point-wise multiplier theorem in [75, Section 3.3.2], and induction that the map

$$(\mu, \eta) \mapsto \partial_{(\mu,\eta)}^{(\alpha,\beta)}[\theta_{\mu,\eta}^* u] = \chi_m^{|\alpha|+\beta} \chi^{\beta} \varpi^{|\alpha|}[\theta_{\mu,\eta}^* \partial_{(x,y)}^{(\alpha,\beta)} u], \quad |\alpha| + \beta \leq k,$$

is separately $C^{k-|\alpha|-\beta}$ -continuous into the space $\mathfrak{F}^s(U, E)$. So the case $k \in \mathbb{N}_0 \cup \{\infty\}$ follows immediately from a well-known fact in multi-variable calculus. When $k = \omega$, the proof is basically the same as that of [26, Proposition 3.2]. \square

Let

$$\mathfrak{Q} := \sum_{|\alpha| \leq l} a_{\alpha} \partial^{\alpha},$$

be an l -th order linear differential operator on U with coefficients $a_{\alpha} : U \rightarrow \mathfrak{L}(E)$, where $\alpha \in \mathbb{N}_0^{m+1}$. We set $\tilde{\mathfrak{G}}_{\mu,\eta} := \theta_{\mu,\eta}^* \mathfrak{Q} \theta_{\mu,\eta}^{\mu,\eta}$.

THEOREM 3.2 Suppose that $a_{\alpha} \in C^{n+k}(O, \mathfrak{L}(E)) \cap BUC^n(U, \mathfrak{L}(E))$. Then

$$[(\mu, \eta) \mapsto \tilde{\mathfrak{G}}_{\mu,\eta}] \in C^k(\mathbb{B}(0, r_0), \mathfrak{L}(\mathfrak{F}^{s+l}(U, E), \mathfrak{F}^s(U, E))),$$

where $\mathfrak{F} \in \{bc, BUC, W_p, H_p\}$, and $s \in [0, n]$.

Proof. By modifying the proof of [26, Proposition 4.1] in an obvious way, one may verify the case of constant coefficients. Now the assertion is a direct consequence of point-wise multiplication results, and Proposition 3.1(c). \square

3.2 Parameter-dependent diffeomorphisms on Ω and Ω_i

By means of $\theta_{\mu,\eta}$, we are now in a position to introduce a family of localized translations on Ω and Ω_i , respectively, by

$$\Theta_{\mu,\eta}(z) = \begin{cases} \psi_{\epsilon,\kappa_c} \circ \theta_{\mu,\eta} \circ \varphi_{\epsilon,\kappa_c}(z), & z \in O_{\epsilon,\kappa_c}, \\ z, & z \in \Omega \setminus O_{\epsilon,\kappa_c}. \end{cases}$$

Thanks to Proposition 3.1(a), $\Theta_{\mu,\eta} \in \text{Diff}^\infty(\Omega) \cap \text{Diff}^\infty(\Omega_i)$ is evident.

There is a universal extension operator $\mathcal{E}_i \in \mathfrak{L}(\mathfrak{F}^s(\Omega_i, E), \mathfrak{F}^s(\Omega, E))$. The restriction operator from $\mathfrak{F}^s(\Omega, E)$ to $\mathfrak{F}^s(\Omega_i, E)$ is denoted by \mathcal{R}_i . Then \mathcal{R}_i is a retraction from $\mathfrak{F}^s(\Omega, E)$ onto $\mathfrak{F}^s(\Omega_i, E)$ with \mathcal{E}_i as a coretraction. When $u \in \mathfrak{F}^s(\Omega \setminus \Sigma, E)$, it is understood that $\mathcal{E}_i u := \mathcal{E}_i(u|_{\Omega_i})$.

For $u \in E^\Omega$, its pull-back and push-forward by $\Theta_{\mu,\eta}$ can be explicitly expressed as

$$\Theta_{\mu,\eta}^* u = \varphi_{\epsilon,\kappa_c}^* \theta_{\mu,\eta}^* \psi_{\epsilon,\kappa_c}^* (\zeta_\epsilon u) + (1_\Omega - \zeta_\epsilon)u, \quad \Theta_*^{\mu,\eta} u = \varphi_{\epsilon,\kappa_c}^* \theta_*^{\mu,\eta} \psi_{\epsilon,\kappa_c}^* (\zeta_\epsilon u) + (1_\Omega - \zeta_\epsilon)u.$$

Analogously, given $u \in E^{\Omega_i}$ of u , we have

$$\Theta_{\mu,\eta}^* u = \mathcal{R}_i \circ \Theta_{\mu,\eta}^* \circ \mathcal{E}_i u, \quad \Theta_*^{\mu,\eta} u = \mathcal{R}_i \circ \Theta_*^{\mu,\eta} \circ \mathcal{E}_i u. \tag{3.5}$$

The construction of $\Theta_{\mu,\eta}$ implies that $\Theta_{\mu,\eta}^*(E^{\Omega_i}) \subset E^{\Omega_i}$, and likewise for $\Theta_*^{\mu,\eta}$. Suppose that $\mathfrak{F} \in \{bc, BC, W_p, H_p\}$. We define

$$\mathfrak{F}_{\text{cp}}^{s,\Omega} := \{u \in \mathfrak{F}^s(\Omega, E) : \text{supp}(u) \subset \psi_{\epsilon,\kappa_c}(\bar{\mathbb{B}}_{5\epsilon_0, 17a/18})\},$$

and

$$\mathfrak{F}_{\text{cp}}^s := \{u \in \mathfrak{F}^s(\mathbb{R}^{m+1}, E) : \text{supp}(u) \subset \bar{\mathbb{B}}_{5\epsilon_0, 17a/18}\}.$$

The analogue of [73, Lemma 3.1] is at our disposal, i.e., it holds that

LEMMA 3.3 $\varphi_{\epsilon,\kappa_c}^* \in \mathfrak{L}(\mathfrak{F}_{\text{cp}}^s, \mathfrak{F}_{\text{cp}}^{s,\Omega})$ with $[\varphi_{\epsilon,\kappa_c}^*]^{-1} = \psi_{\epsilon,\kappa_c}^*$.

Proof. Since [73, (3.2)] still holds with ∇ and g denoting the usual gradient and Euclidean metric in \mathbb{R}^{m+1} , respectively, the proof is essentially the same as that of [73, Lemma 3.1]. \square

By means of Lemma 3.3, with only minor changes to the proofs in [73, Section 3], one can show that $\Theta_{\mu,\eta}$ inherits all the properties of its counterpart therein.

PROPOSITION 3.4 Suppose that $\mathfrak{F} \in \{bc, BC, W_p, H_p\}$. Then

$$\Theta_{\mu,\eta}^* \in \mathfrak{L}(\mathfrak{F}^s(\Omega, E)) \cap \mathfrak{L}(\mathfrak{F}^s(\Omega_i, E)), \quad [\Theta_{\mu,\eta}^*]^{-1} = \Theta_*^{\mu,\eta}, \quad (\mu, \eta) \in \mathbb{B}(0, r_0).$$

Moreover, there exists a constant $M > 0$ such that

$$\|\Theta_{\mu,\eta}^*\|_{\mathfrak{L}(\mathfrak{F}^s(\Omega, E))} + \|\Theta_{\mu,\eta}^*\|_{\mathfrak{L}(\mathfrak{F}^s(\Omega_i, E))} \leq M, \quad (\mu, \eta) \in \mathbb{B}(0, r_0).$$

Proof. The proof is almost the same as that of [73, Proposition 3.3]. For the reader's convenience, we will present a proof based on Lemma 3.3 for the case of Ω_i , and the other case follows in a similar manner. The identity

$$\Theta_{\mu,\eta}^* \circ \Theta_*^{\mu,\eta} = \Theta_*^{\mu,\eta} \circ \Theta_{\mu,\eta}^* = \text{id}_{\mathfrak{F}^s(\Omega, E)} \cap \text{id}_{\mathfrak{F}^s(\Omega_i, E)}$$

is obvious from the definitions of $\Theta_{\mu,\eta}^*$ and $\Theta_*^{\mu,\eta}$ and $\theta_{\mu,\eta}$. In the formula

$$\Theta_{\mu,\eta}^* u = \varphi_{\epsilon,\kappa_c}^* \theta_{\mu,\eta}^* \psi_{\epsilon,\kappa_c}^* (\zeta_\epsilon u) + (1_\Omega - \zeta_\epsilon)u,$$

it can easily be verified that the multiplication operators ζ_ϵ and $(1_\Omega - \zeta_\epsilon)$ are uniformly bounded, i.e.,

$$\|\zeta_\epsilon\|_{\mathfrak{L}(\mathfrak{F}^s(\Omega, E))} + \|(1_\Omega - \zeta_\epsilon)\|_{\mathfrak{L}(\mathfrak{F}^s(\Omega, E))} \leq C_1$$

for some constant C_1 . Lemma 3.3 implies that for some constant C_2

$$\|\varphi_{\epsilon, \kappa_c}^*\|_{\mathfrak{L}(\mathfrak{F}_{\text{cp}}^s, \mathfrak{F}_{\text{cp}}^{s, \Omega})} + \|\psi_{\epsilon, \kappa_c}^*\|_{\mathfrak{L}(\mathfrak{F}_{\text{cp}}^{s, \Omega}, \mathfrak{F}_{\text{cp}}^s)} \leq C_2.$$

It follows from Proposition 3.1(b) that there is a uniform bound C_3 such that

$$\|\theta_{\mu, \eta}^*\|_{\mathfrak{L}(\mathfrak{F}_{\text{cp}}^s)} \leq C_3.$$

The statement then is a direct consequence of the open mapping theorem and (3.5). \square

3.3 Time-dependence

Let $J = [0, T]$, $T > 0$. Assume that $I \subset \mathring{J}$ is an open interval and $t_c \in I$ is a fixed point. Choose ϵ_0 so small that $\mathbb{B}(t_c, 3\epsilon_0) \subset\subset I$. Pick an auxiliary function

$$\xi \in \mathfrak{D}(\mathbb{B}(t_c, 2\epsilon_0), [0, 1]) \quad \text{with} \quad \xi|_{\mathbb{B}(t_c, \epsilon_0)} \equiv 1.$$

The localized temporal translation is defined by

$$\varrho_\lambda(t) := t + \xi(t)\lambda, \quad t \in J \text{ and } \lambda \in \mathbb{R}.$$

For $v : J \times \mathbb{Q}_a \rightarrow E$, the parameter-dependent diffeomorphism can be expressed as

$$\tilde{v}_{\lambda, \mu, \eta}(t, \cdot) := \theta_{\lambda, \mu, \eta}^* v(t, \cdot) = \tilde{T}_{\mu, \eta}(t) \varrho_\lambda^* v(t, \cdot), \quad (\lambda, \mu, \eta) \in \mathbb{B}(0, r_0), \quad (3.6)$$

where $\tilde{T}_{\mu, \eta}(t) := \theta_{\xi(t)\mu, \xi(t)\eta}^*$ for $t \in J$.

Given $v : J \times \mathbb{Q}^m \rightarrow E$, analogously, setting $\tilde{T}_\mu(t) := \theta_{\xi(t)\mu}^*$, we can define

$$\tilde{v}_{\lambda, \mu}(t, \cdot) := \theta_{\lambda, \mu}^* v(t, \cdot) = \tilde{T}_\mu(t) \varrho_\lambda^* v(t, \cdot), \quad (\lambda, \mu) \in \mathbb{B}(0, r_0). \quad (3.7)$$

It is understood that θ_μ in (3.7) is restricted on the hyperplane $y = 0$, appearing as a special case of (3.1).

For any $u \in J \times \Omega \rightarrow E$, or $u \in J \times \Omega_i \rightarrow E$, we define

$$u_{\lambda, \mu, \eta}(t, \cdot) := \Theta_{\lambda, \mu, \eta}^* u(t, \cdot) := T_{\mu, \eta}(t) \varrho_\lambda^* u(t, \cdot), \quad (\lambda, \mu, \eta) \in \mathbb{B}(0, r_0).$$

Here $T_{\mu, \eta}(t) := \Theta_{\xi(t)\mu, \xi(t)\eta}^*$ for $t \in J$. By Proposition 3.4, for every t , $T_{\mu, \eta}(t)$ is invertible.

Let $\Theta_{\lambda, \mu, \eta}^* := [\Theta_{\lambda, \mu, \eta}^*]^{-1}$. Note that $u_{\lambda, \mu, \eta}(0, \cdot) = u(0, \cdot)$.

The following lemma strengthens [26, Lemma 5.1].

LEMMA 3.5 Suppose that $[(\mu, \eta) \mapsto f(\mu, \eta)] \in C^{k+j}(\mathbb{B}(0, r_0), X)$ for $j = 0, 1$ and some Banach space X . Let $F(\mu, \eta)(t) := f(\xi(t)(\mu, \eta))$ with $t \in J$. Then

$$[(\mu, \eta) \mapsto F(\mu, \eta)] \in C^k(\mathbb{B}(0, r_0), C^j(J, X)).$$

Proof. To economize notation, we let $v := (\mu, \eta)$. The case $j = 0$ is proved in [26, Lemma 5.1]. We will only treat the case $k = \omega$ and $j = 1$. The remaining cases follow in a similar way.

Given $v_0 \in \mathbb{B}(0, r_0)$, for every $v \in \mathbb{B}(v_0, r)$ with r small enough

$$f(v) = \sum_{\beta} \frac{1}{\beta!} \partial^{\beta} f(v_0)(v - v_0)^{\beta}, \quad \beta \in \mathbb{N}_0^{m+1},$$

and there exist constants M, R such that

$$\|\partial^{\beta} f(v)\| \leq M \frac{\beta!}{R^{|\beta|}}, \quad v \in \mathbb{B}(v_0, r), \quad \beta \in \mathbb{N}_0^{m+1}.$$

The constants r, M, R depend continuously on v_0 . Thus we can find uniform constants r, M, R for $\bar{\mathbb{B}}^{m+1}(0, |v_0|)$. For each $t \in J$ and $v \in \mathbb{B}(v_0, r)$, the following series

$$\sum_{\beta} \frac{\xi^{|\beta|}(t)}{\beta!} \partial^{\beta} f(\xi(t)v_0)(v - v_0)^{\beta}$$

converges in X and represents $F(v)(t)$. The temporal derivatives can be computed as follows. For $\beta \neq 0$,

$$\begin{aligned} & \left\| \frac{d}{dt} [\xi^{|\beta|}(t) \partial^{\beta} f(\xi(t)v_0)] \right\|_{C(J, X)} \\ & \leq \left\| \xi'(t) |\beta| \xi^{|\beta|-1}(t) \partial^{\beta} f(\xi(t)v_0) + \xi^{|\beta|}(t) \sum_{j=1}^{m+1} \partial^{\beta+e_j} f(\xi(t)v_0)(v_0)_j \right\|_{C(J, X)} \\ & \leq M \|\xi\|_{1, \infty} \left(|\beta| \frac{\beta!}{R^{|\beta|}} + r_0 \sum_{j=1}^{m+1} \frac{(\beta + e_j)!}{R^{|\beta|+1}} \right) \leq M' \frac{\beta!}{(R/C)^{|\beta|}} \end{aligned}$$

for some sufficiently large C . Now the assertion follows right away. □

Suppose that $O_{\epsilon} \subset \mathbb{R}^{m+1}$ is an open subset containing $\psi_{\epsilon, \kappa_c}(\bar{\mathbb{B}}_{3\epsilon_0, 7a/9})$. Put $O_{\epsilon_i} := O_{\epsilon} \cap \Omega_i$.

When $k \in \mathbb{N}_0 \cup \{\infty\}$, we say $u \in C^k(O_{\epsilon_i}, E) \cap \mathfrak{F}^s(\Omega_i, E)$ if $\mathcal{E}_i u \in C^k(O_{\epsilon}, E) \cap \mathfrak{F}^s(\Omega, E)$. By convention, $u \in C^{\omega}(O_{\epsilon_i}, E) \cap \mathfrak{F}^s(\Omega_i, E)$ means that u has an analytic extension $u^e \in C^{\omega}(O_{\epsilon}, E) \cap \mathfrak{F}^s(\Omega, E)$.

PROPOSITION 3.6 Suppose that $(S, V) \in \{(O_{\epsilon}, \Omega), (O_{\epsilon_i}, \Omega_i)\}$.

(a) Suppose that $u \in C^{n+k+j}(S, E) \cap \mathfrak{F}^s(V, E)$ for $j = 0, 1$, where either $s \in [0, n]$ if $\mathfrak{F} \in \{bc, W_p, H_p\}$, or $s = n$ if $\mathfrak{F} = BC$. Here $k \in \mathbb{N}_0 \cup \{\infty\}$ if $(S, V) = (O_{\epsilon_i}, \Omega_i)$. Then we have

$$[(\mu, \eta) \mapsto T_{\mu, \eta} u] \in C^k(\mathbb{B}(0, r_0), C^j(J; \mathfrak{F}^s(V, E))).$$

(b) Suppose that $\mathcal{Q} = \sum_{|\alpha| \leq l} a_{\alpha} \partial^{\alpha}$ with $a_{\alpha} \in C^{n+k+j}(S, \mathfrak{L}(E)) \cap BC^n(V, \mathfrak{L}(E))$ for $j = 0, 1$.

Then

$$[(\mu, \eta) \mapsto T_{\mu, \eta} \mathcal{Q} T_{\mu, \eta}^{-1}] \in C^k\left(\mathbb{B}(0, r_0), C^j\left(J; \mathfrak{L}(\mathfrak{F}^{s+l}(V, E), \mathfrak{F}^s(V, E))\right)\right),$$

for $s \in [0, n]$ if $\mathfrak{F} \in \{BC, W_p, H_p\}$, or $s \in [0, n]$ if $\mathfrak{F} = bc$.

Proof. (a) Modifying the proof of [73, Theorem 3.4] as in that of Proposition 3.4, one can show by means of Proposition 3.1(c) that

$$[(\mu, \eta) \mapsto \Theta_{\mu, \eta}^* u] \in C^k(\mathbb{B}(0, r_0), \mathfrak{F}^s(V, E)).$$

Set $X = \mathfrak{F}^s(V, E)$, $f(\mu, \eta) = \Theta_{\mu, \eta}^* u$. Now (a) is a direct consequence of Lemma 3.5.

(b) When $(S, V) = (\mathcal{O}_{\varepsilon_i}, \Omega_i)$, let $\mathfrak{Q}_i := \sum_{|\alpha| \leq l} (a_\alpha^\varepsilon) \partial^\alpha$ on Ω . Then

$$\mathfrak{Q}_{\mu, \eta} = \mathcal{R}_i \circ \Theta_{\mu, \eta}^* \mathfrak{Q}_i \Theta_{\mu, \eta}^* \circ \mathcal{E}_i.$$

The atlas \mathfrak{A}_ε is real analytically compatible with the Euclidean structure. Similar to the proof of [73, Proposition 3.6], by Theorem 3.2, we can show that

$$[(\mu, \eta) \mapsto \mathfrak{Q}_{\mu, \eta} := \Theta_{\mu, \eta}^* \mathfrak{Q} \Theta_{\mu, \eta}^*] \in C^k(\mathbb{B}(0, r_0), \mathfrak{L}(\mathfrak{F}^{s+l}(V, E), \mathfrak{F}^s(V, E))).$$

Set $X = \mathfrak{L}(\mathfrak{F}^{s+l}(V, E), \mathfrak{F}^s(V, E))$, and $f(\mu, \eta) = \mathfrak{Q}_{\mu, \eta}$. The assertion follows by Lemma 3.5. \square

Remark 3.7. Given any Banach space X , suppose that $u \in C^{n+k}(I, X) \cap \mathfrak{F}^s(J, X)$, where either $s \in [0, n]$ if $\mathfrak{F} \in \{bc, W_p, H_p\}$, or $s = n$ if $\mathfrak{F} = BC$. Following the proofs in [26, Section 3], we can verify that

$$[\lambda \mapsto \varrho_\lambda^* u] \in C^k(\mathbb{B}(0, r_0), \mathfrak{F}^s(J, X)).$$

\square

The localized translation $\Theta_{\mu, \eta}$, being restricted to Σ , induces a family of diffeomorphism $\{\Theta_\mu^* : \mu \in \mathbb{B}(0, r_0)\}$ on function spaces over Σ , which was introduced in [73]. For the reader's convenience, we will briefly state some of its properties herein.

$$\Theta_\mu(q) = \begin{cases} \psi_{\kappa_c}(\theta_\mu(\varphi_{\kappa_c}(q))), & q \in \mathcal{O}_{\kappa_c}, \\ q, & q \in \Sigma \setminus \mathcal{O}_{\kappa_c}. \end{cases}$$

It is evident that $\Theta_\mu \in \text{Diff}^\infty(\Sigma)$ for $\mu \in \mathbb{B}(0, r_0)$ with sufficiently small r_0 . Given $u \in E^\Sigma$, Θ_μ^* and Θ_μ^μ can be explicitly expressed as

$$\Theta_\mu^* u = \varphi_{\kappa_c}^* \theta_\mu^* \psi_{\kappa_c}^*(\zeta u) + (1_\Sigma - \zeta)u, \quad \Theta_\mu^\mu u = \varphi_{\kappa_c}^* \theta_\mu^\mu \psi_{\kappa_c}^*(\zeta u) + (1_\Sigma - \zeta)u,$$

respectively. Given $u \in E^{J \times \Sigma}$, setting $T_\mu(t) := \Theta_{\xi(t)\mu}^*$, we define

$$u_{\lambda, \mu}(t, \cdot) := \Theta_{\lambda, \mu}^* u(t, \cdot) = T_\mu(t) \varrho_\lambda^* u(t, \cdot), \quad (\lambda, \mu) \in \mathbb{B}(0, r_0).$$

Let $\Theta_{\lambda, \mu}^{\lambda, \mu} := [\Theta_{\lambda, \mu}^*]^{-1}$.

A linear operator $\mathfrak{Q} : C^\infty(\Sigma, E) \rightarrow E^\Sigma$ is called a linear differential operator of order l on Σ if in every local chart $(\mathcal{O}_\kappa, \varphi_\kappa)$ with $\kappa \in \mathfrak{R}$, there exists some linear differential operator defined on \mathbb{Q}^m

$$\mathfrak{Q}_\kappa(x, \partial) := \sum_{|\alpha| \leq l} a_\alpha^\kappa(x) \partial^\alpha, \quad \text{with } a_\alpha^\kappa \in \mathfrak{L}(E)^{\mathbb{Q}^m},$$

called the local representation of \mathfrak{Q} in $(\mathcal{O}_\kappa, \varphi_\kappa)$, such that for any $u \in C^\infty(\Sigma, E)$

$$\psi_\kappa^*(\mathfrak{Q}u) = \mathfrak{Q}_\kappa(\psi_\kappa^* u), \quad \kappa \in \mathfrak{R}. \quad (3.8)$$

PROPOSITION 3.8 Let $\mathcal{O} := \mathcal{O}_c \cap \Sigma$.

(a) Suppose that $u \in C^{n+k+j}(\Sigma, E) \cap \mathfrak{F}^s(\Sigma, E)$ for $j = 0, 1$, where either $s \in [0, n]$ if $\mathfrak{F} \in \{bc, W_p, H_p\}$, or $s = n$ if $\mathfrak{F} = BC$. Then we have

$$[\mu \mapsto T_\mu u] \in C^k(\mathbb{B}(0, r_0), C^j(J; \mathfrak{F}^s(\Sigma, E))).$$

(b) Suppose that \mathcal{Q} is a linear differential operator on Σ of order l satisfying that for all $|\alpha| \leq l$ and $\kappa \in \mathfrak{K}$, $a_\alpha^\kappa \in BC^n(\mathbb{Q}^m, \mathfrak{L}(E))$ and $a_\alpha^{\kappa_c} \in C^{n+k+j}(\varphi_{\kappa_c}(\mathcal{O}), \mathfrak{L}(E))$ for $j = 0, 1$. Then for $s \in [0, n]$ if $\mathfrak{F} = BC$, or $s \in [0, n)$ if $\mathfrak{F} = bc$, or $s \in (-\infty, n]$ if $\mathfrak{F} \in \{W_p, H_p\}$

$$[\mu \mapsto T_\mu \mathcal{Q} T_\mu^{-1}] \in C^k(\mathbb{B}(0, r_0), C^j(J; \mathfrak{L}(\mathfrak{F}^{s+l}(\Sigma, E), \mathfrak{F}^s(\Sigma, E)))).$$

Proof. The proofs for (a) and the case $s \geq 0$ in (b) are given in [73, Section 3]. We will only treat the case $k = \omega$ when $s < 0$, the other cases follow similarly. Firstly, we show that for any $s \in (-\infty, 0)$ and $\mathfrak{F}_p \in \{W_p, H_p\}$

$$[\mu \mapsto \theta_\mu^* \partial_j \theta_\mu^\mu] \in C^\omega(\mathbb{B}(0, r_0), \mathfrak{L}(\mathfrak{F}_p^{s+1}(\mathbb{R}^m, E), \mathfrak{F}_p^s(\mathbb{R}^m, E))).$$

(i) $s \leq -1$. On account that with p' denoting the Hölder duality of p ,

$$\partial_j \in \mathfrak{L}(\mathfrak{F}_p^{s+1}(\mathbb{R}^m, E), \mathfrak{F}_p^s(\mathbb{R}^m, E)), \quad \text{and} \quad \mathfrak{F}_p^s(\mathbb{R}^m, E) = (\mathfrak{F}_{p'}^{-s}(\mathbb{R}^m, E))',$$

for every $u \in \mathfrak{F}_p^{s+1}(\mathbb{R}^m, E)$ and $v \in \mathfrak{F}_{p'}^{-s}(\mathbb{R}^m, E)$, we have

$$\begin{aligned} \langle \theta_\mu^* \partial_j \theta_\mu^\mu u, v \rangle &:= -\langle u, \theta_\mu^* \partial_j [\theta_\mu^\mu v | \det(D(\theta_\mu)^{-1})] | \det(D\theta_\mu) \rangle \\ &= -\langle u, \theta_\mu^* \partial_j \theta_\mu^\mu v \theta_\mu^* (\det(D(\theta_\mu)^{-1})) \det(D\theta_\mu) \rangle - \langle u, v \theta_\mu^* \partial_j [\det(D(\theta_\mu)^{-1})] \det(D\theta_\mu) \rangle. \end{aligned}$$

Here $\langle \cdot, \cdot \rangle$ denotes the duality pairing from $\mathfrak{F}_p^s(\mathbb{R}^m, E) \times \mathfrak{F}_{p'}^{-s}(\mathbb{R}^m, E)$ to \mathbb{R} . By [26, Proposition 4.1], we have

$$[\mu \mapsto \theta_\mu^* \partial_j \theta_\mu^\mu] \in C^\omega(\mathbb{B}(0, r_0), \mathfrak{L}(\mathfrak{F}_{p'}^{-s}(\mathbb{R}^m, E), \mathfrak{F}_{p'}^{-s-1}(\mathbb{R}^m, E))).$$

It is a simple matter to check that $\theta_\mu^* (\det(D(\theta_\mu)^{-1})) \det(D\theta_\mu) = 1$. One may compute

$$\begin{aligned} &-\theta_\mu^* \partial_j [\det(D(\theta_\mu)^{-1})] \det(D\theta_\mu) \\ &= -\theta_\mu^* [\partial_j [\det(D(\theta_\mu)^{-1})] \theta_\mu^\mu \det(D\theta_\mu)] = \theta_\mu^* \partial_j \theta_\mu^\mu \det(D\theta_\mu) \theta_\mu^* \det(D(\theta_\mu)^{-1}). \end{aligned}$$

We immediately have for all $k \in \mathbb{N}_0$ that

$$[\mu \mapsto \det(D\theta_\mu)] \in C^\omega(\mathbb{B}(0, r_0), BC^{k+1}(\mathbb{R}^m)).$$

It follows again from [26, Proposition 4.1] that

$$[\mu \mapsto \theta_\mu^* \partial_j \theta_\mu^\mu \det(D\theta_\mu)] \in C^\omega(\mathbb{B}(0, r_0), BC^k(\mathbb{R}^m)).$$

On the other hand, we obtain

$$[\mu \mapsto \theta_\mu^* \det(D(\theta_\mu)^{-1})] = 1 / \det(D\theta_\mu) \in C^\omega(\mathbb{B}(0, r_0), BC^k(\mathbb{R}^m)).$$

Therefore, for every $u \in \mathfrak{F}_p^{s+1}(\mathbb{R}^m, E)$ and $v \in \mathfrak{F}_p^{-s}(\mathbb{R}^m, E)$,

$$[\mu \mapsto \langle \theta_\mu^* \partial_j \theta_\mu^\mu, v \rangle] \in C^\omega(\mathbb{B}(0, r_0)).$$

Now [13, Proposition 1] implies that

$$[\mu \mapsto \theta_\mu^* \partial_j \theta_\mu^\mu] \in C^\omega\left(\mathbb{B}(0, r_0), \mathfrak{L}(\mathfrak{F}_p^{s+1}(\mathbb{R}^m, E), \mathfrak{F}_p^s(\mathbb{R}^m, E))\right).$$

(ii) $s \in (-1, 0)$. [26, Proposition 4.1] and (i) show that

$$\begin{aligned} [\mu \mapsto \theta_\mu^* \partial_j \theta_\mu^\mu] \in C^\omega\left(\mathbb{B}(0, r_0), \mathfrak{L}(\mathfrak{F}_p^{s+2}(\mathbb{R}^m, E), \mathfrak{F}_p^{s+1}(\mathbb{R}^m, E))\right) \\ \cap C^\omega\left(\mathbb{B}(0, r_0), \mathfrak{L}(\mathfrak{F}_p^s(\mathbb{R}^m, E), \mathfrak{F}_p^{s-1}(\mathbb{R}^m, E))\right). \end{aligned}$$

Thus for any $\mu_0 \in \mathbb{B}(0, r_0)$, there exists some constants r_i, M_i and R_i with $i = 1, 2$ such that for all $\mu \in \mathbb{B}^m(\mu_0, r_i)$

$$\left\| \frac{\partial^\alpha}{\partial \mu^\alpha} [\theta_\mu^* \partial_j \theta_\mu^\mu] \right\|_{X_i} \leq M_i \frac{\alpha!}{R_i^{|\alpha|}}, \quad \alpha \in \mathbb{N}_0^m. \quad (3.9)$$

Here $X_1 := \mathfrak{L}(\mathfrak{F}_p^{s+2}, \mathfrak{F}_p^{s+1})$ and $X_2 := \mathfrak{L}(\mathfrak{F}_p^s, \mathfrak{F}_p^{s-1})$. It follows from interpolation theory that $[\mu \mapsto \theta_\mu^* \partial_j \theta_\mu^\mu] \in C^\infty(\mathbb{B}(0, r_0), \mathfrak{L}(\mathfrak{F}_p^{s+1}(\mathbb{R}^m, E), \mathfrak{F}_p^s(\mathbb{R}^m, E)))$. Indeed, we have for $h = 1, 2, \dots, m$ and any $\mu \in \mathbb{B}(0, r_0)$

$$\left\| \frac{\theta_{\mu+te_h}^* \partial_j \theta_{\mu+te_h}^{\mu+te_h} - \theta_\mu^* \partial_j \theta_\mu^\mu}{t} - \frac{\partial}{\partial \mu_h} \theta_\mu^* \partial_j \theta_\mu^\mu \right\|_{X_i} \rightarrow 0, \quad t \rightarrow 0, \quad i = 1, 2.$$

By interpolation theory, the above limits converge when X_i is replaced by $X := \mathfrak{L}(\mathfrak{F}_p^{s+1}(\mathbb{R}^m, E), \mathfrak{F}_p^s(\mathbb{R}^m, E))$. Continuity, or even C^∞ -smoothness, of $\theta_\mu^* \partial_j \theta_\mu^\mu$ in X can be verified analogously.

A similar Cauchy estimate to (3.9) holds for X by interpolation theory, i.e., for any $\mu_0 \in \mathbb{B}(0, r_0)$, there exists some constants $r = \min r_i, M = \max M_i$ and $R = \min R_i$ such that for all $\mu \in \mathbb{B}^m(\mu_0, r)$

$$\left\| \frac{\partial^\alpha}{\partial \mu^\alpha} [\theta_\mu^* \partial_j \theta_\mu^\mu] \right\|_{\mathfrak{L}(\mathfrak{F}_p^{s+1}, \mathfrak{F}_p^s)} \leq M \frac{\alpha!}{R^{|\alpha|}}, \quad \alpha \in \mathbb{N}_0^m.$$

It is well-known that this estimate implies that

$$[\mu \mapsto \theta_\mu^* \partial_j \theta_\mu^\mu] \in C^\omega\left(\mathbb{B}(0, r_0), \mathfrak{L}(\mathfrak{F}_p^{s+1}(\mathbb{R}^m, E), \mathfrak{F}_p^s(\mathbb{R}^m, E))\right).$$

(iii) For $s \in (-\infty, n]$, in view of the proofs for [26, Proposition 4.1, Theorem 4.2], we thus infer that for any linear differential operator $\tilde{\mathcal{Q}} = \sum_{|\alpha| \leq l} a_\alpha \partial^\alpha$, if $a_\alpha \in BC^n(\mathbb{R}^m, \mathfrak{L}(E)) \cap C^{n+k+j}(O, \mathfrak{L}(E))$ for $O := \varphi_{\kappa_c}(O)$, then

$$[\mu \mapsto \theta_\mu^* \tilde{\mathcal{Q}} \theta_\mu^\mu] \in C^\omega\left(\mathbb{B}(0, r_0), \mathfrak{L}(\mathfrak{F}_p^{s+l}(\mathbb{R}^m, E), \mathfrak{F}_p^s(\mathbb{R}^m, E))\right).$$

A similar proof to (i) shows that [73, Lemma 3.1] still holds for Sobolev–Slobodeckii and Bessel potential spaces of negative order. Indeed, let

$$\mathfrak{F}_{\text{cp}}^{s,\Sigma} := \{u \in \mathfrak{F}^s(\Sigma, E) : \text{supp}(u) \subset \psi_{\kappa_c}(\bar{\mathbb{B}}(x_c, 5\varepsilon_0))\},$$

and

$$\mathfrak{F}_{\text{cp}}^{s,\mathbb{R}^m} := \{u \in \mathfrak{F}^s(\mathbb{R}^m, E) : \text{supp}(u) \subset \bar{\mathbb{B}}(x_c, 5\varepsilon_0)\}.$$

Pick $\bar{\mathbb{B}}(x_c, 5\varepsilon_0) \subset \mathring{U} \subset\subset \mathcal{Q}^m$ with U closed, and $\tau \in \mathfrak{D}(\mathring{U}, [0, 1])$ with $\tau|_{\bar{\mathbb{B}}(x_c, 5\varepsilon_0)} \equiv 1$. For $r > 0$, given any $u \in W_{p,\text{cp}}^r$ and $v \in W_{p'}^{-r}(\mathbb{R}^m, E)$, we have

$$\begin{aligned} |\langle u, \varphi_{\kappa_c}^* v \rangle_{\Sigma}| &= |\langle \psi_{\kappa_c}^* u, \tau v \sqrt{|\det G|} \rangle| \\ &\leq M \|\psi_{\kappa_c}^* u\|_{W_p^r(\mathbb{R}^m)} \|v\|_{W_{p'}^{-r}(\mathbb{R}^m)} \leq M \|u\|_{W_p^r(\Sigma)} \|v\|_{W_{p'}^{-r}(\mathbb{R}^m)}. \end{aligned}$$

Here G is the local matrix expression of the metric g , and $\langle \cdot, \cdot \rangle_{\Sigma}$ denotes the duality pairing from $\mathfrak{F}_p^r(\Sigma, E) \times \mathfrak{F}_{p'}^{-r}(\Sigma, E)$ to \mathbb{R} . The ultimate line follows from the point-wise multiplier theorem, see [5, Section 9], and a similar assertion to [73, Lemma 3.1]. Thus the constant M is independent of u and v . It implies that for $r \in \mathbb{R}$

$$\varphi_{\kappa_c}^* \in \mathfrak{L}is(\mathfrak{F}_{\text{cp}}^{r,\mathbb{R}^m}, \mathfrak{F}_{\text{cp}}^{r,\Sigma}), \quad \text{with} \quad [\varphi_{\kappa_c}^*]^{-1} = \psi_{\kappa_c}^*.$$

Modifying the proof [73, Proposition 3.6] in an obvious way and applying Lemma 3.5, we have proved the statement of (b). \square

Recall that $\partial_v = \partial_{v\Sigma}$.

PROPOSITION 3.9 $T_{\mu} \partial_v T_{\mu,\eta}^{-1} = \partial_v$ on $H_p^1(\Omega_i)$.

Proof. It suffices to show that $\Theta_{\mu}^* \partial_v \Theta_{\mu}^{\mu,\eta} = \partial_v$. On account of

$$\psi_{\kappa}^*(\partial_v u) = \psi_{\kappa}^*(v_{\Sigma} \cdot \nabla u) = \partial_{m+1} \psi_{\kappa}^* \mathcal{E}_i u$$

for any $u \in H_p^1(\Omega_i)$, one readily computes

$$\begin{aligned} \Theta_{\mu}^* \partial_v \Theta_{\mu}^{\mu,\eta} u &= \Theta_{\mu}^* \partial_v \varphi_{\varepsilon,\kappa_c}^* \theta_{\mu}^{\mu} \psi_{\varepsilon,\kappa_c}^* \zeta_{\varepsilon} \mathcal{E}_i u + \Theta_{\mu}^* \partial_v (1 - \zeta_{\varepsilon}) u \\ &= \varphi_{\kappa_c}^* \theta_{\mu}^* \psi_{\kappa_c}^* \zeta_{\varepsilon} \partial_v \varphi_{\varepsilon,\kappa_c}^* \theta_{\mu}^{\mu} \psi_{\varepsilon,\kappa_c}^* \zeta_{\varepsilon} \mathcal{E}_i u + (1 - \zeta_{\varepsilon}) \partial_v \varphi_{\varepsilon,\kappa_c}^* \theta_{\mu}^{\mu} \psi_{\varepsilon,\kappa_c}^* \zeta_{\varepsilon} \mathcal{E}_i u + \partial_v (1 - \zeta_{\varepsilon}) u \\ &= \varphi_{\kappa_c}^* \theta_{\mu}^* \psi_{\kappa_c}^* \zeta_{\varepsilon} \varphi_{\kappa_c}^* \partial_{m+1} \theta_{\mu}^{\mu} \psi_{\varepsilon,\kappa_c}^* \zeta_{\varepsilon} \mathcal{E}_i u \\ &\quad + (1 - \zeta_{\varepsilon}) \varphi_{\kappa_c}^* \partial_{m+1} \theta_{\mu}^{\mu} \psi_{\varepsilon,\kappa_c}^* \zeta_{\varepsilon} \mathcal{E}_i u + \partial_v (1 - \zeta_{\varepsilon}) u \\ &= \varphi_{\kappa_c}^* \theta_{\mu}^* \psi_{\kappa_c}^* \zeta_{\varepsilon} \varphi_{\kappa_c}^* \theta_{\mu}^{\mu} \partial_{m+1} \psi_{\varepsilon,\kappa_c}^* \zeta_{\varepsilon} \mathcal{E}_i u + \varphi_{\kappa_c}^* \theta_{\mu}^* \psi_{\kappa_c}^* (1 - \zeta_{\varepsilon}) \varphi_{\kappa_c}^* \theta_{\mu}^{\mu} \partial_{m+1} \psi_{\varepsilon,\kappa_c}^* \zeta_{\varepsilon} \mathcal{E}_i u \\ &\quad + \partial_v (1 - \zeta_{\varepsilon}) u \\ &= \varphi_{\kappa_c}^* \partial_{m+1} \psi_{\varepsilon,\kappa_c}^* \zeta_{\varepsilon} \mathcal{E}_i u + \partial_v (1 - \zeta_{\varepsilon}) u = \partial_v u. \end{aligned}$$

In the above, we have used the fact that $\Theta_{\mu}^* \partial_v (1 - \zeta_{\varepsilon}) u = (1 - \zeta_{\varepsilon}) u$. \square

Recall definitions (3.6) and (3.7). The following proposition shows that the space $\mathbb{E}(J)$ is invariant under the transformation $\Theta_{\lambda,\mu,\eta}^*$ and $\Theta_{\lambda,\mu}^*$, respectively.

PROPOSITION 3.10 $[(\vartheta, h) \mapsto (\Theta_{\lambda, \mu, \eta}^* \vartheta, \Theta_{\lambda, \mu}^* h)] \in \mathfrak{Lis}(\mathbb{E}(J)) \cap \mathfrak{Lis}({}_0\mathbb{E}(J))$ with $(\lambda, \mu, \eta) \in \mathbb{B}(0, r_0)$. Moreover, there exist some $B_{\lambda, \mu, \eta}, B_{\lambda, \mu}$ satisfying

$$\begin{cases} [(\lambda, \mu, \eta) \mapsto B_{\lambda, \mu, \eta}] \in C^\omega(\mathbb{B}(0, r_0), \mathfrak{L}(\mathbb{E}_1(J), \mathbb{F}_1(J))), \\ [(\lambda, \mu) \mapsto B_{\lambda, \mu}] \in C^\omega(\mathbb{B}(0, r_0), \mathfrak{L}(\mathbb{E}_2(J), \mathbb{F}_2(J))), \end{cases}$$

such that

$$\begin{cases} \partial_t [\vartheta_{\lambda, \mu, \eta}] = (1 + \xi' \lambda) \Theta_{\lambda, \mu, \eta}^* \partial_t \vartheta + B_{\lambda, \mu, \eta}(\vartheta_{\lambda, \mu, \eta}), \\ \partial_t [h_{\lambda, \mu}] = (1 + \xi' \lambda) \Theta_{\lambda, \mu}^* \partial_t h + B_{\lambda, \mu}(h_{\lambda, \mu}). \end{cases}$$

In particular, $B_{\lambda, 0, 0} = 0$ and $B_{\lambda, 0} = 0$.

Proof. (i) Following the proof of [26, Proposition 2.4] and interpolation theory, we can show that for any Banach space X , $\varrho_\lambda^* \in \mathfrak{Lis}(\mathfrak{F}(I, X))$ and $[\varrho_\lambda^*]^{-1} = \varrho_\lambda^*$ with $\mathfrak{F} \in \{BC^s, W_p^s\}$. In particular, there exists $M > 0$ such that $\|\varrho_\lambda^*\|_{\mathfrak{L}(\mathfrak{F}(I, X))} \leq M$ for $\lambda \in \mathbb{B}$. A similar estimate as in [26, Lemma 8.3] by using the intrinsic norms of Besov spaces reveals that

$$[(u, \rho) \mapsto (\tilde{u}_{\lambda, \mu, \eta}, \tilde{\rho}_{\lambda, \mu})] \in \mathfrak{Lis}(\mathbb{E}_1(J, \mathbb{R}^{m+1}; E) \times \mathbb{E}_2(J, \mathbb{R}^m; E)).$$

(ii) Observe that

$$(\Theta_{\lambda, \mu, \eta}^* \vartheta, \Theta_{\lambda, \mu}^* h)(0) = (\vartheta, h)(0), \quad \partial_t (\Theta_{\lambda, \mu, \eta}^* \vartheta, \Theta_{\lambda, \mu}^* h)(0) = \partial_t (\vartheta, h)(0), \quad (3.10)$$

whenever the corresponding derivative exists. For any $z \in \Omega \setminus \mathcal{O}_{\epsilon, \kappa_c}$, $T_{\mu, \eta} \vartheta(z) = \vartheta(z)$, we conclude that $\partial_{v_\Omega} T_{\mu, \eta} \vartheta = 0$. For every $t \in J$, $\vartheta(t) \in BC(\Omega)$. It follows from Proposition 3.4 that $T_{\mu, \eta}(t) \vartheta(t) \in BC(\Omega)$, which implies that $\llbracket T_{\mu, \eta}(t) \vartheta(t) \rrbracket = 0$. Moreover, for $(\mu, \eta) \in \mathbb{B}(0, r_0)$

$$\begin{aligned} \|T_{\mu, \eta} \vartheta\|_{\mathbb{E}_1(J)} &= \sum_i \|\mathcal{R}_i \varphi_{\epsilon, \kappa_c}^* \tilde{T}_{\mu, \eta} \psi_{\epsilon, \kappa_c}^* (\zeta_\epsilon \mathcal{E}_i \vartheta) + (1 - \zeta_\epsilon) \vartheta\|_{\mathbb{E}_1(J)} \\ &\leq M \sum_i [\|\tilde{T}_{\mu, \eta} \psi_{\epsilon, \kappa_c}^* (\zeta_\epsilon \mathcal{E}_i \vartheta)\|_{\mathbb{E}_1(J, \mathbb{R}^{m+1}; E)} + \|(1 - \zeta_\epsilon) \mathcal{E}_i \vartheta\|_{\mathbb{E}_1(J)}] \leq M \|\vartheta\|_{\mathbb{E}_1(J)}. \end{aligned}$$

The last line follows from (i), Lemma 3.3 and point-wise multiplier theorem, see [75, Section 3.3.2]. In virtue of [73, Lemma 3.1], one can show similarly that

$$[h \mapsto T_\mu h] \in \mathfrak{L}(\mathbb{E}_2(J)), \quad \mu \in \mathbb{B}(0, r_0).$$

Then it is a direct consequence of the open mapping theorem that

$$[(\vartheta, h) \mapsto (\Theta_{\lambda, \mu, \eta}^* \vartheta, \Theta_{\lambda, \mu}^* h)] \in \mathfrak{Lis}(\mathbb{E}(J)).$$

By (3.10), the statement $[(\vartheta, h) \mapsto (T_{\mu, \eta} \vartheta, T_\mu h)] \in \mathfrak{Lis}({}_0\mathbb{E}(J))$ is immediate.

(iii) The temporal derivative of $\vartheta_{\lambda,\mu,\eta}$ can be computed as follows.

$$\begin{aligned} \partial_t[\vartheta_{\lambda,\mu,\eta}] &= (1 - \varsigma_\epsilon)\partial_t[\varrho_\lambda^*\vartheta] + \sum_i \mathcal{R}_i\varphi_{\epsilon,\kappa_c}^* \partial_t[\tilde{T}_{\mu,\eta}\psi_{\epsilon,\kappa_c}^*(\varsigma_\epsilon\varrho_\lambda^*\mathcal{E}_i\vartheta)] \\ &= (1 - \varsigma_\epsilon)(1 + \xi'\lambda)\varrho_\lambda^*\partial_t\vartheta + \sum_i \left\{ \mathcal{R}_i\varphi_{\epsilon,\kappa_c}^* \tilde{T}_{\mu,\eta}\psi_{\epsilon,\kappa_c}^*(\varsigma_\epsilon\partial_t[\varrho_\lambda^*\mathcal{E}_i\vartheta]) \right. \\ &\quad \left. + \mathcal{R}_i \sum_{j=1}^m \varphi_{\epsilon,\kappa_c}^*(\xi'\chi_m\varpi\mu_j\tilde{T}_{\mu,\eta}\partial_j[\psi_{\epsilon,\kappa_c}^*\varsigma_\epsilon\varrho_\lambda^*\mathcal{E}_i\vartheta]) \right. \\ &\quad \left. + \mathcal{R}_i\varphi_{\epsilon,\kappa_c}^*(\xi'\chi_m\chi\eta\tilde{T}_{\mu,\eta}\partial_{m+1}[\psi_{\epsilon,\kappa_c}^*\varsigma_\epsilon\varrho_\lambda^*\mathcal{E}_i\vartheta]) \right\} \\ &= (1 + \xi'\lambda)\Theta_{\lambda,\mu,\eta}^*\partial_t\vartheta + B_{\lambda,\mu,\eta}(\vartheta_{\lambda,\mu,\eta}). \end{aligned}$$

The last line follows from the definition of $\Theta_{\lambda,\mu,\eta}^*$ and the equality

$$\begin{aligned} &\mathcal{R}_i\varphi_{\epsilon,\kappa_c}^* \tilde{T}_{\mu,\eta}\psi_{\epsilon,\kappa_c}^*(\varsigma_\epsilon\partial_t[\varrho_\lambda^*\mathcal{E}_i\vartheta]) + (1 - \varsigma_\epsilon)(1 + \xi'\lambda)\varrho_\lambda^*\partial_t\vartheta \\ &= \mathcal{R}_i\varphi_{\epsilon,\kappa_c}^* \tilde{T}_{\mu,\eta}\psi_{\epsilon,\kappa_c}^*[\varsigma_\epsilon(1 + \xi'\lambda)\varrho_\lambda^*\partial_t\mathcal{E}_i\vartheta] + (1 - \varsigma_\epsilon)(1 + \xi'\lambda)\varrho_\lambda^*\partial_t\vartheta \\ &= (1 + \xi'\lambda)\Theta_{\lambda,\mu,\eta}^*\partial_t\vartheta. \end{aligned}$$

Given $u \in H_p^1(\Omega \setminus \Sigma)$, the term $B_{\lambda,\mu,\eta}$ can be explicitly written as

$$\begin{aligned} B_{\lambda,\mu,\eta}(u) &= \sum_i \left\{ \sum_{j=1}^m \mathcal{R}_i\varphi_{\epsilon,\kappa_c}^*(\xi'\chi_m\varpi\mu_j\tilde{T}_{\mu,\eta}\partial_j[\psi_{\epsilon,\kappa_c}^*\varsigma_\epsilon T_{\mu,\eta}^{-1}\mathcal{E}_i u]) \right\} \\ &\quad + \mathcal{R}_i\varphi_{\epsilon,\kappa_c}^*(\xi'\chi_m\chi\eta\tilde{T}_{\mu,\eta}\partial_{m+1}[\psi_{\epsilon,\kappa_c}^*\varsigma_\epsilon T_{\mu,\eta}^{-1}\mathcal{E}_i u]) \\ &= \sum_i \left\{ \sum_{j=1}^m \mathcal{R}_i\xi'\varphi_{\epsilon,\kappa_c}^*\chi_m\varphi_{\epsilon,\kappa_c}^*\varpi\mu_j T_{\mu,\eta}\mathcal{Q}^j T_{\mu,\eta}^{-1}\mathcal{E}_i u \right. \\ &\quad \left. + \mathcal{R}_i\xi'\varphi_{\epsilon,\kappa_c}^*\chi_m\varphi_{\epsilon,\kappa_c}^*\chi\eta T_{\mu,\eta}\mathcal{Q}^{m+1} T_{\mu,\eta}^{-1}\mathcal{E}_i u \right\}. \end{aligned}$$

Here \mathcal{Q}^j with $j = 1, \dots, m + 1$ are first order linear differential operators defined on $\Omega \setminus \Sigma$ compactly supported in τ_{a,κ_c} satisfying $\psi_{\epsilon,\kappa_c}^* \mathcal{Q}^j u = \varsigma_\epsilon \partial_j \psi_{\epsilon,\kappa_c}^* u$ for all $u \in H_p^1(\Omega \setminus \Sigma)$. By means of Proposition 3.6(b) and the real analytic compatibility of the atlas \mathfrak{A}_ϵ with the Euclidean structure, it is not hard to check that

$$[(\lambda, \mu, \eta) \mapsto B_{\lambda,\mu,\eta}] \in C^\omega\left(\mathbb{B}(0, r_0), C^1\left(J; \mathfrak{L}(W_p^{s+1}(\Omega \setminus \Sigma), W_p^s(\Omega \setminus \Sigma))\right)\right). \tag{3.11}$$

In [73, Proposition 3.10], it is shown that

$$\partial_t[h_{\lambda,\mu}] = (1 + \xi'\lambda)\Theta_{\lambda,\mu}^*\partial_t h + B_{\lambda,\mu}(h_{\lambda,\mu})$$

with $B_{\lambda,0} = 0$ and

$$[(\lambda, \mu) \mapsto B_{\lambda,\mu}] \in C^\omega\left(\mathbb{B}(0, r_0), C^1\left(J; \mathfrak{L}(W_p^{s+1}(\Sigma), W_p^s(\Sigma))\right)\right). \tag{3.12}$$

This completes the proof. □

With minor modification of the above proof, we immediately have the following similar result.

PROPOSITION 3.11 For $(\lambda, \mu, \eta) \in \mathbb{B}(0, r_0)$ with sufficiently small $r_0 > 0$,

$$[(f, g, q) \mapsto (f_{\lambda, \mu, \eta}, g_{\lambda, \mu}, q_{\lambda, \mu})] \in \mathfrak{Lis}\left(\prod_{i=1}^3 \mathbb{F}_i(J), \prod_{i=1}^3 \mathbb{F}_i(J)\right),$$

and

$$[(f, g, q) \mapsto (f_{\lambda, \mu, \eta}, g_{\lambda, \mu}, q_{\lambda, \mu})] \in \mathfrak{Lis}\left(\prod_{i=1}^3 {}_0\mathbb{F}_i(J), \prod_{i=1}^3 {}_0\mathbb{F}_i(J)\right).$$

The main theorem of this section is

THEOREM 3.12 Suppose that

$$(\vartheta, h) \in BC(J \times \Omega, E) \times BC(J \times \Sigma, E).$$

Then

$$(\vartheta, h) \in C^k(\mathring{J} \times \mathbb{T}_{a/3} \setminus \Sigma, E) \times C^k(\mathring{J} \times \Sigma, E)$$

iff for any $(t_c, z_c) \in \mathring{J} \times \mathbb{T}_{a/3} \setminus \Sigma$, there exists some $r_0 = r_0(t_c, z_c) > 0$ and a corresponding family of parameter-dependent diffeomorphisms $\{(\Theta_{\lambda, \mu, \eta}^*, \Theta_{\lambda, \mu}^*) : (\lambda, \mu, \eta) \in \mathbb{B}(0, r_0)\}$ such that

$$[(\lambda, \mu, \eta) \mapsto (\vartheta_{\lambda, \mu, \eta}, h_{\lambda, \mu})] \in C^k(\mathbb{B}(0, r_0), BC(J \times \Omega, E) \times BC(J \times \Sigma, E)).$$

Proof. The proof follows in a similar manner to that of [73, Theorem 3.5]. \square

4. Regularity of solutions to the Stefan Problem

Throughout the rest of this paper, we assume that $\sigma > 0$ and $d_i, \gamma \in C^{k+2}(0, \infty)$, $\psi_i \in C^{k+3}(0, \infty)$ for $i = 1, 2$ with $k \in \mathbb{N} \cup \{\infty, \omega\}$ such that

$$\kappa_i(u) = -u\psi_i''(u) > 0, \quad d_i(u) > 0, \quad u \in (0, \infty),$$

and

$$\gamma(u) > 0, \quad u \in (0, \infty), \quad \text{or} \quad \gamma \equiv 0.$$

Let $J := [0, T]$. We define

$$\begin{aligned} \mathbb{G}_2(J) &:= H_p^1(J; W_p^{-1/p}(\Sigma)) \cap L_p(J; W_p^{2-1/p}(\Sigma)) \\ \mathbb{G}_3(J) &:= H_p^1(J; W_p^{-1-1/p}(\Sigma)) \cap L_p(J; W_p^{1-1/p}(\Sigma)), \end{aligned}$$

and

$$\begin{aligned} \mathbb{C}_1(J) &:= C(J \times \bar{\Omega}) \cap C(J; C^1(\bar{\Omega}_i)) \\ \mathbb{C}_2(J) &:= C(J; C^3(\Sigma)) \cap C^1(J; C(\Sigma)). \end{aligned}$$

It is shown in [65, Section 3] that

$$\mathbb{E}_1(J) \hookrightarrow \mathbb{C}_1(J), \quad \mathbb{E}_2(J) \hookrightarrow \mathbb{C}_2(J). \quad (4.1)$$

Observe that the constants in these embeddings blow up as $T \rightarrow 0$, however, they are uniform in T if one considers the space ${}_0\mathbb{E}(J)$!

4.1 Linearization at a real analytic temperature and the initial interface

Suppose that $\hat{z}_0 = (\hat{\vartheta}_0, \hat{h}_0) \in \mathbb{F}_4$ satisfies $\hat{\vartheta}_0 > 0$. Given any $\varepsilon > 0$, by the Weierstrass approximation theorem, we can find some

$$\vartheta_A \in C^\omega(\bar{\Omega}), \quad \|\vartheta_A - \hat{\vartheta}_0\|_{BC(\Omega)} \leq \varepsilon.$$

For sufficiently small ε , $\vartheta_A > 0$, and when $\gamma \equiv 0$, $l(\vartheta_A) \neq 0$. Let

$$\begin{aligned} \kappa_A(x) &= \kappa(\vartheta_A(x)), \quad d_A(x) = d(\vartheta_A(x)), \quad l_A(x) = l(\vartheta_A(x)), \quad \sigma_0 = \frac{\sigma}{m}, \\ l_1(t, \cdot) &= \llbracket \psi'(e^{\Delta_\Sigma t} \vartheta_A) \rrbracket, \quad \gamma_1(t, \cdot) = \gamma(e^{\Delta_\Sigma t} \vartheta_A). \end{aligned}$$

For a function $\vartheta \in \mathbb{E}_1(J)$, we do not distinguish $\vartheta|_\Sigma$ from ϑ if the choice is self-evident from the context. Here Δ_Σ denotes the Laplace-Beltrami operator on Σ . Similarly, we define

$$\begin{aligned} \hat{\kappa}_0(x) &= \kappa(\hat{\vartheta}_0(x)), \quad \hat{d}_0(x) = d(\hat{\vartheta}_0(x)), \quad \hat{l}_0(x) = l(\hat{\vartheta}_0(x)), \quad \hat{\gamma}_0 = \gamma(\hat{\vartheta}_0), \\ \hat{l}_1(t, \cdot) &= \llbracket \psi'(e^{\Delta_\Sigma t} \hat{\vartheta}_0) \rrbracket, \quad \hat{\gamma}_1(t, \cdot) = \gamma(e^{\Delta_\Sigma t} \hat{\vartheta}_0). \end{aligned}$$

For $z = (\vartheta, h) \in \mathbb{E}(J)$, we define

$$\begin{aligned} F(z) &= (\kappa_A - \kappa(\vartheta))\partial_t \vartheta + (d(\vartheta) - d_A)\Delta \vartheta - d(\vartheta)M_2(h) : \nabla^2 \vartheta \\ &\quad + d'(\vartheta)(|I - M_1(h)|\nabla \vartheta|^2 - d(\vartheta)(M_3(h)|\nabla \vartheta|) + \kappa(\vartheta)\mathcal{R}(h)\vartheta, \\ G(z) &= -(\llbracket \psi(\vartheta) \rrbracket + \sigma \mathcal{H}(h)) + l_1 \vartheta + \sigma_0 \Delta_\Sigma h + (\gamma(\vartheta)\beta(h) - \gamma_1)\partial_t h, \\ Q(z) &= \llbracket (d(\vartheta) - d_A)\partial_\nu \vartheta \rrbracket + (l_A - l(\vartheta))\partial_t h - (\llbracket d(\vartheta)\nabla \vartheta \rrbracket M_4(h)\nabla_\Sigma h) \\ &\quad + \gamma(\vartheta)\beta(h)(\partial_t h)^2. \end{aligned}$$

Here we have

$$\begin{aligned} M_2(h) &= M_1(h) + M_1^\top(h) - M_1(h)M_1^\top(h), \\ M_3(h) &= (I - M_1(h)) : \nabla M_1(h), \\ M_4(h) &= (I - M_1(h))^\top M_0(h). \end{aligned}$$

Employing the above notations and splitting into the principal linear part and a nonlinear part, we arrive at the following formulation of problem (2.1)

$$\left\{ \begin{array}{ll} \kappa_A \partial_t \vartheta - d_A \Delta \vartheta = F(\vartheta, h) & \text{in } \Omega \setminus \Sigma, \\ \partial_{\nu_\Omega} \vartheta = 0 & \text{on } \partial\Omega, \\ \llbracket \vartheta \rrbracket = 0 & \text{on } \Sigma, \\ l_1(t)\vartheta + \sigma_0 \Delta_\Sigma h - \gamma_1(t)\partial_t h = G(\vartheta, h) & \text{on } \Sigma, \\ l_A \partial_t h - \llbracket d_A(x)\partial_\nu \vartheta \rrbracket = Q(\vartheta, h) & \text{on } \Sigma, \\ \vartheta(0) = \hat{\vartheta}_0, \quad h(0) = \hat{h}_0. & \end{array} \right. \quad (4.2)$$

We assume that \hat{z}_0 further satisfies the compatibility conditions (CC), that is, when $\gamma \equiv 0$

$$\partial_{\nu_\Omega} \hat{\vartheta}_0 = 0, \quad l_1(0)\hat{\vartheta}_0 + \sigma_0 \Delta_\Sigma \hat{h}_0 = G(\hat{z}_0), \quad Q(\hat{z}_0) + \llbracket d_A \partial_\nu \hat{\vartheta}_0 \rrbracket \in W_p^{2-6/p}(\Sigma),$$

and when $\gamma > 0$

$$\partial_{v_\Omega} \hat{\vartheta}_0 = 0, \quad l_A l_1(0) \hat{\vartheta}_0 + l_A \sigma_0 \Delta_\Sigma \hat{h}_0 - \gamma_1(0) \llbracket d_A \partial_v \hat{\vartheta}_0 \rrbracket = \gamma_1(0) Q(\hat{z}_0) + l_A G(\hat{z}_0).$$

In the definition of $(G(\hat{z}_0), Q(\hat{z}_0))$, it is understood that $\partial_t h(0)$ is replaced by

$$\partial_t h(0) = \begin{cases} \frac{1}{\hat{l}_0} (\llbracket \hat{d}_0 \partial_v \hat{\vartheta}_0 \rrbracket - (\llbracket \hat{d}_0 \nabla \hat{\vartheta}_0 \rrbracket | M_4(\hat{h}_0) \nabla_\Sigma \hat{h}_0)), & \gamma \equiv 0, \\ \frac{1}{\beta(\hat{h}_0) \hat{\gamma}_0} (\llbracket \psi(\hat{\vartheta}_0) \rrbracket + \sigma \mathcal{H}(\hat{h}_0)), & \gamma > 0. \end{cases} \quad (4.3)$$

When $\gamma \equiv 0$, we also need to impose the *well-posedness condition* $l(\hat{\vartheta}_0) \neq 0$ on Γ_0 .

4.2 Regularity of a special solution

In this section, we prove the analyticity of the solution to a linearized Stefan problem with initial data \hat{z}_0 , that is, we consider regularity of solutions to the following equation.

$$\left\{ \begin{array}{ll} \kappa_A \partial_t \vartheta - d_A \Delta \vartheta = 0 & \text{in } \Omega \setminus \Sigma, \\ \partial_{v_\Omega} \vartheta = 0 & \text{on } \partial\Omega, \\ \llbracket \vartheta \rrbracket = 0 & \text{on } \Sigma, \\ l_1 \vartheta + \sigma_0 \Delta_\Sigma h - \gamma_1 \partial_t h = g_1 & \text{on } \Sigma, \\ l_A \partial_t h - \llbracket d_A \partial_v \vartheta \rrbracket = q_1 & \text{on } \Sigma, \\ \vartheta(0) = \hat{\vartheta}_0, \quad h(0) = \hat{h}_0. \end{array} \right. \quad (4.4)$$

Here $g_1 := e^{\Delta_\Sigma t} G(\hat{z}_0)$ and $q_1 := e^{\Delta_\Sigma t} Q(\hat{z}_0)$. Recall definition (2.2). We set

$$\mathbb{F}(J) = \{(f, g, q, (\vartheta_0, h_0)) \in \mathbb{F}_1(J) \times \mathbb{F}_2(J) \times \mathbb{F}_3(J) \times \mathbb{F}_4 : \\ (f, g, q, (\vartheta_0, h_0)) \text{ satisfies the linear compatibility conditions } (\mathcal{LCC})\},$$

where $(f, g, q, (\vartheta_0, h_0))$ is said to satisfy the linear compatibility conditions (\mathcal{LCC}) if

$$\begin{aligned} \partial_{v_\Omega} \vartheta_0 = 0, \quad l_1(0) \vartheta_0 + \sigma_0 \Delta_\Sigma h_0 = g(0), \quad q(0) + \llbracket d_A \partial_v \vartheta_0 \rrbracket \in W_p^{2-6/p}(\Sigma), \quad \gamma \equiv 0, \\ \partial_{v_\Omega} \vartheta_0 = 0, \quad l_A l_1(0) \vartheta_0 + l_A \sigma_0 \Delta_\Sigma h_0 - \gamma_1(0) \llbracket d_A \partial_v \vartheta_0 \rrbracket = \gamma_1(0) q(0) + l_A g(0), \quad \gamma > 0. \end{aligned}$$

We equipped $\mathbb{F}(J)$ with the following norm

$$\begin{aligned} & \| (f, g, q, (\vartheta_0, h_0)) \|_{\mathbb{F}(J)} \\ & := \|f\|_{\mathbb{F}_1(J)} + \|g\|_{\mathbb{F}_2(J)} + \|q\|_{\mathbb{F}_3(J)} + \|(\vartheta_0, h_0)\|_{\mathbb{F}_4} + (1 - \text{sgn}(\gamma)) \|q(0) + \llbracket d_A \partial_v \vartheta_0 \rrbracket\|_{W_p^{2-6/p}(\Sigma)}. \end{aligned}$$

(i) Regularity of g_1 and q_1 .

Case 1: $\gamma \equiv 0$. It is not hard to check that

$$l_1, \gamma_1 \in \mathbb{F}_2(J), \quad \kappa_A \in W_p^1(\Omega \setminus \Sigma), \quad d_A \in W_p^{2-2/p}(\Omega \setminus \Sigma), \quad l_A \in W_p^{2-3/p}(\Sigma). \quad (4.5)$$

Indeed, the first term can be obtained as follows. $\vartheta_A|_\Sigma \in W_p^{2-3/p}(\Sigma)$ and the L_p -maximal regularity of Δ_Σ imply that

$$e^{\Delta_\Sigma t} \vartheta_A \in H_p^1(J; W_p^{-1/p}(\Sigma)) \cap L_p(J; W_p^{2-1/p}(\Sigma)).$$

By [55, Proposition 3.2], we infer that $e^{\Delta_\Sigma t} \vartheta_A \in \mathbb{F}_2(J)$. Now the first term in (4.5) follows from the regularity of l, γ . Similar to (4.5), one checks that

$$\hat{l}_1, \hat{\gamma}_1 \in \mathbb{F}_2(J), \quad \hat{\kappa}_0 \in W_p^1(\Omega \setminus \Sigma), \quad \hat{d}_0 \in W_p^{2-2/p}(\Omega \setminus \Sigma), \quad \hat{l}_0 \in W_p^{2-3/p}(\Sigma). \quad (4.6)$$

In [72], an analysis of the structure of the mean curvature operator $\mathcal{H}(h)$ and $\beta(h)$ is obtained, which shows that $\mathcal{H}(h)$ is a rational function in the height function h and its spatial derivatives up to second order, while $\beta(h)$ is a rational function in h and its first order spatial derivatives. On account of the embeddings $W_p^{3-3/p}(\Sigma) \hookrightarrow C(\Sigma)$ and $W_p^{2-3/p}(\Sigma) \hookrightarrow C(\Sigma)$, we conclude from [75, Theorem 2.8.3] and a similar argument to [74, Proposition 2.7] that $W_p^{3-3/p}(\Sigma)$ and $W_p^{2-3/p}(\Sigma)$ are multiplication algebras under point-wise multiplication. Since

$$[x \mapsto x^a] \in C^\omega((0, \infty)), \quad a \in \mathbb{R},$$

well-known results for substitution operators for Sobolev-Slobodeckii spaces imply that

$$\beta(\hat{h}_0) \in W_p^{3-3/p}(\Sigma), \quad \mathcal{H}(\hat{h}_0) \in W_p^{2-3/p}(\Sigma). \quad (4.7)$$

(4.5)–(4.7) then imply

$$G(\hat{z}_0) \in W_p^{2-3/p}(\Sigma), \quad (4.8)$$

and a similar argument yields

$$Q(\hat{z}_0) \in W_p^{1-3/p}(\Sigma). \quad (4.9)$$

Together with the L_p -maximal regularity of Δ_Σ and [55, Proposition 3.2], we conclude from (4.8) and (4.9) that

$$g_1 \in \mathbb{G}_2(J) \hookrightarrow \mathbb{F}_2(J), \quad q_1 \in \mathbb{G}_3(J) \hookrightarrow \mathbb{F}_3(J). \quad (4.10)$$

Case 2: $\gamma > 0$. Based on the discussion in (i), it suffices to show the regularity of $\partial_t h(0)$, which is defined in (4.3). As illustrated in (i), we have

$$\beta(\hat{h}_0) \hat{\gamma}_0, ([\psi(\hat{\vartheta}_0)]) + \sigma \mathcal{H}(\hat{h}_0) \in W_p^{2-3/p}(\Sigma).$$

It implies that $\partial_t h(0) \in W_p^{2-3/p}(\Sigma)$. It yields the desired results, i.e.,

$$g_1 \in \mathbb{F}_2(J), \quad q_1 \in \mathbb{F}_3(J).$$

In virtue of condition (CC) and the definitions of g_1 and q_1 , one checks that condition (LCC) is at our disposal. Therefore, all the compatibility conditions in [65, Theorems 3.3, 3.5] are satisfied, and then there exists a unique solution $z^* = (\vartheta^*, h^*) \in \mathbb{E}(J)$ to the linear system (4.4).

Remark 4.1. The compatibility condition $[[d(\theta_0) \partial_{\nu_{\Gamma_0}} \theta_0]] \in W_p^{2-6/p}(\Gamma_0)$ in Theorem 1.1 implies that

$$[[\hat{d}_0 \partial_\nu \hat{\vartheta}_0]] - ([[\hat{d}_0 \nabla \hat{\vartheta}_0]]) M_4(\hat{h}_0) \nabla_\Sigma \hat{h}_0 \in W_p^{2-6/p}(\Sigma),$$

which is equivalent to

$$([\hat{d}_0 \nabla \hat{\vartheta}_0]|_{\nu_\Sigma} - M_4(\hat{h}_0) \nabla_\Sigma \hat{h}_0) \in W_p^{2-6/p}(\Sigma).$$

From the above discussion, we infer that

$$[\nabla \hat{\vartheta}_0] \in W_p^{2-6/p}(\Sigma, \mathbb{R}^{m+1}).$$

Now based on this observation and (4.6), we conclude that

$$Q(\hat{z}_0) + [d_A \partial_\nu \hat{\vartheta}_0] \in W_p^{2-6/p}(\Sigma).$$

The other two conditions in conditions (CC), i.e., when $\gamma \equiv 0$

$$\partial_{\nu_\Omega} \hat{\vartheta}_0 = 0, \quad l_1(0) \hat{\vartheta}_0 + \sigma_0 \Delta_\Sigma \hat{h}_0 = G(\hat{z}_0),$$

and when $\gamma > 0$

$$\partial_{\nu_\Omega} \hat{\vartheta}_0 = 0, \quad l_A l_1(0) \hat{\vartheta}_0 + l_A \sigma_0 \Delta_\Sigma \hat{h}_0 - \gamma_1(0) [d_A \partial_\nu \hat{\vartheta}_0] = \gamma_1(0) Q(\hat{z}_0) + l_A G(\hat{z}_0).$$

can be easily obtained from the remaining compatibility conditions in Theorems 1.1 and 1.2. Similarly, the compatibility conditions in Theorems 1.1 and 1.2 can also be concluded from conditions (CC). By [65, Theorems 3.1, 3.2], problem (4.2) with initial data \hat{z}_0 has a unique L_p -solution on some possibly small but nontrivial interval $J := [0, T]$, which is denoted by $\hat{z} = (\hat{\vartheta}, \hat{h})$. \square

Next, we apply the parameter-dependent diffeomorphisms $\Theta_{\lambda, \mu, \eta}^*$ and $\Theta_{\lambda, \mu}^*$ to show the analyticity of the solution z^* . We will use the following useful fact that for any time-independent map \mathfrak{N} acting on $\mathfrak{F}^s(\Omega_i, E)$

$$\Theta_{\lambda, \mu, \eta}^* \mathfrak{N} = T_{\mu, \eta} \mathfrak{N} T_{\mu, \eta}^{-1} \Theta_{\lambda, \mu, \eta}^*,$$

and a similar result also holds for $\Theta_{\lambda, \mu}^*$. By Proposition 3.10, we have

$$\begin{aligned} \partial_t [\vartheta_{\lambda, \mu, \eta}^*] &= (1 + \xi' \lambda) \Theta_{\lambda, \mu, \eta}^* \partial_t \vartheta^* + B_{\lambda, \mu, \eta}(\vartheta_{\lambda, \mu, \eta}^*) \\ &= (1 + \xi' \lambda) \Theta_{\lambda, \mu, \eta}^* (d_A / \kappa_A) \Delta \vartheta^* + B_{\lambda, \mu, \eta}(\vartheta_{\lambda, \mu, \eta}^*) \\ &= (1 + \xi' \lambda) (d_{A, \lambda, \mu, \eta} / \kappa_{A, \lambda, \mu, \eta}) T_{\mu, \eta} \Delta T_{\mu, \eta}^{-1} \vartheta_{\lambda, \mu, \eta}^* + B_{\lambda, \mu, \eta}(\vartheta_{\lambda, \mu, \eta}^*), \end{aligned}$$

and either by Proposition 3.9

$$\begin{aligned} \partial_t [h_{\lambda, \mu}^*] &= (1 + \xi' \lambda) \Theta_{\lambda, \mu}^* \partial_t h^* + B_{\lambda, \mu}(h_{\lambda, \mu}^*) \\ &= (1 + \xi' \lambda) \Theta_{\lambda, \mu}^* [(q_1 + [d_A \partial_\nu \vartheta^*]) / l_A] + B_{\lambda, \mu}(h_{\lambda, \mu}^*) \\ &= (1 + \xi' \lambda) \{ (q_{1, \lambda, \mu} / l_{A, \lambda, \mu}) + [(d_{A, \lambda, \mu} / l_{A, \lambda, \mu}) \partial_\nu \vartheta_{\lambda, \mu}^*] \} + B_{\lambda, \mu}(h_{\lambda, \mu}^*), \end{aligned}$$

when $\gamma \equiv 0$, or when $\gamma > 0$, we have

$$\begin{aligned} \partial_t [h_{\lambda, \mu}^*] &= (1 + \xi' \lambda) \Theta_{\lambda, \mu}^* [(l_1 \vartheta^* + \sigma_0 \Delta_\Sigma h^* - g_1) / \gamma_1] + B_{\lambda, \mu}(h_{\lambda, \mu}^*) \\ &= (1 + \xi' \lambda) (l_{1, \lambda, \mu} \vartheta_{\lambda, \mu}^* + \sigma_0 T_\mu \Delta_\Sigma T_\mu^{-1} h_{\lambda, \mu}^* - g_{1, \lambda, \mu}) / (\gamma_{1, \lambda, \mu}) + B_{\lambda, \mu}(h_{\lambda, \mu}^*). \end{aligned}$$

We define a map $\Phi : \mathbb{E}(J) \times \mathbb{B}(0, r_0) \rightarrow \mathbb{F}(J)$: $((\vartheta, h), (\lambda, \mu, \eta)) \mapsto$

$$\begin{cases} \kappa_{A,\lambda,\mu,\eta} \partial_t \vartheta - (1 + \xi' \lambda) d_{A,\lambda,\mu,\eta} T_{\mu,\eta} \Delta T_{\mu,\eta}^{-1} \vartheta - \kappa_{A,\lambda,\mu,\eta} B_{\lambda,\mu,\eta}(\vartheta) & \text{in } \Omega \setminus \Sigma, \\ (1 + \operatorname{sgn}(\gamma) \xi' \lambda) (l_{1,\lambda,\mu} \vartheta + \sigma_0 T_{\mu} \Delta_{\Sigma} T_{\mu}^{-1} h - g_{1,\lambda,\mu}) - \gamma_{1,\lambda,\mu} \partial_t h \\ \quad + \gamma_{1,\lambda,\mu} B_{\lambda,\mu}(h) & \text{on } \Sigma, \\ l_{A,\lambda,\mu} \partial_t h - (1 + \xi' \lambda) (q_{1,\lambda,\mu} + \llbracket d_{A,\lambda,\mu} \partial_v \vartheta \rrbracket) - l_{A,\lambda,\mu} B_{\lambda,\mu}(h) & \text{on } \Sigma, \\ \vartheta(0) - \hat{\vartheta}_0 & \text{in } \Omega \setminus \Sigma, \\ h(0) - \hat{h}_0 & \text{on } \Sigma. \end{cases}$$

Setting $z_{\lambda,\mu,\eta}^* = (\vartheta_{\lambda,\mu,\eta}^*, h_{\lambda,\mu}^*)$, note that $\Phi(z_{\lambda,\mu,\eta}^*, (\lambda, \mu, \eta)) = 0$ for all $(\lambda, \mu, \eta) \in \mathbb{B}(0, r_0)$.

(ii) We need to show that Φ actually maps into $\mathbb{F}(J)$. For simplification, we set $\Phi(z, (\lambda, \mu, \eta)) = (f_2, g_2, q_2, (\vartheta_0, h_0))^T$. It is obvious that based on trace theory of anisotropic Sobolev-Slobodeckii spaces, see [22, Section 2],

$$(\vartheta_0, h_0) \in \mathbb{F}_4.$$

By means of Propositions 3.6 and 3.10, we infer that $f_2 \in \mathbb{F}_1(J)$. For the regularity of the next two terms, we split into two cases as before. When $\gamma \equiv 0$, observe that $\Delta_{\Sigma} \in \mathcal{L}(\mathbb{E}_2(J), \mathbb{F}_2(J))$. Proposition 3.11, (4.5) and (4.10) yield

$$\gamma_{1,\lambda,\mu}, l_{1,\lambda,\mu}, g_{1,\lambda,\mu} \in \mathbb{F}_2(J), \quad q_{1,\lambda,\mu} \in \mathbb{F}_3(J). \quad (4.11)$$

Taking into consideration that $\partial_t h \in \mathbb{F}_3(J)$ and the fact that $\mathbb{F}_2(J), \mathbb{F}_3(J)$ are multiplication algebras, we conclude from Propositions 3.8, 3.10 and (4.11) that

$$g_2 \in \mathbb{F}_2(J), \quad q_2 \in \mathbb{F}_3(J).$$

When $\gamma > 0$, since $\partial_t h \in \mathbb{F}_2(J)$, the desired regularity results clearly hold true. It remains to show condition (\mathcal{LCC}) .

Case 1: $\gamma \equiv 0$. It is immediate that $\partial_{v,\Omega} \vartheta_0 = \partial_{v,\Omega} (\vartheta(0) - \hat{\vartheta}_0) = 0$. One checks that

$$\begin{aligned} l_1(0) \vartheta_0 + \sigma_0 \Delta_{\Sigma} h_0 &= l_1(0) (\vartheta(0) - \hat{\vartheta}_0) + \sigma_0 \Delta_{\Sigma} (h(0) - \hat{h}_0) \\ &= l_{1,\lambda,\mu}(0) \vartheta(0) + \sigma T_{\mu}(0) \Delta_{\Sigma} T_{\mu}^{-1}(0) h(0) - g_{1,\lambda,\mu}(0) = g_2(0), \end{aligned}$$

by recalling $u_{\lambda,\mu}(0, \cdot) = u(0, \cdot)$ for any $u \in E^{\Sigma}$, and

$$\begin{aligned} q_2(0) + \llbracket d_{A,\lambda,\mu}(0) \partial_v \vartheta_0 \rrbracket &= l_{A,\lambda,\mu}(0) \partial_t h(0) - q_{1,\lambda,\mu}(0) - \llbracket d_A \partial_v \vartheta(0) \rrbracket \\ &\quad - l_{A,\lambda,\mu}(0) B_{\lambda,\mu}(h(0)) + \llbracket d_A \partial_v (\vartheta(0) - \hat{\vartheta}_0) \rrbracket \\ &= l_A \partial_t h(0) - q_1(0) - l_A B_{\lambda,\mu}(h(0)) - \llbracket d_A \partial_v \hat{\vartheta}_0 \rrbracket. \end{aligned}$$

By the discussion in (ii), $q_1(0) + \llbracket d_A \partial_v \hat{\vartheta}_0 \rrbracket \in W_p^{2-6/p}(\Sigma)$. Since $W_p^{2-6/p}(\Sigma)$ is a multiplication algebra, (3.12), (4.5) and [65, formula (3.1)] imply that

$$q_2(0) + \llbracket d_{A,\lambda,\mu}(0) \partial_v \vartheta_0 \rrbracket \in W_p^{2-6/p}(\Sigma).$$

Case 2: $\gamma > 0$.

$$\begin{aligned}
 & l_A l_1(0) \vartheta_0 + l_A \sigma_0 \Delta_\Sigma h_0 - \gamma_1(0) \llbracket d_A \partial_v \vartheta_0 \rrbracket \\
 &= l_{A,\lambda,\mu}(0) l_{1,\lambda,\mu}(0) (\vartheta(0) - \hat{\vartheta}_0) + l_{A,\lambda,\mu}(0) \sigma_0 T_\mu(0) \Delta_\Sigma T_\mu^{-1}(0) (h(0) - \hat{h}_0) \\
 &\quad - \gamma_{1,\lambda,\mu}(0) \llbracket d_{A,\lambda,\mu} \partial_v (\vartheta(0) - \hat{\vartheta}_0) \rrbracket \\
 &= l_A l_1(0) \vartheta(0) + l_A \sigma_0 \Delta_\Sigma h(0) - \gamma_1(0) \llbracket d_A \partial_v \vartheta(0) \rrbracket - l_A g_1(0) - \gamma_1(0) q_1(0) \\
 &= l_A g_2(0) + \gamma_1(0) q_2(0).
 \end{aligned}$$

Therefore $\Phi(z, (\lambda, \mu, \eta)) \in \mathbb{F}(J)$.

(iii) Let $w = (u, \rho) \in \mathbb{E}(J)$. The Fréchet derivative of Φ with respect to (ϑ, h) at $(z^*, 0)$ is clearly given by

$$D_1 \Phi(z^*, 0) w = \begin{cases} \kappa_A \partial_t u - d_A \Delta u & \text{in } \Omega \setminus \Sigma, \\ l_1 u + \sigma_0 \Delta_\Sigma \rho - \gamma_1 \partial_t \rho & \text{on } \Sigma, \\ l_A \partial_t \rho - \llbracket d_A \partial_v u \rrbracket & \text{on } \Sigma, \\ u(0) & \text{in } \Omega \setminus \Sigma, \\ \rho(0) & \text{on } \Sigma. \end{cases}$$

It is obvious that

$$D_1 \Phi(z^*, 0) w \in \mathbb{F}_1(J) \times \mathbb{F}_2(J) \times \mathbb{F}_3(J) \times \mathbb{F}_4.$$

Condition (\mathcal{LCC}) can be verified as in (ii) by using [65, formula (3.1)] in the case $\gamma \equiv 0$. We can deduce from [65, Theorems 3.3, 3.5] that

$$D_1 \Phi(z^*, 0) \in \mathcal{L}is(\mathbb{E}(J), \mathbb{F}(J)).$$

(iv) Regularity of $[(\lambda, \mu) \mapsto (g_{1,\lambda,\mu}, q_{1,\lambda,\mu})]$. We express $g_{1,\lambda,\mu}$ as

$$\begin{aligned}
 g_{1,\lambda,\mu} &= \Theta_{\lambda,\mu}^* e^{\Delta_\Sigma t} G(\hat{z}_0) = \Theta_{\lambda,\mu}^* (c + \Delta_\Sigma) e^{\Delta_\Sigma t} (c + \Delta_\Sigma)^{-1} G(\hat{z}_0) \\
 &= T_\mu (c + \Delta_\Sigma) T_\mu^{-1} \Theta_{\lambda,\mu}^* e^{\Delta_\Sigma t} (c + \Delta_\Sigma)^{-1} G(\hat{z}_0).
 \end{aligned}$$

For sufficiently large c , it is well-known that $(c + \Delta_\Sigma)$ is an isomorphism from $W_p^{s+2}(\Sigma)$ to $W_p^s(\Sigma)$ for any $s \in \mathbb{R}$. We consider the solution to

$$\partial_t \rho + \Delta_\Sigma \rho = 0, \quad \rho(0) = (c + \Delta_\Sigma)^{-1} G(\hat{z}_0). \quad (4.12)$$

$\hat{\rho} := e^{\Delta_\Sigma t} (c + \Delta_\Sigma)^{-1} G(\hat{z}_0)$ is the unique solution to (4.12). Furthermore, Proposition 3.10 shows that $\hat{\rho}_{\lambda,\mu}$ satisfies

$$\begin{aligned}
 \partial_t [\hat{\rho}_{\lambda,\mu}] &= (1 + \xi' \lambda) \Theta_{\lambda,\mu}^* \partial_t \hat{\rho} + B_{\lambda,\mu}(\hat{\rho}_{\lambda,\mu}) \\
 &= -(1 + \xi' \lambda) T_\mu \Delta_\Sigma T_\mu^{-1} \hat{\rho}_{\lambda,\mu} + B_{\lambda,\mu}(\hat{\rho}_{\lambda,\mu}).
 \end{aligned}$$

We define the map $\Phi_g : \mathbb{X}_2(J) \times \mathbb{B}(0, r_0) \rightarrow L_p(J; W_p^{2-1/p}(\Sigma)) \times W_p^{4-3/p}(\Sigma)$ by

$$\Phi_g(\rho, (\lambda, \mu)) = \begin{pmatrix} \partial_t \rho + (1 + \xi' \lambda) T_\mu \Delta_\Sigma T_\mu^{-1} \rho - B_{\lambda,\mu}(\rho) \\ \rho(0) - (c + \Delta_\Sigma)^{-1} G(\hat{z}_0) \end{pmatrix}.$$

Here $\mathbb{X}_2(J) := H_p^1(J; W^{2-1/p}(\Sigma)) \cap L_p(J; W^{4-1/p}(\Sigma))$. Note that $\Phi_g(\hat{\rho}_{\lambda,\mu}, (\lambda, \mu)) = (0, 0)^\top$. By the L_p -maximal regularity of Δ_Σ , we immediately have

$$D_1\Phi_g(\hat{\rho}, 0) \in \mathcal{L}\text{is}\left(\mathbb{X}_2(J), L_p(J; W_p^{2-1/p}(\Sigma)) \times W_p^{4-3/p}(\Sigma)\right).$$

We define a bilinear and continuous map

$$T : C(J; \mathcal{L}(W_p^{4-1/p}(\Sigma), W_p^{3-1/p}(\Sigma))) \times \mathbb{X}_2(J) \rightarrow L_p(J; W_p^{2-1/p}(\Sigma))$$

by $(B, u) \mapsto [t \mapsto B(t)u(t)]$. Since T is real analytic, by (3.12), we get

$$[(u, (\lambda, \mu)) \mapsto B_{\lambda,\mu}(u)] \in C^\omega\left(\mathbb{X}_2 \times \mathbb{B}(0, r_0), L_p(J; W_p^{2-1/p}(\Sigma))\right).$$

In virtue of Proposition 3.8(b) and the above bilinear map argument, one gets

$$\Phi_g \in C^\omega\left(\mathbb{X}_2 \times \mathbb{B}(0, r_0), L_p(J; W_p^{2-1/p}(\Sigma)) \times W_p^{4-3/p}(\Sigma)\right).$$

By the implicit function theorem, there exists some $\mathbb{B}(0, r_1) \subset \mathbb{B}(0, r_0)$ such that

$$[(\lambda, \mu) \mapsto \Theta_{\lambda,\mu}^* e^{\Delta_\Sigma t} (c + \Delta_\Sigma)^{-1} G(\hat{z}_0)] \in C^\omega(\mathbb{B}(0, r_1), \mathbb{X}_2).$$

Without loss of generality, we may assume that $r_1 = r_0$. In view of Proposition 3.8(b), we have that for all $s \in \mathbb{R}$

$$[\mu \mapsto T_\mu(c + \Delta_\Sigma)T_\mu^{-1}] \in C^\omega\left(\mathbb{B}(0, r_0), C^1\left(J; \mathcal{L}(W_p^{s+2}(\Sigma), W_p^s(\Sigma))\right)\right).$$

The above bilinear map argument and the embedding $\mathbb{G}_2(J) \hookrightarrow \mathbb{F}_2(J)$ yield

$$[(\lambda, \mu) \mapsto g_{1,\lambda,\mu}] \in C^\omega(\mathbb{B}(0, r_0), \mathbb{F}_2(J)). \tag{4.13}$$

Following a similar discussion, one obtains

$$[(\lambda, \mu) \mapsto q_{1,\lambda,\mu}] \in C^\omega(\mathbb{B}(0, r_0), \mathbb{F}_3(J)). \tag{4.14}$$

(v) Regularity of the map Φ . It follows from Proposition 3.6 that

$$[(\lambda, \mu, \eta) \mapsto (\kappa_{A,\lambda,\mu,\eta}, d_{A,\lambda,\mu,\eta})] \in C^k\left(\mathbb{B}(0, r_0), C^1(J; BC(\Omega))^2\right). \tag{4.15}$$

Proposition 3.8 implies that

$$[(\lambda, \mu) \mapsto (l_{A,\lambda,\mu}, d_{A,\lambda,\mu})] \in C^k\left(\mathbb{B}(0, r_0), C^1(J; BC^1(\Sigma))^2\right). \tag{4.16}$$

Similarly, we have

$$[(\lambda, \mu) \mapsto (l_{1,\lambda,\mu}, \gamma_{1,\lambda,\mu})] \in C^k\left(\mathbb{B}(0, r_0), \left(C^1(J; BC(\Sigma)) \cap C(J; BC^2(\Sigma))\right)^2\right). \tag{4.17}$$

By Proposition 3.6, one concludes that

$$[(\mu, \eta) \mapsto T_{\mu, \eta} \Delta T_{\mu, \eta}^{-1}] \in C^\omega \left(\mathbb{B}(0, r_0), C^1 \left(J; \mathfrak{L}(H_p^2(\Omega \setminus \Sigma), L_p(\Omega)) \right) \right).$$

Combined with (4.15), this yields

$$[(\lambda, \mu, \eta) \mapsto d_{A, \lambda, \mu, \eta} T_{\mu, \eta} \Delta T_{\mu, \eta}^{-1}] \in C^k \left(\mathbb{B}(0, r_0), \mathfrak{L}(\mathbb{E}_1(J), \mathbb{F}_1(J)) \right).$$

It follows again from (4.15) that

$$[(\vartheta, (\lambda, \mu, \eta)) \mapsto \kappa_{A, \lambda, \mu, \eta} \partial_t \vartheta] \in C^k \left(\mathbb{E}_1(J) \times \mathbb{B}(0, r_0), \mathbb{F}_1(J) \right).$$

Together with Proposition 3.10, the above discussion shows that

$$\begin{aligned} [(\vartheta, (\lambda, \mu, \eta)) \mapsto \kappa_{A, \lambda, \mu, \eta} \partial_t \vartheta - (1 + \xi' \lambda) d_{A, \lambda, \mu, \eta} T_{\mu, \eta} \Delta T_{\mu, \eta}^{-1} \vartheta - B_{\lambda, \mu, \eta}(\vartheta)] \\ \in C^k \left(\mathbb{E}_1(J) \times \mathbb{B}(0, r_0), \mathbb{F}_1(J) \right). \end{aligned}$$

Applying (4.13)-(4.17), one can check the regularity of all the other entries of Φ , and thus

$$\Phi \in C^k \left(\mathbb{E}(J) \times \mathbb{B}(0, r_0), \mathbb{F}(J) \right).$$

Now the implicit function theorem yields

$$[(\lambda, \mu, \eta) \mapsto z_{\lambda, \mu, \eta}^*] \in C^k \left(\mathbb{B}(0, r_0), \mathbb{E}(J) \right).$$

As a conclusion of Theorem 3.12, we obtain

$$z^* \in C^k(\mathring{J} \times \mathbb{T}_{a/3} \setminus \Sigma) \times C^k(\mathring{J} \times \Sigma).$$

To attain the regularity of ϑ^* in $\Omega \setminus \bar{\mathbb{T}}_{a/6}$, we study the linear equations

$$\begin{cases} \kappa_1(\vartheta_A) \partial_t \vartheta - d_1(\vartheta_A) \Delta \vartheta = 0 & \text{in } \Omega_1 \setminus \bar{\mathbb{T}}_{a/6} \\ \vartheta = \vartheta^* & \text{on } \partial_1 \mathbb{T}_{a/6} \\ \vartheta(0) = \hat{\vartheta}_0 & \text{in } \Omega_1 \setminus \bar{\mathbb{T}}_{a/6}, \end{cases}$$

and

$$\begin{cases} \kappa_2(\vartheta_A) \partial_t \vartheta - d_2(\vartheta_A) \Delta \vartheta = 0 & \text{in } \Omega_2 \setminus \bar{\mathbb{T}}_{a/6} \\ \partial_{\nu_\Omega} \vartheta = 0 & \text{on } \partial \Omega \\ \vartheta = \vartheta^* & \text{on } \partial_2 \mathbb{T}_{a/6} \\ \vartheta(0) = \hat{\vartheta}_0 & \text{in } \Omega_2 \setminus \bar{\mathbb{T}}_{a/6}, \end{cases}$$

where $\partial_i \mathbb{T}_{a/6} := \Lambda(\Sigma, (-1)^i a/6)$. Since ϑ^* is analytic on $\partial_i \mathbb{T}_{a/6}$, we can obtain the regularity of ϑ^* in these two domains by means of the parameter-dependent diffeomorphism defined in [26], the results in [21, Section 8], and the implicit function theorem as above. To sum up, we have established the analyticity of the solution z^* , i.e.,

$$z^* \in C^k(\mathring{J} \times \Omega \setminus \Sigma) \times C^k(\mathring{J} \times \Sigma). \tag{4.18}$$

4.3 Reduction to zero initial data

Recall that $\hat{z} = (\hat{\vartheta}, \hat{h})$ is the unique L_p solution to problem (4.2) with initial data \hat{z}_0 . We set $\bar{\vartheta} = \hat{\vartheta} - \vartheta^*$, $\bar{h} = \hat{h} - h^*$. Then $\bar{z} = (\bar{\vartheta}, \bar{h})$ is the unique solution to the following equation.

$$\left\{ \begin{array}{ll} \kappa_A \partial_t \vartheta - d_A \Delta \vartheta = F(\vartheta + \vartheta^*, h + h^*) & \text{in } \Omega \setminus \Sigma, \\ \partial_{\nu_\Omega} \vartheta = 0 & \text{on } \partial\Omega, \\ \llbracket \vartheta \rrbracket = 0 & \text{on } \Sigma, \\ l_1 \vartheta + \sigma_0 \Delta_\Sigma h - \gamma_1 \partial_t h = \bar{G}(\vartheta, h; \vartheta^*, h^*) & \text{on } \Sigma, \\ l_A \partial_t h - \llbracket d_A \partial_\nu \vartheta \rrbracket = \bar{Q}(\vartheta, h; \vartheta^*, h^*) & \text{on } \Sigma, \\ \vartheta(0) = 0 & \text{in } \Omega \setminus \Sigma, \\ h(0) = 0 & \text{on } \Sigma. \end{array} \right. \quad (4.19)$$

Here we have set

$$\begin{aligned} \bar{G}(\vartheta, h; \vartheta^*, h^*) &= G(\vartheta + \vartheta^*, h + h^*) - e^{\Delta_\Sigma t} G(\hat{z}_0), \\ \bar{Q}(\vartheta, h; \vartheta^*, h^*) &= Q(\vartheta + \vartheta^*, h + h^*) - e^{\Delta_\Sigma t} Q(\hat{z}_0). \end{aligned}$$

Note that $\bar{G}(0, 0; \hat{\vartheta}_0, \hat{h}_0) = \bar{Q}(0, 0; \hat{\vartheta}_0, \hat{h}_0) = 0$ by construction, which ensures time trace zero at $t = 0$. Let $z_{\lambda, \mu, \eta}^* := (\vartheta_{\lambda, \mu, \eta}^*, h_{\lambda, \mu}^*)$. As in Section 5.1, we compute the temporal derivative of $\bar{z}_{\lambda, \mu, \eta} := (\bar{\vartheta}_{\lambda, \mu, \eta}, \bar{h}_{\lambda, \mu})$ as follows.

$$\begin{aligned} \partial_t [\bar{\vartheta}_{\lambda, \mu, \eta}] &= (1 + \xi' \lambda) (d_{A, \lambda, \mu, \eta} / \kappa_{A, \lambda, \mu, \eta}) T_{\mu, \eta} \Delta T_{\mu, \eta}^{-1} \bar{\vartheta}_{\lambda, \mu, \eta} + B_{\lambda, \mu, \eta}(\bar{\vartheta}_{\lambda, \mu, \eta}) \\ &\quad + F_{\lambda, \mu, \eta}(\bar{z}_{\lambda, \mu, \eta}, z_{\lambda, \mu, \eta}^*) / \kappa_{A, \lambda, \mu, \eta}, \end{aligned}$$

and either

$$\partial_t [\bar{h}_{\lambda, \mu}] = (1 + \xi' \lambda) \llbracket (d_{A, \lambda, \mu} / l_{A, \lambda, \mu}) \partial_\nu \bar{\vartheta} \rrbracket + B_{\lambda, \mu}(\bar{h}_{\lambda, \mu}) + \bar{Q}_{\lambda, \mu}(\bar{z}_{\lambda, \mu, \eta}, z_{\lambda, \mu, \eta}^*) / l_{A, \lambda, \mu},$$

when $\gamma \equiv 0$, or when $\gamma > 0$ we have

$$\begin{aligned} \partial_t [\bar{h}_{\lambda, \mu}] &= (1 + \xi' \lambda) (l_{1, \lambda, \mu} \bar{\vartheta}_{\lambda, \mu} + \sigma_0 T_\mu \Delta_\Sigma T_\mu^{-1} \bar{h}_{\lambda, \mu}) / \gamma_{1, \lambda, \mu} + B_{\lambda, \mu}(\bar{h}_{\lambda, \mu}) \\ &\quad - \bar{G}_{\lambda, \mu}(\bar{z}_{\lambda, \mu, \eta}, z_{\lambda, \mu, \eta}^*) / \gamma_{1, \lambda, \mu}. \end{aligned}$$

For sufficiently small r_0 , $\Theta_{\mu, \eta}(z) = z$ for $(\mu, \eta) \in \mathbb{B}(0, r_0)$ and $z \in \Omega \setminus \mathbb{T}_{a/3}$. Hence, $(\zeta \circ d_\Sigma)_{\lambda, \mu, \eta} = \zeta \circ d_\Sigma$. In virtue of $(\zeta \circ d_\Sigma)_{\lambda, \mu, \eta}(x) = 0$ for $x \notin \Omega \setminus \mathbb{T}_{2a/3}$, and $(h \circ \Pi)_{\lambda, \mu, \eta}(x) = (h_{\lambda, \mu} \circ \Pi)(x)$ for $x \in \mathbb{T}_{2a/3}$, one readily verifies that

$$\begin{aligned} \Theta_{\lambda, \mu, \eta}^* \Upsilon(h) &= (\zeta \circ d_\Sigma)_{\lambda, \mu, \eta} (h_{\lambda, \mu} \circ \Pi)_{\lambda, \mu, \eta} (\nu_\Sigma \circ \Pi)_{\lambda, \mu, \eta} \\ &= (\zeta \circ d_\Sigma) (h_{\lambda, \mu} \circ \Pi) (\nu_\Sigma \circ \Pi)_{\lambda, \mu, \eta}. \end{aligned}$$

In the above expressions, for $z = (\vartheta, h)$,

$$\begin{aligned} F_{\lambda,\mu,\eta}(z, z_{\lambda,\mu,\eta}^*) &= (1 + \xi'\lambda)\Theta_{\lambda,\mu,\eta}^* F(\Theta_{\lambda,\mu,\eta}^{\lambda,\mu,\eta}(z + z_{\lambda,\mu,\eta}^*)) \\ &= (\kappa_{A,\lambda,\mu,\eta} - \kappa(\vartheta + \vartheta_{\lambda,\mu,\eta}^*))[\partial_t(\vartheta + \vartheta_{\lambda,\mu,\eta}^*) - B_{\lambda,\mu,\eta}(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)] \\ &\quad + (1 + \xi'\lambda)\left\{ (d(\vartheta + \vartheta_{\lambda,\mu,\eta}^*) - d_{A,\lambda,\mu,\eta})T_{\mu,\eta}\Delta T_{\mu,\eta}^{-1}(\vartheta + \vartheta_{\lambda,\mu,\eta}^*) \right. \\ &\quad - d(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)M_{2,\lambda,\mu,\eta}(h + h_{\lambda,\mu}^*) : T_{\mu,\eta}\nabla^2 T_{\mu,\eta}^{-1}(\vartheta + \vartheta_{\lambda,\mu,\eta}^*) \\ &\quad + d'(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)|(I - M_{1,\lambda,\mu,\eta}(h + h_{\lambda,\mu}^*))T_{\mu,\eta}\nabla T_{\mu,\eta}^{-1}(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)|^2 \\ &\quad - d(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)(M_{3,\lambda,\mu,\eta}(h + h_{\lambda,\mu}^*)|T_{\mu,\eta}\nabla T_{\mu,\eta}^{-1}(\vartheta + \vartheta_{\lambda,\mu,\eta}^*) \\ &\quad \left. + \kappa(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)\mathcal{R}_{\lambda,\mu,\eta}(h + h_{\lambda,\mu}^*)(\vartheta + \vartheta_{\lambda,\mu,\eta}^*) \right\}, \end{aligned}$$

with

$$\begin{aligned} \Upsilon_{\lambda,\mu,\eta}(h) &= \Theta_{\lambda,\mu,\eta}^* \Upsilon(\Theta_{\lambda,\mu,\eta}^{\lambda,\mu}h) = (\zeta \circ d_{\Sigma})(h \circ \Pi)(v_{\Sigma} \circ \Pi)_{\lambda,\mu,\eta} \\ M_{1,\lambda,\mu,\eta}(h) &= \Theta_{\lambda,\mu,\eta}^* M_1(\Theta_{\lambda,\mu,\eta}^{\lambda,\mu}h) \\ &= \left[(I + (T_{\mu,\eta}\nabla T_{\mu,\eta}^{-1}\Upsilon_{\lambda,\mu,\eta}(h))^{\top})^{-1} (T_{\mu,\eta}\nabla T_{\mu,\eta}^{-1}\Upsilon_{\lambda,\mu,\eta}(h))^{\top} \right]^{\top}, \\ M_{2,\lambda,\mu,\eta}(h) &= \Theta_{\lambda,\mu,\eta}^* M_2(\Theta_{\lambda,\mu,\eta}^{\lambda,\mu}h) \\ &= M_{1,\lambda,\mu,\eta}(h) + (M_{1,\lambda,\mu,\eta}(h))^{\top} - M_{1,\lambda,\mu,\eta}(h)(M_{1,\lambda,\mu,\eta}(h))^{\top}, \\ M_{3,\lambda,\mu,\eta}(h) &= \Theta_{\lambda,\mu,\eta}^* M_3(\Theta_{\lambda,\mu,\eta}^{\lambda,\mu}h) = (I - M_{1,\lambda,\mu,\eta}(h)) : T_{\mu,\eta}\nabla T_{\mu,\eta}^{-1}M_{1,\lambda,\mu,\eta}(h), \\ \mathcal{R}_{\lambda,\mu,\eta}(h)\vartheta &= \Theta_{\lambda,\mu,\eta}^* \mathcal{R}(\Theta_{\lambda,\mu,\eta}^{\lambda,\mu}h)\Theta_{\lambda,\mu,\eta}^{\lambda,\mu,\eta}\vartheta \\ &= \left(T_{\mu,\eta}\nabla T_{\mu,\eta}^{-1}\vartheta | (I + (T_{\mu,\eta}\nabla T_{\mu,\eta}^{-1}\Upsilon_{\lambda,\mu,\eta}(h))^{\top})^{-1} \right. \\ &\quad \left. \left[\partial_t \Upsilon_{\lambda,\mu,\eta}(h) - B_{\lambda,\mu,\eta}(\Upsilon_{\lambda,\mu,\eta}(h)) \right] \right) / (1 + \xi'\lambda), \end{aligned}$$

and

$$\begin{aligned} \bar{G}_{\lambda,\mu}(z, z_{\lambda,\mu,\eta}^*) &= (1 + \xi'\lambda)\Theta_{\lambda,\mu}^* \bar{G}(\Theta_{\lambda,\mu}^{\lambda,\mu}(z + z_{\lambda,\mu,\eta}^*)) \\ &= \{\gamma(\vartheta + \vartheta_{\lambda,\mu}^*)\beta_{\lambda,\mu}(h + h_{\lambda,\mu}^*) - \gamma_{1,\lambda,\mu}\}[\partial_t(h + h_{\lambda,\mu}^*) - B_{\lambda,\mu}(h + h_{\lambda,\mu}^*)] \\ &\quad + (1 + \xi'\lambda)\left\{ -([\psi(\vartheta + \vartheta_{\lambda,\mu}^*)] + \sigma\mathcal{H}_{\lambda,\mu}(h + h_{\lambda,\mu}^*)) + l_{1,\lambda,\mu}(\vartheta + \vartheta_{\lambda,\mu}^*) \right. \\ &\quad \left. + \sigma_0 T_{\mu}\Delta_{\Sigma}T_{\mu}^{-1}(h + h_{\lambda,\mu}^*) - g_{1,\lambda,\mu} \right\}, \end{aligned}$$

with $\mathcal{H}_{\lambda,\mu} = \Theta_{\lambda,\mu}^* \mathcal{H}\Theta_{\lambda,\mu}^{\lambda,\mu}$, and

$$\begin{aligned} M_{0,\lambda,\mu}(h) &= \Theta_{\lambda,\mu}^* M_0(\Theta_{\lambda,\mu}^{\lambda,\mu}h) = (I - hL_{\Sigma,\lambda,\mu})^{-1}, \\ \alpha_{\lambda,\mu}(h) &= \Theta_{\lambda,\mu}^* \alpha(\Theta_{\lambda,\mu}^{\lambda,\mu}h) = M_{0,\lambda,\mu}(h)T_{\mu}\nabla T_{\mu}^{-1}h, \\ \beta_{\lambda,\mu}(h) &= \Theta_{\lambda,\mu}^* \beta(\Theta_{\lambda,\mu}^{\lambda,\mu}h) = (1 + |\alpha_{\lambda,\mu}(h)|^2)^{-1/2}, \end{aligned}$$

and

$$\begin{aligned} \bar{Q}_{\lambda,\mu}(z, z_{\lambda,\mu,\eta}^*) &= (1 + \xi'\lambda)\Theta_{\lambda,\mu}^* \bar{Q}(\Theta_{\lambda,\mu}^{\lambda,\mu}(z + z_{\lambda,\mu,\eta}^*)) \\ &= \gamma(\vartheta + \vartheta_{\lambda,\mu}^*)\beta_{\lambda,\mu}(h + h_{\lambda,\mu}^*)[\partial_t(h + h_{\lambda,\mu}^*) - B_{\lambda,\mu}(h + h_{\lambda,\mu}^*)]^2 / (1 + \xi'\lambda) \\ &\quad + (l_{A,\lambda,\mu} - l(\vartheta + \vartheta_{\lambda,\mu}^*))[\partial_t(h + h_{\lambda,\mu}^*) - B_{\lambda,\mu}(h + h_{\lambda,\mu}^*)] \\ &\quad + (1 + \xi'\lambda)\{[(d(v + \bar{v}_{\lambda,\mu}) - d_{A,\lambda,\mu})\partial_v(\vartheta + \vartheta_{\lambda,\mu}^*)] - q_{1,\lambda,\mu} \\ &\quad - ([d(\vartheta + \vartheta_{\lambda,\mu}^*)T_\mu \nabla T_\mu^{-1}(\vartheta + \vartheta_{\lambda,\mu}^*)]M_{4,\lambda,\mu}(h + h_{\lambda,\mu}^*)T_\mu \nabla_\Sigma T_\mu^{-1}(h + h_{\lambda,\mu}^*))\} \end{aligned}$$

with $M_{4,\lambda,\mu}(h) = \Theta_{\lambda,\mu}^* M_4(\Theta_{\lambda,\mu}^{\lambda,\mu} h) = (I - M_{1,\lambda,\mu}(h))^\top M_{0,\lambda,\mu}(h)$.

Consider the map $\Phi : {}_0\mathbb{E}(J) \rightarrow {}_0\mathbb{F}(J) : ((\vartheta, h), (\lambda, \mu, \eta)) \mapsto$

$$\begin{cases} \kappa_{A,\lambda,\mu,\eta}\partial_t\vartheta - (1 + \xi'\lambda)d_{A,\lambda,\mu,\eta}T_{\mu,\eta}\Delta T_{\mu,\eta}^{-1}\vartheta - \kappa_{A,\lambda,\mu,\eta}B_{\lambda,\mu,\eta}(\vartheta) \\ \quad - F_{\lambda,\mu,\eta}(z, z_{\lambda,\mu,\eta}^*) & \text{in } \Omega \setminus \Sigma, \\ (1 + \operatorname{sgn}(\gamma)\xi'\lambda)(l_{1,\lambda,\mu}\vartheta + \sigma_0 T_\mu \Delta_\Sigma T_\mu^{-1}h) - \gamma_{1,\lambda,\mu}\partial_t h \\ \quad + \gamma_{1,\lambda,\mu}B_{\lambda,\mu}(h) - \bar{G}_{\lambda,\mu}(z, z_{\lambda,\mu,\eta}^*) & \text{on } \Sigma, \\ l_{A,\lambda,\mu}\partial_t h - (1 + \xi'\lambda)([d_{A,\lambda,\mu}\partial_v\vartheta] - l_{A,\lambda,\mu}B_{\lambda,\mu}(h) - \bar{Q}_{\lambda,\mu}(z, z_{\lambda,\mu,\eta}^*)) & \text{on } \Sigma, \end{cases}$$

where $z = (\vartheta, h)$. Note that $\Phi(\bar{z}_{\lambda,\mu}, (\lambda, \mu)) = 0$ for all $(\lambda, \mu) \in \mathbb{B}(0, r_0)$. It is understood that ${}_0\mathbb{F}(J) := \mathbb{F}_1(J) \times {}_0\mathbb{F}_2(J) \times {}_0\mathbb{F}_3(J)$.

(i) First, we shall check that Φ actually maps into ${}_0\mathbb{F}(J)$. Functions in ${}_0\mathbb{F}(J)$ automatically satisfy (\mathcal{LCC}) . One can check the temporal traces without difficulty. Indeed, recalling that $z_{\lambda,\mu,\eta}^*(0, \cdot) = z^*(0, \cdot)$, we have

$$\begin{aligned} \bar{G}_{\lambda,\mu}(z, z_{\lambda,\mu,\eta}^*)(0) &= \Theta_{\lambda,\mu}^* G(z^*)(0) - g_1(0) = 0, \\ \bar{Q}_{\lambda,\mu}(z, z_{\lambda,\mu,\eta}^*)(0) &= \Theta_{\lambda,\mu}^* Q(z^*)(0) - q_1(0) = 0. \end{aligned}$$

It suffices to check regularity of $F_{\lambda,\mu,\eta}(z, z_{\lambda,\mu,\eta}^*)$, $\bar{G}_{\lambda,\mu}(z, z_{\lambda,\mu,\eta}^*)$ and $\bar{Q}_{\lambda,\mu}(z, z_{\lambda,\mu,\eta}^*)$, which will become clear in our argument for the regularity of Φ .

It follows from well-known results for substitution operators for Sobolev spaces and (4.18) that

$$[(\vartheta, (\lambda, \mu, \eta)) \mapsto \kappa(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)] \in C^k\left({}_0\mathbb{E}_1(J) \times \mathbb{B}(0, r_0), C(J, BC(\Omega))\right).$$

Analogous statements hold for $d(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)$, $d'(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)$, $l(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)$, $\gamma(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)$ and $\psi(\vartheta + \vartheta_{\lambda,\mu,\eta}^*)$ as well.

Adopting the notation in Section 2, we introduce an extension operator

$$e_\Pi : \mathbb{E}_2(J) \rightarrow \mathbb{E}_2(J, \mathbb{T}_a) : h \mapsto h \circ \Pi.$$

$((\pi_\kappa \circ \Pi)^2)_{\kappa \in \mathfrak{K}}$ forms a partition of unity for \mathbb{T}_a . For $\mathfrak{F} \in \{W_p, H_p\}$, we define

$$\begin{aligned} \mathfrak{R}_\Pi^c &: \mathfrak{F}^s(\mathbb{T}_a, E) \rightarrow \prod_{\kappa \in \mathfrak{K}} \mathfrak{F}^s(\mathbb{T}_{a,\kappa}, E) : u \mapsto (u\pi_\kappa \circ \Pi)_\kappa, \\ \mathfrak{R}_\Pi &: \prod_{\kappa \in \mathfrak{K}} \mathfrak{F}^s(\mathbb{T}_{a,\kappa}, E) \rightarrow \mathfrak{F}^s(\mathbb{T}_a, E) : (u_\kappa)_\kappa \mapsto \sum_{\kappa \in \mathfrak{K}} u_\kappa \pi_\kappa \circ \Pi. \end{aligned}$$

Then \mathcal{R}_Π is a retraction with \mathcal{R}_Π^c as a coretraction. By using this retraction-coretraction system, it is a simple matter to check that

$$e_\Pi \in \mathcal{L}(\mathbb{E}_2(J), \mathbb{E}_2(J, \mathcal{T}_a)).$$

Extending $\mathcal{V} := (\zeta \circ d_\Sigma)(\nu_\Sigma \circ \Pi)$ to be identically zero outside \mathcal{T}_a , it belongs to $BC^\infty(\Omega, \mathbb{R}^{m+1}) \cap C^\omega(\mathcal{T}_{a/3}, \mathbb{R}^{m+1})$. Proposition 3.6(a) implies

$$[(\lambda, \mu, \eta) \mapsto \mathcal{V}_{\lambda, \mu, \eta}] \in C^\omega(\mathbb{B}(0, r_0), C^1(J, BC^k(\Omega, \mathbb{R}^{m+1})))$$

for all $k \in \mathbb{N}_0$. Proposition 3.8 and (4.18) then yield

$$[(h, (\lambda, \mu)) \mapsto \Upsilon_{\lambda, \mu, \eta}(h + h_{\lambda, \mu}^*)] \in C^\omega({}_0\mathbb{E}_2(J) \times \mathbb{B}(0, r_0), \mathbb{E}_2(J, \Omega; \mathbb{R}^{m+1})).$$

Because $\mathbb{F}_j(J, \Omega)$ with $j = 2, 3$ and $W_p^{1-1/2p}(J; H_p^1(\Omega)) \cap L_p(J; W_p^{3-1/p}(\Omega))$ are Banach algebras, we verify via Proposition 3.6(b) that

$$[(h, (\lambda, \mu, \eta)) \mapsto M_{j, \lambda, \mu, \eta}(h + h_{\lambda, \mu}^*)] \in C^\omega({}_0\mathbb{E}_2(J) \times \mathbb{B}(0, r_0), \mathbb{F}_2(J, \Omega; \mathbb{R}^{(m+1)^2})) \quad (4.20)$$

for $j = 1, 2, 3$. Proposition 3.6(b) and (4.18) lead to

$$[(\vartheta, (\lambda, \mu, \eta)) \mapsto T_{\mu, \eta} \nabla T_{\mu, \eta}^{-1}(\vartheta + \vartheta_{\lambda, \mu, \eta})] \in C^\omega({}_0\mathbb{E}_1(J) \times \mathbb{B}(0, r_0), \mathbb{F}_3(J, \Omega_i; \mathbb{R}^{m+1})).$$

In virtue of Proposition 3.6(b) and (3.11), we get

$$[(z, (\lambda, \mu, \eta)) \mapsto \mathcal{R}_{\lambda, \mu, \eta}(h + h_{\lambda, \mu}^*)(\vartheta + \vartheta_{\lambda, \mu, \eta}^*)] \in C^\omega({}_0\mathbb{E}(J) \times \mathbb{B}(0, r_0), \mathbb{F}_3(J, \Omega_i)).$$

Now it is immediate that

$$[(z, (\lambda, \mu)) \mapsto F_{\lambda, \mu, \eta}(z, z_{\lambda, \mu}^*)] \in C^k({}_0\mathbb{E}(J) \times \mathbb{B}(0, r_0), \mathbb{F}_1(J)).$$

Since $\mathbb{C}_2(J)$ is a multiplication algebra, we have

$$[(h, (\lambda, \mu)) \mapsto M_{0, \lambda, \mu}(h + h_{\lambda, \mu}^*)] \in C^\omega({}_0\mathbb{E}_2(J) \times \mathbb{B}(0, r_0), \mathbb{C}_2(J; T_1^1 \Sigma)).$$

Proposition 3.8 and point-wise multiplier results on Σ then imply that

$$[(h, (\lambda, \mu)) \mapsto \alpha_{\lambda, \mu}(h + h_{\lambda, \mu}^*)] \in C^\omega({}_0\mathbb{E}_2(J) \times \mathbb{B}(0, r_0), \mathbb{F}_2(J; T \Sigma)).$$

Combined with the properties for substitution operators for Sobolev-Slobodeckii spaces, it yields

$$[(h, (\lambda, \mu)) \mapsto \beta_{\lambda, \mu}(h + h_{\lambda, \mu}^*)] \in C^\omega({}_0\mathbb{E}_2(J) \times \mathbb{B}(0, r_0), \mathbb{F}_2(J)).$$

It is shown in [72] that in every coordinate patch $(\mathcal{O}_\kappa, \varphi_\kappa)$, the local expression of the mean curvature operator reads as

$$(\mathcal{H}(h))_\kappa = \beta(h) \frac{P_\kappa(h, \partial_j h, \partial_{jk} h)}{R_\kappa(h, \partial_j h)}.$$

Here P_κ is a polynomial in h and its derivatives up to second order, and R_κ is a polynomial in h and its first order derivatives. Both have $BC^\infty \cap C^\omega$ -coefficients. We again use the fact that $\mathbb{F}_2(J)$ is a Banach algebra. Following a similar argument in the same reference, we get

$$[(h, (\lambda, \mu)) \mapsto \mathcal{H}_{\lambda, \mu}(h + h_{\lambda, \mu}^*)] \in C^\omega({}_0\mathbb{E}_2(J) \times \mathbb{B}(0, r_0), \mathbb{F}_2(J)).$$

By (4.20) and trace theorems of anisotropic Sobolev-Slobodeckii spaces, we infer that

$$[(h, (\lambda, \mu)) \mapsto M_{1, \lambda, \mu}(h + h_{\lambda, \mu}^*)|_\Sigma] \in C^\omega({}_0\mathbb{E}_2(J) \times \mathbb{B}(0, r_0), \mathbb{F}_3(J; \mathbb{R}^{(m+1)^2})).$$

It follows from the point-wise multiplier theorem in [5, Section 9] that

$$[(h, (\lambda, \mu)) \mapsto M_{4, \lambda, \mu}(h + h_{\lambda, \mu}^*)] \in C^\omega({}_0\mathbb{E}_2(J) \times \mathbb{B}(0, r_0), \mathbb{F}_3(J; \mathbb{R}^{(m+1)^2})).$$

To sum up, we conclude that

$$\begin{aligned} [(z, (\lambda, \mu)) \mapsto \bar{G}_{\lambda, \mu}(z, z_{\lambda, \mu}^*)] &\in C^k({}_0\mathbb{E}(J) \times \mathbb{B}(0, r_0), \mathbb{F}_2(J)), \\ [(z, (\lambda, \mu)) \mapsto \bar{Q}_{\lambda, \mu}(z, z_{\lambda, \mu}^*)] &\in C^k({}_0\mathbb{E}(J) \times \mathbb{B}(0, r_0), \mathbb{F}_3(J)). \end{aligned}$$

Taking into account all the above discussion, this leads to

$$\Phi \in C^k({}_0\mathbb{E}(J) \times \mathbb{B}(0, r_0), {}_0\mathbb{F}(J)).$$

(ii) We look at the Fréchet derivative of Φ with respect to ${}_0\mathbb{E}(J)$ at $(\bar{z}, 0)$. To this end, we find it more convenient to consider the following non-linear maps.

$$\begin{aligned} \hat{F}(z) &= (\hat{\kappa}_0 - \kappa(\vartheta))\partial_t \vartheta + (d(\vartheta) - \hat{d}_0)\Delta \vartheta - d(\vartheta)M_2(h) : \nabla^2 \vartheta \\ &\quad + d'(\vartheta)(I - M_1(h))\nabla \vartheta|^2 - d(\vartheta)(M_3(h)|\nabla \vartheta) + \kappa(\vartheta)\mathcal{R}(h)\vartheta, \\ \hat{G}(z) &= -([\psi(\vartheta)] + \sigma \mathcal{H}(h)) + \hat{l}_1 \vartheta + \sigma_0 \Delta_\Sigma h + (\gamma(\vartheta)\beta(h) - \hat{\gamma}_1)\partial_t h, \\ \hat{Q}(z) &= [(d(\vartheta) - \hat{d}_0)\partial_v \vartheta] + (\hat{l}_0 - l(\vartheta))\partial_t h - ([d(\vartheta)\nabla \vartheta]|M_4(h)\nabla_\Sigma h) \\ &\quad + \gamma(v)\beta(h)(\partial_t h)^2, \end{aligned}$$

i.e., replacing ϑ_A by $\hat{\vartheta}_0$ in the definition of F , G and H . With only slight modification of (i), we immediately obtain

$$(\hat{F}, \hat{G}, \hat{Q}) \in C^k(\mathbb{E}(J), \mathbb{F}_1(J) \times \mathbb{F}_2(J) \times \mathbb{F}_3(J)).$$

Recall that $\hat{z} = \bar{z} + z^*$. Letting $w = (u, \rho)$, one computes

$$D_1 \Phi(\bar{z}, 0)w = \begin{cases} \hat{\kappa}_0 \partial_t u - \hat{d}_0 \Delta u - \hat{F}'(\hat{z})w & \text{in } \Omega \setminus \Sigma, \\ \hat{l}_1 u + \sigma_0 \Delta_\Sigma \rho - \hat{\gamma}_1 \partial_t \rho - \hat{G}'(\hat{z})w & \text{on } \Sigma, \\ \hat{l}_0 \partial_t \rho - [\hat{d}_0 \partial_v u] - \hat{Q}'(\hat{z})w & \text{on } \Sigma. \end{cases}$$

A moment of reflection shows that we have the liberty to exchange the coefficients (κ_A, d_A, l_A) , used in the definition of Φ , by $(\hat{\kappa}_0, \hat{d}_0, \hat{l}_0)$. These quantities are only used for the principal linearization, and are then subtracted off in the nonlinearities.

We split $D_1\Phi(\bar{z}, 0)$ into two parts. Let $\hat{\beta}_0 := \beta(\hat{h}_0)$. Define $\mathbb{L}(\hat{\vartheta}_0) : {}_0\mathbb{E}(J) \rightarrow {}_0\mathbb{F}(J)$ by

$$\mathbb{L}(\hat{\vartheta}_0)w = \begin{cases} \hat{\kappa}_0\partial_t u - \hat{d}_0(I - M_2(\hat{h}_0)) : \nabla^2 u & \text{in } \Omega \setminus \Sigma, \\ \hat{l}_1 u + \sigma\mathcal{H}'(\hat{h}_0)\rho - \hat{\gamma}_1\beta'(\hat{h}_0)e^{\Delta\Sigma t}(\partial_t\hat{h}(0))\rho \\ \quad - \hat{\gamma}_1\hat{\beta}_0\partial_t\rho - \gamma'(e^{\Delta\Sigma t}\hat{\vartheta}_0)\hat{\beta}_0e^{\Delta\Sigma t}(\partial_t\hat{h}(0))u & \text{on } \Sigma, \\ \hat{l}_0\partial_t\rho - (\llbracket\hat{d}_0\nabla u\rrbracket|_{\nu\Sigma} - M_4(\hat{h}_0)\nabla_\Sigma\hat{h}_0) & \text{on } \Sigma. \end{cases}$$

and $\mathbb{K}(\hat{z}) : {}_0\mathbb{E}(J) \rightarrow {}_0\mathbb{F}(J)$ by

$$\mathbb{K}(\hat{z}) := (\mathbb{K}_1, \mathbb{K}_2, \mathbb{K}_3)^\top := \mathbb{L}(\hat{\vartheta}_0)w - D_1\Phi(\hat{z}, 0).$$

THEOREM 4.2 Let $J = [0, T]$. Then $\mathbb{L}(\hat{\vartheta}_0) \in \mathfrak{Lis}(\mathbb{E}(J), \mathbb{F}(J))$. In particular, $\mathbb{L}(\hat{\vartheta}_0) \in \mathfrak{Lis}({}_0\mathbb{E}(J), {}_0\mathbb{F}(J))$.

Proof. We first solve the following model problem for the case $\gamma \equiv 0$. Suppose that

$$\kappa_0 \in BUC(\mathbb{R}^m \times \mathbb{R}_+), \quad d_0 \in BUC^1(\mathbb{R}^m), \quad a_0 \in BUC^1(\mathbb{R}^m; \mathbb{R}^m), \quad \kappa_0, d_0 > 0.$$

Moreover, $l_0 \in W_p^{2-6/p}(\mathbb{R}^m)$, $l_2 \in \mathbb{F}_2(J, \mathbb{R}^m)$ satisfy $l_2 l_0 > 0$. The differential operators $-P(x, y) : \nabla^2$ and $-S(x) : \nabla_x^2$ are uniformly elliptic. For any given $(f, g, q, (u_0, \rho_0)) \in \mathbb{F}(J, \mathbb{R}^m \times \mathbb{R}_+)$,

$$\begin{cases} \kappa_0(x, y)\partial_t u(x, y) - P(x, y) : \nabla^2 u(x, y) = f(x, y) & \text{in } \mathbb{R}^m \times \mathbb{R}_+, \\ l_2(t, x)u(x) + S : \nabla_x^2 \rho(x) = g(x) & \text{on } \mathbb{R}^m, \\ l_0(x)\partial_t \rho(x) + d_0(x)\partial_\nu u(x) + a_0(x) \cdot \nabla_x u(x) = q(x) & \text{on } \mathbb{R}^m, \\ u(0) = u_0, \quad \rho(0) = \rho_0, \end{cases}$$

admits a unique solution $(u, \rho) \in \mathbb{E}(J, \mathbb{R}^m \times \mathbb{R}_+)$. We identify \mathbb{R}^m with $\mathbb{R}^m \times \{0\}$. Here $x \in \mathbb{R}^m$, $y \in \mathbb{R}_+$, and the space $\mathbb{F}(J, \mathbb{R}^m \times \mathbb{R}_+)$ is defined by replacing $\Omega \setminus \Sigma$ and Σ by $\mathbb{R}^m \times \mathbb{R}_+$ and \mathbb{R}^m , respectively. The space $\mathbb{E}(J, \mathbb{R}^m \times \mathbb{R}_+)$ is defined analogously. Similar problems have been considered in [22]. It suffices to check the Lopatinskii-Shapiro condition **(LS)** and the asymptotic Lopatinskii-Shapiro condition **(LS_∞⁺)** defined therein. For simplicity, we assume all the coefficients to be constants and set

$$P(\xi) := \xi_e^\top P \xi_e, \quad S(\xi) := \xi^\top S \xi, \quad P_{m+1} := P_{m+1, m+1}$$

and $P_m := (P_{m+1, 1}, \dots, P_{m+1, m})$ with $\xi \in \mathbb{R}^m$ and $\xi_e := (\xi^\top, 0)^\top \in \mathbb{R}^{m+1}$. **(LS)** is satisfied, if for any $\xi \in \mathbb{R}^m$ and $\lambda \in \mathbb{C}_+$ with $|\xi| + |\lambda| \neq 0$, the following ordinary differential equation in \mathbb{R}_+

$$\begin{cases} (\kappa_0\lambda + P(\xi) + 2iP_m \cdot \xi\partial_y - P_{m+1}\partial_y^2)v(y) = 0, & y > 0, \\ l_2v(0) - S(\xi)\delta = 0, \\ l_0\lambda\delta - (ia_0 \cdot \xi + d_0(x)\partial_y)v(0) = 0, \end{cases}$$

has a unique solution $(v, \delta) \in C_0(\mathbb{R}_+; \mathbb{C}) \times \mathbb{C}$. It is clear that the only stable solution to the first line is

$$v(y) = e^{\mu y}v(0), \quad \mu := \frac{\sqrt{-(P_m \cdot \xi)^2 + P_{m+1}\kappa_0\lambda + P_{m+1}P(\xi)} + iP_m \cdot \xi}{P_{m+1}}$$

in the case $\lambda \notin \mathbb{R}$, or, $v \equiv 0$, otherwise. We conclude from the second and third lines that

$$[iS(\xi)a_0 \cdot \xi + S(\xi)\mu d_0 - l_2 l_0 \lambda] \delta = 0.$$

It implies that $\delta = 0$, $v \equiv 0$. (\mathbf{LS}_∞^+) is satisfied, if for any $\xi \in \mathbb{R}^m$ and $\lambda \in \bar{\mathbb{C}}_+$ with $|\xi| + |\lambda| \neq 0$,

$$\begin{cases} (\kappa_0 \lambda + P(\xi) + 2iP_m \cdot \xi \partial_y - P_{m+1} \partial_y^2)v(y) = 0, & y > 0, \\ l_2 v(0) - S(\xi)\delta = 0, \\ (i a_0 \cdot \xi + d_0(x) \partial_y)v(0) = 0, \end{cases}$$

and $|\xi| = 1$, $\lambda \in \bar{\mathbb{C}}_+ \setminus \{0\}$

$$\begin{cases} (\kappa_0 \lambda - P_{m+1} \partial_y^2)v(y) = 0, & y > 0, \\ l_2 v(0) - S(\xi)\delta = 0, \\ l_0 \lambda \delta + d_0(x) \partial_y v(0) = 0, \end{cases}$$

and $|\xi| = 1$, $\lambda \in \bar{\mathbb{C}}_+ \setminus \{0\}$

$$\begin{cases} (\kappa_0 \lambda - P_{m+1} \partial_y^2)v(y) = 0, & y > 0, \\ l_2 v(0) - S(\xi)\delta = 0, \\ d_0(x) \partial_y v(0) = 0, \end{cases}$$

admit unique solutions $(v, \delta) \in C_0(\mathbb{R}_+; \mathbb{C}) \times \mathbb{C}$. One can check in an analogous manner that stable solutions to those equations are trivial. For general coefficients, the problem can be proved by a perturbation argument as in [21]. The rest of the proof now follows from similar arguments to those of [65, Theorems 3.3, 3.5]. \square

In order to prove that $D_1 \Phi(\bar{z}, 0)$ is an isomorphism, we need to control the norm $\|\mathbb{K}(\hat{z}_0)\|_{\mathcal{L}(0, \mathbb{E}(J), 0, \mathbb{F}(J))}$. To this end, we first compute several derivatives related to \mathbb{K} explicitly.

$$\begin{aligned} \hat{F}'(z)w &= (\hat{\kappa}_0 - \kappa(\vartheta)) \partial_t u - \kappa'(\vartheta) u \partial_t \vartheta + (d(\vartheta) - \hat{d}_0) \Delta u + d'(\vartheta) u \Delta \vartheta \\ &\quad - d'(\vartheta) u M_2(h) : \nabla^2 \vartheta - d(\vartheta) M_2'(h) \rho : \nabla^2 \vartheta - d(\vartheta) M_2(h) : \nabla^2 u \\ &\quad - d'(\vartheta) u (M_3(h) |\nabla \vartheta| - d(\vartheta) (M_3'(h) \rho |\nabla \vartheta| - d(\vartheta) (M_3(h) |\nabla u) \\ &\quad + 2d'(\vartheta) (I - M_1(h) \nabla \vartheta | (I - M_1(h)) \nabla u - M_1'(h) \rho \nabla \vartheta) \\ &\quad + d''(\vartheta) u | (I - M_1(h)) \nabla \vartheta|^2 + \kappa(\vartheta) (\nabla u | (I + \nabla \Upsilon(h)^\top)^{-1} \partial_t \Upsilon(h)) \\ &\quad - \kappa(\vartheta) (\nabla \vartheta | (I + \nabla \Upsilon(h)^\top)^{-1} \nabla \Upsilon(\rho)^\top (I + \nabla \Upsilon(h)^\top)^{-1} \partial_t \Upsilon(h)) \\ &\quad + \kappa(\vartheta) (\nabla \vartheta | (I + \nabla \Upsilon(h)^\top)^{-1} \partial_t \Upsilon(\rho)) + \kappa'(\vartheta) u \mathcal{R}(h) \vartheta, \end{aligned}$$

and

$$\begin{aligned} \hat{G}'(z)w &= -([\psi'(\vartheta)])u + \sigma \mathcal{H}'(h) \rho + \hat{l}_1 u + \sigma_0 \Delta_\Sigma \rho + (\gamma(\vartheta) \beta(h) - \hat{\gamma}_1) \partial_t \rho \\ &\quad + (\gamma'(\vartheta) u \beta(h) - \gamma(\vartheta) \beta'(h) \rho) \partial_t h, \end{aligned}$$

and

$$\begin{aligned}\hat{Q}'(z)w &= \llbracket d'(\vartheta)\partial_\nu\vartheta \rrbracket u + \llbracket (d(\vartheta) - \hat{d}_0)\partial_\nu u \rrbracket + (\hat{l}_0 - l(\vartheta))\partial_t\rho - l'(\vartheta)u\partial_t h \\ &\quad - (\llbracket d'(\vartheta)\nabla\vartheta \rrbracket u \mid M_4(h)\nabla_\Sigma h) - (\llbracket d(\vartheta)\nabla u \rrbracket \mid M_4(h)\nabla_\Sigma h) \\ &\quad - (\llbracket d(\vartheta)\nabla\vartheta \rrbracket \mid M_4(h)\nabla_\Sigma\rho) - (\llbracket d(\vartheta)\nabla\vartheta \rrbracket \mid M_4'(h)\rho\nabla_\Sigma h) \\ &\quad + \gamma'(\vartheta)u\beta(h)[\partial_t h]^2 + \gamma(\vartheta)\beta'(h)\rho[\partial_t h]^2 + 2\gamma(\vartheta)\beta(h)\partial_t h\partial_t\rho.\end{aligned}$$

The derivatives of $M_0(h)$, $\alpha(h)$, $\beta(h)$ and $\mathcal{H}(h)$ are given by

$$\begin{aligned}M_0'(h)\rho &= \rho M_0(h)L_\Sigma M_0(h), \quad \alpha'(h)\rho = M_0(h)\nabla_\Sigma\rho + \rho M_0(h)L_\Sigma M_0(h)\alpha(h), \\ \beta'(h)\rho &= -\beta^3(h)(\alpha(h) \mid M_0(h)\nabla_\Sigma\rho + \rho M_0(h)L_\Sigma\alpha(h)), \\ \mathcal{H}'(h)\rho &= \beta(h)\left\{\text{tr}[M_0'(h)\rho(L_\Sigma + \nabla_\Sigma\alpha(h))] + \text{tr}[M_0(h)\nabla_\Sigma\alpha'(h)\rho] \right. \\ &\quad - 2\beta(h)\beta'(h)\rho(M_0(h)\alpha(h) \mid [\nabla_\Sigma\alpha(h)]\alpha(h)) - \beta^2(h)(M_0'(h)\rho\alpha(h) \mid [\nabla_\Sigma\alpha(h)]\alpha(h)) \\ &\quad - \beta^2(h)(M_0(h)\alpha'(h)\rho \mid [\nabla_\Sigma\alpha(h)]\alpha(h)) - \beta^2(h)(M_0(h)\alpha(h) \mid [\nabla_\Sigma\alpha'(h)\rho]\alpha(h)) \\ &\quad \left. - \beta^2(h)(M_0(h)\alpha(h) \mid [\nabla_\Sigma\alpha(h)]\alpha'(h)\rho)\right\}/m + \beta'(h)\rho(\mathcal{H}(h)/\beta(h)).\end{aligned}$$

See [63, formula (32)] for a justification for the last equality.

We will use the following lemma frequently in the sequel.

LEMMA 4.3 There exists a constant C_0 independent of T such that

(a) For all $(v_1, v_2) \in \mathbb{F}_j(J) \times {}_0\mathbb{F}_j(J)$ and $j = 2, 3$,

$$\|v_1 v_2\|_{\mathbb{F}_j(J)} \leq C_0(\|v_1\|_{C(J \times \Sigma)} + \|v_1\|_{\mathbb{F}_j(J)})\|v_2\|_{\mathbb{F}_j(J)}.$$

(b) For all $(v_1, v_2) \in {}_0\mathbb{F}_j(J) \times {}_0\mathbb{F}_j(J)$ and $j = 2, 3$,

$$\|v_1 v_2\|_{{}_0\mathbb{F}_j(J)} \leq C_0\|v_1\|_{{}_0\mathbb{F}_j(J)}\|v_2\|_{{}_0\mathbb{F}_j(J)}.$$

(c) For all $(v_1, v_2) \in \mathbb{F}_3(J) \times {}_0\mathbb{F}_2(J)$

$$\|v_1 v_2\|_{\mathbb{F}_3(J)} \leq C_0\|v_1\|_{\mathbb{F}_3(J)}\|v_2\|_{{}_0\mathbb{F}_2(J)}.$$

Proof. (a) The case $j = 2$ is shown in [60, Lemma 5.5(b)]. For the reader's convenience, we will nevertheless include a proof herein. The case $j = 3$ follows in a similar way. By the retraction-coretraction system defined in Section 2, it suffices to show the estimates for functions on \mathcal{Q}^m , i.e., we assume $(v_1, v_2) \in \mathbb{F}_j(J, \mathcal{Q}^m) \times {}_0\mathbb{F}_j(J, \mathcal{Q}^m)$. We equip $\mathbb{F}_2(J, \mathcal{Q}^m)$ with the norm:

$$\begin{aligned}\|v\|_{\mathbb{F}_2(J, \mathcal{Q}^m)} &= \|v\|_{W_p^{1-1/2p}(J; L_p(\mathcal{Q}^m))} + \|v\|_{L_p(J; W_p^{2-1/p}(\mathcal{Q}^m))}, \\ \|v\|_{L_p(J; W_p^{2-1/p}(\mathcal{Q}^m))} &= \|v\|_{L_p(J; H_p^1(\mathcal{Q}^m))} + \sum_{j=1}^m \|\langle \partial_j v \rangle_{W_p^{1-1/p}(\mathcal{Q}^m)}\|_{L_p(J)}.\end{aligned}$$

Here $\langle \cdot \rangle_{W_p^{1-1/p}(J; X)}$ is the Slobodeckii seminorm of the space $W_p^{1-1/p}(J; X)$ for a Banach space

X. Then,

$$\begin{aligned} \|v_1 v_2\|_{W_p^{1-1/2p}(J;L_p(Q^m))} &\leq C' \left\{ \|v_1\|_{W_p^{1-1/2p}(J;L_p(Q^m))} \|v_2\|_{BC(J \times Q^m)} \right. \\ &\quad \left. + \left(\int_J \int_J \|v_1(s)(v_2(t) - v_2(s))\|_{L_p(Q^m)}^p \frac{1}{|t-s|^{1/2+p}} dt ds \right)^{1/p} \right\} \\ &\leq C' \left\{ \|v_1\|_{W_p^{1-1/2p}(J;L_p(Q^m))} \|v_2\|_{BC(J \times Q^m)} + \|v_1\|_{BC(J \times Q^m)} \langle v_2 \rangle_{W_p^{1-1/2p}(J;L_p(Q^m))} \right\}. \end{aligned}$$

Similarly, one computes

$$\begin{aligned} \|v_1 v_2\|_{L_p(J;W_p^{2-1/p}(Q^m))} &\leq C' \left\{ \|v_1\|_{L_p(J;W_p^{2-1/p}(Q^m))} \|v_2\|_{C(J,BC^1(Q^m))} \right. \\ &\quad \left. + \sum_{j=1}^m \left[\int_J \left(\int_{Q^m} \int_{Q^m} |v_1(t,x)(\partial_j v_2(t,x) - \partial_j v_2(t,y)|^p \frac{1}{|x-y|^{m-1+p}} dx dy \right) dt \right]^{1/p} \right. \\ &\quad \left. + \sum_{j=1}^m \left[\int_J \left(\int_{Q^m} \int_{Q^m} |\partial_j v_1(t,x)(v_2(t,x) - v_2(t,y)|^p \frac{1}{|x-y|^{m-1+p}} dx dy \right) dt \right]^{1/p} \right\}. \end{aligned}$$

We immediately have

$$\begin{aligned} \int_J \left(\int_{Q^m} \int_{Q^m} |v_1(t,x)(\partial_j v_2(t,x) - \partial_j v_2(t,y)|^p \frac{1}{|x-y|^{m-1+p}} dx dy \right) dt \\ \leq \|v_1\|_{BC(J \times Q^m)}^p \|v_2\|_{L_p(J,W_p^{2-1/p}(Q^m))}^p. \end{aligned}$$

The remaining estimate can be carried out as follows.

$$\begin{aligned} &\int_{Q^m} \int_{Q^m} |\partial_j v_1(t,x)(v_2(t,x) - v_2(t,y)|^p \frac{1}{|x-y|^{m-1+p}} dx dy \\ &\leq \int_{Q^m} \int_{Q^m} |\partial_j v_1(t,x)|^p \left(\int_0^1 |(\nabla v_2(t,x + \tau(y-x)))(y-x)| d\tau \right)^p \frac{1}{|x-y|^{m-1+p}} dx dy \\ &\leq C' \int_{Q^m} |\partial_j v_1(t,x)|^p \int_{Q^m} \frac{|x-y|^p}{|x-y|^{m-1+p}} dy dx \|\nabla v_2(t)\|_{BC(Q^m)}^p \\ &\leq C' \|v_1(t)\|_{H_p^1(Q^m)}^p \|v_2(t)\|_{BC^1(Q^m)}^p. \tag{4.21} \end{aligned}$$

Combining these discussions yields

$$\begin{aligned} \|v_1 v_2\|_{L_p(J;W_p^{2-1/p}(Q^m))} &\leq C' \left(\|v_1\|_{BC(J \times Q^m)} + \|v_1\|_{L_p(J;W_p^{2-1/p}(Q^m))} \right) \\ &\quad \left(\|v_2\|_{C(J;BC^1(Q^m))} + \|v_2\|_{L_p(J;W_p^{2-1/p}(Q^m))} \right). \end{aligned}$$

Using the fact that the embedding constant of ${}_0\mathbb{F}_2(J, Q^m) \hookrightarrow C(J; BC^1(Q^m))$ is independent of T yields the asserted result.

(b) is an immediate consequence of (a) and the fact that the embedding constant of ${}_0\mathbb{F}_2(J, Q^m) \hookrightarrow C(J; BC(Q^m))$ is independent of T .

(c) Suppose that $(v_1, v_2) \in \mathbb{F}_3(J, \mathbb{Q}^m) \times {}_0\mathbb{F}_2(J, \mathbb{Q}^m)$. Then

$$\begin{aligned} \|v_1 v_2\|_{W_p^{1/2-1/2p}(J; L_p(\mathbb{Q}^m))} &\leq \|v_1\|_{W_p^{1/2-1/2p}(J; L_p(\mathbb{Q}^m))} \|v_2\|_{BC(J \times \mathbb{Q}^m)} \\ &\quad + \left(\int_J \int_J \|v_1(s)(v_2(t) - v_2(s))\|_{L_p(\mathbb{Q}^m)}^p \frac{1}{|t-s|^{1/2+p/2}} dt ds \right)^{1/p}. \end{aligned}$$

By [55, Lemma 2.5], the comments below [55, formula (3.5)], an analogue of [58, diagram (6.12)] with BUC replaced by C^s , we obtain the embedding result

$${}_0\mathbb{F}_2(J, \mathbb{Q}^m) \hookrightarrow C^s(J; BC(\mathbb{Q}^m))$$

for some $s > 1/2 - 1/2p$ with an embedding constant $M = M(s)$ uniform in T . Thus, we infer that

$$\begin{aligned} \int_J \int_J \|v_1(s)(v_2(t) - v_2(s))\|_{L_p(\mathbb{Q}^m)}^p \frac{1}{|t-s|^{1/2+p/2}} dt ds \\ \leq C \int_J \|v_1(s)\|_{L_p(\mathbb{Q}^m)}^p \int_J \frac{1}{|t-s|^{1/2+p/2-sp}} dt ds \|v_2\|_{C^s(J; BC(\mathbb{Q}^m))}^p \\ \leq C' \|v_1\|_{L_p(J; L_p(\mathbb{Q}^m))}^p \|v_2\|_{C^s(J; BC(\mathbb{Q}^m))}^p. \end{aligned}$$

By an analogous estimate as in (a), see in particular (4.21), one obtains

$$\|v_1 v_2\|_{L_p(J; W_p^{1-1/p}(\mathbb{Q}^m))} \leq C' \|v_1\|_{L_p(J; W_p^{1-1/p}(\mathbb{Q}^m))} \|v_2\|_{C(J; BC^1(\mathbb{Q}^m))}.$$

□

Note that the multiplication constant C_0 in (b) blows up as $T \rightarrow 0$ if ${}_0\mathbb{F}_j(J)$ is replaced by $\mathbb{F}_j(J)$.

We write

$$\begin{aligned} \mathbb{K}_2(\hat{z})w &= \hat{l}_1 u - \llbracket \psi'(\hat{\vartheta}) \rrbracket u + (\gamma(\hat{\vartheta}) - \hat{\gamma}_1) \beta(\hat{h}) \partial_t \rho + \hat{\gamma}_1 (\beta(\hat{h}) - \hat{\beta}_0) \partial_t \rho \\ &\quad - \sigma(\mathcal{H}'(\hat{h}) - \mathcal{H}'(\hat{h}_0)) \rho + (\gamma(\hat{\vartheta}) \beta'(\hat{h}) \partial_t \hat{h} - \hat{\gamma}_1 \beta'(\hat{h}_0) e^{\Delta \Sigma t} (\partial_t \hat{h}(0))) \rho \\ &\quad + \left(\gamma'(\hat{\vartheta}) \beta(\hat{h}) \partial_t \hat{h} - \gamma'(e^{\Delta \Sigma t} \hat{\vartheta}_0) \hat{\beta}_0 e^{\Delta \Sigma t} (\partial_t \hat{h}(0)) \right) u. \end{aligned}$$

Let ε sufficiently small be fixed. Recall that

$$\hat{l}_1 u - \llbracket \psi'(\hat{\vartheta}) \rrbracket u = (\llbracket \psi'(e^{\Delta \Sigma t} \hat{\vartheta}_0) \rrbracket - \psi'(\hat{\vartheta})) u.$$

Due to fact that $\llbracket \psi'(e^{\Delta \Sigma t} \hat{\vartheta}_0) \rrbracket - \psi'(\hat{\vartheta}) \in {}_0\mathbb{F}_2(J)$ and Lemma 4.3(b), by making T small enough, we can achieve that

$$\|\hat{l}_1 u - \llbracket \psi'(\hat{\vartheta}) \rrbracket u\|_{\mathbb{F}_2(J)} \leq \varepsilon \|u\|_{\mathbb{F}_2(J)}.$$

The arguments for the remaining terms in $\mathbb{K}_2(\hat{z})w$ are similar. Thus given any $\varepsilon > 0$, for T small enough, we have

$$\|\mathbb{K}_2(\hat{z})w\|_{\mathbb{F}_2(J)} \leq \varepsilon \|w\|_{{}_0\mathbb{E}(J)}.$$

One can obtain an analogous assertion for $\|\mathbb{K}_3(\hat{z})\|_{\mathfrak{L}(0\mathbb{E}(J), 0\mathbb{F}_3(J))}$. Indeed,

$$\|[(d(\vartheta) - \hat{d}_0)\partial_v u]\|_{\mathbb{F}_3(J)} \leq C_0 \| (d(\vartheta) - \hat{d}_0) \|_{0\mathbb{F}_3(J)} \|[\partial_v u]\|_{0\mathbb{F}_3(J)}.$$

Similar estimates also apply to

$$\|(\hat{l}_0 - l(\hat{\vartheta}))\partial_t \rho\|_{\mathbb{F}_3(J)}, \|([\hat{d}(\hat{h})\nabla u]|M_4(\hat{h})\nabla \hat{h}) - ([\hat{d}_0\nabla u]|M_4(\hat{h}_0)\nabla \hat{h}_0)\|_{\mathbb{F}_3(J)}.$$

It follows from Lemma 4.3(c) that

$$\|[\hat{d}'(\hat{\vartheta})\partial_v \hat{\vartheta}]u\|_{\mathbb{F}_3(J)} \leq C_0 \|[\hat{d}'(\hat{\vartheta})\partial_v \hat{\vartheta}]\|_{\mathbb{F}_3(J)} \|u\|_{0\mathbb{F}_2(J)}.$$

The remaining terms in $\mathbb{K}_3(\hat{z})w$ can be estimates in an analogous way.

For $\|\mathbb{K}_1(\hat{z})\|_{\mathfrak{L}(0\mathbb{E}(J), \mathbb{F}_1(J))}$, one verifies by direct computation that

$$\begin{aligned} \|\mathbb{K}_1(\hat{z})w\|_{\mathbb{F}_1(J)} &= \|\hat{F}'(\hat{z})w + \hat{d}_0 M_2(\hat{h}_0) : \nabla^2 u\|_{\mathbb{F}_1(J)} \\ &\leq C_2 \|w\|_{\mathbb{E}(J)} + C_3 \|w\|_{\mathbb{C}_1(J) \times \mathbb{C}_2(J)}. \end{aligned}$$

Here the constants C_2 and C_3 tend to zero as $T \rightarrow 0$.

PROPOSITION 4.4 Let $p > m + 3$, $\sigma > 0$. Suppose that $d_i \in C^2(0, \infty)$, $\gamma, \psi_i \in C^3(0, \infty)$. Then there exists some constant τ_0 such that given any $T \leq \tau_0$, on $J = [0, T]$, we have

$$\|\mathbb{L}^{-1}(\hat{\vartheta}_0)\|_{\mathfrak{L}(0\mathbb{F}(J), 0\mathbb{E}(J))} \|\mathbb{K}'(\hat{z}_0)\|_{\mathfrak{L}(0\mathbb{E}(J), 0\mathbb{F}(J))} \leq 1/2.$$

It follows from a Neuman series argument that

$$D_1 \Phi(\bar{z}, 0) \in \mathfrak{L}is(0\mathbb{E}(J), 0\mathbb{F}(J)).$$

Employing now the implicit function theorem, we attain

$$[(\lambda, \mu, \eta) \mapsto \bar{z}_{\lambda, \mu, \eta}] \in C^k(\mathbb{B}(0, r_0), 0\mathbb{E}(J)).$$

It follows then from Theorem 3.12 that

$$\bar{z} \in C^k(\mathring{J} \times \mathbb{T}_{a/3} \setminus \Sigma) \times C^k(\mathring{J} \times \Sigma).$$

Remark 4.5. By a similar argument to the proof given at the end of Section 4.2, we can show that

$$\bar{z} \in C^k(\mathring{J} \times \Omega \setminus \Sigma) \times C^k(\mathring{J} \times \Sigma).$$

Together with (4.18), we thus have

$$\hat{z} \in C^k(\mathring{J} \times \Omega \setminus \Sigma) \times C^k(\mathring{J} \times \Sigma).$$

For $k \in \mathbb{N} \cup \{\infty\}$, since the Hanzawa transformation is C^∞ -smooth, the above assertion implies that the solution (θ, Γ) to (1.4) has the same regularity.

But when $k = \omega$, analyticity of the temperature θ , in general, cannot be attained by applying the Hanzawa transformation. \square

REFERENCES

1. ALMGREN, F. & L. WANG, L., Mathematical existence of crystal growth with Gibbs–Thomson curvature effects. *J. Geom. Anal.* **10** (2000), 1–100. [Zb10981.74041](#) [MR1758583](#)
2. ALT, H.W. & CAFFARELLI, L.A., Existence and regularity for a minimum problem with free boundary. *J. Reine Angew. Math.* **325** (1981), 105–144. [Zb10449.35105](#) [MR0618549](#)
3. ALT, H.W. , CAFFARELLI, L.A. & FRIEDMAN, A., Variational problems with two phases and their free boundaries. *Trans. Amer. Math. Soc.* **282** (1984), 431–461. [Zb10844.35137](#) [MR0732100](#)
4. AMANN, H., *Anisotropic function spaces and maximal regularity for parabolic problems*. Part 1. Function spaces. Jindřich Nečas Center for Mathematical Modeling Lecture Notes, **6**. Matfyzpress, Prague, 2009. [Zb11216.35001](#) [MR2907677](#)
5. AMANN, H., Function spaces on singular manifolds. *Math. Nachr.* **286** (2013), 436–475. [Zb11280.46022](#) [MR3048125](#)
6. AMANN, H., Anisotropic function spaces on singular manifolds. arXiv.1204.0606.
7. ANDERSON, D. M. , CERMELLI, P., FRIED, E., GURTIN, M. E. & MCFADDEN, G. B., General dynamical sharp-interface conditions for phase transformations in viscous heat-conducting fluids. *J. Fluid Mech.* **581** (2007), 323–370. [Zb11119.76066](#) [MR2333066](#)
8. ANGENENT, S. B., Analyticity of the interface of the porous media equation after the waiting time. *Proc. Amer. Math. Soc.* **102** (1988), 329–336. [Zb10653.35040](#) [MR0920995](#)
9. ANGENENT, S. B., Parabolic equations for curves on surfaces. I. Curves with p-integrable curvature, *Ann. of Math. (2)* **132** (1990), 451–483 . [Zb10789.58070](#) [MR1078266](#)
10. ATHANASOPOULOS, I., CAFFARELLI, L. & SALSA, S., Regularity of the free boundary in parabolic phase-transition problems. *Acta Math. (2)* **176**, 245–282 (1996). [Zb10891.35164](#) [MR1397563](#)
11. ATHANASOPOULOS, I., CAFFARELLI, L. & SALSA, S., Phase transition problems of parabolic type: flat free boundaries are smooth. *Comm. Pure Appl. Math* **51** (1998), 77–112. [Zb10924.35197](#) [MR1486632](#)
12. BERESTYCKI, H., CAFFARELLI, L. A. & NIRENBERG, L., *Uniform estimates for regularization of free boundary problems*. Analysis and partial differential equations, 567–619, Lecture Notes in Pure and Appl. Math., 122, Dekker, New York, 1990. [Zb10702.35252](#) [MR1044809](#)
13. BROWDER, F. E., Analyticity and partial differential equations I. *Amer. J. Math.* **84** (1962), 666–710. [Zb10124.31004](#) [MR0150463](#)
14. CAFFARELLI, L. A., The regularity of free boundaries in higher dimensions. *Acta Math.* **139** (1977), 155–184. [Zb10386.35046](#) [MR0454350](#)
15. CAFFARELLI, L. A., Some aspects of the one-phase Stefan problem. *Indiana Univ. Math. J.* **27** (1978), 73–77 . [Zb10393.35064](#) [MR0466965](#)
16. CAFFARELLI, L. A., A Harnack inequality approach to the regularity of free boundaries. I. Lipschitz free boundaries are $C^{1+\alpha}$. *Rev. Mat. Iberoamericana* **3** (1987), 139–162 . [Zb10676.35085](#) [MR0990856](#)
17. CAFFARELLI, L. A. & EVANS, L. C., Continuity of the temperature in the two-phase Stefan problem. *Arch. Rational Mech. Anal.* **81** (1983), 199–220. [Zb10516.35080](#) [MR0683353](#)
18. CAFFARELLI, L. A. & FRIEDMAN, A., Continuity of the temperature in the Stefan problem. *Indiana Univ. Math. J.* **28**, (1979), 53–70. [Zb10406.35032](#) [MR0523623](#)
19. CAGINALP, G., An analysis of a phase field model of a free boundary. *Arch. Rational Mech. Anal.* **92** (1986), 205–245. [Zb10608.35080](#) [MR0816623](#)
20. CHALMERS, B., *Principles of Solidification*. Krieger, Huntington, 1977.
21. DENK, R., HIEBER, M. & PRÜSS, J., \mathcal{R} -boundedness, Fourier multipliers and problems of elliptic and parabolic type. *Mem. Amer. Math. Soc.* **166**, (2003). [Zb11274.35002](#) [MR2006641](#)
22. DENK, R., PRÜSS, J. & ZACHER, R., Maximal Lp-regularity of parabolic problems with boundary dynamics of relaxation type. *J. Funct. Anal.* **255** (2008), 3149–3187. [Zb11160.35030](#) [MR2464573](#)

23. DiBENEDETTO, E., Regularity properties of the solution of an n -dimensional two-phase Stefan problem. *Boll. Un. Mat. Ital. Suppl.* (1980), 129–152. [Zb10458.35098](#) [MR0677695](#)
24. DiBENEDETTO, E., Continuity of weak solutions to certain singular parabolic equations. *Ann. Mat. Pura Appl.* **130** (1982), 131–176. [Zb10503.35018](#) [MR0663969](#)
25. DUVAUT, G., Résolution d'un problème de Stefan (fusion d'un bloc de glace à zéro degré). *C. R. Acad. Sci. Paris Sér. A-B* **276** (1973), A1461–A1463. [Zb10258.35037](#) [MR0328346](#)
26. ESCHER, J., PRÜSS, J. & SIMONETT, G., *A new approach to the regularity of solutions for parabolic equations*. Evolution equations, 167–190, Lecture Notes in Pure and Appl. Math. **234**, Dekker, New York, 2003. [Zb11070.35009](#) [MR2073744](#)
27. ESCHER, J., PRÜSS, J. & SIMONETT, G., Analytic solutions for a Stefan problem with Gibbs–Thomson correction. *J. Reine Angew. Math.* **563** (2003), 1–52. [Zb11242.35220](#) [MR2009238](#)
28. ESCHER, J. & SIMONETT, G., Analyticity of the interface in a free boundary problem. *Math. Ann.* **305** (1996), 439–459. [Zb10857.76086](#) [MR1397432](#)
29. ESCHER, J. & SIMONETT, G., A center manifold analysis for the Mullins–Sekerka Model. *J. Differential Equations* **143** (1998), 267–292. [Zb10896.35142](#) [MR1607952](#)
30. FRIEDMAN, A., The Stefan problem in several space variables. *Trans. Amer. Math. Soc.* **133** (1968), 51–87. [Zb10162.41903](#) [MR0227625](#)
31. FRIEDMAN, A., The Stefan problem with small surface tension. *Trans. Amer. Math. Soc.* **328** (1991), 465–515. [Zb10744.35068](#) [MR1040260](#)
32. FRIEDMAN, A. & KINDERLEHRER, D., A one phase Stefan problem. *Indiana Univ. Math. J.* **24** (1974/75), 1005–1035 [Zb10334.49002](#) [MR0385326](#)
33. FRIEDMAN, A. & REITICH, F., *Variational principles and free-boundary problems*. Second edition. Robert E. Krieger Publishing Co., Inc., Malabar, FL 1988. [MR1009785](#)
34. GURTIN, M. E., On the two-phase Stefan problem with interfacial energy and entropy. *Arch. Rational Mech. Anal.* **96**, no. 3, 199–241 (1986). [Zb10654.73008](#) [MR0855304](#)
35. GURTIN, M. E., Multiphase thermomechanics with interfacial structure. I. Heat conduction and the capillary balance law. *Arch. Rational Mech. Anal.* **104** (1988), 195–221. [Zb10723.73016](#) [MR1017288](#)
36. GURTIN, M. E., Toward a nonequilibrium thermodynamics of two-phase materials. *Arch. Rational Mech. Anal.* **100** (1988), 275–312. [Zb10673.73007](#) [MR0918798](#)
37. HADŽIĆ, M., Orthogonality conditions and asymptotic stability in the Stefan problem with surface tension. *Arch. Ration. Mech. Anal.* **203** (2012), 719–745. [Zb11268.80006](#) [MR2928131](#)
38. HADŽIĆ, M. & GUO, Y., Stability in the Stefan problem with surface tension (I). *Comm. Partial Differential Equations* **35** (2010), 201–244. [Zb11195.35303](#) [MR2748623](#)
39. HANZAWA, E. I., Classical solution of the Stefan problem. *Tôhoku Math. J (2)* **33** (1981), 297–335. [Zb10571.35109](#) [MR0633045](#)
40. HARTMAN, P., *Crystal Growth: An Introduction*. North-Holland, 1973.
41. ISHII, M. & TAKASHI, H., *Thermo-fluid dynamics of two-phase flow*. Springer, New York, 2006. [Zb11204.76002](#)
42. KAMENOMOSTSKAJA, S. L., On Stefan's problem. *Mat. Sb. (N.S.)* **53 (95)** 489–514 (1961). [MR0141895](#)
43. KINDERLEHRER, D. & NIRENBERG, L., Regularity in free boundary problems. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **4** (1977), 373–391. [Zb10352.35023](#) [MR0440187](#)
44. KINDERLEHRER, D. & NIRENBERG, L., The smoothness of the free boundary in the one phase Stefan problem. *Comm. Pure Appl. Math.* **31** (1978), 257–282. [Zb10391.35060](#) [MR0480348](#)
45. KINDERLEHRER, D., NIRENBERG, L. & SPRUCK, J., Regularity in elliptic free boundary problems. *J. Analyse Math.* **34** (1978), 86–119 (1979). [Zb10402.35045](#) [MR0531272](#)

46. KINDERLEHRER, D. & STAMPACCHIA, G., *An introduction to variational inequalities and their applications*. Pure and Applied Mathematics, 88. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1980. [Zb10457.35001](#) [MR0567696](#)
47. KOCH, H., Classical solutions to phase transition problems are smooth. *Comm. Partial Differential Equations* **23** (1998), 389–437. [Zb10904.35014](#) [MR1620648](#)
48. LANGER, J. S., Instabilities and pattern formation of crystal growth. *Rev. Mod. Phys.* **52** (1980), 1–28.
49. LUCKHAUS, S., Solutions for the two-phase Stefan problem with the Gibbs–Thomson law for the melting temperature. *European J. Appl. Math.* **1** (1990), 101–111. [Zb10734.35159](#) [MR1117346](#)
50. LUNARDI, A., *Analytic Semigroups and Optimal Regularity in Parabolic Problems*. Birkhäuser Verlag, Basel, 1995. [Zb10816.35001](#) [MR1329547](#)
51. MATANO, H., *Asymptotic behavior of the free boundaries arising in one phase Stefan problems in multi-dimensional spaces*. Lecture Notes in Num. Appl. Anal. **5**, Kinokuniya, Tokyo, 133–151 (1982). [Zb10569.35076](#) [MR0730240](#)
52. MEÏRMANOV, A. M., On the classical solution of the multidimensional Stefan problem for quasilinear parabolic equations. *Mat. Sb. (N.S.)* **112(154)** (1980), 170–192. [MR0585773](#)
53. MEÏRMANOV, A. M., *The Stefan problem*. Translated from the Russian by Marek Niezgodka and Anna Crowley. With an appendix by the author and I. G. Götz. de Gruyter Expositions in Mathematics, 3. Walter de Gruyter & Co., Berlin, 1992. [Zb10751.35052](#) [MR1154310](#)
54. MEÏRMANOV, A. M., The Stefan problem with surface tension in the three-dimensional case with spherical symmetry: nonexistence of the classical solution. *European J. Appl. Math.* **5** (1994), 1–19. [Zb10812.35165](#) [MR1270785](#)
55. MEYRIES, M. & SCHNAUBELT, R., Interpolation, embeddings and traces of anisotropic fractional Sobolev spaces with temporal weights. *J. Funct. Anal.* **262** (2012), 1200–1229. [Zb11250.46022](#) [MR2863860](#)
56. MULLINS, W. W. & SEKERKA, R. F., Morphological stability of a particle growing by diffusion or heat flow. *J. Appl. Phys.* **34** (1963), 323–329.
57. MULLINS, W. W. & SEKERKA, R. F., Stability of a planar interface during solidification of a dilute binary alloy. *J. Appl. Phys.* **35** (1964), 444–451.
58. PRÜSS, J., SAAL, J. & SIMONETT, G., Existence of analytic solutions for the classical Stefan problem. *Math. Ann.* **338** (2007), 703–755. [Zb11130.35136](#) [MR2317935](#)
59. PRÜSS, J., SAAL, J. & SIMONETT, G., Stability of equilibria for the Stefan problem with surface tension. *SIAM J. Math. Anal.* **40** (2008), 675–698. [Zb11157.35502](#) [MR2438781](#)
60. PRÜSS, J. & SIMONETT, G., *Analytic solutions for the two-phase Navier-Stokes equations with surface tension and gravity*. Parabolic problems. Progr. Nonlinear Differential Equations Appl., **80**, Birkhäuser/Springer Basel AG, Basel, 2011. [Zb11247.35207](#) [MR3052594](#)
61. PRÜSS, J. & SIMONETT, G., On the two-phase Navier-Stokes equations with surface tension. *Interfaces Free Bound.* **12** (2010), 311–345. [Zb11202.35359](#) [MR2727674](#)
62. PRÜSS, J. & SIMONETT, G., On the Rayleigh–Taylor instability for the two-phase Navier-Stokes equations. *Indiana Univ. Math. J.* **59** (2010), 1853–1871. [Zb11234.35323](#) [MR2919738](#)
63. PRÜSS, J. & SIMONETT, G., On the manifold of closed hypersurfaces in \mathbb{R}^n . *Discrete Contin. Dyn. Syst.* **33** (2013), 5407–5428. [Zb11274.35434](#) [MR3297714](#)
64. PRÜSS, J. & SIMONETT, G., *Moving interfaces and quasilinear parabolic evolution equations*. Monographs in Mathematics, Birkhäuser Verlag. To appear.
65. PRÜSS, J., SIMONETT, G. & ZACHER, R., Qualitative behavior of solutions for thermodynamically consistent Stefan problems with surface tension. *Arch. Ration. Mech. Anal.* **207** (2013), 611–667. [Zb11269.80004](#) [MR3005325](#)

66. RADKEVICH, E. V., The Gibbs–Thompson correction and conditions for the existence of a classical solution of the modified Stefan problem. *Dokl. Akad. Nauk SSSR* **316** **6**, 1311–1315; translation in *Soviet Math. Dokl.* **43** (1991), 274–278. [Zb10782.35087](#) [MR1122261](#)
67. RODRIGUES, J.-F., The variational inequality approach to the one-phase Stefan problem. *Acta Appl. Math.* **8** (1987), 1–35. [Zb10653.35083](#) [MR0871691](#)
68. RODRIGUES, J.-F., The Stefan problem revisited. In *Mathematical models for phase change problems*, volume 88 of *Internat. Ser. Numer. Math.*, 129–190. Birkhäuser, Basel, 1989. [Zb10699.35265](#) [MR1038069](#)
69. RODRIGUES, J.-F., Variational methods in the Stefan problem. In *Phase transitions and hysteresis (Montecatini Terme, 1993)*, volume 1584 of *Lecture Notes in Math.*, 147–212. Springer, Berlin, 1994. [Zb10819.35154](#) [MR1321833](#)
70. RÖGER, M., Solutions for the Stefan problem with Gibbs–Thomson law by a local minimisation. *Interfaces Free Bound.* **6** (2004), 105–133. [Zb11050.35155](#) [MR2047075](#)
71. RUBINSTEIN, L. I., *The Stefan Problem*. Translations of Mathematical Monographs, Vol. 27, American Mathematical Society, Providence, R.I. 1971. [MR0351348](#)
72. SHAO, Y., Real analytic solutions to the Willmore Flow. Ninth MSU-UAB Conference on Differential Equations and Computational Simulations. *Electron. J. Diff. Eqns., Conf.* **20** (2013), 151–164. [Zb11295.35160](#) [MR3128075](#)
73. SHAO, Y., A family of parameter-dependent diffeomorphisms acting on function spaces over a Riemannian manifold and applications to geometric flows. *NoDEA Nonlinear Differential Equations Appl.* **22** (2015), 45–85. [Zb11322.58016](#) [MR3311893](#)
74. SHAO, Y. & SIMONETT, G., Continuous maximal regularity on uniformly regular Riemannian manifolds. *J. Evol. Equ.* **1** (2014), 211–248. [Zb11295.35161](#) [MR3169036](#)
75. TRIEBEL, H., *Interpolation Theory, Function Spaces, Differential Operators*. North-Holland Publishing Co., Amsterdam-New York, 1978. [Zb10387.46033](#) [MR0503903](#)
76. VISINTIN, A., Remarks on the Stefan problem with surface tension. Boundary value problems for partial differential equations and applications, 455–460, *RMA Res. Notes Appl. Math.*, 29, Masson, Paris (1993). [Zb10803.35173](#) [MR1260478](#)
77. VISINTIN, A., *Models of phase transitions*. Progress in Nonlinear Differential Equations and their Applications. 28. Birkhäuser Boston, Inc., Boston, MA, 1996. [Zb10882.35004](#) [MR1423808](#)
78. ZIEMER, W. P., Interior and boundary continuity of weak solutions of degenerate parabolic equations. *Trans. Amer. Math. Soc.* **271** (1982), 733–748. [Zb10506.35053](#) [MR0654859](#)