

All order asymptotics of hyperbolic knot invariants from non-perturbative topological recursion of A-polynomials

Gaëtan Borot and Bertrand Eynard¹

Abstract. We propose a conjecture to compute the all-order asymptotic expansion of the colored Jones polynomial of the complement of a hyperbolic knot, $J_N(q = e^{\frac{2\pi i}{N}})$ when $N \rightarrow \infty$. Our conjecture claims that the asymptotic expansion of the colored Jones polynomial is a formal wave function of an integrable system whose semiclassical spectral curve \mathcal{C} would be the $SL_2(\mathbb{C})$ character variety of the knot (the A-polynomial), and is formulated in the framework of the topological recursion. It takes as starting point the proposal made recently by Dijkgraaf, Fuji and Manabe (who kept only the perturbative part of the wave function, and found some discrepancies), but it also contains the non-perturbative parts, and solves the discrepancy problem. These non-perturbative corrections are derivatives of Theta functions associated to \mathcal{C} . For a large class of knots, this expansion is still in powers of $1/N$ due to the special properties of A-polynomials. We provide a detailed check of our proposal for the figure-eight knot and the once-punctured torus bundle L^2R . We also present a heuristic argument inspired from the case of torus knots, for which knot invariants can be computed by a matrix model.

Mathematics Subject Classification (2010). 57M27, 14-XX, 15B52, 81Txx.

Keywords. Knot invariants, all-order asymptotics, A-polynomial, A-hat polynomial, topological recursion, non-perturbative effects.

¹ We thank warmly David Cimasoni, Tudor Dimofte, Stavros Garoufalidis, Rinat Kashaev, Marcos Mariño and Don Zagier for fruitful discussions, as well as the organizers of the BIRS workshop “New recursion formulae and integrability for Calabi-Yau spaces” in Banff, where this work was initiated, and the anonymous referee for his useful comments. Gaëtan Borot thanks the participants of the Oberwolfach workshop “Low-dimensional topology and number theory,” whose comments led to improve a later version of the article, and the MPI Bonn for hospitality. Bertrand Eynard thanks the university of Geneva and Stanislav Smirnov for their hospitality, and the Centre de Recherche Mathématiques de Montréal. G.B. benefited from the support of Fonds Européen S16905 (UE7 - CONFRA) and the Swiss NSF (200021_143434). Bertrand Eynard is supported by the ANR project GranMa “Grandes Matrices Aléatoires” ANR-08-BLAN-0311-01 and by the European Science Foundation through the Misgam program.

Contents

1	Introduction	40
2	$SL_2(\mathbb{C})$ -character variety and algebraic geometry on A-spectral curves	46
3	Topological recursion	65
4	Non-perturbative topological recursion	73
5	Application to knot invariants	80
6	Examples	91
7	Heuristics imported from torus knots	111
8	Perspectives	120
A	Diagrammatic representation for the non-perturbative topological recursion	122
B	Some A-spectral curves	127
	References	132

1. Introduction

The asymptotic expansion of the colored Jones polynomial $J_N(\mathfrak{K}, q)$ of a knot \mathfrak{K} when $N \rightarrow \infty$, and more generally of invariants of 3-manifolds, has received much attention recently. The terms of such an asymptotic expansion are also invariants of 3-manifolds, which are interesting for themselves. They are generically called “perturbative invariants.” Many intriguing properties of these expansions have been observed, first in relation with hyperbolic geometry and the volume conjecture [58, 73], then concerning arithmeticity [30], modularity [62] or quantum modularity [89, Examples 4 and 5].

1.1. Solutions of the A-hat recursion relation. Garoufalidis and Lê have shown that the Jones polynomial of a knot $\mathfrak{K} \subseteq \mathbb{S}_3$, denote $J_N(\mathfrak{K}, q)$, is q -holonomic [47]: if we denote by Δ the shift $N \rightarrow N + 1$, there exists an operator $\mathfrak{A}_{\mathfrak{K}}(q^{N/2}, \Delta, q)$ which is polynomial in its three variables, so that

$$\widehat{\mathfrak{A}}_{\mathfrak{K}} \cdot J_N(\mathfrak{K}, q) = 0.$$

More generally, one may consider the (infinite dimensional) space of solutions $\mathcal{J}_{\hbar}(u)$ of the difference equation

$$\widehat{\mathfrak{A}}_{\mathfrak{K}}(e^u, e^{\hbar \partial_u}, e^{2\hbar}) \cdot \mathcal{J}_{\hbar}(u) = 0, \tag{1}$$

where we replaced formally $u = N\hbar$ and $q = e^{2\hbar}$. The Jones polynomial is by construction a solution of equation (1) where u/\hbar is restricted to a discrete set of values. One may also look for solutions of equation (1) among formal power series of the form

$$\mathcal{J}_{\hbar}(u) = \hbar^{\delta/2} \exp\left(\sum_{\chi \geq -1} \hbar^{\chi} J_{\chi}(u)\right). \quad (2)$$

The expectation values of Wilson loops in the representation of dimension N in $\mathrm{SL}_2(\mathbb{C})$ Chern–Simons theory – viewed as a perturbative quantum field theory expanded around a flat connection α with meridian holonomy u and coupling constant $\hbar = i\pi/\text{integer}$ – produce such formal series $\mathcal{J}_{\hbar}^{\mathrm{CS},(\alpha)}(u)$. For some examples of hyperbolic 3-manifolds and a choice of triangulation, it has been observed [30] that the asymptotics of Hikami integral¹ $\mathcal{J}_{\hbar}^{\mathrm{H},(\alpha)}(u)$ (which depend on a choice of integration contour $\gamma_{\alpha} \subseteq \mathbb{C}$) coincides with $\mathcal{J}_{\hbar}^{\mathrm{CS},(\alpha)}(u)$. In other words, $\mathcal{J}_{\hbar}^{\mathrm{H},(\alpha)}$ are also solutions (in those examples) of equation (1).

We would like to propose a third method which we conjecture to provide formal solutions of equation (1) for any hyperbolic 3-manifold \mathfrak{M} with 1 cusp², and relies only on algebraic geometry of the A-polynomial curve (Conjecture 5.5). The representations

$$\pi_1(\partial\mathfrak{M}) \longrightarrow \mathrm{SL}_2(\mathbb{C})$$

which extend to representations

$$\pi_1(\mathfrak{M}) \longrightarrow \mathrm{SL}_2(\mathbb{C})$$

form a set whose algebraic closure is the union of points and curves. The union of those curves can be obtained as the zero locus of a polynomial $\mathfrak{A}(m, l) \in \mathbb{Z}[m, l]$, called the A-polynomial of \mathfrak{M} . The AJ conjecture³ [45] states that

$$\lim_{\hbar \rightarrow 0} \widehat{\mathfrak{A}}_{\hbar}(m, l, e^{2\hbar}) \propto \mathfrak{A}(m, l), \quad (3)$$

where \propto means up to an (irrelevant) polynomial in m . Equation (3) has been checked in numerous examples (it holds for instance for the figure-eight knot) and

¹ The Hikami integral is a finite-dimensional constructed from a triangulation of a hyperbolic 3-manifold and an elimination procedure; however, it is not an invariant of 3-manifolds.

² We identify a 3-manifold with 1 cusp with a compact manifold whose boundary is a torus.

³ Whereas the A-polynomial is defined for any 3-manifold with 1-cusp, the colored Jones polynomial is only defined for knots and links in \mathbb{S}_3 . However, for any knot in integer homology spheres, there is an element in the Habiro ring which “defines” the analog of the colored Jones function for all values of q at roots of unity, and this element is q -holonomic, [48], so allows to define \mathfrak{A} . In this sense, the AJ conjecture make sense for any knot in an integer homology sphere.

has been proved recently for a infinite class of knots [63]. In the light of the AJ conjecture, we can summarize our work by saying that we propose an algorithm to construct formal solutions of equation (1) starting only from the classical limit of the operator $\hat{\mathfrak{A}}$. For the figure-eight knot, we have checked that it gives a correct result for the first few terms.

The final goal would be to identify our series with the genuine all-order asymptotics of invariants of 3-manifolds defined in the realm of quantum topology, like the colored Jones polynomial. This step is subtle because of wild behavior when q is a root of unity, and non trivial Stokes phenomena, as one can already observe in the case of the figure-eight knot, for which rigorous results of Murakami [71] are available (see §5.2). Though not completely predictive on the range of validity in u , the generalized volume conjecture of Gukov asserts that $J_{u/\hbar}(q = e^{\hbar})$ has an asymptotic expansion of the form of equation (2) when $\hbar = i\pi/k$ and k is a integer going to infinity, and the coefficients coincide with those of $\mathcal{J}_{\hbar}^{\text{CS},(\alpha)}(u)$ for some α . In this framework, we can also reformulate our conjecture by saying that our method retrieves the coefficients in the expansion of the colored Jones polynomial (Conjecture 5.6), and will discuss in §5.2 how this statement has to be understood.

1.2. Historical background. Let us describe briefly the origin of our proposals. Twenty years ago, Witten showed in his pioneering article [86] that expectation values of Wilson loops in a Chern–Simons theory with gauge group G on $S^3 \setminus \mathfrak{K}$ where \mathfrak{K} is a knot, compute knot invariants. Moreover, he proposed a correspondence between Chern–Simons theory on a 3-manifold \mathfrak{M} and topological string theory on $T^*\mathfrak{M}$, cf. [87], which has been developed later on [49, 65]. More recently, Bouchard, Klemm, Mariño and Pasquetti [15] suggested⁴ that amplitudes in topological string theory can be computed from the axiomatics of the “topological recursion” developed in [38]. Putting these two ideas together, Dijkgraaf, Fuji and Manabe [26, 27], proposed that the topological recursion’s wave function, applied to the $SL_2(\mathbb{C})$ -character variety of the knot, coincides with $\mathcal{J}_{\hbar}^{\text{CS}}(u)$. However, they kept only the “perturbative part” of the topological recursion’s wave function, and their conjectured formula did not match $\mathcal{J}_{\hbar}^{\text{CS}}(u)$ or $\mathcal{J}_{\hbar}^{\text{H}}(u)$ for the figure-eight knot. They could fix this mismatching problem by introducing additional *ad hoc* constants to all orders. Here, we propose a formula using the thoroughly non-perturbative wave function introduced in [35, 37], which should successfully match $\mathcal{J}_{\hbar}^{\text{CS}}(u)$ without having to introduce additional terms.

⁴ This conjecture has been proved lately [41].

1.3. Short presentation. Let us give a flavor of our construction (all terms will be defined in the body of the article). The geometric component $\mathfrak{A}(m, l) = 0$ of the A-polynomial of \mathfrak{K} has a projective smooth model which is a compact Riemann surface \mathcal{C}_0 of genus g . It is endowed with a point p_c corresponding to the complete hyperbolic metric on $\mathbb{S}_3 \setminus \mathfrak{K}$, and a neighborhood $\tilde{\mathcal{U}} \subseteq \mathcal{C}_0$ of p_c in bijection with a neighborhood $\mathcal{U} \subseteq \mathbb{C}$ of $i\pi$ which parametrizes deformations of the hyperbolic metric of $\mathbb{S}_3 \setminus \mathfrak{K}$. Let p_u the unique point in $\tilde{\mathcal{U}}$ such that $m(p_u) = e^u$. We denote by ι the involution of \mathcal{C}_0 sending (m, l) to $(1/m, 1/l)$. In particular we have $p_{-u} = \iota(p_u)$. In the following, p denotes a point of the curve, and in the comparison with the asymptotics of the colored Jones near $u = i\pi$, one wishes to specialize at $p = p_u$. Let $(\mathcal{A}, \mathcal{B})$ be a symplectic basis of homology on \mathcal{C}_0 . We construct a formal asymptotic series with leading coefficient

$$\tilde{J}_{-1}(p) = \int_o^p \ln l \, d \ln m,$$

and for ⁵ $\chi \geq 1$,

$$\tilde{J}_\chi(u) = \frac{1}{2} \sum_{\ell=1}^\chi \sum_{\substack{2h_j-2+d_j+n_j>0 \\ \sum_j 2h_j-2+d_j+n_j=\chi}} \left(\bigotimes_{j=1}^\ell \frac{\int_{\bullet} \dots \int_{\bullet} \omega_{n_j}^{h_j, (d_j)}}{(2i\pi)^{d_j} d_j! n_j!} \right) \cdot U_{\ell, \mathbf{d}, \bullet} \quad (4)$$

The notation

$$\int_{\bullet}$$

is used for

$$\int_o^p + \int_{\iota(o)}^{\iota(p)}$$

for some basepoint o , ω_n^h are the differentials forms computed by the topological recursion for the spectral curve $(\mathcal{C}_0, \ln m, \ln l)$ with a Bergman kernel normalized on \mathcal{A} -cycles, and $U_{\ell, \mathbf{d}, \bullet}$ is $(0, |\mathbf{d}|)$ tensor which is a sum of terms of the form

$$\bigotimes_{i=1}^s \frac{\nabla^{\otimes (\sum_{j \in \mathcal{J}_i} d_j)} \vartheta_{\bullet} [\mu]_v}{\vartheta_{\bullet} [\mu]_v},$$

⁵ The definition of $\tilde{J}_0(p)$ involve in principle a regularization scheme. We do not attempt to give a definition here, since it is not clear to us which scheme should be chosen to match \tilde{J}_0 with the order 1 term in the asymptotics of the colored Jones – which is expected to be related to the open Reidemeister torsion.

where J_1, \dots, J_s denote a partition of $\{1, \dots, \ell\}$ in s subsets. $\vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]$ denotes the theta function with characteristics $\mu, \nu \in \mathbb{C}^g$ associated to the matrix of periods of \mathcal{C}_0 for the chosen basis of homology. The notations $\vartheta_{\bullet}\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]$ means that we specialize its argument to

$$\mathbf{w} = \mathbf{w}_{\bullet} \equiv \int_{\bullet} d\mathbf{a} + \zeta_{\hbar},$$

where $d\mathbf{a}$ is the vector of holomorphic differentials dual to the \mathcal{A} -cycles, $\zeta_{\hbar} \in \mathbb{C}^g$ is a constant defined in equation (28), and ∇ the gradient acting on the argument \mathbf{w} . Besides, the undotted version of theta means that we specialize to $\mathbf{w} = \zeta_{\hbar}$.

Formula (4) depends strongly on a convention for the basepoint \bullet , and on characteristics $\mu, \nu \in \mathbb{C}^g$. Although we do not know the general prescription for those data needed for comparison with asymptotics of knot invariants, we will see that a natural choice arises for the examples of Section 6. If we change the homology basis $(\mathcal{A}, \mathcal{B})$, we merely obtain the same quantities for a different characteristics μ, ν . The dependence of $J_{-1}(u)$ of the choice of branches for the logarithms will be discussed later. For instance, the first coefficient is given by

$$\begin{aligned} 2 \tilde{j}_1(p) &= \int^{\bullet} \omega_1^1 + \frac{1}{6} \int_{\bullet} \int_{\bullet} \int_{\bullet} \omega_3^0 + \left(\frac{\nabla \vartheta_{\bullet}\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]}{2i\pi \vartheta_{\bullet}\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]} - \frac{\nabla \vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]}{2i\pi \vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]} \right) \oint_{\mathcal{B}} \omega_1^1 \\ &+ \frac{1}{2} \frac{\nabla \vartheta_{\bullet}\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]}{2i\pi \vartheta_{\bullet}\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]} \oint_{\mathcal{B}} \int_{\bullet} \int_{\bullet} \omega_3^0 + \frac{1}{2} \frac{\nabla^{\otimes 2} \vartheta_{\bullet}\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]}{(2i\pi)^2 \vartheta_{\bullet}\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]} \oint_{\mathcal{B}} \oint_{\mathcal{B}} \int_{\bullet} \omega_3^0 \\ &+ \frac{1}{6} \left(\frac{\nabla^{\otimes 3} \vartheta_{\bullet}\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]}{(2i\pi)^3 \vartheta_{\bullet}\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]} - \frac{\nabla^{\otimes 3} \vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]}{(2i\pi)^3 \vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]} \right) \oint_{\mathcal{B}} \oint_{\mathcal{B}} \oint_{\mathcal{B}} \omega_3^0. \end{aligned}$$

In general, ζ_{\hbar} depend on \hbar in a non trivial way, so our series is not a priori a power series in \hbar . It is however a well-defined formal asymptotic series: as we will see, $j_{\chi}(u)$ is actually a function of \hbar , which does not have a power series expansion in powers of \hbar . But A-polynomials of 3-manifolds are very special polynomials: for K-theoretical reasons, ζ_{\hbar} is constant along sequences $\hbar = i\pi/k$ where k ranges over the integers. Hence, $j_{\chi}(u)$ specialized to such subsequences of \hbar , is indeed a function of u only. We also point out that another huge simplification occurs for a certain class of knots (containing the figure-eight knot). Let us denote ι_* , the linear involution induced by ι on the homology of \mathcal{C}_0 . When $\iota_* = -\text{id}$, we have actually $\zeta_{\hbar} = 0$. Thus, our series is always a power series in \hbar (without restriction) in this case.

The proposal of Dijkgraaf, Fuji and Manabe [27] is tantamount to setting

$$U_{\ell, \mathbf{d}, \bullet} \equiv \delta_{\ell, 1} \delta_{\mathbf{d}, \mathbf{0}},$$

and thus miss the theta functions. For the figure-eight knot, as we indicated, $U_{\ell, \mathbf{d}, \bullet}$ contributes as constants in \tilde{j}_χ for $\chi \geq 1$, and their value explain the renormalizations observed by these authors. To summarize, they are due the fact that the geometric component of the character variety is not simply connected.

The leading coefficient $\tilde{j}_{-1}(p_u)$ is known to be related to the complexified volume of M for a family of incomplete hyperbolic metrics parametrized by u . Within our conjecture, the other coefficients $\tilde{j}_\chi(u)$ also acquires a geometric meaning, as primitives of certain meromorphic 1-forms on the $\mathrm{SL}_2(\mathbb{C})$ character variety. The computation of the coefficients with our method is less efficient than making an ansatz like equation (2), plugging into the A-hat recursion relation and solving for the coefficients [30, 91]. However, it underlines the relevance of the geometry of the character variety itself for asymptotics of knot invariants, and also suggests unexpected links between knot theory and other topics in mathematical physics (Virasoro constraints, integrable systems, intersection theory on the moduli space, non-perturbative effects, etc.), via the topological recursion. It also provides a natural framework to discuss the arithmetic properties of perturbative knot invariants, at least when $\iota_* = -\mathrm{id}$.

1.4. Outline. We first review the notions of geometry of the character variety needed to present our construction (Section 2), and the axiomatics of the topological recursion with the definition of the correlators, the partition function and the kernels (Section 3 and 4). We state precisely our conjecture concerning the asymptotic expansion of the Jones polynomial in Section 5, and check it to first orders for the figure-eight knot and the manifold L^2R . Our intuition comes from two other aspects of the topological recursion, namely its relation to integrable systems [13] and to matrix integrals [3, 21, 34]. We give some heuristic motivations in Section 7, by examining the relation of our approach with computation of torus knots invariants from the topological recursion presented in [17]. This section is however independent of the remaining of the text. In Appendix A, we propose a diagrammatic way to write J_χ , which may help reading the formulae, but requires more notations.

2. $\mathrm{SL}_2(\mathbb{C})$ -character variety and algebraic geometry on A-spectral curves

We review standard facts on the $\mathrm{SL}_2(\mathbb{C})$ -character variety of 3-manifolds, especially of hyperbolic 3-manifolds with 1-cusp. A work of reference is [23], where most of the facts presented here are rigorously stated and proved. In Sections 2.8–2.11, we focus on the (irreducible components of the) character variety seen as a compact algebraic curves. In order to prepare the presentation of the topological recursion, we describe some algebraic geometry of the character variety, with the notion of branchpoints, symplectic basis of cycles, theta functions, Bergman kernel, etc.

2.1. A-polynomial and spectral curve. Let \mathfrak{M} be a 3-manifold with one cusp. If R a $\mathrm{SL}_2(\mathbb{C})$ representation of $\pi_1(\mathfrak{M})$ and (m, l) a basis of $\mathbb{Z}^2 \simeq \pi_1(\partial\mathfrak{M}) \subseteq \pi_1(\mathfrak{M})$, $R[m]$ and $R[l]$ can be written in Jordan form

$$R[m] = \begin{pmatrix} m & \star \\ 0 & m^{-1} \end{pmatrix}, \quad R[l] = \begin{pmatrix} l & \star \\ 0 & l^{-1} \end{pmatrix},$$

up to a global conjugation. When \mathfrak{M} is a knot complement in \mathbb{S}_3 , the choice of l and m is canonically the longitude and the meridian around the knot. In general, we continue to call the (arbitrarily chosen) (m, l) *meridian* and *longitude*.

The locus of possible eigenvalues $(m, l) \in \mathbb{C}^\times \times \mathbb{C}^\times$ has been studied in detail in [23]. It is the union of points and curves. In particular, the union of the 1-dimensional components is non empty and its algebraic closure coincides with the zero locus of a polynomial with integer coefficients: $\mathfrak{A}(m, l) = 0$. The latter is uniquely defined up to normalization and is called the A-polynomial of \mathfrak{M} . The A-polynomial is topological invariant of 3-manifolds endowed with a choice of basis of $\pi_1(\partial\mathfrak{M})$, and it contains a lot of geometric information about \mathfrak{M} .

The A-polynomial has many properties, and we shall highlight those we need along the way. The first one is that, since (m, l) and $(1/m, 1/l)$ describe the same representation up to conjugation, the A-polynomial is quasi-reciprocal: there exists integers a, b and a sign ε such that

$$\varepsilon m^a l^b \mathfrak{A}(1/m, 1/l) = \mathfrak{A}(l, m).$$

To simplify, we assume throughout the paper that \mathfrak{M} is a knot complement in a homology sphere, although most of the ideas can be extended to arbitrary 3-manifolds. In particular, this assumption implies [23] that the A-polynomial is actually even in m . We take this property into account by defining $x = m^2$ and $y = l$. The 1-form

$$\phi = \ln l \, d \ln m \tag{5}$$

is related to a notion of volume and will play an important role.

The A-polynomial might not be irreducible. We denote generically $A(m, l)$ an irreducible factor, its zero locus in $(\mathbb{C}^\times)^2$ is denoted \mathcal{C} and called *component*. There always abelian representations with $l = 1$, so $(l - 1)$ is always a factor, it defines the abelian component. The remaining components are called non-abelian, and for any hyperbolic knot, there exist at least one non-abelian component [23, Proposition 2.6]. From now on, we focus on non-abelian components. \mathcal{C} is in general a singular curve, i.e.

$$\mathcal{Z} = \{(m, l) \in (\mathbb{C}^\times)^2 : \partial_m A(m, l) = 0 \text{ and } \partial_l A(m, l) = 0\} \neq \emptyset.$$

However, after a birational transformation $\mathcal{Q}: (m, l) \mapsto (m', l')$, we can obtain a smooth projective model \mathcal{C}_0 of \mathcal{C} . It provides of a compact Riemann surface \mathcal{C}_0 , a finite subset $\mathcal{Z}_0 \subseteq \mathcal{C}_0$, and an isomorphism of Riemann surfaces $\mathcal{C}_0 \setminus \mathcal{Z}_0 \rightarrow \mathcal{C} \setminus \mathcal{Z}$. x and y can then be seen as meromorphic functions on \mathcal{C}_0 .

We refer to the triple (\mathcal{C}_0, x, y) as the *spectral curve* of the component we considered. The examples treated in detail in Section 6 illustrate the method to arrive unambiguously to the spectral curve. Properties of those spectral curves for many example of knots up to 10 crossings are given in Appendix B. In all those cases, we have found that (1) the set of singular points of any component \mathcal{C} is $\mathcal{Z} = \mathcal{C} \cap \{(m, l) = (1, \pm 1)\}$; (2) the irreducible factors of the A-polynomial all have $A \in \mathbb{Z}[m, l]$. We do not know if (1) is a general fact, but there exist examples for which (2) is not true.

2.2. Properties of the A-polynomial. As a polynomial, the A-polynomial of a 3-manifold is very special: it satisfies the Boutroux condition and a quantization condition. These two properties hold for any 3-manifold (and any component of its A-polynomial). They come from a property in K-theory, which is proved in [23, p. 59], and were clarified in [64] Before coming to the K-theoretic point of view, let us describe these properties.

Boutroux condition. We have a *Boutroux property*: for any closed cycle $\Gamma \subseteq \mathcal{C}_0 \setminus \mathcal{Z}_0$,

$$\oint_{\Gamma} \text{Im } \phi = 0.$$

For hyperbolic 3-manifolds, this is related to the existence of a function giving the hyperbolic volume. The Boutroux condition has been underlined in [57] for plane curves of the form $\text{Pol}(x, y) = 0$ endowed with the 1-form $\phi = y dx$. It appears naturally in the asymptotic study of matrix integrals, (bi)orthogonal polynomials and Painlevé transcendents, and is related to a choice of steepest descent

integration contours to apply a saddle-point analysis [8, 9]. Actually, Hikami observed [54] that the A-polynomial can be obtained as the saddle-point condition in integrals of product of quantum dilogarithm constructed from triangulations and related to knot invariants. So, it is not surprising to meet a Boutroux property here.

Quantization condition. The real periods of ϕ are quantized: there exists a positive integer ς such that, for any closed cycle $\Gamma \subseteq \mathcal{C}_0 \setminus \mathcal{Z}_0$ with base point p_0 such that $\text{Ln } m(p_0) = 0, i\pi$,

$$\oint_{\Gamma} \text{Re } \phi \in \frac{2\pi^2}{\varsigma} \cdot \mathbb{Z}. \quad (6)$$

This condition has first been pointed out by Gukov [51] in his formulation of the generalized volume conjecture, as a necessary condition for the $\text{SL}_2(\mathbb{C})$ -Chern–Simons theory to be quantizable. In our framework also, equation (6) implies the existence of a expansion in powers of \hbar for certain quantities. We explain the mechanism in Section 2.10.

2.3. Triangulations and hyperbolic structures on 3-manifolds. By definition, an oriented 3-manifold \mathfrak{M} is *hyperbolic* if it can be endowed with a smooth, complete hyperbolic metrics with finite volume. There exists an infinite number of hyperbolic knots, i.e. knots whose complement in the ambient space is hyperbolic. The A-polynomial of an hyperbolic 3-manifold \mathfrak{M} is closely related to the deformations of the hyperbolic structure on \mathfrak{M} . We review this relation and follow the foundational work of W. Thurston [84] and Neumann and Zagier [78].

Mostow rigidity theorem then states that the metrics in the definition above is unique. \mathfrak{M} is either compact, or has c cusps. We assume that \mathfrak{M} can be decomposed in a set of ideal tetrahedra glued face to face. Ideal means that all vertices of the triangulations are on the cusps. It can then identified with the interior of an oriented, bordered compact 3-manifold, whose boundary consists in c tori. So, its Euler characteristics is 0, and counting reveals that the number of tetrahedra $N_{\mathcal{T}}$ equals the number of distinct edges in the triangulation. And, by construction, the number of vertices is c , the number of cusps.

In an ideal tetrahedron \mathfrak{T} , let us choose an oriented edge e pointing towards a vertex \circ . If we intersect \mathfrak{T} by a horosphere centered at \circ , we obtain a triangle whose sum of angles is π . It is thus similar to some euclidean triangle $T(z_e)$ with vertices 0, 1 and z_e . We choose a representative for which the image of z_e in the tetrahedron belongs to e , and such that $\text{Im } z_e > 0$. z_e is called a *shape parameter*, and we may define in a unique way logarithmic shape parameters $\zeta_e = \text{Ln } z_e$, which are more natural to express geometric conditions.

For a given vertex with incident edges e_1, e_2, e_3 in cyclic order, the shape parameters are $z_{e_1}, 1 - z_{e_1}^{-1}$ and $(1 - z_{e_1})^{-1}$. As a manifestation of the angle sum condition around a Euclidean condition, we have

$$\zeta_{e_1} + \zeta_{e_2} + \zeta_{e_3} = i\pi$$

and in particular, the product of the shape parameters around a vertex is always -1 . The shape parameter of an edge in opposite orientation is $z_{-e} = -z_e^{-1}$. Opposite edges in the tetrahedron have the same shape parameter. Thus, the triangulation depends a priori on N_T shape parameters.

An oriented edge in the ideal triangulation of \mathfrak{M} correspond to the identification of a collection of distinct oriented edges $(e_j)_j$ of the tetrahedra. Since \mathfrak{M} is smooth along edges, we have N_T gluing conditions, which are in general redundant:

$$\sum_j \zeta_{e_j} = 2i\pi, \quad (7)$$

The data of N_T shape parameters satisfying equation (7) fixes a hyperbolic metrics with finite volume for \mathfrak{M} , which in general becomes singular when approaching the vertices of the triangulation. The completion \mathfrak{M}_z of \mathfrak{M} with respect to this metrics is a topological space, which may differs from \mathfrak{M} by addition of set of points at the cusps. \mathfrak{M}_z happens to be a genuine hyperbolic manifold if and only if for any vertex $\bullet \in \{1, \dots, c\}$ in the triangulation

$$\alpha_\bullet \equiv \sum_{e_\bullet} \zeta_{e_\bullet} = 2i\pi. \quad (8)$$

where $\{e_\bullet\}$ is the set of oriented edges of tetrahedra whose image in the triangulation points towards \bullet . It is shown in [78, 84], that the set of solutions of equations (7)–(8) is discrete. Moreover, in the neighborhood of a solution (i.e. of a manifold \mathfrak{M}_{z_0}), the cusp anomalies α_\bullet are local coordinates for the set of solutions of equation (7).

In a triangulated hyperbolic 3-manifold \mathfrak{M}_{z_0} , there is a natural $\mathrm{PSL}_2(\mathbb{C})$ representation of the fundamental group of \mathfrak{M}_{z_0} , namely the holonomy representation. If we assume only $c = 1$ cusp, let us choose two closed paths $\gamma_m, \gamma_l \subseteq \mathfrak{M}$ which are representatives of a meridian and a longitude. Then, the holonomy eigenvalues (m, l) arise such that m^2 (resp. l^2) is the product of the shape parameters of the oriented edges crossed by γ_m (resp. γ_l). The holonomy representation can be lifted [23, p. 71] to a $\mathrm{SL}_2(\mathbb{C})$ representation. The lift is not unique because of a choice of square root, but we always have $l_c = -1$, cf. [19], and we can choose $m_c = -1$. Now, if we waive the completeness requirement for hyperbolic

metrics on \mathfrak{M} , we have a 1-parameter deformation of hyperbolic metrics in the neighborhood of \mathfrak{M}_{z_0} , and (m, l) can be defined in a unique way as continuous (in fact, holomorphic) functions along the deformation. Also, the locus \mathcal{U} of (m, l) achieved by the deformation is included in some 1-dimensional component of the A-polynomial curve: the⁶ geometric component $\mathcal{C}^{\text{geom}}$. $(m_c, l_c) = (-1, 1)$ is very often a singular point in $\mathcal{C}^{\text{geom}}$, but e.g. the germ of l as a function of m is well defined along the deformation. So, there is a unique lift p_c of this point to the smooth model $\mathcal{C}_0^{\text{geom}}$. Logarithmic variables on $\mathcal{C}_0^{\text{geom}}$ are also very useful. We define (u, v) as holomorphic functions on $\mathcal{C}_0^{\text{geom}} \setminus \{\text{singularities and branch cuts}\}$ assuming the initial value $(i\pi, i\pi)$ at p_c , and such that $(m, l) = (e^u, e^v)$. The branches of the logarithm in equation (5) can be unambiguously chosen as

$$\phi = v du.$$

There is a map from the variety \mathfrak{Z} of shape parameters achieved by deformed hyperbolic structures on \mathfrak{M} , to $\mathcal{C}^{\text{geom}}$, which is 1-to-1 at least in a neighborhood $\tilde{\mathcal{U}} \subseteq \mathcal{U}$ of $(m_c, l_c) = (-1, -1)$. If $p \in \tilde{\mathcal{U}}$, we denote $z(p) \in \mathfrak{Z}$ the corresponding point. For a countable set of points $z \in \mathfrak{Z}$ (corresponding to a countable set of points in $\mathcal{C}^{\text{geom}}$), the space \mathfrak{M}_z is not as wild as in the generic case. Indeed, if there exists coprime integers q, q' such that $qu + q'v = i\pi$, \mathfrak{M}_z is a manifold that is just obtained from \mathfrak{M} by performing a (q, q') Dehn filling of the cusp.

2.4. Volume and Chern–Simons invariant. By a standard computation, the volume of an ideal tetrahedron with shape parameter z , endowed with its complete hyperbolic metrics, is given by the Bloch-Wigner dilogarithm $D(z)$, which is a continuous function defined on $\mathbb{C} \setminus \{0, 1\}$ as

$$D(z) = \text{Im Li}_2(z) + \arg(1 - z) \ln |z|. \quad (9)$$

The hyperbolic volume of \mathfrak{M}_z is thus

$$\text{Vol}(\mathfrak{M}_z) = \sum_e D(z_e),$$

and the functional relations satisfied by the dilogarithm ensure that it does not depend on the triangulation.

⁶The *geometric component* is usually defined as the component of the $\text{PSL}_2(\mathbb{C})$ character variety containing the (unique) discrete faithful representation of $\pi_1(\mathfrak{M})$. When projecting to representations of $\pi_1(\partial\mathfrak{M})$ and lifting to $\text{SL}_2(\mathbb{C})$ representations, we may obtain 1 or 2 irreducible factors of the A-polynomial. Here, we chose the one containing $(m_c, l_c) = (-1, -1)$, and we called it geometric component.

Another invariant of hyperbolic 3-manifolds is the *Chern–Simons invariant*. For compact manifolds, it was introduced in [22] and belongs to $\mathbb{R}/(2\pi^2\mathbb{Z})$. For manifolds \mathfrak{M}_z obtained by Dehn surgery on a hyperbolic manifold \mathfrak{M} , this definition was generalized in [68] and the invariant belongs to $\mathbb{R}/\pi^2\mathbb{Z}$. Its definition in terms of a triangulation involves $\sum_e \operatorname{Re} \operatorname{Li}_2(z_e)$ plus a tricky part described in [74]. Note that, a priori, the Chern–Simons invariant $\operatorname{CS}(\mathfrak{M}_z)$ only makes sense when \mathfrak{M}_z is a (smooth) manifold.

From a differential geometry standpoint, [78] and independently [88, Theorem 2] proved that both invariants can be extracted from the function on $\mathcal{C}^{\text{geom}}$,

$$\Phi(p) \equiv \int_o^p \phi = \int_o^p v \, du,$$

and

$$d\Phi(p) = \frac{i}{2} d(\operatorname{Vol}_a + i\operatorname{CS}_a).$$

For a point $p \in \tilde{\mathcal{U}} \subseteq \mathcal{C}^{\text{geom}}$, the volume of $\mathfrak{M}_{z(p)}$ is directly related to the imaginary part of $\Phi(p)$:

$$\operatorname{Vol}(\mathfrak{M}_{z(p)}) - \operatorname{Vol}(\mathfrak{M}_{z(p_c)}) = \operatorname{Vol}_a(p) - 2 \operatorname{Re} u \operatorname{Im} v,$$

and thanks to the Boutroux condition, it does not depend of the path from p_c to p . If we assume that $\mathfrak{M}_{z(p)}$ is the manifold obtained by (q, q') Dehn filling, the real part is related to the Chern–Simons invariant of $\mathfrak{M}_{z(p)}$. The formula involves the conjugate integers (r, r') such that $qr' - q'r = 1$:

$$\operatorname{CS}(\mathfrak{M}_{z(p)}) - \operatorname{CS}(\mathfrak{M}_{z(p_0)}) = \operatorname{CS}_a(p) + \pi(r \operatorname{Im} u + r' \operatorname{Im} v),$$

and thanks to the quantization condition, it does not depend modulo $2\pi^2\mathbb{Z}/\zeta$ of the choice of path from p_c to p . In this article, we call Vol_a the *analytic volume*, and CS_a the *analytic Chern–Simons term*.

Remark 2.1. Even if \mathfrak{M} is not hyperbolic, the primitive of the 1-form $\ln l \, d \ln m$ (i.e. a function Φ so that $d\Phi = \ln l \, d \ln m$) defined over (one of the component of) the A-polynomial curve varieties defines a notion of complexified volume, whose imaginary part is closely related to the notion of volume of a representation.

It is enlightening to understand the volume, the Chern–Simons invariant and the properties raised in §2.2 from the point of view of K-theory. This is the matter of the next two paragraphs.

2.5. Bloch group and hyperbolic geometry. Let \mathbb{K} be a number field or a function field. To fix notations, \mathbb{K}^\times is the multiplicative group of invertible elements of \mathbb{K} , and \mathbb{K}^+ is just \mathbb{K} considered as an additive group. For an abelian group G , the exterior product $\bigwedge_{\mathbb{Z}}^2 G$ is the \mathbb{Z} -module generated by the antisymmetric elements $x \wedge y$ for $x, y \in G$, modulo the relations of compatibility with the group law $(n \cdot x) \wedge y = n(x \wedge y)$. When S is a set, $\mathbb{Z} \cdot S$ is the free \mathbb{Z} -module with basis the elements of S .

The *pre-Bloch group* $P(\mathbb{K})$, cf. [11], is the quotient of $\mathbb{Z} \cdot (\mathbb{K}^\times \setminus \{1\})$ by the relations $[z] + [1 - z] = 0$ and $[z] + [1/z] = 0$ for any $z \in \mathbb{K}^\times \setminus \{0\}$, and the five term relations

$$[z] + [z'] + [1 - zz'] + \left[\frac{1 - z}{1 - zz'} \right] + \left[\frac{1 - z'}{1 - zz'} \right] = 0$$

for any $z, z' \in \mathbb{K}^\times \setminus \{1\}$ such that $zz' \neq 1$. Those combinations appear precisely in the functional relations of the function $D(z)$ of equation (9). Indeed, D induces a well-defined function $D: P(\mathbb{K}) \rightarrow \mathbb{R}$ if we interpret

$$D\left(\xi = \sum_j [z_j]\right) = \sum_j D(z_j).$$

For a hyperbolic manifold \mathfrak{M} with a triangulation, a point p in $\mathcal{C}^{\text{geom}}$ determines shape parameters $\mathbf{z}(p) = (z_e(p))_e$ for the triangulation. We can apply the above construction to a field \mathbb{K} where the functions \mathbf{z} live. It is in general an extension of the field $\mathbb{C}(\mathcal{C}^{\text{geom}})$, of finite degree that we denote d . The element

$$\xi_{\mathbf{z}} = \sum_e [z_e] \in P(\mathbb{K}) \tag{10}$$

is actually independent of the triangulation. Also, the volume is a well-defined function on $\mathcal{C}^{\text{geom}}$, given by $D(\xi_{\mathbf{z}})$.

Up to now, the introduction of the pre-Bloch group has served merely as a rephrasing of §2.4. Neumann and Yang [77] took a step further to reach the Chern–Simons invariant. We introduce the Rogers dilogarithm, which is a multivalued holomorphic function on $\mathbb{C}^\times \setminus \{1\}$:

$$R(z) = \text{Li}_2(z) + \frac{\ln z \ln(1 - z)}{2}.$$

Some computations shows that the diagram below⁷ is well-defined and commutative:

$$\begin{array}{ccc}
 \mathbb{B}(\mathbb{C}) & \xrightarrow{\subseteq} & \mathbb{P}(\mathbb{C}) \xrightarrow{\mu} \bigwedge_{\mathbb{Z}}^2 \mathbb{C}^\times \\
 \hat{\rho} \downarrow & & \downarrow \rho \nearrow e \\
 \mathbb{C}/\mathbb{Q} & \xleftarrow{\dots} & \bigwedge_{\mathbb{Z}}^2 \mathbb{C}^+
 \end{array}
 \quad
 \begin{array}{l}
 \mu([z]) = 2z \wedge (1-z), \\
 \rho([z]) = \frac{\ln z}{2i\pi} \wedge \frac{\ln(1-z)}{2i\pi} + 1 \wedge \frac{R(z)}{2\pi^2}, \\
 e(\zeta \wedge \zeta') = 2e^{2i\pi\zeta} \wedge e^{2i\pi\zeta'}
 \end{array}$$

The *Bloch group* of \mathbb{C} by definition

$$\mathbb{B}(\mathbb{C}) = \ker \mu,$$

and we have $\rho(\mathbb{B}(\mathbb{C})) \subseteq \text{Ker } e$. As a matter of fact,

$$\zeta \mapsto 1 \wedge \zeta \in \bigwedge_{\mathbb{Z}}^2 \mathbb{C}^+$$

induces an isomorphism between \mathbb{C}/\mathbb{Q} and $\text{Ker } e$. Thus, there is a map $\hat{\rho}$ from the Bloch group to \mathbb{C}/\mathbb{Q} . Coming back to hyperbolic geometry: since two edges carry the same shape parameter in each tetrahedron, the element ξ_z defined in equation (10) actually sits in $\mathbb{B}(\mathbb{C}) \subseteq \mathbb{P}(\mathbb{C})$. When \mathfrak{M}_z is a manifold, it was proved in [76] that $\hat{\rho}$ gives the irrational part of the Chern–Simons invariant:

$$\text{CS}(\mathfrak{M}_z) = -2\pi^2 \text{Re } \hat{\rho}(\xi_z) \pmod{8\pi^2\mathbb{Q}}.$$

2.6. K-theory viewpoint. We now review the interpretation of the Boutroux and quantization condition in the context of K-theory, and its relations to hyperbolic geometry.

Symbols. After a classical result of Matsumoto [69, §11], the second K-group $\mathbb{K}_2(\mathbb{K})$ of a field \mathbb{K} is isomorphic to $\bigwedge_{\mathbb{Z}}^2 \mathbb{K}^\times$ modulo the relations $z \wedge (1-z) = 0$. In other words, $\mathbb{K}_2(\mathbb{K}) = \text{coker } \mu/2$, where μ is the morphism introduced in §2.5. The elements of $\mathbb{K}_2(\mathbb{K})$ are usually called *symbols*, and denoted $\{z_1, z_2\}$. When \mathcal{C} is a component of an A-polynomial of a 3-manifold, a theorem [23, p. 61] shows the existence of a integer ζ , that we choose minimal, such that

$$2\zeta \cdot \{m, l\} = 0 \in \mathbb{K}_2(\mathbb{C}(\mathcal{C})). \tag{11}$$

⁷ The factor of 2 is convenient for applications to knot theory in the homology spheres.

Regulators. If $z_1, z_2 \in \mathbb{C}(\mathcal{C})$, and \mathcal{P} denotes the set of zeroes and poles of z_1, z_2 , the *regulator map* is defined as

$$r[z_1, z_2]: H_1(\mathcal{C} \setminus \mathcal{Z}, \mathbb{P}) \longrightarrow \mathbb{C}^\times$$

$$\gamma \longmapsto \exp \left[\frac{1}{2i\pi} \left(\oint_\gamma \ln z_1 d \ln z_2 - \ln z_1(o) \oint_\gamma d \ln z_2 \right) \right].$$

o is a basepoint in γ and given a choice of branch of $\ln z_1$ and $\ln z_2$ at o , the logarithms are analytically continued starting from o along γ . One can show that this definition does not depend on o , on the initial choice of branches for the logarithm, and of the representative z_1, z_2 of the symbol $\{z_1, z_2\}$. Hence, there exists a map

$$r: \mathbf{K}_2(\mathbb{K}) \longrightarrow \varinjlim_{\mathcal{P} \text{ finite}} \text{Hom}(H_1(\mathcal{C} \setminus \mathcal{Z}), \mathbb{C}^\times).$$

If $\{z_1, z_2\}$ is 2ζ -torsion (as in equation (11)), we see that $r[z_1, z_2](\gamma)$ is a $2\zeta^{\text{th}}$ -root of unity for all closed cycles γ . We deduce that, for any closed cycle γ with basepoint o such that $\ln z_1(o) = 0, i\pi$ and the integral is well-defined,

$$\oint_\gamma \ln z_1 d \ln z_2 \in \frac{2\pi^2}{\zeta} \mathbb{Z}. \quad (12)$$

This line of reasoning has been written explicitly in [64]. This can be applied to $\{m, l\}$ for a component of an A-polynomial, and justifies the Boutroux and the quantization condition of §2.2.

Tame symbol and Boutroux condition. Given an algebraic curve \mathcal{C} with two functions z_1, z_2 defined on it, it might not be easy to check if $\{z_1, z_2\}$ is torsion. However, it is elementary to check if there is a local obstruction to being torsion, i.e. if equation (12) holds for all contractible, closed cycles γ in \mathcal{C} . We focus in this paragraph only on the imaginary part of equation (12), which gives rise to the Boutroux condition, and discuss its relation with the tame condition. The reason is that the Boutroux condition already has interesting consequences for the Baker–Akhiezer kernel (§2.10) and thus the construction of Section 4.

This is formalized as follows. To any $z_1, z_2 \in \mathbb{K}^\times$, we can associate the *regulator form*, which is the 1-form

$$\begin{aligned} \eta[z_1, z_2] &= \ln |z_1| (d \arg z_2) - \ln |z_2| (d \arg z_1) \\ &= \text{Im}(\ln z_1 d \ln z_2) - d(\arg z_1 \ln |z_2|). \end{aligned}$$

For any point $p \in \mathcal{C}_0$, let

$$T_p: \mathbf{K}_2(\mathbb{K}) \longrightarrow \mathbb{C}^\times$$

be the map defined by

$$\begin{aligned} T_p(\{z_1, z_2\}) &= (-1)^{\text{ord}_p z_1 \cdot \text{ord}_p z_2} z_1(p)^{\text{ord}_p z_2} z_2(p)^{-\text{ord}_p z_1} \\ &= \exp(i\pi \left(\underset{p}{\text{Res}} \, d \ln z_1 \right) \left(\underset{p}{\text{Res}} \, d \ln z_2 \right) + \underset{p}{\text{Res}} [(d \ln z_2) \ln z_1 - (d \ln z_1) \ln z_2]). \end{aligned} \quad (13)$$

This expression is indeed independent of the representative of $\{z_1, z_2\}$. It is also independent of the branches of the logarithms and of the basepoint to define the integral over a small circle around p . A computation shows

$$\underset{p}{\text{Res}} \, \eta[z_1, z_2] = -i \ln |T_p(\{z_1, z_2\})|,$$

so T_p is closely related to the regulator map $r[z_1, z_2]$ evaluated on a small circle around p . For a given curve, z_1 and z_2 only have a finite number of zeroes and poles, so $T_p(\{z_1, z_2\}) = 1$ except at a finite number of points. Note that the Riemann bilinear identity applied to the meromorphic 1-forms $\frac{dz_1}{z_1}$ and $\frac{dz_2}{z_2}$ implies $\prod_{p \in \mathcal{C}_0} T_p(\{z_1, z_2\}) = 1$. We say that $\{z_1, z_2\}$ is a *weakly tame* symbol if $|T_p(\{z_1, z_2\})| = 1$ for all p , i.e. we define the subgroup

$$\mathbf{K}_2^{\text{w-tame}}(\mathcal{C}) = \bigcap_{p \in \mathcal{C}(\mathbb{C})} \ker |T_p|.$$

It is very easy to check if an element of $\mathbf{K}_2(\mathcal{C}(\mathbb{C}))$ is weakly tame or not, given equation (13), and this provides a local obstruction for the Boutroux condition, and a fortiori for being torsion. Moreover, if there exists an integer ζ_0 such that $T_p(\{z_1, z_2\})$ is a $2\zeta_0^{\text{th}}$ -root of unity for all $p \in \mathcal{C}$, and if $\{z_1, z_2\}$ is torsion, ζ_0 must divide the order of torsion. The *tame group* itself is defined as

$$\mathbf{K}_2^{\text{tame}}(\mathcal{C}) = \bigcap_{p \in \mathcal{C}(\mathbb{C})} \ker T_p.$$

$\eta[z_1, z_2]$ is always closed, since $\eta[z_1, z_2] = \text{Im}(d \ln z_1 \wedge d \ln z_2) = 0$. It is in general not exact, but $\eta[z, 1 - z] = dD(z)$. So, we can illustrate this discussion in the context of hyperbolic 3-manifolds. The shape parameters $(z_e)_e$ sit in an extension \mathbb{K} of $\mathbb{C}(\mathcal{C}^{\text{geom}})$ of some degree ζ , we have at our disposal the element $\xi \in \mathbf{B}(\mathbb{K})$ (see equation (10)) and the symbol $\{m^2, l\} = \sum_e \{z_e, 1 - z_e\}$ is by construction zero in $\mathbf{K}_2(\mathbb{K})$. By coming back to $\mathbf{K}_2(\mathbb{C}(\mathcal{C}^{\text{geom}}))$, one only obtains that $\zeta_0 \cdot \{m^2, l\}$, so $\{m, l\}$ is 2ζ -torsion. Hence $\{m, l\}$ is weakly tame in a trivial way.

2.7. Arithmetics and cusp field. We now come to aspects of the A-polynomial which are relevant to the arithmeticity properties of the perturbative invariants of 3-manifolds. We aim at preparing for a clarification of the arithmetic nature of the invariants defined from the topological recursion in Section 3.1, especially when applied to A-polynomials. Unless precised otherwise, we work in the remaining of this section with any of the irreducible factor of the A-polynomial which is not of the form $(lm^a \pm 1)$, and the corresponding spectral curve (\mathcal{C}_0, u, v) is endowed with a marked point $p_c \in \mathcal{C}_0$ such that $m^2(p_c) = l^2(p_c) = 1$.

We already stated that $m(p_c), l(p_c)$ is a singular point for A . More precisely,

$$A(m, l) \propto (l - l(p_c))^a C\left(\frac{l - l(p_c)}{m - m(p_c)}\right)$$

in the neighborhood of this singularity. If A was a polynomial with integer coefficients, C is also a polynomial with integer coefficients, called the *cusp polynomial*. Although it contains less information than the A-polynomial, it retains some geometric significance and is closely related to the C-polynomial studied by Zhang [90]. We also introduce the *cusp field* \mathbb{F} , which is the splitting field of the cusp polynomial. In particular, at the vicinity of p_c in \mathcal{C}_0 , we have $(l + 1) \sim \gamma(m + 1)$ where γ is a root of C , thus an element of the cusp field.

There are several notions of fields associated to a hyperbolic 3-manifold \mathfrak{M} . In presence of an ideal triangulation of \mathfrak{M} , the *tetrahedron field* is the field generated by the shape parameters of the tetrahedra. From another point of view, \mathfrak{M} can be realized as quotients \mathbb{H}_3/Γ where Γ is a discrete subgroup of $\mathrm{PSL}_2(\mathbb{C})$ of finite covolume. One can define the *invariant trace field*, which is the field generated by the trace of squares of elements of Γ . It is clear that

$$\text{cusp field} \subseteq \text{invariant trace field} \subseteq \text{tetrahedron field}.$$

For hyperbolic knot complements, there are examples where the cusp field is strictly smaller than the tetrahedron field [75], but the invariant trace field coincide with the tetrahedron field after a more general theorem of Neumann and Reid [75, Theorem 2.4]. The numbers produced from the topological recursion will naturally live in the cusp field \mathbb{F} .

Remark 2.2. When \mathcal{C} is the geometric component of an A-polynomial of a triangulated 3-manifold, [20] ensures that the shape parameters are rational functions of l and m . Hence, the order of torsion of $\{m, l\}$ is 2.

2.8. Definition of A-spectral curves. Since we will often use this setting, we give the name *A-spectral curve* (over a field \mathbb{K}) to the object composed by the following data.

- A curve \mathcal{C} defined by an equation of the form $\text{Pol}(m, l) = 0$ (with coefficients in \mathbb{K}), such that $\{m, l\}$ is 2ζ -torsion in $\mathbb{K}_2(\mathcal{C}(\mathbb{K}))$ for some minimal integer ζ (We assume that $\text{Pol}(m, l)$ is irreducible and not proportional to $lm^b \pm 1$ for some integer b).
- A compact Riemann surface \mathcal{C}_0 which is the projective smooth model for \mathcal{C} , and a marked point $p_c \in \mathcal{C}_0$ such that $l(p_c)^2 = m(p_c)^2 = 1$.
- Two holomorphic functions $u = \ln m$ and $v = \ln l$ on

$$\mathcal{C}_0 \setminus \{\text{singularities and branch cuts}\},$$

and the differential form $\phi = v du$.

- We add the technical assumption that the zeroes of v are simple.

One may wonder if all A-spectral curves over \mathbb{K} arise as components of the A-polynomial of some 3-manifold. The answer does not seem to be known. $\mathbb{K}_2^{\text{tame}}(\mathcal{C})$ (and a fortiori $\mathbb{K}_2^{\text{w.tame}}(\mathcal{C})$) for a compact Riemann surface \mathcal{C}_0 of genus $g \geq 1$ defined over \mathbb{Q} is in general not trivial. Part of a conjecture of Beilinson predicts that a certain subgroup of $\mathbb{K}_2^{\text{tame}}(\mathcal{C})$ has rank g . Yet, non zero tame symbols are not easy to exhibit, see for instance [31] where elements in the tame group of some hyperelliptic curves over \mathbb{Q} are constructed.

2.9. Algebraic geometry on the spectral curve. We now come to the study of algebraic geometry on the spectral curve (\mathcal{C}_0, u, v)

Topology, cycles, and holomorphic 1-forms. The curve \mathcal{C}_0 defines a compact Riemann surface of a certain genus g . Actually, the genus can be computed from the polynomial $A(m, l)$ as the dimension g of the space of holomorphic forms, i.e. rational expressions $h(m^2, l)dm$ which are nowhere singular. Let $(\mathcal{A}_j, \mathcal{B}_j)_j$ be a symplectic basis of homology cycles,

$$\mathcal{A}_j \cap \mathcal{A}_{j'} = 0 \quad \mathcal{B}_j \cap \mathcal{B}_{j'} = 0 \quad \mathcal{A}_j \cap \mathcal{B}_{j'} = \delta_{j,j'}, \quad \text{for all } j, j' \in \{1, \dots, g\}. \quad (14)$$

For the moment, we choose an arbitrary basis, and we will have to consider later how objects depend on the basis, i.e to describe the action of the modular group $\text{Sp}_{2g}(\mathbb{Z})$. There is a notion of dual basis of holomorphic forms $(da_j)_j$, characterized by

$$\oint_{\mathcal{A}_{j'}} da_j = \delta_{j,j'}, \quad j \in \{1, \dots, g\}.$$

Then, the *period matrix* is defined as

$$\oint_{\mathcal{B}_{j'}} da_j = \tau_{j,j'},$$

and a classical result states that it is symmetric with positive definite imaginary part. We choose an arbitrary base point o , for example $o = p_c$, and introduce the *Abel map*

$$\begin{aligned} \mathbf{a}: \mathcal{C}_0 &\longrightarrow \mathbb{J} = \mathbb{C}^g / (\mathbb{Z}^g \oplus \tau \mathbb{Z}^g), \\ p &\longmapsto \left(\int_o^p da_1, \dots, \int_o^p da_g \right) \bmod \mathbb{Z}^g \oplus \tau \mathbb{Z}^g. \end{aligned}$$

When $g = 1$, \mathcal{C}_0 is an elliptic curve and \mathbf{a} is an isomorphism. When $g \geq 2$, this is only an immersion.

Theta functions and characteristics. For any $g \times g$ matrix τ which is symmetric with positive definite imaginary part, we can define the theta function

$$\theta(\mathbf{w}|\tau) = \sum_{\mathbf{p} \in \mathbb{Z}^g} e^{i\pi \mathbf{p} \cdot \tau \cdot \mathbf{p} + 2i\pi \mathbf{w} \cdot \mathbf{p}}, \quad \text{for all } \mathbf{w} \in \mathbb{C}^g.$$

Where there is no confusion, we omit to write the dependence in τ . θ is an even, quasi periodic function with respect to the lattice $\mathbb{Z}^g \oplus \tau \mathbb{Z}^g$:

$$\theta(\mathbf{w} + \mathbf{m} + \tau \cdot \mathbf{n}) = e^{-i\pi(2\mathbf{n} \cdot \mathbf{w} + \mathbf{n} \cdot \tau \cdot \mathbf{n})} \theta(\mathbf{w}). \quad (15)$$

We define a gradient ∇ acting implicitly on the variable \mathbf{w} , and a gradient D acting on the variable τ :

$$\nabla \theta = \left(\frac{\partial \theta}{\partial w_1}, \dots, \frac{\partial \theta}{\partial w_g} \right), \quad D \theta = \left(4i\pi \frac{\partial \theta}{\partial \tau_{i,j}} \right)_{1 \leq i, j \leq g}.$$

The theta function is solution to the heat equation

$$4i\pi \partial_{\tau_{j,j'}} \theta = \partial_{w_j} \partial_{w_{j'}} \theta$$

i.e.

$$D \vartheta = \nabla^{\otimes 2} \vartheta.$$

Throughout the article, we are going to use tensor notations, and indicate with a “ \cdot ” the contraction of indices. We consider $\nabla\theta$ and $D\theta$, and more generally $\nabla^{\otimes l}\theta$ (resp. $D^{\otimes l}\theta$) as a l -linear form (resp. a $2l$ -linear form), i.e. a $[0, l]$ tensor. For example, if T is a $[l, 0]$ tensor, we may write

$$\nabla^{\otimes l}\theta \cdot T = \sum_{j_1=1}^g \cdots \sum_{j_l=1}^g \frac{\partial^l \theta}{\partial w_{j_1} \cdots \partial w_{j_l}} T_{j_1, \dots, j_l}.$$

A *half-characteristics* is a vector

$$\mathbf{c} = \frac{1}{2}(\mathbf{n} + \tau \cdot \mathbf{m}) \quad \text{where } \mathbf{n}, \mathbf{m} \in \mathbb{Z}^g.$$

It is said *odd* or *even* depending on the parity of the scalar product $\mathbf{n} \cdot \mathbf{m}$. equation (15) implies that $\theta(\mathbf{c}|\tau)$ and its even-order derivatives vanish at odd half-characteristics, while the odd-order derivatives of $\theta(\mathbf{w})e^{i\pi\mu \cdot \mathbf{w}}$ vanish at even half-characteristics of the form $\frac{1}{2}(\mathbf{n} + \tau \cdot \mathbf{m})$. There is a notation for theta functions whose argument is shifted by a half-characteristics $\mathbf{c} = \nu + \tau \cdot \mu$,

$$\begin{aligned} \vartheta \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right] (\mathbf{w}) &= \sum_{\mathbf{p} \in \mathbb{Z}^g} e^{i\pi(\mathbf{p}+\mu) \cdot \tau \cdot (\mathbf{p}+\mu) + 2i\pi(\mathbf{w}+\nu) \cdot (\mathbf{p}+\mu)} \\ &= e^{i\pi\mu \cdot \tau \cdot \mu + 2i\pi\mu \cdot \nu + 2i\pi\mathbf{w} \cdot \mu} \theta(\mathbf{w} + \nu + \tau \cdot \mu). \end{aligned}$$

Note that we still have

$$\nabla^{\otimes 2} \vartheta \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right] = D \vartheta \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right].$$

Bergman kernel. For us, a *Bergman kernel* is a symmetric $(1, 1)$ form $B(p_1, p_2)$ on $\mathbb{C}_0 \times \mathbb{C}_0$ which has no residues and has no singularities except for a double pole with leading coefficient 1 on the diagonal, i.e., in a local coordinate λ ,

$$B(p_1, p_2) \underset{p_1 \rightarrow p_2}{=} \frac{d\lambda(p_1) \otimes d\lambda(p_2)}{(\lambda(p_1) - \lambda(p_2))^2} + O(1).$$

If we pick up a symplectic basis of homology $(\mathcal{A}, \mathcal{B})$, there is a unique Bergman kernel $B(p_1, p_2)$ which is normalized on the \mathcal{A} -cycles,

$$\oint_{\mathcal{A}_j} B(p_1, \cdot) = 0 \quad \text{for all } j \in \{1, \dots, g\}. \quad (16)$$

Moreover, B is symmetric in p_1 and p_2 and the basis of holomorphic form is retrieved by

$$\oint_{\mathcal{B}_j} B(p_1, \cdot) = 2i\pi \, da_j \quad \text{for all } j \in \{1, \dots, g\}. \quad (17)$$

Any other Bergman kernel takes the form

$$B_\kappa(p_1, p_2) = B(p_1, p_2) + 2i\pi \mathbf{da}(p_1) \cdot \kappa \cdot \mathbf{da}^t(p_2),$$

where κ is a symmetric $g \times g$ matrix of complex numbers and t denotes the transposition. As a matter of fact, B_κ satisfies the relations (16) and (17) if we replace $(\mathcal{A}, \mathcal{B})$ by a symplectic basis of generalized cycles $(\mathcal{A}^\kappa, \mathcal{B}^\kappa)$ defined by

$$\mathcal{A}^\kappa = \mathcal{A} - \kappa \mathcal{B}^\kappa, \quad \mathcal{B}^\kappa = \mathcal{B} - \tau \mathcal{A}.$$

In this formula, \mathcal{A} and \mathcal{B} should be interpreted as column vectors with g rows.

The Bergman kernel normalized of the \mathcal{A} -cycles can always be expressed in terms of theta functions

$$B_0(p_1, p_2) = d_{p_1} d_{p_2} \ln \theta(\mathbf{a}(p_1) - \mathbf{a}(p_2) + \mathbf{c} | \tau),$$

where \mathbf{c} is any non singular odd half-characteristics. Non-singular means that the right hand does not vanish identically when $p_1, p_2 \in \mathcal{C}_0$, and such characteristics exist [70]. Yet, this formula is not very useful for computations when $g \geq 2$. In practice, one may start from the equation $A(m^2, l) = 0$ defining \mathcal{C} and \mathcal{C}_0 , and find “by hand” a Bergman kernel and a basis of holomorphic forms expressed as rational expressions in m^2 and l with rational coefficients. Both methods are illustrated for genus 1 curves in Section 6.2.

Prime form. Let \mathbf{c} be a non singular odd half-characteristics. We introduce a holomorphic 1-form

$$dh_{\mathbf{c}}(p) = \nabla \theta(\mathbf{c}) \cdot \mathbf{da}(p).$$

It is such that its $2g - 2$ zeroes are all double. Then, the *prime form* $E(p_1, p_2)$, cf. [70], is a $(-1/2, -1/2)$ form defined on the universal cover of $\mathcal{C}_0 \times \mathcal{C}_0$,

$$E(p_1, p_2) = \frac{\theta(\mathbf{a}(p_1) - \mathbf{a}(p_2) + \mathbf{c})}{\sqrt{dh_{\mathbf{c}}(p_1)} \otimes \sqrt{dh_{\mathbf{c}}(p_2)}}.$$

It is antisymmetric in p_1 and p_2 , it has a zero if and only if $p_1 = p_2$ in \mathcal{C}_0 , and in a local coordinate λ ,

$$E(p_1, p_2) \Big|_{p_1 \rightarrow p_2} = \frac{\lambda(p_1) - \lambda(p_2)}{\sqrt{d\lambda(p_1)} \otimes \sqrt{d\lambda(p_2)}} + O((\lambda(p_1) - \lambda(p_2))^3).$$

The prime form appears in this article through the formulae

$$\exp \left[\frac{1}{2} \int_{p_2}^{p_1} \int_{p_2}^{p_1} (B_0(p, p') - \frac{du(p) du(p')}{(u(p) - u(p'))^2}) \right] = \frac{u(p_2) - u(p_1)}{E(p_1, p_2) \sqrt{du(p_1) du(p_2)}},$$

$$\exp \left[\int_{p_2}^{p_1} \int_{p_4}^{p_3} B_0(p, p') \right] = \frac{E(p_1, p_3) E(p_2, p_4)}{E(p_1, p_4) E(p_2, p_3)}.$$

Modular transformations. The group $\mathrm{Sp}_{2g}(\mathbb{Z})$ acts on those objects by transformation of the symplectic basis of homology cycles. Let γ be an element of $\mathrm{Sp}_{2g}(\mathbb{Z})$.

- The cycles \mathcal{A} and \mathcal{B} , interpreted as column vectors with g rows, transform by definition as

$$\begin{pmatrix} {}^\gamma \mathcal{A} \\ {}^\gamma \mathcal{B} \end{pmatrix} = M_\gamma \cdot \begin{pmatrix} \mathcal{A} \\ \mathcal{B} \end{pmatrix}, \quad M_\gamma = \begin{pmatrix} d & c \\ b & a \end{pmatrix},$$

where a, b, c, d are $g \times g$ integer matrices. The new basis $({}^\gamma \mathcal{A}, {}^\gamma \mathcal{B})$ is symplectic (see equation (14)) if and only if ${}^t b d$ and ${}^t c a$ are symmetric and ${}^t a d - {}^t c b = 1$. These are indeed the condition under which M_γ belongs to $\mathrm{Sp}_{2g}(\mathbb{Z})$. The dual basis of holomorphic forms, interpreted as a row vector with g columns, transforms as a modular weight -1 vector:

$${}^\gamma da(p) = {}^t(c\tau + d)^{-1} da(p).$$

- The matrix of periods transforms as

$${}^\gamma \tau = (a\tau + b)(c\tau + d)^{-1}.$$

Using the relations defining Sp_{2g} , one can check

$$(c\tau + d)^t (a\tau + b)(c\tau + d)^{-1} = (a\tau + b)^t, \quad (18)$$

so that ${}^\gamma \tau$ is indeed symmetric. We have denoted M^t , the transposed of a matrix M .

- The Bergman kernel $B_{\kappa; \tau}$ defined from the chosen basis of cycles (we have stressed the dependence in τ), transforms as

$${}^\gamma (B_{\kappa; \tau})(p_1, p_2) = B_{{}^\gamma \kappa; {}^\gamma \tau}(p_1, p_2),$$

$${}^\gamma \kappa = c(c\tau + d)^t + (c\tau + d)\kappa(c\tau + d)^t. \quad (19)$$

- The generalized cycles (A^κ, B^κ) on which B_κ is normalized are modular expression of weight 1:

$${}^\nu A^\kappa = (c\tau + d)A^\kappa, \quad {}^\nu B^\kappa = [(c\tau + d)^{-1}]^t B^\kappa.$$

We have used the relation

$$[a - (a\tau + b)(c\tau + d)^{-1}c] = [(c\tau + d)^{-1}]^t$$

which can be deduced from equation (18).

- The theta function transforms as

$$\theta(\mathbf{w} + \Delta_\gamma | {}^\nu \tau) = \Xi_\gamma \sqrt{\det(c\tau + d)} e^{-\mathbf{w} \cdot \tau \cdot \mathbf{w}} \theta((c\tau + d)\mathbf{w} | \tau), \quad (20)$$

where Δ_γ is the half-characteristics

$$\Delta_\gamma = \frac{1}{2}(\text{diag}(ab^t) + \text{diag}(cd^t)\tau)$$

and Ξ_γ a eighth root of unity.

2.10. Baker–Akhiezer spinors. Given a 1-form ω on \mathbb{C}_0 , a complex number $\mathfrak{h} \in \mathbb{C}^\times$, and vectors $\mu, \nu \in \mathbb{C}^g / \mathbb{Z}^g$, we set

$$\begin{aligned} \psi_{\text{BA}}(p_1, p_2) &= \frac{\exp\left(\frac{1}{\mathfrak{h}} \int_{p_2}^{p_1} \omega\right)}{E(p_1, p_2)} \frac{\vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right](\mathbf{a}(p_1) - \mathbf{a}(p_2) + \zeta_{\mathfrak{h}})}{\vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right](\zeta_{\mathfrak{h}})} \\ &= \sqrt{\nabla\theta(\mathbf{c}) \cdot \mathbf{d}\mathbf{a}(p_1)} \otimes \sqrt{\nabla\theta(\mathbf{c}) \cdot \mathbf{d}\mathbf{a}(p_2)} \\ &\quad \exp\left(\int_{p_2}^{p_1} (\mathfrak{h}^{-1}\omega + 2i\pi\mu \cdot \mathbf{d}\mathbf{a})\right) \\ &\quad \frac{\theta(\mathbf{a}(p_1) - \mathbf{a}(p_2) + \zeta + \nu + \tau \cdot \mu)}{\theta(\mathbf{a}(p_1) - \mathbf{a}(p_2) + \mathbf{c}) \theta(\zeta_{\mathfrak{h}} + \nu + \tau \cdot \mu)}, \end{aligned}$$

with

$$\zeta_{\mathfrak{h}} = \text{frac}\left[\frac{1}{2i\pi\mathfrak{h}} \oint_{\mathcal{B}} \omega\right] - \tau \cdot \text{frac}\left[\frac{1}{2i\pi\mathfrak{h}} \oint_{\mathcal{A}} \omega\right].$$

For a vector $\mathbf{w} \in \mathbb{C}^g$, we have denoted $\text{frac}[\mathbf{w}]$ the vector of $[0, 1]^g$ which is equal to \mathbf{w} modulo \mathbb{Z}^g . ψ_{BA} is called a *Baker–Akhiezer spinor*, it is a $(1/2, 1/2)$ -form defined a priori on the universal cover of $\mathcal{C}_0 \times \mathcal{C}_0$, since we have

$$\frac{\psi_{\text{BA}}(p_1 + \mathbf{m}\mathcal{A} + \mathbf{n}\mathcal{B}, p_2)}{\psi_{\text{BA}}(p_1, p_2)} = \exp\left(\left(2i\pi\mu + \mathfrak{h}^{-1} \oint_{\mathcal{A}} \omega\right) \cdot \mathbf{m} + \left(\zeta + 2i\pi(\nu - \mathbf{c}) + \mathfrak{h}^{-1} \oint_{\mathcal{B}} \omega\right) \cdot \mathbf{n}\right).$$

It is regular apart from a simple pole when $p_1 = p_2$:

$$\psi_{\text{BA}}(p_1, p_2) \underset{p_1 \rightarrow p_2}{\sim} \frac{\sqrt{d\lambda(p_1)} \otimes \sqrt{d\lambda(p_2)}}{\lambda(p_1) - \lambda(p_2)},$$

and has an essential singularity when p_1 or p_2 reach a singularity of ω , of the form

$$\psi_{\text{BA}}(p_1, p_2) \propto \exp\left(\frac{1}{\mathfrak{h}} \int_{p_2}^{p_1} \omega\right).$$

Baker–Akhiezer functions have been introduced in [60] to write down some explicit solutions of the KP hierarchy. They can be obtained from the Baker–Akhiezer spinor when ω is a meromorphic 1-form, and by sending p_2 to a pole of ω with an appropriate regularization (see for instance [13]). Modular transformations act on ψ_{BA} only by a change of the vectors μ, ν . We have introduced a normalization constant $\mathfrak{h} \in \mathbb{C}^\times$, to be adjusted later. In general, the ratio involving $\vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right]$ does not have a limit, neither has a power series expansion when $\mathfrak{h} \rightarrow 0$.

But we can say more if we assume the Boutroux and the quantization condition, i.e. that there exists $\varsigma \in \mathbb{N}^*$ such that, for all closed cycles Γ ,

$$\left(\text{Im} \oint_{\Gamma} \omega\right) = 0 \quad \text{and} \quad \left(\text{Re} \oint_{\Gamma} \omega\right) \in \frac{2\pi^2}{\varsigma} \cdot \mathbb{Z}.$$

Let us denote s_A and s_B integer vectors such that

$$\oint_{\mathcal{A}} \omega = \frac{2\pi^2}{\varsigma} s_A \quad \text{and} \quad \oint_{\mathcal{B}} \omega = \frac{2\pi^2}{\varsigma} s_B.$$

It is then natural to consider values of \mathfrak{h}^{-1} belonging to arithmetic subsequences on the imaginary axis:

$$\mathfrak{h} = \frac{i\pi}{k}, \quad k \in \varsigma \cdot \mathbb{Z} + r.$$

Indeed, we find

$$\zeta = \text{frac}\left[\frac{r s_B}{\zeta}\right] - \tau \cdot \text{frac}\left[\frac{r s_A}{\zeta}\right],$$

so that the argument of the theta functions only depend on $r = k \bmod \zeta$. We have

$$\frac{2\pi}{k} \ln |\psi_{\text{BA}}(p_1, p_2)| \underset{k \in \zeta \cdot \mathbb{Z} + r}{\underset{k \rightarrow \infty}{\sim}} \text{Im}\left(\int_{p_2}^{p_1} 2\omega\right),$$

and the Boutroux condition also ensure that

$$\text{Im}\left(\int_{p_2}^{p_1} \omega\right)$$

does not depend on the path of integration between p_1 and p_2 . For a hyperbolic 3-manifold, if we choose $\omega = v du$, the right hand side is $\text{Vol}_a(p_1) - \text{Vol}_a(p_2)$ and this asymptotics is exactly the one involved in the generalized the volume conjecture (see §5.2)

2.11. Branchpoints and local involution. In this article, we reserve the name *ramification points* to points in \mathcal{C}_0 which are zeroes of $du = d \ln m$. The value of m at a ramification point is called a *branchpoint*. We use generically the letter a to denote a ramification point. Since \mathcal{C} is defined by a polynomial equation $A(e^u, e^v) = 0$, we must have $m(a) \neq 0, \infty$. When a is a simple zero of $d \ln m$, we call it a *simple ramification point*, and we can define at least in a neighborhood $\mathcal{U}_a \subseteq \mathcal{C}_0$ of a the *local involution* $p \mapsto \bar{p}$,

$$p, \bar{p} \in \mathcal{U}_a, \quad m(p) = m(\bar{p}) \text{ and } p \neq \bar{p}.$$

Since A is quasi-reciprocal and has real coefficients, the involution

$$\iota: (l, m) \mapsto \left(\frac{1}{m}, \frac{1}{l}\right)$$

and the complex conjugation $*$ act on the set of coordinates $(m(a), l(a))$ of the ramification points, and decompose it into orbits with 2 elements (for an a such that $(l(a), m(a))$ is real or unitary) or 4 elements (in general). *Amphichiral* knot complements admit an orientation reversing automorphism, so that $\iota_m(m, l) = (1/m, l)$ by $\iota_l(m, l) = (m, 1/l)$ are separately symmetries of their A-polynomial. Then at the level of spectral curves, the set of ramification points can be decomposed further into orbits of 2, 4 or 8 elements.

3. Topological recursion

The *topological recursion* associates, to any spectral curve (\mathcal{C}_0, u, v) , a family of symmetric $(1, \dots, 1)$ forms $\omega_n^h(p_1, \dots, p_n)$ on \mathcal{C}_0^n ($n \in \mathbb{N}^*$, $h \in \mathbb{N}$) and a family of numbers F_h ($h \in \mathbb{N}$). These objects have many properties, we shall only mention those we use without proofs. We refer to [40] for a detailed review of the topological recursion. The fact that, here or in topological strings, one encounters spectral curves of the form $\text{Pol}(e^u, e^v) = 0$ rather than $\text{Pol}(u, v) = 0$, does not make a big difference in the formalism.

We assume that all ramification points are simple. This is satisfied for most of the A-polynomials we have studied (see the tables in §B.3). The topological recursion can also be defined when some ramification points are not simple [81, 14], but we do not address this issue here.

3.1. Definitions. Let (\mathcal{C}_0, u, v) be a spectral curve endowed with a basis of cycles (A, B) . Hence, there is a privileged Bergman kernel $B(p_1, p_2)$. To shorten notations, we write

$$dz_1 \cdots dz_n$$

instead of

$$dz_1 \otimes \cdots \otimes dz_n,$$

for a $(1, \dots, 1)$ -form.

Recursion kernel. We introduce the recursion kernel

$$K(p_0, p) = \frac{-\frac{1}{2} \int_{\bar{p}}^p B(\cdot, p_0)}{(v(p) - v(\bar{p}))du(p)}. \tag{21}$$

$K(p_0, p)$ is a 1-form with respect to p_0 globally defined on \mathcal{C}_0 , and a (-1) -form with respect to p which is defined locally around each ramification point.

Differential forms. We define

$$\begin{aligned} \omega_1^0(p) &= v(p)du(p), \\ \omega_2^0(p_1, p_2) &= B(p_1, p_2), \end{aligned}$$

and recursively

$$\begin{aligned} \omega_n^h(p_0, p_I) &= \sum_a \operatorname{Res}_{p \rightarrow a} K(p_0, p) \left[\omega_{n+1}^{h-1}(p, \bar{p}, p_I) + \sum'_{h', J} \omega_{|J|+1}^{h'}(p, p_J) \omega_{n-|J|}^{h-h'}(\bar{p}, p_{I \setminus J}) \right]. \end{aligned} \tag{22}$$

In the left hand side, $I = \{1, \dots, n - 1\}$ and p_I in a $(n - 1)$ -uple of points of \mathbb{C}_0 . For any $J \subseteq I$, p_J is the uple of points indexed by the subset J . In the right hand side, we take the residues at all ramification points, and the \sum' in the right hand side ranges over $h' \in \{0, \dots, h\}$ and all splitting of variables $J \subseteq I$, excluding $(J, h') = (\emptyset, 0)$ and (I, h) . The formula above is a recursion on the *level*

$$\chi = 2h - 2 + n.$$

ω_n^h has a diagrammatic interpretation (Figure 1), it can be written as a sum over graphs with n external legs, h handles, and thus Euler characteristics $-\chi$. However, the weights of the graphs are non local, they involve stacks of $2g + 2 - n$ residues where the ordering matters.




Figure 1. Diagrammatic representation of the topological recursion which defines ω_n^h . Each ω_n^h is represented as a “surface” with h handles and n punctures, i.e. with Euler characteristics $\chi = 2 - 2h - n$. The diagrammatic representation of the topological recursion, is that one can compute ω_n^h with Euler characteristics χ in terms of $\omega_{n'}^{h'}$ with $\chi' = 2 - 2h' - n' > \chi$, by “removing a pair of pants” from the corresponding surface.

Although equation (22) seems to give a special role to the variable p_0 , one can prove (for instance from the diagrammatic representation) that $\omega_n^h(p_0, \dots, p_{n-1})$ is symmetric in p_0, \dots, p_{n-1} . Except maybe ω_1^0 , the ω_n^h are all meromorphic $(1, \dots, 1)$ -forms on \mathbb{C}_0^n , which have no residues and have poles only at the ramification points.

We illustrate the computation at level 1. To write down the residues it is convenient to choose a local coordinate at each ramification point, for instance

$$\lambda_a(p) = \sqrt{m(p) - m(a)} = \sqrt{e^{u(p)} - e^{u(a)}},$$

which has the advantage that

$$\lambda_a(\bar{p}) = -\lambda_a(p).$$

If r is a function or R is a 1-form, we denote

$$\partial^j r(\underline{a}) = \frac{\partial^j r(p)}{\partial \lambda_a(p)^j} \Big|_{p=a} \quad \text{and} \quad \partial^j R(\underline{a}) = \frac{\partial^j \left(\frac{R(p)}{d\lambda_a(p)} \right)}{\partial \lambda_a(p)^j} \Big|_{p=a}.$$

Then, we find

$$\begin{aligned} \omega_3^0(p_0, p_1, p_2) &= \sum_a \operatorname{Res}_{z \rightarrow a} K(p_0, p) [B(p, p_1) B(p_2, \bar{p}) + B(\bar{p}, p_1) B(p, p_2)] \\ &= \sum_a \frac{B(p_0, \underline{a}) B(p_1, \underline{a}) B(p_2, \underline{a})}{2 v'(\underline{a})}. \end{aligned} \tag{23}$$

To write down ω_1^1 , we need to expand

$$\frac{B(p, \bar{p})}{(d\lambda_a(p))^2} \Big|_{p \rightarrow a} = -\frac{1}{4\lambda_a^2(p)} + S_B(a) + o(1).$$

Then, we have

$$\begin{aligned} \omega_1^1(p_0) &= \sum_a \operatorname{Res}_{p \rightarrow a} K(p_0, p) B(p, \bar{p}) \\ &= \sum_a \frac{S_B(a)}{2} B(p_0, \underline{a}) + \frac{1}{48} (v'''(\underline{a}) B(p_0, \underline{a}) - v'(\underline{a}) \partial_2^2 B(p_0, \underline{a})). \end{aligned}$$

Stable free energies. We have already met the abelian function

$$\Phi(p) = \int_o^p \omega_1^0 = \int_o^p v \, du.$$

For $h \geq 2$, we define

$$F_h = \frac{1}{2h-2} \sum_a \operatorname{Res}_{p \rightarrow a} \Phi(p) \omega_1^h(p).$$

Since ω_1^h has no residues, F_h does not depend on the basepoint o . The numbers F_h are called the *stable free energies* of the spectral curve. We are not going to give an explicit definition of the *unstable free energies* F_0 and F_1 . Actually, for the computation of the BA kernels and later the asymptotics of the colored Jones polynomial, it is not necessary to know how to compute the free energies, we only need one of their key property called special geometry (see equation (24)). So, we just state that there exists F_0 and F_1 satisfying equation (24), it is in fact a way to define them.

3.2. Deformation of spectral curves. By abuse of notations, we write

$$F_h = \omega_{n=0}^h,$$

i.e. we consider ω_n^h for all $n, h \in \mathbb{N}$. Unless specified, the properties mentioned below also hold for the unstable free energies. Special geometry expresses the variation of ω_n^h when $\phi = v \, du$ is deformed by addition of a meromorphic 1-form Ω . By form-cycle duality on \mathcal{C}_0 , to any meromorphic 1-form Ω we can associate a cycle Ω^* and a germ of holomorphic function on Ω^* denoted Λ_Ω , such that

$$\Omega(p) = \int_{\Omega^*} \Lambda_\Omega(\cdot) B(\cdot, p).$$

Then, for a smooth family of spectral curves $\mathcal{S}_\alpha = (\mathcal{C}_0, u_\alpha, v_\alpha)$ such that

$$(v_\alpha - v) \, du - (u_\alpha - u) \, dv \underset{\alpha \rightarrow 0}{\sim} \alpha \, \Omega.$$

we have

$$\left. \frac{\partial}{\partial \alpha} \omega_n^h[\mathcal{S}_\alpha](p_I) \right|_{\alpha=0} = \int_{\Omega^*} \Lambda_\Omega(\cdot) \omega_{n+1}^h[\mathcal{S}_{\alpha=0}](\cdot, p_I). \quad (24)$$

Note that from the expression of ω_3^0 in equation (23), one retrieves as a special case the analog of Rauch variational formula [82] for the variation of the Bergman kernel $\omega_2^0 = B$ along any meromorphic deformation.

In this article, deformations by holomorphic 1-forms and by 1-forms with simple poles will play a special role.

Variations of filling fractions. The *filling fractions* are defined by

$$\epsilon_j = \frac{1}{2i\pi} \oint_{\mathcal{A}_j} v \, du.$$

Performing a variation of filling fractions amounts to add to $v \, du$ a holomorphic 1-form, i.e. use the deformation

$$\Omega(p) = 2i\pi \, da_j(p) = \int_{\mathcal{B}_j} B(\cdot, p). \quad (25)$$

We denote $\omega_n^{h,(l)}$ the $[0, l]$ -tensor of l^{th} derivatives of ω_n^h with respect to the filling fractions, and, according to equation (24),

$$\omega_n^{h,(l)}(p_I) = \underbrace{\oint_{\mathcal{B}} \cdots \oint_{\mathcal{B}}}_{l \text{ times}} \omega_{n+l}^h(\cdots, p_I).$$

In particular, the tensor of second derivatives of $F_0 = \omega_0^0$ is the matrix of periods

$$F_0'' = \oint_{\mathcal{B}} \oint_{\mathcal{B}} B = 2i\pi \, \tau.$$

Deformation by simple poles. Given a couple of distinct points (p_1, p_2) , we denote

$$dS_{p_2, p_1}(p) = \int_{p_2}^{p_1} B(\cdot, p). \quad (26)$$

This 1-form is characterized by a simple pole at $p = p_1$ (resp. $p = p_2$) with residue 1 (resp. -1), no other singularities, and vanishing \mathcal{A} -cycle integrals. If we perform an infinitesimal deformation with $\Omega(p) = dS_{p_2, p_1}(p)$, we obtain according to equation (24)

$$\delta\omega_n^h(p_I) = \int_{p_2}^{p_1} \omega_{n+1}^h(\cdot, p_I).$$

3.3. Symplectic invariance. The topological recursion also has nice properties under global transformations of the spectral curve (\mathcal{C}_0, u, v) . To simplify, we consider in this paragraph $(n, h) \neq (0, 0), (0, 1), (1, 0)$, and just mention that the properties below are slightly modified for those cases.

It is very easy to prove from the definitions the following properties.

Property 3.1. *If $v du$ is replaced by $\alpha v du$ for some $\alpha \in \mathbb{C}^\times$, ω_n^h is replaced by $\alpha^{2-2h-n} \omega_n^h$. In particular, the stable free energies F_h are unchanged when (u, v) is replaced by $(-u, v)$ or $(u, -v)$.*

Property 3.2. *If (u, v) is replaced by $(u, v + f(u))$ with f at least a germ of holomorphic function in the neighborhood of the values $u(a)$, ω_n^h are unchanged.*

According to the first property, replacing $m = e^u$ and $l = e^v$ by some of their powers i.e. use $(\pm m^a, \pm l^b)$ instead of (m, l) , only affect the ω_n^h by a scaling factor. The second property tells us that the ω_n^h are the same if we change the signs of m and l , or even replace⁸ l by lm^a for some power a . There is conjecturally a third property concerning the exchange of u and v .

Property 3.3. *If (u, v) is replaced by $(v, -u)$, the F_h are unchanged, and for $n \geq 1$, the cohomology class of ω_n^h is multiplied by the sign $(-1)^n$.*

This has only been proved [39] when u and v are meromorphic function on the curve \mathcal{C}_0 , that is for spectral curves defined by an equation $\text{Pol}(u, v) = 0$. This invariance of the free energies under this exchange has meaningful consequences in random matrix theory and enumerative geometry (see [38, §10.4.1] for an example). Here and in topological strings, we rather have to consider spectral curves of the form $\text{Pol}(e^u, e^v) = 0$. We believe that Property 3.3 survives in this context with a few extra assumptions, although this has not been established yet. For example, within “remodeling the B-model,” it implies the framing independence of the closed topological string sector.

In other words, if Property 3.3 holds, the F_g , and cohomology classes of the ω_n^h up to a sign, are invariant under all the transformations which preserve the symbol $du \wedge dv$. This suggests to consider the F_h and the ω_n^h up to a sign as “symplectic invariants” of the function field $\mathbb{K} = \mathbb{C}(\mathcal{C}_0)$. We have seen in §2.5 that the real part of the primitive of ω_1^0 essentially coincide with the Bloch regulator of the symbol $\{m, l\}$. It would be interesting to investigate the possible meaning of the topological recursion in terms of K-theory of \mathbb{K} .

⁸ This last operation is very useful to lower the degree of m in A-polynomials. For instance, the A-polynomial of the Pretzel(-2, 3, 7)

$$\mathfrak{A}(l, m) = m^{110}l^6 - m^{90}(m^2 - 1)^2l^5 - m^{72}(2m^2 + 1)l^8 + m^{36}(m^2 + 2)l^4 + m^{16}(m^2 - 1)^2l - 1$$

looks simpler if we use the variable $\ell = lm^8$:

$$\mathfrak{A}(l, m) = m^{14}\ell^6 - m^{10}(m^2 - 1)\ell^5 - m^8(2m^2 + 1)\ell^4 + m^4(m^2 + 2)\ell^2 + (m^2 - 1)^2\ell - 1$$

3.4. Deformation of the Bergman kernel. Instead of $B(p_1, p_2)$, we could have used in the definitions (21) and (22) another Bergman kernel,

$$B_\kappa(p_1, p_2) = B(p_1, p_2) + 2i\pi \mathbf{a}(p_1) \cdot \kappa \cdot \mathbf{da}^t(p_2).$$

We denote $\omega_{n|\kappa}^h$ the corresponding objects. We note that they are polynomials of degree $3h - 3 + n$ in κ , and it is not difficult to prove

$$\begin{aligned} \frac{\partial \omega_{n|\kappa}^h(p_I)}{\partial \kappa} &= \frac{1}{2} \frac{1}{2i\pi} \left(\oint_{\mathcal{B}_\kappa} \oint_{\mathcal{B}_\kappa} \omega_{n+2|\kappa}^{h-1}(\cdot, \cdot, p_I) \right. \\ &\quad \left. + \sum_{\substack{J \subseteq I \\ 0 \leq h' \leq h}} \oint_{\mathcal{B}_\kappa} \omega_{|J|+1|\kappa}^h(\cdot, p_J) \oint_{\mathcal{B}_\kappa} \omega_{n-|J|+1|\kappa}^{h-h'}(\cdot, p_{I \setminus J}) \right). \end{aligned} \quad (27)$$

The special geometry (equation (24)) for meromorphic deformations normalized on the \mathcal{A} -cycles still holds for $\omega_{n|\kappa}^h$ at any fixed κ . However, variations of κ and filling fractions are mixed, since the holomorphic forms da_j in equation (25) are defined from $B = B_{\kappa=0}$ and not B_κ . The appropriate formula can be found in [38], it is closely related to ‘‘holomorphic anomaly equations’’ [7], but it will not be used in this article.

3.5. Effect of an involution. The A-polynomial comes with an involution

$$\iota: (m, l) = (1/m, 1/l).$$

It induces an involutive linear map ι_* on the space of holomorphic 1-forms on \mathcal{C}_0 . The g eigenvalues of ι_* are thus ± 1 . By integration, it induces an involutive isomorphism of the Jacobian of the curve, that we denote $\underline{\iota}_*$. The number of $+1$ eigenvalues is the genus of the quotient curve \mathcal{C}_0/ι .

The case $\iota_* = \varepsilon \text{id}$ is of particular interest. When $\varepsilon = 1$, $\underline{\iota}_*$ is a translation by a half-period, and when $\varepsilon = -1$, $\underline{\iota}_*$ is a central symmetry. In these two situations, all admissible Bergman kernels

$$B_\kappa(z_1, z_2) = d_{z_1} d_{z_2} \ln \theta(\mathbf{a}(z_1) - \mathbf{a}(z_2) + \mathbf{c}) + 2i\pi \mathbf{da}(z_1) \cdot \kappa \cdot \mathbf{da}(z_2)$$

are invariant under ι , and so is the recursion kernel $K_\kappa(z_0, z)$. Since the set of ramification points is stable under ι , we can recast the residue formula by choosing

a representative a' in each pair $\{a, \iota(a)\}$ of ramification points

$$\begin{aligned} \omega_{n|\kappa}^h(p_0, p_I) &= \sum_{a'} \operatorname{Res}_{p \rightarrow a'} K_\kappa(p_0, p) E_{n|\kappa}^h(p, z_I) + \operatorname{Res}_{p \rightarrow \iota(a')} K_\kappa(p_0, p) E_{n|\kappa}^h(p, p_I) \\ &= \sum_{a'} \operatorname{Res}_{p \rightarrow a'} [K_\kappa(p_0, p) E_{n|\kappa}^h(p, p_I) + K_\kappa(p_0, \iota(p)) E_{n|\kappa}^h(\iota(p), p_I)] \\ &= \sum_{a'} \operatorname{Res}_{p \rightarrow a'} [K_\kappa(p_0, p) E_{n|\kappa}^h(p, p_I) + K_\kappa(\iota(p_0), p) E_{n|\kappa}^h(\iota(p), p_I)]. \end{aligned}$$

By recursion on $2g - 2 + n \leq 0$, we infer that

$$E_{n|\kappa}^h(\iota(p), p_I) = E_{n|\kappa}^h(p, \iota(p_I))$$

and

$$\omega_{n|\kappa}(p_0, p_I) = \omega_{n|\kappa}^h(\iota(p_0), \iota(p_I)).$$

This result has an interesting corollary when $\iota_* = -\operatorname{id}$: by duality,

$$\iota_* \mathcal{B}_\kappa = -\mathcal{B}_\kappa.$$

Hence we get the property below.

Property 3.4. *If $\iota_* = -\operatorname{id}$, then*

$$\underbrace{\oint_{\mathcal{B}_\kappa} \cdots \oint_{\mathcal{B}_\kappa}}_{d \text{ times}} \omega_{n|\kappa}^h(p_1, \dots, p_{n-d}, \cdot) = 0 \quad \text{when } d \text{ is odd.}$$

In the case $(n, h) = (1, 0)$, since $\ln d \ln m$ is always invariant under ι , we have the next property.

Property 3.5. *If $\iota_* = -\operatorname{id}$, for any closed cycle $\Gamma \subseteq \mathcal{C}_0$, then*

$$\oint_{\Gamma} v \, du = 0$$

As one can see in see the tables in §B.3, $\iota_* = -\operatorname{id}$ is neither rare nor the rule for complement of hyperbolic knots. We observe however that the genus of the quotient \mathcal{C}_0/ι is low compared to the genus of \mathcal{C}_0 : the “simplest” knot we found for which the quotient has not genus 0 is $\mathbf{8}_{21}$. The geometrical significance of these observations from the point of view of knot theory is unclear.

4. Non-perturbative topological recursion

The *perturbative partition* function is usually defined as

$$Z_{\text{pert},\hbar} = \exp\left(\sum_{h \geq 0} \hbar^{2h-2} F_h\right),$$

where F_h are the free energies. However, the genuine partition function of a quantum field theory (like the Chern–Simons theory or topological string theory) should have properties that $Z_{\text{pert},\hbar}$ does not satisfy. For instance, it should be independent of the classical solution chosen to quantize the theory (background independence), and it should have modular properties (e.g. S-duality) whenever this makes sense.

From the topological recursion applied to a spectral curve (\mathcal{C}_0, u, v) , and theta functions, we are going to define a non-perturbative partition function \mathcal{T}_\hbar which implements such properties. Modular transformations correspond here to change of symplectic basis of cycles on \mathcal{C}_0 . Then, one can define non-perturbative “wave functions.” To keep a precise vocabulary, we shall introduce quantities

$$\psi_\hbar^{[n|n]}(p_1, p_2; \dots; p_{2n-1}, p_{2n}),$$

that we call $n|n$ -kernels, which depend on $2n$ points on the curve. In particular, the leading order of $\psi_\hbar^{[1|1]}(p_1, p_2)$ when $\hbar \rightarrow 0$ will be given by the Baker–Akhiezer spinor. We prefer to use a new letter \hbar for the formal parameter. We shall find later that in the application we consider, it must be identified to \hbar defined in terms of the parameter $q = e^{2\hbar}$ in which the colored Jones polynomial is a Laurent polynomial, but this identification might be different when considering other problems.

4.1. Definitions. We use the notations of §2.9. We take as data a spectral curve (\mathcal{C}_0, u, v) endowed with a basis of cycles, we choose two vectors $\mu, \nu \in \mathbb{C}^g$ and we set

$$\tau = \frac{F_0''}{2i\pi},$$

and

$$\zeta = \text{frac}\left[\frac{1}{2i\pi\hbar} \oint_{\mathcal{A}} v \, du\right] - \tau \cdot \text{frac}\left[\frac{1}{2i\pi\hbar} \oint_{\mathcal{B}} v \, du\right]. \quad (28)$$

We give the definitions, which we comment in §4.2.

Partition function. The *non-perturbative partition function* is by definition

$$\mathcal{T}_{\hbar} = \exp\left(\sum_{h \geq 0} \hbar^{2h-2} F_h\right) \mathfrak{S}_{\hbar} \vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right](\zeta_{\hbar}|\tau), \quad (29)$$

where

$$\mathfrak{S}_{\hbar} = \sum_{r \geq 0} \frac{1}{r!} \sum_{\substack{h_j \geq 0, d_j \geq 1 \\ 2h_j - 2 + d_j > 0}} \hbar^{\sum_j 2h_j - 2 + d_j} \bigotimes_{j=1}^r \frac{F_{h_j}^{(d_j)} \cdot \nabla^{\otimes d_j}}{(2i\pi)^{d_j} d_j!}.$$

We may isolate its leading behavior by writing

$$\mathcal{T}_{\hbar} = e^{\hbar^{-2}F_0 + F_1} \vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right](\zeta_{\hbar}|\tau) \widehat{\mathcal{T}}_{\hbar},$$

where now $\lim_{\hbar \rightarrow 0} \widehat{\mathcal{T}}_{\hbar} = 1$. We consider this expression as a formal asymptotic series with parameter $\hbar \rightarrow 0$. The coefficient of \hbar^{χ} in general depend on \hbar , but does not have a power series expansion in \hbar . Thus, it is meaningful to speak of the χ^{th} -order term in the expansion, keeping in mind that this coefficient may also depend on \hbar .

(1|1)-Kernel. In integrable systems, the Sato formula expresses the wave function as Schlesinger transforms of the tau function, which in our language correspond to adding a 1-form with simple poles to $\phi = v du$. Actually, we prefer to work with the *kernel* $\psi_{\hbar}(p_1, p_2)$ which is a function on $\mathcal{C}_0 \times \mathcal{C}_0$, defined as

$$\psi_{\hbar}(p_1, p_2) = \frac{\mathcal{T}_{\hbar}[v du \rightarrow v du + \hbar dS_{p_2, p_1}]}{\mathcal{T}_{\hbar}[v du]}, \quad (30)$$

where dS was defined in equation (26). We introduce shortcut notations

$$\vartheta = \vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right](\zeta_{\hbar}|\tau), \quad \vartheta_{\bullet} = \vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right](\zeta_{\hbar} + \mathbf{a}(p_1) - \mathbf{a}(p_2)|\tau).$$

$\psi_{\hbar}(p_1, p_2)$ can be computed thanks to special geometry:

$$\psi_{\hbar}(p_1, p_2) = \frac{\exp\left(\frac{1}{\hbar} \int_{p_2}^{p_1} v du\right)}{E(p_1, p_2)(du(p_1) du(p_2))^{1/2}} \frac{\mathfrak{N}_{\hbar, p_1, p_2}}{\mathfrak{D}_{\hbar}}. \quad (31)$$

where

$$\mathfrak{N}_{\hbar, p_1, p_2} = \left(\sum_{r \geq 0} \frac{1}{r!} \sum_{\substack{h_j, n_j \geq 0, d_j \geq 1 \\ 2h_j - 2 + d_j + n_j > 0}} \hbar^{\sum_j 2h_j - 2 + d_j + n_j} \bigotimes_{j=1}^r \frac{n_{j, p_1, p_2}}{(2i\pi)^{d_j} d_j! n_j!} \right) \vartheta_{\bullet},$$

$$n_{j,p_1,p_2} = \int_{p_2}^{p_1} \cdots \int_{p_2}^{p_1} \omega_{n_j}^{h_j,(d_j)} \cdot \nabla^{\otimes d_j},$$

and

$$\mathfrak{D}_{\hat{\hbar}} = \left(\sum_{r \geq 0} \frac{1}{r!} \sum_{\substack{h_j \geq 0, d_j \geq 1 \\ 2h_j - 2 + d_j > 0}} \hat{\hbar}^{\sum_j 2h_j - 2 + d_j} \bigotimes_{j=1}^r \frac{F_{h_j}^{(d_j)} \cdot \nabla^{\otimes d_j}}{(2i\pi)^{d_j} d_j!} \right) \vartheta.$$

In the second line of equation (31), all the n_j variables in $\omega_{n_j}^{h_j,(d_j)}$ are integrated over $\int_{p_2}^{p_1}$, and recall from special geometry that

$$\int_{p_2}^{p_1} \cdots \int_{p_2}^{p_1} \omega_n^{h,(d)} = \int_{p_2}^{p_1} \cdots \int_{p_2}^{p_1} \underbrace{\oint_{\mathcal{B}} \cdots \oint_{\mathcal{B}}}_{d \text{ times}} \omega_{n+d}^h.$$

Again, equation (31) should be understood as a formal asymptotic series with parameter $\hat{\hbar} \rightarrow 0$. It can be shown [13] that $\psi_{\hat{\hbar}}(p_1, p_2)$ does not change when p_1 or p_2 goes around an \mathcal{A} or a \mathcal{B} cycle. Since ψ is the ratio of two partition function, the exponential involving the free energies F_h in the numerator of the first line of equation (31) cancels with the same factor present in the denominator. As we claimed earlier, only the expression of ω_n^h is needed to compute the kernel, not the expression of the free energies. We may isolate its leading behavior:

$$\psi_{\hat{\hbar}}(p_1, p_2) = \frac{\psi_{\text{BA}}(p_1, p_2)}{(\text{du}(p_1) \text{du}(p_2))^{1/2}} \hat{\psi}_{\hat{\hbar}}(p_1, p_2),$$

where now $\lim_{\hat{\hbar} \rightarrow 0} \hat{\psi}_{\hat{\hbar}}(p_1, p_2) = 1$.

$n|n$ -kernels. If we perform n successive Schlesinger transformations, we are led to define the $n|n$ -kernels

$$\psi_{\hat{\hbar}}^{[n|n]}(p_1, o_1; \cdots; p_n, o_n) = \frac{\mathcal{T}_{\hat{\hbar}} \left[v \text{ du} \longrightarrow v \text{ du} + \sum_{k=1}^n \hat{\hbar} \text{dS}_{o_k, p_k} \right]}{\mathcal{T}_{\hat{\hbar}} [v \text{ du}]}, \quad (32)$$

which are functions on \mathcal{C}_0^{2n} . Equation (31) has a straightforward generalization:

$$\begin{aligned} & \psi_{\hbar}^{[n|n]}(p_1, o_1; \dots; p_n, o_n) \\ &= \prod_{1 \leq k < k' \leq n} \frac{E(p_k, p_{k'}) E(o_k, o_{k'})}{E(p_k, o_{k'}) E(o_k, p_{k'})} \frac{\exp\left(\frac{1}{\hbar} \int_{\bullet} v \, du\right)}{\prod_{k=1}^n E(p_k, o_k) (du(p_k) \, du(o_k))^{1/2}} \frac{\mathfrak{D}_{\hbar}^{[n|n]}}{\mathfrak{D}_{\hbar}}, \end{aligned} \quad (33)$$

where

$$\begin{aligned} \mathfrak{N}_{\hbar}^{[n|n]} &= \mathfrak{N}_{\hbar}^{[n|n]}(p_1, o_1; \dots; p_n, o_n) \\ &= \left(\sum_{r \geq 0} \frac{1}{r!} \sum_{\substack{h_j, n_j \geq 0, d_j \geq 1 \\ 2h_j - 2 + d_j + n_j > 0}} \hbar^{\sum_j 2h_j - 2 + d_j + n_j} \bigotimes_{j=1}^r \frac{n_{\hbar}^{[n|n]}}{(2i\pi)^{d_j} d_j! n_j!} \right) \vartheta_{\bullet}, \end{aligned}$$

and

$$n_{\hbar}^{[n|n]} = \int_{\bullet} \dots \int_{\bullet} \omega_{n_j}^{h_j, (d_j)} \cdot \nabla^{\otimes d_j}.$$

In this context,

$$\int_{\bullet} = \sum_{i=1}^n \int_{o_i}^{p_i},$$

and ϑ_{\bullet} stands for

$$\vartheta_{\bullet} = \vartheta\left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix}\right] \left(\xi_{\hbar} + \sum_{k=1}^n \mathbf{a}(p_k) - \mathbf{a}(o_k) \mid \tau \right).$$

Diagrammatic representation. In Appendix A, we explain that the formulae for the non perturbative partition function (equation (29)) and the $n|n$ kernels (equation (33)) can be represented as a sum over (maybe disconnected) diagrams. To a given order in \hbar , there is only a finite sum of allowed diagrams. With this formalism, it is easy to re-exponentiate the series above, i.e. to compute the asymptotic series for $\ln \mathfrak{T}_{\hbar}$ or $\ln \hat{\psi}_{\hbar}$: they can be written as sum over connected diagrams.

Special properties for A-spectral curves. When the symbol $\{e^u, e^v\}$ is 2ζ -torsion in $K_2(C(\mathcal{C}))$, the spectral curve satisfies the Boutroux condition and the quantization condition. So, when k is an integer going to infinity along arithmetic subsequences of step ζ and

$$\hbar = \frac{i\pi}{k},$$

the non-perturbative partition function and the $n|n$ -kernels do have an expansion in powers of \hbar , and besides, the factors

$$\exp\left(\frac{1}{\hbar} \int_o^p v \, du\right)$$

do not depend on the path from o to p . This follows from the discussion of §2.10, and especially from the fact that the argument of the theta functions are independent of \hbar along such subsequences. If one consider $\hbar \rightarrow 0$ in a generic way, or if the symbol $\{e^u, e^v\}$ is not torsion and ζ_{\hbar} is not zero for other reasons, the asymptotic expansion features fast oscillations at all orders in \hbar arising from the theta functions and their derivatives.

4.2. Remarks. Equation (29) for the non-perturbative partition function was first derived in [35] as a heuristic formula to compute the asymptotics of matrix integrals, $N = \hbar^{-1}$ playing the role of the matrix size. In [37] it was proved that it has order by order in powers of \hbar^{-1} a property of background independence, and that it transform like a theta function of characteristics $[\mu, \nu]$ under modular transformation. Actually, equation (29) is the result of summing all perturbative partition functions over filling fractions shifted by integer multiples of \hbar . This operation looks very much like the Whitham averaging in integrable systems, and we conjectured (and checked to the first non trivial order) in [13] that \mathcal{T}_{\hbar} is indeed a formal tau function of an integrable system whose times are moduli of the spectral curves. In that article, we also introduced a spinor version of the kernel $\psi_{\hbar}(p_1, p_2)$ (equation (30)), in order to build a wave function in the language of integrable systems. “Wave function” is a generic name for any complex-valued solution of a linear ODE’s or difference equation. $\Psi^{\alpha}(m) \equiv \psi_{\hbar}(p^{\alpha}(m), p_0)$ should be considered as the asymptotic series of a wave function, where p_0 is a point hold fixed in \mathcal{C}_0 . Different branches $p^{\alpha}(m)$ give rise to wave functions with dominant asymptotic behavior in different sectors. Typically, a wave function is a linear combination of $\Psi^{\alpha}(m)$, and thus its asymptotics is subject to the Stokes phenomenon when one goes from one sector to the other. The advantage of introducing the kernels is that the Stokes phenomenon is described by a single object $\psi_{\hbar}(p, p_0)$ with $p \in \mathcal{C}_0$, through the branching structure of the covering $m: \mathcal{C}_0 \rightarrow \hat{\mathcal{C}}$.

One can in principle derive the difference equation satisfied by $\psi_{\hbar}(p, p_0)$ order by order in \hbar , and we expect it to have an expansion in powers of \hbar , no matter if the spectral curve satisfy the Boutroux and the quantization condition. However, a general expression for the resummed difference operator annihilating the wave function (and its $n|n$ counterpart) just from the data of the spectral curve is not

available. Recently, Gukov and Sułkowski [53] have pointed out that, adding some assumption on the form of the answer, allows to reconstruct the full ODE or difference operator from the knowledge of the first orders. In particular in the context of hyperbolic geometry, the A-hat polynomial [45] is expected to appear as one of those operators (cf. §5.3). The A-hat polynomial is known in closed form for many knots, and Dimofte [28] has discussed a procedure to construct the \widehat{A} -polynomial from the A-polynomial. Those observations might give hints towards a general theory for the reconstruction of an exact integrable system whose tau function has precisely an asymptotic given to all order by equation (29) in the limit $\hbar \rightarrow 0$.

4.3. Rewriting in terms of modular quantities. It was proved in [37] that \mathcal{T}_{\hbar} has modular properties. Since the Bergman kernel B_0 is not modular invariant, the ω_n^h are not either modular. Similarly, although the theta function is modular, its derivative are not. So, the modular properties in the expression (29) are not manifest.

From equation (19), one sees that the deformed Bergman kernel B_{κ} is modular invariant if we have chosen $\kappa = \underline{\kappa}(\tau)$ as a function of τ which is quasimodular of weight 2, namely

$$\underline{\kappa}((a\tau + b)(c\tau + d)^{-1}) = c(c\tau + d)^t + (c\tau + d) \cdot \underline{\kappa}(\tau) \cdot (c\tau + d)^t.$$

If this is the case, it is straightforward to deduce from the topological recursions formula that $\omega_{n|\underline{\kappa}(\tau)}^h$ is modular invariant when $2h - 2 + n \geq 0$. It is often easier to compute modular objects than non-modular ones, so imagine that we have computed the $\omega_{n|\underline{\kappa}(\tau)}^h$. We would like to write \mathcal{T}_{\hbar} only in terms of $\omega_{n|\underline{\kappa}(\tau)}^h$. This can be done using equation (27) to express $\omega_n^h \equiv \omega_{n|\kappa=0}^h$ in terms of $\omega_{n|\kappa}^h$. The result (valid for any κ) is

$$\mathcal{T}_{\hbar} = \exp\left(\sum_{h \geq 0} \hbar^{2h-2} F_{h|\kappa}\right) \vartheta\left(\sum_{r \geq 0} \frac{1}{r!} \sum_{\substack{h_j \geq 0, d_j \geq 1 \\ 2h_j - 2 + d_j > 0}} \prod_{j=1}^r \frac{F_{h_j|\kappa}^{(d_j)} \cdot T_{d_j|\kappa}}{(2i\pi)^{d_j} d_j!}\right),$$

where

$$F_{h|\kappa}^{(d)} = \oint_{\mathcal{B}_{\kappa}} \dots \oint_{\mathcal{B}_{\kappa}} \omega_{d|\kappa}^h,$$

and

$$T_{d|\kappa} = \sum_{d'=0}^{\lfloor d/2 \rfloor} \frac{d!(-1)^{d'}(2i\pi)^{d'}}{2^{d'} d'!} \kappa^{\otimes d'} \otimes \frac{\nabla^{\otimes (d-2d')} \vartheta}{\vartheta}. \quad (34)$$

Certain linear combinations of derivatives of theta are modular, and the $T_{d|\underline{\kappa}(\tau)}$ precisely provide such combinations. In fact, the proof that $\mathcal{T}_{\mathfrak{h}}$ is modular given in [37] amounts to prove that $T_{d|\underline{\kappa}(\tau)}$ are modular. In the context of elliptic curves, we shall see in §6.1 that it is natural to choose $\underline{\kappa}(\tau)$ proportional to $E_2(\tau)$.

4.4. Effect of an involution. When the genus of the quotient \mathcal{C}_0/ι is zero, only the terms with even d_j remain in the partition function (equation (29)) and the kernel (equation (31)). In this paragraph, we assume it is the case. The conclusion of §3.5 was that only even order derivatives of theta functions appear in the formulae, since the odd order derivatives are contracted with zero. Then, we may trade $\nabla^{\otimes 2}$ for a derivative with respect to the period matrix:

$$\nabla^{\otimes 2} = 4i\pi \partial_\tau = D.$$

Besides, from Property 3.5 we learn that $\zeta_{\mathfrak{h}} = 0$ for any \mathfrak{h} . Then, the non-perturbative partition function and the non-perturbative $n|n$ kernels happen to be formal power series in \mathfrak{h} . And, in order to compute them, we only have to compute derivatives of Thetanullwerten with respect to the matrix of periods. For instance, the partition function reads

$$\mathcal{T}_{\mathfrak{h}} = \exp\left(\sum_{h \geq 0} \mathfrak{h}^{2h-2} F_h\right) \left(\sum_{r \geq 0} \frac{1}{r!} \sum_{\substack{h_j \geq 0, d'_j \geq 1 \\ 2h_j - 2 + 2d'_j > 0}} \mathfrak{h}^{\sum_j 2h_j - 2 + 2d'_j} \bigotimes_{j=1}^r \frac{F_{h_j}^{(2d'_j)} \cdot D^{\otimes d'_j}}{(2i\pi)^{2d'_j} (2d'_j)!}\right) \vartheta$$

On the other hand, if we compute the *perturbative partition function* with the Bergman kernel B_κ , we find, with help of equation (27),

$$\begin{aligned} Z_{\text{pert}, \mathfrak{h}|\kappa} &\equiv \exp\left(\sum_{h \geq 0} \mathfrak{h}^{2h-2} F_{h|\kappa}\right) \\ &= \exp\left(\sum_{h \geq 0} \mathfrak{h}^{2h-2} F_h\right) \left(\sum_{r \geq 0} \frac{1}{r!} \sum_{\substack{h_j \geq 0, d'_j \geq 1 \\ 2h_j - 2 + 2d'_j > 0}} \mathfrak{h}^{\sum_j 2h_j - 2 + 2d'_j} \prod_{j=1}^r \frac{F_{h_j}^{(2d'_j)} \cdot \kappa^{\otimes d'_j}}{d'_j! 2^{d'_j}}\right). \end{aligned}$$

This expression is very similar to the non-perturbative partition function computed with the Bergman kernel B_0 . More precisely,

$$\mathcal{T}_{\mathfrak{h}} = Z_{\text{pert}, \mathfrak{h}|\kappa}, \tag{35}$$

where

$$\frac{\kappa^{\otimes d'}}{2^{d'} (2i\pi)^{d'} d'!} \mapsto \frac{1}{(2i\pi)^{2d'} (2d')!} \frac{D^{\otimes d'} \vartheta}{\vartheta}.$$

The analogy carries at the level of the kernels. For instance, the *perturbative kernel* computed with B_κ is defined as

$$\psi_{\text{pert}, \mathfrak{h}|\kappa}(p_1, p_2) = \exp \left(\sum_{n \geq 1} \frac{1}{n!} \sum_{h \geq 0} \mathfrak{h}^{2h-2+n} \int_{p_2}^{p_1} \cdots \int_{p_2}^{p_1} \omega_{n|\kappa}^h \right),$$

and we observe that

$$\psi_{\mathfrak{h}}(p_1, p_2) = \frac{\vartheta \bullet}{\vartheta} \psi_{\text{pert}, \mathfrak{h}|\kappa}(p_1, p_2), \quad (36)$$

where

$$\frac{\kappa^{\otimes d'}}{2^{d'} (2i\pi)^{d'} d'!} \mapsto \frac{1}{(2i\pi)^{2d'} (2d')!} \frac{D^{\otimes d'} \vartheta}{\vartheta}.$$

Examples of knots for which $\iota_* = -\text{id}$ can be read off the tables in §B.3. For instance, it happens for the figure eight-knot and the manifold $L^2 R$. These two examples have been studied in [27], where it was proposed that asymptotics of the colored Jones polynomial could be computed from $\psi_{\text{pert}, \mathfrak{h}|\kappa}$, at the price of *ad hoc* renormalizations of $\kappa^{\otimes d'}$ to all orders. This phenomenon is explained by equations (35) and (36), and this explanation is verified on examples in Section 6.

5. Application to knot invariants

Our main conjecture is formulated in §5.4. We first explain the background of Chern–Simons theory and facts about volume conjectures, which allow a better understanding of the identification of parameters and of the complicated Stokes phenomenon when we consider functions in the variable u .

5.1. Generalities on Chern–Simons theory

With compact gauge group. The partition function of Chern–Simons theory of compact gauge group G (and corresponding Lie algebra \mathfrak{g}) in a closed 3-manifold $\overline{\mathfrak{M}}$ is formally the path integral over \mathfrak{g} -connections \mathcal{A} on $\overline{\mathfrak{M}}$, of the Chern–Simons action,

$$Z_{\text{CS}; G}(\overline{\mathfrak{M}}) = \int [\mathcal{D}\mathcal{A}] e^{-\frac{1}{\hbar} S_{\text{CS}}[\mathcal{A}]}, \quad S_{\text{CS}}[\mathcal{A}] = \int_{\overline{\mathfrak{M}}} \frac{1}{4} (\mathcal{A} \wedge d\mathcal{A} + \frac{2}{3} \mathcal{A} \wedge \mathcal{A} \wedge \mathcal{A}).$$

It depends on the Planck constant \hbar .

A way to define properly this integral is to choose a saddle point \mathcal{A}^{cl} of the action S_{CS} , and perform an expansion around \mathcal{A}_{cl} as usual in perturbative quantum field theory. By construction of Chern–Simons theory, the saddle points (also called “classical solutions”) are flat connections on $\overline{\mathcal{M}}$, i.e. those satisfying

$$d\mathcal{A}^{\text{cl}} + \mathcal{A}^{\text{cl}} \wedge \mathcal{A}^{\text{cl}} = 0.$$

However, there are in general many equivalence classes of flat connections, and one wishes the genuine partition function to be a sum over all classes of the perturbative partition functions, with some coefficients α_{cl}

$$Z_{\text{CS}; G}(\overline{\mathcal{M}}) = \sum_{\text{cl}} \alpha_{\text{cl}} Z_G^{\text{cl}}(\overline{\mathcal{M}}).$$

This sum is finite when \hbar assumes a value⁹ of the form

$$\hbar = \frac{i\pi}{K + H^\vee},$$

where K is an integer called *level* and H^\vee is the dual Coxeter number of G . There is actually a rigorous definition of the expectation values of Wilson loops for these values: the so-called Witten–Reshetikhin–Turaev invariants [83, 85].

With complex gauge group. The complexification comes in two steps. We shall be sketchy here and refer to [33] for details. Firstly, one considers a Chern–Simons theory with complex gauge group $G_{\mathbb{C}}$, whose Lie algebra is obtained from \mathfrak{g} by Weyl’s unitary trick, and with the new action $S_{\text{CS}}[\mathcal{A}] + S_{\text{CS}}[\mathcal{A}^*]$. The partition function then admits a decomposition in perturbative blocks, defined by expansion around a $\mathfrak{g}_{\mathbb{C}}$ -valued flat connexion,

$$Z_{G_{\mathbb{C}}} = \sum_{\text{cl}} \alpha_{\text{cl}} Z_{G_{\mathbb{C}}}^{\text{cl}} = \sum_{\text{cl}} \alpha_{\text{cl}} Z_G^{\text{cl}}(Z_G^{\text{cl}}) \quad (37)$$

The partition function is real when \hbar^* and \mathcal{A}^* are the complex conjugates of \hbar and \mathcal{A} . Secondly, at the level of the perturbative blocks, one consider a complexified version of the theory by assuming \mathcal{A} and \mathcal{A}^* independent $\mathfrak{g}_{\mathbb{C}}$ -valued connections. The blocks then have a factorization $Z_{G_{\mathbb{C}}}^{\text{cl}} = \Phi_G^{\text{cl}} \Phi_G^{*,\text{cl}}$. The Φ_G^{cl} are called *holomorphic blocks*, and they will play an important rôle in the following. By construction, they have an expansion in power series of \hbar , whose coefficients can be computed as well-defined sums over Feynman diagrams.

⁹ There exists two conventions for the Planck constant in Chern–Simons theory: either one puts \hbar [28] in the denominator in the exponential, or $2\hbar$ [30], [27]. We adopt the second convention, where $\hbar = 2i\pi/\text{integer}$.

Wilson loops and colored Jones polynomial. The most important observables in Chern–Simons theory are the expectation values of *Wilson loops*: given an oriented loop \mathfrak{K} in $\overline{\mathfrak{M}}$, and a representation R of G , they are defined as

$$\mathcal{W}_{G,R}(\mathfrak{K}, \hbar) = \left\langle \text{Tr}_R \text{P exp} \left(\oint_{\mathfrak{K}} \mathcal{A} \right) \right\rangle,$$

where P is the ordering operation along the loop. \mathcal{L} can be considered as a knot drawn in $\overline{\mathfrak{M}}$, and in fact the expectation value of the Wilson loops is a partition function for the knot complement $\overline{\mathfrak{M}} \setminus \mathfrak{K}$, where the classical solutions are now flat connections on $\overline{\mathfrak{M}} \setminus \mathfrak{K}$ with a meridian holonomy prescribed by R . To be precise, if $\rho = \frac{1}{2} \sum_{\alpha > 0} \alpha$ is the vector of Weyl’s constants, $\Lambda_R = (\lambda_j)_j$ is the highest weight associated to R , we must identify the holonomy eigenvalues to $e^{\hbar(\rho_j + \lambda_j)} = e^{u_j}$.

A foundational result is that the expectation values of Wilson loops define knot invariants. When $G = \text{SU}(2)$ and R is the spin $\frac{N-2}{2}$ representation, which has dimension N and is represented by the Young diagram

$$\lambda_{N-1} = \underbrace{\square \square \cdots \square}_{N-1 \text{ boxes}},$$

the expectation value of the Wilson loop is related to the colored Jones polynomial $J_N(\mathfrak{K}, q)$, with identifications

$$q = e^{2\hbar}, \quad N\hbar = u, \quad J_N(\mathfrak{K}, q) = \frac{\mathcal{W}_{\text{SU}(2), \lambda_{N-1}}(\mathfrak{K}, \hbar)}{\mathcal{W}_{\text{SU}(2), \lambda_{N-1}}(\bigcirc, \hbar)}. \quad (38)$$

The denominator accounts for the normalization of the Jones polynomial, which is 1 for the unknot in \mathbb{S}_3 , denoted \bigcirc . The expectation value of the Wilson loop of the unknot is itself given by

$$\mathcal{W}_{\text{SU}(2), \lambda_{N-1}}(\bigcirc, \hbar) = \frac{q^{N/2} - q^{-N/2}}{q^{1/2} - q^{-1/2}} = \frac{\text{sh } u}{\text{sh } \hbar}.$$

5.2. The volume conjectures. Initially, the Jones polynomial $J_2(\mathfrak{K}, q)$ has been defined in [55] and its colored version $J_N(\mathfrak{K}, q)$ in [83, 85], in the context of quantum groups. J_N is a Laurent polynomial in q with integer coefficients. The number $J_N(\mathfrak{K}, q = e^{2i\pi/N})$ is usually called the Kashaev invariant, and the original volume conjecture is

Conjecture 5.1 ([58]). *For any hyperbolic knot \mathfrak{K} in \mathbb{S}_3 ,*

$$\lim_{N \rightarrow \infty} \frac{2\pi}{N} \ln |J_N(\mathfrak{K}, q = e^{2i\pi/N})| = \text{Vol}(\mathbb{S}_3 \setminus \mathfrak{K}).$$

It was later enhanced by Gukov [51] to include hyperbolic deformations of $\mathbb{S}_3 \setminus \mathfrak{K}$, and subleading terms.

Conjecture 5.2 ([52]). *For any knot \mathfrak{K} in \mathbb{S}_3 , and u in an open domains $\mathcal{U}^{(\alpha)} \subseteq \mathbb{C}$, in the regime $N \rightarrow \infty$, $k \rightarrow \infty$ and $i\pi N/k = u$ kept fixed, the colored Jones has an asymptotic expansion of the form*

$$J_N(q = e^{2\hbar}, \mathfrak{K}) \sim \hbar^{\delta^{(\alpha)}/2} \exp\left(\sum_{\chi \geq -1} \hbar^\chi J_\chi^{(\alpha)}(u)\right) \quad \hbar = i\pi/k \quad (39)$$

- The leading order is a complexified volume

$$J_{-1}^{(\alpha)}(u) = \int^{p_u^{(\alpha)}} \ln l \, d \ln m,$$

where $p_u^{(\alpha)}$ is a point in some component of the A-polynomial such that

$$m(p_u^{(\alpha)}) = e^u$$

(see Remark 2.1 when \mathfrak{K} is not hyperbolic),

- $\delta^{(\alpha)}$ is an integer computed from cohomology, and $J_0^{(\alpha)}(u)$ is related to the Ray-Singer torsion.
- $J_\chi(u)$ for $\chi \geq 1$ are the coefficients in the \hbar -expansion of a certain holomorphic block $\Phi^{(\alpha)}(u)$ for $SL_2(\mathbb{C})$ Chern–Simons theory on \mathfrak{M} with boundary condition specified by u .

The statement about the leading order is called the *generalized volume conjecture* (GVC). The range of validity in u is not obvious, because of resonances and Stokes phenomena, that we attempt to describe in the next paragraph. The Kashaev invariant is retrieved for $u = i\pi$. We first recall two rigorous results about the leading order of the GVC. The first one is due to Garoufalidis and Lê.

Theorem 5.3. [47] *For any knot \mathfrak{K} , if u is nonnegative and small enough, then*

$$\lim_{\substack{N \rightarrow \infty \\ u = N\hbar}} \ln J_N(\mathfrak{K}, q = e^{2\hbar}) = 0,$$

i.e. the GVC holds with the choice of the abelian component.

The second is due to Murakami, who studied the figure-eight knot starting from a closed formula available in this case.

Theorem 5.4 ([71]). *For the figure-eight knot, the expression $\hbar \ln J_N(\mathbf{4}_1, q = e^{2\hbar})$ has the following behavior in the regime $N \rightarrow \infty$, $\hbar \rightarrow 0$ while $N\hbar = u$.*

- *When u is real and $|u| \leq \frac{1}{2} \ln\left(\frac{3+\sqrt{5}}{2}\right)$, or $u \in [0, \frac{i\pi}{6}[$, the lim is 0 (GVC for the abelian component).*
- *When $u = i\pi$ or $u \in]\frac{5i\pi}{6}, \frac{7i\pi}{6}[\notin i\pi\mathbb{Q}$, the lim is given by the GVC for the geometric component.*
- *When $u = \frac{i\pi P}{Q}$ with P, Q coprime integers and $P/Q \in]\frac{5}{6}, \frac{7}{6}[$, the lim is 0 when $N \rightarrow \infty$ along multiples of Q , whereas the lim is given by the GVC for the geometric component if $N \rightarrow \infty$ avoiding multiples of Q .*

We recall that the A-polynomial of the figure-eight knot has two components, one abelian ($l-1$) and one geometric, which intersect at $m^2 = -1$ and $m^2 = \frac{3\pm\sqrt{5}}{2}$. One recognizes in the latter a value of u at which a transition between components occur for the GVC to be valid according to Theorem 5.4. The example of the figure-eight is special in two ways. Firstly, its branchpoints are located at $m^2 = e^{\pm 2i\pi/3}$ and $m^2 = \frac{3\pm\sqrt{5}}{2}$, so two of them coincide with the intersection points. So, we do not see a change of branch within a single component at $\sqrt{\frac{3+\sqrt{5}}{2}}$ but actually a transition to the abelian component, and the behavior around the other branchpoints is beyond the range of validity of Theorem 5.4. Secondly, $\text{CS}_a(p)$ vanishes along the path from p_c to p in the geometric component such that

$$m^2(p) \in \left[\frac{3-\sqrt{5}}{2}, \frac{3+\sqrt{5}}{2} \right] \cup \left\{ e^{i\varphi} : \varphi \in \left[-\frac{2i\pi}{3}, \frac{2i\pi}{3} \right] \right\}.$$

so that Theorem 5.4 is only sensitive to the volume, not to the Chern–Simons part.

5.3. \widehat{A} -polynomial, AJ conjecture and Stokes phenomenon. Garoufalidis and Lê [46] showed that $J_N(\mathfrak{K}, q)$ always satisfy some recurrence relation on N (this was also anticipated in [42]). At the level of the analytic continuation, this turns into the existence of an operator $\widehat{\mathfrak{A}}_{\mathfrak{K}} \in \mathbb{Z}[e^{\frac{\hbar}{2}\partial_u}, e^u, e^{\hbar}]$ so that

$$\widehat{\mathfrak{A}}_{\mathfrak{K}} \cdot J_{u/\hbar}(\mathfrak{K}, q = e^{2\hbar}) = 0. \quad (40)$$

The AJ conjecture [45] states that the limit $\hbar \rightarrow 0$ of $\widehat{\mathfrak{A}}$ coincides with the A-polynomial of \mathfrak{K} up to a factor which is a polynomial in e^u , i.e. the A-polynomial is the semiclassical spectral curve associated to the difference equation equation (40). It has been proved recently in [63] for hyperbolic knots satisfying some technical assumptions and for which the A-polynomial has only a single irreducible factor apart from $(l-1)$.

If we treated equation (40) like an ODE, the leading asymptotic of the colored Jones when $\hbar \rightarrow 0$ would be given naively by a WKB analysis, namely,

$$J_{u/\hbar}(\mathfrak{K}, q = e^{2\hbar}) \sim \exp\left(\frac{1}{\hbar} \int^{p_u} \ln l \, d \ln m\right),$$

where l and m satisfies $\lim_{\hbar \rightarrow 0} \widehat{A}_{\hbar}(m, l) = 0$, and p_u is a point on this curve such that $\ln m(p_u) = u$. At the heuristic level, it explains the appearance of the complexified volume in the leading asymptotics of the colored Jones polynomial, by combining the AJ conjecture and Neumann–Zagier results reviewed in §2.4. Going a step further, we could imagine to introduce a infinite set of times and embed equation (40) (at least perturbatively in the new times) in a system of compatible ODE's, for which we know how to associate quantities satisfying loop equations [6, 12, 5]. Those loop equations have many solutions, and the non-perturbative topological recursion applied to the semiclassical spectral curve provide distinguished solutions as formal asymptotic series in \hbar [13]. This naive approach can be seen as a vague intuition why it is sensible to compare objects computed from the topological recursion to the asymptotics of solutions of the $\widehat{\mathfrak{A}}$ recursion relation, which we attempt to do in §5.4.

Difference equation are of discrete nature, and if we treat it like an ODE we may miss resonance phenomena, which here occur when q is a root of unity. On top of that, we have to take into account the usual Stokes phenomenon, hidden in the specification of the point p_u on the semiclassical spectral curve which projects to $\ln m(p_u) = u$. This choice comes in three part:

- to which component $\mathcal{C}_0^{(\alpha)}$ of the A-polynomial should p_u belong?
- in which sheet of the covering $m^2: \mathcal{C}_0^{(\alpha)} \rightarrow \mathbb{C}$ should p_u belong?
- which determination of the logarithms in $J_{-1}(u)$ should be chosen?

We call the data of such a triple

$$\mathcal{T}^{(\alpha)} = (\mathcal{C}_0^{(\alpha)}, p^{(\alpha)}, \ln)$$

a *determination*. Although we can consider the right-hand side intrinsically as a function of a point p in the universal covering of $\bigcup_{\alpha} \mathcal{C}_0^{(\alpha)}$, it is a non trivial issue to predict for which determination it can be matched to the asymptotics of the left-hand side which is a function of u . The transition between different determinations occurs across *Stokes curves* in the u -complex plane. Although the values of u at which several components intersect, and branchcut structures of the coverings $m^2: \mathcal{C}_0 \rightarrow \mathbb{C}$ represented in the u -plane obviously play a role, we do not know

of a unambiguous algorithm which would give, for any knot, the determination corresponding to each domain and the correct pattern of Stokes curves which separate them. For second order differential equation (i.e. for semiclassical spectral curves having a single component, the form $y^2 = \text{Pol}(x)$), the algorithm yielding the Stokes curves is known [8], but it is not obvious to generalize this construction to curves of the form $\text{Pol}(e^x, e^y) = 0$ and having several components. The only reliable facts are that, for hyperbolic knots, one has to choose:

- for u close to $i\pi$: the determination corresponding to the geometric branch of the geometric component (see §2.3). We call it the *geometric determination*;
- and for u nonnegative and close to 0, the determination corresponding to the abelian component, so that $J_{-1}(u) \equiv 0$.

5.4. Main conjectures. Let \mathfrak{M} is a hyperbolic 3-manifold with 1-cusp, and let us consider an A-spectral curve (\mathcal{C}_0, u, v) coming from an irreducible component of the A-polynomial of \mathfrak{M} . We would like to consider the asymptotics series constructed from the 2|2-kernel introduced in §4.1,

$$\mathcal{J}_{\hbar}^{\text{n.p.TR}}(p) = (\psi_{\hbar}^{[2|2]}(p, o; \iota(p), \iota(o)))^{1/2},$$

depending on a choice of basepoint o and a characteristics $\mu, \nu \in \mathbb{C}^g$. We identified the formal parameter $\hat{\hbar}$ to \hbar . We recall that ι is the involution

$$(m, l) \longrightarrow \left(\frac{1}{m}, \frac{1}{l} \right)$$

defined on \mathcal{C}_0 . Let us recapitulate its properties.

- $\mathcal{J}_{\hbar}^{\text{n.p.TR}}(p)$ is defined as a formal asymptotic series

$$\mathcal{J}_{\hbar}^{\text{n.p.TR}}(p) = \exp \left(\sum_{\chi \geq -1} \hbar^{\chi} \tilde{J}_{\chi}(p) \right).$$

- The leading order is the complexified volume up to a constant

$$\tilde{J}_{-1}(p) = \frac{1}{2} \int_{o, \iota(o)}^{p, \iota(p)} v \, du = \int_o^p v \, du = \frac{i}{2} (\text{Vol}_a(p) + i \text{CS}_a(p)).$$

- For any $\chi \geq 1$, $\tilde{J}_{\chi}(p)$ is a meromorphic function of $p \in \mathcal{C}_0$, which is either independent of \hbar , or is a function of \hbar which does not have a power series expansion when $\hbar \rightarrow 0$. We give in §5.5 its expression up to $\chi = 3$.

- If 2ζ is the order of torsion of the symbol $\{m, l\}$ in $K_2(\mathbb{C}(\mathbb{C}_0))$, for any $\chi \geq 1$, $\tilde{j}_\chi(p)$, seen as a function of \hbar , assumes a constant value on the subsequences $\hbar = \frac{i\pi}{k}$ where k is a integer with fixed congruence modulo ζ .
- When $\iota_* = -\text{id}$, for any $\chi \geq 1$, $\tilde{j}_\chi(p)$ is independent of \hbar .

Conjecture 5.5. *There exists a choice of o and μ, ν such that $\mathcal{J}_\hbar^{\text{n.p. TR}}(p)$ is annihilated by the $\widehat{\mathfrak{A}}$ -operator.*

We also attempt to formulate a stronger version of the conjecture to identify this series with the all-order asymptotics of the colored Jones polynomial.

Conjecture 5.6. *If \mathfrak{M} is the complement of a hyperbolic knot, with a choice of determination as in the GVC (Conjecture 5.2) and keeping the same notations, we have the all-order asymptotic expansion*

$$J_N(\mathfrak{K}, q = e^{2\hbar}) \sim C_\hbar B(u) \mathcal{J}_\hbar^{\text{n.p. TR}}(p_u^{(\alpha)})$$

for a constant C_\hbar independent of u , and a prefactor $B(u)$ independent of \hbar . In other words, for any $\chi \geq 1$, the $J_\chi^{(\alpha)}(u)$ of equation (39) coincide with $\tilde{j}_\chi(p^{(\alpha)}(u))$ up to a constant independent of \hbar and u .

5.5. First few terms. In the comparison to the colored Jones polynomial, there is always an issue of normalization, which is reflected in the prefactors C_\hbar and $B(u)$ that we do not attempt to predict. Thus, the definition of \tilde{j}_0 is irrelevant here, and we refer to [26] for some discussion on the computation of the constant term $j_0(u)$ in the GVC in terms of algebraic geometry on the A-polynomial.

We now write down the general expression for $\tilde{j}_\chi(p)$ for $\chi = 1, 2, 3$, in terms of modular quantities (see Section 4 and in particular §4.3). We first introduce the $[l, 0]$ tensors

$$G_{n|k}^{h,(d)}(p) = \frac{1}{n!} \frac{1}{(2i\pi)^d d!} \underbrace{\int_{o,\iota(o)}^{p,\iota(p)} \cdots \int_{o,\iota(o)}^{p,\iota(p)}}_{n \text{ times}} \underbrace{\oint_{\mathcal{B}_\kappa} \cdots \oint_{\mathcal{B}_\kappa}}_{d \text{ times}} \omega_{n+d|k}^h.$$

and $G_{n|k}^h(p) \equiv G_{n|k}^{h,(0)}(p)$. Then, we have

$$\begin{aligned} 2\tilde{j}_1(p) &= G_{1|\underline{k}(\tau)}^{1,(0)}(p) + G_{3|\underline{k}(\tau)}^{0,(0)}(p) + T_{1,\bullet|\underline{k}(\tau)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) \\ &\quad + T_{2,\bullet|\underline{k}(\tau)} G_{1|\underline{k}(\tau)}^{0,(2)}(p) + (T_{3,\bullet|\underline{k}(\tau)} - T_{3|\underline{k}(\tau)}) G_{0|\underline{k}(\tau)}^{0,(3)}, \end{aligned}$$

$$\begin{aligned}
2 \tilde{j}_2(p) = & G_{2|\underline{k}(\tau)}^{1,(0)}(p) + T_{1,\bullet|\underline{k}(\tau)} G_{1|\underline{k}(\tau)}^{1,(1)}(p) + (T_{2,\bullet|\underline{k}(\tau)} - T_{2|\underline{k}(\tau)}) G_{0|\underline{k}(\tau)}^{1,(2)} \\
& + G_{4|\underline{k}(\tau)}^{0,(0)}(p) + T_{1,\bullet|\underline{k}(\tau)} G_{3|\underline{k}(\tau)}^{0,(1)}(p) + T_{2,\bullet|\underline{k}(\tau)} G_{2|\underline{k}(\tau)}^{0,(2)}(p) \\
& + T_{3,\bullet|\underline{k}(\tau)} G_{1|\underline{k}(\tau)}^{0,(3)}(p) + (T_{4,\bullet|\underline{k}(\tau)} - T_{4|\underline{k}(\tau)}) G_{0|\underline{k}(\tau)}^{0,(4)} \\
& + \frac{1}{2} (V_{\bullet|\underline{k}(\tau)}^{(1,1)} - T_{1|\underline{k}(\tau)}^2) (G_{0|\underline{k}(\tau)}^{1,(1)})^2 + V_{\bullet|\underline{k}(\tau)}^{(1,1)} G_{0|\underline{k}(\tau)}^{0,(1)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) \\
& + V_{\bullet|\underline{k}(\tau)}^{(1,2)} G_{0|\underline{k}(\tau)}^{0,(1)} G_{1|\underline{k}(\tau)}^{0,(2)}(p) + (V_{\bullet|\underline{k}(\tau)}^{(1,3)} - T_{1|\underline{k}(\tau)} T_{3|\underline{k}(\tau)}) G_{0|\underline{k}(\tau)}^{1,(1)} G_{0|\underline{k}(\tau)}^{0,(3)} \\
& + \frac{1}{2} V_{\bullet|\underline{k}(\tau)}^{(1,1)} (G_{2|\underline{k}(\tau)}^{0,(1)}(p))^2 + V_{\bullet|\underline{k}(\tau)}^{(1,2)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) G_{1|\underline{k}(\tau)}^{0,(2)}(p) \\
& + V_{\bullet|\underline{k}(\tau)}^{(1,3)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) G_{0|\underline{k}(\tau)}^{0,(3)} + \frac{1}{2} V_{\bullet|\underline{k}(\tau)}^{(2,2)} (G_{1|\underline{k}(\tau)}^{0,(2)}(p))^2 \\
& + V_{\bullet|\underline{k}(\tau)}^{(2,3)} G_{0|\underline{k}(\tau)}^{0,(2)}(p) G_{0|\underline{k}(\tau)}^{0,(3)} + \frac{1}{2} (V_{\bullet|\underline{k}(\tau)}^{(3,3)} - T_{3|\underline{k}(\tau)}^2) (G_{0|\underline{k}(\tau)}^{0,(3)})^2,
\end{aligned}$$

$$\begin{aligned}
2 \tilde{j}_3(p) = & G_{1|\underline{k}(\tau)}^{2,(0)}(p) + (T_{1,\bullet|\underline{k}(\tau)} - T_{1|\underline{k}(\tau)}) G_{0|\underline{k}(\tau)}^{2,(1)} + G_{3|\underline{k}(\tau)}^{1,(0)}(p) \\
& + T_{1,\bullet|\underline{k}(\tau)} G_{2|\underline{k}(\tau)}^{1,(1)}(p) + T_{2,\bullet|\underline{k}(\tau)} G_{1|\underline{k}(\tau)}^{1,(2)}(p) + (T_{3,\bullet|\underline{k}(\tau)} - T_{3|\underline{k}(\tau)}) G_{0|\underline{k}(\tau)}^{1,(3)} \\
& + G_{5|\underline{k}(\tau)}^{0,(0)}(p) + T_{1,\bullet|\underline{k}(\tau)} G_{4|\underline{k}(\tau)}^{0,(1)} + T_{2,\bullet|\underline{k}(\tau)} G_{3|\underline{k}(\tau)}^{0,(2)}(p) \\
& + T_{3,\bullet|\underline{k}(\tau)} G_{2|\underline{k}(\tau)}^{0,(3)}(p) + T_{4,\bullet|\underline{k}(\tau)} G_{1|\underline{k}(\tau)}^{0,(4)}(p) \\
& + (T_{5,\bullet|\underline{k}(\tau)} - T_{5|\underline{k}(\tau)}) G_{0|\underline{k}(\tau)}^{0,(5)} + V_{\bullet|\underline{k}(\tau)}^{(1,1)} G_{0|\underline{k}(\tau)}^{1,(1)} G_{1|\underline{k}(\tau)}^{1,(1)}(p) \\
& + V_{\bullet|\underline{k}(\tau)}^{(1,2)} G_{0|\underline{k}(\tau)}^{1,(1)} G_{0|\underline{k}(\tau)}^{1,(2)} + V_{\bullet|\underline{k}(\tau)}^{(1,2)} G_{0|\underline{k}(\tau)}^{1,(1)} G_{3|\underline{k}(\tau)}^{0,(1)}(p) \\
& + V_{\bullet|\underline{k}(\tau)}^{(1,2)} G_{0|\underline{k}(\tau)}^{1,(1)} G_{2|\underline{k}(\tau)}^{0,(2)}(p) + V_{\bullet|\underline{k}(\tau)}^{(1,3)} G_{0|\underline{k}(\tau)}^{1,(1)} G_{1|\underline{k}(\tau)}^{0,(3)}(p) \\
& + (V_{\bullet|\underline{k}(\tau)}^{(1,4)} - T_{1|\underline{k}(\tau)} T_{4|\underline{k}(\tau)}) G_{0|\underline{k}(\tau)}^{1,(1)} G_{0|\underline{k}(\tau)}^{0,(4)} + V_{\bullet|\underline{k}(\tau)}^{(1,1)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) G_{1|\underline{k}(\tau)}^{1,(1)}(p) \\
& + V_{\bullet|\underline{k}(\tau)}^{(1,1)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) G_{0|\underline{k}(\tau)}^{1,(2)} + V_{\bullet|\underline{k}(\tau)}^{(1,1)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) G_{3|\underline{k}(\tau)}^{0,(1)}(p) \\
& + V_{\bullet|\underline{k}(\tau)}^{(1,2)} G_{2|\underline{k}(\tau)}^{0,(1)} G_{2|\underline{k}(\tau)}^{0,(2)}(p) + V_{\bullet|\underline{k}(\tau)}^{(1,3)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) G_{1|\underline{k}(\tau)}^{0,(3)}(p) \\
& + (V_{\bullet|\underline{k}(\tau)}^{(1,4)} - T_{1|\underline{k}(\tau)} T_{4|\underline{k}(\tau)}) G_{2|\underline{k}(\tau)}^{0,(1)}(p) G_{0|\underline{k}(\tau)}^{0,(4)} \\
& + V_{\bullet|\underline{k}(\tau)}^{(2,1)} G_{1|\underline{k}(\tau)}^{0,(2)}(p) G_{1|\underline{k}(\tau)}^{1,(1)}(p) + V_{\bullet|\underline{k}(\tau)}^{(2,2)} G_{1|\underline{k}(\tau)}^{0,(2)}(p) G_{0|\underline{k}(\tau)}^{1,(2)} \\
& + V_{\bullet|\underline{k}(\tau)}^{(2,1)} G_{1|\underline{k}(\tau)}^{0,(2)}(p) G_{3|\underline{k}(\tau)}^{0,(1)}(p) + V_{\bullet|\underline{k}(\tau)}^{(2,2)} G_{1|\underline{k}(\tau)}^{0,(2)}(p) G_{2|\underline{k}(\tau)}^{0,(2)}(p)
\end{aligned}$$

$$\begin{aligned}
& + V_{\bullet|\underline{k}(\tau)}^{(2,3)} G_{1|\underline{k}(\tau)}^{0,(2)}(p) G_{1|\underline{k}(\tau)}^{0,(3)}(p) + V_{\bullet|\underline{k}(\tau)}^{(2,4)} G_{1|\underline{k}(\tau)}^{0,(2)}(p) G_{0|\underline{k}(\tau)}^{0,(4)} \\
& + V_{\bullet|\underline{k}(\tau)}^{(3,1)} G_{0|\underline{k}(\tau)}^{0,(3)} G_{1|\underline{k}(\tau)}^{1,(1)}(p) + (V_{\bullet|\underline{k}(\tau)}^{(3,2)} - T_{3|\underline{k}(\tau)} T_{2|\underline{k}(\tau)}) G_{0|\underline{k}(\tau)}^{0,(3)} G_{0|\underline{k}(\tau)}^{1,(2)} \\
& + V_{\bullet|\underline{k}(\tau)}^{(3,1)} G_{0|\underline{k}(\tau)}^{0,(3)} G_{3|\underline{k}(\tau)}^{0,(1)}(p) + V_{\bullet|\underline{k}(\tau)}^{(3,2)} G_{0|\underline{k}(\tau)}^{0,(3)} G_{2|\underline{k}(\tau)}^{0,(2)}(p) \\
& + V_{\bullet|\underline{k}(\tau)}^{(3,3)} G_{0|\underline{k}(\tau)}^{0,(3)} G_{1|\underline{k}(\tau)}^{0,(3)}(p) + (V_{\bullet|\underline{k}(\tau)}^{(3,4)} - T_{3|\underline{k}(\tau)} T_{4|\underline{k}(\tau)}) G_{0|\underline{k}(\tau)}^{0,(3)} G_{0|\underline{k}(\tau)}^{0,(4)} \\
& + \frac{1}{6} (V_{\bullet|\underline{k}(\tau)}^{(1,1,1)} - T_{1|\underline{k}(\tau)}^3) (G_{0|\underline{k}(\tau)}^{1,(1)})^3 + \frac{1}{2} V_{\bullet|\underline{k}(\tau)}^{(1,1,2)} (G_{0|\underline{k}(\tau)}^{1,(1)})^2 G_{2|\underline{k}(\tau)}^{0,(1)}(p) \\
& + \frac{1}{2} V_{\bullet|\underline{k}(\tau)}^{(1,1,2)} (G_{0|\underline{k}(\tau)}^{1,(1)})^2 G_{1|\underline{k}(\tau)}^{0,(2)}(p) \\
& + \frac{1}{2} (V_{\bullet|\underline{k}(\tau)}^{(1,1,3)} - T_{1|\underline{k}(\tau)}^2 T_{3|\underline{k}(\tau)}) (G_{0|\underline{k}(\tau)}^{1,(1)})^2 G_{0|\underline{k}(\tau)}^{0,(3)} \\
& + V_{\bullet|\underline{k}(\tau)}^{(1,1,2)} G_{0|\underline{k}(\tau)}^{1,(1)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) G_{1|\underline{k}(\tau)}^{0,(2)}(p) \\
& + V_{\bullet|\underline{k}(\tau)}^{(1,1,3)} G_{0|\underline{k}(\tau)}^{1,(1)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) G_{0|\underline{k}(\tau)}^{0,(3)} + \frac{1}{2} V_{\bullet|\underline{k}(\tau)}^{(1,2,2)} G_{0|\underline{k}(\tau)}^{1,(1)} (G_{1|\underline{k}(\tau)}^{0,(2)}(p))^2 \\
& + V_{\bullet|\underline{k}(\tau)}^{(1,2,3)} G_{0|\underline{k}(\tau)}^{1,(1)} G_{1|\underline{k}(\tau)}^{0,(2)}(p) G_{0|\underline{k}(\tau)}^{0,(3)} \\
& + \frac{1}{2} (V_{\bullet|\underline{k}(\tau)}^{(1,3,3)} - T_{1|\underline{k}(\tau)} T_{3|\underline{k}(\tau)}^2) G_{0|\underline{k}(\tau)}^{1,(1)} (G_{0|\underline{k}(\tau)}^{0,(3)})^2 \\
& + \frac{1}{6} V_{\bullet|\underline{k}(\tau)}^{(1,1,1)} (G_{2|\underline{k}(\tau)}^{0,(1)}(p))^3 + \frac{1}{2} V_{\bullet|\underline{k}(\tau)}^{(1,1,2)} (G_{2|\underline{k}(\tau)}^{0,(1)}(p))^2 G_{1|\underline{k}(\tau)}^{0,(2)}(p) \\
& + \frac{1}{2} V_{\bullet|\underline{k}(\tau)}^{(1,1,3)} (G_{2|\underline{k}(\tau)}^{0,(1)}(p))^2 G_{0|\underline{k}(\tau)}^{0,(3)} + \frac{1}{2} V_{\bullet|\underline{k}(\tau)}^{(1,2,2)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) (G_{1|\underline{k}(\tau)}^{0,(2)}(p))^2 \\
& + V_{\bullet|\underline{k}(\tau)}^{(1,2,3)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) G_{1|\underline{k}(\tau)}^{0,(2)}(p) G_{0|\underline{k}(\tau)}^{0,(3)} \\
& + \frac{1}{2} V_{\bullet|\underline{k}(\tau)}^{(1,3,3)} G_{2|\underline{k}(\tau)}^{0,(1)}(p) (G_{0|\underline{k}(\tau)}^{0,(3)})^2 \\
& + \frac{1}{6} V_{\bullet|\underline{k}(\tau)}^{(2,2,2)} (G_{1|\underline{k}(\tau)}^{0,(2)}(p))^3 + \frac{1}{6} V_{\bullet|\underline{k}(\tau)}^{(2,3,3)} (G_{1|\underline{k}(\tau)}^{0,(2)}(p))^2 G_{0|\underline{k}(\tau)}^{0,(3)} \\
& + \frac{1}{2} V_{\bullet|\underline{k}(\tau)}^{(2,3,3)} G_{1|\underline{k}(\tau)}^{0,(2)}(p) (G_{0|\underline{k}(\tau)}^{0,(3)})^2 + \frac{1}{6} (V_{\bullet|\underline{k}(\tau)}^{(3,3,3)} - T_{3|\underline{k}(\tau)}^3) (G_{0|\underline{k}(\tau)}^{0,(3)})^3.
\end{aligned}$$

In the formulae above, the $G_{l|\underline{k}(\tau)}^{h,(d)}$ are contracted (from left to right) with the tensors $T_{d,\bullet|\underline{k}(\tau)}$ which were defined in equations (34) and their combinations:

$$V_{2,\bullet|\underline{k}(\tau)}^{(d_1,d_2)} = T_{d_1+d_2,\bullet|\underline{k}(\tau)} - T_{d_1,\bullet|\underline{k}(\tau)} T_{d_2,\bullet|\underline{k}(\tau)}$$

and

$$\begin{aligned}
V_{2, \bullet | \underline{k}(\tau)}^{(d_1, d_2, d_3)} &= T_{d_1 + d_2 + d_3, \bullet | \underline{k}(\tau)} \\
&\quad - (T_{d_1, \bullet | \underline{k}(\tau)} T_{d_2 + d_3, \bullet | \underline{k}(\tau)} + T_{d_2, \bullet | \underline{k}(\tau)} T_{d_1 + d_3, \bullet | \underline{k}(\tau)} \\
&\quad \quad + T_{d_3, \bullet | \underline{k}(\tau)} T_{d_1 + d_2, \bullet | \underline{k}(\tau)}) \\
&\quad + 2T_{d_1, \bullet | \underline{k}(\tau)} T_{d_2, \bullet | \underline{k}(\tau)} T_{d_3, \bullet | \underline{k}(\tau)}.
\end{aligned}$$

They are combination of derivatives of theta functions evaluated at

$$\mathbf{w}_\bullet = \mathbf{a}(p) - \mathbf{a}(o) + \mathbf{a}(\iota(p)) - \mathbf{a}(\iota(o)) + \zeta_\hbar,$$

and the constant ζ is defined in equation (28). When $\iota_* = -\text{id}$, several simplifications occur: the gray terms vanish; $\mathbf{a}(p) - \mathbf{a}(o) + \mathbf{a}(\iota(p)) - \mathbf{a}(\iota(o)) = 0$ and $\zeta_\hbar = 0$, the argument of the ϑ and ϑ_\bullet is always zero, i.e. $\mathbf{w}_\bullet = \mathbf{w} = 0$. In particular, this implies that, for any $\chi \geq 1$, $\tilde{j}_\chi(p)$ is independent of \hbar .

5.6. Comments. We check that Conjecture 5.6 holds for the figure-eight knot in §6.4 up to $o(\hbar^3)$, with o chosen (in a certain sense, as explained later) at a branchpoint, $[\mu, \nu]$ the unique half-integer characteristics with reality properties, and $C_\hbar \equiv 1 + o(\hbar^3)$. This very natural choice of the normalization allows to retrieve the asymptotic expansion of the Kashaev invariant $J_N(\mathbf{4}_1, q = e^{2i\pi/N})$ when $N \rightarrow \infty$, by specializing to $u = i\pi$ and taking the geometric determination.

For the once punctured torus bundle L^2R (a knot complement in lens space), we check in §6.5 up to $o(\hbar^2)$ that

$$\mathcal{J}_\hbar^H(u) \sim C_\hbar B(u) \mathcal{J}_\hbar^{\text{n.p. TR}}(p^{(\omega)}(u)),$$

where $\mathcal{J}_\hbar^H(u)$ is a Hikami-type integral associated to L^2R , and for the RHS, o is chosen at a branchpoint, $[\mu, \nu]$ is the unique half-integer characteristics with reality properties, and we choose the geometric determination. However, the normalization is now

$$C_\hbar = 1 + \frac{\hbar^2}{32} + o(\hbar^2).$$

The free parameters in our conjecture are the basepoint o for computing primitives, and the characteristics $[\mu, \nu]$ of the theta functions. Note that different choices of o affects $\tilde{j}_\chi(p)$ in a non trivial way, since it contains products of primitives. We have not found a general rule to specify neither o , nor μ, ν , and the choices might be also subjected to Stokes jumps regarding the identification to asymptotics of the Jones polynomial. In the examples treated in Section 6, the curve has $g = 1$ and we find natural choices for them. In general, we think that

it must be chosen among even half-integer characteristics, so $\binom{2g+1}{g}$ possibilities are left. Recall that, for hyperelliptic curves, they are in bijection with partitions of the $2g + 2$ Weierstraß points in two sets of $g + 1$ elements. For A-spectral curves of §B.1 that we found to be hyperelliptic,¹⁰ it turned out that they can be represented after birational transformations with rational coefficients $(m, l) \mapsto (X, Y)$, in the form

$$Y^2 = S_1(X)S_2(X),$$

where S_1 and S_2 are polynomials with integer coefficients and of the same degree $g + 1$, hence providing a canonical choice of even half-characteristics, for which $(\vartheta \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right] (0))^8$ computed by Thomae's formula is an integer. This suggests that a deeper study of the $\mathrm{SL}_2(\mathbb{C})$ character variety could entirely fix the appropriate choice of μ, ν .

In such a conjecture, it is natural to identify the Planck constant \hbar of Chern–Simons theory with the parameter \mathfrak{h} of the non-perturbative partition functions of Section 4, since special properties arise on each side when \mathfrak{h} and \hbar assume values of the form $i\pi/\text{integer}$. In the framework of Chern–Simons theory, the expectation value of Wilson loop can be thought as a wave function, hence it is natural to compare them to kernels. The 2-kernel $\psi_{\mathfrak{h}}^{[2|2]}(p_1, p'_1; p_2, p'_2)$ is symmetric by exchange of (p_1, p'_1) with (p_2, p'_2) , so the right hand side is invariant under the involution ι , which is also a property of the holomorphic blocks. We attempt to motivate¹¹ further the precise form of the conjecture in Section 7. We shall see that, for torus knots, $\psi_{\hbar}^{[2|2]}$ *without the power 1/2* appears heuristically in the computation of the colored Jones polynomial. For torus knots, it is known [51, Appendix B] that the Chern–Simons partition function $Z_{\mathrm{SL}_2(\mathbb{C})}$ coincide with $Z_{\mathrm{SU}(2)}$ up to a simple factor. For hyperbolic knots, we rather have equation (37), which incite to identify the holomorphic block with the analytic continuation of $\sqrt{Z_{\mathrm{SL}_2(\mathbb{C})}}$. This may account for the power 1/2 in Conjecture 5.6.

6. Examples

From the point of view adopted in this article, the complexity of hyperbolic 3-manifolds with 1-cusp is measured by the complexity of the algebraic curve defined

¹⁰ On top of curves of genus $g = 1, 2$ which are necessarily hyperelliptic, we found that all genus 3 curves of §B.1 are hyperelliptic, as well as $\mathbf{7}_2, \mathbf{9}_{10}^{(1)}, \mathbf{10}_{146}^{(2)}$ (genus 4), $\mathbf{8}_1$ (genus 5) and $\mathbf{9}_2$ (genus 7). This list is not exhaustive within §B.1, because we could not obtain an answer from MAPLE in reasonable time for curves of high degree.

¹¹ Dijkgraaf, Fuji and Manabe [26] also provided topological string arguments for the identification of parameters in equation (38) and the role of ι .

by the geometric component $\mathcal{C}_0^{\text{geom}}$ of its A-polynomial: to compute $\mathcal{J}_h^{\text{n.p.TR}}(p)$, we need to compute explicitly meromorphic forms (and their primitives) on the curve, as well as values of theta functions and their derivatives. From the tables of A-polynomials of Culler [24, 25], we collected the genus of the A-polynomial components of various knots in §B.1.

The simplest non trivial class of manifolds correspond to those for which $\mathcal{C}_0^{\text{geom}}$ is a genus 1 curve, i.e. an elliptic curve. This happens for the geometric components of the figure 8-knot and the manifold L^2R . The theta and theta derivatives values can be computed in a simple and efficient way thanks to the theory of modular forms (Section 6.1).

The next simplest class corresponds to manifolds for which $\mathcal{C}_0^{\text{geom}}$ is hyperelliptic. In this case there are uniform expressions for a Bergman kernel in terms of the coordinates m^2 and l , and the theta values are well-known in terms of the coordinates of Weierstrass points. For curves of genus $g \geq 2$, in principle, the values of theta derivatives can be related to the theta values via the theory of Siegel modular forms and the work of [10]. The 5_2 knot and the Pretzel $(-2, 3, 7)$ give rise to A-polynomial with a single component, of genus 2 thus hyperelliptic. We leave to a future work explicit computations for A-spectral curves of genus 2 and comparison to the perturbative invariants obtained by other methods.

We observe many times that some components of the A-polynomial of different knots are either the same, or birationally equivalent. For instance, the A-polynomial of the 5_2 and the Pretzel $(-2, 3, 7)$ are birationally equivalent, and one of the two factors of the A-polynomial of the 7_4 coincides with the A-polynomial of the 4_1 . This remark has some interest because values of theta derivatives, which provide the corrective terms to be added to the topological recursion for comparison with the asymptotics of the colored Jones polynomial, only depend on the isomorphism class of \mathcal{C}_0 as a Riemann surface, i.e. only depend on $A(m, l)$ up to birational equivalence.

Since \mathcal{C} is a singular curve, we do not expect a naive inequality between the degree of A or of the invariant trace field (which contains the cusp field), and the genus g . We observe that

- g looks experimentally much lower than the genus of a generic smooth curve of the same degree as the A polynomial and
- the genus of the quotient \mathcal{C}/ι also drops compared to the genus of \mathcal{C} .

It would be interesting to have a interpretation of g as well as those two observations from the point of view of representation theory. In the same vein, we can ask other open questions, here focused on geometric components.

Problem 6.1. *Describe the set of elliptic curves over \mathbb{Q} which are obtained as geometric components of hyperbolic 3-manifolds. Do all elliptic curves arise in that way?*

Problem 6.2. *Characterize the hyperbolic 3-manifolds so that the quotient $\mathcal{C}^{\text{geom}}/\iota$ has genus 0.*

Problem 6.3. *For a given genus g , do an infinite number of non-isomorphic curves of genus g arise as geometric components of a hyperbolic 3-manifold?*

These problems are already interesting in one replaces “geometric component” by the class of “A-spectral curves” defined in equation (2.8).

6.1. Thetanullwerten for elliptic curves. In this section, we give a self-contained presentation to compute the theta functions and their derivatives appearing in Section 4 and the computation of $\mathcal{J}_h^{\text{n.p.TR}}(p)$ for a genus 1 spectral curve. For more details about elliptic modular forms, the reader may consult the recent textbook [18, Chapter 1].

6.1.1. Modular forms and their derivatives. Elliptic curves are characterized by the orbit of their period τ in the upper-half plane \mathbb{H} under the modular group $\text{SL}_2(\mathbb{Z})$. A modular form of weight k for a subgroup $\Gamma \subseteq \text{SL}_2(\mathbb{Z})$ is a by definition a holomorphic function $f : \mathbb{H} \rightarrow \mathbb{C}$ such that $f(\tau) = O(1)$ when $q = e^{2i\pi\tau} \rightarrow 0$, and satisfying

$$f\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^k f(\tau) \quad \text{for all } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma. \quad (41)$$

When the subgroup is not precised, it is understood that equation (41) holds for the full modular group. Obviously, modular forms are 1-periodic functions, so have a Fourier expansion

$$f(\tau) = \sum_{n \geq 0} a_n q^n, \quad q = e^{2i\pi\tau},$$

where only nonnegative indices appear owing to the growth condition when $q \rightarrow 0$. The Eisenstein series

$$E_{2l}(\tau) = \frac{1}{2\zeta(2l)} \sum_{(n,m) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{1}{(n + m\tau)^{2l}}, \quad \text{for all } l \geq 1,$$

provide important examples of modular forms of weight $2l$ when $l \geq 2$. The zeta value in the denominator enforces the normalization $E_{2l}(\tau) = 1 + O(q)$ when $q \rightarrow 0$. We find convenient to absorb a factor of π per unit weight, and introduce non standard notations

$$\tilde{E}_{2l} = \pi^{2l} E_{2l}.$$

It is well known that the ring of modular forms is generated by \tilde{E}_4 and \tilde{E}_6 . Thus, identities between modular forms of a given weight can be proved by checking that only a finite number of their Fourier coefficients match. \tilde{E}_2 fails to be modular, indeed one can show

$$\tilde{E}_2\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^2 \tilde{E}_2(\tau) - 6i\pi c(c\tau + d) \quad \text{for all } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}). \quad (42)$$

Let us define a differentiation operator with an accurate normalization for our purposes:

$$D = 4i\pi \partial_\tau = 2(2i\pi)^2 q \partial_q.$$

Obviously, derivatives of modular forms are not modular forms. If f is a modular form of weight k for some subgroup Γ , then, for all $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$, we rather have

$$(Df)\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^{k+2} (Df)(\tau) + 4i\pi k c(c\tau + d)^{k+1} f(\tau).$$

This behavior is captured by the notion of “quasi-modular forms” and its relation with “non-holomorphic modular forms” [18, Chapter 1]. We adopt however a more pedestrian way. It is easy to check that the combination $\mathfrak{D}_k f = Df + \frac{2}{3}k \tilde{E}_2 f$ is modular of weight $k + 2$. Consequently, the differential closure of the ring of modular forms is generated by E_2 , E_4 and E_6 . The basic relations in the new ring are

$$D\tilde{E}_2 = \frac{2}{3}(\tilde{E}_4 - \tilde{E}_2^2), \quad (43)$$

$$D\tilde{E}_4 = \frac{8}{3}(\tilde{E}_6 - \tilde{E}_2\tilde{E}_4),$$

$$D\tilde{E}_6 = 4(\tilde{E}_4^2 - \tilde{E}_2\tilde{E}_6). \quad (44)$$

Since the vector spaces of modular forms of weight 4, 6 and 8 are 1-dimensional, these relations can be proved by checking from equation (42) that $\mathfrak{D}_1 \tilde{E}_2$ is modular of weight 4, hence of the form $c_4 \tilde{E}_4$, and similarly $\mathfrak{D}_4 \tilde{E}_4 = c_4 \tilde{E}_6$ and $\mathfrak{D}_6 \tilde{E}_6 = c_6 \tilde{E}_6$. Then, one finds c_{2l} by matching the constant Fourier coefficients of the two sides.

6.1.2. Theta functions and their derivatives. In genus 1 there are 3 even characteristics $\frac{1}{2}$, 0 and $\frac{\tau}{2}$. The corresponding theta values are

$$\vartheta_2(\tau) = \sum_{n \in \mathbb{Z}} (-1)^n q^{n^2/2},$$

$$\vartheta_3(\tau) = \sum_{n \in \mathbb{Z}} q^{n^2/2},$$

$$\vartheta_4(\tau) = \sum_{n \in \mathbb{Z} + 1/2} q^{n^2/2},$$

and they satisfy the relation

$$\vartheta_2^4 + \vartheta_4^4 = \vartheta_3^4.$$

The ϑ_i are modular forms of weight $1/2$, but only for a congruence subgroup $\Gamma(2)$ of $\mathrm{SL}_2(\mathbb{Z})$ (this is related to the shift of argument and the eight root of unity in equation (20)). Their fourth powers build a vector modular form of weight 2,

$$\vec{\vartheta}^4(\tau + 1) = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \cdot \vec{\vartheta}^4(\tau), \quad \vec{\vartheta}^4(-1/\tau) = \tau^2 \begin{pmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix} \cdot \vec{\vartheta}^4(\tau). \quad (45)$$

It is possible to build out of ϑ_i^4 expressions which are modular forms, resulting in relations to Eisenstein series upon checking a few Fourier coefficients. As before, we prefer to work with $\tilde{\vartheta}_i = \pi^{1/2} \vartheta_i$, and we obtain

$$\tilde{E}_4 = \tilde{\vartheta}_2^8 + \tilde{\vartheta}_4^8 + \tilde{\vartheta}_2^4 \tilde{\vartheta}_4^4, \quad (46)$$

$$\tilde{E}_6 = -\tilde{\vartheta}_2^{12} - \frac{3}{2} \tilde{\vartheta}_2^8 \tilde{\vartheta}_2^4 + \frac{3}{2} \tilde{\vartheta}_2^4 \tilde{\vartheta}_4^8 + \tilde{\vartheta}_4^{12}. \quad (47)$$

Combining equations (46) and (47) with equations (43) and (44), we obtain after some algebra the basic relations in the differential ring generated by the $\tilde{\vartheta}_i$,

$$\frac{D \tilde{\vartheta}_2}{\tilde{\vartheta}_2} = \frac{1}{3} (-\tilde{\vartheta}_2^4 - 2\tilde{\vartheta}_4^4 - \tilde{E}_2), \quad (48)$$

$$\frac{D \tilde{\vartheta}_3}{\tilde{\vartheta}_3} = \frac{1}{3} (-\tilde{\vartheta}_2^4 + \tilde{\vartheta}_4^4 - \tilde{E}_2),$$

$$\frac{D \tilde{\vartheta}_4}{\tilde{\vartheta}_4} = \frac{1}{3} (2\tilde{\vartheta}_2^4 + \tilde{\vartheta}_4^4 - \tilde{E}_2). \quad (49)$$

From there follows the computation of the d -th derivative of $\tilde{\vartheta}_i$ to all orders, and we observe especially that

$$D^d \tilde{\vartheta}_i / \tilde{\vartheta}_i = 3^{-d} P_d(\tilde{\vartheta}_2^4, \tilde{\vartheta}_4^4, \tilde{E}_2),$$

where P_d is a polynomial with integer coefficients. But we explained in §4.3 how the kernels could be computed in terms of combinations of derivatives which were modular. We immediately see that the appropriate combination must be equal to $3^{-d} P_d(\tilde{\vartheta}_2^4, \tilde{\vartheta}_4^4, 0)$. We are led to define

$$T_{2d;i} = 3^{-d} P_d(\tilde{\vartheta}_2^4, \tilde{\vartheta}_4^4, 0).$$

and we now focus on the computation of those numbers.

6.1.3. Application to elliptic curves. In this paragraph we consider a curve \mathcal{C} defined by an equation $\text{Pol}(m^2, l) = 0$ with integer coefficients, whose smooth model \mathcal{C}_0 is a Riemann surface of genus 1. Alternatively, there exists $x, y \in \mathbb{Q}(m^2, l)$ such that the defining equation of \mathcal{C}_0 can be brought in Weierstraß form

$$y^2 = 4x^3 - g_2x - g_3, \quad g_2, g_3 \in \mathbb{Q}, \quad (50)$$

and g_2, g_3 are called *elliptic invariants*. Up to a multiplicative constant, the unique holomorphic 1-form on \mathcal{C}_0 is

$$z = \frac{dx}{y}.$$

We assume we have chosen \mathcal{A} and \mathcal{B} cycles on the curve, it is not necessary to be precise about this choice as we will see in a moment. If we denote

$$2\varpi_A = \oint_{\mathcal{A}} dz \quad \text{and} \quad 2\varpi_B = \oint_{\mathcal{B}} dz,$$

then the holomorphic 1-form normalized on the \mathcal{A} -cycle is

$$da = \frac{dz}{2\varpi_A}$$

and the period is

$$\tau = \frac{\varpi_B}{\varpi_A}.$$

The curve \mathcal{C}_0 is isomorphic to $\mathbb{C}/(\mathbb{Z} \oplus \tau\mathbb{Z})$, and we can uniformize equations (50) by

$$x = \wp\left(\frac{z}{2\varpi_A} \mid \tau\right) \quad \text{and} \quad y = \wp'\left(\frac{z}{2\varpi_A} \mid \tau\right),$$

where \wp is the Weierstraß function

$$\wp(w \mid \tau) = \frac{1}{w^2} + \sum_{(n,m) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{1}{(w+n+m\tau)^2} - \frac{1}{(n+m\tau)^2}.$$

Let us recall the main properties of $\wp(w \mid \tau)$. It is an even periodic function with periods 1 and τ , which has a double pole with coefficient 1 and first subleading order $O(w^2)$. Its full asymptotic expansion when $w \rightarrow 0$ is

$$\wp(w \mid \tau) = \frac{1}{w^2} + \sum_{j \geq 1} \frac{2^{2j+2}}{(2j)!} \frac{(-1)^j B_{2j+2}}{2j+2} \tilde{E}_{2j+2}(\tau) w^{2j},$$

where B_{2j} are the Bernoulli numbers. The values of the Eisenstein series for \mathcal{C}_0 can be expressed in terms of g_2 and g_3 , by a comparison of the expansion of the left and right hand side of equation (50) when $z \rightarrow 0$:

$$\tilde{E}_4(\tau) = (2\varpi_A)^4 \frac{3g_2}{4}, \quad \tilde{E}_6(\tau) = (2\varpi_A)^6 \frac{27g_3}{8}. \quad (51)$$

Equations (46)-(47) allow in principle the

$$(t_2, t_3 = t_2 + t_4, t_4) = (\vartheta_2^4, \vartheta_3^4, \vartheta_4^4).$$

Yet, the precise choice of the solution depends on the choice of the basis of cycles, i.e. of a representative in the $\mathrm{SL}_2(\mathbb{Z})$ -orbit of τ . The modular group acts on equations (46) and (47) and their solution set according to equation (45). We remind that \tilde{E}_4 and \tilde{E}_6 are modular forms, but since we are dividing by the right power of $(2\varpi_A)$ at the end which is also modular, the point in the $\mathrm{SL}_2(\mathbb{Z})$ -orbit can be chosen arbitrarily as regards \tilde{E}_4 and \tilde{E}_6 . We will actually take the natural value of \tilde{E}_4 and \tilde{E}_6 coming from an equation of the elliptic curve, see equation (51). Those values are preserved by the subgroup Γ_+ of upper triangular matrices with 1's on the diagonal. Actually, Γ_+ is the subgroup preserving the \mathcal{A} -cycle (see §2.9), so ϖ_A is also invariant. Γ_+ acts transitively on the set of solutions of equations (46) and (47): if (t_2, t_3, t_4) is a solution, the others are

$$(-t_4, -t_3, -t_2), \quad (-t_2, t_4, t_3), \quad (-t_3, -t_4, t_2), \quad (t_4, -t_2, -t_3), \quad (t_3, t_2, -t_4). \quad (52)$$

To summarize, any solution of equations (46) and (47) – with \tilde{E}_4 and \tilde{E}_6 values deduced from an equation of the elliptic curve by equation (51) – will give us the fourth powers of the theta values, maybe in the wrong order and with the wrong sign. But the sign does not matter to compute $D^d \tilde{\vartheta}_i / \tilde{\vartheta}_i$ from equations (48)–(49), so the choice of another solution just results in a permutation of $i = 2, 3, 4$, i.e. of the label of the even characteristics.

6.1.4. Arithmetic aspects. The modular discriminant

$$\Delta(\tau) = e^{2i\pi\tau} \prod_{n=1}^{\infty} (1 - e^{2i\pi n\tau})^{24}$$

is another important modular form, of weight 12. In terms of Eisenstein series

$$\Delta(\tau) = \frac{E_4^3(\tau) - E_6^2(\tau)}{1728}.$$

Equivalently, we find its value from equations (46) and (47) or equation (51),

$$\Delta(\tau) = \frac{(2\varpi_A)^{12}}{(2\pi)^{12}} (g_2^3 - 27g_3^2) = \frac{1}{\pi^{12}} \frac{\tilde{\vartheta}_2^8 \tilde{\vartheta}_3^8 \tilde{\vartheta}_4^8}{256}.$$

If we assume g_2 and g_3 rational (and this is so when \mathcal{C}_0 comes from an A-polynomial), we learn from equations (46) and (47) and equation (51) that, for $i = 2, 3, 4$, $\tilde{\vartheta}_i^4 / (2\varpi_A)^2$ are algebraic numbers. Even more, the reality of g_2 and g_3 imply that the complex conjugates (t_2^*, t_3^*, t_4^*) must be in the list of solutions (52), and looking case by case we infer that one of the numbers $\tilde{\vartheta}_i^4 / (2\varpi_A)^4$ is real (if $\Delta > 0$) or pure imaginary (if $\Delta < 0$), while the two others are complex conjugates up to a sign. When $\Delta < 0$, we also have a privileged choice of even characteristics, namely the one for which $\tilde{\vartheta}_{i_0}^4 / 2\varpi_A^2$ is purely imaginary. This would remain true if τ was slightly changed by addition of an imaginary part (the $\tilde{E}_{2j}(\tau)$ would remain real). We deduce that $T_{2d;i_0} / (2\varpi_A)^{2d}$ are real, algebraic numbers. This last statement is also true for all $T_{2d;i} / (2\varpi_A)^{2d}$ when $\Delta > 0$, because the three numbers $\frac{\tilde{\vartheta}_i^4}{(2\varpi_A)^2}$ are real and we can apply directly the formulae (48)–(49).

6.1.5. Examples of A-spectral curves. Among the knots we investigated, we have found five non isomorphic elliptic curves arising as components of A-polynomials. They turn out to have a minimal model with coefficients $0, \pm 1$. We label them by their name in Cremona classification, the equation of their minimal model (l and m^2 are obtained by a birational transformation from x and y), and the knot for which they arise as geometric component.

15A8: $y^2 + xy + y = x^3 + y^2$ (**figure-eight knot**). A set of elliptic invariants is $g_2 = -\frac{1}{12}$ and $g_3 = \frac{161}{216}$, and the discriminant is $\Delta = -15$. We find that $T_{2d;i_0}$ are rational numbers.

$\tilde{\vartheta}_i^4 / (2\varpi_A)^2$	$\frac{7+i\sqrt{15}}{8}$	$\frac{7-i\sqrt{15}}{8}$	$-\frac{i\sqrt{15}}{4}$
$T_{2;i} / (2\varpi_A)^2$	$\frac{-7+3i\sqrt{15}}{24}$	$\frac{-7-3i\sqrt{15}}{24}$	$\frac{7}{12}$
$T_{4;i} / (2\varpi_A)^4$	$\frac{47+21i\sqrt{15}}{96}$	$\frac{47-21i\sqrt{15}}{96}$	$-\frac{47}{48}$
$T_{6;i} / (2\varpi_A)^6$	$\frac{-665+9i\sqrt{15}}{1152}$	$\frac{-665-9i\sqrt{15}}{1152}$	$-\frac{301}{576}$
$T_{8;i} / (2\varpi_A)^8$	$\frac{28375-12999i\sqrt{15}}{13824}$	$\frac{28375+12999i\sqrt{15}}{13824}$	$-\frac{28285}{6912}$

This curve also arises as non-geometric component in $7_4^{(1)}$, $8_{18}^{(1)}$, $8_{18}^{(2)}$, $9_{24}^{(1)}$, $9_{37}^{(1)}$, $9_{49}^{(1)}$, $10_{142}^{(1)}$, $10_{145}^{(1)}$, $10_{146}^{(1)}$, $10_{147}^{(1)}$, $10_{155}^{(1)}$.

14A4: $y^2 + xy + y = x^3 - x$ (**L^2R**). A set of elliptic invariants is $g_2 = -\frac{25}{12}$ and $g_3 = \frac{253}{216}$, and the discriminant is $\Delta = -28$. We find again that $T_{2d;i_0}$ are rational numbers.

$\tilde{\vartheta}_i^4 / (2\varpi_A)^2$	$\frac{11-i\sqrt{7}}{8}$	$\frac{11+i\sqrt{7}}{8}$	$\frac{i\sqrt{7}}{4}$
$T_{2;i} / (2\varpi_A)^2$	$\frac{-11-3i\sqrt{7}}{24}$	$\frac{-11+3i\sqrt{7}}{24}$	$\frac{11}{12}$
$T_{4;i} / (2\varpi_A)^4$	$\frac{71-33i\sqrt{7}}{96}$	$\frac{71+33i\sqrt{7}}{96}$	$-\frac{71}{48}$
$T_{6;i} / (2\varpi_A)^6$	$\frac{-1837-225i\sqrt{7}}{1152}$	$\frac{-1837+225i\sqrt{7}}{1152}$	$\frac{319}{576}$
$T_{8;i} / (2\varpi_A)^8$	$\frac{72583+37509i\sqrt{7}}{13824}$	$\frac{72583-37509i\sqrt{7}}{13824}$	$-\frac{16333}{6912}$

19A3: $y^2 + y = x^3 + x^2 + x$ (**9₃₅**). A set of elliptic invariant is $g_2 = \frac{8}{3}$ and $g_3 = -\frac{1}{27}$, and the discriminant is $\Delta = -19$. We find that $T_{2d;i_0} \in \mathbb{Q}(\alpha)$ with

$$\tilde{\vartheta}_{i_0}^4 / (2\varpi_A)^2 = \frac{2i}{\sqrt{6}}(\alpha^{1/2} + \alpha^{-1/2}), \quad \alpha = \left(\frac{257 + 3\sqrt{57}}{4}\right)^{1/3}.$$

11A3: $y^2 + y = x^3 - x^2$ (**9₄₈**). A set of elliptic invariants is $g_2 = -\frac{4}{3}$ and $g_3 = \frac{19}{27}$, and the discriminant is $\Delta = -11$. We find that $T_{2d;i_0} \in \mathbb{Q}(\alpha)$ with

$$\tilde{\vartheta}_{i_0}^4 / (2\varpi_A)^2 = \frac{i}{\sqrt{3}}(\alpha^{1/2} - \alpha^{-1/2}), \quad \alpha = \left(\frac{329 + 57\sqrt{33}}{2}\right)^{1/3}.$$

43A1: $y^2 + y = x^3 + x^2$ (10₁₃₉). A set of elliptic invariants is $g_2 = -\frac{4}{3}$ and $g_3 = \frac{35}{27}$, and the discriminant is $\Delta = -43$. We find that $T_{2d;i_0} \in \mathbb{Q}(\alpha)$, with

$$\tilde{\vartheta}_{i_0}^4 / (2\varpi_A)^2 = \frac{i}{\sqrt{3}}(\alpha^{1/2} - \alpha^{-1/2}), \quad \alpha = \left(\frac{1193 + 105\sqrt{129}}{2} \right)^{1/3}.$$

In the two first examples (figure-eight knot and L^2R , the fact that $T_{2d;i_0}$ are rational numbers imply that the coefficients $\tilde{j}_\chi(p)$ (for the choice of the characteristics associated to i_0) for $\chi \geq 1$ sit in the same function field as the amplitudes of the topological recursion. On the contrary, for the three last examples, they will sit a priori in an extension by the element α of the function field where the amplitudes of the topological recursion live.

Remark 6.4. The two first elliptic curves do not have complex multiplication, whereas the three last do. We thank Farshid Hajir for pointing us this property.

6.2. Bergman kernel for elliptic curves. In genus 1, there is only one odd characteristics, and the corresponding theta function is

$$\vartheta_1(w|\tau) = i \sum_{n \in \mathbb{Z}} e^{i\pi\tau(n+1/2)^2 + 2i\pi wn}.$$

It satisfies

$$\vartheta_1(w + 1|\tau) = -\vartheta_1(w|\tau)$$

and

$$\vartheta_1(w + \tau) = -e^{-i\pi(2w+\tau)} \vartheta_1(w|\tau).$$

The Bergman kernel normalized on the \mathcal{A} -cycle is

$$B_0(z_1, z_2) = -(\ln \vartheta_1)'' \left(\frac{z}{2\varpi_A} | \tau \right) \frac{dz_1}{2\varpi_A} \otimes \frac{dz_2}{2\varpi_A}.$$

On the other hand, the Weierstraß function provides another natural Bergman kernel

$$B_{\underline{k}(\tau)}(z_1, z_2) = \wp \left(\frac{z_1 - z_2}{2\varpi_A} | \tau \right) \frac{dz_1}{2\varpi_A} \otimes \frac{dz_2}{2\varpi_A}.$$

By “natural,” we mean that it can be written in terms of the coordinates x and y (see equation (50)) thanks to the addition relation for \wp . The result is

$$B_{\underline{k}(\tau)}(z_1, z_2) = \left[\frac{1}{4} \left(\frac{y(z_1) + y(z_2)}{x(z_1) - x(z_2)} \right)^2 - x(z_1) - x(z_2) \right] \frac{dz_1}{2\varpi_A} \otimes \frac{dz_2}{2\varpi_A}. \quad (53)$$

There is a well-known relation between \wp and ϑ_1 :

$$\wp(w|\tau) = -(\ln \vartheta_1)''(w|\tau) - \frac{\tilde{E}_2(\tau)}{3}.$$

In other words, equation (53) is the expression for a Bergman kernel normalized on the $\mathcal{A}_{\underline{\kappa}(\tau)}$ -cycle, with the value

$$\underline{\kappa}(\tau) = -\frac{1}{2i\pi} \frac{\tilde{E}_2(\tau)}{3}.$$

6.3. Application to some degree 2, elliptic A-spectral curves. We assume in this paragraph that the spectral curve $(\mathcal{C}, \ln l, \ln m)$ has a defining equation

$$l = \frac{P_1(m^2) + P_2(m^2)\sqrt{S(m^2)}}{R(m^2)}, \quad (54)$$

where P_1, P_2, R are polynomials and S is a polynomial of degree 4 with simple roots and leading coefficient 1. Also, it admits a smooth model \mathcal{C}_0 of the equation

$$\tilde{l} = \sqrt{S(m^2)}.$$

We also assume that $\{m, l\}$ is 2-torsion (i.e. $\zeta = 1$) and that $\iota_* = -\text{id}$. The spectral curve for the figure-eight knot and L^2R takes this form. Then, many simplifications occur.

6.3.1. Writing the n -forms. First, the ramification points a_i of the spectral curve coincide with the Weierstraß points of \mathcal{C}_0 , i.e. with the roots of S . Besides, the local involutions correspond to changing the sign of the squareroot

$$\sqrt{S(X)} \mapsto -\sqrt{S(X)},$$

and is in fact defined globally on \mathcal{C}_0 . We define a variable z by integration of the holomorphic 1-form

$$dz = \frac{dX}{\sqrt{S(X)}}.$$

One can check

$$\wp\left(\frac{z_1 - z_2}{2\varpi_A}\right) = \frac{1}{4} \frac{(\sqrt{S(X_1)} + \sqrt{S(X_2)})^2}{(X_1 - X_2)^2} - \frac{S'(X_1) - S'(X_2)}{12(X_1 - X_2)} + \frac{(X_1 - X_2)^2}{12}.$$

by matching the behavior of the two sides at their double pole $z_1 = z_2$ up to $\mathcal{O}(z_1 - z_2)$. In particular, we find

$$\wp\left(\frac{z - \bar{z}}{2\varpi_A}\right) = \frac{(S'(X))^2}{16S(X)} - \frac{S''(X)}{12}.$$

which intervenes in the computation of ω_1^1 . We also compute

$$\wp\left(\frac{z - a_i}{2\varpi_A}\right) = \frac{S'(a_i)}{4(X - a_i)} + \frac{S''(a_i)}{24}$$

and

$$\int^X \wp\left(\frac{z' - a_i}{2\varpi_A}\right) dz' = -\frac{\sqrt{S(X)}}{2(X - a_i)} + \frac{1}{24} \int^X \frac{dX S''(X)}{\sqrt{S(X)}}. \quad (55)$$

To avoid cumbersome notations, we use the same letter a_i to denote the image of the ramification point a_i in the Jacobian of \mathcal{C}_0 (on the left hand side) and the value of the X -coordinate at the same ramification point. The second term in equations (55) is odd when we change the sign of the squareroot, so disappear when we integrate from z to \bar{z} :

$$\int_{\bar{z}}^z \wp\left(\frac{z' - a_i}{2\varpi_A}\right) dz' = \frac{\sqrt{S(X)}(1/a_i - a_i)}{2(X - a_i)(X - 1/a_i)}. \quad (56)$$

Besides, one can check by differentiating the two sides of the equality

$$\int_{o, \iota(o)}^{z, \iota(z)} \wp\left(\frac{z - a_i}{2\varpi_A}\right) dz = \frac{\sqrt{S(X)}(1/a_i - a_i)}{2(X - a_i)(X - 1/a_i)} + C_i,$$

where C_i only depends on the basepoint o . This can be checked by differentiation and the fact that both sides of the equality are invariant under ι . If one denotes $\{a_i, \iota(a_i), a_{\bar{i}}, \iota(a_{\bar{i}})\}$ the set of ramification points, one finds that $C_i = 0$ when the basepoint o is chosen as $a_{\bar{i}}$ or $\iota(a_{\bar{i}})$. These formulae allow to complete the computation of the $\omega_{n|\underline{k}(\tau)}^h$, i.e. the topological recursion with the Bergman kernel $B_{\underline{k}(\tau)}$ defined in equation (53). We just need to compute expansion of the quantities in equation (56) at the branchpoints $X \rightarrow a_j$, and then take residues. The coordinate z is convenient for these computations, because Taylor expanding then amounts to differentiating the Weierstraß function with respect its argument. This method yields $\omega_{n|\underline{k}(\tau)}^h$ as a linear combination with rational coefficients of elementary n -forms of the type

$$\bigotimes_{j=1}^n \wp^{(p_j)}\left(\frac{z_j - a_{ij}}{2\varpi_A}\right) \frac{dz_j}{(2\varpi_A)^{p_j+1}}, \quad (57)$$

where p_j are even integers. It is easy to integrate $\omega_{n|\underline{k}(\tau)}^h$ over cycles with such a representation (we have to use equation (56) for terms with some $p_j = 0$). In particular, integrating z_j over

$$\frac{1}{2i\pi} \oint_{\mathcal{B}_{\underline{k}(\tau)}}$$

in equation (57) gives 0 if $p_j > 0$, and replaces the j^{th} -factor by $(2\varpi_A)^{-1}$ if $p_j = 0$.

The map ι_* has a single eigenvalue, which we assumed to be -1 . The discussion of §4.4 applies and can be explicitly checked: we find that the amplitudes

$$G_{n|\underline{k}(\tau)}^{h,(d)} = \frac{1}{n!} \frac{1}{(2i\pi)^d d!} \underbrace{\int_{o,\iota(o)}^{p,\iota(p)} \cdots \int_{o,\iota(o)}^{p,\iota(p)}}_{n \text{ times}} \underbrace{\oint_{\mathcal{B}_{\underline{k}(\tau)}} \cdots \oint_{\mathcal{B}_{\underline{k}(\tau)}}}_{d \text{ times}} \omega_{n+d|\underline{k}(\tau)}^h$$

vanish if d is odd. Then, the computation of $T_{2d;i}$ for d integer as detailed in §6.1 is all we need to evaluate $\mathcal{J}_h^{\text{n.p.TR}}(p)$.

6.3.2. Arithmetic aspects. Since S is palindromic, we can write the results in a more compact form with a variable

$$w = \frac{m^2 + m^{-2}}{2}.$$

We also denote

$$\frac{S(m^2)}{m^4} = \sigma(w).$$

Note that w is a uniformization variable for the quotient \mathcal{C}_0/ι which has genus 0 by assumption. The amplitudes $G_{n|\underline{k}(\tau)}^{h,(d)}$ are rational functions in w and $\sigma(w)$, with coefficients in $\mathbb{Q}[C_i]$. For positive levels, let \mathcal{P}_n^h the set of n -uples (p_1, \dots, p_n) such that the monomial

$$\bigotimes_{j=1}^n \wp\left(\frac{z_j - a_{i_j}}{2\varpi_A}\right) \frac{dz_j}{2\varpi_A}$$

appears in $\omega_n^h(z_1, \dots, z_n)$. At level 1, we have

$$\mathcal{P}_3^0 = \{(2, 2, 2)\}, \quad \mathcal{P}_1^1 = \{(4)\}.$$

It is easy to construct recursively a set $\mathcal{Q}_n^h \supseteq \mathcal{P}_n^h$ from the residue formula equation (22). At level one, we define $\mathcal{Q}_1^1 = \mathcal{P}_1^1$ and $\mathcal{Q}_3^0 = \mathcal{P}_3^0$, and if we know all \mathcal{Q}_n^h at level χ , we use the following rules to define \mathcal{Q}_n^h at level $\chi + 1$.

- If

$$(p, p', p_2, \dots, p_n) \in \mathcal{Q}_{n+1}^{h-1},$$

then

$$(p + p' + 2, p_2, \dots, p_n) \in \mathcal{Q}_n^h.$$

- For all $(n', h') \neq (1, 0)$, $(n + 1, h)$ such that $0 \leq h' \leq h$ and $0 \leq n' \leq n$, if

$$(p, p_2, \dots, p_{n'+1}) \in \mathcal{Q}_{n'+1}^{h'} \quad \text{and} \quad (p', p_{n'+2}, \dots, p_n) \in \mathcal{Q}_{n-n'}^{h-h'},$$

then

$$(p + p' + 2, p_2, \dots, p_n) \in \mathcal{Q}_n^h.$$

- If

$$(p_1, \dots, p_n) \in \mathcal{Q}_n^h$$

and σ is a permutation of n , then

$$(p_{\sigma(1)}, \dots, p_{\sigma(n)}) \in \mathcal{Q}_n^h.$$

We have an inclusion $\mathcal{P}_n^h \subseteq \mathcal{Q}_n^h$, and not an equality, because a monomial with indices (p_1, \dots, p_n) may come from several terms in the residue formula, and there might exist coincidences leading to cancellations. As a consequence

$$G_{n|\underline{k}(\tau)}^{h,(d)} \in (\sigma(w))^{-3r_n^{h,(d)}/2} \cdot \mathbb{Q}[w, C_i],$$

where

$$r_n^{h,(d)} = \max \left\{ \sum_{i=1}^{n-d} (p_{j_i} + 1) : (p_1, \dots, p_n) \in \mathcal{Q}_n^h, j_1, \dots, j_{n-d} \in \{1, \dots, n\} \right\}.$$

We already know $r_3^{0,(0)} = 3$, $r_3^{0,(2)} = 1$, and $r_1^{1,(0)} = 3$, and by recursion one can show

$$r_n^{h,(d)} = 3(2h - 2 + n) + d.$$

Thus, our construction naturally entails

$$\tilde{j}_\chi(w) \in (\sigma(w))^{-3\chi/2} \cdot \mathbb{Q}[w], \quad \text{for all } \chi \geq 1. \quad (58)$$

At the points such that $m^2 = 1$ (in particular the reference point where the hyperbolic metric is complete), we have $w = 1$. From the definition, we see that the cusp field is $\mathbb{F} = \mathbb{Q}[\sqrt{\sigma(1)}]$, and our construction naturally entails

$$\tilde{j}_\chi(w = 1) \in (\sigma(1))^{-3\chi/2} \cdot \mathbb{Q}[C_i] \subseteq \mathbb{F}[C_i], \quad \text{for all } \chi \geq 1. \quad (59)$$

In the examples below, we need $C_i \equiv 0$, so one can forget about C_i in equation (58) and equation (59). In this case, our Conjecture 5.6 predicts that the coefficients of the asymptotic expansion of the Kashaev invariant belong to the cusp field \mathbb{F} .

6.4. 4_1 (figure-eight knot). Apart from the abelian factor $(l - 1)$, the A-polynomial has a unique factor (necessarily the geometric one)

$$\mathfrak{A}(m, l) = l^2 m^4 + l(-m^8 + m^6 + 2m^4 + m^2 - 1) + m^4$$

It defines a curve \mathcal{C}_0 of genus 1. The symbol $\{m, l\}$ is 2-torsion, and $\iota_* = -\text{id}$. The spectral curve can be put in the form of equation (54),

$$l = \frac{m^8 - m^6 - 2m^4 - m^2 + 1 + (m^4 - 1)\sqrt{S(m^2)}}{2m^4}$$

and

$$S(X) = X^4 - 2X^3 - X^2 - 2X + 1,$$

so the results of §6.3 can be applied, and we introduce

$$w = \frac{m^2 + m^{-2}}{2} = \text{ch}(2u),$$

so

$$\frac{S(m^2)}{m^4} = \sigma(w) = 4w^2 - 4w - 3.$$

The curve has 4 ramification points, of coordinates

$$(m^2, l) = \left(\frac{3 \pm \sqrt{5}}{2}, 1\right) \quad \text{and} \quad \left(\frac{-1 \pm i\sqrt{3}}{2}, -1\right).$$

The local involution

$$z \mapsto \bar{z}$$

is defined globally on \mathcal{C}_0 , and it corresponds to

$$(m, l) \mapsto (m, 1/l).$$

Incidentally, this involution coincides with the amphichiral symmetry. The cusp field is $\mathbb{Q}[\sqrt{-3}]$. It is known [66] that the hyperbolic volume of $\mathfrak{M}_u = \mathbb{S}_3 \setminus 4_1$ with cusp angle $2 \text{Im } u$ and $\text{Re } u = 0$, is

$$\text{Vol}(\mathfrak{M}_u) = 2 \left[\Lambda \left(-iu + \frac{\beta(u)}{2} \right) - \Lambda \left(-iu - \frac{\beta(u)}{2} \right) \right],$$

where

$$\beta(u) = \arccos \left(\text{ch}(2u) - \frac{1}{2} \right)$$

and Λ is the Lobachevsky function

$$\Lambda(x) = - \int_0^x |\ln(2 \sin x')| dx'.$$

In particular, it vanishes when $u = \pm 2i\pi/3$, and this value coincide with the u -projection of two of the four branchpoints. Hence, if we denote by a_0 any of these points, we find that

$$\text{Im} \int_{a_0}^p \ln l \, d \ln m = 0$$

is half of the hyperbolic volume with the correct additive constant.




Figure 2. The figure-eight knot.

Amplitudes. We now derive the three first terms of $\mathcal{J}_h^{\text{n.p.TR}}$. We choose to compute the primitive with $C_i \equiv 0$. We computed the ω_n^h up to level 3 (i.e. in the case $2h - 2 + n \leq 3$). We just present here the expression for the non-vanishing amplitudes $G_{n|\underline{k}(\tau)}^{h,(d)}$ up to level 3. The values of $G_{n|\underline{k}(\tau)}^{h,(d=0)}$ for level 1 and 2 coincide¹² with the amplitudes computed in [27, §3.3]:

¹² More precisely, since [27] uses the spectral curve $2v \, du$ (instead of $v \, du$ here), we retrieve their amplitudes $\mathcal{F}^{(h,n)}$ by multiplying our results by $(1/2)^x$.

$$G_{3|\underline{k}(\tau)}^{0,(0)} = \frac{1}{6\sigma^{3/2}(w)}(-12w^2 + 12w - 7),$$

$$G_{1|\underline{k}(\tau)}^{0,(2)} = \frac{2(2\varpi_A)^{-2}}{15\sigma^{1/2}(w)}(-4w + 3),$$

$$G_{1|\underline{k}(\tau)}^{1,(0)} = \frac{1}{90\sigma^{3/2}(w)}(-8w^3 + 44w^2 - 30w - 87),$$

$$G_{4|\underline{k}(\tau)}^{0,(0)} = \frac{1}{3\sigma^3(w)}(16w^5 - 32w^4 + 24w^3 + 44w^2 - 67w + 25),$$

$$G_{2|\underline{k}(\tau)}^{0,(2)} = \frac{2(2\varpi_A)^{-2}}{135\sigma^2(w)}(64w^4 - 232w^3 + 156w^2 + 378w - 243),$$

$$G_{0|\underline{k}(\tau)}^{0,(4)} = -\frac{256(2\varpi_A)^{-4}}{10125},$$

$$G_{2|\underline{k}(\tau)}^{1,(0)} = \frac{1}{1620\sigma^3(w)} \begin{pmatrix} 1280w^6 - 9088w^5 + 13136w^4 + 22176w^3 \\ -17928w^2 - 26352w + 23193 \end{pmatrix},$$

$$G_{0|\underline{k}(\tau)}^{1,(2)} = \frac{-368(2\varpi_A)^{-2}}{10125},$$

$$G_{5|\underline{k}(\tau)}^{0,(0)} = \frac{1}{60\sigma^{9/2}(w)} \begin{pmatrix} -640w^8 + 1600w^7 - 1440w^6 - 9520w^5 + 18184w^4 \\ -3988w^3 - 18542w^2 + 19071w - 5644 \end{pmatrix},$$

$$G_{3|\underline{k}(\tau)}^{0,(2)} = \frac{2(2\varpi_A)^{-2}}{30375\sigma^{7/2}(w)} \begin{pmatrix} -37888w^7 + 283424w^6 - 471088w^5 - 636000w^4 \\ +1368360w^3 + 174906w^2 - 1767663w + 883791 \end{pmatrix},$$

$$G_{1|\underline{k}(\tau)}^{0,(4)} = \frac{4(2\varpi_A)^{-4}}{455625\sigma^{5/2}(w)} \begin{pmatrix} 94208w^5 - 287512w^4 + 76748w^3 \\ 370230w^2 - 112527w - 219564 \end{pmatrix},$$

$$G_{3|\underline{k}(\tau)}^{1,(0)} = \frac{1}{109350\sigma^{9/2}(w)} \mathfrak{E}_1,$$

$$G_{1|\underline{k}(\tau)}^{1,(2)} = \frac{8(2\varpi_A)^{-2}}{455625\sigma^{7/2}(w)} \mathfrak{E}_2,$$

$$G_{1|\underline{k}(\tau)}^{2,(0)} = \frac{1}{2733750\sigma^{9/2}(w)} \begin{pmatrix} 13238272w^9 - 70087552w^8 + 94437312w^7 \\ +49067168w^6 - 177750608w^5 + 952056w^4 \\ -78657516w^3 + 179205966w^2 + 191047329w \\ -219603474 \end{pmatrix}.$$

where

$$\mathfrak{G}_1 = \begin{pmatrix} -249856w^9 + 2901504w^8 - 6701952w^7 - 8240960w^6 \\ +8573472w^5 + 30776112w^4 - 55663848w^3 \\ -12104316w^2 + 71667990w - 34229709 \end{pmatrix}$$

and

$$\mathfrak{G}_2 = \begin{pmatrix} 353792w^7 - 1479360w^6 + 1280256w^5 + 1398544w^4 \\ -1258392w^3 - 1990008w^2 - 10098w + 1832382 \end{pmatrix}$$

First orders of $\mathcal{J}_{\hbar}^{\text{n.p. TR}}$ /comparison to colored Jones. We choose the even-half characteristics $[\mu, \nu]$ leading to real-valued theta derivatives (the last column of the table for the curve labeled 15A8 in §6.1.5 is selected). The reader may recognize in those values the *ad hoc* renormalizations of the constants G^k found by the authors of [27]. We use the general expressions given in §5.5 to compute

$$\begin{aligned} \tilde{J}_1 &= \frac{-1}{12\sigma^{3/2}(w)}(8w^3 - 4w^2 - 10w + 17) \\ &= \frac{-1}{12S^{3/2}(m^2)}(m^{12} - m^{10} - 2m^8 + 15m^6 - 2m^4 - m^2 + 1), \end{aligned}$$

$$\begin{aligned} \tilde{J}_2 &= \frac{2}{\sigma^3(w)}(8w^3 - 4w^2 - 10w + 7) \\ &= \frac{2m^6}{S^3(m^2)}(m^{12} - m^{10} - 2m^8 + 5m^6 - 2m^4 - m^2 + 1), \end{aligned}$$

and

$$\begin{aligned} \tilde{J}_3 &= \frac{1}{90\sigma^{9/2}(w)} \begin{pmatrix} 256w^8 - 512w^7 - 8704w^6 + 2048w^5 + 29792w^4 \\ -46928w^3 + 1272w^2 + 49164w - 27469 \end{pmatrix} \\ &= \frac{m^2}{90S^{9/2}(m^2)} \mathfrak{G}_3, \end{aligned}$$

where

$$\mathfrak{G}_3 = \begin{pmatrix} m^{32} - 4m^{30} - 128m^{28} + 36m^{26} + 1074m^{24} - 5630m^{22} \\ +5782m^{20} + 7484m^{18} - 18311m^{16} + 7484m^{14} + 5782m^{12} \\ -5630m^{10} + 1074m^8 + 36m^6 - 128m^4 - 4m^2 + 1 \end{pmatrix},$$

These coefficients are exactly those found in [30, Section 4.2.1] for a power series solution of the A-hat recursion relation of the figure-eight knot, so we have checked our Conjecture 5.5 up to $o(\hbar^3)$, with

$$C_{\hbar} = 1 + o(\hbar^3).$$

These authors as well as [27] also find the same coefficients in the asymptotic expansion of a Hikami-type integral $J_h^H(u)$ associated to the figure-eight knot. This is believed to be the correct asymptotic expansion for the colored Jones polynomial in the GVC near $i\pi$.

Specialization to $u = i\pi$ /comparison to Kashaev invariant. We recall that the complete hyperbolic point correspond to $w = 1$. The coefficients $J_\chi(w = 1)$ belong to $\mathbb{Q}[\sqrt{-3}]$. We find for the first orders

$$\begin{aligned} \tilde{j}_1(w = 1) &= -\frac{11}{12} (-3)^{-3/2}, \\ \tilde{j}_2(w = 1) &= 2 (-3)^{-3}, \\ \tilde{j}_3(w = 1) &= -\frac{1081}{90} (-3)^{-9/2}. \end{aligned}$$

This is in agreement with the asymptotic expansion of the Kashaev invariant, proved in [30] with help of numerics of numerical computations to be

$$\begin{aligned} J_N(\mathfrak{K}, q = e^{2i\pi/N}) \\ \underset{N \rightarrow \infty}{=} \frac{N^{3/2}}{3^{1/4}} \exp\left(\frac{N \text{Vol}(\mathbf{4}_1)}{2\pi} + \frac{11\epsilon_N}{12} + 2\epsilon_N^2 - \frac{1081\epsilon_N^3}{90} + O(\epsilon_N^4)\right), \end{aligned}$$

where

$$\epsilon_N = \frac{i\pi}{(-3)^{3/2}N}.$$

6.5. Once-punctured torus bundle L^2R . If $\mathfrak{t} = (\mathbb{R}^2/\mathbb{Z}^2) \setminus \{0\}$ denotes the once-punctured torus, the 3-manifold L^2R is defined as $\mathfrak{t} \times [0, 1]/\sim$ where the equivalence relation is generated by

$$(x, 0) \sim (L \circ L \circ R(x), 1), \quad L(x) = x + 1, R(x) = \frac{x}{x + 1}.$$

This manifold is hyperbolic [50, 80], . Apart from the abelian factor $(l - 1)$, the A -polynomial has a unique factor (necessarily the geometric one)

$$\mathfrak{A}(m, l) = l^2 m^4 + l(-m^6 + 2m^4 + 2m^2 - 1) + m^2.$$

It defines a curve of genus 1. The symbol $\{m, l\}$ is 2-torsion, and $l_* = -\text{id}$. The spectral curve can be put in the form equation (54):

$$l = \frac{m^6 - 2m^4 - 2m^2 + 1 + (m^2 - 1)\sqrt{S(m^2)}}{2m^4},$$

$$S(X) = X^4 - 2X^3 - 5X^2 - 2X + 1.$$

so the results of §6.3 can be applied, and we introduce

$$w = \frac{m^2 + m^{-2}}{2} = \text{ch}(2u), \quad \frac{S(m^2)}{m^4} = \sigma(w) = 4w^2 - 4w - 7.$$

The curve has 4 ramification points, of coordinates

$$(m^2, l) = \left(\frac{1 + 2\sqrt{2} \pm \sqrt{5 + 4\sqrt{2}}}{2}, \frac{1 + \sqrt{2} \pm (\sqrt{2} - 1)\sqrt{5 + 4\sqrt{2}}}{2} \right)$$

and

$$\left(\frac{1 - 2\sqrt{2} \pm i\sqrt{4\sqrt{2} - 5}}{2}, \frac{1 - \sqrt{2} \pm i(\sqrt{2} + 1)\sqrt{4\sqrt{2} - 5}}{2} \right).$$

The local involution $z \mapsto \bar{z}$ is defined globally of \mathcal{C}_0 . The cusp field is $\mathbb{Q}[\sqrt{-7}]$.

Amplitudes. We again choose to compute all primitives with $C_i \equiv 0$:

$$G_{3|\mathbb{K}(\tau)}^{0,(0)} = -\frac{1}{24\sigma^{3/2}(w)}(8w^3 + 36w^2 + 6w + 19)$$

$$G_{1|\mathbb{K}(\tau)}^{0,(2)} = \frac{(2\varpi_A)^{-2}}{14\sigma^{1/2}(w)}(-6w + 7),$$

$$G_{1|\mathbb{K}(\tau)}^{1,(0)} = \frac{1}{168\sigma^{3/2}(w)}(40w^3 - 12w^2 - 210w - 217),$$

$$G_{4|\mathbb{K}(\tau)}^{0,(0)} = \frac{1}{192\sigma^3(w)} \begin{pmatrix} 64w^6 + 832w^5 - 144w^4 + 3168w^3 \\ + 1532w^2 - 2060w + 1257 \end{pmatrix},$$

$$G_{2|\mathbb{K}(\tau)}^{0,(2)} = \frac{(2\varpi_A)^{-2}}{294\sigma^2(w)}(144w^4 - 816w^3 + 952w^2 + 1988w - 931),$$

$$G_{0|\mathbb{K}(\tau)}^{0,(4)} = -\frac{27(2\varpi_A)^{-4}}{1372},$$

$$G_{2|\mathbb{K}(\tau)}^{1,(0)} = \frac{1}{28224\sigma^3(w)} \begin{pmatrix} 7872w^6 - 116544w^5 + 341968w^4 + 841120w^3 \\ - 443884w^2 - 350644w + 556003 \end{pmatrix},$$

$$G_{0|\mathbb{K}(\tau)}^{1,(2)} = -\frac{57(2\varpi_A)^{-2}}{2744}.$$

First orders of $\mathcal{J}_\hbar^{\text{n.p. TR}}$ /comparison to colored Jones. We choose the even-half characteristics $[\mu, \nu]$ leading to real-valued theta derivatives (the last column of the table for the curve labeled 14A4 in §6.1.5 is selected):

$$\tilde{j}_1(w) = \frac{1}{24\sigma^{3/2}(w)}(-40w^3 + 44w^2 - 14w - 127),$$

$$\tilde{j}_2(w) = \frac{1}{128\sigma^3(w)} \begin{pmatrix} -64w^6 + 192w^5 + 1168w^4 + 3488w^3 \\ -2300w^2 - 2996w + 2071 \end{pmatrix}.$$

This is in agreement with the results of [27], and we recognize again their *ad hoc* renormalizations in the values of theta derivatives. These authors have computed the asymptotic expansion of a Hikami type integral $\mathcal{J}_\hbar^{\text{H}}(u)$ associated to L^2R :

$$\mathcal{J}_\hbar^{\text{H},\pm}(u) = \hbar^{\delta/2} \exp \left(\sum_{\chi \geq -1} J_\chi^{\text{H},(\pm)}(u) \right),$$

where \pm indicates the dependence of the integration contour. We have

$$\tilde{j}_1(w_u^\pm) = J_1^{\text{H},\pm}(u), \quad \tilde{j}_2(w_u^\pm) + \frac{1}{128} = J_2^{\text{H},\pm}(u)$$

Hence, a version of Conjecture 5.6 holds up to $o(\hbar^2)$, if the colored Jones on the LHS is replaced by $\mathcal{J}_\hbar^{\text{H},\pm}(u)$ holds. The appropriate normalization constant is

$$C_\hbar = 1 + \frac{\hbar^2}{128} + o(\hbar^2).$$

7. Heuristics imported from torus knots

Let (P, Q) be coprime integers. The A-polynomial of the torus knot $\mathfrak{K}_{P,Q}$ contains a non-abelian component of the form $A(m, l) = lm^PQ + 1$. Since the corresponding spectral curve does not have branchpoints, its partition function and kernels are ill-defined, so our conjecture for the Jones polynomial cannot be correct as such for torus knots for the non-abelian branch. Nevertheless, we shall see heuristically how the shape of our conjecture for any expectation value of Wilson loops arises in the case of torus knots. It is only at the end of this derivation, when we specialize to the Jones polynomial, that one discovers that the A-polynomial should be replaced by the germ of one of its deformation for the conjecture to be meaningful.

7.1. Matrix model for torus knots

7.1.1. General case Thanks to toric symmetry, the red expectation values of Wilson loops of $\mathfrak{K}_{P,Q}$ can be computed by localization [4, 56, 61, 86], and the sum over flat connections on $S_3 \setminus \mathfrak{K}_{P,Q}$ can be written as a matrix-like integral

$$\mathcal{W}_{G,R}(\mathfrak{K}_{P,Q}, \hbar) = \frac{1}{Z_{P,Q}} \int dX e^{-\frac{1}{\hbar} \frac{X^2}{4PQ}} \prod_{\alpha>0} 4 \operatorname{sh}\left(\frac{\alpha \cdot X}{2P}\right) \operatorname{sh}\left(\frac{\alpha \cdot X}{2Q}\right) \chi_R(e^X), \quad (60)$$

where $\alpha > 0$ are the positive roots of G , χ_R is the character of the representation R , and the normalization constant $Z_{P,Q}$ is

$$Z_{P,Q} = \int_{\mathbb{R}^n} dX e^{-\frac{1}{\hbar} \frac{X^2}{4PQ}} \prod_{\alpha>0} 4 \operatorname{sh}\left(\frac{\alpha \cdot X}{2P}\right) \operatorname{sh}\left(\frac{\alpha \cdot X}{2Q}\right). \quad (61)$$

This can be written even more explicitly, using Weyl's formula for the characters

$$\chi_R(e^X) = \frac{\sum_{\mathbf{w} \in \operatorname{Weyl}(G)} \epsilon(\mathbf{w}) e^{\mathbf{w}(\rho + \Lambda_R) \cdot X}}{\prod_{\alpha>0} 2 \operatorname{sh}\left(\frac{\alpha \cdot X}{2}\right)},$$

where

$$\rho = \frac{1}{2} \sum_{\alpha>0} \alpha$$

is the vector of Weyl's constants, Λ_R is the highest weight associated to R .

7.1.2. SU(n) case. For $\operatorname{SU}(n)$, the positive roots are

$$\alpha_{i,j} = \mathbf{e}_i - \mathbf{e}_j, \quad i < j,$$

where $\mathbf{e}_i = (0, \dots, 0, 1, 0, \dots, 0)$ with 1 in the i^{th} position, and

$$\rho = \sum_{i=1}^n \left(\frac{n+1}{2} - i \right) \mathbf{e}_i.$$

The Weyl group is the symmetric group \mathfrak{S}_n . Irreducible representations R are in correspondence with Young diagrams with n rows, or partitions

$$\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0),$$

and we have

$$\Lambda_R = (\lambda_1, \dots, \lambda_n).$$

The character associated to the representation indexed by λ is the Schur polynomial

$$\begin{aligned} s_\lambda(e^{\mathbf{X}}) &= \frac{\sum_{\sigma \in \mathfrak{S}_n} \epsilon(\sigma) \prod_{i=1}^n e^{X_{\sigma(i)}(\lambda_i - i + \frac{n+1}{2})}}{\prod_{1 \leq i < j \leq n} 2 \operatorname{sh}\left(\frac{X_i - X_j}{2}\right)} \\ &= \frac{\det(e^{H_i X_j})}{\prod_{1 \leq i < j \leq n} 2 \operatorname{sh}\left(\frac{X_i - X_j}{2}\right)} \prod_{i=1}^n e^{(\frac{n+1}{2} - c)X_i}, \end{aligned}$$

where

$$H_i = \lambda_i - i + c$$

and c is an arbitrary constant. From Harish–Chandra formula,

$$s_\lambda(e^{\mathbf{X}}) = \frac{\Delta(\mathbf{H})\Delta(\mathbf{X}) \prod_{i=1}^n e^{(\frac{n+1}{2} - c)X_i}}{\prod_{1 \leq i < j \leq n} 2 \operatorname{sh}\left(\frac{X_i - X_j}{2}\right)} \int_{\mathbf{U}(n)} dU e^{\operatorname{Tr} \mathbf{H} \mathbf{U} \mathbf{X} \mathbf{U}^\dagger},$$

where dU is the Haar measure on $\mathbf{U}(n)$ with total mass 1,

$$\mathbf{H} = \operatorname{diag}(H_1, \dots, H_n), \quad \mathbf{X} = \operatorname{diag}(X_1, \dots, X_n),$$

and

$$\Delta(\mathbf{X}) = \prod_{1 \leq i < j \leq n} (X_i - X_j)$$

is the Vandermonde determinant. Thus, equation (60) gives

$$\begin{aligned} &\mathcal{W}_{\mathbf{SU}(n), R}(\mathfrak{K}_{P, Q}, \hbar) \\ &= \frac{\Delta(\mathbf{H})}{(PQ)^{n(n-1)/2} Z_{P, Q}} \int_{\mathbb{R}^n \times \mathbf{U}(n)} d\mathbf{X} dU (\Delta(\mathbf{X}))^2 \\ &\quad e^{-\frac{1}{\hbar} \operatorname{Tr} \frac{\mathbf{X}^2}{4PQ} + (\frac{n+1}{2} - c) \operatorname{Tr} \mathbf{X} + \operatorname{Tr} \mathbf{H} \mathbf{U} \mathbf{X} \mathbf{U}^\dagger} \\ &\quad e^{\mathcal{V}(\mathbf{X}/P) + \mathcal{V}(\mathbf{X}/Q) - \mathcal{V}(\mathbf{X})}, \end{aligned} \quad (62)$$

where we have defined the potential $\mathcal{V}(\mathbf{X})$ as

$$e^{\mathcal{V}(\mathbf{X})} = \frac{\prod_{1 \leq i < j \leq n} 2 \operatorname{sh}\left(\frac{X_i - X_j}{2}\right)}{\Delta(\mathbf{X})}.$$

Note that $\mathcal{V}(X)$ is invariant under translation of X by a matrix proportional to the identity matrix $\mathbf{1}_n$. We may include the factors $i = j$ since they are equal to 1, and rewrite

$$\begin{aligned} \mathcal{V}(X) &= \text{Tr} \ln \left[\frac{2 \text{sh}(\mathbf{1}_n \otimes X - X \otimes \mathbf{1}_n)}{\mathbf{1}_n \otimes X - X \otimes \mathbf{1}_n} \right] \\ &= \frac{1}{2} \sum_{l, m \geq 1} (-1)^m \frac{B_{l+m}}{l+m} \frac{\text{Tr} X^l}{l!} \frac{\text{Tr} X^m}{m!}, \end{aligned} \quad (63)$$

where B_l the l^{th} Bernoulli number. To get rid of the linear term in the exponential in equation (62), we shift

$$X \mapsto X + \hbar P Q \left(\frac{n+1}{2} - c \right) \mathbf{1}_n.$$

The contour of integration \mathbb{R}^n is shifted to $\mathcal{C}_{R, \text{fi}}^n$ accordingly:

$$\mathcal{C}_{R, \text{fi}} = \mathbb{R} + \hbar P Q \left(\frac{n+1}{2} - c \right).$$

We define the normal matrix $M = U X U^\dagger$, and the invariant measure on the space $\mathcal{H}_n(\mathcal{C}_{\hbar, R})$ of normal matrices with eigenvalues on the contour $\mathcal{C}_{R, \hbar}$ is

$$dM = \frac{\text{Vol}[\text{U}(n)]}{(2\pi)^n n!} (\Delta(X))^2 dX dU.$$

Therefore,

$$\begin{aligned} &\mathcal{W}_{\text{SU}(n), R}(\mathfrak{K}_{P, Q}, \hbar) \\ &= \frac{\text{Vol}[\text{U}(n)]}{(2\pi)^n n!} \frac{e^{\hbar P Q n(n+1)/2 - c)^2/2}}{(PQ)^{n(n-1)} Z_{P, Q}} \Delta(\mathbf{H}) e^{2PQ\hbar((n+1)/2 - c) \text{Tr} \mathbf{H}}, \quad (64) \\ &\int_{\mathcal{H}_n(\mathcal{C}_{R, \hbar})} dM e^{-\frac{1}{\hbar} \text{Tr} \frac{M^2}{4PQ}} e^{\mathcal{V}(M/P) + \mathcal{V}(M/Q) - \mathcal{V}(M)}. \end{aligned}$$

The Vandermonde of the H 's is related to the dimension of the representation

$$\dim R = \frac{\Delta(\mathbf{H})}{\prod_{i=1}^n (H_i + n - c)!},$$

and the trace of \mathbf{H} is related to the number of boxes in λ ,

$$\text{Tr} \mathbf{H} = |\lambda| + n \left(c - \frac{n+1}{2} \right).$$

So far, the constant c was arbitrary, in particular it can depend on λ . The choice

$$c = -\frac{|\lambda|}{n} + \frac{n+1}{2}$$

allows to have $\text{Tr } \mathbf{H} = 0$, and we now stick to it. The normalization constant $Z_{P,Q}$ is computed with the same steps for the trivial representation R_\emptyset of $\text{SU}(n)$.

$$\lambda_i^\emptyset = 0 \quad H_i^\emptyset = -i + \frac{n+1}{2} \quad \text{for all } i \in \{1, \dots, n\}.$$

Thus,

$$\begin{aligned} & \mathcal{W}_{\text{SU}(n), R}(\mathfrak{R}_{P,Q}, \hbar) \\ &= D_R \frac{\int_{\mathcal{H}_n(\mathbb{C}_{R,\hbar})} dM e^{-\frac{1}{\hbar} \text{Tr} \frac{M^2}{4PQ}} e^{\mathcal{V}(M/P) + \mathcal{V}(M/Q) - \mathcal{V}(M)} e^{\text{Tr } \mathbf{H} M}}{\int_{\mathcal{H}_n(\mathbb{R})} dM e^{-\frac{1}{\hbar} \text{Tr} \frac{M^2}{4PQ}} e^{\mathcal{V}(M/P) + \mathcal{V}(M/Q) - \mathcal{V}(M)} e^{\text{Tr } \mathbf{H}^\emptyset M}}, \end{aligned} \quad (65)$$

where the multiplicative constant is given by

$$D_R = e^{\frac{\hbar PQ |\lambda|^2}{n}} \frac{\prod_{j=1}^n \left(H_j + \frac{n-1}{2} + \frac{|\lambda|}{n} \right)!}{\prod_{j=1}^{n-1} j!} \dim R.$$

The integral in the numerator is similar to that in the denominator, except for the *external field* \mathbf{H} (resp. \mathbf{H}^\emptyset) encoding the highest weight associated to R (resp. to the trivial representation). We also shifted the contour from \mathbb{R} to

$$\mathbb{C}_{R,\hbar} = \mathbb{R} + \frac{2\hbar PQ}{n} |\lambda|.$$

Since the discussion to come remains at a formal level, we shall move it back to \mathbb{R} , and assume the range of integration to be the space of Hermitian matrices $\mathcal{H}_n(\mathbb{R})$ in equation (65).

7.2. Computation from the topological recursion

7.2.1. Principle. For matrix integrals (with or without external potential) of the form equation (64), we have [38]

$$Z = \mathcal{J}_\hbar(\mathbb{S}_{n,\hbar}), \quad (66)$$

where

$$\mathcal{S}_{n,\hbar} = (\mathcal{C}, x, y)$$

is the spectral curve of the matrix integral, which in general depend on \hbar and the size of the matrix n , and \mathcal{T}_{\hbar} is the non-perturbative partition function defined in Section 4. We have define somewhat arbitrarily¹³ equation (66) means that the asymptotic expansion of the left hand side is given by the right hand side (which we defined as a formal asymptotic series). Adding an external field in the form $e^{\text{Tr} \mathbf{H} \mathbf{M}}$ amounts (see for instance [79]) to modify the spectral curve by addition of simple poles $p_1, \dots, p_n \in \mathcal{C}$ to x with residue \hbar with respect to dy , and such that

$$y(p_j) = \hbar H_j,$$

and some other simple poles $o_1, \dots, o_n \in \mathcal{C}$ with residue $-\hbar$,

$$(\mathcal{C}, x, y) \mapsto \left(\mathcal{C}, x + \hbar \sum_{j=1}^n \frac{dS_{o_j, p_j}}{dy}, y \right).$$

Similarly, we denote p_j^\emptyset the poles associated to the external field \mathbf{H}^\emptyset , i.e.

$$y(p_j^\emptyset) = \hbar \left(-i + \frac{n+1}{2} \right).$$

We thus find

$$\mathcal{W}_{\text{SU}(n), R}(\mathcal{R}_{P, Q}, \hbar) = D_R \frac{\mathcal{T}_{\hbar} \left(\mathcal{C}, x + \hbar \sum_{j=1}^n dS_{o_j, p_j} / dy, y \right)}{\mathcal{T}_{\hbar} \left(\mathcal{C}, x + \hbar \sum_{j=1}^n dS_{o_j, p_j^\emptyset} / dy, y \right)}.$$

Then, the symplectic invariance (see §3.3) allows to exchange¹⁴ the role of x and y :

$$\mathcal{W}_{\text{SU}(n), R}(\mathcal{R}_{P, Q}, \hbar) = D_R \frac{\mathcal{T}_{\hbar} \left(\mathcal{C}, y, x + \hbar \sum_{j=1}^n dS_{o_j, p_j} / dy \right)}{\mathcal{T}_{\hbar} \left(\mathcal{C}, y, x + \hbar \sum_{j=1}^n dS_{o_j, p_j^\emptyset} / dy \right)}.$$

¹³ This line of reasoning does not tell us the scale of \hbar , because changing \hbar to $\alpha \hbar$ amounts to rescaling $y dx$ to $\alpha y dx$, and we are not precise enough to identify y and x to (some multiple of) the meridian and longitude eigenvalues m and l of knot theory. Outside Section 7, the quantization condition satisfied by A-spectral curves provided a good argument in favor of this choice.

¹⁴ This exchange is possible when the set of zeroes of dy is not empty, i.e. when the spectral curve (\mathcal{C}, y, x) has at least one ramification point.

Now, the simple poles are added to the second function in the spectral curve, and we recognize the $n|n$ -kernel described in §4.1, for the spectral curve

$$\widehat{\mathcal{S}}_{n,\hbar} = (\mathbb{C}, y, x)$$

for some basepoint o :

$$\mathcal{W}_{\text{SU}(n),R}(\mathfrak{R}_{P,Q}, \hbar) = D_R \frac{\psi_{\hbar}^{[n|n]}(p_1, o_1; \dots; p_n, o_n)}{\psi_{\hbar}^{[n|n]}(p_1^{\emptyset}, o_1; \dots; p_n^{\emptyset}, o_n)}. \quad (67)$$

7.2.2. SU(2) case: Jones polynomial. For $\text{SU}(n = 2)$ and the representation R associated to $(\lambda_1, \lambda_2) = (N - 1, 0)$, we retrieve the colored Jones polynomial (see equation (38)). It is thus computed from the 2-kernel with points p_1 and p_2 of projections,

$$y(p_1) = 2\hbar H_1 = N\hbar, \quad y(p_2) = 2\hbar H_2 = -N\hbar, \quad (68)$$

and, for the trivial representation,

$$y(p_1^{\emptyset}) = \hbar, \quad y(p_2^{\emptyset}) = -\hbar.$$

This leads to

$$J_N(\mathfrak{R}_{P,Q}, q = e^{2\hbar}) = N e^{\hbar P Q (N-1)^2} \frac{\text{sh } \hbar}{\text{sh } N\hbar} \frac{\psi_{\hbar}^{[2|2]}(p_1, p_2)}{\psi_{\hbar}^{[2|2]}(p_1^{\emptyset}, p_2^{\emptyset})}.$$

and we insist that the kernels are computed for the spectral curve of the matrix integral after exchange of x and y .

7.2.3. Spectral curve for the torus knots. Let P', Q' be integers such that $P'Q - Q'P = 1$. The spectral curve $\mathcal{S}_{n,\hbar} = (\mathbb{C}, x, y)$ of the matrix integral in equation (61) was derived in [17], in the regime when $n\hbar$ is of order 1,

$$\mathcal{S}: e^{-(P+Q)y} - e^{-n\hbar} e^{-(P'x+Qy)} - e^{-(Q'x+Py)} + e^{n\hbar} e^{-(P'+Q')x} = 0. \quad (69)$$

This curve can be uniformized with a variable $z \in \mathbb{C} \equiv \mathbb{C}$:

$$e^x = e^{(P-Q)n\hbar} e^{-Pz} \left(\frac{e^{2n\hbar} - e^z}{1 - e^z} \right)^Q$$

$$e^y = e^{(P'-Q'-1/Q)n\hbar} e^{-P'z} \left(\frac{e^{2n\hbar} - e^z}{1 - e^z} \right)^{Q'}.$$

hence its genus is 0, and there is no theta function in the definition of its partition function and kernels. It was argued in [17] that the topological recursion for this curve reproduces the torus knots invariants.

Here, we are interested in the regime where \hbar is small and n is fixed (and in particular $n = 2$). If we keep x and y of order 1, the curve is trivial:

$$e^{Py} = e^{P'x}.$$

in the sense that it does not have ramification points. So, the partition function \mathcal{T}_{\hbar} of this spectral curve is ill-defined. Actually, when the limit spectral curve is ill-defined, the information about the unstable terms (i.e. the terms decaying with $\hbar \rightarrow 0$) are actually obtained contained in the blow-up of $\mathcal{S}_{n,\hbar}$ at its basepoint o . It is realized by setting

$$x = \sqrt{2n\hbar} \tilde{x} \quad \text{and} \quad y = \sqrt{2n\hbar} \tilde{y},$$

with \tilde{x} and \tilde{y} of order 1, and retaining the first non trivial order in equation (69) when $n\hbar \rightarrow 0$, we find

$$PQ \tilde{y}^2 - (P'Q + PQ')\tilde{x}\tilde{y} + P'Q' \tilde{x}^2 + 1 = 0. \quad (70)$$

The formulae (66) and (67) are expected to be correct if applied to the spectral curve of equation (70) at least for the terms of order $o(1)$ when $\hbar \rightarrow 0$. The non decaying terms are rather given by the limit spectral curve itself, and thus are trivial. This is in agreement with the fact [59] that there is no exponential growth of the Jones polynomial of torus knots (they are not hyperbolic), in other words $\tilde{j}_{-1} \equiv 0$.

7.3. General mechanism. We expect that the mechanism described for torus knots complements is general (see for instance the conjecture in [61, Section 6]), and our proposal should essentially compute expectation values of Wilson loops for more general 3-manifold \mathfrak{M} , provided the spectral curve is well-chosen. A scenario would be that

$$\mathcal{W}_{G,R}(\partial\mathfrak{M}, \hbar) = \frac{\int dV \chi_R(V) e^{S(V)}}{\int dV e^{S(V)}}, \quad (71)$$

where V consists of eigenvalues of the holonomy operator

$$\text{P exp} \oint_{\partial\mathfrak{M}} \mathcal{A},$$

and $S(V)$ the effective action for V after integrating out the other degrees of freedom against the Chern–Simons action. If a formula like (71) holds, the steps

of §7.1.2 can be repeated to find

$$\mathcal{W}_{\text{SU}(n)}(\partial\mathfrak{M}, \hbar/2) \propto \frac{\int dM e^{S(M)} e^{\text{Tr} \mathbf{H} M}}{\int dM e^{S(M)}},$$

where

$$M = U X U^\dagger, \quad U \in \text{SU}(n),$$

and

$$H_i = \lambda_i - i + c,$$

for some constant c . Then, if the integral

$$\int dM e^{S(M)}$$

is a Tau function with spectral curve $\mathcal{S}_{n,\hbar} = (\mathcal{C}, x, y)$, we would find again

$$\mathcal{W}_{\text{SU}(n)}(\partial\mathfrak{M}, \hbar) \propto \frac{\mathcal{T}_{\hbar}(\mathcal{C}, y, x + \hbar \sum_{i=1}^n dS_{o,p_i}/dy)}{\mathcal{T}_{\hbar}(\mathcal{C}, y, x + \hbar \sum_{i=1}^n dS_{o,p_i^\emptyset}/dy)} \propto \frac{\psi_{\hbar}^{[n|n]}(p_1, \dots, p_n)}{\psi_{\hbar}^{[n|n]}(p_1^\emptyset, \dots, p_n^\emptyset)},$$

where the $n|n$ -kernel is computed for the spectral curve $\widehat{\mathcal{S}}_{n,\hbar} = (\mathcal{C}, y, x)$ after the exchange $x \leftrightarrow y$. For hyperbolic manifolds, one expects to find as spectral curve a deformation of the A-polynomial (or at least of subcomponents of it), which reduces to the A-polynomial in the $\text{SU}(n = 2)$ case, i.e. to $m \propto e^{2y}$, $l \propto e^x$ and $A(m, l) = 0$. This is plausible because it is known that the A-polynomial can be obtained as the saddle point equation obtained by elimination from the Neumann–Zagier potential [54]. In our argument, we see then from equation (68) that the points p_1 and p_2 needed to compute the Jones polynomial have m -projection

$$\ln m(p_1) = N\hbar + \text{cte}, \quad \ln m(p_2) = -N\hbar + \text{cte},$$

and one recognizes¹⁵ (up to constant shift) the identification between the hyperbolic structure parameter $m = e^u$ and the quantum group parameter $q = e^{2\hbar}$ and N appearing in equation (38).

¹⁵ Again, the argument for the overall scaling of \hbar does not come from the discussion of this Section.

8. Perspectives

We have constructed a formal asymptotic series $\mathcal{J}_\hbar^{\text{n.p.}, \text{TR}}(p)$ depending on a point on the A-polynomial curve of a hyperbolic 3-manifold with 1-cusp, which has interesting properties *per se*. It depends on a choice of characteristics $\mu, \nu \in \mathbb{C}^g$ (which might be restricted to even-half characteristics) and basepoint o for the computation of iterated primitives. Provided an accurate choice is made for those data, we have conjectured that it computes the asymptotic expansion of the colored Jones polynomial, discarding roots of unity. We have made a non-trivial check to first orders for the figure-eight knot. A weaker conjecture is that our series is a formal solution of the A-hat recursion relation. We made a closely related check to first orders for the L^2R . We think that working on the A-hat recursion relation satisfied by the colored Jones polynomial is a good approach in an attempt to prove our conjecture (or a slight modification of it).

The intuition behind our construction comes from the theory of integrable systems and its relations to loop equations. $\mathcal{J}_\hbar^{\text{n.p.}, \text{TR}}(p)$ was defined formally by introducing an infinite number of infinitesimal deformations of the A-polynomial curve, and one may wonder if this can be interpreted as an integrable perturbation of the Wess–Zumino–Witten CFT.

The main interest of our proposal is rather structural than computational. Although we do have an algorithm, it requires the use of a basis of meromorphic forms which behaves well under integration, and the computation of theta functions and derivatives, so is less efficient than other methods. Yet, the expression in terms of the topological recursion suggests possible connections between knot theory and respectively integrable systems [13, 40], intersection theory on the moduli space [36] and enumerative geometry, which deserve further investigation.

This conjecture gives also a framework for the study of the arithmetic properties of perturbative knot invariants. It would be particularly interesting to compare our predictions for the expansion of the Kashaev invariant (i.e. at the complete hyperbolic point) to those of [29] obtained by a gluing procedure.

When the quotient of (a component of) the A-polynomial curve by the involution

$$\iota(m, l) = \left(\frac{1}{m}, \frac{1}{l} \right)$$

is a genus $g_\iota = 0$ curve, $\mathcal{J}_\hbar^{\text{n.p.}, \text{TR}}$ is a formal power series, although it takes into account non-perturbative effects. When this property does not hold, it is rather an asymptotic series which contains fast oscillations to all orders when $\hbar \rightarrow 0$. However, the K-theoretical properties of the A-polynomial imply that, if we specialize \hbar to sequences $i\pi/k$ with k an integer of fixed congruence, we retrieve a formal

power series. It would be interesting to know if this wild behavior can be seen in the asymptotics of the colored Jones polynomial for knots such that $g_i \neq 0$. The simplest examples of this kind we know are 8_{21} and the Pretzel $(-2, 3, 9)$, and are currently under investigation. Actually, earlier experiments on the asymptotics of the colored Jones have been performed to our knowledge only for knots with $g_i = 0$. If Conjecture 5.6 contains some part of truth, new phenomena may be discovered. Else, one would need to understand how it should be modify to preserve the matching for 4_1 and $L^2 R$.

Our conjectures could be generalized in several directions.

- One may wonder if the $n|n$ kernels can be identified to asymptotics of other relevant knot invariants. Note also that Hirota equations imply determinantal formulae [13]

$$\psi_{\hbar}^{[n|n]}(p_1, q_1; \dots; p_n, q_n) \propto \det_{i,j} \psi_{\hbar}^{[1|1]}(p_i, q_j),$$

where \propto means equality up to a factor involving prime forms. A naive guess, inspired by Section 7, would be to compare $\psi_{\hbar}^{[n|n]}$ to expectation values of Wilson loops in a Chern–Simons theory with $SU(n|n)$ gauge group in the limit of large representations. The rescaled size of the representation would be in correspondence (see §5.1) in the case of $SU(n)$ with points p_1, \dots, p_{2n} on the A-polynomial curve.

- One may wish to study the asymptotics of the colored Jones when $q \rightarrow \zeta_d$ (a root of unity, instead of $q \rightarrow 1$ here). It is natural to propose a conjecture similar to 5.6, with the curve of equation $\lim_{q \rightarrow \zeta_d} \hat{\mathfrak{A}}(e^u, e^v, q)$ replacing the A-polynomial.
- One may wish to study the a - (or Q -)deformation of the knot invariants, considered recently in [2]. In the regime when $q \rightarrow 1$ but keeping a finite, this amounts to study asymptotics for gauge groups of large rank. We guess that the non-topological recursion for the a -deformed A-polynomial will come into play.
- And, at the top of the hierarchy, one may consider the categorified knot invariants, which results from another deformation with a variable t [32]. These knot invariants can be seen as generating series of BPS invariants. They are conjectured to be annihilated by an operator $\mathfrak{A}(e^u, e^{\hbar \partial_u}, a, q, t)$, called the super-A-hat polynomial, which is explicitly known in a few examples [43, 44]. In this case, although a t -deformed spectral curve can be defined, we think that a “deformed topological recursion” should be used in order to compute something meaningful about their asymptotics. This intuition

is based on the fact that Schur polynomial have to be replaced by Macdonald polynomials under this deformation, and ongoing work suggests that the analysis of the matrix model of equation (60) requires a deformation of the topological recursion.

Although the identification for the colored Jones polynomial ($N\hbar = \ln m(p_u)$) was rather simple, appropriate and non-trivial “mirror maps” (like in topological strings [1, 15]) could be necessary to make any of those generalizations effective.

In yet another direction, the (generalized) volume conjecture can also be formulated for links with L components. The $\mathrm{SL}_n(\mathbb{C})$ character variety has local complex dimension $(n-1)L$ at a generic point [67]. It is a challenging problem to reduce – if only possible – the asymptotics of quantum invariants of 3-manifolds with L cusps to algebraic geometry on this variety.

APPENDIXES

A. Diagrammatic representation for the non-perturbative topological recursion

A.1. Non-perturbative partition function. The non-perturbative Tau function $\mathcal{T}_{\hat{\mathfrak{h}}}$ was defined in equation (29). We had

$$\mathcal{T}_{\hat{\mathfrak{h}}} = e^{\hat{\mathfrak{h}}^{-2}F_0 + F_1} \vartheta_{[\nu]^\mu}^{[\mu]}(\zeta|\tau) \hat{\mathcal{T}}_{\hat{\mathfrak{h}}},$$

with

$$\hat{\mathcal{T}}_{\hat{\mathfrak{h}}} = \exp\left(\sum_{h \geq 2} \hat{\mathfrak{h}}^{2h-2} F_h\right) \frac{1}{\vartheta_{[\nu]^\mu}^{[\mu]}(\zeta|\tau)} \left\{ 1 + \sum_{r \geq 1} \frac{1}{r!} \sum_{\substack{h_j \geq 0, d_j \geq 1 \\ 2h_j - 2 + d_j > 0}} \hat{\mathfrak{h}}^{\sum_j 2h_j - 2 + d_j} \bigotimes_{j=1}^r \frac{F_{h_j}^{(d_j)} \cdot \nabla^{\otimes d_j}}{(2i\pi)^{d_j} d_j!} \right\} \vartheta_{[\nu]^\mu}^{[\mu]}(\zeta|\tau). \quad (72)$$

We also recall that, owing to special geometry, the k^{th} derivative of ω_n^h with respect to filling fractions is

$$\omega_n^{h,(k)} = \overbrace{\oint_{\mathbb{B}} \dots \oint_{\mathbb{B}}}^{k \text{ integrals}} \omega_{n+k}^h.$$

For $2 - 2h - n < 0$, we represent ω_n^h by a surface with h handles and n legs, and we represent $\nabla^{\otimes k} \vartheta / \vartheta$ by a black vertex with k legs.

Then, with those diagrammatic notations, equation (72) is represented as a sum of graphs. Each graph has exactly one black vertex, whose legs are attached to the legs of a product of ω_n^h 's, such that all legs are paired,


$$\widehat{\mathcal{T}}_{\text{fi}} = 1 + \sum \frac{\hbar^{-\chi_{\text{Euler}}}}{\# \text{Aut}} \text{weight} \left(\begin{array}{c} \text{Diagram with a black vertex connected to three punctured surfaces} \end{array} \right),$$

where χ_{Euler} is sum of the Euler characteristics of all punctured surfaces of the graph (each of them having a negative Euler characteristics), and $\# \text{Aut} \in \mathbb{N}^*$ is the symmetry factor of the graph.


A.2. Logarithm of the non-perturbative partition function. Note that the generating function for the derivatives of $\ln \vartheta$, is related to the generating function for the derivatives of ϑ , by keeping the cumulants. If we represent $\nabla^{\otimes k} \ln \vartheta$ by a white vertex with k legs, we have that the black vertex is the sum of all possible products of white vertices having the same legs.

$$\frac{\nabla^{\otimes k} \vartheta \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right]}{\vartheta \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right]} = \text{Black vertex with } k \text{ legs} \quad \nabla^{\otimes k} \ln \vartheta \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right] = \text{White vertex with } k \text{ legs}.$$

and



This means that the diagrammatic representation of $\widehat{\mathcal{T}}_{\mathfrak{h}}$, in terms of white vertices is a sum of all graphs, not necessarily connected, whose vertices are either surfaces with handles and punctures, or white vertices, and whose edges connect the punctures to white vertices. Then the logarithm of $\widehat{\mathcal{T}}_{\mathfrak{h}}$ has the same diagrammatic representation, but keeping only connected graphs,

$$\ln \widehat{\mathcal{T}}_{\mathfrak{h}} = \sum \frac{\mathfrak{h}^{-\chi_{\text{Euler}}}}{\# \text{Aut}} \text{weight} \left(\begin{array}{c} \text{Diagrammatic representation of connected graphs} \end{array} \right),$$


where χ_{Euler} is the sum of the Euler characteristics of the punctured surfaces appearing in the graph.

A.3. $n|n$ kernels. We take the example of the $2|2$ kernel since it is the one which can be compared to the Jones polynomial. It was defined in equation (32) by

$$\psi_{\mathfrak{h}}^{[2|2]}(p_1, o_1; p_2, o_2) = \frac{\mathcal{T}_{\mathfrak{h}}[v \, du \longrightarrow v \, du + \mathfrak{h} \, dS_{o_1, p_1} + \mathfrak{h} \, dS_{o_2, p_2}]}{\mathcal{T}_{\mathfrak{h}}[v \, du]}.$$

Note that, when we add in the numerator simple poles p_1, o_1, p_2, o_2 to the spectral curve, we obtain

$$\omega_n^h \rightarrow \sum_{r=0}^{\infty} \frac{\mathfrak{h}^r}{r!} \int_{o_1, o_2}^{\overbrace{p_1, p_2}^{r \text{ times}}} \dots \int_{o_1, o_2}^{\overbrace{p_1, p_2}^{r \text{ times}}} \omega_{n+r}^h, \tag{73}$$

and thus

$$\omega_n^{h, (d)} \rightarrow \sum_{r=0}^{\infty} \frac{\mathfrak{h}^r}{r!} \oint_{\mathcal{B}} \dots \oint_{\mathcal{B}} \int_{o_1, o_2}^{\overbrace{p_1, p_2}^{d \text{ times}}} \dots \int_{o_1, o_2}^{\overbrace{p_1, p_2}^{r \text{ times}}} \omega_{n+r+d}^h. \tag{74}$$

Note that the formula for $\mathcal{T}_{\mathfrak{h}}$ involves only

$$F_h^{(d)} = \omega_0^{h,(d)}, \quad \text{with } 2 - 2h - d < 0.$$

In particular, equations (73) and (74) do not yield the terms with $h = 0$ and $d = 1, 2$, but they produce all the other terms. Besides, the argument of the theta function and its matrix of periods is also shifted by the addition of simple poles in the spectral curve. We obtain

$$\vartheta \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right] (\mathbf{w} | \tau) \longrightarrow \vartheta \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right] (\mathbf{w}' | \tau'),$$

$$\mathbf{w}' = \mathbf{w} + \mathbf{a}(p_1) - \mathbf{a}(o_1) + \mathbf{a}(p_2) - \mathbf{a}(o_2) + \sum_{r \geq 2} \frac{\mathfrak{h}^{r-1}}{r!} \oint_{\mathcal{B}} \overbrace{\int_{o_1, o_2}^{p_1, p_2} \dots \int_{o_1, o_2}^{p_1, p_2}}^{r \text{ times}} \omega_{r+1}^0,$$


and

$$\tau' = \tau + \sum_{r \geq 1} \frac{\mathfrak{h}^r}{r!} \oint_{\mathcal{B}} \oint_{\mathcal{B}} \overbrace{\int_{o_1, o_2}^{p_1, p_2} \dots \int_{o_1, o_2}^{p_1, p_2}}^{r \text{ times}} \omega_{r+2}^0.$$

and its Taylor expansion in \mathfrak{h} generates the terms with $h = 0, d = 1, 2$ which were missing in the expansion of $\omega_0^{h,(d)}$. Eventually, taking into account the finite shift appearing in equation (75), the vertices are now associated with derivatives of theta functions evaluated at

$$\mathbf{w}_\bullet = \mathbf{a}(p_1) - \mathbf{a}(o_1) + \mathbf{a}(p_2) - \mathbf{a}(o_2) + \zeta,$$

namely



Thus, the result can be represented diagrammatically as follows:

$$\begin{aligned} & \psi_{\mathfrak{h}}^{[2][2]}(p_1, o_1; p_2, o_2) \\ &= \frac{e^c E(p_1, p_2) E(o_1, o_2)}{E(p_1, o_1) E(p_2, o_2) E(p_1, o_2) E(o_1, p_2)} \frac{\vartheta_{\bullet} \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right]}{\vartheta_{\bullet} \left[\begin{smallmatrix} \mu \\ \nu \end{smallmatrix} \right]} \frac{\hat{\psi}_{\mathfrak{h}}^{[2][2]}(p_1, o_1; p_2, o_2)}{\hat{\mathcal{T}}_{\mathfrak{h}}}, \end{aligned}$$

where

$$\epsilon = \frac{1}{\hbar} \left(\int_{o_1}^{p_1} v \, du + \int_{o_2}^{p_2} v \, du \right)$$

with

$$\ln \hat{\psi}_{\hbar}^{[2|2]}(p_1, o_1; p_2, o_2) = \sum \frac{\hbar^{-\chi_{\text{Euler}}}}{\# \text{Aut}} \text{weight} \left(\begin{array}{c} \text{Diagram} \end{array} \right), \quad (76)$$

where now some legs attached to surfaces are not contracted with legs of white vertices. Those open legs are associated to

$$\int_{o_1, o_2}^{p_1, p_2} = \int_{o_1}^{p_1} + \int_{o_2}^{p_2},$$

or, more generally,

$$\int_{o_1}^{p_1} + \dots + \int_{o_n}^{p_n},$$

if we wanted to compute the $n|n$ kernels. At each order χ , the sum consists of a finite number of connected graphs whose sum of Euler characteristics of surfaces is χ_{Euler} . Note that a surface with n punctures and h handles is associated to an ω_n^h , which can itself be expressed as a sum over pants decomposition (or equivalently skeleton graphs) of that surface (see Figure 1 and for a more detailed description [40, Section 3]).

A.4. Perturbative knot invariants to first orders. The central object in our conjecture concerning the asymptotics of the colored Jones was

$$\ln \hat{\psi}_{\hbar}^{[2|2]}(p, o; \iota(p), \iota(o)) = 2 \sum_{\chi \geq 0} \hbar^{-\chi} \tilde{j}_{\chi}(p),$$

and from equation (76) they acquire a diagrammatic representation. When $\iota_* = -\text{id}$, $\omega_n^{h,(d)}$ vanish whenever d is odd, so the only graphs with non-zero weight are those where each surface is contracted with an even number of legs incident to a white vertex. We give below the two first orders in diagrams in this case:

$$\begin{aligned}
 2\tilde{j}_1(p) &= \text{diagram 1} + \frac{1}{6} \text{diagram 2} + \frac{1}{2} \text{diagram 3} + \frac{1}{2} \text{diagram 4}, \\
 2\tilde{j}_2(p) &= \frac{1}{24} \text{diagram 5} + \frac{1}{2} \text{diagram 6} + \frac{1}{4} \text{diagram 7} + \frac{1}{2} \text{diagram 8} \\
 &+ \frac{1}{24} \text{diagram 9} + \frac{1}{8} \text{diagram 10} + \frac{1}{2} \text{diagram 11} + \text{diagram 12} \\
 &+ \frac{1}{6} \text{diagram 13} + \frac{1}{8} \text{diagram 14} + \frac{1}{2} \text{diagram 15} + \frac{1}{2} \text{diagram 16} \\
 &+ \frac{1}{4} \text{diagram 17} + \frac{1}{2} \text{diagram 18} + \frac{1}{24} \text{diagram 19},
 \end{aligned}$$

and we can compare this diagrammatic representation to the (black terms in) the expressions given in §5.4.

B. Some A-spectral curves

B.1. In Rolfsen classification. We present list of hyperbolic knot complement sorted by genus of their components, which is exhaustive up to 8 crossings (knots which do not appear have components of higher genus). Some knots with 9 and 10 crossings knots and once-punctured torus bundle (knot complement in lens spaces) were included. We also added $m129(3, 0)$, which is studied in [16], and is the orbifold obtained by $(0, 3)$ Dehn filling on the first cusp of $m129$. When the

A-polynomial have several components which are not of the form $(lm^a + b)$, we indicate their label (α) in exponent.

genus	knot complements
0	$10_{152}^{(1)}$
1	4_1 $7_4^{(1)}$ $8_{18}^{(1)}$ $8_{18}^{(2)}$ $8_{18}^{(3)}$ $9_{24}^{(1)}$ $9_{35}^{(1)}$ $9_{37}^{(1)}$ $9_{48}^{(1)}$ $9_{49}^{(1)}$ 10_{139} $10_{142}^{(1)}$ $10_{145}^{(1)}$ $10_{146}^{(1)}$ $10_{147}^{(1)}$ $10_{155}^{(1)}$ L^2R $m129(0,3)$
2	5_2 Pretzel $(-2, 3, 7)$ $7_4^{(2)}$ $7_7^{(1)}$ $8_5^{(1)}$ $9_{37}^{(2)}$ $10_{136}^{(1)}$ $10_{154}^{(1)}$ $10_{160}^{(1)}$ $10_{163}^{(1)}$
3	6_1 $7_7^{(2)}$ $8_5^{(2)}$ $9_{35}^{(2)}$ $9_{47}^{(1)}$ $9_{48}^{(2)}$ LR^3
4	7_2 8_{21} $9_{10}^{(1)}$ $9_{23}^{(1)}$ 9_{46} $9_{49}^{(2)}$ $10_{61}^{(1)}$ $10_{145}^{(2)}$ $10_{146}^{(2)}$
5	8_1 8_{20} $9_{10}^{(2)}$ $9_{17}^{(1)}$ $9_{41}^{(1)}$ $9_{47}^{(2)}$ $10_{142}^{(2)}$ $10_{144}^{(1)}$
6	$8_{16}^{(1)}$ $9_{17}^{(2)}$ $9_{31}^{(1)}$ $10_{136}^{(2)}$ $10_{152}^{(2)}$
7	6_3 8_2 9_2 $9_{16}^{(1)}$ $9_{41}^{(2)}$
8	$9_{37}^{(3)}$ 9_{42} 10_1 $10_{61}^{(2)}$ $10_{138}^{(1)}$ $10_{138}^{(2)}$ 10_{140} $10_{141}^{(2)}$
9	7_5 8_3 8_4 $8_{16}^{(2)}$ 9_3 $9_{10}^{(1)}$ $9_{16}^{(2)}$ $9_{28}^{(1)}$ $10_{62}^{(1)}$ $10_{154}^{(2)}$ $10_{155}^{(2)}$
10	10_2 10_{125} 10_{132}
11	9_4 $9_{23}^{(11)}$
12	7_6 8_{15} $9_{38}^{(1)}$
13	7_3 8_{11} 9_{43} $10_{144}^{(2)}$
14	8_7 9_5
15	8_9 8_{10} 9_6 10_3 $10_{62}^{(2)}$ 10_{128} $10_{146}^{(3)}$ 10_{161} 10_{162}
16	8_6 $9_{28}^{(16)}$ $9_{29}^{(1)}$ 10_4
17	$9_{31}^{(2)}$ 10_8
18	10_{126} 10_{143}
19	8_8 9_{44}
20	9_9 10_5 $10_{160}^{(2)}$
21	8_{12} 9_7 $9_{29}^{(2)}$
22	10_{153}
23	7_4 8_{13} $10_{147}^{(2)}$
24	9_8
25	9_{11} 10_{133}
26	8_{14} $9_{38}^{(2)}$ 9_{46}
28	9_{12}

B.2. In tetrahedron census. We present an exhaustive list of hyperbolic 3-manifolds triangulated with atmost 6 tetrahedra. Those which are complement of a knot with atmost ten crossings were rather included in Figure B.1. For all the knots in this table, we observe that the A-polynomial has a single component.

genus	knot complements
2	$k3_1$
5	$k4_3$ $k4_4$ $k5_{11}$
6	$k5_1$
7	$k6_{32}$
8	$k5_5$ $k5_{10}$ $k5_{14}$ $k5_{15}$ $k5_{16}$ $k6_2$ $k6_{33}$
9	$k5_4$ $k5_6$ $k6_{42}$
10	$k5_7$ $k6_{19}$
11	$k5_{18}$ $k6_5$ $k6_{39}$
12	$k5_{13}$ $k5_{17}$
13	$k6_3$ $k6_{22}$ $k6_{31}$ $k6_{38}$
14	$k6_{10}$ $k6_{11}$ $k6_{14}$ $k6_{29}$
15	$k6_4$ $k6_6$
16	$k6_{16}$
17	$k6_{15}$ $k6_{17}$ $k6_{25}$ $k6_{27}$
18	$k6_7$ $k6_{18}$
20	$k6_{13}$ $k6_{37}$
21	$k6_8$ $k6_{12}$
22	$k6_9$ $k6_{30}$
24	$k6_{21}$ $k6_{35}$

B.3. Properties. We present properties of spectral curves for various knots. Each block collect equivalent curves modulo birational transformations. Note that the transformation $l \mapsto C l^\pm m^a$ implies $\omega_n^h \mapsto (\pm)^n \omega_n^h$. The column g gives the genus of the curve, and the column g_l gives the genus of the quotient curve \mathcal{C}/ι , i.e. the number of +1 eigenvalues of ι_* . In the column H, we indicate if ι coincide or not with the hyperelliptic involution. If this is the case, we necessarily have $\iota_* = -id$. $|\{a\}|$ indicates the number of ramification points. They are all simple, except when we indicate with a superscript +1 the presence of one extra ramification point of order 3 at $(m, l) = (-1, 1)$. We then indicate the minimal positive integer ζ such that $2\zeta \cdot \{m, l\} = 0 \in \text{in } K_2(\mathcal{C})$. When the knot is amphichiral and when the component is stable under $\alpha(m, l) = (1/m, l)$, we indicate the number of +1 eigenvalues of the induced map α_* in homology. We put question marks when we could not obtain the answer in a reasonable time with MAPLE.

knot	g	g_i	H	$ \{a\} $	ζ	amphicheiral	α_*
4_1	1	0	yes	4	1	yes	0
$7_4^{(1)}$	deduced from 4_1 by $l \rightarrow lm^4$				1	no	
$8_{18}^{(1)}$	1	0	no	4	2	yes	
$8_{18}^{(2)}$	deduced from $8_{18}^{(1)}$ by $l \rightarrow l^{-1}$				2	yes	
$9_{24}^{(1)}$	idem 4_1				1	no	
$9_{37}^{(1)}$	idem 4_1				1	no	
$9_{49}^{(1)}$	idem 4_1 with $l \rightarrow lm^8$				1	no	
$10_{142}^{(1)}$	idem 4_1 with $l \rightarrow lm^{12}$				1	no	
$10_{145}^{(1)}$	idem 4_1 with $l \rightarrow -lm^{-2}$				2	no	
$10_{146}^{(1)}$	idem 4_1 with $l \rightarrow -lm^{-6}$				2	no	
$10_{147}^{(1)}$	idem 4_1 with $l \rightarrow -l^{-1}m^2$				2	no	
$10_{155}^{(1)}$	idem $8_{18}^{(1)}$ with $l \rightarrow l^{-1}m^{-4}$				2	no	
$8_{18}^{(3)}$	1	0	?	6	1	yes	0
$9_{35}^{(1)}$	1	0	yes	4^{+1}	2	no	
$9_{48}^{(1)}$	1	0	no	2^{+1}	1	no	
10_{139}	1	0	yes	6	2	no	
L^2R	1	0	yes	4	1	no	
5_2	2	0	yes	8	2	no	
$7_7^{(1)}$	idem 5_2 with $l \rightarrow lm^{-4}$				2	no	
$8_5^{(1)}$	idem 5_2 with $l \rightarrow l^{-1}m^{-12}$				2	no	
$10_{154}^{(1)}$	2	0	yes	8	2	no	
$10_{160}^{(1)}$	idem 5_2 with $l \rightarrow lm^8$				2	no	
$10_{163}^{(1)}$	idem 5_2 with $l \rightarrow lm^{-8}$				2	no	
$P(-2, 3, 7)$	idem $10_{154}^{(1)}$ with $l \rightarrow l^{-1}m^{-26}$				2	no	
$k3_1$	idem $10_{154}^{(1)}$ with $l \rightarrow -lm^{-26}$				2	no	
$7_4^{(2)}$	2	0	yes	6	2	no	
$9_{37}^{(2)}$	idem $7_4^{(2)}$ with $l \rightarrow l^{-1}m^{-4}$				2	no	
$10_{136}^{(1)}$	2	0	yes	6	2	no	

knot	g	g_i	$ \{a\} $	ζ	amphicheiral	α_*
$\mathbf{6}_1$	3	0	12	1	no	
$\mathbf{9}_{35}^{(2)}$	idem $\mathbf{6}_1$ with $l \rightarrow lm^4$			1	no	
$\mathbf{9}_{47}^{(1)}$	idem $\mathbf{6}_1$ with $l \rightarrow lm^8$			1	no	
$\mathbf{9}_{48}^{(2)}$	idem $\mathbf{6}_1$ with $l \rightarrow l^{-1}m^{-4}$			1	no	
$\mathbf{7}_7^{(2)}$	3	0	8	2	no	
$\mathbf{8}_5^{(2)}$	3	0	14	2	no	
LR^3	3	0	10	1	no	
$\mathbf{7}_2$	4	0	16	2	no	
$\mathbf{8}_{21}$	4	1	12	2	no	
$\mathbf{9}_{10}$	4	0	12	2	no	
$\mathbf{9}_{23}^{(1)}$	4	0	12	2	no	
$\mathbf{9}_{46}$	4	1	10	1	no	
$\mathbf{9}_{49}^{(2)}$	4	1	2	2	no	
$\mathbf{10}_{61}^{(1)}$	4	0	16	2	no	
$\mathbf{10}_{146}^{(2)}$	idem $\mathbf{10}_{61}^{(2)}$ with $l \rightarrow lm^{-8}$			2	no	
$\mathbf{10}_{145}^{(2)}$	4	1	14	2	no	
$\mathbf{8}_1$	5	0	20	1	no	
$\mathbf{8}_{20}$	5	1	12	2	no	
$\mathbf{9}_{17}^{(1)}$	5	0	18	2	no	
$\mathbf{10}_{144}^{(1)}$	idem $\mathbf{9}_{17}^{(2)}$ with $l \rightarrow l^{-1}m^8$			1	no	
$\mathbf{10}_{142}^{(2)}$	5	1	18	2	no	
$\mathbf{k4}_3$	5	0	16	2	no	
$\mathbf{k4}_4$	5	0	18	2	no	
$\mathbf{k5}_{11}$	5	0	14	1	no	
$\mathbf{10}_{136}^{(2)}$	6	2	16	2	no	
$\mathbf{6}_3$	7	1	12	2	yes	3
$\mathbf{9}_{16}^{(1)}$	idem $\mathbf{6}_3$ with $l \rightarrow lm^{12}$			1	no	
$\mathbf{9}_2$	7	0	24	2	no	

References

- [1] M. Aganagic, A. Klemm, M. Mariño, and C. Vafa, The topological vertex. *Comm. Math. Phys.* **254** (2005), no. 2, 425–478. [MR 2117633](#) [Zbl 1114.81076](#)
- [2] M. Aganagic and C. Vafa, Large n duality, mirror symmetry, and a q -deformed A-polynomial for knots. Preprint 2012. [arXiv:1204.4709](#) [hep-th]
- [3] J. Ambjørn and Y. Makeenko, Properties of loop equations for the Hermitian matrix model and for two-dimensional gravity. *Mod. Phys. Lett. A* **5** (1990), no. 22, 1753–1763. [Zbl Makeenko](#)
- [4] C. Beasley, Localization for Wilson loops in Chern–Simons theory. *Adv. Theor. Math. Phys.* **17** (2013), no. 1, 1–240. [MR 3066489](#) [Zbl 1290.81057](#)
- [5] M. Bergère, G. Borot, and B. Eynard, Rational differential systems, loop equations, and application to the q th Reductions of KP. *Ann. Henri Poincaré*, online first, January 2015.
- [6] M. Bergère and B. Eynard, Determinantal formulae and loop equations. Preprint 2009. [arXiv:0901.3273](#) [math-ph]
- [7] M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, Kodaira–Spencer theory of gravity and exact results for quantum string amplitudes. *Comm. Math. Phys.* **165** (1994), no. 2, 311–428. [MR 1301851](#) [Zbl 0815.53082](#)
- [8] M. Bertola, Boutroux curves with external field: equilibrium measures without a variational problem. *Anal. Math. Phys.* **1** (2011), no. 2-3, 167–211. [MR 2853811](#) [Zbl 1259.33021](#)
- [9] M. Bertola and M. Mo, Commuting difference operators, spinor bundles and the asymptotics of orthogonal polynomials with respect to varying complex weights. *Adv. Math.* **220** (2009), no. 1, 154–218. [MR 2462838](#) [Zbl 1153.39027](#)
- [10] D. Bertrand and W. Zudilin, On the transcendence degree of the differential field generated by Siegel modular forms. *J. Reine Angew. Math.* **554** (2003), 47–68. [MR 1952168](#) [Zbl 1130.11020](#)
- [11] S. Bloch, Applications of the dilogarithm function in algebraic K-theory and algebraic geometry. In M. Nagata (ed.), *Proceedings of the International Symposium on Algebraic Geometry*. A Taniguchi Symposium. Held at Kyoto University, Kyoto, January 10–14, 1977. Kinokuniya Book Store Co., Ltd., Tokyo, 103–114. [MR 0578856](#) [MR 0578850](#) (collection) [Zbl 0416.18016](#) [Zbl 0394.00009](#) (collection)
- [12] G. Borot and B. Eynard, Tracy–Widom GUE law and symplectic invariants. Preprint 2010. [arXiv:1011.1418](#) [nlin.SI]
- [13] G. Borot and B. Eynard, Geometry of spectral curves and all order dispersive integrable system. *SIGMA Symmetry Integrability Geom. Methods Appl.* **8** (2012), Paper 100, 53 pp. [MR 3007259](#) [Zbl 1270.14017](#)
- [14] V. Bouchard, J. Hutchinson, P. Loliencar, M. Meiers, and M. Rupert, A generalized topological recursion for arbitrary ramification. *Ann. Henri Poincaré* **15** (2014), no. 1, 143–169. [MR 3147410](#) [Zbl 1291.81212](#)

- [15] V. Bouchard, A. Klemm, M. Mariño, and S. Pasquetti, Remodeling the B-model. *Comm. Math. Phys.* **287** (2009), 117–178. [MR 2480744](#) [Zbl 1178.81214](#)
- [16] D. Boyd, F. Rodriguez-Villegas, and N. Dunfield, Mahler’s measure and the dilogarithm (II). Preprint 2003. [arXiv:math/0308041](#) [math.NT]
- [17] A. Brini, B. Eynard, and M. Mariño, Torus knots and mirror symmetry. *Ann. Henri Poincaré* **13** (2012), no. 8, 1873–1910. [MR 2994763](#) [Zbl 1256.81086](#)
- [18] J. Bruinier, G. van der Geer, G. Harder, and D. Zagier, *The 1 – 2 – 3 of modular forms*. Lectures from the Summer School on Modular Forms and their Applications held in Nordfjordeid, June 2004. Edited by K. Ranestad. Universitext. Springer, Berlin, 2008. [MR 2385372](#) [Zbl 1197.11047](#)
- [19] D. Calegari, Real places and torus bundles. *Geom. Dedicata* **118** (2006), 209–227. [MR 2239457](#) [Zbl 05046847](#)
- [20] A. Champanerkar, *A-polynomials and Bloch invariants of hyperbolic 3-manifolds*. PhD Thesis. Columbia University, New York, N.Y., 2003 [MR 2704573](#) <http://www.math.csi.cuny.edu/~abhjit/research.html>
- [21] L. Chekhov and B. Eynard, Hermitean matrix model free energy: Feynman graph technique for all genera. *J. High Energy Phys.* **2006** (2006), no. 3, article id. 014.
- [22] S. Chern and J. Simons, Some cohomology classes in principal fiber bundles and their application to Riemannian geometry. *Proc. Nat. Acad. Sci. U.S.A.* **68** (1971), 791–794. [MR 0279732](#) [Zbl 0209.25401](#)
- [23] D. Cooper, M. Culler, H. Gillet, D. Long, and P. Shalen, Plane curves associated to character varieties of 3-manifolds. *Invent. Math.* **118** (1994), no. 1, 47–84. [MR 1288467](#) [Zbl 0842.57013](#)
- [24] M. Culler, A-polynomials in the tetrahedral census of knots. http://www.indiana.edu/~knotinfo/descriptions/tetrahedral_census_name.html
- [25] M. Culler, Personal webpage, <http://homepages.math.uic.edu/~culler>
- [26] R. Dijkgraaf and H. Fuji, The volume conjecture and topological strings. *Fortsch. Phys.* **57** (2009), no. 9 825–856. [MR 2567388](#) [Zbl 1210.81081](#)
- [27] R. Dijkgraaf, H. Fuji, and M. Manabe, The volume conjecture, perturbative knot invariants, and recursion relations for topological strings. *Nucl. Phys. B* **849** (2011), no. 1, 166–211. [MR 2795276](#) [Zbl 1215.81082](#)
- [28] T. Dimofte, Quantum Riemann surfaces in Chern–Simons theory. *Adv. Theor. Math. Phys.* **17** (2013), no. 3, 479–599. [MR 3250765](#) [Zbl 1304.81143](#)
- [29] T. Dimofte and S. Garoufalidis, The quantum content of the gluing equations. *Geom. Topol.* **17** (2013), no. 3, 1253–1315. [MR 3073925](#) [Zbl 1283.57017](#)
- [30] T. Dimofte, S. Gukov, J. Lennels, and D. Zagier, Exact results for perturbative Chern–Simons theory with complex gauge group. *Commun. Number Theory Phys.* **3** (2009), no. 2, 363–443. [MR 2551896](#) [Zbl 1214.81151](#)

- [31] T. Dokchitser, R. de Jeu, and D. Zagier, Numerical verification of Beilinson's conjecture for K_2 of hyperelliptic curves. *Compos. Math.* **142** (2006), no. 2, 339–373. [MR 2218899](#) [Zbl 1186.11037](#)
- [32] N. Dunfield, S. Gukov, and J. Rasmussen, The superpolynomial for knot homologies. *Experiment. Math.* **15** (2006), no. 2, 129–160. [MR 2253002](#) [Zbl 1118.57012](#)
- [33] S. Elitzur, G. Moore, A. Schwimmer, and N. Seiberg, Remarks on the canonical quantization of the Chern–Simons–Witten theory. *Nucl. Phys. B* **326** (1989), no. 1, 108–134. [MR 1025431](#)
- [34] B. Eynard, All genus correlation functions for the Hermitian 1-matrix model. *J. High Energy Phys.* **2004** (2004), no. 11, article id. 031.
- [35] B. Eynard, Large N expansion of convergent matrix integrals, holomorphic anomalies, and background independence. *J. High Energy Phys.* **2009** (2009), no. 3, article id. 003.
- [36] B. Eynard, Invariants of spectral curves and intersection theory of moduli spaces of complex curves. *Commun. Number Theory Phys.* **8** (2014), no. 3, 541–588. [MR 3282995](#) [Zbl 06394371](#)
- [37] B. Eynard and M. Mariño, A holomorphic and background independent partition function for matrix models and topological strings. *J. Geom. Phys.* **61** (2011), no. 7, 1181–1202. [MR 2788322](#) [Zbl 1215.81084](#)
- [38] B. Eynard and N. Orantin, Invariants of algebraic curves and topological expansion. *Comm. Number Theory and Physics* **1** (2007), no. 2, 347–452. [MR 2346575](#) [Zbl 1161.14026](#)
- [39] B. Eynard and N. Orantin, Topological expansion of mixed correlations in the Hermitian 2 matrix model and $x-y$ symmetry of the F_g invariants. *J. Phys. A* **41** (2008), no. 1, Article id. 015203. [MR 2450700](#) [Zbl 1134.81040](#)
- [40] B. Eynard and N. Orantin, Topological recursion in enumerative geometry and random matrices. *J. Phys. A* **42** (2009), no. 29, Article id. 293001. [MR 2519749](#) [Zbl 1177.82049](#)
- [41] B. Eynard and N. Orantin, Computation of open Gromov–Witten invariants for toric Calabi–Yau 3-folds by topological recursion, a proof of the BKMP conjecture. Preprint 2012, [arXiv:1205.1103](#) [math-ph]
- [42] C. Frohman, R. Gelca, and W. Lofaro, The A-polynomial from the noncommutative viewpoint. *Trans. Amer. Math. Soc.* **354** (2002), no. 2, 735–747. [MR 1862565](#) [Zbl 0980.57002](#)
- [43] H. Fuji, S. Gukov, and P. Sułkowski, Super-A-polynomial for knots and BPS states. *Nuclear Phys. B* **867** (2013), no. 2, 506–546. [MR 2992793](#) [Zbl 1262.81170](#)
- [44] H. Fuji, S. Gukov, and P. Sułkowski Volume conjecture: refined and categorified. *Adv. Theor. Math. Phys.* **16** (2012), no. 6, 1669–1777. [MR 3065082](#) [Zbl 1282.57016](#)

- [45] S. Garoufalidis, On the characteristic and deformation varieties of a knot. In C. Gordon and Y. Rieck (eds.), *Proceedings of the Casson Fest*. Papers from the 28th University of Arkansas Spring Lecture Series “The Andrews–Curtis and the Poincaré Conjectures” held in Fayetteville, AR, April 10–12, 2003 and from the Conference on the Topology of Manifolds of Dimensions 3 and 4 held in honor of the 60th birthday of A. Casson at the University of Texas, Austin, TX, May 19–21, 2003. *Geometry & Topology Monographs*, 7. Geometry & Topology Publications, Coventry, 2004, 291–309. [MR 2172488](#) [MR 2169266](#) (collections) [Zbl 1080.57014](#) [Zbl 1080.57014](#) (collections)
- [46] S. Garoufalidis and T. Lê, The colored Jones function is q -holonomic. *Geom. Topol.* **9** (2005), 1253–1293. [MR 2174266](#) [Zbl 1078.57012](#)
- [47] S. Garoufalidis and T. Lê, On the volume conjecture for small angles. Preprint 2005. [arXiv:0502163](#) [math.GT]
- [48] S.+ Garoufalidis and T. Lê, Is the Jones polynomial of a knot really a polynomial? *J. Knot Theory Ramifications* **15** (2006), no. 8, 983–1000. [MR 2275093](#) [Zbl 1126.57007](#)
- [49] R. Gopakumar and C. Vafa, On the gauge theory/geometry correspondence. *Adv. Theor. Math. Phys.* **3** (1999), no. 5, 1415–1443. [MR 1796682](#) [Zbl 0972.81135](#)
- [50] F. Guéritaud and D. Futer, On canonical triangulations of once-punctured torus bundles and two-bridge link complements. *Geom. Topology* With an appendix by David Futer. *Geom. Topol.* **10** (2006), 1239–1284. [MR 2255497](#) [Zbl 1130.57024](#)
- [51] S. Gukov, Three-dimensional quantum gravity, Chern–Simons theory, and the A-polynomial. *Comm. Math. Phys.* **255** (2005), no. 3, 577–627. [MR 2134725](#) [Zbl 1115.57009](#)
- [52] S. Gukov and H. Murakami, $SL_2(\mathbb{C})$ Chern–Simons theory and the asymptotic behavior of the colored Jones polynomial. *Lett. Math. Phys.* **86** (2008), no. 2–3, 79–98. [MR 2465747](#) [Zbl 1183.57012](#)
- [53] S. Gukov and P. Sułkowski, A-polynomial, B-model, and quantization. *J. High Energy Phys.* **2012** (2012), no. 2, article id. 070. [MR 2996110](#)
- [54] K. Hikami, Generalized volume conjecture and the A-polynomials: the Neumann–Zagier potential function as a classical limit of the partition function. *J. Geom. Phys.* **57** (2007), 1895–1940. [MR 2330673](#) [Zbl 1139.57013](#)
- [55] V. F. R. Jones, A polynomial invariant for knots via von Neumann algebras. *Bull. Amer. Math. Soc. (N.S.)* **12** (1985), no. 1 103–111. [MR 0766964](#) [Zbl 0564.57006](#)
- [56] J. Källén, Cohomological localization of Chern–Simons theory. *J. High Energy Phys.* **2011** 2011, no. 8, article id. 008. [MR 2876080](#) [Zbl 1298.81347](#)
- [57] A. Kapaev, Monodromy approach to the scaling limits in the isomonodromy systems. *Teoret. Mat. Fiz.* **137** (2003), no. 3, 393–407. In Russian. English transl., *Theoret. and Math. Phys.* **137** (2003), no. 3, 1691–1702. [MR 2084149](#) [Zbl 1178.34112](#)
- [58] R. Kashaev, The hyperbolic volume of knots from the quantum dilogarithm. *Lett. Math. Phys.* **39** (1997), no. 3, 269–275. [MR 1434238](#) [Zbl 0876.57007](#)

- [59] R. Kashaev and O. Tirkkonen, Proof of the volume conjecture for torus knots. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **269** (2000), *Vopr. Kvant. Teor. Polya i Stat. Fiz.* 16, 262–268, 370. In Russian. English transl., *J. Math. Sci. (N. Y.)* **115** (2003), no. 1, 2033–2036. [MR 1805865](#) [Zbl 1030.57021](#)
- [60] I. Krichever, Methods of algebraic geometry in the theory of nonlinear equations. *Usp. Mat. Nauk.* **32** (1977), no. 6(198), 183–208. In Russian. English transl., *Russ. Math. Surv.* **32**(1977), no. 6, 185–213. [Zbl 0372.35002](#) [Zbl 0386.35002](#) (transl.)
- [61] R. Lawrence and L. Rozansky, Witten–Reshetikhin–Turaev invariants of Seifert manifolds. *Comm. Math. Phys.* **205** (1999), no. 2, 287–314. [MR 1712599](#) [Zbl 0966.57017](#)
- [62] R. Lawrence and D. Zagier, Modular forms and quantum invariants of 3-manifolds. *Asian J. Math.* **3** (1999), 93–108. Sir Michael Atiyah: a great mathematician of the twentieth century. [MR 1701924](#) [Zbl 1024.11028](#)
- [63] T. Lê and A. Tran, On the AJ conjecture for knots. Preprint 2011. [arXiv:1111.525](#) [math.GT]
- [64] W. Li and Q. Wang, On the generalized volume conjecture and regulator. *Comm. Contemporary Math.* **10** (2008), suppl. 1, 1023–1032. [MR 2468376](#) [Zbl 1221.57021](#)
- [65] M. Mariño, Chern–Simons theory and topological strings. *Rev. Mod. Phys.* **77** (2005), no. 2, 675–720. [MR 2168778](#) [Zbl 1205.81013](#)
- [66] A. Mednykh and A. Vesnin, Hyperbolic volumes of Fibonacci manifolds. *Sibirsk. Mat. Zh.* **36** (1995), no. 2, 266–277, i. In Russian. English transl., *Siberian Math. J.* **36** (1995), no. 2, 235–245. [MR 1340395](#) [Zbl 0865.57012](#)
- [67] P. Menal-Ferre and J. Porti, Twisted cohomology for hyperbolic three manifolds. *Osaka J. Math.* (2012), no. 3, 741–769. [MR 2993065](#) [Zbl 1255.57018](#)
- [68] R. Meyerhoff, *The Chern–Simons invariant for hyperbolic 3-Manifolds*. Ph.D. thesis. Princeton University, Princeton, N.Y., 1981. [MR 2631318](#)
- [69] J. Milnor, *Introduction to algebraic K-theory*. Annals of Mathematics Studies, 72. Princeton University Press, Princeton, N.Y., and University of Tokyo Press, Tokyo, 1971. [MR 0349811](#) [Zbl 0237.18005](#)
- [70] D. Mumford, *Tata lectures on Theta*. II. Jacobian theta functions and differential equations. With the collaboration of C. Musili, M. Nori, E. Previato, M. Stillman, and H. Umemura. Progress in Mathematics, 43. Birkhäuser Boston, Boston, MA, 1984. [MR 0742776](#) [Zbl 0549.14014](#)
- [71] H. Murakami, Some limits of the colored Jones polynomials of the figure-eight knot. *Kyungpook Math. J.* **44** (2004), no. 3, 369–383. [MR 2095421](#) [Zbl 1079.57011](#)
- [72] H. Murakami, An introduction to the volume conjecture and its generalizations. *Acta Math. Vietnam.* **33** (2008), no. 3, 219–253. [MR 2501844](#) [Zbl 1179.57023](#)
- [73] H. Murakami and J. Murakami, The colored Jones polynomials and the simplicial volume of a knot. *Acta Math.* **186** (2001), no. 1, 85–104. [MR 1828373](#) [Zbl 0983.57009](#)

- [74] W. Neumann, Combinatorics of triangulations and the Chern–Simons invariant for hyperbolic 3-manifolds. In B. Apanasov, W. D. Neumann, A. W. Reid, and L. Siebenmann (eds.), *Topology '90*. Papers from the Research Semester in Low-dimensional Topology held at Ohio State University, Columbus, Ohio, February–June 1990. Ohio State University Mathematical Research Institute Publications, 1. Walter de Gruyter & Co., Berlin, 1992, 243–271. [MR 1184415](#) [MR 1184397](#) (collection) [Zbl 0768.57006](#) [Zbl 0747.00024](#) (collection)
- [75] W. Neumann and A. Reid, Arithmetic of hyperbolic manifolds. In B. Apanasov, W. D. Neumann, A. W. Reid, and L. Siebenmann (eds.), *Topology '90*. Papers from the Research Semester in Low-dimensional Topology held at Ohio State University, Columbus, Ohio, February–June 1990. Ohio State University Mathematical Research Institute Publications, 1. Walter de Gruyter & Co., Berlin, 1992, 273–310. [MR 1184416](#) [MR 1184397](#) (collection) [Zbl 0777.57007](#) [Zbl 0747.00024](#) (collection)
- [76] W. Neumann and J. Yang, Rationality problems for K-theory and Chern–Simons invariants of hyperbolic 3-manifolds. *Enseign. Math.* (2) **41** (1995), no. 3–4, 281–296. [MR 1365848](#) [Zbl 0861.57022](#)
- [77] W. Neumann and J. Yang, Bloch invariants of hyperbolic 3-manifolds. *Duke Math. J.* **96** (1999), no. 1, 29–59. [MR 1663915](#) [Zbl 0943.57008](#)
- [78] W. Neumann and D. Zagier, Volumes of hyperbolic three-manifolds. *Topology* **24** (1985), no. 3, 307–332. [MR 0815482](#) [Zbl 0589.57015](#)
- [79] N. Orantin, Chain of matrices, loop equations and topological recursion. In G. Akemann, J. Baik, and Ph. Di Francesco (eds), *The Oxford handbook of random matrix theory*. Oxford University Press, Oxford, 2011, Chapter 16, 329–352. [MR 2932636](#) [MR 2920518](#) (collection) [Zbl 1236.15073](#) [Zbl 1225.15004](#) (collection)
- [80] J.-P. Otal, Le théorème d'hyperbolisation pour les variétés fibrées de dimension 3. *Astérisque* 235, 1996. [MR 0855.57003](#) [Zbl 0855.57003](#)
- [81] A. Prats-Ferrer, New recursive residue formulas for the topological expansion of the Cauchy matrix model. *J. High Energy Phys.* **2010** (2010), no. 10, article id. 090. [MR 2780511](#) [Zbl 1291.81309](#)
- [82] H. Rauch, Weierstrass points, branch points, and moduli of Riemann surfaces. *Comm. Pure Appl. Math.* **12** (1959), 543–560. [MR 0110799](#) [Zbl 0091.07301](#)
- [83] N. Reshetikhin and V. Turaev, Ribbon graphs and their invariants derived from quantum groups. *Comm. Math. Phys.* **127** (1990), no. 1, 1–26. [MR 1036112](#) [Zbl 0768.57003](#)
- [84] W. Thurston, *The geometry and topology of 3-manifolds*. Princeton Lecture Notes, 1980. Chapter 4. <http://library.msri.org/books/gt3m>
- [85] V. Turaev, The Yang–Baxter equation and invariants of links. *Invent. Math.* **92** (1988), no. 3, 527–553. [MR 0939474](#) [Zbl 0648.57003](#)
- [86] E. Witten, Quantum field theory and the Jones polynomial. *Comm. Math. Phys.* **121** (1989), no. 3, 351. [MR 0990772](#) [Zbl 0667.57005](#)

- [87] E. Witten, Chern–Simons gauge theory as a string theory. In H. Hofer, C. H. Taubes, A. Weinstein, and E. Zehnder (eds.), *The Floer memorial volume*. Progress in Mathematics, 133. Birkhäuser, Basel, 1995, 637–678. [MR 1362846](#) [MR 1362819](#) (collection) [Zbl 0844.58018](#) [Zbl 0824.00019](#) (collection)
- [88] T. Yoshida, The η -invariant of hyperbolic 3-manifolds. *Invent. Math.* **81** (1985), 473–514. [MR 0807069](#) [Zbl 0594.58012](#)
- [89] D. Zagier, Quantum modular forms. In E. Blanchard, D. Ellwood, M. Khalkhali, M. Marcolli, H. Moscovici, and S. Popa (eds.), *Quanta of maths*. Including papers from the Conference on Non Commutative Geometry held in honor of A. Connes in Paris, March 29–April 6, 2007. Clay Mathematics Proceedings, 11. American Mathematical Society, Providence, R.I., and Clay Mathematics Institute, Cambridge, MA, 2010, 659–675. [MR 2757599](#) [MR 2724742](#) (collection) [Zbl](#) [Zbl 1206.00042](#) (collection)
- [90] X. Zhang, The C -polynomial of a knot. *Topology App.* **139** (2004), no. 1-3, 185–198. [MR 2051104](#) [Zbl 1054.57017](#)
- [91] S. Zhu, The higher order terms in asymptotic expansion of color Jones polynomials. Preprint 2011. [arXiv:1104.0403](#) [math.GN]

Received September 13, 2012

Gaëtan Borot, Département de Mathématiques, Université de Genève, 1211 Genève, Switzerland

e-mail: gborot@mpim-bonn.mpg.de

Bertrand Eynard, Institut de Physique Théorique, CEA Saclay, 91191 Gif-sur-Yvette, France

e-mail: bertrand.eynard@cea.fr