

Fully extended r -spin TQFTs

Nils Carqueville and Lóránt Szegedy

Abstract. We prove the r -spin cobordism hypothesis in the setting of (weak) 2-categories for every positive integer r : the 2-groupoid of 2-dimensional fully extended r -spin TQFTs with given target is equivalent to the homotopy fixed points of an induced Spin_2^r -action. In particular, such TQFTs are classified by fully dualisable objects together with a trivialisation of the r th power of their Serre automorphisms. For $r = 1$, we recover the oriented case (on which our proof builds), while ordinary spin structures correspond to $r = 2$.

To construct examples, we explicitly describe Spin_2^r -homotopy fixed points in the equivariant completion of any symmetric monoidal 2-category. We also show that every object in a 2-category of Landau–Ginzburg models gives rise to fully extended spin TQFTs and that half of these do not factor through the oriented bordism 2-category.

1. Introduction and summary

The spin group Spin_n in dimension n is by definition the double cover of the group of rotations SO_n in Euclidean space \mathbb{R}^n . A spin structure on an n -dimensional oriented manifold is a lift of its tangent bundle along the covering $\mathrm{Spin}_n \rightarrow \mathrm{SO}_n$. Such geometric structures and their close cousins in Lorentzian geometry are fundamental in theoretical physics, since, e.g., electrons are classically modelled as sections of spin bundles.

More generally, for any continuous group homomorphism $\xi: G \rightarrow \mathrm{GL}_n$, a tangential structure on an n -dimensional manifold M is a principal G -bundle on M together with a bundle map to the frame bundle of M that is compatible with ξ (see Section 2.1 for details). The case of spin structures is precisely when ξ is the covering map $\mathrm{Spin}_n \rightarrow \mathrm{SO}_n$ post-composed with the inclusion $\mathrm{SO}_n \subset \mathrm{GL}_n$; in the case of orientations, ξ is just that inclusion, while in the case of framings, ξ is the inclusion of the trivial group into GL_n .

Given the relevance of spin structures in physics and the motivation to study functorial topological quantum field theories (TQFTs) as a means to gain insight into physics, it is natural to consider spin TQFTs. These are (higher) symmetric monoidal functors on (higher) categories of bordisms with prescribed spin structures. The case

of closed spin TQFTs in dimension $n = 2$ was first considered in [1, 7, 33, 37], and in [1, 39], it was classified¹ in terms of “closed Λ_2 -Frobenius algebras” (see Section 2.2 for the definition). Such algebraic structures formalise the relation between topological Neveu–Schwarz and Ramond sectors, examples of which can be obtained as a \mathbb{Z}_2 -graded version of the centre construction of [30]. In particular, there is a (1|1)-dimensional example in $\text{Vect}_{\mathbb{C}}^{\mathbb{Z}_2}$ whose associated TQFT computes the Arf invariant of spin surfaces. Not many other explicit examples have been studied in the literature, and all previously known classes of examples are constructed from semisimple algebraic data.

In the setting of symmetric monoidal (∞, n) -categories, fully extended TQFTs with G -structure are widely believed to be classified by homotopy fixed points of a G -action (induced from the G -action on framed bordisms) on the maximal ∞ -subgroupoids of fully dualisable objects in the target (∞, n) -categories. This is described in significant, yet non-exhaustive, detail in [31]. To our knowledge, this general version of the cobordism hypothesis, originally put forward in [3], is established as a theorem only up to a completion of the extended proof sketch in [31], or up to a conjecture on the relation between factorisation homology and adjoints; see [2, Conjecture 1.2].

On the other hand, in dimension $n = 2$ and in the setting of (weak) 2-categories, the cobordism hypothesis for the framed and oriented case was proved explicitly in [34] and [25–27], respectively: for any symmetric monoidal 2-category \mathcal{B} , the 2-groupoid of fully extended framed TQFTs $\text{Bord}_{2,1,0}^{\text{fr}} \rightarrow \mathcal{B}$ is equivalent to the maximal sub-2-groupoid $(\mathcal{B}^{\text{fd}})^{\times}$ of fully dualisable objects in \mathcal{B} , while fully extended oriented TQFTs $\text{Bord}_{2,1,0}^{\text{or}} \rightarrow \mathcal{B}$ are described by SO_2 -homotopy fixed points. The latter are objects of a 2-groupoid

$$[(\mathcal{B}^{\text{fd}})^{\times}]^{\text{SO}_2}$$

and correspond to pairs (α, λ) , where $\alpha \in \mathcal{B}^{\text{fd}}$ and $\lambda: S_{\alpha} \cong 1_{\alpha}$ is a trivialisation of the Serre automorphism of α . In Sections 3.1.4 and 3.3.1–3.3.3, we recall the notions just mentioned, in particular how the Serre automorphism $S_{\alpha}: \alpha \rightarrow \alpha$, defined in (3.4), corresponds to one full rotation of frames.

r -spin cobordism hypothesis. In the present paper, we classify fully extended spin TQFTs valued in an arbitrary symmetric monoidal 2-category \mathcal{B} (Section 3), and we construct a number of examples (Section 4). More precisely, we consider r -spin TQFTs for any positive integer r . Recall that while for $n \geq 3$, the double cover $\text{Spin}_n \rightarrow \text{SO}_n$ is also the universal cover, this is not true for $n \leq 2$. Hence, there

¹In fact, [39, Theorem 5.2.1] provides a classification of open/closed r -spin TQFTs, of which the closed case for $r = 2$ discussed here is a special case.

is less reason to single out double covers of SO_2 and instead consider the r -fold cover $\mathrm{Spin}_2^r \rightarrow \mathrm{SO}_2$ for all $r \in \mathbb{Z}_{\geq 1}$ ². Note that necessarily

$$\mathrm{Spin}_2^r \cong \mathrm{SO}_2$$

as groups, and that by definition,

$$\mathrm{Spin}_2 = \mathrm{Spin}_2^2 \quad \text{and} \quad \mathrm{Spin}_2^1 = \mathrm{SO}_2.$$

Following [38], in Section 3.2, we describe a 2-category $\mathrm{Bord}_{2,1,0}^{r\text{-spin}}$ of bordisms with r -spin structure related to $\xi: \mathrm{Spin}_2^r \rightarrow \mathrm{SO}_2 \hookrightarrow \mathrm{GL}_2$, and in Section 3.3.3, we construct a 2-category $2D^r(\mathcal{B}^{\mathrm{fd}})$ whose objects are pairs (α, θ) , where $\alpha \in \mathcal{B}^{\mathrm{fd}}$ and $\theta: S_\alpha^r \cong 1_\alpha$. Then, we prove Lemma 3.18 and Theorem 3.19.

Theorem (r -spin cobordism hypothesis). *Let \mathcal{B} be a symmetric monoidal 2-category, and let $r \in \mathbb{Z}_{\geq 1}$. The 2-groupoid of fully extended r -spin TQFTs valued in \mathcal{B} is equivalent to the homotopy fixed points $[(\mathcal{B}^{\mathrm{fd}})^\times]^{\mathrm{Spin}_2^r}$. This in turn is equivalent to $2D^r((\mathcal{B}^{\mathrm{fd}})^\times)$, and under these equivalences, we have*

$$\begin{aligned} \mathrm{Fun}^{\mathrm{sm}}(\mathrm{Bord}_{2,1,0}^{r\text{-spin}}, \mathcal{B}) &\xrightarrow{\cong} [(\mathcal{B}^{\mathrm{fd}})^\times]^{\mathrm{Spin}_2^r} \xrightarrow{\cong} 2D^r((\mathcal{B}^{\mathrm{fd}})^\times) \\ \mathcal{Z} &\longmapsto (\mathcal{Z}(+), S_{\mathcal{Z}(+)}^r \cong 1_{\mathcal{Z}(+)}) \end{aligned}$$

Put differently, (fully) extended r -spin TQFTs are classified by what they assign to the positively framed point $+ \in \mathrm{Bord}_{2,1,0}^{r\text{-spin}}$ together with a trivialisation of the r th power of the associated Serre automorphism. The main ingredients of the proof are a generators-and-relations presentation of $\mathrm{Bord}_{2,1,0}^{r\text{-spin}}$, inspired by the work [27], and an explicit description of r -spin bordisms in terms of holonomies, following [36].

Examples. The choice of target 2-category \mathcal{B} is essential for extended TQFTs. To broaden the class of known r -spin TQFTs, in Section 4, we explicitly describe Spin_2^r -homotopy fixed points in the “equivariant completion” $\mathcal{B}_{\mathrm{eq}}$ of any given symmetric monoidal 2-category \mathcal{B} . As introduced in [16] and reviewed in Section 4.1, objects in $\mathcal{B}_{\mathrm{eq}}$ are pairs (α, A) , where $\alpha \in \mathcal{B}$ and $A \in \mathcal{B}(\alpha, \alpha)$ is endowed with the structure of a Δ -separable Frobenius algebra, while 1- and 2-morphisms are bimodules and bimodule maps. We show the following proposition (see Corollary 4.9, and (4.1) for the definition of the Nakayama automorphism $\gamma_A: A \rightarrow A$).

Proposition. *Let $(\alpha, A) \in \mathcal{B}_{\mathrm{eq}}$ be such that $\alpha \in \mathcal{B}^{\mathrm{fd}}$ as well as $S_\alpha^r \cong 1_\alpha$ and $\gamma_A^r = 1_A$ in \mathcal{B} . Then, there is an r -spin TQFT:*

$$\begin{aligned} \mathcal{Z}: \mathrm{Bord}_{2,1,0}^{r\text{-spin}} &\longrightarrow \mathcal{B}_{\mathrm{eq}} \\ + &\longmapsto (\alpha, A). \end{aligned}$$

²For $n = 1$, we have $\mathrm{SO}_1 = \{1\}$, and its r -fold cover is the unique map $\{1, \dots, r\} \rightarrow \{1\}$.

Moreover, in Section 4.1.5, we explain how to compute the invariants such that TQFTs associate to r -spin surfaces, by explicitly constructing the closed Λ_r -Frobenius algebras which classify the underlying non-extended TQFTs.

An advantage of considering \mathcal{B}_{eq} -valued (as opposed to \mathcal{B} -valued, for a given \mathcal{B}) TQFTs is as follows. As explained in Remark 3.28, r -spin TQFTs valued in a pivotal 2-category \mathcal{B} cannot detect all r -spin structures if $r \geq 3$. However, the equivariant completion \mathcal{B}_{eq} of a pivotal 2-category \mathcal{B} is itself not pivotal.

As a specific example of a target \mathcal{B} , in Section 4.2, we consider the symmetric monoidal 2-category \mathcal{LG} of Landau–Ginzburg models, constructed in [13, 15]. (Examples of extended 2-spin TQFTs were first considered in [22].) Objects of \mathcal{LG} are “potentials” $W \in \mathbb{k}[x_1, \dots, x_n]$ that describe isolated singularities, and Hom categories are homotopy categories of matrix factorisations. In [13], it was observed that every object in \mathcal{LG} is fully dualisable and that precisely those potentials $W(x_1, \dots, x_n)$ that depend on an even number of variables give rise to fully extended oriented TQFTs. Moreover, these oriented TQFTs indeed extend the closed TQFTs associated to the (generically non-semisimple) Jacobi algebras Jac_W to the point. In light of the r -spin cobordism hypothesis proved in Section 3, it is straightforward to extend these results as follows (Theorem 4.17)³.

Theorem. *Every object $W(x_1, \dots, x_n) \in \mathcal{LG}$ gives rise to an extended 2-spin TQFT valued in \mathcal{LG} . These TQFTs factor through the oriented bordism 2-category iff n is even.*

Explicitly, the 2-spin TQFT associated to an object $W \in \mathcal{LG}$ with an odd number of variables consists of the even Neveu–Schwarz sector $\text{Jac}_W \in \text{Vect}_{\mathbb{k}} \subset \text{Vect}_{\mathbb{k}}^{\mathbb{Z}_2}$ and the odd Ramond sector $\text{Jac}_W[1] \in \text{Vect}_{\mathbb{k}}^{\mathbb{Z}_2}$, together with the structure maps described in general in Section 3.1.5. Moreover, in Example 4.18, we illustrate how to apply our results on equivariant completion (Section 4.1) to a variant of

$$\mathcal{B} = \mathcal{LG}$$

and explicitly compute the invariants of r -spin tori in the simplest non-trivial (and novel) example.

Examples not treated in this paper. We close this introductory section with a few comments on potential further applications of the r -spin cobordism hypothesis. Besides the 2-categories $\text{Alg}_{\mathbb{k}}$ and \mathcal{LG} (as well as their variants with additional \mathbb{Z}_2 -, \mathbb{Z} - or \mathbb{Q} -gradings), it is natural to consider the 2-category $\mathcal{V}\text{ar}$ of [11] of smooth

³In the case of Landau–Ginzburg models, the vector space of automorphisms of the identity 1-morphism on any object is 1-dimensional. Hence, the choice of trivialisation of the square of the Serre automorphism is unique up to a non-zero scalar.

projective varieties and derived categories (see also Example 3.3), which appears in the study of B -twisted sigma models. The 2-category $\mathcal{V}\text{ar}$ has a natural symmetric monoidal structure [5]. As explained in [11, 31], the Serre automorphism S_X of $X \in \mathcal{V}\text{ar}$ can be identified with the Serre functor of the derived category associated to X .

In [29], Kuznetsov constructs “fractional Calabi–Yau categories” \mathcal{A}_X as the admissible subcategories of semiorthogonal decompositions of derived categories of certain varieties $X \in \mathcal{V}\text{ar}$. This means in particular that \mathcal{A}_X is a triangulated category with suspension functor Σ such that \mathcal{A}_X has a Serre functor S which satisfies $S^q \cong \Sigma^p$ for some $p, q \in \mathbb{Z}$ with $q \neq 0$. It follows that the orbit category \mathcal{A}_X/\mathbb{Z} has a Serre functor whose $(p - q)$ -th power is trivialisable; see, e.g., [21, Theorem 5.14].

It is tempting to expect that some of the fractional Calabi–Yau categories constructed in [29, Section 4] classify $(p - q)$ -spin TQFTs whose target is $\mathcal{V}\text{ar}$ up to the \mathbb{Z} -action quotiented out in orbit categories. This is possible only if one can identify the Serre functor of \mathcal{A}_X/\mathbb{Z} with the Serre automorphism of some other object in the target 2-category. More generally, we could work in the 2-category of smooth and proper triangulated differential graded categories described in [4, Appendix A]. In this setting, both the geometric constructions of [29] and the representation theoretic examples of fractional Calabi–Yau categories in [21, Section 6] may lead to interesting r -spin TQFTs.

2. Non-extended r -spin TQFTs

In this section, we review the classification of non-extended closed r -spin and framed TQFTs following [39], to which we refer for details. We recall the relevant categories of 2-dimensional bordisms as well as the notion of a closed Λ_r -Frobenius algebra, and we state the main classification result: 2-dimensional r -spin and framed ($r = 0$) TQFTs are equivalent to closed Λ_r -Frobenius algebras in the target category.

2.1. Framed and r -spin TQFTs

By a *surface*, we mean a 2-dimensional compact smooth manifold. Let G be a topological group, and let

$$\xi: G \longrightarrow \text{GL}_2$$

be a continuous group homomorphism, and recall that the frame bundle $F\Sigma \rightarrow \Sigma$ of a surface Σ is a principal GL_2 -bundle. A G -structure (more precisely: a *tangential structure for $\xi: G \rightarrow \text{GL}_2$*) on Σ is a principal G -bundle $\pi: P \rightarrow \Sigma$ together with a

bundle map q intertwining the group actions via ξ :

$$\begin{array}{ccc} P & \xrightarrow{q} & F\Sigma \\ & \searrow \pi & \swarrow \\ & \Sigma & \end{array}$$

A *map of surfaces with G -structure* is a bundle map which is a local diffeomorphism of the underlying surfaces. Such a map is called a *diffeomorphism* if its underlying map of surfaces is a diffeomorphism, and an *isomorphism of G -structures* if the underlying map of surfaces is the identity.

We will consider the following tangential structures:

- A *framing* is a tangential structure for the trivial group:

$$\star \longrightarrow \mathrm{GL}_2. \quad (2.1)$$

- An *orientation* is a tangential structure for the inclusion:

$$\mathrm{SO}_2 \simeq \mathrm{GL}_2^+ \hookrightarrow \mathrm{GL}_2, \quad (2.2)$$

where GL_2^+ is the subgroup of elements in GL_2 with positive determinant.

- For $r \in \mathbb{Z}_{\geq 0}$, an *r -spin structure* is a tangential structure:

$$\widetilde{\mathrm{GL}}_2^{+r} \xrightarrow{p^r} \mathrm{GL}_2^+ \hookrightarrow \mathrm{GL}_2, \quad (2.3)$$

where $p^r: \widetilde{\mathrm{GL}}_2^{+r} \rightarrow \mathrm{GL}_2^+$ is the r -fold covering for $r > 0$, while for $r = 0$, it is the universal cover.

By a trivial r -spin structure on a surface Σ , we mean an r -spin structure isomorphic to the r -spin structure with trivial bundles

$$P = \Sigma \times \widetilde{\mathrm{GL}}_2^{+r}, \quad F\Sigma = \Sigma \times \mathrm{GL}_2$$

and trivial bundle map $q^{(+)} = \mathrm{id}_\Sigma \times p^r$ (positive orientation) or $q^{(-)} = \mathrm{id}_\Sigma \times (T \circ p^r)$ (negative orientation), where T is composition with the matrix $\begin{pmatrix} +1 & 0 \\ 0 & -1 \end{pmatrix} \in \mathrm{GL}_2$.

Remark 2.1. A 1-spin structure is the same as an orientation, and a 2-spin structure is usually called a *spin structure*. Moreover, we can identify framings with 0-spin structures by noting that the fibres of a 0-spin bundle are contractible; see [37, Proposition 2.2]. This is consistent with the fact that, for any $r \in \mathbb{Z}_{\geq 0}$, an r -spin structure is a \mathbb{Z}_r -bundle over the oriented frame bundle.

Let $r \in \mathbb{Z}_{\geq 0}$. There is a symmetric monoidal category of 2-dimensional r -spin bordisms $\text{Bord}_{2,1}^{r\text{-spin}}$ as follows. An object S is a 1-dimensional closed manifold s embedded in a cylinder $s \times (-1, 1)$, together with an r -spin structure on the cylinder. For an object S , we write

$$S^{(+)} := s \times [0, 1) \quad \text{and} \quad S^{(-)} := s \times (-1, 0]$$

with the restricted r -spin structures. The morphisms of $\text{Bord}_{2,1}^{r\text{-spin}}$ are diffeomorphism classes of r -spin bordisms: for $S, S' \in \text{Bord}_{2,1}^{r\text{-spin}}$, an r -spin bordism $S \rightarrow S'$ is a compact surface Σ with r -spin structure, together with a boundary parametrisation map $S^{(+)} \sqcup S'^{(-)} \hookrightarrow \Sigma$, i.e., a map of r -spin surfaces that identifies the boundary of Σ with the 1-dimensional embedded manifolds

$$s \times \{0\} \subset S \quad \text{and} \quad s' \times \{0\} \subset S'.$$

Finally, a diffeomorphism of r -spin bordisms is a diffeomorphism of r -spin surfaces compatible with the boundary parametrisations. We usually refer to a morphism in $\text{Bord}_{2,1}^{r\text{-spin}}$ by a bordism that represents it.

A particular class of r -spin bordisms are *deck transformation bordisms*. These are cylinders whose boundary parametrisations are given by deck transformations of the r -spin bundle on the source or target object.

The composition of morphisms in $\text{Bord}_{2,1}^{r\text{-spin}}$ is given by gluing along boundary parametrisations; hence, the unit morphisms are given by cylinders with trivial boundary parametrisations. Taking disjoint unions endows $\text{Bord}_{2,1}^{r\text{-spin}}$ with its standard symmetric monoidal structure. In light of Remark 2.1, we write

$$\text{Bord}_{2,1}^{\text{fr}} = \text{Bord}_{2,1}^{0\text{-spin}}, \quad \text{Bord}_{2,1}^{\text{or}} = \text{Bord}_{2,1}^{1\text{-spin}}.$$

Definition 2.2. Let $\mathbb{C}c$ be a symmetric monoidal category. A (closed) r -spin TQFT valued in $\mathbb{C}c$ is a symmetric monoidal functor

$$\mathfrak{Z}: \text{Bord}_{2,1}^{r\text{-spin}} \longrightarrow \mathbb{C}c.$$

The case of 2-dimensional closed spin TQFTs ($r = 2$) was first described and classified in [1], including concrete examples in terms of Clifford algebras viewed as objects in the category of super vector spaces

$$\mathbb{C}c = \text{Vect}_{\mathbb{C}}^{\mathbb{Z}_2}.$$

Spin TQFTs were further discussed from the perspective of extended TQFTs in [22], and spin state sum constructions were given in [7, 33]. TQFTs with r -spin structure for arbitrary r were introduced in [32] and further studied in [37]. The classification of general r -spin TQFTs appears in [39], in terms of the algebraic structures we review next.

2.2. Classification in terms of closed Λ_r -Frobenius algebras

A closed Λ_r -Frobenius algebra C in a symmetric monoidal category \mathcal{C} consists of a collection of objects $C_a \in \mathcal{C}$ for all $a \in \mathbb{Z}_r$ as well as morphisms

$$\begin{aligned} \mu_{a,b} &= \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} : C_a \otimes C_b \longrightarrow C_{a+b-1}, & \eta_1 &= \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} : 1 \longrightarrow C_1, \\ \Delta_{a,b} &= \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} : C_{a+b+1} \longrightarrow C_a \otimes C_b, & \varepsilon_{-1} &= \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} : C_{-1} \longrightarrow 1 \end{aligned}$$

for all $a, b \in \mathbb{Z}_r$. The Nakayama automorphisms of C are

$$N_a := \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} : C_a \longrightarrow C_a \quad (2.4)$$

for all $a \in \mathbb{Z}_r$. These data by definition satisfy the following conditions:

(co)associativity:

$$\begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} = \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array}, \quad \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} = \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array}, \quad (2.5)$$

(co)unitality :

$$\begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} = \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array}, \quad \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} = \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array}, \quad (2.6)$$

Frobenius relation:

$$\begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} = \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array}, \quad (2.7)$$

commutativity:

$$N_b^{1-a} \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} = \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} = \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array}, \quad (2.8)$$

twist relations:

$$N_a^a = 1_{C_a}, \quad \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array} = \begin{array}{c} \bullet \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \bullet \end{array}, \quad (2.9)$$

$$\text{deck transformation relations: } N_a^r = 1_{C_a}. \quad (2.10)$$

A map of closed Λ_r -Frobenius algebras $\varphi: C \rightarrow D$ is a collection of morphisms $\varphi_a: C_a \rightarrow D_a$ preserving the structure morphisms. Analogously to the case of ordinary Frobenius algebras, maps of closed Λ_r -Frobenius algebras are always isomorphisms.

Example 2.3. (i) One class of closed Λ_r -Frobenius algebras in a given symmetric monoidal category \mathcal{C} can be constructed from ordinary Frobenius algebras A in \mathcal{C} whose ordinary Nakayama automorphism γ_A satisfies $\gamma_A^r = 1_A$ (see Section 4.1.1 and (4.1) below for details). Indeed, as explained in [37] and [39, Section 4.2], the construction of commutative Frobenius algebras as the centres of certain types of non-commutative Frobenius algebras in [30, Section 2.7] is naturally the special case of $r = 1$ of a construction of “ \mathbb{Z}_r -graded centre” for any $r \in \mathbb{Z}_{\geq 0}$.

(ii) In the category $\text{Bord}_{2,1}^{r\text{-spin}}$, r -spin circles, pair of pants, cups, and caps naturally assemble into a closed Λ_r -Frobenius algebra C . The precise presentation is given in [39, Section 5.1] in terms of a combinatorial description of r -spin structures. In particular, it follows from [39, equation (5.2)] that the Nakayama automorphisms of C are deck transformation bordisms.

The closed Λ_r -Frobenius algebra C of Example 2.3 (ii) is not just any example. As proven in [39, Theorem 5.2.1], $\text{Bord}_{2,1}^{r\text{-spin}}$ is generated as a symmetric monoidal category by the data of C , subject to relations given by the defining properties (2.5)–(2.10). This implies the following theorem.

Theorem 2.4 ([39, Corollary 5.2.2]). *There is an equivalence of symmetric monoidal groupoids between the groupoid of r -spin TQFTs with target \mathcal{C} and the groupoid of closed Λ_r -Frobenius algebras in \mathcal{C} .*

For this reason, we will refer to the objects C_a of a closed Λ_r -Frobenius algebra in any given symmetric monoidal category \mathcal{C} (not necessarily equivalent to $\text{Bord}_{2,1}^{r\text{-spin}}$) as the a -th circle spaces. The a -th circle space in $\text{Bord}_{2,1}^{r\text{-spin}}$ is simply the circle with “framing number” a , and we denote it by S_a^1 . Below in Sections 3.1.5, 4.1.5 and 4.2, we will use Theorem 2.4 to construct examples of closed r -spin TQFTs beyond those mentioned in Section 2.1.

2.3. Computing invariants

The above classification theorem provides a way to compute invariants of r -spin surfaces from r -spin TQFTs in terms of the algebraic data of a closed Λ_r -Frobenius algebra. As the number of diffeomorphism classes of r -spin structures on a connected oriented surface of genus $g \geq 2$ is, if non-zero, either one (r odd) or two (r even), we

are mainly interested in surfaces of genera 0 and 1; see, e.g., [40, Section 3] and the references therein for a detailed account.

The sphere S^2 admits an r -spin structure only if $r \in \{1, 2\}$, in which case it is unique up to isomorphism; hence, the torus T^2 is of most interest. Any torus with r -spin structure can be presented in terms of the closed Λ_r -Frobenius algebra in $\text{Bord}_{2,1}^{r\text{-spin}}$ as

$$T(a, b) := \begin{array}{c} \text{---} a \cdot a \\ \circ \\ \text{---} N_{-a}^{1-b} \\ \bullet \\ \text{---} -a \cdot a \\ \circ \end{array} \in \text{End}_{\text{Bord}_{2,1}^{r\text{-spin}}}(\emptyset)$$

for some $a, b \in \mathbb{Z}_r$. Moreover, as shown in [39, Proposition 4.1.4], the r -spin torus $T(a, b)$ is diffeomorphic to the r -spin torus $T(\gcd(a, b, r), 0)$, and in fact, diffeomorphism classes of r -spin tori are in bijection with divisors of r . Hence, we write

$$T(d) = T(d, 0)$$

for the class of r -spin tori corresponding to the divisor d .

Proposition 2.5 ([39, Proposition 4.1.4]). *The invariant of the r -spin torus $T(d)$ computed by a \mathcal{C} -valued closed r -spin TQFT \mathcal{Z} classified by a closed Λ_r -Frobenius algebra C is the quantum dimension of C_d :*

$$\mathcal{Z}(T(d)) = \dim(C_d) = \text{ev}_{C_d} \circ b_{C_{-d}, C_d} \circ \text{coev}_{C_d},$$

where b is the braiding of \mathcal{C} .

3. Fully extended r -spin TQFTs

In this section, we describe fully extended r -spin TQFTs and prove the corresponding cobordism hypothesis in the 2-categorical setting. In Section 3.1, we recall some aspects of symmetric monoidal 2-categories \mathcal{B} , their Serre automorphisms, and we construct canonical closed Λ_0 -Frobenius algebras. Section 3.2 describes the 2-category $\text{Bord}_{2,1,0}^{r\text{-spin}}$ of r -spin bordisms. Then, in Section 3.3, we define the 2-groupoid of fully extended r -spin TQFTs $\text{Bord}_{2,1,0}^{r\text{-spin}} \rightarrow \mathcal{B}$ and explain that it is equivalent to the 2-groupoid of Spin_2^r -homotopy fixed points.

3.1. Dualisability in symmetric monoidal 2-categories


In this section, we present our notational conventions for dualisability in symmetric monoidal 2-categories. Moreover, we construct a closed Λ_0 -Frobenius algebra (in the sense of Section 2.2) for every fully dualisable object.

For complete definitions, we refer to [8, 34, 38] and references therein; with an eye towards examples in Section 4, below we mostly use the same conventions as in [13, Section 2].

3.1.1. Conventions for 2-categories. By a 2-category we mean a (possibly non-strict) bicategory \mathcal{B} in the sense of [38, Appendix A.1]. For objects $\alpha, \beta \in \mathcal{B}$, we denote the category of 1-morphisms $\alpha \rightarrow \beta$ by $\mathcal{B}(\alpha, \beta)$; for 1-morphisms $X, Y \in \mathcal{B}(\alpha, \beta)$, we write $\text{Hom}_{\mathcal{B}}(X, Y)$, or simply $\text{Hom}(X, Y)$, for the set of 2-morphisms $X \rightarrow Y$. Horizontal and vertical compositions are denoted by \otimes and \circ , respectively:

$$\begin{aligned} \otimes: \mathcal{B}(\beta, \gamma) \times \mathcal{B}(\alpha, \beta) &\longrightarrow \mathcal{B}(\alpha, \gamma) \\ (X', X) &\longmapsto X' \otimes X, \\ \circ: \text{Hom}(Y, Z) \times \text{Hom}(X, Y) &\longrightarrow \text{Hom}(X, Z) \\ (\psi, \varphi) &\longmapsto \psi \circ \varphi. \end{aligned}$$

We read string diagrams from right to left and from bottom to top. For instance, for 1-morphisms $X \in \mathcal{B}(\alpha, \beta)$, $X' \in \mathcal{B}(\beta, \gamma)$, and $V \in \mathcal{B}(\alpha, \gamma)$, a 2-morphism $\varphi \in \text{Hom}(X' \otimes X, V)$ is represented by



However, sometimes, we will suppress object labels in string diagrams, as, e.g., in (3.1) below.

3.1.2. Adjoints. Let \mathcal{B} be a 2-category. A 1-morphism $X \in \mathcal{B}(\alpha, \beta)$ has a *left adjoint* if there exists a 1-morphism ${}^\dagger X \in \mathcal{B}(\beta, \alpha)$ together with *adjunction 2-morphisms*

$$\begin{aligned} \begin{array}{c} \text{---} \curvearrowright \text{---} \\ \text{ }^\dagger X \quad X \\ \text{---} \curvearrowleft \text{---} \\ X \quad \text{ }^\dagger X \end{array} &= \text{ev}_X: \text{ }^\dagger X \otimes X \longrightarrow 1_\alpha, \\ \begin{array}{c} \text{---} \curvearrowleft \text{---} \\ X \quad \text{ }^\dagger X \\ \text{---} \curvearrowright \text{---} \\ \text{ }^\dagger X \quad X \end{array} &= \text{coev}_X: 1_\beta \longrightarrow X \otimes \text{ }^\dagger X \end{aligned} \quad (3.1)$$

such that the Zorro moves

$$\begin{array}{c} \text{U-shaped curve with arrows} \end{array} = \begin{array}{c} | \end{array} = 1_X, \quad \begin{array}{c} \text{Inverted U-shaped curve with arrows} \end{array} = \begin{array}{c} | \end{array} = 1_{\dagger X}$$

are satisfied. Similarly, a *right adjoint* for X consists of $X^\dagger \in \mathcal{B}(\beta, \alpha)$ with

$$\begin{array}{c} \text{Loop with } X \text{ and } X^\dagger \end{array} = \tilde{\text{ev}}_X: X \otimes X^\dagger \longrightarrow 1_\beta, \\
 \begin{array}{c} \text{Loop with } X^\dagger \text{ and } X \end{array} = \widetilde{\text{coev}}_X: 1_\alpha \longrightarrow X^\dagger \otimes X$$

that satisfy analogous Zorro moves.

If $X, Y \in \mathcal{B}(\alpha, \beta)$ have left and right adjoints (with chosen adjunction maps), then we write

$$\dagger\varphi := \begin{array}{c} \text{Diagram with } \dagger X \text{ and } \dagger Y \end{array}, \quad \varphi^\dagger := \begin{array}{c} \text{Diagram with } X^\dagger \text{ and } Y^\dagger \end{array}$$

for the left and right adjoints of $\varphi \in \text{Hom}(X, Y)$, respectively. We call \mathcal{B} *pivotal* if every 1-morphism X comes with chosen left and right adjunction data such that

$$\dagger X = X^\dagger, \quad \dagger\varphi = \varphi^\dagger$$

for all 2-morphisms φ , and

$$\begin{array}{c} \text{Complex diagram with } X^\dagger, Y^\dagger, \text{ and } (Y \otimes X)^\dagger \end{array} = \begin{array}{c} \text{Complex diagram with } X^\dagger, Y^\dagger, \text{ and } (Y \otimes X)^\dagger \end{array}$$

for all composable 1-morphisms X, Y .

3.1.3. Symmetric monoidal structure. Let \mathcal{B} be a 2-category. A monoidal structure on \mathcal{B} consists of a 2-functor


$$\square: \mathcal{B} \times \mathcal{B} \longrightarrow \mathcal{B}$$

called monoidal product, a unit object

$$1 \in \mathcal{B},$$

a pseudonatural transformation $a: \square \circ (\square \times \text{Id}_{\mathcal{B}}) \rightarrow \square \circ (\text{Id}_{\mathcal{B}} \times \square)$ called associator, a weak inverse a^- for a , as well as unitors, 2-unitors, and a pentagonator (which we will usually suppress), subject to the coherence axioms in [38, Section 2.3].

Viewing a monoidal 2-category \mathcal{B} as a 3-category with a single object and using the strictification results of [23, 24], we can use the 3-dimensional graphical calculus of [6, 41]. For this, we extend our diagrammatic conventions by reading 3-dimensional diagrams from front to back. For instance, for 1-morphisms $X \in \mathcal{B}(\varepsilon \square \delta, \alpha)$, $Y \in \mathcal{B}(\gamma \square \zeta, \varepsilon)$, $Z \in \mathcal{B}(1, \zeta \square \delta)$, $X' \in \mathcal{B}(\beta, \alpha)$, and $Y' \in \mathcal{B}(\gamma, \beta)$, the diagram




represents a 2-morphism $\varphi \in \text{Hom}(X' \otimes Y', X \otimes (Y \square 1_\delta) \otimes (1_\gamma \square Z))$; compare [12, Section 3.1.2].

Let \mathcal{B} be a monoidal 2-category. Writing $\tau: \mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B} \times \mathcal{B}$ for the strict 2-functor that acts as $(\zeta, \xi) \mapsto (\xi, \zeta)$ on objects, 1- and 2-morphisms, a symmetric braided structure on \mathcal{B} consists of a pseudonatural transformation


$$b: \square \longrightarrow \square \circ \tau$$

called braiding, a weak inverse b^- for b , and an invertible modification $\sigma: 1_\square \rightarrow b^- \circ b$, as well as two further invertible modifications between compositions of a , a^- , b , b^- , subject to the coherence axioms of [38, Section 2.3].



The braiding b consists of 1-morphism components




for all $\alpha, \alpha' \in \mathcal{B}$ and of 2-morphism components $b_{X,Y}: (Y \square X) \otimes b_{\alpha,\beta} \rightarrow b_{\alpha',\beta'} \otimes (X \square Y)$ for all $X \in \mathcal{B}(\alpha, \alpha')$ and $Y \in \mathcal{B}(\beta, \beta')$. Graphically, the 2-morphism components are depicted as




3.1.4. Duality and Serre automorphism. Let \mathcal{B} be a symmetric monoidal 2-category. An object $\alpha \in \mathcal{B}$ has a *dual* if there exists an object $\alpha^\#$ together with *adjunction 1-morphisms*



and *cuspidal 2-isomorphisms*



$$= c_l: (\tilde{e}v_\alpha \square 1_\alpha) \otimes (1_\alpha \square \widetilde{\text{coev}}_\alpha) \longrightarrow 1_\alpha, \quad (3.2)$$



$$= c_r: (1_{\alpha^\#} \square \tilde{e}v_\alpha) \otimes (\widetilde{\text{coev}}_\alpha \square 1_{\alpha^\#}) \longrightarrow 1_{\alpha^\#}. \quad (3.3)$$


More precisely, these data witness $\alpha^\#$ as the *right dual* of α . Using the symmetric braiding of \mathcal{B} , the object $\alpha^\#$ is also the left dual of α , with adjunction maps

$$\text{ev}_\alpha = \tilde{e}v_\alpha \otimes b_{\alpha^\#, \alpha}, \quad \text{coev}_\alpha = b_{\alpha^\#, \alpha} \otimes \widetilde{\text{coev}}_\alpha.$$

If $\alpha, \beta \in \mathcal{B}$ have duals $\alpha^\#, \beta^\#$ with chosen adjunction 1-morphisms


$$\tilde{e}v_\alpha, \quad \widetilde{\text{coev}}_\alpha, \quad \tilde{e}v_\beta, \quad \widetilde{\text{coev}}_\beta,$$

the associated dual $X^\#$ of a 1-morphism $X \in \mathcal{B}(\alpha, \beta)$ is



$$\hat{=} X^\# \in \mathcal{B}(\beta^\#, \alpha^\#).$$

For another 1-morphism $Y \in \mathcal{B}(\alpha, \beta)$ and a 2-morphism $\varphi \in \text{Hom}(X, Y)$, its dual $\varphi^\#$ is




An object α in a symmetric monoidal 2-category \mathcal{B} is called *fully dualisable* if it has a dual $\alpha^\#$ such that the 1-morphisms $\tilde{e}v_\alpha, \widetilde{\text{co}e}v_\alpha$ have both left and right adjoints (as in Section 3.1.2). The full sub-2-category of fully dualisable objects is denoted by \mathcal{B}^{fd} , and we call \mathcal{B} fully dualisable if $\mathcal{B} \cong \mathcal{B}^{\text{fd}}$.


Convention 3.1. Whether or not a symmetric monoidal 2-category \mathcal{B} is fully dualisable is a property of \mathcal{B} . If it is fully dualisable, we will assume that we have chosen explicit duality data $(\alpha^\#, \tilde{e}v_\alpha, \widetilde{\text{co}e}v_\alpha)$ and adjunction data $(\dagger \tilde{e}v_\alpha, \text{ev}_{\tilde{e}v_\alpha}, \text{coev}_{\tilde{e}v_\alpha}), (\tilde{e}v_\alpha^\dagger, \tilde{e}v_{\tilde{e}v_\alpha}, \widetilde{\text{co}e}v_{\tilde{e}v_\alpha}), (\dagger \widetilde{\text{co}e}v_\alpha, \text{ev}_{\widetilde{\text{co}e}v_\alpha}, \text{coev}_{\widetilde{\text{co}e}v_\alpha}),$ and $(\widetilde{\text{co}e}v_\alpha^\dagger, \tilde{e}v_{\widetilde{\text{co}e}v_\alpha}, \widetilde{\text{co}e}v_{\widetilde{\text{co}e}v_\alpha})$ for all $\alpha \in \mathcal{B}$. Put differently, we then view \mathcal{B} as “fully dualised”.

As shown in [34], the adjunction 1-morphisms $\tilde{e}v_\alpha, \widetilde{\text{co}e}v_\alpha$ of a fully dualisable object α do not only have left and right adjoints, but these again have left and right adjoints, and so on infinitely. The relations between multiple adjoints are negotiated by the *Serre automorphism*:

$$S_\alpha = (1_\alpha \square \tilde{e}v_\alpha) \otimes (b_{\alpha,\alpha} \square 1_{\alpha^\#}) \otimes (1_\alpha \square \tilde{e}v_\alpha^\dagger)$$



with inverse

$$S_\alpha^{-1} = (1_\alpha \square \tilde{e}v_\alpha) \otimes (b_{\alpha,\alpha} \square 1_{\alpha^\#}) \otimes (1_\alpha \square \dagger \tilde{e}v_\alpha)$$


The general result [34, Theorem 3.9] on multiple adjoints in \mathcal{B}^{fd} implies, in particular,


$$\begin{aligned} \tilde{e}v_\alpha^\dagger &\cong (S_\alpha \square 1_{\alpha^\#}) \otimes b_{\alpha^\#, \alpha} \otimes \widetilde{\text{coev}}_\alpha, \\ \dagger \tilde{e}v_\alpha &\cong (S_\alpha^{-1} \square 1_{\alpha^\#}) \otimes b_{\alpha^\#, \alpha} \otimes \widetilde{\text{coev}}_\alpha. \end{aligned} \quad (3.6)$$

Let $(\mathcal{B}^{\text{fd}})^\times$ be the maximal sub-2-groupoid of \mathcal{B}^{fd} . Then, as shown in [27, Proposition 2.8], for all $X \in (\mathcal{B}^{\text{fd}})^\times(\alpha, \beta)$, there are 2-morphisms

$$S_X: X \otimes S_\alpha \longrightarrow S_\beta \otimes X,$$

which together with the components S_α assemble into a pseudonatural transformation $\text{Id}_{(\mathcal{B}^{\text{fd}})^\times} \rightarrow \text{Id}_{(\mathcal{B}^{\text{fd}})^\times}$. This can be slightly generalised as follows.

Proposition 3.2. *Let \mathcal{B} be a symmetric monoidal pivotal 2-category such that \mathcal{B}^{fd} has adjoints for all 1-morphisms. Then, the Serre automorphisms S_α together with the 2-morphisms (expressed in terms of the graphical calculus of [6] for symmetric monoidal pivotal 2-categories)*

$$S_X =$$


for all $X \in \mathcal{B}^{\text{fd}}(\alpha, \beta)$

form a pseudonatural transformation $S: \text{Id}_{\mathcal{B}^{\text{fd}}} \rightarrow \text{Id}_{\mathcal{B}^{\text{fd}}}$.

Proof. If $X \in \mathcal{B}^{\text{fd}}(\alpha, \beta)$ has a quasi-inverse X^{-1} , then X^{-1} is isomorphic to the (chosen) adjoint X^\dagger , and we have $(X^\#)^{-1} \cong (X^{-1})^\# \cong (X^\dagger)^\#$. Substituting this into the proof of [27, Proposition 2.8], we find that specifying S_X amounts to filling the diagram

$$\begin{array}{ccccccc}
 \alpha \cong \alpha \square 1 & \xrightarrow{1_\alpha \square \tilde{e}v_\alpha^\dagger} & \alpha \square \alpha \square \alpha^\# & \xrightarrow{b(\alpha, \alpha) \square 1_{\alpha^\#}} & \alpha \square \alpha \square \alpha^\# & \xrightarrow{1_\alpha \square \tilde{e}v_\alpha} & \alpha \square 1 \cong \alpha \\
 \downarrow X & & \downarrow X \square X \square (X^\dagger)^\# & & \downarrow X \square X \square (X^\dagger)^\# & & \downarrow X \\
 \beta \cong \beta \square 1 & \xrightarrow{1_\beta \square \tilde{e}v_\beta^\dagger} & \beta \square \beta \square \beta^\# & \xrightarrow{b(\beta, \beta) \square 1_{\beta^\#}} & \beta \square \beta \square \beta^\# & \xrightarrow{1_\beta \square \tilde{e}v_\beta} & \beta \square 1 \cong \beta
 \end{array}$$

This is precisely what the expression of S_X in (3.7) does. \blacksquare


Example 3.3. We sketch a few fully dualisable symmetric monoidal 2-categories that appear in connection with 2-dimensional TQFT:

- (i) There is a 2-category $\text{Bord}_{2,1,0}^{\text{fr}}$ of 2-framed points, 1-dimensional bordisms and 2-dimensional bordism classes, that we review below in Section 3.2. The Serre automorphism S_+ of the positively framed point $+$ $\in \text{Bord}_{2,1,0}^{\text{fr}}$ generates an action of $\pi_1(\text{SO}_2) \cong \mathbb{Z}$ and corresponds to a twist of the interval over the point $+$.
- (ii) State sum models: for \mathbb{k} field, there is a 2-category $\text{Alg}_{\mathbb{k}}^{\text{fd}}$ of separable \mathbb{k} -algebras, bimodules, and bimodule maps [31, 38]. The Serre automorphism of $A \in \text{Alg}_{\mathbb{k}}^{\text{fd}}$ is the A - A -bimodule $\text{Hom}_{\mathbb{k}}(A, \mathbb{k})$.
- (iii) Landau–Ginzburg models: there is a 2-category \mathcal{LG} of isolated singularities, matrix factorisations, and their maps up to homotopy [13, 15], which we briefly review in Section 4.2 below. The Serre automorphism of $W \in \mathcal{LG}$ is isomorphic to 1_W up to a shift.
- (iv) B-twisted sigma models: there is a 2-category \mathcal{Var} of smooth projective varieties, Fourier–Mukai kernels and Ext groups [5, 11]. The Serre automorphism of $X \in \mathcal{Var}$ is given by tensoring with the canonical line bundle of X shifted by $-\dim(X)$.
- (v) Topologically twisted models: there is a 2-category $\text{DGSat}_{\mathbb{k}}$ of essentially small, smooth, proper, and triangulated differential graded \mathbb{k} -categories and their derived categories of bimodules [4, Appendix A]. The Serre automorphism of $\mathcal{C} \in \text{DGSat}_{\mathbb{k}}$ is given in terms of the \mathbb{k} -linear dual composed with the canonical trace functor associated to \mathcal{C} . The 2-categories of parts (iii) and (iv) are equivalent to sub-2-categories of $\text{DGSat}_{\mathbb{k}}$.

3.1.5. A Frobenius algebra. Let \mathcal{B} be a symmetric monoidal 2-category. For a fixed fully dualisable object $\alpha \in \mathcal{B}^{\text{fd}}$ with $\alpha^{\#\#} = \alpha$, we now consider the a -th circle spaces

$$\begin{aligned} C_a^\alpha &:= \tilde{e}\tilde{v}_\alpha \otimes (1_\alpha \square S_{\alpha^\#}^{1-a}) \otimes \dagger \tilde{e}\tilde{v}_\alpha \\ &\cong \tilde{e}\tilde{v}_\alpha \otimes (S_{\alpha^\#}^{1-a} \square 1_{\alpha^\#}) \otimes \dagger \tilde{e}\tilde{v}_\alpha \in \mathcal{B}(1, 1) \quad \text{for all } a \in \mathbb{Z}, \end{aligned} \quad (3.8)$$

where the isomorphism in (3.8) is induced by



which in turn is the cusp isomorphism (3.2) combined with $S_\alpha^\# \cong S_{\alpha^\#}$.

If \mathcal{B} is the 2-category $\text{Alg}_{\mathbb{k}}^{\text{fd}}$ of Example 3.3 (ii), then for an algebra $A \in \text{Alg}_{\mathbb{k}}^{\text{fd}}$ the zeroth and first circle spaces are the zeroth Hochschild homology and cohomology of A , respectively: $C_0^A \cong \text{HH}_0(A)$ and $C_1^A \cong \text{HH}^0(A)$. If \mathcal{B} is the 2-category $\mathcal{L}\mathcal{G}$ of Example 3.3 (iii), then the circle spaces of a given object are (shifts of) the associated Jacobi algebra, as we will explain in Section 4.2 below. In the following, we will sometimes treat the isomorphism in (3.8) as an identity, and we usually drop the index “ α ” in C_a^α .

Next, we set

$$\eta_1 := \begin{array}{c} \tilde{e}\tilde{v}_\alpha \\ \text{cup} \\ \dagger \tilde{e}\tilde{v}_\alpha \\ \text{COEV}_{\tilde{e}\tilde{v}_\alpha} \end{array} : 1_1 \longrightarrow C_1, \quad (3.9)$$

$$\varepsilon_{-1} := \begin{array}{c} \tilde{e}\tilde{v}_{\tilde{e}\tilde{v}_\alpha} \\ \text{cap} \\ \tilde{e}\tilde{v}_\alpha \\ \text{S}_{\alpha^\#}^2 \end{array} : C_{-1} \longrightarrow 1_1, \quad (3.10)$$

$$\mu_{a,b} := \begin{array}{c} \begin{array}{c} \tilde{e}v_\alpha \quad S_{\alpha^\#}^{2-a-b} \quad \dagger \tilde{e}v_\alpha \\ \downarrow \\ \tilde{e}v_\alpha \quad S_{\alpha^\#}^{1-a} \quad \dagger \tilde{e}v_\alpha \quad \tilde{e}v_\alpha \quad S_{\alpha^\#}^{1-b} \quad \dagger \tilde{e}v_\alpha \\ \downarrow \\ \text{ev}_{\tilde{e}v_\alpha} \end{array} \\ : C_a \otimes C_b \longrightarrow C_{a+b-1}, \end{array} \quad (3.11)$$

$$\Delta_{a,b} := \begin{array}{c} \begin{array}{c} \tilde{e}v_\alpha \quad S_{\alpha^\#}^{1-a} \quad \dagger \tilde{e}v_\alpha \quad \tilde{e}v_\alpha \quad S_{\alpha^\#}^{1-b} \quad \dagger \tilde{e}v_\alpha \\ \downarrow \\ \tilde{e}v_\alpha \quad S_{\alpha^\#}^{-2} \quad \dagger \tilde{e}v_\alpha \\ \downarrow \\ \text{coev}_{\tilde{e}v_\alpha} \\ \downarrow \\ \tilde{e}v_\alpha \quad S_{\alpha^\#}^{-1-a} \\ \downarrow \\ \tilde{e}v_\alpha \quad S_{\alpha^\#}^{-a-b} \quad \dagger \tilde{e}v_\alpha \end{array} \\ : C_{a+b+1} \longrightarrow C_a \otimes C_b \end{array} \quad (3.12)$$

for all $a, b \in \mathbb{Z}$, where in the expressions for ε_{-1} and $\Delta_{a,b}$, we use the isomorphisms

$$(S_\alpha^2 \square 1_{\alpha^\#}) \otimes \dagger \tilde{e}v_\alpha \longrightarrow \tilde{e}v_\alpha^\dagger, \quad (1_\alpha \square S_{\alpha^\#}^{-2}) \otimes \tilde{e}v_\alpha^\dagger \longrightarrow \dagger \tilde{e}v_\alpha \quad (3.13)$$

obtained from (3.6). The above data have a familiar structure. (Recall the definition of closed Λ_0 -Frobenius algebras in Section 2.2.)

Proposition 3.4. *The data $\{C_a\}_{a \in \mathbb{Z}}$ and $\eta_1, \varepsilon_{-1}, \{\mu_{a,b}, \Delta_{a,b}\}_{a,b \in \mathbb{Z}}$ have the properties of a closed Λ_0 -Frobenius algebra in the symmetric monoidal category $\mathcal{B}(1, 1)$.*

Proof. The fact that $\eta_1, \varepsilon_{-1}, \{\mu_{a,b}, \Delta_{a,b}\}_{a,b \in \mathbb{Z}}$ satisfy the (co)associativity, (co)unitality, and Frobenius conditions is straightforward to check using the diagrammatic calculus for monoidal 2-categories. The remaining defining relations (2.8)–(2.10) of a closed Λ_0 -Frobenius algebra are more difficult to verify directly. Instead, we use the

framed cobordism hypothesis (see Theorem 3.8 below) to argue indirectly: to $\alpha \in \mathcal{B}^{\text{fd}}$ corresponds a symmetric monoidal functor $\mathcal{Z}: \text{Bord}_{2,1,0}^{\text{fr}} \rightarrow \mathcal{B}$ with $\mathcal{Z}(+) = \alpha$ such that the data $\eta_1, \varepsilon_{-1}, \{\mu_{a,b}, \Delta_{a,b}\}_{a,b \in \mathbb{Z}}$ are the images under \mathcal{Z} of 2-morphisms in $\text{Bord}_{2,1,0}^{\text{fr}}$. The latter in turn are generators of the framed bordism 1-category and satisfy all the relations of a closed Λ_0 -Frobenius algebras, which follows from the special case $r = 0$ of [39, Theorem 5.2.1]. Hence, also their images $\eta_1, \varepsilon_{-1}, \{\mu_{a,b}, \Delta_{a,b}\}_{a,b \in \mathbb{Z}}$ in \mathcal{B} satisfy these relations. ■

Together with the r -spin cobordism hypothesis proved in Section 3.3 below, this implies the following.

Corollary 3.5. *For $r \in \mathbb{Z}_{\geq 1}$, every isomorphism $S_\alpha^r \cong 1_\alpha$ induces a closed Λ_r -Frobenius algebra structure on $\{C_a\}_{a \in \{0,1,\dots,r-1\}}$.*

Proof. Combine [39, Theorem 5.2.1] for $r \in \mathbb{Z}_{\geq 1}$ with Theorem 3.19 below. This in particular guarantees the existence of isomorphisms $C_a \cong C_{a+r}$ for all $a \in \mathbb{Z}$. ■

3.2. The 2-category of r -spin bordisms

3.2.1. 2-categories of bordisms with tangential structure. Here, we briefly recall 2-categories of bordisms with G -structure. For more background and details, we refer to [38, Sections 3.1–3.3].

We begin by fixing conventions for double categories. A *double category* \mathbb{D} consists of a category of objects \mathbb{D}_0 , a category of horizontal morphisms \mathbb{D}_1 , unit horizontal morphisms, a composition functor, and natural transformations that implement associativity and unitality of the composition. The morphisms of \mathbb{D}_1 are called 2-morphisms. The *horizontal 2-category* of a double category \mathbb{D} is the 2-category consisting of the objects of \mathbb{D}_0 , horizontal 1-morphisms and 2-morphisms between parallel 1-morphisms.

We continue with a sketch of the double category of bordisms with tangential structure for a chosen group homomorphism $\xi: G \rightarrow \text{GL}_2$, which we denote Bord^G . A (*2-*)*halo* of a d -dimensional manifold ($d \leq 2$) is loosely speaking a stratified 2-dimensional manifold in which the d -manifold is embedded. We will not need the precise definition, but we refer to Figure 3.1 for illustrative examples of the cases $d = 0$ and $d = 1$. A *d -bordism* between two $(d - 1)$ -dimensional manifolds with haloes S and T is a d -dimensional compact manifold M together with an embedding $S \sqcup T \hookrightarrow M$ that identifies the boundary of M with the $(d - 1)$ -dimensional manifold underlying the halo $S \sqcup T$. We write $M: S \rightarrow T$. A *diffeomorphism* of such a bordism is a diffeomorphism which is compatible with the boundary parametrisation maps.

The objects of $(\text{Bord}^G)_0$ are compact 0-dimensional manifolds with haloes with G -structure, and morphisms are diffeomorphisms of these haloes with G -structure.




Figure 3.1. 0- and 1-dimensional manifolds (below) and their (vertical) inclusions into 2-haloes (above). The horizontal embeddings give a 1-bordism $M: S \rightarrow T$.

The objects of $(\text{Bord}^G)_1$ are 1-bordisms with G -structure (recall Section 2.1), and its morphisms are diffeomorphism classes of 2-bordisms with G -structure. The composition functor is given by gluing of bordisms.

The double categories Bord^G are symmetric monoidal via the disjoint union. The 2-category of bordisms with G -structure $\text{Bord}_{2,1,0}^G$ is defined to be the horizontal 2-category of Bord^G . We will use the notation

$$\text{Bord}_{2,1,0}^{\text{fr}}, \quad \text{Bord}_{2,1,0}^{\text{or}}, \quad \text{Bord}_{2,1,0}^{r\text{-spin}}$$

for the 2-categories of framed, oriented, and r -spin bordisms, respectively. By [42, Theorem 1.1], these categories inherit a symmetric monoidal structure from the respective double categories.

3.2.2. Functors from group homomorphisms. Consider the following commutative diagram of homomorphisms of topological groups:

$$\begin{array}{ccc} G & \xrightarrow{\lambda} & G' \\ & \searrow \xi & \swarrow \xi' \\ & & \text{GL}_2 \end{array}$$

For a G -structure (P, q) on a surface Σ , the group homomorphism λ induces a G' -structure on Σ via the associated bundle construction:

$$\begin{array}{ccc} P \times_{\lambda} G' & \xrightarrow{q\lambda} & F\Sigma \\ & \searrow \pi_{\lambda} & \swarrow \\ & & \Sigma \end{array}$$

This construction is compatible with gluing of bordisms with tangential structure and with disjoint union. Hence, it gives rise to symmetric monoidal functors of double categories and of 2-categories:

$$\mathrm{Bord}^G \xrightarrow{\lambda} \mathrm{Bord}^{G'}, \quad \mathrm{Bord}_{2,1,0}^G \xrightarrow{\Lambda\lambda} \mathrm{Bord}_{2,1,0}^{G'}. \quad (3.14)$$

The group homomorphisms in (2.1)–(2.3) fit into the commutative diagram

$$\begin{array}{ccccc} \star & \xleftarrow{\tilde{\lambda}} & \widetilde{\mathrm{GL}}_2^{+r} & \xrightarrow{\lambda := p^r} & \mathrm{GL}_2^+ \\ & \searrow & \downarrow \iota \circ p^r & \swarrow \iota & \\ & & \mathrm{GL}_2 & & \end{array} \quad (3.15)$$

and induce symmetric monoidal functors:

$$\mathrm{Bord}_{2,1,0}^{\mathrm{fr}} \xrightarrow{\tilde{\Lambda} := \Lambda\tilde{\lambda}} \mathrm{Bord}_{2,1,0}^{r\text{-spin}} \xrightarrow{\Lambda := \Lambda p^r} \mathrm{Bord}_{2,1,0}^{\mathrm{or}},$$

where we use the notations of (3.14) and (3.15). The functor $\tilde{\Lambda}$ assigns to a framed manifold the manifold with the trivial r -spin structure corresponding to the orientation induced by the framing, and Λ assigns to a haloed r -spin surface the haloed surface with the underlying orientation.

3.3. Fully extended r -spin TQFTs

In this section, we consider 2-dimensional extended TQFTs with tangential structure and the cobordism hypothesis for r -spin structures, $r \in \mathbb{Z}_{\geq 0}$. For this, we first recall the framed cobordism hypothesis, homotopy group actions on 2-categories, and their homotopy fixed points. The latter are expected to describe TQFTs with tangential structures, as is known to be the case for oriented (or equivalently: 1-spin) TQFTs. After a review of earlier results in the oriented case, we give a presentation of all r -spin bordism 2-categories in terms of fully dualisable objects and prove the r -spin cobordism hypothesis (for 2-categories, not for $(\infty, 2)$ -categories).

Definition 3.6. Let \mathcal{B} be a symmetric monoidal 2-category. A *fully extended 2-dimensional TQFT with G -structure valued in \mathcal{B}* is a symmetric monoidal functor

$$\mathcal{Z}: \mathrm{Bord}_{2,1,0}^G \longrightarrow \mathcal{B}.$$

We write $\mathrm{Fun}^{\mathrm{sm}}(\mathrm{Bord}_{2,1,0}^G, \mathcal{B})$ for the symmetric monoidal 2-groupoid of fully extended TQFTs with G -structure and values in \mathcal{B} .

3.3.1. The framed cobordism hypothesis. Denote by $2D^0$ the symmetric monoidal 2-category freely generated by a single 2-dualisable object $+$; cf. [34,38]. Our slightly ambiguous notation for the generating object in $2D^0$ draws justification from the following fact.

Theorem 3.7 ([34, Theorem 7.1]). *There is an equivalence of symmetric monoidal 2-categories*

$$\begin{aligned} \iota^0: 2D^0 &\longrightarrow \text{Bord}_{2,1,0}^{\text{fr}} \\ + &\longmapsto +, \end{aligned} \tag{3.16}$$

sending the object $+$ in $2D^0$ to the positively framed (halo of a) point $+$.

The framed cobordism hypothesis classifies framed fully extended TQFTs in terms of fully dualisable objects.

Theorem 3.8 ([34, Theorem 8.1]). *The 2-groupoid of framed fully extended TQFTs with target \mathcal{B} is equivalent to the core of the 2-category of fully dualisable objects in \mathcal{B} as a symmetric monoidal 2-groupoid:*

$$\text{Fun}^{\text{sm}}(\text{Bord}_{2,1,0}^{\text{fr}}, \mathcal{B}) \cong (\mathcal{B}^{\text{fd}})^{\times}.$$

3.3.2. Homotopy G -actions on 2-categories. In order to state the cobordism hypothesis with orientation and more generally with r -spin structure, we will need the notion of homotopy action of a group on a 2-category, as well as its fixed points.

Let G be a topological group. The *homotopy action of G* on a symmetric monoidal 2-category \mathcal{B} is a monoidal functor

$$\rho: \prod_{\leq 2}(G) \longrightarrow \text{Aut}^{\text{sm}}(\mathcal{B})$$

from the fundamental 2-groupoid of G to the 2-category of symmetric monoidal autoequivalences of \mathcal{B} . On $\prod_{\leq 2}(G)$, the monoidal structure comes from the group structure on G ; on Aut^{sm} , it is the composition of functors. Equivalently, a homotopy G -action on \mathcal{B} is a functor

$$\begin{aligned} \text{B} \prod_{\leq 2}(G) &\longrightarrow \{\text{symmetric monoidal 2-categories}\} \\ \star &\longmapsto \mathcal{B} \end{aligned} \tag{3.17}$$

from the delooping of $\prod_{\leq 2}(G)$ to the 3-category of symmetric monoidal 2-categories.

For $G = \text{GL}_2^+ \simeq \text{SO}_2$, the fundamental 2-groupoid is equivalent to the 2-groupoid $\text{B}\mathbb{Z}$ with a single object \star with automorphism group, the free abelian group on a single generator \mathbb{Z} , and only identity 2-morphisms. Below we will identify $\prod_{\leq 2}(\text{GL}_2^+)$ with the 2-groupoid $\text{B}\mathbb{Z}$.

To define an action ρ of GL_2^+ on a 2-category, we only need to specify the value of ρ on the generator $-1 \in \mathbb{Z}$. Recall from Proposition 3.2 the Serre automorphism $S: \mathrm{Id}_{\mathcal{B}^{\mathrm{fd}}} \rightarrow \mathrm{Id}_{\mathcal{B}^{\mathrm{fd}}}$. We define the homotopy action of GL_2^+ on fully dualisable objects as follows:

$$\begin{aligned} \rho: \prod_{\leq 2} (\mathrm{GL}_2^+) &\longrightarrow \mathrm{Aut}^{\mathrm{sm}}(\mathcal{B}^{\mathrm{fd}}) \\ \star &\longmapsto \mathrm{Id}_{\mathcal{B}^{\mathrm{fd}}} \\ \mathbb{Z} \ni -1 &\longmapsto S \\ 1 &\longmapsto 1. \end{aligned} \tag{3.18}$$

Similarly, the homotopy action of the r -spin group is defined through the r th power of the Serre automorphism:

$$\begin{aligned} \rho^r: \prod_{\leq 2} (\widetilde{\mathrm{GL}}_2^{+r}) &\longrightarrow \mathrm{Aut}^{\mathrm{sm}}(\mathcal{B}^{\mathrm{fd}}) \\ \star &\longmapsto \mathrm{Id}_{\mathcal{B}^{\mathrm{fd}}} \\ \mathbb{Z} \ni -1 &\longmapsto S^r \\ 1 &\longmapsto 1. \end{aligned}$$

Note that this action is the GL_2^+ -action (3.18) composed with the functor induced from the covering map $p^r: \widetilde{\mathrm{GL}}_2^{+r} \rightarrow \mathrm{GL}_2^+$ in (2.3).

3.3.3. Presentations of r -spin bordism 2-categories. The 2-category of *homotopy fixed points* \mathcal{B}^G of a homotopy action ρ as in (3.17) is defined to be the 2-category of natural transformations of functors of 3-categories:

$$\mathcal{B}^G = \mathrm{Nat}(\Delta_\star, \rho),$$

where the constant functor $\Delta_\star: \mathbf{B} \prod_{\leq 2} (G) \rightarrow \{\mathrm{sym. mon. 2-cat.}\}$ sends the unique object in $\mathbf{B} \prod_{\leq 2} (G)$ to the 2-category \star with a single object and only identity morphisms, see [26, Remark 3.11–3.14]. It is expected that \mathcal{B}^G is the 3-limit of the functor (3.17), but we are not aware of a rigorous development of the theory of 3-limits.

By the cobordism hypothesis, it is expected that 2-dimensional fully extended TQFTs with G -structure and target \mathcal{B} are classified by homotopy fixed points of a G -action on $(\mathcal{B}^{\mathrm{fd}})^\times$, originating from the G -action on $\mathrm{Bord}_{2,1,0}^{\mathrm{fr}}$. To our knowledge, there is no complete proof for arbitrary G available in the literature, but in the case of orientations this is a known theorem.

Theorem 3.9 ([25, Corollary 5.9]). *The 2-groupoid of oriented fully extended TQFTs with target \mathcal{B} is equivalent to the 2-groupoid of homotopy fixed points of the SO_2 -action on the core of fully dualisable objects in \mathcal{B} :*

$$\mathrm{Fun}^{\mathrm{sm}}(\mathrm{Bord}_{2,1,0}^{\mathrm{or}}, \mathcal{B}) \cong [(\mathcal{B}^{\mathrm{fd}})^\times]^{\mathrm{SO}_2}.$$

The proof in [25] of this uses the presentation of $\text{Bord}_{2,1,0}^{\text{or}}$ from [38], which is not in terms of 2-dualisability data. We also mention that the equivalence as stated is one of 2-groupoids, but later we will see that this can be extended to an equivalence of symmetric monoidal 2-groupoids.

Let $n \in \mathbb{Z}_{\geq 1}$. Given a symmetric monoidal 2-category \mathcal{B} , we define a 2-category $2\text{D}^n(\mathcal{B}^{\text{fd}})$. For $n = 1$, this reduces to the 2-category in [27, Theorem 4.3] which is equivalent to the SO_2 -homotopy fixed points of $(\mathcal{B}^{\text{fd}})^{\times}$. Later, we will consider the case $n = r$ for r -spin TQFTs with $r \geq 2$.

- Objects of $2\text{D}^n(\mathcal{B}^{\text{fd}})$ are pairs (α, θ) , where $\alpha \in \mathcal{B}^{\text{fd}}$, and $\theta: S_{\alpha}^n \rightarrow 1_{\alpha}$ is a 2-isomorphism in \mathcal{B}^{fd} .
- A 1-morphism $(\alpha, \theta) \rightarrow (\alpha', \theta')$ in $2\text{D}^n(\mathcal{B}^{\text{fd}})$ is a 1-morphism $X: \alpha \rightarrow \alpha'$ in \mathcal{B}^{fd} such that the following diagram commutes:

$$\begin{array}{ccc}
 X \otimes S_{\alpha}^n & \xrightarrow{1_X \otimes \theta} & X \otimes 1_{\alpha} \\
 \downarrow S_X^n & & \searrow \cong \\
 S_{\alpha'}^n \otimes X & \xrightarrow{\theta' \otimes 1_X} & 1_{\alpha'} \otimes X \\
 & & \nearrow \cong \\
 & & X
 \end{array} \quad (3.19)$$

- A 2-morphism $X \rightarrow Y$ in $2\text{D}^n(\mathcal{B}^{\text{fd}})$ is a 2-morphism $X \rightarrow Y$ in \mathcal{B}^{fd} .
- Composition and units of $2\text{D}^n(\mathcal{B}^{\text{fd}})$ are induced from \mathcal{B}^{fd} .

To keep the cases $n = 1$ and $n \neq 1$ separate, for a given object $(\alpha, \theta) \in 2\text{D}^n(\mathcal{B}^{\text{fd}})$, we write $\lambda := \theta$ if $n = 1$, and for $n = r \notin \{0, 1\}$, we write $\vartheta := \theta$.

Theorem 3.10 ([27, Theorem 4.3]). *There is an equivalence of 2-categories:*

$$[\mathcal{B}^{\text{fd}}]^{\text{SO}_2} \cong 2\text{D}^1(\mathcal{B}^{\text{fd}}).$$

In the following, we will determine a presentation of $\text{Bord}_{2,1,0}^{\text{or}}$ and $\text{Bord}_{2,1,0}^{r\text{-spin}}$ in terms of 2-dualisability data. The results are collected in Theorems 3.14 and 3.17, but first we need some preparation.

Lemma 3.11. *Let G be a topological group, and let $\xi: G \rightarrow \text{GL}_2$ be a continuous group homomorphism.*

- (i) *Every object in $\text{Bord}_{2,1,0}^G$ is isomorphic to a disjoint union of points with trivial G -structure.*
- (ii) *Every object in $\text{Bord}_{2,1,0}^G$ is fully dualisable.*

Proof. Every connected component c of the underlying manifold of an object in $\text{Bord}_{2,1,0}^G$ is contractible; hence, the G -structure on c is trivialisable. The mapping cylinder for a trivialisisation gives an isomorphism in $\text{Bord}_{2,1,0}^G$. This proves part (i).

To prove part (ii), consider the commutative diagram of group homomorphisms:

$$\begin{array}{ccc}
 \star & \xrightarrow{\text{incl.}} & G \\
 & \searrow & \swarrow \xi \\
 & & \text{GL}_2
 \end{array}$$

from which we get the induced functor Λ_{incl} as in (3.14). Composing this with the functor in (3.16) provides a symmetric monoidal functor:

$$2D^0 \xrightarrow{\iota^0} \text{Bord}_{2,1,0}^{\text{fr}} \xrightarrow{\Lambda_{\text{incl.}}} \text{Bord}_{2,1,0}^G.$$

This composition sends $+\in 2D^0$ to the haloed point with trivial G -structure, and symmetric monoidality implies that the image of $+$ is fully dualisable. The claim of part (i) then completes the proof. ■

A deck transformation on an r -spin surface (P, q, Σ) is an automorphism of the r -spin structure (P, q) which permutes the elements of each fibre of the \mathbb{Z}_r -bundle $q: P \rightarrow F\Sigma$. We also refer to an r -spin bordism as a deck transformation if it is a mapping cylinder of a deck transformation. The 1-morphism components S_p for $p \in \text{Bord}_{2,1,0}^{r\text{-spin}}$ of the Serre functor on $\text{Bord}_{2,1,0}^{r\text{-spin}}$ are isomorphic to deck transformations ([17, Remark 1.3.1]):

For later use, we recall from Example 2.3 (ii) the relation between deck transformations and Nakayama automorphisms.

Lemma 3.12. *The Nakayama automorphisms $N_a: C_a \rightarrow C_a$ of the closed Λ_r -Frobenius algebra in $\text{Bord}_{2,1,0}^{r\text{-spin}}(\emptyset, \emptyset)$ are deck transformations.*

Another way to express this relation is as follows.

Lemma 3.13. *There are invertible modifications:*

$$\lambda: S \xrightarrow{\cong} 1_{\text{Id}_{\text{Bord}_{2,1,0}^{\text{or}}}} \quad \text{and} \quad \vartheta: S^r \xrightarrow{\cong} 1_{\text{Id}_{\text{Bord}_{2,1,0}^{r\text{-spin}}}}.$$

Proof. In $\text{Bord}_{2,1,0}^{\text{or}}$, the 1-morphism components of the Serre automorphism S are diffeomorphic to the identity, and mapping cylinders of these diffeomorphisms assemble into the modification λ . In $\text{Bord}_{2,1,0}^{r\text{-spin}}$, the r th power of the 1-morphism components of S are diffeomorphic to the r th power of a deck transformation, which in turn is isomorphic to the identity, thus providing ϑ . ■

This motivates the following definition of a symmetric monoidal 2-category $2\mathcal{D}^n$ via generators and relations for every $n \in \mathbb{Z}_{\geq 1}$. The generators of $2\mathcal{D}^n$ are the objects 1- and 2-morphisms of $2\mathcal{D}^0$ (cf. Section 3.3.1) together with additional 2-morphisms:

$$\theta_\alpha: S_\alpha^n \longrightarrow 1_\alpha, \quad \theta_\alpha^{-1}: 1_\alpha \longrightarrow S_\alpha^n$$

for all $\alpha \in 2\mathcal{D}^n$. The relations of $2\mathcal{D}^n$ are

- the relations of $2\mathcal{D}^0$,
- $\theta_\alpha \circ \theta_\alpha^{-1} = 1_{1_\alpha}$ and $\theta_\alpha^{-1} \circ \theta_\alpha = 1_{S_\alpha^n}$,
- the commutativity of the diagram

$$\begin{array}{ccc} X \otimes S_\alpha^n & \xrightarrow{1_X \otimes \theta_\alpha} & X \otimes 1_\alpha \\ \downarrow S_X^n & & \searrow \cong \\ S_{\alpha'}^n \otimes X & \xrightarrow{\theta_{\alpha'} \otimes 1_X} & 1_{\alpha'} \otimes X \\ & & \nearrow \cong \\ & & X \end{array} \quad (3.21)$$

for all $\alpha, \alpha' \in 2\mathcal{D}^0$ and $X \in 2\mathcal{D}^0(\alpha, \alpha')$.

We note that the condition in (3.21) expresses the naturality of S . Furthermore, the θ_α are components of an invertible modification $\theta: S^n \rightarrow 1_{\text{Id}_{2\mathcal{D}^n}}$. For $n = 1$, we write $\lambda_\alpha := \theta_\alpha$, and for $n = r$, we write $\vartheta_\alpha := \theta_\alpha$.

Theorems 3.9 and 3.10 together with the 3-categorical Yoneda lemma [10, Theorem 2.12] imply the following theorem.

Theorem 3.14. *There is an equivalence of symmetric monoidal 2-categories:*

$$\iota^1: 2\mathcal{D}^1 \xrightarrow{\cong} \text{Bord}_{2,1,0}^{\text{or}}.$$

Proof. We have a chain of equivalences:

$$\text{Fun}^{\text{sm}}(\text{Bord}_{2,1,0}^{\text{or}}, \mathcal{B}) \cong [(\mathcal{B}^{\text{fd}})^{\times}]^{\text{SO}_2} \cong 2\mathcal{D}^1((\mathcal{B}^{\text{fd}})^{\times}) \cong \text{Fun}^{\text{sm}}(2\mathcal{D}^1, \mathcal{B}), \quad (3.22)$$

all natural in \mathcal{B} . The first equivalence is from Theorem 3.9, the second is from Theorem 3.10. To explain the last equivalence, we will define functors

$$\begin{array}{ccc} & F & \\ \text{Fun}^{\text{sm}}(2\mathcal{D}^1, \mathcal{B}) & \xrightarrow{\quad} & 2\mathcal{D}^1((\mathcal{B}^{\text{fd}})^{\times}) \\ & F^{-1} & \end{array}$$

and then show that F^{-1} is in fact a weak inverse of F .

The functor F is the evaluation on the generating object $+ \in 2\mathcal{D}^1$ and the corresponding generating 2-morphism λ_+ . In detail,

- for a functor $Y \in \text{Fun}^{\text{sm}}(2\mathcal{D}^1, \mathcal{B})$, we set $F(Y) := (Y(+), Y(\lambda_+))$;
- for a natural transformation $f: Y \rightarrow Y'$, we set

$$F(f) := (f_+: (Y(+), Y(\lambda_+)) \rightarrow (Y'(+), Y'(\lambda_+)));$$

- for a modification $\varphi: f \rightarrow f'$, we set $F(\varphi) := (\varphi_+: f_+ \rightarrow f'_+)$.

We need to show that F indeed lands in $2\mathcal{D}^1((\mathcal{B}^{\text{fd}})^{\times})$; i.e., the diagram (3.19) commutes for $X = f_+$ and $n = 1$. The 2-morphism component of f for $U: \alpha \rightarrow \alpha'$ is $f_U: f_{\alpha'} \otimes Y(U) \rightarrow Y'(U) \otimes f_{\alpha}$. Substituting $\alpha = \alpha'$ and $U = S_{\alpha}$, we get

$$f_{S_{\alpha}}: f_{\alpha} \otimes S_{Y(\alpha)} \rightarrow S_{Y'(\alpha)} \otimes f_{\alpha},$$

where we used monoidality of Y to obtain $Y(S_{\alpha}) \cong S_{Y(\alpha)}$, etc. The key observation is that by functoriality and monoidality of f we have $f_{S_{\alpha}} = S_{f_{\alpha}}$. Naturality, of f implies that (3.19) then indeed commutes:

$$\begin{array}{ccc} f_{\alpha} \otimes S_{Y(\alpha)} & \xrightarrow{1_{f_{\alpha}} \otimes Y(\lambda_{\alpha})} & f_{\alpha} \otimes 1_{Y(\alpha)} & \xrightarrow{\cong} & f_{\alpha} \\ f_{S_{\alpha}} = S_{f_{\alpha}} \downarrow & & & \searrow \cong & \\ S_{Y'(\alpha)} \otimes f_{\alpha} & \xrightarrow{1_{Y'(\alpha)} \otimes f_{\alpha}} & 1_{Y'(\alpha)} \otimes f_{\alpha} & \xrightarrow{\cong} & f_{\alpha} \end{array} \quad (3.23)$$

Now, we construct the functor F^{-1} . Since $2\mathcal{D}^1$ is defined in terms of generators and relations, in order to define a symmetric monoidal functor $Y: 2\mathcal{D}^1 \rightarrow \mathcal{B}$ it is enough to specify the value of Y on the generating objects and morphisms. Then, one needs to check that relations in $2\mathcal{D}^1$ are sent to relations in \mathcal{B} . The same holds for defining natural transformations and modifications, where we only need to specify their components on generators.

- For $(\alpha, \theta) \in 2D^1((\mathcal{B}^{\text{fd}})^\times)$, we let $F^{-1}(\alpha, \theta)$ be the functor Y with values $Y(+) = \alpha$ and $Y(\lambda_+) = \theta$. This determines the value of Y on all of $2D^1$.
- For $X: (\alpha, \theta) \rightarrow (\alpha', \theta')$, we let $F^{-1}(X)$ to be the monoidal natural transformation $f: Y \rightarrow Y'$ with 1-morphism component $f_+ = X$. This determines all 1- and 2-morphism components of f .
- For $\phi: X \rightarrow X'$, we let $F^{-1}(\phi)$ be the modification $\varphi: f \rightarrow f'$ with 2-morphism component $\varphi_+ = \phi$. This determines all 2-morphism components of φ .

We need to check that F^{-1} is well defined. First, we note that $Y = F^{-1}(\alpha, \theta)$ is indeed a functor $2D^1 \rightarrow \mathcal{B}$ because the relations in $2D^0$ are satisfied by definition of Y , and so are the relations on λ and its inverse. By symmetric monoidality, Y sends S_+ in $2D^1$ to S_α in $2D^1((\mathcal{B}^{\text{fd}})^\times)$; hence, naturality is satisfied as well. $F^{-1}(X)$ is a natural transformation, since its components satisfy (3.23). For modifications, there are no further conditions to check.

By construction, we have

$$F \circ F^{-1} = \text{Id}.$$

Moreover, we also have $F^{-1} \circ F \cong \text{Id}$, since two functors out of $2D^1$ are isomorphic if they agree on generators [38, Theorem 2.78]. This shows that F is an equivalence.

Finally, we observe that the functor ι^1 respects the symmetric monoidal structures on $2D^1$ and $\text{Bord}_{2,1,0}^{\text{or}}$ and hence, it can canonically be promoted to a symmetric monoidal functor. Thus, the claim follows from the 3-categorical Yoneda lemma of [10, Theorem 2.12]. ■

By looking at the chain of equivalences in (3.22), we can read off the value of ι^1 on generators of the 2-category $2D^1$:

- ι^1 sends $+ \in 2D^1$ together with its 2-dualisability data to the positively oriented point with its 2-dualisability data (cf. Lemma 3.11);
- $\iota^1(\lambda_+) = \lambda_+$ from Lemma 3.13.

Remark 3.15. Using the symmetric monoidal equivalence ι^1 , one also obtains a symmetric monoidal structure on the equivalence in Theorem 3.9.

In line with the above presentation of ι^1 , we now define a symmetric monoidal functor

$$\iota^r: 2D^r \longrightarrow \text{Bord}_{2,1,0}^{r\text{-spin}} \quad (3.24)$$

on generators of $2D^r$ for $r \in \mathbb{Z}_{\geq 2}$ as follows:

- ι^r sends $+ \in 2D^r$, together with its 2-dualisability data, to the point with trivial r -spin structure in $\text{Bord}_{2,1,0}^{r\text{-spin}}$, with its 2-dualisability data as in Lemma 3.11;
- $\iota^r(\vartheta_+) = \vartheta_+$ from Lemma 3.13.

Proposition 3.16. *We have a strictly commutative diagram of symmetric monoidal functors:*

$$\begin{array}{ccccc}
 \text{Bord}_{2,1,0}^{\text{fr}} & \xrightarrow{\tilde{\Lambda}} & \text{Bord}_{2,1,0}^{r\text{-spin}} & \xrightarrow{\Lambda} & \text{Bord}_{2,1,0}^{\text{or}} \\
 \uparrow \iota^0 & & \uparrow \iota^r & & \uparrow \iota^1 \\
 2\text{D}^0 & \xrightarrow{\tilde{K}} & 2\text{D}^r & \xrightarrow{K} & 2\text{D}^1
 \end{array}$$

where the functors \tilde{K} and K by definition act as the identity on objects as well as on 1- and 2-morphism generators of 2D^0 , while on the other generators, we have

$$K \left(S_+^r \begin{array}{c} \xleftarrow{\vartheta_+} \\ \xrightarrow{\vartheta_+^{-1}} \end{array} 1_+ \right) = \left(S_+^r \begin{array}{c} \xleftarrow{\lambda_+^r} \\ \xrightarrow{\lambda_+^{-r}} \end{array} 1_+ \begin{array}{c} \xleftarrow{\cong} \\ \xrightarrow{\cong} \end{array} 1_+ \right). \quad (3.25)$$

By Theorems 3.7 and 3.14, the functors ι^0 and ι^1 are equivalences. In Section 3.3.4 below, we will prove the following theorem.

Theorem 3.17. *The functor in (3.24) is an equivalence for all $r \in \mathbb{Z}_{\geq 1}$,*

$$\iota^r: 2\text{D}^r \xrightarrow{\cong} \text{Bord}_{2,1,0}^{r\text{-spin}}.$$

We also have the analogous statement of Theorem 3.10, which follows immediately by applying [27, Theorem 4.3] to the natural transformation S^r .

Lemma 3.18. *The homotopy fixed points of the r -spin action on \mathcal{B}^{fd} are*

$$[\mathcal{B}^{\text{fd}}]^{\text{Spin}_2^r} \cong 2\text{D}^r(\mathcal{B}^{\text{fd}}).$$

Theorem 3.17 and Lemma 3.18 imply the 2-categorical cobordism hypothesis with r -spin structure.

Theorem 3.19. *The 2-groupoid of fully extended r -spin TQFTs with target \mathcal{B} is equivalent to the homotopy fixed points of the r -spin action:*

$$\text{Fun}^{\text{sm}}(\text{Bord}_{2,1,0}^{r\text{-spin}}, \mathcal{B}) \cong [(\mathcal{B}^{\text{fd}})^{\times}]^{\text{Spin}_2^r}.$$

Proof. We have a chain of equivalences:

$$\text{Fun}^{\text{sm}}(\text{Bord}_{2,1,0}^{r\text{-spin}}, \mathcal{B}) \cong \text{Fun}^{\text{sm}}(2\text{D}^r, \mathcal{B}) \cong 2\text{D}^r((\mathcal{B}^{\text{fd}})^{\times}) \cong [(\mathcal{B}^{\text{fd}})^{\times}]^{\text{Spin}_2^r}.$$

The first equivalence is from Theorem 3.17, the last equivalence is from Lemma 3.18, and the proof of the second equivalence is completely analogous to the $n = 1$ case in the proof of Theorem 3.14. \blacksquare

Remark 3.20. The proof of the oriented cobordism hypothesis in [25] (Theorem 3.9) uses the presentation of $\text{Bord}_{2,1,0}^{\text{or}}$ of [38], which is not in terms of 2-dualisability data, and a direct computation of the SO_2 -homotopy fixed points $[(\mathcal{B}^{\text{fd}})^{\times}]^{\text{SO}_2}$ (Theorem 3.10). In order to prove the r -spin cobordism hypothesis (Theorem 3.19), we need a presentation of $\text{Bord}_{2,1,0}^{r\text{-spin}}$ (Theorem 3.17) in terms of 2-dualisability data and a direct computation of Spin_2^r -homotopy fixed points $[(\mathcal{B}^{\text{fd}})^{\times}]^{\text{Spin}_2^r}$ (Lemma 3.18).

It is straightforward to check the following factorisation of r -spin TQFTs.

Proposition 3.21. *Let $\alpha \in \mathcal{B}^{\text{fd}}$ and $k < r$ be such that there are 2-isomorphisms $\varphi: S_\alpha^k \rightarrow 1_\alpha$ and $\psi: S_\alpha^r \rightarrow 1_\alpha$. Then,*

- (i) *there is a 2-isomorphism $\phi: S_\alpha^g \rightarrow 1_\alpha$, where $g = \text{gcd}(k, r)$;*
- (ii) *the diagram*

$$\begin{array}{ccc}
 \text{Bord}_{2,1,0}^{r\text{-spin}} & \xrightarrow{\Lambda_{r,g}} & \text{Bord}_{2,1,0}^{g\text{-spin}} \\
 & \searrow \mathcal{Z}_\psi & \swarrow \mathcal{Z}_\phi \\
 & \mathcal{B} &
 \end{array}$$

commutes up to a natural isomorphism, where $\Lambda_{r,g}$ is the functor from (3.14) for the group homomorphism $\widetilde{\text{GL}}_2^{+r} \rightarrow \widetilde{\text{GL}}_2^{+g}$, while \mathcal{Z}_ψ and \mathcal{Z}_ϕ denote the r -spin and g -spin TQFTs from Theorem 3.19 corresponding to $(\alpha, \psi) \in 2D^r(\mathcal{B}^{\text{fd}})$ and $(\alpha, \phi) \in 2D^g(\mathcal{B}^{\text{fd}})$, respectively.

Remark 3.22. Let us assume that the adjoints of 1-morphisms in \mathcal{B} satisfy $X^\dagger = \dagger X$, which is, for example, the case when \mathcal{B} is pivotal. Then, by the definition of the Serre automorphism (3.4) and its inverse (3.5), we have $S = S^{-1}$. Hence, under this assumption, r -spin TQFTs with target \mathcal{B} factorise through oriented TQFTs (r odd), or through 2-spin TQFTs (r even).

3.3.4. Proof of the r -spin cobordism hypothesis. Here, we prove Theorem 3.17. To do this, we will check the conditions listed in the following Whitehead-type theorem for the functor ι^r in (3.24).

Theorem 3.23 ([38, Theorem 2.25]). *A functor $F: \mathcal{C} \rightarrow \mathcal{B}$ of 2-categories is an equivalence iff it is essentially surjective on objects and 1-morphisms and fully faithful on 2-morphisms.*

Lemma 3.24. *The functor ι^r is essentially surjective on objects.*

Proof. This follows from Lemma 3.11 (i). ■

Lemma 3.25. *The functor ι^r is essentially surjective on 1-morphisms.*

Proof. For every connected component c of a 1-morphism $P \rightarrow FX$ in $\text{Bord}_{2,1,0}^{r\text{-spin}}$, we obtain an element $\delta(c) \in \mathbb{Z}_r$ as follows. If the 1-manifold of which c is a halo is closed, set $\bar{c} := c$; otherwise, let \bar{c} be the r -spin surface with closed embedded 1-manifold defined by identifying the two boundary points of the embedded 1-manifold in c via the boundary parametrisation maps. This identification is possible by choosing a trivialisation of the r -spin structures of the objects parametrising the boundary of c . Consider a curve $\Gamma: S^1 \rightarrow \bar{c}$ parametrising the 1-manifold in \bar{c} and its lift $\tilde{\Gamma}: S^1 \rightarrow F\bar{c}$ to the frame bundle defined by picking at every point a tangent vector to Γ and another vector so that the induced orientation agrees with the orientation underlying the r -spin structure of c . This lift is unique up to homotopy. There is a unique lift $\hat{\Gamma}: S^1 \rightarrow P|_{\bar{c}}$ of $\tilde{\Gamma}$ after fixing it at one point, as the fibres are discrete. We define $\delta(c) \in \mathbb{Z}_r$ to be the holonomy of $\hat{\Gamma}$, which only depends on c ; for more details on this construction, we refer to [36] or [37, Section 5.2].

Recall that, by (3.20), S_+ is isomorphic to a deck transformation with holonomy $-1 \in \mathbb{Z}_r$. If $c = \bar{c}$, then

$$c \cong C_{\delta(c)} \in \text{Bord}_{2,1,0}^{r\text{-spin}}(\emptyset, \emptyset)$$

from (3.8). Otherwise, c is isomorphic either to the endomorphism $S_{\pm}^{-\delta(c)}$, or to $S_{\pm}^{-\delta(c)}$ pre- or post-composed with one of the adjunction 1-morphisms of $+$; for example,

$$c \cong \tilde{e}v_+ \circ (S_+^{-\delta(c)} \sqcup 1_-). \quad \blacksquare$$

Lemma 3.26. *The functor ι^r is full on 2-morphisms.*

Proof. Let $X, X': \alpha \rightarrow \alpha'$ be parallel 1-morphisms in $2D^r$, and let $\Sigma: \iota^r(X) \rightarrow \iota^r(X')$ be a 2-morphism in $\text{Bord}_{2,1,0}^{r\text{-spin}}$. Without loss of generality, we can assume that Σ is connected of genus g . We write $\Lambda(\Sigma)$ for the oriented surface underlying Σ . Hence, the r -spin structure on Σ is represented by a bundle $P \rightarrow \Lambda(\Sigma)$ and a \mathbb{Z}_r -bundle map $q: P \rightarrow F\Lambda(\Sigma)$.

The strategy of our proof is as follows. We describe the r -spin structure on Σ up to diffeomorphisms of r -spin surfaces with underlying diffeomorphism of surfaces the identity, which we refer to as *isomorphisms of r -spin structures*. Then, we consider a decomposition of the oriented surface $\Lambda(\Sigma)$ suitable for our description of r -spin structures. Finally, we lift the oriented generators to r -spin generators and restore the r -spin structure up to isomorphism in the above sense. Therefore, the r -spin surface we build from the generators is in particular diffeomorphic to Σ , thus representing the same 2-morphism in $\text{Bord}_{2,1,0}^{r\text{-spin}}$.

Step 1. Following [36], we describe the r -spin structure of Σ in terms of holonomies along curves in the underlying oriented surface $\Lambda(\Sigma)$ in the \mathbb{Z}_r -bundle $q: P \rightarrow F\Lambda(\Sigma)$.




Figure 3.2. Curves on a closed surface.

Step 1.1. If $\Lambda(\Sigma)$ is a closed surface (i.e., if $\alpha = \alpha' = \emptyset$ and $X = 1_\emptyset = X'$), then r -spin structures up to isomorphism on $\Lambda(\Sigma)$ are in (non-canonical) bijection with

$$H^1(\Lambda(\Sigma), \mathbb{Z}_r) \cong \mathbb{Z}_r^{2g}.$$

The latter bijection is given by picking simple closed curves in $\Lambda(\Sigma)$ which represent a basis of $H_1(\Lambda(\Sigma))$. For each handle, we choose two curves a_k, b_k which intersect at precisely one point and which do not intersect with the curves associated to the other handles; see Figure 3.2.

Step 1.2. If $\Lambda(\Sigma)$ is not a closed surface, we introduce a new surface $\tilde{\Sigma}$ and additional curves on $\Lambda(\Sigma)$.

Step 1.2.1 (definition of $\tilde{\Sigma}$). We define the new oriented surface $\tilde{\Sigma}$ using the boundary parametrisation maps. Let $\partial_i, i \in \{1, \dots, |\pi_0(\partial\Sigma)|\}$, denote the parametrised boundary components of $\Lambda(\Sigma)$, which may be circles or intervals. We arbitrarily single out the component ∂_1 , and we choose a connected subset U_j of an open neighbourhood of ∂_1 for each remaining boundary component $\partial_j, j \in \{2, \dots, |\pi_0(\partial\Sigma)|\}$, so that the U_j are pairwise disjoint. Furthermore, we choose a connected subset V_j of an open neighbourhood of ∂_j in each remaining boundary component ($j \neq 1$). We illustrate such choices in Figure 3.3.

Using the boundary parametrisation maps, we glue $U_j \cap \partial_1$ to $V_j \cap \partial_j$. Finally, we retract each remaining boundary component to a single point. The surface $\tilde{\Sigma}$ obtained in this way is a closed surface, whose genus is the sum of g and the number of closed parametrised boundary components of $\Lambda(\Sigma)$, with a point \tilde{p}_i removed for each parametrised boundary component ∂_i ; see Figure 3.4.

Step 1.2.2 (additional curves). We extend our collection of curves in $\Lambda(\Sigma)$ by defining curves in $\tilde{\Sigma}$: we pick a simple closed curve \tilde{u}_j for each $\partial_j, j \in \{2, \dots, |\pi_0(\partial\Sigma)|\}$, encircling \tilde{p}_j , and not intersecting with each other, as well as a simple closed curve \tilde{d}_i for each boundary component “parallel” along the boundary curve, see Figure 3.4 (b).




Figure 3.3. An example of an oriented surface $\Lambda(\Sigma)$ with non-empty boundary, together with choices of open neighbourhoods near parametrised boundary components (U_i and V_j) and their identification (from Step 1.2.1) along the boundary indicated by thick grey lines. The curves u_j (from Step 1.2.2) are shown in blue.

Then, we obtain the following curves in $\Lambda(\Sigma)$ corresponding the curves in $\tilde{\Sigma}$; see Figure 3.5 (a):

- two closed curves $a_k, b_k, k \in \{1, \dots, g\}$, for each handle, $k \in \{1, \dots, g\}$,
- one curve d_i for each boundary component $\partial_i, i \in \{1, \dots, |\pi_0(\partial\Sigma)|\}$,
- one curve u_j for each boundary component $\partial_j, j \in \{2, \dots, |\pi_0(\partial\Sigma)|\}$.

Since near each U_j and V_j the r -spin surface is trivial, we obtain an r -spin structure on $\tilde{\Sigma}$. Also note that the set of r -spin structures on $\Lambda(\Sigma)$ with prescribed r -spin structure near the boundary and the set of r -spin structures on $\tilde{\Sigma}$ with prescribed r -spin structure near its punctures are in bijection by construction. These sets are in bijection with the set

$$\{\delta \in H^1(\tilde{\Sigma}, \mathbb{Z}_r) \mid \delta(d_i) = x_i \text{ for all } i; \delta(u_j) = y_j \text{ for all } j \text{ with } \partial_j \not\cong S^1\},$$

where x_i is the holonomy along d_i and y_j is the holonomy along u_j , which are fixed by the boundary parametrisation.




Figure 3.4. (a) The surface $\tilde{\Sigma}$ obtained from $\Lambda(\Sigma)$ as in Figure 3.3 in Step 1.2.1, and the additional curves \tilde{u}_j from Step 1.2.2. (b) The surface $\tilde{\Sigma}$ with all the curves $\tilde{a}_k, \tilde{b}_k, \tilde{u}_j, \tilde{d}_i$ from Step 1.2.2.




Figure 3.5. (a) All curves on $\Lambda(\Sigma)$ obtained in Step 1.2.2. (b) Only those curves on $\Lambda(\Sigma)$ whose images in $\tilde{\Sigma}$ form a minimal generating set of $\pi_1(\tilde{\Sigma})$.

In order to describe the r -spin structure on Σ , it is enough to remember the holonomies along a set of curves that generate $\pi_1(\tilde{\Sigma})$. Therefore, we reduce the set of curves a_k, b_k, d_i, u_j by discarding the curves u_j with $\partial_j \not\cong S^1$. The remaining curves are illustrated in Figure 3.5 (b).

Step 2. We pick a decomposition of $\Lambda(\Sigma)$ into oriented generators so that

- for each handle we have the decomposition as in Figure 3.6 (a);
- for each boundary component we have the decomposition as in Figure 3.6 (b).

Note that we can require that 1-morphism components of the Serre automorphism only appear at the boundary of generating 2-morphisms in $\text{Bord}_{2,1,0}^{\text{or}}$ and not in their interior, as in $\text{Bord}_{2,1,0}^{\text{or}}$ there is a trivialisation of the Serre automorphism; cf. Lemma 3.13. We furthermore require that the curves a_k, b_k, u_j, d_i cross the generating 2-morphisms as in Figure 3.7.




Figure 3.6. Decomposition of a surface into oriented generators. (a) Decomposition of a handle. (b) Decomposition near a closed parametrised boundary component.

Step 3. Recall from Lemma 3.12 that the Nakayama automorphisms $N_a: C_a \rightarrow C_a$ are deck transformations. We lift the oriented generators to r -spin generators by inserting (i.e., by replacing a neighbourhood of a_k and d_j with)

- $N_{\delta(a_k)}^{1-\delta(b_k)}$ at the intersection of a_k and b_k ; see Figure 3.7 (a),
- $N_{\delta(d_j)}^{-\delta(u_j)}$ at the intersection of d_j and u_j ; see Figure 3.7 (b),
- $\mu_{-\delta(a_k), \delta(a_k)}$ and $\Delta_{-\delta(a_k), \delta(a_k)}$ from (3.11)–(3.12) at the saddles crossed by b_k ,
- identity 2-morphisms where no intersections occur.

The r -spin structure given by this construction has the same holonomies along the above mentioned curves as the r -spin structure of Σ . Note that a full circle along a positively oriented simple closed loop, where no insertions of $N_{\delta(a_k)}$ appear, contributes +1 to the holonomy. Hence, the two r -spin structures are isomorphic, and thus, the two r -spin surfaces represent the same 2-morphisms. ■

Lemma 3.27. *The functor ι^r is faithful on 2-morphisms.*

Proof. Let $\sigma, \sigma' \in 2D^r(\alpha, \beta)(X, Y)$. Assume that $\iota^r(\sigma) = \iota^r(\sigma')$, and that these are connected bordisms. By Proposition 3.16, we have

$$(\Lambda \circ \iota^r)(\sigma) = (\iota^1 \circ K)(\sigma),$$

and analogously for σ' . Hence, since ι^1 is an equivalence (Theorem 3.14), we have

$$K(\sigma) = K(\sigma').$$

This means that there is a sequence of relations in $2D^1$ relating $K(\sigma)$ and $K(\sigma')$.




Figure 3.7. Insertion of the Nakayama automorphism at the intersection of curves.

By (3.25), the numbers of λ and λ^{-1} in $K(\sigma)$ and $K(\sigma')$ are each divisible by r . Using the coherence theorem for 2-categories, we can bundle together the relations in the sequence involving λ in tuples of r . These can be lifted to relations in $2D^r$ via (3.25). Noting that all other relations in $2D^1$ and $2D^r$ are the same (to wit, those of $2D^0$) and that they can hence also be lifted, it follows that

$$\sigma = \sigma'. \quad \blacksquare$$

3.3.5. Computing invariants of r -spin bordisms with closed boundary. With the r -spin cobordism hypothesis at hand, we can describe the closed Λ_r -Frobenius algebra which classifies the non-extended r -spin TQFT associated to a fully extended r -spin TQFT. In particular, we can explicitly describe the values of the non-extended r -spin TQFT on r -spin surfaces with closed boundary. For convenience, we also present the corresponding results for framed TQFTs.

Recall from Sections 2 and 3.2 that

$$\text{Bord}_{2,1}^{\text{fr}} \cong \text{Bord}_{2,1,0}^{\text{fr}}(\emptyset, \emptyset) \quad \text{and} \quad \text{Bord}_{2,1}^{r\text{-spin}} \cong \text{Bord}_{2,1,0}^{r\text{-spin}}(\emptyset, \emptyset).$$

Consider the fully extended framed and r -spin TQFTs:

$$\begin{aligned} Y: \text{Bord}_{2,1,0}^{\text{fr}} &\longrightarrow \mathcal{B} & \text{and} & & Z: \text{Bord}_{2,1,0}^{r\text{-spin}} &\longrightarrow \mathcal{B} \\ + &\longmapsto \alpha & & & + &\longmapsto \alpha \\ \vartheta_+ &\longmapsto (\vartheta: S_\alpha^r \xrightarrow{\cong} 1_\alpha). \end{aligned}$$

The corresponding non-extended TQFTs

$$Y|: \text{Bord}_{2,1}^{\text{fr}} \longrightarrow \mathcal{B}(1,1) \quad \text{and} \quad Z|: \text{Bord}_{2,1}^{r\text{-spin}} \longrightarrow \mathcal{B}(1,1)$$

$$S_a^1 \longmapsto C_a^\alpha \qquad S_a^1 \longmapsto C_a^\alpha$$

are classified by the closed Λ_0 - and Λ_r -Frobenius algebras in Proposition 3.4 and Corollary 3.5, respectively.

In particular, the invariants assigned to the framed and r -spin tori $T(d)$ introduced in Section 2.3 are quantum dimensions of the circle spaces in $\mathcal{B}(1,1)$:

$$T(d) \longmapsto \dim(C_d^\alpha).$$

Remark 3.28. (i) Assume that, as in Remark 3.22, the left and right adjoints of 1-morphisms in the target 2-category \mathcal{B} agree. In this case, we effectively have oriented TQFTs (r odd) with all the circle spaces being isomorphic, or 2-spin TQFTs (r even) with

$$C_a^\alpha \cong C_{a+2}^\alpha$$

for every $a \in \mathbb{Z}$. Accordingly, the invariants associated to framed and r -spin tori may take at most two distinct values if left and right adjoints agree in \mathcal{B} .

(ii) For oriented and 2-spin surfaces, there already exist TQFTs which compute complete invariants (the oriented TQFT of [35] with target $\text{Vect}_{\mathbb{k}}$ computed from the relative Euler characteristic and the 2-spin TQFT of [1, 37] with target $\text{Vect}_{\mathbb{k}}^{\mathbb{Z}_2}$ computing the Arf invariant). For $r > 2$, TQFTs with pivotal 2-categories as targets cannot distinguish all r -spin structures, but other targets may allow for more interesting r -spin TQFTs.

4. Examples

By the main result of the previous section, constructing extended r -spin TQFTs amounts to finding fully dualisable objects whose Serre automorphisms are such that their r th power is trivialisable. In Section 4.1 we increase our chances to find such objects by passing from a given target 2-category to its “equivariant completion”, where we translate the condition on the Serre automorphism to a condition on the Nakayama automorphism of certain Frobenius algebras (Corollary 4.9), and we study the associated circle spaces and hence torus invariants in detail (Section 4.1.5). Then in Section 4.2 we show that every object in the 2-category of Landau–Ginzburg models \mathcal{LG} gives rise to an extended 2-spin TQFT, and we illustrate how to do computations in the equivariant completion of \mathcal{LG} and its variants.

4.1. Equivariant completion


In this section, we consider the representation 2-category \mathcal{B}_{eq} of certain Frobenius algebras internal to a given symmetric monoidal pivotal 2-category \mathcal{B} . In particular, we explicitly determine the Serre automorphisms and circle spaces associated to objects in \mathcal{B}_{eq} , from which invariants of extended r -spin TQFTs with values in \mathcal{B}_{eq} can be computed with the help of Theorem 3.23. We stress that even if the original 2-category \mathcal{B} is pivotal, its completion \mathcal{B}_{eq} need not be pivotal, which in light of Remark 3.28 is a desired feature.

Throughout this section, we fix a symmetric monoidal pivotal 2-category \mathcal{B} which satisfies the condition (*) below. (The symmetric monoidal structure will not be relevant before Section 4.1.3.)

4.1.1. Equivariant completion of a 2-category. A Δ -separable Frobenius algebra on an object $\alpha \in \mathcal{B}$ consists of

$$A \in \mathcal{B}(\alpha, \alpha), \quad \mu_A = \text{cup}, \quad \eta_A = \text{cap}, \quad \Delta_A = \text{cup}, \quad \varepsilon_A = \text{cap}$$

such that



Recall, e.g., from [16, Section 2.2], the notions of (bi)modules and (bi)module maps over the underlying algebra (A, μ_A, η_A) . If X is a right A -module and Y is a left A -module, then the *relative tensor product* $X \otimes_A Y$ is the coequaliser of the canonical maps $X \otimes A \otimes Y \rightrightarrows X \otimes Y$. Since A is a Δ -separable Frobenius algebra, the map

$$\pi_A^{X,Y} = \text{diagram of a cup with a cap inside, connected to lines X and Y}$$

is an idempotent. If $\pi_A^{X,Y}$ splits, then $X \otimes_A Y$ can be identified with $\text{Im}(\pi_A^{X,Y})$; see, e.g., [16, Lemma 2.3]. Hence, we will make the following assumption.

- (*) For all Δ -separable Frobenius algebras A on all objects of \mathcal{B} , the idempotents $\pi_A^{X,Y}$ split for all modules X, Y , and we choose adjunction data for A such that $\dagger A = A^\dagger$ as well as $\dagger\mu = \mu^\dagger$ and $\dagger\Delta = \Delta^\dagger$.

Thus, we have splitting maps

$$X \otimes_A Y \begin{array}{c} \xrightarrow{\iota_A^{X,Y}} \\ \xleftarrow{\vartheta_A^{X,Y}} \end{array} X \otimes Y$$

with

$$\iota_A^{X,Y} \circ \vartheta_A^{X,Y} = \pi_A^{X,Y} \quad \text{and} \quad \vartheta_A^{X,Y} \circ \iota_A^{X,Y} = 1_{X \otimes_A Y}.$$

Note that every Frobenius algebra is self-dual, so there always exist adjunction data such that $\dagger A = A^\dagger$. The conditions

$$\dagger\mu = \mu^\dagger \quad \text{and} \quad \dagger\Delta = \Delta^\dagger$$

automatically hold if \mathcal{B} is pivotal, but we do not make this stronger assumption on \mathcal{B} .

Definition 4.1. The *equivariant completion* \mathcal{B}_{eq} of \mathcal{B} is the 2-category whose

- objects are pairs (α, A) with $\alpha \in \mathcal{B}$ and $A \in \mathcal{B}(\alpha, \alpha)$, a Δ -separable Frobenius algebra;
- 1-morphisms $(\alpha, A) \rightarrow (\beta, B)$ are 1-morphisms $\alpha \rightarrow \beta$ in \mathcal{B} together with a B - A -bimodule structure;
- 2-morphisms are bimodule maps in \mathcal{B} ;
- horizontal composition is the relative tensor product, and $1_{(\alpha, A)}$ is A with its canonical A - A -bimodule structure;
- vertical composition and unit 2-morphisms are induced from \mathcal{B} .

Equivariant completion was introduced in [16] in connection with generalised orbifold constructions of oriented TQFTs. The attribute “equivariant” derives from the fact that an action $\rho: \underline{G} \rightarrow \mathcal{B}(\alpha, \alpha)$ of a finite group G (viewed as a discrete monoidal category \underline{G}) gives rise to a Δ -separable Frobenius algebra structure on

$$A_G := \bigoplus_{g \in G} \rho(g)$$

if $\mathcal{B}(\alpha, \alpha)$ has finite direct sums and that G -equivariantisation can be described in terms of categories of A_G -modules.

The assignment $\mathcal{B} \mapsto \mathcal{B}_{\text{eq}}$ is a completion in the sense that $(\mathcal{B}_{\text{eq}})_{\text{eq}} \cong \mathcal{B}_{\text{eq}}$; see [16, Proposition 4.2]. Equivariant completion is the same as (unital and counital) “condensation completion” in dimension 2, as introduced in [20] for arbitrary dimension in the context of fully extended framed TQFTs and topological orders.

4.1.2. Adjoints. Recall that we assume that every Δ -separable Frobenius algebra $A \in \mathcal{B}(\alpha, \alpha)$ comes with chosen adjunction data such that $\dagger A = A^\dagger$. Hence, we can define the *Nakayama automorphism* and its inverse as follows:

$$\gamma_A = \left\langle \begin{array}{c} | \\ \text{---} \text{---} \text{---} \\ | \end{array} \right\rangle, \quad \gamma_A^{-1} = \left\langle \begin{array}{c} | \\ \text{---} \text{---} \text{---} \\ | \end{array} \right\rangle. \tag{4.1}$$

Since we also assume that $\dagger \mu = \mu^\dagger$ and $\dagger \Delta = \Delta^\dagger$, the Nakayama automorphism is a map of the underlying algebra and coalgebra structures of A , and A is a symmetric Frobenius algebra iff $\gamma_A = 1_A$; see, e.g., [19].

Remark 4.2. Let \mathcal{C} be a symmetric monoidal 1-category with left duals, and let $A \in \mathcal{C}$ be a Frobenius algebra. We can endow \mathcal{C} with right duals by setting

$$X^\dagger := \dagger X, \quad \tilde{ev}_X := ev_X \circ b_{X^\dagger, X}, \quad \widetilde{coev}_X := b_{X, X^\dagger} \circ coev_X$$

using the braiding b . In this case, the inverse Nakayama automorphism of A is

$$\gamma_A^{-1} = \left\langle \begin{array}{c} | \\ \text{---} \text{---} \text{---} \\ | \end{array} \right\rangle.$$

Comparing this to the definition (2.4) of the Nakayama automorphisms of a closed Λ_0 -Frobenius algebra C , we see that the conventions for these two different Nakayama structures γ and N , for two different algebraic entities A and C , respectively, are not maximally aligned.

Given a B - A -bimodule $X \in \mathcal{B}(\alpha, \beta)$ together with algebra automorphisms $\varphi: A \rightarrow A$ and $\psi: B \rightarrow B$, the ψ - φ -twisted bimodule ${}_\psi X_\varphi$ is given by

$$\begin{array}{c} \psi X_\varphi \\ | \\ \text{---} \text{---} \text{---} \\ | \\ B \psi X_\varphi A \end{array} = \begin{array}{c} X \\ | \\ \text{---} \text{---} \text{---} \\ | \\ B X A \end{array},$$

where the unlabelled vertices on the right-hand side correspond to the original bimodule structure on X .

If the 2-category \mathcal{B} has adjoints for 1-morphisms, then, as shown in [16, Proposition 4.2], its equivariant completion \mathcal{B}_{eq} inherits this property, by twisting with Nakayama automorphisms. This implies that even if \mathcal{B} is pivotal, \mathcal{B}_{eq} typically is not.

Proposition 4.3. *Let \mathcal{B} be a 2-category, and let $X \in \mathcal{B}_{\text{eq}}((\alpha, A), (\beta, B))$ be such that the underlying 1-morphism $X: \alpha \rightarrow \beta$ in \mathcal{B} has left and right adjoints $\dagger X$ and X^\dagger , respectively. Then, X also has left and right adjoints:*

$${}^*X = \gamma_A^{-1}(\dagger X), \quad X^* = (X^\dagger)_{\gamma_B} \quad (4.2)$$

in \mathcal{B}_{eq} , witnessed by the adjunction maps

$$\begin{aligned} \text{ev}_X^{\mathcal{B}_{\text{eq}}} &= \text{diagram} \circ \iota_B^{\dagger X, X}, & \text{coev}_X^{\mathcal{B}_{\text{eq}}} &= \vartheta_A^{X, \dagger X} \circ \text{diagram} \\ \tilde{\text{ev}}_X^{\mathcal{B}_{\text{eq}}} &= \text{diagram} \circ \iota_A^{X, X^\dagger}, & \tilde{\text{coev}}_X^{\mathcal{B}_{\text{eq}}} &= \vartheta_B^{X^\dagger, X} \circ \text{diagram} \end{aligned}$$

As a consistency check, we recall that

are the canonical A -actions on $\dagger A$, and the A - A -bimodule structure on A^\dagger is obtained as the mirror images of the above diagrams. These actions agree by assumption on A . From this, it is straightforward to verify that

$$\begin{aligned} \text{diagram}_1: \gamma_A A &\xrightarrow{\cong} \dagger A, & \text{diagram}_2: A^\dagger &\xrightarrow{\cong} A_{\gamma_A^{-1}} \end{aligned} \quad (4.3)$$

are bimodule maps. Since $\dagger A = A^\dagger$ by assumption, it follows that

$$\gamma_A: A_{\gamma_A^{-1}} \xrightarrow{\cong} \gamma_A A \quad (4.4)$$

in \mathcal{B}_{eq} . Moreover, the special case $X = A = B$ in (4.2) reads

$${}^*A = \gamma_A^{-1}(\dagger A) \cong \gamma_A^{-1}(\gamma_A A) \cong A \quad \text{and} \quad A^* = (A^\dagger)_{\gamma_A} \cong (A_{\gamma_A^{-1}})_{\gamma_A} \cong A$$

in \mathcal{B}_{eq} , which is consistent with $1_{(\alpha, A)} = A$.


4.1.3. Symmetric monoidal structure. As explained in [42, Corollary 6.12], the equivariant completion \mathcal{B}_{eq} is the horizontal 2-category of a symmetric monoidal double category \mathbb{B}_{eq} , which satisfies the conditions under which the symmetric monoidal structure of \mathbb{B}_{eq} is passed on to \mathcal{B}_{eq} .

Proposition 4.4. *\mathcal{B}_{eq} has a symmetric monoidal structure induced from \mathcal{B} .*


Here, we collect the ingredients of the symmetric monoidal structure on \mathcal{B}_{eq} in graphical presentation, using the conventions of Section 3.1. The monoidal product on objects $(\alpha, A), (\alpha', A') \in \mathcal{B}_{\text{eq}}$ is given by

$$(\alpha, A) \square^{\mathcal{B}_{\text{eq}}} (\alpha', A') = (\alpha \square^{\mathcal{B}} \alpha', A \square^{\mathcal{B}} A'),$$

where the Δ -separable Frobenius structure on $A \square A' \equiv A \square^{\mathcal{B}} A'$ is as follows:



It follows that the Nakayama automorphism of $A \square A'$ factorises with respect to \square ,



On 1-morphisms $X \in \mathcal{B}_{\text{eq}}((\alpha, A), (\beta, B))$ and $X' \in \mathcal{B}_{\text{eq}}((\alpha', A'), (\beta', B'))$, the monoidal product is $X \square^{\mathcal{B}} X'$ with the left $(A \square A')$ - and right $(B \square B')$ -action induced from \mathcal{B} . On 2-morphisms, we have


$$\square^{\mathcal{B}_{\text{eq}}} = \square^{\mathcal{B}}.$$

From now on, we will denote the monoidal product of both \mathcal{B} and \mathcal{B}_{eq} simply by \square .


The braiding in \mathcal{B}_{eq} has 1-morphism components

$$b_{(\alpha, A), (\alpha', A')}^{\mathcal{B}_{\text{eq}}} = (A' \square A) \otimes b_{\alpha, \alpha'}^{\mathcal{B}} \cong b_{\alpha, \alpha'}^{\mathcal{B}} \otimes (A \square A')$$

for $(\alpha, A), (\alpha', A') \in \mathcal{B}_{\text{eq}}$, with left $(A' \square A)$ -action given by $\mu_{A' \square A}$ and right $(A \square A')$ -action given by



The 2-morphism components of $b^{\mathcal{B}_{\text{eq}}}$ are




for all $X \in \mathcal{B}_{\text{eq}}((\alpha, A), (\beta, B))$ and $X' \in \mathcal{B}_{\text{eq}}((\alpha', A'), (\beta', B'))$.

Remark 4.5. It follows that $\mathcal{B} \subset \mathcal{B}_{\text{eq}}$ is an embedding of symmetric monoidal 2-categories. In particular, the braiding components of 1-morphisms in $\mathcal{B}_{\text{eq}}((1, 1_1), (1, 1_1))$ are those in $\mathcal{B}(1, 1)$.

4.1.4. Duality and Serre automorphism. If $\alpha \in \mathcal{B}$ is dualisable with duality data $(\alpha^\#, \tilde{e}v_\alpha, \widetilde{\text{co}e}v_\alpha)$, then every object $(\alpha, A) \in \mathcal{B}_{\text{eq}}$ is dualisable with

$$\begin{aligned} (\alpha, A)^\# &= (\alpha^\#, A^\#), \\ \tilde{e}v_{(\alpha, A)} &= \tilde{e}v_\alpha \otimes (A \square 1_{\alpha^\#}) \cong \tilde{e}v_\alpha \otimes (1_\alpha \square A^\#), \\ \widetilde{\text{co}e}v_{(\alpha, A)} &= (1_{\alpha^\#} \square A) \otimes \widetilde{\text{co}e}v_\alpha \cong (A^\# \square 1_\alpha) \otimes \widetilde{\text{co}e}v_\alpha, \end{aligned}$$

where we used the isomorphism



induced by the inverse cusp isomorphism c_1^{-1} in (3.2), and similarly with $\widetilde{\text{co}e}v_{(\alpha, A)}$ and c_r^{-1} . These isomorphisms together with μ_A also give the above adjunction morphisms their bimodule structures.

The Frobenius algebra structure $(A^\#, \mu_{A^\#}, \eta_{A^\#}, \Delta_{A^\#}, \varepsilon_{A^\#})$ on $A^\#$ is by definition $(A^\#, \mu_A^\#, \eta_A^\#, \Delta_A^\#, \varepsilon_A^\#)$ up to cusp isomorphisms as needed. We illustrate this with the multiplication

$$\mu_{A^\#} = \begin{array}{c} \alpha^\# \\ \bullet \\ c_r \\ \text{---} \\ \mu_A \\ \text{---} \\ \bullet \\ c_r^{-1} \\ \text{---} \\ A^\# \end{array} \equiv \begin{array}{c} | \\ \text{---} \\ | \end{array} \quad (4.5)$$

where the last expression is shorthand for the defining Gray diagram in the middle. Note that there is a canonical isomorphism $A \cong A^{\#\#}$ in \mathcal{B} (see [6, Figure 32]), which we leave implicit. Similarly, we denote the other structure maps of $A^{\#}$ as

$$\eta_{A^{\#}} = \downarrow \circlearrowleft, \quad \Delta_{A^{\#}} = \begin{array}{c} \diagup \quad \diagdown \\ \circlearrowright \\ \downarrow \end{array}, \quad \varepsilon_{A^{\#}} = \downarrow \circlearrowright. \quad (4.6)$$

The enveloping algebra of A is

$$A^e = A \square A^{\#}.$$

The cusp isomorphisms together with μ_A give a canonical right A^e -module structure on $\widetilde{\text{ev}}_{(\alpha, A)}$ and a left $(A^e)^{\#}$ -module structure on $\widetilde{\text{coev}}_{(\alpha, A)}$.

Lemma 4.6. *If $\alpha \in \mathcal{B}$ is dualisable, then for $(\alpha, A) \in \mathcal{B}_{\text{eq}}$ we have that*

$$\gamma_{A^{\#}} = (\gamma_A^{\#})^{-1}$$

up to cusp isomorphisms in \mathcal{B} .

Proof. Up to cusp isomorphisms, $\widetilde{\text{ev}}_{A^{\#}}: A^{\#} \otimes (A^{\#})^{\dagger} \rightarrow 1_{\alpha^{\#}}$ agrees with $\text{ev}_A^{\#}$,

$$\text{ev}_A^{\#} = \begin{array}{c} \circlearrowleft \\ \downarrow \quad \downarrow \\ A^{\#} \quad \dagger A^{\#} \end{array} \equiv \begin{array}{c} \circlearrowleft \\ \downarrow \quad \downarrow \\ A^{\#} \quad (A^{\#})^{\dagger} \end{array} = \widetilde{\text{ev}}_{A^{\#}},$$

and similarly for the coevaluations,

$$\widetilde{\text{coev}}_{A^{\#}} = \begin{array}{c} A^{\#} \quad (A^{\dagger})^{\#} \\ \circlearrowright \\ \downarrow \quad \downarrow \\ A^{\#} \quad \dagger A^{\#} \end{array} \equiv \begin{array}{c} A^{\#} \quad \dagger A^{\#} \\ \circlearrowright \\ \downarrow \quad \downarrow \\ A^{\#} \quad \dagger A^{\#} \end{array} = \text{coev}_{A^{\#}}.$$

Hence, together with (4.5) and (4.6), we find

$$\gamma_{A^{\#}}^{-1} = \begin{array}{c} \downarrow \quad \downarrow \\ \circlearrowleft \quad \circlearrowright \\ \downarrow \quad \downarrow \\ \downarrow \quad \downarrow \\ \circlearrowright \quad \circlearrowleft \\ \downarrow \quad \downarrow \\ \downarrow \quad \downarrow \end{array} = \begin{array}{c} \downarrow \quad \downarrow \\ \downarrow \quad \downarrow \\ \downarrow \quad \downarrow \\ \downarrow \quad \downarrow \\ \downarrow \quad \downarrow \\ \downarrow \quad \downarrow \end{array} = \gamma_{A^{\#}}^{\#}. \quad \blacksquare$$

We now turn to full dualisability, which is another property that is compatible with equivariant completion.

Proposition 4.7. *Let $\alpha \in \mathcal{B}$ be fully dualisable. Then, every $(\alpha, A) \in \mathcal{B}_{\text{eq}}$ is fully dualisable.*

Proof. If $\tilde{e}v_\alpha, \widetilde{\text{coev}}_\alpha$ witness $\alpha^\#$ as a dual of α in \mathcal{B} , then

$$\tilde{e}v_{(\alpha, A)} = \tilde{e}v_\alpha \otimes (A \square 1_{\alpha^\#}), \quad \widetilde{\text{coev}}_{(\alpha, A)} = (1_{\alpha^\#} \square A) \otimes \widetilde{\text{coev}}_\alpha$$


witness $(\alpha^\#, A^\#)$ as a dual of (α, A) in \mathcal{B}_{eq} . Moreover, by Proposition 4.3, these adjunction 1-morphisms have adjoints themselves since $A \in \mathcal{B}(\alpha, \alpha)$ has adjoints thanks to its Frobenius algebra structure, while $\tilde{e}v_\alpha, \widetilde{\text{coev}}_\alpha$ have adjoints by assumption. \blacksquare

In particular, according to (4.2), we have

$$\tilde{e}v_{(\alpha, A)}^\star = (\tilde{e}v_{(\alpha, A)}^\dagger)_{\gamma_{1_1}} = [\tilde{e}v_\alpha \otimes (A \square 1_{\alpha^\#})]^\dagger \cong (A^\dagger \square 1_{\alpha^\#}) \otimes \tilde{e}v_\alpha^\dagger.$$

Hence, the Serre automorphism of $(\alpha, A) \in \mathcal{B}_{\text{eq}}$ is

$$S_{(\alpha, A)} = (A \square \tilde{e}v_{(\alpha, A)}) \otimes (b_{(\alpha, A), (\alpha, A)} \square A^\#) \otimes (A \square \tilde{e}v_{(\alpha, A)}^\star)$$



Applying the 2-isomorphisms $b_{A, A}$ and $b_{A^\dagger, 1_\alpha}^{-1}$, the inner A -line can be moved to the right and the A^\dagger -line can be moved to the left, respectively. Alternatively, the A^\dagger -line can be moved to the left of the $b_{\alpha, \alpha}$ -line with the help of two cusp isomorphisms and then to the right by $b_{A^\dagger, 1_\alpha}$. Thus, we have shown the following proposition.

Proposition 4.8. *Let $\alpha \in \mathcal{B}^{\text{fd}}$. Then, $(\alpha, A) \in \mathcal{B}_{\text{eq}}^{\text{fd}}$ and*

$$S_{(\alpha, A)} \cong A^\dagger \otimes S_\alpha \otimes A \cong A \otimes S_\alpha \otimes A^\dagger. \quad (4.7)$$

Below we will frequently not display $A = 1_{(\alpha, A)}$ and simply write

$$S_{(\alpha, A)} \cong A^\dagger \otimes S_\alpha \cong S_\alpha \otimes A^\dagger.$$

Corollary 4.9. *Let $r \in \mathbb{Z}_{\geq 1}$, $\alpha \in \mathcal{B}^{\text{fd}}$, and $(\alpha, A) \in \mathcal{B}_{\text{eq}}$ such that $S_\alpha^r \cong 1_\alpha$ and $\gamma_A^r = 1_A$ in \mathcal{B} . Then, there is an r -spin TQFT*

$$\begin{aligned} \mathcal{Z}: \text{Bord}_{2,1,0}^{r\text{-spin}} &\longrightarrow \mathcal{B}_{\text{eq}} \\ + &\longmapsto (\alpha, A). \end{aligned}$$

Proof. Combining (4.7) with $A^\dagger \cong A_{\gamma_A^{-1}}$, Lemma 3.18 and Theorem 3.19, we see that any choice of isomorphism $S_\alpha^r \cong 1_\alpha$ determines a Spin_2^r -homotopy fixed point in $(\mathcal{B}^{\text{fd}})^\times$. ■

4.1.5. A Frobenius algebra. For $(\alpha, A) \in \mathcal{B}_{\text{eq}}^{\text{fd}}$ and $a \in \mathbb{Z}$, the a -th circle space (recall (3.8)) is

$$C_a^{(\alpha, A)} \cong \tilde{\text{eV}}_{(\alpha, A)} \otimes_{A^e} (S_{(\alpha, A)}^{1-a} \square 1_{(\alpha, A)^\#}) \otimes_{A^e} {}^* \tilde{\text{eV}}_{(\alpha, A)}. \quad (4.8)$$

By Proposition 4.3, we have

$${}^* \tilde{\text{eV}}_{(\alpha, A)} \cong \gamma_{A^e}^{-1}(\dagger A \square 1_{\alpha^\#}) \otimes \dagger \tilde{\text{eV}}_\alpha \cong (1_\alpha \square \gamma_A^{-1}(\dagger A^\#)_{\gamma_A}) \otimes \tilde{\text{eV}}_\alpha,$$

and by Proposition 4.8 together with (4.3) and $\dagger A = A^\dagger$, we have

$$S_{(\alpha, A)}^{1-a} \equiv S_{(\alpha, A)}^{\otimes A(1-a)} \cong \gamma_A^{1-a} A \otimes S_\alpha^{1-a} \otimes A. \quad (4.9)$$

Our next goal is to explicitly describe the closed Λ_0 -Frobenius structure on the circle spaces $C_a^{(\alpha, A)}$ in \mathcal{B}_{eq} . This means that we will determine the (co)multiplication and (co)unit of (3.9)–(3.12) of $C_a^{(\alpha, A)}$ directly in terms of data in \mathcal{B} . In doing so, we will frequently use the fact that the 2-morphism

$$S_{(\alpha, A)}^{x+y} \cong \begin{array}{c} \text{Diagram} \end{array}, \quad x, y \in \mathbb{Z}, \quad (4.10)$$

The diagram shows a 2-morphism between two Frobenius structures. On the left, the Frobenius structure is $S_{(\alpha, A)}^x \otimes S_{(\alpha, A)}^y$. It consists of a green vertical line labeled $\gamma_A^x A$ on the left, a blue vertical line labeled S_α^x in the middle, and a green vertical line labeled $\gamma_A^y A$ on the right. A green arc connects the top of the left line to the top of the right line, passing over the blue line. A blue arc connects the top of the middle line to the top of the right line, passing under the green arc. On the right, the Frobenius structure is $S_{(\alpha, A)}^{x+y}$. It consists of a single green vertical line labeled $\gamma_A^{x+y} A$ on the left, a blue vertical line labeled S_α^{x+y} in the middle, and a green vertical line labeled A on the right. A green arc connects the top of the left line to the top of the right line, passing over the blue line. A blue arc connects the top of the middle line to the top of the right line, passing under the green arc. The two structures are connected by a 2-morphism represented by a green arc on the left and a blue arc on the right.

in \mathcal{B} induces the identity

$$S_{(\alpha,A)}^x \otimes_A S_{(\alpha,A)}^y \rightarrow S_{(\alpha,A)}^{x+y},$$

up to the isomorphism in (4.9). From now on, we will no longer display all A -lines in diagrams that represent 2-morphisms in \mathcal{B}_{eq} , since

$$A = 1_{(\alpha,A)}.$$

Accordingly, we abbreviate (4.10) and its inverse as

$$\begin{array}{c}
 S_{(\alpha,A)}^{x+y} \\
 | \\
 \text{[Green Square]} \\
 \swarrow \searrow \\
 S_{(\alpha,A)}^x \quad S_{(\alpha,A)}^y
 \end{array}
 =
 \begin{array}{c}
 S_{(\alpha,A)}^{x+y} \\
 | \\
 \text{[Green Square]} \\
 \text{[Blue Arc } \gamma_A^y \text{]} \\
 | \quad | \\
 S_{(\alpha,A)}^x \quad S_{(\alpha,A)}^y
 \end{array},$$

$$\begin{array}{c}
 S_{(\alpha,A)}^x \quad S_{(\alpha,A)}^y \\
 \swarrow \searrow \\
 \text{[Green Square]} \\
 | \\
 S_{(\alpha,A)}^{x+y}
 \end{array}
 =
 \begin{array}{c}
 S_{(\alpha,A)}^x \quad S_{(\alpha,A)}^y \\
 | \quad | \\
 \text{[Green Square]} \quad \text{[Green Square]} \\
 \text{[Blue Arc } \gamma_A^{-y} \text{]} \\
 | \\
 S_{(\alpha,A)}^{x+y}
 \end{array}.
 \tag{4.11}$$

With the above preparations, we can present the isomorphism (3.13) in the case of the equivariant completion.

Lemma 4.10. *Let $\alpha \in \mathcal{B}^{\text{fd}}$. Then, for any $(\alpha, A) \in \mathcal{B}_{\text{eq}}$, there are mutually inverse isomorphisms*

$$(1_{(\alpha,A)} \square S_{(\alpha,A)\#}^2) \otimes_{A^e} {}^* \tilde{\text{eV}}_{(\alpha,A)} \xrightleftharpoons[f'_A]{f_A} \tilde{\text{eV}}_{(\alpha,A)}^*$$


given by

$$\begin{aligned}
 f_A = & \left(\begin{array}{c} \alpha \\ \alpha^\# \\ (A^\#)^\dagger \\ S_{\alpha^\#}^2 \\ S_{(\alpha, A)^\#}^2 \\ \gamma_A^{-1}(\dagger A^\#)\gamma_A \\ \tilde{e}v_\alpha^\dagger \\ \dagger \tilde{e}v_\alpha \end{array} \right), \\
 f'_A = & \left(\begin{array}{c} \alpha \\ \alpha^\# \\ S_{(\alpha, A)^\#}^2 \\ (A^\dagger)^\# \\ S_{\alpha^\#}^{-2} \\ \dagger A^\# \\ \tilde{e}v_\alpha \\ \tilde{e}v_\alpha^\dagger \end{array} \right). \tag{4.12}
 \end{aligned}$$

Proof. Repeated use of (4.3) together with standard manipulations of string diagrams for Δ -separable Frobenius algebras shows that f_A, f'_A are indeed bimodule maps, and that $f_A \circ f'_A = 1_{\dagger A^\#}$. Since their source and target are isomorphic, it follows that f_A and f'_A indeed represent mutually inverse 2-morphisms in \mathcal{B}_{eq} . ■

We have now expressed all the ingredients of the closed Λ_0 -Frobenius structure on $\{C_a^{(\alpha, A)}\}$ in \mathcal{B}_{eq} directly in terms of data in \mathcal{B} . With this, the Nakayama automorphisms $N_a^{(\alpha, A)}$ can be computed.


Proposition 4.11. *Let $\alpha \in \mathcal{B}^{\text{fd}}$. Then, for any $(\alpha, A) \in \mathcal{B}_{\text{eq}}$, we have*




Proof. Our task is to compute $N_a^{(\alpha, A)}$ as defined in (2.4) in \mathcal{B}_{eq} . We will first compute

$$\varepsilon_{-1} \circ \mu_{a, -a} \quad \text{and} \quad \Delta_{a, -a} \circ \eta_1$$


in \mathcal{B}_{eq} , starting with $\varepsilon_{-1} \circ \mu_{a, -a}$. Using the notation introduced in (4.11), we have



and inserting the expression for f_A in (4.12) into (3.10), we have



In the composition $\varepsilon_{-1} \circ \mu_{a, -a}$, we first use



where here and below we suppress S_α -strands. This expression cancels with another subdiagram of $\varepsilon_{-1} \circ \mu_{a,-a}$, leaving

$$\begin{aligned}
 \varepsilon_{-1} \circ \mu_{a,-a} &\equiv \text{Diagram 1} \\
 &= \text{Diagram 2} \quad . \quad (4.13)
 \end{aligned}$$

The second step uses the properties of Δ -separable Frobenius algebras and (4.1); moreover, here and below we suppress \tilde{e}_α and its adjoints (as they are only spectators)

in our string diagram manipulations). Analogously, we arrive at

$$\Delta_{a,-a} \circ \eta_1 \equiv \begin{array}{c} A^\# \quad S_{(\alpha,A)^\#}^{-a+1} (A^\#)^\dagger \quad A^\# \quad S_{(\alpha,A)^\#}^{+a+1} \dagger A^\# \\ \begin{array}{c} \text{Diagram with four vertical lines and a central loop structure. The left two lines are labeled } A^\# \text{ and } S_{(\alpha,A)^\#}^{-a+1} (A^\#)^\dagger. \text{ The right two lines are labeled } A^\# \text{ and } S_{(\alpha,A)^\#}^{+a+1} \dagger A^\#. \text{ A central loop structure connects the lines, with vertices labeled } \gamma_{A^\#}^{1-a} \text{ and } \gamma_{A^\#}. \end{array} \\ \end{array} \quad (4.14)$$


Combining (4.13) with (4.14) into (2.4), we employ the relation

$$\begin{array}{c} \text{Diagram 1: Two vertical lines with a loop between them. A dot labeled } \varphi \text{ is on the right line.} \\ = \\ \text{Diagram 2: Two vertical lines with a loop between them. A dot labeled } \varphi^\dagger \text{ is on the left line.} \end{array}$$

to see that $N_a^{(\alpha,A)}$ is given by pre-composing the twist

$$\begin{array}{c} C_a^{(\alpha,A)} \\ \text{Diagram: A vertical line with a loop on its right side.} \\ C_a^{(\alpha,A)} \end{array} \quad (4.15)$$

with (where we continue to suppress $\tilde{e}\nu_\alpha$)



Repeatedly using the defining properties of Δ -separable Frobenius algebras as well the properties of the Nakayama automorphism $\gamma_{A^\#}$ collected in Section 4.1.2, a straightforward but lengthy computation shows that the above string diagram is equal to

(4.16)

Putting $\tilde{e}v_\alpha$, $\tilde{e}v_\alpha^\dagger$, S_α^{1-a} back in, a final application of the isomorphisms (4.4) and (4.9) allows us to identify (4.16) with

Post-composing with (4.15) thus completes the proof. ■

Combining Proposition 4.11 with the isomorphisms (4.7) and (4.9), we obtain closed Λ_r -Frobenius algebras from Δ -separable Frobenius algebras A on fully dualisable objects if the r th power of the Nakayama automorphism of A is the identity.

Corollary 4.12. *If for $r \in \mathbb{Z}_{\geq 1}$ there is an isomorphism $S_\alpha^r \cong 1_\alpha$ in \mathcal{B}^{fd} and for $(\alpha, A) \in \mathcal{B}_{\text{eq}}$ we have*

$$\gamma_A^r = 1_A,$$

then there is an induced closed Λ_r -Frobenius algebra structure on

$$\{C_a^{(\alpha, A)}\}_{a \in \{0, 1, \dots, r-1\}}.$$

If \mathcal{B} has internal Homs, as is the case in the examples related to TQFTs of state sum, sigma model, and Landau–Ginzburg type, then the computation of both $C_a^{(\alpha, A)}$ and $N_a^{(\alpha, A)}$ can be simplified. For ease of presentation, we further assume that

$$S_\alpha \cong 1_\alpha;$$

in Section 4.2 below, we will see how this restriction can be lifted in practice.

Lemma 4.13. *Let $\alpha \in \mathcal{B}^{\text{fd}}$ with $S_\alpha \cong 1_\alpha$, and assume that for $(\alpha, A) \in \mathcal{B}_{\text{eq}}$, we have*

$$C_a^{(\alpha, A)} \cong \mathcal{B}(1_1, C_a^{(\alpha, A)}).$$

Then

$$C_a^{(\alpha, A)} \cong \left\{ \varphi \in \mathcal{B}(1_\alpha, A) \mid \gamma_A^{1-a} \begin{array}{c} \circ \\ \uparrow \\ \circ \\ \downarrow \\ \circ \\ \downarrow \\ \circ \end{array} \varphi = \downarrow \varphi \right\}, \quad (4.17)$$


and $N_a^{(\alpha, A)}$ corresponds to post-composition with γ_A .

Note that the above result further elucidates the relation between the two different notions of “Nakayama morphism” N and γ .

Proof of Lemma 4.13. We have

$$\begin{aligned} C_a^{(\alpha, A)} &\cong \mathcal{B}_{\text{eq}}(1_{(1, 1_1)}, C_a^{(\alpha, A)}) \\ &\cong \mathcal{B}_{\text{eq}}(\tilde{\mathbf{e}}\tilde{\mathbf{v}}_{(\alpha, A)}, \tilde{\mathbf{e}}\tilde{\mathbf{v}}_{(\alpha, A)} \otimes_{A^e} (S_{(\alpha, A)}^{1-a} \square 1_{\alpha^\#})) \\ &\cong \mathcal{B}_{\text{eq}}(A, S_{(\alpha, A)}^{1-a}) \\ &\cong \mathcal{B}_{\text{eq}}(A, \gamma_A^{1-a} A) \\ &\cong \left\{ \varphi \in \mathcal{B}(1_\alpha, A) \mid \gamma_A^{1-a} \begin{array}{c} \circ \\ \uparrow \\ \circ \\ \downarrow \\ \circ \end{array} \varphi = \downarrow \varphi \right\}. \end{aligned}$$

In the second step, we used (4.8) and adjunction for $\tilde{e}_{\nu(\alpha,A)}$; the third step is the isomorphism



the fourth step is (4.9) together with the assumption $S_\alpha \cong 1_\alpha$; the fifth step is a standard computation with Δ -separable Frobenius algebras along the lines of [9, Section 3.2]. ■

4.2. Landau–Ginzburg models

In this section, we briefly review the 2-category of Landau–Ginzburg models \mathcal{LG} and note that every object in \mathcal{LG} gives rise to an extended 2-spin TQFT. Then, we apply the results of Section 4.1 to a closely related 2-category $\mathcal{LG}^{\bullet/2}$ and consider the simplest non-trivial example.

Recall from [15] that, for every fixed field \mathbb{k} , there is a 2-category \mathcal{LG} whose objects are pairs $(\mathbb{k}[x_1, \dots, x_n], W)$, where $n \in \mathbb{Z}_{\geq 0}$ and $W = 0 \in \mathbb{k}$ if $n = 0$, while for $n > 0$, $W \in \mathbb{k}[x] \equiv \mathbb{k}[x_1, \dots, x_n]$ is such that the *Jacobi algebra*

$$\text{Jac}_W = \mathbb{k}[x_1, \dots, x_n] / (\partial_{x_1} W, \dots, \partial_{x_n} W)$$

is finite-dimensional over \mathbb{k} . We refer to such polynomials W as *potentials*. The Hom categories of \mathcal{LG} are idempotent completions of homotopy categories of finite-rank matrix factorisations. Hence, up to technicalities with idempotents (which will not be relevant to our discussions below), a 1-morphism $(\mathbb{k}[x], W) \rightarrow (\mathbb{k}[z], V)$ is a free \mathbb{Z}_2 -graded $\mathbb{k}[x, z]$ -module $X = X^0 \oplus X^1$ together with an odd $\mathbb{k}[x, z]$ -linear endomorphism $d_X: X \rightarrow X$ such that $d_X^2 = (V - W) \cdot 1_X$. The Hom sets of 2-morphisms $(X, d_X) \rightarrow (X', d_{X'})$ consist of the even cohomology classes of the differential defined on \mathbb{Z}_2 -homogeneous maps as

$$\begin{aligned} \delta_{X, X'}: \text{Hom}_{\mathbb{k}[x, z]}(X, X') &\longrightarrow \text{Hom}_{\mathbb{k}[x, z]}(X, X') \\ \zeta &\longmapsto d_{X'} \circ \zeta - (-1)^{|\zeta|} \zeta \circ d_X \end{aligned} \quad (4.18)$$

and extended linearly to all of $\text{Hom}_{\mathbb{k}[x, z]}(X, X')$.

Given $(X, d_X): (\mathbb{k}[x], W_1) \rightarrow (\mathbb{k}[y], W_2)$ and $(Y, d_Y): (\mathbb{k}[y], W_2) \rightarrow (\mathbb{k}[z], W_3)$, their horizontal composition is $(Y \otimes_{\mathbb{k}[y]} X, d_Y \otimes 1_X + 1 \otimes d_X)$, and the unit 1-morphism of $(\mathbb{k}[x_1, \dots, x_n], W)$ is $1_W = (I_W, d_{I_W})$ with

$$I_W = \bigwedge \left(\bigoplus_{i=1}^n \mathbb{k}[x, x'] \cdot \theta_i \right), \quad d_{I_W} = \sum_{i=1}^n (\partial_{[i]}^{x', x} W \cdot \theta_i + (x'_i - x_i) \cdot \theta_i^*), \quad (4.19)$$

where $\{\theta_i\}$ is a chosen $\mathbb{k}[x', x]$ -basis of $\mathbb{k}[x, x']^{\oplus n}$, and

$$\partial_{[i]}^{x', x} W = \frac{W(x_1, \dots, x_{i-1}, x'_i, \dots, x'_n) - W(x_1, \dots, x_i, x'_{i+1}, \dots, x'_n)}{x'_i - x_i}.$$

A straightforward computation shows that $\text{End}(1_W) \cong \text{Jac}_W$ in $\mathcal{L}\mathcal{G}$.

Every 1-morphism $X \equiv (X, d_X) \in \mathcal{L}\mathcal{G}((\mathbb{k}[x_1, \dots, x_n], W), (\mathbb{k}[z_1, \dots, z_m], V))$ has a left adjoint ${}^\dagger X$ and a right adjoint X^\dagger , and ${}^\dagger X \cong X^\dagger$ iff $m = n \bmod 2$. The associated adjunction 2-morphisms are explicitly known; see [15, Theorem 6.11], or [14] for a concise review.

The 2-category $\mathcal{L}\mathcal{G}$ has a natural monoidal structure, with the monoidal product on objects given by $(\mathbb{k}[x], W) \square (\mathbb{k}[z], V) = (\mathbb{k}[x, z], W + V)$, while on 1- and 2-morphisms, it is basically $\otimes_{\mathbb{k}}$; see [13, Section 2.2]. Hence, $1 := (\mathbb{k}, 0)$ is the unit object. Every $(\mathbb{k}[x], W) \in \mathcal{L}\mathcal{G}$ has a (left and right) dual $(\mathbb{k}[x], W)^\# = (\mathbb{k}[x], -W)$ whose associated adjunction 1-morphisms have 1_W as their underlying matrix factorisation; see [13, Proposition 2.6]. Thus, as every 1-morphism in $\mathcal{L}\mathcal{G}$ has an adjoint, every object in $\mathcal{L}\mathcal{G}$ is fully dualisable.

The monoidal 2-category $\mathcal{L}\mathcal{G}$ has a symmetric braiding, whose 1-morphism components $b_{V, W}$ are given by 1_{V+W} (up to a reordering of variables), while the 2-morphism components are compositions of canonical module isomorphisms and structure maps of the underlying 2-category $\mathcal{L}\mathcal{G}$. For details, we refer to [13, Section 2.3]. In summary, we have the following theorem.

Theorem 4.14 ([13, 15]). *For every field \mathbb{k} , the 2-category of Landau–Ginzburg models $\mathcal{L}\mathcal{G}$ has a symmetric monoidal structure such that $\mathcal{L}\mathcal{G} = \mathcal{L}\mathcal{G}^{\text{fd}}$.*

Remark 4.15. A variant of $\mathcal{L}\mathcal{G}$ is the symmetric monoidal 2-category $\mathcal{L}\mathcal{G}^{\bullet/2}$, which is defined analogously to $\mathcal{L}\mathcal{G}$, but 2-morphisms are given by both even and odd cohomologies of the differentials $\delta_{X, X'}$ in (4.18), but with classes $-\zeta$ and $+\zeta$ identified. This ad hoc \mathbb{Z}_2 -quotient allows to stay within the realm of 2-categories (as opposed to super 2-categories) while allowing odd 2-morphisms; compare [28] and [13, Remark 3.11 (ii)].

Theorem 4.14 also holds for $\mathcal{L}\mathcal{G}^{\bullet/2}$, i.e.,

$$\mathcal{L}\mathcal{G}^{\bullet/2} = (\mathcal{L}\mathcal{G}^{\bullet/2})^{\text{fd}}.$$

The Serre automorphism of $W \equiv (\mathbb{k}[x_1, \dots, x_n], W)$ was computed in [13, Lemma 3.8] to be

$$S_W \cong 1_W[n], \quad (4.20)$$

where $[n]$ denotes the n -fold application of the shift functor [1], which sends a matrix factorisation $(X^0 \oplus X^1, d_X)$ to $(X^1 \oplus X^0, -d_X)$. It follows that [2] is the identity functor, and one finds that $\text{Hom}(1_W, 1_W[n]) \cong \delta_{n,0 \bmod 2} \cdot \text{Jac}_W$ in $\mathcal{L}\mathcal{G}$, as $\mathcal{L}\mathcal{G}$ has only even cohomology classes as 2-morphisms, while $\text{Hom}(1_W, 1_W[n]) \cong \text{Jac}_W[n]$ is purely odd in $\mathcal{L}\mathcal{G}^{\bullet/2}$ if n is odd. As a consequence, $(\mathbb{k}[x_1, \dots, x_n], W)$ determines an extended oriented TQFT with values in $\mathcal{L}\mathcal{G}$ iff n is even, and it determines an extended oriented TQFT with values in $\mathcal{L}\mathcal{G}^{\bullet/2}$ for every value of n .

Remark 4.16. As shown in [13, Section 3], fully extended oriented TQFTs with values in $\mathcal{L}\mathcal{G}$ are indeed extensions of closed Landau–Ginzburg models to the point: Applying the cobordism hypothesis of [25, 38] to the duality data of an object $(\mathbb{k}[x], W)$ in $\mathcal{L}\mathcal{G}$ or in $\mathcal{L}\mathcal{G}^{\bullet/2}$, one recovers the (non-semisimple) commutative Frobenius algebra Jac_W from the circle, the pair of pants, and the disc.

As an immediate consequence of Theorem 3.19, Lemma 3.18, (4.20), and the isomorphism $\text{Aut}(1_W) \cong \mathbb{k}$, we find that every potential depending on an odd number of variables gives rise to a proper extended spin TQFT.

Theorem 4.17. *Every object $W \equiv (\mathbb{k}[x_1, \dots, x_n], W) \in \mathcal{L}\mathcal{G}$ gives rise to a unique-up-to-isomorphism extended 2-spin TQFT valued in $\mathcal{L}\mathcal{G}$. These TQFTs factor through the oriented bordism 2-category iff n is even.*

It is straightforward to compute that

$$C_a^W \equiv C_a^{(\mathbb{k}[x_1, \dots, x_n], W)} \cong \text{Jac}_W[n \cdot (1 - a)] \quad \text{in } \mathcal{L}\mathcal{G}(1, 1) \cong \text{vect}_{\mathbb{k}}^{\mathbb{Z}_2}$$

for $a \in \{0, 1\}$. Hence, these circle spaces are the zeroth Hochschild homology and cohomology, respectively, of the differential graded category of matrix factorisations, first computed in [18]. Moreover, for the Nakayama automorphisms, we have

$$N_a^W = (-1)^{n \cdot (1-a)} \cdot 1_{C_a^W}.$$

We now turn to the equivariant completion of $\mathcal{L}\mathcal{G}^{\bullet/2}$ to look for r -spin TQFTs that can detect more r -spin structures than oriented TQFTs. One type of example+ of Δ -separable Frobenius algebras with trivialisable r th power of its Serre automorphism is the algebra A_G mentioned in Section 4.1.1, in the case $G = \mathbb{Z}_r$.

Recall from [16, Section 7.1] that if a finite group G acts on $\mathbb{k}[x_1, \dots, x_n]$ such that a given $W \in \mathbb{k}[x_1, \dots, x_n]$ is invariant, this induces a Δ -separable Frobenius structure on

$$A_G := \bigoplus_{g \in G} g(1_W),$$

where the g -twisted matrix factorisation $g(1_W)$ is obtained from (4.19) by replacing $x'_i \mapsto g^{-1}(x'_i)$. Its Nakayama automorphism is

$$\gamma_{A_G} = \sum_{g \in G} \det(g)^{-1} \cdot 1_{g(1_W)}, \quad (4.21)$$

where $\det(g)$ is the determinant of the g -action on x_1, \dots, x_n ; cf. [9, Section 3.1].

Example 4.18. For $r \in \mathbb{Z}_{\geq 3}$, we consider $(\mathbb{k}[x], x^r) \in \mathcal{L}\mathcal{G}^{\bullet/2}$ with the \mathbb{Z}_r -action $\mathbb{Z}_r \rightarrow \text{Aut}_{\mathbb{k}}(\mathbb{k}[x])$, $1 \mapsto (x \mapsto \xi \cdot x)$, where

$$\xi := e^{2\pi i/r}.$$

Hence, $W := x^r$ is invariant, and we have

$$g(1_W) = (\mathbb{k}[x] \cdot 1) \oplus (\mathbb{k}[x] \cdot \theta)$$

with

$$d_{g(1_W)} = \frac{x'^r - x^r}{\xi^{-g}x' - x} \cdot \theta + (\xi^{-g}x' - x) \cdot \theta^*$$

for $g \in \{0, 1, \dots, r-1\}$. Setting

$$A_{\mathbb{Z}_r} = \bigoplus_{g \in G} g(1_W),$$

we have an object

$$((\mathbb{k}[x], x^r), A_{\mathbb{Z}_r}) \in (\mathcal{L}\mathcal{G}^{\bullet/2})_{\text{eq}}.$$

For $g = 0$, we have

$$\text{Hom}(1_W, 1_W) \cong \mathbb{k} \cdot \{1, x, \dots, x^{r-2}\}$$

as a purely even \mathbb{Z}_2 -graded vector space, while for $g \neq 0$ one finds that [9, Appendix 2]

$$\text{Hom}(1_W, g(1_W)) \cong \mathbb{k} \cdot \left(\frac{x'^r - x^r}{(x' - x)(\xi^{-g}x' - x)} \cdot \theta + \theta^* \right) [1]$$

is a purely odd, 1-dimensional \mathbb{Z}_2 -graded vector space. Moreover, by (4.20) we have $S_W \cong 1_W$ in $\mathcal{L}\mathcal{G}^{\bullet/2}$, and according to (4.21), the Nakayama automorphism of $A_{\mathbb{Z}_r}$ is

$$\gamma_{A_{\mathbb{Z}_r}} = \sum_{g=0}^{r-1} \xi^{-g} \cdot 1_{g(1_W)}. \quad (4.22)$$

We will use Lemma 4.13 to compute the circle spaces

$$C_a \equiv C_a^{((\mathbb{k}[x], x^r), A_{\mathbb{Z}_r})}.$$

Hence, we have to identify the image of the projector

$$\mathrm{Hom}(1_W, A_{\mathbb{Z}_r}) \longrightarrow \mathrm{Hom}(1_W, A_{\mathbb{Z}_r})$$

$$\downarrow \varphi \longmapsto \gamma_{A_{\mathbb{Z}_r}}^{1-a} \cdot \left(\text{diagram} \right).$$

Expanding $\varphi: 1_W \rightarrow A_{\mathbb{Z}_r}$ as $\sum_{g=0}^{r-1} \varphi_g$ with $\varphi_g: 1_W \rightarrow {}_g(1_W)$ and using (4.22), a variant of (4.17) reads

$$C_a \cong \bigoplus_{g=0}^{r-1} \left\{ \downarrow \varphi_g \left| \frac{1}{r} \sum_{h=0}^{r-1} \xi^{-h(1-a)} \left(\text{diagram} \right) = \downarrow \varphi_g \right. \right\} [1-a], \quad (4.23)$$

where we used (4.20) and the isomorphism $\mathrm{Hom}^i(X, Y[1]) \cong \mathrm{Hom}^{i+1}(X, Y)$ in $\mathcal{L}\mathcal{G}^{\bullet/2}$. A direct computation along the lines of [9, Appendix 2] then reveals that

$$\xi^{-h(1-a)} \left(\text{diagram} \right) \cong \begin{cases} \xi^{h(a-1)-hj} \cdot \varphi_g & \text{if } g = 0 \text{ and } \varphi_g = x^j, \\ \xi^{ha} \cdot \varphi_g & \text{if } g \neq 0. \end{cases}$$

Hence, the summand for $g = 0$ in (4.23) is 0 for $a = 0$, and equal to the 1-dimensional \mathbb{Z}_2 -graded vector space $\mathbb{k} \cdot x^{a-1}[1-a]$ otherwise, while the summands for $g \neq 0$ contribute only if $a = 0$, namely, a term $\mathrm{Hom}(1_W, {}_g(1_W))[1] \cong \mathbb{k}[2] = \mathbb{k}$:

$$C_a \cong \delta_{a \geq 1} \cdot \mathbb{k}[1-a] \oplus \delta_{a,0} \bigoplus_{g=1}^{r-1} \mathbb{k}. \quad (4.24)$$

It follows that the quantum dimension of C_a (as an object in $\mathcal{L}\mathcal{G}(1, 1) \cong \mathrm{vect}^{\mathbb{Z}_2}$, i.e., as a super vector space) is $r - 1$ for $a = 0$, $+1$ for $a \in \mathbb{Z} \setminus \{0\}$ even, and -1 for a odd. However, in $\mathcal{L}\mathcal{G}^{\bullet/2}$ the 2-morphisms $(+1) \cdot 1_{C_a}$ and $(-1) \cdot 1_{C_a}$ are identical. Recalling Proposition 2.5, this means that the $(\mathcal{L}\mathcal{G}^{\bullet/2})_{\mathrm{eq}}$ -valued TQFT associated to $A_{\mathbb{Z}_r}$ can only distinguish two r -spin structures on the torus (for $r \neq 2$), while there are as many non-diffeomorphic r -spin structures on T^2 as there are divisors of r .

We emphasise that this example works for arbitrary $r \geq 3$, whereas the example computing the Arf invariant mentioned in Remark 3.28 (ii) is defined only for r even.

The computational techniques used in Example 4.18 can analogously be applied to more involved examples. For instance, there are \mathbb{Z}_r -actions on $\mathbb{k}[x_1, x_2]$ which leave $W = x_1^r + x_2^{2r}$ invariant, and the associated $(\mathcal{L}\mathcal{G}^{\bullet/2})_{\mathrm{eq}}$ -valued TQFTs may detect more than two r -spin structures on the torus. We leave such computations as well as

the application of the theory developed in Section 4.1 to the 2-category of \mathbb{Q} -graded Landau–Ginzburg models $\mathcal{L}\mathcal{G}^{\text{gr}}$ (see [15] or [13, Section 2.5]) to future work.

Acknowledgements. We are grateful to Tobias Dyckerhoff, Lukas Müller, Ingo Runkel, and Christopher Schommer-Pries for helpful discussions.

Funding. N.C. is supported by the DFG Heisenberg Programme.

References

- [1] P. S. Aspinwall, T. Bridgeland, A. Craw, M. R. Douglas, M. Gross, A. Kapustin, G. W. Moore, G. Segal, B. Szendrői, and P. M. H. Wilson, *Dirichlet branes and mirror symmetry*. Clay Math. Monogr. 4, American Mathematical Society, Providence, RI, 2009 Zbl 1188.14026 MR 2567952
- [2] D. Ayala and J. Francis, The cobordism hypothesis. 2017, arXiv:1705.02240
- [3] J. C. Baez and J. Dolan, [Higher-dimensional algebra and topological quantum field theory](#). *J. Math. Phys.* **36** (1995), no. 11, 6073–6105 Zbl 0863.18004 MR 1355899
- [4] M. Ballard, D. Favero, and L. Katzarkov, [A category of kernels for equivariant factorizations and its implications for Hodge theory](#). *Publ. Math. Inst. Hautes Études Sci.* **120** (2014), 1–111 Zbl 1401.14086 MR 3270588
- [5] P. Banks, Extended TQFTs and algebraic geometry. 2020, arXiv:2011.02394
- [6] J. Barrett, C. Meusburger, and G. Schaumann, Gray categories with duals and their diagrams. 2012, arXiv:1211.0529
- [7] J. W. Barrett and S. O. G. Tavares, [Two-dimensional state sum models and spin structures](#). *Comm. Math. Phys.* **336** (2015), no. 1, 63–100 Zbl 1326.81185 MR 3322367
- [8] J. Bénabou, [Introduction to bicategories](#). In *Reports of the Midwest Category Seminar*, pp. 1–77, Lecture Notes in Math. 47, Springer, Berlin, 1967 Zbl 1375.18001 MR 220789
- [9] I. Brunner, N. Carqueville, and D. Plencner, [Orbifolds and topological defects](#). *Comm. Math. Phys.* **332** (2014), no. 2, 669–712 Zbl 1300.81070 MR 3257659
- [10] L. Buhné, *Topics in three-dimensional descent theory*. Ph.D. thesis, University of Hamburg, 2015, <https://ediss.sub.uni-hamburg.de/handle/ediss/6382>, visited on 15 October 2023
- [11] A. Căldăraru and S. Willerton, The Mukai pairing. I. A categorical approach. *New York J. Math.* **16** (2010), 61–98 Zbl 1214.14013 MR 2657369
- [12] N. Carqueville, C. Meusburger, and G. Schaumann, [3-dimensional defect TQFTs and their tricategories](#). *Adv. Math.* **364** (2020), article no. 107024, 58 Zbl 1441.81125 MR 4065261
- [13] N. Carqueville and F. Montiel Montoya, [Extending Landau–Ginzburg models to the point](#). *Comm. Math. Phys.* **379** (2020), no. 3, 955–977 Zbl 1445.18016 MR 4163357
- [14] N. Carqueville and D. Murfet, [A toolkit for defect computations in Landau–Ginzburg models](#). In *String-Math 2012*, pp. 239–250, Proc. Sympos. Pure Math. 90, American Mathematical Society, Providence, RI, 2015 Zbl 1356.81195 MR 3409797

- [15] N. Carqueville and D. Murfet, [Adjunctions and defects in Landau–Ginzburg models](#). *Adv. Math.* **289** (2016), 480–566 Zbl [1353.18004](#) MR [3439694](#)
- [16] N. Carqueville and I. Runkel, [Orbifold completion of defect bicategories](#). *Quantum Topol.* **7** (2016), no. 2, 203–279 Zbl [1360.18007](#) MR [3459961](#)
- [17] C. L. Douglas, C. Schommer-Pries, and N. Snyder, [Dualizable tensor categories](#). *Mem. Amer. Math. Soc.* **268** (2020), no. 1308, vii+88 Zbl [1514.57001](#) MR [4254952](#)
- [18] T. Dyckerhoff, [Compact generators in categories of matrix factorizations](#). *Duke Math. J.* **159** (2011), no. 2, 223–274 Zbl [1252.18026](#) MR [2824483](#)
- [19] J. Fuchs and C. Stigner, On Frobenius algebras in rigid monoidal categories. *Arab. J. Sci. Eng. Sect. C Theme Issues* **33** (2008), no. 2, 175–191 Zbl [1185.18007](#) MR [2500035](#)
- [20] D. Gaiotto and T. Johnson-Freyd, Condensations in higher categories. 2019, arXiv:[1905.09566](#)
- [21] J. Grant, [Serre functors and graded categories](#). *Algebr. Represent. Theor.* (2022), DOI [10.1007/s10468-022-10151-4](#)
- [22] S. Gunningham, [Spin Hurwitz numbers and topological quantum field theory](#). *Geom. Topol.* **20** (2016), no. 4, 1859–1907 Zbl [1347.81070](#) MR [3548460](#)
- [23] N. Gurski, [Coherence in three-dimensional category theory](#). Cambridge Tracts in Math. 201, Cambridge University Press, Cambridge, 2013 Zbl [1314.18002](#) MR [3076451](#)
- [24] P. Guthmann, The tricategory of formal composites and its strictification. 2019, arXiv:[1903.05777](#)
- [25] J. Hesse, [Group actions on bicategories and topological quantum field theories](#). Ph.D. thesis, University of Hamburg, 2017, <https://ediss.sub.uni-hamburg.de/handle/ediss/7302>, visited on 15 October 2023
- [26] J. Hesse, C. Schweigert, and A. Valentino, Frobenius algebras and homotopy fixed points of group actions on bicategories. *Theory Appl. Categ.* **32** (2017), article no. 18, 652–681 Zbl [1377.18003](#) MR [3649192](#)
- [27] J. Hesse and A. Valentino, The Serre automorphism via homotopy actions and the cobordism hypothesis for oriented manifolds. *Cah. Topol. Géom. Différ. Catég.* **60** (2019), no. 2, 194–236 Zbl [1427.57023](#) MR [3931946](#)
- [28] M. Khovanov and L. Rozansky, [Matrix factorizations and link homology](#). *Fund. Math.* **199** (2008), no. 1, 1–91 Zbl [1145.57009](#) MR [2391017](#)
- [29] A. Kuznetsov, [Calabi–Yau and fractional Calabi–Yau categories](#). *J. Reine Angew. Math.* **753** (2019), 239–267 Zbl [1440.14092](#) MR [3987870](#)
- [30] A. D. Lauda and H. Pfeiffer, [State sum construction of two-dimensional open-closed topological quantum field theories](#). *J. Knot Theory Ramifications* **16** (2007), no. 9, 1121–1163 Zbl [1148.57039](#) MR [2375819](#)
- [31] J. Lurie, On the classification of topological field theories. In *Current developments in mathematics, 2008*, pp. 129–280, International Press of Boston, Somerville, MA, 2009 Zbl [1180.81122](#) MR [2555928](#)
- [32] S. Novak, [Lattice topological field theories in two dimensions](#). Ph.D. thesis, University of Hamburg, 2015, <https://ediss.sub.uni-hamburg.de/handle/ediss/6467>, visited on 15 October 2023

- [33] S. Novak and I. Runkel, [State sum construction of two-dimensional topological quantum field theories on spin surfaces](#). *J. Knot Theory Ramifications* **24** (2015), no. 5, article no. 1550028, 84 Zbl [1320.57035](#) MR [3349955](#)
- [34] P. Pstragowski, *On dualizable objects in monoidal bicategories, framed surfaces and the cobordism hypothesis*. Master's thesis, University of Bonn, 2014, arXiv:[1411.6691](#)
- [35] F. Quinn, [Lectures on axiomatic topological quantum field theory](#). In *Geometry and quantum field theory (Park City, UT, 1991)*, pp. 323–453, IAS/Park City Math. Ser. 1, American Mathematical Society, Providence, RI, 1995 Zbl [0901.18002](#) MR [1338394](#)
- [36] O. Randal-Williams, [Homology of the moduli spaces and mapping class groups of framed, \$r\$ -Spin and Pin surfaces](#). *J. Topol.* **7** (2014), no. 1, 155–186 Zbl [1322.55006](#) MR [3180616](#)
- [37] I. Runkel and L. Szegedy, [Topological field theory on \$r\$ -spin surfaces and the Arf-invariant](#). *J. Math. Phys.* **62** (2021), no. 10, article no. 102302, 34 Zbl [1480.81114](#) MR [4321198](#)
- [38] C. J. Schommer-Pries, *The classification of two-dimensional extended topological field theories*. ProQuest LLC, Ann Arbor, MI, 2009 MR [2713992](#)
- [39] W. H. Stern and L. Szegedy, [Topological field theories on open-closed \$r\$ -spin surfaces](#). *Topology Appl.* **312** (2022), article no. 108062, 40 Zbl [1501.57026](#) MR [4392318](#)
- [40] L. Szegedy, *State-sum construction of two-dimensional functorial field theories*. Ph.D. thesis, Universität Hamburg, 2018, <https://ediss.sub.uni-hamburg.de/handle/ediss/7848>, visited on 15 October 2023
- [41] T. Trimble, *Surface diagrams*. <https://ncatlab.org/toddtrimble/published/Surface+diagrams>, visited on 15 October 2023
- [42] L. Wester Hansen and M. Shulman, *Constructing symmetric monoidal bicategories functorially*. 2019, arXiv:[1910.09240](#)

Received 2 September 2021.

Nils Carqueville

Fakultät für Physik, Universität Wien, Boltzmanngasse 5, 1090 Wien, Austria;
nils.carqueville@univie.ac.at

Lóránt Szegedy

Institute for Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria;
lorant.szegedy@posteo.net