



Marc Hoyois · Joachim Jelisiejew · Denis Nardin · Burt Totaro ·
Maria Yakerson

The Hilbert scheme of infinite affine space and algebraic K-theory

Received August 10, 2021; revised October 30, 2021

Abstract. We study the Hilbert scheme $\mathrm{Hilb}_d(\mathbf{A}^\infty)$ from an $\mathbf{A}^1$ -homotopical viewpoint and obtain applications to algebraic K-theory. We show that the Hilbert scheme $\mathrm{Hilb}_d(\mathbf{A}^\infty)$ is $\mathbf{A}^1$ -equivalent to the Grassmannian of $(d - 1)$ -planes in $\mathbf{A}^\infty$. We then describe the $\mathbf{A}^1$ -homotopy type of $\mathrm{Hilb}_d(\mathbf{A}^n)$ in a certain range, for n large compared to d . For example, we compute the integral cohomology of $\mathrm{Hilb}_d(\mathbf{A}^n)(\mathbb{C})$ in a range. We also deduce that the forgetful map $\mathcal{F}\mathrm{Flat} \rightarrow \mathrm{Vect}$ from the moduli stack of finite locally free schemes to that of finite locally free sheaves is an $\mathbf{A}^1$ -equivalence after group completion. This implies that the moduli stack $\mathcal{F}\mathrm{Flat}$, viewed as a presheaf with framed transfers, is a model for the effective motivic spectrum kgl representing algebraic K-theory. Combining our techniques with the recent work of Bachmann, we obtain Hilbert scheme models for the kgl -homology of smooth proper schemes over a perfect field.

Keywords. Hilbert scheme, K-theory, pure Tate motive

1. Introduction

In this paper we analyze the Hilbert scheme of points from the $\mathbf{A}^1$ -homotopical perspective, yielding topological information about the Hilbert scheme as well as new geometric models for algebraic K-theory. For simplicity, schemes in the introduction are assumed

Marc Hoyois: Fakultät für Mathematik, Universität Regensburg, 93040 Regensburg, Germany;
marc.hoyois@ur.de

Joachim Jelisiejew: Faculty of Mathematics, Informatics and Mechanics, University of Warsaw,
02-097 Warszawa, Poland; jjelisiejew@mimuw.edu.pl

Denis Nardin: Fakultät für Mathematik, Universität Regensburg, 93040 Regensburg, Germany;
denis.nardin@ur.de

Burt Totaro: Mathematics Department, UCLA, Los Angeles, CA 90095-1555, USA;
totaro@math.ucla.edu

Maria Yakerson: Institute for Mathematical Research, ETH Zürich, 8092 Zürich, Switzerland;
maria.yakerson@math.ethz.ch

Mathematics Subject Classification (2020): Primary 14C05; Secondary 14F42, 19E08

to be over a perfect field k , even though many of our results hold over an arbitrary base scheme.

The Hilbert scheme $\mathrm{Hilb}_d(X)$ classifies zero-dimensional degree d closed subschemes of a fixed scheme X . For X a smooth surface, the Hilbert scheme is a smooth variety of dimension $2d$. It has led to rich developments in algebraic geometry, geometric representation theory and string theory [8, 24, 27, 51]. For higher-dimensional varieties X , the picture is more obscure: the Hilbert scheme is singular (for all $d > 3$), has many irreducible components (for large d), and almost arbitrary singularities (for $\dim X \geq 16$) [38, 40, 47]. Many natural questions remain open [1].

Until now, little has been known about the topology of $\mathrm{Hilb}_d(X)$ even for X an affine or projective space. We have an explicit basis for the cohomology of $\mathrm{Hilb}_d(\mathbf{A}^2)$ and its motive is pure Tate, thanks to the Białyński-Birula decomposition for a generic $\mathbf{G}_m$ -action on $\mathbf{A}^2$; see [33, Appendix A] or [13]. For $\mathrm{Hilb}_d(\mathbf{A}^n)$ with $n > 2$ this method fails due to the presence of singularities and the motive is not known. The only general results on the topology of those schemes were that $\mathrm{Hilb}_d(\mathbf{P}^n)$ and $\mathrm{Hilb}_d(\mathbf{A}^n)$ are connected, and $\mathrm{Hilb}_d(\mathbf{P}^n)$ is simply connected [29, 32].

We show that the situation simplifies after stabilization with respect to n . We consider the ind-scheme $\mathrm{Hilb}_d(\mathbf{A}^\infty) = \mathrm{colim}_n \mathrm{Hilb}_d(\mathbf{A}^n)$ and construct an $\mathbf{A}^1$ -equivalence¹ in Theorem 2.1:

$$\mathrm{Hilb}_d(\mathbf{A}^\infty) \simeq \mathrm{Gr}_{d-1}(\mathbf{A}^\infty). \quad (1.1)$$

This implies that for any oriented cohomology theory A^* on k -schemes in the sense of [12, Example 2.1.4], such as l -adic cohomology, integral cohomology over the complex numbers, motivic cohomology, homotopy invariant K-theory, algebraic cobordism, etc., we have

$$A^*(\mathrm{Hilb}_d(\mathbf{A}^\infty)) \simeq A^*(\mathrm{Spec} k)[[c_1, \dots, c_{d-1}]], \quad |c_i| = i.$$

We also prove stability results, describing the $\mathbf{A}^1$ -homotopy type of $\mathrm{Hilb}_d(\mathbf{A}^n)$ in a certain range, for n large compared to d ; see Theorem 8.2 and Corollary 8.3. For example, over the complex numbers, the homomorphism on integral cohomology

$$H^*(\mathrm{Gr}_{d-1}(\mathbf{A}^\infty)(\mathbf{C}), \mathbf{Z}) \simeq \mathbf{Z}[c_1, \dots, c_{d-1}] \rightarrow H^*(\mathrm{Hilb}_d(\mathbf{A}^n)(\mathbf{C}), \mathbf{Z})$$

is an isomorphism in degrees at most $2n - 2d + 2$.

Remark 1.2. One can also consider the Hilbert scheme $\mathrm{Hilb}_d(\mathbf{A}^N)$ where

$$\mathbf{A}^N = \mathrm{Spec} \mathbf{Z}[x_1, x_2, \dots],$$

which contains the ind-scheme $\mathrm{Hilb}_d(\mathbf{A}^\infty)$ as a subfunctor. We will show that the inclusion $\mathrm{Hilb}_d(\mathbf{A}^\infty) \subset \mathrm{Hilb}_d(\mathbf{A}^N)$ is an $\mathbf{A}^1$ -equivalence; see Proposition 4.2.

¹Here and further in the Introduction, we say that a map of presheaves is an $\mathbf{A}^1$ -equivalence if it becomes an $\mathbf{A}^1$ -equivalence when restricted to affine schemes; in particular, such a map is a motivic equivalence.

To prove (1.1), we consider the stack $\mathcal{F}\text{Flat}_d$ of finite flat degree d algebras and the stack $\mathcal{V}\text{ect}_d$ of rank d locally free sheaves. The forgetful maps

$$\text{Hilb}_d(\mathbf{A}^\infty) \rightarrow \mathcal{F}\text{Flat}_d, \quad \text{Gr}_d(\mathbf{A}^\infty) \rightarrow \mathcal{V}\text{ect}_d$$

are shown to be $\mathbf{A}^1$ -equivalences; see Propositions 4.2 and 4.7. For any ring R and R -module M the R -module $R \oplus M$ has a commutative unital R -algebra structure with $M^2 = 0$, called the *square-zero extension*. This construction defines a morphism

$$\alpha: \mathcal{V}\text{ect}_{d-1} \rightarrow \mathcal{F}\text{Flat}_d.$$

We show that α is an $\mathbf{A}^1$ -homotopy equivalence, with inverse β sending an R -algebra A to the R -module $A/(R \cdot 1_A)$. The $\mathbf{A}^1$ -homotopy between $\alpha\beta$ and the identity is obtained using the Rees algebra, making $\mathcal{F}\text{Flat}_d$ a cone over $\mathcal{V}\text{ect}_{d-1}$. On the technical level, this is the key point to avoid dealing with the singularities of $\mathcal{F}\text{Flat}_d$.

The stacks $\mathcal{F}\text{Flat} = \coprod_d \mathcal{F}\text{Flat}_d$ and $\mathcal{V}\text{ect} = \coprod_d \mathcal{V}\text{ect}_d$ are commutative monoids under direct sum. While the maps α and β do not preserve this structure, we employ them together with McDuff–Segal’s group completion theorem to deduce that the forgetful map $\mathcal{F}\text{Flat} \rightarrow \mathcal{V}\text{ect}$ is an $\mathbf{A}^1$ -equivalence after group completion; see Theorem 3.1. As a consequence, we get a new description of algebraic K-theory in terms of Hilbert schemes: on affine schemes, there is an $\mathbf{A}^1$ -equivalence

$$K \simeq \underline{\mathbf{Z}} \times \text{Hilb}_\infty(\mathbf{A}^\infty); \quad (1.3)$$

see Corollary 4.5. Here, K is the presheaf of K-theory spaces, $\underline{\mathbf{Z}}$ is the constant sheaf with value $\mathbf{Z}$, and $\text{Hilb}_\infty(\mathbf{A}^\infty)$ is the colimit of the maps $\text{Hilb}_d(\mathbf{A}^\infty) \rightarrow \text{Hilb}_{d+1}(\mathbf{A}^{\infty+1})$.

Furthermore, we discuss consequences of these computations for stable motivic homotopy theory. These results are part of the theory of framed transfers, developed in [15, 19, 66] and other works. Some of the main results of this theory are as follows: every generalized motivic cohomology theory acquires a unique structure of framed transfers [15, Theorem 3.5.12], and certain cohomology theories acquire a universality property with respect to their transfers. The big picture of various cohomology theories and the corresponding transfers is given in [36, Section 1.1]. For example, the algebraic cobordism spectrum MGL represents the universal cohomology theory with finite syntomic transfers, i.e., with pushforwards along finite flat locally complete intersection morphisms. This is expressed by the equivalence of ∞ -categories

$$\text{Mod}_{\text{MGL}}(\mathbf{SH}(k)) \simeq \mathbf{SH}^{\text{fsyn}}(k),$$

where $\mathbf{SH}^{\text{fsyn}}(k)$ is the ∞ -category of motivic spectra with finite syntomic transfers [14, Theorem 4.1.3].

In Section 5 we work towards an analogous universal property for algebraic K-theory. We observe that the forgetful map $\mathcal{F}\text{Flat} \rightarrow \mathcal{V}\text{ect}$ is a morphism of presheaves with framed transfers in the sense of [15], and the framed suspension spectrum of $\mathcal{V}\text{ect}$ is the effective motivic K-theory spectrum kgl . We therefore obtain an equivalence of motivic spectra

$$\Sigma_{\text{T,fr}}^\infty \mathcal{F}\text{Flat} \simeq \text{kgl};$$

see Theorem 5.4. Bachmann employs it in [3] to obtain an equivalence of symmetric monoidal ∞ -categories

$$\mathrm{Mod}_{\mathrm{kgl}}(\mathbf{SH}(k))[1/e] \simeq \mathbf{SH}^{\mathrm{fflat}}(k)[1/e],$$

where e is the exponential characteristic of k . In that sense, kgl represents the universal generalized motivic cohomology theory with finite locally free transfers.

Some of our results generalize from $\mathbf{A}_k^\infty$ to $\mathbf{A}_X^\infty = X \times \mathbf{A}^\infty$ for a smooth separated k -scheme X . Answering a question by Rahul Pandharipande, we analyze the Hilbert scheme of points of $\mathbf{A}_X^\infty$ and obtain a motivic equivalence

$$\Omega_T^\infty(\mathrm{kgl} \otimes \Sigma_T^\infty X_+) \simeq \mathrm{Hilb}(\mathbf{A}_X^\infty)^{\mathrm{gp}}$$

for X smooth and proper, thus providing a geometric model for the kgl -homology of X ; see Corollary 5.7. Here, the $\mathcal{E}_\infty$ -structure on the Hilbert scheme is given by “taking disjoint unions of closed subschemes”. As we explain in Section 4, this equivalence is analogous to Segal’s model for ko -homology of a topological space [59, Section 1]. Indeed, the ind-scheme $\mathrm{Hilb}(\mathbf{A}_X^\infty)$ can be thought of as a geometric version of Segal’s configuration space of points “labeled” by vector spaces: it has a canonical map to the symmetric power $\mathrm{Sym}(X)$, whose fiber over a k -rational point of the form $\sum_i d_i [x_i]$ is $\mathbf{A}^1$ -equivalent to the stack $\prod_i \mathrm{Vect}_{d_i-1}$.

We proceed with investigating further the connection between algebraic cobordism and algebraic K-theory. In [14], it is shown that the algebraic cobordism spectrum MGL is the framed suspension spectrum of the substack $\mathcal{FSyn} \subset \mathcal{FFlat}$ of finite syntomic schemes. We show in Section 6 that the inclusion $\mathcal{FSyn} \subset \mathcal{FFlat}$ induces the standard orientation map $\mathrm{MGL} \rightarrow \mathrm{kgl}$ upon taking framed suspension spectra. Combining the results of this paper with [5, Theorem 1.1], we obtain the following commutative diagram in $\mathrm{PSh}(\mathrm{Sm}_k)$, where the symbol $\simeq$ denotes a motivic equivalence and $+$ a motivic equivalence up to Quillen’s plus construction:

$$\begin{array}{ccccc}
 \mathbf{Z} \times \mathrm{Hilb}_\infty^{\mathrm{lci}}(\mathbf{A}^\infty) & \hookrightarrow & \mathbf{Z} \times \mathrm{Hilb}_\infty(\mathbf{A}^\infty) & & \mathbf{Z} \times \mathrm{Gr}_\infty(\mathbf{A}^\infty) \\
 \downarrow \simeq & & \downarrow \simeq & & \downarrow \simeq \\
 \mathbf{Z} \times \mathcal{FSyn}_\infty & \hookrightarrow & \mathbf{Z} \times \mathcal{FFlat}_\infty & \xrightarrow{\simeq} & \mathbf{Z} \times \mathrm{Vect}_\infty \\
 \downarrow + & & \downarrow \simeq & & \downarrow \simeq \\
 \mathcal{FSyn}^{\mathrm{gp}} & \longrightarrow & \mathcal{FFlat}^{\mathrm{gp}} & \xrightarrow{\simeq} & \mathrm{Vect}^{\mathrm{gp}} \\
 \downarrow \simeq & & & & \downarrow \simeq \\
 \Omega_T^\infty \mathrm{MGL} & \longrightarrow & & & \Omega_T^\infty \mathrm{KGL}
 \end{array}$$

Here $\mathrm{Hilb}_\infty(\mathbf{A}^\infty) = \mathrm{colim}_{d,n} \mathrm{Hilb}_d(\mathbf{A}^n)$ is the stabilization of the Hilbert scheme of affine space, and $\mathrm{Hilb}_\infty^{\mathrm{lci}}(\mathbf{A}^\infty)$ is the local complete intersection locus. The bottommost arrow in the diagram is the standard orientation of KGL , and all the other horizontal arrows are the obvious forgetful morphisms. Moreover, the bottom rectangle is one of

commutative monoids in presheaves with framed transfers (in particular, of presheaves of $\mathcal{E}_\infty$ -ring spaces).

One reason for our interest in $\mathrm{Hilb}_d(\mathbf{A}^\infty)$ is a conjecture of Mike Hopkins stating that the motive of $\Omega_T^\infty \mathrm{MGL}$ is pure Tate. As observed in [5, Corollary 1.7], this conjecture is equivalent to showing that $\mathrm{Hilb}_\infty^{\mathrm{lci}}(\mathbf{A}^\infty)$ has a pure Tate motive. While this problem remains open, we prove in Section 7 that $\mathrm{Hilb}_3^{\mathrm{lci}}(\mathbf{A}^\infty)$ has a pure Tate motive. In contrast, $\mathrm{Hilb}_3^{\mathrm{lci}}(\mathbf{A}^n)$ is not pure Tate for finite n [5, Remark 1.8].

Notation and terminology. Sch denotes the category of schemes. If S is a scheme, Sch_S denotes the category of S -schemes, Sm_S that of smooth S -schemes, $\mathbf{H}(S)$ the ∞ -category of motivic spaces over S , and $\mathbf{SH}(S)$ that of motivic spectra [6, Sections 2.2 and 4.1]. The *Tate sphere* $\mathbf{T}$ in $\mathbf{H}(S)$ is defined as $\mathbf{A}^1/(\mathbf{A}^1 - 0) \simeq \Sigma \mathbf{G}_m \simeq (\mathbf{P}^1, 1)$.

We only consider the Hilbert scheme of points in this paper. That is, for a morphism of schemes $X \rightarrow S$ and any S -scheme T ,

$$\mathrm{Hilb}_d(X/S)(T) = \left\{ \begin{array}{l} Z \subset X_T \text{ closed subscheme such that} \\ Z \rightarrow T \text{ is finite locally free of degree } d \end{array} \right\}.$$

This defines a scheme over S whenever X is affine or quasi-projective over S [64, Remark 0B9B] (more generally, it is an algebraic space over S when X is separated over S [28, Theorem A (i)]). We denote by $\mathrm{Hilb}_d^{\mathrm{lci}}(X/S)$ the open subscheme of $\mathrm{Hilb}_d(X/S)$ given by the locus of local complete intersections. We write $\mathrm{Hilb}(X/S) = \coprod_{d \geq 0} \mathrm{Hilb}_d(X/S)$. We sometimes omit S from the notation when it is clear from the context.

We denote by $\mathrm{PSh}(\mathcal{C})$ the ∞ -category of presheaves of spaces on $\mathcal{C}$ and by $\mathrm{PSh}_\Sigma(\mathcal{C}) \subset \mathrm{PSh}(\mathcal{C})$ the full subcategory of presheaves that transform finite sums into finite products [44, Section 5.5.8]. If $\mathcal{C}$ is a suitable category of schemes, $L_{\mathbf{A}^1}: \mathrm{PSh}(\mathcal{C}) \rightarrow \mathrm{PSh}(\mathcal{C})$ (resp. $L_{\mathrm{mot}}: \mathrm{PSh}(\mathcal{C}) \rightarrow \mathrm{PSh}(\mathcal{C})$) is the localization functor onto the full subcategory of $\mathbf{A}^1$ -invariant presheaves (resp. of $\mathbf{A}^1$ -invariant Nisnevich sheaves). A morphism f in $\mathrm{PSh}(\mathcal{C})$ is called an $\mathbf{A}^1$ -*equivalence* (resp. a *motivic equivalence*) if $L_{\mathbf{A}^1}(f)$ (resp. $L_{\mathrm{mot}}(f)$) is an equivalence. Thus, if $\mathcal{C} = \mathrm{Sm}_S$, a motivic equivalence means an isomorphism in the ∞ -category of motivic spaces $\mathbf{H}(S)$; this should not be confused with an isomorphism in the derived category of motives, $\mathbf{DM}(S)$, which is a weaker condition. Explicitly, a morphism $f: \mathcal{X} \rightarrow \mathcal{Y}$ in $\mathrm{PSh}(\mathcal{C})$ (e.g., a morphism of stacks) is an $\mathbf{A}^1$ -equivalence if for every scheme T the morphism $\mathcal{X}(T \times \mathbf{A}^\bullet) \rightarrow \mathcal{Y}(T \times \mathbf{A}^\bullet)$ induces an equivalence on geometric realizations.

2. The stacks $\mathcal{F}\mathrm{Flat}_d$ and Vect_{d-1} are $\mathbf{A}^1$ -equivalent

We say that a morphism of schemes $p: Y \rightarrow X$ is *finite locally free* if it is finite, flat, and of finite presentation, or equivalently if it is affine and $p_* \mathcal{O}_Y$ is a locally free $\mathcal{O}_X$ -module of finite rank. We denote by $\mathrm{Vect}(X)$ the groupoid of finite locally free $\mathcal{O}_X$ -modules and by $\mathcal{F}\mathrm{Flat}(X)$ the groupoid of finite locally free X -schemes, or equivalently of finite locally free $\mathcal{O}_X$ -algebras. For $d \geq 0$, write $\mathcal{F}\mathrm{Flat}_d \subset \mathcal{F}\mathrm{Flat}$ for the subfunctor of finite locally

free commutative $\mathcal{O}_X$ -algebras of rank d , and $\mathcal{V}ect_d \subset \mathcal{V}ect$ for the subfunctor of vector bundles of rank d , which is the classifying stack of the group scheme GL_d .

The presheaves of groupoids $\mathcal{V}ect_d$ and $\mathcal{F}lat_d$ are algebraic stacks of finite type over $\mathbf{Z}$. In particular, Poonen observed that $\mathcal{F}lat_d$ is the quotient stack by GL_d of the affine scheme Y_d of commutative algebra structures on $\mathbf{A}^d$, with $\mathbf{A}^d$ viewed as a free module of rank d over $\mathbf{Z}$ [54, Definition 3.2]. The singularities of the stack $\mathcal{F}lat_d$ are known to be almost arbitrarily bad: namely, $\coprod_{d \geq 0} \mathcal{F}lat_d$ satisfies Murphy's law up to retraction [40, after Theorem 1.3].

For $d \geq 1$, we define morphisms $\alpha: \mathcal{V}ect_{d-1} \rightarrow \mathcal{F}lat_d$ and $\beta: \mathcal{F}lat_d \rightarrow \mathcal{V}ect_{d-1}$ as follows. Given a vector bundle $\mathcal{V}$ of rank $d-1$ over a scheme X , let $\alpha(\mathcal{V})$ be the sheaf of $\mathcal{O}_X$ -algebras $\mathcal{O}_X \oplus \mathcal{V}$, with $\mathcal{V}$ a square-zero ideal. Given a sheaf of algebras $\mathcal{A}$ which is locally free of rank d over X , let $\beta(\mathcal{A})$ be $\mathcal{A}/\mathcal{O}_X$, which is a vector bundle of rank $d-1$. (Indeed, the global section 1 of $\mathcal{A}$ is nonzero at every point of X , so locally we can complete it to a basis of $\mathcal{A}$ and find that $\mathcal{O}_X \rightarrow \mathcal{A} \rightarrow \mathcal{A}/\mathcal{O}_X$ is locally split exact.)

Theorem 2.1. *For each $d \geq 1$, the morphism $\alpha: \mathcal{V}ect_{d-1} \rightarrow \mathcal{F}lat_d$ in $\mathbf{PSh}(\mathbf{Sch})$ is an $\mathbf{A}^1$ -equivalence, with inverse β up to $\mathbf{A}^1$ -homotopy.*

The proof uses a degeneration of commutative algebras which we formulate in terms of the Rees algebra construction (Lemma 2.2). Poonen considered a closely related degeneration [54, Proposition 7.1].

The following is standard when R is a field, a recent reference being [58, Remark 2.6.5].

Lemma 2.2. *Let R be a commutative ring, and A an associative R -algebra with an increasing filtration by R -submodules $A_0 \subset A_1 \subset \cdots$ with $1 \in A_0$, $A_i \cdot A_j \subset A_{i+j}$, and $A = \bigcup_i A_i$. Assume that the R -modules $\mathrm{gr}_i(A) = A_i/A_{i-1}$ are flat for all $i \geq 0$ (where $A_{-1} = 0$). Let*

$$\mathrm{Rees}(A) = \bigoplus_{i \geq 0} A_i t^i \subset A[t]$$

be the corresponding Rees algebra. Then $\mathrm{Rees}(A)$ is a flat $R[t]$ -algebra with

$$\mathrm{Rees}(A)/(t) \simeq \mathrm{gr}_*(A) \quad \text{and} \quad \mathrm{Rees}(A)/(t-1) \simeq A.$$

If $\mathrm{gr}_i(A)$ is finite locally free as an R -module for each i and zero for i sufficiently large, then $\mathrm{Rees}(A)$ is finite locally free as an $R[t]$ -module.

Proof. It is immediate that $\mathrm{Rees}(A)/(t) \simeq \mathrm{gr}_*(A)$ and $\mathrm{Rees}(A)/(t-1) \simeq A$. The conclusions about flatness are also straightforward; we only write out the proof in the case needed below, where $\mathrm{gr}_i(A)$ is finite locally free as an R -module for each i and zero for i sufficiently large. In this case, $\mathrm{gr}_i(A)$ is projective as an R -module for each i [64, Lemma 00NX], and so each inclusion $A_{i-1} \rightarrow A_i$ has an R -linear splitting. These splittings determine an isomorphism $\mathrm{Rees}(A) \simeq \mathrm{gr}_*(A)[t]$ of $R[t]$ -modules. Here $\mathrm{gr}_*(A)$ is a finite locally free R -module, and so $\mathrm{Rees}(A)$ is a finite locally free $R[t]$ -module. ■

Proof of Theorem 2.1. The composition $\mathcal{V}ect_{d-1} \rightarrow \mathcal{F}Flat_d \rightarrow \mathcal{V}ect_{d-1}$ is isomorphic to the identity. It remains to show that the composition $\mathcal{F}Flat_d \rightarrow \mathcal{V}ect_{d-1} \rightarrow \mathcal{F}Flat_d$ is $\mathbf{A}^1$ -homotopic to the identity. We define an explicit $\mathbf{A}^1$ -homotopy as follows.

Let R be a ring and let A be a finite locally free R -algebra of rank d . The point is that A has a canonical increasing filtration given by $A_0 = R \cdot 1_A$ and $A_1 = A$. The R -modules $gr_i(A)$ are locally free, and so Lemma 2.2 applies. The associated Rees algebra $\text{Rees}(A)$ is locally free of rank d over $R[t]$, with $\text{Rees}(A)/(t) \simeq gr_*(A)$ and $\text{Rees}(A)/(t-1) \simeq A$. Here $gr_*(A)$ is the square-zero extension $R \oplus A/(R \cdot 1)$.

The operation $A \mapsto \text{Rees}(A)$ is functorial in A and hence defines a morphism of stacks over $\mathbf{Z}$,

$$\mathbf{A}^1 \times \mathcal{F}Flat_d \rightarrow \mathcal{F}Flat_d.$$

At $t = 1$ in $\mathbf{A}^1$, this is the identity, while at $t = 0$ it is isomorphic to the composition $\mathcal{F}Flat_d \rightarrow \mathcal{V}ect_{d-1} \rightarrow \mathcal{F}Flat_d$, sending $\mathcal{A}$ to the square-zero extension $\mathcal{O}_X \oplus \mathcal{A}/(\mathcal{O}_X \cdot 1_{\mathcal{A}})$. This is the desired $\mathbf{A}^1$ -homotopy. $\blacksquare$

Remark 2.3. The stacks $\mathcal{V}ect$ and $\mathcal{F}Flat_{\geq 1}$ are the Zariski sheafifications of the disjoint unions $\coprod_{d \geq 0} \mathcal{V}ect_d$ and $\coprod_{d \geq 1} \mathcal{F}Flat_d$ in $\text{PSh}(\text{Sch})$. Since disjoint unions and sheafification preserve $\mathbf{A}^1$ -homotopies, Theorem 2.1 implies that $\alpha: \mathcal{V}ect \rightarrow \mathcal{F}Flat_{\geq 1}$ is an $\mathbf{A}^1$ -equivalence in $\text{PSh}(\text{Sch})$, with inverse β up to $\mathbf{A}^1$ -homotopy.

3. $\mathbf{A}^1$ -equivalence between the group completions of the stacks $\mathcal{F}Flat$ and $\mathcal{V}ect$

The direct sum and tensor product of $\mathcal{O}_X$ -modules define an $\mathcal{E}_{\infty}$ -semiring structure on the stack $\mathcal{V}ect$. Similarly, the disjoint union and cartesian product of schemes define an $\mathcal{E}_{\infty}$ -semiring structure on $\mathcal{F}Flat$. In order to reduce the technical burden here, we postpone the formal construction of these semirings to Section 5. We denote by $\mathcal{V}ect^{\text{gp}}$ and $\mathcal{F}Flat^{\text{gp}}$ the corresponding group completions, which are presheaves of $\mathcal{E}_{\infty}$ -ring spaces on the category of schemes.

Forgetting the algebra structure on a finite locally free $\mathcal{O}_X$ -algebra defines a morphism of $\mathcal{E}_{\infty}$ -semirings $\eta: \mathcal{F}Flat \rightarrow \mathcal{V}ect$. The main result of this section is the following theorem:

Theorem 3.1. *The map $\eta^{\text{gp}}: \mathcal{F}Flat^{\text{gp}} \rightarrow \mathcal{V}ect^{\text{gp}}$ is an $\mathbf{A}^1$ -equivalence.*

Let $\sigma: \mathcal{F}Flat \rightarrow \mathcal{F}Flat$ be the map that adds a disjoint point over the base, and let $\tau: \mathcal{V}ect \rightarrow \mathcal{V}ect$ be the map that adds a trivial line bundle. We denote by $\mathcal{F}Flat^{\text{st}}$ the colimit of the sequence

$$\mathcal{F}Flat \xrightarrow{\sigma} \mathcal{F}Flat \xrightarrow{\sigma} \mathcal{F}Flat \rightarrow \cdots.$$

We similarly define $\mathcal{V}ect^{\text{st}}$ as the colimit of the sequence

$$\mathcal{V}ect \xrightarrow{\tau} \mathcal{V}ect \xrightarrow{\tau} \mathcal{V}ect \rightarrow \cdots.$$

The maps $\mathcal{F}\text{Flat} \rightarrow \mathcal{F}\text{Flat}^{\text{gp}}, \mathcal{A} \mapsto \mathcal{A} - n\mathcal{O}$, for $n \geq 0$ form a cone over the above sequence, which induces a canonical map $\mathcal{F}\text{Flat}^{\text{st}} \rightarrow \mathcal{F}\text{Flat}^{\text{gp}}$. There is similarly a canonical map $\mathcal{V}\text{ect}^{\text{st}} \rightarrow \mathcal{V}\text{ect}^{\text{gp}}$.

Note that there is a commutative square

$$\begin{array}{ccc} \mathcal{F}\text{Flat} & \xrightarrow{\eta} & \mathcal{V}\text{ect} \\ \sigma \downarrow & & \downarrow \tau \\ \mathcal{F}\text{Flat} & \xrightarrow{\eta} & \mathcal{V}\text{ect} \end{array}$$

inducing a map $\eta^{\text{st}}: \mathcal{F}\text{Flat}^{\text{st}} \rightarrow \mathcal{V}\text{ect}^{\text{st}}$ in the colimit. We shall deduce Theorem 3.1 from the following variant, which does not involve group completion:

Theorem 3.2. *The map $\eta^{\text{st}}: \mathcal{F}\text{Flat}^{\text{st}} \rightarrow \mathcal{V}\text{ect}^{\text{st}}$ is an $\mathbf{A}^1$ -equivalence.*

To prove Theorem 3.2, we consider the algebraic stack $\mathcal{F}\text{Flat}^{\text{mrk}}$ of finite locally free schemes with a distinguished point. More precisely, for every scheme X , the groupoid $\mathcal{F}\text{Flat}^{\text{mrk}}(X)$ is given by

$$\mathcal{F}\text{Flat}^{\text{mrk}}(X) = \{(f, s) \mid f: Z \rightarrow X \text{ finite locally free, } s: X \rightarrow Z \text{ a section of } f\}.$$

We will refer to the section s as the *marking* and to such pairs (f, s) as *marked schemes*. There is a forgetful map $\theta: \mathcal{F}\text{Flat}^{\text{mrk}} \rightarrow \mathcal{F}\text{Flat}$ that discards the marking (which is the universal finite locally free family).

Let $\mathcal{F}\text{Flat}^{\text{nu}}$ be the stack of finite locally free sheaves of nonunital commutative algebras. There is an equivalence

$$\mathcal{F}\text{Flat}^{\text{nu}} \simeq \mathcal{F}\text{Flat}^{\text{mrk}}$$

sending a nonunital algebra $\mathcal{A}$ to its unitization $\mathcal{O} \oplus \mathcal{A}$, with inverse sending an augmented algebra to its augmentation ideal. Let $v: \mathcal{V}\text{ect} \rightarrow \mathcal{F}\text{Flat}^{\text{nu}}$ be the functor sending a finite locally free sheaf $\mathcal{V}$ to $\mathcal{V}$ regarded as a nonunital algebra with zero multiplication.

The following table summarizes the various maps we will use in the proof of Theorem 3.2, some of which are introduced below:

Name	Description
$\eta: \mathcal{F}\text{Flat} \rightarrow \mathcal{V}\text{ect}$	forgets the algebra structure
$\theta: \mathcal{F}\text{Flat}^{\text{mrk}} \rightarrow \mathcal{F}\text{Flat}$	forgets the marking
$\tau: \mathcal{V}\text{ect} \rightarrow \mathcal{V}\text{ect}$	adds a trivial line bundle
$\sigma: \mathcal{F}\text{Flat} \rightarrow \mathcal{F}\text{Flat}$	adds a disjoint point
$\sigma^{\text{mrk}}: \mathcal{F}\text{Flat}^{\text{mrk}} \rightarrow \mathcal{F}\text{Flat}^{\text{mrk}}$	adds a disjoint point without changing the marking
$\varepsilon: \mathcal{F}\text{Flat}^{\text{mrk}} \rightarrow \mathcal{F}\text{Flat}^{\text{mrk}}$	adds a new tangent direction at the marked point
$v: \mathcal{V}\text{ect} \rightarrow \mathcal{F}\text{Flat}^{\text{nu}}$	equips a module with zero multiplication
$\alpha: \mathcal{V}\text{ect} \rightarrow \mathcal{F}\text{Flat}$	forms the trivial square-zero extension
$\pi: \mathcal{F}\text{Flat}^{\text{nu}} \rightarrow \mathcal{V}\text{ect}$	forgets the algebra structure

Proposition 3.3. *The map $v: \mathcal{V}\text{ect} \rightarrow \mathcal{F}\text{Flat}^{\text{nu}}$ is an $\mathbf{A}^1$ -equivalence.*

Proof. Let $\pi: \mathcal{F}\text{Flat}^{\text{nu}} \rightarrow \mathcal{V}\text{ect}$ be the forgetful functor. Then $\pi \circ \nu$ is the identity, and the map

$$\mathcal{F}\text{Flat}^{\text{nu}} \rightarrow \text{Maps}(\mathbf{A}^1, \mathcal{F}\text{Flat}^{\text{nu}}), \quad \mathcal{A} \mapsto t\mathcal{A}[t],$$

is an $\mathbf{A}^1$ -homotopy from $\nu \circ \pi$ to the identity of $\mathcal{F}\text{Flat}^{\text{nu}}$. ■

Proposition 3.4. *The map $\theta: \mathcal{F}\text{Flat}^{\text{mrk}} \rightarrow \mathcal{F}\text{Flat}_{\geq 1}$ is an $\mathbf{A}^1$ -equivalence.*

Proof. We have a commutative square

$$\begin{array}{ccc} \mathcal{V}\text{ect} & \xrightarrow{\alpha} & \mathcal{F}\text{Flat}_{\geq 1} \\ \nu \downarrow & & \uparrow \theta \\ \mathcal{F}\text{Flat}^{\text{nu}} & \xrightarrow{\simeq} & \mathcal{F}\text{Flat}^{\text{mrk}} \end{array}$$

The map α is an $\mathbf{A}^1$ -equivalence by Remark 2.3 and the map ν is an $\mathbf{A}^1$ -equivalence by Proposition 3.3. By the 2-out-of-3 property, θ is an $\mathbf{A}^1$ -equivalence. ■

Let $\sigma^{\text{mrk}}: \mathcal{F}\text{Flat}^{\text{mrk}} \rightarrow \mathcal{F}\text{Flat}^{\text{mrk}}$ be the map that adds a disjoint point over the base without changing the marking. We then have a commutative square

$$\begin{array}{ccc} \mathcal{F}\text{Flat}^{\text{mrk}} & \xrightarrow{\theta} & \mathcal{F}\text{Flat} \\ \downarrow \sigma^{\text{mrk}} & & \downarrow \sigma \\ \mathcal{F}\text{Flat}^{\text{mrk}} & \xrightarrow{\theta} & \mathcal{F}\text{Flat} \end{array}$$

inducing in the colimit a map

$$\theta^{\text{st}}: \mathcal{F}\text{Flat}^{\text{mrk, st}} \rightarrow \mathcal{F}\text{Flat}^{\text{st}}.$$

Corollary 3.5. *The map $\theta^{\text{st}}: \mathcal{F}\text{Flat}^{\text{mrk, st}} \rightarrow \mathcal{F}\text{Flat}^{\text{st}}$ is an $\mathbf{A}^1$ -equivalence.*

Proof. This follows immediately from Proposition 3.4 and the observation that the inclusion $\mathcal{F}\text{Flat}_{\geq 1} \subset \mathcal{F}\text{Flat}$ becomes an equivalence after stabilization along σ . ■

Lemma 3.6. *Let S be a scheme and let $Z \rightarrow X$ and $Z \rightarrow Y$ be morphisms between finite locally free S -schemes with $Z \rightarrow X$ a closed immersion. Then the pushout $X \sqcup_Z Y$ in the category of schemes exists and is finite locally free over S .*

Proof. The pushout exists and is affine over S by [64, Tag 0E25]. Now, local freeness translates into an algebraic statement: if $M \rightarrow P$ and $N \rightarrow P$ are morphisms of finite locally free modules with $M \rightarrow P$ surjective, then $M \times_P N$ is finite locally free. This is true, since the module $M \times_P N$ is an extension of the locally free module $\ker(M \rightarrow P)$ by the locally free module N . ■

Proof of Theorem 3.2. Consider the finite locally free $\mathbf{Z}[t]$ -scheme

$$\mathcal{R} = \text{Spec } \mathbf{Z}[x, t]/((x - t)x)$$

whose fibers over 0 and 1 are $\mathcal{R}_0 = \operatorname{Spec} \mathbf{Z}[x]/(x^2)$ and $\mathcal{R}_1 = \operatorname{Spec} \mathbf{Z} \sqcup \operatorname{Spec} \mathbf{Z}$. There is a section $i: \operatorname{Spec} \mathbf{Z}[t] \rightarrow \mathcal{R}$ sending x to 0, which defines an element $(\mathcal{R}, i)$ in $\mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}}(\mathbf{Z}[t])$.

Let $\varepsilon: \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}} \rightarrow \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}}$ be the map that sends $(f: Z \rightarrow S, s: S \rightarrow Z)$ to $Z \sqcup_S (\mathcal{R}_0)_S$ with section the canonical map $S \rightarrow Z \sqcup_S (\mathcal{R}_0)_S$. The map ε is well-defined by Lemma 3.6. In plain words, the transformation ε adds a new tangent direction at the marked point of the finite scheme. Similarly, let $H: \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}} \rightarrow \operatorname{Maps}(\mathbf{A}^1, \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}})$ be the map that sends $(f: Z \rightarrow S, s: S \rightarrow Z)$ to the pushout $\mathbf{A}_Z^1 \sqcup_{\mathbf{A}_S^1} \mathcal{R}_S$. It is then clear that $H_0 = \varepsilon$ and $H_1 = \sigma^{\operatorname{mrk}}$, i.e., H is an $\mathbf{A}^1$ -homotopy from ε to $\sigma^{\operatorname{mrk}}$.

We now have commutative squares

$$\begin{array}{ccc} \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t} & \xrightarrow{\nu} & \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{nu}} \simeq \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}} \\ \downarrow \tau & & \downarrow \varepsilon \\ \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t} & \xrightarrow{\nu} & \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{nu}} \simeq \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}} \end{array} \quad \begin{array}{ccc} \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}} & \xrightarrow{\theta} & \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t} \xrightarrow{\eta} \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t} \\ \downarrow \sigma^{\operatorname{mrk}} & & \downarrow \sigma \\ \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}} & \xrightarrow{\theta} & \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t} \xrightarrow{\eta} \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t} \end{array}$$

with the following properties:

- (1) $\nu: \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t} \rightarrow \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{nu}}$ is an $\mathbf{A}^1$ -equivalence, by Proposition 3.3;
- (2) there is an $\mathbf{A}^1$ -homotopy $H: \varepsilon \rightsquigarrow \sigma^{\operatorname{mrk}}$;
- (3) $\theta^{\operatorname{st}}: \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}, \operatorname{st}} \rightarrow \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{st}}$ is an $\mathbf{A}^1$ -equivalence, by Corollary 3.5;
- (4) the $\mathbf{A}^1$ -homotopy $\eta \circ \theta \circ H \circ \nu$ is constant if we identify its endpoints using the isomorphism

$$\eta \circ \theta \circ \nu \circ \tau \simeq \tau \circ \tau \stackrel{\operatorname{swap}}{\simeq} \tau \circ \tau \simeq \tau \circ \eta \circ \theta \circ \nu.$$

Assertion (4) is a straightforward verification from the definition of the $\mathbf{A}^1$ -homotopy H . Assertion (2) allows us to define a map $\nu^{\operatorname{st}}: L_{\mathbf{A}^1} \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}^{\operatorname{st}} \rightarrow L_{\mathbf{A}^1} \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{mrk}, \operatorname{st}}$, which is an equivalence by (1). Assertion (4) implies that $L_{\mathbf{A}^1} (\eta^{\operatorname{st}} \circ \theta^{\operatorname{st}}) \circ \nu^{\operatorname{st}} \simeq L_{\mathbf{A}^1} \tau^{\operatorname{st}}$, where $\tau^{\operatorname{st}}: \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}^{\operatorname{st}} \rightarrow \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}^{\operatorname{st}}$ is the action of $\mathcal{O} \in \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}$ on the $\mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}$ -module $\mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}^{\operatorname{st}}$. Note that τ^{st} is *not* an equivalence, but it is an $\mathbf{A}^1$ -equivalence by [5, Proposition 5.1] since the cyclic permutation of $\mathcal{O}^3$ is $\mathbf{A}^1$ -homotopic to the identity. From this and (3) we deduce that η^{st} is an $\mathbf{A}^1$ -equivalence. $\blacksquare$

Proof of Theorem 3.1. Let $\mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}[-1]$ and $\mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}[-1]$ be the commutative monoids obtained from $\mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}$ and $\mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}$ by inverting $\mathcal{O}$. We have a commutative square

$$\begin{array}{ccc} \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}^{\operatorname{st}} & \longrightarrow & \mathcal{F}\mathcal{F}\mathcal{a}\mathcal{t}[-1] \\ \downarrow & & \downarrow \\ \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}^{\operatorname{st}} & \longrightarrow & \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}[-1] \end{array}$$

The cyclic permutation of $\mathcal{O}^{\oplus 3}$ becomes homotopic to the identity in $L_{\mathbf{A}^1} \mathcal{V}\mathcal{e}\mathcal{c}\mathcal{t}$, since it is a product of elementary matrices. It thus follows from [5, Proposition 5.1] that the lower horizontal map is an $\mathbf{A}^1$ -equivalence. By Theorem 3.2, the left vertical map is an $\mathbf{A}^1$ -equivalence. It then follows from [5, Proposition 5.1] that the upper horizontal map is also an $\mathbf{A}^1$ -equivalence. Hence, the right vertical map is an $\mathbf{A}^1$ -equivalence.

Since the functor $L_{\mathbf{A}^1}$ commutes with group completion [34, Lemma 5.5], we deduce that $\mathcal{F}\text{Flat}^{\text{gp}} \rightarrow \text{Vect}^{\text{gp}}$ is an $\mathbf{A}^1$ -equivalence. ■

Remark 3.7. It follows from Theorem 3.2 and [5, Proposition 5.1] that the cyclic permutation of three points in $\mathcal{F}\text{Flat}$ becomes $\mathbf{A}^1$ -homotopic to the identity in $\mathcal{F}\text{Flat}^{\text{st}}$. In fact, one can directly show that it is $\mathbf{A}^1$ -homotopic to the identity in $\mathcal{F}\text{Flat}$ as follows. Let $X = \mathbf{Z}[x, y, z]/(xy, xz, yz)$ be three affine lines glued together at the origin. There is a finite flat morphism $X \rightarrow \mathbf{A}^1 = \text{Spec } \mathbf{Z}[t]$ given by $t \mapsto x + y + z$, which is an $\mathbf{A}^1$ -homotopy between three points and the finite flat $\mathbf{Z}$ -scheme

$$X_0 = \text{Spec } \mathbf{Z}[x, y, z]/(xy, xz, yz, x + y + z).$$

The cyclic group C_3 acts on $X/\mathbf{A}^1$ by permuting the three lines, so it remains to show that the induced automorphism c of X_0 is $\mathbf{A}^1$ -homotopic to the identity. Note that X_0 is a square-zero extension $\text{Spec } \mathbf{Z} \oplus V$ where $V = \mathbf{Z}\{x, y, z\}/\mathbf{Z}\{x + y + z\}$. Hence the group $\text{GL}(V)$ acts on X_0 , and the matrix of c in the basis (x, y) of V is $\begin{pmatrix} 0 & 1 \\ 1 & -1 \end{pmatrix}$, which has determinant 1. We conclude using the fact that $(L_{\mathbf{A}^1} \text{SL}_2)(\mathbf{Z})$ is connected, since $\text{SL}_2(\mathbf{Z})$ is generated by elementary matrices of row additions.

4. Consequences for the Hilbert scheme of $\mathbf{A}^\infty$

Fix a base scheme S . For a separated S -scheme X we write $\text{Hilb}(\mathbf{A}_X^\infty/S)$ for the colimit $\text{colim}_n \text{Hilb}(\mathbf{A}_X^n/S)$ taken in $\text{PSh}(\text{Sch}_S)$. For a set E , we also consider $\text{Hilb}(\mathbf{A}_X^E/S)$, the Hilbert scheme of points of the scheme $\mathbf{A}_X^E = \text{Spec } \mathbf{Z}[x_e \mid e \in E] \times X$ over S .

Let us recall a useful lemma from [15]. A presheaf $\mathcal{F}: \text{Sch}_S^{\text{op}} \rightarrow \text{Spc}$ satisfies *closed gluing (for affine schemes)* if it sends the empty scheme to the point and it sends pushouts of closed embeddings (of affine schemes) to pullbacks [15, Definition A.2.1]. Presheaves parametrizing embeddings often satisfy closed gluing.

We shall say that a morphism f in $\text{PSh}(\text{Sch}_S)$ is a *universal $\mathbf{A}^1$ -equivalence on affine schemes* if any base change of f in $\text{PSh}(\text{Sch}_S)$ is an $\mathbf{A}^1$ -equivalence on affine schemes.

Lemma 4.1. (1) *Let $\mathcal{F}: \text{Sch}_S^{\text{op}} \rightarrow \text{Spc}$ be a presheaf of spaces that satisfies closed gluing for affine schemes and assume that for every affine scheme $\text{Spec } A$ over S and finitely generated ideal $I \subset A$ the morphism*

$$\mathcal{F}(\text{Spec } A) \rightarrow \mathcal{F}(\text{Spec } A/I)$$

is surjective on π_0 . Then $\mathcal{F}$ is $\mathbf{A}^1$ -contractible on affine schemes.

(2) *Let $f: \mathcal{F} \rightarrow \mathcal{G}$ be a map of presheaves of spaces on Sch_S such that for every affine scheme $T \in \text{Sch}_S$ the fiber over every T -point $T \rightarrow \mathcal{G}$ satisfies closed gluing for affine schemes (as a presheaf on Sch_T). Suppose that for every affine scheme $\text{Spec } A$ over S and every finitely generated ideal $I \subset A$ the morphism*

$$\mathcal{F}(\text{Spec } A) \rightarrow \mathcal{F}(\text{Spec } A/I) \times_{\mathcal{G}(\text{Spec } A/I)} \mathcal{G}(\text{Spec } A)$$

is surjective on π_0 . Then f is a universal $\mathbf{A}^1$ -equivalence on affine schemes.

Proof. (1) We will use the same strategy as in the proof of [15, Lemma 2.3.22]. Recall that $(L_{\mathbf{A}^1} \mathcal{F})(\mathrm{Spec} A)$ is the geometric realization of the simplicial space

$$[n] \mapsto \mathcal{F}(\Delta_A^n) = \mathcal{F}(\mathrm{Spec} A[t_0, \dots, t_n] / (\sum_i t_i - 1)).$$

Since $\mathcal{F}$ satisfies closed gluing, [15, Lemma A.2.6] implies that

$$\mathrm{Maps}(\partial \Delta^n, \mathcal{F}(\Delta_A^\bullet)) \simeq \mathcal{F}(\partial \Delta_A^n)$$

where $\partial \Delta_A^n$ is the zero locus of $t_0 \cdots t_n$ in Δ_A^n . In particular, the map

$$\mathrm{Maps}(\Delta^n, \mathcal{F}(\Delta_A^\bullet)) \rightarrow \mathrm{Maps}(\partial \Delta^n, \mathcal{F}(\Delta_A^\bullet))$$

is surjective on π_0 . Therefore the geometric realization of $\mathcal{F}(\Delta_A^\bullet)$ is contractible [45, Lemma A.5.3.7].

(2) This follows from (1) by the universality of colimits. Namely, any presheaf $\mathcal{H}$ on affine schemes in Sch_S can be written as a colimit of representables T , and any base change $\mathcal{F} \times_{\mathcal{G}} \mathcal{H} \rightarrow \mathcal{H}$ of f is correspondingly the colimit of the projection maps $\mathcal{F} \times_{\mathcal{G}} T \rightarrow T$. By the assumption and (1), $\mathcal{F} \times_{\mathcal{G}} T$ is $\mathbf{A}^1$ -contractible on affine schemes, as a presheaf on Sch_T . Since $\mathbf{A}^1$ -equivalences on affine schemes are preserved by the forgetful functor $\mathrm{PSh}(\mathrm{Sch}_T) \rightarrow \mathrm{PSh}(\mathrm{Sch}_S)$ and by colimits, we conclude that $\mathcal{F} \times_{\mathcal{G}} \mathcal{H} \rightarrow \mathcal{H}$ is an $\mathbf{A}^1$ -equivalence on affine schemes. ■

For an S -scheme X , let $h_S^{\mathrm{fflat}}(X)$ (resp. $h_S^{\mathrm{fsyn}}(X)$) be the presheaf of groupoids that sends an S -scheme U to the groupoid of spans $U \leftarrow Z \rightarrow X$ where $Z \rightarrow U$ is finite locally free (resp. finite syntomic) and the morphisms are over S . In particular, $h_S^{\mathrm{fflat}}(S)$ is the presheaf $\mathcal{FFlat}$ restricted to S -schemes. There is a forgetful map $\mathrm{Hilb}(\mathbf{A}_X^\infty/S) \rightarrow h_S^{\mathrm{fflat}}(X)$ sending a closed subscheme $Z \hookrightarrow \mathbf{A}_X^\infty \times_S U$ to the span $U \leftarrow Z \rightarrow X$.

Proposition 4.2. *Let X be a separated S -scheme.*

- (1) *The forgetful map $\mathrm{Hilb}(\mathbf{A}_X^\infty/S) \rightarrow h_S^{\mathrm{fflat}}(X)$ is a universal $\mathbf{A}^1$ -equivalence on affine schemes.*
- (2) *If E is an infinite set and $X \rightarrow S$ is affine, the forgetful map $\mathrm{Hilb}(\mathbf{A}_X^E/S) \rightarrow h_S^{\mathrm{fflat}}(X)$ is a universal $\mathbf{A}^1$ -equivalence on affine schemes.*

Proof. The presheaves $\mathrm{Hilb}(\mathbf{A}_X^\infty/S)$, $\mathrm{Hilb}(\mathbf{A}_X^E/S)$, and $h_S^{\mathrm{fflat}}(X)$ all satisfy closed gluing, since they are ind-representable by algebraic stacks (alternatively, one can verify closed gluing for each of them by hand using closed gluing for finite locally free sheaves [17, Théorème 2.2 (iv)]).

So Lemma 4.1 (2) applies. Suppose $U = \mathrm{Spec} A$ is an affine scheme over S and let $U_0 = \mathrm{Spec} A/I$ for some finitely generated ideal $I = (i_1, \dots, i_r)$. Let $U \leftarrow Z = \mathrm{Spec} B \rightarrow X$ be a span in $h_S^{\mathrm{fflat}}(X)(U)$ and let $Z_0 = Z \times_U U_0$. To prove (1), we must show that every closed embedding $Z_0 \hookrightarrow \mathbf{A}_{X \times_S U_0}^r$ over $X \times_S U_0$ can be extended to a closed embedding $Z \hookrightarrow \mathbf{A}_{X \times_S U}^{r+m}$ over $X \times_S U$, for some $m \geq 0$:

$$\begin{array}{ccc}
Z_0 & \hookrightarrow & \mathbf{A}_{X \times_S U_0}^r \xrightarrow{\text{inj}} \mathbf{A}_{X \times_S U_0}^{r+m} \\
\downarrow & & \downarrow \\
Z & \dashrightarrow & \mathbf{A}_{X \times_S U}^{r+m} \\
& \searrow & \swarrow \\
& X \times_S U &
\end{array}$$

Here, inj is the canonical map adding zero coordinates.

We construct the embedding $Z \rightarrow \mathbf{A}_{X \times_S U}^{r+m} = \mathbf{A}^{r+m} \times X \times_S U$ as follows. The map to $X \times_S U$ is already given, and we lift the map $Z_0 \rightarrow \mathbf{A}^r$ to any map $Z \rightarrow \mathbf{A}^r$. It remains to fix m and the map $Z \rightarrow \mathbf{A}^m$. Such a map corresponds to a tuple of global functions on Z .

Let $i_1, \dots, i_r$ be generators of the ideal $I \subset A$ and let $b_1, \dots, b_e$ be generators of the A -module B . Let $m = re$ and take the m -tuple of functions given by all possible products $\{i_{r'}b_{e'}\}_{r' \leq r, e' \leq e}$. By assumption, the scheme Z is finite over U , hence over $\mathbf{A}_U^{r+m}$. Since X is separated over S , the scheme Z is finite over $\mathbf{A}_{X \times_S U}^{r+m}$ as well [23, Remark 9.11]. Therefore to prove that $f: Z \rightarrow \mathbf{A}_{X \times_S U}^{r+m}$ is a closed immersion it remains to show that $f^\# : \mathcal{O}_{\mathbf{A}_{X \times_S U}^{r+m}} \rightarrow f_*(\mathcal{O}_Z)$ is a surjection of sheaves. Both sheaves are $\mathcal{O}_{\mathbf{A}_{X \times_S U}^{r+m}}$ -modules of finite type, and so we can use Nakayama's lemma and check the condition on the fibers over the image of any point of Z . The statement is obvious for the points of Z_0 , since we know that $Z_0 \rightarrow \mathbf{A}_{X \times_S U_0}^{r+m}$ is a closed immersion. Every point $z \in Z - Z_0$ has a distinguished open neighborhood in Z where one of the generators of I , say i_s , is invertible. But then the image of $f^\#$ contains $i_s b_1, \dots, i_s b_e$ which generate $\mathcal{O}_{Z,z}$ as an $\mathcal{O}_{\mathbf{A}_{X \times_S U}^{r+m}, f(z)}$ -module.

Therefore $f^\#$ is surjective at z as well. This concludes the proof of (1).

We now prove (2). By the assumption, $X \times_S U$ is affine; denote it by $\text{Spec } C$. We must then prove that every generating family $(\bar{b}_e)_{e \in E}$ of the C -algebra B/IB can be lifted to a generating family $(b_e)_{e \in E}$ of B . Since B/IB is a finitely generated C -algebra, there exists a finite subset $E' \subset E$ such that $(\bar{b}_e)_{e \in E'}$ generates B/IB ; to construct it, fix a set of finitely many generators of B/IB over C and then pick E' indexing the elements of $(\bar{b}_e)_{e \in E}$ that appear in expressions for these generators. Choose arbitrary lifts $b_e \in B$ of $\bar{b}_e$ for $e \in E'$.

Since B is a finite C -algebra and I is finitely generated, IB is a finite C -module. Let $h_1, \dots, h_m$ be generators of IB as a C -module. Note that the elements h_i and b_e for $e \in E'$ generate B as a C -algebra. Since E is infinite, there exist distinct elements $e_i \in E - E'$ for $1 \leq i \leq m$. For every such i , choose an element r_i lifting $\bar{b}_{e_i}$ in the C -subalgebra of B generated by b_e for $e \in E'$. Then $b_{e_i} := r_i + h_i$ is also a lift of $\bar{b}_{e_i}$, and the elements b_{e_i} and b_e for $e \in E'$ generate B as a C -algebra. Choosing the remaining lifts b_e arbitrarily, we thus obtain a generating family $(b_e)_{e \in E}$ as desired. ■

Remark 4.3. The separatedness assumption in Proposition 4.2 is necessary for the following reason. Let X be an affine line with doubled origin and $i: \mathbf{A}^1 \hookrightarrow X$ be one of the inclusions. View $\mathbf{A}^1$ as a degree 1 family over itself. Then there does not exist a closed immersion $\mathbf{A}^1 \hookrightarrow \mathbf{A}_X^{m+1}$ whose projection is i .

Proposition 4.2 implies that for every affine S -scheme X the inclusion $\mathrm{Hilb}_d(\mathbf{A}_X^\infty/S) \subset \mathrm{Hilb}_d(\mathbf{A}_X^N/S)$ is an $\mathbf{A}^1$ -equivalence on affine schemes, so we concentrate on the former space.

Corollary 4.4. *The forgetful map $\mathrm{Hilb}(\mathbf{A}^\infty) \rightarrow \mathcal{F}\mathrm{Flat}$ is a universal $\mathbf{A}^1$ -equivalence on affine schemes.*

Proof. Take $X = S = \mathrm{Spec} \mathbf{Z}$ in Proposition 4.2. ■

Corollary 4.5. *The forgetful map*

$$\mathbf{Z} \times \mathrm{Hilb}_\infty(\mathbf{A}^\infty) \rightarrow \mathbf{K}$$

is an $\mathbf{A}^1$ -equivalence on affine schemes, where $\mathbf{K}$ is the presheaf of $\mathbf{K}$ -theory spaces. In particular, if R is a regular noetherian commutative ring, then

$$\mathbf{K}(R) \simeq \mathbf{Z}^{\mathrm{Spec} R} \times |\mathrm{Hilb}_\infty(\mathbf{A}^\infty)(\Delta_R^\bullet)|.$$

Proof. This map factors as

$$\mathbf{Z} \times \mathrm{Hilb}_\infty(\mathbf{A}^\infty) \rightarrow \mathcal{F}\mathrm{Flat}^{\mathrm{st}} \rightarrow \mathcal{V}\mathrm{ect}^{\mathrm{st}} \rightarrow \mathbf{K}.$$

The first map is an $\mathbf{A}^1$ -equivalence on affine schemes by Corollary 4.4, and the second map is an $\mathbf{A}^1$ -equivalence by Theorem 3.2. The third map is also an $\mathbf{A}^1$ -equivalence on affine schemes; see for example [5, Example 5.2]. The last statement follows because $\mathbf{K} \simeq L_{\mathbf{A}^1} \mathbf{K}$ on regular noetherian schemes. ■

Corollary 4.6. *Let X be a separated S -scheme. Then the forgetful map $\mathrm{Hilb}^{\mathrm{lci}}(\mathbf{A}_X^\infty/S) \rightarrow h_S^{\mathrm{fsyn}}(X)$ is a universal $\mathbf{A}^1$ -equivalence on affine schemes.*

Proof. We have a tautological pullback square

$$\begin{array}{ccc} \mathrm{Hilb}^{\mathrm{lci}}(\mathbf{A}_X^\infty/S) & \longrightarrow & h_S^{\mathrm{fsyn}}(X) \\ \downarrow & & \downarrow \\ \mathrm{Hilb}(\mathbf{A}_X^\infty/S) & \longrightarrow & h_S^{\mathrm{fflat}}(X) \end{array}$$

so the claim follows from Proposition 4.2 (1). ■

It is well known that the forgetful map $\mathrm{Gr}_d(\mathbf{A}^\infty) \rightarrow \mathcal{V}\mathrm{ect}_d$ is an $\mathbf{A}^1$ -equivalence on affine schemes. We give a proof in Proposition 4.7 below, since we could not locate a reference. In fact, we consider the following slightly more general situation, which is the linearized version of Proposition 4.2. For $X \rightarrow S$ an affine morphism of schemes and $\mathcal{F}$ a quasi-coherent sheaf on X , the scheme $\mathrm{Quot}_d(X/S, \mathcal{F})$ parametrizes quotients of $\mathcal{F}$ that are finite locally free of rank d over S [25, Theorem 4.7]. We denote by $\mathcal{C}\mathrm{oh}_d(X/S)$ the presheaf of groupoids on S -schemes parametrizing quasi-coherent sheaves on X that are finite locally free of rank d over S (which is an algebraic stack at least when $X \rightarrow S$ is of

finite type [28, Section 4]). There is a forgetful map $\text{Quot}_d(X/S, \mathcal{F}) \rightarrow \text{Coh}_d(X/S)$ that for $X = S$ becomes $\text{Gr}_d(\mathcal{F}) \rightarrow \text{Vect}_d$. We define

$$\text{Quot}_d(X/S, \mathcal{O}_X^\infty) = \text{colim}_n \text{Quot}_d(X/S, \mathcal{O}_X^n).$$

Proposition 4.7. *Let S be a scheme, X an affine S -scheme, and $d \geq 0$. Then the forgetful map*

$$\text{Quot}_d(X/S, \mathcal{O}_X^\infty) \rightarrow \text{Coh}_d(X/S)$$

is a universal $\mathbf{A}^1$ -equivalence on affine schemes.

Proof. By Lemma 4.1 (2), it suffices to show that for any affine scheme $Y = \text{Spec } A$ and any finitely presented closed subscheme $Y_0 = \text{Spec } A/I$, the map

$$\text{Quot}_d(X/S, \mathcal{O}_X^\infty)(Y) \rightarrow \text{Quot}_d(X/S, \mathcal{O}_X^\infty)(Y_0) \times_{\text{Coh}_d(X/S)(Y_0)} \text{Coh}_d(X/S)(Y)$$

is surjective. Concretely, if $Y \times_S X = \text{Spec } B$, we must show that for any B -module M that is finite locally free of rank d over A and any generators $(\bar{x}_1, \dots, \bar{x}_r)$ of M/IM as a B/IB -module, we can find generators $(x_1, \dots, x_{r+s})$ of M as a B -module such that $x_1, \dots, x_r$ lift $\bar{x}_1, \dots, \bar{x}_r$ and $x_{r+1}, \dots, x_{r+s}$ live in IM . This is indeed possible, since we can take any lifts $x_1, \dots, x_r$ and any family $(x_{r+1}, \dots, x_{r+s})$ of generators of the finitely generated B -module IM . ■

Remark 4.8. As in Proposition 4.2, one can also show that the map $\text{Quot}_d(X/S, \mathcal{O}_X^E) \rightarrow \text{Coh}_d(X/S)$ is a universal $\mathbf{A}^1$ -equivalence on affine schemes for any infinite set E .

There is a canonical map $\text{Gr}_{d-1}(\mathbf{A}^n) \rightarrow \text{Hilb}_d(\mathbf{A}^n)$ sending a surjection $\mathcal{O}_T^n \rightarrow \mathcal{E}$ to the surjection $\text{Sym}_{\mathcal{O}_T}(\mathcal{O}_T^n) \rightarrow \mathcal{O}_T \oplus \mathcal{E}$, where $\mathcal{O}_T \oplus \mathcal{E}$ is the square-zero extension of $\mathcal{O}_T$ by $\mathcal{E}$.

Corollary 4.9. *The map $\text{Gr}_{d-1}(\mathbf{A}^\infty) \rightarrow \text{Hilb}_d(\mathbf{A}^\infty)$ is an $\mathbf{A}^1$ -equivalence on affine schemes.*

Proof. There is a commutative square

$$\begin{array}{ccc} \text{Gr}_{d-1}(\mathbf{A}^\infty) & \longrightarrow & \text{Hilb}_d(\mathbf{A}^\infty) \\ \downarrow & & \downarrow \\ \text{Vect}_{d-1} & \xrightarrow{\alpha} & \mathcal{F}\text{Flat}_d \end{array}$$

The vertical maps are $\mathbf{A}^1$ -equivalences on affine schemes by Propositions 4.7 and 4.2, and the lower horizontal map is an $\mathbf{A}^1$ -equivalence by Theorem 2.1. ■

In [59, Section 1], Segal constructs for a compact topological space X its “labeled” configuration space $F(X_+)$, which is an $\mathcal{E}_\infty$ -space whose points are finite sets of points of X , labeled by jointly orthogonal real vector spaces embedded into $\mathbf{R}^\infty$. The space $F(X_+)$ has a canonical forgetful map to the total symmetric power of X , which sends $\{V_x\}_{x \in S}$ to $\sum_{x \in S} \dim(V_x) \cdot [x]$, where S is a finite set of points of X .

For a smooth quasi-projective variety X over a field k , the ind-scheme $\mathrm{Hilb}(\mathbf{A}_X^\infty)$ can be thought of as a geometric analogue of Segal's labeled configuration space of X , with vector spaces replaced by finite algebras. To see this, recall the Hilbert–Chow morphism [9, Theorem 2.16]

$$\rho_X: \mathrm{Hilb}_d(X) \rightarrow \mathrm{Sym}^d(X),$$

where $\mathrm{Sym}^d(X)$ is the quotient of X^d by the naturally acting symmetric group on d letters. On points, the morphism ρ_X sends a finite closed subscheme $Z \subset X$ to the support of Z counted with multiplicities. Fix another quasi-projective k -scheme Y and consider the composition

$$\rho_{X \times Y/X} := \mathrm{pr}_1 \circ \rho_{X \times Y}: \mathrm{Hilb}_d(X \times Y) \rightarrow \mathrm{Sym}^d(X \times Y) \rightarrow \mathrm{Sym}^d(X).$$

For $x \in X$ a closed point, denote by $\mathrm{Hilb}_d(X \times Y, x)$ the fiber $\rho_{X \times Y/X}^{-1}(d[x])$. This is the scheme parametrizing subschemes of $X \times Y$ of degree d supported on $\{x\} \times Y$. By construction [9, Theorem 2.16, (2.34)], for every quasi-projective X and Y the morphism $\rho_{X \times Y/X}$ factors through $h_d^{\mathrm{flat}}(X)$.

Lemma 4.10. *Let $\Gamma = \sum_{i=1}^r \mu_i[x_i]$ be a k -point of $\mathrm{Sym}^d(X)$, where $x_1, \dots, x_r \in X$ are nonsingular k -points. Then the fiber $\rho_{X \times Y/X}^{-1}(\Gamma)$ is isomorphic to*

$$\prod_{i=1}^r \mathrm{Hilb}_{\mu_i}(\mathbf{A}^{\dim_{x_i} X} \times Y, 0).$$

Proof. First let us reduce to the case where $r = 1$. Choose for every point x_i an open neighborhood U_i that does not contain the other points. Then for every k -scheme T and a family $Z \subset X \times Y \times T$ corresponding to a T -point of the fiber we can write Z as

$$Z \simeq \bigsqcup_i Z \times_X U_i,$$

where $Z \times_X U_i$ is supported on $\{x_i\} \times Y \times T$. Therefore

$$\rho_{X \times Y/X}^{-1}(\Gamma) \simeq \prod_i \rho_{X \times Y/X}^{-1}(\mu_i[x_i]).$$

For the rest of the proof we assume that $\Gamma = d[x]$, so that the fiber is equal to $\mathrm{Hilb}_d(X \times Y, x)$. We need to prove

$$\mathrm{Hilb}_d(X \times Y, x) \simeq \mathrm{Hilb}_d(\mathbf{A}^{\dim_x X} \times Y, 0).$$

To do so we will prove a stronger statement. Let $e = d \cdot \dim_x(X)$ and let

$$X_{x,e} = \mathrm{Spec} \mathcal{O}_{X,x} / \mathfrak{m}_{X,x}^e$$

be the e -thickening of $x \in X$. We will prove

$$\mathrm{Hilb}_d(X \times Y, x) \simeq \mathrm{Hilb}_d(X_{x,e} \times Y, x).$$

The claim then follows from the fact that $X_{x,e} \simeq \mathbf{A}_{0,e}^{\dim_x X}$ since x is a nonsingular point of X . Without loss of generality we can replace X by an open neighborhood of x and we thus assume X is affine.

Summing up, we need to show that for every k -scheme T and every T -family $Z \subset X \times Y \times T$ of degree d supported on $\{x\} \times Y \times T$, the subscheme Z in fact lies in $X_{x,e} \times Y \times T$. It is enough to show that the d -th power of every $f \in \mathfrak{m}_{X,x}$ vanishes when pulled back to Z . Shrinking X further if necessary, we may view such an f as a map $f: X \rightarrow \mathbf{A}^1$ with $f(x) = 0$. Consider the composition

$$f' = \mathrm{Sym}^d(f) \circ \rho_{X \times Y/X}: \mathrm{Hilb}_d(X \times Y) \rightarrow \mathrm{Sym}^d(X) \rightarrow \mathrm{Sym}^d(\mathbf{A}^1) \simeq \mathbf{A}^d,$$

where the last isomorphism is given by elementary symmetric polynomials and maps $d[0]$ to $0 \in \mathbf{A}^d$. Since $f(x) = 0$, the map f' restricted to $\mathrm{Hilb}_d(X \times Y, x)$ is zero.

Let $T \rightarrow \mathrm{Hilb}_d(X \times Y)$ be a map classifying a family $Z \subset X \times Y \times T$. By the description of the Hilbert–Chow morphism in terms of linearized determinants (see [9, Corollary 2.18] and [39, Equation 2.4, p. 9]), the restriction of f' to T computes exactly the non-leading coefficients of the characteristic polynomial of the $\mathcal{O}_T$ -linear endomorphism of $\mathcal{O}_Z$ given by multiplication by $f \in \mathcal{O}(X)$. Applying this to a family Z in the fiber $\mathrm{Hilb}_d(X \times Y, x)$, we see that all these coefficients are 0 and so $f^d = 0$ on Z by the Cayley–Hamilton theorem. ■

Let $\mathrm{Hilb}_d(\mathbf{A}^n \times \mathbf{A}^\infty, 0)$ be the ind-scheme $\mathrm{colim}_m \mathrm{Hilb}_d(\mathbf{A}^n \times \mathbf{A}^m, 0)$. We deduce from Lemma 4.10 that for a smooth quasi-projective n -dimensional k -scheme X , the fiber of

$$\rho_{\mathbf{A}^\infty/X}: \mathrm{Hilb}_d(X \times \mathbf{A}^\infty) \rightarrow \mathrm{Sym}^d(X) \quad (4.11)$$

over a k -point $\sum_{i=1}^r \mu_i [x_i]$ is isomorphic to $\prod_{i=1}^r \mathrm{Hilb}_{\mu_i}(\mathbf{A}^n \times \mathbf{A}^\infty, 0)$.

The restrictions of the forgetful maps $\mathrm{Hilb}_d(\mathbf{A}^n \times \mathbf{A}^m) \rightarrow \mathcal{F}\mathrm{Flat}_d$ induce a forgetful map

$$\mathrm{Hilb}_d(\mathbf{A}^n \times \mathbf{A}^\infty, 0) \rightarrow \mathcal{F}\mathrm{Flat}_d.$$

Lemma 4.12. *The forgetful map $\mathrm{Hilb}_d(\mathbf{A}^n \times \mathbf{A}^\infty, 0) \rightarrow \mathcal{F}\mathrm{Flat}_d$ is an $\mathbf{A}^1$ -equivalence on affine schemes.*

Proof. As remarked before Lemma 4.10, the map $\rho_{\mathbf{A}^n \times \mathbf{A}^\infty/\mathbf{A}^n}$ factors through $h_d^{\mathrm{fflat}}(\mathbf{A}^n)$. Denote by $h_d^{\mathrm{fflat}}(\mathbf{A}^n, 0) \hookrightarrow h_d^{\mathrm{fflat}}(\mathbf{A}^n)$ the fiber over $d[0] \in \mathrm{Sym}^d(\mathbf{A}^n)$ of the induced map. By Proposition 4.2 (1) we know that

$$\mathrm{Hilb}_d(\mathbf{A}^n \times \mathbf{A}^\infty, 0) \rightarrow h_d^{\mathrm{fflat}}(\mathbf{A}^n, 0)$$

is an $\mathbf{A}^1$ -equivalence on affine schemes. It remains to prove that the forgetful map

$$\alpha': h_d^{\mathrm{fflat}}(\mathbf{A}^n, 0) \rightarrow \mathcal{F}\mathrm{Flat}_d$$

is an $\mathbf{A}^1$ -equivalence. The map α' has a natural section $\beta': \mathcal{F}\mathrm{Flat}_d \rightarrow h_d^{\mathrm{fflat}}(\mathbf{A}^n, 0)$ that equips a family with a constant map to $0 \in \mathbf{A}^n$. The monoid $(\mathbf{A}^1, \cdot)$ acts on $\mathbf{A}^n$ and so induces a map

$$\mathbf{A}^1 \times h_d^{\mathrm{fflat}}(\mathbf{A}^n) \rightarrow h_d^{\mathrm{fflat}}(\mathbf{A}^n) \quad (4.13)$$

that sends $(t, U \leftarrow Z \xrightarrow{\varphi} \mathbf{A}^n)$ to $U \leftarrow Z \xrightarrow{t\varphi} \mathbf{A}^n$. Since $d[0]$ is an $\mathbf{A}^1$ -fixed point, the map (4.13) restricts to a map $\mathbf{A}^1 \times h_d^{\text{fflat}}(\mathbf{A}^n, 0) \rightarrow h_d^{\text{fflat}}(\mathbf{A}^n, 0)$ which gives an $\mathbf{A}^1$ -homotopy $\beta'\alpha' \simeq \text{id}_{\mathcal{F}\text{Flat}_d}$. This shows that β' is an inverse of α' up to $\mathbf{A}^1$ -homotopy. ■

Corollary 4.14. *Let X be a smooth quasi-projective k -scheme. Then the fiber of the map*

$$\rho_{\mathbf{A}_X^\infty/X}: \text{Hilb}(\mathbf{A}_X^\infty) \rightarrow \text{Sym}(X)$$

over a k -point $\sum_i d_i [x_i] \in \text{Sym}(X)$ is $\mathbf{A}^1$ -equivalent on affine schemes to $\prod_i \mathcal{F}\text{Flat}_{d_i}$, and hence also to $\prod_i \text{Vect}_{d_i-1}$.

Proof. Combine Lemma 4.10, Lemma 4.12, and Theorem 2.1. ■

5. The effective motivic K-theory spectrum

Fix a base scheme S . Let $\mathbf{Corr}^{\text{fflat}}(\text{Sch}_S)$ be the $(2, 1)$ -category whose objects are S -schemes and whose morphisms are spans $X \leftarrow Z \rightarrow Y$ with $Z \rightarrow X$ finite locally free. It is semiadditive and has a symmetric monoidal structure given by the cartesian product of S -schemes. We equip the ∞ -category $\text{PSh}_\Sigma(\mathbf{Corr}^{\text{fflat}}(\text{Sch}_S))$ with the induced symmetric monoidal structure given by Day convolution. Recall that commutative monoids for the Day convolution are precisely right-lax symmetric monoidal functors [22, Proposition 2.12]. The unit of this symmetric monoidal structure is the presheaf $\mathcal{F}\text{Flat}_S$ (the restriction of $\mathcal{F}\text{Flat}$ to Sch_S), which therefore has a unique structure of commutative monoid.

We claim that the presheaf $\mathcal{V}\text{ect}_S$ also has a structure of commutative monoid in $\text{PSh}_\Sigma(\mathbf{Corr}^{\text{fflat}}(\text{Sch}_S))$, with transfers given by the pushforward of finite locally free sheaves and multiplication given by the tensor product. To see this, we apply the symmetric monoidal unfurling construction of [7, Corollary 7.8.1] to the category of finite locally free sheaves fibered over Sch_S^{op} to obtain a right-lax symmetric monoidal functor $\mathcal{V}\text{ect}_S: \mathbf{Corr}^{\text{fflat}}(\text{Sch}_S)^{\text{op}} \rightarrow \text{Spc}$.² The forgetful map $\eta: \mathcal{F}\text{Flat}_S \rightarrow \mathcal{V}\text{ect}_S$ can then be promoted to a morphism of commutative monoids in $\text{PSh}_\Sigma(\mathbf{Corr}^{\text{fflat}}(\text{Sch}_S))$, namely the unique such morphism.

Let $\mathbf{Corr}^{\text{fr}}(\text{Sch}_S)$ be the symmetric monoidal ∞ -category of framed correspondences constructed in [15, Section 4]. Its objects are S -schemes and its morphisms are spans

$$\begin{array}{ccc} & Z & \\ f \swarrow & & \searrow g \\ X & & Y \end{array}$$

²The unfurling construction in *loc. cit.* requires a symmetric monoidal Waldhausen bicartesian fibration over $(\text{Sch}_S^{\text{op}}, \text{fflat})$ as input. The symmetric monoidal bicartesian fibration is here the fibered category of finite locally free sheaves with the external tensor product. One can equip it with the minimal Waldhausen structure, whose cofibrations are summand inclusions of locally free sheaves, but in fact the Waldhausen structure is not used to construct the unfurling.

where f is finite syntomic and equipped with a trivialization of its cotangent complex $\mathcal{L}_f$ in the K-theory space $K(Z)$. There is a symmetric monoidal forgetful functor $\mathbf{Corr}^{\text{fr}}(\text{Sch}_S) \rightarrow \mathbf{Corr}^{\text{flat}}(\text{Sch}_S)$ [15, Section 4.3.15], allowing us to regard $\eta: \mathcal{F}\text{Flat}_S \rightarrow \mathcal{V}\text{ect}_S$ as a morphism of commutative monoids in $\text{PSh}_\Sigma(\mathbf{Corr}^{\text{fr}}(\text{Sch}_S))$. Below we also denote by $\mathcal{F}\text{Flat}_S$ and $\mathcal{V}\text{ect}_S$ the restriction of these presheaves to $\mathbf{Corr}^{\text{fr}}(\text{Sm}_S)$, which is the full subcategory of $\mathbf{Corr}^{\text{fr}}(\text{Sm}_S)$ spanned by the smooth S-schemes.

Recall that any presheaf with framed transfers on Sm_S gives rise to a motivic spectrum via the functor

$$\Sigma_{\mathbf{T}, \text{fr}}^\infty: \text{PSh}_\Sigma(\mathbf{Corr}^{\text{fr}}(\text{Sm}_S)) \rightarrow \mathbf{SH}^{\text{fr}}(S) \simeq \mathbf{SH}(S),$$

where the equivalence is [35, Theorem 18].

Let $\beta: (\mathbf{P}^1, \infty) \rightarrow \mathcal{V}\text{ect}^{\text{gp}}$ be the morphism corresponding to the formal difference of locally free sheaves $\mathcal{O}_{\mathbf{P}^1} - \mathcal{O}_{\mathbf{P}^1}(-1)$. For any scheme S , β induces a canonical element in $\pi_{2,1}(\Sigma_{\mathbf{T}, \text{fr}}^\infty \mathcal{V}\text{ect}_S)$. Following [34, Section 3], we shall say that a $\mathcal{V}\text{ect}_S^{\text{gp}}$ -module $\mathcal{F}$ in some presentable ∞ -category with an action of $\text{PSh}_\Sigma(\mathbf{Corr}^{\text{fr}}(\text{Sm}_S))$ is β -periodic if the morphism

$$\mathcal{F} \rightarrow \Omega_{\mathbf{P}^1} \mathcal{F}$$

induced by β is an equivalence. For example, algebraic K-theory (as a pointed presheaf on Sm_S) is β -periodic, by the projective bundle formula.

Proposition 5.1. *For any scheme S , there is an equivalence of $\mathcal{E}_\infty$ -ring spectra*

$$\text{KGL}_S \simeq (\Sigma_{\mathbf{T}, \text{fr}}^\infty \mathcal{V}\text{ect}_S)[\beta^{-1}] \quad \text{in } \mathbf{SH}(S) \simeq \mathbf{SH}^{\text{fr}}(S).$$

Proof. By [34, Proposition 3.2], the functors $\Omega_{\mathbf{T}}^\infty$ and $\Omega_{\mathbf{T}, \text{fr}}^\infty$ restrict to equivalences of symmetric monoidal ∞ -categories of β -periodic modules

$$\begin{aligned} \text{P}_\beta \text{Mod}_{\mathcal{V}\text{ect}^{\text{gp}}}(\mathbf{H}(S)_*) &\simeq \text{P}_\beta \text{Mod}_{\mathcal{V}\text{ect}^{\text{gp}}}(\mathbf{SH}(S)), \\ \text{P}_\beta \text{Mod}_{\mathcal{V}\text{ect}^{\text{gp}}}(\mathbf{H}^{\text{fr}}(S)) &\simeq \text{P}_\beta \text{Mod}_{\mathcal{V}\text{ect}^{\text{gp}}}(\mathbf{SH}^{\text{fr}}(S)). \end{aligned}$$

Under the former equivalence, the $\mathcal{E}_\infty$ -ring spectrum KGL_S corresponds to the pointed motivic space $\Omega^\infty \text{KH}$ [11, Proposition 2.14], which is the β -periodization of $\text{L}_{\text{mot}} \mathcal{V}\text{ect}^{\text{gp}}$ [34, Proposition 4.10].

For $\mathcal{F}$ a $\mathcal{V}\text{ect}^{\text{gp}}$ -module in either $\mathbf{H}(S)_*$ or $\mathbf{H}^{\text{fr}}(S)$, its β -periodization $\text{P}_\beta \mathcal{F}$ is computed as a sequential colimit of the presheaves $\Omega_{\mathbf{T}}^n \mathcal{F}$ [34, Theorem 3.8 and Lemma 4.9]. Since the forgetful functor $\mathbf{H}^{\text{fr}}(S) \rightarrow \mathbf{H}(S)_*$ commutes with sequential colimits and with $\Omega_{\mathbf{T}}^n$, the square

$$\begin{array}{ccc} \text{Mod}_{\mathcal{V}\text{ect}^{\text{gp}}}(\mathbf{H}^{\text{fr}}(S)) & \xrightarrow{\text{P}_\beta} & \text{P}_\beta \text{Mod}_{\mathcal{V}\text{ect}^{\text{gp}}}(\mathbf{SH}^{\text{fr}}(S)) \\ \downarrow & & \downarrow \\ \text{Mod}_{\mathcal{V}\text{ect}^{\text{gp}}}(\mathbf{H}(S)_*) & \xrightarrow{\text{P}_\beta} & \text{P}_\beta \text{Mod}_{\mathcal{V}\text{ect}^{\text{gp}}}(\mathbf{SH}(S)) \end{array}$$

of right-lax symmetric monoidal functors commutes. The commutativity of this square on the framed motivic space $\text{L}_{\text{mot}} \mathcal{V}\text{ect}^{\text{gp}}$ gives the desired equivalence. $\blacksquare$

Let $\mathrm{kgl}_S \in \mathbf{SH}(S)^{\mathrm{veff}}$ denote the very effective cover of KGL_S [62, Definition 5.5]. We shall only consider the spectrum kgl_S when S is regular over a field (i.e., regular and equicharacteristic), in which case it coincides with the effective cover of KGL_S (this is equivalent to the vanishing of the negative K-theory of fields, by the characterization of very effectivity in terms of homotopy sheaves [2, Section 3]).

Corollary 5.2. *If S is regular over a field, there is an equivalence of $\mathcal{E}_\infty$ -ring spectra*

$$\mathrm{kgl}_S \simeq \Sigma_{\mathrm{T},\mathrm{fr}}^\infty \mathcal{V}\mathrm{ect}_S \quad \text{in } \mathbf{SH}(S) \simeq \mathbf{SH}^{\mathrm{fr}}(S).$$

Proof. By Theorem 5.1, there is a canonical $\mathcal{E}_\infty$ -map $\Sigma_{\mathrm{T},\mathrm{fr}}^\infty \mathcal{V}\mathrm{ect}_S \rightarrow \mathrm{kgl}_S$. The assertion that it is an equivalence is local on S , so we may assume S affine. By Popescu's theorem [64, Tag 07GC], S is then a cofiltered limit of smooth k -schemes for some perfect field k . Since pro-smooth base change preserves very effective covers, we can assume that S is the spectrum of a perfect field k . In this case, by the motivic recognition principle [15, Theorem 3.5.14], the very effective cover of a motivic spectrum E is $\Sigma_{\mathrm{T},\mathrm{fr}}^\infty \Omega_{\mathrm{T},\mathrm{fr}}^\infty(E)$, and for X a framed motivic space we have $\Omega_{\mathrm{T},\mathrm{fr}}^\infty \Sigma_{\mathrm{T},\mathrm{fr}}^\infty X \simeq X^{\mathrm{gp}}$. Since $L_{\mathrm{mot}} \mathcal{V}\mathrm{ect}^{\mathrm{gp}}$ is already β -periodic on regular schemes (as it agrees with the K-theory presheaf K), we deduce that $\Sigma_{\mathrm{T},\mathrm{fr}}^\infty \mathcal{V}\mathrm{ect}_k$ is the very effective cover of $(\Sigma_{\mathrm{T},\mathrm{fr}}^\infty \mathcal{V}\mathrm{ect}_k)[\beta^{-1}]$, so the corollary follows from Proposition 5.1. ■

Remark 5.3. By [6, Proposition B.3], Corollary 5.2 holds for a finite-dimensional scheme S if and only if, for every $s \in S$, the canonical map $s^*(\mathrm{kgl}_S) \rightarrow \mathrm{kgl}_{\kappa(s)}$ is an equivalence. One expects this to be true at least if S is smooth over a Dedekind scheme.

Theorem 5.4. *If S is regular over a field, there is an equivalence of $\mathcal{E}_\infty$ -ring spectra*

$$\mathrm{kgl}_S \simeq \Sigma_{\mathrm{T},\mathrm{fr}}^\infty \mathcal{F}\mathcal{F}\mathrm{lat}_S \quad \text{in } \mathbf{SH}(S) \simeq \mathbf{SH}^{\mathrm{fr}}(S).$$

Proof. Combine Corollary 5.2 and Theorem 3.1. ■

Remark 5.5. One can lift the Bott element $\beta: (\mathbf{P}^1, \infty) \rightarrow \mathcal{V}\mathrm{ect}^{\mathrm{gp}}$ to $\mathcal{F}\mathcal{F}\mathrm{lat}^{\mathrm{gp}}$ (for example, by taking the formal difference of square-zero extensions $\mathcal{O} \oplus \mathcal{O} - \mathcal{O} \oplus \mathcal{O}(-1)$). Combining Proposition 5.1 and Theorem 3.1, we then have an equivalence of $\mathcal{E}_\infty$ -ring spectra

$$\mathrm{KGL}_S \simeq (\Sigma_{\mathrm{T},\mathrm{fr}}^\infty \mathcal{F}\mathcal{F}\mathrm{lat}_S)[\beta^{-1}]$$

for any scheme S .

Let $\mathbf{H}^{\mathrm{fflat}}(S) \subset \mathrm{PSh}_\Sigma(\mathbf{Corr}^{\mathrm{fflat}}(\mathrm{Sm}_S))$ be the full subcategory spanned by the $\mathbf{A}^1$ -invariant Nisnevich sheaves, and let $\mathbf{SH}^{\mathrm{fflat}}(S)$ be the ∞ -category of $\mathbf{T}$ -spectra in $\mathbf{H}^{\mathrm{fflat}}(S)_*$. Theorem 5.4 formally leads to an adjunction

$$\mathrm{Mod}_{\mathrm{kgl}}(\mathbf{SH}(S)) \rightleftarrows \mathbf{SH}^{\mathrm{fflat}}(S),$$

where the left adjoint is symmetric monoidal. In [3, Corollary 4.3], Bachmann proves that this adjunction induces an equivalence

$$\mathrm{Mod}_{\mathrm{kgl}}(\mathbf{SH}(k))[1/e] \simeq \mathbf{SH}^{\mathrm{fflat}}(k)[1/e],$$

for k a perfect field of exponential characteristic e .

The results in [3] (which in turn depend on Theorem 5.4) have the following consequence.

Proposition 5.6. *Let k be a perfect field of exponential characteristic e and X a smooth k -scheme. Then there is an equivalence*

$$\Omega_T^\infty(\mathrm{kgl} \otimes \Sigma_T^\infty X_+)[1/e] \simeq L_{\mathrm{Zar}} L_{\mathbf{A}^1} h^{\mathrm{fflat}}(X)^{\mathrm{gp}}[1/e].$$

If moreover $\mathrm{kgl} \otimes \Sigma_T^\infty X_+$ is a dualizable kgl -module (for example if X is proper), there is an equivalence

$$\Omega_T^\infty(\mathrm{kgl} \otimes \Sigma_T^\infty X_+) \simeq L_{\mathrm{Zar}} L_{\mathbf{A}^1} h^{\mathrm{fflat}}(X)^{\mathrm{gp}}.$$

Proof. By the cancellation theorem for finite locally free correspondences [3, Theorem 3.5 and Proposition 2.13], for every $\mathcal{F} \in \mathbf{H}^{\mathrm{fflat}}(k)$ there is a natural equivalence

$$\mathcal{F}^{\mathrm{gp}} \simeq \Omega_T^\infty \Sigma_T^\infty \mathcal{F}$$

in $\mathbf{H}^{\mathrm{fflat}}(k)$. Moreover, if $\mathcal{F} \in \mathrm{PSh}_\Sigma(\mathbf{Corr}^{\mathrm{fflat}}(\mathrm{Sm}_k))^{\mathrm{gp}}$, then $L_{\mathrm{mot}} \mathcal{F} \simeq L_{\mathrm{Zar}} L_{\mathbf{A}^1} \mathcal{F}$ by [15, Theorem 3.4.11]. The first statement now follows from the equivalence of ∞ -categories $\mathrm{Mod}_{\mathrm{kgl}}(\mathbf{SH}(k))[1/e] \simeq \mathbf{SH}^{\mathrm{fflat}}(k)[1/e]$ [3, Corollary 4.3]. Without inverting e , the unit of the adjunction $\mathrm{Mod}_{\mathrm{kgl}}(\mathbf{SH}(k)) \rightleftarrows \mathbf{SH}^{\mathrm{fflat}}(k)$ is an equivalence on the unit object [3, Theorem 4.2], hence on any dualizable object. This implies the second statement. ■

Corollary 5.7. *Let k be a perfect field and X a smooth separated k -scheme. Suppose that $\mathrm{kgl} \otimes \Sigma_T^\infty X_+$ is a dualizable kgl -module (for example X is proper). Then there is an equivalence*

$$\Omega_T^\infty(\mathrm{kgl} \otimes \Sigma_T^\infty X_+) \simeq (L_{\mathrm{Zar}} L_{\mathbf{A}^1} \mathrm{Hilb}(\mathbf{A}_X^\infty))^{\mathrm{gp}}.$$

Proof. Combine Propositions 5.6 and 4.2. ■

In light of Corollary 4.14, we can think of Corollary 5.7 as a geometric analogue of Segal's result [59, Proposition 1.1], expressing ko -homology of a topological space X as the homotopy groups of the labeled configuration space $F(X)$ (see Section 4). Here ko is the connective cover of the real topological K-theory spectrum KO .

Remark 5.8. Let k be a perfect field and $\mathrm{HZ} \in \mathbf{SH}(k)$ the motivic Eilenberg–Mac Lane spectrum over k . Since $s_0(\mathbf{1}) \simeq s_0(\mathrm{kgl}) \simeq \mathrm{HZ}$ [43], there is a *unique* morphism of $\mathcal{E}_\infty$ -ring spectra $\mathrm{kgl} \rightarrow \mathrm{HZ}$. Let $\mathrm{Cor}(k)$ be Voevodsky's category of finite correspondences over k [46, Definition 1.5]. By [15, Section 5.3], there is a symmetric monoidal functor

$$\mathbf{Corr}^{\mathrm{fflat}}(\mathrm{Sm}_k^{\mathrm{ft,sep}}) \rightarrow \mathrm{Cor}(k)$$

sending a span $X \xleftarrow{f} Z \xrightarrow{g} Y$ to the cycle $(f, g)_*[Z]$ on $X \times Y$. It induces an adjunction

$$\mathbf{SH}^{\mathrm{fflat}}(k) \rightleftarrows \mathbf{DM}(k),$$

where the left adjoint is symmetric monoidal. Recall that HZ is by definition the image of the unit object of $\mathbf{DM}(k)$ by the forgetful functor $\mathbf{DM}(k) \rightarrow \mathbf{SH}(k)$. The unit of the

above adjunction therefore induces the (unique) morphism of $\mathcal{E}_\infty$ -ring spectra $\mathrm{kgl} \rightarrow \mathrm{HZ}$, and we obtain a commutative square of adjunctions

$$\begin{array}{ccc} \mathrm{Mod}_{\mathrm{kgl}}(\mathbf{SH}(k)) & \xrightleftharpoons{\quad} & \mathrm{Mod}_{\mathrm{HZ}}(\mathbf{SH}(k)) \\ \downarrow \uparrow & & \downarrow \uparrow \\ \mathbf{SH}^{\mathrm{fflat}}(k) & \xrightleftharpoons{\quad} & \mathbf{DM}(k) \end{array}$$

The vertical adjunctions are known to be equivalences after inverting the exponential characteristic e (see [57] and [37, Theorem 5.8]). For $X \in \mathrm{Sm}_k^{\mathrm{ft}, \mathrm{sep}}$, the canonical map

$$\Omega_{\mathrm{T}}^\infty(\mathrm{kgl} \otimes \Sigma_{\mathrm{T}}^\infty X_+) \rightarrow \Omega_{\mathrm{T}}^\infty(\mathrm{HZ} \otimes \Sigma_{\mathrm{T}}^\infty X_+) \quad (5.9)$$

can thus be identified, after inverting e (or if X is proper), with the motivic localization of the map

$$h^{\mathrm{fflat}}(X)^{\mathrm{gp}} \rightarrow h^{\mathrm{Cor}}(X),$$

where $h^{\mathrm{Cor}}(X)$ is the presheaf represented by X on $\mathrm{Cor}(k)$. Furthermore, if X is quasi-projective, there is a commutative diagram

$$\begin{array}{ccc} \mathrm{Hilb}(\mathbf{A}_X^\infty) & \xrightarrow{\rho_{\mathbf{A}_X^\infty/X}} & \mathrm{Sym}(X)^{\mathrm{gp}} \\ \downarrow & & \uparrow \\ h^{\mathrm{fflat}}(X) & \longrightarrow & h^{\mathrm{Cor}}(X) \end{array}$$

where $\rho_{\mathbf{A}_X^\infty/X}$ is the Hilbert–Chow morphism (4.11), $\mathrm{Hilb}(\mathbf{A}_X^\infty) \rightarrow h^{\mathrm{fflat}}(X)$ is a motivic equivalence (Proposition 4.2), and $h^{\mathrm{Cor}}(X) \rightarrow \mathrm{Sym}(X)^{\mathrm{gp}}$ is an isomorphism after inverting e [63, Theorem 6.8]. Altogether, the map (5.9) corresponds, up to inverting e , to the group completion of the Hilbert–Chow morphism $\rho_{\mathbf{A}_X^\infty/X}: \mathrm{Hilb}(\mathbf{A}_X^\infty) \rightarrow \mathrm{Sym}(X)$.

6. Comparison with algebraic cobordism

Let $\mathcal{FSyn}_S$ be the presheaf with framed transfers on Sm_S associating to X the groupoid of finite syntomic X -schemes. The forgetful map $\mathcal{FSyn}_S \rightarrow \mathrm{Vect}_S$ induces a map of $\mathcal{E}_\infty$ -ring spectra

$$\Sigma_{\mathrm{T}, \mathrm{fr}}^\infty \mathcal{FSyn}_S \rightarrow \Sigma_{\mathrm{T}, \mathrm{fr}}^\infty \mathrm{Vect}_S.$$

By [14, Theorem 3.4.1], $\Sigma_{\mathrm{T}, \mathrm{fr}}^\infty \mathcal{FSyn}_S \simeq \mathrm{MGL}_S$ is the algebraic cobordism spectrum of Voevodsky. By Proposition 5.1, $(\Sigma_{\mathrm{T}, \mathrm{fr}}^\infty \mathrm{Vect}_S)[\beta^{-1}]$ is identified with KGL_S , so we get an $\mathcal{E}_\infty$ -map $\mathrm{MGL}_S \rightarrow \mathrm{KGL}_S$. The goal of this section is to prove that this map is an $\mathcal{E}_\infty$ -refinement of the usual orientation of KGL_S .

Let PMGL_S be the spectrum $\bigoplus_{n \in \mathbb{Z}} \Sigma^{2n, n} \mathrm{MGL}_S$ with the $\mathcal{E}_\infty$ -ring structure from [6, Theorem 16.19]. The commutative monoid structure on PMGL_S in the homotopy category $\mathbf{hSH}(S)$ is determined by that of MGL_S , and it classifies *periodic oriented ring spectra*, i.e., oriented ring spectra E with a given unit in $\pi_{2,1}(E)$.

By [14, Theorem 3.4.1], there is an equivalence of $(\mathbf{Z}$ -graded) $\mathcal{E}_\infty$ -ring spectra

$$(\Sigma_{\mathbf{T}, \text{fr}}^\infty \mathcal{FQSm}_S)[u^{-1}] \simeq \text{PMGL}_S, \quad (6.1)$$

where $u \in \pi_{2,1}(\Sigma_{\mathbf{T}, \text{fr}}^\infty \mathcal{FQSm}_S^1) \simeq \pi_{2,1}(\Sigma^{2,1} \text{MGL}_S)$ is the suspension of the unit of MGL_S . Here, $\mathcal{FQSm}_S(X)$ is the ∞ -groupoid of finite quasi-smooth derived X -schemes, with the $\mathcal{E}_\infty$ -semiring structure given by the sum and the product of X -schemes. Recall that a morphism of derived schemes $f: Y \rightarrow X$ is *quasi-smooth* if it is locally of finite presentation with perfect cotangent complex $\mathcal{L}_f$ of Tor-amplitude ≤ 1 , and *finite* if the underlying morphism of classical schemes is finite. Recall also that $f: Y \rightarrow X$ is quasi-smooth if and only if, locally on Y , it is the vanishing locus of finitely many functions on a smooth X -scheme [41, Section 2.3.13]; in particular, quasi-smooth morphisms are locally of finite Tor-amplitude.

Under the equivalence (6.1), Thom classes of vector bundles have the following geometric description. If $\mathcal{E}$ is a locally free sheaf of rank n over a smooth S -scheme X , its Thom class in $\text{MGL}_X^{2n,n}(\mathbf{V}(\mathcal{E}))$ is represented by the zero section $X \hookrightarrow \mathbf{V}(\mathcal{E})$ in $\mathcal{FQSm}_S^n(\mathbf{V}(\mathcal{E}))$. Indeed, this holds by construction of the equivalence $\Sigma_{\mathbf{T}, \text{fr}}^\infty \mathcal{FQSm}_S^n \simeq \Sigma^{2n,n} \text{MGL}_S$; cf. [14, Construction 3.1.1].

Proposition 6.2. *Let S be a scheme. The $\mathcal{E}_\infty$ -map $\text{MGL}_S \rightarrow \text{KGL}_S$ induced by $\mathcal{FSyn}_S \rightarrow \text{Vect}_S$ is the standard orientation of KGL_S .*

Proof. By [45, Theorem 6.1.3.2], the structure sheaf of any finite quasi-smooth S -scheme is a perfect $\mathcal{O}_S$ -module, since quasi-smooth morphisms are locally of finite presentation and locally of finite Tor-amplitude. There is therefore a commutative square of commutative monoids in presheaves with framed transfers

$$\begin{array}{ccc} \mathcal{FSyn}_S & \longrightarrow & \mathcal{FQSm}_S \\ \downarrow & & \downarrow \\ \text{Vect}_S & \longrightarrow & \text{Perf}_S \end{array}$$

where $\text{Perf}_S(X)$ is the ∞ -groupoid of perfect complexes on X and the right vertical morphism $\mathcal{FQSm}_S \rightarrow \text{Perf}_S$ forgets the algebra structure. By Proposition 5.1, there is an $\mathcal{E}_\infty$ -map of presheaves with framed transfers $\text{Vect}_S \rightarrow \Omega_{\mathbf{T}, \text{fr}}^\infty \text{KGL}_S$, which factors through $L_{\text{Nis}} \text{Vect}_S^{\text{gp}}$ (since Nisnevich sheafification is compatible with framed transfers [15, Proposition 3.2.4]). On the other hand, the algebraic K-theory presheaf $\mathbf{K}$ has finite locally free transfers induced by the pushforward functors between the ∞ -categories of perfect complexes, which extend the pushforwards of vector bundles. Since $\text{Vect}^{\text{gp}} \rightarrow \mathbf{K}$ is a Nisnevich-local equivalence, we deduce that the $\mathcal{E}_\infty$ -map $\text{Vect}_S \rightarrow L_{\text{Nis}} \text{Vect}_S^{\text{gp}}$ factors through Perf_S in $\text{PSh}_\Sigma(\mathbf{Corr}^{\text{fr}}(\text{Sm}_S))$. Hence, the composite $\mathcal{FSyn}_S \rightarrow \text{Vect}_S \rightarrow \Omega_{\mathbf{T}, \text{fr}}^\infty \text{KGL}_S$ factors through $\mathcal{FQSm}_S$. By adjunction and [14, Theorem 3.4.1], we obtain a sequence of $\mathcal{E}_\infty$ -maps

$$\text{MGL}_S \rightarrow \bigoplus_{n \geq 0} \Sigma^{2n,n} \text{MGL}_S \rightarrow \text{KGL}_S.$$

Recall that the standard periodic orientation of KGL_S is such that the Thom class of a finite locally free sheaf $\mathcal{E}$ over X is represented by the perfect $\mathcal{O}_{V(\mathcal{E})}$ -module $\mathcal{O}_X$, which is precisely the image of the zero section $X \hookrightarrow V(\mathcal{E})$ by the forgetful map $\mathcal{FQSm}_S(V(\mathcal{E})) \rightarrow \mathcal{P}erf_S(V(\mathcal{E}))$. Hence, the map $\Sigma^{2n,n}\mathrm{MGL}_S \rightarrow \mathrm{KGL}_S$ constructed above preserves Thom classes of locally free sheaves of rank n . In particular, $\Sigma^{2,1}\mathrm{MGL}_S \rightarrow \mathrm{KGL}_S$ sends u to the Bott element β , which is the Thom class of $\mathcal{O}_S$, and we obtain an induced $\mathcal{E}_\infty$ -map

$$\mathrm{PMGL}_S \rightarrow \mathrm{KGL}_S.$$

This map preserves Thom classes by construction, i.e., it induces the standard periodic orientation of KGL_S . $\blacksquare$

Remark 6.3. For $S = \mathrm{Spec} \mathbf{C}$, the $\mathcal{E}_\infty$ -map $\mathrm{PMGL}_S \rightarrow \mathrm{KGL}_S$ constructed above realizes to the folklore $\mathcal{E}_\infty$ -map $\mathrm{MUP} \rightarrow \mathrm{KU}$ alluded to in [26, p. 1841]. In particular, it differs from the Gepner–Snaith $\mathcal{E}_\infty$ -map $\mathrm{PMGL}_S \rightarrow \mathrm{KGL}_S$ constructed in [21], which uses a different $\mathcal{E}_\infty$ -ring structure on PMGL_S [6, Remark 16.20].

Corollary 6.4. *Suppose that S is regular over a field. Under the equivalence $\mathrm{kg}l_S \simeq \Sigma_{\mathrm{T},\mathrm{fr}}^\infty \mathcal{F}\mathrm{Flat}_S$ of Theorem 5.4, the standard orientation $\mathrm{MGL}_S \rightarrow \mathrm{kg}l_S$ is $\Sigma_{\mathrm{T},\mathrm{fr}}^\infty$ of the forgetful map $\mathcal{F}\mathrm{Syn}_S \rightarrow \mathcal{F}\mathrm{Flat}_S$.*

7. The Hilbert scheme of finite lci schemes of degree 3

The open subset of lci subschemes in the Hilbert scheme has a rich homotopy type. In particular, in positive characteristic, the limiting space $\mathrm{Hilb}_\infty^{\mathrm{lci}}(\mathbf{A}^\infty)$ has the $\mathbf{A}^1$ -homotopy type of the unit component of the infinite loop space associated to algebraic cobordism, $\Omega_{\mathrm{T}}^\infty \mathrm{MGL}$ [5, Theorem 1.2]. In characteristic zero, these two spaces at least have isomorphic motives in the derived category of motives, $\mathbf{DM}(k)$. Hopkins conjectured that the motive of $\Omega_{\mathrm{T}}^\infty \mathrm{MGL}$ is pure Tate, but that remains open. By definition, a pure Tate motive is a direct sum of the motives $\mathbf{Z}\{a\} := \mathbf{Z}(a)[2a]$ for integers a .

A possible approach to Hopkins’ conjecture is through the space $\mathrm{Hilb}_\infty^{\mathrm{lci}}(\mathbf{A}^\infty)$. In that direction, we have the following result.

Theorem 7.1. *The motive of $\mathrm{Hilb}_3^{\mathrm{lci}}(\mathbf{A}^\infty)$ over $\mathbf{Z}$ is pure Tate in $\mathbf{DM}(\mathbf{Z})$. The Chow ring of $\mathrm{Hilb}_3^{\mathrm{lci}}(\mathbf{A}^\infty)$ is*

$$\mathbf{Z}[c_1, c_2]/(c_2(-2c_1^2 + 9c_2)),$$

where $|c_i| = i$.

Remark 7.2. As observed in [5, Remark 1.8], the motive of $\mathrm{Hilb}_3^{\mathrm{lci}}(\mathbf{A}^n)$ is *not* pure Tate for any finite $n \geq 2$.

Here, $\mathbf{DM}(\mathbf{Z})$ denotes Spitzweck’s ∞ -category of motives over $\mathrm{Spec} \mathbf{Z}$ [61, Chapter 9]. If S is a regular scheme of dimension ≤ 1 , then the ∞ -category $\mathbf{DM}(S)$ categorifies Bloch–Levine motivic cohomology, in the sense that the presheaf

$X \mapsto \text{Maps}_{\mathbf{DM}(S)}(M(X), \mathbf{Z}\{n\})$ on smooth S -schemes is the Zariski sheafification of Bloch's cycle complex $X \mapsto z^n(X, *)$. By [20, Theorem 1.2 (5)], we have

$$\pi_0 \text{Maps}_{\mathbf{DM}(S)}(\mathbf{Z}\{a\}, \mathbf{Z}\{b\}) = \begin{cases} \mathbf{Z} & \text{if } a = b, \\ \text{Pic}(S) & \text{if } a + 1 = b, \\ 0 & \text{otherwise.} \end{cases}$$

In particular, the homotopy category of pure Tate motives over $\text{Spec } \mathbf{Z}$ is equivalent to the category of free graded abelian groups. For $X \in \text{PSh}(\text{Sm}_S)$, we will write

$$\text{CH}^a(X) = H^{2a}(X, \mathbf{Z}(a)) = \pi_0 \text{Maps}_{\mathbf{DM}(S)}(M(X), \mathbf{Z}\{a\}).$$

These are the classical Chow groups when S is semilocal and X is a smooth S -scheme of finite type (since Bloch's cycle complex is already a Zariski sheaf in this case [42, Theorem 1.7]). By [61, Proposition 10.1] and [52, Proposition 6.2], we have a canonical isomorphism

$$\text{CH}^*(\text{BGL}_n) \simeq \mathbf{Z}[c_1, \dots, c_n]$$

with c_i in degree i .

Proof of Theorem 7.1. Let $X_n = \text{Hilb}_3(\mathbf{A}^n)$ over $\mathbf{Z}$, and let $Y_n \subset X_n$ be the closed complement of $\text{Hilb}_3^{\text{lc}}(\mathbf{A}^n)$. Let $X = \text{colim}_n X_n$ and $Y = \text{colim}_n Y_n$; we want to describe the motive of $X - Y = \text{colim}_n (X_n - Y_n)$. For an algebraically closed field k , every non-lci finite k -scheme of degree 3 is isomorphic to $\text{Spec } k[x, y]/(x, y)^2$ [53, Table 1], and any two embeddings into $\mathbf{A}_k^n$ differ by an affine transformation. It follows that the obvious morphism $f: W_n := \text{Gr}_2(\mathbf{A}^n) \times \mathbf{A}^n \rightarrow X_n$ is injective on k -points with image $Y_n(k)$ for every algebraically closed field k . In particular, f is quasi-finite; by definition, it suffices to check this over fields, and then it suffices to prove this over algebraically closed fields by descent [64, Tag 02VI].

Also, f is proper, as it is the restriction over $\text{Hilb}_3(\mathbf{A}^n)$ of a $\text{Gr}_2(\mathbf{A}^n)$ -bundle over $\mathbf{P}^n$ mapping to $\text{Hilb}_3(\mathbf{P}^n)$. Since f is proper and quasi-finite, it is finite [64, Tag 02OG]. It is also unramified, meaning that the sheaf of relative differentials Ω_{W_n/X_n}^1 is zero. Indeed, the sheaf is coherent, and so this can be checked over fields by Nakayama's lemma. After reducing further to an algebraically closed base field, one can check that f is injective on tangent spaces using the calculation below of the tangent space to the Hilbert scheme X_n ; so f is unramified. Since f is finite, unramified, and injective on points over algebraically closed fields, f is a closed immersion [64, Tag 04DG]. That is, Y_n with reduced scheme structure is isomorphic to $\text{Gr}_2(\mathbf{A}^n) \times \mathbf{A}^n$.

By Corollary 4.9, the inclusion $Y \rightarrow X$ is a motivic equivalence. Both Y_n and X_n are smooth over $\mathbf{Z}$, with Y_n of codimension 4 in X_n . So we have a cofiber sequence

$$M(X - Y) \rightarrow M(X) \rightarrow M(Y)\{4\}$$

in $\mathbf{DM}(\mathbf{Z})$, where $M(Y)\{4\}$ and $M(X)$ are pure Tate. So, to show that $M(X - Y)$ is pure Tate as well, it suffices to show that the morphism $M(X) \rightarrow M(Y)\{4\}$ is the projection

onto a summand. A morphism between pure Tate motives is determined by what it does on Chow groups CH^* , so it suffices to show that the pushforward homomorphism

$$\mathrm{CH}^{*-4}(Y) \rightarrow \mathrm{CH}^*(X)$$

is a split injection of abelian groups. Since the restriction $\mathrm{CH}^*(X) \rightarrow \mathrm{CH}^*(Y)$ is an isomorphism, it suffices to show that the composed map

$$\mathrm{CH}^{*-4}(Y) \rightarrow \mathrm{CH}^*(X) \rightarrow \mathrm{CH}^*(Y)$$

is split injective. This composition is multiplication by the top Chern class of the normal bundle, $c_4(N_{Y/X})$ [18, Corollary 6.3]. So it suffices to show that this class is nonzero modulo every prime number p in $\mathrm{CH}^*(Y) = \mathrm{CH}^*(\mathrm{BGL}_2) = \mathbf{Z}[c_1, c_2]$. (Indeed, then multiplication by $c_4(N_{Y/X})$ is injective modulo p for every prime number p , and (looking at each graded piece) a homomorphism of finitely generated free abelian groups which is injective modulo every prime is split injective.) We will show that $c_4(N_{Y/X})$ is $c_2(-2c_1^2 + 9c_2)$, which will complete the proof.

Let U_n be the open subset of X_n of degree 3 subschemes not contained in any affine line. Then Y_n is contained in U_n , and so the normal bundle of Y_n in X_n can also be viewed as its normal bundle in U_n . The point is that each point of U_n , viewed as a degree 3 subscheme of $\mathbf{A}^n$, spans an affine plane, and so U_n is fibered over the scheme of affine planes in $\mathbf{A}^n$ with fiber U_2 . More precisely, let Aff_n be the group scheme of affine automorphisms of $\mathbf{A}^n$, and let $A(m, n)$ be the scheme of affine embeddings $\mathbf{A}^m \hookrightarrow \mathbf{A}^n$. Then Aff_n acts on X_n , the subschemes U_n and Y_n are Aff_n -invariant, and the closed immersion $Y_n \hookrightarrow U_n$ can be identified with

$$Y_2 \times_{\mathrm{Aff}_2} A(2, n) \hookrightarrow U_2 \times_{\mathrm{Aff}_2} A(2, n).$$

Hence, the normal bundle of $Y_n \hookrightarrow X_n$ is extended from the Aff_2 -equivariant normal bundle of $Y_2 \hookrightarrow U_2$. In other words, it is the pullback of N_{Y_2/X_2} by the projection

$$Y_n \simeq Y_2 \times_{\mathrm{Aff}_2} A(2, n) \rightarrow [Y_2/\mathrm{Aff}_2].$$

Since $Y_2 \simeq \mathbf{A}^2$, there is an equivalence of stacks $[Y_2/\mathrm{Aff}_2] \simeq \mathrm{BGL}_2$. Moreover, the induced map $Y = \mathrm{colim}_n Y_n \rightarrow \mathrm{BGL}_2$ is a motivic equivalence, inducing the canonical motivic equivalence $\mathrm{Gr}_2 \rightarrow \mathrm{BGL}_2$ (by Proposition 4.7). So our goal is to compute the top Chern class of the GL_2 -representation given by the (rank 4) normal bundle to Y_2 in $X_2 = \mathrm{Hilb}_3(\mathbf{A}^2)$ at the $\mathbf{Z}$ -point $S = \mathrm{Spec} \mathbf{Z}[x, y]/(x^2, xy, y^2) \subset \mathbf{A}^2$ in Y_2 .

Classically [60, Section 3.2.1] the tangent bundle to $\mathrm{Hilb}_3(\mathbf{A}^2)$ at $[S]$ is

$$H^0(S, N_{S/\mathbf{A}^2}) = \mathrm{Hom}_{\mathcal{O}_S}(I/I^2, \mathcal{O}_S),$$

where $I = (x^2, xy, y^2) \subset \mathbf{Z}[x, y]$ and $\mathcal{O}_S = \mathbf{Z}[x, y]/I$. By direct inspection, this $\mathbf{Z}$ -module is free of rank 6 and isomorphic to

$$\mathrm{Hom}_{\mathbf{Z}}(\mathbf{Z}\{x^2, xy, y^2\}, \mathbf{Z}\{x, y\}) = \mathrm{Hom}(S^2(V^*), V^*),$$

where $V = \mathbf{A}^2$, so V^* is the space of linear functions on $\mathbf{A}^2$, with basis x, y .

To compute the Chern classes of this representation of GL_2 , we can work over $\mathbf{C}$, so GL_2 becomes linearly reductive. The Clebsch–Gordan formula implies that the tangent space of $X_2 = \mathrm{Hilb}_3(\mathbf{A}^2)$ at $[S]$ is $V \oplus (S^3(V) \otimes \det(V)^*)$ as a GL_2 -representation [10, Proposition II.5.5]. In particular, its only 2-dimensional subrepresentation is V , which thus coincides with the tangent space to Y_2 . So the normal space to Y_2 in X_2 at $[S]$ is $S^3(V) \otimes \det(V)^*$ as a representation of GL_2 . The top Chern class c_4 of $S^3(V) \otimes \det(V)^*$ is $c_2(-2c_1^2 + 9c_2)$ in $\mathrm{CH}^*(\mathrm{BGL}_2) = \mathbf{Z}[c_1, c_2]$, as one checks by the splitting principle (the fact that the Chow ring of BGL_2 injects into that of $\mathrm{BG}_{\mathrm{m}}^2$). That is what we want. ■

8. Stability theorems for the Hilbert scheme

We have shown that $\mathrm{Hilb}_d(\mathbf{A}^\infty)$ has the $\mathbf{A}^1$ -homotopy type of BGL_{d-1} . We now prove corresponding stability theorems for the inclusion $\mathrm{Hilb}_d(\mathbf{A}^n) \subset \mathrm{Hilb}_d(\mathbf{A}^\infty)$. There are several notions of connectivity available in $\mathbf{A}^1$ -homotopy theory, and we shall prove that the inclusion $\mathrm{Hilb}_d(\mathbf{A}^n) \subset \mathrm{Hilb}_d(\mathbf{A}^\infty)$ is highly connected in various senses (Theorem 8.2). In particular, we show that it induces an isomorphism in motivic cohomology in weights $\leq n - d + 1$ (Corollary 8.3).

Remark 8.1. We use the convention that, for $n \geq -1$, a morphism f in Spc is n -connected if its fibers are n -connected (meaning that $\pi_i = 0$ for $i \leq n$). It follows, in particular, that f is surjective on π_0 . In the older literature, such a map would be called $(n + 1)$ -connected. Our numbering is becoming more common, as in [49, Definition 6.55] or [56, Remark 1.6].

We will say that a morphism $f: Y \rightarrow X$ in $\mathrm{PSh}(\mathrm{Sm}_S)$ is $\mathbf{A}^1$ -*n-connected* if $L_{\mathrm{mot}}(f)$ is n -connected as a morphism of Nisnevich sheaves (i.e., n -connected on Nisnevich stalks). We will frequently use Morel’s $\mathbf{A}^1$ -connectivity theorem [49, Theorem 6.38], which states that if k is a perfect field and $X \in \mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_k)$ is n -connected, then X is $\mathbf{A}^1$ - n -connected. Recall also that a motivic spectrum is *very n-effective* if it can be obtained using colimits and extensions from the image of the functor $\Sigma_{\mathbf{T}}^n \Sigma_{\mathbf{T}}^\infty$.

Theorem 8.2. *Let $n \geq d \geq 0$.*

(1) *The morphism*

$$\mathrm{Hilb}_d(\mathbf{A}^n)(\mathbf{C}) \rightarrow \mathrm{Hilb}_d(\mathbf{A}^\infty)(\mathbf{C})$$

is $(2n - 2d + 2)$ -connected and the morphism

$$\mathrm{Hilb}_d(\mathbf{A}^n)(\mathbf{R}) \rightarrow \mathrm{Hilb}_d(\mathbf{A}^\infty)(\mathbf{R})$$

is $(n - d)$ -connected.

(2) *If k is a field, the morphism*

$$\Sigma \mathrm{Hilb}_d(\mathbf{A}_k^n) \rightarrow \Sigma \mathrm{Hilb}_d(\mathbf{A}_k^\infty)$$

is $\mathbf{A}^1$ -($n - d + 1$)-connected. Here, ΣX denotes the pushout $ \sqcup_X *$ in $\mathrm{PSh}(\mathrm{Sm}_k)$.*

(3) If k is a field, the cofiber of

$$\Sigma_{\mathbf{T}}^{\infty} \mathrm{Hilb}_d(\mathbf{A}_k^n)_+ \rightarrow \Sigma_{\mathbf{T}}^{\infty} \mathrm{Hilb}_d(\mathbf{A}_k^{\infty})_+$$

in $\mathbf{SH}(k)$ is very $(n - d + 2)$ -effective.

The proof of Theorem 8.2 is given at the end of this section.

Corollary 8.3. *Let k be a field and A an abelian group.*

(1) *The homomorphism of motivic cohomology groups*

$$H^*(\mathrm{BGL}_{d-1}, A(a)) \rightarrow H^*(\mathrm{Hilb}_d(\mathbf{A}^n), A(a))$$

is an isomorphism for all $a \leq n - d + 1$.

(2) *Let $i \geq 0$. The homomorphism of Milnor–Witt motivic cohomology groups*

$$H^{2a+i}(\mathrm{BGL}_{d-1}, \tilde{A}(a)) \rightarrow H^{2a+i}(\mathrm{Hilb}_d(\mathbf{A}^n), \tilde{A}(a))$$

is an isomorphism if $a + i \leq n - d$ and injective if $a + i = n - d + 1$.

Proof. Both statements follow from Theorem 8.2 (3) and the following facts: if $E \in \mathbf{SH}(k)$ is very r -effective, then $H^*(E, A(a)) = 0$ for $a < r$ and $H^{2a+i}(E, \tilde{A}(a)) = 0$ for $i \geq 0$ and $a + i < r$. The former is because the 1-effective cover of the motivic Eilenberg–MacLane spectrum HA is zero, since motivic cohomology vanishes in negative weights. The latter is because the very 1-effective cover of $H\tilde{A}$ is zero, since in negative weights Milnor–Witt motivic cohomology is given by the cohomology of the Witt sheaf, which vanishes in negative degrees [4, Chapter 2, Example 6.1.5]. ■

Remark 8.4. In Theorem 8.2 (2, 3) and in Corollary 8.3, we regard $\mathrm{Hilb}_d(\mathbf{A}_k^n)$ as a presheaf on smooth k -schemes. Since $\mathrm{Hilb}_d(\mathbf{A}_k^n)$ is in fact a scheme, one can also define its motivic cohomology and Milnor–Witt motivic cohomology more intrinsically by working in $\mathbf{SH}(\mathrm{Hilb}_d(\mathbf{A}_k^n))$. By standard arguments using resolution of singularities or alterations, these two ways of defining the cohomology of nonsmooth schemes are known to agree with $\mathbf{Z}[1/e]$ -linear coefficients, where e is the exponential characteristic of k .

Let $\mathcal{Z}_d \rightarrow \mathcal{F}\mathrm{Flat}_d$ be the universal finite locally free scheme of degree d and let $\mathcal{V}_d(\mathbf{A}^n)$ be the vector bundle over $\mathcal{F}\mathrm{Flat}_d$ defined by

$$\mathcal{V}_d(\mathbf{A}^n) = \mathrm{Hom}_{\mathcal{F}\mathrm{Flat}_d}(\mathcal{Z}_d, \mathbf{A}_{\mathcal{F}\mathrm{Flat}_d}^n).$$

Then $\mathrm{Hilb}_d(\mathbf{A}^n) \subset \mathcal{V}_d(\mathbf{A}^n)$ is the open substack of closed immersions $\mathcal{Z}_d \rightarrow \mathbf{A}_{\mathcal{F}\mathrm{Flat}_d}^n$. The key geometric input to all our stability theorems is the following observation:

Lemma 8.5. *The closed complement of $\mathrm{Hilb}_d(\mathbf{A}^n)$ in $\mathcal{V}_d(\mathbf{A}^n)$ has codimension at least $n - d + 2$ in every fiber over $\mathcal{F}\mathrm{Flat}_d$.*

Proof. Let k be a field and $S = \mathrm{Spec} R$ a finite k -scheme of degree d . We must show that the closed complement of $\mathrm{Emb}_k(S, \mathbf{A}_k^n)$ in $\mathrm{Hom}_k(S, \mathbf{A}_k^n) \simeq \mathbf{A}_k^{nd}$ has codimension at

least $n - d + 2$. A k -morphism $S \rightarrow \mathbf{A}_k^n$ can be viewed as a k -algebra homomorphism $k[x_1, \dots, x_n] \rightarrow R$, or as a k -linear map $k\{x_1, \dots, x_n\} \rightarrow R$. Such a morphism $S \rightarrow \mathbf{A}_k^n$ is an embedding if and only if the corresponding homomorphism $k[x_1, \dots, x_n] \rightarrow R$ is surjective. In particular, $\text{Emb}_k(S, \mathbf{A}_k^n)$ contains the open subset of k -linear maps $k\{x_1, \dots, x_n\} \rightarrow R$ such that $k\{x_1, \dots, x_n\} \rightarrow R/(k \cdot 1)$ is surjective. The subset of non-surjective linear maps $k^n \rightarrow k^r$ has codimension $n - r + 1$; indeed, the space of all linear maps has dimension nr , whereas any nonsurjective map lands in some hyperplane in k^r (corresponding to a point in $\mathbf{P}^{r-1}$), and so the subset of nonsurjective maps has dimension $n(r - 1) + (r - 1) = nr - (n - r + 1)$. Thus the complement of $\text{Emb}_k(S, \mathbf{A}_k^n)$ in $\text{Hom}_k(S, \mathbf{A}_k^n)$ has codimension at least $n - (d - 1) + 1 = n - d + 2$, as we want. ■

Remark 8.6. The bound in Lemma 8.5 is optimal for $n \geq d - 2$, the worst case being the square-zero extension algebra

$$R = k[x_1, \dots, x_{d-1}]/(x_i x_j : \text{all } i, j)$$

over a field k . In that case, a k -algebra homomorphism $k[x_1, \dots, x_n] \rightarrow R$ is surjective if and only if the k -linear map $k\{x_1, \dots, x_n\} \rightarrow R/(k \cdot 1)$ is surjective. As a result, the subset of nonsurjective k -algebra homomorphisms $k[x_1, \dots, x_n] \rightarrow R$ has codimension equal to $n - (d - 1) + 1 = n - d + 2$.

Proposition 8.7. *Let k be a subfield of $\mathbf{C}$, X an algebraic stack of finite type over k , $V \rightarrow X$ a vector bundle, and $r \geq 0$. Let $U \subset V$ be an open substack whose complement has codimension at least r in every fiber of $V \rightarrow X$. Then $U(\mathbf{C}) \rightarrow X(\mathbf{C})$ is $(2r - 2)$ -connected in the classical topology. If moreover $k \subset \mathbf{R}$ and X is an algebraic space, then $U(\mathbf{R}) \rightarrow X(\mathbf{R})$ is $(r - 2)$ -connected in the classical topology.*

Proof. Let $X_\bullet \rightarrow X$ be a smooth hypercover where each X_i is an affine scheme. Then $X(\mathbf{C})$ is the colimit of $X_\bullet(\mathbf{C})$ in Spc . If X is an algebraic space, we can choose $X_\bullet \rightarrow X$ to be a Nisnevich hypercover by [55, Proposition 5.7.6], in which case also $X(\mathbf{R})$ is the colimit of $X_\bullet(\mathbf{R})$ in Spc . Since simplicial colimits preserve connectivity, we can assume that X is an affine scheme.

We give the rest of the proof for $\mathbf{C}$, but exactly the same argument applies to $\mathbf{R}$. Let $0 \leq j < 2r$; we need to show that $\pi_j U(\mathbf{C}) \rightarrow \pi_j X(\mathbf{C})$ is surjective (for any base points u_0 in $U(\mathbf{C})$ and x_0 in $X(\mathbf{C})$), and that $\pi_{j-1} U(\mathbf{C}) \rightarrow \pi_{j-1} X(\mathbf{C})$ is injective. Let h be a continuous map $S^j \rightarrow X(\mathbf{C})$; we will show that h has arbitrarily small perturbations that lift to $U(\mathbf{C})$. Although the perturbations we construct need not keep the base point fixed, they will send the base point s_0 of S^j near u_0 in $U(\mathbf{C})$; this implies the surjectivity of $\pi_j(U(\mathbf{C}), u_0) \rightarrow \pi_j(X(\mathbf{C}), x_0)$, since $U(\mathbf{C})$ is locally contractible.

By the triangulation of real semialgebraic sets, there are semialgebraic triangulations of S^j and $X(\mathbf{C})$ [30, Section 1]. By simplicial approximation, after subdividing the triangulation of S^j , we can perturb h to a simplicial map (still called h), which in particular is semialgebraic. By [16, Theorem 6], we can perturb h to a $\mathbf{C}^1$ semialgebraic map. (This makes sense even though $X(\mathbf{C})$ is singular, using local embeddings of $X(\mathbf{C})$ into affine space. Any two embeddings are equivalent by complex analytic maps, and so the notion

of “ C^1 ” is well-defined.) The pullback of a C^1 vector bundle on $X(C)$ along h is a C^1 vector bundle on S^j , for example by considering transition functions: a composite of C^1 maps is C^1 . As a result, the pullback of the map $U(C) \rightarrow X(C)$ to S^j is an open subset of a C^1 semialgebraic real vector bundle $p: W \rightarrow S^j$, with complement a semialgebraic closed subset $Z \subset W$ that has real codimension at least $2r$ in every fiber. It follows that Z has real codimension at least $2r$ in W . Here W is a C^1 manifold, and Z is a finite union of (possibly noncompact) C^1 submanifolds of real codimension at least $2r$. (The smooth locus of Z is a (possibly noncompact) C^1 manifold, and its complement has lower dimension; so we can decompose Z by induction on dimension.) Choose a C^1 section e of the vector bundle $W \rightarrow S^j$ such that $e(s_0) = u_0$. Since $j < 2r$, the transversality theorem for C^1 manifolds [31, Theorem 3.2.1] implies that e can be perturbed to a C^1 -section disjoint from Z . That is, the map $h: S^j \rightarrow X(C)$ lifts to $U(C)$, as we want. The same argument (transversality applied to the pullback vector bundle) also proves, for $j < 2r$, that every continuous map $S^{j-1} \rightarrow U(C)$ such that the composed map to $X(C)$ extends to D^j has arbitrarily small perturbations that extend to a continuous map $D^j \rightarrow U(C)$. So $U(C) \rightarrow X(C)$ is $(2r - 2)$ -connected, as we want. ■

Lemma 8.8. *Let k be a perfect field, $f: Y \rightarrow X$ a morphism in $\text{PSh}(\text{Sm}_k)$, and $n \geq -1$. If f is $\mathbf{A}^1$ - n -connected, then $\text{cofib}(f)$ is $\mathbf{A}^1$ -($n + 1$)-connected. The converse holds if X and Y are $\mathbf{A}^1$ -1-connected.*

Proof. Let C be the cofiber of $L_{\text{mot}}(f)$ in $\text{Shv}_{\text{Nis}}(\text{Sm}_k)$. The class of n -connected maps is closed under cobase change [44, Corollary 6.5.1.17]. Therefore, if $L_{\text{mot}}(f)$ is n -connected, then $* \rightarrow C$ is n -connected, which is equivalent to C being $(n + 1)$ -connected [44, Proposition 6.5.1.20]. By the $\mathbf{A}^1$ -connectivity theorem, C is $\mathbf{A}^1$ -($n + 1$)-connected. Conversely, suppose that X and Y are $\mathbf{A}^1$ -1-connected and that C is $\mathbf{A}^1$ -($n + 1$)-connected. Let F be the fiber of $L_{\text{mot}}(f)$. We will prove that C is in fact $(n + 1)$ -connected, which implies (by considering the Nisnevich stalks) that F is n -connected [67, Theorem IV.7.13]. Note that C is 1-connected since $L_{\text{mot}}(X)$ and $L_{\text{mot}}(Y)$ are. By induction, we may assume that C is n -connected and F is $(n - 1)$ -connected. By Blakers–Massey, the canonical map $F \rightarrow \Omega C$ is then n -connected. In particular, $\pi_{n+1}(C)$ is isomorphic to $\pi_n(F)$, hence is strictly $\mathbf{A}^1$ -invariant. By [49, Corollary 6.60], we have $\pi_{n+1}(C) \simeq \pi_{n+1}L_{\text{mot}}(C) = 0$, as desired. ■

Lemma 8.9. *Let k be a perfect field, X a smooth k -scheme, and $Z \subset X$ a closed subscheme of codimension $\geq r$. Then $\Sigma(X/(X - Z))$ is $\mathbf{A}^1$ - r -connected.*

Proof. We can assume X quasi-compact. If Z is smooth, then $X/(X - Z)$ is $\mathbf{A}^1$ -($r - 1$)-connected by the purity isomorphism [50, Section 3, Theorem 2.23]. In general, the result is trivial if $r = 0$. If $r \geq 1$, then $X/(X - Z)$ is $\mathbf{A}^1$ -connected by [48, Lemma 6.1.4]. We can therefore assume $r \geq 2$. Since k is perfect, there exists a filtration

$$\emptyset = Z_0 \subset Z_1 \subset \cdots \subset Z_n = Z$$

by closed subschemes such that $Z_i - Z_{i-1}$ is smooth. We prove the result by induction on the length n of the filtration, the case $n = 0$ being trivial. We therefore assume that

$\Sigma(X/(X - Z_{n-1}))$ is $\mathbf{A}^1$ - r -connected. Since $\Sigma(X/(X - Z))$ and $\Sigma((X - Z_{n-1})/(X - Z))$ are $\mathbf{A}^1$ -1-connected, Lemma 8.8 and the cofiber sequence

$$\frac{X - Z_{n-1}}{X - Z} \rightarrow \frac{X}{X - Z} \rightarrow \frac{X}{X - Z_{n-1}}$$

imply that the morphism $\Sigma((X - Z_{n-1})/(X - Z)) \rightarrow \Sigma(X/(X - Z))$ is $\mathbf{A}^1$ -($r - 1$)-connected. Since the source is $\mathbf{A}^1$ - r -connected (by the smooth case), it follows that $\Sigma(X/(X - Z))$ is $\mathbf{A}^1$ - r -connected. ■

Proposition 8.10. *Let k be a perfect field, $X: \mathrm{Sm}_k^{\mathrm{op}} \rightarrow \mathrm{Spc}$ a presheaf, $V \rightarrow X$ a vector bundle, and $r \geq 0$. Let $U \subset V$ be an open subpresheaf such that, for every finite field extension k'/k and every $\alpha \in X(k')$, the closed complement of $\alpha^*(U)$ in $\alpha^*(V)$ has codimension at least r .*

- (1) *The morphism $\Sigma^2 U \rightarrow \Sigma^2 X$ is $\mathbf{A}^1$ - r -connected. If moreover U and X are $\mathbf{A}^1$ -connected, then the morphism $\Sigma U \rightarrow \Sigma X$ is $\mathbf{A}^1$ -($r - 1$)-connected.*
- (2) *The cofiber of $\Sigma_T^\infty U_+ \rightarrow \Sigma_T^\infty X_+$ in $\mathbf{SH}(k)$ is very r -effective.*

Proof. (1) By Lemma 8.8, it suffices to show that $\Sigma \mathrm{cofib}(U \rightarrow X)$ is $\mathbf{A}^1$ - r -connected. Colimits of pointed objects preserve connectivity, hence $\mathbf{A}^1$ -connectivity by the $\mathbf{A}^1$ -connectivity theorem, so we are reduced by universality of colimits to the case $X \in \mathrm{Sm}_k$. Since $V \rightarrow X$ is an $\mathbf{A}^1$ -equivalence and $V - U$ has codimension $\geq r$ in V , the result follows from Lemma 8.9.

(2) As in the proof of (1), we can assume $X \in \mathrm{Sm}_k$. Let Z be the closed complement of U in V . If Z is smooth, the claim follows from the purity isomorphism. In general, since k is perfect, one can stratify Z by smooth subschemes of codimension $\geq r$ in V (cf. the proof of Lemma 8.9), thereby reducing to the case Z smooth. ■

Lemma 8.11. *Let k be a field and $n \geq d - 1 \geq 0$. Then $\mathrm{Hilb}_d(\mathbf{A}_k^n)$ is $\mathbf{A}^1$ -connected.*

In fact, $\mathrm{Hilb}_d(\mathbf{A}_k^n)$ is $\mathbf{A}^1$ -connected for all n and d , at least when k is infinite, as Totaro shows by a more elaborate argument [65, Theorem 6.1]. Since $\mathbf{A}^1$ -connectedness implies connectedness, this result recovers Hartshorne's theorem that $\mathrm{Hilb}_d(\mathbf{P}^n)$ is connected [29] (although Hartshorne's theorem also applies to nonconstant Hilbert polynomials).

Proof. By [48, Lemma 6.1.3] and [50, Section 2, Corollary 3.22], it suffices to show that $\mathrm{Hilb}_d(\mathbf{A}^n)(k)$ is nonempty and that any two points of $\mathrm{Hilb}_d(\mathbf{A}^n)(F)$ can be connected by a chain of affine lines over F , for every separable finitely generated field extension F/k . It is clear that $\mathrm{Hilb}_d(\mathbf{A}^n)(k)$ is nonempty.

Fix coordinates $\mathbf{A}_F^n = \mathrm{Spec} F[x_1, \dots, x_n]$. Consider an F -point $[A] \in \mathrm{Hilb}_d(\mathbf{A}^n)(F)$ corresponding to a surjection $\pi: F[x_1, \dots, x_n] \rightarrow A$. We claim that π can be connected by a chain of affine lines to a surjection such that the images of $1, x_1, \dots, x_{d-1}$ are linearly independent. We prove this by adjusting π as necessary. If $1, x_1, \dots, x_{i-1}$ are linearly independent while $1, x_1, \dots, x_i$ are dependent, then

$$\pi|_{\hat{\cdot}: F[x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n]} \rightarrow A$$

is a surjection. For $a \in A$ consider the $F[t]$ -algebra homomorphism $\rho: F[x_1, \dots, x_n, t] \rightarrow A[t]$ defined by $\rho(x_j) = \pi(x_j)$ for $j \neq i$ and $\rho(x_i) = ta + (1-t)\pi(x_i)$. For every $\lambda \in \bar{F}$, the map ρ_λ is surjective since it extends $\pi|_i$. Hence, ρ itself is surjective and it defines an $\mathbf{A}^1$ -homotopy from $\pi = \rho_0$ to ρ_1 in $\text{Hilb}_d(\mathbf{A}^n)$. If we choose $a \in A - F\{\pi(1), \pi(x_1), \dots, \pi(x_{i-1})\}$, then the images of $1, x_1, \dots, x_i$ by ρ_1 are linearly independent. Continuing by induction we prove the claim. Having π such that $1, x_1, \dots, x_{d-1}$ span A , we use another $\mathbf{A}^1$ -homotopy to ensure $\pi(x_j) = 0$ for $j \geq d$.

As in the proof of Theorem 2.1, consider the algebra $\text{Rees}(A) := F \oplus tA[t] \subset A[t]$ which is finite locally free over $F[t]$. Define an $F[t]$ -algebra homomorphism

$$\tilde{\pi}: F[x_1, \dots, x_n, t] \rightarrow \text{Rees}(A)$$

by $\tilde{\pi}(x_i) = t\pi(x_i)$ for every i . Since $1, x_1, \dots, x_{d-1}$ span A as an F -vector space, the image of $\tilde{\pi}$ contains t as well as tA , and so $\tilde{\pi}$ is surjective. Geometrically, $\tilde{\pi}$ corresponds to a morphism $\tilde{\pi}: \mathbf{A}_F^1 \rightarrow \text{Hilb}_d(\mathbf{A}^n)$ that links $[A] = \tilde{\pi}(1)$ with $\tilde{\pi}(0)$. The latter F -point corresponds to an embedding of Spec of the square-zero extension $F \oplus F^{d-1}$ at the origin of the $(d-1)$ -dimensional subspace $x_d = \dots = x_n = 0$ in $\mathbf{A}^n$. In particular, it is independent of $[A]$, which proves that every F -point can be connected to a fixed one by a chain of affine lines over F , as we want. ■

Proof of Theorem 8.2. In (2) and (3), we may replace the field k by a perfect subfield, since the conclusions are preserved by essentially smooth base change. Recall that the forgetful map $\text{Hilb}_d(\mathbf{A}^\infty) \rightarrow \mathcal{F}\text{Flat}_d$ is a motivic equivalence (Corollary 4.4). For the connectivity on complex points and for (2) and (3), we apply Propositions 8.7 and 8.10 with $X = \mathcal{F}\text{Flat}_d$, $V = \mathcal{V}_d(\mathbf{A}^n)$, $U = \text{Hilb}_d(\mathbf{A}^n)$, and $r = n - d + 2$, see Lemma 8.5. For (2), note that both $\text{Hilb}_d(\mathbf{A}^n)$ and $\mathcal{F}\text{Flat}_d$ are $\mathbf{A}^1$ -connected by Lemma 8.11. To prove the connectivity on real points, we pull back the above situation to the scheme $X_m = \text{Hilb}_d(\mathbf{A}^m)$. By Proposition 8.7, $(U \times_X X_m)(\mathbf{R}) \rightarrow X_m(\mathbf{R})$ is $(n-d)$ -connected. Taking the colimit over m proves the result, using the fact that $\text{Hilb}_d(\mathbf{A}^\infty) \rightarrow \mathcal{F}\text{Flat}_d$ is a universal $\mathbf{A}^1$ -equivalence on affine schemes (Corollary 4.4). ■

Remark 8.12. Note that the assumptions of Propositions 8.7 and 8.10 are preserved by any base change $X' \rightarrow X$. For example, by considering the open substack $\mathcal{F}\text{Syn}_d \subset \mathcal{F}\text{Flat}_d$, we see that Theorem 8.2 holds with Hilb_d replaced by $\text{Hilb}_d^{\text{lci}}$.

Acknowledgments. We would like to thank Tom Bachmann, Elden Elmanto, Rahul Pandharipande, Bjorn Poonen, Oliver Röndigs, Vladimir Sosnilo, Markus Spitzweck, Ravi Vakil, Andrzej Weber, and the referees for discussions and suggestions. This work was partly initiated during Yakerson's visit to the Institute of Mathematics of Polish Academy of Sciences, and then continued during Jelisiejew's visit to the University of Regensburg. We would like to express our gratitude to both institutions for their hospitality. We also thank the organizers of the Southern California Algebraic Geometry Seminar and the ETH Zürich Algebraic Geometry and Moduli Zoominar.

Funding. M.H., D.N., and M.Y. were supported by SFB 1085 "Higher invariants".

J.J. was supported by NCN grant 2017/26/D/ST1/00913 and by the START fellowship of the Foundation for Polish Science. B.T. was supported by NSF grant DMS-2054553.

References

- [1] American Institute of Mathematics Problem List: Components of Hilbert schemes. <http://aimpl.org/hilbertschemes> (2010)
- [2] Bachmann, T.: [The generalized slices of Hermitian K-theory](#). J. Topol. **10**, 1124–1144 (2017) Zbl [1453.14065](#) MR [3743071](#)
- [3] Bachmann, T.: [Cancellation theorem for motivic spaces with finite flat transfers](#). Doc. Math. **26**, 1121–1144 (2021) Zbl [1483.14039](#) MR [4324462](#)
- [4] Bachmann, T., Calmès, B., Déglise, F., Fasel, J., Østvær, P. A.: Milnor–Witt motives. [arXiv:2004.06634](#) (2020), to appear in Mem. Amer. Math. Soc.
- [5] Bachmann, T., Elmanto, E., Hoyois, M., Khan, A. A., Sosnilo, V., Yakerson, M.: [On the infinite loop spaces of algebraic cobordism and the motivic sphere](#). Épijournal Géom. Algébrique **5**, art. 9, 13 pp. (2021) Zbl [1478.14043](#) MR [4275258](#)
- [6] Bachmann, T., Hoyois, M.: [Norms in motivic homotopy theory](#). Astérisque **425**, ix+207 pp. (2021) Zbl [07403459](#) MR [4288071](#)
- [7] Barwick, C., Glasman, S., Shah, J.: [Spectral Mackey functors and equivariant algebraic K-theory, II](#). Tunis. J. Math. **2**, 97–146 (2020) Zbl [1461.18009](#) MR [3933393](#)
- [8] Beauville, A.: [Variétés Kähleriennes dont la première classe de Chern est nulle](#). J. Differential Geom. **18**, 755–782 (1984) (1983) Zbl [0537.53056](#) MR [730926](#)
- [9] Bertin, J.: The punctual Hilbert scheme: an introduction. In: Geometric methods in representation theory. I, Séminaires et Congrès 24, Société Mathématique de France, Paris, 1–102 (2012) Zbl [1327.13026](#) MR [3202701](#)
- [10] Bröcker, T., tom Dieck, T.: Representations of compact Lie groups. Graduate Texts in Mathematics 98, Springer, New York (1995) Zbl [0874.22001](#) MR [1410059](#)
- [11] Cisinski, D.-C.: [Descente par éclatements en K-théorie invariante par homotopie](#). Ann. of Math. (2) **177**, 425–448 (2013) Zbl [1264.19003](#) MR [3010804](#)
- [12] Déglise, F.: Orientation theory in arithmetic geometry. In: K-Theory—Proceedings of the International Colloquium, Mumbai, 2016, Hindustan Book Agency, New Delhi, 239–347 (2018) Zbl [1451.14067](#) MR [3930052](#)
- [13] Ellingsrud, G., Strømme, S. A.: [On the homology of the Hilbert scheme of points in the plane](#). Invent. Math. **87**, 343–352 (1987) Zbl [0625.14002](#) MR [870732](#)
- [14] Elmanto, E., Hoyois, M., Khan, A. A., Sosnilo, V., Yakerson, M.: [Modules over algebraic cobordism](#). Forum Math. Pi **8**, art. e14, 44 pp. (2020) Zbl [1458.14027](#) MR [4190058](#)
- [15] Elmanto, E., Hoyois, M., Khan, A. A., Sosnilo, V., Yakerson, M.: [Motivic infinite loop spaces](#). Cambridge J. Math. **9**, 431–549 (2021) Zbl [07422194](#) MR [4325285](#)
- [16] Fernando, J. F., Ghiloni, R.: [Differentiable approximation of continuous semialgebraic maps](#). Selecta Math. (N.S.) **25**, art. 46, 30 pp. (2019) Zbl [1472.14067](#) MR [3984104](#)
- [17] Ferrand, D.: [Conducteur, descente et pincement](#). Bull. Soc. Math. France **131**, 553–585 (2003) Zbl [1058.14003](#) MR [2044495](#)
- [18] Fulton, W.: [Intersection theory](#). 2nd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete 2, Springer, Berlin (1998) Zbl [0885.14002](#) MR [1644323](#)
- [19] Garkusha, G., Panin, I.: [Framed motives of algebraic varieties \(after V. Voevodsky\)](#). J. Amer. Math. Soc. **34**, 261–313 (2021) Zbl [1491.14034](#) MR [4188819](#)
- [20] Geisser, T.: [Motivic cohomology over Dedekind rings](#). Math. Z. **248**, 773–794 (2004) Zbl [1062.14025](#) MR [2103541](#)
- [21] Gepner, D., Snaith, V.: [On the motivic spectra representing algebraic cobordism and algebraic K-theory](#). Doc. Math. **14**, 359–396 (2009) Zbl [1232.55010](#) MR [2540697](#)
- [22] Glasman, S.: [Day convolution for \$\infty\$ -categories](#). Math. Res. Lett. **23**, 1369–1385 (2016) Zbl [1375.18041](#) MR [3601070](#)

- [23] Görtz, U., Wedhorn, T.: [Algebraic geometry I](#). Advanced Lectures in Mathematics, Vieweg + Teubner, Wiesbaden (2010) Zbl [1213.14001](#) MR [2675155](#)
- [24] Göttsche, L.: Hilbert schemes of points on surfaces. In: Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), Higher Education Press, Beijing, 483–494 (2002) Zbl [1055.14005](#) MR [1957058](#)
- [25] Gustavsen, T. S., Laksov, D., Skjelnes, R. M.: [An elementary, explicit, proof of the existence of Quot schemes of points](#). Pacific J. Math. **231**, 401–415 (2007) Zbl [1152.14012](#) MR [2346502](#)
- [26] Hahn, J., Yuan, A.: [Exotic multiplications on periodic complex bordism](#). J. Topol. **13**, 1839–1852 (2020) Zbl [1461.55007](#) MR [4186145](#)
- [27] Haiman, M.: [Hilbert schemes, polygraphs and the Macdonald positivity conjecture](#). J. Amer. Math. Soc. **14**, 941–1006 (2001) Zbl [1009.14001](#) MR [1839919](#)
- [28] Hall, J., Rydh, D.: [General Hilbert stacks and Quot schemes](#). Michigan Math. J. **64**, 335–347 (2015) Zbl [1349.14013](#) MR [3359029](#)
- [29] Hartshorne, R.: [Connectedness of the Hilbert scheme](#). Inst. Hautes Études Sci. Publ. Math. **29**, 5–48 (1966) Zbl [0171.41502](#) MR [213368](#)
- [30] Hironaka, H.: Triangulations of algebraic sets. In: Algebraic geometry (Arcata, CA, 1974), Proceedings of Symposia in Pure Mathematics 29, American Mathematical Society, Providence, RI, 165–185 (1975) Zbl [0332.14001](#) MR [374131](#)
- [31] Hirsch, M. W.: Differential topology. Graduate Texts in Mathematics 33, Springer, New York (1976) Zbl [0356.57001](#) MR [1336822](#)
- [32] Horrocks, G.: [Sheaves on projective space invariant under the unitriangular group](#). Invent. Math. **10**, 108–118 (1970) Zbl [0194.51901](#) MR [269664](#)
- [33] Hoskins, V., Pepin Lehalleur, S.: [On the Voevodsky motive of the moduli space of Higgs bundles on a curve](#). Selecta Math. (N.S.) **27**, art. 11, 37 pp. (2021) Zbl [1470.14014](#) MR [4222601](#)
- [34] Hoyois, M.: [Cdh descent in equivariant homotopy K-theory](#). Doc. Math. **25**, 457–482 (2020) Zbl [1453.14068](#) MR [4124487](#)
- [35] Hoyois, M.: [The localization theorem for framed motivic spaces](#). Compos. Math. **157**, 1–11 (2021) Zbl [1455.14042](#) MR [4215649](#)
- [36] Hoyois, M., Jelisiejew, J., Nardin, D., Yakerson, M.: [Hermitian K-theory via oriented Gorenstein algebras](#). J. Reine Angew. Math. **793**, 105–142 (2022) Zbl [1520.14046](#) MR [4513165](#)
- [37] Hoyois, M., Kelly, S., Østvær, P. A.: [The motivic Steenrod algebra in positive characteristic](#). J. Eur. Math. Soc. **19**, 3813–3849 (2017) Zbl [1386.14087](#) MR [3730515](#)
- [38] Iarrobino, A.: [Reducibility of the families of 0-dimensional schemes on a variety](#). Invent. Math. **15**, 72–77 (1972) Zbl [0227.14006](#) MR [301010](#)
- [39] Iversen, B.: [Linear determinants with applications to the Picard scheme of a family of algebraic curves](#). Lecture Notes in Mathematics 174, Springer, Berlin (1970) Zbl [0205.50802](#) MR [292835](#)
- [40] Jelisiejew, J.: [Pathologies on the Hilbert scheme of points](#). Invent. Math. **220**, 581–610 (2020) Zbl [1473.14007](#) MR [4081138](#)
- [41] Khan, A. A., Rydh, D.: Virtual Cartier divisors and blow-ups. arXiv:[1802.05702v2](#) (2019)
- [42] Levine, M.: Techniques of localization in the theory of algebraic cycles. J. Algebraic Geom. **10**, 299–363 (2001) Zbl [1077.14509](#) MR [1811558](#)
- [43] Levine, M.: [The homotopy coniveau tower](#). J. Topol. **1**, 217–267 (2008) Zbl [1154.14005](#) MR [2365658](#)
- [44] Lurie, J.: [Higher topos theory](#). Annals of Mathematics Studies 170, Princeton University Press, Princeton, NJ (2009) Zbl [1175.18001](#) MR [2522659](#)
- [45] Lurie, J.: Spectral algebraic geometry. <http://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf> (2018)

- [46] Mazza, C., Voevodsky, V., Weibel, C.: Lecture notes on motivic cohomology. Clay Mathematics Monographs 2, American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA (2006) Zbl 1115.14010 MR 2242284
- [47] Miller, E., Sturmfels, B.: [Combinatorial commutative algebra](#). Graduate Texts in Mathematics 227, Springer, New York (2005) Zbl 1090.13001 MR 2110098
- [48] Morel, F.: [The stable \$\mathbb{A}^1\$ -connectivity theorems](#). K-Theory **35**, 1–68 (2005) Zbl 1117.14023 MR 2240215
- [49] Morel, F.: [\$\mathbb{A}^1\$ -algebraic topology over a field](#). Lecture Notes in Mathematics 2052, Springer, Heidelberg (2012) Zbl 1263.14003 MR 2934577
- [50] Morel, F., Voevodsky, V.: [\$\mathbb{A}^1\$ -homotopy theory of schemes](#). Inst. Hautes Études Sci. Publ. Math. **90**, 45–143 (2001) (1999) Zbl 0983.14007 MR 1813224
- [51] Nakajima, H.: [Lectures on Hilbert schemes of points on surfaces](#). University Lecture Series 18, American Mathematical Society, Providence, RI (1999) Zbl 0949.14001 MR 1711344
- [52] Naumann, N., Spitzweck, M., Østvær, P. A.: [Chern classes, K-theory and Landweber exactness over nonregular base schemes](#). In: Motives and algebraic cycles, Fields Institute Communications 56, American Mathematical Society, Providence, RI, 307–317 (2009) Zbl 1190.14022 MR 2562464
- [53] Poonen, B.: [Isomorphism types of commutative algebras of finite rank over an algebraically closed field](#). In: Computational arithmetic geometry, Contemporary Mathematics 463, American Mathematical Society, Providence, RI, 111–120 (2008) Zbl 1155.13015 MR 2459993
- [54] Poonen, B.: [The moduli space of commutative algebras of finite rank](#). J. Eur. Math. Soc. **10**, 817–836 (2008) Zbl 1151.14011 MR 2421162
- [55] Raynaud, M., Gruson, L.: [Critères de platitude et de projectivité. Techniques de “platification” d’un module](#). Invent. Math. **13**, 1–89 (1971) Zbl 0227.14010 MR 308104
- [56] Rezk, C.: Proof of the Blakers–Massey theorem. <https://faculty.math.illinois.edu/~rezk/freudenthal-and-blakers-massey.pdf> (2015)
- [57] Röndigs, O., Østvær, P. A.: [Modules over motivic cohomology](#). Adv. Math. **219**, 689–727 (2008) Zbl 1180.14015 MR 2435654
- [58] Schedler, T.: Deformations of algebras in noncommutative geometry. In: Noncommutative algebraic geometry, Mathematical Sciences Research Institute Publications 64, Cambridge University Press, Cambridge, 71–165 (2016) Zbl 1386.14057 MR 3618473
- [59] Segal, G.: [K-homology theory and algebraic K-theory](#). In: K-theory and operator algebras (Athens, GA, 1975), Lecture Notes in Mathematics 575, Springer, Berlin, 113–127 (1977) Zbl 0363.55002 MR 515311
- [60] Sernesi, E.: Deformations of algebraic schemes. Grundlehren der mathematischen Wissenschaften 334, Springer, Berlin (2006) Zbl 1102.14001 MR 2247603
- [61] Spitzweck, M.: [A commutative \$\mathbb{P}^1\$ -spectrum representing motivic cohomology over Dedekind domains](#). Mém. Soc. Math. France (N.S.) **110** (2018) Zbl 1408.14081 MR 3865569
- [62] Spitzweck, M., Østvær, P. A.: [Motivic twisted K-theory](#). Algebr. Geom. Topol. **12**, 565–599 (2012) Zbl 1282.14040 MR 2916287
- [63] Suslin, A., Voevodsky, V.: [Singular homology of abstract algebraic varieties](#). Invent. Math. **123**, 61–94 (1996) Zbl 0896.55002 MR 1376246
- [64] The Stacks Project Authors: The Stacks project. <http://stacks.math.columbia.edu> (2020)
- [65] Totaro, B.: [Torus actions, Morse homology, and the Hilbert scheme of points on affine space](#). Épijournal Géom. Algébrique **5**, art. 12, 14 pp. (2021) Zbl 1470.14094 MR 4315780
- [66] Voevodsky, V.: Notes on framed correspondences. <http://www.math.ias.edu/vladimir/files/framed.pdf> (2001)
- [67] Whitehead, G. W.: Elements of homotopy theory. Graduate Texts in Mathematics 61, Springer, New York (1978) Zbl 0406.55001 MR 516508