

Exploring the unknown: The work of Louis Nirenberg on partial differential equations

Tristan Rivière

Abstract. Partial differential equations are central objects in the mathematical modeling of natural and social sciences (sound propagation, heat diffusion, thermodynamics, electromagnetism, elasticity, fluid dynamics, quantum mechanics, population growth, finance, etc.). They were for a long time restricted only to the study of natural phenomena or questions pertaining to geometry, before becoming over the course of time, and especially in the last century, a field in itself.

The second half of the 20th century was the “golden age” of the exploration of partial differential equations from a theoretical perspective. Louis Nirenberg, starting in the early 1950s, played a major role in the growth of this fundamental area of human knowledge and his name is associated with many of the milestones in the study of PDEs.

Louis Nirenberg passed away in January 2020 and his memory as a mathematician and colleague is a special invitation to review the development of the field of PDEs through the work of its leading pioneers.

Dedicated to the memory of Louis Nirenberg

1. Introduction

Louis Nirenberg used to qualify the field of *partial differential equations* as being “messy” (and also often acknowledged his special taste for what he called this “messiness”), which is probably a reference to the “intrinsic diversity” of the field. We would like to illustrate the pertinence of this quote by contradiction and by presenting the original attempts made mostly in the 19th century to see PDEs as a whole and the limits and inadequacies by which this approach has been confronted.

1.1. A general existence result: The Cauchy–Kowalevski theorem

Perhaps the first general systematic study of partial differential equations goes back to the work of Augustin-Louis Cauchy and his existence theorem for *quasilinear* first order PDEs with real analytic data.

2020 *Mathematics Subject Classification.* Primary 35-xx; Secondary 53Axx, 58Exx, 58Jxx, 70Gxx, 70H20, 76D03, 76D05.

Keywords. Partial differential equations, geometric analysis, calculus of variations.

Theorem 1.1 (Cauchy (1842)). *Let $(A^j)_{j=1,\dots,n-1}$ be a family of $n-1$ real analytic maps from a neighborhood of $(0,0)$ in $\mathbb{R}^n \times \mathbb{R}^m$ into the space of real $m \times m$ matrices and let f be a real analytic map into \mathbb{R}^m . Then there exists a unique real analytic solution u in a neighborhood of the origin to the following system:*

$$\begin{cases} \partial_{x_n} u = \sum_{j=1}^{n-1} A^j(x, u) \partial_{x_j} u + f(x, u), \\ u(x_1, \dots, x_{n-1}, 0) \equiv 0. \end{cases}$$

In 1874, Sofia Kowalevski (or Kowalevskaya), apparently unaware of Cauchy's work, proved in her thesis a general non-linear version of the previous result. Let us present the theorem called nowadays *Cauchy–Kowalevski theorem* in the particular case of *second order non-linear scalar equations*

$$F(x, u, \partial u, \partial^2 u) = 0, \quad \text{where } \partial u := (\partial_{x_i} u)_{i=1,\dots,n} \text{ and } \partial^2 u := (\partial_{x_i x_j}^2 u)_{i,j=1,\dots,n}.$$

Here, $F(x, p, q, r)$ is a real analytic function of all the entries $p \in \mathbb{R}$, $q = (q_i)_{i=1,\dots,n} \in \mathbb{R}^n$ and $r = (r_{ij})_{i,j=1,\dots,n} \in \mathbb{R}^{n^2}$. To that purpose, we introduce the notion of *characteristic direction*. A direction $X = (\xi_i)_{i=1,\dots,n} \in \mathbb{R}^n$ is called *characteristic* at (x, p, q, r) if

$$\sum_{i,j}^n \frac{\partial F}{\partial r_{ij}}(x, p, q, r) \xi_i \xi_j = 0.$$

Given a function u defined in a neighborhood of a point x_0 , a hypersurface $f(x) = 0$ is called *non-characteristic* at x_0 if $\xi := \nabla f(x_0)$ is not a characteristic direction at $(x_0, u(x_0), \partial u(x_0), \partial^2 u(x_0))$. The Cauchy–Kowalevski theorem requires that *initial data*, the so called *Cauchy data*, be given on a *non-characteristic* hyper-surface. Observe that if the PDE has the form

$$F(x, u, \partial u, \partial^2 u) = \partial_{x_n x_n}^2 u - f(x, u, \partial u, \partial^2 u),$$

where f does not depend on r_{nn} , the surface $x_n = 0$ is automatically *non characteristic*. We can now state the *Cauchy–Kowalevski theorem* in that particular case.

Theorem 1.2 (Cauchy–Kowalevski (1874)). *Let $f(x, p, q, r)$ be a real analytic function of its variables, and assume that f is independent of r_{nn} . Let $u_0(x_1, \dots, x_{n-1})$ and $u_1(x_1, \dots, x_{n-1})$ be two real analytic functions defined in the neighborhood of the origin 0. Then there is a unique real analytic solution defined in a neighborhood of the origin for the problem*

$$\begin{cases} \partial_{x_n x_n}^2 u = f(x, u, \partial u, \partial^2 u), \\ u = u_0 \text{ and } \partial_{x_n} u = u_1 \text{ on } x_n = 0. \end{cases}$$

1.2. Some inadequacies of the Cauchy–Kowalevski theory

The Cauchy–Kowalevski theorem requires an analytic framework. Its historical proof consists of an argument based on the convergence of power series. Only the analyticity assumptions with respect to x_n could be relaxed. The question of whether there could be more solutions for the same analytic data (in the C^∞ class, for instance) has stimulated much research and, although there are uniqueness theorems for some classes of linear PDEs, there are also counter-examples to uniqueness; see [55]. The general question remains to be settled.

If one seeks global solutions, which are expected to exist in physical problems, there is an “intrinsic” need to relax the analytic framework, since singularities can appear in “finite time” (the time variable here being x_n) even though all data are analytic. Consider, for instance, the Cauchy solution to

$$\begin{cases} \partial_{x_2} u = u \partial_{x_1} u, \\ u(x_1, 0) = -\frac{x_1}{1+x_1^2}. \end{cases}$$

The gradient on each level set of $u = u_0 = -x_0/(1+x_0^2)$ for any $u_0 \in (-1, 1)$ is constant, hence each level set is made of straight segments leaving the point $(x_0, 0)$ in the direction given by the vector $(1, u_0)$. These segments have to meet at points where u thus necessarily ceases to be continuous.

In her thesis, Sofia Kowalevski illustrated the need for the assumption that the initial surface be *non-characteristic* by the following example. Consider locally an analytic solution of

$$\begin{cases} \partial_{x_2} u = \partial_{x_1 x_1}^2 u, \\ u(x_1, 0) = \frac{1}{1-x_1}. \end{cases}$$

An elementary computation gives the explicit value of the coefficients of the Taylor series expansion for the solution, which happens not to converge at the origin. Hence, near the origin, there is no analytic solution to that equation with the given initial data. The reason why Cauchy’s theorem does not apply in this case, is that $x_2 = 0$ is *characteristic* for the equation. However this equation is nothing but the *heat equation* modeling in particular the diffusion of heat in an homogeneous material starting from the data $u(x_1, 0)$. A non-existence result certainly seems counter-intuitive! It suggests that one should leave the analytic framework imposed by Cauchy and Kowalevski.

The conclusion at this stage is that the analysis of PDEs *cannot be captured in a single theory* having simply to do with the convergence of power series.

2. Local solvability

2.1. The notion of local solvability and Lewy's counterexample

The first attempt to go beyond the Cauchy–Kowalevski theory would be to give up uniqueness requirements and look at “germs” of PDEs at a point. We are thus led to considering the linear framework and ask whether one can enlarge the class of possible solutions from the analytic class to the C^∞ class, or even to the space of distributions¹ \mathcal{D}' . Is a linear PDE always *locally* solvable? Precisely, consider $a_\alpha(x)$ to be a family of C^∞ complex coefficients defined in a neighborhood of the origin in \mathbb{R}^n and indexed by $\alpha = (\alpha_1, \dots, \alpha_n)$, where $\alpha_j \in \mathbb{N}$ and $|\alpha| := \sum \alpha_j \leq p$ for some $p \in \mathbb{N}^*$. Given any $f \in C_0^\infty(\mathbb{R}^n, \mathbb{C})$, does there exist locally near the origin a complex-valued solution (if need be, even in the very weak sense of distributions) to

$$L u = \sum_{\alpha=(\alpha_1 \dots \alpha_n)} a_\alpha(x) \frac{\partial^{|\alpha|} u}{\partial^{\alpha_1} x_1 \dots \partial^{\alpha_n} x_n} = f ?$$

If for a given L and any f , the answer to the question is “yes”, the PDE is called *locally solvable* at 0. Around 1955, Leon Ehrenpreis and Bernard Malgrange proved independently the *local solvability* of any linear PDE with constant coefficients $a_\alpha(x) \equiv a_\alpha$; see [25–28, 54]. Using Laurent Schwartz’s theory of tempered distributions such a PDE can be converted into a convolution equation, and the problem is reduced to a division problem in function algebra. Encouraged by this result, the conjecture asserting that any linear operator L should be *locally solvable* became notorious in the PDE community. That was until Hans Lewy came up in 1957 with a spectacular counter-example, namely: there exists a non-analytic function $f \in C^\infty$ such that the PDE

$$\frac{\partial u}{\partial x_1} + i \frac{\partial u}{\partial x_2} - 2i(x_1 + i x_2) \frac{\partial u}{\partial x_3} = f$$

has no C^1 solution in any neighborhood of the origin in \mathbb{R}^3 ; see [50]. This counter-example triggered an intense research activity which involved many prominent analysts of that time (such as Lars Hörmander, Louis Nirenberg, François Trèves, Yuri Vladimirovich Egorov, and so on). They were investigating which necessary and sufficient conditions would ensure *local solvability*.

2.2. The Nirenberg–Trèves local solvability condition

In three fundamental papers, Louis Nirenberg and François Trèves (see [63–65]) in 1963 proposed a condition on the *principal symbol of the PDE*

$$p_m(x, \xi) := \sum_{|\alpha|=p} a_\alpha(x) \prod_{j=1}^n \xi_j^{\alpha_j}$$

that should be necessary and sufficient for local solvability, the so called *(P) condition*.

¹We return to this notion of weak solutions introduced by Laurent Schwartz later in this presentation.

The bi-characteristics of a real function $A(x, \xi)$ are the curves $(x(t), \xi(t))$ that solve the Hamilton–Jacobi equation

$$\frac{dx}{dt} = \nabla_{\xi} A(x, \xi) \quad \text{and} \quad \frac{d\xi}{dt} = -\nabla_x A(x, \xi).$$

On such a curve, $A(x(t), \xi(t))$ is constant, and the curves for which this constant is zero are called the *null bi-characteristics* of A . The *Nirenberg–Trèves (P) condition* reads as follows:

On every null bi-characteristic of $\Re p_m$ the function $\Im p_m$ does not change sign, (P)

where $\Re p_m$ and $\Im p_m$ denote respectively the real and imaginary parts of the principal symbol $p_m(x, \xi)$. The necessity and sufficiency of the *Nirenberg–Trèves (P) condition* for the *local solvability* was established for a growing number of cases in successive works by Nirenberg and Trèves themselves, then by Beals and Fefferman [5], Hörmander [40], until a very recent result of Nils Dencker in 2006 (see [21]), which establishes the sufficiency of the generalized *Nirenberg–Trèves (P) condition* for general *pseudo-differential operators*², the so called *Nirenberg–Trèves (ψ) condition*.

3. Cauchy problems and global solvability for linear PDEs

3.1. The notion of a Cauchy problem

Understanding the local solvability of a PDE is certainly one important question, but one might argue that its relevance should not come into play in physical applications where solutions are expected to be *globally* defined on a given subdomains of spacetime.

The question of global solvability is traditionally coupled with that of uniqueness, and together they form what is called a *Cauchy problem*. A *Cauchy problem*, or *well-posed problem*, consists of a linear PDE L , of a function space E in which the data (the input) makes sense, of a function space F to which the expected solution (the output – also called the “unknown”) should belong, with the requirements that:

- (i) there exists exactly one solution in F of the PDE L for any given data in the function space E ;
- (ii) the dependence of the solution on the data is continuous from E into F .

²This generalized notion of differential operators was introduced by Louis Nirenberg in collaboration with Joseph J. Kohn in [46], and by Lars Hörmander in [39] a few years after the introduction by Calderón and Zygmund of the theory of *singular integral operators*. This revival of the Calderón Zygmund theory has probably been strongly stimulated by the proof of the index theorem by Michael Atiyah and Isadore Singer in 1962. The author of this presentation would like to apologize at this stage to the mathematicians from an important branch of the PDE community for certainly not giving enough credit to the pioneered work of Kohn and Nirenberg which is a milestone in the modern theory of *micro-local analysis*. As Louis Nirenberg used to say, mathematics is very much about taste and individual enthusiasm ...

Finding appropriate *Cauchy problems* for linear PDEs has generated a tremendous amount of mathematical activities in the last century. It also gives the “asymptotes”, or “constraints at the horizon”, in order to solve many *non-linear* problems, as we shall see.

One can illustrate the notion of *Cauchy problem* by looking at examples of *ill-posed problems*. One of them deals with finding a holomorphic extension in the 2 dimensional disc, that we will denote D^2 , of a prescribed C^1 boundary data. In other words, given $\phi \in E := C^1(\partial D^2, \mathbb{C})$, one seeks $u \in F := C^1(\bar{D}^2, \mathbb{C})$ satisfying

$$\begin{cases} Lu := \bar{\partial}u = \frac{1}{2} \left(\frac{\partial u}{\partial x_1} + i \frac{\partial u}{\partial x_2} \right) = 0 & \text{in } D^2, \\ u = \phi & \text{on } \partial D^2. \end{cases}$$

This is an ill-posed problem, because not every $\phi \in C^1(\partial D^2, \mathbb{C})$ admits a C^1 holomorphic extension in D^2 (take for instance $\phi(\theta) = e^{-i\theta}$). The ill-posedness can however be thwarted by replacing $E := C^1(\partial D^2, \mathbb{C})$ with the subspace of C^1 functions that are L^2 -orthogonal to $e^{-ik\theta}$ for any $k \in \mathbb{N}^*$.

3.2. The fragmentation of the analysis of PDEs

The search for Cauchy problems has imposed a fragmentation of the field of PDEs into multiple areas of analysis which have often grown independently of each other. This is due to the very different behaviors of various differential operators. The analysis of PDEs cannot be encapsulated into a single theory, and it is intrinsically split. It is in fact a field that might seem disorderly from the outside. But this “messiness”, which sometimes discourages young vocations, is in the very nature of PDEs and it is the source of an infinite diversity of phenomena, arguments, and results. One may nevertheless attempt to put some order in this diversity, by singling out three main families of operators that we briefly present in the framework of *linear second order scalar equations* with constant coefficients:

$$Lu := \sum_{i,j=1}^n a_{ij} \frac{\partial^2 u}{\partial x_i \partial x_j} + \sum_{i=1}^n b_i \frac{\partial u}{\partial x_i} + c u$$

with the symmetric principal symbol: $a_{ij} = a_{ji}$.

(i) The *elliptic operators* are those for which L has no characteristic direction:

$$\forall \xi \neq 0 \quad \sum_{i,j=1}^n a_{ij} \xi_i \xi_j > 0.$$

The archetype of the elliptic operator is the *Laplacian operator*

$$Lu := \Delta u = \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i \partial x_i}.$$

The *Laplace equation* $\Delta u = 0$, also called the *potential equation*, arises in the physics of gravitational and electromagnetic fields. The field is generated by the potential u satisfying $\Delta u = f$, where f denotes either the mass or the charge density.

(ii) The *parabolic operators* are those for which one direction is singled out and does not appear in the second derivatives. They have the form

$$Lu := \frac{\partial u}{\partial t} - \sum_{i,j=1}^{n-1} a_{ij} \frac{\partial^2 u}{\partial x_i \partial x_j} - \sum_{i=1}^{n-1} b_i \frac{\partial u}{\partial x_i} - c u,$$

where the (a_{ij}) operator for the remaining variables is (negative) elliptic:

$$\forall \xi \neq 0 \quad \sum_{i,j=1}^{n-1} a_{ij} \xi_i \xi_j > 0.$$

The model case of a parabolic operator is the *heat operator*

$$Lu := \frac{\partial u}{\partial t} - \Delta u = \frac{\partial u}{\partial t} - \sum_{i=1}^{n-1} \frac{\partial^2 u}{\partial x_i \partial x_i}.$$

The *heat equation* $\partial_t u - \Delta u = 0$ is a fundamental tool in the modeling of diffusion (heat, combustion, chemical reactions, and so on).

(iii) The *hyperbolic operators* are those for which the matrix (a_{ij}) has signature $(+, -, \dots, -)$. The prototype is the *wave operator*

$$Lu := \frac{\partial^2 u}{\partial t^2} - \Delta u = \frac{\partial^2 u}{\partial t^2} - \sum_{i=1}^{n-1} \frac{\partial^2 u}{\partial x_i \partial x_i}.$$

The *wave equation* $\partial_{t^2}^2 u - \Delta u = 0$ is a fundamental equation arising in the modeling of propagation of waves in gases and fluids.

This (overly) simplified classification leaves out many equations, including ones that are relevant to physical phenomena, such as the Schrödinger equation, Korteweg–deVries water-wave equation, Navier–Stokes, and so on. Nonetheless, understanding how the three basic families differ from one another constitutes a first step in the study of “hybrid” and of more complicated PDEs. Parabolic equations can be understood as elliptic equations with time propagation, and thus these two families enjoy many similar properties, such as smoothing effect and infinite speed propagation. Hyperbolic equations on the other hand are very different, and they involve, for examples, finite speed propagation and the “transport” of singularities. The work of Louis Nirenberg has a barycenter whose location lies closer to the first two families, and this is why we shall mostly restrict this discussion to *elliptic* and *parabolic* PDEs.

3.3. The birth of Cauchy problems: The Dirichlet principle

Peter Gustav Lejeune Dirichlet was a German mathematician of the first half of the 19th century. His interest for physics led him to formulate a mathematical problem related to potential theory, a field which finds its roots in Newtonian mechanics. This problem, named after him by Riemann, is formulated as follows:

A bounded domain empty of charge being given as well as the value of the potential at its boundary, show mathematically that it produces a unique electromagnetic field in the interior of the domain.

In his lecture³ on potential theory in Göttingen in 1857, Dirichlet gave a more precise mathematical formulation of the problem. Namely, show that for any bounded connected (smooth) domain Ω in \mathbb{R}^3 and for any continuous differentiable function ϕ on $\partial\Omega$, there exists a unique continuously differentiable function u equal to ϕ on $\partial\Omega$ and minimizing the following integral, now called *Dirichlet energy*:

$$E(u) := \int_{\Omega} |\nabla u|^2 dx^3.$$

Such a function satisfies the Laplace equation

$$\begin{cases} \Delta u = 0 & \text{in } \Omega, \\ u = \phi & \text{on } \partial\Omega. \end{cases} \quad (3.1)$$

The claims made by Dirichlet in his lecture were not completely settled, but they were however used in the following years by Bernhard Riemann in two-dimensional complex analysis. This situation created some trouble in the mathematical community and the Dirichlet principle came under attack, in particular in the work of Karl Weierstrass, who produced examples of coercive Lagrangians (akin to the Dirichlet energy), which do not possess minimizers for some boundary data. The revival of the Dirichlet principle came in the early 20th century in the work of David Hilbert who solved it rigorously in some special cases and devoted the 20th of his famous list of 23 problems to this principle. It is hard to find papers of the beginning of the 20th century on PDE without any mention of the *Dirichlet principle*. A detailed description of the steps towards the full resolution of Dirichlet's assertions, although very instructive, would take us on a too long detour. The resolution of Dirichlet's problem gave rise to numerous ad hoc tools from functional analysis for solving PDEs, such as the notion of *Sobolev spaces*, *distributions*, *weak solutions*, and so on. Let us also mention the names of Henri Léon Lebesgue, Leonida Tonelli, Beppo Levi, Sergei Sobolev, Laurent Schwartz.

The modern way to solve the Dirichlet problem goes as follows. One introduces the so called *Sobolev Space* of Lebesgue measurable functions u on the bounded domain Ω whose Schwartz distributional derivatives lie in the *Hilbert space* of square integrable functions:

$$W^{1,2}(\Omega, \mathbb{R}) = \{u \in L^2; \partial_{x_i} u \in L^2 \text{ for } i = 1, \dots, 3\},$$

where the distributional derivatives of u are defined by duality with compactly supported smooth functions

$$\forall \phi \in C_0^\infty(\Omega, \mathbb{R}) \quad \langle \partial_{x_i} u, \phi \rangle := - \int_{\Omega} u \frac{\partial \phi}{\partial x_i} dx^3.$$

³published after his death by F. Grube in 1876; see [22, 23].

This space is equipped with the scalar product for which it is complete, namely,

$$(u, v) := \int_{\Omega} \sum_{i=1}^3 \partial_{x_i} u \partial_{x_i} v + u v \, dx^3.$$

The Hilbert Sobolev space $W^{1,2}$ has favorable analytical properties, such as *weak sequential pre-compactness* (from any sequence with uniformly bounded norm one can extract a subsequence converging weakly – in duality with smooth functions – to a function in $W^{1,2}$). Furthermore, the Dirichlet energy $E(u)$ is *weakly sequentially lower semi-continuous* and for any weakly converging sequence u_k with limit u_{∞} , there holds

$$\liminf_{k \rightarrow +\infty} E(u_k) \geq E(u_{\infty}).$$

One proves that smooth functions are dense in this *Hilbert space*. Moreover, the map which to a smooth function in $\bar{\Omega}$ assigns its restriction to the boundary $\partial\Omega$ is continuous from $W^{1,2}(\Omega, \mathbb{R})$ into the space of square integrable functions on the boundary $\partial\Omega$, and therefore extends continuously to a well defined map from $W^{1,2}(\Omega, \mathbb{R})$ into $L^2(\partial\Omega)$. These facts combined together give the existence of a minimizer of the Dirichlet energy for any weak boundary data (L^2 function ϕ) which admits a $W^{1,2}$ extension inside Ω . This subspace of $L^2(\partial\Omega)$ is denoted by $H^{1/2}(\partial\Omega)$. The use of distribution theory makes it possible to assert that any such minimizer u satisfies the Laplace equation in a weak sense:

$$\forall \phi \in C_0^{\infty}(\Omega, \mathbb{R}) \quad \langle \Delta u, \phi \rangle := \int_{\Omega} u \Delta \phi \, dx^3 = 0.$$

A fundamental lemma of Laurent Schwartz states that such a function u has to be analytic in the interior of the domain Ω and is the unique solution to (3.1). This full resolution of the Cauchy problem for the Dirichlet question (i.e. the *Dirichlet Problem*) can be extended without much difficulty in the case where “inside charges” are present, and one shows that for any smooth function f in Ω and any smooth function ϕ in $\partial\Omega$, there is a unique solution – smooth inside Ω – to the inhomogeneous problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = \phi & \text{on } \partial\Omega. \end{cases} \quad (3.2)$$

3.4. The Agmon–Douglis–Nirenberg elliptic Cauchy problems in the Banach L^p spaces

Numerous problems from classical mechanics, quantum mechanics, chemistry, biology, and geometry can be described by means of elliptic non-linear PDEs of the form

$$-\Delta u = f(x, u, \nabla u), \quad (3.3)$$

where the highest-order term is given by the Laplace operator or more generally by an *elliptic operator* which has no null characteristic. Considering this PDE on a bounded

smooth domain Ω with zero boundary condition, it is tempting to view it as a perturbation of (3.4) and use the inverse of the Dirichlet problem, which was given in the previous section for $\phi = 0$ and which we denote by $(-\Delta)_0^{-1}$. Then (3.3) becomes

$$u = (-\Delta)_0^{-1} f(x, u, \nabla u).$$

The idea behind this reformulation is ultimately to use a fixed point argument for solving the PDE. This is done in the same vein as for proving Cauchy–Lipschitz–Picard existence theorem for ODEs of the form $\dot{y} = f(t, y(t))$. One writes the solution as an integral

$$y(t) = y(t_0) + \int_{t_0}^t f(s, y(s)) ds,$$

before applying a fixed point argument. The difficulty is to find the right space in which to work. In the ODE case, for smooth f , the integral operator is very explicit and one shows that the space of Lipschitz functions gives an ad hoc framework for the fixed point argument to work. In the case of PDEs, one must first have a thorough understanding of the operator $(-\Delta)_0^{-1}$ and how it acts on elements of various Banach spaces.

In two fundamental papers published in 1959 and in 1964, Schmel Agmon, Avron Douglis, and Louis Nirenberg solved the elliptic Cauchy problems and the invertibility of elliptic operators of arbitrary orders in domains; see [2, 3]. They worked in the context of Banach L^p spaces. They obtained a series of optimal results that opened the way in exploring not only linear but also non-linear PDEs, which were beforehand completely out of reach. We give below the simplest example of a boundary problem solved by the Agmon–Douglis–Nirenberg theory of Cauchy problems in Banach spaces bearing in mind that the theory applies to a very wide family of problems.

Theorem 3.1 (Agmon–Douglis–Nirenberg (1959)). *Let Ω be a smooth bounded domain of \mathbb{R}^n and $p \in (1, +\infty)$. For any $f \in L^p(\Omega)$, there exists a unique solution u to*

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (3.4)$$

such that the distributional second and first derivatives of u lie in $L^p(\Omega)$, and moreover

$$\int_{\Omega} \sum_{|\alpha| \leq 2} |\partial^\alpha u|^p \leq C(\Omega, p) \int_{\Omega} |f|^p,$$

where $C(\Omega, p)$ is a positive constant independent of $f \in L^p(\Omega)$. \square

With such an estimate at hand, it becomes possible to use the operator $(-\Delta)_0^{-1}$ and a fixed point argument in the Banach spaces L^p so as to obtain a solution to non-linear PDEs. A major piece of the puzzle was brought in by Alberto Calderón and Antoni Zygmund in 1952. The “singular integral theory” asserts in particular⁴ that, for Schwartz

⁴It is striking that the trace of the Hessian matrix encodes the whole Hessian matrix, in the L^p space perspective.

functions⁵ u , the mapping

$$\Delta u \rightarrow (\partial_{x_i x_j}^2 u)_{i,j=1,\dots,n}$$

is continuous for the $L^p(\mathbb{R}^n)$ norm into $L^p(\mathbb{R}^n)$ for any $p \in (1, +\infty)$; see [17, 18]. The Agmon–Douglis–Nirenberg L^p theory is parallel to a previous theory developed by the Polish mathematician Juliusz Schauder around 1930, which involves the more regular Hölder spaces⁶ instead of L^p or Sobolev spaces.

4. Inequalities and a priori estimates

4.1. Gagliardo–Nirenberg interpolation inequalities

The field of PDEs is structured by *inequalities*. They are the “power horses” of the field. In the mid 1950s, functional analysis was already rich in inequalities: Hölder, Minkowski, Poincaré, Poincaré–Wirtinger, Young, Hausdorff–Young, Hardy, Hardy–Littlewood, etc., and the more recent Sobolev inequalities. Deep scaling considerations (a common trait in Louis Nirenberg’s work) led Louis Nirenberg around 1959 (see [61, 62]), and independently Emilio Gagliardo (see [34]), to discover a large family of inequalities for Sobolev norms “lying” between two others. As an illustration, we have that for any $p \in [1, +\infty]$, any $r \in [1, +\infty]$, and any $n \in \mathbb{N}^*$, there exists a constant $C > 0$ such that for all Schwartz functions $u \in \mathcal{S}(\mathbb{R}^n)$, there holds

$$\left[\int_{\mathbb{R}^n} \sum_{|\alpha|=1} |\partial^\alpha u|^q dx^n \right]^{1/q} \leq C \left[\int_{\mathbb{R}^n} |u|^p dx^n \right]^{1/2p} \left[\int_{\mathbb{R}^n} \sum_{|\alpha| \leq 2} |\partial^\alpha u|^r dx^n \right]^{1/2r}, \quad (4.1)$$

where $q^{-1} = 2^{-1}(p^{-1} + r^{-1})$ and dx^n denotes the Lebesgue measure on \mathbb{R}^n . In the family of *Gagliardo–Nirenberg interpolation inequalities*, one also finds the following one that we shall use later on. For any $1 \leq p \leq q < +\infty$ and any $n \in \mathbb{N}^*$, there exists $C > 0$ such that for all Schwartz functions $u \in \mathcal{S}(\mathbb{R}^n)$ there holds

$$\left[\int_{\mathbb{R}^n} |u|^q dx^n \right]^{1/q} \leq C \left[\int_{\mathbb{R}^n} |u|^p dx^n \right]^{t/p} \left[\int_{\mathbb{R}^n} \sum_{|\alpha| \leq 1} |\partial^\alpha u|^n dx^n \right]^{(1-t)/n}, \quad (4.2)$$

⁵The space of Schwartz function in \mathbb{R}^n is the space of C^∞ functions whose derivatives of arbitrary order decrease faster than any polynomial at infinity. It is preserved by the Fourier transform.

⁶These spaces were discovered by Otto Ludwig Hölder in his dissertation in 1882. He had been asked to characterize the optimal regularity of the charge distribution f that would ensure that the potential u solving (3.4) have continuous second derivatives. The continuity of f is not enough and he discovered that the existence of $\alpha > 0$ such that

$$\sup_{x \neq y} \frac{|f(x) - f(y)|}{|x - y|^\alpha} < +\infty \quad (3.5)$$

is a sufficient condition for u to be C^2 . This norm is denoted $\|f\|_{C^{0,\alpha}}$.

where $t = p q^{-1}$. The discovery of these estimates, together with the pioneering works of Marcel Riesz, Olaf Thorin, and Józef Marcinkiewicz, lies at the origin of an important subfield of functional analysis developed by Jacques-Louis Lions, Jaak Peetre (see [53]) and Alberto Calderón ([16]), and called *interpolation theory*.

4.2. The use of Gagliardo–Nirenberg inequalities for proving a priori estimates

The notion of a priori estimates is central in PDEs. Roughly speaking, it consists in finding an estimate of the form

$$\|u\|_E < +\infty,$$

for some suitable Banach space E , all the while assuming we have a solution u of some PDE, but prior to having actually proved the existence of such a solution. In concrete situations, looking at a given non-linear PDE problem, one establishes such an a priori bound in order to perform one of the numerous available analytical methods in order to finally prove the existence of a solution satisfying that bound: fixed point argument in a perturbative approach, continuity method⁷, topological techniques (e.g. Leray–Schauder theory), functional analysis approaches (e.g. monotone operator theory, Hille Yosida), successive approximation (e.g. Galerkin method, convex integration), penalization approaches (e.g. elliptization or viscosity method), variational approaches (e.g. minimization, min-max methods, Morse theory).

Gagliardo–Nirenberg inequalities are mostly used to control non-linear terms in PDEs, and to establish *a priori estimates*. There are countless applications for these inequalities. In order to illustrate their might in dealing with non-linearities, the author of these notes is cherry-picking a subject dear to his heart: the Dirichlet problem for maps taking values into submanifolds, also called *harmonic map* or *vectorial Dirichlet problem*. This problem has many applications in geometry (in minimal surface theory and in complex geometry – it is used to describe the Teichmüller space of 2-dimensional surfaces) as well as in physics (for instance the Dirichlet problem for maps taking values into the sphere is the main mathematical object of the Ericksen–Leslie model of liquid crystals).

We consider exactly the same problem as the one posed by Dirichlet, that is to find critical points of the Dirichlet energy E on a smooth bounded domain Ω for some boundary condition, but this time under the additional constraint that the “unknown” u takes values in a given closed sub-manifold $N^n \subset \mathbb{R}^m$ of the Euclidean space. Having fixed $\phi \in C^1(\partial\Omega, N^n)$, we look for a critical point u of E with the constraint that $u \in N^n$ everywhere. This constraint generates a new equation generalizing the Laplace equation (3.1)

⁷This method, probably first used by Serge Bernstein in 1905, consists in “deforming” continuously a given PDE problem to one which is directly solvable, in such a way as to preserve the *a priori estimates*. The continuity method applied to elliptic non-linear problems is roughly done as follows. Once the *a priori estimate* is established, one linearizes the problem with respect to the deformation parameter, then calls upon the linear Agmon–Douglis–Nirenberg L^p theory or the Schauder $C^{0,\alpha}$ theory, and finally uses the local inversion theorem to ensure the “continuation” of the existence of the solution as the deformation parameter evolves.

and one shows that the problem is equivalent to

$$\begin{cases} \Delta u \in T_{u(x)}N^n & \text{in } \Omega, \\ u = \phi & \text{on } \partial\Omega, \end{cases} \quad (4.3)$$

where $T_{u(x)}N^n$ denotes the tangent space to N^n in \mathbb{R}^m at the point $u(x)$. The equation $\Delta u \in T_{u(x)}N^n$ is a natural generalization of the Laplace equation previously obtained for the unconstrained *Dirichlet problem*. Indeed, it says that the tangential component of Δu to the constraint is zero, which simply implies that $\Delta u \equiv 0$ when there is no constraint. The problem (4.3) can be recast as an elliptic non-linear equation of the form

$$\begin{cases} -\Delta u = f(u, \nabla u) & \text{in } \Omega, \\ u = \phi & \text{on } \partial\Omega. \end{cases} \quad (4.4)$$

This PDE is said to be *semi-linear*, because the term involving the highest order derivatives is linear. In contrast with the original Dirichlet problem (3.1), one can show that (4.4) sometimes has more than one solution (and sometimes has infinitely many of them! or even no solutions at all for the general form (4.4)).

To what extent is this non-linear Dirichlet problem degenerate, and what sufficient conditions would ensure that (4.4) is a well posed *Cauchy problem*? Taking two solutions u and v , the difference $u - v$ satisfies

$$u - v = (-\Delta)_0^{-1}(f(u, \nabla u) - f(v, \nabla v)). \quad (4.5)$$

In order to proceed to a “contraction mapping argument” we need to find a function space E in which the difference of the non-linearities $(-\Delta)_0^{-1}(f(u, \nabla u) - f(v, \nabla v))$ is controlled by the norm of the difference $u - v$ in the same space, weighted by a constant $k < 1$:

$$\|(-\Delta)_0^{-1}(f(u, \nabla u) - f(v, \nabla v))\|_E \leq k\|u - v\|_E. \quad (4.6)$$

Such an inequality, also called an *a priori estimate* – since we have not yet settled the question of existence for either u or v – offers the possibility to absorb the non-linear right-hand side of (4.5) into the linear left-hand side $u - v$, and in turn deduce that $u = v$.

The *Gagliardo–Nirenberg interpolation inequalities* are essential to prove such *a priori estimates* as (4.6). Our example, picked among countless others, was chosen to illustrate the possibility to solve Cauchy problems for non-linear PDEs. It is however representative of the power of the Gagliardo–Nirenberg inequalities.

Among the various inequalities pertaining to *vectorial Dirichlet problems*, there is a particularly elegant one⁸, which, in combination with the Agmon–Douglis–Nirenberg L^p theory, enables us to show the well-posedness of the *vectorial Dirichlet problem* for two dimensional domains under small energy assumptions. Namely, there exists a constant

⁸This inequality, which is a particular case of (4.2), also appears in the work of Olga Ladyzenskaya.

$C(\Omega) > 0$ such that for any smooth function w supported in the two dimensional domain Ω the following inequality holds:

$$\int_{\Omega} |w|^4 dx^2 \leq C(\Omega) \int_{\Omega} |w|^2 dx^2 \int_{\Omega} |\nabla w|^2 dx^2. \quad (4.7)$$

This inequality was centrally used by Michael Struwe in the framework of the *vectorial Dirichlet problem* to obtain *a priori estimates* that imply the existence and uniqueness for the corresponding flow, also called *harmonic map flow*, in two dimensions; see [70].

5. The John–Nirenberg BMO space: When elasticity meets harmonic analysis

The analysis of PDEs has evolved and keeps evolving in close “partnership” with the development of functional analysis and function space theory. Many linear and non-linear problems in PDE have stimulated the introduction of new function spaces – we have already outlined above the importance of the Sobolev spaces for solving the Dirichlet problem. The converse is also true: knowledge and properties of certain function spaces can trigger a new understanding of PDE problems.

In 1961, the mathematician – and colleague of Louis Nirenberg at the Courant Institute in New York – Fritz John, was studying a rigidity problem from elasticity [41]. The strain exerted on a perfect elastic solid can be measured by the distance of the gradient of the resulting deformation with respect to the orthogonal group. In relation with this fundamental notion in elasticity, he asked the following question:

Is it true that if the gradient of a transformation f from Euclidean space \mathbb{R}^n into itself is “close to” the group of rotations at every point, then it is globally close to one single rotation?

By “close to”, it was originally meant in the L^∞ norm. This rigidity question finds its origin in a work of the mathematician Arthur Korn from 1914 [47]. Extending Korn’s results and now celebrated inequalities, Kurt Friedrichs⁹ proved in an important work in 1947 (see [33]) that the gradient of a deformation is everywhere antisymmetric if and only if it is constant, provided that suitable boundary conditions, which exclude rigid motions, are imposed. This result is an infinitesimal version of the question asked by John.

F. John first gave a relatively straightforward counter-example to the L^∞ version of the question, but he also proved that on every ball $B_r(x) \subset \mathbb{R}^n$ of arbitrary center x and

⁹The name Kurt Otto Friedrichs is strongly associated with the Courant Institute for having contributed to its development since its very beginning in the mid 1930s.

radius $r > 0$, there exists a rotation $R_{x,r}$ such that

$$\frac{1}{|B_r(x)|} \int_{B_r(x)} |\nabla f - R_{x,r}| dx^n \leq C \|\text{dist}(\nabla f, SO(n))\|_\infty,$$

where $|B_r(x)|$ is the volume of the ball $B_r(x)$. Thus, John was proving that although the gradient of such a deformation could not be close to one single rotation globally, it is in average in the L^1 -norm at any scale close to a rotation $R_{x,r}$ that possibly depends on the ball.

In a subsequent collaboration [42], which has since become a milestone in analysis, Fritz John and Louis Nirenberg systematically studied the sub-space of locally integrable functions, called the space of functions of *Bounded Mean Oscillation* (BMO), whose elements satisfy

$$\sup_{x \in \mathbb{R}^n, r > 0} \frac{1}{|B_r(x)|} \int_{B_r(x)} |u - \bar{u}_{x,r}| dx^n < +\infty,$$

where $\bar{u}_{x,r}$ is the average of u on the ball $B_r(x)$. They proved that this space is strictly larger than L^∞ , the space of globally bounded functions (e.g. $\log|x| \in \text{BMO} \setminus L^\infty$), but that it is smaller than L^p_{loc} for any $p < +\infty$. Precisely following an ingenious decomposition of \mathbb{R}^n of *Calderón–Zygmund type* they proved the existence of $\alpha_n > 0$ such that for any ball $B \subset \mathbb{R}^n$

$$\int_B \exp\left[\alpha_n \frac{|u - \bar{u}|}{\|u\|_{\text{BMO}}}\right] dx^n \leq C_n |B|,$$

where \bar{u} is the average of u on B . The existence of such a bound for any ball is proved to uniquely characterize the BMO space.

The BMO space, which naturally arose in the context of elasticity in 1960, was apparently unknown to functional analysts. It was therefore a big surprise to discover, after the remarkable work of Elias Stein and Charles Fefferman in 1972 ([32]), that BMO was the Banach dual of a famous space introduced in complex function theory some 40 years earlier by Friedrich Riesz [67] and named “Hardy space” after a famous work by Godfrey Hardy from 1915 [36]. Historically, the Hardy space \mathcal{H}^1 was defined in the context of holomorphic functions on the disc D^2 : it is made of the traces on the circle S^1 of holomorphic functions f such that

$$\lim_{r \rightarrow 1^-} \int_0^{2\pi} |f(re^{i\theta})| d\theta < +\infty.$$

Note that this integral is an increasing function of the parameter r for any holomorphic function f . The Hardy space was later extended to a much broader context of real function space theory. The dual spaces \mathcal{H}^1 and BMO play a fundamental role in PDEs. Empirically, one could say that they are the “natural replacements” for L^1 and L^∞ . These two spaces are not compatible with Calderón–Zygmund theory, and in fact the Agmon–Douglis–Nirenberg results do not hold either for L^1 nor for L^∞ . In contrast, \mathcal{H}^1 and BMO are well behaved in these theories.

It is unfortunately beyond the scope of this presentation to expose in its full glory the usefulness of the duality $\mathcal{H}^1 - \text{BMO}$ in the analysis of PDEs. It plays a central role for instance in a theory called *integrability by compensation*, where some non-linear quantities appear in the form of products with such algebraic structures that improved integrability might be deduced. This has frequently been used in fluid mechanics as well as in geometry.

We content ourselves with mentioning one application. Using some straightforward integration by parts, one proves the following *Gagliardo–Nirenberg interpolation inequality*: for any $n \in \mathbb{N}^*$ there exists $C_n > 0$ such that for all Schwartz functions u in $\mathcal{S}(\mathbb{R}^n)$ one has

$$\int_{\mathbb{R}^n} \sum_{|\alpha|=1} |\partial^\alpha u|^4 dx^n \leq C_n \|u\|_\infty^2 \int_{\mathbb{R}^n} \sum_{|\alpha|=2} |\partial^\alpha u|^2 dx^n.$$

Using more sophisticated arguments (Littlewood-Paley decomposition of tempered distributions) the author of these notes in collaboration with Yves Meyer prove a “slight” improvement of this inequality (see [56]) by replacing the L^∞ norm of u with its BMO-norm, namely:

$$\int_{\mathbb{R}^n} \sum_{|\alpha|=1} |\partial^\alpha u|^4 dx^n \leq C_n \|u\|_{\text{BMO}}^2 \int_{\mathbb{R}^n} \sum_{|\alpha|=2} |\partial^\alpha u|^2 dx^n.$$

This improved inequality entering in a larger class of inequalities by David Adams and Michael Fraizer (see [1]) has been crucially used in [56] for proving a partial regularity result for *stationary Yang–Mills fields*, which was obtained independently by Terence Tao and Gang Tian; see [71].

6. The maximum principle

6.1. Nirenberg’s strong maximum principle for parabolic equations

It would be impossible to speak about the work of Louis Nirenberg without mentioning the *maximum principle*. The contrast between the immense range of applications of this principle and the simplicity of the heuristic idea behind it is amazing.

In one dimension, the maximum principle states that a continuously twice differentiable function on the segment $[0, 1]$ satisfying

$$u'' \geq 0 \quad (\text{resp.}, u'' \leq 0) \quad \text{on } [0, 1]$$

achieves its maximal (respectively, minimal) value on the boundary of the segment, i.e. on $\{0\} \cup \{1\}$.

In higher dimension, the maximum principle was known to Gauss since 1839 for solutions of the Laplace equation, owing to the mean value theorem for harmonic functions:

A solution to the Laplace equation on a bounded smooth domain achieves its extremal values on the boundary of the domain.

This formulation requires u to solve a PDE, and it is only at the beginning of the 20th century that, after successive contributions by Charles Émile Picard (1905) [66], Serge Bernstein (1910) [8] and Leon Lichtenstein (1924) [52], the idea of a general principle for elliptic partial differential *inequalities* emerged.

In a seminal five-page long paper published in 1927 [37], Eberhard Hopf opened the way to a wide range of applications of this principle by proving a general *strong version of the maximum principle* for C^2 solutions to the following linear elliptic second order inequalities on a smooth bounded domain $\Omega \subset \mathbb{R}^n$:

$$Lu := \sum_{i,j=1}^n a_{ij}(x) \frac{\partial^2 u}{\partial x_i \partial x_j} + \sum_{i=1}^n b_i(x) \frac{\partial u}{\partial x_i} + c(x)u \geq 0, \quad (6.1)$$

where $a_{ij}(x)$ is a map into uniformly positive definite symmetric matrices, and the coefficient a_{ij} as well as b_i and c are bounded in $L^\infty(\Omega)$, with moreover $c(x) \leq 0$ in Ω . A C^2 function u satisfying (6.1) is called a *sub-solution of L* (and a *super-solution of L* if $Lu \leq 0$).

As in the 1-dimensional case, one formulates the *weak version of the maximum principle* as follows:

$$Lu \geq 0 \text{ in } \Omega \Rightarrow \max_{x \in \Omega} u(x) \leq \max_{x \in \partial\Omega} u(x),$$

whereas the *strong version of the maximum principle* discovered by Eberhard Hopf asserts that

$$Lu \geq 0 \text{ in } \Omega \quad \text{and} \quad \exists x_0 \in \Omega \quad \text{s.t.} \quad \max_{x \in \Omega} u(x) = u(x_0) \Rightarrow u(x) \equiv u(x_0) \text{ in } \Omega.$$

In the early 1950s, a weak version of this principle was known to hold for *parabolic operators* of the type $\partial_t u - Lu$. In 1953, Louis Nirenberg proved the corresponding strong version; see [59].

6.2. The notion of “barriers”, the “moving planes” method, and the Gidas–Ni–Nirenberg symmetry principle

The heuristic idea behind the strong maximum principle, at least in the simpler elliptic framework, has an interesting geometric representation. We say that a linear elliptic operator L satisfies the strong maximum principle if the following holds. Let any pair of hyper-surfaces realized by two graphs of respectively a sub-solution u and super-solution v over a bounded domain, with one of them sitting above the other (i.e. $u \geq v$) be given. The strong maximum principle says that if they touch at some *interior* point, then the two hyper-surfaces are necessarily identical.

A *sub-solution* satisfying $Lu \geq 0$ is then said to be a “barrier” with respect to a *super solution* satisfying $Lv \leq 0$, and vice versa. This geometric interpretation is an incentive

to manufacture *barriers* with respect to *solutions*, *sub-solutions*, and to *super-solutions* in order to prove *pointwise inequalities* via the maximum principle. This fruitful technique has become a classic in analysis, where it is known as a *comparison argument*. Devising suitable *barriers* is nothing short of being an art in itself. It requires deep intuition and thorough experience of the problems considered. Louis Nirenberg was notoriously known, as it has been reported by his collaborators, for being a master in conceiving barrier solutions.

The geometric interpretation of the maximum principle in both its strong and weak formulations was probably first used in the 1955 work of Alexander Danilowitsch Alexandrov; see [4] for a later publications of his original ideas that he presented for the first time in Zürich in 1955 according to Heinz Hopf [38]. He used a *comparison argument* between the solution itself and some reflections of it in order to prove that embedded constant mean curvature closed surfaces in \mathbb{R}^3 are necessarily isometric to a dilation of the unit sphere S^2 . The relevance of the maximum principle and of elliptic theory in the resolution of this geometric problem is apparent in the equation satisfied by the graph u of constant mean curvature H , namely:

$$\operatorname{div} \left[\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right] = 2H.$$

Following an important paper by James Serrin [69], Louis Nirenberg in collaboration with Basilis Gidas and Wei Ming Ni, converted Alexandrov’s original idea for constant mean curvature surfaces into a general method, as beautiful as it is efficient, nowadays known as the “moving planes method”; see [35]. With it, one can prove symmetry results (either with respect to a given direction, or full rotational symmetry) and uniqueness results for positive solutions to semi-linear scalar equations of the form

$$-\Delta u = f(u). \tag{6.2}$$

These symmetry and uniqueness results are of utmost importance, since they extend to a non-linear framework a fundamental principle in quantum mechanics and in spectral theory; stating that the ground state of the Laplace operator, which is necessarily positive,

$$-\Delta u = \lambda_1 u$$

enjoys special symmetries and has multiplicity one (i.e. it is unique – Krein–Rutman theorem).

This method and the ensuing symmetry results have important applications to diverse areas of science: the study of ground states of non-linear Schrödinger models in quantum mechanics, the vortex theory of Onsager in thermodynamics, turbulence in statistical physics, phase-transitions in Van der Waals fluids, the Yamabe problem in differential geometry (which is concerned with finding constant scalar curvature metrics in a given conformal class), etc.

The moving planes method consists in comparing an arbitrary solution u for the semilinear equation (6.2) with its successive reflections u_λ across a continuous family of parallel hyper-planes. These reflections u_λ are used as barrier functions for u . A key ingredient of the method is the strong version of the maximum principle and a refinement of it discovered by E. Hopf in the 1950s, and now known as “Hopf lemma”. It states that at maximum points on the boundary, the outward normal derivative of a *sub-solution* is strictly positive¹⁰ unless the *sub-solution* is identically constant.

Later on, Louis Nirenberg in collaboration with Henri Berestycki introduced a new method, still based on the *strong maximum principle*, and called the “sliding method”; see, for instance, [6]. This new method was devised to prove various pointwise estimates and asymptotic behaviors for solutions in infinite cylindrical domains to semi-linear equations of the form (6.2), as well as for their parabolic counterparts. Its extension to the case of finite cylinder was then proposed by Luis Caffarelli. The *sliding method* has numerous important applications to traveling fronts problems in the mathematical modelling of combustion and flame propagation. Its novel idea consists in comparing the solution with its translations along the axis of the cylinder, rather than using the images by successive symmetries of the solution as *barriers*. It requires the use of a version of the maximum principle on “narrow domains” where the sign of the 0th order coefficient $c(x)$ is not required to be controlled due to S. R. Srinivasa Varadhan; see [7].

6.3. The Dirichlet problem for non-linear second order elliptic equations

We have stressed the importance of *a priori estimates* for solving the Dirichlet problem of semi-linear equations. In order to prove such estimates, we have combined the Agmon–Douglis–Nirenberg L^p -theory for boundary-value problems along with the Gagliardo–Nirenberg estimates in various Banach spaces. For many scalar equations of elliptic type which are more non-linear and also more degenerate than semi-linear equations, the maximum principle is an additional tool that can be added into the mix to reach the desired estimates.

In a series of five fundamental papers written in collaboration with Luis Caffarelli and Joel Spruck, Louis Nirenberg identified the *maximum principle* as a fundamental device to obtain *a priori* estimates, and to solve the Dirichlet problem for “highly” non-linear PDEs known as *fully non-linear PDEs*; see [10, 12–15].

An example of such an equation is the *Monge–Ampère* equation, which appears in problems related to optimal transport as well as in geometric problems of prescribed curvature.

Theorem 6.1 (Caffarelli–Nirenberg–Spruck (1984)). *Let Ω be a strictly convex domain. For any smooth and positive function f in $\bar{\Omega}$, and for any smooth function ϕ in $\bar{\Omega}$, there*

¹⁰This strict positivity of the normal outward derivative at the maximal point was already proved by Leon Lichtenstein in 1923 in [52] for exact solutions of $Lu = 0$ in 2 dimensions.

exists a unique strictly convex C^∞ solution u to

$$\begin{cases} \det \nabla^2 u = f & \text{in } \Omega, \\ u = \phi & \text{on } \partial\Omega. \end{cases}$$

The strict convexity requirement imposed on the solution forces ellipticity on the problem. To see this, let $F(\partial^2 u) := \log \det \nabla^2 u$. Let $(u^{ij})_{i,j=1,\dots,n}$ denote the inverse of the Hessian matrix of u . Assuming that u is convex, the matrix (u^{ij}) is symmetric and positive definite. Hence,

$$\sum_{i,j}^n \frac{\partial F}{\partial r_{ij}}(r) \xi_i \xi_j = \sum_{i,j} u^{ij} \xi_i \xi_j > 0,$$

which is tantamount to ellipticity.

The core of the argument involves several steps. First, one establishes *a priori estimates* for the Hölder norm of the solution (assumed to exist). These estimates follow from the classical elliptic Agmon–Douglis–Nirenberg L^p -theory or Schauder theory for Hölder spaces, provided that $F(\partial^2 u)$ is *uniformly elliptic*:

$$\lambda |\xi|^2 \leq \frac{\partial F}{\partial r_{ij}} \xi_i \xi_j \leq \Lambda |\xi|^2,$$

for some constants $0 < \lambda < \Lambda$, and also provided that the modulus of continuity of $\partial_r F(\partial^2 u)$ can be controlled up to the boundary. In other words, the task is to prove that $\partial^2 u$ is uniformly bounded and uniformly continuous up to the boundary. This is done with the help of the maximum principle and comparison arguments. Eventually, the authors reach the *a priori estimate*¹¹

$$\|\partial^2 u\|_{C^{0,\alpha}(\bar{\Omega})} \leq C < +\infty, \quad (6.3)$$

where $C^{0,\alpha}$ is the Hölder norm defined in (3.5). Once (6.3) is established, the final part of the argument relies on a *continuity method*, where one interpolates ϕ with the determinant of the Hessian matrix of a function which coincides with ϕ on the boundary.

In these highly intricate works, the *maximum principle* is showing its full quintessence and potential to obtain hidden *a priori estimates*. The core of the argument involves the sophisticated construction of *barrier functions* with which the solution u and its successive derivatives (or finite differences) are compared.

These five papers by Caffarelli, Nirenberg, Spruck and also by Joseph Kohn for one of them, have stimulated a tremendous amount of research activity on *fully non-linear equations* since their publications. These equations have an immense range of applications in many fields of science, including material sciences, finance, and computer vision.

¹¹This bound was independently obtained by Nikolai Vladimirovich Krylov [48, 49], using works of Lawrence Craig Evans [30, 31], and Neil Trudinger [72, 73].

The original ideas of Nirenberg et al. have influenced the development of a whole branch of analysis, called *viscosity theory for PDEs*, where the *maximum principle* plays a central and decisive role. The viscosity theory for PDEs was introduced by Lawrence Craig Evans [29] and by Michael Crandall and Pierre-Louis Lions [20].

7. Solving problems from geometry

The analysis of PDEs and differential geometry are by essence intimately intertwined. The central roles played by the Laplace operator and by the $\bar{\partial}$ -operator in Riemann surface theory, constitutes the simplest illustration of this connection. The second half of the 20th century saw a dramatic acceleration of the transfer of techniques from non-linear PDEs to the resolution of problems that seem a priori confined to geometry. A spectacular example of the might of the PDE approach in geometry is the recent proof of the Poincaré conjecture by Grigori Perelman, which heavily relies on the parabolic Ricci flow devised by Richard Hamilton.

7.1. Nirenberg's resolution of the Weyl problem

The taste for geometry and the influence of geometric questions are manifest in Louis Nirenberg's work. He is among the pioneers who introduced elaborate analysis tools for solving questions pertaining to embeddings, tensors, curvature, complex structures, etc. His doctoral work itself dealt with geometry, and, following the invitation from his advisors James Stoker and Kurt Friedrichs, he solved a problem originally posed by Hermann Weyl:

Given a metric of positive Gauss curvature on the 2-dimensional sphere, does there exist an isometric embedding of this sphere into a convex surface of \mathbb{R}^3 ?

Nirenberg answered this question assuming the given metric is four times continuously differentiable¹² in [60]. In this first work, the general “philosophy” we have discussed is already present, and it will remain recurrent in Louis Nirenberg's papers: one looks for *a priori estimates*, and combine them with suitable *continuity methods* that leave the *a priori estimates* unchanged along the deformation. The link with the previous section is made by writing the PDE satisfied by a graph $(x, u(x))$ of Gaussian curvature $K(x)$ at $(x, u(x))$:

$$\det(\partial^2 u) = K(1 + |\partial u|^2)^2.$$

Its associated Dirichlet problem was the subject of the last part of the first paper of the aforementioned series by L. Caffarelli, L. Nirenberg, and J. Spruck.

¹²Several mathematicians had already considered this question. Hans Lewy in the case of analytic metrics, and more generally Aleksei Pogorelov who solved the problem with the help of different methods.

7.2. The Nirenberg problem

The original “Nirenberg problem” can be stated as follows:

For which function K on S^2 does there exist a metric which is pointwise conformal to the canonical metric g_0 , and such that K is the Gauss curvature of that metric?

By “pointwise conformal”, we mean the existence of a function u on S^2 such that $g = e^{2u} g_0$. Stated differently, the *Nirenberg problem* consists in establishing the existence (or lack thereof) of a solution to the Liouville equation

$$-\Delta_{g_0} u = e^{2u} K - 1,$$

where Δ_{g_0} is the *negative Laplace Beltrami* operator for g_0 , the canonical metric of S^2 . This simply formulated question has brought forth an enormous amount of work since the early 1970s. Not only because it is the “simplest” instance of a wide range of similar questions (higher dimension, different curvature tensor, scalar curvature, Q -curvature, σ_k -curvatures, fractional curvatures, etc.), but also because it gives rise to the major issues faced by conformal geometric analysts in their study of “critical non-linear PDEs”, such as *concentration of compactness phenomena*, *Palais–Smale sequences*, *Morse theory*, inter alia. These issues appear as well in many celebrated problems: the Yamabe problem, harmonic map theory, Yang–Mills equations, constant mean curvature surfaces, Willmore surfaces to name a few. The apparent simplicity of the Nirenberg problem fosters the universal difficulties arising in conformal geometric analysis.

It would be much beyond the scope of the present report to give a detailed account of the various arguments and creativity which have flourished in the quest for solving Nirenberg’s problem. This problem is moreover at the origin of the whole branch of analysis devoted to the study of critical elliptic problems which is central in conformal differential geometry. We content ourselves to close the presentation of the *Nirenberg Problem* with mentioning that not every choice for K gives rise to a solution. This is seen, for example, using the Gauss–Bonnet theorem, and the now well-known Kazdan–Warner necessary condition. Let us also mention an important sufficient condition for the existence due to Alice Sun-Yung Chang and Paul Yang [19]. Let K be positive on S^2 with only non-degenerate critical points and with at least two local maxima. If in addition $\Delta_{g_0} K > 0$ at all saddle points of K , then Nirenberg’s problem has a solution.

7.3. The Newlander–Nirenberg complex Fröbenius theorem

Louis Nirenberg made important contributions to complex geometry and complex analysis. Once again, PDE techniques lie at the heart of the approach he favored to tackle various geometric questions.

An important one deals with the *integrability of almost complex structures*. The question goes as follows. Let a map J from \mathbb{R}^{2n} into the space of real-valued $n \times n$ matrices be given. Assume it satisfies everywhere the condition

$$J^2 = -I_n,$$

where I_n is the $n \times n$ identity matrix.

Does there exist a local diffeomorphism w in a neighborhood of any point which transports the endomorphism J into multiplication by i everywhere, after the canonical identification $\mathbb{R}^{2n} \simeq \mathbb{C}^n$?

For $n = 1$, the question amounts to solving locally a Beltrami equation

$$\partial_{\bar{z}} w = \mu \partial_z w.$$

This is successfully achieved by introducing the singular integral operator associated to the inverse of the Cauchy–Riemann operator $\bar{\partial}^{-1}$ and by using an elementary fixed point argument.

The general case $n > 1$ is much more involved and leads to an overdetermined system of coupled linear Cauchy–Riemann type PDEs of the form

$$\partial_{\bar{z}_j} w = \sum_{k=1}^n \mu_{jk} \partial_{z_k} w, \quad (7.1)$$

where z_j are complex coordinates for the complex structure given by the value of J at the point in the neighborhood of which we are working. The system being overdetermined (indeed, differentiating in \bar{z}_k the j th equation has to give the same result as differentiating in \bar{z}_j the k -th equation, namely: $\partial_{\bar{z}_j} \partial_{\bar{z}_k} w = \partial_{\bar{z}_k} \partial_{\bar{z}_j} w$) there must be a structural constraint on the system for it to be *locally solvable*. A necessary condition was discovered by Albert Nijenhuis in his 1951 doctoral thesis, and independently by Paulette Libermann in [51], as well as by Beno Eckmann and Alfred Frölicher in [24]¹³. It can be stated as follows. We consider the complexification of the tangent space to \mathbb{R}^{2n} at each point, and we call a $(0, 1)$ vector any vector of the complexified tangent space and of the form

$$X - iJX,$$

where X is a real vector. The necessary condition says that the space of $(0, 1)$ vectors has to be stable under bracket operation. Louis Nirenberg and his student A. Newlander succeeded in proving that this *complex Fröbenius* type condition is in fact sufficient for the local solvability in the C^∞ framework.

Note that, unlike in the one-dimensional case, applying the inverse of the Cauchy–Riemann operator $\partial_{\bar{z}_j}$ to the right-hand side of (7.1) does not enable an iteration argument in any classical Banach space (Hölder, Sobolev). Indeed, an integration in z_j is applied to a derivative in z_k of w , for $k \neq j$: this leads nowhere.

Nirenberg and Newlander in [58] had the idea to recast the problem in terms of the inverse $z(w)$ of the diffeomorphism $w(z)$. It solves a PDE system of the form

$$\partial_{\bar{w}_l} z = F(w) \partial_{w_l} z.$$

¹³In these works, using the Cauchy–Kowalevski approach, this condition is also shown to be sufficient in the analytic framework.

Although this problem is now becoming non-linear (!) unlike the original one which was linear, it is this time possible to implement a scheme similar to the one devised in the one-dimensional case, since differentiation occurs with respect to only one of the independent variables in each equation. This problem is solvable in classical Hölder spaces.

We have decided to end this presentation with this remarkable work, which can but trigger the admiration of any mathematician, even beyond the field of partial differential equations, and this more than 58 years after its original publication!

8. Conclusion

At the end of these notes, one feels somewhat frustrated to have only presented one part of the prolific and monumental work of Louis Nirenberg. Many important contributions have been omitted such as the analyticity of solutions to analytic PDEs (in collaboration with Charles Morrey [57]), the regularity of free-boundary problems (in collaboration with David Kinderlehrer and Joel Spruck [43–45]), the analysis of solutions to non-linear elliptic equations involving critical Sobolev exponents (in collaboration with Haïm Brezis [9]), the partial regularity of solutions to the Navier–Stokes equation (in collaboration with Luis Caffarelli and Robert Kohn [11]) which to this day remains the optimal step towards solving the Millennium problem. The author of these notes also apologizes to the numerous collaborators of Louis Nirenberg whose joint works with him have not been mentioned here.

Louis Nirenberg’s scientific endeavor is an exemplary reminder to all of us that research is first and foremost a collective venture, in which debating, discussing, and exchanging ideas play a decisive role.

It is the result of no coincidence that Louis Nirenberg made his professional home at the Courant Institute at New York University, a prestigious institution that has fostered since its very creation a unique laboratory for the free exchange of scientific ideas.

Although there are still many important theoretical questions to answer¹⁴, the analysis of partial differential equations is nowadays mostly aimed at better understanding other fields of science, with applications in geometry, physics, mechanics, chemistry, biology, social sciences, technology. These developments, and the ones to come, anchor their roots on the immense efforts deployed in the last century by human intelligence in this area of mathematics. Mathematical knowledge is however not only made of an accumulation of truths and results confined to papers and books, and transmitted in this form to future generations. A large and immaterial share of mathematical knowledge resides in the “living part” of mathematics, in mathematicians themselves, with their intuitions, their hesitations, their perseverance, and most importantly with their esthetical quest and search for beauty (as surprising as it may sound to non-mathematicians!). Hermann Weyl once said:

¹⁴pertaining, inter alia, to very non-linear PDEs, degenerate PDEs, non-local PDEs, rough data, PDEs in connection with stochastics, and much more.

My work always tried to unite the truth with the beautiful, but when I had to choose one or the other, I usually chose the beautiful.

We do not know whether Louis Nirenberg would have had agreed with this quote, but we would nonetheless like to pay tribute to his memory for the beautiful mathematics he has produced all along his life and has generously shared with us all for so many years.

Acknowledgments. The author is very grateful to the numerous colleagues who sent him remarks and comments about this presentation which was written for a big part on the occasion of the award of the Abel prize to Louis Nirenberg in May 2015 in Oslo.

A condensed version of this article, based on the Abel lecture the author delivered in Oslo in May 2015, appeared in [68] and he would like to take this occasion to thank Frank Morgan for his input on the text at the time.

A very special thanks goes to Sylvia Serfaty and Alain Sznitman as well as to the anonymous referees for their careful reading, corrections and suggestions that contributed a lot to the improvement of these notes.

References

- [1] D. R. Adams and M. Frazier, Composition operators on potential spaces. *Proc. Amer. Math. Soc.* **114** (1992), no. 1, 155–165 Zbl [0753.47045](#) MR [1076570](#)
- [2] S. Agmon, A. Douglis, and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I. *Comm. Pure Appl. Math.* **12** (1959), 623–727 Zbl [0093.10401](#) MR [125307](#)
- [3] S. Agmon, A. Douglis, and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. II. *Comm. Pure Appl. Math.* **17** (1964), 35–92 Zbl [0123.28706](#) MR [162050](#)
- [4] A. D. Alexandrov, A characteristic property of spheres. *Ann. Mat. Pura Appl. (4)* **58** (1962), 303–315 Zbl [0107.15603](#) MR [143162](#)
- [5] R. Beals and C. Fefferman, On local solvability of linear partial differential equations. *Ann. of Math. (2)* **97** (1973), 482–498 Zbl [0256.35002](#) MR [352746](#)
- [6] H. Berestycki and L. Nirenberg, On the method of moving planes and the sliding method. *Bol. Soc. Brasil. Mat. (N.S.)* **22** (1991), no. 1, 1–37 Zbl [0784.35025](#) MR [1159383](#)
- [7] H. Berestycki, L. Nirenberg, and S. R. S. Varadhan, The principal eigenvalue and maximum principle for second-order elliptic operators in general domains. *Comm. Pure Appl. Math.* **47** (1994), no. 1, 47–92 Zbl [0806.35129](#) MR [1258192](#)
- [8] S. Bernstein, Sur la généralisation du problème de Dirichlet. *Math. Ann.* **69** (1910), no. 1, 82–136 Zbl [41.0427.02](#) MR [1511579](#)
- [9] H. Brezis and L. Nirenberg, Positive solutions of non-linear elliptic equations involving critical Sobolev exponents. *Comm. Pure Appl. Math.* **36** (1983), no. 4, 437–477 Zbl [0541.35029](#) MR [709644](#)
- [10] L. Caffarelli, J. J. Kohn, L. Nirenberg, and J. Spruck, The Dirichlet problem for non-linear second-order elliptic equations. II. Complex Monge–Ampère, and uniformly elliptic, equations. *Comm. Pure Appl. Math.* **38** (1985), no. 2, 209–252 Zbl [0598.35048](#) MR [780073](#)

- [11] L. Caffarelli, R. Kohn, and L. Nirenberg, Partial regularity of suitable weak solutions of the Navier–Stokes equations. *Comm. Pure Appl. Math.* **35** (1982), no. 6, 771–831
Zbl [0509.35067](#) MR [673830](#)
- [12] L. Caffarelli, L. Nirenberg, and J. Spruck, The Dirichlet problem for non-linear second-order elliptic equations. I. Monge–Ampère equation. *Comm. Pure Appl. Math.* **37** (1984), no. 3, 369–402
Zbl [0598.35047](#) MR [739925](#)
- [13] L. Caffarelli, L. Nirenberg, and J. Spruck, The Dirichlet problem for non-linear second-order elliptic equations. III. Functions of the eigenvalues of the Hessian. *Acta Math.* **155** (1985), no. 3–4, 261–301
Zbl [0654.35031](#) MR [806416](#)
- [14] L. Caffarelli, L. Nirenberg, and J. Spruck, non-linear second order elliptic equations. IV. Star-shaped compact Weingarten hypersurfaces. In *Current topics in partial differential equations*, pp. 1–26, Kinokuniya, Tokyo, 1986
Zbl [0672.35027](#) MR [1112140](#)
- [15] L. Caffarelli, L. Nirenberg, and J. Spruck, non-linear second-order elliptic equations. V. The Dirichlet problem for Weingarten hypersurfaces. *Comm. Pure Appl. Math.* **41** (1988), no. 1, 47–70
Zbl [0672.35028](#) MR [917124](#)
- [16] A.-P. Calderón, Intermediate spaces and interpolation. *Studia Math. (Ser. Specjalna) Zeszyt 1* (1963), 31–34
Zbl [0124.31803](#) MR [0147896](#)
- [17] A. P. Calderon and A. Zygmund, On the existence of certain singular integrals. *Acta Math.* **88** (1952), 85–139
Zbl [0047.10201](#) MR [52553](#)
- [18] A. P. Calderón and A. Zygmund, On singular integrals. *Amer. J. Math.* **78** (1956), 289–309
Zbl [0072.11501](#) MR [84633](#)
- [19] S.-Y. A. Chang and P. C. Yang, Prescribing Gaussian curvature on S^2 . *Acta Math.* **159** (1987), no. 3–4, 215–259
Zbl [0636.53053](#) MR [908146](#)
- [20] M. G. Crandall and P.-L. Lions, Viscosity solutions of Hamilton–Jacobi equations. *Trans. Amer. Math. Soc.* **277** (1983), no. 1, 1–42
Zbl [0599.35024](#) MR [690039](#)
- [21] N. Dencker, The resolution of the Nirenberg–Treves conjecture. *Ann. of Math. (2)* **163** (2006), no. 2, 405–444
Zbl [1104.35080](#) MR [2199222](#)
- [22] P. G. L. Dirichlet, *Vorlesungen über die im umgekehrten Verhältniss des Quadrats der Entfernung wirkenden Kräfte*, edited by F. Grube. Teubner, Leipzig, 1876
Zbl [08.0619.01](#)
- [23] P. G. L. Dirichlet, *G. Lejeune Dirichlet’s Werke. 2 Vols.*, edited by L. Kronecker and L. Fuchs. First published 1889–1897. Reprinted by Chelsea Publishing Company, New York, 1969
Zbl [0212.00801](#)
- [24] B. Eckmann and A. Frölicher, Sur l’intégrabilité des structures presque complexes. *C. R. Acad. Sci. Paris* **232** (1951), 2284–2286
Zbl [0042.40503](#) MR [42202](#)
- [25] L. Ehrenpreis, Solution of some problems of division. I. Division by a polynomial of derivation. *Amer. J. Math.* **76** (1954), 883–903
Zbl [0098.08401](#) MR [68123](#)
- [26] L. Ehrenpreis, Solution of some problems of division. II. Division by a punctual distribution. *Amer. J. Math.* **77** (1955), 286–292
Zbl [0064.11504](#) MR [70048](#)
- [27] L. Ehrenpreis, Solution of some problems of division. III. Division in the spaces \mathcal{D}' , \mathcal{H} , \mathcal{O}_A . *Amer. J. Math.* **78** (1956), 685–715
Zbl [0072.32801](#) MR [83690](#)
- [28] L. Ehrenpreis, Solution of some problems of division. IV. Invertible and elliptic operators. *Amer. J. Math.* **82** (1960), 522–588
Zbl [0098.08401](#) MR [119082](#)
- [29] L. C. Evans, On solving certain non-linear partial differential equations by accretive operator methods. *Israel J. Math.* **36** (1980), no. 3–4, 225–247
Zbl [0454.35038](#) MR [597451](#)
- [30] L. C. Evans, Classical solutions of fully non-linear, convex, second-order elliptic equations. *Comm. Pure Appl. Math.* **35** (1982), no. 3, 333–363
Zbl [0469.35022](#) MR [649348](#)

- [31] L. C. Evans, Classical solutions of the Hamilton–Jacobi–Bellman equation for uniformly elliptic operators. *Trans. Amer. Math. Soc.* **275** (1983), no. 1, 245–255 MR [678347](#)
- [32] C. Fefferman and E. M. Stein, H^p spaces of several variables. *Acta Math.* **129** (1972), no. 3–4, 137–193 Zbl [0257.46078](#) MR [447953](#)
- [33] K. O. Friedrichs, On the boundary-value problems of the theory of elasticity and Korn’s inequality. *Ann. of Math. (2)* **48** (1947), 441–471 Zbl [0029.17002](#) MR [22750](#)
- [34] E. Gagliardo, Ulteriori proprietà di alcune classi di funzioni in più variabili. *Ricerche Mat.* **8** (1959), 24–51 Zbl [0199.44701](#) MR [109295](#)
- [35] B. Gidas, W. M. Ni, and L. Nirenberg, Symmetry and related properties via the maximum principle. *Comm. Math. Phys.* **68** (1979), no. 3, 209–243 Zbl [0425.35020](#) MR [544879](#)
- [36] G. Hardy, On the mean value of the modulus of an analytic function, *Proc. London Math. Soc.* **14** (1915) 269–277 Zbl [45.1331.03](#)
- [37] E. Hopf, Elementare Bemerkungen über die Lösungen partieller Differentialgleichungen zweiter Ordnung vom elliptischen Typus. *Sitzungsberichte Akad. Berlin* (1927), 147–152 Zbl [53.0454.02](#)
- [38] H. Hopf, *Differential geometry in the large*. Lecture Notes in Math. 1000, Springer, Berlin, 1983 Zbl [0526.53002](#) MR [707850](#)
- [39] L. Hörmander, Pseudo-differential operators. *Comm. Pure Appl. Math.* **18** (1965), 501–517 Zbl [0125.33401](#) MR [180740](#)
- [40] L. Hörmander, Pseudodifferential operators of principal type. In *Singularities in boundary value problems (Proc. NATO Adv. Study Inst., Maratea, 1980)*, pp. 69–96, NATO Adv. Study Inst. Ser. C: Math. Phys. Sci. 65, Reidel, Boston, MA, 1981 Zbl [0459.35096](#) MR [617227](#)
- [41] F. John, Rotation and strain. *Comm. Pure Appl. Math.* **14** (1961), 391–413 Zbl [0102.17404](#) MR [138225](#)
- [42] F. John and L. Nirenberg, On functions of bounded mean oscillation. *Comm. Pure Appl. Math.* **14** (1961), 415–426 Zbl [0102.04302](#) MR [131498](#)
- [43] D. Kinderlehrer and L. Nirenberg, Regularity in free boundary problems. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **4** (1977), no. 2, 373–391 Zbl [0352.35023](#) MR [440187](#)
- [44] D. Kinderlehrer, L. Nirenberg, and J. Spruck, Regularity in elliptic free boundary problems. *J. Analyse Math.* **34** (1978), 86–119 Zbl [0402.35045](#) MR [531272](#)
- [45] D. Kinderlehrer, L. Nirenberg, and J. Spruck, Regularity in elliptic free boundary problems. II. Equations of higher order. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **6** (1979), no. 4, 637–683 Zbl [0425.35097](#) MR [563338](#)
- [46] J. J. Kohn and L. Nirenberg, An algebra of pseudo-differential operators. *Comm. Pure Appl. Math.* **18** (1965), 269–305 Zbl [0171.35101](#) MR [176362](#)
- [47] A. Korn, Über die Lösung des Grundproblemens der Elastizitätstheorie. *Math. Ann.* **75** (1914), no. 4, 497–544 Zbl [45.1015.01](#) MR [1511808](#)
- [48] N. V. Krylov, Degenerate non-linear elliptic equations (Russian). *Mat. Sb. (N.S.)* **120(162)** (1983), no. 3, 311–330 Zbl [0525.35038](#) MR [691980](#)
- [49] N. V. Krylov, Degenerate non-linear elliptic equations. II (Russian). *Mat. Sb. (N.S.)* **121(163)** (1983), no. 2, 211–232 Zbl [0538.35037](#) MR [703325](#)
- [50] H. Lewy, An example of a smooth linear partial differential equation without solution. *Ann. of Math. (2)* **66** (1957), 155–158 Zbl [0078.08104](#) MR [88629](#)
- [51] P. Libermann, Problèmes d’équivalence relatifs à une structure presque complexe sur une variété à quatre dimensions. *Acad. Roy. Belgique. Bull. Cl. Sci. (5)* **36** (1950), 742–755 Zbl [0041.50102](#) MR [0040791](#)

- [52] L. Lichtenstein, Neue Beiträge zur Theorie der linearen partiellen Differentialgleichungen zweiter Ordnung vom elliptischen Typus. *Math. Z.* **20** (1924), no. 1, 194–212
Zbl [50.0325.01](#) MR [1544673](#)
- [53] J.-L. Lions and J. Peetre, Sur une classe d'espaces d'interpolation. *Inst. Hautes Études Sci. Publ. Math.* (1964), no. 19, 5–68 Zbl [0148.11403](#) MR [165343](#)
- [54] B. Malgrange, Existence et approximation des solutions des équations aux dérivées partielles et des équations de convolution. *Ann. Inst. Fourier (Grenoble)* **6** (1955/56), 271–355
Zbl [0071.09002](#) MR [86990](#)
- [55] G. Métivier, Counterexamples to Hölmgren's uniqueness for analytic non-linear Cauchy problems. *Invent. Math.* **112** (1993), no. 1, 217–222 Zbl [0794.35026](#) MR [1207483](#)
- [56] Y. Meyer and T. Rivière, A partial regularity result for a class of stationary Yang–Mills fields in high dimension. *Rev. Mat. Iberoamericana* **19** (2003), no. 1, 195–219 Zbl [1127.35317](#)
MR [1993420](#)
- [57] C. B. Morrey, Jr. and L. Nirenberg, On the analyticity of the solutions of linear elliptic systems of partial differential equations. *Comm. Pure Appl. Math.* **10** (1957), 271–290
Zbl [0082.09402](#) MR [89334](#)
- [58] A. Newlander and L. Nirenberg, Complex analytic coordinates in almost complex manifolds. *Ann. of Math. (2)* **65** (1957), 391–404 Zbl [0079.16102](#) MR [88770](#)
- [59] L. Nirenberg, A strong maximum principle for parabolic equations. *Comm. Pure Appl. Math.* **6** (1953), 167–177 Zbl [0050.09601](#) MR [55544](#)
- [60] L. Nirenberg, The Weyl and Minkowski problems in differential geometry in the large. *Comm. Pure Appl. Math.* **6** (1953), 337–394 Zbl [0051.12402](#) MR [58265](#)
- [61] L. Nirenberg, On elliptic partial differential equations. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)* **13** (1959), 115–162 Zbl [0088.07601](#) MR [109940](#)
- [62] L. Nirenberg, An extended interpolation inequality. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)* **20** (1966), 733–737 Zbl [0163.29905](#) MR [208360](#)
- [63] L. Nirenberg and F. Trèves, Solvability of a first order linear partial differential equation. *Comm. Pure Appl. Math.* **16** (1963), 331–351 Zbl [0117.06104](#) MR [163045](#)
- [64] L. Nirenberg and F. Trèves, On local solvability of linear partial differential equations. I. Necessary conditions. *Comm. Pure Appl. Math.* **23** (1970), 1–38 Zbl [0191.39103](#) MR [264470](#)
- [65] L. Nirenberg and F. Trèves, On local solvability of linear partial differential equations. II. Sufficient conditions. *Comm. Pure Appl. Math.* **23** (1970), 459–509 Zbl [0208.35902](#)
MR [264471](#)
- [66] É. Picard, *Traité d'analyse. Tome II. Les Grands Classiques Gauthier-Villars, Jacques Gabay, Sceaux*, 1991 MR [1188643](#)
- [67] F. Riesz, Über die Randwerte einer analytischen Funktion. *Math. Z.* **18** (1923), no. 1, 87–95
Zbl [49.0225.01](#) MR [1544621](#)
- [68] T. Rivière, Exploring the unknown: the work of Louis Nirenberg on partial differential equations. *Notices Amer. Math. Soc.* **63** (2016), no. 2, 120–125 Zbl [1338.35006](#) MR [3409063](#)
- [69] J. Serrin, A symmetry problem in potential theory. *Arch. Rational Mech. Anal.* **43** (1971), 304–318 Zbl [0222.31007](#) MR [333220](#)
- [70] M. Struwe, On the evolution of harmonic mappings of Riemannian surfaces. *Comment. Math. Helv.* **60** (1985), no. 4, 558–581 Zbl [0595.58013](#) MR [826871](#)
- [71] T. Tao and G. Tian, A singularity removal theorem for Yang–Mills fields in higher dimensions. *J. Amer. Math. Soc.* **17** (2004), no. 3, 557–593 Zbl [1086.53043](#) MR [2053951](#)

- [72] N. S. Trudinger, Elliptic equations in nondivergence form. In *Miniconference on partial differential equations (Canberra, 1981)*, pp. 1–16, Proc. Centre Math. Anal. Austral. Nat. Univ. 1, Austral. Nat. Univ., Canberra, 1982 Zbl [0555.35040](#) MR [758449](#)
- [73] N. S. Trudinger, Fully non-linear, uniformly elliptic equations under natural structure conditions. *Trans. Amer. Math. Soc.* **278** (1983), no. 2, 751–769 Zbl [0518.35036](#) MR [701522](#)

Received 17 April 2020.

Tristan Rivière

Department of Mathematics, ETH Zürich, 8093 Zürich, Switzerland; riviere@math.ethz.ch