

The Artin monoid Cayley graph

Rachael Boyd, Ruth Charney, Rose Morris-Wright, and Sarah Rees

Abstract. In this paper, we investigate properties of the Artin monoid Cayley graph. This is the Cayley graph of an Artin group A_Γ with respect to the (infinite) generating set given by the associated Artin monoid A_Γ^+ . In a previous paper, the first three authors introduced a monoid Deligne complex and showed that this complex is contractible for all Artin groups. In this paper, we show that the Artin monoid Cayley graph is quasi-isometric to a modification of the Deligne complex for A_Γ obtained by coning off translates of the monoid Deligne complex. We then address the question of when the monoid Cayley graph has infinite diameter. We conjecture that this holds for all Artin groups of infinite type. We give a set of criteria that imply infinite diameter, and using existing solutions to the word problem for large type Artin groups and 3-free Artin groups, we prove that the conjecture holds for any Artin group containing a 3-generator subgroup of one of these two types.

1. Introduction

Artin groups (also known as Artin–Tits groups) form a large class of groups closely associated to Coxeter groups. They arise naturally in algebraic geometry, topology, and representation theory. The classical examples of such groups are braid groups, whose associated Coxeter groups are the symmetric groups. Braid groups have been extensively studied using both combinatorial and geometric methods. While many of these techniques can be generalised to all spherical type Artin groups (that is, those whose associated Coxeter group is finite), Artin groups associated to infinite Coxeter groups present a much bigger challenge and remain largely mysterious. Recently, there has been a major push to find new geometric approaches to study these groups. In this paper, we investigate some new combinatorial approaches.

To specify an Artin group, we begin with a labelled graph Γ with a finite vertex set

$$X = \{x_1, \dots, x_n\}$$

and edge set E such that each edge $\{x_i, x_j\} \in E$ is labelled by an integer

$$m_{i,j} = m_{j,i} \geq 2.$$

2020 Mathematics Subject Classification. Primary 20F36; Secondary 20F55, 20M32, 20F65.

Keywords. Artin monoids, Artin groups.

We then define the *Artin group*, A_Γ , to be the group with presentation

$$A_\Gamma = \left\langle X \mid \underbrace{x_i x_j x_i \cdots}_{\text{length } m_{i,j}} = \underbrace{x_j x_i x_j \cdots}_{\text{length } m_{i,j}} \forall \{x_i, x_j\} \in E \right\rangle.$$

If there is no edge in Γ between x_i and x_j in X , we set $m_{i,j} = \infty$ and there is no relation between x_i and x_j in the presentation. The associated Coxeter group W_Γ is the group whose presentation is the same as A_Γ with the added relations $x_i^2 = e$ for all i , where e is the identity element.

If we allow only the positive powers of the generators, then the presentation above gives rise to a monoid, A_Γ^+ . As it turns out, questions that seem intractable for Artin groups are often easier to solve in the monoid case. For example, because all of the relations in the presentation preserve the length of a word, there is an easily computable length function for any given monoid element. Thus, the word problem in the monoid is easily solved. Moreover, there are nice normal forms for the elements of the monoid [3, 12]. However, solving the word problem in Artin groups is much more difficult, and for many Artin groups, it is not even known whether such solutions exist.

By a theorem of Paris [14], the natural map from A_Γ^+ to the submonoid of A_Γ consisting of positive words in X is an isomorphism. So, it is natural to ask if and how one might use information about the monoid to understand the Artin group. In the case of a spherical type Artin group, a connection between the group and the monoid is obtained via the existence of a Garside element. Namely, there is an element Δ in the Artin monoid (corresponding to the longest element in the Coxeter group) such that for any $g \in A_\Gamma$, $g\Delta^n$ is represented by a positive word in A_Γ^+ for sufficiently large n . In addition, Δ^2 lies in the center of A_Γ . It follows that most questions about the behaviour of the Artin group can be translated into questions about the Artin monoid. The existence of this Garside element is key to our understanding of spherical type Artin groups. (We refer the reader to [6, 11] for more details about Garside groups and Garside elements.)

For infinite type Artin groups, there is no Garside element and no clear way to translate information from the Artin monoid to the Artin group. In this paper, we begin to address this problem by investigating the Cayley graph $\text{Cay}(A_\Gamma, M)$ of the Artin group A_Γ with respect to the (infinite) generating set $M := A_\Gamma^+$. We call this the *monoid Cayley graph* for A_Γ . Geodesics in this Cayley graph correspond to minimal length multi-fractions, $g = a_1 a_2^{-1} \cdots a_k^{\pm 1}$, $a_i \in A_\Gamma^+$, representing an element $g \in A_\Gamma$. Understanding these geodesics is a first step towards defining normal forms for elements of A_Γ and offers inductive approaches to answer some questions about these groups. In the case of a spherical type Artin group, $\text{Cay}(A_\Gamma, M)$ is neither interesting nor useful as the Garside structure implies that this graph has diameter two; we can

reach any element by travelling along an edge labelled by a single monoid element followed by an edge labelled (forward or backward) by a power of Δ . We conjecture that in all other cases, that is for all infinite type Artin groups, $\text{Cay}(A_\Gamma, M)$ has infinite diameter. It is this question that we address in this paper. Surprisingly, it is not at all obvious.

The first step is to observe that if A_T is a special subgroup of A_Γ (that is, the subgroup generated by a subset $T \subset X$), then the inclusion of the monoid Cayley graph of A_T into the monoid Cayley graph of A_Γ is an isometric embedding (see Proposition 2.2). Hence, to show that $\text{Cay}(A_\Gamma, M)$ has infinite diameter it suffices to find a special subgroup for which this holds. Thus, while we focus primarily on 3-generator Artin groups, our results extend to any Artin group containing one of these groups as a special subgroup. Our main result is the following theorem.

Theorem 6.1. *Let $\text{Cay}(A_\Gamma, M)$ denote the Cayley graph of A_Γ with respect to the generating set $M := A_\Gamma^+$. Then, $\text{Cay}(A_\Gamma, M)$ has infinite diameter providing one of the following holds.*

- (1) Γ contains a pair of vertices not connected by an edge (or equivalently, there exists i, j with $m_{i,j} = \infty$).
- (2) Γ contains a triangle with all labels ≥ 3 .
- (3) Γ contains a triangle with no label equal to 3 and at most one label is equal to 2.

Note that the only 3-generator Artin groups of infinite type not covered by this theorem are those with edge labels $\{2, 3, n\}$ with $n \geq 6$. After posting this paper, we were informed by Arye Juhasz that he has a proof of the $\{2, 3, n\}$ case. Together with our results, this leaves open only the case of Artin groups all of whose 3-generator special subgroups are spherical type. The proof of the theorem involves constructing a sequence of geodesics of increasing length in $\text{Cay}(A_\Gamma, M)$. To do this, we first establish some explicit criteria on words in the generators X , which allow us to construct such geodesics. Using solutions to the word problem established by Holt and the fourth author in the case of large type Artin groups [9] and by Blasco–García, Cumplido, and the third author in the case of 3-free Artin groups [1], we then show that these criteria hold for any 3-generator Artin group satisfying one of the conditions above.

While the proof of the main theorem involves combinatorial arguments, the original motivation came from geometric questions. A geometric construction that has played a key role in the study of Artin groups is the Deligne complex \mathcal{D}_Γ , which we describe in Section 3. A central conjecture about Artin groups, known as the $K(\pi, 1)$ -conjecture, can be reduced to proving that the Deligne complex is contractible. In 1972, Deligne [7] proved that this conjecture holds for spherical type Artin groups

using the combinatorial information given by the Garside structure. The conjecture has subsequently been proved using more geometric arguments for some special classes of infinite type Artin groups (see the survey paper by Paris [15]) but the general conjecture has remained unapproachable. In a paper by the first three authors, we introduce an analogue of the Deligne complex for the monoid, called the monoid Deligne complex \mathcal{D}_Γ^+ , and prove that this complex is contractible for *all* Artin monoids. The monoid Deligne complex embeds in the full Deligne complex and its translates cover all of \mathcal{D}_Γ . Our attempt to better understand this covering, and how these translates intersect, led us to the study of the monoid Cayley graph.

There is a direct connection between the geometry of the monoid Cayley graph and that of the Deligne complex. Since \mathcal{D}_Γ^+ is contractible, attaching a cone to each translate of \mathcal{D}_Γ^+ in \mathcal{D}_Γ does not change its topology. In Section 3, we define the “coned-off Deligne complex”, $C\mathcal{D}_\Gamma$, and prove the following theorem.

Theorem 3.2. *For any infinite type Artin group A_Γ , $C\mathcal{D}_\Gamma$ is quasi-isometric to $\text{Cay}(A_\Gamma, M)$.*

The paper is organised as follows. In Section 2, we introduce the monoid Cayley graph $\text{Cay}(A_\Gamma, M)$ and prove that the monoid Cayley graph of a special subgroup isometrically embeds into the monoid Cayley graph of A_Γ . In Section 3, we investigate the coned-off Deligne complex. Sections 4 and 5 contain the proof of the main theorem.

2. The monoid Cayley graph

In this section, we introduce some basic notation and discuss the relationship between the monoid Cayley graph of an Artin group and the monoid Cayley graphs of special subgroups.

A standard tool in geometric group theory for studying a finitely generated group G is the assignment of a metric to G . To do this, one chooses a finite generating set S and considers the Cayley graph of G with respect to S , denoted by $\text{Cay}(G, S)$. This is the graph whose vertices are the elements of G and edges join two vertices g and h when they differ by a single element of S , that is, $h = gs$ for some $s \in S$. One can then define a metric on G by taking the distance from g_1 to g_2 to be the length of a shortest path from g_1 to g_2 ; in this Cayley graph; each such path is labelled by a word over S (string over $S \cup S^{-1}$). Different choices of (finite) generating sets give rise to quasi-isometric metrics, so this metric is “coarsely” well-defined. For an Artin group A_Γ with its standard generating set X , we denote this Cayley graph by $\text{Cay}(A_\Gamma, X)$.

In this paper, we will consider a different type of Cayley graph, namely, the Cayley graph with respect to an infinite generating set M . As defined above, M consists of all

elements in the Artin monoid A_Γ^\pm . The vertices of this Cayley graph, $\text{Cay}(A_\Gamma, M)$, are again the elements of A_Γ , but two vertices g, h are now joined by an edge whenever $h = gm$ for some $m \in M$. Note that these edges come with an orientation from g to h . However, paths in this Cayley graph may follow edges in either direction. Thus, a path from the identity vertex e to a vertex g corresponds to a word over M (string over $M \cup M^{-1}$) representing g . Given a word w over the standard generating set X , we can subdivide w into maximal subwords which are all positive or all negative. This determines a word over M representing the same element of A_Γ . We will refer to the length of this word, viewed as a path in $\text{Cay}(A_\Gamma, M)$, as the *monoidal length* of w and denote it by $|w|_M$.

We are interested in understanding the geometry of $\text{Cay}(A_\Gamma, M)$. Here are a few easy observations. First, if A_Γ is a spherical type Artin group with Garside element Δ , then any $g \in A_\Gamma$ can be translated into the monoid by multiplying by a sufficiently high power of Δ , that is, there exists n such that

$$g\Delta^n = m \in M.$$

Thus, there is a path from the identity vertex e to g consisting of two edges, the edge from e to m , followed by the (reverse of) the edge from g to m . It follows that in this case, the diameter of $\text{Cay}(A_\Gamma, M)$ is two. Thus, the monoid Cayley graph is of little interest in the spherical type case. We will restrict our attention to infinite type Artin groups.

Conjecture 2.1. *For any infinite type Artin group, $\text{Cay}(A_\Gamma, M)$ has infinite diameter.*

This conjecture will be the main focus of this paper. Studying the diameter involves understanding geodesics in this Cayley graph. These are of interest in their own right. Note that any two vertices connected by a path all of whose edges have the same orientation will also be connected by a single edge, since the product of monoid elements is again a monoid element. Thus, any geodesic from e to g in $\text{Cay}(A_\Gamma, M)$ must be an alternating path, or, in other words, an expression of the form $g = a_1 a_2^{-1} a_3 \cdots a_k^{\pm 1}$ (or $g = a_1^{-1} a_2 a_3^{-1} \cdots a_k^{\pm 1}$) with $a_i \in M$.

In this paper, we will focus on the case of 3-generator Artin groups. However, as we will now see, this has immediate implications for higher rank Artin groups. Let A_Γ be an Artin group with standard generating set X . A *special subgroup* of A_Γ is a subgroup A_T generated by a subset $T \subseteq X$. By a theorem of van der Lek [16], this group is itself an Artin group with defining graph the full subgraph of Γ spanned by T .

Proposition 2.2. *Let A_Γ be an Artin group with generating set X and let $T \subset X$ be any non-empty subset. Let M_T be the positive monoid in A_T . There is a simplicial retraction of $\text{Cay}(A_\Gamma, M)$ onto $\text{Cay}(A_T, M_T)$. Hence, the natural inclusion of*

$\text{Cay}(A_T, M_T)$ into $\text{Cay}(A_\Gamma, M)$ is an isometric embedding, that is, it maps geodesics to geodesics.

Proof. This follows from work of Godelle and Paris [8] in which they construct a retraction of the Salvetti complex of A_Γ to the Salvetti complex of A_T . Subsequently, Charney and Paris [5] studied the universal cover of this retraction in more detail. The 1-skeleton of the universal cover of the Salvetti complex of A_Γ is precisely the Cayley graph of A_Γ with respect to the standard generating set X . Edges come with a preferred orientation, namely, the orientation corresponding to the positive power of a generator in X . In the proof of [5, Theorem 1.2], they give a precise description of this retraction: the retraction either collapses an edge to a single point or maps it to an edge with the same orientation. It follows that it takes any strictly positive path, corresponding to a monoid element in A_Γ , to a strictly positive path corresponding to a monoid element in A_T . Thus, it induces a retraction of $\text{Cay}(A_\Gamma, M)$ onto $\text{Cay}(A_T, M_T)$ taking each edge, labelled by an element of M , to either a single vertex or a single edge labelled by an element of M_T . In particular, if γ is an edge path in $\text{Cay}(A_\Gamma, M)$ connecting two vertices of $\text{Cay}(A_T, M_T)$, then its image under the retraction is an edge path of the same or shorter length in $\text{Cay}(A_T, M_T)$ between these vertices. It follows that geodesic paths in $\text{Cay}(A_T, M_T)$ must also be geodesic in $\text{Cay}(A_\Gamma, M)$. ■

Corollary 2.3. *If A_Γ contains a special subgroup A_T whose monoid Cayley graph has infinite diameter, then the same holds for the monoid Cayley graph of A_Γ .*

As an example, let $T = \{s, t\}$ be two generators not connected by an edge, so A_T is the free group on $\{s, t\}$. It is easy to see that the word $st^{-1}st^{-1} \dots st^{-1}$ of length $2m$ is geodesic in $\text{Cay}(A_T, M_T)$. It then follows from the corollary that for any Γ whose underlying graph is not a clique $\text{Cay}(A_\Gamma, M)$ has infinite diameter.

3. Relationship to the monoid Deligne complex

In this section, we introduce the coned-off Deligne complex (which we note here is different from the coned-off Deligne complex defined in the paper of Martin and Przytycki [10]) and show that the resulting complex is quasi-isometric to $\text{Cay}(A_\Gamma, M)$.

We first give a bit of history. Associated to any Coxeter group W_Γ is a complex hyperplane complement on which W_Γ acts freely. It is known that the quotient of this space by W_Γ has fundamental group A_Γ . The $K(\pi, 1)$ -conjecture states that, in fact, this quotient space is a $K(A_\Gamma, 1)$ -space. In 1972, Deligne [7] proved that this holds for spherical type Artin groups by constructing a simplicial complex homotopy equivalent to the universal cover of this hyperplane complement and proving that this simplicial complex is contractible. In [4], the second author and M. Davis constructed a slight

variation on this complex for infinite-type Artin groups, now known as the Deligne complex, which plays an analogous role for infinite-type Artin groups. The $K(\pi, 1)$ -conjecture is equivalent to the conjecture that the Deligne complex is contractible for all A_Γ . In [4], the conjecture was proved in some special cases (such as for FC type) and more recently, it was proved for affine type Artin groups by Paolini and Salvetti [13], but it remains open in general.

To define the Deligne complex, first note that a special subgroup A_T of an infinite type Artin group A_Γ may have either spherical type or infinite type. The Deligne complex \mathcal{D}_Γ is the geometric realisation of the partially ordered set \mathcal{P} of cosets gA_T , where A_T is a spherical type special subgroup and the ordering is given by inclusion. (Note that we allow $T = \emptyset$ and define $gA_\emptyset = \{g\}$.) There are two natural piecewise Euclidean metrics that we can put on \mathcal{D}_Γ . The first is obtained by using the simplicial structure and declaring each simplex to be a standard Euclidean simplex with all edges of length 1. We denote the Deligne complex with this metric by $\mathcal{D}_\Gamma^\Delta$. An alternate metric is obtained by viewing \mathcal{D}_Γ as a cube complex. Here, the cubes correspond to intervals $[gA_T, gA_R]$ in the poset \mathcal{P} and the metric assigns each k -cube to be isometric to the unit cube I^k in \mathbb{R}^k . These two metrics on \mathcal{D}_Γ are quasi-isometric since the maximal dimension of cubes in the Deligne complex is finite, so viewing the cube as a union of simplices distorts the metric by a bounded amount. We will use the simplicial structure and metric defined on $\mathcal{D}_\Gamma^\Delta$ as it is more convenient for our arguments.

We can also define an analogue of this space for the Artin monoid. Namely, we define $\mathcal{D}_\Gamma^{\Delta+}$ to be the geometric realization of the subposet of \mathcal{P} consisting of cosets mA_T , where $m \in A_\Gamma^+ = M$. In [2], the first three authors prove that $\mathcal{D}_\Gamma^{\Delta+}$ is contractible for all infinite type A_Γ . The complex $\mathcal{D}_\Gamma^{\Delta+}$ naturally embeds into $\mathcal{D}_\Gamma^\Delta$. There is an action of A_Γ on $\mathcal{D}_\Gamma^\Delta$, via left multiplication on the cosets $g \cdot mA_T = gmA_T$. We call the image of $\mathcal{D}_\Gamma^{\Delta+}$ under the action of $g \in A_\Gamma$ a *translate* of $\mathcal{D}_\Gamma^{\Delta+}$ and denote it by $g\mathcal{D}_\Gamma^{\Delta+}$.

Definition 3.1. Define the *coned-off Deligne complex* $C\mathcal{D}_\Gamma$ to be the simplicial Deligne complex $\mathcal{D}_\Gamma^\Delta$ with a ‘‘cone point’’ added for each translate of $\mathcal{D}_\Gamma^{\Delta+}$ in the following sense: for all $g \in A_\Gamma$, add

- (1) a vertex v_g ,
- (2) an edge between every vertex in $g\mathcal{D}_\Gamma^{\Delta+}$ and the vertex v_g ,
- (3) a $p + 1$ simplex $v_g * \sigma_p$ for every p -simplex $\sigma_p \in g\mathcal{D}_\Gamma^{\Delta+}$.

This is again a simplicial complex, and we put the standard metric on each simplex such that all edges have length 1.

One motivation for defining this coned-off complex is that we know that the monoid Deligne complex is contractible, as are its translates. Therefore, coning them off

does not change the homotopy type, i.e.,

$$C \mathcal{D}_\Gamma \simeq \mathcal{D}_\Gamma^\Delta.$$

We now define a map from vertices in $\text{Cay}(A_\Gamma, M)$ to vertices in $C \mathcal{D}_\Gamma$ as follows:

$$f : \text{Cay}(A_\Gamma, M) \rightarrow C \mathcal{D}_\Gamma; \quad g \mapsto v_g.$$

Theorem 3.2. *The above map f is a quasi-isometry, i.e., $\text{Cay}(A_\Gamma, M) \simeq_{q.i.} C \mathcal{D}_\Gamma$.*

Proof. First, note that because the translates of $\mathcal{D}_\Gamma^{\Delta+}$ by A_Γ cover $\mathcal{D}_\Gamma^\Delta$, any point in $\mathcal{D}_\Gamma^\Delta$ is in some translate $g\mathcal{D}_\Gamma^{\Delta+}$, and therefore at most distance one from the cone point for that translate, v_g , in $C \mathcal{D}_\Gamma$. Thus, every point in $C \mathcal{D}_\Gamma$ is at most distance one from a point in $f(\text{Cay}(A_\Gamma, M))$.

First, we claim that if g and h are joined by an edge in $\text{Cay}(A_\Gamma, M)$ then $f(g) = v_g$ and $f(h) = v_h$ are joined by an edge path of length 2 in $C \mathcal{D}_\Gamma$. Without loss of generality, we can assume that the edge is directed from h to g so that $g = h \cdot m$ for some $m \in M$. Now, $f(g) = v_g$ is joined by an edge to gA_\emptyset , and $gA_\emptyset = h \cdot mA_\emptyset$ is in $h\mathcal{D}_\Gamma^{\Delta+}$, so it is joined by an edge to $v_h = f(h)$. We have therefore constructed a path of length 2 from $f(g)$ to $f(h)$ as claimed.

Since any geodesic in $\text{Cay}(A_\Gamma, M)$ is a path of such edges, for x and y in $\text{Cay}(A_\Gamma, M)$ we deduce that

$$d_{C \mathcal{D}_\Gamma}(f(x), f(y)) \leq 2 \cdot d_{\text{Cay}}(x, y).$$

In particular, this means that, for any $A \in \mathbb{R}$ such that $A \geq 2$, we have

$$d_{C \mathcal{D}_\Gamma}(f(x), f(y)) \leq A \cdot d_{\text{Cay}}(x, y).$$

It remains to show that

$$d_{\text{Cay}}(x, y) \leq A \cdot d_{C \mathcal{D}_\Gamma}(f(x), f(y))$$

for some A .

For the following argument, refer to Figure 1. Start with a geodesic γ between $f(x)$ and $f(y)$ in $C \mathcal{D}_\Gamma$. We can deform γ into an edge path γ' from x to y contained in the same simplices as γ . Since the maximal dimension of the simplices in $\mathcal{D}_\Gamma^\Delta$ is finite, there exists some constant $B \geq 1$, depending only on this dimension such that $B \text{lg}(\gamma) \geq \text{lg}(\gamma')$, where $\text{lg}(\cdot)$ denotes the length of the path in $C \mathcal{D}_\Gamma$. Label the edges of γ' by e_1, \dots, e_k . Each edge e_i of γ' is either an edge to or from a cone point or an edge in $\mathcal{D}_\Gamma^\Delta$. Since the translates of $\mathcal{D}_\Gamma^{\Delta+}$ cover $\mathcal{D}_\Gamma^\Delta$, any edge in $\mathcal{D}_\Gamma^\Delta$ lies in some translate of $\mathcal{D}_\Gamma^{\Delta+}$. Without loss of generality, we may assume that the only edges with cone point vertices are the first and last (with vertices v_x and v_y), as otherwise we can




Figure 1. The edge path γ' in $C\mathcal{D}_\Gamma$ joining v_x to v_y , the corresponding modified edge path in $C\mathcal{D}_\Gamma$ (with γ' shown in grey) and the path in $\text{Cay}(A_\Gamma, M)$ from x to y .

apply this proof to every section of the edge path between cone point vertices. Then, except for e_1 and e_k , every edge lies in a translate of $\mathcal{D}_\Gamma^{\Delta+}$ by an element $g_i \in A_\Gamma$ for $2 \leq i \leq k-1$. We replace the edge e_i with the two edges d_i^1, d_i^2 , with d_i^1 travelling from the initial vertex of e_i to the cone point v_{g_i} , and d_i^2 from the cone point to the terminal vertex of e_i . Edges e_i and e_{i+1} meet at a vertex in the overlap of the translates $g_i\mathcal{D}_\Gamma^{\Delta+}$ and $g_{i+1}\mathcal{D}_\Gamma^{\Delta+}$; i.e., the vertex satisfies $g_im_ia_i = g_{i+1}m_{i+1}a_i$ for m_i and m_{i+1} in M . Since A_T is of spherical type, this means in particular that there exist $a_i, b_i \in A_T^+$ such that

$$g_im_ia_ib_i^{-1} = g_{i+1}m_{i+1} \implies g_im_ia_i = g_{i+1}m_{i+1}b_i.$$

In $\text{Cay}(A_\Gamma, M)$, the vertices g_i and g_{i+1} are therefore joined by an edge path of length at most 2. Similarly, the vertices x and g_2 are joined by an edge path of length 2, as are the vertices g_{k-1} and y . This is because, for example, v_x is joined by the edge e_1 to a vertex which also lies in the edge e_2 , and thus in $g_2\mathcal{D}_\Gamma^{\Delta+}$. So, this vertex satisfies $xm_1A_T = g_2m_2A_T$ for m_1 and m_2 in M , and as before, we can argue that this corresponds to an edge path of length 2. From the modified path in $C\mathcal{D}_\Gamma$, we

are therefore able to construct a path between x and y in $\text{Cay}(A_\Gamma, M)$, as shown in Figure 1. It follows that

$$d_{\text{Cay}(A_\Gamma, M)}(x, y) \leq 2k - 2 \leq 2k = 2 \lg(\gamma'),$$

and hence,

$$d_{\text{Cay}(A_\Gamma, M)}(x, y) \leq 2B \lg(\gamma) = 2B \cdot d_{\mathcal{C}_{\mathcal{D}_\Gamma}}(f(x), f(y)).$$

Taking

$$A = 2B \geq 2,$$

this completes the proof. ■

4. Criteria for infinite diameter

In this section, we will introduce two criteria which an Artin group may or may not satisfy and prove that if an Artin group A_Γ satisfies both of these criteria, then, as a consequence, $\text{Cay}(A_\Gamma, M)$ has infinite diameter. To illustrate the criteria, we will include some simple examples of Artin groups which satisfy them and Artin groups which do not. Further examples appear in Section 5, where we prove that large type and 3-free Artin groups of infinite type satisfy the two criteria.

Assuming an Artin group A_Γ satisfies the two criteria, our strategy to prove that $\text{Cay}(A_\Gamma, M)$ has infinite diameter is as follows. We start with a particular sequence of words $w_1, w_2, \dots, w_n, \dots$ over the standard generating set X , for which $|w_n|_M$ is equal to n . This sequence exists due to Criterion 2. Using Criterion 1, we now show that when we premultiply w_n by a word of monoidal length less than n the word we obtain cannot reduce to the empty word and hence cannot represent the identity. It follows that w_n is geodesic in $\text{Cay}(A_\Gamma, M)$, and so the sequence of words $\{w_n\}$ represents a sequence of geodesics of increasing length in $\text{Cay}(A_\Gamma, M)$. This shows that $\text{Cay}(A_\Gamma, M)$ has infinite diameter.

4.1. Criterion 1. A_Γ has preserved signed suffixes

Definition 4.1. Suppose that A_Γ is an Artin group with generating set X . Let w be a geodesic word over X that ends in a positive (resp., negative) letter, and let u denote the longest positive (resp., negative) proper suffix of w . Let a be a single positive (resp., negative) generator. We say that A_Γ has *preserved positive (resp., negative) suffixes* if we can guarantee the existence of a word for the element of the group given by aw that is geodesic in X and also has u as a suffix. If A_Γ has both preserved positive and negative suffixes, then we say that the group has *preserved signed suffixes*.

For example, free groups have preserved signed suffixes. In a free group, if w is a geodesic with respect to the standard generators (i.e., a freely reduced word) and aw is not, this means that w must begin with the letter a^{-1} . A geodesic representative for the group element given by aw can thus be obtained via free cancellation of aa^{-1} , and this leaves the remainder of w (in particular, the longest positive suffix u) unchanged.

Similarly, the free abelian group \mathbb{Z}^n has preserved signed suffixes. If w is a geodesic word with respect to the standard generators and aw is not, this means that w can be factored as

$$w = w_1 a^{-1} w_2,$$

and a geodesic word for the group element given by aw can be obtained by allowing a to commute with each individual letter of w_1 to obtain the word $w_1 a a^{-1} w_2$. After the final cancellation of aa^{-1} , a geodesic is obtained and the suffix w_2 is unchanged. The longest positive suffix of w must also be a suffix of w_2 , so this positive suffix is preserved. A symmetric argument shows that \mathbb{Z}^n also has preserved negative suffixes and so has preserved signed suffixes.

In our results, we use the solution to the word problem for large type Artin groups given in [9] and for 3-free Artin groups given in [1]. In Section 5, we explain in more detail how these solutions to the word problem guarantee that large type and 3-free Artin groups have preserved signed suffixes.

4.2. Criterion 2. A_Γ contains an alternating blocking sequence $\{\alpha_n\}$

In this section, we will describe the criterion required to construct the desired sequence of words $\{w_n\}$.

Definition 4.2. Let A_Γ be an Artin group with generating set X . Let u be a word and $x \in X \cup X^{-1}$. We say that (u, x) is a *blocking pair* if, given any geodesic word over X of the form wu , it follows that the word wux is also a geodesic word over X .

Intuitively, the subword u is blocking any letters from w from cancelling out the letter x .

For example, in a free group, if $a, x \in X$ (both positive) then (a, x) is a blocking pair. In a free abelian group, if $a, x \in X$ and $a \neq x$, then (a, x) is not a blocking pair because the word $x^{-1}a$ is geodesic, but the word $x^{-1}ax$ is not.

We now use blocking pairs to build a sequence of words over X that label arbitrarily long paths in $\text{Cay}(A_\Gamma, M)$.

Definition 4.3. Suppose that $\{\alpha_n\}$ is an infinite sequence of words over X , where α_i is a nonempty positive word over X if i is odd and α_i is a nonempty negative word if i is even. We say that the sequence $\{\alpha_n\}$ is an *alternating blocking sequence* if for

any letter x in α_n there exists some subword u of $\alpha_{n-1}\alpha_n$ immediately preceding x such that (u, x) is a blocking pair.

Our second criterion states that A_Γ must contain such an alternating blocking sequence.

Remark 4.4. If A_Γ is of spherical type, then A_Γ has no alternating blocking sequence. For suppose that u is any positive word and that $x^{-1} \in X^{-1}$. Then, (u, x^{-1}) cannot be a blocking pair. To prove this, we first recall the following facts about spherical type Artin groups from Garside theory: the square of the Garside element Δ^2 is in the center of A_Γ and satisfies that, for all $x \in X$, x is a right divisor of Δ . Now, consider the word $\Delta^2 u$, which represents an element of the monoid and so is a geodesic word over X . On the other hand, $\Delta^2 u x^{-1}$ represents the same group element as $u \Delta^2 x^{-1}$ which is not freely reduced (since x is a right divisor of Δ^2). Therefore, (u, x) is not a blocking pair. It follows that no alternating sequence of words $\{\alpha_n\}$ can satisfy the conditions for an alternating blocking sequence.

4.3. Sufficiency of these criteria

In this section, we prove that if an Artin group A_Γ satisfies Criterion 1 and Criterion 2, then the monoid Cayley graph $\text{Cay}(A_\Gamma, M)$ has infinite diameter.

Theorem 4.5. *Let A_Γ be an Artin group with standard generating set X and positive monoid M . Suppose that A_Γ has preserved signed suffixes, and that A_Γ contains an alternating blocking sequence of words $\{\alpha_n\}$. Then, the monoid Cayley graph $\text{Cay}(A_\Gamma, M)$ has infinite diameter.*

Proof. Let $w_n = \alpha_1 \cdots \alpha_n$. We will show that the words $\{w_n\}$ label geodesic paths of length n in $\text{Cay}(A_\Gamma, M)$. The claim that $\text{Cay}(A_\Gamma, M)$ is infinite diameter follows.

Consider some arbitrary word $v_m = \beta_m \cdots \beta_2 \beta_1$ where β_i is a positive word over X if i is odd and β_i is a negative word over X if i is even. We allow the possibility that β_1 is the empty word, but for $i > 1$, we assume that β_i is non-empty; hence, v_m labels a path of length m or $m - 1$ in $\text{Cay}(A_\Gamma, M)$.

To show that w_n is a geodesic word over M , it is sufficient to show that for all v_m such that $m \leq n$, the word $v_m w_n$ does not represent the identity element in A_Γ . Because the identity is represented by the empty word regardless of which generating set we use, we do this by showing that $v_m w_n$ regarded as a word over X cannot be reduced to the empty word. This follows from the below claim.

Claim. If $m \leq n$, then the group element represented by the word $v_m w_n$ has a geodesic representative over X ending in $\alpha_m \cdots \alpha_{n-1} \alpha_n$.

We prove this claim via induction on m .

Base case. Assume that $m = 1$. We prove that for all n , $v_1 w_n$ is a geodesic word over X , and is therefore the representative we require. If $n = 1$, then $v_1 w_1 = \beta_1 \alpha_1$ is a positive word over X and so must be geodesic over X ; it is therefore a geodesic with suffix α_1 as required.

Now, we build the word $v_1 w_n$ and see that it is geodesic. We suppose that $j > 1$ and that $v_1 w_{j-1}$ is a geodesic word and build the word $v_1 w_j$ from $v_1 w_{j-1}$ by appending one letter of $\alpha_j = x_{j,1} x_{j,2} \cdots x_{j,k_j}$ at a time and using induction on the number of letters we have appended. As the base case, we append the first letter $x_{j,1}$. Then, the fact that $\{\alpha_n\}$ is an alternating blocking sequence guarantees the existence of some suffix $u_{j,1}$ of the word $v_1 w_{j-1}$ such that $(u_{j,1}, x_{j,1})$ is a blocking pair. Since we assumed that $v_1 w_{j-1}$ is a geodesic word over X , the blocking pair condition implies that $v_1 w_{j-1} x_{j,1}$ must also be a geodesic word over X . Now, assume the inductive hypothesis that $v_1 w_{j-1} x_{j,1} \cdots x_{j,i-1}$ is a geodesic word over X and append the letter $x_{j,i}$. Again, since $\{\alpha_n\}$ is an alternating blocking sequence, there exists some suffix $u_{j,i}$ of the word $v_1 w_{j-1} x_{j,1} \cdots x_{j,i-1}$ such that $(u_{j,i}, x_{j,i})$ is a blocking pair. Since $v_1 w_{j-1} x_{j,1} \cdots x_{j,i-1}$ is a geodesic word over X , the blocking pair condition implies that $v_1 w_{j-1} x_{j,1} \cdots x_{j,i}$ must also be a geodesic word over X . Iterating this method to append $\alpha_2 \cdots \alpha_n$ to $v_1 w_1 = v_1 \alpha_1$, we see at each step a letter $x_{j,i}$ is appended and the word remains geodesic over X . Therefore, $v_1 w_n$ is a geodesic word over X with suffix $\alpha_1 \cdots \alpha_n$.

Inductive hypothesis. Suppose that $m > 1$. We assume that for all $m \leq n$, the element of A_Γ represented by $v_{m-1} w_n$ can be represented by a geodesic word ξ over X that ends with $\alpha_{m-1} \cdots \alpha_n$.

Inductive step. Consider the case $n = m$. By the inductive hypothesis the group element represented by $v_{m-1} w_m$ is represented by a geodesic word ξ over X that ends with $\alpha_{m-1} \alpha_m$. We assume that m is odd, and note that a symmetric argument will work for the m even case. Recall the word v_m satisfies $v_m = \beta_m v_{m-1}$, and hence, $v_m w_n = \beta_m v_{m-1} w_n = \beta_m \xi$. Suppose that $\beta_m = b_{l_m} \cdots b_2 b_1$; i.e., β_m is of length l_m over X . Since m is odd, α_m and β_m are both positive words over X , so each $b_i \in X$. Starting with b_1 , we append these letters to the left of ξ one at a time. The group A_Γ has preserved signed suffixes, so when we append the positive letter b_1 to the left of the geodesic word ξ , we can obtain a new geodesic that ends in the same positive suffix as ξ , namely, the suffix α_m . Repeating this process for each b_i , we obtain a geodesic word over X which represents the element $v_m w_m$ and has suffix α_m .

For $n > m$, we repeat the argument from the base case, in which we started with the geodesic representative of the group element given by $v_1 w_{j-1}$ and built one for $v_1 w_j$. In this case, we start with the geodesic word we just constructed, which represents the element given by $v_m w_m$ and ends with α_m . As in the base case we append the letters $x_{j,i}$ one at a time on the right, for $m < j \leq n$, and due to the fact that $\{\alpha_n\}$ is an

alternating blocking sequence, the word remains a geodesic over X at each step. After we have appended all the $x_{j,i}$ we obtain a geodesic representative of $v_m w_n$ ending in $\alpha_m \cdots \alpha_n$ as desired. ■

Remark 4.6. In the above proof, we allowed the possibility that $|v_n|_M = |w_n|_M$ providing the last word β_1 of v_n has the same sign as the first word α_1 of w_n . Thus, we have also shown that the element of A_Γ represented by w_n cannot be represented by a geodesic v_n^{-1} in $\text{Cay}(A_\Gamma, M)$ beginning with a word of the opposite sign.

5. The large type and 3-free case

In this section, we show that large type Artin groups and 3-free Artin groups that are not of spherical type satisfy the criteria in Section 4 that we need in order to apply Theorem 4.5.

An Artin group is said to have large type if all the associated parameters m_{ij} are at least 3 (possibly infinite). A 3-free Artin group is an Artin group for which none of the parameters m_{ij} are equal to 3.

Lemma 5.1. *Let A_Γ be a 3-free Artin group of infinite type with generating set X . Then, either some $m_{ij} = \infty$, or Γ contains a triangle with at most one edge labelled 2.*

Proof. Assume that A_Γ has no $m_{ij} = \infty$. Since A_Γ is infinite type, it must contain at least 3 generators, that is, $|X| \geq 3$. We will prove by induction on $|X|$ that Γ contains a triangle with at most one edge labelled 2. Suppose that $X = \{a, b, c\}$. If a commutes with both b and c , then $A_\Gamma \cong \mathbb{Z} \times A_{\{b,c\}}$ which contradicts our assumption that A_Γ is infinite type.

Now, suppose that $X = \{x_1, \dots, x_n\}$, $n > 3$. Since A_Γ is infinite type it must contain an edge connecting some x_i, x_j with $m_{ij} \neq 2$. If every triangle containing this edge has the other two edges labelled 2, then x_i and x_j commute with all other generators x_k . Thus,

$$A_\Gamma \cong A_R \times A_T,$$

where $R = \{x_i, x_j\}$ and $T = X \setminus R$. Since A_R is spherical type, A_T must be infinite type, so by induction, the subgraph spanned by T contains a triangle with at most one edge labelled 2. ■

From now on, we will assume that our Artin groups A_Γ are of infinite type.

A solution to the word problem for large type Artin groups is given in [9] and we will rely heavily on the consequences of this algorithm. The algorithm works by successive manipulations of subwords of a non-geodesic word until a free cancellation

is possible. In [1], the algorithm is modified to apply to 3-free Artin groups by the addition of specific commutations between two generators.

We will say that two geodesics words are *equivalent* if they represent the same element of A_Γ . We will use the following key takeaway from these papers to prove Criterion 1.

Proposition 5.2. *Properties of rightward reducing sequences [9, Proposition 3.3], [1, Proposition 5.1]*

Let A_Γ be a large type or 3-free Artin group, w be a geodesic word over X , and x be a letter in $x \in X \cup X^{-1}$ such that wx is not geodesic. Then, there is a suffix ζ of w and an algorithm called a rightward reducing sequence (RRS) such that when the RRS is applied to w , ζ is transformed into an equivalent word ζ' that ends in x^{-1} . Furthermore, if ζ begins in a positive (resp., negative) letter, then ζ' ends in a positive (resp., negative) letter.

5.1. Criterion 1: Preserving signed suffixes

Lemma 5.3. *Let A_Γ be an Artin group of large type or 3-free type. Then, A_Γ has preserved signed suffixes.*

Proof. Suppose that w is a geodesic word over X in A_Γ , with longest positive suffix u , and suppose that a is a positive generator from X .

If aw is geodesic, then u is also a suffix of the geodesic word aw .

If aw is non-geodesic but not freely reduced, then the first letter of w must be a^{-1} and this cancellation must produce a geodesic representative of aw that has u as a suffix.

So, now suppose that aw is freely reduced but not geodesic. Factor aw as aw_1w_2 , where aw_1 is the largest prefix of aw that is geodesic. Denote by x the first letter of w_2 . By Proposition 5.2, there is an RRS applied to a suffix ζ of aw_1 that results in a new geodesic ending in x^{-1} . This suffix ζ must start with the positive letter a , as otherwise the RRS could be applied to w_1 and w would not be geodesic. So, $\zeta = aw_1$, and after performing the RRS algorithm, we get a word ζ' equivalent to aw_1 . By Proposition 5.2, ζ' ends in a positive letter, since a was positive. We know that this letter is x^{-1} , so we conclude that x is a negative letter. This means that u , as a positive suffix of aw , must be a proper suffix of w_2 .

Note also that w is geodesic so that any word representing the same group element as aw must have length at least $|w|_X - 1$. For suppose to the contrary that there were a shorter word ξ representing the same element as aw . Then, $a^{-1}\xi$ is a word representing the same element as w but with length smaller than $|w|_X$, contradicting w being geodesic. This implies that, applying a single RRS on aw followed by the free cancellation between x^{-1} and x , we obtain a geodesic representative of the same

group element as aw . As explained in the previous paragraph, except for the cancellation of the first letter, the suffix w_2 has been unaffected by this algorithm; thus, we have constructed a geodesic representative of aw that ends in u . ■

5.2. Criterion 2: Constructing alternating blocking sequences

In this section, we will explain how to construct an alternating blocking sequence for large type and 3-free Artin groups with at least 3 generators. Once this sequence is constructed, we may apply Theorem 4.5 and Corollary 2.3 to conclude that any infinite type Artin group A_Γ that is either large type or 3-free has infinite diameter monoid Cayley graph $\text{Cay}(A_\Gamma, M)$.

To prove Criterion 2, we rely on the following:

Proposition 5.4. *For suffixes of equivalent geodesics, we have the following:*

- (1) *As per [9, Proposition 4.5], suppose that A_Γ is a large type Artin group and that $a, b \in X$ are distinct generators. Given a geodesic word $wb^\pm a^\pm$, any equivalent geodesic word must end in either a^\pm or b^\pm .*
- (2) *Suppose that A_Γ is a 3-free Artin group and $a, b, c \in X$ are distinct generators such that $m_{a,b} > 3$ and $m_{b,c} > 3$*
 - (a) *Given a geodesic word $wb^\pm a^\pm$, any equivalent geodesic must end in a^\pm , b^\pm or a letter that commutes with a .*
 - (b) *Given a geodesic word $wb^\pm c^\pm b^\pm a^\pm$, any equivalent geodesic must end in a^\pm or a letter that commutes with a .*

Part (2) above of the above proposition is new. We will prove Criterion 2 is satisfied for both large type and 3-free Artin groups assuming this result and return to the proof of part (2) at the end of this section.

Lemma 5.5. *Let A_Γ be an Artin group of large type. Then, A_Γ contains an alternating blocking sequence.*

Proof. Suppose that A_Γ is a large type Artin group and that $a, b, c \in X$ are three distinct positive generators. We define an alternating blocking sequence by

$$\begin{aligned} \alpha_1 &= ab, & \alpha_2 &= c^{-1}a^{-1}, & \alpha_3 &= bc, \\ \alpha_4 &= a^{-1}b^{-1}, & \alpha_5 &= ca, & \alpha_6 &= b^{-1}c^{-1}, \end{aligned}$$

and for $i > 6$, $\alpha_i = \alpha_{i-6}$.

This is a sequence of words in the standard generators such that α_i is a positive word when i is odd and a negative word when i is even. It remains to verify that for any letter x in α_i there exists some subword u of $\alpha_{i-1}\alpha_i$ immediately preceding x

such that (u, x) is a blocking pair. Without loss of generality, we assume that $i = 2 \pmod 6$ as other arguments are symmetric. Thus, it remains to show that both (ab, c^{-1}) and (bc^{-1}, a^{-1}) are blocking pairs.

Suppose to the contrary that, for some word w , wab is geodesic, but $wabc^{-1}$ is not. Then, by Proposition 5.2, the word wab admits a rightward reducing sequence that transforms it to a word w' such that the final letter of w' is c . However, since w' is a geodesic representative of wab , this contradicts part (1) of Proposition 5.4, which requires that every geodesic representative of wab must end with a letter in the set $\{a, b, a^{-1}, b^{-1}\}$. Hence, (ab, c^{-1}) is a blocking pair. Notice that in the above argument we used only the fact that a, b, c were distinct generators in a large type Artin group, so an identical argument can be used to show that (bc^{-1}, a^{-1}) is a blocking pair. This shows that the given sequence $\{\alpha_i\}$ is an alternating blocking sequence. ■

Lemma 5.6. *Let A_Γ be a 3-free Artin group of infinite type such that no $m_{ij} = \infty$. Then, A_Γ contains an alternating blocking sequence.*

Proof. By Lemma 5.1, Γ contains three generators a, b, c such that at most one of the relations is a commutation. We have also already covered large type Artin groups, so we can assume that

$$m_{a,b} > 3, \quad m_{b,c} > 3, \quad m_{a,c} = 2.$$

We define an alternating sequence of words by

$$\alpha_i = abc \quad \text{if } i \text{ is odd,} \quad \alpha_i = b^{-1}a^{-1}b^{-1}c^{-1} \quad \text{if } i \text{ is even.}$$

This sequence alternates between positive and negative words in the standard generators, and so, it remains to show that for any letter x in α_i there exists some subword u of $\alpha_{i-1}\alpha_i$ immediately preceding x such that (u, x) is a blocking pair. Assuming without loss of generality that i is even, it follows as in the proof of Lemma 5.5 that

$$(abc, b^{-1}), \quad (cb^{-1}, a^{-1}), \quad (bcb^{-1}a^{-1}, b^{-1}), \quad \text{and} \quad (a^{-1}b^{-1}, c^{-1})$$

are all blocking pairs, using part (2) of Proposition 5.4 in place of part (1). Note that in some cases part (2) of Proposition 5.4 is applied to the three generators ordered as c, b, a rather than as a, b, c , which is valid because of the symmetry in the triple $\{m_{a,b}, m_{a,c}, m_{b,c}\}$. ■

We conclude this section with the proof of part (2) of Proposition 5.4, for which we require more technical elements of the solution to the word problem as given in [1]. Note that we essentially follow the notation from [1], except that we have used subwords ξ_i here where subwords w_i are used in [1], in order to avoid confusion with our previously defined sequence w_n .

Recall that given a word u we can write

$$u = x_1^{k_1} x_2^{k_2} \cdots x_n^{k_n},$$

where we assume that the word is freely reduced and $x_i \neq x_{i+1}$. We call n the syllable length of this word.

Remark 5.7. To prove part (2) of Proposition 5.4, we will use the following key facts from [1]. These facts follow directly from the way the algorithm for the solution to the word problem is defined in that paper. Suppose that a word w can be changed to a new word w' via a rightward reducing sequence. Then, w has a suffix $\xi_1 \xi_2 \cdots \xi_{k+1}$ satisfying the following properties:

- (F1) For $1 \leq i \leq k$, the first i steps of the algorithm involve a modification to $\xi_1 \cdots \xi_i$ and leave the suffix $\xi_{i+1} \cdots \xi_{k+1}$ unaffected.
- (F2) The final $(k+1)^{\text{st}}$ step in the algorithm involves only commutations of individual letters. Thus, if w' ends in a letter x , then either x commutes with all the letters in ξ_{k+1} , or ξ_{k+1} is the empty word, in which case the algorithm only involves k steps.
- (F3) The 3-free condition, together with [1, Lemma 3.11], implies that the syllable length of ξ_i must be at least 2 for all $i \leq k$.
- (F4) If ξ_i has $x^\pm y^\pm$ as a suffix and $m_{x,y} > 3$, then after the application of the i th step in the algorithm, the word created has suffix $\mu \xi_{i+1} \cdots \xi_{k+1}$, where μ is a word in x, y of syllable length 2.
- (F5) The subword ξ_i cannot end in $x^\pm y^\pm z^\pm$, where $m_{y,z} > 3$ and $m_{x,y} > 3$.

Proof of Proposition 5.4 (2). To prove part (a), suppose that u is a geodesic word ending in the letter x and representing the same group element as $wb^\pm a^\pm$. Then, $wb^\pm a^\pm x^{-1}$ is not geodesic and it follows from Proposition 5.2 that either $wb^\pm a^\pm x^{-1}$ is not freely reduced, in which case, $x = a^\pm$ and we are done, or the word $wb^\pm a^\pm$ can be changed via a rightward reducing sequence to a new word u' ending in x .

Our goal is to show that one of the cases $x = a^\pm$, $x = b^\pm$, or $m_{x,a} = 2$ holds. We consider a factorisation of $wb^\pm a^\pm$ with suffix $\xi_1 \xi_2 \cdots \xi_{k+1}$. If ξ_{k+1} is non-empty, then it ends in a^\pm , and then by (F2), x must commute with a . Otherwise, ξ_{k+1} is empty. Then, by (F3), ξ_k has $b^\pm a^\pm$ as a suffix, and, by (F4), after the k^{th} and final step in the algorithm, the result is a word that ends in a^\pm or b^\pm .

Now, we prove part (b). As for part (a), we consider a word u ending in x and equivalent to $wb^\pm c^\pm b^\pm a^\pm$. If $wb^\pm c^\pm b^\pm a^\pm x^{-1}$ is not geodesic, then either $wb^\pm c^\pm b^\pm a^\pm x^{-1}$ is not freely reduced and $x = a^\pm$ or we can use a rightward reducing sequence to obtain a new word u' ending in x and equivalent to $wb^\pm c^\pm b^\pm a^\pm$ by

Proposition 5.2. We will show that the suffix $b^\pm c^\pm b^\pm a^\pm$ implies that there is no such sequence.

We have already shown in the proof of part (a) that if ξ_{k+1} is non-empty then x must commute with a , and if ξ_{k+1} is empty, then $b^\pm a^\pm$ must be a suffix of ξ_k . We need now to consider further the case where ξ_{k+1} is empty, and consider the two cases where c^\pm is a part of ξ_k and where c^\pm is a part of ξ_{k-1} . In the first case, $c^\pm b^\pm a^\pm$ is a suffix of the word ξ_k , but then, since

$$m_{b,c} > 3 \quad \text{and} \quad m_{a,b} > 3,$$

by (F5) the algorithm cannot proceed to achieve our word ending in x , a contradiction.

If, on the other hand, c^\pm is part of ξ_{k-1} , then ξ_{k-1} must have $b^\pm c^\pm$ as a suffix. Then, (F4) implies that after the $(k-1)^{st}$ step in the algorithm the word still ends in $\mu b^\pm a^\pm$, where μ is a word in b, c of syllable length 2, that is, $\mu b^\pm a^\pm$ is either of the form

$$c^i b^j a^\pm \quad \text{or} \quad b^j c^i b^\pm a^\pm.$$

In either case, (F5) implies that the algorithm cannot proceed, which is again a contradiction. ■

6. Conclusions

Combining Theorem 4.5 with Lemmas 5.3, 5.5, and 5.6, we conclude that large type and 3-free Artin groups (of infinite type) have infinite diameter monoid Cayley graphs. Together with Corollary 2.3, we thus obtain our main theorem.

Theorem 6.1. *Let $\text{Cay}(A_\Gamma, M)$ denote the Cayley graph of A_Γ with respect to the generating set $M := A_\Gamma^\pm$. Then, $\text{Cay}(A_\Gamma, M)$ has infinite diameter providing one of the following holds:*

- (1) Γ contains a pair of vertices not connected by an edge (or equivalently, there exists i, j with $m_{i,j} = \infty$).
- (2) Γ contains a triangle with all labels ≥ 3 .
- (3) Γ contains a triangle with no label equal to 3, and at most one label is equal to 2.

Recall that Conjecture 2.1 states that $\text{Cay}(A_\Gamma, M)$ has infinite diameter for all infinite type Artin groups. To complete the proof of this conjecture, it remains to consider the case in which all 3-generator special subgroup of A_Γ are either of spherical type or correspond to triangles labelled $(2, 3, n)$, $n \geq 6$. We believe that the latter case can be solved by arguments similar to those used in the large type and 3-free cases.

Acknowledgements. The authors would like to thank Derek Holt for helpful conversations and suggestions, and the anonymous referee for helpful comments. The authors would also like to thank ICERM for hosting the 2022 semester on Braids where some discussions for this project took place.

Funding. The first author was supported by EPSRC Fellowship no. EP/V043323/1 and EP/V043323/2.

References

- [1] R. Blasco-García, M. Cumplido, and R. Morris-Wright, The word problem is solvable for 3-free Artin groups in quadratic time. 2022, arXiv:2204.03523
- [2] R. Boyd, R. Charney, and R. Morris-Wright, A Deligne complex for Artin monoids. *J. Algebra* **607** (2022), no. part B, 53–78 Zbl 1517.20056 MR 4441310
- [3] E. Brieskorn and K. Saito, Artin-Gruppen und Coxeter-Gruppen. *Invent. Math.* **17** (1972), 245–271 Zbl 0243.20037 MR 323910
- [4] R. Charney and M. W. Davis, The $K(\pi, 1)$ -problem for hyperplane complements associated to infinite reflection groups. *J. Amer. Math. Soc.* **8** (1995), no. 3, 597–627 Zbl 0833.51006 MR 1303028
- [5] R. Charney and L. Paris, Convexity of parabolic subgroups in Artin groups. *Bull. Lond. Math. Soc.* **46** (2014), no. 6, 1248–1255 Zbl 1308.20037 MR 3291260
- [6] P. Dehornoy, F. Digne, E. Godelle, D. Krammer, and J. Michel, *Foundations of Garside theory*. EMS Tracts Math. 22, EMS Press, Zürich, 2015 Zbl 1370.20001 MR 3362691
- [7] P. Deligne, Les immeubles des groupes de tresses généralisés. *Invent. Math.* **17** (1972), 273–302 Zbl 0238.20034 MR 422673
- [8] E. Godelle and L. Paris, $K(\pi, 1)$ and word problems for infinite type Artin-Tits groups, and applications to virtual braid groups. *Math. Z.* **272** (2012), no. 3-4, 1339–1364 Zbl 1300.20045 MR 2995171
- [9] D. F. Holt and S. Rees, Artin groups of large type are shortlex automatic with regular geodesics. *Proc. Lond. Math. Soc. (3)* **104** (2012), no. 3, 486–512 Zbl 1275.20034 MR 2900234
- [10] A. Martin and P. Przytycki, Acylindrical actions for two-dimensional Artin groups of hyperbolic type. *Int. Math. Res. Not. IMRN* **2022** (2022), no. 17, 13099–13127 Zbl 07582349 MR 4475273
- [11] J. McCammond, An introduction to Garside structures. 2005, <https://web.math.ucsb.edu/~jon.mccammond/papers/intro-garside.pdf>
- [12] J. Michel, A note on words in braid monoids. *J. Algebra* **215** (1999), no. 1, 366–377 Zbl 0937.20017 MR 1684142
- [13] G. Paolini and M. Salvetti, Proof of the $K(\pi, 1)$ conjecture for affine Artin groups. *Invent. Math.* **224** (2021), no. 2, 487–572 Zbl 07355938 MR 4243019
- [14] L. Paris, Artin monoids inject in their groups. *Comment. Math. Helv.* **77** (2002), no. 3, 609–637 Zbl 1020.20026 MR 1933791

- [15] L. Paris, *$K(\pi, 1)$ conjecture for Artin groups*. *Ann. Fac. Sci. Toulouse Math. (6)* **23** (2014), no. 2, 361–415 Zbl [1306.20040](#) MR [3205598](#)
- [16] H. van der Lek, *The homotopy type of complex hyperplane complements*. Ph.D. thesis, Katholieke Universiteit te Nijmegen, Nijmegen, 1983

Received 31 March 2023; revised 12 July 2023.

Rachael Boyd

School of Mathematics and Statistics, University of Glasgow, Glasgow G12 8QQ, UK;
rachael.boyd@glasgow.ac.uk

Ruth Charney

Department of Mathematics, Brandeis University, 415 South St., Waltham, MA 02421, USA;
charney@brandeis.edu

Rose Morris-Wright

Department of Mathematics, Middlebury College, 14 Old Chapel Rd, Middlebury, VT 05753, USA; rmorriswright@middlebury.edu

Sarah Rees

School of Mathematics, Statistics and Physics, University of Newcastle, Newcastle NE1 7RU, UK; sarah.rees@newcastle.ac.uk