

Quadratic automaton algebras and intermediate growth

Natalia Iyudu and Stanislav Shkarin

Abstract. We present an example of a quadratic algebra given by three generators and three relations, which is automaton (the set of normal words forms a regular language) and such that its ideal of relations does not possess a finite Gröbner basis with respect to any choice of generators and any choice of a well-ordering of monomials compatible with multiplication. This answers a question of Ufnarovski.

Another result is a simple example (4 generators and 7 relations) of a quadratic algebra of intermediate growth.

Mathematics Subject Classification (2010). 17A45, 16A22.

Keywords. Automaton algebras, word combinatorics, regular language, Gröbner basis, quadratic algebras, Koszul algebras, finitely presented algebras, Hilbert series, Gelfand–Kirillov dimension.

1. Introduction

In the first part of the paper we consider automaton algebras, that is algebras which could be associated to a finite graph. This is a property of an algebra, formulated in terms of language of its normal words, that is in terms of words combinatorics. This approach to the study of algebra (its linear basis, Hilbert series, growth, structural properties) is very powerful, widely known, see, for example, book of collective author M. Lothaire [8] or A. Belov, V. Borisenko, V. Latyshev [1], or book of M. Sapir [12], and fruitfully connects algebraic study with symbolic dynamics.

The class of automaton algebras was introduced by Ufnarovski [15] (see also his book [14, Chapter 5]). He asked if every automaton algebra admits a finite Gröbner basis in its ideal of relations with respect to some choice of generators and an ordering on monomials. It is a very natural question, and many people tried to figure it out, however without a success. The difficulty here lies in the fact that there are infinitely many generating sets as well as infinitely many admissible orderings on monomials to deal with.

We argue, using mod p specialization of coefficients of initial algebra over the field of characteristic zero. On the way we prove quite interesting statement, allowing

to capture the fact of existence of finite Gröbner basis. It says that if algebra over a field of characteristic zero did have a finite Gröbner bases, then after specialization mod p its growth can't go up (thus stays the same) for all but finitely many p . Then the difficulty of considering all possible orderings transforms to consideration of all possible mod p specializations, and this turns out to be more tractable, we are able to follow the the pattern of the Gröbner basis for any p . We prove that in a particular example, namely, the algebra with three generators x, y, z , and relations

$$xx - xz - 2zz = 0, \quad yz = 0, \quad xy = 0,$$

which is automaton and has a polynomial growth, Hilbert series $H(t) = \frac{1}{(1-t)^3}$ in characteristic zero and two, in any odd finite characteristic becomes an algebra of exponential growth.

Thus, the first main result of this paper is an example of a quadratic automaton algebra A given by three generators and three relations such that the ideal of relations of A does not possess a finite Gröbner basis with respect to any choice of generators and any choice of a well-ordering on monomials compatible with multiplication. A finite Gröbner basis does not exist after any extension of the ground field as well.

The second our result is a simple example of quadratic algebra (4 generators, 7 relations) with intermediate growth.

One example of finitely presented cubic algebra of intermediate growth was mentioned in a note added in proof of the paper by Shearer [13]. There was no explanation why the Hilbert series is as stated, and it was elaborated in paper by Kobayashi [6], where he also constructs other examples and applies them to the word problem in monoids. Another example was due to Ufnarovskii [16], who proves that the universal enveloping of the (infinite dimensional) Witt Lie algebra is finitely presented and has intermediate growth. The paper [5] by Kirillov, Kontsevich and Molev contains a number of results on algebras of intermediate growth both in the associative and Lie setting together with yet another example of universal enveloping with intermediate growth. All these examples were not quadratic. Recently, Koçak [7] used the idea suggested by Zelmanov, that one can take a Veronese subalgebra of (a slightly modified to make it degree graded) universal enveloping of the Witt algebra, to obtain quadratic algebra with intermediate growth. One drawback of his example is its size: it is presented by 14 generators and 96 quadratic relations. Literally, the presentation of this quadratic algebras in [7] has two and a half pages of relations. So we thought, that example of quadratic algebra with 4 generators and 7 relations, with intermediate growth, could be of interest.

Now we formulate results more precisely. We start by recalling relevant definitions. All algebras we deal with are associative. Throughout the paper, \mathbb{K} is a field. If A is a unital \mathbb{K} -algebra and $X \subset A$ generates A (as a unital algebra), then A can be naturally interpreted as the quotient of the free algebra $\mathbb{K}\langle X \rangle$ on generators X by the ideal I consisting of all $f \in \mathbb{K}\langle X \rangle$ vanishing when considered

as elements of A . The ideal I is known as the *ideal of relations* of A . A generating set X together with a set $F \subseteq I$ generating I as a (two-sided) ideal is known as a *presentation* of A . An algebra is called *finitely presented* if X and F can be chosen finite. In other words, a finitely presented algebra is an algebra isomorphic to a quotient of $\mathbb{K}\langle X \rangle$ with finite X by a finitely generated ideal. The set of all words (including the empty word 1) in the alphabet X is denoted $\langle X \rangle$. A well-ordering \leq on $\langle X \rangle$ is said to be *compatible with multiplication* if

$$1 \leq u \text{ for all } u \in \langle X \rangle \text{ and } u \leq v \implies uw \leq vw, \quad wu \leq wv \text{ for all } u, v, w \in \langle X \rangle.$$

If we fix a well-ordering \leq on $\langle X \rangle$ compatible with multiplication, we can talk of the leading monomial \overline{f} of a non-zero $f \in \mathbb{K}\langle X \rangle$ (=the biggest with respect to \leq monomial, which features in f with non-zero coefficient). A subset G of an ideal I in $\mathbb{K}\langle X \rangle$ is called a *Gröbner basis of I* if $0 \notin G$, G generates I as an ideal and for each non-zero $f \in I$, there is $g \in G$ such that \overline{g} is a subword of \overline{f} . Such a subset is by no means unique: for one, $I \setminus \{0\}$ fits the bill. However, a couple of extra conditions pinpoint G . Namely, if we additionally assume that a Gröbner basis G satisfies

- for every two distinct $f, g \in G$, \overline{f} is not a subword of any monomial featuring in g ;
- every $f \in G$ is monic: the \overline{f} -coefficient in f equals 1,

then G becomes unique. Such a basis is called the *reduced Gröbner basis of I* . Note that I possesses a finite Gröbner basis if and only if its reduced Gröbner basis is finite.

The non-commutative Buchberger algorithm [2] applied to the set of defining relations yields the reduced Gröbner basis for the ideal of relations of any finitely presented algebra. One of the problems though is that (unlike for the commutative case) the procedure does not have to terminate in finitely many steps. What is even worse, there is no *a-priory* way to say if it does. Furthermore, everything is highly sensitive to the choice of the generators and the ordering. The words $u \in \langle X \rangle$, which have no leading monomials of elements of the ideal I of relations of A as subwords, are called *normal words* for A . It is easy to see that normal words form a linear basis in A . Clearly, if G is a Gröbner basis for I , then a word $u \in \langle X \rangle$ is normal if and only if it has no leading monomials of elements of G as subwords.

Recall that if X is a finite set, then a subset L of $\langle X \rangle$ forms a *regular language* if there is a finite quiver (loops and multiple edges are allowed) with each arrow marked by a letter from X , while some vertices are marked by symbols S and/or T such that L consists (exactly) of the words that can be read from paths in this quiver starting at a vertex marked by S and terminating at a vertex marked by T . Ufnarovski [14] has introduced the class of automaton algebras in the following way.

Definition 1.1. A finitely generated algebra A is called an *automaton algebra* if there exists a finite generating set X for A such that with respect to some well-ordering

on $\langle X \rangle$ compatible with multiplication, the set NW of normal words for A forms a regular language.

There is correspondence between the set of normal words NW , and the set LM of leading monomials of the reduced Gröbner basis of the ideal of relations of A , namely each of them can be reconstructed from another. Namely, NW is the set of all monomials which does not contain elements of LM , as subwords, and LM , are all those words in which all subwords are from NW , but they themselves are not. Hence NW is a regular language if and only if the set LM is a regular language. This means that NW and LM are interchangeable in the definition of automaton algebras. In particular, an algebra, whose ideal of relations possesses a finite Gröbner basis (with respect to some choice of a generating set and an ordering on monomials), is an automaton algebra. On the other hand, there are examples [9, 14] of automaton algebras with infinite Gröbner bases. The question mentioned above is whether this may be rectified by a clever choice of generators and of an ordering.

Question 1.2. *Given an automaton algebra A , is it possible to find a finite generating set X and an ordering on $\langle X \rangle$ compatible with multiplication such that the ideal of relations of A has a finite Gröbner basis?*

Theoretically speaking, there is one more degree of freedom here. Replacing the ground field \mathbb{K} by a bigger field (a one containing \mathbb{K} as a subfield) preserves the automaton property, but it yields extra opportunities for a finite Gröbner basis to exist. Replacing \mathbb{K} by a bigger field \mathbb{F} amounts to replacing a finitely presented \mathbb{K} -algebra A by $A_{\mathbb{F}} = \mathbb{F} \otimes_{\mathbb{K}} A$ treated as an \mathbb{F} -algebra. Clearly, any presentation of A as a \mathbb{K} -algebra works as a presentation of $A_{\mathbb{F}}$ as an \mathbb{F} -algebra (but not vice versa).

Note that automaton algebras share a lot of properties with algebras possessing a finite Gröbner basis in the ideal of relations [3, 9, 10, 14, 15]. For instance, the Hilbert series of each automaton algebra is rational (hence intermediate growth is impossible), nil automaton algebras are finite dimensional, an automaton algebra of exponential growth contains a subalgebra isomorphic to the free algebra on two generators, etc. This was another evidence for the conjecture that these two classes coincide. It is also quite clear, that not every quadratic algebra is automaton. This could be deduced, for instance, from the example by Shearer [13] of a quadratic algebra (= a finitely presented algebra with all defining relations being homogeneous of degree 2), whose Hilbert series is not rational. Shearer produced this example answering a well-known question of Govorov [4] whether every finitely presented algebra has rational Hilbert series.

Definition 1.3. We say that a finitely presented \mathbb{K} -algebra A is a *finite Gröbner basis algebra (FGB-algebra for short)*, if there exists a field \mathbb{F} containing \mathbb{K} as a subfield such that there is a finite Gröbner basis in the ideal of relations of $A_{\mathbb{F}}$ for some choice of a finite generating set X and of a well-ordering on $\langle X \rangle$ compatible

with multiplication. Otherwise, we say that A is an *infinite Gröbner basis algebra* (IGB-algebra for short).

Our main objective is to provide examples of finitely presented algebras, which are IGB and automaton. Our method allows to construct unlimited number of such examples. We give two: one quadratic on three generators given by three relations and one cubic on two generators given by two relations.

Theorem 1.4. *Let \mathbb{K} be an arbitrary field of characteristic zero. Let A be the \mathbb{K} -algebra given by generators x, y, z and relations xy, yz and $x^2 - xz - 2z^2$ and B be the \mathbb{K} -algebra given by generators x, y and relations y^3 and $x^2y - yx^2 - yxy$. Then both A and B are automaton IGB-algebras.*

The advantage of A is in being quadratic, while the advantage of B is that it is generated by just two elements. The proof is based on the fact (Theorem 2.4) that the existence of finite Gröbner basis implies that the growth of algebra and any its specialization mod p coincide (for almost all p). Then the example of algebra without finite Gröbner basis, is an algebra with integer coefficients and finite Gelfand–Kirillov dimension, which becomes infinite when coefficients are written mod p , for all prime p except from 2. The growth is determined via calculation of Gröbner bases and Hilbert series.

Before proceeding, we would like to make one more comment. Ufnarovskii [14, 15] demonstrated that a quadratic algebra given by two (homogeneous degree 2) relations is automaton. What one can actually observe is that it is always an FGB-algebra. On the other hand, one easily checks that any quadratic algebra on two generators is an FGB-algebra as well. Thus, as far as quadratic algebras are concerned, our example is minimal: we could not possibly have less than three generators or less than three relations.

1.1. Growth of finitely generated algebras. Part of our argument is based on the growth of finitely generated algebras. We recall relevant concepts. Let A be a finitely generated unital \mathbb{K} -algebra and X be a finite generating set for A . Let $A^{(n)} = A_X^{(n)}$ be a linear subspace of A spanned by the products of up to n elements of X (we include 1, which is the product of zero elements of X). Clearly, $A^{(n)}$ form a filtration on A . The power series

$$H_A(t) = H_A^X(t) = \sum_{n=0}^{\infty} a_n t^n$$

with $a_0 = 1$ and $a_n = a_n(X) = \dim A^{(n)}/A^{(n-1)}$ for $n \geq 2$ is called the *Hilbert series* of A (with respect to the generating set X). If we know the normal words for A with respect to some well-ordering on $\langle X \rangle$, we know the Hilbert series as well: a_n is the number of normal words of degree n . Note that if we assume that each element of X has degree 1 and that each defining relation is homogeneous, then

the ideal of relations I and therefore A itself are degree graded. In this case a_n becomes the dimension of the space A_n of homogeneous elements of A of degree n (by convention, zero has any degree we like). The series

$$P_A(t) = P_A^X(t) = \frac{H_A^X(t)}{1-t} = \sum_{n=0}^{\infty} \alpha_n t^n$$

with $\alpha_n = \alpha_n(X) = a_0 + \cdots + a_n = \dim A^{(n)}$ is called the *Poincaré series* of A (with respect to the generating set X).

Let A be a finitely generated unital \mathbb{K} -algebra. Obviously, the Hilbert and Poincaré series H_A^X and $P_A^X(t) = \sum \alpha_n(X)t^n$ depend on the choice of the generating set X . However, some of their features do not depend on this choice. Indeed, let X and Y be two finite generating sets of A . Since X and Y are finite, we can choose $m \in \mathbb{N}$ such that $X \subset A_Y^{(m)}$ and $Y \subset A_X^{(m)}$. Then

$$\alpha_n(X) \leq \alpha_{mn}(Y) \quad \text{and} \quad \alpha_n(Y) \leq \alpha_{mn}(X) \quad \text{for all } n \in \mathbb{Z}_+. \quad (1.1)$$

In particular, it follows that the limits

$$\text{GKdim}(A) = \limsup_{n \rightarrow \infty} \frac{\ln \alpha_n(X)}{\ln n} \in [0, \infty] \quad (1.2)$$

and

$$\varkappa(A) = \limsup_{n \rightarrow \infty} \frac{\ln \ln \alpha_n(X)}{\ln n} \in [0, 1] \quad (1.3)$$

do not depend on the choice of X .

Definition 1.5. The number $\text{GKdim}(A)$ defined in (1.2) is known as the *Gelfand–Kirillov dimension* of A . Algebras of finite Gelfand–Kirillov dimension are said to have *polynomial growth*. On the other hand, if

$$\limsup_{n \rightarrow \infty} \frac{\ln \alpha_n(X)}{n} > 0,$$

we say that A has *exponential growth*. A finitely generated algebra A which has infinite Gelfand–Kirillov dimension and is not of exponential growth is said to have *intermediate growth*.

Remark 1.6. Again, for an algebra to have exponential growth does not depend on the choice of a finite generating set X . Note that while the limit in the above display does depend on X , it being positive does not according to (1.1). Obviously, every algebra with exponential growth has infinite Gelfand–Kirillov dimension. Note also that polynomial and exponential growths are related to the number $\varkappa(A)$ defined in (1.3):

$$\begin{aligned} &\text{if } A \text{ has polynomial growth, then } \varkappa(A) = 0; \\ &\text{if } A \text{ has exponential growth, then } \varkappa(A) = 1. \end{aligned} \quad (1.4)$$

We use growth arguments in the proof of Theorem 1.4. The second result of this paper is the following example.

Theorem 1.7. *Let \mathbb{K} be an arbitrary field and C be the \mathbb{K} -algebra given by four generators x, y, z, u and seven quadratic relations $xu - yz, yu - zx, zu - uz, yy, yx, xy$ and xx . Then C has intermediate growth.*

In Section 2 we describe a way to verify that a finitely presented algebra does not possess a finite Gröbner basis in its ideal of relations for any choice of generators and ordering. We also do the preparatory work for applying the general result to particular algebras of Theorem 1.4. We study the Gröbner bases of members of two varieties of finitely presented algebras containing the algebras A and B of Theorem 1.4 in Section 3, which culminates in the proof of Theorem 1.4. We prove Theorem 1.7 in Section 4 and make few extra remarks in the final Section 5.

2. The IGB-criterion

Everywhere below we use the following notation. The symbol \mathbb{Q} denotes the field of rational numbers, \mathbb{Z} stands for the ring of integers, \mathbb{N} is the set of positive integers, while $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$ is the set of non-negative integers. If p is a prime number then $\mathbb{Z}_p = \mathbb{Z}/p\mathbb{Z}$ is the p -element field. We denote by π_p the canonical map $\pi_p : \mathbb{Z} \rightarrow \mathbb{Z}_p$: $\pi_p(n)$ is the congruence class of n modulo p . We naturally extend π_p to a ring homomorphism

$$\pi_p : \mathbb{Z}\langle x_1, \dots, x_n \rangle \rightarrow \mathbb{Z}_p\langle x_1, \dots, x_n \rangle$$

(we use the same symbol to denote the extension) by acting on coefficients. That is, for $f \in \mathbb{Z}\langle x_1, \dots, x_n \rangle$, $\pi_p(f)$ is obtained from f by replacing each coefficient of f by its congruence class modulo p . We would like to introduce the following technical notions.

Definition 2.1. Let \mathbb{K} be a field and R_0 be a subring of \mathbb{K} . We say that R is a *finite ring extension of R_0 in \mathbb{K}* if R is a subring of \mathbb{K} , $R_0 \subseteq R$ and there is a finite set $S \subset R$ such that the set $R_0 \cup S$ generates R as a ring.

Definition 2.2. We say that a map $A \mapsto \gamma(A)$, which assigns to a finitely generated algebra A a number $\gamma(A) \in [0, \infty]$ is a *growth measuring function* if the following conditions are satisfied:

- $\gamma(A) = \gamma(A_{\mathbb{F}})$ for every field \mathbb{F} containing the ground field \mathbb{K} of A as a subfield;
- if A and B are two finitely generated algebras (possibly over different fields) and there exist finite generating sets X and Y for A and B respectively such that the corresponding Poincaré series $P_A^X = \sum \alpha_n t^n$ and $P_B^Y = \sum \beta_n t^n$ satisfy $\alpha_n \leq \beta_n$ for all $n \in \mathbb{Z}_+$, then $\gamma(A) \leq \gamma(B)$.

Remark 2.3. Note that the Gelfand–Kirillov dimension $\text{GKdim}(A)$ as well as $\varkappa(A)$ defined in (1.3) are growth measuring functions. The dimension $\dim A$ of a finitely generated \mathbb{K} -algebra A as a \mathbb{K} -vector space is a growth measuring function as well. One can come up with quite a few other examples.

Theorem 2.4. *Let (X, F) be a finite presentation of a \mathbb{K} -algebra A and let R_0 be a unital subring of \mathbb{K} such that each $f \in F$ belongs to $R_0\langle X \rangle$. That is, all coefficients of each $f \in F$ are in R_0 . Assume also that we have a collection of non-zero ring homomorphisms $\rho_\alpha : R_0 \rightarrow \mathbb{K}_\alpha$ labelled by α from some index set Λ such that each \mathbb{K}_α is a field. We extend every ρ_α to a ring homomorphism $\rho_\alpha : R_0\langle X \rangle \rightarrow \mathbb{K}_\alpha\langle X \rangle$ by the original ρ_α acting on coefficients. For each $\alpha \in \Lambda$, let A_α be the \mathbb{K}_α -algebra defined by the presentation (X, F_α) , where $F_\alpha = \rho_\alpha(F)$. Finally, assume that γ is a growth measuring function and the following conditions are satisfied:*

- (1) $\gamma(A_\alpha) > \gamma(A)$ for every $\alpha \in \Lambda$;
- (2) for every field \mathbb{F} containing \mathbb{K} as a subfield and for every finite ring extension R of R_0 in \mathbb{F} , there exists $\alpha \in \Lambda$ such that ρ_α extends to a ring homomorphism $\rho'_\alpha : R \rightarrow \mathbb{F}_\alpha$, where \mathbb{F}_α is some field containing \mathbb{K}_α as a subfield.

Then A is an IGB-algebra.

Proof. Assume the contrary: A is an FGB-algebra. Then there is a field \mathbb{F} containing \mathbb{K} as a subfield such that $A_\mathbb{F}$ has a finite generating set Y and there is a well-ordering on $\langle Y \rangle$ compatible with multiplication for which the ideal of relations of $A_\mathbb{F}$ corresponding to the generating set Y has finite reduced Gröbner basis $G = \{g_1, \dots, g_t\}$. If we replace the original field \mathbb{K} by \mathbb{F} , nothing really changes except for A being replaced by $A_\mathbb{F}$: (X, F) is still a presentation of $A_\mathbb{F}$ as an \mathbb{F} -algebra, $\gamma(A_\mathbb{F}) = \gamma(A)$, A_α do not change at all and (2) remains true. Thus without loss of generality, we can assume that $\mathbb{F} = \mathbb{K}$ and $A_\mathbb{F} = A$. Let $X = \{x_1, \dots, x_n\}$, $Y = \{y_1, \dots, y_m\}$ and $F = \{f_1, \dots, f_s\}$. Since both X and Y are generating sets in A , there exist $\widehat{x}_j \in \mathbb{K}\langle t_1, \dots, t_m \rangle$ and $\widehat{y}_k \in \mathbb{K}\langle s_1, \dots, s_n \rangle$ such that $x_j = \widehat{x}_j(y_1, \dots, y_m)$ in A for each $1 \leq j \leq n$ and $y_k = \widehat{y}_k(x_1, \dots, x_n)$ in A for each $1 \leq k \leq m$. It follows that

$$\begin{aligned} x_j - \widehat{x}_j(\widehat{y}_1(x_1, \dots, x_n), \dots, \widehat{y}_m(x_1, \dots, x_n)) \\ = \sum_i a_{i,j} u_{i,j} f_{r(i,j)}(x_1, \dots, x_n) v_{i,j} \text{ in } \mathbb{K}\langle X \rangle \end{aligned} \quad (2.1)$$

for $1 \leq j \leq n$, where the sum in the right-hand side is finite, $u_{i,j}, v_{i,j} \in \langle X \rangle$ and $a_{i,j} \in \mathbb{K}^*$.

Since each g_k is a relation for A with respect to the generating set Y , $g_k(y_1, \dots, y_m) = 0$ in A and therefore $g_k(\widehat{y}_1(x_1, \dots, x_n), \dots, \widehat{y}_m(x_1, \dots, x_n)) = 0$ in A . This happens precisely when

$$g_k(\widehat{y}_1(x_1, \dots, x_n), \dots, \widehat{y}_m(x_1, \dots, x_n)) = \sum_i c_{i,k} q_{i,k} f_{j(i,k)}(x_1, \dots, x_n) w_{i,k} \text{ in } \mathbb{K}\langle X \rangle, \quad (2.2)$$

for $1 \leq k \leq t$, where the sum in the right-hand side is finite, $q_{i,k}, w_{i,k} \in \langle X \rangle$ and $c_{i,k} \in \mathbb{K}^*$.

Now consider the finite set $S \subset \mathbb{K}$ comprising all (non-zero) coefficients of all g_k , all (non-zero) coefficients of all \widehat{x}_j and all \widehat{y}_k , all coefficients $a_{i,j}$ from (2.1) and all coefficients $c_{i,k}$ from (2.2). Let R be the subring of \mathbb{K} generated by $R_0 \cup S$. Clearly R is a finite ring extension of R_0 in $\mathbb{K} = \mathbb{F}$. By (2), there is $\alpha \in \Lambda$, a field \mathbb{F}_α containing \mathbb{K}_α as a subfield and a ring homomorphism $\rho'_\alpha : R \rightarrow \mathbb{F}_\alpha$ such that $\rho'_\alpha|_{R_0} = \rho_\alpha$. Let $B_\alpha = (A_\alpha)_{\mathbb{F}_\alpha}$. That is, B_α is obtained from A_α by extending the ground field from \mathbb{K}_α to \mathbb{F}_α . Hence $\gamma(B_\alpha) = \gamma(A_\alpha)$ and therefore $\gamma(B_\alpha) > \gamma(A)$ by (1). Note that as an \mathbb{F}_α -algebra, B_α is presented by (X, F_α) .

As we have done with ρ_α , we can treat ρ'_α as a homomorphism from $R\langle X \rangle$ to $\mathbb{F}_\alpha\langle X \rangle$. The way we defined R yields that both (2.1) and (2.2) consist of equalities between elements of $R\langle X \rangle$. Since $\rho'_\alpha(f_r) = \rho_\alpha(f_r)$ is a relation for B_α with respect to the generating set X , after the application of ρ'_α , the right-hand sides of both (2.1) and (2.2) vanish as elements of B_α . Consider $\rho'_\alpha(\widehat{y}_k)(x_1, \dots, x_n)$ for $1 \leq k \leq m$ as elements of B_α and denote them by y_k again. Applying ρ'_α to both sides of (2.1), we see that

$$x_j - \rho'_\alpha(\widehat{x}_j)(y_1, \dots, y_m) = 0 \text{ in } B_\alpha \text{ for } 1 \leq j \leq n.$$

It follows that $Y = \{y_1, \dots, y_m\}$ is a generating set for B_α . Now applying ρ'_α to both sides of (2.2), we see that

$$\rho'_\alpha(g_k)(y_1, \dots, y_m) = 0 \text{ in } B_\alpha \text{ for } 1 \leq k \leq t. \quad (2.3)$$

Since ρ_α is non-zero, R_0 is unital and ρ'_α extends ρ_α , we have $\rho'_\alpha(1) = 1$. Keeping $\langle Y \rangle$ equipped with the same ordering \leq for which G is the reduced Gröbner basis of the ideal of relations of A , we see that the equality $\rho'_\alpha(1) = 1$ implies that the leading monomial of each g_k is the same as the leading monomial of $\rho'_\alpha(g_k)$. By (2.3), each $\rho'_\alpha(g_k)$ is a relation for B_α with respect to the generating set Y . It follows that the normal words for A with respect to $(\langle Y \rangle, \leq)$, span B_α as an \mathbb{F}_α -vector space. Hence the Poincaré series $P_A = \sum s_n t^n$ of A and $P_{B_\alpha} = \sum r_n t^n$ of B_α with respect to the generating set Y satisfy $r_n \leq s_n$ for all $n \in \mathbb{Z}_+$. Thus $\gamma(B_\alpha) \leq \gamma(A)$, which contradicts the earlier inequality $\gamma(B_\alpha) > \gamma(A)$. \square

The only growth measuring function we shall apply Theorem 2.4 with is the Gelfand–Kirillov dimension GKdim. We gave Theorem 2.4 in its full generality on the off-chance that using it with other growth measuring functions may happen in the future. In the proof of Theorem 1.4, we are going to apply Theorem 2.4 in the particular case $R_0 = \mathbb{Z}$. In order to do this smoothly we need the following commutative lemma. We also need two well-known facts. First,

$$\text{a subgroup of a finitely generated abelian group is finitely generated.} \quad (2.4)$$

This easily follows from the fact that every finitely generated abelian group is isomorphic to a direct product of finitely many cyclic groups. Our second tool is the weak Hilbert Nullstellensatz:

$$\begin{aligned} &\text{if } \mathbb{K} \text{ is a field, } \mathbb{F} \text{ is an algebraically closed field containing } \mathbb{K} \text{ as a subfield} \\ &\text{and } I \text{ is a proper ideal in the polynomial } \mathbb{K}\text{-algebra } \mathbb{K}[x_1, \dots, x_n], \text{ then} \quad (2.5) \\ &\text{there is } (a_1, \dots, a_n) \in \mathbb{F}^n \text{ such that } f(a_1, \dots, a_n) = 0 \text{ for all } f \in I. \end{aligned}$$

Lemma 2.5. *Let R be a non-zero unital commutative finitely generated ring such that the group $(R, +)$ is torsion-free. Let E be the set of all prime numbers p such that $1 = py$ for some $y \in R$. Then E is finite.*

Proof. Let $\{x_1, \dots, x_n\}$ be a generating set for R . Then R is naturally isomorphic to $\mathbb{Z}[t_1, \dots, t_n]/I$, where I is the ideal consisting of all f such that $f(x_1, \dots, x_n) = 0$ in R . Since $(R, +)$ is torsion-free, there is a unique ideal J in the \mathbb{Q} -algebra $\mathbb{Q}[t_1, \dots, t_n]$ such that $I = \mathbb{Z}[t_1, \dots, t_n] \cap J$. Since R is unital, I is proper and therefore J is proper. Since the field \mathbb{A} of algebraic (over \mathbb{Q}) numbers is algebraically closed and contains \mathbb{Q} , (2.5) provides $\alpha_1, \dots, \alpha_n \in \mathbb{A}$ such that $f(\alpha_1, \dots, \alpha_n) = 0$ for every $f \in J$. Now we have

$$p \in E \implies \text{there exists } f \in \mathbb{Z}[t_1, \dots, t_n] \text{ such that } f(\alpha_1, \dots, \alpha_n) = \frac{1}{p}. \quad (2.6)$$

Pick $k \in \mathbb{N}$ such that $\beta_j = k\alpha_j$ is an algebraic integer for $1 \leq j \leq n$. For instance, as k we can take the product of leading coefficients of the minimal polynomials for α_j . Obviously,

$$\begin{aligned} &\text{for each } f \in \mathbb{Z}[t_1, \dots, t_n], \text{ there is } g \in \mathbb{Z}[t_1, \dots, t_n] \\ &\text{such that } f(\alpha_1, \dots, \alpha_n) = \frac{g(\beta_1, \dots, \beta_n)}{k^m}, \end{aligned} \quad (2.7)$$

where m is the degree of f . Combining (2.6) and (2.7), we get

$$p \in E \implies \begin{aligned} &\text{there exist } g \in \mathbb{Z}[t_1, \dots, t_n] \text{ and } m \in \mathbb{N} \\ &\text{such that } g(\beta_1, \dots, \beta_n) = \frac{k^m}{p}. \end{aligned} \quad (2.8)$$

Now consider the set

$$G = \{g(\beta_1, \dots, \beta_n) : g \in \mathbb{Z}[t_1, \dots, t_n]\}.$$

Obviously, G is a subgroup of $(\mathbb{A}, +)$. Since each β_j is an algebraic integer, $\beta_j^{m_j}$ is a linear combination with integer coefficients of $1, \beta_j, \dots, \beta_j^{m_j-1}$, where m_j is the degree of β_j . Hence the set G does not change if in its definition we use only $g \in \mathbb{Z}[t_1, \dots, t_n]$ such that $\deg_{t_j} g < m_j$ for $1 \leq j \leq n$. Hence the group $(G, +)$ is finitely generated. By (2.4), the group $(G \cap \mathbb{Q}, +)$ is also finitely generated. Since every finitely generated subgroup of $(\mathbb{Q}, +)$ is cyclic, $G \cap \mathbb{Q}$ as an additive group is generated by a fraction $\frac{a}{b}$ with $a \in \mathbb{Z}, b \in \mathbb{N}$. By (2.8), for every $p \in E$, there is $r_p \in \mathbb{Z}_+$ such that $\frac{k^{r_p}}{p} \in G \cap \mathbb{Q}$. On the other hand, a number of the form $\frac{k^m}{p}$ with p being a prime and $m \in \mathbb{Z}_+$ can only belong to $G \cap \mathbb{Q}$ if p divides $N = kb$. Since N has only finitely many prime factors, E is finite. \square

Corollary 2.6. *Let R be a non-zero unital commutative finitely generated ring such that the group $(R, +)$ is torsion-free. Let E be the set of all prime numbers p such that there is no non-zero ring homomorphisms from R to the algebraic closure \mathbb{F}_p of the field \mathbb{Z}_p . Then E is finite.*

Proof. Let $S = \{s_1, \dots, s_n\}$ be generators of R . Consider the ideal I in $\mathbb{Z}[t_1, \dots, t_n]$ consisting of all f such that $f(s_1, \dots, s_n) = 0$. It is clear that $p \in E$ if $\pi_p(I) = \mathbb{Z}_p[t_1, \dots, t_n]$. On the other hand, if $\pi_p(I) \neq \mathbb{Z}_p[t_1, \dots, t_n]$, (2.5) provides $\alpha_1, \dots, \alpha_n \in \mathbb{F}_p$ such that $f(\alpha_1, \dots, \alpha_n) = 0$ whenever $f \in I$. Now the map $s_j \mapsto \alpha_j$ extends to a homomorphism from R to \mathbb{F}_p yielding $p \notin E$. Thus E is exactly the set of primes p for which $\pi_p(I) = \mathbb{Z}_p[t_1, \dots, t_n]$. The latter equality is equivalent to $1 \in \pi_p(I)$, which in turn, is equivalent to $1 = py$ in R for some $y \in R$. By Lemma 2.5, E is finite. \square

Now we state and prove a particular case of Theorem 2.4 corresponding to $R_0 = \mathbb{Z}$ and the growth measuring function γ being the Gelfand–Kirillov dimension GKdim .

Theorem 2.7. *Let A be an algebra over a field \mathbb{K} of characteristic zero given by a finite presentation (X, F) such that each defining relation (=an element of F) has integer coefficients. For each prime number p , let A_p be the \mathbb{Z}_p -algebra defined by the presentation (X, F_p) , where $F_p = \pi_p(F) = \{\pi_p(f) : f \in F\}$. Assume also that $\text{GKdim}(A_p) > \text{GKdim}(A)$ for infinitely many prime numbers p . Then A is an IGB-algebra.*

Proof. We apply Theorem 2.4 with $R_0 = \mathbb{Z}$ and the homomorphisms ρ_α being $\pi_p : \mathbb{Z} \rightarrow \mathbb{Z}_p$ with the index set Λ being the infinite set of all prime numbers p for which $\text{GKdim}(A_p) > \text{GKdim}(A)$. Thus condition (1) of Theorem 2.4 with $\gamma = \text{GKdim}$ is automatically satisfied. In order to complete the proof, it remains to verify condition (2) of Theorem 2.4. Let R be a finite ring extension of \mathbb{Z} in a

field \mathbb{F} containing \mathbb{K} as a subfield. Then R is a finitely generated ring and $(R, +)$ is torsion-free. Since Λ is infinite, Corollary 2.6 provides $p \in \Lambda$ for which there exists a non-zero ring homomorphism $\varphi : R \rightarrow \mathbb{F}_p$, where \mathbb{F}_p is the algebraic closure of \mathbb{Z}_p . Since $\varphi \neq 0$, we must have $\varphi(1) = 1$, which implies that φ extends π_p . Thus condition (2) of Theorem 2.4 is satisfied as well and the result follows. \square

3. Two varieties of finitely presented algebras

In this section we compute reduced Gröbner bases of some finitely presented algebras. The algebras we deal with are degree graded. The graded case is much better than the general one in terms of computing the Gröbner basis since the elements of the reduced Gröbner basis are homogeneous and the degree n elements arise only from resolving the degree n overlaps of the leading monomials (aka obstructions, aka ambiguities) of elements computed so far. Hence the Buchberger algorithm climbs up the degrees in this case: for each given n , after finitely many steps we may be sure that no new members of the Gröbner basis of degree n will appear. In the non-graded case we do not have even this modest luxury.

Example 3.1. Let \mathbb{K} be a field of characteristic other than 2, $b \in \mathbb{K}$ is such that $b^2 \neq 1$, $b \neq 0$ and $a = \frac{b^2-1}{4}$. Consider the \mathbb{K} -algebra A given by generators x, y, z and relations xy, yz and $xx - xz - azz$. We equip monomials with the left-to-right degree-lexicographical ordering assuming $x > y > z$. If $s = \frac{1-b}{1+b}$ has infinite order in the group \mathbb{K}^* , then the leading monomials of the members of the reduced Gröbner basis of the ideal of relations of A are $yz, xz^j x$ and $xz^j y$ with $j \in \mathbb{Z}_+$. If s has finite order m in \mathbb{K}^* , then the leading monomials are $yz, xz^j x$ and $xz^j y$ with $0 \leq j \leq m-2, xz^m$ and $z^m(xz^{m-1})^j y$ for $j \in \mathbb{Z}_+$. Furthermore,

$$H_A(t) = \frac{1}{(1-t)^3}$$

if s has infinite order and

$$H_A(t) = \frac{1-t^m-t^{m+1}}{(1-t)^2(1-t-t^m)}$$

if s has order $m \in \mathbb{N}$.

Proof. Assume that $k \in \mathbb{N}$ and $s^j \neq 1$ for $1 \leq j \leq k$. A direct computation shows that the yz together with

$$p_{j+1}xz^jx - \frac{p_{j+2}}{2}xz^{j+1} + 2ap_jz^{j+1}x - ap_{j+1}z^{j+2}, \quad p_{j+1}xz^jy + 2ap_jz^{j+1}y$$

for $0 \leq j \leq k$, where $p_j = (1+b)^j - (1-b)^j$, comprise (up to normalization) all members of degree up to $k+2$ of the reduced Gröbner basis of the ideal of

relations of A . Indeed, relations in the above display for $j = 0$ are (up to a non-zero scalar multiple) the defining relations $x^2 - xz - az^2$ and xy and we proceed inductively. The degree $j + 2$ overlaps $xz^{j-1}x^2 = (xz^{j-1}x)x = xz^{j-1}(xx)$ and $xz^{j-1}xy = (xz^{j-1}x)y = xz^{j-1}(xy)$ produce (after reduction) the two relations in the above display, while all other overlaps of degree $j + 2$ resolve.

If s has infinite order, this works for every k and we are furnished with the complete reduced Gröbner basis. The leading monomials of the members of the basis are yz , $xz^m x$ and $xz^m y$ for $m \in \mathbb{Z}_+$. The corresponding normal words are $z^j y^m$ and $z^j y^m xz^n$ with $j, m, n \in \mathbb{Z}_+$. Counting the number of normal words of any given degree, we arrive to

$$H_A(t) = \frac{1}{(1-t)^3}.$$

Now assume that the order of s in \mathbb{K}^* is $k + 1$ with $k \in \mathbb{N}$ (note that $s \neq 1$ since $b \neq 0$). The above display together with yz still provides all members of degree up to $k + 2$ of the reduced Gröbner basis of the ideal of relations of A . However, the degree $k + 2$ elements are now up to a scalar multiple $z^{k+1}y$ and $xz^{k+1} - z^{k+1}x$. From this point on the pattern of the basis changes dramatically. The overlaps $xz^{2k+1}(xz^k)^j y = [xz^{k+1}]z^k(xz^k)^j y = xz^k[z^{k+1}(xz^k)^j y]$ yield the monomial relation $z^{k+1}(xz^k)^{j+1}y$ for every $j \in \mathbb{Z}_+$ while all other overlaps of degrees $\geq k + 3$ resolve. Thus the leading monomials of the members of the reduced Gröbner basis of the ideal of relations of A are yz , $xz^j x$ and $xz^j y$ with $0 \leq j \leq m - 2$, xz^m and $z^m(xz^{m-1})^j y$ for $j \in \mathbb{Z}_+$, where $m = k + 1$ is the order of s . The corresponding normal words are

$$\begin{aligned} z^j & \text{ with } j \in \mathbb{Z}_+, \\ z^j y^r & \text{ with } j, r \in \mathbb{Z}_+, j < m, \\ z^r (xz^{m-1})^j xz^t & \text{ with } r, j, t \in \mathbb{Z}_+, t < m, r \geq m, \\ z^r y^{m_0} xz^{m-1} y^{m_1} xz^{m-1} y^{m_2} x \dots xz^{m-1} y^{m_j}, \\ z^r y^{m_0} xz^{m-1} y^{m_1} xz^{m-1} y^{m_2} x \dots xz^{m-1} y^{m_j} xz^t & \text{ with } j, r, t, m_0, \dots, m_j \in \mathbb{Z}_+, \\ & r, t < m. \end{aligned}$$

Counting the number of normal words of given degree, we arrive to

$$H_A = \frac{1 - t^m - t^{m+1}}{(1-t)^2(1-t-t^m)}. \quad \square$$

Example 3.2. Let \mathbb{K} be an arbitrary field and $a \in \mathbb{K}^*$. Consider the \mathbb{K} -algebra B given by generators x, y and relations y^3 and $x^2y - ayx^2 - yxy$. We equip monomials with the left-to-right degree-lexicographical ordering assuming $x > y$. If $1 + a + \dots + a^k \neq 0$ for all $k \in \mathbb{N}$, then the leading monomials of the members of the

reduced Gröbner basis of the ideal of relations of B are y^3 , x^2y and $y^2(xy)^jxy^2$ with $j \in \mathbb{Z}_+$. If $k \in \mathbb{N}$ is the minimal positive integer for which $1 + a + \dots + a^k = 0$, then the leading monomials are y^3 , x^2y and $y^2(xy)^jxy^2$ with $0 \leq j < k$ (finite Gröbner basis in this case). In the first case

$$H_B(t) = \frac{1}{(1+t)(1-t)^3},$$

while in the second case

$$H_B(t) = \frac{1 - t^{2k+3}}{(1-t)^2(1-t^2-t^{2k+3})}.$$

Proof. Whatever the case, a direct computation shows that apart for defining relations the only other degree ≤ 5 member of the reduced Gröbner basis is the monomial y^2xy^2 , which arises from the overlap $x^2y^3 = (x^2y)y^2 = x^2(y^3)$. If $1 + a = 0$, the defining relations together with y^2xy^2 form the reduced Gröbner basis. If $1 + a \neq 0$, apart from the above three members, the only degree ≤ 7 member of the reduced Gröbner basis is the monomial y^2xyxy^2 , which arises from the overlap $x^2y^2xy^2 = (x^2y)yxy^2 = x^2(y^2xy^2)$. If $1 + a + a^2 = 0$, the process ends (we have the complete basis). Otherwise, apart from the above four members, the only degree ≤ 9 member of the reduced Gröbner basis is the monomial $y^2xyxyxy^2$, which arises from the overlap $x^2y^2xyxy^2 = (x^2y)yxyxy^2 = x^2(y^2xyxy^2)$. This pattern goes on indefinitely (one can give an easy inductive argument as well). This takes care of the Gröbner bases in all cases and of their leading monomials.

Now if $1 + a + \dots + a^k \neq 0$ for all $k \in \mathbb{N}$, then as we have just seen, the leading monomials of the members of the reduced Gröbner basis of the ideal of relations of B are y^3 , x^2y and $y^2(xy)^jxy^2$ with $j \in \mathbb{Z}_+$. Hence the normal words of B are $y^\varepsilon(xy)^mxy^2(xy)^jx^s$ with $\varepsilon \in \{0, 1\}$, $m, j, s \in \mathbb{Z}_+$ and $y^{\text{GKdim}}(xy)^jx^s$ with $\text{GKdim} \in \{0, 1, 2\}$ and $j, s \in \mathbb{Z}_+$. Counting normal words of given degree yields

$$H_B(t) = \frac{1}{(1+t)(1-t)^3}.$$

If $k \in \mathbb{N}$ is the minimal positive integer for which $1 + a + \dots + a^k = 0$, then as we have already seen, the leading monomials of the members of the reduced Gröbner basis of the ideal of relations of B are y^3 , x^2y and $y^2(xy)^jxy^2$ with $0 \leq j < k$. Now we have a finite Gröbner basis and applying standard techniques [14], one gets

$$H_B(t) = \frac{1 - t^{2k+3}}{(1-t)^2(1-t^2-t^{2k+3})}. \quad \square$$

Now we plug $a = 2$ into Example 3.1.


Lemma 3.3. *Let \mathbb{K} be a field and A be the quadratic \mathbb{K} -algebra given by generators x, y, z and relations xy, yz and $x^2 - xz - 2z^2$. If $\text{char } \mathbb{K} = 0$, then A is an automaton*

algebra and satisfies $\text{GKdim}(A) = 3$. If $\text{char } \mathbb{K}$ is a prime $p \geq 5$, then A has exponential growth and therefore $\text{GKdim}(A) = \infty$.

Proof. We equip monomials with the left-to-right degree-lexicographical ordering assuming $x > y > z$. If $\text{char } \mathbb{K} \notin \{2, 3\}$, A is the algebra from Example 3.1 with $a = 2$, $b = -3$ and $s = -2$. If $\text{char } \mathbb{K} = 0$, s has infinite order in \mathbb{K}^* . By Example 3.1,

$$H_A(t) = \frac{1}{(1-t)^3}$$

(hence $\text{GKdim}(A) = 3$) and the leading monomials of the members of the reduced Gröbner basis of the ideal of relations of A are yz , xz^jx and xz^jy with $j \in \mathbb{Z}_+$. They are easily seen to form a regular language. Indeed, the corresponding graph is



Hence A is an automaton algebra and $\text{GKdim}(A) = 3$ provided $\text{char } \mathbb{K} = 0$. If $\text{char } \mathbb{K} = p \geq 5$, then by Example 3.1,

$$H_A(t) = \frac{1 - t^m - t^{m+1}}{(1-t)^2(1-t-t^m)},$$

where $m = m(p)$ is the order of -2 in \mathbb{Z}_p^* if $p \geq 5$. The latter shows that A has exponential growth. Indeed, the Poincaré series of A is

$$P_A(t) = \frac{1 - t^m - t^{m+1}}{(1-t)^3(1-t-t^m)} = \sum r(n)t^n$$

and an easy calculus exercise shows that

$$\lim_{n \rightarrow \infty} \frac{\ln r(n)}{n} = \ln \frac{1}{c_m} > 0,$$

where c_m is the only zero of $1 - t - t^m$ in the interval $(0, 1)$. □


Next we plug in $a = 1$ into Example 3.2.

Lemma 3.4. *Let \mathbb{K} be a field and B be the cubic \mathbb{K} -algebra given by generators x, y and relations y^3 and $x^2y - yx^2 - yxy$. If $\text{char } \mathbb{K} = 0$, then B is an automaton algebra and satisfies $\text{GKdim}(B) = 3$. If $\text{char } \mathbb{K} \neq 0$, then B has exponential growth and therefore $\text{GKdim}(B) = \infty$.*

Proof. We equip monomials with the left-to-right degree-lexicographical ordering assuming $x > y$. If $\text{char } \mathbb{K} = 0$, Example 3.2 yields that

$$H_B(t) = \frac{1}{(1+t)(1-t)^3}$$

(hence $\text{GKdim}(B) = 3$) and that the leading monomials of the members of the reduced Gröbner basis of the ideal of relations of B are y^3 , x^2y and $y^2(xy)^jxy^2$ with $j \in \mathbb{Z}_+$. They are easily seen to form a regular language. Indeed, the corresponding graph is



Hence B is an automaton algebra and $\text{GKdim}(B) = 3$ provided $\text{char } \mathbb{K} = 0$. If $\text{char } \mathbb{K} = p$ with p being a prime number, then k in Example 3.1 equals $p - 1$ and we have

$$H_B(t) = \frac{1 - t^{2p+2}}{(1-t)^2(1-t^2-t^{2p+2})}.$$

This shows that B has exponential growth. Indeed, the Poincaré series of B is

$$P_B(t) = \frac{1 - t^{2p+2}}{(1-t)^3(1-t^2-t^{2p+2})} = \sum r(n)t^n$$

and an easy calculus exercise shows that

$$\lim_{n \rightarrow \infty} \frac{\ln r(n)}{n} = \ln \frac{1}{c_p} > 0,$$

where c_p is the only zero of the polynomial $1 - t^2 - t^{2p+2}$ in the interval $(0, 1)$. \square

3.1. Proof Theorem 1.4. The combination of Lemmas 3.3, 3.4 and Theorem 2.7 yield Theorem 1.4. Indeed, by Lemmas 3.3 and 3.4, A and B from Theorem 1.4 are automaton algebras. The same lemmas also imply that $\text{GKdim}(A) = \text{GKdim}(B) = 3$, while $\text{GKdim}(A_p) = \text{GKdim}(B_p) = \infty$ for every prime number $p \geq 5$. By Theorem 2.7 then A and B are IGB-algebras, which completes the proof of Theorem 1.4.

4. Proof of Theorem 1.7

Let \mathbb{K} be an arbitrary field and C be the \mathbb{K} -algebra given by four generators x, y, z, u and seven quadratic relations $xu - yz, yu - zx, zu - uz, yy, yx, xy$ and xx . We have

to show that C has intermediate growth. We equip the monomials in x, y, z, u with the left-to-right degree-lexicographical ordering assuming $x > y > z > u$. First, we find the reduced Gröbner basis in the ideal of relations of C . Degree 3 overlaps $xyu = (xy)u = x(yu)$ and $yyu = (yy)u = y(yu)$ produce two monomial relations yzx and xzx , while all other degree 3 overlaps of the leading monomials of defining relations resolve. Thus we have exactly two degree three members of the reduced Gröbner basis: yzx and xzx . Degree 4 overlaps $yzxu = (yzx)u = yz(xu)$ and $xzxu = (xzx)u = xz(xu)$ produce two monomial relations $yzyz$ and $xzxz$, while all other degree 4 overlaps resolve. Thus we have exactly two degree 4 members of the reduced Gröbner basis: $yzyz$ and $xzxz$. Degree 5 overlaps $yzyzu = (yzyz)u = yzy(zu)$ and $xzxcz = (xzcxz)u = xzx(zu)$ produce (after reduction) two monomial relations yz^2xz and xz^2xz , while all other degree 5 overlaps resolve. Thus we have exactly two degree 5 members of the reduced Gröbner basis: yz^2xz and xz^2xz . This pattern is easily seen to go on:

the reduced Gröbner basis in the ideal of relations of C consists of
 $xu - yz, yu - zx, zu - uz, yx, xx, yz^{k+1}xz^k, xz^{k+1}xz^k, yz^k yz^k$ (4.1)
 and $xz^k yz^k$ for $k \in \mathbb{Z}_+$.

Indeed, one can use the following inductive argument. If $n \geq 5$ is even, we have $n = 2k + 2$ with $k > 1$. The degree $n + 1$ overlaps $yz^k yz^k u = (yz^k yz^k)u = yz^k yz^{k-1}(zu)$ and $xz^k yz^k u = (xz^k yz^k)u = xz^k yz^{k-1}(zu)$ produce (after reduction) $yz^{k+1}xz^k$ and $xz^{k+1}xz^k$. All other overlaps of degree $n + 1$ resolve. It is especially easy to see since an overlap of two monomial relations always resolves. If $n \geq 5$ is odd, we have $n = 2k + 1$ with $k > 1$. The degree $n + 1$ overlaps $yz^k xz^{k-1}u = (yz^k xz^{k-1})u = yz^k xz^{k-2}(zu)$ and $xz^k xz^{k-1}u = (xz^k xz^{k-1})u = xz^k xz^{k-2}(zu)$ produce (after reduction) $yz^k yz^k$ and $xz^k yz^k$. All other overlaps of degree $n + 1$ resolve. This argument verifies (4.1), from which we immediately have

the leading monomials of the reduced Gröbner basis in the ideal
 of relations of C are $xu, yu, zu, yx, xx, yz^{k+1}xz^k, xz^{k+1}xz^k,$ (4.2)
 $yz^k yz^k$ and $xz^k yz^k$ for $k \in \mathbb{Z}_+$.

Now we can easily describe the corresponding normal words:

the normal words for C are $u^j z^k$ and $u^j z^k s_0 z^{k_1} s_1 z^{k_2} s_2 \dots z^{k_m} s_m z^{k_{m+1}},$
 where $s_r \in \{x, y\}, j, k, m, k_0, \dots, k_{m+1} \in \mathbb{Z}_+, k_1 > k_2 > \dots > k_{m+1}$ (4.3)
 and $k_r > k_{r+1} + 1$ whenever $1 \leq r \leq m$ and $s_r = x$.

Let $p(n)$ be the number of normal words for C of degree at most n . That is, the Poincaré series of C is

$$P_C(t) = \sum p(n)t^n.$$

According to (1.4), in order to show that C has intermediate growth, it suffices to prove that

$$\lim_{n \rightarrow \infty} \frac{\ln \ln p(n)}{\ln n} = \frac{1}{2}. \quad (4.4)$$

Let $n \in \mathbb{N}$ and m be the largest integer not exceeding \sqrt{n} . By (4.3), the words

$$s_1 z^{2m-2} s_2 z^{2m-4} s_3 z^{2m-6} \dots z^4 s_{m-1} z^2 s_m \text{ with } s_j \in \{x, y\}$$

are normal words for C of degree $\leq n$. Since we listed 2^m words, we have $p(n) \geq 2^m$. Since $\sqrt{n} - 1 < m \leq \sqrt{n}$, it follows that

$$\liminf_{n \rightarrow \infty} \frac{\ln p(n)}{\sqrt{n}} \geq \ln 2 \implies \liminf_{n \rightarrow \infty} \frac{\ln \ln p(n)}{\ln n} \geq \frac{1}{2}. \quad (4.5)$$

Next, by (4.3), the number $a(n)$ of normal words of C of degree n increases as a function of n . Since $p(n) = a(0) + \dots + a(n)$, we have $p(n) \leq (n+1)a(n)$. If $b(n)$ is the number of normal words for C of degree n starting with x or y (we include the empty word 1 as well), (4.3) implies that

$$a(n) = (n+1)b(0) + nb(1) + \dots + 2b(n-1) + b(n).$$

Since $b(n)$ increases as well, $a(n) \leq \frac{n(n+1)}{2}b(n) \leq (n+1)^2b(n)$. Hence

$$p(n) \leq (n+1)a(n) \leq (n+1)^3b(n).$$

By (4.3), $b(n)$ does not exceed the number $\varphi(n)$ of degree n words of the form $s_1 z^{k_1} s_2 z^{k_2} s_3 \dots z^{k_{m-1}} s_m z^{k_m}$, where $s_j \in \{x, y\}$, $m \in \mathbb{N}$, $k_1, \dots, k_m \in \mathbb{Z}_+$ and $k_1 > k_2 > \dots > k_m$. Now one can easily see that the numbers $\varphi(n)$ have the following generating function:

$$\sum_{n=0}^{\infty} \varphi(n) t^n = \prod_{n=1}^{\infty} (1 + 2t^n). \quad (4.6)$$

Note that the infinite product in the right-hand side of (4.6) converges uniformly on compact subsets of the open unit disk \mathbb{D} of the complex plane \mathbb{C} . Thus we can treat it as a holomorphic function Φ on \mathbb{D} . By the Cauchy formula

$$\varphi(n) = \frac{1}{2\pi i} \oint_{\Gamma} \frac{\Phi(z) dz}{z^{n+1}}, \quad (4.7)$$

where Γ is any closed contour inside \mathbb{D} which encircles 0 once counterclockwise. Now we specify Γ to be the circle Γ_n centered at zero of radius $r_n = 1 - \frac{1}{\sqrt{n}}$ (assume $n \geq 2$). First, observe that

$$|\Phi(z)| \leq \prod_{k=1}^{\infty} (1 + 2r_n^k)$$

for $z \in \Gamma_n$. Hence,

$$\ln |\Phi(z)| \leq \sum_{k=1}^{\infty} \ln(1 + 2r_n^k) \leq 2 \sum_{k=1}^{\infty} r_n^k = \frac{2r_n}{1 - r_n} \leq 2\sqrt{n} \text{ for } z \in \Gamma_n.$$

Hence

$$|\Phi(z)| \leq e^{2\sqrt{n}} \text{ for } z \in \Gamma_n.$$

Using the L'Hopital rule or the Taylor expansion of $\ln(1 + t)$, one easily shows that

$$\lim_{n \rightarrow \infty} \frac{r_n^{-n}}{e^{\sqrt{n}}} = \sqrt{e}.$$

Using the above two displays and (4.7) with $\Gamma = \Gamma_n$ and estimating the integral from above by the product of the length of the path by the maximum of the absolute value of the function under the integral, we see that there exists a constant $a > 0$ such that

$$\varphi(n) = |\varphi(n)| \leq r_n^{-n} \max\{|\Phi(z)| : z \in \Gamma_n\} \leq ae^{3\sqrt{n}} \text{ for all } n.$$

Since $p(n) \leq (n + 1)a(n) \leq (n + 1)^3b(n) \leq (n + 1)^3\varphi(n) \leq a(n + 1)^3e^{3\sqrt{n}}$, we have

$$\limsup_{n \rightarrow \infty} \frac{\ln p(n)}{\sqrt{n}} \leq e^3 \implies \limsup_{n \rightarrow \infty} \frac{\ln \ln p(n)}{\ln n} \leq \frac{1}{2}. \tag{4.8}$$

The inequalities (4.5) and (4.8) imply (4.4), which completes the proof of Theorem 1.7.

5. Concluding remarks

Remark 5.1. One could apply similar arguments starting with other than in Examples 3.1 or 3.2 varieties of finitely presented algebras. What one needs is many (impossible to cover by a single non-trivial Zariski closed set) algebras with exceptionally fast growth rate compared to that of generic members of the variety. We went for the ones we considered to be the simplest among other suitable varieties.

Remark 5.2. Since the algebra A in Theorem 1.4 is quadratic, one may ask whether it is Koszul and whether Koszulity has any bearing on the question we considered. We refer to the book [11] for the definition and the properties of Koszul algebras. Well, A is Koszul. This can be verified directly by computing the Koszul complex of A and showing that it is exact. Thus Koszulity does not save the day: a Koszul automaton algebra can still be an IGB-algebra.

Remark 5.3. It so happens that all examples of automaton IGB algebras we were able to produce (including those we did not mention in this text) have non-trivial zero divisors. We have no idea of why it happens this way. This prompts us to ask the following question. Unfortunately, we have no arguments to back either positive or negative answer.

Question 5.4. *Does there exist an automaton IGB-algebra A , which is also a domain (= has no non-trivial zero divisors)?*

Remark 5.5. Another reason we include Theorem 1.7 is that, as it is apparent from its proof, the ideal of relations of the algebra C with respect to the original generators and the left-to-right degree-lexicographical ordering assuming $x > y > z > u$ is very nice (although infinite) in a sense that it follows a clear pattern and can be written in full. On the other hand, the algebra C is not an automaton algebra because it has intermediate growth, see [14].

Remark 5.6. More subtle considerations than in the proof of Theorem 1.7 allow to show that the sequence $\frac{\ln p(n)}{\sqrt{n}}$ converges to a positive limit. What we actually did in the proof is to demonstrate that its lower limit is at least $\ln 2$ and its upper limit is at most e^3 .

Remark 5.7. As we have mentioned in the introduction, a quadratic algebra with either at most two generators or at most two relations is an FGB-algebra and therefore can not have intermediate growth. Thus a quadratic algebra of intermediate growth has at least 3 generators and at least 3 linearly independent quadratic relations. Our example in Theorem 1.7 has 4 generators and 7 quadratic relations. So, potentially, there is still room for improvement. In particular, the following question is open.

Question 5.8. *Does there exist a quadratic algebra A of intermediate growth presented by three generators and some homogeneous degree 2 relations?*

We do have an argument showing that if such an algebra A exists, its space of quadratic relations must have dimension either 4 or 5. That is, A must be presented by 3 generators and either 4 or 5 linearly independent quadratic relations.

Acknowledgements. We are grateful to IHES and MPIM for hospitality, support, and excellent research atmosphere. This work is funded by the ERC grant 320974, and partially supported by the project PUT9038. We would also like to express our gratitude to the anonymous referee for numerous comments, which helped to improve the presentation of the paper.

References

- [1] A. Belov, V. Borisenko, and V. Latyshev, Monomial algebras, Algebra 4, *Modern Math. and Appl.*, VINITI, **26** (2002), 35–2014; *J. Math. Sci. (New York)*, **87** (1997), 3463–3575. [Zbl 0927.16018](#) [MR 1604202](#)
- [2] G. Bergman, The diamond lemma for ring theory, *Adv. Math.*, **29** (1978), 178–218. [Zbl 0326.16019](#) [MR 506890](#)
- [3] F. Cedó and J. Okniński, Gröbner bases for quadratic algebras of skew type, *Proc. Edinburgh Math. Soc.*, **55** (2012), 387–401. [Zbl 1348.20062](#) [MR 2928501](#)

- [4] V. Govorov, Graded algebras, *Math. Notes*, **12** (1972), 552–556. [Zbl 0239.16002](#) [MR 318199](#)
- [5] A. Kirillov, M. Kontsevich, and A. Molev, Algebras of intermediate growth, *Selecta Mathematica Sovietica*, **9** (1990), 137–153. [Zbl 0705.17013](#) [MR 753873](#)
- [6] Y. Kobayashi, A finitely presented monoid which has solvable word problem but has no regular complete presentation, *Theoret. Comput. Sci.*, **146** (1995), 321–329. [Zbl 0873.68105](#) [MR 1343236](#)
- [7] D. Koçak, Finitely presented quadratic algebras of intermediate growth, *Algebra Discrete Math.*, **20** (2015), 69–88. [Zbl 1350.16017](#) [MR 3431952](#)
- [8] M. Lothaire, *Combinatorics on Words*, Encyclopedia of Mathematics and its Applications, 17, Addison-Wesley, Reading, MA, 1983. [Zbl 0514.20045](#) [MR 675953](#)
- [9] J. Månsson and P. Nordbeck, Regular Gröbner bases, *J. Symb. Comp.*, **33** (2002), 163–181. [Zbl 1011.16017](#) [MR 1879379](#)
- [10] J. Månsson and P. Nordbeck, A generalized Ufnarovski graph, *Appl. Algebra Engrg. Comm. Comput.*, **16** (2005), 293–306. [Zbl 1092.68058](#) [MR 2233759](#)
- [11] A. Polishchuk and L. Positselski, *Quadratic algebras*, University Lecture Series, 37, American Mathematical Society, Providence, RI, 2005. [Zbl 1145.16009](#) [MR 2177131](#)
- [12] M. Sapir, *Combinatorial algebra: syntax and semantics*, Springer, 2014. [Zbl 1319.05001](#) [MR 3243545](#)
- [13] J. Shearer, A graded algebra with a non-rational Hilbert series, *J. Algebra*, **62** (1980), 228–231. [Zbl 0436.16001](#) [MR 561125](#)
- [14] V. Ufnarovski, Combinatorial and asymptotic methods in algebra, in *Algebra*, VI, 1–196, Enciclopedia Math. Sci., 57, Springer, Berlin, 1995. [Zbl 0826.16001](#) [MR 1360005](#)
- [15] V. Ufnarovski, On the use of graphs for calculating the basis, growth and Hilbert series of associative algebras, *Math. Sb.*, **180** (1989), 1548–1560. [Zbl 0685.16007](#) [MR 1034428](#)
- [16] V. Ufnarovski, Poincaré series of graded algebras, *Math. Notes*, **27** (1980), 12–18. [Zbl 0446.16034](#) [MR 562474](#)

Received 11 June, 2017

N. Iyudu, School of Mathematics, The University of Edinburgh,
James Clerk Maxwell Building, The King’s Buildings,
Peter Guthrie Tait Road, Edinburgh EH9 3FD, UK
E-mail: niyudu@exseed.ed.ac.uk

S. Shkarin, Department of Pure Mathematics, Queen’s University Belfast,
University Road, Belfast BT7 1NN, UK
E-mail: s.shkarin@qub.ac.uk