

Iterated and mixed discriminants

Alicia Dickenstein, Sandra Di Rocco, and Ralph Morrison

Abstract. Classical work by Salmon and Bromwich classified singular intersections of two quadric surfaces. The basic idea of these results was already pursued by Cayley in connection with tangent intersections of conics in the plane and used by Schäfli for the study of hyperdeterminants. More recently, the problem has been revisited with similar tools in the context of geometric modeling and a generalization to the case of two higher dimensional quadric hypersurfaces was given by Ottaviani. We propose and study a generalization of this question for systems of Laurent polynomials with support on a fixed point configuration.

In the non-defective case, the closure of the locus of coefficients giving a non-degenerate multiple root of the system is defined by a polynomial called the *mixed discriminant*. We define a related polynomial called the multivariate *iterated discriminant*, generalizing the classical Schäfli method for hyperdeterminants. This iterated discriminant is easier to compute and we prove that it is always divisible by the mixed discriminant. We show that tangent intersections can be computed via iteration if and only if the singular locus of a corresponding dual variety has sufficiently high codimension. We also study when point configurations corresponding to Segre–Veronese varieties and to the lattice points of planar smooth polygons, have their iterated discriminant equal to their mixed discriminant.

1. Introduction

Let K be an algebraically closed field of characteristic zero and $A \subset \mathbb{Z}^n$ a finite lattice subset. A (Laurent) polynomial $p = \sum_{a \in A} c_a x^a \in K[x_1, \dots, x_n]$ with support on the point configuration A is called an *A-polynomial*.

Classical work by Salmon [25] and Bromwich [1] classified singular intersections of two quadric surfaces, corresponding to the case of two A -polynomials where A consists of the lattice points in the dilated simplex $2\Delta_3$ in \mathbb{R}^3 . The basic idea of these results was already pursued by Cayley in connection with tangent intersections of conics in \mathbb{C}^2 . More recently, the problem has been revisited with similar tools in [15], in the context of geometric modeling with focus on the real case; and in [29], where these techniques are used to classify singular Darboux cyclides. These are surfaces in




Figure 1. Transverse (left) and non-transverse (right) hyperbolas.

3-space that are the projection of the intersection of two quadrics in dimension four. A generalization to the case of two higher dimensional quadric hypersurfaces is given in [24].

Consider two space quadrics, given in matrix form by

$$p_i = [1 \quad x_1 \quad x_2 \quad x_3] M_i \begin{bmatrix} 1 \\ x_1 \\ x_2 \\ x_3 \end{bmatrix}, \quad i = 0, 1. \quad (1)$$

For generic matrices $M_i \in K^{4 \times 4}$, the intersection ($p_1 = p_2 = 0$) describes a non-singular curve of degree 4. The non-generic intersections are described in [16, 26] in the following way. Consider the pencil of quadrics given by $p_0 + tp_1$. Using the Schäftli decomposition method, the existence of a tangential intersection can be studied by considering the zero locus of the following polynomial in the entries of M_0, M_1 :

$$D_{4, \Delta_1}(\det(M_0 + tM_1)), \quad (2)$$

where D_{4, Δ_1} is the univariate discriminant of the degree 4 polynomial $\det(M_0 + tM_1)$, considered as a polynomial in t . For generic matrices this is a polynomial of degree 6 in its entries (that is, in the coefficients of p_0, p_1), and it vanishes whenever the polynomial $\det(M_0 + tM_1)$ does not have four simple roots. To classify the different singular intersections, they then studied the Segre characteristics arising from the Jordan normal form of the matrix $M_0 + tM_1$.

In this paper, we propose and study a generalization of this approach for any support A .

We consider equation (2) to be an *iterated process*, as we are computing the discriminant of a discriminant. Factorizations of iterated discriminants and resultants for

polynomials of three variables where studied in [4]. Our aim is to define and study an *iterated discriminant* generalizing Schäfli's method for hyperdeterminants, and to show when tangent intersections can be computed via iteration.

The theory of A -discriminants was introduced in [16] and has been extensively studied both from a geometric and a computational viewpoint [10, 12, 13, 18]. Denote by $X_A \subset \mathbb{P}^{|A|-1}$ the projective variety defined as the closed image of the monomial embedding given by the A -monomials. The dual variety $X_A^\vee \subset \mathbb{P}^{|A|-1^\vee}$ is the closure of the coefficient vectors of the A -polynomials p whose zero-locus ($p = 0$) has a singular point $x \in (K^*)^n$ with non-zero coordinates. Equivalently, the dual variety is the closure of the hyperplane sections of X_A which are singular at a point with non-zero coordinates. The expected codimension of X_A^\vee is one and when this is the case we say that A is *non-defective*. When A is non-defective the irreducible polynomial $D_A \in \mathbb{Z}[(c_a)_{a \in A}]$ defining (up to sign) the dual variety: $X_A^\vee = (D_A = 0)$, is called the *A -discriminant* [16]. We will use the notation $D_A((c_a)_{a \in A}) = D_A(p)$.

If $n = 1$ and $A = \{0, 1, 2\}$, then $p = c_2x^2 + c_1x + c_0$ and $D_A(p) = c_1^2 - 4c_0c_2$ is the classical discriminant of a degree two polynomial. More generally, $D_{\{0,1,\dots,\delta\}}$ coincides with the classical discriminant of univariate polynomials of degree δ . This $D_{0,1,\dots,\delta}$ is a polynomial of degree $2(\delta - 1)$ in the coefficients, which we denote by $D_{\delta\Delta_1}$. The case of multi-linear polynomials (i.e., tensors) corresponds to the case in which the convex hull of A equals the product $\Delta_{n_1} \times \dots \times \Delta_{n_l}$ where Δ_s denotes the unit simplex of dimension s . This multivariate A -discriminant is also referred to as the hyperdeterminant of size $(n_1 + 1) \times \dots \times (n_l + 1)$ (see [16, Chapter 14]). This is a classical object defined originally by Cayley [7].

Note that for a quadratic polynomial p with associated matrix M as in (1), that is for A consisting of the lattice points in $2\Delta_3$, the existence of a singular point in ($p = 0$) implies that the linear forms given by its partial derivatives vanish and so $\det(M) = 0$. Indeed, $D_A(p) = \det(M)$ (up to an integer factor). This suggests that an iterated discriminant should be connected to the notion of discriminant for a system of polynomials. This notion is called the *mixed discriminant* [6, 9, 16], which is a natural generalization of the classical A -discriminant.

Given $r + 1$ finite configurations $A_0, \dots, A_r \subset \mathbb{Z}^n$, and a system of A_i -polynomials p_0, \dots, p_r ,

$$p_0 = p_1 = \dots = p_r = 0, \quad p_i = \sum_{a \in A_i} c_{i,a}x^a, \tag{3}$$

we call an isolated solution $x \in (K^*)^n$ a *non-degenerate multiple root* for the system (3) if the $r + 1$ gradient vectors $\nabla_x p_i(x)$, $i = 0, \dots, r$ are linearly dependent but any subset of r of them is linearly independent. The associated *mixed discriminantal variety* is the closure of the locus of coefficients for which the system has a non-degenerate multiple root. If this variety is a hypersurface, it is defined by a single

irreducible polynomial which we call the mixed discriminant, denoted MD_{A_0, \dots, A_r} . If it is not a hypersurface, we call the system *defective* and set $MD_{A_0, \dots, A_r} = 1$.

Observe that when $r = 0$, $MD_{0, A_0} = D_{A_0}$ equals the A_0 discriminant. In fact, in the non-defective case, mixed discriminants are special cases of discriminants of a single polynomial. This was settled in [16], but without the hypothesis of non-degeneracy of the common multiple root and in [6] for the case $r + 1 = n$. Given A_0, \dots, A_r , the associated Cayley configuration $C = C(A_0, \dots, A_r) \subset \mathbb{Z}^{n+r}$ is the union of the lifted configurations $e_i \times A_i \in \mathbb{Z}^{n+r}$ for $i = 0, \dots, r$, where $e_0 = 0$ and e_i is the standard i th basis vector in \mathbb{Z}^r for $i \geq 1$. As sparse discriminants are affine invariants of lattice configurations [16], we could equivalently consider $C \subset \mathbb{Z}^{n+r+1}$, where now e_0, \dots, e_r denote the canonical basis in \mathbb{Z}^{r+1} . We introduce $r + 1$ new variables $\lambda_0, \dots, \lambda_r$ and encode the initial system by one auxiliary C -polynomial:

$$P_\lambda = \lambda_0 p_0 + \dots + \lambda_r p_r \in K[\lambda_0, \dots, \lambda_r, x_1, \dots, x_n].$$

We will denote both this polynomial and its tuple of coefficients by P_λ , where $\lambda = (\lambda_0, \dots, \lambda_r)$. In Proposition 3.3 we prove that when C is non-defective,

$$MD(A_0, \dots, A_r)(p_0, \dots, p_r)$$

can be computed as $D_C(P_\lambda)$ for any r .

This characterization leads to the following definition of *multivariate iterated discriminant* of order r . In the present paper we consider the case when $A_0 = \dots = A_r = A$ and use the notation $MD_{r,A} := MD_{A, \dots, A}$. Notice that $D_A(P_\lambda)$ is a homogeneous polynomial of degree $\deg(D_A)$ in $\lambda_0, \dots, \lambda_r$.

Definition 1.1. Given $A \subset \mathbb{Z}^n$ non-defective, denote by d the codimension of the singular locus of the dual variety X_A^v . Given $r \geq 0$, the *multivariate iterated discriminant of order r* is the polynomial $ID_{r,A}$ on the coefficients of $(r + 1)$ A -polynomials p_0, \dots, p_r defined by

$$\begin{cases} ID_{r,A}(p_0, \dots, p_r) := D_{\delta_A \Delta_r}(D_A(P_\lambda)), & \text{if } d \geq r, \\ ID_{r,A}(p_0, \dots, p_r) := 0, & \text{otherwise.} \end{cases}$$

It is worth noting that in the classical case of $r = 0$, all these polynomials coincide by definition:

$$MD_{0,A} = ID_{0,A} = D_A \quad \text{and} \quad D_{\delta_{\Delta_0}}(D_A(\lambda p_A)) = D_A.$$

The latter equality is a consequence of the fact that the discriminant (in the variable λ) of the monomial $D_A \lambda^\delta$ is the coefficient D_A (see [19]). Moreover, when A consists of the vertices of a simplex, $ID_{r,A}$ coincides with the hyperdeterminant Schafli decomposition [16, Chapter 14].

Our main results give a precise relation between $MD_{r,A}$ and $ID_{r,A}$. The advantage of relating $MD_{r,A}$ with $ID_{r,A}$ is that the latter polynomial is much easier to compute. We show that in the non-defective case $MD_{r,A}$ is always an irreducible factor of $ID_{r,A}$, as a consequence of biduality (see Section 4). Therefore, if $ID_{r,A}(p_0, \dots, p_r) \neq 0$, we get a certificate that the intersection $(p_0 = \dots = p_r)$ is smooth. When A is non-defective, we denote by $\text{sing}(X_A^\vee)$ the subscheme of X_A^\vee defined by the ideal generated by the partial derivatives of D_A . We show that $ID_{r,A}$ can have other irreducible factors given by the Chow forms Ch_{Y_k} of the higher dimensional irreducible components of the schematic singular locus of the dual variety X_A^\vee . We recall the notion of Chow forms at the beginning of Section 4. Theorem 4.4 and Proposition 4.5 imply the following theorem.

Theorem. *Assume $A \subset \mathbb{Z}^n$ is non-defective and let $r \in \mathbb{Z}$ with $0 \leq r \leq \dim(X_A)$. Then, the mixed discriminant $MD_{r,A}$ always divides the iterated discriminant $ID_{r,A}$. Moreover,*

- (1) *If $\text{codim}_{X_A^\vee}(\text{sing}(X_A^\vee)) > r$, then $ID_{r,A} = MD_{r,A}$.*
- (2) *If $\text{codim}_{X_A^\vee}(\text{sing}(X_A^\vee)) = r$, then $ID_{r,A} = MD_{r,A} \prod_{i=1}^{\ell} \text{Ch}_{Y_i}^{\mu_i}$, where Y_1, \dots, Y_ℓ are the irreducible components of $\text{sing}(X_A^\vee)$ of codimension r , with respective multiplicities $\mu_i \geq 2$.*
- (3) *If $\text{codim}_{X_A^\vee}(\text{sing}(X_A^\vee)) < r$, then $ID_{r,A} = 0$.*

The paper is organized as follows. In Section 2 we present some examples that motivate the theory of iterated discriminants. In Section 3 we present material on mixed discriminants and Cayley configurations.

In Section 4 we develop the theory of iterated discriminants and prove our main results Theorem 4.4 and Proposition 4.5. We prove in Proposition 4.8 that the multiplicities μ_k in Theorem 4.4 are at least two. Very few is known in general about these multiplicities, except for the homogeneous case of three variables studied in [4] and the general results in [22]. Based on this evidence and some examples we computed, we state Conjecture 4.9. The difficulty in determining these multiplicities relies in the fact that for general point configurations A , a complete description of the components of the singular locus of the dual varieties X_A and their codimensions is out of reach for the moment. By a result of Katz ([20, Proposition 3.4]), it is expected that the codimension one components correspond to the double point locus (the closure of those hypersurfaces with two different non-degenerate singular points) and the cusp locus (the closure of those hypersurfaces having a single degenerate singular point with an A_2 -singularity). The case of hyperdeterminants has been exhaustively described in [28, Theorem 0.5], where it is shown that in the non-defective case only one irreducible component with codimension one can exist, or there could be several irreducible components of codimension one of both types. Already the univariate

sparse case poses some challenges [11]. Even the particular case of the existence of a cusp component with codimension one when D_A corresponds to the mixed discriminant of two planar configurations, recently studied in [23], is not trivial. A general approach to describe the irreducible components (and much more information) via the computation of tropical fans and characteristic classes is developed in [14].

In Section 5 we ask more broadly when mixed and iterated discriminants are equal, for products of scaled simplices, that is, when X_A is a Segre–Veronese variety. The case of Segre varieties was solved in [28], via a careful study of the singularities of hyperdeterminant varieties. As a corollary of our results, we show in Proposition 5.2 that the iterated method to characterize singular complete intersections for $r + 1$ hypersurfaces of the same degree $d > 1$ in \mathbb{P}^n gives the corresponding mixed discriminant if and only if $r = 1$ and $d = 2$ (the case of two quadric hypersurfaces already found in [24, Theorem 8.2]).

Our Conjecture 5.3 is the following, with notation as in Section 5:

Conjecture. *The equality $\deg(ID_{r,A_\ell,d,k}) = \deg(MD_{r,A_\ell,d,k})$ holds if and only if*

$$\mathbb{P}^r(1) \times \mathbb{P}^{k_1}(d_1) \times \cdots \times \mathbb{P}^{k_\ell}(d_\ell)$$

is of one of the following cases:

- (1) $\mathbb{P}^r \times \mathbb{P}^m \times \mathbb{P}^m$, $m \geq 1$, $r = 1, 2$,
- (2) $(\mathbb{P}^1)^4$,
- (3) $\mathbb{P}^1 \times \mathbb{P}^n(2)$.

A partial answer is given in Theorem 5.6 and Proposition 5.2.

Finally, in Section 6 we analyze the case of plane curves. Theorem 6.3 shows that for planar configurations A consisting of the lattice points of a smooth polygon, the only case where $MD_{1,A}$ equals $ID_{1,A}$ are the known cases in which the polygon is the unit square (the bilinear case) or $2\Delta_2$, the standard triangle of size 2. This implies that in all other cases, the singularities of the discriminant locus have codimension one; that is, there are “many” different types of singular hypersurfaces defined by A -polynomials. A factorization of the iterated discriminants gives all components of the singular locus of codimension one.

2. Motivating examples

In this section we present some motivating examples that we abstract in the paper. The first two correspond to two classical cases in which the iterated discriminant actually computes the mixed discriminant. The last two are the simplest cases which already show the occurrence of other factors of the iterated discriminant.

Example 2.1. Let $A = \{(0, 0), (1, 0), (0, 1), (1, 1)\}$ be the vertices of the unit cube and let $f = c_{00} + c_{10}x_1 + c_{01}x_2 + c_{11}x_1x_2$ be an A -polynomial. In this case, $D_A(f) = c_{00}c_{11} - c_{10}c_{01}$ is a polynomial of degree 2, which equals the determinant of the matrix

$$\begin{pmatrix} c_{00} & c_{01} \\ c_{10} & c_{11} \end{pmatrix}.$$

In case $r + 1 = 2$, the mixed discriminant associated with two A -polynomials

$$p_0 = c_{00}^1 + c_{10}^1x_1 + c_{01}^1x_2 + c_{11}^1x_1x_2, \quad p_1 = c_{00}^2 + c_{10}^2x_1 + c_{01}^2x_2 + c_{11}^2x_1x_2,$$

is the following degree four irreducible polynomial, which is the hyperdeterminant of format $2 \times 2 \times 2$ (see [16, pp. 475–479]):

$$\begin{aligned} MD_{1,A}(p_0, p_1) &= c_{00}^2 c_{11}^2 - 2c_{00}^2 c_{01}^2 c_{10}^1 c_{11}^1 - 2c_{00}^2 c_{10}^2 c_{01}^1 c_{11}^1 - 2c_{00}^2 c_{11}^2 c_{00}^1 c_{11}^1 \\ &+ 4c_{00}^2 c_{11}^2 c_{01}^1 c_{10}^1 + 2c_{00}^2 c_{00}^1 c_{11}^2 - 4c_{00}^2 c_{01}^1 c_{10}^1 c_{11}^1 + c_{01}^2 c_{10}^2 + 4c_{01}^2 c_{10}^2 c_{00}^1 c_{11}^1 \\ &- 2c_{01}^2 c_{10}^2 c_{01}^1 c_{10}^1 - 2c_{01}^2 c_{11}^2 c_{00}^1 c_{10}^1 + 2c_{01}^2 c_{00}^1 c_{10}^1 c_{11}^1 + c_{10}^2 c_{01}^2 \\ &- 2c_{10}^2 c_{11}^2 c_{00}^1 c_{01}^1 + 2c_{10}^2 c_{00}^1 c_{01}^1 c_{11}^1 + c_{11}^2 c_{00}^2 c_{10}^2 - 2c_{11}^2 c_{00}^2 c_{11}^1 \\ &+ c_{00}^1 c_{11}^2. \end{aligned}$$

It vanishes at (p_0, p_1) with respective coefficient vectors $(1, 1, -2, -1)$, $(1, 1, -3, -2)$, corresponding to the tangent hyperbolas in Figure 1.

One form of computing D_A is as the iterated discriminant $ID_{1,A}$. Write

$$\det \begin{pmatrix} c_{00}^1 + \lambda c_{00}^2 & c_{01}^1 + \lambda c_{01}^2 \\ c_{10}^1 + \lambda c_{10}^2 & c_{11}^1 + \lambda c_{11}^2 \end{pmatrix} = \Delta_0 + \Delta_1 \lambda + \Delta_2 \lambda^2,$$

and then compute

$$MD_{1,A}(c^1, c^2) = \Delta_1^2 - 4\Delta_0\Delta_2,$$

as the univariate resultant of the degree 2 polynomial $\Delta_0 + \Delta_1 \lambda + \Delta_2 \lambda^2$ in λ with coefficients in $\mathbb{Z}[c^1, c^2]$. This compact formula is the simplest case of Schäfli's formula to compute the mixed discriminant $MD_{1,A}$.

Example 2.2. Let us consider again the case discussed in the introduction corresponding to the singular intersections of two quadric surfaces p_0, p_1 in three-space. We display their common support A as the columns of the following 3×10 matrix:

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 2 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 2 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 2 \end{bmatrix}.$$

We also display the corresponding Cayley configuration $C = \Delta_1 \times A$ as the columns of the following 5×20 -matrix:

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 2 & 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 2 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 2 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 2 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 2 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 2 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix}.$$

In this case, we know that $(X_C)^v$ is a hypersurface by [8]. Thus we have that the polynomial $MD_{1,A}(p_0, p_1)$ cuts out the closure of the locus of coefficients for which the two quadrics lie tangent to one another at a point and it can be computed via the discriminant D_C by Proposition 3.3. It can be also computed as the iterated discriminant in (2). This polynomial can be studied through tropical discriminants as in [10].

Moreover, one can compute the univariate discriminant $D_{4\Delta_1}$ of a degree 4 polynomial as the discriminant of its cubic resolvent from Galois theory. Let Δ_i denote the coefficient λ^i in $\det(M_0 + tM_1)$. Then

$$MD_{1,A}(p_0, p_1) = \frac{4p^3 - q^2}{27},$$

where

$$\begin{aligned} p &= 12\Delta_4\Delta_0 - 3\Delta_3\Delta_1 + \Delta_2^2, \\ q &= 72\Delta_4\Delta_2\Delta_0 + 9\Delta_3\Delta_2\Delta_1 - 27\Delta_4\Delta_1^2 - 27\Delta_0\Delta_3^2 - 2\Delta_2^3. \end{aligned}$$


This gives a compact and feasible way of computing the mixed discriminant $MD_{1,A}$ we are interested in. In fact, expanding this expression in terms of the coefficients of p_0, p_1 is beyond the capabilities of the excellent Computer Algebra system Macaulay2 [17] in a standard computer, because it is a polynomial of degree 24 which has degree 12 in both the coefficients of p_0 and p_1 . Note that a general polynomial of bidegree (12, 12) in two groups of 10 variables has more than $4 \cdot 10^{11}$ monomials!

The general case is hinted in the following simple examples.

Example 2.3. Consider the two-dimensional configuration

$$A = \{(0, 0), (1, 0), (2, 0), (0, 1), (1, 1)\}$$

corresponding to the first Hirzebruch surface \mathbb{F}_1 .



Given a generic polynomial f with support A :

$$f(x, y) = a_0 + a_1x + a_2x^2 + y(b_0 + b_1x),$$

the A -discriminant coincides with the resultant of the two univariate polynomials $a_0 + a_1x + a_2x^2$ and $b_0 + b_1x$ and thus is equal to the degree 3 polynomial

$$D_A(f) = a_0b_1^2 - a_1b_0b_1 + a_2b_0^2.$$

The mixed discriminant $MD_{1,A}$ has degree 8, while the iterated discriminant $ID_{2,A}$ has degree $2 \cdot 3 \cdot (3 - 1)^1 = 12$ by (16). There is another irreducible factor that we explain in Theorem 4.4 and compute in Example 4.6.

Example 2.4. We now consider the case of a univariate polynomial of degree 3 with $A = \{0, 1, 2, 3\}$ and $r = 1$. Given two cubic polynomials p_0, p_1 depending on a variable x , their mixed discriminant equals the discriminant of the Cayley configuration

$$C = \{(0, 0), (0, 1), (0, 2), (0, 3), (1, 0), (1, 1), (1, 2), (1, 3)\}$$

at the polynomial $p_0 + tp_1$ in one more variable t . In fact, $D_C(p_0 + tp_1)$ equals the univariate resultant $\text{Res}_{3,3}(p_0, p_1)$. This resultant can be computed as the determinant of the associated Sylvester matrix and therefore has degree 6 in the vectors of coefficients of p_0, p_1 . Since the discriminant D_A of a cubic univariate polynomial has degree 4, the iterated discriminant $ID_{1,A} = D_{4\Delta_1}(D_A(p_0 + tp_1))$ instead has degree $2 \cdot 4 \cdot 3 = 24$ according to (16). It has another irreducible factor of degree 6 raised to the third power, which is the Chow form of the singular locus of $D_A = 0$ corresponding to degree 3 polynomials with a triple root (a degenerate multiple root), predicted by Theorem 4.4.

3. The mixed discriminant and the discriminant of the Cayley configuration

In this section we show in Proposition 3.3 that the mixed discriminant $MD(A_0, \dots, A_r)$ coincides in the non-defective case with the discriminant of the associated Cayley configuration D_C , thus generalizing [6, Theorem 2.1]. Note that when C is defective these varieties need not coincide, as shown in [6, Example 2.2]. We also characterize, in Proposition 3.4, non-defectivity of C when all A_i are equal. The latter result relies on a classical criterion by Katz, stated as Lemma 3.1 below, and is a simple consequence of [27, Theorem 0.1].

Recall that for a projective variety X , the *dual defect* of X is defined to be

$$\text{def}(X) := \text{codim}(X^\vee) - 1, \tag{4}$$

where X^\vee is the dual variety consisting of singular hyperplane sections to X . In particular, if the dual variety is a hypersurface as expected, then the dual defect is equal to 0 and X is said to be non-defective. When $X = X_A$ for some finite lattice configuration A , we also say that A is non-defective. In this context, we have the following lemma due to Katz.

Lemma 3.1 ([20]). *Let $A \subset \mathbb{Z}^n$ be a lattice configuration with $|A| = N + 1$. Let $H_p(f)$ denote the Hessian matrix of an A -polynomial f . Then*

$$\text{codim } X_A^\vee = 1 + \min_f (\text{corank}(H_u(f)))$$

where u is a general point and f varies among the polynomials with support in A vanishing at u .

In particular, $\text{codim } X_A^\vee = 1$ implies that polynomials vanishing at a general point u together with their partial derivatives have Hessian of maximal rank.

Observe that Lemma 3.1 is equivalent to saying that in the non-defective case, the closure of the singular A -polynomials coincides with the closure of the nodal A -polynomials, that is, polynomials only admitting non-degenerate multiple roots.

Corollary 3.2. *If A is a non-defective finite lattice configuration, then*

$$X_A^\vee = \overline{\left\{ f \in \mathbb{P}^{N^\vee} : f(u) = 0, \frac{\partial f}{\partial x_i}(u) = 0 \text{ and } \det(H(f))(u) \neq 0 \text{ for some } u \in (K^*)^n \right\}}.$$

Proof. The inclusion “ \subseteq ” follows by definition and the inclusion “ \supseteq ” follows by Lemma 3.1. ■

Let us now consider the Cayley configuration C associated to $r + 1$ finite lattice configurations A_0, \dots, A_r in \mathbb{Z}^n . We remark that the r in [16] corresponds to our $r - 1$. We use the following notation: $(\lambda, x) = (\lambda_0, \dots, \lambda_r, x_1, \dots, x_n)$. A polynomial f with support on C has the form

$$f = \sum_0^r \lambda_i p_i,$$

where p_i are A_i -polynomials in the variables x . Consider the Jacobian matrix

$$[\nabla_x(p_0)(u) \quad \cdots \quad \nabla_x(p_r)(u)]$$

of p_0, \dots, p_r at u . Notice that $f \in X_C^\vee$ if there exists $(\lambda, u) \in (\mathbb{C}^\nu)^{r+1} \times (\mathbb{C}^\nu)^n$ such that

$$p_0(u) = \cdots = p_r(u) = 0 \quad \text{and} \quad \lambda \in \ker [\nabla_x(p_0)(u) \quad \cdots \quad \nabla_x(p_r)(u)]^T;$$

or equivalently, if $\sum_{i=0}^r \lambda_i \nabla_x(p_i)(u) = 0$ and thus the gradients are linearly dependent. In particular,

$$\text{rank}([\nabla_x(p_0)(u) \ \cdots \ \nabla_x(p_r)(u)]) \leq r.$$

We will now prove that the locus where the rank is exactly r characterizes the dual variety X_C^\vee , assuming it is a hypersurface.

Proposition 3.3. *Let A_0, \dots, A_r and C as above and assume that C is non-defective. Then,*

$$MD(A_0, \dots, A_r)(p_0, \dots, p_r) = D_C\left(\sum_{i=0}^r \lambda_i p_i\right),$$

where p_i are A_i -polynomials for $i = 0, \dots, r$ and $(\lambda_0, \dots, \lambda_r)$ are variables.

Proof. Let ϕ_f be the tuple of coefficients of f . Corollary 3.2 implies that

$$X_C^\vee = \overline{\left\{ \phi_f : f(\lambda, u) = 0, p_{A_i}(u) = 0, i = 0, \dots, r, \right. \\ \left. \sum_{j=0}^r \lambda_j \frac{\partial p_{A_j}}{\partial x_i}(u) = 0 \text{ and } \det(H(f))(\lambda, u) \neq 0 \right\}},$$

where $(\lambda, u) \in (\mathbb{C}^\nu)^{r+1} \times (\mathbb{C}^\nu)^n$. Here, $H(f)(\lambda, u)$ means the following: as f is homogeneous in the variables λ and $\lambda \in (\mathbb{C}^\nu)^{r+1}$, we assume that $\lambda_0 = 1$ and that $(\lambda_1, \dots, \lambda_r)$ are its affine coordinates. Thus, $H(f)(\lambda, u)$ is the Hessian of f with respect to the variables $(\lambda_1, \dots, \lambda_r, x_1, \dots, x_n)$. This Hessian matrix is of the form

$$H_{(u,\lambda)}(\phi_f) = \begin{bmatrix} \sum \lambda_i H(p_{A_i})(u) & \begin{bmatrix} \nabla(p_{A_1})(u) \\ \vdots \\ \nabla(p_{A_r})(u) \end{bmatrix}^T \\ \begin{bmatrix} \nabla(p_{A_1})(u) \\ \vdots \\ \nabla(p_{A_r})(u) \end{bmatrix} & 0 \end{bmatrix}.$$

It follows that if $\phi_f \in X_C^\vee$ and $\det H(f)(\lambda, u) \neq 0$, which happens for generic points in X_C^\vee by Corollary 3.2, then

$$\text{rank} \begin{bmatrix} \nabla(p_{A_1})(u) \\ \vdots \\ \nabla(p_{A_r})(u) \end{bmatrix} = r,$$

that is, the gradients of the polynomials $p_{A_i}(u)$ for $i \neq 0$ form a matrix of rank r , that is, they are linearly independent. This happens similarly for the gradients of

any subset of r polynomials $p_{A_j}(u)$. Moreover, this is exactly the condition implying that ϕ_f belongs to the mixed-discriminantal variety $MD(A_0, \dots, A_r) = 0$ which we denote by X_{MD} . It follows that $X_C^\vee \subseteq X_{MD}$ and that X_{MD} is also a hypersurface, i.e., $MD(A_0, \dots, A_r) \neq 1$.

The reverse inclusion follows essentially from the definition. In fact, if $\phi_f \in X_{MD}$ is generic, then there is a common zero $u \in (C^\vee)^n$ of p_{A_0}, \dots, p_{A_r} and a linear dependency $\sum \lambda_i \nabla(p_{A_i})(u) = 0$ with all $\lambda_i \neq 0$, because all the maximal minors in the matrix

$$\begin{bmatrix} \nabla(p_{A_0})(u) \\ \vdots \\ \nabla(p_{A_r})(u) \end{bmatrix}$$

are assumed to be non-zero. It follows that $\phi_f \in X_C^\vee$. ■

Notice that if $A_0 = A_1 = \dots = A_r = A$ then $C = \{e_0, \dots, e_r\} \times A$, which is usually written as $C = \Delta_r \times A$. Following [27], we define the following quantity associated to a projective variety X :

$$\mu(X) = \dim(X) + \text{def}(X), \tag{5}$$

where the defect of X has been defined in (4). We end this section with the following result about non-defectivity.

Proposition 3.4. *Let A be a non-defective finite lattice configuration. Then, the associated Cayley configuration $C = \Delta_r \times A$ is non-defective if and only if $r \leq \dim(X_A)$.*

Proof. Note that X_C equals the Segre embedding of $\mathbb{P}^r \times X_A$. We can then use [28, Theorem 0.1], which says that

$$\mu(\mathbb{P}^r \times X_A) = \max(r + \dim(X_A), r + \text{def}(\mathbb{P}^r), \dim(X_A) + \text{def}(X_A)).$$

According to (5), we have that $\mu(X_C) = r + \dim(X_A) + \text{def}(X_C)$. Since $\text{def}(\mathbb{P}^r) = r$, and by hypothesis $\text{def}(X_A) = 0$, we get that

$$\mu(X_C) = \max(r + \dim(X_A), 2r, \dim(X_A)).$$

When $r \leq \dim(X_A)$, we get that $\mu(X_C) = r + \dim(X_A)$, which implies that

$$\text{def}(X_C) = 0.$$

On the other side, when $r > \dim(X_A)$, we have that $\mu(X_C) = 2r$, and so

$$\text{def}(X_C) = r - \dim(X_A) > 0. \quad \blacksquare$$

4. The multivariate iterated discriminant

In the remainder of the paper we will consider the case $A_i = A$ for $i = 0, \dots, r$. In order to establish an iterated process for the mixed discriminant it is convenient to consider the *geometric iterated discriminant* $JD_{r,A}$ introduced in Definition 4.2 below. In Proposition 4.5 we prove that this polynomial coincides with the iterated discriminant $ID_{r,A}$ from Definition 1.1. It implies that Theorem 4.4, which can be considered the main result of this paper, also holds for $ID_{r,A}$, as stated in the Introduction.

Recall that given an irreducible and reduced projective variety $Y \subset \mathbb{P}^N$ of codimension s , its Chow form Ch_Y is defined as follows. Consider linear subspaces of dimension ℓ in \mathbb{P}^N , $L \in \text{Gr}(\ell + 1, N + 1)$. If $s \geq \ell + 1$, any generic L will not intersect Y . The irreducible subvariety

$$\{L \in \text{Gr}(\ell + 1, N + 1) : L \cap Y \neq \emptyset\}$$

parametrizing the exceptional intersection locus, has codimension $(s - \ell)$ in $\text{Gr}(\ell + 1, N + 1)$. In case $\ell = s - 1$, the defining polynomial is denoted by Ch_Y and it is called the *Chow form* of Y [16, p. 99].

We also need to recall two classical facts which will be used in the proof of our main Theorem 4.4.

Remark 4.1. Given a finite lattice configuration A and a generic singular hyperplane section of X_A , we can recover the intersection point by means of the gradient of the discriminant D_A . Precisely,

- (1) as we are assuming that $\text{char}(K) = 0$, if a regular point H in the dual variety X_A^\vee is tangent to X_A at a regular point y_H , then this projective point is unique and $y_H = \nabla D_A(H)$ (see [16, Theorem 1.1, Chapter 1]). This is referred to as *biduality*.
- (2) when $X_A^\vee \subset (\mathbb{P}^N)^\vee$ is a hypersurface, biduality implies that the Gauss map $\gamma: X_A^\vee \dashrightarrow \mathbb{P}^n$ is defined by $H \mapsto \nabla D_A(H) = y_H$ and the closure of its image equals X_A .

Let $A = \{a_0, \dots, a_N\} \subset \mathbb{Z}^n$ be a lattice configuration. We will assume henceforth that A is non-defective and that D_A is a homogeneous polynomial of degree $\delta > 0$.

Given A -polynomials

$$p_i = \sum_{j=0}^N c_{ij} x^{a_j}, \quad i = 0, \dots, r,$$

we also denote by $(p_0, \dots, p_r) \in (\mathbb{P}^{(r+1)(N+1)-1})^\vee$ the vector of their coefficients.

For any $\lambda = (\lambda_0, \dots, \lambda_r) \in \mathbb{P}^r$, we write

$$P_\lambda := \lambda_0 p_0 + \dots + \lambda_r p_r \in (\mathbb{P}^N)^\nu.$$

Definition 4.2. Consider the incidence variety

$$\Sigma = \left\{ ((p_0, p_1, \dots, p_r), \lambda) \in (\mathbb{P}^{(r+1)(N+1)-1})^\nu \times \mathbb{P}^r : \sum_j c_{ij} \frac{\partial D_A}{\partial c_j}(P_\lambda) = 0, \right. \\ \left. i = 0, \dots, r \right\}. \quad (6)$$

Let $\pi: \Sigma \rightarrow (\mathbb{P}^{(r+1)(N+1)-1})^\nu$ be the linear projection onto the first factor. The *r-multivariate iterated dual scheme* $\pi(\Sigma)$ is defined by the projective elimination ideal

$$\pi I = (I : m^\infty) \cap \mathbb{C}[c], \quad (7)$$

where $\mathbb{C}[c]$ is the ring of polynomials in the variables c_{ij} , m is the irrelevant ideal of \mathbb{P}^r , and I is the ideal

$$I = \left\langle \sum_j c_{ij} \frac{\partial D_A}{\partial c_j}(P_\lambda), i = 0, \dots, r \right\rangle.$$

When $\pi(\Sigma)$ has codimension one, we denote by $JD_{r,A} \in \mathbb{Z}[c]$ a generator (unique up to multiplication by a non-zero constant) of the union of the codimension one components of πI and we call it the *geometric iterated discriminant*.

Notice that the projection is in general not irreducible; see, for instance, Example 2.3. We will see in Proposition 4.5 below that the geometric iterated discriminant $JD_{r,A}$ coincides with the more naive definition of the iterated discriminant $ID_{r,A}$ from Definition 1.1.

Let $(p, \lambda) = ((p_0, \dots, p_r), (\lambda_0, \dots, \lambda_r)) \in \Sigma$. In order to understand the projection π we consider two auxiliary maps

$$\phi: \Sigma \rightarrow X_A^\nu, \quad T: (\mathbb{P}^{(r+1)(N+1)-1})^\nu \rightarrow \text{Gr}(r+1, N+1)$$

such that

$$\begin{array}{ccc} \Sigma & \xrightarrow{\phi} & X_A^\nu \\ \downarrow & & \\ \pi(\Sigma) & \dashrightarrow^{T} & \text{Gr}(r+1, N+1) \end{array}$$

defined by $\phi(p, \lambda) = P_\lambda$ and $T(p_0, \dots, p_r) = T_p$, where we denote by T_p the projective linear span of p_0, \dots, p_r .

Lemma 4.3. *Let $p = (p_0, \dots, p_r) \in (\mathbb{P}^{(r+1)(N+1)-1})^\nu$ such that $JD_{r,A}(p) = 0$. Then, T_p is tangent to X_A^ν at some point ξ .*

Proof. If $JD_{r,A}(p_0, \dots, p_r) = 0$, then there exists λ such that $(p, \lambda) \in \Sigma$; let $\xi = P_\lambda$. Consider the equalities

$$0 = \sum_j c_{ij} \frac{\partial D_A}{\partial c_j}(P_\lambda), \quad i = 0, \dots, r. \quad (8)$$

Note that these equations equal the derivatives with respect to $\lambda_0, \dots, \lambda_r$ of the composed function $(f, \mu) \rightarrow D_A(\sum_{i=0}^r \mu_i f_i)$ at the point (p, λ) . The Euler relation implies that $D_A(P_\lambda) = 0$, and thus $P_\lambda \in X_A^\nu$. Moreover, (8) implies that each p_i lies in $T_{X^\nu, \xi}$, which is equivalent to $T_p \subset T_{X^\nu, \xi}$. ■

Recall that we denote by $\text{sing}(X_A^\nu)$ the subscheme of X_A^ν defined by the ideal generated by the partial derivatives of D_A .

Theorem 4.4. *Assume $A \subset \mathbb{Z}^n$ is non-defective and let $r \in \mathbb{Z}$ with $0 \leq r \leq \dim(X_A)$. Then, the mixed discriminant $MD_{r,A}$ always divides the geometric iterated discriminant $JD_{r,A}$. Moreover,*

- (1) *if $\text{codim}_{X_A^\nu}(\text{sing}(X_A^\nu)) > r$, then $JD_{r,A} = MD_{r,A}$.*
- (2) *if $\text{codim}_{X_A^\nu}(\text{sing}(X_A^\nu)) = r$, then $JD_{r,A} = MD_{r,A} \prod_{i=k}^\ell \text{Ch}_{Y_k}^{\mu_k}$, where Y_1, \dots, Y_ℓ are the irreducible components of $\text{sing}(X_A^\nu)$ of maximal dimension r , with respective multiplicities $\mu_k \geq 2$.*
- (3) *if $\text{codim}_{X_A^\nu}(\text{sing}(X_A^\nu)) < r$, then $\pi(\Sigma) = (\mathbb{P}^{(r+1)(N+1)-1})^\nu$ and $JD_{r,A} = 0$.*

Proof. As already observed, in the classical case of $r = 0$, we have

$$ID_{0,A} = MD_{0,A} = D_A.$$

Note also that by Propositions 3.4 and 3.3, $\deg(MD_{r,A}) > 0$ and it is irreducible.

Observe that the map ϕ is surjective since for any $F \in X_A^\nu$,

$$F = \phi\left(F, \dots, F, \frac{1}{(r+1)}, \dots, \frac{1}{(r+1)}\right),$$

and we have

$$\left(F, \dots, F, \frac{1}{(r+1)}, \dots, \frac{1}{(r+1)}\right) \in \Sigma.$$

The rational map T is defined over the open dense subset $U_T = \{p : T_p \simeq \mathbb{P}^r\}$ of all p with linear span of projective dimension r . Notice also that T is surjective and that for each $H \in \text{Gr}(r+1, N+1)$, the fiber $T^{-1}(H)$ has dimension $(r+1)^2 - 1$.

Let $\Sigma^\circ = \phi^{-1}((X_A^\vee)_{\text{reg}})$ and $\Sigma' = \phi^{-1}(\text{sing}(X_A^\vee))$; that is, let

$$\Sigma^\circ = \{(p, \lambda) \in \Sigma : P_\lambda \in (X_A^\vee)_{\text{reg}}\}, \quad \Sigma' = \{(p, \lambda) \in \Sigma : P_\lambda \in \text{sing}(X_A^\vee)\}.$$

It follows that $\pi(\Sigma) = \pi(\Sigma^\circ) \cup \pi(\Sigma')$.

We claim that $\overline{\pi(\Sigma^\circ)} \subseteq V(MD_{r,A})$. In fact, take a generic point $(p, \lambda) \in \pi(\Sigma^\circ)$. We can then assume that not only $P_\lambda \in (X_A^\vee)_{\text{reg}}$, but also there is a unique regular point $y = (x^{m_0} : \dots : x^{m_N}) \in X_A$ with $x \in (K^*)^n$ such that

$$P_\lambda(x) = 0, \quad \frac{\partial P_\lambda}{\partial x_i}(x) = 0, \quad i = 1, \dots, n.$$

By Remark 4.1,

$$y = \left(\frac{\partial D_A}{\partial c_0}(P_\lambda) : \dots : \frac{\partial D_A}{\partial c_n}(P_\lambda) \right).$$

The equations $\frac{\partial P_\lambda}{\partial x_i}(x) = 0$ for $i = 1, \dots, n$ mean that $\sum \lambda_i \nabla(p_i)(x) = 0$. Moreover, $p_i(x) = 0$ for all i , because

$$p_i(x) = \sum_j c_{ij} y_j = k \sum_j c_{ij} \frac{\partial D_A}{\partial c_j}(P_\lambda) = 0, \quad i = 0, \dots, r \quad (9)$$

for some $k \in K^*$ such that $y = k \nabla D_A(P_\lambda)$. This implies that $MD_{r,A}(p_0, \dots, p_r) = 0$.

We now show that $V(MD_{r,A}) \subseteq \pi(\Sigma)$. Let (p_0, \dots, p_r) be a generic element in the zero locus of $MD_{r,A}$. Then there exists $(u, \lambda) \in (K^*)^{n+r+1}$ such that

$$p_0(u) = \dots = p_r(u) = 0, \quad \left[\nabla p_0(u) \quad \dots \quad \nabla p_r(u) \right] \begin{bmatrix} \lambda_0 \\ \vdots \\ \lambda_r \end{bmatrix} = 0.$$

We claim that $(p, \lambda) \in \Sigma$. If $P_\lambda \in \text{sing}(X_A^\vee)$, then

$$\frac{\partial D_A}{\partial c_j}(P_\lambda) = 0,$$

and thus $(p_0, \dots, p_r) \in \pi(\Sigma)$. If instead $P_\lambda \in (X_A^\vee)_{\text{reg}}$ is generic, then biduality gives $\nabla D_A(P_\lambda) = y$ with $y = (u^{m_0} : \dots : u^{m_N})$, and thus

$$\sum_j c_{ij} \frac{\partial D_A}{\partial c_j}(P_\lambda) = p_i(u) = 0$$

as in (9), implying again that $(p_0, \dots, p_r) \in \pi(\Sigma)$. We have then proved that

$$\overline{\pi(\Sigma^\circ)} \subseteq V(MD_{r,A}) \subseteq \pi(\Sigma). \quad (10)$$

Consider the non-embedded primary components of the ideal $\langle \frac{\partial D_A}{\partial c_j}, j = 0, \dots, N \rangle$ defining the singular locus of X_A^v . Correspondingly, we consider the decomposition into irreducible components $\text{sing}(X_A^v) = \bigcup Y_k$. Define

$$V_k = \{H \in \text{Gr}(r+1, N+1) : H \cap Y_k \neq \emptyset\}, \quad \Sigma_k = \phi^{-1}(Y_k). \quad (11)$$

Recall that $\text{codim}_{\text{Gr}(r+1, N+1)}(V_k) = \max\{0, \text{codim}_{\mathbb{P}^N}(Y_k) - r\}$.

Assume that $\text{codim}_{X_A^v}(\text{sing}(X_A^v)) > r$. Then $\text{codim}_{\text{Gr}(r+1, N+1)}(V_k) \geq 2$ for all k . It follows that for all i ,

$$\begin{aligned} \text{codim}_{\mathbb{P}^{(r+1)(N+1)-1^v}}(\pi(\Sigma_i)) &= \text{codim}_{\mathbb{P}^{(r+1)(N+1)-1^v}}(\overline{\pi(\Sigma_i) \cap U_T}) \\ &\geq \text{codim}_{\mathbb{P}^{(r+1)(N+1)-1^v}}(\overline{T^{-1}(V_i)}) \geq 2. \end{aligned}$$

The containment in equation (10) then implies that $\pi(\Sigma)$ is of codimension one and set-theoretically coincides with $V(MD_{r,A})$. If $\text{codim}_{X_A^v}(\text{sing}(X_A^v)) = r$, then

$$\text{codim}_{\text{Gr}(r+1, N+1)}(V_k) = 1,$$

and thus by definition $\pi(\Sigma_k) = V(\text{Ch}_{Y_k}^{\mu_k})$ for some integer exponents μ_k . We prove that these multiplicities are at least equal to 2 in Proposition 4.8 below.

As the mixed discriminant $MD_{r,A}$ is irreducible, it remains to show that the multiplicity of $MD_{r,A}$ in $ID_{r,A}$ is equal to 1. For that, it is enough to show that there exists $(p_0^v, \dots, p_r^v) \in V(MD_{r,A})$ and λ^v such that $(p^v, \lambda^v) \in \Sigma$ and $d\pi((p^v, \lambda^v))$ has maximal rank.

We start by choosing a point $\xi \in \text{reg}(X_A^v)$ such that $\text{rank}(H) = n+1$, where $H = \text{Hess}(D_A)(\xi)$. Notice that $H = \text{Jac}(\gamma)(\xi)$, where $\gamma: X_A^v \dashrightarrow X_A$ is the Gauss map defined as $\gamma(y) = \nabla D_A(y)$, which in affine coordinates has generic rank equal to $n = \dim(X_A)$. Up to a change of coordinates, H can be assumed to be of the form

$$H = \begin{bmatrix} I_{n+1} & 0 \\ 0 & 0 \end{bmatrix}. \quad (12)$$

Consider $Z = \{M \in \text{Gr}(r+1, N+1) : \xi \in M \subset T_{X_A^v, \xi}\}$. Note that for every $p \in T^{-1}(Z)$, we have $\xi \in T_p \subset T_{X_A^v, \xi}$ and $\dim(T_p) = r$ as $p \in U_T$. It follows that there exists a unique λ^* such that $(p, \lambda^*) \in \Sigma$ and $\phi(p, \lambda^*) = \xi$. Consider the $(r+1) \times (N+1)$ matrix C_p whose i th row corresponds to the coefficients of p_i . Without loss of generality, the matrix C_p can be assumed to be of the form $[I_{r+1}, C]$.

The matrix M of the lifted linear map

$$\pi: (\mathbb{C}^{(r+1)(N+1)})^v \times \mathbb{C}^{r+1} \rightarrow (\mathbb{C}^{(r+1)(N+1)})^v$$

is equal to

$$M = [I_{(r+1)(N+1)}, 0],$$

where 0 denotes the zero matrix of size $(r+1)(N+1) \times (r+1)$. Let us call g_0, \dots, g_r the defining equations of Σ in (6). It follows that $d\pi(p, \lambda^*)$ is of maximal rank if and only if the square $(r+1)(N+1) + (r+1)$ -matrix M' with upper rows consisting of the Jacobian of g_0, \dots, g_n with respect to the variables $(c_{01}, \dots, c_{0N}, \dots, c_{r0}, \dots, c_{rN}, \lambda_0, \dots, \lambda_r)$ evaluated at (p, λ^*) and lower rows given by the matrix M , has maximal rank. But given the form of M , this is equivalent to the fact that the $(r+1) \times (r+1)$ submatrix H^* at the right upper corner of M' has maximal rank $r+1$. Recall from the proof of Lemma 4.3 that g_0, \dots, g_r equal the derivatives with respect to $\lambda_0, \dots, \lambda_r$ of the composed function

$$(f, \lambda) \rightarrow D_A \left(\sum_{i=0}^r \lambda_i f_i \right).$$

Then, H^* consists of the Hessian matrix with respect to the λ -variables of this composed function. Therefore, we have that

$$H^* = C_p H C_p^t.$$

Recall that we assume that $r \leq n$, and thus $r+1 \leq n+1$. Given the form of the coefficient matrix C_p and of the Hessian matrix H in (12), we deduce that H^* is of maximal rank because it is the identity matrix I_{r+1} .

Assume that $\text{codim}_{X_A^v}(\text{sing}(X_A^v)) < r$. Then $\text{codim}_{\mathbb{P}^N}(Y_k) < r+1$ for all k . The assumption also implies that any element of the Grassmannian belongs to V_k for all k (defined in (11)), and that

$$\pi(\Sigma') \cap U_T = T^{-1}(\text{Gr}(r+1, N+1)) = U_T.$$

It follows that

$$\mathbb{P}^{(r+1)(N+1)-1} = \overline{U_T} = \overline{\pi(\Sigma_i) \cap U_T} = \pi(\Sigma_i) \subset \pi(\Sigma),$$

and thus $\pi(\Sigma) = (\mathbb{P}^{(r+1)(N+1)-1})^v$. ■

The following Proposition 4.5 explains the name geometric iterated discriminant: we show that under the hypotheses of Theorem 4.4, the polynomial $JD_{r,A}$ in Definition 4.2 equals the polynomial $ID_{r,A}$ in Definition 1.1, and thus when it is non-zero it can be computed as a discriminant of a discriminant.

Recall that, given a natural number d , we denote by $d\Delta_r$ in \mathbb{R}^{r+1} the lattice configuration given by the integer points in the dilated unit simplex d times, and by $D_{d\Delta_r}$ the associated discriminant. For any homogeneous polynomial $H = H(\lambda_0, \dots, \lambda_r)$ of degree d , the discriminant of H equals, up to constant, the resultant of its partial derivatives:

$$D_{d\Delta_r}(H) = \text{Res}_{d-1} \left(\frac{\partial H}{\partial \lambda_0}, \dots, \frac{\partial H}{\partial \lambda_r} \right), \quad (13)$$

where Res_{d-1} denotes the homogeneous resultant associated to $r + 1$ homogeneous polynomials of degree $d - 1$ (see [16, Proposition 1.7, Chapter 13]). Moreover, the following universal property is proved in [19] (see [2, Theorem 3.8] for an English concise version). Let $G_0, \dots, G_r \in \mathbb{Z}[u][\lambda_0, \dots, \lambda_r]$ have degree $d - 1$ with generic coefficients u :

$$G_i(u, \lambda) = \sum_{|\alpha|=d_i} u_{i,\alpha} \lambda^\alpha.$$

Denote by I_G the ideal $\langle G_0, \dots, G_r \rangle \subset \mathbb{Z}[u][\lambda_0, \dots, \lambda_r]$ generated by G_0, \dots, G_r . Then, $\text{Res}(G_0, \dots, G_r)$ is a generator of the (generic) projective elimination ideal

$$\pi I_G = (I_G : m^\infty) \cap \mathbb{Z}[u]. \quad (14)$$

In particular, for any variable λ_i and any $N > \sum_i d_i - n$, it holds that

$$\lambda_i^N \text{Res}_{d-1}(G_0, \dots, G_r) \in \langle G_0, \dots, G_r \rangle. \quad (15)$$

Thus, such an equality holds for any specialization of the coefficients u in a ring.

Let A be a non-defective configuration with

$$\text{codim}(\text{sing}(X_A^v)) \geq r.$$

Call $\delta = \deg(D_A)$ and for a choice of A -polynomials p_0, \dots, p_r consider the evaluation

$$D_A(P_\lambda) = D_A\left(\sum_{i=0}^r \lambda_i p_i\right),$$

which is either zero or a homogeneous polynomial in $\lambda = (\lambda_0, \dots, \lambda_r)$ of degree δ .

Proposition 4.5. *Under the hypotheses of Theorem 4.4, the following equality holds:*

$$JD_{r,A} = ID_{r,A}.$$

Moreover, when $\text{codim}_{X_A^v}(\text{sing}(X_A^v)) \geq r$, the degree of the iterated discriminant is equal to

$$\deg(ID_{r,A}) = (r + 1)\delta(\delta - 1)^r. \quad (16)$$

Proof. By Theorem 4.4 and Definition 1.1, we can assume that

$$\text{codim}_{X_A^v}(\text{sing}(X_A^v)) \geq r.$$

Let $(p_0^0, \dots, p_r^0, \lambda^0)$ be a point in the incidence variety Σ defined in (6). Note that for any $i = 0, \dots, r$, we have that

$$0 = \sum_j c_{ij} \frac{\partial D_A}{\partial c_j} \left(\sum_{i=0}^r \lambda_i^0 p_i^0 \right) = \frac{\partial}{\partial \lambda_i} D_A \left(\sum_{i=0}^r \lambda_i p_i^0 \right) (\lambda^0).$$

Then,

$$ID_{r,A} = D_{\delta\Delta_r} \left(D_A \left(\sum_{i=0}^r \lambda_i p_i^0 \right) \right) = 0.$$

As we pointed out in (13), this homogeneous discriminant equals, up to constant, the resultant

$$\text{Res}_{\delta-1} \left(\frac{\partial}{\partial \lambda_0} D_A \left(\sum_{i=0}^r \lambda_i p_i^0 \right), \dots, \frac{\partial}{\partial \lambda_r} D_A \left(\sum_{i=0}^r \lambda_i p_i^0 \right) \right).$$

It then follows that if $D_{\delta\Delta_r}(D_A(\sum_{i=0}^r \lambda_i p_i^0)) = 0$, then there exists $\lambda^0 \in \mathbb{P}^r$ which is a common zero of all these partial derivatives. We deduce from Equation (15) that for any ring R containing the coefficients of $D_A(P_\lambda)$ and for any $i = 0, \dots, r$, the iterated discriminant $D_{\delta\Delta_r}(P_\lambda)$ lies in the ideal generated by the partial derivatives

$$\frac{\partial}{\partial \lambda_j} D_A \left(\sum_{i=0}^r \lambda_i p_i^0 \right), \quad j = 0, \dots, r,$$

in each localization $R[\lambda_0, \dots, \lambda_r]_{\lambda_i}$. Moreover, we have that the ideal πI in (7) is the specialization of the ideal πI_G in (14). Then, $ID_{r,A} = JD_{r,A}$, as claimed.

To see that equation (16) holds, recall that $\deg(D_A) = \delta$ and so the degree of $D_A(\sum_{i=0}^r \lambda_i p_i)$ in the coefficients of p_0, \dots, p_r as well as in the λ variables is equal to δ . On the other side, the degree of $D_{\delta\Delta_r}$ is equal to $(r+1)\delta^r$. ■

4.1. The exponents in Theorem 4.4

We first present two examples that illustrate Theorem 4.4 with $r+1=2$. In the first one, the singular locus has codimension $r+1=2$, which implies a factor (with multiplicity 2) of the iterated discriminant. In the second one, the singular locus has codimension bigger than 2, which implies equality between $MD_{1,A}$ and $ID_{2,A}$.

Example 4.6 (Example 2.3, continued). Consider again the two-dimensional configuration corresponding to the first Hirzebruch surface \mathbb{F}_1 :

$$A = \{(0, 0), (1, 0), (2, 0), (0, 1), (1, 1)\}.$$

Given a generic A -polynomial $f(x, y) = a_0 + a_1x + a_2x^2 + y(b_0 + b_1x)$, we saw that $D_A(f) = a_0b_1^2 - a_1b_0b_1 + a_2b_0^2$. The ideal defining the singular locus S of X_A^y is generated by

$$b_0^2, b_0b_1, b_1^2, -a_1b_1 + 2a_2b_0, 2a_0b_1 - a_1, b_0.$$

This ideal has multiplicity 2 and its radical is generated by b_0, b_1 . In this case, $ID_{2,A}$ has another irreducible factor Ch_S of degree 2 coming from the Chow form of S , to the second power:

$$ID_{2,A} = MD_{1,A} \cdot \text{Ch}_S^2,$$

where $\text{Ch}_S((a_0, a_1, a_2, b_0, b_1), (A_0, A_1, A_2, B_0, B_1)) = B_0b_1 - B_1b_0$.

Example 4.7. Let X_A be the Segre embedding of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, so D_A is the hyperdeterminant of format $(2, 2, 2)$ of degree 4, whose singular locus has codimension greater than 2 by [28]. Take $r = 2$, so that $MD_{1,A}$ equals the discriminant of the hyperdeterminant of format $(2, 2, 2, 2)$ (corresponding to the Segre embedding of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$). In this case, $MD_{1,A}$ equals the iterated discriminant $ID_{2,A}$ and thus has degree $2 \cdot 4 \cdot (4 - 1)^1 = 24$.

This is the only known case of polynomials of degree bigger than 2 for which the iterated and the mixed discriminants coincide.

We have not completely identified the exponents μ_k occurring in the factorization of the iterated discriminant in Theorem 4.4 by the difficulties expressed in the Introduction, where we gave the only references to the literature we are aware of, but the following proposition shows that these exponents are strictly bigger than 1.

Proposition 4.8. *With notation and assumptions as in Theorem 4.4, the exponents μ_k in item (2) satisfy $\mu_k \geq 2$ for all k .*

Proof. To simplify the notation, assume that

$$\text{codim}_{X_A^v}(\text{sing}(X_A^v)) = r = 1, \quad \text{sing}(X_A^v) = Y, \quad JD_{r,A} = MD_{r,A} \text{Ch}_Y^\mu.$$

Considering a generic point (p_0, p_1) such that $MD(p_0, p_1) \neq 0$ and $\text{Ch}_Y(p_0, p_1) = 0$, we have that

$$\begin{aligned} \frac{\partial ID}{\partial c_{i,a}}(p_0, p_1) &= \mu \text{Ch}_Y^{\mu-1}(p_0, p_1) MD_{1,A}(p_0, p_1) \frac{\partial \text{Ch}_Y}{\partial c_{i,a}}(p_0, p_1) \\ &\quad + \text{Ch}_Y^\mu(p_0, p_1) \frac{\partial MD}{\partial c_{i,a}}(p_0, p_1) \end{aligned}$$

for all $a \in A$ and $i = 0, 1$. By genericity, we may assume that $\frac{\partial \text{Ch}_Y}{\partial c_{i,a}}(p_0, p_1) \neq 0$ for some $c_{i,a}$. We can conclude that if $\mu = 1$, then

$$\frac{\partial ID}{\partial c_{i,a}}(p_0, p_1) \neq 0$$

for some $c_{i,a}$, and thus that $\frac{\partial ID}{\partial c_{i,a}}(p_0, p_1) = 0$ for all $c_{i,a}$ would imply $\mu \geq 2$. We will prove that this is the case.

Recall that $\text{Ch}_Y(p_0, p_1) = 0$ implies that the line spanned by p_0, p_1 intersects Y at some point which we denote by $p_0 + \lambda^* p_1$. This means that

$$D_A(p_0 + \lambda^* p_1) = \frac{\partial D_A}{\partial c_a}(p_0 + \lambda^* p_1) = 0$$

for all $a \in A$. Let

$$D_A(p_0 + \lambda p_1) = \sum_0^\delta \Gamma_j(c_{0,a}, c_{1,a})_{a \in A} \lambda^j$$

and recall that $ID_{r,A}(p_0, \dots, p_r) := D_\delta(D_A(p_0 + \lambda p_1))$ by Theorem 4.5. It follows that

$$\frac{\partial ID}{\partial c_{i,a}}(p_0, p_1) = \sum_1^\delta \frac{\partial D_\delta}{\partial \Gamma_j}(D_A(p_0 + \lambda^* p_1)) \frac{\partial \Gamma_j}{\partial c_{i,a}}(p_0, p_1).$$

Observe that

$$\begin{aligned} \frac{\partial D_A(p_0 + \lambda^* p_1)}{\partial c_{0,a}} &= \frac{\partial D_A}{\partial c_a}(p_0 + \lambda^* p_1) = 0, \\ \frac{\partial D_A(p_0 + \lambda^* p_1)}{\partial c_{1,a}} &= \lambda^* \frac{\partial D_A}{\partial c_a}(p_0 + \lambda^* p_1) = 0. \end{aligned}$$

Moreover,

$$0 = \frac{\partial D_A(p_0 + \lambda^* p_1)}{\partial c_{i,a}} = \sum_{j=0}^\delta \frac{\partial \Gamma_j}{\partial c_{i,a}}(p_0, p_1) (\lambda^*)^j.$$

Recall that if $\frac{\partial D_\delta}{\partial \Gamma_j}(D_A(p_0 + \lambda^* p_1)) \neq 0$ for some j , then by biduality

$$\frac{\partial D_\delta}{\partial \Gamma_j}(D_A(p_0 + \lambda^* p_1)) = (\lambda^*)^j,$$

which would conclude the proof. If otherwise, $\frac{\partial D_\delta}{\partial \Gamma_j}(D_A(p_0 + \lambda^* p_1)) = 0$ for all j , then the assertion is also true. \blacksquare

Iterated discriminants with respect to one variable appear frequently in the study and applications of the Cylindrical Algebraic Decomposition proposed by Collins in 1975, and this lead to try to describe the singularities of discriminant hypersurfaces. In particular, the best detailed study is done by Busé and Mourrain in [4, Theorem 6.8, Corollary 6.9] for homogeneous polynomials of three variables (or more, but iterating twice the computation of a discriminant with respect to one of the variables) using resultants, with proofs that cannot be extended for general configurations A .

An interesting subsequent work is the paper by Lazard and McCallum [22]. Again, they consider polynomials f in variables (x, y, z_1, \dots, z_m) and univariate iterated

discriminants in x and y (thinking of f in the ring $k[z_1, \dots, z_m][x, y]$) with rather elementary techniques. They identify the factors of their iterated discriminants but do not identify the exponents in general. However, they prove a series of very nice general and useful results, in particular, Proposition 9 about the regular points of the discriminant (which is a version of biduality), and Propositions 10 through 14 about the singular points, that could be used to identify the exponents μ_k in particular cases.

Based on the computation of different examples (see, for instance, Examples 2.3 and Example 4.6), and the results in [4] and [22] that we mentioned, we see some evidence of the following.

Conjecture 4.9. *The multiplicity μ_k are equal to 2 if Y_k is a component of codimension one corresponding to the closure of the locus of those p for which there are two different non-degenerate multiple roots (the double point locus), while μ_k equals 3 when Y_k is a component of codimension one corresponding to the locus of those p for which there is a degenerate multiple root (the cusp locus).*

5. Comparing mixed and iterated discriminants

In this section we consider the case when A equals the lattice points in a cartesian product of dilates of standard simplices: $d_1 \Delta_{k_1} \times \dots \times d_\ell \Delta_{k_\ell}$, for some $\ell \geq 1$. In other words we investigate Segre–Veronese varieties $X_A = \mathbb{P}^{k_1}(d_1) \times \dots \times \mathbb{P}^{k_\ell}(d_\ell)$.

The symbol $\mathbb{P}^k(d)$ denotes the Veronese embedding of degree d in dimension k , i.e., the variety \mathbb{P}^k embedded in $\mathbb{P}^{\binom{k+d}{d}-1}$ by the global sections of the line bundle $\mathcal{O}_{\mathbb{P}^k}(d)$. We occasionally denote $\mathbb{P}^k(1)$ by \mathbb{P}^k .

The symbol $\mathbb{P}^{k_1}(d_1) \times \dots \times \mathbb{P}^{k_\ell}(d_\ell)$ denotes the Segre embedding of the above defined Veronese embeddings, more precisely the variety $\mathbb{P}^{k_1} \times \dots \times \mathbb{P}^{k_\ell}$ embedded via the global sections of the line bundle $\pi_1^* \mathcal{O}_{\mathbb{P}^{k_1}}(d_1) \otimes \dots \otimes \pi_\ell^* \mathcal{O}_{\mathbb{P}^{k_\ell}}(d_\ell)$, where π_i denotes the i th projection $\pi_i: \mathbb{P}^{k_1} \times \dots \times \mathbb{P}^{k_\ell} \rightarrow \mathbb{P}^{k_i}$. These are toric embeddings corresponding to the configurations of lattice points of the polytopes $d_1 \Delta_{k_1} \times \dots \times d_\ell \Delta_{k_\ell}$.

When $d_i = 1$ we recover the case of hyperdeterminants, which has been completely solved in [28]. In Proposition 5.2, we show that when $\ell = 1$ there is equality if and only if $r = 1$ and $d_1 = 2$. We then conjecture that these are all the possible cases (see Conjecture 5.3), that is, in all other cases the singularities of the discriminantal locus have codimension one in the dual variety. We conclude with Theorem 5.6, which covers the case in which all $d_i > 1$.

To determine when the iterated and mixed discriminants of Segre–Veronese varieties are equal, we start with the following lemma, which allows us to compute the degree of MD_{r,d,Δ_n} , that is, the case in which we consider $(r + 1)$ polynomials of

degree d in n variables. Recall that when $r \leq n$, we know by Proposition 3.3 that the mixed discriminant equals the discriminant of the Cayley configuration given by the lattice points in the product of simplices $\Delta_r \times d\Delta_n$.

Lemma 5.1. *If $r \leq n$, then $\deg(MD_{r,d\Delta_n}) = (n+1)\binom{n}{r}d^r(d-1)^{n-r}$.*

Proof. We will use [16, Theorem 2.4, Chapter 13], which tells us that this degree is equal to the coefficient of the monomial $x^r y^n$ in the expansion of

$$\begin{aligned} S(x, y) &= \frac{1}{((1+x)(1+y) - x(1+y) - dy(1+x))^2} \\ &= \frac{1}{(1 - (d-1)y - dxy)^2}. \end{aligned}$$

We may write

$$S(x, y) = \left(\frac{1}{1-q}\right)^2 = \sum_{n \geq 0} (n+1)q^n,$$

where $q = (d-1)y + dxy = y((d-1) + dx)$.

Since

$$q^n = \left(\sum_{j=0}^n \binom{n}{j} (d-1)^{n-j} d^j x^j\right) y^n,$$

we have

$$S(x, y) = \sum_{n \geq 0} \sum_{j=0}^n (n+1) \binom{n}{j} (d-1)^{n-j} d^j x^j y^n. \quad (17)$$

From this expansion, we see that the coefficient of $x^r y^n$ is equal to

$$(n+1) \binom{n}{r} (d-1)^{n-r} d^r$$

when $r \leq n$, and is equal to 0 if $r > n$. This completes the proof. \blacksquare

Proposition 5.2. *Let $1 < d$ and $1 \leq r \leq n$. Then $MD_{r,d\Delta_n} = ID_{r,d\Delta_n}$ if and only if $r = 1$ and $d = 2$.*

The fact that this equality holds in the case of $r = 1$ and $d = 2$ was shown in [24, Theorem 8.2]. Although we include this in our proof for completeness, the main contribution of this result is that equality does not hold in any other case.

Proof. For any $d > 1$, the configuration of lattice points in $d\Delta_n$ is non-defective [3] and as $r \leq n$, it is enough to check that

$$\deg(MD_{r,d\Delta_n}) = \deg(ID_{r,d\Delta_n})$$

by Propositions 3.3 and 3.4.

From Lemma 5.1, we know that

$$\deg(MD_{r,d\Delta_n}) = (n+1) \binom{n}{r} d^r (d-1)^{n-r}.$$

By Proposition 4.5, we also know that

$$\deg(ID_{r,d\Delta_n}) = (n+1)(d-1)^n (r+1)((n+1)(d-1)^n - 1)^r.$$

To determine when these are equal, we will consider the ratio of the two degrees, both of which are non-zero for $d > 1$. We have

$$\begin{aligned} \frac{\deg(ID_{r,d\Delta_n})}{\deg(MD_{r,d\Delta_n})} &= \frac{(n+1)(d-1)^n (r+1)((n+1)(d-1)^n - 1)^r}{(n+1) \binom{n}{r} d^r (d-1)^{n-r}} \\ &= \frac{(d-1)^r (r+1)((n+1)(d-1)^n - 1)^r}{\binom{n}{r} d^r}. \end{aligned}$$

For any $d > 1$, we have

$$(n+1)(d-1)^n - 1 = n(d-1)^n + (d-1)^n - 1 \geq n(d-1)^n,$$

with equality if and only if $d = 2$. Thus the numerator satisfies

$$\begin{aligned} (d-1)^r (r+1)((n+1)(d-1)^n - 1)^r &\geq (d-1)^r (r+1)(n(d-1)^n)^r \\ &= (d-1)^{r(n+1)} (r+1)n^r. \end{aligned}$$

Since $\binom{n}{r} \leq \frac{n^r}{r!}$, we have

$$\begin{aligned} \frac{\deg(ID_{r,d\Delta_n})}{\deg(MD_{r,d\Delta_n})} &\geq \frac{(d-1)^{r(n+1)} (r+1)n^r}{\frac{n^r}{r!} d^r} \\ &= \frac{(d-1)^{r(n+1)} (r+1)!}{d^r} = \left(\frac{(d-1)^{n+1}}{d} \right)^r \cdot (r+1)!. \end{aligned}$$

If $d = 2$, then this ratio is

$$\frac{(r+1)!}{2^r} = \frac{(r+1)!}{2^r} \geq 1,$$

with equality if and only if $r = 1$. If $d > 2$, then

$$(d-1)^{n+1} \geq (d-1)^2 \geq 2(d-1) = 2d-2 > d.$$

Thus

$$\frac{(d-1)^{n+1}}{d} > 1,$$

and so

$$\frac{\deg(ID_{r,d\Delta_n})}{\deg(MD_{r,d\Delta_n})} > (r+1)!.$$

Thus, except possibly in the case of $d = 2$ and $r = 1$, we have

$$\deg(ID_{r,d\Delta_n}) > \deg(MD_{r,d\Delta_n}).$$

To see that $d = 2$ and $r = 1$ gives $\deg(MD_{r,d\Delta_n}) = \deg(ID_{r,d\Delta_n})$, note that in this case the ratio of the degrees is

$$\frac{(2-1)^1(1+1)((n+1)(2-1)^n-1)^1}{\binom{n}{1}2^1} = \frac{2(n+1-1)}{2n} = 1. \quad \blacksquare$$

Geometrically, Proposition 5.2 shows that for $\mathbb{P}^r(1) \times \mathbb{P}^n(d)$ the associated mixed discriminant is equal to the iterated discriminant only when $r = 1$ and $d = 2$. Note that we do not consider the case $d = 1$ because this case is defective.

It is natural to consider the same question for any product-of-simplices

$$\mathbb{P}^r(1) \times \mathbb{P}^{k_1}(d_1) \times \cdots \times \mathbb{P}^{k_\ell}(d_\ell).$$

Setting $d = (d_1, \dots, d_\ell)$ and $k = (k_1, \dots, k_\ell)$, let $A_{\ell,d,k}$ denote the configuration corresponding to $\mathbb{P}^{k_1}(d_1) \times \cdots \times \mathbb{P}^{k_\ell}(d_\ell)$.

We conjecture the following:

Conjecture 5.3. *We have $ID_{r,A_{\ell,d,k}} = MD_{r,A_{\ell,d,k}}$ if and only if*

$$\mathbb{P}^r(1) \times \mathbb{P}^{k_1}(d_1) \times \cdots \times \mathbb{P}^{k_\ell}(d_\ell)$$

is of one of the following forms:

- (1) $\mathbb{P}^r \times \mathbb{P}^m \times \mathbb{P}^m$, $m \geq 1$, $r = 1, 2$,
- (2) $(\mathbb{P}^1)^4$,
- (3) $\mathbb{P}^1 \times \mathbb{P}^n(2)$.

This conjecture was inspired by the question posed in [16, Chapter 14, p. 479], which coincides with the above conjecture when $d_i = 1$ for all i . Their conjecture (and thus our conjecture in this special case) was proved in [28]. Note that Proposition 5.2 implies that Conjecture 5.3 is true when $\ell = 1$, which puts us into case (3).

To study our conjecture in general, the following theorem giving the degree of the mixed discriminant $MD_{r,A_{\ell,d,k}}$ will be useful. Let $B = B_\ell$ be the set of all non-empty subsets $\Omega \subset \{0, 1, \dots, \ell\}$. For each $\Omega \in B$, let

$$d_\Omega = \sum_{j \in \Omega} d_j.$$

Let $\delta(\Omega) \in \mathbb{Z}_+^{\ell+1}$ be the characteristic vector of Ω . For every $\kappa = (r, k_1, \dots, k_\ell) \in \mathbb{Z}_+^{r+1}$, let $\mathcal{P}(\kappa)$ denote the set of all partitions of κ into a sum of vectors $\delta(\Omega)$; in other words, $\mathcal{P}(\kappa)$ is the set of all non-negative integral vectors $(m_\Omega)_{\Omega \in B}$ such that

$$\sum_{\Omega \in B} m_\Omega \delta(\Omega) = \kappa.$$

Theorem 5.4 ([16, Theorem 13.2.5]). *The degree of $MD_{r,A_\ell,d,k}$ is given by*

$$\sum_{(m_\Omega) \in \mathcal{P}(\kappa)} \left(1 + \sum_{\Omega \in B} m_\Omega\right)! \prod_{\Omega \in B} \frac{(d_\Omega - 1)^{m_\Omega}}{m_\Omega!}.$$

Note that any partition using the vector $\delta(\{0\}) = (1, 0, \dots, 0)$ will not contribute to this sum, since $d_{\{0\}} - 1 = 0$. Letting C be the set of all non-empty subsets of $\{1, \dots, \ell\}$ and letting $k = (k_1, \dots, k_\ell)$ as before, we have that

$$\deg(ID_{r,A_\ell,d,k}) = (r + 1)\delta(\delta - 1)^r,$$

where

$$\delta = \sum_{(m_\Omega) \in \mathcal{P}(k)} \left(1 + \sum_{\Omega \in C} m_\Omega\right)! \prod_{\Omega \in C} \frac{(d_\Omega - 1)^{m_\Omega}}{m_\Omega!}.$$

When it is clear from context, we will abbreviate $\deg(MD_{r,A_\ell,d,k})$ as $\deg(MD)$ and $\deg(ID_{r,A_\ell,d,k})$ as $\deg(ID)$.

Example 5.5. Let us compare the degrees of the mixed and the iterated discriminant when $r = 1$, $\ell = 2$, and $k_1 = k_2 = 1$. To compute the degree of the mixed discriminant, we consider all partitions of $\kappa = (1, 1, 1)$. We may discount any partition with the vector $(1, 0, 0)$, as this partition would contribute a term of 0 to $\deg(MD)$. Thus, the only relevant partitions are

- $(1, 1, 1)$,
- $(1, 1, 0) + (0, 0, 1)$, and
- $(1, 0, 1) + (0, 1, 0)$.

The contributions from these terms to $\deg(MD)$ are

- $2! \cdot \frac{(d_1+d_2)^1}{1!} = 2(d_1 + d_2)$,
- $3! \cdot \frac{d_1^1 \cdot (d_2-1)^1}{1! \cdot 1!} = 6d_1(d_2 - 1)$, and
- $3! \cdot \frac{d_2^1 \cdot (d_1-1)^1}{1! \cdot 1!} = 6d_2(d_1 - 1)$,

respectively. (Note that some of these contributions will be zero if one or both of d_1 and d_2 are equal to 1.) Adding these gives

$$\begin{aligned} \deg(MD) &= 2(d_1 + d_2) + 6d_1(d_2 - 1) + 6d_2(d_1 - 1) \\ &= 12d_1d_2 - 4d_1 - 4d_2 = 4(3d_1d_2 - d_1 - d_2). \end{aligned}$$

To compute $\deg(ID)$, we must consider the partitions of $k = (1, 1)$. There are only two: $(1, 1)$ and $(1, 0) + (0, 1)$. The contributions of these to δ are

- $2! \cdot \frac{(d_1+d_2-1)!}{1!} = 2(d_1 + d_2 - 1)$ and
- $3! \cdot \frac{(d_1-1)!(d_2-1)!}{1! \cdot 1!} = 6(d_1 - 1)(d_2 - 1)$,

respectively. Thus

$$\delta = 2(d_1 + d_2 - 1) + 6(d_1 - 1)(d_2 - 1) = 6d_1d_2 - 4d_1 - 4d_2 + 4.$$

It follows that

$$\deg(ID) = 2\delta(\delta - 1) = 2(6d_1d_2 - 4d_1 - 4d_2 + 4)(6d_1d_2 - 4d_1 - 4d_2 + 3).$$

We will now argue that $\deg(ID) > \deg(MD)$, unless $d_1 = d_2 = 1$. First we perform the change of variables $d_1 = d'_1 + 1$ and $d_2 = d'_2 + 1$, to remove some of the negatives. This gives

$$\deg(MD) = 4(3d'_1d'_2 + 2d'_1 + 2d'_2 + 1)$$

and

$$\deg(ID) = 2(6d'_1d'_2 + 2d'_1 + 2d'_2 + 2)(6d'_1d'_2 + 2d'_1 + 2d'_2 + 1).$$

Now, if either d_1 or d_2 is greater than 1, then $(6d'_1d'_2 + 2d'_1 + 2d'_2 + 1)$ is at least 3, meaning that

$$\deg(ID) \geq 6(6d'_1d'_2 + 2d'_1 + 2d'_2 + 2) = 36d'_1d'_2 + 12d'_1 + 12d'_2 + 12.$$

This is certainly greater than

$$\deg(MD) = 12d'_1d'_2 + 8d'_1 + 8d'_2 + 4,$$

since d'_1 and d'_2 are non-negative. So, in this case $\deg(ID) > \deg(MD)$. If we do have $d_1 = d_2 = 1$, then $\deg(MD) = 4 = \deg(ID)$. This equality was predicted by case (1) of Conjecture 5.3.

We will now prove that Conjecture 5.3 holds in the case that $r = 1$ and $d_i > 1$ for all i .

Theorem 5.6. *Suppose $d_i > 1$ for all i . Then the only case where $MD_{1,A_{\ell,d,k}} = ID_{1,A_{\ell,d,k}}$ is when $\ell = 1$ and $d_1 = 2$.*

Proof. This proposition holds when $\ell = 1$ by Proposition 5.2, and when $\ell = 2$ and $k_1 = k_2 = 1$ by Example 5.5. Thus it suffices to prove that we have

$$\deg(MD_{1,A_{\ell,d,k}}) < \deg(ID_{1,A_{\ell,d,k}})$$

for $\ell = 2$ with $(k_1, k_2) \neq (1, 1)$, and for $\ell \geq 3$.

First we consider how partitions of $\kappa = (1, k_1, \dots, k_\ell)$ relate to partitions of $k = (k_1, \dots, k_\ell)$. Each partition of κ gives rise to a partition of k simply by deleting the first coordinate and grouping together vectors that are now identical. Note that no partition (m_Ω) contributing to $\deg(MD)$ uses the vector $(1, 0, \dots, 0)$. Also, exactly one vector in each partition of κ is of the form $(1, *, \dots, *)$. Call the support of this vector $\Psi((m_\Omega))$, or simply Ψ when the context is clear. Note that $m_\Psi = 1$. Let Ξ denote $\Psi \setminus \{0\}$. Isolating Ψ and Ξ , we may write

$$\begin{aligned} \deg(MD) &= \sum_{(m_\Omega) \in \mathcal{P}(\kappa)} (1 + \sum_{\Omega \in B} m_\Omega)! \prod_{\Omega \in B} \frac{(d_\Omega - 1)^{m_\Omega}}{m_\Omega!} \\ &= \sum_{(m_\Omega) \in \mathcal{P}(\kappa)} (1 + \sum_{\Omega \in B} m_\Omega)! \cdot \frac{(d_\Psi - 1)^{m_\Psi}}{m_\Psi!} \cdot \frac{(d_\Xi - 1)^{m_\Xi}}{m_\Xi!} \prod_{\Omega \in B, \Omega \neq \Psi, \Xi} \frac{(d_\Omega - 1)^{m_\Omega}}{m_\Omega!} \\ &= \sum_{(m_\Omega) \in \mathcal{P}(\kappa)} (1 + \sum_{\Omega \in B} m_\Omega)! \cdot (d_\Psi - 1) \cdot \frac{(d_\Xi - 1)^{m_\Xi}}{m_\Xi!} \prod_{\Omega \in B, \Omega \neq \Psi, \Xi} \frac{(d_\Omega - 1)^{m_\Omega}}{m_\Omega!}. \end{aligned}$$

Given (m_Ω) a partition of κ , let (n_Ω) be the corresponding partition of k . So, if the term in $\deg(MD)$ coming from (m_Ω) is

$$(1 + \sum_{\Omega \in B} m_\Omega)! \cdot (d_\Psi - 1) \cdot \frac{(d_\Xi - 1)^{m_\Xi}}{m_\Xi!} \prod_{\Omega \in B, \Omega \neq \Psi, \Xi} \frac{(d_\Omega - 1)^{m_\Omega}}{m_\Omega!},$$

then the term in δ coming from (n_Ω) is

$$(1 + \sum_{\Omega \in B} m_\Omega)! \cdot \frac{(d_\Xi - 1)^{m_\Xi + 1}}{(m_\Xi + 1)!} \prod_{\Omega \in B, \Omega \neq \Psi, \Xi} \frac{(d_\Omega - 1)^{m_\Omega}}{m_\Omega!}.$$

Note that $d_\Xi = d_\Psi - 1$. We know that $d_\Xi \neq 1$ by our assumption that $d_i > 1$ for all i , so the change in factor between these two contributions is

$$\frac{d_\Xi - 1}{d_\Xi(m_\Xi + 1)}.$$

Since $d_\Xi > 1$, this is at least $\frac{1}{2(m_\Xi + 1)}$. In general, $m_\Omega \leq \max\{k_i\}$ for any Ω ; since the vector $\delta(\Psi)$ also appears in the partition of κ , we in fact have

$$m_\Xi \leq \max\{k_i\} - 1.$$

So, $m_\Xi + 1 \leq \max\{k_i\}$. It follows that $\frac{1}{2(m_\Xi + 1)}$ is greater than or equal to $\frac{1}{2 \max\{k_i\}}$. So, passing from a partition of κ to a partition of k , the corresponding term in δ is at least $\frac{1}{2 \max\{k_i\}}$ times the corresponding term in $\deg(MD)$.

Now we consider how many partitions of κ give rise to the same partition of k . Given a partition (n_Ω) of k , all relevant partitions of κ that map to it can be constructed

by choosing a single vector used in (n_Ω) , and appending a 1 to the 0th coordinate. Thus, the number of partitions of κ mapping to (n_Ω) is equal to the number of distinct vectors used in (n_Ω) . The number of distinct vectors in this partition can be bounded by $k_1 + \dots + k_\ell$, since this is the total sum of all the entries of all the vectors used. Thus, we have that

$$\delta \geq \frac{1}{2 \max\{k_i\}(k_1 + \dots + k_\ell)} \deg(MD).$$

It follows that

$$\begin{aligned} \deg(ID) &= 2\delta(\delta - 1) \geq 2 \frac{1}{2 \max\{k_i\}(k_1 + \dots + k_\ell)} \deg(MD) \cdot (\delta - 1) \\ &= \frac{\delta - 1}{\max\{k_i\}(k_1 + \dots + k_\ell)} \cdot \deg(MD). \end{aligned}$$

To show that $\deg(ID) > \deg(MD)$, it remains to show that

$$\delta - 1 > \max\{k_i\}(k_1 + \dots + k_\ell).$$

First, rewrite

$$\begin{aligned} \delta &= \sum_{(m_\Omega) \in \mathcal{P}(k)} (1 + \sum_{\Omega \in C} m_\Omega)! \prod_{\Omega \in C} \frac{(d_\Omega - 1)^{m_\Omega}}{m_\Omega!} \\ &= \sum_{(m_\Omega) \in \mathcal{P}(k)} (1 + \sum_{\Omega \in C} m_\Omega) \cdot \frac{(\sum_{\Omega \in C} m_\Omega)!}{\prod_{\Omega \in C} m_\Omega!} \prod_{\Omega \in C} (d_\Omega - 1)^{m_\Omega} \\ &= \sum_{(m_\Omega) \in \mathcal{P}(k)} (1 + \sum_{\Omega \in C} m_\Omega) \cdot \binom{\sum_{\Omega \in C} m_\Omega}{m_{\Omega_1}, \dots, m_{\Omega_t}} \prod_{\Omega \in C} (d_\Omega - 1)^{m_\Omega}. \end{aligned}$$

For any partition of k , we have that $\sum_{\Omega \in C} m_\Omega \geq \max\{k_i\}$, since k_i vectors (counted with multiplicity) must have non-zero i th coordinate. Moreover, a multinomial coefficient

$$\binom{a_1 + a_2 + \dots + a_s}{a_1, a_2, \dots, a_s}$$

can be rewritten as the product

$$\binom{a_1}{a_1} \binom{a_1 + a_2}{a_2} \dots \binom{a_1 + a_2 + \dots + a_s}{a_s},$$

so it is at least as large as

$$\binom{a_1 + a_2 + \dots + a_s}{a_s}.$$

Of course, we may reorder the a_i 's in any way we desire. So, as long as some a_i satisfies $0 < a_i < a_1 + \cdots + a_s$, we have

$$\begin{aligned} \binom{a_1 + a_2 + \cdots + a_s}{a_1, a_2, \dots, a_s} &\geq \binom{a_1 + a_2 + \cdots + a_s}{a_i} \\ &\geq \binom{a_1 + a_2 + \cdots + a_s}{1} = (a_1 + \cdots + a_s). \end{aligned}$$

This means that if (m_Ω) is a partition of k that uses at least two different vectors, we have

$$\binom{\sum_{\Omega \in C} m_\Omega}{m_{\Omega_1}, \dots, m_{\Omega_t}} \geq \sum_{\Omega \in C} m_\Omega \geq \max\{k_i\}.$$

Finally, the product $\prod_{\Omega \in C} (d_\Omega - 1)^{m_\Omega}$ is greater than or equal to 1. Thus, every partition of k that uses at least distinct two vectors contributes at least $(1 + \max\{k_i\}) \cdot \max\{k_i\}$ to δ . We will now argue that there are at least ℓ such partitions of k .

To do this, we split into two cases: where $\ell = 2$, and where $\ell \geq 3$. If $\ell = 2$ and $(k_1, k_2) \neq (1, 1)$, then there are indeed at least two such partitions of $k = (k_1, k_2)$. For instance, we could use $(1, 1) + (k_1 - 1)(1, 0) + (k_2 - 1)(0, 1)$ and $k_1(1, 0) + k_2(0, 1)$. Both do indeed use at least two distinct vectors since at least one of $k_1 - 1$ and $k_2 - 1$ is non-zero.

Assume now $\ell \geq 3$. We can construct a partition of k that uses at least two vectors by choosing any $0 - 1$ vector with support size at least 2 and at most $\ell - 1$, and then completing the partition by using standard basis vectors. The condition on the support size guarantees that at least one other vector will be used, and that this new standard basis vector has not already been used. There are $2^\ell - 2 - \ell$ such initial vectors, which is greater than or equal to ℓ since $\ell \geq 3$. Thus, at least ℓ partitions of k contribute at least $(1 + \max\{k_i\}) \cdot \max\{k_i\}$ to δ .

Note that $\ell \max\{k_1\} \geq k_1 + \cdots + k_\ell$. It follows that

$$\begin{aligned} \delta &\geq \ell(1 + \max\{k_i\}) \cdot \max\{k_i\} \geq \ell \max\{k_i\} \cdot \max\{k_i\} + \ell \\ &> \ell \max\{k_i\} \cdot \max\{k_i\} + 1 \geq \max\{k_i\}(k_1 + \cdots + k_\ell) + 1. \end{aligned}$$

Equivalently,

$$\delta - 1 > \max\{k_i\}(k_1 + \cdots + k_\ell).$$

This implies that $\deg(ID) > \deg(MD)$, as desired. ■

6. Curves in the plane

In this section we will determine when the mixed and iterated discriminants associated to a planar configuration are equal. Let $A = P \cap \mathbb{Z}^2$, where P is a smooth lattice

polygon of dimension 2. Let v_A , p_A , and V_A denote the normalized area $\text{area}_{\mathbb{Z}}(P)$ (that is, twice its Euclidean area), the lattice perimeter (that is, the number of points in A on the edges of P), and the number of vertices of P , respectively. It is well known [16] that in this smooth case the degree δ_A of D_A equals

$$\delta_A = 3v_A - 2p_A + V_A.$$

The degree of the mixed discriminant can be computed from [6, Corollary 3.15] as

$$\begin{aligned} \deg(MD(A, A)) &= 2(\text{area}_{\mathbb{Z}}(2P) - \text{area}_{\mathbb{Z}}(P) - p_A) \\ &= 2(4v_A - v_A - p_A) = 6v_A - 2p_A. \end{aligned}$$

We can reformulate these equations in terms of the number of interior lattice points of P . Let i_A denote the number of interior lattice points of P . Then we know by Pick's theorem that

$$v_A = 2i_A + p_A - 2,$$

which can be rewritten as $v_A - p_A = 2i_A - 2$. This allows us to write

$$\begin{aligned} \deg(MD(A, A)) &= 6v_A - 2p_A = 4v_A + 2(v_A - p_A) \\ &= 4v_A + 4i_A - 4 = 4(v_A + i_A - 1). \end{aligned}$$

and

$$\begin{aligned} \delta_A &= 3v_A - 2p_A + V_A \\ &= v_A + 2(v_A - p_A) + V_A = v_A + 4(i_A - 1) + V_A. \end{aligned}$$

Example 6.1. Let $A = ((0, 0), (2, 0), (0, 2))$. Let us verify that $\deg(MD(A, A)) = \deg(ID_{1,A})$, as implied by Proposition 5.2. We have $v_A = 4$, $i_A = 0$, and $V_A = 3$. This gives us

$$\deg(MD(A, A)) = 4(v_A + i_A - 1) = 4(4 - 0 - 1) = 12$$

and

$$\delta_A = v_A + 4(i_A - 1) + V_A = 4 + 4(0 - 1) + 3 = 3.$$

This means that $\deg(ID_{1,A}) = 2\delta_A(\delta_A - 1) = 2 \cdot 3 \cdot 2 = 12 = \deg(MD(A, A))$.

Example 6.2. Assume $A = \text{conv}((0, 0), (1, 0), (0, 1), (1, 1))$. Let us verify that

$$\deg(MD(A, A)) = \deg(ID_{1,A}),$$

as implied by Example 2.1. We have $v_A = 2$, $i_A = 0$, and $V_A = 4$. This gives us

$$\deg(MD(A, A)) = 4(v_A + i_A - 1) = 4(2 - 0 - 1) = 4$$

and

$$\delta_A = v_A + 4(i_A - 1) + V_A = 2 + 4(0 - 1) + 4 = 2.$$

This means that $\deg(ID_{1,A}) = 2\delta_A(\delta_A - 1) = 2 \cdot 2 \cdot 1 = 4 = \deg(MD(A, A))$.

It turns out that these two examples are the only smooth polygons P where the iterated and the mixed discriminants associated to the configuration of lattice points in P coincide.

Theorem 6.3. *The only smooth polygons P with an associated discriminant without singularities in codimension bigger than 1 are the known cases of the triangle $2\Delta_2$ and the unit square.*

Proof. Assume P is such a polygon, $A = P \cap \mathbb{Z}^2$ and $\delta_A = \deg(D_A)$. Using our formulas for $MD(A, A)$ and δ_A , we have that

$$4(v_A + i_A - 1) = 2(v_A + 4(i_A - 1) + V_A)(v_A + 4(i_A - 1) + V_A - 1),$$

which is equivalent to

$$2(v_A + i_A - 1) = (v_A + 4(i_A - 1) + V_A)(v_A + 4(i_A - 1) + V_A - 1).$$

Suppose for the sake of contradiction that $i_A > 0$. Then

$$v_A + i_A - 1 \leq v_A + 4(i_A - 1) < v_A + 4(i_A - 1) + V_A.$$

Now, if a, b, c, d are positive real numbers with $ab = cd$, then $b < c$ implies $a > d$. This means that

$$2 > v_A + 4(i_A - 1) + V_A - 1 \geq v_A + V_A - 1 \geq v_A + 2.$$

In other words, $v_A < 0$, a contradiction. Thus we know that $i_A = 0$.

Setting $i_A = 0$ reduces our equation to

$$2(v_A - 1) = (v_A + V_A - 4)(v_A + V_A - 5).$$

By a classification result due to [21] and presented again in [5], all convex lattice polygons with no interior lattice points are equivalent to either the triangle

$$2\Delta_2 = \text{conv}((0, 0), (2, 0), (0, 2)),$$

or to a polygon of the form

$$\text{conv}((0, 0), (0, 1), (a, 0), (b, 1)),$$

where $a \geq b \geq 0$ and $a \geq 1$. These polygons are illustrated in Figure 2. All of these polygons have either three or four vertices. So, we must have $V_A = 3$ or $V_A = 4$.




Figure 2. All lattice polygons with no interior lattice points.

If $V_A = 3$, our equation becomes

$$2(v_A - 1) = (v_A - 1)(v_A - 2).$$

This means that either $v_A = 1$, or $2 = v_A - 2$; that is, $v_A = 1$ or $v_A = 4$. If $v_A = 1$, the only possibility for P is the primitive lattice triangle of normalized 1; but this gives a degenerate system, and so is removed from our consideration. If $v_A = 4$, the only possibilities of P are

$$\text{conv}((0, 0), (2, 0), (0, 2)) \quad \text{and} \quad \text{conv}((0, 0), (4, 0), (0, 1)).$$

The second polygon is not smooth, so the only possible triangle is

$$\text{conv}((0, 0), (2, 0), (0, 2)).$$

If $V_A = 4$, our equation becomes

$$2(v_A - 1) = v_A(v_A - 1).$$

This means that either $v_A = 1$ (which is impossible with $V_A = 4$), or that $v_A = 2$. The only polygon with 4 vertices and area 2 is the square

$$\text{conv}((0, 0), (1, 0), (0, 1), (1, 1)).$$

Thus we have shown that

$$\text{conv}((0, 0), (2, 0), (0, 2)) \quad \text{and} \quad \text{conv}((0, 0), (1, 0), (0, 1), (1, 1))$$

are the only possibilities for P . Having already verified that

$$\deg(MD(A, A)) = 2\delta_A(\delta_A - 1)$$

for both of these polygons, this completes the proof. ■

Acknowledgments. We thank Frédéric Bihan, Laurent Busé, Carlos D’Andrea, Bernard Mourrain, and Giorgio Ottaviani for helpful discussions and references to previous work in this direction.

Funding. The first author acknowledges the support of UBACYT 20020170100048-BA and CONICET PIP 11220200100182, Argentina. The second author acknowledges supported by KTH and Williams College. The third author acknowledges support by ICERM and VR grants NT:2014-4763, NT:2018-03688. All three authors acknowledge support by the Knut and Alice Wallenberg foundation.

References

- [1] T. J. I. Bromwich, *Quadratic forms and their classification by means of invariant-factors*. Cambridge Tracts in Math. 3, Hafner Publishing, New York, 1971 Zbl 37.0134.01 MR 0349723
- [2] L. Busé, Elimination theory in codimension one and applications. Notes of lectures given at the CIMPA-UNESCO-IRAN school in Zanjan, Iran, July 9–22, 2005, <https://hal.inria.fr/inria-00077120>.
- [3] L. Busé and J.-P. Jouanolou, [On the discriminant scheme of homogeneous polynomials](#). *Math. Comput. Sci.* **8** (2014), no. 2, 175–234 Zbl 1302.13028 MR 3224627
- [4] L. Busé and B. Mourrain, [Explicit factors of some iterated resultants and discriminants](#). *Math. Comp.* **78** (2009), no. 265, 345–386 Zbl 1200.13042 MR 2448711
- [5] W. Castryck, [Moving out the edges of a lattice polygon](#). *Discrete Comput. Geom.* **47** (2012), no. 3, 496–518 Zbl 1237.52002 MR 2891244
- [6] E. Cattani, M. A. Cueto, A. Dickenstein, S. Di Rocco, and B. Sturmfels, [Mixed discriminants](#). *Math. Z.* **274** (2013), no. 3–4, 761–778 Zbl 1273.13051 MR 3078246
- [7] A. Cayley, On the theory of linear transformations. *Cambridge. Mat. J.* **4** (1845), no. 23, 193–209
- [8] S. Di Rocco, [Linear toric fibrations](#). In *Combinatorial algebraic geometry*, pp. 119–147, Lecture Notes in Math. 2108, Springer, Cham, 2014 Zbl 1326.14002 MR 3329088
- [9] A. Dickenstein, I. Z. Emiris, and A. Karasoulou, [Plane mixed discriminants and toric Jacobians](#). In *SAGA—Advances in ShApes, Geometry, and Algebra*, pp. 105–121, Geom. Comput. 10, Springer, Cham, 2014 Zbl 1331.65047 MR 3289657
- [10] A. Dickenstein, E. M. Feichtner, and B. Sturmfels, [Tropical discriminants](#). *J. Amer. Math. Soc.* **20** (2007), no. 4, 1111–1133 Zbl 1166.14033 MR 2328718
- [11] A. Dickenstein, M. I. Herrero, and L. F. Tabera, [Arithmetics and combinatorics of tropical Severi varieties of univariate polynomials](#). *Israel J. Math.* **221** (2017), no. 2, 741–777 Zbl 1401.14236 MR 3704934
- [12] A. Dickenstein, J. M. Rojas, K. Rusek, and J. Shih, [Extremal real algebraic geometry and \$\mathcal{A}\$ -discriminants](#). *Mosc. Math. J.* **7** (2007), no. 3, 425–452, 574 Zbl 1137.14044 MR 2343141

- [13] A. Esterov, [Newton polyhedra of discriminants of projections](#). *Discrete Comput. Geom.* **44** (2010), no. 1, 96–148 Zbl [1273.14105](#) MR [2639821](#)
- [14] A. Esterov, [Characteristic classes of affine varieties and Plücker formulas for affine morphisms](#). *J. Eur. Math. Soc. (JEMS)* **20** (2018), no. 1, 15–59 Zbl [1472.14006](#) MR [3743235](#)
- [15] R. T. Farouki, C. Neff, and M. A. O’Conner, Automatic parsing of degenerate quadric-surface intersections. *ACM Trans. Graph.* **8** (1989), no. 3, 174–203 Zbl [0746.68093](#)
- [16] I. M. Gel’fand, M. M. Kapranov, and A. V. Zelevinsky, [Discriminants, resultants, and multidimensional determinants](#). Math. Theory Appl., Birkhäuser, Boston, MA, 1994 Zbl [0827.14036](#) MR [1264417](#)
- [17] D. R. Grayson and M. E. Stillman, Macaulay2, a software system for research in algebraic geometry, <http://www.math.uiuc.edu/Macaulay2/>.
- [18] E. Gross, H. A. Harrington, Z. Rosen, and B. Sturmfels, [Algebraic systems biology: a case study for the Wnt pathway](#). *Bull. Math. Biol.* **78** (2016), no. 1, 21–51 Zbl [1356.92038](#) MR [3452313](#)
- [19] J.-P. Jouanolou, [Le formalisme du résultant](#). *Adv. Math.* **90** (1991), no. 2, 117–263 Zbl [0747.13007](#) MR [1142904](#)
- [20] N. M. Katz, [Pinceaux de Lefschetz: Theoreme d’existence](#). In *Groupes de Monodromie en Géométrie Algébrique*, pp. 212–253, Lect. Notes Math. 340, Springer, Berlin-Heidelberg, 1973. Zbl [0284.14006](#)
- [21] R. Koelman, The number of moduli of families of curves on a toric surface. PhD thesis, Katholieke Universiteit de Nijmegen, 1991.
- [22] D. Lazard and S. McCallum, [Iterated discriminants](#). *J. Symbolic Comput.* **44** (2009), no. 9, 1176–1193 Zbl [1206.13030](#) MR [2532164](#)
- [23] I. Nikitin, [Bivariate systems of polynomial equations with roots of high multiplicity](#). *J. Algebra Appl.* **22** (2023), no. 1, article no. 2350014 Zbl [7637623](#) MR [4526174](#)
- [24] G. Ottaviani, [Introduction to the hyperdeterminant and to the rank of multidimensional matrices](#). In *Commutative algebra*, pp. 609–638, Springer, New York, 2013 Zbl [1276.14078](#) MR [3051388](#)
- [25] G. Salmon, *A treatise on the analytic geometry of three dimensions*. W. Metcalfe and Son, Cambridge, 1882
- [26] L. Schläfli, *Gesammelte mathematische Abhandlungen. Band II*. Birkhäuser, Basel, 1953 Zbl [0051.24102](#) MR [0053873](#)
- [27] J. Weyman and A. Zelevinsky, [Multiplicative properties of projectively dual varieties](#). *Manuscripta Math.* **82** (1994), no. 2, 139–148 Zbl [0839.14039](#) MR [1256155](#)
- [28] J. Weyman and A. Zelevinsky, [Singularities of hyperdeterminants](#). *Ann. Inst. Fourier (Grenoble)* **46** (1996), no. 3, 591–644 Zbl [0853.14001](#) MR [1411723](#)
- [29] M. Zhao, X. Jia, C. Tu, B. Mourrain, and W. Wang, [Enumerating the morphologies of non-degenerate Darboux cyclides](#). *Comput. Aided Geom. Design* **75** (2019), article no. 101776 Zbl [07137433](#) MR [4014752](#)

Received 5 February 2021; revised 19 December 2022.

Alicia Dickenstein

Department of Mathematics, FCEN, University of Buenos Aires and IMAS (UBA-CONICET), Ciudad Universitaria, Pab. I, C1428EGA Buenos Aires, Argentina; alidick@dm.uba.ar

Sandra Di Rocco

School of Engineering Sciences, KTH Royal Institute of Technology, 10044 Stockholm, Sweden; dirocco@kth.se

Ralph Morrison

Department of Mathematics, Williams College, Williamstown, MA 01267, USA; 10rem@williams.edu