

On the dimension spectra of infinite conformal iterated function systems

Tushar Das and David Simmons

Abstract. The dimension spectrum of a conformal iterated function system (CIFS) is the set of all Hausdorff dimensions of its various subsystem limit sets. This brief note provides two constructions – (i) a compact perfect set that cannot be realized as the dimension spectrum of a CIFS; and (ii) a similarity IFS whose dimension spectrum has zero Hausdorff dimension, and thus is not uniformly perfect – which resolve questions posed by Choussionis, Leykekhman and Urbański, and goes on to provoke fresh conjectures and questions regarding the topological and metric properties of IFS dimension spectra.

1. Introduction

The study of iterated function systems (IFSes), which began in earnest in the early 1980s, increased in popularity during the renaissance following Benoit Mandelbrot's seminal work *Les objets fractals* [32] and his invention of the word *fractal* to describe “a mathematical set or concrete object whose form is extremely irregular and/or fragmented at all scales”. Leading researchers who worked on IFS theory and developed a variety of extensions include Bandt, Barnsley, Dekking, Falconer, Graf, Hata, Hutchinson, Mauldin, Schief, Simon, Solomyak and Urbański; see [2, 4, 11, 14, 18, 24, 33, 35, 40, 41] for a small sample of their seminal research. For applications in engineering and science, e.g., in computer graphics, image processing, wavelets, probabilistic growth models and stochastic dynamical systems, see [1, 12, 13, 27, 30, 31].

Several pioneering works focussed on IFSes consisting of finitely many Euclidean similarities; afterwards the theory was extended to handle systems with infinitely many maps (called infinite IFSes) that were conformal. Mauldin and Urbański were among the pioneers of this extension of IFS theory, first to the study of infinite conformal iterated function systems (CIFSes), and then to their generalizations, namely conformal graph directed Markov systems (CGDMSes); see [33, 35]. The CIFS and CGDMS limit sets model several among the intensively studied fractals arising from

2020 *Mathematics Subject Classification.* 37D35, 28A80, 37B10.

Keywords. Hausdorff dimension, iterated function system, conformal dynamics, IFS, CIFS, dimension spectrum, Texan conjecture, thermodynamic formalism, transfer operator.

either side of Sullivan’s dictionary [36, 42] (see also [10, Table 1]); namely, certain Julia sets associated with holomorphic and meromorphic iteration, as well as certain Fuchsian and Kleinian limit sets associated with actions of discrete subgroups of isometries of hyperbolic (negatively curved) spaces.

In particular, the CIFS framework is perfectly suited to encode various sets that appear naturally at the interface of dynamical systems, fractal geometry and Diophantine approximation. For instance, one can encode real numbers via their continued fraction expansions and this leads to the Gauss continued fraction IFS, which is a prime example of an infinite CIFS that comprises of the Möbius maps $x \mapsto 1/(a+x)$ for $a \in \mathbb{N}$. Given any subset $A \subset \mathbb{N}$, let Λ_A denote the set of all irrationals $x \in [0, 1]$ whose continued fraction partial quotients all lie in A . Then, Λ_A may be expressed as the limit set of the subsystem of the Gauss IFS that comprises the maps $x \mapsto 1/(a+x)$ for $a \in A$; see [21]. The elements of Λ_A when A is a finite set are known as *bounded type* numbers. For instance, studying $E_{\{1,2\}}$ relates to the problem of finding rational numbers of given denominator having all partial quotients equal to 1 or 2. The Hausdorff dimensions of such sets have continued to be intensely investigated since several decades; see, e.g., [5, 26, 34, 37, 38] and the references therein.

It was conjectured independently by Hensley [20] and by Mauldin and Urbański [34] in the 1990s that the set $\{\dim_{\mathbb{H}}(\Lambda_A) : A \subset \mathbb{N} \text{ with } |A| < \infty\}$ is a dense subset of the interval $[0, 1]$. This conjecture, dubbed the *Texan conjecture* by Jenkinson [25], was resolved in the affirmative in 2006 by Kesseböhmer and Zhu [29]. The research surrounding the Texan conjecture gave birth to the study of topological and metric features of the dimension spectrum of an infinite CIFS.

Understanding the geometry and topology of the IFS and GDMS dimension spectra has since presented researchers with several challenges; see [7, 8, 15–17, 28]. In their recent papers [7, 8], Chousionis, Leykekhman and Urbański leverage the thermodynamic formalism to commence a careful study of the dimension spectra of finitely irreducible CGDMSes as well as for continued fractions with coefficients restricted to infinite subsets of natural numbers. In particular, they provide a positive answer to the analogue of the Texan conjecture for complex continued fractions [7, Theorem 1.4]. They proved that the dimension spectrum of every infinite CIFS satisfying the open set condition is compact and perfect, and conjectured that every such set may be realized as the dimension spectrum of a similarity IFS. They also asked whether there exists an IFS whose dimension spectrum is not uniformly perfect. This short note resolves both these questions posed by Chousionis, Leykekhman and Urbański in [7] regarding the dimension spectrum of infinite CIFSes, and concludes with some fresh conjectures and research directions in this seam.

Notation. In this note, $\mathbb{N} := \{1, 2, 3, \dots\}$. We write $x \asymp y$ to mean that x and y are *multiplicatively comparable*, i.e., there exists $C > 0$ such that $1/C \leq x/y \leq C$.

We use $\Theta(x)$ to denote any positive quantity multiplicatively comparable to x . We use $x \gg y$ to mean that for every $c > 1$ we have that x is eventually bigger than cy . We write $\dim_{\text{H}}(X)$ to denote the Hausdorff dimension of a set $X \subset \mathbb{R}^n$; and write $\rho(T)$ to denote the spectral radius of an operator T . We simplify notation by writing $\dim_{\text{H}}(A) := \dim_{\text{H}}(\Lambda_A)$ for the Hausdorff dimension of subsystem limit sets (see Definitions 2.1 and 2.4).

2. Definitions and statement of results

The definition of a CIFS appears in several places in the literature; see, for example, [7, Remark 3.2]¹.

Definition 2.1. Fix $d \in \mathbb{N}$. A collection $\mathcal{U} = (u_a)_{a \in E}$ of self-maps of \mathbb{R}^d is a *conformal iterated function system* (CIFS) on \mathbb{R}^d if:

1. E is a countable (finite or infinite) index set, which is referred to as an *alphabet*.
2. $X \subset \mathbb{R}^d$ is a nonempty compact set which is equal to the closure of its interior.
3. For all $a \in E$, $u_a(X) \subset X$.
4. $V \subset \mathbb{R}^d$ is an open connected bounded set such that $\text{dist}(X, \mathbb{R}^d \setminus V) > 0$.
5. For each $a \in E$, u_a is a conformal homeomorphism from V to an open subset of V .
6. (*Cone condition*)

$$\inf_{x \in X} \inf_{r \in (0,1)} \frac{\lambda(X \cap B(x, r))}{r^d} > 0,$$

where λ denotes the Lebesgue measure on \mathbb{R}^d .

7. (*Open set condition, OSC*) For all $a \in E$, the collection $(u_a(\text{int}(X)))_{a \in E}$ is disjoint, where $\text{int}(X)$ denotes the interior of the set X .
8. (*Uniform contraction*)

$$\sup_{a \in E} \sup |u'_a| < 1,$$

and, if E is infinite,

$$\limsup_{a \in E} |u'_a| = 0.$$

9. (*Bounded distortion property*) For all $n \in \mathbb{N}$, $\omega \in E^n$ and $x, y \in V$,

$$|u'_\omega(x)| \asymp |u'_\omega(y)|.$$

¹The original definition may be found in [33, §2, pp. 6–7].

Definition 2.2. Given a countable alphabet E as above, we denote by E^n the set of all words of length n formed using this alphabet, and by E^* the set of all finite words formed using this alphabet. In other words,

$$E^* = \bigcup_{n=0}^{\infty} E^n.$$

If $\omega \in E^* \cup E^{\mathbb{N}}$, i.e., ω is either a finite or infinite word formed using the alphabet E , then we denote subwords of ω by

$$\omega_{n+1}^{n+r} := (\omega_{n+i})_{i=1}^r \in E^r.$$

If $\omega \in E^n$ is a finite word, then we define

$$u_{\omega}(x) := u_{\omega_1} \circ \cdots \circ u_{\omega_n}(x).$$

The *coding map* of the CIFS $\mathcal{U} = (u_a)_{a \in E}$ is the map $\pi : E^{\mathbb{N}} \rightarrow X$ defined by the formula

$$\pi(\omega) = \lim_{n \rightarrow \infty} u_{\omega_1^n}(x_0) = \lim_{n \rightarrow \infty} u_{\omega_1} \circ \cdots \circ u_{\omega_n}(x_0),$$

where $x_0 \in X$ is an arbitrary point. By the uniform contraction hypothesis, $\pi(\omega)$ exists and is independent of the choice of x_0 . The *limit set* of the CIFS is the image of $E^{\mathbb{N}}$ under the coding map, and will be denoted by $\Lambda = \Lambda_E := \pi(E^{\mathbb{N}})$.

Note that the uniform contraction hypothesis implies that the coding map is always Hölder continuous, assuming that the metric on $E^{\mathbb{N}}$ is given by the formula

$$d(\omega, \tau) = \lambda^{|\omega \wedge \tau|},$$

where $\lambda \in (0, 1)$ and $\omega \wedge \tau$ is the longest word which is an initial segment of both ω and τ .

The class of CIFSes consisting of similarities has been studied particularly intensively. We give the definition below in the basic case where $d = 1$.

Definition 2.3. Let E be a countable alphabet, as above. A *similarity iterated function system* (SIFS) on \mathbb{R} is a uniformly contracting and uniformly bounded collection of similarities $\mathcal{U} = \mathcal{U}_E := (u_a)_{a \in E}$ indexed by E . We write each similarity $u_a : \mathbb{R} \rightarrow \mathbb{R}$ as $u_a(x) = \lambda_a x + b_a$ for $0 < |\lambda_a| < 1$ and $b_a \in \mathbb{R}$. A collection $(u_a)_{a \in E}$ of similarities is *uniformly contracting* or *uniformly bounded* if

$$\sup_{a \in E} |\lambda_a| < 1 \quad \text{or} \quad \sup_{a \in E} |b_a| < \infty,$$

respectively. To guarantee that our SIFS \mathcal{U} is a CIFS as defined above, we assume that \mathcal{U} satisfies the *open set condition* (OSC), i.e., there exists an open set $W \subset \mathbb{R}$, whose

closure satisfies the cone condition, such that the collection $(u_a(W))_{a \in E}$ is a disjoint collection of subsets of W . Note that the OSC assumption implies that the collection of similarities are uniformly bounded, i.e., that

$$\sup_{a \in E} |b_a| < \infty,$$

and also that $\lim_{a \in E} |\lambda_a| = 0$ (by taking the Lebesgue measure of the inclusion $\bigcup_{a \in E} u_a(W) \subset W$).

As above, the *limit set* of $\mathcal{U} = \mathcal{U}_E$ is the image of the *coding map* $\pi : E^{\mathbb{N}} \rightarrow \mathbb{R}$ defined by

$$\pi(\omega) = \lim_{n \rightarrow \infty} u_{\omega_1^n}(x_0) = \lim_{n \rightarrow \infty} u_{\omega_1} \circ \cdots \circ u_{\omega_n}(x_0),$$

and will be denoted $\Lambda = \Lambda_E := \pi(E^{\mathbb{N}})$. Note that given any SIFS (not necessarily satisfying the OSC) the uniformly contracting and uniformly bounded condition implies that π is defined. When we write that \mathcal{U} is an SIFS, we assume, as is common [7, Remark 3.2], that the OSC is satisfied.

Definition 2.4. Given an SIFS or CIFS $\mathcal{U} = \mathcal{U}_E$ we will be interested in sub-CIFSes or sub-SIFSes (called *subsystems*) formed by restrictions of \mathcal{U} to various subsets of the original alphabet E . Given $A \subset E$, the corresponding subsystem, coding map and limit set are denoted by \mathcal{U}_A , π_A and Λ_A , respectively.

Definition 2.5. The (*Hausdorff*) *dimension spectrum* of a CIFS $\mathcal{U} = (u_a)_{a \in E}$ is defined as

$$\text{DimSpec}(\mathcal{U}) := \{\dim_{\text{H}}(\Lambda_A) : A \subset E\}.$$

Chousionis, Leykekhman and Urbański proved [7, Theorem 1.2] that the dimension spectrum of an infinite CIFS is compact and perfect. They went on to conjecture [7, Conjecture 1.3] that every compact perfect set $K \subset [0, \infty)$ can be the dimension spectrum of a CIFS. Note that by taking a one-element subset of the alphabet, we get a subsystem whose limit set is a singleton and thus of Hausdorff dimension zero. Thus, $0 \in \text{DimSpec}(\mathcal{U})$ for all iterated function systems \mathcal{U} . Hence, their original conjecture should be reformulated to only consider compact perfect sets containing zero. Our first result shows that their (reformulated) conjecture was too optimistic.

Theorem 2.6. *There exists a compact and perfect set $K \subset [0, 1]$ such that $0 \in K$ and $\text{DimSpec}(\mathcal{U}) \neq K$ for all CIFSes \mathcal{U} on \mathbb{R} .*

The proof of Theorem 2.6 shows that it remains true if \mathbb{R} is replaced by \mathbb{R}^d for any $d \in \mathbb{N}$.

In [7], Chousionis, Leykekhman and Urbański recognized that their conjecture had “room for many partial results and open questions”. They asked, in particular,

whether there exists an IFS whose dimension spectrum is not uniformly perfect. Our second result answers their question in the affirmative.

Theorem 2.7. *There exists an infinite SIFS on \mathbb{R} whose dimension spectrum has Hausdorff dimension zero. In particular, the dimension spectrum is not uniformly perfect.*

Before we present the proofs of these theorems in the following two sections, let us briefly remark on the salient ideas. The idea of Theorem 2.7 is to construct an SIFS \mathcal{U} whose contraction ratios form a quickly decreasing sequence, and then simply calculate an upper bound on the Hausdorff dimension of $\text{DimSpec}(\mathcal{U})$.

On the other hand, the basic idea of the proof of Theorem 2.6 is to exploit some geometric structure of the set $\text{DimSpec}(\mathcal{U})$. Namely, if F_1 and F_2 are two subsets of the alphabet E , then for each $b \in E \setminus F_1 \cup F_2$, we have a quadruple of points

$$\begin{aligned}\delta_1 &= \delta(F_1), & \delta'_1 &= \delta(F_1 \cup \{b\}), \\ \delta_2 &= \delta(F_2), & \delta'_2 &= \delta(F_2 \cup \{b\}),\end{aligned}$$

where $\delta(F) := \dim_{\text{H}}(\Lambda_F)$. We show that these four points satisfy the relation

$$(\delta'_1 - \delta_1)^C \leq \delta'_2 - \delta_2 \leq (\delta'_1 - \delta_1)^{1/C},$$

where C is a constant. We then construct a set K where this relation is satisfied for only finitely many δ'_1 and δ'_2 , given fixed δ_1 and δ_2 .

3. Proof of Theorem 2.6

To simplify the notation in the sequel, we will write $\dim_{\text{H}}(A) := \dim_{\text{H}}(\Lambda_A)$ for all sets $A \subset E$.

Let $A = \{0, 1\}$, and let $(\sigma_n)_{n \geq 1}$ be an enumeration of A^* such that the map $n \mapsto |\sigma_n|$ is nondecreasing. Next, let $g(\sigma_n) := 4^{-n!}$, and let $f : A^{\mathbb{N}} \rightarrow [0, 1]$ be defined by

$$f(\omega) := \sum_{n: \omega_{n+1}=1} g(\omega_1^n).$$

We define $K := f(A^{\mathbb{N}})$. The map f is continuous, and thus K is compact.

The map f is injective. Indeed, suppose $\omega, \tau \in A^{\mathbb{N}}$ are distinct. Without loss of generality, we may take them to be of the form $\omega = \omega_1^n 0 \omega_{n+2}^{\infty}$ and $\tau = \omega_1^n 1 \tau_{n+2}^{\infty}$. It follows from the properties of $(\sigma_n)_{n \geq 1}$ and g that $g(\omega_1^{n+k}) \leq 4^{-k} g(\omega_1^n)$ for all k . Therefore, we have that

$$f(\tau) - f(\omega) \geq g(\omega_1^n) - \sum_{m>n} g(\omega_1^m) \geq g(\omega_1^n) \left[1 - \sum_{k>0} 4^{-k} \right] = \frac{2}{3} g(\omega \wedge \tau) > 0.$$

Now, since K is the continuous injective image of a perfect space, it itself is thus perfect.

Next, observe that

$$f(\tau) - f(\omega) \leq g(\tau_1^n) + \sum_{m>n} g(\tau_1^m) \leq g(\tau_1^n) \left[1 + \sum_{k>0} 4^{-k} \right] = \frac{4}{3} g(\omega \wedge \tau).$$

The previous two calculations show that

$$f(\tau) - f(\omega) \asymp g(\omega \wedge \tau). \quad (1)$$

We now want to prove that $\text{DimSpec}(\mathcal{U}) \neq K$ for all CIFSes \mathcal{U} on \mathbb{R} . So let $\mathcal{U} = (u_a)_{a \in E}$ be an infinite CIFS on \mathbb{R} with alphabet E and, by way of contradiction, suppose that $\text{DimSpec}(\mathcal{U}) = K$. Next, for a given $F \subseteq E$ with $2 \leq \#(F) < \infty$, we estimate how much the Hausdorff dimension of F increases after we add an extra symbol $b \in E \setminus F$. Note that similar results, though only with upper bounds rather than two-sided bounds, may be found in [19, Theorem 3.3] and [7, Theorem 5.4].

Claim 3.1. *For every $F \subseteq E$ with $2 \leq \#(F) < \infty$ and for every $b \in E \setminus F$, we have*

$$\dim_{\text{H}}(F \cup \{b\}) = \delta + \Theta(D_b^\delta),$$

where $\delta = \dim_{\text{H}}(F)$ and $D_b = \|u'_b\| := \sup |u'_b|$, and the constant associated with Θ depends on the subset $F \subseteq E$.

Proof. Recall from Definition 2.1 of a CIFS that $X \subset \mathbb{R}^d$ is a nonempty compact set which is equal to the closure of its interior. Let $C(X)$ denote the Banach space of continuous functions from X to $(0, \infty)$, and let $L : C(X) \rightarrow C(X)$ denote the Perron–Frobenius operator of the CIFS $(u_a)_{a \in F}$, i.e.,

$$Lf(x) := \sum_{a \in F} |u'_a(x)|^\delta f \circ u_a(x).$$

Then there exists a positive continuous map $g : X \rightarrow (0, \infty)$ such that $Lg = g$, by [35, Theorem 6.1.2]. Now, fix $b \in E \setminus F$ and let

$$L'f(x) := Lf(x) + |u'_b(x)|^\delta f \circ u_b(x).$$

Then, the logarithm of the spectral radius of L' is $\log \rho(L') = P(F \cup \{b\}, \delta)$, the pressure of the CIFS $(u_a)_{a \in F \cup \{b\}}$ evaluated at δ ; see [35, Theorem 2.4.3, Theorem 2.4.6, and p. 29]. To estimate this, we compute $L'g$. Now, by the bounded distortion property (see Definition 2.1) and since g is bounded from above and below on the compact set X , we have

$$(L' - L)g(x) = |u'_b(x)|^\delta g \circ u_b(x) \asymp D_b^\delta g(x),$$

and thus,

$$L'g = (1 + \Theta(D_b^\delta))g.$$

Since g and L' are both positive, this tells us² that the spectral radius satisfies

$$\exp(P(F \cup \{b\}, \delta)) = \rho(L') = 1 + \Theta(D_b^\delta).$$

Thus, we have that

$$P(F \cup \{b\}, \delta) \asymp D_b^\delta.$$

On the other hand, from Bowen's formula, we know that $P(F \cup \{b\}, s) = 0$, where $s = \dim_{\text{H}}(F \cup \{b\})$; see [35, Theorem 4.2.11]. Moreover, the negative derivative of pressure satisfies

$$-P'(F \cup \{b\}, \cdot) \asymp 1 \tag{2}$$

on $[\delta, \infty)$, independently of b . Indeed, the lower bound in (2) follows from direct calculation, while the upper bound in (2) follows from the convexity of pressure [35, Proposition 4.2.8(b)] together with the fact that $P(F \cup \{b\}, 0) = \log(\#(F) + 1)$ is independent of b , and that $\delta > 0$ since $\#(F) \geq 2$. It thus follows that $s = \delta + \Theta(D_b^\delta)$. This concludes the proof of Claim 3.1. \blacksquare

Next, we consider $F_1, F_2 \subset E$ such that $2 \leq |F_i| < \infty$, $\delta_i := \dim_{\text{H}}(F_i) > 0$, and $\delta_2 > \delta_1$; for instance, we could take $F_1 = \{a_1, a_2\}$ and $F_2 = \{a_1, a_2, a_3\}$, where $(a_n)_{n \geq 1}$ is an enumeration of E .

Now fix some $b \in E \setminus (F_1 \cup F_2)$. Since we assumed $\text{DimSpec}(U) = K = f(A^{\mathbb{N}})$, there exist $\omega, \omega', \tau, \tau' \in A^{\mathbb{N}}$ such that

$$\begin{aligned} f(\omega) &= \dim_{\text{H}}(F_1) = \delta_1, & f(\omega') &= \dim_{\text{H}}(F_1 \cup \{b\}), \\ f(\tau) &= \dim_{\text{H}}(F_2) = \delta_2, & f(\tau') &= \dim_{\text{H}}(F_2 \cup \{b\}). \end{aligned}$$

By applying Claim 3.1 twice, we have

$$f(\omega') - f(\omega) \asymp D^{f(\omega)} \quad \text{and} \quad f(\tau') - f(\tau) \asymp D^{f(\tau)},$$

where $D = D_b = \|u'_b\|$. Now, let $\omega'' := \omega \wedge \omega'$ and $\tau'' := \tau \wedge \tau'$. Then, using (1), we have

$$\begin{aligned} g(\omega'') &\asymp f(\omega') - f(\omega) \asymp D^{\delta_1}, \\ g(\tau'') &\asymp f(\tau') - f(\tau) \asymp D^{\delta_2}, \end{aligned}$$

²Though this is potentially standard to functional analysts, we include a brief argument for the reader's convenience: If $Lg \geq \lambda g$, then $\|L^n\| \gtrsim \|L^n g\| \geq \|\lambda^n g\| = \lambda^n \|g\|$ and thus, $\rho(L) \geq \lambda$, and if $Lg \leq \lambda g$, then for all f , we have $\|L^n f\| \leq \|f\| \cdot \|g^{-1}\| \cdot \|L^n g\| \leq \|f\| \cdot \|g^{-1}\| \cdot \lambda^n \|g\|$ and thus, $\|L^n\| \lesssim \lambda^n$, yielding $\rho(L) \leq \lambda$.

and thus,

$$g(\tau'') \asymp g(\omega'')^{\delta_2/\delta_1}.$$

Rewriting this using the definition of g , we see that

$$4^{n!} \asymp 4^{sm!},$$

where $s = \delta_2/\delta_1$, $\sigma_n = \omega''$ and $\sigma_m = \tau''$.

Note that since $\lim_{b \in E} \|u'_b\| = 0$, Claim 3.1 implies that $\lim_{b \in E} f(\omega') = f(\omega)$ or, equivalently, that $\lim_{b \in E} |\omega''| = \infty$. Thus, n and m will both become arbitrarily large as b ranges over E . If $n > m$ for infinitely many $b \in E$, then $4^{n!} \geq 4^{nm!} \gg 4^{sm!}$, a contradiction. Similarly, if $m \geq n$ for infinitely many $b \in E$, then since $s = \delta_2/\delta_1 > 1$, we have $4^{m!} \geq 4^{n!} \gg 4^{s^{-1}n!}$, another contradiction. This concludes the proof of Theorem 2.6.

4. Proof of Theorem 2.7

Let $\mathcal{U} = (u_a)_{a \in E}$ be a collection of similarities satisfying the OSC such that for all $a \in E := \mathbb{N}$,

$$|u'_a| = 2^{-a^2}.$$

(The precise choice of these similarities does not matter as long as they satisfy the OSC.) Let $A := \{0, 1\}$, and let $f : A^* \cup A^{\mathbb{N}} \rightarrow \text{DimSpec}(\mathcal{U})$ be defined by the formula $f(\tau) := \dim_{\mathbb{H}}(\Lambda_{A_\tau})$, where $A_\tau := \{a \in E : \tau_a = 1\}$.

Next we prove an analogue of Claim 3.1, where the associated constant is now independent of the unperturbed limit set.

Claim 4.1. *Fix two elements $a_1, a_2 \in E$, and define $\tilde{A}^* := \{\omega \in A^* : \omega_{a_1} = \omega_{a_2} = 1\}$ and $\tilde{A}^{\mathbb{N}} := \{\omega \in A^{\mathbb{N}} : \omega_{a_1} = \omega_{a_2} = 1\}$. Then, for all $\omega \in \tilde{A}^*$, we have*

$$f(\omega 1) = f(\omega 0) + \Theta(2^{-|\omega|^2 f(\omega)}),$$

where the associated constants may depend on a_1, a_2 .

Proof. Let $\omega \in \tilde{A}^*$. Then it follows from Bowen's formula [35, Theorem 4.2.11] that

$$\sum_{a: \omega_n=1} 2^{-a^2 f(\omega 0)} = 1,$$

and that

$$\sum_{a: \omega_n=1} 2^{-a^2 f(\omega 1)} + 2^{-(|\omega|+1)^2 f(\omega 1)} = 1.$$

Hence,

$$\begin{aligned} 2^{-(|\omega|+1)^2 f(\omega 1)} &= \sum_{a:\omega_n=1} 2^{-a^2 f(\omega 0)} - \sum_{a:\omega_n=1} 2^{-a^2 f(\omega 1)} \\ &= (f(\omega 1) - f(\omega 0)) \cdot \sum_{a:\omega_n=1} \ln(2) 2^{-a^2 \xi} a^2, \end{aligned}$$

where $\xi \in (f(\omega 0), f(\omega 1))$. It then follows that

$$2^{-(|\omega|+1)^2 f(\omega 1)} \asymp f(\omega 1) - f(\omega 0),$$

since

$$0 < 2^{-a_1^2} a_1^2 + 2^{-a_2^2} a_2^2 \leq \sum_{a:\omega_n=1} 2^{-a^2 \xi} a^2 \leq \sum_{a \in E} 2^{-a^2 s} a^2 < \infty,$$

where $s := \dim_{\text{H}}(\{a_1, a_2\}) > 0$. This concludes the proof of Claim 4.1. \blacksquare

It then follows from Claim 4.1 that for all $a_1, a_2 \in E$ and for all $\omega, \tau \in \tilde{A}^*$, we have

$$|f(\omega) - f(\tau)| = O(2^{-|\omega \wedge \tau|^2 f(\omega)}) = O(2^{-s|\omega \wedge \tau|^2}),$$

where $s = \dim_{\text{H}}(\{a_1, a_2\})$. Thus, for each $n \in \mathbb{N}$, the set $f(\tilde{A}^{\mathbb{N}})$ can be covered by $\#(A)^n = 2^n$ balls of radius $O(2^{-sn^2})$. This implies that the box dimension, and thus that Hausdorff dimension, of $f(\tilde{A}^{\mathbb{N}})$ is zero. By the countable stability of the Hausdorff dimension,

$$\dim_{\text{H}}(\text{DimSpec}(\mathcal{U})) = f(A^{\mathbb{N}}) = 0.$$

This concludes the proof of Theorem 2.7.

5. Conjectures and future work

Our investigations of topological and metric properties of the dimension spectra of various conformal iterated function systems led to the following conjectures.

Conjecture 5.1. The only sets $K \subset [0, \infty)$ such that both K and its mirror image $\text{sup}(K) - K$ are dimension spectra of CIFs are intervals, i.e., $K = [0, \lambda]$ for some $\lambda > 0$.

Let us recall the definition of local Hausdorff dimension in the setting of a metric space.

Definition 5.2. Let X be a metric space and let $F \subset X$. For all $x \in X$, we define the *local Hausdorff dimension* of F at x as

$$\dim_x(F) := \inf \{ \dim_{\text{H}}(F \cap B(x, \varepsilon)) : \varepsilon > 0 \}.$$

The monotonicity of the Hausdorff dimension implies that the infimum in the definition above is actually a limit as ε tends to zero, i.e.,

$$\dim_x(F) = \lim_{\varepsilon \rightarrow 0} \dim_{\text{H}}(F \cap B(x, \varepsilon)).$$

Conjecture 5.3. Let $F \subset [0, \infty)$ be the dimension spectrum of a CIFS. The map $x \mapsto \dim_x(F)$ restricted to $x \in F$ is a continuous, weakly decreasing (i.e., nonincreasing) function.

Conjecture 5.4. Let $F \subset [0, \infty)$ be the dimension spectrum of a CIFS. Then, one of the following three mutually exclusive scenarios holds:

- Type I: F is equal to the union of finitely many intervals.
- Type II: F has zero Hausdorff dimension.
- Type III: The local Hausdorff dimension satisfies $\dim_x(F) = \min(1, c/x)$ for all $x \in F$, for some $0 < c < \sup(F)$; i.e., the graph of the function $F \ni x \mapsto \dim_x(F)$ is a horizontal line followed by a hyperbola.

Remark 5.5. For each of the three scenarios in Conjecture 5.4, there exists a set K exemplifying the scenario, which can be realized as the dimension spectrum of a SIFS. Indeed, this observation led us to Conjecture 5.4. Theorem 2.7 provides an example of Type II. We leave it as an exercise for the interested reader to verify that the dimension spectrum of any SIFS whose similarities have contraction ratios $1/2, 1/4, 1/8, \dots$ is of Type I, and similarly that the spectrum of one whose similarities have contraction ratios $1/3, 1/3, 1/9, 1/27, \dots$ is of Type III.

Remark 5.6. If Conjecture 5.4 is true, then for each $F \subset [0, \infty)$ that is a dimension spectrum of a CIFS, the local Hausdorff dimension satisfies $\dim_x(F) = \min(1, c/x)$ for all $x \in F$, for some $0 \leq c \leq \sup(F)$. The cases where $c = \sup(F)$ and $c = 0$ correspond to Types I and II, respectively.

In general, it appears difficult to distinguish between sets that can be SIFS or CIFS dimension spectra and those that cannot. In particular, it would be interesting to understand when an SIFS dimension spectrum could be realized as that of a CIFS that is not an SIFS, and vice versa.

The study of finite SIFSes with overlaps has witnessed several breakthroughs in the last decade; see [22, 23]. It would be interesting to know whether dimension spectra behave differently in the absence of the OSC. For instance, recall that Chouisionis,

Leykekhman and Urbański proved [7, Theorem 1.2] that the dimension spectrum of an infinite conformal iterated function system satisfying the open set condition is compact and perfect. However, this theorem is false for some systems that satisfy all conditions of being a CIFS except for the OSC. Indeed, take the family of maps $\mathcal{U} = \{u_a : \mathbb{R} \rightarrow \mathbb{R}\}_{a \in E}$ defined by $u_a(x) := (1/2)x + a$ for $a \in E := \mathbb{Q} \cap [0, 1]$. Then, for any $F \subseteq E$, the dimension of Λ_F is either 0 or 1 depending on whether or not $\#(F) \geq 2$, and thus, $\text{DimSpec}(\mathcal{U}) = \{0, 1\}$.

Given that a finite alphabet CIFS will have a finite dimension spectrum, it is natural to ask the following inverse problem.

Question 5.7. If D is a finite subset of the nonnegative real numbers containing zero does there exist a finite CIFS whose dimension spectrum is D ?

Beyond similarity and conformal IFSes, the dimension spectra of affine IFSes remain unanalyzed. It may be fruitful to first focus on *infinitely generated versions* of certain well-studied classes of finitely generated affine or other non-conformal IFSes; see, e.g., [3, 6, 9, 39]. See [28] for some recent progress in this direction.

In a different direction, rather than focussing on solely the Hausdorff dimension spectra, the study of spectra of other fractal dimensions – such as packing dimension, box dimension, and Assouad dimension – also awaits investigation.

Acknowledgements. The authors discussed the results of this research with Vasileios Chousionis and Mariusz Urbański at the American Institute of Mathematics (AIM) in March 2018, where they were collaborating via the [SQuaREs program](#). We thank the AIM staff for nurturing this outstanding collective research opportunity and for providing us with excellent working conditions. We thank Balázs Bárány for stimulating discussions, as well as two anonymous referees for their valuable suggestions that helped clarify certain infelicities, improved the exposition of our proofs, and led to the inclusion of Question 5.7 in Section 5.

Funding. David Simmons is supported by a 2018 Royal Society University Research Fellowship, URF\R1\180649.

References

- [1] C. Bandt, M. F. Barnsley, R. Devaney, K. J. Falconer, V. Kannan, and P. B. Vinod Kumar (eds.), *Fractals, wavelets, and their applications*. Springer Proc. Math. Stat. 92, Springer, Cham, 2014 Zbl [1300.28001](#) MR [3280210](#)
- [2] C. Bandt and S. Graf, Self-similar sets. VII. A characterization of self-similar fractals with positive Hausdorff measure. *Proc. Amer. Math. Soc.* **114** (1992), no. 4, 995–1001 Zbl [0823.28003](#) MR [1100644](#)

- [3] K. Barański, Hausdorff dimension of the limit sets of some planar geometric constructions. *Adv. Math.* **210** (2007), no. 1, 215–245 MR [2298824](#)
- [4] M. Barnsley, *Fractals everywhere*. Academic Press, Inc., Boston, MA, 1988
Zbl [0691.58001](#) MR [977274](#)
- [5] R. T. Bumby, Hausdorff dimensions of Cantor sets. *J. Reine Angew. Math.* **331** (1982), 192–206 Zbl [0468.10032](#) MR [647383](#)
- [6] J. Chen and Y. Pesin, Dimension of non-conformal repellers: a survey. *Nonlinearity* **23** (2010), no. 4, R93–R114 Zbl [1201.37026](#) MR [2602012](#)
- [7] V. Chousionis, D. Leykekhman, and M. Urbański, The dimension spectrum of conformal graph directed Markov systems. *Selecta Math. (N.S.)* **25** (2019), no. 3, Paper No. 40, 74
Zbl [1421.37013](#) MR [3960790](#)
- [8] V. Chousionis, D. Leykekhman, and M. Urbański, On the dimension spectrum of infinite subsystems of continued fractions. *Trans. Amer. Math. Soc.* **373** (2020), no. 2, 1009–1042
Zbl [1437.37032](#) MR [4068257](#)
- [9] T. Das and D. Simmons, The Hausdorff and dynamical dimensions of self-affine sponges: a dimension gap result. *Invent. Math.* **210** (2017), no. 1, 85–134 Zbl [1387.37026](#)
MR [3698340](#)
- [10] T. Das, D. Simmons, and M. Urbański, Dimension rigidity in conformal structures. *Adv. Math.* **308** (2017), 1127–1186 Zbl [1405.37047](#) MR [3600084](#)
- [11] F. M. Dekking, Recurrent sets. *Adv. in Math.* **44** (1982), no. 1, 78–104 Zbl [0495.51017](#)
MR [654549](#)
- [12] M. Dekking, J. Lévy Véhel, E. Lutton, and C. Tricot (eds.), *Fractals: theory and applications in engineering*. Springer, London, 1999 Zbl [0936.00006](#) MR [1726363](#)
- [13] J. L. Encarnação, H.-O. Peitgen, G. Sakas, and G. Englert (eds.), *Fractal Geometry and Computer Graphics*. Beiträge zur Graphischen Datenverarbeitung, Springer, Berlin, Heidelberg, 1992
- [14] K. Falconer, *Fractal geometry: mathematical foundations and applications*. John Wiley & Sons, Ltd., Chichester, 1990 Zbl [0689.28003](#) MR [1102677](#)
- [15] A. Ghenciu, S. Munday, and M. Roy, The Hausdorff dimension spectrum of conformal graph directed Markov systems and applications to nearest integer continued fractions. *J. Number Theory* **175** (2017), 223–249 Zbl [1375.11057](#) MR [3608189](#)
- [16] A. E. Ghenciu, Parabolic iterated function systems with applications to the backward continued fractions. *Far East J. Dyn. Syst.* **9** (2007), no. 1, 75–91 Zbl [1130.37006](#)
MR [2300398](#)
- [17] A. E. Ghenciu, Gauss-like continued fraction systems and their dimension spectrum. *Real Anal. Exchange* **34** (2009), no. 1, 17–28 Zbl [1179.37013](#) MR [2527119](#)
- [18] S. Graf, R. D. Mauldin, and S. C. Williams, The exact Hausdorff dimension in random recursive constructions. *Mem. Amer. Math. Soc.* **71** (1988), no. 381, x+121
Zbl [0641.60003](#) MR [920961](#)
- [19] S.-M. Heinemann and M. Urbański, Hausdorff dimension estimates for infinite conformal IFSs. *Nonlinearity* **15** (2002), no. 3, 727–734 Zbl [1015.37019](#) MR [1901102](#)

- [20] D. Hensley, A polynomial time algorithm for the Hausdorff dimension of continued fraction Cantor sets. *J. Number Theory* **58** (1996), no. 1, 9–45 Zbl [0858.11039](#) MR [1387719](#)
- [21] D. Hensley, *Continued fractions*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2006 Zbl [1161.11028](#) MR [2351741](#)
- [22] M. Hochman, Some problems on the boundary of fractal geometry and additive combinatorics. In *Recent developments in fractals and related fields*, pp. 129–174, Trends Math., Birkhäuser/Springer, Cham, 2017 Zbl [1383.28002](#) MR [3775463](#)
- [23] M. Hochman, Dimension theory of self-similar sets and measures. In *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. III. Invited lectures*, pp. 1949–1972, World Sci. Publ., Hackensack, NJ, 2018 Zbl [1447.28007](#) MR [3966837](#)
- [24] J. E. Hutchinson, Fractals and self-similarity. *Indiana Univ. Math. J.* **30** (1981), no. 5, 713–747 Zbl [0598.28011](#) MR [625600](#)
- [25] O. Jenkinson, On the density of Hausdorff dimensions of bounded type continued fraction sets: the Texan conjecture. *Stoch. Dyn.* **4** (2004), no. 1, 63–76 Zbl [1089.28006](#) MR [2069367](#)
- [26] O. Jenkinson and M. Pollicott, Rigorous effective bounds on the Hausdorff dimension of continued fraction Cantor sets: a hundred decimal digits for the dimension of E_2 . *Adv. Math.* **325** (2018), 87–115 Zbl [1386.30039](#) MR [3742587](#)
- [27] P. E. T. Jorgensen, *Analysis and probability: wavelets, signals, fractals*. Graduate Texts in Mathematics 234, Springer, New York, 2006 Zbl [1104.42001](#) MR [2254502](#)
- [28] N. Jurga, Dimension spectrum of infinite self-affine iterated function systems. *Selecta Math. (N.S.)* **27** (2021), no. 3, Paper No. 49 Zbl [1469.28007](#) MR [4273646](#)
- [29] M. Kesseböhmer and S. Zhu, Dimension sets for infinite IFSs: the Texan conjecture. *J. Number Theory* **116** (2006), no. 1, 230–246 Zbl [1085.37018](#) MR [2197868](#)
- [30] H. Kunze, D. La Torre, F. Mendivil, and E. R. Vrscay, *Fractal-based methods in analysis*. Springer, New York, 2012 Zbl [1237.28002](#) MR [3014680](#)
- [31] J. Lévy Véhel and E. Lutton (eds.), *Fractals in Engineering*. Springer, London, 2005
- [32] B. Mandelbrot, *Les objets fractals*. Nouvelle Bibliothèque Scientifique, Flammarion, Paris, 1975 Zbl [0900.00018](#) MR [0462040](#)
- [33] R. D. Mauldin and M. Urbański, Dimensions and measures in infinite iterated function systems. *Proc. London Math. Soc. (3)* **73** (1996), no. 1, 105–154 Zbl [0852.28005](#) MR [1387085](#)
- [34] R. D. Mauldin and M. Urbański, Conformal iterated function systems with applications to the geometry of continued fractions. *Trans. Amer. Math. Soc.* **351** (1999), no. 12, 4995–5025 Zbl [0940.28009](#) MR [1487636](#)
- [35] R. D. Mauldin and M. Urbański, *Graph directed Markov systems*. Cambridge Tracts in Mathematics 148, Cambridge University Press, Cambridge, 2003 Zbl [1033.37025](#) MR [2003772](#)
- [36] C. T. McMullen, The classification of conformal dynamical systems. In *Current developments in mathematics, 1995 (Cambridge, MA)*, pp. 323–360, Int. Press, Cambridge, MA, 1994 Zbl [0908.30028](#) MR [1474980](#)

- [37] C. G. Moreira, Geometric properties of the Markov and Lagrange spectra. *Ann. of Math.* (2) **188** (2018), no. 1, 145–170 Zbl [1404.11095](#) MR [3815461](#)
- [38] C. G. T. d. A. Moreira, Dynamical systems, fractal geometry and Diophantine approximations. In *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. I. Plenary lectures*, pp. 731–757, World Sci. Publ., Hackensack, NJ, 2018 Zbl [1451.37006](#) MR [3966744](#)
- [39] H. W. J. Reeve, Infinite non-conformal iterated function systems. *Israel J. Math.* **194** (2013), no. 1, 285–329 Zbl [1301.37012](#) MR [3047072](#)
- [40] A. Schief, Separation properties for self-similar sets. *Proc. Amer. Math. Soc.* **122** (1994), no. 1, 111–115 Zbl [0807.28005](#) MR [1191872](#)
- [41] K. Simon, B. Solomyak, and M. Urbański, Invariant measures for parabolic IFS with overlaps and random continued fractions. *Trans. Amer. Math. Soc.* **353** (2001), no. 12, 5145–5164 Zbl [0987.37075](#) MR [1852098](#)
- [42] D. Sullivan, Conformal dynamical systems. In *Geometric dynamics (Rio de Janeiro, 1981)*, pp. 725–752, Lecture Notes in Math. 1007, Springer, Berlin, 1983 Zbl [0524.58024](#) MR [730296](#)

Received 20 September 2020.

Tushar Das

Department of Mathematics & Statistics, University of Wisconsin – La Crosse, 1725 State Street, La Crosse, WI 54601, USA; tdas@uwlax.edu

David Simmons

Department of Mathematics, University of York, Heslington, York YO10 5DD, UK; david9550@gmail.com