

The pointwise behavior of Riemann's function

Frederik Broucke and Jasson Vindas

Abstract. We present a new and simple method for the determination of the pointwise Hölder exponent of Riemann's function $\sum_{n=1}^{\infty} n^{-2} \sin(\pi n^2 x)$ at every point of the real line. In contrast to earlier approaches, where wavelet analysis and the theta modular group were needed for the analysis of irrational points, our method is direct and elementary, being only based on the following tools from number theory and complex analysis: the evaluation of quadratic Gauss sums, the Poisson summation formula, and Cauchy's theorem.

1. Introduction

According to an account of Weierstrass, Riemann would have suggested the function

$$f(x) = \sum_{n=1}^{\infty} \frac{\sin(n^2 \pi x)}{n^2} \quad (1.1)$$

as an example of a function which is continuous but nowhere differentiable. In 1916, Hardy [12] proved, based on earlier work by him and Littlewood [13], that Riemann's function f is not differentiable in a certain subset of \mathbb{R} that contains every irrational point. This seemed to confirm the nowhere differentiability conjecture, but, on the contrary, Gerver [10] showed in 1970 that f is actually differentiable at any rational number of the form $(2r + 1)/(2s + 1)$, $r, s \in \mathbb{Z}$. His results [10, 11] in combination with Hardy's ones imply that Riemann's function is not differentiable at any other real number. Gerver's proofs are elementary, but difficult and long. Simpler proofs were found later by Smith in 1972 [23] and Itatsu in 1981 [15] (see also [14, 18]). They provided more precise information about the pointwise behavior of Riemann's "non-differentiable" function, which in particular gives the pointwise Hölder exponent [17] at any rational point. This left open the determination of the exact pointwise regularity of Riemann's function at the irrationals.

2020 Mathematics Subject Classification. Primary 42A16; Secondary 26A16, 26A27, 11F30, 11J70, 42A55.

Keywords. Riemann "non-differentiable" function, pointwise Hölder exponent, quadratic Gaussian sums, fractional integrals of modular forms.

Duistermaat [7] used a variant of Itatsu's approach to exhibit explicit dependence of the O -constants on the analyzed rational point in the Smith–Itatsu asymptotic formulas. His error terms were strong enough to enable him to find an upper bound for the pointwise Hölder exponent at every irrational point. His upper bound depends on approximation properties of the irrational number by certain continued fractions. The problem of finding the pointwise Hölderian regularity of Riemann's function at irrational points was finally solved by Jaffard [16] in 1996, who showed that Duistermaat's upper bound was sharp, that is, it is exactly equal to the pointwise Hölder exponent. Jaffard's proof is indirect and non-elementary.

A generalization of Jaffard's result to fractional integrals of modular forms has recently been obtained by Pastor [20, Theorem 2.3(2)]. As Jaffard's, his proof is of Tauberian nature and makes use of wavelet analysis and the action of subgroups of the modular group.

We refer to the articles [2, 3, 8, 9] for other recently revealed fascinating properties of Riemann's function.

The purpose of this paper is to provide a new and self-contained approach for the determination of the pointwise Hölder exponent of Riemann's function at every point. Our arguments are direct and lead to completely elementary and fairly short proofs that only rely on the following tools: the evaluation of quadratic Gauss sums, the Poisson summation formula, and Cauchy's integral theorem. We highlight that the main novelty in our treatment is the use of a simple contour integral formula for the pointwise analysis at *irrational* numbers. To the best of our knowledge, this is the first explicit proof in the literature that avoids passing through the wavelet transform for the computation of the pointwise Hölder exponent at each irrational point.

Our method can be sketched as follows. For the sake of convenience, we work with a rescaled and complex version of Riemann's function, namely,

$$\phi(z) = \sum_{n=1}^{\infty} \frac{1}{2\pi i n^2} e(n^2 z), \quad (1.2)$$

where we use the notation $e(z)$ for $e^{2\pi i z}$ and $z = x + iy$ with $y \geq 0$. The pointwise properties of Riemann's original function can easily be deduced from those of ϕ . We are interested in the computation of the pointwise Hölder exponent

$$\alpha(x) = \sup\{\alpha > 0: \phi(x+h) = P_x(h) + O_x(|h|^\alpha) \text{ for some polynomial } P_x\}. \quad (1.3)$$

Restricting the complex variable z to the upper half-plane, one has

$$\phi'(z) = \frac{1}{2}(\theta(z) - 1),$$

where θ stands for the Jacobi theta function, namely, $\theta(z) = \sum_{n \in \mathbb{Z}} e(n^2 z)$. Therefore, for each $x \in \mathbb{R}$, we obtain the basic identity

$$\phi(x+h) - \phi(x) + \frac{1}{2}h = \frac{1}{2} \lim_{y \rightarrow 0^+} \int_{iy}^{h+iy} \theta(x+z) dz, \quad (1.4)$$

a formula that was already employed by Itatsu for $x = 0$.

We will exploit the formula (1.4) for the analysis of both rational and irrational numbers x . Itatsu and Duistermaat used (1.4) at $x = 0$ and then transformation properties (under the theta modular group) to study all rational points. We take a different path, in the spirit of Smith, and use the Poisson summation formula to study the boundary behavior of $\theta(x+z)$. This directly gives an exact expression for the limit of the integral in (1.4) when x is rational that yields an asymptotic series and that we shall discuss in Section 3. Approximating x by the n th convergent $r_n = p_n/q_n$ in its continued fraction expansion when x is irrational and using our exact formula for $\theta(r_n+z)$, one generates sufficiently good bounds for $\theta(x+z)$. The next key step in our method is to use Cauchy's theorem to transform (1.4) into

$$\phi(x+h) - \phi(x) + \frac{1}{2}h = -\frac{1}{2} \int_{\Gamma} \theta(x+z) dz, \quad (1.5)$$

where Γ is the part of the counterclockwise oriented boundary of the rectangle with vertices $0, h, h+i|h|$, and $h+i|h|$ that lies in the (open) upper half-plane. In Section 4 we shall combine the crucial formula (1.5) with our bounds for $\theta(x+z)$ to give a lower bound for $\alpha(x)$, and hence to obtain a new and simpler proof of Jaffard's theorem.

We would like to point out that a straightforward modification of our arguments from Section 4 can also be used to deduce Pastor's result [20, Theorem 2.3(2)] without having to resort on Tauberian theorems for the wavelet transform. We briefly sketch this in Section 5.

We close this introduction by mentioning other generalizations of the function ϕ . Its fractional integrals

$$\phi_a(x) = \sum_{n=1}^{\infty} \frac{e(n^2 x)}{n^a}$$

are covered by the results of Jaffard [16] and Pastor only when $a > 1$. For $a \leq 1$, ϕ_a is no longer a continuous function. Interestingly, Seuret and Ubis [22] have performed a pointwise regularity analysis of the Fourier series ϕ_a for values of the parameter $a \in (1/2, 1]$ in terms of L^2 -type local Hölder exponents. Another important generalization of ϕ is obtained when one replaces the frequency n^2 by a polynomial $P(n)$ and considers

$$\phi_{P,a}(x) = \sum_{n=1}^{\infty} \frac{e(P(n)x)}{n^a}, \quad a > 1.$$

When the degree k of P is larger than 2, the analysis becomes much more difficult: one loses the underlying modularity and Poisson summation only seems to yield good control in intervals $|x - p/q| \ll q^{-k}$, which appear to be too small to obtain the pointwise regularity at the irrationals. Only partial results are known, but more can be said about the local behavior “on average” by considering the spectrum of singularities $d_{\phi_{P,a}}(\alpha)$, which is defined as the Hausdorff dimension of the set of points x at which the Hölder exponent of $\phi_{P,a}$ equals α . We refer to [5, 6] for investigations in this direction.

2. Preliminaries: Quadratic Gauss sums

The following exponential sums naturally arise in the analysis of ϕ at rational points.

Definition 2.1. Let q, p, m be integers with $(p, q) = 1$. The quadratic Gauss sum $S(q, p)$ and the generalized quadratic Gauss sum $S(q, p, m)$ are defined as

$$S(q, p) = \sum_{j=1}^q e\left(\frac{pj^2}{q}\right) \quad \text{and} \quad S(q, p, m) = \sum_{j=1}^q e\left(\frac{pj^2 + mj}{q}\right).$$

The quadratic Gauss sums were already evaluated by Gauss (see e.g. [1, Section 9.10] or [19, Section 9.3]).

Theorem 2.2. Suppose p and q are positive integers with $(p, q) = 1$. For odd n , define

$$\varepsilon_n = \begin{cases} 1 & \text{if } n \equiv 1 \pmod{4}, \\ i & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

Then

$$S(q, p) = \begin{cases} \varepsilon_q \left(\frac{p}{q}\right) \sqrt{q} & \text{if } q \text{ is odd,} \\ 0 & \text{if } q \equiv 2 \pmod{4}, \\ (1 + i) \overline{\varepsilon_p} \left(\frac{q}{p}\right) \sqrt{q} & \text{if } q \equiv 0 \pmod{4}. \end{cases}$$

Here, $\left(\frac{p}{q}\right)$ is the Jacobi symbol (see [1, 19]).

The generalized quadratic Gauss sums $S(q, p, m)$ can be related to $S(q, p)$ as follows. Let p^* be the multiplicative inverse of $p \pmod{q}$. Suppose first that $m \equiv 2m' \pmod{q}$ for some m' . Then we can complete the square to get

$$S(q, p, m) = \sum_{j=1}^q e\left(\frac{p(j + p^*m')^2}{q}\right) e\left(-\frac{p^*m'^2}{q}\right) = e\left(-\frac{p^*m'^2}{q}\right) S(q, p). \quad (2.1)$$

If there is no such m' , then q is even and m odd. In this case we have

$$\begin{aligned} S(4q, p) &= \sum_{j=1}^{2q} e\left(\frac{p(2j + p^*m)^2}{4q}\right) + \sum_{j=1}^{2q} e\left(\frac{p(2j)^2}{4q}\right) \\ &= 2e\left(\frac{p^*m^2}{4q}\right)S(q, p, m) + 2S(q, p), \end{aligned}$$

since $2j + p^*m$ runs over all odd residues mod $4q$ when j runs over $\{1, \dots, 2q\}$. Therefore,

$$S(q, p, m) = \begin{cases} \frac{1}{2}e\left(-\frac{p^*m^2}{4q}\right)S(4q, p) & \text{if } q \equiv 2 \pmod{4}, \\ 0 & \text{if } q \equiv 0 \pmod{4}. \end{cases} \quad (2.2)$$

3. Behavior at rational points

In this section we deduce an asymptotic expansion for ϕ at every rational number. We first prove a simple but crucial lemma that describes the behavior of θ near rationals¹. This lemma will be used again in Section 4 to derive bounds for θ near irrational points. For a complex number $z \neq 0$, we define $z^{-1/2}$ via the principal branch of the logarithm continuously extended to the negative real axis from the upper half-plane, i.e., $\arg(z) \in (-\pi, \pi]$. Accordingly, our convention is thus $t^{1/2} = i|t|^{1/2}$ for $t < 0$, which simplifies the writing of some formulas below.

Lemma 3.1. *Suppose $1 \leq p \leq q$, $(p, q) = 1$ and $\text{Im } z > 0$. Then*

$$\theta\left(\frac{p}{q} + z\right) = \frac{e^{\pi i/4}}{q\sqrt{2}}z^{-1/2}\left(S(q, p) + 2\sum_{m=1}^{\infty} S(q, p, m)\exp\left(-\frac{i\pi m^2}{2q^2z}\right)\right).$$

Proof. Rearranging terms according to their value mod q , we write

$$\theta\left(\frac{p}{q} + z\right) = \sum_{n \in \mathbb{Z}} e\left(\frac{pn^2}{q}\right)e(n^2z) = \sum_{j=1}^q e\left(\frac{pj^2}{q}\right) \sum_{n \in j+q\mathbb{Z}} e(n^2z).$$

For fixed z , the function $f_z: \mathbb{R} \rightarrow \mathbb{C}$, $f_z: t \mapsto e(zt^2)$ has Fourier transform

$$\hat{f}_z(u) = \int_{-\infty}^{\infty} f_z(t)e^{-2\pi iut} dt = \frac{e^{\pi i/4}}{\sqrt{2}}z^{-1/2}\exp\left(-\frac{i\pi u^2}{2z}\right).$$

An application of the well-known Poisson summation formula then yields

$$\theta\left(\frac{p}{q} + z\right) = \frac{e^{\pi i/4}}{q\sqrt{2}}z^{-1/2}\sum_{j=1}^q e\left(\frac{pj^2}{q}\right)\sum_{m \in \mathbb{Z}} e\left(\frac{mj}{q}\right)\exp\left(-\frac{i\pi m^2}{2q^2z}\right)$$

¹In the language of modular forms, this lemma gives the expansion of θ at the cusps.

$$\begin{aligned}
 &= \frac{e^{\pi i/4}}{q\sqrt{2}} z^{-1/2} \sum_{m \in \mathbb{Z}} S(q, p, m) \exp\left(-\frac{i\pi m^2}{2q^2 z}\right) \\
 &= \frac{e^{\pi i/4}}{q\sqrt{2}} z^{-1/2} \left(S(q, p) + 2 \sum_{m=1}^{\infty} S(q, p, m) \exp\left(-\frac{i\pi m^2}{2q^2 z}\right) \right). \quad \blacksquare
 \end{aligned}$$

Define the “twisted” ϕ -function

$$\phi_{q,p}(z) = \sum_{m=1}^{\infty} \frac{S(q, p, m)}{2\pi i m^2} e(m^2 z).$$

Lemma 3.1 allows us to give a short proof of the following theorem, essentially due to Smith [23] and Itatsu [15] (cf. [7, 14]; a generalization of Theorem 3.2 is given in [5, Corollary 2.4]).

Theorem 3.2. *Let p and q be integers, $q \geq 1$, $(p, q) = 1$. Then*

$$\phi(p/q + h) = \phi(p/q) + C_{p/q}^- |h|_-^{1/2} + C_{p/q}^+ |h|_+^{1/2} - h/2 + R_{q,p}(h),$$

where $C_{p/q}^{\pm}$ are given by

$$C_{p/q}^- = \frac{e^{3\pi i/4}}{q\sqrt{2}} S(q, p) \quad \text{and} \quad C_{p/q}^+ = \frac{e^{\pi i/4}}{q\sqrt{2}} S(q, p), \quad (3.1)$$

and $R_{q,p}(h)$ satisfies the estimate $R_{q,p}(h) \ll q^{3/2} |h|^{3/2}$. Furthermore, $C_{p/q}^- = C_{p/q}^+ = 0$ (and hence ϕ is differentiable at p/q) if and only if $q \equiv 2 \pmod{4}$.

Proof. Suppose $y > 0$. By equation (1.4),

$$\phi\left(\frac{p}{q} + h + iy\right) = \phi\left(\frac{p}{q} + iy\right) + \frac{1}{2} \int_{iy}^{h+iy} \theta\left(\frac{p}{q} + \zeta\right) d\zeta - \frac{1}{2}h.$$

Using Lemma 3.1 and integrating by parts,

$$\begin{aligned}
 \int_{iy}^{h+iy} \theta\left(\frac{p}{q} + \zeta\right) d\zeta &= \frac{e^{\pi i/4}}{q\sqrt{2}} \left(S(q, p) \left[2\zeta^{1/2} \right]_{iy}^{h+iy} \right. \\
 &\quad \left. + 2 \int_{iy}^{h+iy} \zeta^{-1/2} (4q^2 \zeta^2) \left(\phi_{q,p}\left(-\frac{1}{4q^2 \zeta}\right) \right)' d\zeta \right) \\
 &= \frac{2e^{\pi i/4}}{q\sqrt{2}} \left(S(q, p) \left[\zeta^{1/2} \right]_{iy}^{h+iy} + \left[4q^2 \zeta^{3/2} \phi_{q,p}\left(-\frac{1}{4q^2 \zeta}\right) \right]_{iy}^{h+iy} \right. \\
 &\quad \left. - 6q^2 \int_{iy}^{h+iy} \zeta^{1/2} \phi_{q,p}\left(-\frac{1}{4q^2 \zeta}\right) d\zeta \right).
 \end{aligned}$$

All the occurring functions have continuous extensions to \mathbb{R} . Letting $y \rightarrow 0^+$ we obtain the desired result, with the constants $C_{p/q}^\pm$ as in (3.1) and with

$$R_{q,p}(h) = -4q \frac{e^{-3\pi i \operatorname{sgn} h/4}}{\sqrt{2}} \phi_{q,p} \left(-\frac{1}{4q^2 h} \right) |h|^{3/2} - 6q \frac{e^{\pi i/4}}{\sqrt{2}} \int_0^h t^{1/2} \phi_{q,p} \left(-\frac{1}{4q^2 t} \right) dt. \quad (3.2)$$

The fact that the coefficients $C_{p/q}^\pm$ are both zero if and only if $q \equiv 2 \pmod{4}$ is an immediate consequence of Theorem 2.2. The bound $R_{q,p}(h) \ll q^{3/2} |h|^{3/2}$ easily follows since $\phi_{q,p} \ll \sqrt{q}$, in view of (2.1), (2.2), and Theorem 2.2. ■

Iterating the integration by parts procedure, we obtain a full asymptotic series for the remainder $R_{q,p}$. Indeed, for any $K \in \mathbb{N}$,

$$R_{q,p}(h) = -\frac{e^{-3\pi i \operatorname{sgn} h/4}}{\sqrt{2}} \sum_{k=0}^K a_k q^{2k+1} \phi_{q,p}^{(-k)} \left(-\frac{1}{4q^2 h} \right) e^{k\pi i(1-\operatorname{sgn} h)/2} |h|^{k+3/2} - \frac{e^{\pi i/4}}{\sqrt{2}} (K+3/2) a_K q^{2K+1} \int_0^h t^{K+1/2} \phi_{q,p}^{(-K)} \left(-\frac{1}{4q^2 t} \right) dt,$$

where²

$$a_k = (-1)^k 4^{k+1} \prod_{j=1}^k (j+1/2)$$

and $\phi_{q,p}^{(-k)}$ stands for the k th-order primitive

$$\phi_{q,p}^{(-k)}(x) = \sum_{m=1}^{\infty} \frac{S(q,p,m)}{(2\pi i m^2)^{k+1}} e(m^2 x).$$

A similar asymptotic series was obtained by Duistermaat in [7].

Inspecting the k th term in this asymptotic series, we see that it is of the form $|h|^{3/2+k} g_k^\pm(|h|^{-1})$, where $\pm = \operatorname{sgn} h$ and where the functions g_k^\pm are $4q^2$ -periodic with zero mean and global Hölder regularity $1/2+k$. One readily verifies that $R_{q,p}$ is a so-called *trigonometric chirp* at 0 of type $(3/2, 1)$ and of regularity $1/2$. The latter refines a theorem of Jaffard and Meyer [17, Theorem 7.1] for Riemann's function; see [17, p. 73] for the precise definition of a trigonometric chirp. The prototypical example of a trigonometric chirp at 0 of type (α, β) , $\alpha > -1$, $\beta > 0$ is the function $|x|^\alpha \sin(|x|^{-\beta})$.

Using the explicit expression for $S(q,p)$ given by Theorem 2.2, we can exhibit the behavior of $\operatorname{Re}(\phi(p/q+h) - \phi(p/q))$ in a precise fashion, which we summarize

²For $k=0$, the product equals 1 in accordance with the empty product convention.

$q \bmod 4$	$p \bmod 4$	$h < 0$	$h > 0$
1	any	$-\left(\frac{p}{q}\right) \frac{1}{2\sqrt{q}} \sqrt{ h } + O_q(h)$	$\left(\frac{p}{q}\right) \frac{1}{2\sqrt{q}} \sqrt{h} + O_q(h)$
3	any	$-\left(\frac{p}{q}\right) \frac{1}{2\sqrt{q}} \sqrt{ h } + O_q(h)$	$-\left(\frac{p}{q}\right) \frac{1}{2\sqrt{q}} \sqrt{h} + O_q(h)$
2	any	$-\frac{1}{2}h + O(q^{3/2} h ^{3/2})$	$-\frac{1}{2}h + O(q^{3/2}h^{3/2})$
0	1	$-\left(\frac{q}{p}\right) \frac{1}{\sqrt{q}} \sqrt{ h } + O_q(h)$	$-\frac{1}{2}h + O(q^{3/2}h^{3/2})$
0	3	$-\frac{1}{2}h + O(q^{3/2} h ^{3/2})$	$\left(\frac{q}{p}\right) \frac{1}{\sqrt{q}} \sqrt{h} + O_q(h)$

Table 1. Behavior of $\operatorname{Re}(\phi(p/q + h) - \phi(p/q)) = \sum_{n=1}^{\infty} \frac{\sin(2\pi n^2 x)}{2\pi n^2}$

in Table 1. Note that at some rational points the function $\operatorname{Re} \phi$ has a (finite) left (resp. right) derivative, but an infinite right (resp. left) derivative. The table displays the behavior of the 1-periodic function $\operatorname{Re}(\phi(p/q + h) - \phi(p/q))$. By rescaling $\operatorname{Re} \phi$ by a factor of $1/2$, we obtain the well-known regularity of the 2-periodic Riemann’s function f (1.1) at rational points.

Corollary 3.3. *Suppose $r = p/q$ is rational. If p and q are both odd, then f is differentiable at r and its Hölder exponent at r is $3/2$; otherwise the Hölder exponent of f at r equals $1/2$.*

4. Behavior at irrational points

We now investigate the behavior of ϕ at irrational points ρ . Unlike in the rational case, we will not be able to derive an asymptotic formula for ϕ near ρ . Instead, we will determine the Hölder exponent $\alpha(\rho)$ introduced in (1.3).

We need some preparation in order to state the formula for $\alpha(\rho)$. Denote the n th convergent in the continued fraction expansion of ρ by $r_n = p_n/q_n$, where $(p_n, q_n) = 1$. The quality of the approximation of ρ by r_n is quantified by the number τ_n , which is defined via the relation

$$|\rho - r_n| = \left(\frac{1}{q_n}\right)^{\tau_n}.$$

Let $(r_{n_k})_k$ be the subsequence³ of convergents r_{n_k} with $q_{n_k} \not\equiv 2 \pmod{4}$, and set

$$\tau(\rho) := \limsup_{k \rightarrow \infty} \tau_{n_k}. \quad (4.1)$$

Theorem 4.1. *Let ρ be irrational. The Hölder exponent $\alpha(\rho)$ of ϕ at ρ is given by*

$$\alpha(\rho) = \frac{1}{2} + \frac{1}{2\tau(\rho)}. \quad (4.2)$$

The same result also holds for the Hölder exponent at ρ of $\operatorname{Re} \phi$ and $\operatorname{Im} \phi$.

The rest of this section is devoted to the proof of Theorem 4.1, which consists of two parts, namely, establishing the two inequalities \geq and \leq in (4.2). Let us first recall some basic properties of continued fractions (we refer to [21] for proofs and more advanced properties). The continued fractions have the following properties: for every $n \in \mathbb{N}$, $\tau_n > 2$, consecutive convergents r_n and r_{n+1} lie on different sides of ρ , $|\rho - r_{n+1}| < |\rho - r_n|$, and

$$p_{n+1}q_n - p_nq_{n+1} = (-1)^n. \quad (4.3)$$

Since $|r_n - r_{n+1}| = 1/(q_nq_{n+1})$, we have

$$|\rho - r_n| \leq \frac{1}{q_nq_{n+1}} \leq 2|\rho - r_n|,$$

so

$$\left(\frac{1}{q_n}\right)^{\tau_n-1} \leq \frac{1}{q_{n+1}} \leq 2\left(\frac{1}{q_n}\right)^{\tau_n-1}. \quad (4.4)$$

Note also that in view of (4.3), we have that q_n and q_{n+1} are never both congruent to 2 modulo 4.

4.1. The lower bound for $\alpha(\rho)$

The lower bound for $\alpha(\rho)$ was first found by Jaffard [16] by means of Tauberian arguments involving the continuous wavelet transform. To estimate the wavelet transform, Jaffard deduced bounds⁴ for the theta function near the irrational number ρ . We will present a simple proof of these bounds, using Lemma 3.1. Furthermore, we will show how these bounds directly furnish the lower bound for $\alpha(\rho)$, without needing to pass through the wavelet transform.

³Using a basic property of continued fractions (see (4.3) below), it is readily seen that this is an infinite subsequence.

⁴Recently, Pastor has extended such bounds to modular forms, see Lemma 5.2 below.

Comparing the sum with an integral, we immediately obtain the following estimate:

$$\sum_{n=1}^{\infty} e(n^2 z) \ll y^{-1/2} \quad (4.5)$$

for $z = x + iy$ with $y > 0$. Here and below ‘ \ll ’ stands for Vinogradov’s notation.

Proposition 4.2. *Let $z = x + iy$ with $y > 0$. For each $\varepsilon > 0$ there exists a $\delta = \delta(\rho, \varepsilon) > 0$ such that for $|z| < \delta$, $y > 0$ the following bound holds:*

$$\theta(\rho + z) \ll |z|^{\frac{1}{2\tau(\rho)} - \varepsilon - \frac{1}{2}} + y^{-1/2} |z|^{\frac{1}{2\tau(\rho)} - \varepsilon}. \quad (4.6)$$

Proof. We first derive bounds for θ near a rational p/q with $(p, q) = 1$. By Lemma 3.1,

$$\theta\left(\frac{p}{q} + \zeta\right) = \frac{e^{\pi i/4}}{q\sqrt{2}} \zeta^{-1/2} \sum_{m \in \mathbb{Z}} S(q, p, m) \exp\left(-\frac{\pi i}{2q^2 \zeta} m^2\right).$$

By the results obtained in Section 2, we have that $S(q, p, m) \ll \sqrt{q}$. Estimating via (4.5),

$$\theta\left(\frac{p}{q} + \zeta\right) \ll \frac{|S(q, p)|}{q|\zeta|^{1/2}} + \frac{\sqrt{q}|\zeta|^{1/2}}{(\operatorname{Im} \zeta)^{1/2}}. \quad (4.7)$$

Let N be such that $n \geq N$ implies $\tau_n \leq \tau(\rho) + \varepsilon'$ whenever $q_n \not\equiv 2 \pmod{4}$, and where ε' is such that $1/(2\tau(\rho) + 2\varepsilon') = 1/(2\tau(\rho)) - \varepsilon$. Set $\delta := 2|\rho - r_N|$. For z with $|z| \leq \delta$, let n be the unique integer larger than N such that $2|\rho - r_{n+1}| < |z| \leq 2|\rho - r_n|$, and set $\zeta = z + (\rho - r_{n+1})$. Then

$$\frac{1}{2}|z| \leq |\zeta| \leq \frac{3}{2}|z|, \quad \operatorname{Im} \zeta = \operatorname{Im} z = y. \quad (4.8)$$

Suppose first that $q_n \not\equiv 2 \pmod{4}$. We then apply (4.7) with $p = p_{n+1}$, $q = q_{n+1}$. For the second term, we use (4.4) to see that

$$\sqrt{q_{n+1}} \leq q_n^{\frac{\tau_n - 1}{2}} = |\rho - r_n|^{\frac{1}{2\tau_n} - \frac{1}{2}} \leq \sqrt{2}|z|^{\frac{1}{2\tau_n} - \frac{1}{2}}.$$

Since $1/(2\tau_n) \geq 1/(2\tau(\rho)) - \varepsilon$ (because $q_n \not\equiv 2 \pmod{4}$) and $|\zeta| \asymp |z|$, this second term is of the desired order. The first term vanishes if $q_{n+1} \equiv 2 \pmod{4}$, while otherwise we have

$$\frac{1}{\sqrt{q_{n+1}}} = |\rho - r_{n+1}|^{\frac{1}{2\tau_{n+1}}} \leq |z|^{\frac{1}{2\tau(\rho)} - \varepsilon},$$

so this first term is also of the desired order.

Suppose now that $q_n \equiv 2 \pmod{4}$. We then apply (4.7) with $p = p_n$, $q = q_n$ and get

$$\theta(\rho + z) = \theta\left(\frac{p_n}{q_n} + \left(\zeta + \frac{p_{n+1}}{q_{n+1}} - \frac{p_n}{q_n}\right)\right) \ll \frac{\sqrt{q_n}}{\sqrt{y}} \left| \zeta + \frac{(-1)^n}{q_{n+1} q_n} \right|^{1/2}$$

$$= \frac{1}{\sqrt{q_{n+1}}\sqrt{y}} |\zeta q_n q_{n+1} + (-1)^n|^{1/2} \ll \frac{1}{\sqrt{q_{n+1}}\sqrt{y}}.$$

In the first estimate we employed (4.3), while in the last estimate we used that $|\zeta| \ll |\rho - r_n| \leq 1/(q_n q_{n+1})$. Since $q_n \equiv 2 \pmod{4}$, we have that $q_{n+1} \not\equiv 2 \pmod{4}$ (by (4.3)), so that we can bound $(q_{n+1})^{-1/2}$ like before. ■

Using this bound for θ near ρ , we now deduce the lower bound for $\alpha(\rho)$. Pick $\varepsilon > 0$ arbitrarily, and use Proposition 4.2 to find a $\delta > 0$ such that the bound (4.6) holds for $|z| < \delta$. Suppose $|h| < \delta/\sqrt{2}$. We use again (1.4), so that

$$\phi(\rho + h) - \phi(\rho) = -\frac{1}{2}h + \frac{1}{2} \lim_{y \rightarrow 0^+} \int_{iy}^{h+iy} \theta(\rho + z) dz.$$

By Cauchy's theorem, the limit of this integral equals

$$\int_0^{i|h|} \theta(\rho + z) dz + \int_{i|h|}^{h+i|h|} \theta(\rho + z) dz - \int_h^{h+i|h|} \theta(\rho + z) dz =: I_1 + I_2 + I_3.$$

Using the bounds (4.6), we get

$$\begin{aligned} I_1 &\ll \int_0^{|h|} y^{-\frac{1}{2} + \frac{1}{2\tau(\rho)} - \varepsilon} \ll |h|^{\frac{1}{2} + \frac{1}{2\tau(\rho)} - \varepsilon}, \\ I_2 &\ll |h|^{-\frac{1}{2} + \frac{1}{2\tau(\rho)} - \varepsilon} \cdot |h| = |h|^{\frac{1}{2} + \frac{1}{2\tau(\rho)} - \varepsilon}, \\ I_3 &\ll |h|^{-\frac{1}{2} + \frac{1}{2\tau(\rho)} - \varepsilon} \cdot |h| + |h|^{\frac{1}{2\tau(\rho)} - \varepsilon} \int_0^{|h|} y^{-1/2} dy \ll |h|^{\frac{1}{2} + \frac{1}{2\tau(\rho)} - \varepsilon}. \end{aligned}$$

Since ε was arbitrary, $\alpha(\rho) \geq 1/2 + 1/(2\tau(\rho))$. A fortiori, this lower bound also holds for the Hölder exponent at ρ of the real and imaginary part of ϕ .

4.2. The upper bound for $\alpha(\rho)$

An upper bound for the Hölder exponent at ρ can be obtained from the expansion of ϕ at rationals, and was first done by Duistermaat [7, Proposition 5.2]. For the sake of being self-contained, we repeat his proof here.

Let $\varepsilon > 0$ be arbitrary, and let $(r_l)_l$ be a subsequence of $(r_n)_n$ with the properties that $q_l \not\equiv 2 \pmod{4}$ and that $\tau_l \geq \tau(\rho) - \varepsilon$. We will construct a sequence of points $(h_l)_l$ such that $h_l \rightarrow 0$ and $\phi(\rho + h_l) - \phi(\rho)$ is bounded from below by a constant multiple of $|h_l|^{1/2 + 1/(2(\tau(\rho) - \varepsilon))}$. We will do this by exploiting the square root behavior of ϕ in r_l . Set

$$x_l = \lambda|\rho - r_l|, \tag{4.9}$$

where λ is a fixed positive constant, independent of l , to be determined later. Using Theorem 3.2 and Theorem 2.2 we see that

$$|\phi(r_l + x_l) - \phi(r_l)| \geq \frac{x_l^{1/2}}{\sqrt{2q_l}} - \frac{1}{2}x_l + O((q_l x_l)^{3/2}),$$

where the big- O constant is absolute (independent of q_l). By equation (4.9), and using that $|\rho - r_l| \leq q_l^{-2}$, this is at least

$$\frac{\sqrt{\lambda}}{\sqrt{q_l}} |\rho - r_l|^{1/2} \left(\frac{1}{\sqrt{2}} - \frac{1}{2\sqrt{q_l}} \sqrt{\lambda} + O(\lambda) \right).$$

If we now fix a $\lambda > 0$ sufficiently small, then for q_l sufficiently large

$$|\phi(r_l + x_l) - \phi(r_l)| \gg \frac{\sqrt{\lambda}}{\sqrt{q_l}} |\rho - r_l|^{1/2}.$$

Using that $q_l^{-(\tau(\rho)-\varepsilon)} \geq |\rho - r_l|$, we get

$$|\phi(r_l + x_l) - \phi(r_l)| \gg |\rho - r_l|^{\frac{1}{2} + \frac{1}{2(\tau(\rho)-\varepsilon)}} \asymp |x_l|^{\frac{1}{2} + \frac{1}{2(\tau(\rho)-\varepsilon)}}.$$

Finally, since $|\phi(\rho) - \phi(r_l)|$ and $|\phi(\rho) - \phi(r_l + x_l)|$ are not both smaller than the term $|\phi(r_l + x_l) - \phi(r_l)|/2$, we can take $h_l = r_l - \rho$ or $h_l = r_l + x_l - \rho$ such that $|\phi(\rho) - \phi(\rho + h_l)|$ is maximal, and we get

$$|\phi(\rho) - \phi(\rho + h_l)| \gg |h_l|^{\frac{1}{2} + \frac{1}{2(\tau(\rho)-\varepsilon)}}, \quad h_l \rightarrow 0.$$

Since ε was arbitrary, this shows that $\alpha(\rho) \leq 1/2 + 1/(2\tau(\rho))$.

With a small modification, the above argument shows that the same upper bound also holds for the Hölder exponent at ρ of the real and imaginary part of ϕ . Indeed, using the same notation as above, we now define x_l by setting $|x_l| = \lambda|\rho - r_l|$ and by choosing the sign of x_l so that $r_l + x_l$ lies on the side where the square root behavior is present (see Table 1 for $\text{Re } \phi$; for $\text{Im } \phi$ one can make a similar table).

5. Behavior at irrationals for fractional integrals of modular forms

As we indicated in the introduction, our ideas can also be applied to simplify the proof of a theorem of Pastor on the pointwise Hölder exponents of modular forms at the irrationals.

We adopt here similar notations as in Pastor's paper [20]. We let g be a modular form of weight k for a subgroup Γ of finite index of $\mathrm{SL}(2, \mathbb{Z})$, let⁵

$$g(z) = \sum_{n=0}^{\infty} c_n e(nz)$$

be its Fourier expansion at ∞ , and define for $a > 0$ its a -fractional integral as

$$g_a(z) := \sum_{n=1}^{\infty} \frac{c_n}{n^a} e(nz),$$

which converges uniformly for $z = x \in \mathbb{R}$ if $a > k$ in general, and for $a > k/2$ whenever g is a cusp form. Let $\beta(\rho)$ be the pointwise Hölder exponent of g_a at an irrational number ρ . It was shown in [4] that $\beta(\rho) = a - k/2$ if g is a cusp form. The corresponding computation of $\beta(\rho)$ when g is not a cusp form is one of the main results from [20]. In the remainder of this section we assume that g is *not a cusp form* and that $a > k$.

Theorem 5.1 ([20, Theorem 2.3(2)]). *Let ρ be irrational and let⁶*

$$\tau(\rho) := \sup \left\{ \tau : \left| \rho - \frac{p}{q} \right| \ll \frac{1}{q^\tau} \text{ for infinitely many noncuspidal rationals } \frac{p}{q} \right\}. \quad (5.1)$$

Then, the Hölder exponent $\beta(\rho)$ of g_a at ρ is given by

$$\beta(\rho) = a - k \left(1 - \frac{1}{\tau(\rho)} \right).$$

Note that the case of Riemann's (complex) function (1.2) corresponds to g being the theta function, which is modular of weight $1/2$ and for which the noncuspidal rationals are precisely those rationals p/q with $q \not\equiv 2 \pmod{4}$.

In order to prove Theorem 5.1, Pastor uses a Tauberian argument involving the wavelet transform, in combination with bounds of g near rationals. In this section, we show that the evaluation of the Hölder exponent $\beta(\rho)$ follows directly from these bounds of g by using basic complex analysis. The bounds on g are expressed in the following lemma of Pastor, which extends Proposition 4.2 to general modular forms, and which we shall use as a black box.

⁵For simplicity, we assume (without loss of generality) that the modular form is holomorphic at the cusp at ∞ and that this cusp has width 1.

⁶One always has $\tau(\rho) \geq 2$.

Lemma 5.2 ([20, Lemma 3.3]). *Let ρ be irrational and $\tau = \tau(\rho)$ be given by (5.1). For each $\varepsilon > 0$, the following bounds hold:*

$$g(x + iy) \ll y^{\frac{k}{\tau} - \varepsilon - k} + y^{-k} |x + iy - \rho|^{\frac{k}{\tau} - \varepsilon} \quad \text{for } 0 < y < 1/2, \quad (5.2)$$

and

$$g(\rho + iy) \gg y^{-k + \frac{k}{\tau} + \varepsilon} \quad \text{infinitely often as } y \rightarrow 0^+. \quad (5.3)$$

We slightly adapt our method from Section 4.1 in order to obtain the lower bound for $\beta(\rho)$. The starting point is the following integral representation for g_a ([20, Eq. (3.3)]):

$$g_a(x) = \frac{(2\pi)^a}{i^a \Gamma(a)} \int_x^{x+i\infty} (z-x)^{a-1} (g(z) - c_0) dz.$$

Let J be the integer such that $a - (1 - 1/\tau)k \in (J, J + 1]$. Then by taking ε small enough in (5.2) from Lemma 5.2, one sees that the integrals⁷

$$\frac{(2\pi)^a}{i^a \Gamma(a)} \int_\rho^{\rho+i\infty} (-1)^j \binom{a-1}{j} (z-\rho)^{a-1-j} (g(z) - c_0) dz =: \frac{g_a^{(j)}(\rho)}{j!}, \quad j = 1, \dots, J,$$

converge absolutely. Let now h be a small number. By Cauchy's theorem we have

$$\begin{aligned} & \frac{i^a \Gamma(a)}{(2\pi)^a} \left(g_a(\rho + h) - g_a(\rho) - \sum_{j=1}^J \frac{g_a^{(j)}(\rho)}{j!} h^j \right) \\ &= \int_{\rho+h}^{\rho+ih} (z-\rho-h)^{a-1} (g(z) - c_0) dz \\ & - \int_\rho^{\rho+ih} \sum_{j=0}^J (-1)^j \binom{a-1}{j} (z-\rho)^{a-1-j} h^j (g(z) - c_0) dz \\ & + \int_{\rho+ih}^{\rho+i\infty} \left((z-\rho-h)^{a-1} - \sum_{j=0}^J (-1)^j \binom{a-1}{j} (z-\rho)^{a-1-j} h^j \right) (g(z) - c_0) dz. \end{aligned}$$

Applying Taylor's theorem and the bounds provided by Lemma 5.2 (and the exponential decay of $g(z) - c_0$ as $z \rightarrow \infty$), one readily sees that this is bounded by a constant times $|h|^{a - (1 - \frac{1}{\tau})k - \varepsilon}$, showing that $\beta(\rho) \geq a - (1 - 1/\tau)k$.

It might also be possible to apply a variant of Duistermaat's method from Section 4.2 in combination with an approximate functional equation to deduce the upper bound for $\beta(\rho)$. This would however require to exhibit explicit bounds in terms of

⁷Note that a is not necessarily an integer. As customary, $\binom{a-1}{j} = (a-1) \cdots (a-j)/j!$.

the involved parameters of the error term in e.g. [20, Theorem 2.5]. We rather choose a shorter path and shall show the upper bound via a simple Abelian argument that involves the maximum modulus principle and Pastor's oscillation bound (5.3). We now distinguish two cases according to whether a is a positive integer or not.

Assume first that a is a positive integer. Let β be such that $g_a(\rho + h) = P_\rho(h) + O(|h|^\beta)$ for a certain polynomial P_ρ . We have

$$g(z) = c_0 + \frac{a!}{(2\pi i)^{a+1}} \oint \frac{g_a(\zeta)}{(\zeta - z)^{a+1}} d\zeta, \quad \text{for } \text{Im } z > 0,$$

where the integral is over a counterclockwise oriented circle with centre z in the upper half-plane. Consider now the function $(z - \rho)^{-\beta} (g_a(z) - P_\rho(z - \rho))$. It is holomorphic for $\text{Im } z > 0$, has a continuous extension to $\{z: \text{Im } z \geq 0 \text{ and } z \neq \rho\}$, it is $O(|z - \rho|^{-\beta})$ when $z \rightarrow \rho$, and bounded on $\mathbb{R} \setminus \{\rho\}$. Hence, by the Phragmén–Lindelöf theorem⁸, it is bounded on $\text{Im } z > 0$. Let now z be close to ρ . Integrating over a circle of radius $|z - \rho|/2$, we get

$$g(z) = c_0 + \frac{a!}{(2\pi i)^{a+1}} \oint \frac{g_a(\zeta) - P_\rho(\zeta - \rho)}{(\zeta - z)^{a+1}} d\zeta + O(1) \ll 1 + |z - \rho|^{\beta - a}, \quad \text{as } z \rightarrow \rho.$$

Comparing this to (5.3) from Lemma 5.2, we see that $\beta \leq a - k(1 - 1/\tau)$.

Suppose now that a is not a positive integer, and let $N = \lceil a \rceil$. We have

$$g(z) = c_0 + \frac{1}{i^N (2\pi)^a \Gamma(N - a)} \int_0^\infty t^{N-a} g_a^{(N)}(z + it) \frac{dt}{t}.$$

If we again assume that $g_a(\rho + h) = P_\rho(h) + O(|h|^\beta)$, then by the first part we have $g_a^{(N)}(z) \ll |z - \rho|^{\beta - N} + 1$ as $z \rightarrow \rho$. Hence,

$$\begin{aligned} g(\rho + iy) &= c_0 + \frac{1}{i^N (2\pi)^a \Gamma(N - a)} \int_0^1 t^{N-a} g_a^{(N)}(\rho + i(y + t)) \frac{dt}{t} + O(1) \\ &\ll \int_0^1 t^{N-a} (y + t)^{\beta - N} \frac{dt}{t} + 1 \ll y^{\beta - a}, \quad \text{as } y \rightarrow 0, \end{aligned}$$

provided that $\beta < a$, which we may assume without loss of generality. Comparing again with (5.3) this yields $\beta \leq a - k(1 - 1/\tau)$.

Funding. F. Broucke was partly supported by the Ghent University through the BOF-grant 01J04017 and partly by a postdoctoral fellowship (grant number 12ZZH23N)

⁸We apply here the classical Phragmén–Lindelöf principle after a transformation $\zeta = -1/(z - \rho)$ to functions holomorphic in the open sector $0 < \arg \zeta < \pi$ with continuous extension to the boundary.

of the Research Foundation – Flanders. J. Vindas was partly supported by Ghent University through the BOF-grant 01J04017 and by the Research Foundation–Flanders through the FWO-grant G067621N.

References

- [1] T. M. Apostol, *Introduction to analytic number theory*. Undergraduate Texts in Mathematics, Springer, New York-Heidelberg, 1976 Zbl 0335.10001 MR 434929
- [2] V. Banica and L. Vega, [Riemann’s non-differentiable function and the binormal curvature flow](#). *Arch. Ration. Mech. Anal.* **244** (2022), no. 2, 501–540 Zbl 1506.35153 MR 4408174
- [3] A. Boritchev, D. Eceizabarrena, and V. V. Da Rocha, [Intermittency of Riemann’s non-differentiable function through the fourth-order flatness](#). *J. Math. Phys.* **62** (2021), no. 9, article no. 093101 Zbl 1500.76046 MR 4309227
- [4] F. Chamizo, I. Petrykiewicz, and S. Ruiz-Cabello, [The Hölder exponent of some Fourier series](#). *J. Fourier Anal. Appl.* **23** (2017), no. 4, 758–777 Zbl 1376.42004 MR 3685989
- [5] F. Chamizo and A. Ubis, [Some Fourier series with gaps](#). *J. Anal. Math.* **101** (2007), 179–197 Zbl 1211.42003 MR 2346544
- [6] F. Chamizo and A. Ubis, [Multifractal behavior of polynomial Fourier series](#). *Adv. Math.* **250** (2014), 1–34 Zbl 1293.42002 MR 3122161
- [7] J. J. Duistermaat, Self-similarity of “Riemann’s nondifferentiable function”. *Nieuw Arch. Wisk. (4)* **9** (1991), no. 3, 303–337 Zbl 0760.26009 MR 1166151
- [8] D. Eceizabarrena, [Geometric differentiability of Riemann’s non-differentiable function](#). *Adv. Math.* **366** (2020), article no. 107091 Zbl 1437.26005 MR 4072795
- [9] D. Eceizabarrena, [On the Hausdorff dimension of Riemann’s non-differentiable function](#). *Trans. Amer. Math. Soc.* **374** (2021), no. 11, 7679–7713 Zbl 1481.26003 MR 4328680
- [10] J. Gerver, [The differentiability of the Riemann function at certain rational multiples of \$\pi\$](#) . *Amer. J. Math.* **92** (1970), 33–55 Zbl 0203.05904 MR 265525
- [11] J. Gerver, [More on the differentiability of the Riemann function](#). *Amer. J. Math.* **93** (1971), 33–41 Zbl 0228.26008 MR 276422
- [12] G. H. Hardy, [Weierstrass’s non-differentiable function](#). *Trans. Amer. Math. Soc.* **17** (1916), no. 3, 301–325 Zbl 46.0401.03 MR 1501044
- [13] G. H. Hardy and J. E. Littlewood, [Some problems of diophantine approximation](#). *Acta Math.* **37** (1914), no. 1, 193–239 Zbl 45.0305.03 MR 1555099
- [14] M. Holschneider and P. Tchamitchian, [Pointwise analysis of Riemann’s “nondifferentiable” function](#). *Invent. Math.* **105** (1991), no. 1, 157–175 Zbl 0741.26004 MR 1109624
- [15] S. Itatsu, [Differentiability of Riemann’s function](#). *Proc. Japan Acad. Ser. A Math. Sci.* **57** (1981), no. 10, 492–495 Zbl 0501.26004 MR 640257
- [16] S. Jaffard, [The spectrum of singularities of Riemann’s function](#). *Rev. Mat. Iberoamericana* **12** (1996), no. 2, 441–460 Zbl 0889.26005 MR 1402673

- [17] S. Jaffard and Y. Meyer, [Wavelet methods for pointwise regularity and local oscillations of functions](#). *Mem. Amer. Math. Soc.* **123** (1996), no. 587 Zbl 0873.42019 MR 1342019
- [18] E. Mohr, [Wo ist die Riemannsche Funktion \$\sum_{n=1}^{\infty} \frac{\sin n^2 x}{n^2}\$ nichtdifferenzierbar?](#) *Ann. Mat. Pura Appl. (4)* **123** (1980), 93–104 Zbl 0456.26003 MR 581926
- [19] H. L. Montgomery and R. C. Vaughan, [Multiplicative number theory. I. Classical theory](#). Cambridge Studies in Advanced Mathematics 97, Cambridge University Press, Cambridge, 2006 Zbl 1142.11001 MR 2378655
- [20] C. Pastor, [On the regularity of fractional integrals of modular forms](#). *Trans. Amer. Math. Soc.* **372** (2019), no. 2, 829–857 Zbl 1440.42178 MR 3968789
- [21] W. M. Schmidt, [Diophantine approximation](#). Lecture Notes in Mathematics 785, Springer, Berlin, 1980 Zbl 0421.10019 MR 568710
- [22] S. Seuret and A. Ubis, [Local \$L^2\$ -regularity of Riemann's Fourier series](#). *Ann. Inst. Fourier (Grenoble)* **67** (2017), no. 5, 2237–2264 Zbl 1412.42008 MR 3732690
- [23] A. Smith, [The differentiability of Riemann's functions](#). *Proc. Amer. Math. Soc.* **34** (1972), 463–468 Zbl 0254.26013 MR 308337

Received 9 November 2022.

Frederik Broucke

Department of Mathematics: Analysis, Logic and Discrete Mathematics, Ghent University, Krijgslaan 281, 9000 Gent, Belgium; fabrouck.broucke@ugent.be

Jasson Vindas

Department of Mathematics: Analysis, Logic and Discrete Mathematics, Ghent University, Krijgslaan 281, 9000 Gent, Belgium; jasson.vindas@ugent.be