



Singular orthotropic functionals with nonstandard growth conditions

Pierre Bousquet, Lorenzo Brasco and Chiara Leone

Abstract. We pursue the study of a model convex functional with orthotropic structure and nonstandard growth conditions, this time focusing on the sub-quadratic case. We prove that bounded local minimizers are locally Lipschitz. No restrictions on the ratio between the highest and the lowest growth rates are needed. The result holds also in presence of a non-autonomous lower order term, under sharp integrability assumptions. Finally, we prove higher differentiability of bounded local minimizers as well.

1. Introduction

1.1. Overview

In this paper, we expand on the gradient regularity theory for minimizers of functionals from the calculus of variations, having an *orthotropic* structure, in the *nonstandard growth* case. This may be seen as a follow up of our previous papers [10] and [12].

Specifically, for an open set $\Omega \subseteq \mathbb{R}^N$ and a set of exponents $1 < p_1 \leq \dots \leq p_N$, we take the anisotropic Sobolev space $W_{\text{loc}}^{1,\mathbf{p}}(\Omega)$, defined by

$$W_{\text{loc}}^{1,\mathbf{p}}(\Omega) = \{u \in L_{\text{loc}}^1(\Omega) : u_{x_i} \in L_{\text{loc}}^{p_i}(\Omega), i = 1, \dots, N\}.$$

Given a function $f \in L_{\text{loc}}^1(\Omega)$, we consider the following functional:

$$\mathfrak{F}_{\mathbf{p}}(u, \Omega') := \sum_{i=1}^N \frac{1}{p_i} \int_{\Omega'} |u_{x_i}|^{p_i} dx - \int_{\Omega'} f u dx,$$

for $u \in W_{\text{loc}}^{1,\mathbf{p}}(\Omega) \cap L_{\text{loc}}^\infty(\Omega)$ and $\Omega' \Subset \Omega$.

In the superquadratic case, i.e., for $2 \leq p_1 \leq \dots \leq p_N < \infty$, and for $f \equiv 0$, it has been recently proved in [10] that any local minimizer $U \in W_{\text{loc}}^{1,\mathbf{p}}(\Omega) \cap L^\infty(\Omega)$ is such that

$$\nabla U \in L_{\text{loc}}^\infty(\Omega, \mathbb{R}^N) \quad \text{and} \quad |U_{x_i}|^{(p_i-2)/2} U_{x_i} \in W_{\text{loc}}^{1,2}(\Omega), \quad \text{for } i = 1, \dots, N.$$

2020 *Mathematics Subject Classification*: Primary 35B65; Secondary 35J75, 49K20.

Keywords: Nonstandard growth conditions, singular elliptic equations, Lipschitz regularity, Sobolev regularity, orthotropic functionals.

The main goal of this paper is to address the same kind of regularity issues, again for *bounded* local minimizers, this time in the subquadratic case

$$1 < p_1 \leq \dots \leq p_N \leq 2.$$

However, we will obtain some regularity results which actually hold in the full range $1 < p_1 \leq \dots \leq p_N < \infty$, see the next section for more details.

We recall that $u \in W_{\text{loc}}^{1,p}(\Omega) \cap L_{\text{loc}}^\infty(\Omega)$ is a *local minimizer* of \mathfrak{F}_p in Ω if

$$\mathfrak{F}_p(u, \Omega') \leq \mathfrak{F}_p(\varphi, \Omega'), \quad \text{for every } \varphi - u \in W_0^{1,p}(\Omega') \cap L^\infty(\Omega') \text{ and every } \Omega' \Subset \Omega.$$

Here we denote by $W_0^{1,p}(\Omega')$ the completion of $C_0^\infty(\Omega')$ with respect to the norm

$$\varphi \in C_0^\infty(\Omega') \mapsto \sum_{i=1}^N \|\varphi_{x_i}\|_{L^{p_i}(\Omega')}.$$

By convexity of \mathfrak{F}_p , we have that u is a local minimizer if and only if it is a local weak solution in $W_{\text{loc}}^{1,p}(\Omega) \cap L_{\text{loc}}^\infty(\Omega)$ of the quasilinear equation

$$-\sum_{i=1}^N (|u_{x_i}|^{p_i-2} u_{x_i})_{x_i} = f, \quad \text{in } \Omega.$$

This can be seen as a particular instance of elliptic equation in the wide context of the Musielak–Orlicz spaces, see [18] for a comprehensive study on the subject.

We emphasize the fact that in this paper we just consider *bounded* minimizers u . As a consequence, we discard a priori all the counterexamples to regularity arising in the literature related to nonstandard growth variational problems, see [38, 40, 49]. For completeness, we mention that the boundedness of minimizers in this setting has already been extensively studied, see [37] for the homogeneous case $f \equiv 0$, and [20, 21] for the non-homogeneous one, see also [25–28, 44]. Moreover, we ignore the problem of the *existence* of a minimizer in $W^{1,p}$, for which we would also need to assume that f belongs to a suitable dual Sobolev space. Here instead, we assume *a priori* to have a bounded minimizer u and focus on identifying sharp conditions on the function f needed to obtain its Lipschitz continuity and higher differentiability.

The main feature of all our regularity results will be that *we do not need to impose any restriction on the ratio p_N/p_1 .*

We refer the reader to our previous papers [10–12] for an introduction to the realm of gradient regularity for minimizers of orthotropic functionals (see also [31] for an approach based on viscosity methods). We just recall here that already for the standard growth case $p_1 = p_N = p$, the superquadratic case $p > 2$ is much more involved than the case of the model functional

$$\mathfrak{L}(u; \Omega') = \int_{\Omega'} |\nabla u|^p dx, \quad \text{for } u \in W_{\text{loc}}^{1,p}(\Omega) \text{ and } \Omega' \Subset \Omega,$$

as far as the regularity of the gradient of local minimizers is concerned. On the other hand, the subquadratic case $1 < p < 2$ is simpler, in a sense: the Lipschitz continuity for

minimizers of orthotropic functionals is a consequence of a general result due to Fonseca and Fusco Theorem 2.2 in [36]), as observed in the introductions of [11] and [12].

In particular, it seems natural to try to adapt the techniques used in [36] since they allow to establish the Lipschitz regularity for the subquadratic case when $p_1 = p_N = p < 2$. However, we stress that in the case $p_1 \neq p_N$, our functional pertains to the class of variational problems with nonstandard growth conditions, following the terminology of Marcellini in [50, 51]. Then it couples in a nontrivial way the difficulties coming from the two situations: orthotropic structure and nonstandard growth conditions. Thus, even if our proof of the Lipschitz regularity is inspired from Theorem 2.2 in [36], nontrivial adaptations and intermediate results will be needed.

Finally, it is worth recalling that, in spite of a large number of papers and contributions on anisotropic problems (including for example [6, 7, 15, 17, 34, 35, 41–43, 53, 57]), a complete gradient regularity theory is still missing, even for the case of orthotropic structures. Moreover, we recall that the basic regularity for local minimizers of \mathfrak{F}_p (i.e., $C^{0,\alpha}$ estimates, Harnack inequalities, and an extension of the De Giorgi's regularity theory) is still not fully understood, see however [1, 3] and [48] for some results.

1.2. Main results

Our first result is an higher integrability statement, which is valid without any restriction on the exponents p_i . As we will see in a while, this will be instrumental to the two main regularity results of this paper. In what follows, we use the notation

$$(1.1) \quad \mathcal{G}_0(\nabla u) = \left(\sum_{i=1}^N \frac{1}{p_i} |u_{x_i}|^{p_i} - 1 \right)_+ + 1,$$

where $(\cdot)_+$ stands for the positive part. This function naturally arises from the principal part of \mathfrak{F}_p . It encodes in a natural way the full summability information for each component of the gradient. A similar idea has been considered for example in the papers [2, 23, 24], dealing with the so-called *double phase problems*.

Proposition 1.1 (Higher integrability: general growth). *Let $1 < p_1 \leq \dots \leq p_N < \infty$ and let $f \in L_{\text{loc}}^\gamma(\Omega)$, for some $\gamma \geq 2$. Then for every local minimizer $U \in W_{\text{loc}}^{1,p}(\Omega) \cap L_{\text{loc}}^\infty(\Omega)$, we have*

$$\mathcal{G}_0(\nabla U) \in L_{\text{loc}}^\gamma(\Omega).$$

Moreover, for every ball $B_R(x_0) \Subset \Omega$ such that $B_{4R}(x_0) \Subset \Omega$, as well, we have

$$(1.2) \quad \int_{B_{R/2}(x_0)} \mathcal{G}_0(\nabla U)^\gamma dx \leq \Gamma_1 + \Gamma_2 \int_{B_R(x_0)} \mathcal{G}_0(\nabla U) dx,$$

for two constants $\Gamma_1, \Gamma_2 > 0$ which depend only on $N, p_N, p_1, \gamma, R, \|f\|_{L^\gamma(B_{4R}(x_0))}$ and $\|U\|_{L^\infty(B_{4R}(x_0))}$.

Theorem L (Lipschitz regularity: subquadratic growth). *Let $1 < p_1 \leq \dots \leq p_N \leq 2$ and let $f \in L_{\text{loc}}^\gamma(\Omega)$, for some $\gamma > N$. Then every local minimizer $U \in W_{\text{loc}}^{1,p}(\Omega) \cap L_{\text{loc}}^\infty(\Omega)$ is locally Lipschitz continuous. Moreover, for every ball $B_R(x_0) \Subset \Omega$ such that $B_{4R}(x_0) \Subset \Omega$,*

with $R < 1$, we have

$$\begin{aligned} & \|\mathcal{G}_0(\nabla U)\|_{L^\infty(B_{R/4}(x_0))} \\ & \leq C \left[\frac{1}{R^{N\gamma/(\gamma-N)}} \left(\Gamma_1 + \Gamma_2 \int_{B_R(x_0)} \mathcal{G}_0(\nabla U) dx \right)^{N/(\gamma-N)} + \|f\|_{L^\gamma(B_R(x_0))}^{N\gamma/(\gamma-N)} \right] \\ & \quad \cdot \|\mathcal{G}_0(\nabla U)\|_{L^1(B_R(x_0))}. \end{aligned}$$

for some $C = C(N, p_N, p_1, \gamma) > 0$ and for the same constants Γ_1, Γ_2 as in (1.2).

Observe that the assumption $\gamma > N$ is sharp (in the scale of Lebesgue spaces) to obtain the Lipschitz continuity of U . Actually, this is already true when $p_1 = \dots = p_N = 2$. It is a remarkable fact that, even in the orthotropic case with nonstandard growth conditions, this universal assumption on f still leads to Lipschitz continuity. We refer the reader to [4] for a wide class of variational problems (not including orthotropic structures, however) where this same condition is known to guarantee Lipschitz continuity of local minimizers.

Theorem S (Sobolev regularity: subquadratic growth). *Let $1 < p_1 \leq \dots \leq p_N \leq 2$ and let*

$$(1.3) \quad f \in L_{\text{loc}}^{1+2/p_1}(\Omega).$$

Then for every local minimizer $U \in W_{\text{loc}}^{1,\mathbf{p}}(\Omega) \cap L_{\text{loc}}^\infty(\Omega)$, we have

$$\mathcal{W}_i := |U_{x_i}|^{(p_i-2)/2} U_{x_i} \in W_{\text{loc}}^{1,2}(\Omega), \quad \text{for } i = 1, \dots, N,$$

and

$$U_{x_i} \in W_{\text{loc}}^{1,\mathbf{p}}(\Omega), \quad \text{for } i = 1, \dots, N.$$

Moreover, for every ball $B_R(x_0) \Subset \Omega$ such that $B_{4R}(x_0) \Subset \Omega$, we have for every $i = 1, \dots, N$,

$$\begin{aligned} & \sum_{i=1}^N \int_{B_{R/4}(x_0)} |\nabla \mathcal{W}_i|^2 dx \\ & \leq \frac{C}{R^2} \left(\Gamma_1 + \Gamma_2 \int_{B_R(x_0)} \mathcal{G}_0(\nabla U) dx \right) + CR^{2/p_1-1} \int_{B_R(x_0)} |f|^{1+2/p_1} dx, \end{aligned}$$

and

$$\|\nabla U_{x_i}\|_{L^{p_i}(B_{R/4}(x_0), \mathbb{R}^N)} \leq \frac{2}{p_i} \|U_{x_i}\|_{L^{p_i}(B_{R/4}(x_0))}^{(2-p_i)/2} \|\nabla \mathcal{W}_i\|_{L^2(B_{R/4}(x_0), \mathbb{R}^N)},$$

for some $C = C(N, p_N, p_1) > 0$ and for the same constants Γ_1, Γ_2 as in (1.2), corresponding to the choice $\gamma = (p_1 + 2)/p_1$.

1.3. Comparison with known results

In the homogeneous case $f = 0$, Proposition 1.1 can be obtained as a consequence of Lemma 4.2 in [46]. In the superquadratic case $p_1 \geq 2$ and still for $f \equiv 0$, an alternate proof can also be found in Proposition 6.1 of [10]. We present here a new proof that takes into account the presence of the forcing term f . Our argument is certainly more elementary than the one in [10], and arguably more natural in our setting than the one in [46], in

the sense that it strongly relies on some tools that will be repeatedly used in the proofs of the other main results, see the next section for further comments.

Theorem L is the counterpart for the subquadratic case of our previous result Theorem 1.1 in [10], which deals with the superquadratic case. We shall explain in the next section why the two situations require different arguments. We point out that experts in the field may recognize Theorem L (and Theorem 1.1 in [10], as well) as a particular case of the main result in [46], at least in the homogeneous case $f \equiv 0$. However, it turns out that the proof of Proposition 2.1 in [46] is affected by a crucial flaw, we refer the reader to Remark 1.4 in [10] for a detailed discussion on this delicate point. In any case, it is fair to admit that some other parts of Lieberman's paper [46] have been an important source of inspiration for the proof of Proposition 1.1.

In Corollary 3.4 of [33], the authors prove the local Lipschitz continuity of local minimizers (not a priori bounded) of the following functional:

$$(1.4) \quad \sum_{i=1}^N \frac{1}{p_i} \int (1 + |u_{x_i}|^2)^{p_i/2} dx, \quad \text{for } \frac{2N}{N+2} < p_1 \leq \dots \leq p_N \leq 2.$$

Observe that such a functional has an orthotropic structure, with nonstandard subquadratic growth conditions, exactly as our \mathfrak{F}_p . However, it should be noticed that the functional (1.4) is *neither degenerate nor singular*: this is the crucial difference with our case. Indeed, the Hessian of the function

$$\tilde{G}(z) := \sum_{i=1}^N \frac{1}{p_i} (1 + z_i^2)^{p_i/2}, \quad \text{for } z = (z_1, \dots, z_N) \in \mathbb{R}^N,$$

satisfies

$$(p_1 - 1)(1 + |z|^2)^{(p_1-2)/2} |\xi|^2 \leq \langle D^2 \tilde{G}(z) \xi, \xi \rangle \leq |\xi|^2, \quad \text{for } z, \xi \in \mathbb{R}^N.$$

This property fails to be satisfied by our functional, where the integrand is given by

$$(1.5) \quad G(z) = \sum_{i=1}^N \frac{1}{p_i} |z_i|^{p_i}, \quad \text{for } z \in \mathbb{R}^N.$$

Even worse, in contrast with the general framework of [33], in our situation there are no continuous functions $h_1, h_2: [0, +\infty) \rightarrow [0, +\infty)$ such that

$$h_1(|z|) |\xi|^2 \leq \langle D^2 G(z) \xi, \xi \rangle \leq h_2(|z|) |\xi|^2, \quad \text{for } z, \xi \in \mathbb{R}^N,$$

even for large values of z . Indeed, $D^2 G(z)$ is given by the diagonal matrix

$$D^2 G(z) = \begin{bmatrix} (p_1 - 1) |z_1|^{p_1-2} & & \\ & \ddots & \\ & & (p_N - 1) |z_N|^{p_N-2} \end{bmatrix},$$

and each entry on the diagonal blows-up as the corresponding component of z vanishes.

As for Theorem S, we observe that this may be seen as a generalization of the following classical result for the p -Laplacian: for $1 < p < 2$, any local weak solution $u \in W_{\text{loc}}^{1,p}(\Omega)$ of

$$-\operatorname{div}(|\nabla u|^{p-2}\nabla u) = f, \quad \text{in } \Omega,$$

belongs to $W_{\text{loc}}^{2,p}(\Omega)$, provided $f \in L_{\text{loc}}^{p'}(\Omega)$ (see for example [30]). The reader may notice that for $p_1 = p_N = p$, our assumption (1.3) boils down to

$$f \in L_{\text{loc}}^{\gamma}(\Omega), \quad \text{for } \gamma \geq 1 + \frac{2}{p}.$$

Since for $1 < p < 2$ we have $1 + 2/p < p'$, this is a weaker requirement when compared with the classical result recalled above. This is not surprising, since we are now assuming that u is a priori bounded. Such an assumption is responsible for this new feature. In the standard growth case, this has been recently observed in Theorem 1.2 of [22].

Higher differentiability of local minimizers is a well-studied problem: for the specific case of orthotropic functionals with subquadratic nonstandard growth, some prior results can be found for example in Theorem 3 of [6], Corollary 1 of [5], and Theorem 2 of [17].

Finally, in the superquadratic case $p_1 \geq 2$, as already recalled, the counterpart of Theorem S has been obtained in Corollary 7.1 of [10], for $f \equiv 0$. In the case of a right-hand side $f \not\equiv 0$, some results have been obtained in Theorem 1.1 of [14] and Corollary 2 of [53] (see also Proposition 3.11 in [55] for an approach based on convex duality, leading to global estimates on the torus).

Remark 1.2 (On the C^1 regularity). In dimension $N = 2$, the C^1 regularity of a Lipschitz local minimizer essentially follows from Theorem 1.1 in [29], both in the cases $p_2 \leq 2$ and $p_1 \geq 2$, provided that $f \equiv 0$. This assertion is detailed in [8], where the ‘‘mixed’’ case $p_1 \leq 2 \leq p_2$ is considered as well. For a slightly different approach, see [47] when $p_1 \geq 2$ and [54] when $p_1 = p_2 < 2$: these references still deal with the case $f \equiv 0$. In the non-homogeneous case, the strategy followed in [9] (and originally written for $p_1 = p_2$ and $f \equiv 0$) could be adapted to more general situations, provided f satisfies suitable differentiability and summability conditions. In any case, the C^1 regularity of local minimizers when $N \geq 3$ is entirely open, even for $p_1 = p_N$ and $f \equiv 0$.

1.4. Structure of the proofs

The proofs of Proposition 1.1, Theorem L and Theorem S are based on a classical three steps strategy. We first approximate our local minimizer $U \in W_{\text{loc}}^{1,p}(\Omega) \cap L_{\text{loc}}^{\infty}(\Omega)$ by a sequence of minimizers $\{u^{\varepsilon}\}_{\varepsilon>0}$ of regularized functionals $\mathfrak{F}_{p,\varepsilon}$ having good smoothing properties. We next obtain uniform a priori estimates on these minimizers. Finally, we pass to the limit in order to transfer these a priori bounds to U .

The first step is usually quite easy: it is sufficient to perturb the initial functional by adding some uniformly convex ε -perturbation and possibly smooth out the coefficients, for example by replacing f with its mollification f^{ε} . This regularization strategy allows to avoid the usual difference quotient method, and the technicalities that go with its use in the nonstandard growth setting. However, here we face a first difficulty: remember that we are not assuming f to be in the correct dual Sobolev space. This also entails that we

do not have a good a priori L^∞ estimate at our disposal. Thus, such an approximation has to be handled with great care. We circumvent this technical difficulty by adding a nonlinear lower term in the regularized functional, which forces the minimizers u^ε to be bounded, with a uniform L^∞ bound which only depends on the local L^∞ norm of U (see Lemma 2.4). This is a technical aspect of the proof, which we believe to have its own interest. We note that this approach has features in common with [16].

The core of the matter is next to establish the a priori estimates for the gradient of u^ε , the minimizer of the regularized functional $\mathfrak{F}_{p,\varepsilon}$. As for the estimates leading to Proposition 1.1 and Theorem L, these are achieved by means of Moser-type schemes: a *slow one* and a *fast one*, respectively.

The cornerstone of these schemes is a Caccioppoli inequality for power functions of the gradients (see Proposition 3.1). In a simplified way, for every $\alpha \geq 0$ this reads as

$$(1.6) \quad \sum_{i=1}^N \int |u_{x_i}|^{p_i-2} |(G(\nabla u)^{(\alpha+1)/2})_{x_i}|^2 dx \lesssim (\alpha+1)^2 \int G(\nabla u)^{\alpha+2-2/p_N} dx,$$

where G is the same function as in (1.5). Here, for simplicity, we put $f \equiv 0$ and write u in place of u^ε .

Observe that on the left-hand side of (1.6) we have a weighted gradient of a power of $G(\nabla u)$: the weights $|u_{x_i}|^{p_i-2}$ are the typical feature of degenerate/singular orthotropic functionals. The main difficulty in getting regularity results out of this estimate is precisely due to their presence. In contrast with the Caccioppoli inequality previously obtained in Lemma 3.1 of [10] to handle the superquadratic case $p_1 \geq 2$, now these weights do not pop-up on the right-hand side. This is a crucial ingredient of the estimate: indeed, no control from above would be possible on $|u_{x_i}|^{p_i-2}$ if $p_i < 2$.

Not surprisingly, the proof of (1.6) relies on the differentiated Euler–Lagrange equation, which is nothing but the equation solved by the components of ∇u^ε . In a nutshell, the idea to reach such an estimate not containing the nasty weights $|u_{x_i}|^{p_i-2}$ on the right-hand side is that of using an integration by parts trick: this permits to trade the presence of the term D^2G , with the more tractable one DG . This idea is certainly not new in the context of singular variational problems: it goes back at least to [52], and has then become standard in the field.

However, as natural as this idea may appear, its technical implementation in our context needs some efforts: in particular, a careful choice of the test functions for the differentiated equation has to be done. Such a choice must reflect the algebraic structure of the operator, in a sense. Without entering too much into the details, we refer to the proof of Proposition 3.1 below. The choice of the correct test functions here has been suggested to us by [46], even if our choice seems to be simpler and more natural.

The Caccioppoli inequality (1.6) is first used in the proof of the higher integrability result of Proposition 1.1. More specifically, it permits to obtain a self-improving estimate of the type

$$(1.7) \quad \int G(\nabla u)^{\beta+1} dx \lesssim \|u\|_{L^\infty}^2 \beta^2 \int G(\nabla u)^{\beta+1-2/p_N} dx.$$

This is the *slow* Moser iteration scheme we were referring to above: by iterating (1.7) a finite number of times, we can conclude that $G(\nabla u)$ (and thus ∇u itself) can be estimated

in L^q , for every finite $q \geq 1$. Observe that the *additive* integrability gain at each step and the presence of the factor β^2 on the right-hand side make the previous scheme not suitable for being iterated infinitely many times. This explains why we cannot reach the limiting case $\nabla u \in L^\infty$ with this approach.

Estimates like (1.7) are quite typical in regularity theory, both in the contexts of standard and nonstandard growth problems (among others, see for example Proposition 3.1 in [32] and Theorem 3 in [19], respectively). Usually, they are obtained by coupling an integration by parts with a Caccioppoli inequality for the gradient, like the one (1.6) at our disposal. The L^∞ bound on the solution is used to treat the solution itself as a constant in the estimates.

We stress here that in this part of the proofs we do not need the restriction $p_N \leq 2$. Thus, in particular, we can extend and simplify the higher integrability result we previously obtained¹ in Proposition 4.3 of [10].

With the aid of (1.7), in the case $p_N \leq 2$ we can transpose to our situation the typical *absorption trick* which lies at the basis of the Lipschitz estimate for the standard p -Laplacian (see for example [32], Section 3). Up to some nontrivial technical issues, this consists in observing that when $G(\nabla u) \geq 1$, we have

$$|u_{x_i}|^{p_i-2} \gtrsim G(\nabla u)^{(p_i-2)/p_i}, \quad \text{since } p_i - 2 \leq 0,$$

and thus

$$(1.8) \quad \sum_{i=1}^N \int |u_{x_i}|^{p_i-2} |(G(\nabla u)^{(\alpha+1)/2})_{x_i}|^2 dx \gtrsim \sum_{i=1}^N \int |(G(\nabla u)^{(\alpha+2)/2-1/p_i})_{x_i}|^2 dx.$$

The weights $|u_{x_i}|^{p_i-2}$ have then been absorbed into a suitable power function of the gradient. In this sense, in the case $p_N \leq 2$, the presence of the weights $|u_{x_i}|^{p_i-2}$ on the left-hand side of (1.7) helps, more than it hurts.

At this point, by joining (1.8) and (1.7), the orthotropic nature of the problem completely disappears and we simply fall into the realm of nonstandard growth problems. A standard application of the Sobolev inequality makes then possible to launch a standard Moser iterative scheme (i.e., a *fast* one, with a *multiplicative* gain of integrability at each step). This permits to reach an L^∞ - L^q estimate on $G(\nabla u)$, after infinitely many iterations. This is not the end of the story. Indeed, we still have to pay attention to a detail which is quite typical of the nonstandard growth case: the exponent q in this a priori estimate could be too large. However, this preliminary estimate can be “rectified” by combining the higher integrability result of Proposition 1.1 together with an interpolation trick which decreases the initial integrability requirement on $G(\nabla u)$. We then finally get a L^∞ - L^1 estimate on $G(\nabla u)$, as desired.

In contrast to Proposition 1.1 and Theorem L, the proof of Theorem S does not rely on the Caccioppoli inequality of Proposition 3.1. The proof follows the same idea as in

¹There is however a subtle detail here: the result in [10] was obtained through a complicate self-improving iterative scheme (inspired from that of Theorem 1.1 in [7]), which *was not* of Moser-type. Actually, this was much more sophisticated and could be roughly described as follows: improvement of integrability of $N - 1$ components of the gradient entails that the missing one improves its integrability as well.

the case of the result for the familiar p -Laplacian, for the case $1 < p < 2$: we test the differentiated equation with the gradient itself u_{x_k} and perform an integration by parts as in Naumann's trick [52]. Again, this permits to avoid using the undesired upper bound on the Hessian D^2G . In order to conclude, one has to control from above terms of the form

$$u_{x_k x_k} |u_{x_i}|^{p_i-1}.$$

Observe that for every $k \neq i$, the two terms are completely decoupled. However, when $p_i \leq 2$, we can simply estimate this term from above by Young's inequality:

$$u_{x_k x_k} |u_{x_i}|^{p_i-1} \lesssim |u_{x_k}|^{p_k-2} |u_{x_k x_k}|^2 + |u_{x_k}|^{2-p_k} |u_{x_i}|^{2(p_i-1)}.$$

The first term is absorbed on the left-hand side, while the second term can be estimated from above by means of an integrability estimate (here we rely again on the information provided by Proposition 1.1). This explains why we require $p_N \leq 2$ in the statement of Theorem S.

1.5. Plan of the paper

In Section 2, we present the approximation scheme and some basic material used all along the paper. Section 3 contains the crucial Caccioppoli-type inequality for the gradient (Proposition 3.1). The latter is exploited in Section 4 to perform the slow Moser iteration leading to the higher integrability estimate needed in Proposition 1.1. The Lipschitz bound related to Theorem L is proved in Section 5, while Section 6 is devoted to the proof of the higher differentiability estimates corresponding to Theorem S. Then, in Section 7, we eventually prove our three main results by passing to the limit in the approximation scheme. Finally, for completeness, we include in Appendix A the proof of a maximum principle ensuring the uniform boundedness of the approximating sequence.

2. Preliminaries

In this section, we fix $1 < p_1 \leq p_2 \leq \dots \leq p_N < \infty$.

2.1. Some auxiliary functions

For every $i = 1, \dots, N$ and $\varepsilon > 0$, we define

$$(2.1) \quad g_{i,\varepsilon}(t) = \frac{1}{p_i} (\varepsilon + t^2)^{p_i/2}, \quad \text{for every } t \in \mathbb{R}.$$

Lemma 2.1 (Sub-quadratic case). *Let $1 < p_i \leq 2$. Then for every $t \in \mathbb{R}$, we have*

$$(2.2) \quad (p_i - 1) (\varepsilon + t^2)^{(p_i-2)/2} \leq g''_{i,\varepsilon}(t) \leq (\varepsilon + t^2)^{(p_i-2)/2},$$

$$(2.3) \quad g_{i,\varepsilon}(t) \leq \frac{1}{p_i} (\varepsilon^{p_i/2} + g'_{i,\varepsilon}(t) t),$$

$$(2.4) \quad |g'_{i,\varepsilon}(t)|^2 \leq \frac{p_i}{p_i - 1} g''_{i,\varepsilon}(t) g_{i,\varepsilon}(t).$$

Proof. The second derivative of $g_{i,\varepsilon}$ is given by

$$\begin{aligned} g''_{i,\varepsilon}(t) &= (\varepsilon + t^2)^{(p_i-2)/2} + (p_i - 2) (\varepsilon + t^2)^{(p_i-4)/2} t^2 \\ &= (\varepsilon + t^2)^{(p_i-2)/2} \left(1 + (p_i - 2) \frac{t^2}{\varepsilon + t^2} \right). \end{aligned}$$

In particular, by using that $p_i - 2 \leq 0$, we easily get (2.2).

We also observe that

$$\begin{aligned} g_{i,\varepsilon}(t) &= \frac{1}{p_i} (\varepsilon + t^2)^{p_i/2} = \frac{1}{p_i} (\varepsilon + t^2)^{(p_i-2)/2} \varepsilon + \frac{1}{p_i} (\varepsilon + t^2)^{(p_i-2)/2} t^2 \\ &\leq \frac{\varepsilon^{p_i/2}}{p_i} + \frac{g'_{i,\varepsilon}(t)}{p_i} t, \end{aligned}$$

which proves (2.3).

Finally, (2.4) follows by writing

$$|g'_{i,\varepsilon}(t)|^2 = (\varepsilon + t^2)^{(p_i-2)/2} (\varepsilon + t^2)^{(p_i-2)/2} t^2 \leq (\varepsilon + t^2)^{(p_i-2)/2} (\varepsilon + t^2)^{p_i/2},$$

and then using the definition of $g_{i,\varepsilon}$ and the lower bound in (2.2). ■

Lemma 2.2 (Super-quadratic case). *Let $p_i > 2$. Then for every $t \in \mathbb{R}$, we have*

$$(2.5) \quad (\varepsilon + t^2)^{(p_i-2)/2} \leq g''_{i,\varepsilon}(t) \leq (p_i - 1) (\varepsilon + t^2)^{(p_i-2)/2},$$

$$(2.6) \quad g_{i,\varepsilon}(t) \leq \frac{2^{(p_i-2)/2}}{p_i} (\varepsilon^{p_i/2} + g'_{i,\varepsilon}(t) t),$$

$$(2.7) \quad |g'_{i,\varepsilon}(t)|^2 \leq p_i g''_{i,\varepsilon}(t) g_{i,\varepsilon}(t).$$

Proof. The proofs of (2.5) and (2.7) are similar to those of (2.2) and (2.4) respectively, and we omit them. In order to prove (2.6), we use the convexity of the map $\tau \mapsto |\tau|^{p_i/2}$. This implies

$$(2.8) \quad g_{i,\varepsilon}(t) = \frac{1}{p_i} (\varepsilon + t^2)^{p_i/2} \leq \frac{2^{(p_i-2)/2}}{p_i} (\varepsilon^{p_i/2} + |t|^{p_i}).$$

We then observe that

$$|t|^{p_i} = |t|^{p_i-2} t^2 \leq (\varepsilon + t^2)^{(p_i-2)/2} t^2 = g'_{i,\varepsilon}(t) t.$$

By combining the last two inequalities, we get (2.6). ■

We also define the function

$$(2.9) \quad G_\varepsilon(z) := \sum_{i=1}^N g_{i,\varepsilon}(z_i) = \sum_{i=1}^N \frac{1}{p_i} (\varepsilon + z_i^2)^{p_i/2},$$

which will play a crucial role in our estimates. The next result holds without any restriction on p_i .

Lemma 2.3. *For every $z = (z_1, \dots, z_N) \in \mathbb{R}^N$ and every $i = 1, \dots, N$, we have*

$$(2.10) \quad |g'_{i,\varepsilon}(z_i)| \leq p_N^{(p_N-1)/p_N} G_\varepsilon(z)^{(p_i-1)/p_i}.$$

Proof. By recalling the definition of both $g_{i,\varepsilon}$ and G_ε , we have

$$\begin{aligned} |g'_{i,\varepsilon}(z_i)| &\leq (\varepsilon + |z_i|^2)^{(p_i-1)/2} = (p_i g_{i,\varepsilon}(z_i))^{(p_i-1)/p_i} \\ &\leq \left(\sum_{k=1}^N p_k g_{k,\varepsilon}(z_k) \right)^{(p_i-1)/p_i} \leq p_N^{(p_i-1)/p_i} G_\varepsilon(z)^{(p_i-1)/p_i}. \end{aligned}$$

By using that $p_N > 1$ and that $p_i \leq p_N$, we can estimate the last term from above as claimed. \blacksquare

2.2. Regularized problems

We will use an approximation scheme which is similar to that already used in our previous papers, starting from [11], Section 2. We want to consider local minimizers of the following convex integral functional:

$$\mathfrak{F}_p(u; \Omega') = \sum_{i=1}^N \frac{1}{p_i} \int_{\Omega'} |u_{x_i}|^{p_i} dx - \int_{\Omega'} f u dx, \quad u \in W_{\text{loc}}^{1,p}(\Omega) \cap L_{\text{loc}}^\infty(\Omega), \quad \Omega' \Subset \Omega.$$

The function f is assumed to belong to $L_{\text{loc}}^1(\Omega)$. In the rest of the paper, we fix $U \in W_{\text{loc}}^{1,p}(\Omega) \cap L_{\text{loc}}^\infty(\Omega)$ a bounded local minimizer of \mathfrak{F}_p . We also fix a ball

$$B \Subset \Omega \quad \text{such that} \quad 2B \Subset \Omega \text{ as well.}$$

Here by λB we denote the ball concentric with B , scaled by a factor $\lambda > 0$. We set

$$\varepsilon_0 = \min \left\{ 1, \frac{\text{radius of } B}{2} \right\} > 0.$$

For every $0 < \varepsilon \leq \varepsilon_0$, we denote by ϱ_ε the usual Friedrichs mollifier, supported in a ball of radius ε centered at the origin. For every $x \in \overline{B}$, we then define

$$U^\varepsilon(x) = U * \varrho_\varepsilon(x) \quad \text{and} \quad f^\varepsilon(x) = f * \varrho_\varepsilon(x).$$

Finally, we set

$$M := \|U\|_{L^\infty(2B)},$$

and take ζ^ε to be a C^∞ function of one variable, such that

$$0 \leq (\zeta^\varepsilon)' \leq 1 \quad \text{and} \quad \zeta^\varepsilon(t) = \begin{cases} M, & \text{if } t \geq M + \varepsilon, \\ -M, & \text{if } t \leq -M - \varepsilon, \end{cases}$$

and such that

$$\lim_{\varepsilon \rightarrow 0} \|\zeta^\varepsilon - \zeta\|_{L^\infty(\mathbb{R})} = 0, \quad \text{where } \zeta(t) = \max\{\min\{t, M\}, -M\}, \text{ for } t \in \mathbb{R}.$$

By recalling the definition (2.1) of $g_{i,\varepsilon}$, we then define the regularized functional

$$\mathfrak{F}_{\mathbf{p},\varepsilon}(v; B) = \sum_{i=1}^N \int_B g_{i,\varepsilon}(v_{x_i}) dx - \int_B f^\varepsilon \zeta^\varepsilon(v) dx.$$

Lemma 2.4 (Existence and regularity of a minimum for $\mathfrak{F}_{\mathbf{p},\varepsilon}$). *For every $0 < \varepsilon \leq \varepsilon_0$, the problem*

$$(2.11) \quad \min \{ \mathfrak{F}_{\mathbf{p},\varepsilon}(v; B) : v - U^\varepsilon \in W_0^{1,\mathbf{p}}(B) \},$$

admits a solution u^ε , which belongs to $C^\infty(\bar{B})$. Moreover, for every $0 < \varepsilon \leq \varepsilon_0$, we have

$$\|u^\varepsilon\|_{L^\infty(B)} \leq M + \varepsilon.$$

Proof. We first show that we can apply Theorem 9.2 in [56] and get existence of a solution to

$$(2.12) \quad \min \{ \mathfrak{F}_{\mathbf{p},\varepsilon}(v; B) : v - U^\varepsilon \in W_0^{1,\infty}(B) \}.$$

For this, we check the required assumptions. We first claim that, for every $z, \xi \in \mathbb{R}^N$,

$$(2.13) \quad \sum_{i=1}^N g''_{i,\varepsilon}(z_i) \xi_i^2 \geq \nu_\varepsilon (1 + |z|^2)^\tau |\xi|^2, \quad \text{with} \quad \begin{cases} \nu_\varepsilon := \min \{ p_1 - 1, \varepsilon^{(p_N-2)/2} \} > 0, \\ \tau := \min \{ 0, (p_1-2)/2 \} > -1/2. \end{cases}$$

From (2.2) and (2.5), we get

$$\sum_{i=1}^N g''_{i,\varepsilon}(z_i) \xi_i^2 \geq \sum_{i=1}^N \min \{ p_i - 1, 1 \} (\varepsilon + z_i^2)^{(p_i-2)/2} \xi_i^2.$$

Then we observe that if $p_i \leq 2$, then $p_1 \leq 2$, and we can write

$$\min \{ p_i - 1, 1 \} (\varepsilon + z_i^2)^{(p_i-2)/2} \geq (p_1 - 1) (1 + |z|^2)^{(p_1-2)/2} = (p_1 - 1) (1 + |z|^2)^\tau.$$

If instead $p_i > 2$, then

$$\min \{ p_i - 1, 1 \} (\varepsilon + z_i^2)^{(p_i-2)/2} \geq \varepsilon^{(p_N-2)/2} \geq \varepsilon^{(p_N-2)/2} (1 + |z|^2)^\tau.$$

Thus, in both cases, (2.13) holds.

As for the lower order term, observe that the smooth function $h^\varepsilon(x, u) := f^\varepsilon(x) \zeta^\varepsilon(u)$ satisfies

$$\frac{\partial h^\varepsilon}{\partial u}(x, u) = 0, \quad \text{for every } (x, u) \in B \times (\mathbb{R} \setminus (-M - \varepsilon, M + \varepsilon)).$$

Finally, the uniform convexity of B and the smoothness of U^ε entail that the latter satisfies the bounded slope condition.

Then Theorem 9.2 in [56] yields the existence of a solution to (2.12). Since all the data are smooth, Theorem 9.3 in [56] implies that $u^\varepsilon \in C^\infty(\bar{B})$. We claim that u^ε is a solution

of (2.11), as well. Indeed, by using Theorem 1.1 in [13], for every $v \in U^\varepsilon + W_0^{1,\mathbf{p}}(B)$, we can infer the existence of $\{v^k\}_{k \in \mathbb{N}} \subset U^\varepsilon + W_0^{1,\infty}(B)$ converging to v in $W^{1,p_1}(B)$ and such that

$$\lim_{k \rightarrow \infty} \sum_{i=1}^N \int_B g_{i,\varepsilon}(v_{x_i}^k) dx = \sum_{i=1}^N \int_B g_{i,\varepsilon}(v_{x_i}) dx.$$

By the dominated convergence theorem and the uniform boundedness of ζ^ε , we also have

$$\lim_{k \rightarrow \infty} \int_B f^\varepsilon \zeta^\varepsilon(v^k) dx = \int_B f^\varepsilon \zeta^\varepsilon(v) dx.$$

This proves that there is no Lavrentiev phenomenon for $\mathfrak{F}_{\mathbf{p},\varepsilon}$, that is,

$$\mathfrak{F}_{\mathbf{p},\varepsilon}(u^\varepsilon; B) = \min_{U^\varepsilon + W_0^{1,\infty}(B)} \mathfrak{F}_{\mathbf{p},\varepsilon} = \inf_{U^\varepsilon + W_0^{1,\mathbf{p}}(B)} \mathfrak{F}_{\mathbf{p},\varepsilon}.$$

Thus, we get that u^ε solves (2.11), as well. Finally, the claimed L^∞ estimate readily follows from Lemma A.1 in the Appendix. \blacksquare

The smooth minimizer u^ε satisfies the Euler–Lagrange equation

$$(2.14) \quad \sum_{i=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) \varphi_{x_i} dx = \int f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) \varphi dx, \quad \text{for every } \varphi \in W_0^{1,\mathbf{p}}(B).$$

For every $k = 1, \dots, N$, one can insert test functions of the form φ_{x_k} , with $\varphi \in C^2$ compactly supported in B . By integrating by parts, we then get the equation for the partial derivatives of u^ε : for every $k = 1, \dots, N$,

$$(2.15) \quad \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) u_{x_k x_i}^\varepsilon \varphi_{x_i} dx = \int (f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon))_{x_k} \varphi dx.$$

As usual, by a density argument, the equation can be tested by any $\varphi \in W_0^{1,\mathbf{p}}(B)$.

The first ingredient of our recipe is a simple a priori estimate, which is essentially the same as in Lemma 2.1 of [10]: the only difference is the presence of the non-autonomous and nonlinear term $f^\varepsilon \zeta^\varepsilon(v)$, together with a slight modification of the function $g_{i,\varepsilon}$.

Lemma 2.5 (Basic energy estimate). *For every $0 < \varepsilon \leq \varepsilon_0$, the following uniform estimate holds:*

$$\sum_{i=1}^N \frac{1}{p_i} \int_B |u_{x_i}^\varepsilon|^{p_i} dx \leq C \left(\sum_{i=1}^N \frac{1}{p_i} \int_{2B} |U_{x_i}|^{p_i} dx + \varepsilon^{p_1/2} + \|f\|_{L^1(2B)} \right),$$

for some $C = C(N, B, \mathbf{p}, M) > 0$.

Proof. By testing the minimality of u^ε against U^ε , we obtain

$$\sum_{i=1}^N \int_B g_{i,\varepsilon}(u_{x_i}^\varepsilon) dx \leq \sum_{i=1}^N \int_B g_{i,\varepsilon}(U_{x_i}^\varepsilon) dx - \int_B f^\varepsilon(\zeta^\varepsilon(U^\varepsilon) - \zeta^\varepsilon(u^\varepsilon)) dx.$$

The convexity of the function $g_{i,\varepsilon}$ allows to apply Jensen's inequality in connection with the fact that U^ε is defined by a convolution. This gives

$$\int_B g_{i,\varepsilon}(U_{x_i}^\varepsilon) dx \leq \int_B g_{i,\varepsilon}(U_{x_i}) * \varrho_\varepsilon dx \leq \int_{2B} g_{i,\varepsilon}(U_{x_i}) dx.$$

By using also that $g_{i,\varepsilon}(t) \geq |t|^{p_i}/p_i$ and the 1-Lipschitz character of ζ^ε , we get

$$\begin{aligned} \sum_{i=1}^N \frac{1}{p_i} \int_B |u_{x_i}^\varepsilon|^{p_i} dx &\leq \sum_{i=1}^N \int_{2B} g_{i,\varepsilon}(U_{x_i}) dx + \int_B |f^\varepsilon| |U^\varepsilon - u^\varepsilon| dx \\ &\leq \sum_{i=1}^N \int_{2B} g_{i,\varepsilon}(U_{x_i}) dx + C_M \|f\|_{L^1(2B)}, \end{aligned}$$

where C_M is a positive constant which only depends on M . We finally rely on (2.8) when $p_i > 2$, or the subadditivity of $t \mapsto |t|^{p_i/2}$ when $p_i \leq 2$, to obtain

$$(2.16) \quad g_{i,\varepsilon}(t) \leq \frac{\max\{1, 2^{(p_i-2)/2}\}}{p_i} (\varepsilon^{p_i/2} + |t|^{p_i}).$$

This concludes the proof. ■

In view of our goal, we need a convergence result for the minimizers $\{u^\varepsilon\}_{0 < \varepsilon < \varepsilon_0}$. This is the content of the next lemma, which is an extension of Lemma 2.2 in [10].

Lemma 2.6 (Convergence to a minimizer). *With the same notation as above, we have*

$$(2.17) \quad \lim_{\varepsilon \rightarrow 0} \left[\|u^\varepsilon - U\|_{L^q(B)} + \sum_{i=1}^N \|(u^\varepsilon - U)_{x_i}\|_{L^{p_i}(B)} \right] = 0,$$

for every $1 \leq q < \infty$.

Proof. The proof goes as in Lemma 2.2 of [10]. We repeat the argument, since this gives us the occasion to fix some missing details in [10]. By using the uniform estimate of Lemma 2.5 and the definition of U^ε , we get that $\{u^\varepsilon - U^\varepsilon\}_{0 < \varepsilon \leq \varepsilon_0}$ is a bounded family in $W_0^{1,p}(B)$. Thanks to Lemma 2.4, we also have that $\{u^\varepsilon - U^\varepsilon\}_{0 < \varepsilon \leq \varepsilon_0}$ is a bounded family in $L^\infty(B)$. From these two facts, we can infer the existence of an infinitesimal sequence $\{\varepsilon_k\}_{k \in \mathbb{N}} \subset (0, \varepsilon_0]$ such that $\{u^{\varepsilon_k} - U^{\varepsilon_k}\}_{k \in \mathbb{N}}$ converges weakly in $W_0^{1,p}(B)$ and almost everywhere to a function $\phi \in W_0^{1,p}(B) \cap L^\infty(B)$, in the sense that

$$\lim_{k \rightarrow \infty} \int_B (u^{\varepsilon_k} - U^{\varepsilon_k}) \varphi dx = \int_B \phi \varphi dx, \quad \text{for every } \varphi \in L^1(B),$$

and

$$\lim_{k \rightarrow \infty} \int_B (u^{\varepsilon_k} - U^{\varepsilon_k})_{x_i} \varphi dx = \int_B \phi_{x_i} \varphi dx, \quad \text{for every } \varphi \in L^{p_i'}(B), i = 1, \dots, N.$$

By recalling that U^{ε_k} has been constructed by convolution, we also have that it converges strongly in $W^{1,p}(B)$ and almost everywhere to U . This permits to conclude that $\{u^{\varepsilon_k}\}_{k \in \mathbb{N}}$

converges weakly and almost everywhere to $u := \phi + U$. We need to prove that actually $u = U$. With this aim, we test the minimality of each u^{ε_k} against the function U^{ε_k} . Thus, by lower semicontinuity of the L^{p_i} norms, we can infer

$$\begin{aligned}
 \sum_{i=1}^N \frac{1}{p_i} \int_B |u_{x_i}|^{p_i} dx &\leq \liminf_{k \rightarrow \infty} \sum_{i=1}^N \frac{1}{p_i} \int_B |u_{x_i}^{\varepsilon_k}|^{p_i} dx \leq \liminf_{k \rightarrow \infty} \sum_{i=1}^N \frac{1}{p_i} \int_B g_{i,\varepsilon_k}(u_{x_i}^{\varepsilon_k}) dx \\
 &\leq \lim_{k \rightarrow \infty} \sum_{i=1}^N \frac{1}{p_i} \int_B g_{i,\varepsilon_k}(U_{x_i}^{\varepsilon_k}) dx - \int_B f^{\varepsilon_k} (\zeta^{\varepsilon_k}(U^{\varepsilon_k}) - \zeta^{\varepsilon_k}(u^{\varepsilon_k})) dx \\
 (2.18) \quad &= \sum_{i=1}^N \frac{1}{p_i} \int_B |U_{x_i}|^{p_i} dx - \int_B f(\zeta(U) - \zeta(u)) dx.
 \end{aligned}$$

Observe that for the convergence of the lower order term, we used that f^{ε_k} converges strongly in $L^1(B)$, that U^{ε_k} and u^{ε_k} are equibounded in $L^\infty(B)$ and converge almost everywhere to U and u respectively, and that ζ^{ε_k} converges uniformly to the Lipschitz function ζ . This shows that

$$\sum_{i=1}^N \frac{1}{p_i} \int_B |u_{x_i}|^{p_i} dx - \int_{\Omega'} f \zeta(u) dx \leq \sum_{i=1}^N \frac{1}{p_i} \int_B |U_{x_i}|^{p_i} dx - \int_{\Omega'} f \zeta(U) dx.$$

The L^∞ -boundedness of u^ε proved in Lemma 2.4, gives that $\|u\|_{L^\infty(B)} \leq M$. A similar estimate holds for U by assumption. Since $\zeta(t) = t$ for every $t \in [-M, M]$, one gets

$$\mathfrak{F}_p(u; B) \leq \mathfrak{F}_p(U; B).$$

By the strict convexity of the functional \mathfrak{F}_p , the minimizer must be unique and thus we get $u = U$, as desired.

In order to prove (2.17), we can adapt the argument of Lemma 2.2 in [10]. By (2.18), we get

$$(2.19) \quad \lim_{k \rightarrow \infty} \sum_{i=1}^N \frac{1}{p_i} \int_B |u_{x_i}^{\varepsilon_k}|^{p_i} dx = \sum_{i=1}^N \frac{1}{p_i} \int_B |U_{x_i}|^{p_i} dx.$$

For every $i = 1, \dots, N$, we rely on the lower semicontinuity of the L^{p_i} norm to get

$$\liminf_{k \rightarrow \infty} \int_B |u_{x_i}^{\varepsilon_k}|^{p_i} dx \geq \int_B |U_{x_i}|^{p_i} dx,$$

In connection with (2.19), this implies that

$$\lim_{k \rightarrow \infty} \int_B |u_{x_i}^{\varepsilon_k}|^{p_i} dx = \int_B |U_{x_i}|^{p_i} dx.$$

The convergence of the norms, in conjunction with the weak convergence, permits to infer that $(u_{x_i}^{\varepsilon_k})_{k \in \mathbb{N}}$ converges to U_{x_i} in $L^{p_i}(B)$ for every $i = 1, \dots, N$ (see for example Theorem 2.11 in [45]).

Moreover, since $\{u^{\varepsilon_k}\}_{k \in \mathbb{N}}$ is bounded by M and converges almost everywhere in B to U , the dominated convergence theorem implies that $(u^{\varepsilon_k})_{k \in \mathbb{N}}$ converges to U in $L^q(B)$ for every $1 \leq q < \infty$.

Finally, we observe that we can repeat this argument with any subsequence of the original family $\{u^\varepsilon\}_{\varepsilon > 0}$. Thus the above limit holds true for the whole family $\{u^\varepsilon\}_{0 < \varepsilon \leq \varepsilon_0}$ instead of $\{u^{\varepsilon_k}\}_{k \in \mathbb{N}}$ and (2.17) follows. ■

The following technical result is classical in the regularity theory. This is taken from Lemma 6.1 in [39], and we state it here for the reader's convenience.

Lemma 2.7. *Let $0 < r < R$ and let $Z: [r, R] \rightarrow [0, \infty)$ be a bounded function. Assume that for $r \leq s < t \leq R$, we have*

$$Z(s) \leq \frac{\mathcal{A}}{(t-s)^{\alpha_0}} + \frac{\mathcal{B}}{(t-s)^{\beta_0}} + \mathcal{C} + \vartheta Z(t),$$

with $\mathcal{A}, \mathcal{B}, \mathcal{C} \geq 0$, $\alpha_0 \geq \beta_0 > 0$ and $0 \leq \vartheta < 1$. Then we have

$$Z(r) \leq \left(\frac{1}{(1-\lambda)^{\alpha_0}} \frac{\lambda^{\alpha_0}}{\lambda^{\alpha_0} - \vartheta} \right) \left[\frac{\mathcal{A}}{(R-r)^{\alpha_0}} + \frac{\mathcal{B}}{(R-r)^{\beta_0}} + \mathcal{C} \right],$$

where λ is any number such that $\vartheta^{1/\alpha_0} < \lambda < 1$.

3. Caccioppoli-type inequalities for the gradient

Throughout this section, we assume that $1 < p_1 \leq \dots \leq p_N$, without any further restriction. In what follows, we will use the function

$$(3.1) \quad \mathcal{G}_\varepsilon(z) = ((G_\varepsilon(z) - 1)_+ + 1), \quad \text{for every } z \in \mathbb{R}^N,$$

where G_ε is the same function as in (2.9).

Proposition 3.1 (Caccioppoli inequality for power functions of the gradient). *Let $1 < p_1 \leq p_2 \leq \dots \leq p_N < \infty$. For every $\alpha \geq 0$ and every non-negative $\eta \in C_0^2(B)$, we have*

$$(3.2) \quad \begin{aligned} & \sum_{i=1}^N \int g''_{i,\varepsilon}(u^\varepsilon_{x_i}) |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+1)/2})_{x_i}|^2 \eta^2 dx \\ & \leq C(\alpha+1)^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha+2-2/p_N} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ & \quad + C(\alpha+1)^2 \int |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\alpha \eta^2 dx, \end{aligned}$$

for some $C = C(N, p_1, p_N) > 0$.

Proof. We are going to use a trick based on integration by parts, taken from Theorem 1 in [52] (see also [36]). This permits to circumvent the use of the upper bound on the

Hessian of the function G_ε . We start by fixing $k \in \{1, \dots, N\}$ and inserting in (2.15) the test function

$$\varphi = F(G_\varepsilon(\nabla u^\varepsilon)) g'_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta^2,$$

where F is a non-negative Lipschitz continuous monotone non-decreasing function, that will be specified later on. This is a feasible test function, thanks to the regularity of u^ε . Thus we get

$$\begin{aligned} & \sum_{i=1}^N \int g''_{i,\varepsilon}(u^\varepsilon_{x_i}) u^\varepsilon_{x_k x_i} (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_i} g'_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta^2 dx \\ & + \sum_{i=1}^N \int g''_{i,\varepsilon}(u^\varepsilon_{x_i}) u^\varepsilon_{x_k x_i} F(G_\varepsilon(\nabla u^\varepsilon)) (g'_{k,\varepsilon}(u^\varepsilon_{x_k}))_{x_i} \eta^2 dx \\ (3.3) \quad & = -2 \sum_{i=1}^N \int g''_{i,\varepsilon}(u^\varepsilon_{x_i}) u^\varepsilon_{x_k x_i} F(G_\varepsilon(\nabla u^\varepsilon)) g'_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta \eta_{x_i} dx \\ & + \int (f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon))_{x_k} F(G_\varepsilon(\nabla u^\varepsilon)) g'_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta^2 dx. \end{aligned}$$

We observe that

$$g''_{i,\varepsilon}(u^\varepsilon_{x_i}) u^\varepsilon_{x_k x_i} = (g'_{i,\varepsilon}(u^\varepsilon_{x_i}))_{x_k}.$$

Then by integrating by parts on the right-hand side² of (3.3), we obtain

$$\begin{aligned} & \sum_{i=1}^N \int g''_{i,\varepsilon}(u^\varepsilon_{x_i}) u^\varepsilon_{x_k x_i} (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_i} g'_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta^2 dx \\ & + \sum_{i=1}^N \int g''_{i,\varepsilon}(u^\varepsilon_{x_i}) u^\varepsilon_{x_k x_i} F(G_\varepsilon(\nabla u^\varepsilon)) (g'_{k,\varepsilon}(u^\varepsilon_{x_k}))_{x_i} \eta^2 dx \\ (3.4) \quad & = 2 \sum_{i=1}^N \int g'_{i,\varepsilon}(u^\varepsilon_{x_i}) (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_k} g'_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta \eta_{x_i} dx \\ & + 2 \sum_{i=1}^N \int g'_{i,\varepsilon}(u^\varepsilon_{x_i}) F(G_\varepsilon(\nabla u^\varepsilon)) (g'_{k,\varepsilon}(u^\varepsilon_{x_k}))_{x_k} \eta \eta_{x_i} dx \\ & + 2 \sum_{i=1}^N \int g'_{i,\varepsilon}(u^\varepsilon_{x_i}) F(G_\varepsilon(\nabla u^\varepsilon)) g'_{k,\varepsilon}(u^\varepsilon_{x_k}) (\eta_{x_k} \eta_{x_i} + \eta \eta_{x_i x_k}) dx \\ & + \int (f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon))_{x_k} F(G_\varepsilon(\nabla u^\varepsilon)) g'_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta^2 dx. \end{aligned}$$

This is valid for every $k = 1, \dots, N$, we then take the sum over k .

²This is the trick in [52] (and [36]) mentioned above.

On the left-hand side, the first term then becomes

$$\begin{aligned}
& \sum_{i,k=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) u_{x_k x_i}^\varepsilon (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_i} g'_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx \\
&= \sum_{i,k=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_i} (g_{k,\varepsilon}(u_{x_k}^\varepsilon))_{x_i} \eta^2 dx \\
&= \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_i} \left(\sum_{k=1}^N g_{k,\varepsilon}(u_{x_k}^\varepsilon) \right)_{x_i} \eta^2 dx \\
&= \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_i} (G_\varepsilon(\nabla u^\varepsilon))_{x_i} \eta^2 dx \\
&= \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) F'(G_\varepsilon(\nabla u^\varepsilon)) |(G_\varepsilon(\nabla u^\varepsilon))_{x_i}|^2 \eta^2 dx.
\end{aligned}$$

For the second term of the left-hand side in (3.4), we observe that

$$\begin{aligned}
& \sum_{i,k=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) u_{x_k x_i}^\varepsilon F(G_\varepsilon(\nabla u^\varepsilon)) (g'_{k,\varepsilon}(u_{x_k}^\varepsilon))_{x_i} \eta^2 dx \\
&= \sum_{i,k=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2 F(G_\varepsilon(\nabla u^\varepsilon)) g''_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx,
\end{aligned}$$

and this is non-negative, since each $g_{k,\varepsilon}$ is convex and $F \geq 0$. We thus obtain

$$\begin{aligned}
& \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) F'(G_\varepsilon(\nabla u^\varepsilon)) |(G_\varepsilon(\nabla u^\varepsilon))_{x_i}|^2 \eta^2 dx \\
&+ \sum_{i,k=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2 F(G_\varepsilon(\nabla u^\varepsilon)) g''_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx \\
(3.5) \quad &= 2 \sum_{i,k=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_k} g'_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta \eta_{x_i} dx \\
&+ 2 \sum_{i,k=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) F(G_\varepsilon(\nabla u^\varepsilon)) (g'_{k,\varepsilon}(u_{x_k}^\varepsilon))_{x_k} \eta \eta_{x_i} dx \\
&+ 2 \sum_{i,k=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) F(G_\varepsilon(\nabla u^\varepsilon)) g'_{k,\varepsilon}(u_{x_k}^\varepsilon) (\eta_{x_k} \eta_{x_i} + \eta \eta_{x_i x_k}) dx \\
&+ \sum_{k=1}^N \int (f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon))_{x_k} F(G_\varepsilon(\nabla u^\varepsilon)) g'_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx.
\end{aligned}$$

By (2.14), the second term of the right-hand side can be written as

$$(3.6) \quad -2 \sum_{i=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) F(G_\varepsilon(\nabla u^\varepsilon)) f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) \eta \eta_{x_i} dx.$$

By an integration by parts and (2.14) again, the last term on the right-hand side of (3.5) is equal to

$$\begin{aligned} & - \sum_{k=1}^N \int f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_k} g'_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx \\ & \quad + \int |f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon)|^2 F(G_\varepsilon(\nabla u^\varepsilon)) \eta^2 dx \\ & \quad - 2 \sum_{k=1}^N \int f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) F(G_\varepsilon(\nabla u^\varepsilon)) g'_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta \eta_{x_k} dx. \end{aligned}$$

We observe that the third term in the above sum is equal to the quantity in (3.6). So (3.5) is equivalent to

$$\begin{aligned} \mathcal{J}_1 + \mathcal{J}_2 & := \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) F'(G_\varepsilon(\nabla u^\varepsilon)) |(G_\varepsilon(\nabla u^\varepsilon))_{x_i}|^2 \eta^2 dx \\ & \quad + \sum_{i,k=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2 F(G_\varepsilon(\nabla u^\varepsilon)) g''_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx \\ & = 2 \sum_{i,k=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_k} g'_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta \eta_{x_i} dx \\ & \quad - 4 \sum_{i=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) F(G_\varepsilon(\nabla u^\varepsilon)) f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) \eta \eta_{x_i} dx \\ & \quad + 2 \sum_{i,k=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) F(G_\varepsilon(\nabla u^\varepsilon)) g'_{k,\varepsilon}(u_{x_k}^\varepsilon) (\eta_{x_k} \eta_{x_i} + \eta \eta_{x_i x_k}) dx \\ & \quad - \sum_{k=1}^N \int f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_k} g'_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx \\ & \quad + \int |f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon)|^2 F(G_\varepsilon(\nabla u^\varepsilon)) \eta^2 dx \\ (3.7) \quad & =: 2 \mathcal{J}_3 + 4 \mathcal{F}_1 + 2 \mathcal{J}_4 + \mathcal{F}_2 + \mathcal{F}_3. \end{aligned}$$

We first estimate \mathcal{J}_3 : by Young's inequality, we have for every $\tau > 0$,

$$\begin{aligned}
|\mathcal{J}_3| &\leq \int \sum_{k=1}^N \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right) (F'(G_\varepsilon(\nabla u^\varepsilon)) |(G_\varepsilon(\nabla u^\varepsilon))_{x_k} g'_{k,\varepsilon}(u_{x_k}^\varepsilon)|) \eta |\nabla \eta| dx \\
&\leq \frac{1}{2\tau} \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 F'(G_\varepsilon(\nabla u^\varepsilon)) \left(\sum_{k=1}^N \frac{(g'_{k,\varepsilon}(u_{x_k}^\varepsilon))^2}{g''_{k,\varepsilon}(u_{x_k}^\varepsilon)} \right) |\nabla \eta|^2 \\
&\quad + \frac{\tau}{2} \int \sum_{k=1}^N F'(G_\varepsilon(\nabla u^\varepsilon)) |(G_\varepsilon(\nabla u^\varepsilon))_{x_k}|^2 g''_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx \\
&\leq \frac{1}{2\tau} \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 F'(G_\varepsilon(\nabla u^\varepsilon)) \left(\sum_{k=1}^N \frac{(g'_{k,\varepsilon}(u_{x_k}^\varepsilon))^2}{g''_{k,\varepsilon}(u_{x_k}^\varepsilon)} \right) |\nabla \eta|^2 dx + \frac{\tau}{2} \mathcal{J}_1.
\end{aligned}$$

By taking $\tau = 1/2$, we can absorb the term \mathcal{J}_1 on the right-hand side and obtain from (3.7) the following:

$$\begin{aligned}
(3.8) \quad \frac{1}{2} \mathcal{J}_1 + \mathcal{J}_2 &\leq 2 \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 F'(G_\varepsilon(\nabla u^\varepsilon)) \left(\sum_{k=1}^N \frac{(g'_{k,\varepsilon}(u_{x_k}^\varepsilon))^2}{g''_{k,\varepsilon}(u_{x_k}^\varepsilon)} \right) |\nabla \eta|^2 dx \\
&\quad + 4 \mathcal{F}_1 + 2 \mathcal{J}_4 + \mathcal{F}_2 + \mathcal{F}_3.
\end{aligned}$$

The term

$$\mathcal{J}_4 = \sum_{i,k=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) F(G_\varepsilon(\nabla u^\varepsilon)) g'_{k,\varepsilon}(u_{x_k}^\varepsilon) (\eta_{x_k} \eta_{x_i} + \eta \eta_{x_i x_k}) dx$$

is easier to handle: we simply have

$$|\mathcal{J}_4| \leq \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 F(G_\varepsilon(\nabla u^\varepsilon)) (|\nabla \eta|^2 + \eta |D^2 \eta|) dx.$$

In conclusion, from (3.8) we get

$$\begin{aligned}
(3.9) \quad \mathcal{J}_1 + 2 \mathcal{J}_2 &\leq 4 \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 F'(G_\varepsilon(\nabla u^\varepsilon)) \left(\sum_{k=1}^N \frac{(g'_{k,\varepsilon}(u_{x_k}^\varepsilon))^2}{g''_{k,\varepsilon}(u_{x_k}^\varepsilon)} \right) |\nabla \eta|^2 dx \\
&\quad + 8 \mathcal{F}_1 + 4 \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 F(G_\varepsilon(\nabla u^\varepsilon)) (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\
&\quad + 2 \mathcal{F}_2 + 2 \mathcal{F}_3.
\end{aligned}$$

We now treat the three terms containing f^ε : we start from

$$\begin{aligned}
\mathcal{F}_1 &= - \sum_{i=1}^N \int g'_{i,\varepsilon}(u^\varepsilon_{x_i}) F(G_\varepsilon(\nabla u^\varepsilon)) f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) \eta \eta_{x_i} dx \\
&\leq \int \sum_{i=1}^N |g'_{i,\varepsilon}(u^\varepsilon_{x_i})| F(G_\varepsilon(\nabla u^\varepsilon)) |f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon)| \eta |\nabla \eta| dx \\
&\leq \frac{1}{2} \int \left(\sum_{i=1}^N g'_{i,\varepsilon}(u^\varepsilon_{x_i}) \right)^2 F(G_\varepsilon(\nabla u^\varepsilon)) |\nabla \eta|^2 dx \\
&\quad + \frac{1}{2} \int |f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon)|^2 F(G_\varepsilon(\nabla u^\varepsilon)) \eta^2 dx.
\end{aligned}$$

The last term coincides with $\frac{1}{2} \mathcal{F}_3$ while the first term is bounded from above (up to a multiplicative constant) by the third term on the right-hand side of (3.9). Using also that $\|(\zeta^\varepsilon)'\|_{L^\infty(\mathbb{R})} \leq 1$, we thus get, from (3.9),

$$\begin{aligned}
\mathcal{J}_1 + 2 \mathcal{J}_2 &\leq 4 \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u^\varepsilon_{x_i})| \right)^2 F'(G_\varepsilon(\nabla u^\varepsilon)) \left(\sum_{k=1}^N \frac{(g'_{k,\varepsilon}(u^\varepsilon_{x_k}))^2}{g''_{k,\varepsilon}(u^\varepsilon_{x_k})} \right) |\nabla \eta|^2 dx \\
(3.10) \quad &+ 8 \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u^\varepsilon_{x_i})| \right)^2 F(G_\varepsilon(\nabla u^\varepsilon)) (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\
&+ 6 \int |f^\varepsilon|^2 F(G_\varepsilon(\nabla u^\varepsilon)) \eta^2 dx + 2 \mathcal{F}_2.
\end{aligned}$$

The last term \mathcal{F}_2 contains second order derivatives of u^ε that should be absorbed on the left-hand side. We proceed similarly as for \mathcal{J}_3 and estimate it as follows:

$$\begin{aligned}
\mathcal{F}_2 &= - \sum_{k=1}^N \int f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) (F(G_\varepsilon(\nabla u^\varepsilon)))_{x_k} g'_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta^2 dx \\
&\leq \sum_{k=1}^N \int |f^\varepsilon| (F'(G_\varepsilon(\nabla u^\varepsilon)) |(G_\varepsilon(\nabla u^\varepsilon))_{x_k} g'_{k,\varepsilon}(u^\varepsilon_{x_k})|) \eta^2 dx \\
&\leq \frac{1}{2\tau} \sum_{k=1}^N \int |f^\varepsilon|^2 F'(G_\varepsilon(\nabla u^\varepsilon)) \frac{(g'_{k,\varepsilon}(u^\varepsilon_{x_k}))^2}{g''_{k,\varepsilon}(u^\varepsilon_{x_k})} \eta^2 dx \\
&\quad + \frac{\tau}{2} \sum_{k=1}^N \int F'(G_\varepsilon(\nabla u^\varepsilon)) |(G_\varepsilon(\nabla u^\varepsilon))_{x_k}|^2 g''_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta^2 dx \\
&= \frac{1}{2\tau} \sum_{k=1}^N \int |f^\varepsilon|^2 F'(G_\varepsilon(\nabla u^\varepsilon)) \frac{(g'_{k,\varepsilon}(u^\varepsilon_{x_k}))^2}{g''_{k,\varepsilon}(u^\varepsilon_{x_k})} \eta^2 dx + \frac{\tau}{2} \mathcal{J}_1.
\end{aligned}$$

thanks to Young's inequality. Here, as always, $\tau > 0$ is arbitrary.

By inserting this estimate in (3.10) and choosing $\tau = 1/2$, we obtain

$$\begin{aligned} \mathcal{J}_1 + 4\mathcal{J}_2 &\leq 8 \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 F'(G_\varepsilon(\nabla u^\varepsilon)) \left(\sum_{k=1}^N \frac{(g'_{k,\varepsilon}(u_{x_k}^\varepsilon))^2}{g''_{k,\varepsilon}(u_{x_k}^\varepsilon)} \right) |\nabla \eta|^2 dx \\ &\quad + 16 \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 F(G_\varepsilon(\nabla u^\varepsilon)) (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ &\quad + 12 \int |f^\varepsilon|^2 F(G_\varepsilon(\nabla u^\varepsilon)) \eta^2 dx + 4 \sum_{k=1}^N \int |f^\varepsilon|^2 F'(G_\varepsilon(\nabla u^\varepsilon)) \frac{(g'_{k,\varepsilon}(u_{x_k}^\varepsilon))^2}{g''_{k,\varepsilon}(u_{x_k}^\varepsilon)} \eta^2 dx. \end{aligned}$$

We observe that if we set

$$(3.11) \quad \delta_k = \max \left\{ \frac{p_k}{p_k - 1}, p_k \right\} \quad \text{and} \quad \bar{\delta} = \max_{k=1, \dots, N} \delta_k = \max \left\{ \frac{p_1}{p_1 - 1}, p_N \right\},$$

by (2.4) or (2.7), we have

$$(3.12) \quad \sum_{k=1}^N \frac{(g'_{k,\varepsilon}(u_{x_k}^\varepsilon))^2}{g''_{k,\varepsilon}(u_{x_k}^\varepsilon)} \leq \sum_{k=1}^N \delta_k g_{k,\varepsilon}(u_{x_k}^\varepsilon) \leq \bar{\delta} G_\varepsilon(\nabla u^\varepsilon).$$

Thus we have obtained

$$\begin{aligned} \mathcal{J}_1 + 4\mathcal{J}_2 &\leq 8\bar{\delta} \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 F'(G_\varepsilon(\nabla u^\varepsilon)) G_\varepsilon(\nabla u^\varepsilon) |\nabla \eta|^2 dx \\ &\quad + 16 \int \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 F(G_\varepsilon(\nabla u^\varepsilon)) (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ (3.13) \quad &\quad + 12 \int |f^\varepsilon|^2 F(G_\varepsilon(\nabla u^\varepsilon)) \eta^2 dx + 4\bar{\delta} \int |f^\varepsilon|^2 F'(G_\varepsilon(\nabla u^\varepsilon)) G_\varepsilon(\nabla u^\varepsilon) \eta^2 dx. \end{aligned}$$

We now use Lemma 2.3 to estimate from above the right-hand side. Thus, from (3.13), we get

$$\begin{aligned} (3.14) \quad &\mathcal{J}_1 + 4\mathcal{J}_2 \\ &\leq 8\bar{\delta} (p_N^{(p_N-1)/p_N})^2 \int \left(\sum_{i=1}^N G_\varepsilon(\nabla u^\varepsilon)^{(p_i-1)/p_i} \right)^2 F'(G_\varepsilon(\nabla u^\varepsilon)) G_\varepsilon(\nabla u^\varepsilon) |\nabla \eta|^2 dx \\ &\quad + 16 (p_N^{(p_N-1)/p_N})^2 \int \left(\sum_{i=1}^N G_\varepsilon(\nabla u^\varepsilon)^{(p_i-1)/p_i} \right)^2 F(G_\varepsilon(\nabla u^\varepsilon)) (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ &\quad + 12\bar{\delta} \int |f^\varepsilon|^2 [F(G_\varepsilon(\nabla u^\varepsilon)) + F'(G_\varepsilon(\nabla u^\varepsilon)) G_\varepsilon(\nabla u^\varepsilon)] \eta^2 dx. \end{aligned}$$

By recalling the definition (3.1) of \mathcal{E}_ε , we observe that

$$\mathcal{E}_\varepsilon(\nabla u^\varepsilon) \geq G_\varepsilon(\nabla u^\varepsilon) \quad \text{and} \quad \mathcal{E}_\varepsilon(\nabla u^\varepsilon) \geq 1.$$

Thus, in particular, we get

$$\sum_{i=1}^N G_\varepsilon(\nabla u^\varepsilon)^{(p_i-1)/p_i} \leq N \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(p_N-1)/p_N},$$

and from (3.14), we obtain

$$\begin{aligned} \mathcal{J}_1 + 4\mathcal{J}_2 &\leq 8\bar{\delta} (Np_N^{(p_N-1)/p_N})^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2(p_N-1)/p_N+1} F'(G_\varepsilon(\nabla u^\varepsilon)) |\nabla \eta|^2 dx \\ &\quad + 16(Np_N^{(p_N-1)/p_N})^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2(p_N-1)/p_N} F(G_\varepsilon(\nabla u^\varepsilon)) (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ (3.15) \quad &+ 12\bar{\delta} \int |f^\varepsilon|^2 [F(G_\varepsilon(\nabla u^\varepsilon)) + F'(G_\varepsilon(\nabla u^\varepsilon)) G_\varepsilon(\nabla u^\varepsilon)] \eta^2 dx. \end{aligned}$$

In order to conclude, we now make the choice

$$F(t) = ((t-1)_+ + 1)^\alpha, \quad \text{with } \alpha > 0.$$

We observe that

$$(3.16) \quad |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon))_{x_i}|^2 \leq |(G_\varepsilon(\nabla u^\varepsilon))_{x_i}|^2,$$

thanks to the definition of \mathcal{G}_ε .

It follows that

$$\begin{aligned} F'(G_\varepsilon(\nabla u^\varepsilon)) |(G_\varepsilon(\nabla u^\varepsilon))_{x_i}|^2 &\geq \alpha \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha-1} |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon))_{x_i}|^2 \\ &= \frac{4\alpha}{(\alpha+1)^2} |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+1)/2})_{x_i}|^2, \end{aligned}$$

and thus

$$\begin{aligned} \mathcal{J}_1 &= \sum_{i=1}^N \int g''_{i,\varepsilon}(u^\varepsilon_{x_i}) F'(G_\varepsilon(\nabla u^\varepsilon)) |(G_\varepsilon(\nabla u^\varepsilon))_{x_i}|^2 \eta^2 dx \\ &\geq \frac{4\alpha}{(\alpha+1)^2} \sum_{i=1}^N \int g''_{i,\varepsilon}(u^\varepsilon_{x_i}) |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+1)/2})_{x_i}|^2 \eta^2 dx. \end{aligned}$$

From (3.15) we get

$$\begin{aligned} &\frac{4\alpha}{(\alpha+1)^2} \sum_{i=1}^N \int g''_{i,\varepsilon}(u^\varepsilon_{x_i}) |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+1)/2})_{x_i}|^2 \eta^2 dx + 4\mathcal{J}_2 \\ (3.17) \quad &\leq C(\alpha+1) \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha+2 \frac{p_N-1}{p_N}} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ &\quad + C(\alpha+1) \int |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\alpha \eta^2 dx, \end{aligned}$$

for some $C = C(N, p_1, p_N) > 0$.

We are only left with estimating \mathcal{J}_2 from below: recall that we have

$$\begin{aligned} \mathcal{J}_2 &= \sum_{i,k=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2 F(G_\varepsilon(\nabla u^\varepsilon)) g''_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx \\ &= \sum_{i,k=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\alpha g''_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx. \end{aligned}$$

We now observe that by (3.16) and through some lengthy though elementary computations, we get

$$\begin{aligned} |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/2})_{x_i}|^2 &= \frac{1}{4 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)} |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)_{x_i})|^2 \leq \frac{1}{4 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)} |(G_\varepsilon(\nabla u^\varepsilon)_{x_i})|^2 \\ &= \frac{1}{4 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)} \left| \left(\sum_{k=1}^N g_{k,\varepsilon}(u_{x_k}^\varepsilon) \right)_{x_i} \right|^2 = \frac{1}{4 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)} \left| \sum_{k=1}^N g'_{k,\varepsilon}(u_{x_k}^\varepsilon) u_{x_k x_i}^\varepsilon \right|^2 \\ &\leq \frac{N}{4 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)} \sum_{k=1}^N |g'_{k,\varepsilon}(u_{x_k}^\varepsilon)|^2 |u_{x_k x_i}^\varepsilon|^2. \end{aligned}$$

We then apply (2.4) or (2.7) on the last term, so to get

$$|(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/2})_{x_i}|^2 \leq \frac{N}{4 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)} \sum_{k=1}^N \delta_k g''_{k,\varepsilon}(u_{x_k}^\varepsilon) g_{k,\varepsilon}(u_{x_k}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2,$$

where δ_k is the same quantity defined in (3.11). We further observe that

$$\frac{g_{k,\varepsilon}(u_{x_k}^\varepsilon)}{\mathcal{G}_\varepsilon(\nabla u^\varepsilon)} \leq \frac{G_\varepsilon(\nabla u^\varepsilon)}{\mathcal{G}_\varepsilon(\nabla u^\varepsilon)} \leq 1.$$

This discussion leads us to

$$\frac{4}{N} \frac{1}{\delta} |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/2})_{x_i}|^2 \leq \sum_{k=1}^N g''_{k,\varepsilon}(u_{x_k}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2.$$

By inserting this inequality in \mathcal{J}_2 , we get

$$\begin{aligned} \mathcal{J}_2 &\geq \frac{4}{N} \frac{1}{\delta} \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\alpha |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/2})_{x_i}|^2 \eta^2 dx \\ &= \frac{4}{(\alpha+1)^2} \frac{1}{N \delta} \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+1)/2})_{x_i}|^2 \eta^2 dx. \end{aligned}$$

Finally, we can use this estimate in (3.17), so as to get the desired conclusion for $\alpha > 0$. The limit case $\alpha = 0$ can now be simply obtained by taking the limit α goes to 0 in the previously obtained estimate, since the relevant constant remains bounded. \blacksquare

Proposition 3.2 (Slow Moser's iteration). *Let $1 < p_1 \leq p_2 \leq \dots \leq p_N < \infty$. For every $\vartheta \geq 2/p'_N$ and every non-negative function $\eta \in C_0^2(B)$, we have*

$$(3.18) \quad \begin{aligned} & \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\vartheta+2/p_N} \eta^2 dx \\ & \leq \frac{C^\vartheta}{\vartheta} \int \eta^2 dx + C \|u^\varepsilon\|_{L^\infty(B)}^2 \vartheta^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\vartheta (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ & \quad + C^\vartheta \vartheta^\vartheta \|u^\varepsilon\|_{L^\infty(B)}^{\vartheta+2/p_N} \int |f^\varepsilon|^{\vartheta+2/p_N} \eta^2 dx, \end{aligned}$$

for some $C = C(N, p_1, p_N) > 0$.

Proof. We start by taking $\beta \geq 1$ and writing

$$\begin{aligned} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx &= \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta ((G_\varepsilon(\nabla u^\varepsilon) - 1)_+ + 1) \eta^2 dx \\ &\leq \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx + \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \sum_{k=1}^N g_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx. \end{aligned}$$

We observe that if we set

$$\sigma_k = \max\{1, 2^{(p_k-2)/2}\} \quad \text{and} \quad \bar{\sigma} = \max_{k=1,\dots,N} \sigma_k = \max\{1, 2^{(p_N-2)/2}\},$$

using (2.3) or (2.6) on the second integral, we get

$$\begin{aligned} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx &\leq \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx + \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \sum_{k=1}^N \sigma_k \frac{\varepsilon^{p_k/2}}{p_k} \eta^2 dx \\ &\quad + \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \sum_{k=1}^N \sigma_k \frac{g'_{k,\varepsilon}(u_{x_k}^\varepsilon)}{p_k} u_{x_k}^\varepsilon \eta^2 dx. \end{aligned}$$

By recalling that $g'_{k,\varepsilon}(t) t \geq 0$ and using that $0 < \varepsilon \leq 1$, we get

$$\begin{aligned} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx &\leq \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx + \bar{\sigma} \frac{N}{p_1} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx \\ &\quad + \frac{\bar{\sigma}}{p_1} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \sum_{k=1}^N g'_{k,\varepsilon}(u_{x_k}^\varepsilon) u_{x_k}^\varepsilon \eta^2 dx. \end{aligned}$$

In the last term, using the product rule and equation (2.14), we obtain

$$\begin{aligned} & \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx \\ & \leq \left(1 + \bar{\sigma} \frac{N}{p_1}\right) \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx - \frac{\bar{\sigma}}{p_1} \sum_{k=1}^N \int (\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta)_{x_k} g'_{k,\varepsilon}(u_{x_k}^\varepsilon) u^\varepsilon \eta^2 dx \\ & \quad - \frac{2\bar{\sigma}}{p_1} \sum_{k=1}^N \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta g'_{k,\varepsilon}(u_{x_k}^\varepsilon) u^\varepsilon \eta \eta_{x_k} dx + \frac{\bar{\sigma}}{p_1} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) u^\varepsilon \eta^2 dx. \end{aligned}$$

By using that u^ε is bounded and that $0 \leq (\cdot^\varepsilon)' \leq 1$, we get

$$\begin{aligned}
 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx &\leq \left(1 + \bar{\sigma} \frac{N}{p_1}\right) \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx \\
 &+ \frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \sum_{k=1}^N \int |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta)_{x_k}| |g'_{k,\varepsilon}(u^\varepsilon_{x_k})| \eta^2 dx \\
 (3.19) \quad &+ \frac{2\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \sum_{k=1}^N \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta |g'_{k,\varepsilon}(u^\varepsilon_{x_k})| \eta |\eta_{x_k}| dx \\
 &+ \frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta |f^\varepsilon| \eta^2 dx.
 \end{aligned}$$

Now, (2.10) together with the fact that $\mathcal{G}_\varepsilon \geq G_\varepsilon$ and $\mathcal{G}_\varepsilon \geq 1$ entail

$$\begin{aligned}
 &\frac{2\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \sum_{k=1}^N \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta |g'_{k,\varepsilon}(u^\varepsilon_{x_k})| \eta |\eta_{x_k}| dx \\
 &\leq \frac{2\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} N p_N^{(p_N-1)/p_N} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1-1/p_N} \eta |\nabla \eta| dx \\
 &\leq \tau \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx \\
 &+ \frac{1}{\tau} (N p_N^{(p_N-1)/p_N})^2 \left(\frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1}\right)^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1-2/p_N} |\nabla \eta|^2 dx,
 \end{aligned}$$

where in the last inequality we applied Young's inequality.

By choosing $\tau = 1/2$, we can absorb the term containing $\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1}$, and we get from (3.19) the following:

$$\begin{aligned}
 &\frac{1}{2} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx \\
 (3.20) \quad &\leq \left(1 + \frac{\bar{\sigma} N}{p_1}\right) \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx \\
 &+ \frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \sum_{k=1}^N \int |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta)_{x_k}| |g'_{k,\varepsilon}(u^\varepsilon_{x_k})| \eta^2 dx \\
 &+ 2(N p_N^{(p_N-1)/p_N})^2 \left(\frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1}\right)^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1-2/p_N} |\nabla \eta|^2 dx \\
 &+ \frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta |f^\varepsilon| \eta^2 dx.
 \end{aligned}$$

For the second term on the right-hand side, we use Young's inequality: for every $\tau > 0$,

$$\begin{aligned}
& \frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \sum_{k=1}^N \int |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta)_{x_k} | |g'_{k,\varepsilon}(u^\varepsilon_{x_k})| \eta^2 dx \\
&= \frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \sum_{k=1}^N \beta \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta-1} |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon))_{x_k} | |g'_{k,\varepsilon}(u^\varepsilon_{x_k})| \eta^2 dx \\
&\leq \left(\frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \right)^2 \frac{\beta^2}{2\tau} \sum_{k=1}^N \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta-2} |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon))_{x_k} |^2 g''_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta^2 dx \\
&\quad + \frac{\tau}{2} \sum_{k=1}^N \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \frac{(g'_{k,\varepsilon}(u^\varepsilon_{x_k}))^2}{g''_{k,\varepsilon}(u^\varepsilon_{x_k})} \eta^2 dx \\
&= \frac{2}{\tau} \left(\frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \right)^2 \sum_{k=1}^N \int |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta/2})_{x_k} |^2 g''_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta^2 dx \\
&\quad + \frac{\tau}{2} \sum_{k=1}^N \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \frac{(g'_{k,\varepsilon}(u^\varepsilon_{x_k}))^2}{g''_{k,\varepsilon}(u^\varepsilon_{x_k})} \eta^2 dx.
\end{aligned}$$

We also notice that by (3.12) we have

$$\sum_{k=1}^N \frac{(g'_{k,\varepsilon}(u^\varepsilon_{x_k}))^2}{g''_{k,\varepsilon}(u^\varepsilon_{x_k})} \leq \bar{\delta} \mathcal{G}_\varepsilon(\nabla u^\varepsilon).$$

Thus from (3.20), we get

$$\begin{aligned}
& \frac{1}{2} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx \leq \left(1 + \frac{\bar{\sigma} N}{p_1} \right) \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx \\
&\quad + \frac{2}{\tau} \left(\frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \right)^2 \sum_{k=1}^N \int |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta/2})_{x_k} |^2 g''_{k,\varepsilon}(u^\varepsilon_{x_k}) \eta^2 dx \\
&\quad + \frac{\tau \bar{\delta}}{2} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx \\
&\quad + 2 \left(N p_N^{(p_N-1)/p_N} \right)^2 \left(\frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \right)^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1-2/p_N} |\nabla \eta|^2 dx \\
&\quad + \frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta |f^\varepsilon| \eta^2 dx.
\end{aligned}$$

By choosing $\tau = 1/(2\bar{\delta})$, we can absorb again the term containing $\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1}$ on the right-hand side and obtain

$$\begin{aligned} \frac{1}{4} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx &\leq \left(1 + \frac{\bar{\sigma} N}{p_1}\right) \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx \\ &\quad + 4\bar{\delta} \left(\frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1}\right)^2 \sum_{k=1}^N \int |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta/2})_{x_k}|^2 g''_{k,\varepsilon}(u_{x_k}^\varepsilon) \eta^2 dx \\ &\quad + 2(Np_N^{(p_N-1)/p_N})^2 \left(\frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1}\right)^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1-2/p_N} |\nabla \eta|^2 dx \\ &\quad + \frac{\bar{\sigma} \|u^\varepsilon\|_{L^\infty(B)}}{p_1} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta |f^\varepsilon| \eta^2 dx. \end{aligned}$$

On the right-hand side, we now use the Caccioppoli inequality (3.2) with $\alpha = \beta - 1 \geq 0$, so to get

$$\begin{aligned} &\sum_{k=1}^N \int g''_{k,\varepsilon}(u_{x_k}^\varepsilon) |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta/2})_{x_k}|^2 \eta^2 dx \\ &\leq C\beta^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1-2/p_N} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx + C\beta^2 \int |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta-1} \eta^2 dx, \end{aligned}$$

for some $C = C(N, p_1, p_N) > 0$. This finally gives

$$\begin{aligned} &\int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx \\ &\leq C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx + C \|u^\varepsilon\|_{L^\infty(B)}^2 \beta^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1-2/p_N} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ &\quad + C \|u^\varepsilon\|_{L^\infty(B)}^2 \beta^2 \int |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta-1} \eta^2 dx + C \|u^\varepsilon\|_{L^\infty(B)} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta |f^\varepsilon| \eta^2 dx, \end{aligned}$$

for some $C = C(N, p_1, p_N) > 0$. On the first term on the right-hand side, we can use Young's inequality:

$$C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta \eta^2 dx \leq \frac{\tau \beta}{\beta + 1} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx + \frac{C\beta^{+1}}{\tau \beta (\beta + 1)} \int \eta^2 dx.$$

By choosing $\tau = 1/2$, we can re-absorb the term $\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1}$. This gives

$$\begin{aligned} &\frac{1}{2} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx \\ &\leq \frac{2^\beta C\beta^{+1}}{\beta + 1} \int \eta^2 dx + C \|u^\varepsilon\|_{L^\infty(B)}^2 \beta^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1-2/p_N} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ &\quad + C \|u^\varepsilon\|_{L^\infty(B)}^2 \beta^2 \int |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta-1} \eta^2 dx + C \|u^\varepsilon\|_{L^\infty(B)} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta |f^\varepsilon| \eta^2 dx. \end{aligned}$$

We proceed in a similar way for the two terms containing f^ε . By using Young's inequality with exponents

$$\frac{\beta + 1}{\beta - 1}, \frac{\beta + 1}{2} \quad \text{and} \quad \frac{\beta + 1}{\beta}, \beta + 1,$$

respectively, we get

$$\begin{aligned} & C \|u^\varepsilon\|_{L^\infty(B)}^2 \beta^2 \int |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta-1} \eta^2 dx + C \|u^\varepsilon\|_{L^\infty(B)} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\beta |f^\varepsilon| \eta^2 dx \\ & \leq \tau \frac{\beta-1}{\beta+1} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx + \frac{2\tau^{\frac{1-\beta}{2}} C^{\frac{\beta+1}{2}}}{\beta+1} \|u^\varepsilon\|_{L^\infty(B)}^{\beta+1} \beta^{\beta+1} \int |f^\varepsilon|^{\beta+1} \eta^2 dx \\ & \quad + \tau \frac{\beta}{\beta+1} \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\beta+1} \eta^2 dx + \frac{C^{\beta+1}}{\tau^\beta (\beta+1)} \|u^\varepsilon\|_{L^\infty(B)}^{\beta+1} \int |f^\varepsilon|^{\beta+1} \eta^2 dx. \end{aligned}$$

By choosing $\tau = 1/8$, we can absorb again the terms containing the power $\beta + 1$ of $\mathcal{G}_\varepsilon(\nabla u^\varepsilon)$. This finally leads to the estimate (3.18), up to renaming $\vartheta = \beta + 1 - 2/p_N$. This concludes the proof. \blacksquare

4. Uniform higher integrability

In this section, we establish a higher integrability estimate for ∇u^ε , which will eventually lead to the result of Proposition 1.1. Throughout this section, we shall assume that $1 < p_1 \leq \dots \leq p_N$, without any further restriction. Up to some technical facts, the proof of the next result is simply based on iterating the estimate of Proposition 3.2, starting from

$$\vartheta_0 = \frac{2}{p'_N}.$$

However, some care is required in order to get an a priori estimate only depending on the L^1 norm of \mathcal{G}_ε : observe indeed that $\vartheta_0 > 1$ in the case $p_N > 2$. For this purpose, we further need an interpolation trick.

Proposition 4.1. *Let $1 < p_1 \leq p_2 \leq \dots \leq p_N < \infty$. Then for every pair of concentric balls $B_r \Subset B_R \Subset B$ and every $\gamma \geq 2$, we have*

$$\int_{B_r} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\gamma dx \leq \Gamma_1 + \Gamma_2 \int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx,$$

for two constants $\Gamma_1, \Gamma_2 > 0$ depending on

$$N, p_N, p_1, \gamma, R, R-r, \|f^\varepsilon\|_{L^\gamma(B_R)} \text{ and } \|u^\varepsilon\|_{L^\infty(B)}.$$

Proof. We take $\gamma \geq 2$ and define the sequence of exponents

$$\vartheta_0 = \frac{2}{p'_N}, \quad \vartheta_{i+1} = \vartheta_i + \frac{2}{p_N} = \frac{2}{p'_N} + (i+1) \frac{2}{p_N} = 2 + i \frac{2}{p_N}, \quad \text{for } i \in \mathbb{N}.$$

We set

$$(4.1) \quad i_0 = \max \left\{ i \in \mathbb{N} : i \leq \frac{p_N}{2} (\gamma - 2) \right\}.$$

This in particular implies that

$$\vartheta_{i_0+1} \leq \gamma < \vartheta_{i_0+2}.$$

We now need to distinguish various cases, according to the values of p_N and γ , allowing to iteratively refine estimate (3.18).

Case A.1. Here we assume that

$$(4.2) \quad p_N \leq 2 \quad \text{and} \quad \frac{p_N}{2} (\gamma - 2) \in \mathbb{N}.$$

This is the simplest case: we get the estimate by iterating Proposition 3.2 with exponents $\vartheta = \vartheta_i$ and a suitable sequence of shrinking balls.

More precisely, we fix B_r and B_R as in the statement, and define the sequence of decreasing radii

$$r_i = R - i \frac{R - r}{i_0 + 1}, \quad \text{for } i = 0, \dots, i_0 + 1.$$

Accordingly, we take a cut-off function $\eta_i \in C_0^2(B_{r_i})$ for $i = 0, \dots, i_0$, such that

$$0 \leq \eta_i \leq 1, \quad \eta_i \equiv 1 \text{ on } B_{r_{i+1}}, \quad |\nabla \eta_i|^2 + |D^2 \eta_i| \leq \frac{C(i_0 + 1)^2}{(R - r)^2}.$$

By applying (3.18) with $\vartheta = \vartheta_i$, $\eta = \eta_i$, and using the properties of the cut-off function, we get

$$\begin{aligned} \int_{B_{r_{i+1}}} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\vartheta_{i+1}} dx &\leq \frac{C^{\vartheta_i}}{\vartheta_i} |B_R| + \frac{C \vartheta_i^2 (i_0 + 1)^2}{(R - r)^2} \|u^\varepsilon\|_{L^\infty(B)}^2 \int_{B_{r_i}} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\vartheta_i} dx \\ &\quad + C^{\vartheta_i} \vartheta_i^{\vartheta_i} \|u^\varepsilon\|_{L^\infty(B)}^{\vartheta_{i+1}} \int_{B_{r_i}} |f^\varepsilon|^{\vartheta_{i+1}} dx, \end{aligned}$$

for a constant $C = C(N, p_1, p_N) > 0$. By using Hölder's inequality on the right-hand side, we also get

$$(4.3) \quad \begin{aligned} \int_{B_{r_{i+1}}} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\vartheta_{i+1}} dx &\leq \frac{C^{\vartheta_i}}{\vartheta_i} |B_R| + \frac{C \vartheta_i^2 (i_0 + 1)^2}{(R - r)^2} \|u^\varepsilon\|_{L^\infty(B)}^2 \int_{B_{r_i}} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\vartheta_i} dx \\ &\quad + C^{\vartheta_i} \vartheta_i^{\vartheta_i} |B_R|^{1-\vartheta_{i+1}/\gamma} \|u^\varepsilon\|_{L^\infty(B)}^{\vartheta_{i+1}} \left(\int_{B_R} |f^\varepsilon|^\gamma dx \right)^{\vartheta_{i+1}/\gamma}. \end{aligned}$$

Starting from $i = 0$ and iterating (4.3) from 0 to i_0 , we get

$$(4.4) \quad \int_{B_r} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\vartheta_{i_0+1}} dx \leq \mathcal{M} + D_0 \int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx,$$

where we set, for notational simplicity, and for every natural number $0 \leq k \leq i_0$,

$$D_k = \left[\prod_{i=k}^{i_0} \frac{C \vartheta_i^2 (i_0 + 1)^2}{(R - r)^2} \|u^\varepsilon\|_{L^\infty(B)}^2 \right] = \left[\frac{C(i_0 + 1)^2}{(R - r)^2} \|u^\varepsilon\|_{L^\infty(B)}^2 \right]^{i_0-k+1} \prod_{i=k}^{i_0} \vartheta_i^2,$$

while

$$\mathcal{M} = \sum_{i=0}^{i_0} \left(\frac{C^{\vartheta_i}}{\vartheta_i} |B_R| + C^{\vartheta_i} \vartheta_i^{\vartheta_i} |B_R|^{1-\vartheta_{i+1}/\gamma} (\|u^\varepsilon\|_{L^\infty(B)} \|f\|_{L^\gamma(B_R)})^{\vartheta_{i+1}} \right) D_{i+1},$$

with the notational agreement that $D_{i_0+1} = 1$. Their precise expression is not very important, but we point out that D_0 and \mathcal{M} depend only on

$$N, p_N, p_1, \gamma, R, R - r, \|f_\varepsilon\|_{L^\gamma(B_R)} \text{ and } \|u_\varepsilon\|_{L^\infty(B)}.$$

Thanks to the assumption (4.2) and recalling that $\mathcal{G}_\varepsilon \geq 1$, we get

$$\vartheta_{i_0+1} = \gamma \quad \text{and} \quad \int_{B_R} \mathcal{G}_\varepsilon (\nabla u^\varepsilon)^{2/p'_N} dx \leq \int_{B_R} \mathcal{G}_\varepsilon (\nabla u^\varepsilon) dx,$$

thus the desired conclusion follows from (4.4).

Case A.2. Here we assume that

$$p_N \leq 2 \quad \text{and} \quad \frac{p_N}{2} (\gamma - 2) \notin \mathbb{N}.$$

In light of the second assumption, we have

$$\vartheta_{i_0+1} < \gamma < \vartheta_{i_0+2},$$

where i_0 is the index defined in (4.1). Thus in this case, we need an extra step of the iteration, by suitably adapting the choice of the exponent ϑ . We take $B_r \Subset B_\varrho \Subset B_R$, where

$$\varrho = \frac{R + r}{2},$$

and define the sequence of decreasing radii

$$r_i = R - i \frac{R - \varrho}{i_0 + 1}, \quad \text{for } i = 0, \dots, i_0 + 1.$$

We take a cut-off function $\eta_i \in C_0^2(B_{r_i})$ for $i = 0, \dots, i_0$, such that

$$0 \leq \eta_i \leq 1, \quad \eta_i \equiv 1 \text{ on } B_{r_{i+1}}, \quad |\nabla \eta_i|^2 + |D^2 \eta_i| \leq \frac{C(i_0 + 1)^2}{(R - \varrho)^2}.$$

By proceeding as above, we now get

$$(4.5) \quad \int_{B_\varrho} \mathcal{G}_\varepsilon (\nabla u^\varepsilon)^{\vartheta_{i_0+1}} dx \leq \mathcal{M} + D_0 \int_{B_R} \mathcal{G}_\varepsilon (\nabla u^\varepsilon)^{2/p'_N} dx.$$

The last term can be estimated from above, by using again that $2/p'_N \leq 1$. However, in order to reach the desired exponent $\gamma > \vartheta_{i_0+1}$, we need to apply (3.18) once more. We take a cut-off function $\eta \in C_0^2(B_\varrho)$ such that

$$0 \leq \eta \leq 1, \quad \eta \equiv 1 \text{ on } B_r, \quad |\nabla \eta|^2 + |D^2 \eta| \leq \frac{C}{(\varrho - r)^2}.$$

By applying (3.18) with³

$$\vartheta = \gamma - \frac{2}{p_N},$$

³Observe that such a choice is feasible, since $\gamma - 2/p_N \geq 2/p'_N$ if and only if $\gamma \geq 2$.

and the cut-off function above, we get

$$\begin{aligned} \int_{B_r} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\gamma dx &\leq C|B_R| + C\|u^\varepsilon\|_{L^\infty(B)}^2 \int_{B_\varrho} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\gamma-2/p_N} dx \\ &\quad + C\|u^\varepsilon\|_{L^\infty(B)}^\gamma \int_{B_R} |f^\varepsilon|^\gamma dx. \end{aligned}$$

On the right-hand side, the term containing $\mathcal{G}_\varepsilon(\nabla u^\varepsilon)$ is under control, since by construction,

$$\gamma - \frac{2}{p_N} < \vartheta_{i_0+2} - \frac{2}{p_N} = \vartheta_{i_0+1}.$$

Thus we have

$$\int_{B_\varrho} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\gamma-2/p_N} dx \leq \int_{B_\varrho} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\vartheta_{i_0+1}} dx,$$

and the last term can be estimated by (4.5).

Case B. Here we assume that $p_N > 2$. The proof goes exactly as before, so, for every $r < R$ with $B_R \Subset B$, we certainly have

$$\int_{B_r} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\gamma dx \leq \Gamma_1 + \Gamma_2 \int_{B_{(R+r)/2}} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx,$$

but now the major difference is that we need to estimate the integral on the right-hand side. Indeed, in this case $2/p'_N > 1$ and we cannot directly assure that this term is bounded, uniformly in ε . We need to use an interpolation trick to get a reverse L^{2/p'_N} - L^1 estimate on $\mathcal{G}_\varepsilon(\nabla u^\varepsilon)$. We denote

$$\varrho = \frac{R+r}{2},$$

and we observe that

$$1 < \frac{2}{p'_N} < 2,$$

thus by interpolation in Lebesgue spaces, we get for every $\varrho \leq s < t \leq R$,

$$(4.6) \quad \int_{B_s} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx \leq \left(\int_{B_s} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right)^{2/p_N} \left(\int_{B_s} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^2 dx \right)^{(p_N-2)/p_N}.$$

The L^2 norm on the right-hand side can in turn be estimated by means of (3.18), observing that

$$2 = \frac{2}{p'_N} + \frac{2}{p_N}.$$

By taking $\vartheta = 2/p'_N$ and a cut-off function $\eta \in C_0^2(B_t)$ such that

$$0 \leq \eta \leq 1, \quad \eta \equiv 1 \text{ on } B_s, \quad |\nabla \eta|^2 + |D^2 \eta| \leq \frac{C}{(t-s)^2},$$

we thus get, from (3.18),

$$\begin{aligned}
\int_{B_s} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^2 dx &\leq C|B_R| + \frac{C}{(t-s)^2} \|u^\varepsilon\|_{L^\infty(B)}^2 \int_{B_t} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx \\
&\quad + C \|u^\varepsilon\|_{L^\infty(B)}^2 \int_{B_R} |f^\varepsilon|^2 dx \\
&\leq C|B_R| + \frac{C}{(t-s)^2} \|u^\varepsilon\|_{L^\infty(B)}^2 \int_{B_t} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx \\
&\quad + C|B_R|^{1-2/\gamma} \|u^\varepsilon\|_{L^\infty(B)}^2 \left(\int_{B_R} |f^\varepsilon|^\gamma dx \right)^{2/\gamma},
\end{aligned}$$

for $C = C(N, p_1, p_N) > 0$. We insert the above inequality into (4.6) and use the subadditivity of $\tau \mapsto \tau^{(p_N-2)/p_N}$ to get

$$\begin{aligned}
\int_{B_s} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx &\leq \left(\int_{B_s} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right)^{2/p_N} (C|B_R|)^{(p_N-2)/p_N} \\
&\quad + \left(\int_{B_s} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right)^{2/p_N} \left(\frac{C}{(t-s)^2} \|u^\varepsilon\|_{L^\infty(B)}^2 \right)^{\frac{p_N-2}{p_N}} \left(\int_{B_t} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx \right)^{\frac{p_N-2}{p_N}} \\
&\quad + \left(\int_{B_s} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right)^{2/p_N} (C|B_R|^{1-2/\gamma} (\|u^\varepsilon\|_{L^\infty(B)} \|f^\varepsilon\|_{L^\gamma(B_R)})^2)^{\frac{p_N-2}{p_N}}.
\end{aligned}$$

For the second term on the right-hand side, we apply Young's inequality with conjugate exponents $p_N/2$ and $p_N/(p_N-2)$. We get

$$\begin{aligned}
\int_{B_s} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx &\leq \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right)^{2/p_N} (C|B_R|)^{(p_N-2)/p_N} \\
&\quad + \frac{2}{p_N} \int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \left(\frac{C}{(t-s)^2} \|u^\varepsilon\|_{L^\infty(B)}^2 \right)^{(p_N-2)/2} + \frac{p_N-2}{p_N} \int_{B_t} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx \\
&\quad + \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right)^{2/p_N} (C|B_R|^{1-2/\gamma} (\|u^\varepsilon\|_{L^\infty(B)} \|f^\varepsilon\|_{L^\gamma(B_R)})^2)^{(p_N-2)/p_N}.
\end{aligned}$$

We now use Lemma 2.7 with the choices

$$\begin{aligned}
Z(s) &= \int_{B_s} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx, \quad \vartheta = \frac{p_N-2}{p_N}, \\
\mathcal{A} &= \frac{2}{p_N} \int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx (C \|u^\varepsilon\|_{L^\infty(B)}^2)^{(p_N-2)/2}, \quad \alpha_0 = p_N - 2, \quad \mathcal{B} = 0,
\end{aligned}$$

and

$$\begin{aligned}
\mathcal{C} &= \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right)^{2/p_N} \\
&\quad \cdot \left[(C|B_R|)^{(p_N-2)/p_N} + (C|B_R|^{1-2/\gamma} (\|u^\varepsilon\|_{L^\infty(B)} \|f^\varepsilon\|_{L^\gamma(B_R)})^2)^{(p_N-2)/p_N} \right],
\end{aligned}$$

in order to absorb the penultimate integral. This permits to conclude that

$$\begin{aligned} \int_{B_\varepsilon} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p'_N} dx &\leq \tilde{C} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right)^{2/p_N} (C|B_R|)^{(p_N-2)/p_N} \\ &+ \tilde{C} \frac{2}{p_N} \frac{1}{(R-r)^{p_N-2}} \int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \left(C \|u^\varepsilon\|_{L^\infty(B)}^2 \right)^{(p_N-2)/2} \\ &+ \tilde{C} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right)^{2/p_N} (|B_R|^{1-2/\gamma} (\|u^\varepsilon\|_{L^\infty(B)} \|f^\varepsilon\|_{L^\gamma(B_R)})^2)^{(p_N-2)/p_N}. \end{aligned}$$

This gives the claimed $L^{2/p'_N}-L^1$ estimate on $\mathcal{G}_\varepsilon(\nabla u^\varepsilon)$. The desired conclusion now easily follows. We leave the details to the reader. \blacksquare

Remark 4.2 (Quality of the constants). For future references, it is important to notice that the two constants Γ_1 and Γ_2 in the previous statement are uniformly bounded from above, whenever there exists a constant $C \geq 1$ such that

$$\|f^\varepsilon\|_{L^\gamma(B_R)} + \|u^\varepsilon\|_{L^\infty(B)} \leq C, \quad \text{for every } 0 < \varepsilon \leq \varepsilon_0,$$

and

$$R - r \geq \frac{1}{C}.$$

On the contrary, we see from the proof above that

$$\lim_{\gamma \rightarrow \infty} \Gamma_i = +\infty, \quad \text{for } i = 1, 2.$$

5. Uniform Lipschitz bound

We now establish a local L^∞ estimate for ∇u^ε : this time, this will lead to Theorem L.

Proposition 5.1. *Let $1 < p_1 \leq \dots \leq p_N \leq 2$ and $0 < \varepsilon \leq \varepsilon_0$. Then for every pair of concentric balls $B_r \Subset B_R \Subset B$, $R < 1$, and every $q > N$, we have*

$$\begin{aligned} &\|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^\infty(B_r)} \\ &\leq C \left[\frac{1}{(R-r)^{Nq/(q-N)}} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^q dx \right)^{N/(q-N)} + \|f^\varepsilon\|_{L^q(B_R)}^{Nq/(q-N)} \right] \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^1(B_R)}, \end{aligned}$$

for some $C = C(N, p_N, p_1, q) > 0$.

Proof. We will use a Moser's iteration scheme, in order to get the claimed estimate. By (2.2), for every $i = 1, \dots, N$, we have

$$\begin{aligned} g''_{i,\varepsilon}(u^\varepsilon_{x_i}) &\geq (p_i - 1) (\varepsilon + |u^\varepsilon_{x_i}|^2)^{(p_i-2)/2} = (p_i - 1) (p_i g_{i,\varepsilon}(u^\varepsilon_{x_i}))^{(p_i-2)/p_i} \\ &\geq (p_i - 1) (p_i G_\varepsilon(\nabla u^\varepsilon))^{(p_i-2)/p_i} \geq (p_i - 1) (p_i \mathcal{G}_\varepsilon(\nabla u^\varepsilon))^{(p_i-2)/p_i}, \end{aligned}$$

thanks to the fact that $p_i \leq 2$, for every $i = 1, \dots, N$. We can further estimate the last term from below as follows:

$$g''_{i,\varepsilon}(u^\varepsilon_{x_i}) \geq (p_1 - 1) (p_N)^{(p_1-2)/p_1} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(p_1-2)/p_1}.$$

By using this lower bound in (3.2), we get

$$\begin{aligned}
 & \sum_{i=1}^N \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(p_1-2)/p_1} |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+1)/2})_{x_i}|^2 \eta^2 dx \\
 (5.1) \quad & \leq C(\alpha+1)^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha+2-2/p_N} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\
 & \quad + C(\alpha+1)^2 \int |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\alpha \eta^2 dx,
 \end{aligned}$$

for some $C = C(N, p_1, p_N) > 0$. With simple algebraic manipulations, for every $\alpha \geq 0$ we have

$$\begin{aligned}
 & \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(p_1-2)/p_1} |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+1)/2})_{x_i}|^2 \\
 & = \left(\frac{\alpha+1}{2}\right)^2 \frac{1}{\left(\frac{\alpha+2}{2} - \frac{1}{p_1}\right)^2} |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+2)/2-1/p_1})_{x_i}|^2 \geq |(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+2)/2-1/p_1})_{x_i}|^2.
 \end{aligned}$$

Thus from (5.1) we obtain

$$\begin{aligned}
 & \int |\nabla (\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+2)/2-1/p_1})|^2 \eta^2 dx \\
 (5.2) \quad & \leq C(\alpha+1)^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha+2-2/p_N} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\
 & \quad + C(\alpha+1)^2 \int |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\alpha \eta^2 dx,
 \end{aligned}$$

for some $C = C(N, p_1, p_N) > 0$. By adding on both sides the term

$$\int (\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+2)/2-1/p_1})^2 |\nabla \eta|^2 dx,$$

and using again that $\mathcal{G}_\varepsilon \geq 1$, we then obtain from (5.2),

$$\begin{aligned}
 & \int |\nabla ((\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+2)/2-1/p_1}) \eta)|^2 dx \\
 (5.3) \quad & \leq C(\alpha+1)^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha+2-2/p_N} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\
 & \quad + C(\alpha+1)^2 \int |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\alpha \eta^2 dx,
 \end{aligned}$$

possibly for a different constant $C = C(N, p_1, p_N) > 0$. Let us suppose for simplicity that $N \geq 3$. The case $N = 2$ can be treated with minor modifications. On the left-hand side of (5.3), we then use Sobolev's inequality in $W^{1,2}(\mathbb{R}^N)$. This gives

$$\begin{aligned}
 & \left(\int (\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+2)/2-1/p_1} \eta)^{2^*} dx \right)^{2/2^*} \\
 (5.4) \quad & \leq C(\alpha+1)^2 \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha+2-2/p_N} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\
 & \quad + C(\alpha+1)^2 \int |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\alpha \eta^2 dx,
 \end{aligned}$$

with some new constant $C = C(N, p_1, p_N) > 0$. We choose $\eta \in C_0^2(B_R)$ to be a cut-off function such that

$$0 \leq \eta \leq 1, \quad \eta \equiv 1 \text{ on } B_r, \quad |\nabla \eta|^2 + |D^2 \eta| \leq \frac{C}{(R-r)^2}.$$

Thus we obtain, from (5.4),

$$\begin{aligned} \left(\int_{B_r} \left(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+2)/2-1/p_1} \right)^{2^*} dx \right)^{2/2^*} &\leq C \frac{(\alpha+1)^2}{(R-r)^2} \int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha+2-2/p_N} dx \\ &\quad + C(\alpha+1)^2 \int_{B_R} |f^\varepsilon|^2 \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\alpha dx. \end{aligned}$$

We now take an exponent $q > N$. By Hölder's inequality on the last term, we deduce that

$$\begin{aligned} (5.5) \quad &\left(\int_{B_r} \left(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+2)/2-1/p_1} \right)^{2^*} dx \right)^{2/2^*} \\ &\leq C \frac{(\alpha+1)^2}{(R-r)^2} \int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha+2-2/p_N} dx \\ &\quad + C(\alpha+1)^2 \|f^\varepsilon\|_{L^q(B_R)}^2 \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha \frac{q}{q-2}} dx \right)^{(q-2)/q}. \end{aligned}$$

Before proceeding further, we rely on Hölder's inequality to get

$$\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha+2-2/p_N} dx \leq \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha \frac{q}{q-2}} dx \right)^{(q-2)/q} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{q/p'_N} dx \right)^{2/q}.$$

Moreover, by recalling that $\mathcal{G}_\varepsilon \geq 1$, we have

$$\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha q/(q-2)} \leq \left(\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\alpha+2-2/p_1} \right)^{q/(q-2)}.$$

By using these two facts in (5.5), we obtain

$$\begin{aligned} (5.6) \quad &\left(\int_{B_r} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+2-2/p_1) \frac{2^*}{2}} dx \right)^{2/2^*} \\ &\leq C(\alpha+1)^2 \left[\frac{1}{(R-r)^2} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{q/p'_N} dx \right)^{2/q} + \|f^\varepsilon\|_{L^q(B_R)}^2 \right] \\ &\quad \times \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(\alpha+2-2/p_1) \frac{q}{q-2}} dx \right)^{(q-2)/q}. \end{aligned}$$

We now set

$$\theta = \alpha + 2 - \frac{2}{p_1},$$

so that from (5.6), we get

$$\begin{aligned} (5.7) \quad &\left(\int_{B_r} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\frac{2^*}{2} \theta} dx \right)^{2/(2^* \theta)} \\ &\leq (C\theta^2)^{1/\theta} \left[\frac{1}{(R-r)^2} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{q/p'_N} dx \right)^{2/q} + \|f^\varepsilon\|_{L^q(B_R)}^2 \right]^{1/\theta} \\ &\quad \times \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\theta \frac{q}{q-2}} dx \right)^{(q-2)/(q\theta)}. \end{aligned}$$

We define the sequence of exponents through the following recursive relation:

$$\theta_0 = 1, \quad \theta_{i+1} \frac{q}{q-2} = \frac{2^*}{2} \theta_i, \quad \text{for } i \in \mathbb{N};$$

that is,⁴

$$\theta_{i+1} = \left(\frac{2^*}{2} \frac{q-2}{q} \right) \theta_i = \left(\frac{2^*}{2} \frac{q-2}{q} \right)^{i+1}, \quad \text{for } i \in \mathbb{N}.$$

We also define the classical sequence of shrinking radii

$$r_i = r + \frac{R-r}{2^i}, \quad \text{for } i \in \mathbb{N}.$$

With this notation, from (5.7) we get

$$\begin{aligned} (5.8) \quad & \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^{\frac{q}{q-2}\theta_{i+1}}(B_{r_{i+1}})} \\ & \leq \left[\frac{C}{(R-r)^2} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{q/p'_N} dx \right)^{2/q} + C \|f^\varepsilon\|_{L^q(B_R)}^2 \right]^{1/\theta_i} \\ & \quad \times (4^i \theta_i^2)^{1/\theta_i} \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^{\frac{q}{q-2}\theta_i}(B_{r_i})}. \end{aligned}$$

By starting from $i = 0$ and iterating (5.8) n times, we get

$$\begin{aligned} (5.9) \quad & \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^{\frac{q}{q-2}\theta_{n+1}}(B_{r_{n+1}})} \\ & \leq \left[\frac{C}{(R-r)^2} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{q/p'_N} dx \right)^{2/q} + C \|f^\varepsilon\|_{L^q(B_R)}^2 \right]^{\sum_{i=0}^n \frac{1}{\theta_i}} \\ & \quad \times \prod_{i=0}^n (4^i \theta_i^2)^{1/\theta_i} \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^{\frac{q}{q-2}}(B_R)}. \end{aligned}$$

By observing that

$$\lim_{n \rightarrow \infty} \prod_{i=0}^n (4^i \theta_i^2)^{1/\theta_i} =: C_{N,q} < +\infty,$$

and

$$\lim_{n \rightarrow \infty} \sum_{i=0}^n \frac{1}{\theta_i} = \sum_{i=0}^{\infty} \left(\frac{2q}{2^*(q-2)} \right)^i = \frac{N}{2} \frac{q-2}{q-N},$$

if we take the limit as n goes to ∞ in (5.9), we end up with

$$\begin{aligned} \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^\infty(B_r)} & \leq C \left[\frac{1}{(R-r)^2} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{q/p'_N} dx \right)^{2/q} + \|f^\varepsilon\|_{L^q(B_R)}^2 \right]^{\frac{N}{2} \frac{q-2}{q-N}} \\ & \quad \times \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^{q/(q-2)}(B_R)}, \end{aligned}$$

⁴Observe that

$$\frac{2q}{2^*(q-2)} < 1 \iff q > N,$$

and the latter holds true, in view of our assumption.

for some $C = C(N, p_1, p_N, q) > 0$. The previous estimate holds for every $r < R$ such that $B_R \Subset B$. Thus, we can now use a standard interpolation trick to rectify it and replace the $L^{q/(q-2)}$ norm on the right-hand side by the L^1 norm.

This goes as follows: we first observe that

$$\|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^{q/(q-2)}(B_R)} \leq (\|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^\infty(B_R)})^{2/q} (\|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^1(B_R)})^{(q-2)/q}.$$

Then by using Young's inequality with exponents $q/2$ and $q/(q-2)$, we get

$$\begin{aligned} & \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^\infty(B_r)} \\ & \leq \frac{q-2}{q} C^{q/(q-2)} \left[\frac{1}{(R-r)^2} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{q/p'_N} dx \right)^{2/q} + \|f^\varepsilon\|_{L^q(B_R)}^2 \right]^{\frac{N}{2} \frac{q}{q-2}} \\ & \quad \times \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^1(B_R)} + \frac{2}{q} \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^\infty(B_R)}. \end{aligned}$$

We now take s, t such that $r \leq s < t \leq R$. The previous estimate is valid by replacing r with s and R with t . Thus, with some simple algebraic manipulations, we get

$$\begin{aligned} & \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^\infty(B_s)} \\ & \leq \frac{q-2}{q} \frac{C^{\frac{q}{q-2}}}{2^{\frac{q(N-2)+2N}{2(N-q)}}} \left[\frac{1}{(t-s)^{\frac{Nq}{q-N}}} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{q/p'_N} dx \right)^{\frac{N}{q-N}} + \|f^\varepsilon\|_{L^q(B_R)}^{\frac{Nq}{q-N}} \right] \\ & \quad \times \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^1(B_R)} + \frac{2}{q} \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^\infty(B_t)}. \end{aligned}$$

By relying once again on Lemma 2.7, from the last estimate we get

$$\begin{aligned} & \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^\infty(B_r)} \\ & \leq \tilde{C} \left[\frac{1}{(R-r)^{\frac{Nq}{q-N}}} \left(\int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{q/p'_N} dx \right)^{\frac{N}{q-N}} + \|f^\varepsilon\|_{L^q(B_R)}^{\frac{Nq}{q-N}} \right] \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^1(B_R)}. \end{aligned}$$

By finally using that

$$\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{q/p'_N} \leq \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^q,$$

we eventually conclude the proof. \blacksquare

6. Uniform higher differentiability

At last, we prove a Sobolev-type regularity result for (some nonlinear function of) ∇u^ε , which eventually will permit to establish Theorem S.

Proposition 6.1. *Let $1 < p_1 \leq \dots \leq p_N \leq 2$. For $0 < \varepsilon \leq \varepsilon_0$ and $i = 1, \dots, N$, we set*

$$(6.1) \quad \mathcal{V}_{i,\varepsilon} = V_{i,\varepsilon}(u_{x_i}^\varepsilon), \quad \text{with } V_{i,\varepsilon}(t) = \int_0^t \sqrt{g''_{i,\varepsilon}(\tau)} d\tau.$$

Then for every non-negative $\eta \in C_0^2(B)$ and every $\gamma \geq 2$, we have⁵

$$(6.2) \quad \sum_{i=1}^N \int |\nabla \mathcal{V}_{i,\varepsilon}|^2 \eta^2 dx \leq C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1+2(1/p_1-1/p_N)} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ + C \left(\int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{\frac{2-p_1}{p_1} \frac{\gamma}{\gamma-2}} \eta^2 dx \right)^{(\gamma-2)/\gamma} \left(\int |f^\varepsilon|^\gamma \eta^2 dx \right)^{2/\gamma},$$

for some $C = C(N, p_N, p_1) > 0$.

Proof. We start by fixing $k \in \{1, \dots, N\}$ and inserting in the differentiated equation (2.15) the test function $\varphi = u_{x_k}^\varepsilon \eta^2$. Thus we get

$$\sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2 \eta^2 dx = -2 \sum_{i=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) u_{x_k x_i}^\varepsilon u_{x_k}^\varepsilon \eta \eta_{x_i} dx \\ + \int (f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon))_{x_k} u_{x_k}^\varepsilon \eta^2 dx.$$

For the first term of the right-hand side, we use the same trick as in the proof of Proposition 3.1: we observe that

$$g'_{i,\varepsilon}(u_{x_i}^\varepsilon) u_{x_k x_i}^\varepsilon = (g'_{i,\varepsilon}(u_{x_i}^\varepsilon))_{x_k},$$

and then integrate by parts. We integrate by parts the term $(f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon))_{x_k}$, as well. This yields

$$(6.3) \quad \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2 \eta^2 dx \\ = 2 \sum_{i=1}^N \int g'_{i,\varepsilon}(u_{x_i}^\varepsilon) (u_{x_k}^\varepsilon \eta \eta_{x_i})_{x_k} dx - \int f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) (u_{x_k}^\varepsilon \eta^2)_{x_k} dx \\ = \int u_{x_k x_k}^\varepsilon \left(2 \sum_{i=1}^N g'_{i,\varepsilon}(u_{x_i}^\varepsilon) \eta \eta_{x_i} - f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) \eta^2 \right) dx \\ + 2 \int u_{x_k}^\varepsilon \left(\sum_{i=1}^N g'_{i,\varepsilon}(u_{x_i}^\varepsilon) (\eta_{x_k} \eta_{x_i} + \eta \eta_{x_i x_k}) - \eta \eta_{x_k} f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) \right) dx.$$

By Young's inequality and the fact that $0 \leq (\zeta^\varepsilon)' \leq 1$, we can estimate the first term of the right-hand side as follows:

$$\int u_{x_k x_k}^\varepsilon \left(2 \sum_{i=1}^N g'_{i,\varepsilon}(u_{x_i}^\varepsilon) \eta \eta_{x_i} - f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) \eta^2 \right) dx \\ \leq \frac{1}{2} \int g''_{k,\varepsilon}(u_{x_k}^\varepsilon) |u_{x_k x_k}^\varepsilon|^2 \eta^2 dx + 4 \int \frac{1}{g''_{k,\varepsilon}(u_{x_k}^\varepsilon)} \left(\left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 |\nabla \eta|^2 + |f^\varepsilon|^2 \eta^2 \right) dx.$$

⁵When $\gamma = 2$, the last term is simply $C \|\mathcal{G}_\varepsilon \eta^2\|_{L^\infty}^{(2-p_1)/p_1} \|f^\varepsilon \eta\|_{L^2}^2$.

We use (2.2) to estimate $1/g''_{k,\varepsilon}$ on the right-hand side. On account of this inequality, we get

$$\begin{aligned} \int u_{x_k x_k}^\varepsilon \left(2 \sum_{i=1}^N g'_{i,\varepsilon}(u_{x_i}^\varepsilon) \eta \eta_{x_i} - f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) \eta^2 \right) dx &\leq \frac{1}{2} \int g''_{k,\varepsilon}(u_{x_k}^\varepsilon) |u_{x_k x_k}^\varepsilon|^2 \eta^2 dx \\ &+ \frac{4}{p_k - 1} \int (\varepsilon + (u_{x_k}^\varepsilon)^2)^{(2-p_k)/2} \left(\left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 |\nabla \eta|^2 + |f^\varepsilon|^2 \eta^2 \right) dx. \end{aligned}$$

Inserting the above inequality into (6.3) and absorbing the Hessian term of the right-hand side into the left-hand side, one gets

$$\begin{aligned} &\sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2 \eta^2 dx \\ (6.4) \quad &\leq \frac{8}{p_k - 1} \int (\varepsilon + (u_{x_k}^\varepsilon)^2)^{(2-p_k)/2} \left(\left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 |\nabla \eta|^2 + |f^\varepsilon|^2 \eta^2 \right) dx \\ &+ 4 \int u_{x_k}^\varepsilon \left(\sum_{i=1}^N g'_{i,\varepsilon}(u_{x_i}^\varepsilon) (\eta_{x_k} \eta_{x_i} + \eta \eta_{x_i x_k}) - \eta \eta_{x_k} f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) \right) dx. \end{aligned}$$

We now estimate the last term as follows:

$$\begin{aligned} &\left| \int u_{x_k}^\varepsilon \left(\sum_{i=1}^N g'_{i,\varepsilon}(u_{x_i}^\varepsilon) (\eta_{x_k} \eta_{x_i} + \eta \eta_{x_i x_k}) - \eta \eta_{x_k} f^\varepsilon(\zeta^\varepsilon)'(u^\varepsilon) \right) dx \right| \\ &\leq \int |u_{x_k}^\varepsilon| |f^\varepsilon| \eta |\nabla \eta| dx + \int |u_{x_k}^\varepsilon| \sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| (|\nabla \eta|^2 + \eta |D^2 \eta|) dx. \end{aligned}$$

Observe that we used again that $0 \leq (\zeta^\varepsilon)' \leq 1$. We apply Young's inequality on the last term to obtain

$$\begin{aligned} &\int |u_{x_k}^\varepsilon| \sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ &\leq \frac{1}{2} \int |u_{x_k}^\varepsilon|^{2-p_k} \left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\ &\quad + \frac{1}{2} \int |u_{x_k}^\varepsilon|^{p_k} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx. \end{aligned}$$

By further using that $p_k \leq 2$, we have

$$|u_{x_k}^\varepsilon|^{2-p_k} \leq (\varepsilon + (u_{x_k}^\varepsilon)^2)^{(2-p_k)/2} = p_k^{(2-p_k)/p_k} g_{k,\varepsilon}(u_{x_k}^\varepsilon)^{(2-p_k)/p_k}.$$

It follows from the above inequality and (6.4) that

$$\begin{aligned} & \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |u_{x_k x_i}^\varepsilon|^2 \eta^2 dx \\ & \leq C \int g_{k,\varepsilon}(u_{x_k}^\varepsilon)^{(2-p_k)/p_k} \left(\left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 |\nabla \eta|^2 + |f^\varepsilon|^2 \eta^2 \right) dx \\ & \quad + C \int |u_{x_k}^\varepsilon|^{p_k} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx + C \int |u_{x_k}^\varepsilon| |f^\varepsilon| \eta |\nabla \eta| dx, \end{aligned}$$

for some $C = C(p_1, p_N) > 0$. Then take the sum over $k = 1, \dots, N$. This gives

$$\begin{aligned} & \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |\nabla u_{x_i}^\varepsilon|^2 \eta^2 dx \\ (6.5) \quad & \leq C \int \sum_{k=1}^N g_{k,\varepsilon}(u_{x_k}^\varepsilon)^{(2-p_k)/p_k} \left(\left(\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \right)^2 |\nabla \eta|^2 + |f^\varepsilon|^2 \eta^2 \right) dx \\ & \quad + C \int \sum_{k=1}^N |u_{x_k}^\varepsilon|^{p_k} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx + C \int \sum_{k=1}^N |u_{x_k}^\varepsilon| |f^\varepsilon| \eta |\nabla \eta| dx. \end{aligned}$$

By Lemma 2.3, we have

$$\sum_{i=1}^N |g'_{i,\varepsilon}(u_{x_i}^\varepsilon)| \leq p_N^{(p_N-1)/p_N} \sum_{i=1}^N G_\varepsilon(\nabla u^\varepsilon)^{(p_i-1)/p_i} \leq N p_N^{(p_N-1)/p_N} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(p_N-1)/p_N}.$$

Moreover, by the definitions of $g_{k,\varepsilon}$, G_ε , and \mathcal{G}_ε , it is easily seen that

$$\begin{aligned} & \sum_{k=1}^N g_{k,\varepsilon}(u_{x_k}^\varepsilon)^{(2-p_k)/p_k} \leq C \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(2-p_1)/p_1}, \\ & \sum_{k=1}^N |u_{x_k}^\varepsilon|^{p_k} \leq \sum_{k=1}^N p_k g_{k,\varepsilon}(u_{x_k}^\varepsilon) \leq C \mathcal{G}_\varepsilon(\nabla u^\varepsilon) \quad \text{and} \quad \sum_{k=1}^N |u_{x_k}^\varepsilon| \leq C \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/p_1}, \end{aligned}$$

where all the constants depend only on N , p_1 and p_N . From (6.5), we get

$$\begin{aligned} & \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |\nabla u_{x_i}^\varepsilon|^2 \eta^2 dx \\ & \leq C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1+2(1/p_1-1/p_N)} |\nabla \eta|^2 dx + C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(2-p_1)/p_1} |f^\varepsilon|^2 \eta^2 dx \\ & \quad + C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon) (|\nabla \eta|^2 + \eta |D^2 \eta|) dx + C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/p_1} |f^\varepsilon| \eta |\nabla \eta| dx, \end{aligned}$$

for some $C = C(N, p_1, p_N) > 0$. Since

$$1 \leq 1 + 2 \left(\frac{1}{p_1} - \frac{1}{p_N} \right) \quad \text{and} \quad \mathcal{G}_\varepsilon(\nabla u^\varepsilon) \geq 1,$$

the third term can be absorbed in the first one, up to increasing C if necessary:

$$\begin{aligned}
 & \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |\nabla u_{x_i}^\varepsilon|^2 \eta^2 dx \\
 (6.6) \quad & \leq C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1+2(1/p_1-1/p_N)} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\
 & \quad + C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(2-p_1)/p_1} |f^\varepsilon|^2 \eta^2 dx + C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/p_1} |f^\varepsilon| \eta |\nabla \eta| dx.
 \end{aligned}$$

In the last term, we write

$$\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/p_1} |f^\varepsilon| \eta |\nabla \eta| = (\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/2} |\nabla \eta|) (\mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/p_1-1/2} |f^\varepsilon| \eta),$$

and use Young's inequality:

$$\begin{aligned}
 & \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1/p_1} |f^\varepsilon| \eta |\nabla \eta| dx \leq \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon) |\nabla \eta|^2 dx + \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p_1-1} |f^\varepsilon|^2 \eta^2 dx \\
 & \leq \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1+2(1/p_1-1/p_N)} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx + \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{2/p_1-1} |f^\varepsilon|^2 \eta^2 dx,
 \end{aligned}$$

where in the last line we have used again that $\mathcal{G}_\varepsilon \geq 1$. Inserting this estimate in (6.6), we obtain

$$\begin{aligned}
 & \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |\nabla u_{x_i}^\varepsilon|^2 \eta^2 dx \leq C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{1+2(1/p_1-1/p_N)} (|\nabla \eta|^2 + \eta |D^2 \eta|) dx \\
 & \quad + C \int \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^{(2-p_1)/p_1} |f^\varepsilon|^2 \eta^2 dx.
 \end{aligned}$$

On the left-hand side, we use the definition (6.1) of $\mathcal{V}_{i,\varepsilon}$ which gives that

$$|\nabla \mathcal{V}_{i,\varepsilon}|^2 = g''_{i,\varepsilon}(u_{x_i}^\varepsilon) |\nabla u_{x_i}^\varepsilon|^2.$$

This yields the desired conclusion when the exponent γ in the statement of Proposition 6.1 is equal to 2. When $\gamma > 2$, we only need to apply the Hölder inequality to the last term of the right-hand side with the exponents $\gamma/(\gamma-2)$ and $\gamma/2$. The proof is complete. ■

7. Proofs of the main results

We finally establish the three results presented in the introduction by relying on the relevant a priori estimates that we have obtained in the previous sections. We thus fix a ball $B_{4R}(x_0) \Subset \Omega$ as in the statements of Proposition 1.1, Theorem L and Theorem S: we are going to use the results of the previous sections, with the choice $B = B_{2R}(x_0)$.

We will use the functions \mathcal{S}_0 and \mathcal{S}_ε , defined by (1.1) and (3.1). Moreover, we will omit to indicate the centers of the balls, which will always be x_0 .

7.1. Proof of Proposition 1.1

In this section, we assume that $1 < p_1 \leq \dots \leq p_N < \infty$ and $f \in L^{\gamma}_{\text{loc}}(\Omega)$ for some $\gamma \geq 2$. We consider the ball $B_R \Subset B$. Then for every $0 < \varepsilon < \varepsilon_0$,

$$\|f^\varepsilon\|_{L^\gamma(B_R)} \leq \|f\|_{L^\gamma(2B)}.$$

Using also that $\|u^\varepsilon\|_{L^\infty(B)} \leq M + 1$, Proposition 4.1 and Remark 4.2 imply that for every $0 < \varepsilon < \varepsilon_0$ we have

$$(7.1) \quad \int_{B_{R/2}} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\gamma dx \leq \Gamma_1 + \Gamma_2 \int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx,$$

for two constants $\Gamma_1, \Gamma_2 > 0$ which do not depend on ε , but only on

$$N, p_N, p_1, \gamma, R, \|f\|_{L^\gamma(2B)} \text{ and } M = \|U\|_{L^\infty(2B)}.$$

In particular, by using that⁶

$$(7.2) \quad \mathcal{G}_0(z) \leq \mathcal{G}_\varepsilon(z) \leq C(\varepsilon^{p_1/2} + \mathcal{G}_0(z)), \quad \text{for every } z \in \mathbb{R}^N,$$

for some $C = C(N, p_N, p_1) > 0$, we can infer

$$\int_{B_{R/2}} \mathcal{G}_0(\nabla u^\varepsilon)^\gamma dx \leq \Gamma_1 + C\Gamma_2 \int_{B_R} (\varepsilon^{p_1/2} + \mathcal{G}_0(\nabla u^\varepsilon)) dx.$$

In view of Lemma 2.6, there exists an infinitesimal sequence $\{\varepsilon_k\}_{k \in \mathbb{N}}$ such that

$$(u^{\varepsilon_k}, \nabla u^{\varepsilon_k}) \text{ converges to } (U, \nabla U) \quad \text{a. e. in } B.$$

We then take the limit on both sides of the estimate above and use Fatou's lemma on the left. We get

$$(7.3) \quad \int_{B_{R/2}} \mathcal{G}_0(\nabla U)^\gamma dx \leq \Gamma_1 + C\Gamma_2 \liminf_{k \rightarrow \infty} \int_{B_R} \mathcal{G}_0(\nabla u^{\varepsilon_k}) dx.$$

By Lemma 2.6, the functions $u^{\varepsilon_k}_{x_i}$ converge to U_{x_i} in $L^{p_i}(B)$. Hence, the continuity of the map $v \in L^{p_i}(B) \mapsto |v|^{p_i} \in L^1(B)$ implies that

$$(7.4) \quad \lim_{\varepsilon \rightarrow 0} \|\mathcal{G}_0(\nabla u^\varepsilon) - \mathcal{G}_0(\nabla U)\|_{L^1(B)} = 0.$$

By using this result in (7.3), we obtain

$$\int_{B_{R/2}} \mathcal{G}_0(\nabla U)^\gamma dx \leq \Gamma_1 + C\Gamma_2 \int_{B_R} \mathcal{G}_0(\nabla U) dx.$$

This concludes the proof, up to renaming the constant Γ_2 .

⁶The upper bound simply follows from (2.16), with standard algebraic manipulations.

7.2. Proof of Theorem L

In this section, we assume that $1 < p_1 \leq \dots \leq p_N \leq 2$ and $f \in L^{\gamma}_{\text{loc}}(\Omega)$ for some $\gamma > N$. In particular, $\gamma \geq 2$ and thus we can rely on Proposition 4.1.

We introduce the ball $B_R \Subset B$ as before. By Proposition 5.1 applied with $B_{R/4}$ and $B_{R/2}$, for every $\varepsilon \in (0, \varepsilon_0)$, we have

$$\begin{aligned} \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^\infty(B_{R/4})} &\leq C \left[\left(\frac{4}{R}\right)^{N\gamma/(\gamma-N)} \left(\int_{B_{R/2}} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\gamma dx \right)^{N/(\gamma-N)} \right. \\ &\quad \left. + \|f^\varepsilon\|_{L^\gamma(B_R)}^{N\gamma/(\gamma-N)} \right] \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^1(B_R)}, \end{aligned}$$

for some $C = C(N, p_N, p_1, \gamma) > 0$. On the right-hand side, we can apply (7.1), in order to estimate the term containing $\mathcal{G}_\varepsilon^\gamma$. This yields

$$\begin{aligned} \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^\infty(B_{R/4})} &\leq C \left[\left(\frac{4}{R}\right)^{N\gamma/(\gamma-N)} \left(\Gamma_1 + \Gamma_2 \int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right)^{N/(\gamma-N)} \right. \\ &\quad \left. + \|f^\varepsilon\|_{L^\gamma(B_R)}^{N\gamma/(\gamma-N)} \right] \|\mathcal{G}_\varepsilon(\nabla u^\varepsilon)\|_{L^1(B_R)}. \end{aligned}$$

We now take the same infinitesimal sequence $\{\varepsilon_k\}_{k \in \mathbb{N}}$ as in the proof of Proposition 1.1. By using again (7.2), the lower semicontinuity of the L^∞ norm with respect to almost everywhere convergence, the property (7.4) and the fact that f^{ε_k} is defined from f by convolution with a smooth kernel, the limit as k goes to ∞ gives

$$\begin{aligned} \|\mathcal{G}_0(\nabla U)\|_{L^\infty(B_{R/4})} &\leq C \left[\left(\frac{4}{R}\right)^{N\gamma/(\gamma-N)} \left(\Gamma_1 + \Gamma_2 \int_{B_R} \mathcal{G}_0(\nabla U) dx \right)^{N/(\gamma-N)} \right. \\ &\quad \left. + \|f\|_{L^\gamma(B_R)}^{N\gamma/(\gamma-N)} \right] \|\mathcal{G}_0(\nabla U)\|_{L^1(B_R)}, \end{aligned}$$

possibly for a different $C = C(N, p_N, p_1, \gamma) > 0$. This completes the proof.

7.3. Proof of Theorem S

We assume that $1 < p_1 \leq \dots \leq p_N \leq 2$ and f satisfies (1.3). We can suppose that $p_1 < 2$, otherwise the result is well known. We set, for notational simplicity,

$$\gamma = 1 + \frac{2}{p_1}.$$

Consider the ball $B_R \Subset B$, and let $\eta \in C_0^\infty(B_{R/2})$ be such that

$$\eta \equiv 1 \text{ on } B_{R/4}, \quad 0 \leq \eta \leq 1, \quad \text{and} \quad |\nabla \eta|^2 + |D^2 \eta| \leq \frac{C_0}{R^2},$$

for some C_0 which depends only on N . The choice of γ entails the following estimates:

$$1 + 2 \left(\frac{1}{p_1} - \frac{1}{p_N} \right) < \gamma \quad \text{and} \quad \frac{2 - p_1}{p_1} \frac{\gamma}{\gamma - 2} = \gamma.$$

Then, as a consequence of (6.2), we have

$$\begin{aligned} \sum_{i=1}^N \int_{B_{R/4}} |\nabla \mathcal{V}_{i,\varepsilon}|^2 dx &\leq \frac{C}{R^2} \int_{B_{R/2}} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\gamma dx \\ &\quad + C \left(\int_{B_{R/2}} \mathcal{G}_\varepsilon(\nabla u^\varepsilon)^\gamma dx \right)^{(\gamma-2)/\gamma} \|f^\varepsilon\|_{L^\gamma(B_R)}^2, \end{aligned}$$

where we also used that $\mathcal{G}_\varepsilon \geq 1$, by definition. We now rely again on (7.1), to estimate the terms containing $\mathcal{G}_\varepsilon^\gamma$. This estimate and Young's inequality with exponents $\gamma/2$ and $\gamma/(\gamma-2)$ give

$$(7.5) \quad \sum_{i=1}^N \int_{B_{R/4}} |\nabla \mathcal{V}_{i,\varepsilon}|^2 dx \leq \frac{C}{R^2} \left(\Gamma_1 + \Gamma_2 \int_{B_R} \mathcal{G}_\varepsilon(\nabla u^\varepsilon) dx \right) + CR^{\gamma-2} \|f^\varepsilon\|_{L^\gamma(B_R)}^\gamma,$$

possibly for a different constant $C = C(N, p_N, p_1) > 0$. From this estimate, we deduce that the family $\nabla \mathcal{V}_{i,\varepsilon}$ is uniformly bounded in $L^2(B_{R/4})$. Moreover, by (2.2) we have

$$(7.6) \quad \sqrt{g''_{i,\varepsilon}(t)} \leq (\varepsilon + t^2)^{(p_i-2)/4} \leq |t|^{(p_i-2)/2}, \quad \text{for } t \neq 0.$$

Thus, by recalling the definition of $\mathcal{V}_{i,\varepsilon}$, we get

$$\int_{B_{R/4}} |\mathcal{V}_{i,\varepsilon}|^2 dx = \int_{B_{R/4}} |V_{i,\varepsilon}(u_{x_i}^\varepsilon)|^2 dx \leq \left(\frac{2}{p_i} \right)^2 \int_{B_{R/4}} |u_{x_i}^\varepsilon|^{p_i} dx,$$

and the latter is uniformly bounded, thanks to Lemma 2.5.

Thus, by taking the same infinitesimal sequence $\{\varepsilon_k\}_{k \geq 1}$ as in the proof of Proposition 1.1, we have obtained that $\{\mathcal{V}_{i,\varepsilon_k}\}_{k \in \mathbb{N}}$ is a bounded sequence in $W^{1,2}(B_{R/4})$. By appealing to the Rellich–Kondrašov theorem, we can infer its convergence to a function $\mathcal{V}_i \in W^{1,2}(B_{R/4})$, weakly in $W^{1,2}(B_{R/4})$ and strongly in $L^2(B_{R/4})$ (up to a subsequence). By the lower semicontinuity of the L^2 norm, (7.2) and (7.4), we get from (7.5) that

$$\sum_{i=1}^N \int_{B_{R/4}} |\nabla \mathcal{V}_i|^2 dx \leq \frac{C}{R^2} \left(\Gamma_1 + \Gamma_2 \int_{B_R} \mathcal{G}_0(\nabla U) dx \right) + CR^{\gamma-2} \|f\|_{L^\gamma(B_R)}^\gamma,$$

possibly for a different $C = C(N, p_N, p_1) > 0$. We claim that for every $1 \leq i \leq N$ and for almost every $x \in B_{R/4}$, we have

$$(7.7) \quad \mathcal{V}_i(x) = \frac{2}{p_i} \sqrt{p_i - 1} |U_{x_i}(x)|^{(p_i-2)/2} U_{x_i}(x).$$

Indeed, take $x \in B_{R/4}$ such that $u_{x_i}^{\varepsilon_k}(x)$ converges to $U_{x_i}(x)$, and such that $|U_{x_i}(x)| < +\infty$. Observe that the collection of these points has full measure in $B_{R/4}$. We then set

$$M_i(x) = \sup_{k \in \mathbb{N}} |u_{x_i}^{\varepsilon_k}(x)|,$$

which is finite, by construction. Then, for every $k \geq 0$ we have

$$\begin{aligned}
& \left| \mathcal{V}_{i,\varepsilon_k}(x) - \frac{2}{p_i} \sqrt{p_i - 1} |U_{x_i}(x)|^{(p_i-2)/2} U_{x_i}(x) \right| \\
&= \left| V_{i,\varepsilon_k}(u_{x_i}^{\varepsilon_k}(x)) - \frac{2}{p_i} \sqrt{p_i - 1} |U_{x_i}(x)|^{(p_i-2)/2} U_{x_i}(x) \right| \\
&\leq \left| \int_0^{u_{x_i}^{\varepsilon_k}(x)} \left(\sqrt{g_{i,\varepsilon_k}''(\tau)} - \sqrt{(p_i - 1)|\tau|^{p_i-2}} \right) d\tau \right| \\
&\quad + \frac{2}{p_i} \sqrt{p_i - 1} \left| |u_{x_i}^{\varepsilon_k}(x)|^{(p_i-2)/2} u_{x_i}^{\varepsilon_k}(x) - |U_{x_i}(x)|^{(p_i-2)/2} U_{x_i}(x) \right| \\
&\leq \int_0^{M_i(x)} \left| \sqrt{g_{i,\varepsilon_k}''(\tau)} - \sqrt{(p_i - 1)|\tau|^{p_i-2}} \right| d\tau \\
&\quad + \frac{2}{p_i} \sqrt{p_i - 1} \left| |u_{x_i}^{\varepsilon_k}(x)|^{(p_i-2)/2} u_{x_i}^{\varepsilon_k}(x) - |U_{x_i}(x)|^{(p_i-2)/2} U_{x_i}(x) \right|.
\end{aligned}$$

Thanks to (7.6), one can apply the dominated convergence to conclude that the first term in the right-hand side converges to 0 when k goes to $+\infty$. By also using that $u_{x_i}^{\varepsilon_k}(x)$ converges to $U_{x_i}(x)$, we finally get (7.7).

By using the chain rule in Sobolev spaces, we also obtain that $U_{x_i} \in W^{1,p_i}(B_{R/4})$ and satisfies the estimate claimed in the statement of Theorem S. The proof is complete.

A. A weak maximum principle

Let $G: \mathbb{R}^N \rightarrow [0, +\infty)$ be a convex function such that $G(z) > G(0)$ for every $z \in \mathbb{R}^N \setminus \{0\}$. Let $\zeta: \mathbb{R} \rightarrow \mathbb{R}$ be a Lipschitz function, with the following property: there exists $M > 0$ such that

$$\zeta(t) = \begin{cases} M, & \text{if } t \geq M, \\ -M, & \text{if } t \leq -M. \end{cases}$$

Given a ball $B \subset \mathbb{R}^N$, $f \in L^1(B)$ and $U \in W^{1,1}(B) \cap L^\infty(B)$ such that

$$\|U\|_{L^\infty(B)} \leq M \quad \text{and} \quad \int_B G(\nabla U) dx < +\infty,$$

we consider the functional

$$\mathcal{F}(v) = \int_B [G(\nabla v) + f\zeta(v)] dx, \quad \text{for every } v \in U + W_0^{1,1}(B).$$

Lemma A.1. *If u is a minimum of \mathcal{F} , then $\|u\|_{L^\infty(B)} \leq M$.*

Proof. We want to test the minimality of u against the truncated function

$$v := \max\{-M, \min\{u, M\}\}.$$

By construction, we still have $v \in U + W_0^{1,1}(B)$, and by minimality of u , we get

$$\mathcal{F}(u) \leq \mathcal{F}(v).$$

By the properties of ζ , we have

$$\begin{aligned} \mathcal{F}(v) &= \int_{\{|u| \leq M\}} [G(\nabla u) + f\zeta(u)] dx + \int_{\{|u| > M\}} G(0) dx \\ &\quad + \int_{\{u > M\}} f\zeta(M) dx + \int_{\{u < -M\}} f\zeta(-M) dx. \end{aligned}$$

Hence, using that $\zeta(u) = \zeta(M)$ when $u \geq M$ and $\zeta(u) = \zeta(-M)$ when $u \leq -M$, by comparing the last two displays, we get

$$\int_{\{|u| > M\}} G(\nabla u) dx \leq \int_{\{|u| > M\}} G(0) dx, \quad \text{that is,} \quad \int_{\{|u| > M\}} [G(\nabla u) - G(0)] dx \leq 0.$$

Thanks to the properties of G , we deduce that $\nabla u = 0$ almost everywhere on the set $\{|u| > M\}$. It follows that $\nabla u = \nabla v$ almost everywhere, and since $u = v = U$ on ∂B , this implies that $u = v$ almost everywhere in B . In particular, $|u| \leq M$ almost everywhere in B . ■

Acknowledgements. This work has been finalized during a staying of P.B. and L.B. at the *Institute of Applied Mathematics and Mechanics* of the University of Warsaw, in July 2022. Iwona Chlebicka and Anna Zatorska-Goldstein are gratefully acknowledged for their kind invitation and the nice working atmosphere provided during the whole staying. C.L. is a member of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

Funding. This work was partially supported by the projects FAR 2019 and FAR 2020 of the University of Ferrara.

References

- [1] Baldelli L., Ciani S., Skrypnik I. and Vespri V.: [A note on the point-wise behaviour of bounded solutions for a non-standard elliptic operator](#). To appear on *Discrete Contin. Dyn. Syst. Ser. S*, doi: 10.3934/dcdss.2022143.
- [2] Baroni, P., Colombo, M. and Mingione, G.: [Regularity for general functionals with double phase](#). *Calc. Var. Partial Differential Equations* **57** (2018), no. 2, article no. 62, 48 pp.
- [3] Baroni, P., Di Castro, A. and Palatucci, G.: [Intrinsic geometry and De Giorgi classes for certain anisotropic problems](#). *Discrete Contin. Dyn. Syst. Ser. S* **10** (2017), no. 4, 647–659.
- [4] Beck, L. and Mingione, G.: [Lipschitz bounds and nonuniform ellipticity](#). *Comm. Pure Appl. Math.* **73** (2020), no. 5, 944–1034.
- [5] Bhattacharya, T. and Leonetti, F.: Some remarks on the regularity of minimizers of integrals with anisotropic growth. *Comment. Math. Univ. Carolin.* **34** (1993), no. 4, 597–611.
- [6] Bhattacharya, T. and Leonetti, F.: [On improved regularity of weak solutions of some degenerate, anisotropic elliptic systems](#). *Ann. Mat. Pura Appl. (4)* **170** (1996), 241–255.
- [7] Bildhauer, M., Fuchs, M. and Zhong, X.: [A regularity theory for scalar local minimizers of splitting-type variational integrals](#). *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **6** (2007), no. 3, 385–404.

- [8] Bousquet, P.: Another look to the orthotropic functional in the plane. In *Bruno Pini mathematical analysis Seminar 2020*, pp. 1–29. Bruno Pini Math. Anal. Semin. 11, Univ. Bologna, Alma Mater Stud., Bologna, 2021.
- [9] Bousquet, P. and Brasco, L.: C^1 regularity of orthotropic p -harmonic functions in the plane. *Anal. PDE* **11** (2018), no. 4, 813–854.
- [10] Bousquet, P. and Brasco, L.: Lipschitz regularity for orthotropic functionals with nonstandard growth conditions. *Rev. Mat. Iberoam.* **36** (2020), no. 7, 1989–2032.
- [11] Bousquet, P., Brasco, L. and Julin, V.: Lipschitz regularity for local minimizers of some widely degenerate problems. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **16** (2016), no. 4, 1235–1274.
- [12] Bousquet, P., Brasco, L., Leone, C. and Verde, A.: On the Lipschitz character of orthotropic p -harmonic functions. *Calc. Var. Partial Differential Equations* **57** (2018), no. 3, article no. 88, 33 pp.
- [13] Bousquet, P., Mariconda, C. and Treu, G.: On the Lavrentiev phenomenon for multiple integral scalar variational problems. *J. Funct. Anal.* **266** (2014), no. 9, 5921–5954.
- [14] Brasco, L., Leone, C., Pisante, G. and Verde, A.: Sobolev and Lipschitz regularity for local minimizers of widely degenerate anisotropic functionals. *Nonlinear Anal.* **153** (2017), 169–199.
- [15] Canale, A., D’Ottavio, A., Leonetti, F. and Longobardi, M.: Differentiability for bounded minimizers of some anisotropic integrals. *J. Math. Anal. Appl.* **253** (2001), no. 2, 640–650.
- [16] Carozza, M., Kristensen, J. and Passarelli di Napoli, A.: Higher differentiability of minimizers of convex variational integrals. *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **28** (2011), no. 3, 395–411.
- [17] Cavaliere, P., D’Ottavio, A., Leonetti, F. and Longobardi, M.: Differentiability for minimizers of anisotropic integrals. *Comment. Math. Univ. Carolin.* **39** (1998), no. 4, 685–696.
- [18] Chlebicka, I., Gwiazda, P., Świerczewska-Gwiazda, A. and Wróblewska-Kamińska, A.: *Partial differential equations in anisotropic Musielak–Orlicz spaces*. Springer Monographs in Mathematics, Springer, Cham, 2021.
- [19] Choe, H. J.: Interior behaviour of minimizers for certain functionals with nonstandard growth. *Nonlinear Anal.* **19** (1992), no. 10, 933–945.
- [20] Cianchi, A.: Local boundedness of minimizers of anisotropic functionals. *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **17** (2000), no. 2, 147–168.
- [21] Cianchi, A.: Symmetrization in anisotropic elliptic problems. *Comm. Partial Differential Equations* **32** (2007), no. 4–6, 693–717.
- [22] Clop, A., Gentile, A. and Passarelli di Napoli, A.: Higher differentiability results for solutions to a class of non-homogeneous elliptic problems under sub-quadratic growth conditions. *Bull. Math. Sci.* **13** (2023), no. 2, article no. 2350008, 55 pp.
- [23] Colombo, M. and Mingione, G.: Bounded minimisers of double phase variational integrals. *Arch. Ration. Mech. Anal.* **218** (2015), no. 1, 219–273.
- [24] Colombo, M. and Mingione, G.: Regularity for double phase variational problems. *Arch. Ration. Mech. Anal.* **215** (2015), no. 2, 443–496.
- [25] Cupini, G., Marcellini, P. and Mascolo, E.: Regularity under sharp anisotropic general growth conditions. *Discrete Contin. Dyn. Syst. Ser. B* **11** (2009), no. 1, 66–86.
- [26] Cupini, G., Marcellini, P. and Mascolo, E.: Local boundedness of solutions to quasilinear elliptic systems. *Manuscripta Math.* **137** (2012), no. 3–4, 287–315.

- [27] Cupini, G., Marcellini, P. and Mascolo, E.: [Local boundedness of minimizers with limit growth conditions](#). *J. Optim. Theory Appl.* **166** (2015), no. 1, 1–22.
- [28] Cupini, G., Marcellini, P. and Mascolo, E.: [Regularity of minimizers under limit growth conditions](#). *Nonlinear Anal.* **153** (2017), 294–310.
- [29] De Silva, D. and Savin, O.: [Minimizers of convex functionals arising in random surfaces](#). *Duke Math. J.* **151** (2010), no. 3, 487–532.
- [30] de Thélin, F.: [Local regularity properties for the solutions of a nonlinear partial differential equation](#). *Nonlinear Anal.* **6** (1982), no. 8, 839–844.
- [31] Demengel, F.: [Lipschitz interior regularity for the viscosity and weak solutions of the pseudo \$p\$ -Laplacian equation](#). *Adv. Differential Equations* **21** (2016), no. 3-4, 373–400.
- [32] DiBenedetto, E.: [\$C^{1+\alpha}\$ local regularity of weak solutions of degenerate elliptic equations](#). *Nonlinear Anal.* **7** (1983), no. 8, 827–850.
- [33] Eleuteri, M., Marcellini, P., Mascolo, E. and Perrotta, S.: [Local Lipschitz continuity for energy integrals with slow growth](#). *Ann. Mat. Pura Appl. (4)* **201** (2022), no. 3, 1005–1032.
- [34] Esposito, L., Leonetti, F. and Mingione, G.: [Regularity for minimizers of functionals with \$p\$ - \$q\$ growth](#). *NoDEA Nonlinear Differential Equations Appl.* **6** (1999), no. 2, 133–148.
- [35] Esposito, L., Leonetti, F. and Mingione, G.: [Sharp regularity for functionals with \$\(p, q\)\$ growth](#). *J. Differential Equations* **204** (2004), no. 1, 5–55.
- [36] Fonseca, I. and Fusco, N.: Regularity results for anisotropic image segmentation models. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **24** (1997), no. 3, 463–499.
- [37] Fusco, N. and Sbordone, C.: [Some remarks on the regularity of minima of anisotropic integrals](#). *Comm. Partial Differential Equations* **18** (1993), no. 1-2, 153–167.
- [38] Giaquinta, M.: [Growth conditions and regularity, a counterexample](#). *Manuscripta Math.* **59** (1987), no. 2, 245–248.
- [39] Giusti, E.: *Direct methods in the calculus of variations*. World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
- [40] Hong, M. C.: Some remarks on the minimizers of variational integrals with nonstandard growth conditions. *Boll. Un. Mat. Ital. A (7)* **6** (1992), no. 1, 91–101.
- [41] Kolodñ, Ī. M.: The boundedness of generalized solutions of parabolic differential equations. *Vestnik Moskov. Univ. Ser. I Mat. Meh.* **26** (1971), no. 5, 25–31.
- [42] Korolëv, A. G.: [On the boundedness of generalized solutions of elliptic differential equations](#). *Russ. Math. Surv.* **38** (1983), no. 2, 186–187.
- [43] Leonetti, F.: [Higher integrability for minimizers of integral functionals with nonstandard growth](#). *J. Differential Equations* **112** (1994), no. 2, 308–324.
- [44] Leonetti, F. and Mascolo, E.: [Local boundedness for vector valued minimizers of anisotropic functionals](#). *Z. Anal. Anwend.* **31** (2012), no. 3, 357–378.
- [45] Lieb E. H., Loss M.: *Analysis*. Second edition. Graduate Studies in Mathematics 14, American Mathematical Society, Providence, RI, 2001.
- [46] Lieberman, G. M.: [Gradient estimates for anisotropic elliptic equations](#). *Adv. Differential Equations* **10** (2005), no. 7, 767–812.
- [47] Lindqvist, P. and Ricciotti, D.: [Regularity for an anisotropic equation in the plane](#). *Nonlinear Anal.* **177** (2018), 628–636.

- [48] Liskevich, V. and Skrypnik, I. I.: Hölder continuity of solutions to an anisotropic elliptic equation. *Nonlinear Anal.* **71** (2009), no. 5-6, 1699–1708.
- [49] Marcellini P.: Un exemple de solution discontinue d'un problème variationnel dans le cas scalaire. Preprint no. 11 Ist. Mat. Univ. Firenze, 1987. Available at <http://web.math.unifi.it/users/marcellin/lavori/reprints>.
- [50] Marcellini, P.: Regularity of minimizers of integrals of the calculus of variations with non-standard growth conditions. *Arch. Rational Mech. Anal.* **105** (1989), no. 3, 267–284.
- [51] Marcellini, P.: Regularity and existence of solutions of elliptic equations with p, q -growth conditions. *J. Differential Equations* **90** (1991), no. 1, 1–30.
- [52] Naumann, J.: Interior integral estimates on weak solutions of certain degenerate elliptic systems. *Ann. Mat. Pura Appl. (4)* **156** (1990), 113–125.
- [53] Pagano, A.: Higher integrability for minimizers of variational integrals with nonstandard growth. *Ann. Univ. Ferrara Sez. VII (N.S.)* **39** (1993), 1–17 (1995).
- [54] Ricciotti, D.: Regularity of the derivatives of p -orthotropic functions in the plane for $1 < p < 2$. *Ann. Acad. Sci. Fenn. Math.* **44** (2019), no. 2, 1093–1099.
- [55] Santambrogio, F.: Regularity via duality in calculus of variations and degenerate elliptic PDEs. *J. Math. Anal. Appl.* **457** (2018), no. 2, 1649–1674.
- [56] Stampacchia, G.: On some regular multiple integral problems in the calculus of variations. *Comm. Pure Appl. Math.* **16** (1963), 383–421.
- [57] Ural'tseva, N. N. and Urdaletova, A. B.: The boundedness of the gradients of generalized solutions of degenerate nonuniformly elliptic quasilinear equations. *Vest. Leningr. Univ. Mat.* **16** (1984), 263–270.

Received November 21, 2022; revised August 22, 2023. Published online November 17, 2023.

Pierre Bousquet

Institut de Mathématiques de Toulouse, UMR5219, Université de Toulouse;
CNRS, UPS, 31062 Toulouse Cedex 9, France;
pierre.bousquet@math.univ-toulouse.fr

Lorenzo Brasco

Dipartimento di Matematica e Informatica, Università degli Studi di Ferrara,
Via Machiavelli 30, 44121 Ferrara, Italy;
lorenzo.brasco@unife.it

Chiara Leone

Dipartimento di Matematica “R. Caccioppoli”, Università degli Studi di Napoli Federico II,
Via Cinthia, Complesso Universitario di Monte S. Angelo, 80126 Napoli, Italy;
chiara.leone@unina.it