

On some permutable products of supersoluble groups

Manuel J. Alejandro, A. Ballester-Bolinches,
John Cossey and M. C. Pedraza-Aguilera

Abstract

It is well known that a group $G = AB$ which is the product of two supersoluble subgroups A and B is not supersoluble in general. Under suitable permutability conditions on A and B , we show that for any minimal normal subgroup N both AN and BN are supersoluble. We then exploit this to establish some sufficient conditions for G to be supersoluble.

1. Introduction

All groups considered in this paper will be finite. It is well known that, even for a saturated formation \mathcal{F} , a group that is the product of two subgroups in \mathcal{F} need not to be in \mathcal{F} . The structure of such products and conditions which ensure that the product is in \mathcal{F} have been widely studied for the classes \mathcal{N} of nilpotent groups and \mathcal{U} of supersoluble groups.

The behaviour of minimal normal subgroups of factorized groups has been an important source of information about their structure. Our interest is in extending the following theorem of Stonehewer [12]:

Let G be a finite group which can be written as the product $G = AB$ of two nilpotent subgroups A and B . If N is a minimal normal subgroup of G , then either AN or BN is nilpotent.

Unfortunately, this result is not true if we replace nilpotent by supersoluble in the statement ($PSL(2, 7)$ can be written as the product of two supersoluble subgroups). One of the main purposes of this paper is to

2000 Mathematics Subject Classification: 20D10, 20D35, 20D40.

Keywords: finite groups, products, subnormality, supersolubility.

find conditions which allow us to establish a supersoluble version of Stonehewer's Theorem. We then exploit these results to obtain the supersolubility of some products of supersoluble groups.

Much recent work on products of groups and supersolubility has focussed on products when the factors satisfy extra permutability conditions. The first step in this direction was taken by Baer [4].

Let $G = AB$ be the product of the supersoluble normal subgroups A and B of G . If the derived subgroup G' of G is nilpotent, then G is supersoluble.

Many generalizations of this theorem have been obtained. The aim of most of them has been to weaken the normality hypotheses, replacing them by permutability conditions instead. Following Carocca [7], we will say that:

A group G is said to be the totally permutable product (t.p.p.) of the subgroups A and B if $G = AB$ and every subgroup of A permutes with every subgroup of B .

The product $G = AB$ is said to be mutually permutable (m.p.p.) if A permutes with every subgroup of B and B permutes with every subgroup of A .

Every totally permutable product is clearly mutually permutable, but the converse does not hold. It is true, however, for mutually permutable products $G = AB$ such that $A \cap B = 1$ [7, Proposition 3.5].

Asaad and Shaalan obtained in [3] results about totally and mutually permutable products of supersoluble groups. For totally permutable products, they obtained the following:

If G is the totally permutable product of two supersoluble subgroups A and B , then G is supersoluble.

Moreover they proved that this result is not valid for mutually permutable products, but the following generalization [3, Theorem 3.1] of Baer's result is true:

If G is the mutually permutable product of two supersoluble subgroups A and B , and the derived subgroup G' of G is nilpotent, then G is supersoluble.

Finally, this last result remains valid if the nilpotency of G' is replaced by the nilpotency of one of the factors [3, Theorem 3.2].

In the last part of this paper some generalizations of these results will be naturally obtained from our study.

In recent years, some weaker versions of the concepts of totally and mutually permutable products have been introduced. Permutability of each factor with some specific families of subgroups of the other has often been analyzed. In this context, we study the structure of mutually sn -permutable products:

Definition Let A and B be two subgroups of a group G such that $G = AB$.

- a) We say that $G = AB$ is the totally sn -permutable product of A and B if every subnormal subgroup of A permutes with every subnormal subgroup of B .
- b) We say that $G = AB$ is the mutually sn -permutable product of A and B if A permutes with every subnormal subgroup of B and B permutes with every subnormal subgroup of A .

Carocca showed [8, Theorem 6] that every group which can be factorized as a mutually sn -permutable product of two soluble groups is soluble as well. In the study of factorized groups in which each factor permutes with a specific family of subgroups of the other, the embedding of the intersection of the factors plays a very important role. In Section 2 we analyze this embedding and apply it to obtain an alternative proof of Carocca's result.

Since the purpose of this paper is to investigate the structure of a group $G = AB$ which is either a totally sn -permutable product or a mutually sn -permutable product of two supersoluble subgroups A and B , we shall assume in the sequel that all groups are not only finite but also soluble, except of course in the statement and proof of Carocca's Theorem.

We obtain a supersoluble version of Stonehewer's Theorem by assuming that the product is mutually sn -permutable. In fact, we have the following stronger result:

Theorem A Let $G = AB$ be the mutually sn -permutable product of the supersoluble subgroups A and B . If N is a minimal normal subgroup of G , then both AN and BN are supersoluble.

This theorem turns out to be useful in the study of the supersolubility of sn -permutable products of supersoluble groups. It also allowed us to prove an important structural theorem for mutually permutable products of supersoluble groups (see [1]).

It is not true in general that a group which is the totally sn -permutable product of two supersoluble subgroups is supersoluble, as we shall see in the last section of this paper. However this result of Beidleman, Galoppo, Heineken and Manfredino ([6]) gives a sufficient condition for this property to be satisfied:

Theorem 1 Let $G = AB$ be a finite group which is the totally sn -permutable product of the subgroups A and B . If A is supersoluble and B is nilpotent, then G is supersoluble.

Note that the proof in [6] does not require the finiteness of G . On the other hand, this last Theorem confirms that the result of Asaad and Shaalan holds in the case of a totally sn -permutable product (instead of a totally

permutable product) if we assume the nilpotency of one of the factors. We will see that a proof of this result in the finite case follows easily from Theorem A, as does the following Theorem:

Theorem B. *Let $G = AB$ be the mutually sn-permutable product of the subgroups A and B , where A is supersoluble and B is nilpotent. If B permutes with each Sylow subgroup of A , then the group G is supersoluble.*

The hypothesis on B in the above theorem is essential in order to get supersolubility as we will show by means of an example in the last section of this paper.

Moreover, we prove that the second theorem of Asaad and Shaalan remains true if G is a mutually sn-permutable product without any additional requirement.

Theorem C. *Let $G = AB$ be the mutually sn-permutable product of the supersoluble subgroups A and B . If the derived subgroup G' of G is nilpotent, then G is supersoluble.*

Notice that the group in Theorem C is nilpotent-by-abelian. Hence a natural question is whether or not a metanilpotent group which is the mutually sn-permutable product of two supersoluble subgroups is supersoluble. The second example in Section 5 answers that question negatively, but additional assumptions allow us to get supersolubility:

Theorem D. *Let $G = AB$ be the mutually sn-permutable product of the supersoluble subgroups A and B . Assume that G is metanilpotent. If*

$$(|A/A^N|, |B/B^N|) = 1$$

then G is supersoluble.

2. Some properties of mutually sn-permutable products

Lemma 1 *Let $G = AB$ be a mutually sn-permutable product. Let $I = A \cap B$ and let L be a subnormal subgroup of A such that $I \leq L$. Then LB is again a mutually sn-permutable product.*

Proof. It is obvious that B permutes with every subnormal subgroup of L . On the other hand, if we take R a subnormal subgroup of B , then we have that $LR = LIR = L(A \cap B)R = (A \cap LB)R = AR \cap LB = RA \cap BL = R(A \cap BL) = R(A \cap B)L = RL$. Therefore, we have that LB is a mutually sn-permutable product. ■

Corollary 1 *If $G = AB$ is a mutually sn-permutable product such that $A \cap B = 1$, then $G = AB$ is a totally sn-permutable product.*

Proof. This is a straightforward consequence of the previous lemma. ■

Lemma 2 *Let $G = AB$ be a mutually sn -permutable product of the subgroups A and B . If N is a minimal normal subgroup of G , then either $A \cap N = B \cap N = 1$ or $N = (A \cap N)(B \cap N)$. In this second situation, the product $(A \cap N)(B \cap N)$ is mutually sn -permutable.*

Proof. Clearly $A \cap N$ is a normal subgroup of A and hence $X = (A \cap N)B$ is a subgroup of G such that $N \cap X = N \cap (A \cap N)B = (A \cap N)(B \cap N)$. Now since $N \cap X$ is a normal subgroup of X , B normalizes $N \cap X = (A \cap N)(B \cap N)$.

Arguing similarly, $Y = A(B \cap N)$ is a subgroup of G such that $Y \cap N = A(B \cap N) \cap N = (A \cap N)(B \cap N)$

As before, Y (and hence A) normalizes $Y \cap N = (A \cap N)(B \cap N)$. We can conclude that $(A \cap N)(B \cap N)$ is a normal subgroup of G , and the minimality of N implies that either $A \cap N = B \cap N = 1$ or $N = (A \cap N)(B \cap N)$.

If this last situation holds, then given any subnormal subgroup A_0 of $A \cap N$ we have that A_0 is subnormal in A as well and thus $A_0B \leq G$. But then $A_0(B \cap N) = A_0B \cap N \leq G$. A similar argument shows that $A \cap N$ permutes with every subnormal subgroup of $B \cap N$, and consequently the product $(A \cap N)(B \cap N)$ is mutually sn -permutable. ■

Following the notation used in [9, III.4 and IV.5], we define:

Definition. *Let G be a group, and let \mathcal{X} be a class of groups. We say that a maximal subgroup M is \mathcal{X} -normal in G if $G/\text{Core}_G(M) \in \mathcal{X}$.*

If \mathcal{F} is a saturated formation, a subgroup H of G is said to be \mathcal{F} -subnormal in G if $H = G$ or there exists a chain of subgroups

$$H = H_0 < H_1 < \dots < H_n = G$$

such that H_{i-1} is a maximal \mathcal{F} -normal subgroup of H_i , for each $i = 1, 2, \dots, n$.

\mathcal{U} -subnormality will appear in our context. An interesting connection which will link this embedding property with subnormal permutability can be stated in the following terms:

Theorem 2 *Let G be a group. If H is a subgroup of G such that H permutes with every subnormal subgroup of G , then H is \mathcal{U} -subnormal in G .*

Proof. We assume that the theorem is false and let G be a minimal counterexample. There exists a subgroup H of G which permutes with every subnormal subgroup of G but is not \mathcal{U} -subnormal in G . Let N be a minimal normal subgroup of G . Clearly HN/N permutes with every subnormal subgroup of G/N . The minimality of G yields that HN/N is \mathcal{U} -subnormal in G/N and therefore HN is \mathcal{U} -subnormal in G as a consequence of [10, Lemma 1.1].

Since N is a minimal normal subgroup of the soluble group G , then N must be an elementary abelian p -group, for some prime p . But H permutes with every subnormal subgroup of G and consequently also with every subgroup of N . In particular, there exists a series

$$H < HN_1 < \cdots < HN_r < HN$$

such that $|HN : HN_r| = |HN_{j+1} : HN_j| = |HN_1 : H| = p$, for each $j = 1, \dots, r - 1$. This clearly implies that H is a \mathcal{U} -subnormal subgroup of HN . Hence we have that H is \mathcal{U} -subnormal in G , a contradiction. ■

As a consequence of the above Theorem and [5, Lemma 3.1], we have that if H permutes with every subnormal subgroup of a group G , then the supersoluble residual of H is subnormal in G .

Theorem 3 [8, Theorem 6] *Let $G = AB$ be the mutually sn -permutable product of the subgroups A and B of G . If both A and B are soluble, then G is also soluble.*

Proof. Let us assume that the theorem is false and choose a minimal counterexample, G say. Arguing by induction, we know that G must have a unique minimal normal subgroup, which is non-abelian, N say. Using Lemma 2, we know that either $N = (A \cap N)(B \cap N)$ or $A \cap N = B \cap N = 1$. On the other hand, applying Theorem 2, $A \cap B$ is an \mathcal{U} -subnormal subgroup of A and B .

Take now a minimal normal subgroup N_1 of A and a minimal normal subgroup N_2 of B , respectively. Then it is clear that $N \leq N_1^G \cap N_2^G \leq N_1 B \cap A N_2 = [N_1(A \cap B)]N_2$. Moreover we can suppose that $\text{Core}_A(A \cap B) = \text{Core}_B(A \cap B) = 1$, otherwise we would take N_1 or N_2 contained in $A \cap B$ and N would be contained in A or B , a contradiction.

Let us assume firstly that $N = (A \cap N)(B \cap N)$. If $N \neq G$, then the solubility of G/N follows from the minimality of G . But N is again a mutually sn -permutable product of two soluble subgroups by Lemma 2, and therefore N itself is soluble. It follows that G is soluble, a contradiction. Consequently we can assume that $N = G$ and hence G must be a simple group. Furthermore by the above arguments $N \leq N_1(A \cap B)N_2$. Consequently $G = N_1(A \cap B)N_2$ and hence $A = N_1(A \cap B)$. Notice that N_1 is a minimal normal subgroup of A . Hence $A \cap B$ is an \mathcal{U} -subnormal maximal subgroup of A with $\text{Core}_A(A \cap B) = 1$. Therefore A is supersoluble and N_1 is a cyclic group of prime order, p say. Arguing in an analogous way we can assume also that B is supersoluble and N_2 is a cyclic group of prime order, q say. Since $\text{Core}_G(A) = \text{Core}_G(B) = 1$, we have that G is isomorphic to subgroups of the symmetric groups of degree p and q respectively.

Consequently $p = q$. Moreover, $|N_1N_2| = p^2$, that is, N_1N_2 is an abelian subgroup of G . Recall that $G = N_1(A \cap B)N_2$. This means that N_1 is a normal subgroup of G , a contradiction.

Therefore we may assume for the rest of the proof that $A \cap N = B \cap N = 1$. Take, as above, minimal normal subgroups N_1 of A and N_2 of B . It is clear that N_1 and N_2 are a p -group and a q -group, respectively, for primes p and q . Now if there exists a prime $r \neq p, r \neq q$ such that r divides $|N|$, then every Sylow r -subgroup of $A \cap B$, say H , will be a Sylow r -subgroup of $N_1(A \cap B)N_2$ as well. But therefore $H \cap N$ is a Sylow r -subgroup of N . We reach a contradiction if we note that $H \cap N \leq A \cap B \cap N = 1$. Hence N is a $\{p, q\}$ -group, and this implies the solubility of N by Burnside's Theorem [9, Theorem I.2]. This fact leads to our final contradiction since G/N is also soluble by induction. ■

3. The “supersoluble version” of Stonehewer’s theorem

Proof of Theorem A

Assume the result is not true and let G be a minimal counterexample. Let N be a minimal normal subgroup of G such that either AN or BN is not supersoluble. We know by Theorem 3 that G must be soluble, and thus N is an elementary abelian p -group for some prime p . We prove that N is the unique normal minimal subgroup of G .

Let M denote a minimal normal subgroup of G with $N \neq M$. First the group G/M satisfies the hypotheses of the theorem and hence the minimal choice of G implies that both ANM/M and BNM/M are supersoluble. Therefore, the supersoluble residual $(AN)^u$ of AN is contained in M . Moreover, AN/N is supersoluble. Hence $(AN)^u \leq N$. Consequently $(AN)^u \leq M \cap N = 1$, and AN is supersoluble. Analogously, since BNM/M is supersoluble, we can deduce that BN is supersoluble, a contradiction. Consequently we may assume that $N = Soc(G)$ is the unique minimal normal subgroup of G . Suppose that $N \leq A$. Then every subgroup of $N/(B \cap N)$ is normal in $BN/(B \cap N)$. This implies that BN is supersoluble, a contradiction. Consequently N is neither contained in A nor in B .

We shall see next that p must be the largest prime dividing $|G|$. Let q be this largest prime, and assume $q \neq p$. We can assume that q divides $|B|$. In such case, B has a unique Sylow q -subgroup, B_q say. Note that AB_q is a subgroup of G . Now $1 \neq B_q^G = B_q^{BA} = B_q^A \leq AB_q$ and thus N is contained in B_q^G . This clearly implies that $N \leq A$, a contradiction.

Using Lemma 2, we know that either $N = (A \cap N)(B \cap N)$ or $A \cap N = B \cap N = 1$. We consider these two possible situations separately.

Let us assume first that $N = (A \cap N)(B \cap N)$. Note that we can assume that both $A \cap N$ and $B \cap N$ are non-trivial proper subgroups of N .

We know that either AN or BN is not supersoluble. We assume without loss of generality that $Z = AN$ is not supersoluble. Let A_1 be a supersoluble projector of Z containing A (see [9; III, 3]). Note that $Z = A_1Z^{\mathcal{U}}$ and $1 = (A_1 \cap Z^{\mathcal{U}}) \neq Z^{\mathcal{U}} \leq N$ (see [9; IV, 5.18]). Let x be a non-trivial element of $Z^{\mathcal{U}}$. We can find elements n_1 of $N \cap A$ and n_2 of $N \cap B$ such that $x = n_1n_2$ and $o(n_i) = p$ for $i = 1, 2$. If $\langle n_2 \rangle \cap A_1 \neq 1$ then $n_2 \in A_1$ and thus $x \in Z^{\mathcal{U}} \cap A_1 = 1$, a contradiction. Consequently $\langle n_2 \rangle \cap A_1 = 1$. Moreover, $A_1 = A(N \cap A_1)$. Since $\langle n_2 \rangle$ permutes with both A and $N \cap A_1$, it follows that $\langle n_2 \rangle$ permutes with A_1 . Therefore $X = A_1\langle n_2 \rangle$ is a subgroup of G . Note that

$$N \cap X = N \cap A_1\langle n_2 \rangle = \langle n_2 \rangle(N \cap A_1)$$

is a normal subgroup of X as is $N \cap A_1$. If we consider the series $N \cap A_1 \trianglelefteq \langle n_2 \rangle(N \cap A_1) \trianglelefteq X$, we can deduce that $X/N \cap A_1$ is a supersoluble group. Hence $X^{\mathcal{U}}$ is contained in $A_1 \cap Z^{\mathcal{U}} = 1$ and consequently X is supersoluble. This is impossible since A_1 is a supersoluble projector of X .

Hence we can assume that $A \cap N = B \cap N = 1$. We now prove:

Given any non-trivial element x of N , there must exist a subgroup N_x of N such that $x \in N_x$, $N_x \trianglelefteq AN_x$ and AN_x is supersoluble.

In order to prove this statement, note that if A_p and B_p denote the Sylow p -subgroups of A and B respectively, then $N \leq A_pB_p$ applying [2; Corollary 1.3.3]. Consequently we can write $x = ab$ where $a \in A_p$ and $b \in B_p$. Note that $\langle b \rangle$ is a subnormal subgroup of B and therefore it permutes with A . Let $Y = A\langle b \rangle$, and write $N_x = N \cap Y$. Clearly $x \in N_x \leq N$ and N_x is a normal subgroup of $Y = A(N \cap Y) = AN_x$. We shall be done if Y is supersoluble. Assume on the contrary that Y is not supersoluble and take a supersoluble projector A_1 of Y with $A \leq A_1$ (see [9; III, 3]). Let now β be the smallest non-negative integer such that $b^{p^\beta} \in A_1$. Since $Y = A\langle b \rangle = A_1\langle b \rangle$ and Y is not supersoluble, we have that $\beta > 0$. Moreover, $A_1 \cap \langle b \rangle = \langle b^{p^\beta} \rangle$. Write $T = A_1\langle b^{p^{\beta-1}} \rangle = A_1T^{\mathcal{U}}$. Note that A_1 is a supersoluble projector of T , and that $T^{\mathcal{U}}$ is an abelian group since $T^{\mathcal{U}} \leq Y^{\mathcal{U}}$, and $T^{\mathcal{U}} \cap A_1 = 1$. Now

$$|T^{\mathcal{U}}| = |T : A_1| = |\langle b^{p^{\beta-1}} \rangle : \langle b^{p^\beta} \rangle| = p$$

and therefore T is supersoluble, a contradiction. Hence we have proved the above statement.

Assume now that AN is not supersoluble. There must exist, in such case, a non-cyclic chief factor K/L of AN , with $K \leq N$. Let $x \in K \setminus L$. Following

the above claim, we know that there must exist a subgroup N_x of N in the described situation. Note that N_x is in fact a normal subgroup of AN . Now N_xL/L is a normal subgroup of AN/L and K/L is a minimal normal of AN/L . Consequently $1 \neq xL \in N_xL/L \cap K/L$. Thus $N_xL/L \cap K/L \neq 1$ and hence $K/L \leq N_xL/L \leq AN_x/L$ which is supersoluble. We conclude that K/L is a cyclic group, our final contradiction. ■

4. Main consequences

The following lemma will be used in the proofs of this section.

Lemma 3 *Let G be a primitive group and let N be its unique minimal normal subgroup. Assume that G/N is supersoluble. If N is a p -group, where p is the largest prime dividing $|G|$, then $N = F(G) = O_p(G)$ is a Sylow p -subgroup of G .*

Proof. It is well known that $N = F(G) = O_p(G)$ since $O_q(G) = 1$ for every prime $q \neq p$. Therefore we only need to show that N is a Sylow p -subgroup of G .

Since G is a primitive soluble group, we can write $G = NM$, where M is a maximal subgroup of G , $N \cap M = 1$ and $C_G(N) = N$.

Note that $O_p(M) = 1$ by [9, Theorem A.15.6.b]. Clearly $M \cong G/N$ is supersoluble. On the other hand, a Sylow p -subgroup of M is normal in M and consequently p does not divide the order of M . This completes the proof. ■

We begin by studying a minimal counterexample to claim the following:

“If $G = AB$ is the mutually sn-permutable product of the supersoluble subgroups A and B , then G is supersoluble.”

Let $G = AB$ be a minimal counterexample to the above property. G must be a primitive soluble group, and hence there exists a unique minimal normal subgroup N of G , such that $N = C_G(N)$. Let p be the prime dividing $|N|$. By using Theorem A, we know that both AN and BN are supersoluble.

We show that p must be the largest prime dividing $|G|$. Let q be this largest prime, and assume $q \neq p$. We can assume without loss of generality that q divides $|AN|$. The supersoluble group AN must have a unique Sylow q -subgroup, $(AN)_q$ say, and clearly $(AN)_q$ centralizes N . This implies that $(AN)_q = 1$ since $C_G(N) = N$, a contradiction. Thus p must be the largest prime dividing $|G|$ and so we have the structure given by Lemma 3.

Next we consider a minimal counterexample to the following claim:

“Let $G = AB$ be the mutually sn-permutable product of the subgroups A and B , where A is supersoluble and B is nilpotent. Then G is supersoluble.”

Consider such a minimal counterexample, G say. As before, $G = AB$ is a primitive soluble group with a unique minimal normal subgroup N . N is a self-centralizing p -group for a prime p . By the argument above and by Lemma 3, p must be the largest prime dividing G , N is the unique Sylow p -subgroup of G and by Theorem A both AN and BN are supersoluble.

We shall see now that N must be contained in A . Note that if $N \leq B$, then the nilpotency of B implies that B is a p -group. In such case $G = AN$ and G is supersoluble, a contradiction. Consequently we can assume that N is not contained in B . Consider now a Hall p' -subgroup $B_{p'}$ of B . Since $B_{p'}$ is a normal subgroup of B , it permutes with A and thus $AB_{p'}$ is a subgroup of G . If $B_{p'} \neq 1$ then $1 \neq B_{p'}^G \leq AB_{p'}$ and hence $N \leq A$, as desired. Hence $B_{p'} = 1$ and thus B must be a p -group. In such case $B \leq N$ and hence $G = AB = AN$ and G is supersoluble, a contradiction. Thus $A \cap N = N$ and therefore $N \leq A$, as we wanted to prove.

At this point we easily obtain the following consequences:

Proof of Theorem 1 [6, Theorem 2]

From the argument above, we can assume that $N \leq A$, where N denotes the unique minimal normal subgroup of G , N an elementary abelian Sylow p -subgroup of G . Denote by N_1 a minimal normal subgroup of A such that $N_1 \leq N$. Then N_1 permutes with $B_{p'}$ and is centralized by B_p . Hence $N \leq N_1^G \leq N_1 B_{p'}$ and so $N \leq N_1$. Consequently $N = N_1$ and N is a cyclic group of prime order, the final contradiction. ■

Proof of Theorem B

As B permutes with every Sylow subgroup of A , we can consider the subgroup of G , $A_{p'}B$, where $A_{p'}$ denotes a Hall p' -subgroup of A . Notice that if $A_{p'} = 1$, we would have $A = N$, and $G = BN$ is supersoluble by Theorem A, a contradiction. If $G = A_{p'}B$, then $N \leq B$ and B is a p -group, a contradiction. Hence $A_{p'}B < G$ and $G = NA_{p'}B$. On the other hand, $N \cap A_{p'}B = N \cap B$ is a normal subgroup of $A_{p'}B$. Hence $N \cap B$ is a normal subgroup of G , $N = N \cap B \leq B$ and consequently $N \cap B = 1$. Therefore B is a p' -group. Take now N_1 a minimal normal subgroup of A with $N_1 \leq N$. Then $N \leq N_1^G \leq N_1 B \leq G$. Thus $N = N_1$, a cyclic group of prime order, the final contradiction. ■

Proof of Theorem C

Assume the result is not true and let G be a minimal counterexample. It is clear that $G' \neq 1$ and that G is a primitive group. Let N be the unique minimal normal subgroup. By Lemma 3, we have $N = F(G)$ and so $G' \leq F(G) = N$. From AN and BN are supersoluble (by Theorem A) and normal ($G' \leq AN$, $G' \leq BN$) we can apply Baer's Theorem to give G supersoluble, a contradiction. ■

Proof of Theorem D

Assume that the Theorem is false and take a minimal counterexample $G = AB$. We have that G is a primitive soluble group. Write, as usual, $G = NM$, where N is the unique minimal normal subgroup of G , M is a maximal subgroup of G , $N \cap M = 1$ and $C_G(N) = N$. We also know that N is a p -group for some prime p , and by Lemma 3 and the metanilpotence of G we have $N = F(G) = O_p(G) = G^N$ and N is a Sylow p -subgroup of G . Consequently we may assume that $M \cong G/N$ is nilpotent. Note that $O_p(M) = 1$ by [9, Theorem A.15.6.b].

Let now $A_{p'}$ and $B_{p'}$ be Hall p' -subgroups of A and B , respectively, such that $A_{p'}B_{p'}$ is a Hall p' -subgroup of G (see [2, Theorem 1.3.2]). Replacing M by some conjugate if necessary, we can assume $A_{p'}B_{p'} = M$. The nilpotency of M and the fact that $\pi(|A|) \cap \pi(|B|) \subseteq \{p\}$ allows us to write $M = A_{p'} \times B_{p'}$.

Take now the subgroup $T = BN$ of G . Note that T is supersoluble according to Theorem A. Using the local definition of the class of supersoluble groups, we obtain that $T/O_{p'}(T)$ is an abelian group of exponent dividing $p - 1$.

Now N is self-centralizing in T and hence $O_{p'}(T) = 1$. It follows that $O_{p'}(T) = O_p(T) = O_p(BN) = N$. This implies that $BN/N \cong B/B \cap N \cong B_{p'}$ is an abelian group of exponent dividing $p - 1$. We could reproduce the same argument for A and obtain thus that both $A_{p'}$ and $B_{p'}$ are abelian groups of exponent dividing $p - 1$.

But $M = A_{p'} \times B_{p'}$, and hence M is also abelian of exponent dividing $p - 1$. But N is an irreducible and faithful module for M , and consequently its dimension must equal one. We conclude that N is cyclic, the final contradiction. ■

5. Some counterexamples and remarks

It is not true in general that a group which is the totally sn -permutable product of two supersoluble subgroups is supersoluble, as the following example shows.

Example. Let V be a faithful irreducible module for S , the symmetric group of degree 3, over the field of 7 elements. It is easy to check that V has dimension 2. Let G be the semidirect product of V and S , and let $A = VS_3$, $B = VS_2$, where S_p is a Sylow p -subgroup of S . It is again easy to check that A and B are supersoluble, G is the totally sn -permutable product of A and B , but G is not supersoluble.

Note that the hypothesis on B in Theorem B is essential in order to get the supersolubility of G as the following example illustrates.

Example. Consider $X = \langle a, b : a^4 = 1 = b^2, a^b = a^{-1} \rangle$, the dihedral group of order 8, and let $V = \langle v_1, v_2 \rangle$ be a $GF(5)$ -space of dimension 2. Then V can be considered as an X -module with the following action:

$$\begin{aligned} v_1^a &= 3v_1 & v_1^b &= v_2 \\ v_2^a &= 2v_2 & v_2^b &= v_1 \end{aligned}$$

Take now $G = [V]X$, the corresponding semidirect product, and consider the following subgroups of G :

$$A = V\langle a \rangle ; \quad B = \langle v_1v_2 \rangle \times \langle b \rangle$$

Note that $G = AB$. It is clear that A is supersoluble, B is nilpotent and the nilpotent residual of G is precisely V , which is abelian. It is easy to see that G is the mutually sn -permutable product of the subgroups A and B , but G is not supersoluble.

References

- [1] ALEJANDRE, MANUEL J., BALLESTER-BOLINCHES, A., COSSEY, JOHN: Permutable products of supersoluble groups. To appear in *J. Algebra*.
- [2] AMBERG, B., FRANCIOSI, S. AND DE GIOVANNI, F.: *Products of groups*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1992.
- [3] ASAAD, M., SHAALAN, A.: On the supersolubility of finite groups. *Arch. Math.* **53** (1989), no. 4, 318–326.
- [4] BAER, R.: Classes of finite groups and their properties. *Illinois J. Math.* **1** (1957), 115–187.
- [5] BALLESTER-BOLINCHES, A., DOERK, K. AND PÉREZ-RAMOS, M. D.: On the lattice of \mathcal{F} -subnormal groups. *J. Algebra* **148** (1992), no. 1, 42–52.
- [6] BEIDLEMAN, J. C., GALOPPO, A., HEINEKEN, H. AND MANFREDINO, M.: On certain products of soluble groups. *Forum Math.* **13** (2001), 569–580.
- [7] CAROCCA, A.: p -supersolvability of factorized finite groups. *Hokkaido Math. J.* **21** (1992), 395–403.
- [8] CAROCCA, A.: On factorized finite groups in which certain subgroups of the factors permute. *Arch. Math.* **71** (1998), 257–261.
- [9] DOERK, K. AND HAWKES, T. O.: *Finite soluble groups*. De Gruyter Expositions in Mathematics 4. Walter de Gruyter, Berlin, 1992.
- [10] FÖRSTER, P.: Finite groups all whose subgroups are \mathcal{F} -subnormal or \mathcal{F} -subabnormal. *J. Algebra* **103** (1986), no. 1, 285–293.

- [11] KEGEL, O. H.: Sylow-Gruppen und Subnormalteiler endlicher Gruppen. *Math. Z.* **78** (1962), 205–221.
- [12] STONEHEWER, S. E.: On finite nilpotent groups. *J. Pure Appl. Algebra* **88** (1993), no. 1-3, 239–244.

Recibido: 13 de mayo de 2002

Manuel J. Alejandro
 Centro de Investigación Operativa
 Universidad Miguel Hernández
 Avda. del Ferrocarril s/n
 03202 Elche (Spain)
 m.alejandro@umh.es

A. Ballester-Bolinches
 Departament d'Àlgebra
 Universitat de València
 Dr. Moliner 50
 46100 Burjassot, València (Spain)
 Adolfo.Ballester@uv.es

John Cossey
 Mathematics Department
 School of Mathematical Sciences
 Australian National University
 0200 Canberra (Australia)
 John.Cossey@maths.anu.edu.au

M. C. Pedraza-Aguilera
 Escuela Universitaria de Informática
 Departamento de Matemática Aplicada
 Universidad Politécnica de Valencia
 Camino de Vera, s/n
 46071 Valencia (Spain)
 mpedraza@mat.upv.es

This paper was partly done while the first author was a visitor at the Australian National University. He is grateful to the Department of Mathematics for its warm hospitality while this investigation was carried out. The second and the fourth authors are supported by Proyecto PB 97-0674-C02-02 of DGICYT, MEC, Spain. The third author wishes to thank the Centro de Investigación Operativa of the Universidad Miguel Hernández for its support of this research.