

# A note on the existence of $H$ -bubbles via perturbation methods

Veronica Felli

## Abstract

We study the problem of existence of surfaces in  $\mathbb{R}^3$  parametrized on the sphere  $\mathbb{S}^2$  with prescribed mean curvature  $H$  in the perturbative case, i.e. for  $H = H_0 + \varepsilon H_1$ , where  $H_0$  is a nonzero constant,  $H_1$  is a  $C^2$  function and  $\varepsilon$  is a small perturbation parameter.

## 1. Introduction

In this paper we are interested in the existence of  $H$ -bubbles, namely of  $\mathbb{S}^2$ -type parametric surfaces in  $\mathbb{R}^3$  with prescribed mean curvature  $H$ . This geometrical problem is motivated by some models describing capillarity phenomena and has the following analytical formulation: given a function  $H \in C^1(\mathbb{R}^3)$ , find a smooth nonconstant function  $\omega : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  which is conformal as a map on  $\mathbb{S}^2$  and solves the problem

$$(P_H) \quad \begin{cases} \Delta\omega = 2H(\omega)\omega_x \wedge \omega_y, & \text{in } \mathbb{R}^2, \\ \int_{\mathbb{R}^2} |\nabla\omega|^2 < +\infty, \end{cases}$$

where

$$\begin{aligned} \omega_x &= \left( \frac{\partial\omega_1}{\partial x}, \frac{\partial\omega_2}{\partial x}, \frac{\partial\omega_3}{\partial x} \right), & \omega_y &= \left( \frac{\partial\omega_1}{\partial y}, \frac{\partial\omega_2}{\partial y}, \frac{\partial\omega_3}{\partial y} \right), \\ \Delta\omega &= \omega_{xx} + \omega_{yy}, & \nabla\omega &= (\omega_x, \omega_y), \end{aligned}$$

and  $\wedge$  denotes the exterior product in  $\mathbb{R}^3$ .

Brezis and Coron [4] proved that for constant nonzero mean curvature  $H(u) \equiv H_0$  the only nonconstant solutions are spheres of radius  $|H_0|^{-1}$ .

---

*2000 Mathematics Subject Classification:* 53A10, 35J50, 35B20.

*Keywords:* H-surfaces, nonlinear elliptic systems, perturbative methods.

While the Plateau and the Dirichlet problems have been largely studied both for  $H$  constant and for  $H$  nonconstant (see [3, 4, 10, 12, 13, 14, 15, 16]), problem  $(P_H)$  in the case of nonconstant  $H$  has been investigated only recently, see [5, 6, 7]. In [5] Caldirolì and Musina proved the existence of  $H$ -bubbles with minimal energy under the assumptions that  $H \in C^1(\mathbb{R}^3)$  satisfies

- (i)  $\sup_{u \in \mathbb{R}^3} |\nabla H(u + \xi) \cdot u| < 1$  for some  $\xi \in \mathbb{R}^3$ ,
- (ii)  $H(u) \rightarrow H_\infty$  as  $|u| \rightarrow \infty$  for some  $H_\infty \in \mathbb{R}$ ,
- (iii)  $c_H = \inf_{\substack{u \in C_c^1(\mathbb{R}^2, \mathbb{R}^3) \\ u \neq 0}} \sup_{s > 0} \mathcal{E}_H(su) < \frac{4\pi}{3H_\infty^2}$

where

$$\mathcal{E}_H(u) = \frac{1}{2} \int_{\mathbb{R}^2} |\nabla u|^2 + 2 \int_{\mathbb{R}^2} Q(u) \cdot u_x \wedge u_y$$

and  $Q : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  is any vector field such that  $\operatorname{div} Q = H$ .

The perturbative method introduced by Ambrosetti and Rabinowitz [1, 2] was used in [7] to treat the case in which  $H$  is a small perturbation of a constant, namely

$$H(u) = H_\varepsilon(u) = H_0 + \varepsilon H_1(u),$$

where  $H_0 \in \mathbb{R} \setminus \{0\}$ ,  $H_1 \in C^2(\mathbb{R}^3)$ , and  $\varepsilon$  is a small real parameter. This method allows to find critical points of a functional  $f_\varepsilon$  of the type  $f_\varepsilon(u) = f_0(u) - \varepsilon G(u)$  in a Banach space by studying a finite dimensional problem. More precisely, if the unperturbed functional  $f_0$  has a finite dimensional manifold of critical points  $Z$  which satisfies a nondegeneracy condition, it is possible to prove, for  $|\varepsilon|$  sufficiently small, the existence of a smooth function  $\eta_\varepsilon(z) : Z \rightarrow (T_z Z)^\perp$  such that any critical point  $\bar{z} \in Z$  of the function

$$\Phi_\varepsilon : Z \rightarrow \mathbb{R}, \quad \Phi_\varepsilon(z) = f_\varepsilon(z + \eta_\varepsilon(z))$$

gives rise to a critical point  $u_\varepsilon = \bar{z} + \eta_\varepsilon(\bar{z})$  of  $f_\varepsilon$ , i.e. the perturbed manifold  $Z_\varepsilon := \{z + \eta_\varepsilon(z) : z \in Z\}$  is a natural constraint for  $f_\varepsilon$ . Furthermore  $\Phi_\varepsilon$  can be expanded as

$$(1.1) \quad \Phi_\varepsilon(z) = b - \varepsilon \Gamma(z) + o(\varepsilon) \quad \text{as } \varepsilon \rightarrow 0$$

where  $b = f_0(z)$  and  $\Gamma$  is the Melnikov function defined as the restriction of the perturbation  $G$  on  $Z$ , namely  $\Gamma = G|_Z$ . For problem  $(P_{H_\varepsilon})$ ,  $\Gamma$  is given by

$$\Gamma : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad \Gamma(p) = \int_{|p-q| < \frac{1}{|H_0|}} H_1(q) dq.$$

In [7] Caldiroli and Musina studied the functional  $\Gamma$  giving some existence results in the perturbative setting for problem  $(P_{H_\varepsilon})$ . They prove that for  $|\varepsilon|$  small there exists a smooth  $H_\varepsilon$ -bubble if one of the following conditions holds

- 1)  $H_1$  has a nondegenerate stationary point and  $|H_0|$  is large;
- 2)  $\max_{p \in \partial K} H_1(p) < \max_{p \in K} H_1(p)$  or  $\min_{p \in \partial K} H_1(p) > \min_{p \in K} H_1(p)$  for some non-empty compact set  $K \subset \mathbb{R}^3$  and  $|H_0|$  is large;
- 3)  $H_1 \in L^r(\mathbb{R}^3)$  for some  $r \in [1, 2]$ .

They prove that critical points of  $\Gamma$  give rise to solutions to  $(P_{H_\varepsilon})$  for  $\varepsilon$  sufficiently small. Precisely the assumption that  $H_0$  is large required in cases 1) and 2) ensures that if  $H_1$  is not constant then  $\Gamma$  is not identically constant. If we let this assumption drop, it may happen that  $\Gamma$  is constant even if  $H_1$  is not. This fact produces some loss of information because the first order expansion (1.1) is not sufficient to deduce the existence of critical points of  $\Phi_\varepsilon$  from the existence of critical points of  $\Gamma$ . Instead of studying  $\Gamma$  we perform a direct study of  $\Phi_\varepsilon$  which allows us to prove some new results. In the first one, we assume that  $H_1$  vanishes at  $\infty$  and has bounded gradient, and prove the existence of a solution without branch points. Let us recall that a branch point for a solution  $\omega$  to  $(P_H)$  is a point where  $\nabla\omega = 0$ , i.e. a point where the surface parametrized by  $\omega$  fails to be immersed.

**Theorem 1.1** *Let  $H_0 \in \mathbb{R} \setminus \{0\}$ ,  $H_1 \in C^2(\mathbb{R}^3)$  such that*

$$(H1) \quad \lim_{|p| \rightarrow \infty} H_1(p) = 0;$$

$$(H2) \quad \nabla H_1 \in L^\infty(\mathbb{R}^3, \mathbb{R}^3).$$

*Let  $H_\varepsilon = H_0 + \varepsilon H_1$ . Then for  $|\varepsilon|$  sufficiently small there exists a smooth  $H_\varepsilon$ -bubble without branch points.*

With respect to case 1) of [7] we require neither nondegeneracy of critical points of  $H_1$  nor largeness of  $H_0$ . With respect to case 2) we do not assume that  $H_0$  is large; on the other hand our assumption (H1) implies 2). Moreover we do not assume any integrability condition of type 3). With respect to the result proved in [5], we have the same kind of behavior of  $H_1$  at  $\infty$  (see (ii) and assumption (H1)) but we do not need any assumption of type (iii); on the other hand in [5] it is not required that the prescribed curvature is a small perturbation of a constant.

The following results give some conditions on  $H_1$  in order to have two or three solutions.

**Theorem 1.2** *Let  $H_0 \in \mathbb{R} \setminus \{0\}$ ,  $H_1 \in C^2(\mathbb{R}^3)$  such that (H1), (H2),*

- (H3) *Hess  $H_1(p)$  is positive definite for any  $p \in B_{1/|H_0|}(0)$ ,*
- (H4)  *$H_1(p) > 0$  in  $B_{1/|H_0|}(0)$ ,*

*hold. Then for  $|\varepsilon|$  sufficiently small there exist at least three smooth  $H_\varepsilon$ -bubbles without branch points.*

**Remark 1.3** *If we assume (H1), (H2), and, instead of (H3) – (H4), that  $H_1(0) > 0$  and Hess  $H_1(0)$  is positive definite, then we can prove that for  $|H_0|$  sufficiently large and  $|\varepsilon|$  sufficiently small there exist at least three smooth  $H_\varepsilon$ -bubbles without branch points.*

**Theorem 1.4** *Let  $H_0 \in \mathbb{R} \setminus \{0\}$ ,  $H_1 \in C^2(\mathbb{R}^3)$  such that (H1) and (H2) hold. Assume that there exist  $p_1, p_2 \in \mathbb{R}^3$  such that*

$$(H5) \quad \int_{B(p_1, 1/|H_0|)} H_1(\xi) d\xi > 0 \quad \text{and} \quad \int_{B(p_2, 1/|H_0|)} H_1(\xi) d\xi < 0.$$

*Then for  $|\varepsilon|$  sufficiently small there exist at least two smooth  $H_\varepsilon$ -bubbles without branch points.*

**Remark 1.5** *If we assume (H1), (H2), and, instead of (H5), that there exist  $p_1, p_2 \in \mathbb{R}^3$  such that  $H_1(p_1) > 0$  and  $H_1(p_2) < 0$ , then we can prove that for  $|H_0|$  sufficiently large and  $|\varepsilon|$  sufficiently small there exist at least two smooth  $H_\varepsilon$ -bubbles without branch points.*

The present paper is organized as follows. In Section 2 we introduce some notation and recall some known facts whereas Section 3 is devoted to the proof of Theorems 1.1, 1.2, and 1.4.

## 2. Notation and known facts

In the sequel we will take  $H_0 = 1$ ; this is not restrictive since we can do the change  $H_1(u) = H_0 \tilde{H}_1(H_0 u)$ . Hence we will always write

$$H_\varepsilon(u) = 1 + \varepsilon H(u),$$

where  $H \in C^2(\mathbb{R}^3)$ . Let us denote by  $\omega$  the function  $\omega : \mathbb{R}^2 \rightarrow \mathbb{S}^2$  defined as

$$\omega(x, y) = (\mu(x, y)x, \mu(x, y)y, 1 - \mu(x, y)), \quad \text{where } \mu(x, y) = \frac{2}{1 + x^2 + y^2}.$$

Note that  $\omega$  is a conformal parametrization of the unit sphere and solves

$$(2.1) \quad \begin{cases} \Delta \omega = 2 \omega_x \wedge \omega_y & \text{on } \mathbb{R}^2 \\ \int_{\mathbb{R}^2} |\nabla \omega|^2 < +\infty. \end{cases}$$

Problem (2.1) has in fact a family of solutions of the form  $\omega \circ \phi + p$  where  $p \in \mathbb{R}^3$  and  $\phi$  is any conformal diffeomorphism of  $\mathbb{R}^2 \cup \{\infty\}$ . For  $s \in (1, +\infty)$ , we will set  $L^s := L^s(\mathbb{S}^2, \mathbb{R}^3)$ , where any map  $v \in L^s$  is identified with the map  $v \circ \omega : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  which satisfies

$$\int_{\mathbb{R}^2} |v \circ \omega|^s \mu^2 = \int_{\mathbb{S}^2} |v|^s.$$

We will use the same notation for  $v$  and  $v \circ \omega$ . By  $W^{1,s}$  we denote the Sobolev space  $W^{1,s}(\mathbb{S}^2, \mathbb{R}^3)$  endowed (according to the above identification) with the norm

$$\|v\|_{W^{1,s}} = \left[ \int_{\mathbb{R}^2} |\nabla v|^s \mu^{2-s} \right]^{1/s} + \left[ \int_{\mathbb{R}^2} |v|^s \mu^2 \right]^{1/s}.$$

If  $s'$  is the conjugate exponent of  $s$ , i.e.  $1/s + 1/s' = 1$ , the duality product between  $W^{1,s}$  and  $W^{1,s'}$  is given by

$$\langle v, \varphi \rangle = \int_{\mathbb{R}^2} \nabla v \cdot \nabla \varphi + \int_{\mathbb{R}^2} v \cdot \varphi \mu^2 \quad \text{for any } v \in W^{1,s} \text{ and } \varphi \in W^{1,s'}.$$

Let  $Q$  be any smooth vector field on  $\mathbb{R}^3$  such that  $\operatorname{div} Q = H$ . The energy functional associated to problem

$$(P_\varepsilon) \quad \begin{cases} \Delta u = 2(1 + \varepsilon H(u)) u_x \wedge u_y, & \text{in } \mathbb{R}^2, \\ \int_{\mathbb{R}^2} |\nabla u|^2 < +\infty, \end{cases}$$

is given by

$$\mathcal{E}_\varepsilon(u) = \frac{1}{2} \int_{\mathbb{R}^2} |\nabla u|^2 + 2\mathcal{V}_1(u) + 2\varepsilon \mathcal{V}_H(u), \quad u \in W^{1,3},$$

where

$$\mathcal{V}_H(u) = \int_{\mathbb{R}^2} Q(u) \cdot u_x \wedge u_y$$

has the meaning of an algebraic volume enclosed by the surface parametrized by  $u$  with weight  $H$  (it is independent of the choice of  $Q$ ); in particular

$$\mathcal{V}_1(u) = \frac{1}{3} \int_{\mathbb{R}^2} u \cdot u_x \wedge u_y.$$

In [7], Caldiroli and Musina studied some regularity properties of  $\mathcal{V}_H$  on the space  $W^{1,3}$ . In particular they proved the following properties.

a) For  $H \in C^1(\mathbb{R}^3)$ , the functional  $\mathcal{V}_H$  is of class  $C^1$  on  $W^{1,3}$  and the Fréchet differential of  $\mathcal{V}_H$  at  $u \in W^{1,3}$  is given by

$$(2.2) \quad d\mathcal{V}_H(u)\varphi = \int_{\mathbb{R}^2} H(u) \varphi \cdot u_x \wedge u_y \quad \text{for any } \varphi \in W^{1,3}$$

and admits a unique continuous and linear extension on  $W^{1,3/2}$  defined by (2.2). Moreover for every  $u \in W^{1,3}$  there exists  $\mathcal{V}'_H(u) \in W^{1,3}$  such that

$$(2.3) \quad \langle \mathcal{V}'_H(u), \varphi \rangle = \int_{\mathbb{R}^2} H(u) \varphi \cdot u_x \wedge u_y \quad \text{for any } \varphi \in W^{1,3/2}.$$

b) For  $H \in C^2(\mathbb{R}^3)$ , the map  $\mathcal{V}'_H : W^{1,3} \rightarrow W^{1,3}$  is of class  $C^1$  and

$$(2.4) \quad \begin{aligned} \langle \mathcal{V}''_H(u) \cdot \eta, \varphi \rangle &= \int_{\mathbb{R}^2} H(u) \varphi \cdot (\eta_x \wedge u_y + u_x \wedge \eta_y) + \int_{\mathbb{R}^2} (\nabla H(u) \cdot \eta) \varphi \cdot (u_x \wedge u_y) \\ &\text{for any } u, \eta \in W^{1,3} \text{ and } \varphi \in W^{1,3/2}. \end{aligned}$$

Hence for all  $u \in W^{1,3}$ ,  $\mathcal{E}'_\varepsilon(u) \in W^{1,3}$  and for any  $\varphi \in W^{1,3/2}$

$$\langle \mathcal{E}'_\varepsilon(u), \varphi \rangle = \int_{\mathbb{R}^2} \nabla u \cdot \nabla \varphi + 2 \int_{\mathbb{R}^2} \varphi \cdot u_x \wedge u_y + 2\varepsilon \int_{\mathbb{R}^2} H(u) \varphi \cdot u_x \wedge u_y.$$

As remarked in [7], critical points of  $\mathcal{E}_\varepsilon$  in  $W^{1,3}$  give rise to bounded weak solutions to  $(P_\varepsilon)$  and hence by the regularity theory for  $H$ -systems (see [9]) to classical conformal solutions which are  $C^{3,\alpha}$  as maps on  $\mathbb{S}^2$ .

The unperturbed problem, i.e.  $(P_\varepsilon)$  for  $\varepsilon = 0$ , has a 9-dimensional manifold of solutions given by

$$Z = \{R\omega \circ L_{\lambda,\xi} + p : R \in SO(3), \lambda > 0, \xi \in \mathbb{R}^2, p \in \mathbb{R}^3\}$$

where  $L_{\lambda,\xi}z = \lambda(z - \xi)$  (see [11]). In [11] the nondegeneracy condition  $T_u Z = \ker \mathcal{E}''_0(u)$  for any  $u \in Z$  (where  $T_u Z$  denotes the tangent space of  $Z$  at  $u$ ) is proved (see also [8]).

As observed in [7], in performing the finite dimensional reduction, the dependence on the 6-dimensional conformal group can be neglected since any  $H$ -system is conformally invariant. Hence we look for critical points of  $\mathcal{E}_\varepsilon$  constrained on a three-dimensional manifold  $Z_\varepsilon$  just depending on the translation variable  $p \in \mathbb{R}^3$ .

### 3. Proof of Theorem 1.1

We start by constructing a perturbed manifold which is a natural constraint for  $\mathcal{E}_\varepsilon$ .

**Lemma 3.1** *Assume  $H \in C^2(\mathbb{R}^3) \cap L^\infty(\mathbb{R}^3)$  and  $\nabla H \in L^\infty(\mathbb{R}^3, \mathbb{R}^3)$ . Then there exist  $\varepsilon_0 > 0$ ,  $C_1 > 0$ , and a  $C^1$  map  $\eta : (-\varepsilon_0, \varepsilon_0) \times \mathbb{R}^3 \rightarrow W^{1,3}$  such that for any  $p \in \mathbb{R}^3$  and  $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$*

$$(3.1) \quad \mathcal{E}'_\varepsilon(\omega + p + \eta(\varepsilon, p)) \in T_\omega Z,$$

$$(3.2) \quad \eta(\varepsilon, p) \in (T_\omega Z)^\perp,$$

$$(3.3) \quad \int_{\mathbb{S}^2} \eta(\varepsilon, p) = 0,$$

$$(3.4) \quad \|\eta(\varepsilon, p)\|_{W^{1,3}} \leq C_1 |\varepsilon|.$$

Moreover if we assume that the limit of  $H$  at  $\infty$  exists and

$$(3.5) \quad \lim_{|p| \rightarrow \infty} H(p) = 0$$

we have that  $\eta(\varepsilon, p)$  converges to 0 in  $W^{1,3}$  as  $|p| \rightarrow \infty$  uniformly with respect to  $|\varepsilon| < \varepsilon_0$ .

**Proof.** Let us define the map

$$F = (F_1, F_2) : \mathbb{R} \times \mathbb{R}^3 \times W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3 \rightarrow W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3$$

given by

$$\langle F_1(\varepsilon, p, \eta, \lambda, \alpha), \varphi \rangle = \langle \mathcal{E}'_\varepsilon(\omega + p + \eta), \varphi \rangle - \sum_{i=1}^6 \lambda_i \int_{\mathbb{R}^2} \nabla \varphi \cdot \nabla \tau_i + \alpha \cdot \int_{\mathbb{S}^2} \varphi,$$

for all  $\varphi \in W^{1,3/2}$  and

$$F_2(\varepsilon, p, \eta, \lambda, \alpha) = \left( \int_{\mathbb{R}^2} \nabla \eta \cdot \nabla \tau_1, \dots, \int_{\mathbb{R}^2} \nabla \eta \cdot \nabla \tau_6, \int_{\mathbb{S}^2} \eta \right)$$

where  $\tau_1, \dots, \tau_6$  are chosen in  $T_\omega Z$  such that

$$\int_{\mathbb{R}^2} \nabla \tau_i \cdot \nabla \tau_j = \delta_{ij} \quad \text{and} \quad \int_{\mathbb{S}^2} \tau_i = 0 \quad i, j = 1, \dots, 6$$

so that  $T_\omega Z$  is spanned by  $\tau_1, \dots, \tau_6, e_1, e_2, e_3$ . It has been proved by Caldiroli and Musina [7] that  $F$  is of class  $C^1$  and that the linear continuous operator

$$\begin{aligned} \mathcal{L} : W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3 &\rightarrow W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3 \\ \mathcal{L} &= \frac{\partial F}{\partial(\eta, \lambda, \alpha)}(0, p, 0, 0, 0) \end{aligned}$$

i.e.

$$\langle \mathcal{L}_1(v, \mu, \beta), \varphi \rangle = \langle \mathcal{E}_0''(\omega) \cdot v, \varphi \rangle - \sum_{i=1}^6 \mu_i \int_{\mathbb{R}^2} \nabla \varphi \cdot \tau_i - \beta \int_{\mathbb{S}^2} \varphi$$

for all  $\varphi \in W^{1,3/2}$  and

$$\mathcal{L}_2(v, \mu, \beta) = \left( \int_{\mathbb{R}^2} \nabla v \cdot \nabla \tau_1, \dots, \int_{\mathbb{R}^2} \nabla v \cdot \nabla \tau_6, \int_{\mathbb{S}^2} v \right)$$

is invertible. Caldiroli and Musina applied the Implicit Function Theorem to solve the equation  $F(\varepsilon, p, \eta, \lambda, \alpha) = 0$  locally with respect to the variables  $\varepsilon, p$ , thus finding a  $C^1$ -function  $\eta$  on a neighborhood  $(-\varepsilon_0, \varepsilon_0) \times B_R \subset \mathbb{R} \times \mathbb{R}^3$  satisfying (3.1), (3.2), and (3.3). We will use instead the Contraction Mapping Theorem, which allows to prove the existence of such a function  $\eta$  globally on  $\mathbb{R}^3$ , thanks to the fact that the operator  $\mathcal{L}$  does not depend on  $p$  and hence it is invertible uniformly with respect to  $p \in \mathbb{R}^3$ .

We have that  $F(\varepsilon, p, \eta, \lambda, \alpha) = 0$  if and only if  $(\eta, \lambda, \alpha)$  is a fixed point of the map  $T_{\varepsilon, p}$  defined as

$$T_{\varepsilon, p}(\eta, \lambda, \alpha) = -\mathcal{L}^{-1}F(\varepsilon, p, \eta, \lambda, \alpha) + (\eta, \lambda, \alpha).$$

To prove the existence of  $\eta$  satisfying (3.1), (3.2), and (3.3), it is enough to prove that  $T_{\varepsilon, p}$  is a contraction in some ball  $B_\rho(0)$  with  $\rho = \rho(\varepsilon) > 0$  independent of  $p$ , whereas the regularity of  $\eta(\varepsilon, p)$  follows from the Implicit Function Theorem.

We have that if  $\|\eta\|_{W^{1,3}} \leq \rho$

$$\begin{aligned} & \|T_{\varepsilon, p}(\eta, \lambda, \alpha)\|_{W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3} \\ & \leq C_2 \|F(\varepsilon, p, \eta, \lambda, \alpha) - \mathcal{L}(\eta, \lambda, \alpha)\|_{W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3} \\ & \leq C_2 \|\mathcal{E}'_\varepsilon(\omega + p + \eta) - \mathcal{E}_0''(\omega)\eta\|_{W^{1,3}} \\ & \leq C_2 (\|\mathcal{E}'_0(\omega + \eta) - \mathcal{E}_0''(\omega)\eta\|_{W^{1,3}} + 2|\varepsilon| \|\mathcal{V}'_H(\omega + p + \eta)\|_{W^{1,3}}) \\ & \leq C_2 \left( \int_0^1 \|\mathcal{E}_0''(\omega + t\eta) - \mathcal{E}_0''(\omega)\|_{W^{1,3/2}} dt \|\eta\|_{W^{1,3}} \right. \\ & \quad \left. + 2|\varepsilon| \|\mathcal{V}'_H(\omega + p + \eta)\|_{W^{1,3}} \right) \\ & \leq C_2 \rho \sup_{\|\eta\|_{W^{1,3}} \leq \rho} \|\mathcal{E}_0''(\omega + \eta) - \mathcal{E}_0''(\omega)\|_{W^{1,3/2}} \\ (3.6) \quad & \quad + 2C_2 |\varepsilon| \sup_{\|\eta\|_{W^{1,3}} \leq \rho} \|\mathcal{V}'_H(\omega + p + \eta)\|_{W^{1,3}} \end{aligned}$$

where  $C_2 = \|\mathcal{L}^{-1}\|_{\mathcal{L}(W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3)}$ .



For  $(\eta_1, \lambda_1, \alpha_1), (\eta_2, \lambda_2, \alpha_2) \in B_\rho(0) \subset W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3$  we have

$$\begin{aligned}
 & \frac{\|T_{\varepsilon,p}(\eta_1, \lambda_1, \alpha_1) - T_{\varepsilon,p}(\eta_2, \lambda_2, \alpha_2)\|_{W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3}}{C_2 \|\eta_1 - \eta_2\|_{W^{1,3}}} \\
 & \leq \frac{\|\mathcal{E}'_\varepsilon(\omega + p + \eta_1) - \mathcal{E}'_\varepsilon(\omega + p + \eta_2) - \mathcal{E}''_0(\omega)(\eta_1 - \eta_2)\|_{W^{1,3}}}{C_2 \|\eta_1 - \eta_2\|_{W^{1,3}}} \\
 & \leq \int_0^1 \|\mathcal{E}''_\varepsilon(\omega + p + \eta_2 + t(\eta_1 - \eta_2)) - \mathcal{E}''_0(\omega)\|_{W^{1,3/2}} dt \\
 & \leq \int_0^1 \|\mathcal{E}''_0(\omega + p + \eta_2 + t(\eta_1 - \eta_2)) - \mathcal{E}''_0(\omega)\|_{W^{1,3/2}} dt \\
 & \quad + 2|\varepsilon| \int_0^1 \|\mathcal{V}''_H(\omega + p + \eta_2 + t(\eta_1 - \eta_2))\|_{W^{1,3/2}} dt \\
 & \leq \sup_{\|\eta\|_{W^{1,3}} \leq 3\rho} \|\mathcal{E}''_0(\omega + \eta) - \mathcal{E}''_0(\omega)\|_{W^{1,3/2}} + 2|\varepsilon| \sup_{\|\eta\|_{W^{1,3}} \leq 3\rho} \|\mathcal{V}''_H(\omega + p + \eta)\|_{W^{1,3/2}}.
 \end{aligned}$$

From (2.3), (2.4), and the Hölder inequality it follows that there exists a positive constant  $C_3$  such that for any  $\eta \in W^{1,3}$ ,  $p \in \mathbb{R}^3$

$$(3.7) \quad \|\mathcal{V}'_H(\omega + p + \eta)\|_{W^{1,3}} \leq C_3 \left[ \left( \int_{\mathbb{R}^2} |H(\omega + p + \eta)|^{3/2} |\nabla \omega|^3 \mu^{-1} \right)^{2/3} + \|\eta\|_{W^{1,3}}^2 \right]$$

and

$$\begin{aligned}
 (3.8) \quad \|\mathcal{V}''_H(\omega + p + \eta)\|_{W^{1,3/2}} & \leq C_3 \left[ \left( \int_{\mathbb{R}^2} |H(\omega + p + \eta)|^2 |\nabla(\omega + \eta)|^2 \right)^{1/2} \right. \\
 & \quad \left. + \left( \int_{\mathbb{R}^2} |\nabla H(\omega + p + \eta)|^{3/2} |\nabla(\omega + \eta)|^3 \mu^{-1} \right)^{2/3} \right].
 \end{aligned}$$

Choosing  $\rho_0 > 0$  such that

$$C_2 \sup_{\|\eta\|_{W^{1,3}} \leq 3\rho_0} \|\mathcal{E}''_0(\omega + \eta) - \mathcal{E}''_0(\omega)\|_{W^{1,3/2}} < \frac{1}{2}$$

and  $\varepsilon_0 > 0$  such that

$$(3.9) \quad 8C_2 C_3 \varepsilon_0 \|H\|_{L^\infty(\mathbb{R}^3)} \|\omega\|_{W^{1,3}}^2 < \min \left\{ 1, \rho_0, \frac{1}{8C_2 C_3 \varepsilon_0} \right\},$$

$$(3.10) \quad \sup_{\substack{\|\eta\|_{W^{1,3}} \leq \rho_0 \\ p \in \mathbb{R}^3}} \|\mathcal{V}'_H(\omega + p + \eta)\|_{W^{1,3}} < \frac{\rho_0}{6\varepsilon_0 C_2},$$

$$(3.11) \quad \sup_{\substack{\|\eta\|_{W^{1,3}} \leq 3\rho_0 \\ p \in \mathbb{R}^3}} \|\mathcal{V}''_H(\omega + p + \eta)\|_{W^{1,3/2}} < \frac{1}{8\varepsilon_0 C_2},$$

we obtain that  $T_{\varepsilon,p}$  maps the ball  $\overline{B_{\rho_0}(0)}$  into itself for any  $|\varepsilon| < \varepsilon_0$ ,  $p \in \mathbb{R}^3$ , and is a contraction there.

Hence it has a unique fixed point  $(\eta(\varepsilon, p), \lambda(\varepsilon, p), \alpha(\varepsilon, p)) \in \overline{B_{\rho_0}(0)}$ . From (3.6) we have that the following property holds

$$(*) \quad T_{\varepsilon, p} \text{ maps a ball } \overline{B_{\rho}(0)} \subset W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3 \text{ into itself whenever } \rho \leq \rho_0 \\ \text{and } \rho > 4|\varepsilon|C_2 \sup_{\|\eta\|_{W^{1,3}} \leq \rho} \|\mathcal{V}'_H(\omega + p + \eta)\|_{W^{1,3}}.$$

In particular let us set

$$(3.12) \quad \rho_\varepsilon = 5|\varepsilon|C_2 \sup_{\substack{\|\eta\|_{W^{1,3}} \leq \rho_0 \\ p \in \mathbb{R}^3}} \|\mathcal{V}'_H(\omega + p + \eta)\|_{W^{1,3}}.$$

In view of (3.10) and (3.12), we have that for any  $|\varepsilon| < \varepsilon_0$  and for any  $p \in \mathbb{R}^3$

$$\rho_\varepsilon \leq \rho_0 \quad \text{and} \quad \rho_\varepsilon > 4|\varepsilon|C_2 \sup_{\|\eta\|_{W^{1,3}} \leq \rho_\varepsilon} \|\mathcal{V}'_H(\omega + p + \eta)\|_{W^{1,3}}$$

so that, due to (\*),  $T_{\varepsilon, p}$  maps  $\overline{B_{\rho_\varepsilon}(0)}$  into itself. From the uniqueness of the fixed point we have that for any  $|\varepsilon| < \varepsilon_0$  and  $p \in \mathbb{R}^3$

$$(3.13) \quad \|(\eta(\varepsilon, p), \lambda(\varepsilon, p), \alpha(\varepsilon, p))\|_{W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3} \leq \rho_\varepsilon \leq C_1|\varepsilon|$$

for some positive constant  $C_1$  independent of  $p$  and hence  $\|\eta(\varepsilon, p)\|_{W^{1,3}} \leq \rho_\varepsilon \leq C_1|\varepsilon|$  thus proving (3.4). Assume now (3.5) and set for any  $p \in \mathbb{R}^3$

$$\rho_p = 8C_2C_3\varepsilon_0 \left( \int_{\mathbb{R}^2} \sup_{|q-p| \leq 1+C_0} |H(q)|^{3/2} |\nabla \omega|^3 \mu^{-1} \right)^{2/3}$$

where  $C_0$  is a positive constant such that  $\|u\|_{L^\infty} \leq C_0\|u\|_{W^{1,3}}$  for any  $u \in W^{1,3}$ . From (3.9) we have that

$$\rho_p < \min \left\{ 1, \rho_0, \frac{1}{8C_2C_3\varepsilon_0} \right\}.$$

Hence, due to (3.7), we have that for  $|\varepsilon| < \varepsilon_0$  and  $\|\eta\|_{W^{1,3}} \leq \rho_p$

$$4|\varepsilon|C_2 \|\mathcal{V}'_H(\omega + p + \eta)\|_{W^{1,3}} \\ \leq 4\varepsilon_0C_2C_3 \left( \int_{\mathbb{R}^2} \sup_{|q-p| \leq 1+C_0} |H(q)|^{3/2} |\nabla \omega|^3 \mu^{-1} \right)^{2/3} + 4\varepsilon_0C_2C_3\rho_p^2 < \rho_p.$$

From (\*) and the uniqueness of the fixed point, we deduce that

$$\|\eta(\varepsilon, p)\|_{W^{1,3}} \leq \rho_p$$

for any  $|\varepsilon| < \varepsilon_0$  and  $p \in \mathbb{R}^3$ . On the other hand, since  $H$  vanishes at  $\infty$ , by the definition of  $\rho_p$  we have that  $\rho_p \rightarrow 0$  as  $|p| \rightarrow \infty$ , hence

$$\lim_{|p| \rightarrow \infty} \eta(\varepsilon, p) = 0 \quad \text{in } W^{1,3} \text{ uniformly for } |\varepsilon| < \varepsilon_0.$$

The proof of Lemma 3.1 is now complete. ■

**Remark 3.2** *The map  $\eta$  given in Lemma 3.1 satisfies*

$$\langle \mathcal{E}'_\varepsilon(\omega + p + \eta(\varepsilon, p)), \varphi \rangle - \sum_{i=1}^6 \lambda_i(\varepsilon, p) \int_{\mathbb{R}^2} \nabla \varphi \cdot \nabla \tau_i + \alpha(\varepsilon, p) \cdot \int_{\mathbb{S}^2} \varphi, \quad \forall \varphi \in W^{1,3/2}$$

where  $(\eta(\varepsilon, p), \lambda(\varepsilon, p), \alpha(\varepsilon, p)) \in \overline{B_{\rho_\varepsilon}(0)} \subset W^{1,3} \times \mathbb{R}^6 \times \mathbb{R}^3$  being  $\rho_\varepsilon$  given in (3.12), hence

$$\begin{aligned} & \int_{\mathbb{R}^2} \nabla(\omega + \eta(\varepsilon, p)) \cdot \nabla \varphi + 2 \int_{\mathbb{R}^2} \varphi \cdot (\omega + \eta(\varepsilon, p))_x \wedge (\omega + \eta(\varepsilon, p))_y \\ & + 2\varepsilon \int_{\mathbb{R}^2} H(\omega + p + \eta(\varepsilon, p)) \varphi \cdot (\omega + \eta(\varepsilon, p))_x \wedge (\omega + \eta(\varepsilon, p))_y \\ & = \sum_{i=1}^6 \lambda_i(\varepsilon, p) \int_{\mathbb{R}^2} \nabla \varphi \cdot \nabla \tau_i - \alpha(\varepsilon, p) \cdot \int_{\mathbb{S}^2} \varphi, \quad \forall \varphi \in W^{1,3/2}, \end{aligned}$$

i.e.  $\eta(\varepsilon, p)$  satisfies the equation

$$\Delta \eta(\varepsilon, p) = F(\varepsilon, p)$$

where

$$\begin{aligned} F(\varepsilon, p) &= 2(\omega + \eta(\varepsilon, p))_x \wedge (\omega + \eta(\varepsilon, p))_y - 2\omega_x \wedge \omega_y + \lambda(\varepsilon, p) \cdot \Delta \tau \\ & - \alpha(\varepsilon, p) \mu^2 + 2\varepsilon H(\omega + p + \eta(\varepsilon, p)) (\omega + \eta(\varepsilon, p))_x \wedge (\omega + \eta(\varepsilon, p))_y \end{aligned}$$

in  $\mathbb{R}^2$ . Since  $F(\varepsilon, p) \in L^{3/2}$  and, in view of (3.4) and (3.13),  $F(\varepsilon, p) \rightarrow 0$  in  $L^{3/2}$  as  $\varepsilon \rightarrow 0$  uniformly with respect to  $p$ , by regularity we have that

$$\eta(\varepsilon, p) \in W^{2,3/2} \quad \text{and} \quad \eta(\varepsilon, p) \rightarrow 0 \quad \text{in} \quad W^{2,3/2}$$

hence, by Sobolev embeddings,  $F(\varepsilon, p) \in L^3$  and  $F(\varepsilon, p) \rightarrow 0$  in  $L^3$  as  $\varepsilon \rightarrow 0$  uniformly with respect to  $p$ . Again by regularity

$$\eta(\varepsilon, p) \in W^{2,3} \quad \text{and} \quad \eta(\varepsilon, p) \rightarrow 0 \quad \text{in} \quad W^{2,3}$$

hence  $\eta(\varepsilon, p) \in C^{1,1/3}$  and

$$(3.14) \quad \eta(\varepsilon, p) \rightarrow 0 \quad \text{in} \quad C^{1,1/3} \quad \text{as} \quad \varepsilon \rightarrow 0 \quad \text{uniformly with respect to } p.$$

For any  $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$ , let us define the perturbed manifold

$$Z_\varepsilon := \{\omega + p + \eta(\varepsilon, p) : p \in \mathbb{R}^3\}.$$

From [7], we have that  $Z_\varepsilon$  is a natural constraint for  $\mathcal{E}_\varepsilon$ , namely any critical point  $p \in \mathbb{R}^3$  of the functional

$$\Phi_\varepsilon : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad \Phi_\varepsilon(p) = \mathcal{E}_\varepsilon(\omega + p + \eta(\varepsilon, p))$$

gives rise to a critical point  $u_\varepsilon = \omega + p + \eta(\varepsilon, p)$  of  $\mathcal{E}_\varepsilon$ .

**Proposition 3.3** *Assume  $H \in C^2(\mathbb{R}^3)$ ,  $\nabla H \in L^\infty(\mathbb{R}^3, \mathbb{R}^3)$ , and*

$$\lim_{|p| \rightarrow \infty} \bar{H}(p) = 0.$$

*Then for any  $|\varepsilon| < \varepsilon_0$*

$$\lim_{|p| \rightarrow \infty} \Phi_\varepsilon(p) = \text{const} = \mathcal{E}_0(\omega).$$

**Proof.** We have that

$$\begin{aligned}
\Phi_\varepsilon(p) &= \mathcal{E}_\varepsilon(\omega + p + \eta(\varepsilon, p)) \\
&= \mathcal{E}_0(\omega + p + \eta(\varepsilon, p)) + 2\varepsilon \mathcal{V}_H(\omega + p + \eta(\varepsilon, p)) \\
&= \frac{1}{2} \int_{\mathbb{R}^2} |\nabla \omega|^2 + \frac{1}{2} \int_{\mathbb{R}^2} |\nabla \eta(\varepsilon, p)|^2 + \int_{\mathbb{R}^2} \nabla \omega \cdot \nabla \eta(\varepsilon, p) \\
&\quad + \frac{2}{3} \int_{\mathbb{R}^2} (\omega + p + \eta(\varepsilon, p)) \cdot (\omega + \eta(\varepsilon, p))_x \wedge (\omega + \eta(\varepsilon, p))_y \\
&\quad + 2\varepsilon [\mathcal{V}_H(\omega + p) + \langle \mathcal{V}'_H(\omega + p), \eta(\varepsilon, p) \rangle + o(\|\eta(\varepsilon, p)\|_{W^{1,3}})] \\
&= \frac{1}{2} \int_{\mathbb{R}^2} |\nabla \omega|^2 + \frac{2}{3} \int_{\mathbb{R}^2} \omega \cdot \omega_x \wedge \omega_y + \frac{1}{2} \int_{\mathbb{R}^2} |\nabla \eta(\varepsilon, p)|^2 \\
&\quad + \int_{\mathbb{R}^2} \nabla \omega \cdot \nabla \eta(\varepsilon, p) + \frac{2}{3} \int_{\mathbb{R}^2} \omega \cdot (\omega_x \wedge \eta(\varepsilon, p)_y + \eta(\varepsilon, p)_x \wedge \omega_y) \\
&\quad + \frac{2}{3} \int_{\mathbb{R}^2} \omega \cdot \eta(\varepsilon, p)_x \wedge \eta(\varepsilon, p)_y \\
&\quad + \frac{2}{3} \int_{\mathbb{R}^2} \eta(\varepsilon, p) \cdot (\omega + \eta(\varepsilon, p))_x \wedge (\omega + \eta(\varepsilon, p))_y \\
(3.15) \quad &+ 2\varepsilon \mathcal{V}_H(\omega + p) + 2\varepsilon \langle \mathcal{V}'_H(\omega + p), \eta(\varepsilon, p) \rangle + 2\varepsilon o(\|\eta(\varepsilon, p)\|_{W^{1,3}})
\end{aligned}$$

where we have used the fact that

$$\int_{\mathbb{R}^2} p \cdot u_x \wedge u_y = 0 \quad \forall p \in \mathbb{R}^3, u \in W^{1,3},$$

(see [7], Lemma A.3). Notice that from Lemma 3.1 we have that

$$(3.16) \quad \int_{\mathbb{R}^2} |\nabla \eta(\varepsilon, p)|^2 \leq \sqrt[3]{4\pi} \|\eta(\varepsilon, p)\|_{W^{1,3}}^2 \xrightarrow{|p| \rightarrow \infty} 0,$$

$$(3.17) \quad \left| \int_{\mathbb{R}^2} \nabla \omega \cdot \nabla \eta(\varepsilon, p) \right| \leq \sqrt[6]{4\pi} \left( \int_{\mathbb{R}^2} |\nabla \omega|^2 \right)^{1/2} \|\eta(\varepsilon, p)\|_{W^{1,3}} \xrightarrow{|p| \rightarrow \infty} 0,$$

and, by the Hölder inequality and Lemma 3.1,

$$(3.18) \quad \left| \int_{\mathbb{R}^2} \omega \cdot (\omega_x \wedge \eta(\varepsilon, p)_y + \eta(\varepsilon, p)_x \wedge \omega_y) \right| \leq 2 \|\omega\|_{W^{1,3}}^2 \|\eta(\varepsilon, p)\|_{W^{1,3}} \xrightarrow{|p| \rightarrow \infty} 0,$$

$$(3.19) \quad \left| \int_{\mathbb{R}^2} \omega \cdot (\eta(\varepsilon, p)_x \wedge \eta(\varepsilon, p)_y) \right| \leq \|\omega\|_{W^{1,3}} \|\eta(\varepsilon, p)\|_{W^{1,3}}^2 \xrightarrow{|p| \rightarrow \infty} 0,$$

and

$$(3.20) \quad \left| \int_{\mathbb{R}^2} \eta(\varepsilon, p) \cdot (\omega + \eta(\varepsilon, p))_x \wedge (\omega + \eta(\varepsilon, p))_y \right| \leq \|\omega + \eta(\varepsilon, p)\|_{W^{1,3}}^2 \|\eta(\varepsilon, p)\|_{W^{1,3}} \xrightarrow{|p| \rightarrow \infty} 0.$$

Moreover the Gauss-Green Theorem yields

$$\mathcal{V}_H(\omega + p) = - \int_{B_1} H(\xi + p) d\xi$$

so that by the Dominated Convergence Theorem we have that

$$(3.21) \quad \lim_{|p| \rightarrow \infty} \mathcal{V}_H(\omega + p) = 0.$$

From (2.3), Hölder's inequality, and Lemma 3.1, we have that

$$(3.22) \quad \begin{aligned} |\langle \mathcal{V}'_H(\omega + p), \eta(\varepsilon, p) \rangle| &= \left| \int_{\mathbb{R}^2} H(\omega + p) \eta(\varepsilon, p) \cdot \omega_x \wedge \omega_y \right| \\ &\leq \|H\|_{L^\infty(\mathbb{R}^3)} \|\omega\|_{W^{1,3}}^2 \|\eta(\varepsilon, p)\|_{W^{1,3}} \xrightarrow{|p| \rightarrow \infty} 0. \end{aligned}$$

From (3.15)–(3.22), it follows that

$$\lim_{|p| \rightarrow \infty} \Phi_\varepsilon(p) = \frac{1}{2} \int_{\mathbb{R}^2} |\nabla \omega|^2 + \frac{2}{3} \int_{\mathbb{R}^2} \omega \cdot \omega_x \wedge \omega_y = \mathcal{E}_0(\omega).$$

The proposition is thereby proved. ■

**Proof of Theorem 1.1.** As already observed at the beginning of Section 2, it is not restrictive to take  $H_0 = 1$ . From Proposition 3.3 it follows that for  $|\varepsilon| < \varepsilon_0$  either  $\Phi_\varepsilon$  is constant (and hence we have infinitely many critical points) or it has a global maximum or minimum point. In any case  $\Phi_\varepsilon$  has a critical point. Since  $Z_\varepsilon$  is a natural constraint for  $\mathcal{E}_\varepsilon$ , we deduce the existence of a critical point of  $\mathcal{E}_\varepsilon$  for  $|\varepsilon| < \varepsilon_0$  and hence of a solution to  $(P_\varepsilon)$ . The  $H_\varepsilon$ -bubble  $\omega_\varepsilon$  found in this way is of the form  $\omega + p^\varepsilon + \eta(\varepsilon, p^\varepsilon)$  for some  $p^\varepsilon \in \mathbb{R}^3$  where  $\eta$  is as in Lemma 3.1. Remark 3.2 yields that  $\omega_\varepsilon$  is closed in  $C^{1,1/3}(\mathbb{S}^2, \mathbb{R}^3)$ -norm to the manifold  $\{\omega + p : p \in \mathbb{R}^3\}$  for  $\varepsilon$  small. Since  $\omega$  has no branch points, we deduce that  $\omega_\varepsilon$  has no branch points. ■

To prove Theorems 1.2 and 1.4, we need the following expansion for  $\Phi_\varepsilon$ :

$$(3.23) \quad \Phi_\varepsilon(p) = \mathcal{E}_0(\omega) - 2\varepsilon\Gamma(p) + O(\varepsilon^2)$$

as  $\varepsilon \rightarrow 0$  uniformly in  $p \in \mathbb{R}^3$  (see [7]).

**Proof of Theorem 1.2.** Let  $\varepsilon > 0$  small. Assumption (H4) implies that  $\Gamma(0) > 0$  and hence from (3.23) we have that for  $\varepsilon$  small

$$\Phi_\varepsilon(0) < \mathcal{E}_0(\omega),$$

whereas from assumption (H3) we have that  $\text{Hess}\Gamma(0)$  is positive definite so that  $\Gamma$  has a strict local minimum in 0 and hence from (3.23)  $\Phi_\varepsilon$  has a strict local maximum in  $B_r(0)$  for some  $r > 0$  such that

$$\Phi_\varepsilon(p) < \Phi_\varepsilon(0) - c_\varepsilon < \mathcal{E}_0(\omega)$$

for  $|p| = r$ , where  $c_\varepsilon$  is some positive constant depending on  $\varepsilon$ . In particular  $\Phi_\varepsilon$  has a mountain pass geometry. Moreover by Theorem 1.1,

$$\Phi_\varepsilon(p) \rightarrow \mathcal{E}_0(\omega) \quad \text{as } |p| \rightarrow \infty,$$

and so  $\Phi_\varepsilon$  must have a global minimum point. If the minimum point and the mountain pass point coincide then  $\Phi_\varepsilon$  has infinitely many critical points. Otherwise  $\Phi_\varepsilon$  has at least three critical points: a local maximum point, a global minimum point, and a mountain pass. If  $\varepsilon < 0$  we find the inverse inequalities and hence we find that  $\Phi_\varepsilon$  has a local minimum point, a global maximum point, and a mountain pass. As a consequence  $(P_\varepsilon)$  has at least three solutions provided  $|\varepsilon|$  is sufficiently small. ■

As observed in Remark 1.3, if  $H_1(0) > 0$  and  $\text{Hess}H_1(0)$  is positive definite, by continuity we have that for  $H_0$  sufficiently large  $\Gamma(0) > 0$  and  $\text{Hess}\Gamma(0)$  is positive definite, so that we can still prove the existence of three solutions arguing as above.

**Proof of Theorem 1.4.** Assumption (H5) implies that  $\Gamma(p_1) > 0$  and  $\Gamma(p_2) < 0$ . Since

$$\Phi_\varepsilon(p) = \mathcal{E}_0(\omega) + 2\varepsilon(-\Gamma(p) + o(1)) \quad \text{as } \varepsilon \rightarrow 0,$$

we have for  $\varepsilon$  sufficiently small

$$\Phi_\varepsilon(p_1) < \mathcal{E}_0(\omega) \quad \text{and} \quad \Phi_\varepsilon(p_2) > \mathcal{E}_0(\omega)$$

if  $\varepsilon > 0$  and the inverse inequalities if  $\varepsilon < 0$ . Since by Theorem 1.1

$$\Phi_\varepsilon(p) \rightarrow \mathcal{E}_0(\omega) \quad \text{as } |p| \rightarrow \infty,$$

we conclude that  $\Phi_\varepsilon$  must have a global maximum point and a global minimum point in  $\mathbb{R}^3$ . Since  $Z_\varepsilon$  is a natural constraint for  $\mathcal{E}_\varepsilon$ , we deduce the existence of two critical points of  $\mathcal{E}_\varepsilon$  for  $|\varepsilon|$  sufficiently small and hence of two solutions to  $(P_\varepsilon)$ . ■

As observed in Remark 1.5, if  $H_1(p_1) > 0$  and  $H_1(p_2) < 0$ , by continuity we have that for  $H_0$  sufficiently large  $\Gamma(p_1) > 0$  and  $\Gamma(p_2) < 0$ , so that we can still prove the existence of two solutions arguing as above.

## References

- [1] AMBROSETTI, A. AND BADIALE, M.: Homoclinics: Poincaré-Melnikov type results via a variational approach. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **15** (1998), no. 2, 233–252.
- [2] AMBROSETTI, A. AND BADIALE, M.: Variational perturbative methods and bifurcation of bound states from the essential spectrum. *Proc. Roy. Soc. Edinburgh Sect. A* **128** (1998), 1131–1161.
- [3] BREZIS, H. AND CORON, J. M.: Multiple solutions of  $H$ -systems and Rellich’s conjecture. *Comm. Pure Appl. Math.* **37** (1984), no. 2, 149–187.
- [4] BREZIS, H. AND CORON, J. M.: Convergence of solutions of  $H$ -systems or how to blow bubbles. *Arch. Rational Mech. Anal.* **89** (1985), no. 1, 21–56.
- [5] CALDIROLI, P. AND MUSINA, R.: Existence of minimal  $H$ -bubbles. *Commun. Contemp. Math.* **4** (2002), no. 2, 177–210.
- [6] CALDIROLI, P. AND MUSINA, R.: Existence of  $H$ -bubbles in a perturbative setting. *Rev. Mat. Iberoamericana* **20** (2004), no. 2, 611–626.
- [7] CALDIROLI, P. AND MUSINA, R.:  $H$ -bubbles in a perturbative setting: the finite-dimensional reduction method. *Duke Math. J.* **122** (2004), no. 3, 457–484.
- [8] CHANILLO, S. AND MALCHIODI, A.: Asymptotic Morse theory for the equation  $\Delta u = 2u_x \wedge u_y$ . *Comm. Anal. Geom.* **13** (2005), no. 1, 187–251.
- [9] GRÜTER, M.: Regularity of weak  $H$ -surfaces. *J. Reine Angew. Math.* **329** (1981), 1–15.
- [10] HILDEBRANDT, S.: Randwertprobleme für Flächen mit vorgeschriebener mittlerer Krümmung und Anwendungen auf die Kapillaritätstheorie. I: Fest vorgegebener Rand. *Math. Z.* **112** (1969), 205–213.
- [11] ISOBE, T.: On the asymptotic analysis of  $H$ -systems, I: asymptotic behavior of large solutions. *Adv. Differential Equations* **6** (2001), no. 5, 513–546.
- [12] JAKOBOWSKY, N.: A perturbation result concerning a second solution to the Dirichlet problem for the equation of the prescribed mean curvature. *J. Reine Angew. Math.* **457** (1994), 1–21.
- [13] JAKOBOWSKY, N.: Multiple surfaces of non-constant mean curvature. *Math. Z.* **217** (1994), no. 3, 497–512.

- [14] STEFFEN, K.: On the existence of surfaces with prescribed mean curvature and boundary. *Math. Z.* **146** (1976), no. 2, 113–135.
- [15] STRUWE, M.: Large H-surface via the mountain-pass lemma. *Math. Ann.* **270** (1985), no. 3, 441–459.
- [16] STRUWE, M.: *Plateau's problem and the calculus of variations*. Mathematical Notes **35**. Princeton University Press, Princeton, NJ, 1988.

*Recibido:* 29 de enero de 2003.

Veronica Felli  
Dipartimento di Matematica e Applicazioni  
Università di Milano-Bicocca  
Via Cozzi 53  
20125 Milano, Italy  
`veronica.felli@unimib.it`

---

The author wishes to thank Professor A. Ambrosetti and Professor R. Musina for many helpful suggestions. Supported by MIUR, national project “Variational Methods and Nonlinear Differential Equations”.