

# Dyadic BMO on the bidisk

Óscar Blasco and Sandra Pott

## Abstract

We give several new characterizations of the dual of the dyadic Hardy space  $H^{1,d}(\mathbb{T}^2)$ , the so-called dyadic BMO space in two variables and denoted  $\text{BMO}_{\text{prod}}^d$ . These include characterizations in terms of Haar multipliers, in terms of the “symmetrised paraproduct”  $\Lambda_b$ , in terms of the rectangular BMO norms of the iterated “sweeps”, and in terms of nested commutators with dyadic martingale transforms. We further explore the connection between  $\text{BMO}_{\text{prod}}^d$  and John-Nirenberg type inequalities, and study a scale of rectangular BMO spaces.

## 1. Introduction

Throughout the paper  $\mathcal{D}$  denotes the set of dyadic intervals in the unit circle  $\mathbb{T}$ . In the case of the bicircle  $\mathbb{T}^2$ ,  $\mathcal{D}_1$  denotes the dyadic intervals in the first,  $\mathcal{D}_2$  the dyadic intervals in the second variable. We write  $\mathcal{R} = \mathcal{D}_1 \times \mathcal{D}_2$  for the dyadic rectangles,  $|I|$  for the length of  $I \in \mathcal{D}$  and  $|R|$  for the area of  $R \in \mathcal{R}$ ,  $(h_I)_{I \in \mathcal{D}}$  stands for the Haar basis in  $L^2(\mathbb{T})$  and  $(h_R)_{R \in \mathcal{R}}$  for the product Haar basis of  $L^2(\mathbb{T}^2)$ .

Here

$$h_I(t) = \frac{1}{|I|^{1/2}} (\chi_{I^+}(t) - \chi_{I^-}(t))$$

for each dyadic interval  $I \in \mathcal{D}$ , where  $I^-$  denotes the left half of  $I$ , and  $I^+$  denotes the right half of  $I$ . For each dyadic rectangle  $R = I \times J \in \mathcal{R}$ ,  $h_R$  is defined by

$$h_R(s, t) = h_I(s)h_J(t).$$

We denote by  $\mathcal{H}_{00}$  the space of all functions in  $L^2(\mathbb{T}^2)$  which have a finite expansion in the product Haar basis.

---

*2000 Mathematics Subject Classification:* 42B30, 47B35.

*Keywords:* BMO on the bidisk, Carleson measures, Haar multipliers.

Given  $g \in L^2(\mathbb{T})$ , we use the notation  $g_I = \langle g, h_I \rangle$  and  $m_I g = \frac{1}{|I|} \int_I g(t) dt$ . Similarly, given  $f \in L^2(\mathbb{T}^2)$ , we use the notation  $f_R = \langle f, h_R \rangle$ ,  $f_I(s) = \langle f(\cdot, s), h_I \rangle$ ,  $m_I f(s) = \frac{1}{|I|} \int_I f(t, s) dt$ ,  $f_J(t) = \langle f(t, \cdot), h_J \rangle$  and  $m_J f(t) = \frac{1}{|J|} \int_J f(t, s) ds$ . Therefore

$$f(t, s) = \sum_{R \in \mathcal{R}} f_R h_R(t, s) = \sum_{I \in \mathcal{D}} f_I(s) h_I(t) = \sum_{J \in \mathcal{D}} f_J(t) h_J(s).$$

Let  $P_I g = (g - m_I g) \chi_I$  for  $g \in L^2(\mathbb{T})$ . Observe that  $P_I$  is the orthogonal projection on the subspace spanned by the Haar functions  $h_{I'}$ ,  $I' \in \mathcal{D}$ ,  $I' \subseteq I$ . If  $g = \sum_{I \in \mathcal{D}} g_I h_I$ , then

$$(1.1) \quad P_I g = \sum_{I' \in \mathcal{D}, I' \subseteq I} h_{I'} g_{I'}.$$

Similarly, for each measurable set  $\Omega \subseteq \mathbb{T}^2$ , let  $P_\Omega$  be the orthogonal projection on the subspace spanned by the Haar functions  $h_{R'}$ ,  $R' \in \mathcal{R}$ ,  $R' \subseteq \Omega$ . In particular, for each dyadic rectangle  $R = I \times J \in \mathcal{R}$  and for  $f = \sum_{R' \in \mathcal{R}} h_{R'} f_{R'} \in L^2(\mathbb{T}^2)$ , we have

$$P_R f = P_I \otimes P_J f = \sum_{R' \in \mathcal{R}, R' \subseteq R} h_{R'} f_{R'}.$$

It is easy to see that for  $R \in \mathcal{R}$  and  $f \in L^2(\mathbb{T}^2)$ ,

$$(1.2) \quad P_R f = (f - m_I f - m_J f + m_{I \times J}) \chi_{I \times J}.$$

Recall that  $g \in L^2(\mathbb{T})$  is said to belong to dyadic BMO, to be denoted  $\text{BMO}^d(\mathbb{T})$ , if

$$\sup_{I \in \mathcal{D}} \left( \frac{1}{|I|} \int_I |g(t) - m_I g|^2 dt \right)^{1/2} < \infty.$$

By John-Nirenberg's lemma, this is equivalent to

$$\sup_{I \in \mathcal{D}} \left( \frac{1}{|I|} \int_I |g(t) - m_I g|^p dt \right)^{1/p} < \infty$$

for any  $1 \leq p < \infty$ .

Hence  $g \in \text{BMO}^d(\mathbb{T})$  if and only if there exists a constant  $C$  such that for all  $I \in \mathcal{D}$

$$\sum_{I' \in \mathcal{D}, I' \subseteq I} |g_{I'}|^2 \leq C |I|^{1/2},$$

or equivalently

$$\sup_{I \in \mathcal{D}} \frac{1}{|I|^{1/p}} \|P_I g\|_p < \infty$$

for  $1 \leq p < \infty$ .

The space BMO appears in many different contexts. We shall use that  $\text{BMO}^d(\mathbb{T}) = (H^{1,d}(\mathbb{T}))^*$  where  $H^{1,d}$  is defined in terms of the dyadic square functions  $\mathcal{S}$ ,

$$\mathcal{S}g = \left( \sum_{I \in \mathcal{D}} \frac{\chi_I}{|I|} |g_I|^2 \right)^{1/2}.$$

That is,

$$H^{1,d}(\mathbb{T}) = \{g \in L^1(\mathbb{T}) : \mathcal{S}g \in L^1(\mathbb{T})\}.$$

Using Carleson measures, this gives rise to a description of  $\text{BMO}^d$  in terms of symbols  $g$  for which the dyadic paraproduct  $\pi_g$ ,

$$\pi_g(f) = \sum_{I \in \mathcal{D}} g_I m_I f h_I$$

or its adjoint operator  $\Delta_g$ ,  $\Delta_g(f) = \sum_{I \in \mathcal{D}} g_I f_I \frac{\chi_I}{|I|}$ , is bounded on  $L^2(\mathbb{T})$  (or equivalently, on  $L^p(\mathbb{T})$  for  $1 < p < \infty$ ).

The situation in two variables it is rather different and much more delicate. One main reason for the difficulties encountered in the multi-variable theory is the failure of the naive generalization of the Carleson Embedding Theorem to several variables (see [C], [Fef]). The reader is referred to [ChFef2] for an overview on the theory and an outline of the main differences.

Several new results (e. g. [FS] and [PS]) further exhibit the differences between certain BMO spaces on the polydisk defined by multi-variable versions of the different yet equivalent characterizations of  $\text{BMO}(\mathbb{T})$ .

A function  $f \in L^2(\mathbb{T}^2)$  is said to belong to the rectangular dyadic BMO space, to be denoted  $\text{BMO}_{\text{rect}}^d$ , if

$$(1.3) \quad \sup_{R=I \times J \in \mathcal{R}} \left( \frac{1}{|R|} \int_R |f(t, s) - m_I f(s) - m_J f(t) + m_{I \times J} f|^2 dt ds \right)^{1/2} < \infty.$$

Or equivalently,

$$\|\varphi\|_{\text{BMO}_{\text{rect}}^d} = \sup_{R \in \mathcal{R}} \frac{1}{|R|^{1/2}} \|P_R \varphi\|_2.$$

We will also consider a  $p$ -version of the dyadic rectangular norm for  $1 \leq p < \infty$ ,

$$(1.4) \quad \|\varphi\|_{\text{BMO}_{\text{rect},p}^d} = \sup_{R \in \mathcal{R}} \frac{1}{|R|^{1/p}} \|P_R \varphi\|_p.$$

Here,  $\|\cdot\|_{\text{BMO}_{\text{rect}}^d} = \|\cdot\|_{\text{BMO}_{\text{rect},2}^d}$ . In the one-variable case, the corresponding norms are of course all equivalent because of John-Nirenberg's lemma.

Let us start by defining  $BMO_{\text{prod}}^d(\mathbb{T}^2)$  as the dual of  $H^{1,d}(\mathbb{T}^2)$ , the space of functions  $f \in L^1(\mathbb{T}^2)$  such that  $\mathcal{S}(f) \in L^1(\mathbb{T}^2)$ , where

$$\mathcal{S}(f) = \left( \sum_{R \in \mathcal{R}} |f_R|^2 h_R^2 \right)^{1/2}.$$

Although  $BMO_{\text{prod}}^d(\mathbb{T}^2)$  cannot be characterized by (1.3) [Fef], it was shown by Bernard in the dyadic case [Be] and also by Chang and R. Fefferman in a continuous version [ChFef1] that  $BMO_{\text{prod}}^d(\mathbb{T}^2)$  can also be described as the space of functions  $\varphi \in L^2(\mathbb{T}^2)$  for which there exists  $C > 0$  such that

$$(1.5) \quad \|\varphi\|_{\text{prod}} = \sup_{\Omega \subset \mathbb{T}^2} \frac{1}{|\Omega|^{1/2}} \|P_{\Omega} \varphi\|_2 < \infty,$$

where the supremum is taken over all measurable sets  $\Omega \subseteq \mathbb{T}^2$ . This immediately implies  $BMO_{\text{prod}}^d \subseteq BMO_{\text{rect},2}$ .

The connection between both spaces can be also seen from the description of  $BMO_{\text{prod}}^d$  in terms of the boundedness of the dyadic paraproduct in two variables, defined by  $\pi_b^{(1,2)}(f) = \sum_{R \in \mathcal{R}} b_R m_R f h_R$ .

It follows from Chang’s generalization of the Carleson Embedding Theorem (see [Ch]) that  $b \in BMO_{\text{prod}}^d$  if and only if the double paraproduct  $\pi_b^{(1,2)}$  is bounded on  $L^2(\mathbb{T}^2)$ . In our paper the following fact will be rather crucial:

$$(1.6) \quad \|\varphi\|_{\text{prod}} \approx \|\pi_{\varphi}^{(1,2)}\|.$$

An similar characterization for  $BMO_{\text{rect}}^d$  was proved in [PS], Proposition 3.3.1, namely that  $b \in BMO_{\text{rect}}^d$  if and only if  $\pi_b^{(1,2)}$  maps  $L^2(\mathbb{T}) \hat{\otimes} L^2(\mathbb{T})$  boundedly into  $L^2(\mathbb{T}^2)$ , where  $L^2(\mathbb{T}) \hat{\otimes} L^2(\mathbb{T})$  stands for the projective tensor product. This also implies that  $BMO_{\text{prod}}^d \subsetneq BMO_{\text{rect}}^d$  (see [Fef] for an alternative approach).

We shall try to better understand the difference between both spaces. Two approaches are used to this end. First we observe that John-Nirenberg type inequalities do not hold in  $BMO_{\text{rect}}^d$ , in the sense that the 2-norm in the definition of  $BMO_{\text{rect}}^d$  cannot be replaced by any other  $p$ -norm. This solves a question left open in [FS]. Secondly, we analyse the behaviour of the sweep of functions in the  $BMO_{\text{prod}}^d$  and in the  $BMO_{\text{rect},p}^d$  spaces.

Our main new tool will be characterizations of  $BMO_{\text{prod}}^d$  in terms of Haar multipliers. Recall that sequence of functions  $(\phi_R)_{R \in \mathcal{R}}$  is called a Haar multiplier (see (2.16) or [Per]) on  $L^p(\mathbb{T}^2)$ , if the map

$$f = \sum_{R \in \mathcal{R}} f_R h_R(t, s) \mapsto \sum_{R \in \mathcal{R}} \phi_R(t, s) f_R h_R(t, s)$$

defines a bounded operator on  $L^p(\mathbb{T}^2)$ .

We shall say that  $b \in \text{BMO}_{\text{mult}}^d$  if  $\{(P_R b)\}_{R \in \mathcal{R}}$  defines a Haar multiplier on  $L^2(\mathbb{T}^2)$ .

Using the characterization of  $\text{BMO}_{\text{prod}}^d$  in terms of dyadic paraproducts, we observe that  $b \in \text{BMO}_{\text{prod}}^d$  if the operator  $\Delta_b = (\pi_b^{(1,2)})^*$  given by  $\Delta_b(f) = \sum_R b_R h_R f_R h_R$  defines a bounded operator on  $L^2(\mathbb{T}^2)$ . Hence  $b \in \text{BMO}_{\text{prod}}^d$  if and only if  $(b_R h_R)_{R \in \mathcal{R}}$  is a Haar multiplier on  $L^2(\mathbb{T}^2)$ .

On the other hand, letting the Haar multiplier  $(P_{R'} b)_{R' \in \mathcal{R}}$  act on  $h_R$ , we see that  $\frac{1}{|R|^{1/2}} \|P_{R'} b\|_2 \leq \|b\|_{\text{mult}}$ , implying that  $\text{BMO}_{\text{mult}}^d \subseteq \text{BMO}_{\text{rect}}^d$ .

We shall get a description of  $\text{BMO}_{\text{mult}}^d$  in terms of the boundedness of the operator

$$\Lambda_b = \pi_b^{(1,2)} + (\pi_b^{(1,2)})^* + \Delta_{\pi_b} + (\Delta_{\pi_b})^*,$$

where  $\Delta_{\pi_b}$  (see Definition 2.4) is an operator combining the one-variable paraproduct  $\pi$  and its adjoint. This will allow us to prove that  $\text{BMO}_{\text{mult}}^d(\mathbb{T}^2) = \text{BMO}_{\text{prod}}^d(\mathbb{T}^2)$  (see Theorem 2.8).

On the other hand,  $\text{BMO}_{\text{rect}}^d$  can also be described using  $\Lambda_b$ . We show that  $\text{BMO}_{\text{rect}}^d$  can be characterized in terms of ‘‘average boundedness’’ of  $\Lambda_b$  or in terms of its boundedness from  $L^2(\mathbb{T}) \hat{\otimes} L^2(\mathbb{T})$  into  $L^2(\mathbb{T}^2)$ .

The paper is divided into three sections. The first one is devoted to the introduction of the space  $\text{BMO}_{\text{mult}}^d$  and the proof of some of its properties. We see that  $\text{BMO}_{\text{mult}}^d$  can be characterized as the space of symbols  $b$  for which the operator  $\Lambda_b$  is bounded, and that this space coincides with  $\text{BMO}_{\text{prod}}^d$ .

Section 3 deals with results on sweep functions. We prove the following formula connecting the boundedness of  $\pi_b^{(1,2)}$  and  $\Lambda_{S_b}$  (see Lemma 3.2):

$$(1.7) \quad \pi_b^{(1,2)*} \pi_b^{(1,2)} = \Lambda_{S_b} + D_b,$$

where  $D_b$  is bounded if  $b \in \text{BMO}_{\text{rect},2}^d$ .

This allows us to see that  $b \in \text{BMO}_{\text{prod}}^d$  if and only if  $S_b \in \text{BMO}_{\text{prod}}^d$ . We also obtain a characterization of  $\text{BMO}_{\text{prod}}^d$  in terms of nested commutators with dyadic martingale transforms, sharpening a result from [PS].

We further use the formula (1.7) to quantify the difference between the BMO spaces we have considered, and to get a characterization of  $\text{BMO}_{\text{prod}}^d$  relying only upon the  $\text{BMO}_{\text{rect}}^d$  norm of the  $n$ -fold sweeps.

Finally, in the last section, we apply the results from the previous ones together with interpolation to study the scale of spaces  $\text{BMO}_{\text{rect},p}^d$  introduced in (1.4) and show that these spaces are pairwise distinct. As a corollary, we obtain that  $\text{BMO}_{\text{prod}}^d \subsetneq \bigcap_{p \geq 1} \text{BMO}_{\text{rect},p}^d$ .

## 2. BMO via Haar multipliers

**Definition 2.1** We shall say that  $b \in \text{BMO}_{\text{mult}}^d$  if  $\{(P_R b)\}_{R \in \mathcal{R}}$  defines a Haar multiplier on  $L^2(\mathbb{T}^2)$ , i.e. there exists  $C > 0$  such that

$$\left\| \sum_{R \in \mathcal{R}} P_R b f_R h_R \right\|_2 \leq C \|f\|_2$$

for all  $f \in L^2(\mathbb{T}^2)$ . We define  $\|b\|_{\text{mult}}$  as the norm of the corresponding operator.

Let us start by pointing out some simple facts about this space.

Given  $I \in \mathcal{D}$  we write  $P_I$  for the operator on  $L^2(\mathbb{T})$  given by (1.1), and  $\tilde{P}_I = P_I \otimes \text{id}$  for the corresponding projection on  $L^2(\mathbb{T}^2)$ ,

$$\tilde{P}_I(f)(t, s) = \sum_{I' \in \mathcal{D}, I' \subseteq I} h_{I'}(t) f_{I'}(s).$$

Similarly, given  $J \in \mathcal{D}_2$ , we write  $\tilde{P}_J$  for  $\text{id} \otimes P_J$ .

Of course,  $\tilde{P}_I(f)(t, s) = P_I(f(s, \cdot))(t)$  and  $P_R f = \tilde{P}_J(\tilde{P}_I f)$  for  $R = I \times J$ .

### Proposition 2.2

$$(2.1) \quad L^\infty(\mathbb{T}^2) \subset \text{BMO}_{\text{mult}}^d(\mathbb{T}^2)$$

$$(2.2) \quad \text{BMO}^d(\mathbb{T}) \otimes \text{BMO}^d(\mathbb{T}) \subseteq \text{BMO}_{\text{mult}}^d(\mathbb{T}^2)$$

**Proof.** Using (1.2), one easily obtains the following formula:

$$\sum_{R \in \mathcal{R}} P_R b f_R h_R = fb - \sum_{I \in \mathcal{D}} (m_I b) f_I h_I - \sum_{J \in \mathcal{D}} (m_J b) f_J h_J + \sum_{R \in \mathcal{R}} m_R b f_R h_R.$$

Now (2.1) follows from this expression together with

$$\left\| \sum_{J \in \mathcal{D}} (m_J b) f_J h_J \right\|_{L^2(\mathbb{T}^2)}^2 = \sum_{J \in \mathcal{D}} \|m_J b f_J\|_{L^2(\mathbb{T})}^2,$$

since

$$\left\| \sum_{J \in \mathcal{D}} (m_J b) f_J h_J \right\|_{L^2(\mathbb{T}^2)}^2 \leq \|b\|_\infty^2 \sum_{J \in \mathcal{D}} \|f_J\|_{L^2(\mathbb{T})}^2 = \|b\|_\infty^2 \|f\|_2^2,$$

and the trivial estimates for the terms  $bf$  and  $\sum_{R \in \mathcal{R}} m_R b f_R h_R$ .

To see (2.2), note first that for  $b_1 \in \text{BMO}^d$  and  $f \in L^2(\mathbb{T})$ ,

$$(2.3) \quad \sum_{I \in \mathcal{D}} P_I b_1 f_I h_I = (\pi_{b_1} + \Delta_{b_1})f = (\pi_{b_1} + (\pi_{b_1})^*)f.$$

Therefore  $(P_I b_1)_{I \in \mathcal{D}}$  defines a bounded Haar multiplier on  $L^2(\mathbb{T})$ .

Now let  $b(t, s) = b_1(t)b_2(s)$  with  $b_1, b_2 \in \text{BMO}^d(\mathbb{T})$ . Then  $P_R(b) = P_I(b_1)P_J(b_2)$  and therefore

$$\sum_{R \in \mathcal{R}} P_R b f_R h_R = \sum_{I \in \mathcal{D}} P_I b_1 \left( \sum_{J \in \mathcal{D}} P_J b_2 f_J h_J \right)_I h_I.$$

This yields

$$\begin{aligned} \left\| \sum_{R \in \mathcal{R}} P_R b f_R h_R \right\|_2^2 &= \int_{\mathbb{T}^2} \left| \sum_{I \in \mathcal{D}} P_I b_1(t) \left( \sum_{J \in \mathcal{D}} P_J b_2 f_J h_J \right)_I(s) h_I(t) \right|_2^2 dt ds \\ &\leq C \|b_1\|_{\text{BMO}}^2 \int_{\mathbb{T}} \sum_{I \in \mathcal{D}} \left| \sum_{J \in \mathcal{D}} (P_J b_2 f_J h_J)_I(s) \right|^2 ds \\ &\leq C^2 \|b_1\|_{\text{BMO}}^2 \|b_2\|_{\text{BMO}}^2 \sum_{I \in \mathcal{D}} \sum_{J \in \mathcal{D}} |f_{I \times J}|^2 \end{aligned}$$

with some absolute constant  $C > 0$ . ■

As announced in the introduction, we first relate this space to  $\text{BMO}_{\text{prod}}^d$ . For this purpose we introduce the dyadic paraproducts in two variables (see [PS]):

**Definition 2.3** Given  $b \in L^2(\mathbb{T}^2)$ , we write

$$\pi_b^{(1,2)}(f) = \sum_{R \in \mathcal{R}} b_R m_R f h_R$$

and

$$\Delta_b^{(1,2)}(f) = (\pi_b^{(1,2)})^*(f) = \sum_{R \in \mathcal{R}} b_R f_R \frac{\chi_R}{|R|}.$$

The formula

$$(2.4) \quad \langle \pi_b^{(1,2)}(f), g \rangle = \langle f, \Delta_b^{(1,2)}(g) \rangle = \int_{\mathbb{T}^2} b \left( \sum_{R \in \mathcal{R}} m_R(f) \bar{g}_R h_R \right) dt ds$$

for  $f, g \in \mathcal{H}_{00}$  completely describe the action of the operators  $\pi_b^{(1,2)}$  and  $\Delta_b^{(1,2)}$ .

Let us now define the following mixed operators (see [PS]).

**Definition 2.4** Given  $b \in L^2(\mathbb{T}^2)$ , we define the operators  $\pi_{\Delta_b}$  and  $\Delta_{\pi_b}$  by

$$(2.5) \quad \langle \pi_{\Delta_b}(f), g \rangle = \langle f, \Delta_{\pi_b}(g) \rangle = \int_{\mathbb{T}^2} b \left( \sum_{I \times J \in \mathcal{R}} m_I(f_J) m_J(\bar{g}_I) h_{I \times J} \right) dt ds$$

for  $f, g \in \mathcal{H}_{00}$ .

We write

$$\Lambda_b = \pi_b^{(1,2)} + \Delta_b^{(1,2)} + \Delta_{\pi_b} + \pi_{\Delta_b}.$$

Clearly we have the following expressions:

$$(2.6) \quad \pi_b^{(1,2)}(f)(t, s) = \sum_{I \in \mathcal{D}} \pi_{b_I}(m_I f)(s) h_I(t)$$

$$(2.7) \quad \Delta_b^{(1,2)}(f)(t, s) = \sum_{I \in \mathcal{D}} \Delta_{b_I}(f_I)(s) h_I^2(t)$$

$$(2.8) \quad \Delta_{\pi_b}(f)(t, s) = \sum_{I \in \mathcal{D}} \pi_{b_I}(f_I)(s) h_I^2(t)$$

$$(2.9) \quad \pi_{\Delta_b}(f)(t, s) = \sum_{I \in \mathcal{D}} \Delta_{b_I}(m_I f)(s) h_I(t).$$

**Lemma 2.5** *Let  $R = I \times J \in \mathcal{R}$  and denote  $R_+ = I^+ \times J^+ \cup I^- \times J^-$  and  $R_- = I^+ \times J^- \cup I^- \times J^+$ . Then*

$$(2.10) \quad \pi_b^{(1,2)}(h_R) = (P_{R_+}(b) + P_{R_-}(b))h_R = (P_{R_+}(b) - P_{R_-}(b))|R|^{-1/2}$$

**Proof.** Using that  $m_{R'}(h_R) \neq 0$  only if  $I' \subsetneq I$  and  $J' \subsetneq J$  and that in this case  $m_{R'}(h_R) = h_R(x_{R'})$ , where  $x_{R'} = (t_{I'}, s_{J'})$  is the center of  $R$ , we obtain that

$$\pi_b^{(1,2)}(h_R) = \sum_{I' \subsetneq I, J' \subsetneq J} b_{R'} h_R(x_{R'}) h'_{R'}.$$

Observe that  $h_R(x_{R'}) = h_R(t, s) = \frac{1}{|R|^{1/2}}$  for  $R' \subset R_+$  and  $(t, s) \in R^+$ . Similarly  $h_R(x_{R'}) = h_R(t, s) = -\frac{1}{|R|^{1/2}}$  for  $R' \subset R_-$  and  $(t, s) \in R^-$ . This gives (2.10). ■

**Corollary 2.6** *Let  $b \in L^2(\mathbb{T}^2)$ . Then  $b \in \text{BMO}_{\text{prod}}^d$  if and only if  $(P_{R_+}(b) + P_{R_-}(b))h_R)_{R \in \mathcal{R}}$  is a Haar multiplier on  $L^2(\mathbb{T}^2)$ .*

**Lemma 2.7**  $\Lambda_b(f) = \sum_{R \in \mathcal{R}} P_R(b) f_R h_R.$

**Proof.** Note that for  $\phi, g \in \mathcal{H}_{00}$ , we have

$$(2.11) \quad \phi g = \pi_\phi(g) + \Delta_\phi(g) + \pi_g(\phi).$$

As in (2.3), one obtains

$$(2.12) \quad \sum_{J \in \mathcal{D}} P_J(\phi) g_J h_J = \pi_\phi(g) + \Delta_\phi(g).$$



Combining (2.12) with the formulas in (2.6)- (2.9) we get

$$\begin{aligned}
 \Lambda_b(f)(t, s) &= \sum_{I \in \mathcal{D}} (\pi_{b_I} + \Delta_{b_I})(m_I f)(s) h_I(t) + \sum_{I \in \mathcal{D}} (\pi_{b_I} + \Delta_{b_I})(f_I)(s) h_I^2(t) \\
 &= \sum_{I \in \mathcal{D}} \sum_{J \in \mathcal{D}} P_J(b_I)(s) (m_I f)_J h_J(s) h_I(t) + \sum_{I \in \mathcal{D}} \sum_{J \in \mathcal{D}} P_J(b_I)(s) (f_I)_J h_J(s) h_I^2(t) \\
 &= \sum_{J \in \mathcal{D}} \left( \sum_{I \in \mathcal{D}} (\tilde{P}_J b)_I(s) m_I(f_J) h_I(t) \right) h_J(s) + \sum_{J \in \mathcal{D}} \left( \sum_{I \in \mathcal{D}} (\tilde{P}_J b)_I(s) (f_J)_I h_I^2(t) \right) h_J(s) \\
 &= \sum_{J \in \mathcal{D}} \left( \pi_{\tilde{P}_J b(s, \cdot)}(f_J) \right) (t) h_J(s) + \sum_{J \in \mathcal{D}} \left( \Delta_{\tilde{P}_J b(s, \cdot)}(f_J) \right) (t) h_J(s) \\
 &= \sum_{I \times J \in \mathcal{R}} \tilde{P}_I(\tilde{P}_J(b))(t, s) f_{I \times J} h_{I \times J}(t, s) \\
 &= \sum_{R \in \mathcal{R}} P_R(b)(t, s) f_R h_R(t, s). \quad \blacksquare
 \end{aligned}$$

We now are ready to prove our characterization of  $BMO_{\text{prod}}^d$  in terms of Haar multipliers.

**Theorem 2.8**  $BMO_{\text{prod}}^d = BMO_{\text{mult}}^d$ .

**Proof.** To see that  $BMO_{\text{prod}}^d \subseteq BMO_{\text{mult}}^d$ , it suffices to see that the boundedness of  $\pi_b^{(1,2)}$  implies the boundedness of  $\Delta_{\pi_b}$ . This was proved in [PS], we include here a proof for the sake of completeness.

By (2.5) and the characterization of  $BMO_{\text{prod}}^d$  as the dual of  $H^{1,d}$ , the space of functions with integrable square function, we simply need to show that

$$F = \sum_{I \times J \in \mathcal{R}} m_I(f_J) m_J(g_I) h_{I \times J}(t, s)$$

belongs to  $H^{1,d}$ . Note that

$$\mathcal{S}(F)(t, s) = \left( \sum_{I \times J \in \mathcal{R}} |m_I(f_J)|^2 |m_J(g_I)|^2 h_{I \times J}^2(t, s) \right)^{1/2}.$$

Therefore

$$\begin{aligned}
 \mathcal{S}(F)(t, s) &\leq \left( \sum_{J \in \mathcal{D}} \sum_{I \in \mathcal{D}} (g_I^*(s))^2 h_J^2(s) (f_J^*(t))^2 h_I^2(t) \right)^{1/2} \\
 &= \left( \sum_{I \in \mathcal{D}} (g_I^*(s))^2 h_I^2(t) \right)^{1/2} \left( \sum_{J \in \mathcal{D}} (f_J^*(t))^2 h_J^2(s) \right)^{1/2},
 \end{aligned}$$

and hence

$$\begin{aligned} \int_{\mathbb{T}^2} \mathcal{S}(F)(t, s) dt ds &\leq \\ &\leq \left( \int_{\mathbb{T}^2} \sum_{I \in \mathcal{D}} (g_I^*(s))^2 h_I^2(t) dt ds \right)^{1/2} \left( \int_{\mathbb{T}^2} \sum_{J \in \mathcal{D}} (f_J^*(t))^2 h_J^2(s) dt ds \right)^{1/2} \\ &= \left( \sum_{I \in \mathcal{D}} \int_{\mathbb{T}} (g_I^*(s))^2 ds \right)^{1/2} \left( \sum_{J \in \mathcal{D}} \int_{\mathbb{T}} (f_J^*(t))^2 dt \right)^{1/2} \\ &\leq C \left( \sum_{I \in \mathcal{D}} \|g_I\|_2^2 \right)^{1/2} \left( \sum_{J \in \mathcal{D}} \|f_J\|_2^2 \right)^{1/2} = C \|g\|_2 \|f\|_2. \end{aligned}$$

To prove the reverse inclusion  $\text{BMO}_{\text{mult}}^d \subseteq \text{BMO}_{\text{prod}}^d$ , we shall use the characterization of  $\text{BMO}_{\text{prod}}^d$  given in (1.6).

It is clear that for each measurable set  $\Omega$ , we have  $P_\Omega(b) = P_\Omega(\pi_b^{(1,2)}(\chi_\Omega))$ . We shall show now that

$$P_\Omega(\pi_b^{(1,2)}(\chi_\Omega)) = P_\Omega(\Lambda_b(\chi_\Omega)).$$

Let  $R \in \mathcal{R}$  and  $R \subseteq \Omega$ . Then by (2.10)

$$\langle \Delta_b^{(1,2)}(\chi_\Omega), h_R \rangle = \langle \chi_\Omega, \pi_b^{(1,2)}(h_R) \rangle = |R|^{-1/2} \langle \chi_\Omega, P_{R^+} b - P_{R^-} b \rangle = 0.$$

This shows that  $P_\Omega(\Delta_b^{(1,2)}(\chi_\Omega)) = 0$ .

On the other hand, we also have for  $R = I \times J \subseteq \Omega$  that

$$\pi_{\Delta_b}(h_R) = \sum_{I' \subsetneq I} b_{I' \times J} m_{I'}(h_{I'}) \chi_J h_{I'}.$$

Using that  $\langle \chi_\Omega, \chi_J h_{I'} \rangle = 0$  for all  $I' \subsetneq I$ , we obtain  $P_\Omega(\pi_{\Delta_b}(\chi_\Omega)) = 0$ .

Similarly,  $P_\Omega(\Delta_{\pi_b}(\chi_\Omega)) = 0$ . Finally,

$$\begin{aligned} \|P_\Omega(b)\| &= \|P_\Omega(\pi_b^{(1,2)}(\chi_\Omega))\| = \|P_\Omega(\Lambda_b(\chi_\Omega))\| \\ &\leq \|\Lambda_b(\chi_\Omega)\| \leq \|\Lambda_b\| \|\Omega\|^{1/2}. \end{aligned}$$

■

As a consequence of Thm 2.8, we can sharpen Thm 7.7.2 from [PS] and characterize  $\text{BMO}_{\text{prod}}^d$  in terms of the boundedness of nested commutators with dyadic martingale transforms. This can be understood as a dyadic analogue of the characterization of the continuous product BMO space  $\text{BMO}_{\text{prod}}$  as the space of functions for which the nested commutator

$$[H_1, [H_2, b]] : L^2(\mathbb{T}^2) \rightarrow L^2(\mathbb{T}^2)$$

is bounded, where  $H_1$  resp.  $H_2$  denote the Hilbert transform in the first resp. second variable on  $L^2(\mathbb{T}^2)$ . The latter was proved in [FS] and [LF].

Let  $\Sigma_1, \Sigma_2$  be the spaces of all sequences of signs indexed by the elements of  $\mathcal{D}_1, \mathcal{D}_2$ ,  $\Sigma_1 = \{0, 1\}^{\mathcal{D}_1}$ ,  $\Sigma_2 = \{0, 1\}^{\mathcal{D}_2}$ , and let  $d\sigma_1$  denote the natural product probability measure on  $\Sigma_1$ , which assigns measure  $2^{-n}$  to each cylindrical set of length  $n$ . Let  $d\sigma_2$  denote the corresponding measure on  $\Sigma_2$ . Let  $\Sigma = \Sigma_1 \times \Sigma_2$ , with  $d\sigma$  denoting the product measure, and  $\mathcal{R} = \mathcal{D}_1 \times \mathcal{D}_2$  as before.

For  $\sigma_1 = (\sigma_1(I))_{I \in \mathcal{D}_1} \in \Sigma_1$ ,  $\sigma_2 = (\sigma_2(J))_{J \in \mathcal{D}_2} \in \Sigma_2$ , let  $T_{\sigma_1}, T_{\sigma_2}$  denote the dyadic martingale transforms

$$T_{\sigma_1} : L^2(\mathbb{T}^2) \rightarrow L^2(\mathbb{T}^2), \quad f = \sum_{I \times J \in \mathcal{R}} f_{I \times J} h_{I \times J} \mapsto \sum_{I \times J \in \mathcal{R}} \sigma_1(I) f_{I \times J} h_{I \times J},$$

$$T_{\sigma_2} : L^2(\mathbb{T}^2) \rightarrow L^2(\mathbb{T}^2), \quad f = \sum_{I \times J \in \mathcal{R}} f_{I \times J} h_{I \times J} \mapsto \sum_{I \times J \in \mathcal{R}} \sigma_2(J) f_{I \times J} h_{I \times J}.$$

**Theorem 2.9** *Let  $b \in L^2(\mathbb{T}^2)$ . Then the following are equivalent:*

- (i)  $b \in \text{BMO}_{\text{prod}}^d$
- (ii) *The nested commutators*

$$(2.13) \quad [T_{\sigma_1}, [T_{\sigma_2}, b]] : L^2(\mathbb{T}^2) \rightarrow L^2(\mathbb{T}^2)$$

*are uniformly bounded for all  $\sigma_1 \in \Sigma_1, \sigma_2 \in \Sigma_2$ .*

- (iii) *The nested commutators  $[T_{\sigma_1}, [T_{\sigma_2}, b]] : L^2(\mathbb{T}^2) \rightarrow L^2(\mathbb{T}^2)$  are bounded on average, in the sense that the map*

$$\Phi_b : L^2(\mathbb{T}^2) \rightarrow L^2(\Sigma_1 \times \Sigma_2 \times \mathbb{T}^2), f \mapsto [T_{\sigma_1}, [T_{\sigma_2}, b]]f$$

*is bounded.*

*In this case, we have*

$$(2.14) \quad \|b\|_{\text{BMO}_{\text{prod}}^d} \approx \|\Lambda_b\| \leq \|\Phi_b\| \leq \sup_{\sigma_1 \in \Sigma_1, \sigma_2 \in \Sigma_2} \|[T_{\sigma_1}, [T_{\sigma_2}, b]]\| \leq 4\|\Lambda_b\|.$$

**Proof.** We use the ideas of the proofs of Thm 3.4, Cor 4.1 in [GPTV], adapted to the two-variable case, and of Thm 7.7.2 in [PS].

From [PS, p. 493], we know that

$$[T_{\sigma_1}, [T_{\sigma_2}, b]] = [T_{\sigma_1}, [T_{\sigma_2}, \Lambda_b]].$$

Therefore

$$\sup_{\sigma_1 \in \Sigma_1, \sigma_2 \in \Sigma_2} \|[T_{\sigma_1}, [T_{\sigma_2}, b]]\| \leq 4\|\Lambda_b\|.$$

The second inequality in (2.14) is obvious.

Finally, for  $f \in L^2(\mathbb{T}^2)$  one has

$$\begin{aligned}
 \|\Phi_b f\|^2 &= \int \int_{\Sigma_1 \times \Sigma_2} \|[T_{\sigma_1}, [T_{\sigma_2}, b]]f\|_{L^2(\mathbb{T}^2)}^2 d\sigma_1 d\sigma_2 \\
 &= \int \int_{\Sigma_1 \times \Sigma_2} \left\| \sum_{I \times J \in \mathcal{R}} \sigma_1(I)\sigma_2(J)[\tilde{P}_I, [\tilde{P}_J, b]]f \right\|_{L^2(\mathbb{T}^2)}^2 d\sigma_1 d\sigma_2 \\
 &= \sum_{I \times J \in \mathcal{R}} \|[ \tilde{P}_I, [\tilde{P}_J, b] ]f\|_{L^2(\mathbb{T}^2)}^2 \\
 (2.15) \quad &= \sum_{I \times J \in \mathcal{R}} \|[ \tilde{P}_I, [\tilde{P}_J, \Lambda_b] ]f\|_{L^2(\mathbb{T}^2)}^2 \\
 &= \sum_{I \times J \in \mathcal{R}} \|(\tilde{P}_I \tilde{P}_J \Lambda_b - \tilde{P}_I \Lambda_b \tilde{P}_J - \tilde{P}_J \Lambda_b \tilde{P}_I + \Lambda_b \tilde{P}_J \tilde{P}_I)f\|_{L^2(\mathbb{T}^2)}^2 \\
 &\geq \sum_{I \times J \in \mathcal{R}} \|\tilde{P}_I \tilde{P}_J \Lambda_b f\|^2 = \|\Lambda_b f\|^2,
 \end{aligned}$$

since  $\tilde{P}_I \Lambda_b \tilde{P}_I = 0$  and  $\tilde{P}_J \Lambda_b \tilde{P}_J = 0$ . This proves the first inequality in (2.14). ■

The martingale transformation approach is also interesting in the study of  $\text{BMO}_{\text{rect}}^d$ . Although  $\Lambda_b$  is in general not bounded for  $b \in \text{BMO}_{\text{rect}}^d$ , the space  $\text{BMO}_{\text{rect}}^d$  can be characterized in terms of “average boundedness” of  $\Lambda_b$ , and also in terms of the boundedness of  $\Lambda_b$  from  $L^2(\mathbb{T}) \hat{\otimes} L^2(\mathbb{T})$  into  $L^2(\mathbb{T}^2)$ . For  $\sigma = (\sigma_1, \sigma_2) \in \Sigma$ , let  $T_\sigma = T_{\sigma_1} T_{\sigma_2} : L^2(\mathbb{T}^2) \rightarrow L^2(\mathbb{T}^2)$ .

**Theorem 2.10** *For  $\varphi \in \mathcal{H}_{00}$ ,  $\|\varphi\|_{\text{BMO}_{\text{rect}}^d}$  is equal to the norm of the operator*

$$\Psi_\varphi : L^2(\mathbb{T}^2) \rightarrow L^2(\mathbb{T}^2 \times \Sigma), f \mapsto \Lambda_\varphi T_\sigma f.$$

**Proof.** Let  $f \in L^2(\mathbb{T}^2)$  and  $\varphi \in \mathcal{H}_{00}$ . From Lemma 2.7 we have

$$(2.16) \quad \Lambda_\varphi f = \sum_{R \in \mathcal{R}} P_R \varphi f_R h_R.$$

Thus

$$\begin{aligned}
 \int_\Sigma \int_{\mathbb{T}^2} \|\Lambda_\varphi T_\sigma f\|^2 dt ds d\sigma &= \int_{\mathbb{T}^2} \int_\Sigma \left\| \sum_{R \in \mathcal{R}} \sigma(R)(P_R \varphi)(t, s) f_R h_R(t, s) \right\|^2 d\sigma dt ds \\
 &= \int_{\mathbb{T}^2} \sum_{R \in \mathcal{R}} |f_R|^2 \frac{\chi_R(t, s)}{|R|} |(P_R \varphi)(t, s)|^2 dt ds \\
 &= \sum_{R \in \mathcal{R}} |f_R|^2 \frac{1}{|R|} \|P_R \varphi\|^2.
 \end{aligned}$$

Thus the operator norm of  $\Psi_\varphi$  is  $\sup_{R \in \mathcal{R}} \frac{1}{|R|^{1/2}} \|P_R \varphi\| = \|\varphi\|_{\text{BMO}_{\text{rect}}^d}$ . ■

**Proposition 2.11** *If  $b \in \text{BMO}_{\text{rect}}^d$  then  $\Lambda_b$  maps  $L^2(\mathbb{T}) \hat{\otimes} L^2(\mathbb{T})$  into  $L^2(\mathbb{T}^2)$ .*

**Proof.** Assume  $f(t, s) = f_1(t)f_2(s)$  with  $\|f_1\| = \|f_2\| = 1$ . Then we have

$$\sum_{R \in \mathcal{R}} P_R b f_R h_R = \sum_{I \in \mathcal{D}} \tilde{P}_I \left( \sum_{J \in \mathcal{D}} \tilde{P}_J b(f_2)_J h_J \right) (t) (f_1)_I h_I(t)$$

Writing  $g(t, s) = \sum_{J \in \mathcal{D}} P_J(b(t, \cdot))(s)(f_2)_J h_J(s)$ , we obtain

$$\left\| \sum_{R \in \mathcal{R}} P_R b f_R h_R \right\|_2^2 = \int_{\mathbb{T}} \int_{\mathbb{T}} \left| \sum_{I \in \mathcal{D}} P_I(g(\cdot, s))(t) (f_1)_I h_I(t) \right|^2 dt ds.$$

Now let us consider  $g$  as a function in  $t$  taking values in the Hilbert space  $L^2(\mathbb{T})$ . Recall that as in the scalar case, the Haar multiplier norm of  $(P_I g)_{I \in \mathcal{D}}$  is controlled by the vector  $\text{BMO}^d(\mathbb{T})$  norm of  $g$  given by

$$\sup_{I \in \mathcal{D}} \frac{1}{|I|^{1/2}} \|P_I g\|_{L^2(\mathbb{T}, L^2(\mathbb{T}))}.$$

Thus

$$\sup_{\|f_1\|=1} \int_{\mathbb{T}} \left\| \sum_{I \in \mathcal{D}} P_I(g(\cdot, s))(t) (f_1)_I h_I(t) \right\|_{L^2(\mathbb{T})}^2 dt \leq C \sup_I \frac{1}{|I|} \|P_I g\|_{L^2(\mathbb{T}, L^2(\mathbb{T}))}^2.$$

Notice now that

$$P_I \left( \sum_{J \in \mathcal{D}} P_J(b(t, \cdot))(s)(f_2)_J h_J(s) \right) = \sum_{J \in \mathcal{D}} P_{I \times J}(b)(t, s)(f_2)_J h_J(s).$$

On the other hand, applying a corresponding argument to the function  $(\tilde{P}_I b)(t, s) = \sum_{J \in \mathcal{D}} (P_{I \times J} b)(t, s) h_J(s)$  understood as a function in  $s$  which takes values in  $L^2(\mathbb{T})$ , we obtain for  $\|f_2\|_2 = 1$

$$\begin{aligned} \|P_I g\|_{L^2(\mathbb{T}, L^2(\mathbb{T}))}^2 &= \int_{\mathbb{T}^2} \left| \sum_{J \in \mathcal{D}} P_{I \times J}(b)(t, s)(f_2)_J h_J(s) \right|^2 dt ds \\ &= \int_{\mathbb{T}} \left\| \sum_{J \in \mathcal{D}} P_{I \times J}(b)(\cdot, s)(f_2)_J h_J(s) \right\|_{L^2(\mathbb{T})}^2 ds \\ &\leq C \sup_J \frac{1}{|J|} \|P_{I \times J}(b)\|_{L^2(\mathbb{T}, L^2(\mathbb{T}))}^2 \leq C \|I\| \|b\|_{\text{BMO}_{\text{rect}}^d}^2. \end{aligned}$$

This finishes the proof of the proposition. ■

### 3. Sweeps of functions in BMO

Let us now recall that the (dyadic) sweep of a function  $\varphi \in L^2(\mathbb{T}^2)$  is defined as follows:

$$S_\varphi = \sum_{R \in \mathcal{R}} |\varphi_R|^2 \frac{\chi_R}{|R|},$$

i.e.  $S_\varphi = \mathcal{S}(\varphi)^2$ .

We list some properties of the sweep which will be relevant for our purposes, the proofs of which are elementary and left to the reader.

**Proposition 3.1**

- (i)  $S_\varphi(t, s) = \sum_{I \in \mathcal{D}} S_{\varphi_I}(s) \frac{\chi_I(t)}{|I|}$ .
- (ii)  $S_\varphi = \Delta_\varphi^{(1,2)}(\varphi)$ .
- (iii)  $P_\Omega(S_\varphi) = P_\Omega(S_{P_\Omega \varphi})$ .
- (iv) If  $p > \frac{1}{2}$  then  $\varphi \in L^{2p}(\mathbb{T}^2)$  if and only if  $S_\varphi \in L^p(\mathbb{T}^2)$ .
- (v) If  $S_\varphi \in L^\infty$  then  $\varphi \in \text{BMO}_{\text{prod}}^d$ .
- (vi)  $\|S_\varphi\|_2 \leq C \|\varphi\|_{\text{BMO}_{\text{prod}}^d} \|\varphi\|_2$ .

Here it is the basic result relating the boundedness of  $\pi_b^{(1,2)}$  and  $\Lambda_{S_b}$ .

**Lemma 3.2** *Let  $b \in \mathcal{H}_{00}$ . Then*

$$\pi_b^{(1,2)*} \pi_b^{(1,2)} = \Lambda_{S_b} + D_b,$$

where  $D_b$  is a linear operator on  $L^2(\mathbb{T}^2)$  with  $\|D_b\| \leq C \|b\|_{\text{BMO}_{\text{rect}}^d}^2$ , and  $C > 0$  is an absolute constant.

**Proof.** Let  $R = I \times J, R' = I' \times J' \in \mathcal{R}$ .

First, observe that

$$\begin{aligned} (3.1) \quad & \left\langle \pi_b^{(1,2)*} \pi_b^{(1,2)} h_R, h_{R'} \right\rangle \\ &= \left\langle \sum_{I'' \times J'' \in \mathcal{D}_1 \times \mathcal{D}_2} h_{I'' \times J''} b_{I'' \times J''} m_{I'' \times J''}(h_R), \sum_{I'' \times J'' \in \mathcal{D}_1 \times \mathcal{D}_2} h_{I'' \times J''} b_{I'' \times J''} m_{I'' \times J''}(h_{R'}) \right\rangle \\ &= \sum_{I'' \times J'' \in \mathcal{D}_1 \times \mathcal{D}_2, I'' \subsetneq I, J'' \subsetneq J} |b_{I'' \times J''}|^2 m_{I''}(h_I) m_{I''}(h_{I'}) m_{J''}(h_J) m_{J''}(h_{J'}). \end{aligned}$$

We now do a kind of triangular truncation with respect to the indices  $I, I', J, J'$ .

(i)  $I \supsetneq I', J \supsetneq J'$ .

$$\begin{aligned} \left\langle \pi_{S_b}^{(1,2)} h_R, h_{R'} \right\rangle &= \langle S_b, h_{R'} \rangle m_{R'}(h_R) \\ &= \left\langle \sum_{I'' \times J'' \in \mathcal{D}_1 \times \mathcal{D}_2} \frac{\chi_{I'' \times J''}}{|I''||J''|} |b_{I'' \times J''}|^2, h_{R'} \right\rangle m_{R'}(h_R) \\ &= \sum_{I'' \times J'' \in \mathcal{D}_1 \times \mathcal{D}_2} |b_{I'' \times J''}|^2 m_{I''}(h_{I'}) m_{J''}(h_{J'}) m_{I'}(h_I) m_{J'}(h_J). \end{aligned}$$

This is nonzero only if  $I' \subsetneq I$  and  $J' \subsetneq J$ . In this case, we get contributions only for  $I'' \subsetneq I'$  and  $J'' \subsetneq J'$ , and the expression agrees with (3.1).

(ii)  $I \subsetneq I', J \subsetneq J'$ . Observe that

$$\langle \Delta_{S_b}^{(1,2)} h_R, h_{R'} \rangle = \langle h_R, \pi_{S_b}^{(1,2)} h_{R'} \rangle = \langle \pi_{S_b}^{(1,2)} h_{R'}, h_R \rangle.$$

As shown above, this equals  $\langle \pi_b^{(1,2)*} \pi_b^{(1,2)} h_{R'}, h_R \rangle$  if  $I' \supsetneq I$  and  $J' \supsetneq J$ , and is 0 otherwise.

(iii)  $I \supsetneq I', J \subsetneq J'$ .

$$\begin{aligned} \left\langle \pi_{\Delta_{S_b}} h_R, h_{R'} \right\rangle &= \left\langle \sum_{I'' \times J'' \in \mathcal{D}_1 \times \mathcal{D}_2} S_{b_{I'' \times J''}} h_{I''} \frac{\chi_{J''}}{|J''|} m_{I''}(h_{R_{J''}}), h_{R'} \right\rangle \\ &= \left\langle \sum_{I'' \in \mathcal{D}_1} S_{b_{I'' \times J}} h_{I''} \frac{\chi_J}{|J|} m_{I''}(h_I), h_{R'} \right\rangle \\ &= S_{b_{I' \times J}} m_{I'}(h_J) m_J(h_{I'}) = \langle S_b, h_{I' \times J} \rangle m_{I'}(h_I) m_J(h_{J'}) \\ &= \left\langle \sum_{I'' \times J'' \in \mathcal{D}_1 \times \mathcal{D}_2} \frac{\chi_{I'' \times J''}}{|I''||J''|} |b_{I'' \times J''}|^2, h_{I' \times J} \right\rangle m_{I'}(h_I) m_J(h_{J'}) \\ &= \sum_{I'' \times J'' \in \mathcal{D}_1 \times \mathcal{D}_2} |b_{I'' \times J''}|^2 m_{I''}(h_{I'}) m_{J''}(h_J) m_{I'}(h_I) m_J(h_{J'}). \end{aligned}$$

This is nonzero only for  $I' \subsetneq I$  and  $J' \supsetneq J$ . In this case, the sum has only contributions for  $I'' \subsetneq I'$  and  $J'' \subsetneq J$ , and agrees with (3.1).

(iv)  $I' \supsetneq I$  and  $J' \subsetneq J$ . Note that  $\langle \Delta_{\pi_{S_b}} h_R, h_{R'} \rangle = \langle \pi_{\Delta_{S_b}} h_{R'}, h_R \rangle$ . As shown above, this is only nonzero for  $I' \supsetneq I$  and  $J' \subsetneq J$ , and agrees with (3.1) in this case.

(v)  $I' = I$  or  $J = J'$ . Let  $f \in L^2(\mathbb{T}^2)$ . Then

$$\begin{aligned} & \sum_{I \in \mathcal{D}_1} \sum_{J, J' \in \mathcal{D}_2} \langle \pi_b^{(1,2)*} \pi_b^{(1,2)} h_{I \times J} f_{I \times J}, h_{I \times J'} f_{I \times J'} \rangle \\ &= \sum_{I \in \mathcal{D}_1} \frac{1}{|I|} \sum_{I'' \subsetneq I} \sum_{J'' \in \mathcal{D}_2} |b_{I'' \times J''}|^2 f_{I \times J} f_{I \times J'} m_{J''}(h_J) m_{J''}(h_{J'}) \\ &= \sum_{I \in \mathcal{D}_1} \|\pi_{b^I} f_I\|^2, \end{aligned}$$

where for each  $I$ ,  $f_I$  stands for the one-variable function

$$\sum_{J \in \mathcal{D}_2} h_J f_{I \times J},$$

and  $b^I$  for the function

$$\sum_{J \in \mathcal{D}_2} h_J \frac{1}{|I|^{1/2}} \left( \sum_{I'' \subsetneq I} |b_{I'' \times J}|^2 \right)^{1/2}.$$

It is easy to see that

$$\|b^I\|_{\text{BMO}^d} \leq \|b\|_{\text{BMO}_{\text{rect}}^d} \quad \text{for all } I \in \mathcal{D}_1.$$

Thus the above sum is bounded by  $c \|b\|_{\text{BMO}_{\text{rect}}^d}^2 \|f\|^2$ , where  $c$  is an absolute constant.

The same estimate holds for the terms corresponding to  $J = J'$ .

Now we have counted the terms corresponding to  $I = I'$ ,  $J = J'$  twice and need to estimate them separately. Let  $f \in L^2(\mathbb{T}^2)$ . Then

$$\begin{aligned} (3.2) \quad & \sum_{I \in \mathcal{D}_1} \sum_{J \in \mathcal{D}_2} \langle \pi_b^{(1,2)*} \pi_b^{(1,2)} h_{I \times J} f_{I \times J}, h_{I \times J} f_{I \times J} \rangle \\ &= \sum_{I \in \mathcal{D}_1} \sum_{J \in \mathcal{D}_2} \frac{1}{|I||J|} \sum_{I'' \subsetneq I, J'' \subsetneq J} |b_{I'' \times J''}|^2 |f_{I \times J}|^2 \leq \|b\|_{\text{BMO}_{\text{rect}}^d}^2 \|f\|^2. \end{aligned}$$

Defining  $D_b$  now by

$$\begin{aligned} (3.3) \quad \langle D_b f, f \rangle &= \sum_{I \times J, I' \times J' \in \mathcal{R}, I' = I} \langle \pi_b^{(1,2)*} \pi_b^{(1,2)} h_{I \times J} f_{I \times J}, h_{I' \times J'} f_{I' \times J'} \rangle + \\ &+ \sum_{I \times J, I' \times J' \in \mathcal{R}, I' \neq I, J' = J} \langle \pi_b^{(1,2)*} \pi_b^{(1,2)} h_{I \times J} f_{I \times J}, h_{I' \times J'} f_{I' \times J'} \rangle, \end{aligned}$$

we obtain the statement of the lemma. ■



Now we are ready to state the main result of this section.

**Theorem 3.3** *Let  $b \in \text{BMO}_{\text{rect},2}^d$ . Then  $b \in \text{BMO}_{\text{prod}}^d$  if and only if  $S_b \in \text{BMO}_{\text{prod}}^d$ .*

**Proof.** We will first show that there exist  $C > 0$  such that

$$(3.4) \quad \|S_b\|_{\text{prod}} \leq C \|b\|_{\text{prod}}^2.$$

Indeed, by Chang’s Theorem [Ch], [ChFef2] it is sufficient to show that there exists a constant  $C > 0$  with

$$\|P_\Omega S_b\|_2 \leq C \|b\|_{\text{prod}}^2 |\Omega|^{1/2}$$

for all  $\Omega \subseteq \mathbb{T}^2$  measurable (see (1.5)). Using Proposition 3.1, we obtain

$$(3.5) \quad \|P_\Omega S_b\|_2 = \|P_\Omega S_{P_\Omega b}\|_2 \leq \|S_{P_\Omega b}\|_2 \leq \|P_\Omega b\|_{\text{prod}} \|P_\Omega b\|_2 \leq \|b\|_{\text{prod}}^2 |\Omega|^{1/2}$$

For the converse, assume that  $S_b \in \text{BMO}_{\text{prod}}^d$ . Then  $\Lambda_{S_b}$  is bounded by Theorem 2.8. Now Lemma 3.2 finishes the proof. ■

**Remark.** The first implication can also be shown with the John-Nirenberg Theorem for product BMO, which was proved in [ChFef1] (for a dyadic version, see [T]).

The sweep can be understood as a bilinear map. For  $f, g \in \mathcal{H}_{00}$ , let

$$S_{f,g} = \sum_{R \in \mathcal{R}} \frac{\chi_R}{|R|} f_R g_R,$$

so  $S_f = S_{f,\bar{f}}$ .

**Corollary 3.4**  $S : \text{BMO}_{\text{prod}}^d \times \text{BMO}_{\text{prod}}^d \rightarrow \text{BMO}_{\text{prod}}^d$  is bounded.

**Proof.** The Cauchy-Schwarz inequality gives the pointwise inequality

$$S_{f,g} \leq (S_f)^{1/2} (S_g)^{1/2} \quad \text{for } f, g \in \mathcal{H}_{00}.$$

Let  $\Omega \subseteq \mathbb{T}^2$  be measurable. Using an adaption of 3.1(iii), we see that

$$\begin{aligned} \|P_\Omega S_{f,g}\|_2 &= \|P_\Omega S_{P_\Omega f, P_\Omega g}\|_2 \leq \|S_{P_\Omega f, P_\Omega g}\|_2 \leq \|(S_{P_\Omega f})^{1/2} (S_{P_\Omega g})^{1/2}\|_2 \\ &\leq \|S_{P_\Omega f}\|_2^{1/2} \|S_{P_\Omega g}\|_2^{1/2} \leq |\Omega|^{1/2} \|f\|_{\text{prod}} \|g\|_{\text{prod}} \end{aligned}$$

by (3.5). ■

Another application of Lemma 3.2 yields the following result.

**Theorem 3.5** *Let  $\|\cdot\|_*$  be an positive homogeneous function of degree 1 on  $\mathcal{H}_{00}$  such that*

- (i) *There exists  $c > 0$  such that  $\|\cdot\|_{\text{BMO}_{\text{rect}}^d} \leq c\|\cdot\|_*$*
- (ii) *There exists  $k > 0$  such that  $\|S_\varphi\|_* \leq k\|\varphi\|_*^2$ .*

*Then there exists a constant  $\tilde{C}$  such that for all  $\varphi \in \mathcal{H}_{00}$ ,*

$$\|\varphi\|_{\text{BMO}_{\text{prod}}^d} \leq \tilde{C}\|\varphi\|_*.$$

**Proof.** Throughout this proof we simply write  $\pi_\varphi = \pi_\varphi^{(1,2)}$ . From Lemma 3.2 we have

$$\pi_\varphi^* \pi_\varphi = \pi_{S_\varphi}^{(1,2)} + \pi_{S_\varphi}^{(1,2)*} + \pi_{\Delta_{S_\varphi}} + \pi_{\Delta_{S_\varphi}}^* + D_\varphi,$$

with  $\|D_\varphi\| \leq C\|\varphi\|_{\text{BMO}_{\text{rect}}^d}^2$ .

Let

$$E_n = \text{span}\{h_{I \times J} : I \in \mathcal{D}_1, J \in \mathcal{D}_2, |I|, |J| \geq 2^{-n}\},$$

let  $P_n$  be the orthogonal projection onto  $E_n$  in  $L^2(\mathbb{T}^2)$ , and let

$$c(n) = \sup\{\|\pi_\varphi|_{E_n}\|, \|\varphi\|_* \leq 1\}.$$

A trivial estimate shows that  $c(n) < \infty$  for each  $n \in \mathbb{N}$ . For  $n \in \mathbb{N}$  and  $\varepsilon > 0$ , choose  $f_n \in E_n$  and  $\varphi \in \mathcal{H}_{00}$  with  $\|\varphi\|_* = 1$ ,  $\|f_n\| = 1$  and  $\|\pi_\varphi f_n\| \geq (1 - \varepsilon)c(n)$ . Then

$$\begin{aligned} (3.6) \quad (1 - \varepsilon)^2 c(n)^2 &\leq \|\pi_\varphi f_n\|^2 = \langle \pi_\varphi^* \pi_\varphi f_n, f_n \rangle \\ &= \langle \pi_{S_\varphi}^{(1,2)} f_n, f_n \rangle + \langle \pi_{S_\varphi}^{(1,2)*} f_n, f_n \rangle + \\ &\quad + \langle \pi_{\Delta_{S_\varphi}} f_n, f_n \rangle + \langle \pi_{\Delta_{S_\varphi}}^* f_n, f_n \rangle + \langle D_\varphi f_n, f_n \rangle. \end{aligned}$$

By definition of  $c(n)$ , the first two terms can be estimated by

$$c(n)\|S_\varphi\|_* \leq c(n)k.$$

For the next two terms, we have to remark that that

$$(3.7) \quad \langle \pi_{\Delta_{S_\varphi}} f_n, f_n \rangle = \langle \pi_{\Delta_{P_n S_\varphi}} f_n, f_n \rangle \leq \|\pi_{\Delta_{P_n S_\varphi}}\| \leq \tilde{c}\|\pi_{P_n S_\varphi}^{(1,2)}\| \leq \tilde{c}\|\pi_{S_\varphi}^{(1,2)}|_{E_n}\|$$

Here, we use as in the proof of Thm 2.8 that there exists a constant  $\tilde{c}$  such that  $\|\pi_{\Delta_b}\| \leq \tilde{c}\|\pi_b^{(1,2)}\|$  for all  $b \in \mathcal{H}_{00}$  (see [PS], Thm 7.7.2).

The last term is easily controlled by

$$\langle D_\varphi f_n, f_n \rangle \leq C\|\varphi\|_{\text{BMO}_{\text{rect}}^d}^2.$$

Altogether, we obtain that

$$(3.8) \quad \|\pi_\varphi^{(1,2)}|_{E_n}\|^2 \leq 4\tilde{c}\|\pi_{S_\varphi}^{(1,2)}|_{E_n}\| + C\|\varphi\|_{\text{BMO}_{\text{rect}}^d}^2.$$

With

$$\|\varphi\|_{\text{BMO}_{\text{rect}}^d} \leq c\|\varphi\|_*$$

and

$$\|\pi_{S_\varphi}^{(1,2)}|_{E_n}\| \leq c(n)\|S_\varphi\|_* \leq kc(n)\|\varphi\|_*^2,$$

it follows that

$$(3.9) \quad (1 - \varepsilon)^2 c(n)^2 \leq 4\tilde{c}kc(n) + c^2C$$

Thus

$$c(n) \leq \sqrt{4\tilde{c}^2k^2 + Cc^2} + 2\tilde{c}k.$$

With

$$\tilde{C} = \sqrt{4\tilde{c}^2k^2 + Cc^2} + 2\tilde{c}k,$$

it follows that  $\|\pi_\varphi\| \leq \tilde{C}\|\varphi\|_*$ . ■

We can now characterize  $\text{BMO}_{\text{prod}}^d$  in terms of the  $\text{BMO}_{\text{rect},2}^d$ -norm.

**Theorem 3.6** *Let  $\varphi \in \text{BMO}_{\text{rect}}^d(\mathbb{T}^2)$ . Then  $\varphi \in \text{BMO}_{\text{prod}}^d$  if and only if  $(\|S_\varphi^{(n)}\|_{\text{BMO}_{\text{rect}}^d}^{1/2^n})_{n \in \mathbb{N}}$  is bounded, where  $S_\varphi^{(n)}$  is the  $n$ -fold sweep of  $\varphi$ , defined recursively by  $S_\varphi^{(n)} = S_{S_\varphi}^{(n-1)}$ .*

**Proof.** By Theorem 3.3, we have for each  $n \in \mathbb{N}$

$$\|S_\varphi^{(n)}\|_{\text{BMO}_{\text{rect}}^d} \leq \|S_\varphi^{(n)}\|_{\text{BMO}_{\text{prod}}^d} \leq C \cdot C^2 \cdots C^{2^{n-1}} \|\varphi\|_{\text{BMO}_{\text{prod}}^d}^{2^n} \leq C^{2^n} \|\varphi\|_{\text{BMO}_{\text{prod}}^d}^{2^n},$$

and consequently

$$\|S_\varphi^{(n)}\|_{\text{BMO}_{\text{rect}}^d}^{1/2^n} \leq C\|\varphi\|_{\text{BMO}_{\text{prod}}^d}.$$

Conversely, the map

$$\varphi \mapsto \sup_{n \in \mathbb{N}} \|S_\varphi^{(n)}\|_{\text{BMO}_{\text{rect}}^d}^{1/2^n}$$

clearly defines a positive homogeneous function on  $\mathcal{H}_0$  with satisfies conditions in Theorem 3.5. ■

Another consequence of Theorem 3.5 is

**Corollary 3.7**  *$S$  does not map  $\text{BMO}_{\text{rect}}^d \times \text{BMO}_{\text{rect}}^d$  boundedly into  $\text{BMO}_{\text{rect}}^d$ .*

**Proof.** We know that the  $\|\cdot\|_{\text{prod}}$  norm cannot be controlled by the  $\|\cdot\|_{\text{rect}}$  norm. So Condition (ii) in Theorem 3.5 cannot hold, and in particular  $S$  does not map  $\text{BMO}_{\text{rect}}^d \times \text{BMO}_{\text{rect}}^d$  boundedly into  $\text{BMO}_{\text{rect}}^d$ . ■

### 4. The scale $BMO_{\text{rect},p}^d$

Recall that for  $1 \leq p < \infty$ , a function  $\varphi \in L^2(\mathbb{T}^2)$  is said to belong to  $BMO_{\text{rect},p}^d$  if

$$\|\varphi\|_{\text{rect},p} = \sup_{R \in \mathcal{R}} \frac{1}{|R|^{1/p}} \|P_R \varphi\|_p < \infty.$$

Note that  $BMO_{\text{rect},p_2}^d \subseteq BMO_{\text{rect},p_1}^d$  for  $p_1 \leq p_2$ .

The reader should also be aware that functions in  $BMO_{\text{rect},p}^d$  are actually in  $L^p(\mathbb{T}^2)$ , due to the identities  $m_I(f) = m_I(P_{I \times \mathbb{T}} f)$  and  $m_J(f) = m_J(P_{\mathbb{T} \times J} f)$ .

The following proposition characterizes the behaviour of the  $BMO_{\text{rect},p}^d$  norms under the sweep.

**Proposition 4.1** *Let  $p > \frac{1}{2}$  and let  $C_p = \|\mathcal{S}\|_{L^{2p} \rightarrow L^{2p}}$ . Then*

$$\|S_\varphi\|_{\text{rect},p} \leq 4C_p^2 \|\varphi\|_{\text{rect},2p}^2.$$

**Proof.** Since  $P_R(S_\varphi) = P_R(S_{P_R \varphi})$  and  $\|P_R(g)\|_p \leq 4\|g\|_p$ , we obtain

$$\|S_\varphi\|_{\text{rect},p} \leq 4 \sup_{R \in \mathcal{R}} \frac{1}{|R|^{1/p}} \|S_{P_R(\varphi)}\|_p \leq 4C_p^2 \sup_{R \in \mathcal{R}} \frac{1}{|R|^{1/p}} \|P_R(\varphi)\|_{2p}^2.$$

This gives the result. ■

It is known that  $BMO_{\text{prod}}^d \subsetneq BMO_{\text{rect},2}^d$ . Indeed, this is basically the content of Carleson’s original counterexample [C] (for the continuous case, see [Fef]). As pointed out in [Fef], the example in [C] implies that  $BMO_{\text{rect},2}^d \not\subseteq L^4(\mathbb{T}^2)$ .

We shall improve this by showing that actually  $BMO_{\text{prod}}^d \subsetneq BMO_{\text{rect},p}^d$  for all  $p$ . We will show that for any  $p_2 > p_1 \geq 1$ ,  $BMO_{\text{rect},p_1}^d \not\subseteq L^{p_2}(\mathbb{T}^2)$  and therefore in particular  $BMO_{\text{rect},p_1}^d \not\subseteq BMO_{\text{rect},p_2}^d$ . For the case  $p_1 = 1, p_2 = 2$ , this answers a question posed in [FS].

As a corollary, we show that

$$BMO_{\text{prod}}^d \subsetneq \bigcap_{p \geq 1} BMO_{\text{rect},p}^d.$$

**Theorem 4.2** *Let  $p \geq 2$ . Then  $BMO_{\text{prod}}^d \subseteq BMO_{\text{rect},p}^d$ .*

Moreover

$$(4.1) \quad \|\varphi\|_{\text{rect},p} \leq C \|\varphi\|_{\text{prod}}^{1-2/p} \|\varphi\|_{\text{rect},2}^{2/p}$$

**Proof.** Let us first show that  $\text{BMO}_{\text{prod}}^d \subseteq L^p(\mathbb{T}^2)$  and

$$(4.2) \quad \|\varphi\|_p \leq C \|\varphi\|_{\text{prod}}^{1-2/p} \|\varphi\|_2^{2/p}.$$

For  $p = 2^k$ ,  $k \in \mathbb{N}$ , we shall prove (4.2) by induction.

It is obvious for  $k = 1$ . For  $k = 2$  we have

$$(4.3) \quad \|\varphi\|_4^2 = \|S_\varphi\|_2 = \|\Delta_\varphi^{(1,2)}(\varphi)\|_2 \leq C \|\varphi\|_{\text{prod}} \|\varphi\|_2.$$

Assume it holds for  $p_k = 2^k$ .

$$\|\varphi\|_{p_{k+1}}^2 = \|S_\varphi\|_{p_k} \leq C \|S_\varphi\|_{\text{prod}}^{1-2/p_k} \|S_\varphi\|_2^{2/p_k}.$$

Now from (3.4) and (4.3) we obtain

$$\|\varphi\|_{p_{k+1}} \leq C \|\varphi\|_{\text{prod}}^{1-2/p_k} \|\varphi\|_4^{2/p_k} \leq C \|\varphi\|_{\text{prod}}^{1-2/p_{k+1}} \|\varphi\|_2^{2/p_{k+1}}.$$

Now the general case follows by interpolation.

Given  $p > 2$  and  $p \neq 2^k$  for any  $k \in \mathbb{N}$ , find  $m \in \mathbb{N}$  such that  $2^{m-1} < p < 2^m$ .

Write

$$\frac{1}{p} = \frac{1 - \theta_m}{2^{m-1}} + \frac{\theta_m}{2^m}.$$

Now apply the previous case combined with

$$\|\varphi\|_p \leq \|\varphi\|_{2^{m-1}}^{1-\theta_m} \|\varphi\|_{2^m}^{\theta_m}.$$

Let us use (4.2) to obtain the desired estimate for the  $\text{BMO}_{\text{rect},p}^d$ -norm. Given  $R \in \mathcal{R}$  we have

$$\|P_R \varphi\|_p \leq C \|P_R \varphi\|_{\text{prod}}^{1-2/p} \|P_R \varphi\|_2^{2/p} \leq C \|\varphi\|_{\text{prod}}^{1-2/p} \|\varphi\|_{\text{rect},2}^{2/p} |R|^{1/p}.$$

This finishes the proof. ■

**Proposition 4.3** *Let  $2 < p$ . There exists  $\phi \in \text{BMO}_{\text{rect},2}^d \setminus L^p(\mathbb{T}^2)$ .*

*In particular  $\text{BMO}_{\text{prod}}^d \subseteq \text{BMO}_{\text{rect},p}^d \subsetneq \text{BMO}_{\text{rect},2}^d$ .*

**Proof.** We shall find a sequence  $\varphi_N$  such that  $\sup_N \|\varphi_N\|_{\text{rect},2} < \infty$  but  $\sup_N \|\varphi_N\|_p = \infty$ . A standard argument then gives the existence of  $\phi$ .

From Carleson's construction [C] we know that for each  $N \in \mathbb{N}$  there exists a collection of dyadic rectangles  $\Phi_N$  such that

$$(4.4) \quad \sum_{R \in \Phi_N} |R| = 1$$

$$(4.5) \quad \left| \bigcup_{R \in \Phi_N} R \right| < \frac{1}{N}$$

$$(4.6) \quad \sum_{R \in \Phi_N, R \subseteq R'} |R| \leq C |R'|, \quad R' \in \mathcal{R}$$

where  $C$  is a constant independent of  $N$ .

Defining

$$\varphi_N = \sum_{R \in \Phi_N} |R|^{1/2} h_R$$

we have that

$$\|\varphi_N\|_2 = 1, \quad \|\varphi_N\|_{\text{rect},2} \leq C$$

but, since  $\text{supp}(\varphi_N) \subset \cup_{R \in \Phi_N} R$ ,

$$\|\varphi_N\|_p \geq \left| \bigcup_{R \in \Phi_N} R \right|^{\frac{1}{p} - \frac{1}{2}} \geq N^{\frac{1}{2} - \frac{1}{p}}. \quad \blacksquare$$

We now can answer in the negative the above mentioned question of C. Sadosky and S. Ferguson posed in [FS].

**Corollary 4.4** *There exists  $\phi \in \text{BMO}_{\text{rect},1}^d \setminus \bigcup_{p>1} L^p(\mathbb{T}^2)$ . In particular, for each  $p > 1$ ,  $\text{BMO}_{\text{rect},p}^d \subsetneq \text{BMO}_{\text{rect},1}^d$ , and the norms  $\|\cdot\|_{\text{rect},1}$  and  $\|\cdot\|_{\text{rect},p}$  are not equivalent.*

**Proof.** We use the sequence of functions  $(\varphi_n)_{n \in \mathbb{N}}$  with

$$\|\varphi_n\|_{\text{rect},2} \leq C \text{ and } \|\varphi_n\|_p \geq n^{1/2-1/p}$$

for each  $n \in \mathbb{N}$ ,  $p > 2$  from Proposition 4.3.

Define

$$\phi = \sum_{n=1}^{\infty} \frac{1}{n^2} S_{\varphi_{2^n}}.$$

Then  $\phi \in \text{BMO}_{\text{rect},1}$  by Proposition 4.1, but  $\|S_{\varphi_{2^n}}\|_p \approx \|\varphi_{2^n}\|_{2p}^2 \geq 2^{n(1-1/p)}$  for each  $n \in \mathbb{N}$ ,  $p > 1$  and consequently  $\phi \notin \bigcup_{p>1} L^p(\mathbb{T}^2)$ .  $\blacksquare$

To differentiate the spaces  $\text{BMO}_{\text{rect},p}^d$  and  $\text{BMO}_{\text{prod}}^d$  we shall introduce the following coefficients.

**Definition 4.5** *Let  $E_n = \text{span}\{h_{I \times J} : I \in \mathcal{D}_1, J \in \mathcal{D}_2, |I|, |J| \geq 2^{-n}\}$  and let  $P_n$  be the orthogonal projection onto  $E_n$  in  $L^2(\mathbb{T}^2)$ .*

*For each  $q \geq 1$  and each  $n \in \mathbb{N}$*

$$c(n, q) = \sup\{\|\pi_\varphi\| : \varphi \in E_n, \|\varphi\|_{\text{rect},q} \leq 1\},$$

*and for  $p \geq q$ ,*

$$a(n, p, q) = \sup\{\|\varphi\|_{\text{rect},p} : \varphi \in E_n, \|\varphi\|_{\text{rect},q} \leq 1\},$$

We first analyse the behaviour of these constants. Of course we have

$$\begin{aligned}
 (4.7) \quad & c(n, p_2) \leq c(n, p_1), \quad p_1 \leq p_2 \\
 (4.8) \quad & a(n, p, q_1) \leq a(n, p, q_2), \quad q_1 \leq q_2 \\
 (4.9) \quad & a(n, p_2, q) \leq a(n, p_1, q), \quad p_1 \leq p_2.
 \end{aligned}$$

If  $p \geq q$ , clearly

$$(4.10) \quad c(n, q) \leq a(n, p, q)c(n, p).$$

Let us now extend Theorem 4.2.

**Theorem 4.6** *Let  $p > q \geq 2$  and  $\varphi \in \text{BMO}_{\text{prod}}^d$ . If  $q \leq 2^k \leq p$  for some  $k \in \mathbb{N}$  then*

$$(4.11) \quad \|\varphi\|_{\text{rect}, p} \leq C_{p,q} \|\varphi\|_{\text{prod}}^{1-q/p} \|\varphi\|_{\text{rect}, q}^{q/p}.$$

*In particular, for  $p > q_1 \geq q_2 \geq 2$  we have*

$$(4.12) \quad a(n, p, q_2) \leq C_p c(n, q_2)^{1-q_1/p} a(n, q_1, q_2)^{q_1/p}.$$

**Proof.** We shall see first that

$$(4.13) \quad \|\varphi\|_p \leq C_{p,q} \|\varphi\|_{\text{prod}}^{1-\theta} \|\varphi\|_q^\theta$$

for the above values of  $\theta = q/p$ . We do this in several steps.

First suppose that  $q = 2^n$  for some  $n \in \mathbb{N}$ . Theorem 4.2 gives the case  $n = 1$ . Assume that the result is true for  $n \geq 2$ , and let us consider the case  $q = 2^{n+1}$ .

Let  $p > 2^{n+1}$ . Applying the induction assumption to  $S_\varphi$  for  $p/2$ , we get

$$\begin{aligned}
 \|\varphi\|_p^2 & \approx \|S_\varphi\|_{p/2} \\
 & \leq C \|S_\varphi\|_{\text{prod}}^{1-2^{n+1}/p} \|S_\varphi\|_{2^n}^{2^{n+1}/p} \\
 & \leq C \|\varphi\|_{\text{prod}}^{2(1-2^{n+1}/p)} \|\varphi\|_{2^{n+1}}^{2^{n+2}/p}.
 \end{aligned}$$

This shows that

$$\|\varphi\|_p \leq C \|\varphi\|_{\text{prod}}^{1-2^{n+1}/p} \|\varphi\|_{2^{n+1}}^{2^{n+1}/p}.$$

Let us now proceed to the general case. We may assume that  $q < 2^k \leq p$  for some  $k \in \mathbb{N}$ . We can apply the previous case for  $n = k$  together with interpolation. Writing

$$\frac{1}{2^k} = \frac{1-\alpha}{q} + \frac{\alpha}{p},$$

we obtain

$$\begin{aligned} \|\varphi\|_p &\leq C\|\varphi\|_{\text{prod}}^{1-2^k/p}\|\varphi\|_{2^k}^{2^k/p} \\ &\leq C\|\varphi\|_{\text{prod}}^{1-2^k/p}(\|\varphi\|_q^{1-\alpha}\|\varphi\|_p^\alpha)^{2^k/p} \end{aligned}$$

Consequently

$$\|\varphi\|_p^{1-\alpha 2^k/p} \leq C\|\varphi\|_{\text{prod}}^{1-2^k/p}\|\varphi\|_q^{(1-\alpha)2^k/p}.$$

Note that

$$(1 - \alpha)\frac{2^k}{q} = 1 - \alpha\frac{2^k}{p}.$$

Hence we get with  $\theta = q/p$  that

$$\|\varphi\|_p \leq C\|\varphi\|_{\text{prod}}^{1-\theta}\|\varphi\|_q^\theta.$$

To finish the proof, note that for each  $R \in \mathcal{R}$ ,

$$\begin{aligned} \|P_R\varphi\|_p &\leq C_{p,q}\|P_R\varphi\|_{\text{prod}}^{1-q/p}\|P_R\varphi\|_q^{q/p} \\ &\leq C_{p,q}\|\varphi\|_{\text{prod}}^{1-q/p}\|\varphi\|_{\text{rect},q}^{q/p}|R|^{1/p}. \end{aligned}$$

■

Let us now establish a further connection between the constants introduced in 4.5.

**Theorem 4.7** *There exist  $K_1 > 0$  and  $K_2 > 0$  such that for all  $n \in \mathbb{N}$  and  $p \geq 1$*

$$c^2(n, 2p) \leq K_1 C_p^2 c(n, p) + K_2,$$

where  $C_p = \|\mathcal{S}\|_{L^{2p} \rightarrow L^{2p}}$ .

**Proof.** Write

$$\pi_\varphi^* \pi_\varphi = \Lambda_{S_\varphi} + D_\varphi,$$

with  $\|D_\varphi\| \leq C\|\varphi\|_{\text{BMO}_{\text{rect}}^d}^2$  as above.

For  $n \in \mathbb{N}$ ,  $p \geq 1$  and  $\varepsilon > 0$ , choose  $f_n \in L^2(\mathbb{T}^2)$  and  $\varphi \in E_n$  with  $\|\varphi\|_{\text{rect},2p} = 1$ ,  $\|f_n\|_2 = 1$  and  $\|\pi_\varphi f_n\|_2 \geq (1 - \varepsilon)c(n, 2p)$ . Then

$$(1 - \varepsilon)^2 c(n, 2p)^2 \leq \|\pi_\varphi f_n\|_2^2 = \langle \pi_\varphi^* \pi_\varphi f_n, f_n \rangle = \langle \Lambda_{S_\varphi} f_n, f_n \rangle + \langle D_\varphi f_n, f_n \rangle.$$

Therefore, we obtain that

$$(4.14) \quad (1 - \varepsilon)^2 c(n, 2p)^2 \leq \|\Lambda_{S_\varphi}\| + C\|\varphi\|_{\text{BMO}_{\text{rect}}^d}^2.$$



Since  $\|\varphi\|_{\text{rect},2p} = 1$ , Proposition 4.1 implies  $\|S_\varphi\|_{\text{rect},p} \leq 4C_p^2$ . Therefore, since  $\|\varphi\|_{\text{rect},2} \leq \|\varphi\|_{\text{rect},2p}$ , it follows that

$$(4.15) \quad (1 - \varepsilon)^2 c(n, 2p)^2 \leq 4C_p^2 c(n, p) + C.$$

Using (4.10) we get the second part. ■

**Corollary 4.8** *Let  $p \geq 1$ . Then  $\text{BMO}_{\text{prod}}^d \subsetneq \text{BMO}_{\text{rect},p}^d$ .*

**Proof.** Observe first that Proposition 4.3 implies that

$$(4.16) \quad \lim_{n \rightarrow \infty} c(n, 2) = \infty.$$

This shows that  $\text{BMO}_{\text{prod}}^d \subsetneq \text{BMO}_{\text{rect},p}^d$  for any  $1 \leq p \leq 2$ .

On the other hand, the estimates (4.10) and (4.12) in the case  $q = q_1 = q_2$  imply that if  $p > q \geq 2$  with  $q \leq 2^k \leq p$  for some  $k$ ,

$$(4.17) \quad c^{q/p}(n, q) \leq Cc(n, p),$$

where  $C$  is independent of  $n$ .

Hence  $\text{BMO}_{\text{prod}}^d = \text{BMO}_{\text{rect},p}^d$  for some  $p > 2$  would imply  $\sup_n c(n, p) < \infty$  and therefore  $\sup_n c(n, 2) < \infty$ , contradicting (4.16). ■

The particular case  $p = 4$  means that a question left open in [PS] can be answered in the negative. There, it was asked whether the condition

$$(4.18) \quad \begin{aligned} \|(\pi_b^{(1,2)})^* \pi_b^{(1,2)} h_{I'} f\|^2 &= \frac{1}{|I'|} \left\| \sum_{I \times J \in \mathcal{D}_1 \times \mathcal{D}_2, I \subsetneq I'} \frac{\chi_{I \times J}}{|I||J|} |b_{IJ}|^2 m_J f \right\|_{L^2(\mathbb{T}^2)}^2 \\ &\leq C \|f\|_{L^2(\mathbb{T})} \quad (f \in L^2(\mathbb{T}), I' \in \mathcal{D}_1) \end{aligned}$$

((27) and (28) in [PS]) already implies that  $b \in \text{BMO}_{\text{prod}}^d$ . Note that  $f$  here denotes a function in the second variable. We know from Prop 4.1 that  $b \in \text{BMO}_{\text{rect},4}$  implies  $S_b \in \text{BMO}_{\text{rect}}^d$ . By Lemma 3.2,

$$\|(\pi_b^{(1,2)})^* \pi_b^{(1,2)} h_{I'} f\| = \|(\Lambda_{S_b} + D_b) h_{I'} f\|,$$

where  $D_b$  is bounded on  $L^2(\mathbb{T}^2)$  and  $\Lambda_{S_b}$  maps  $L^2(\mathbb{T}) \hat{\otimes} L^2(\mathbb{T})$  boundedly into  $L^2(\mathbb{T}^2)$  by Prop 2.11. Thus  $b \in \text{BMO}_{\text{rect},4}$  implies (4.18). This condition is therefore not sufficient for  $b \in \text{BMO}_{\text{prod}}^d$ .

As pointed out in [PS], this has also consequences for the study of operator-valued Carleson measures, in the sense that a certain vector BMO condition of the sweep of an operator-valued measure does not imply boundedness of the corresponding vector Carleson embedding.

We can further show that even the intersection of all  $\text{BMO}_{\text{rect},p}^d$  spaces is still bigger than  $\text{BMO}_{\text{prod}}^d$ .

**Corollary 4.9**

$$\text{BMO}_{\text{prod}}^d \subsetneq \bigcap_{p \geq 1} \text{BMO}_{\text{rect},p}^d .$$

**Proof.** Obviously

$$\bigcap_{p \geq 1} \text{BMO}_{\text{rect},p}^d = \bigcap_{p \in \mathbb{N}} \text{BMO}_{\text{rect},p}^d .$$

With the locally convex topology defined by the increasing sequence of seminorms  $(\|\cdot\|_{\text{rect},p})_{p \in \mathbb{N}}$ , the latter is a metrizable locally convex linear space. Since each of the  $\text{BMO}_{\text{rect},p}^d$  is complete in  $\|\cdot\|_{\text{rect},p}$ ,  $\bigcap_{p \in \mathbb{N}} \text{BMO}_{\text{rect},p}^d$  is complete in this topology and therefore a Fréchet space. We know from Theorem 4.2 that

$$\text{BMO}_{\text{prod}}^d \subseteq \bigcap_{p \geq 1} \text{BMO}_{\text{rect},p}^d ,$$

and that the embedding is continuous with respect to the norm topology on  $\text{BMO}_{\text{prod}}^d$  and the locally convex topology on  $\bigcap_{p \geq 1} \text{BMO}_{\text{rect},p}^d$ . Let us assume towards a contradiction that the embedding is surjective. Then the open mapping theorem implies that the locally convex topology on  $\bigcap_{p \geq 1} \text{BMO}_{\text{rect},p}^d$  is normable with  $\|\cdot\|_{\text{prod}}$  and therefore contains a nonempty open neighbourhood of 0 which is bounded with respect to  $\|\cdot\|_{\text{prod}}$ . Since the family  $(\|\cdot\|_{\text{rect},p})_{p \in \mathbb{N}}$  is increasing, this means that there exists  $p \in \mathbb{N}$  and  $\varepsilon > 0$  such that  $\|b\|_{\text{prod}} < 1$  whenever  $\|b\|_{\text{rect},p} < \varepsilon$ , in contradiction to Corollary 4.8. ■

We will now separate the  $\text{BMO}_{\text{rect},p}^d$  spaces. Note that for Corollary 4.4 means that

$$(4.19) \quad \lim_{n \rightarrow \infty} a(n, p, 1) = \infty$$

for all  $p > 1$ .

Let us see that this holds in general.

**Corollary 4.10** *Let  $p > q \geq 1$ . Then  $\text{BMO}_{\text{rect},p}^d \subsetneq \text{BMO}_{\text{rect},q}^d$ .*

**Proof.** We have to show that

$$\lim_{n \rightarrow \infty} a(n, p, q) = \infty .$$

It suffices to prove

$$\lim_{n \rightarrow \infty} a(n, q + \varepsilon, q) = \infty$$

for sufficiently small  $\varepsilon$ . For fixed  $q > 1$ , choose  $\varepsilon > 0$  and  $k \in \mathbb{N}$  such that  $q < q + \varepsilon < 2^k \leq 2q$ .

Using Theorem 4.7, (4.10) and (4.17), we obtain constants  $C_1, C_2$  and  $C_3$  independent of  $n$  such that

$$\begin{aligned} c^2(n, 2q) &\leq C_1 c(n, q) \leq C_1 C_2 a(n, q + \varepsilon, q) c(n, q + \varepsilon) \\ &\leq C_1 C_2 C_3 a(n, q + \varepsilon, q) c(n, 2q)^{\frac{2p}{p+\varepsilon}}. \end{aligned}$$

This shows that

$$c(n, 2q)^{\frac{2\varepsilon}{p+\varepsilon}} \leq C a(n, q + \varepsilon, q)$$

where  $C$  is independent of  $n$ . Now the result follows from Corollary 4.8. ■

**Acknowledgement.** We thank C. Sadosky for fruitful discussions.

## References

- [Be] BERNARD, A.: Espaces  $H^1$  de martingales à deux indices. Dualité avec les martingales de type “BMO” (French). *Bull. Sci. Math. (2)* **103** (1979), no. 3, 297–303.
- [C] CARLESON, L.: A counterexample for measures bounded on  $H^p$  for the bi-disc. Mittag-Leffler Report No. **7**, 1974.
- [Ch] CHANG, S.-Y. A.: Carleson measure on the bi-disc. *Ann. of Math. (2)* **109** (1979), no. 3, 613–620.
- [ChFef1] CHANG, S.-Y. A. AND FEFFERMAN, R.: A continuous version of duality of  $H^1$  with BMO on the bidisc. *Ann. of Math. (2)* **112** (1980), no. 1, 179–201.
- [ChFef2] CHANG, S.-Y. A. AND FEFFERMAN, R.: Some recent developments in Fourier analysis and  $H^p$  theory on product domains. *Bull. Amer. Math. Soc. (N. S.)* **12** (1985), 1–43.
- [Fef] FEFFERMAN, R.: Bounded mean oscillation on the polydisc. *Ann. of Math. (2)* **110** (1979), 395–406.
- [FS] FERGUSON, S. AND SADOSKY, C.: Characterizations of bounded mean oscillation on the polydisk in terms of Hankel operators and Carleson measures. *J. Anal. Math.* **81** (2000), 239–267.
- [GPTV] GILLESPIE, T. A., POTT, S., TREIL, S. AND VOLBERG, A.: Logarithmic growth for matrix martingale transforms. *J. London Math. Soc. (3)* **64** (2001), 624–636.
- [LF] LACEY, M. AND FERGUSON, S.: A characterization of product BMO by commutators. *Acta Math.* **189** (2002), no. 2, 143–160.
- [Per] PEREYRA, C. M.: Lecture notes on dyadic harmonic analysis. In *Second Summer School in Analysis and Mathematical Physics (Cuernavaca, 2000)*, 1–60. Contemp. Math. **289**. Amer. Math. Soc., Providence, 2001.

- [PS] POTT, S. AND SADOSKY, C.: Bounded mean oscillation on the bidisk and Operator BMO. *J. Funct. Anal.* **189** (2002), 475–495.
- [T] TAO, T.: Dyadic product  $H^1$ , BMO and Carleson’s Counterexample. Short stories, <http://www.math.ucla.edu/~tao/>

*Recibido:* 7 de febrero de 2003

*Revisado:* 2 de octubre de 2003

Óscar Blasco  
Department of Mathematics  
Universitat de Valencia  
Burjassot 46100 (Valencia), Spain  
[Oscar.Blasco@uv.es](mailto:Oscar.Blasco@uv.es)

Sandra Pott  
Department of Mathematics  
University of Glasgow  
Glasgow G12 8QW, UK  
[sp@maths.gla.ac.uk](mailto:sp@maths.gla.ac.uk)

---

The first author gratefully acknowledges support by the Spanish grants PR2001-0134, BFM2002-04013 and by the LMS. The second author acknowledges support by EPSRC and by the Nuffield Foundation.