

# Construction of multi-soliton solutions for the $L^2$ -supercritical gKdV and NLS equations

Raphaël Côte, Yvan Martel and Frank Merle

## Abstract

Multi-soliton solutions, i.e. solutions behaving as the sum of  $N$  given solitons as  $t \rightarrow +\infty$ , were constructed for the  $L^2$  critical and subcritical (NLS) and (gKdV) equations in previous works (see [23], [16] and [20]). In this paper, we extend the construction of multi-soliton solutions to the  $L^2$  supercritical case both for (gKdV) and (NLS) equations, using a topological argument to control the direction of instability.

## 1. Introduction

### 1.1. The generalized KdV equation

We consider the generalized Korteweg-de Vries equations:

$$(\text{gKdV}) \quad u_t + (u_{xx} + u^p)_x = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R},$$

where  $p \geq 2$  is an integer. See Section 3.1 for more general nonlinearities.

Recall that the Cauchy problem for (gKdV) in the energy space  $H^1$  has been solved by Kenig, Ponce and Vega [14]: for all  $u_0 \in H^1(\mathbb{R})$ , there exist  $T = T(\|u_0\|_{H^1}) > 0$  and a solution  $u \in \mathcal{C}([0, T], H^1(\mathbb{R}))$  to (gKdV) satisfying  $u(0) = u_0$ , unique in some sense. Moreover, if  $T_1$  denotes the maximal time of existence for  $u$ , then either  $T_1 = +\infty$  (global solution) or  $T_1 < \infty$  and then  $\|u(t)\|_{H^1} \rightarrow \infty$  as  $t \uparrow T_1$  (blow-up solution).

---

*2000 Mathematics Subject Classification:* 35Q51, 35Q53, 35Q55.

*Keywords:* Multi-solitons, generalized Korteweg-de Vries equation, nonlinear Schrödinger equation, instability, supercritical problem.

For such solutions, the mass and energy are conserved:

$$(1.1) \quad \int u^2(t) = \int u^2(0) \quad (L^2 \text{ mass}),$$

$$(1.2) \quad E(u(t)) = \frac{1}{2} \int u_x^2(t) - \frac{1}{p+1} \int u^{p+1}(t) = E(u(0)) \quad (\text{energy}).$$

Now, we define  $Q \in H^1$ ,  $Q > 0$  the unique solution (up to translations) to

$$Q_{xx} + Q^p = Q, \quad \text{i.e.} \quad Q(x) = \left( \frac{p+1}{2 \cosh^2(\frac{p-1}{2}x)} \right)^{\frac{1}{p-1}}.$$

Let  $Q_{c_0}(x) = c_0^{\frac{1}{p-1}} Q(\sqrt{c_0}x)$  and let

$$R_{c_0, x_0}(t, x) = c_0^{\frac{1}{p-1}} Q(\sqrt{c}(x - c_0t - x_0))$$

be the family of soliton solution of the (gKdV) equation.

It is well-known that the stability properties of a soliton  $R_{c_0, x_0}$  depend on the sign of  $\frac{d}{dc} \int Q_c^2|_{c=c_0}$ . Since  $\int Q_c^2 = c^{\frac{5-p}{2(p-1)}} \int Q^2$ , we distinguish the following three cases:

- For  $p < 5$  ( $L^2$  subcritical case), solitons are stable and asymptotically stable in  $H^1$  in some suitable sense: see Cazenave and Lions [3], Weinstein [30], Grillakis, Shatah and Straus [12], for orbital stability, and Pego and Weinstein [27], Martel and Merle [17] for asymptotic stability.
- In the  $L^2$  critical case, i.e.  $p = 5$ , solitons are unstable, and blow up occur for a large class of solutions initially arbitrarily close to a soliton, see Martel and Merle [18], [19].
- In the case  $p > 5$  ( $L^2$  supercritical case), solitons are unstable (see Grillakis, Shatah and Straus [12] and Bona, Souganidis and Strauss [2]).

Now, we focus on multi-soliton solutions. Given  $2N$  parameters defining  $N$  solitons with different speeds,

$$(1.3) \quad 0 < c_1 < \dots < c_N, \quad x_1, \dots, x_N \in \mathbb{R},$$

we call multi-soliton a solution  $u(t)$  to (gKdV) such that

$$(1.4) \quad \lim_{t \rightarrow +\infty} \left\| u(t) - \sum_{j=1}^N R_{c_j, x_j}(t) \right\|_{H^1} = 0.$$

Let us recall known results on multi-solitons:

- For  $p = 2$  and  $3$  (KdV and mKdV), multi-solitons are well-known to exist for any set of parameters (1.3), as a consequence of the inverse

scattering method. Moreover, these special explicit solutions describe the elastic collision of the solitons (see e.g. Miura [24]).

- In the  $L^2$ -subcritical and critical cases, i.e. for (gKdV) with  $p \leq 5$  (or for some more general nonlinearities under the stability assumption  $\frac{d}{dc} \int Q_{c|c=c_j}^2 > 0$  for all  $j$ ), Martel [16] constructed multi-solitons for any set of parameters (1.3). The proof of this result follows the strategy of Merle [23] (compactness argument) and relies on monotonicity properties developed in [17] (see also [21]). Recall that Martel, Merle and Tsai [21] proved stability and asymptotic stability of a sum of  $N$  solitons for large time for the subcritical case. A refined version of the stability result of [21] shows that for a given set of parameters, there exists a *unique* multi-soliton soliton satisfying (1.4), see Theorem 1 in [16].

In the present paper, we extend the multi-soliton existence result to the  $L^2$ -supercritical case, i.e., in a situation where solitons are known to be unstable.

**Theorem 1** (Existence of multi-solitons for  $L^2$ -supercritical (gKdV)). *Let  $p > 5$ . Let  $0 < c_1 < \dots < c_N$  and  $x_1, \dots, x_N \in \mathbb{R}$ . There exist  $T_0 \in \mathbb{R}$ ,  $C, \sigma_0 > 0$ , and a solution  $u \in \mathcal{C}([T_0, \infty), H^1)$  to (gKdV) such that*

$$\forall t \in [T_0, \infty), \quad \left\| u(t) - \sum_{j=1}^N R_{c_j, x_j}(t) \right\|_{H^1} \leq C e^{-\sigma_0^{3/2} t}.$$

**Remark 1.** As in the subcritical case, the proof of Theorem 1 is based on a compactness argument and on some large time uniform estimates, however, it also involves an additional topological argument to control an instable direction of the linearized operator around each  $Q_{c_j}$ . The proof relies decisively on the introduction of  $L^2$  eigenfunctions of the linearized operator, constructed by Pego and Weinstein [26] by ODE techniques. Note that in [26], the existence of such eigenfunctions for  $Q_{c_0}$  is proved to be equivalent to  $\frac{d}{dc} \int Q_{c|c=c_0}^2 < 0$ .

It is possible that other methods of construction work for some range of parameters  $0 < c_1 < \dots < c_N$ , but due to the instable directions, the use of such a topological argument is probably necessary to treat the general case (1.3).

Finally, note that the solution  $u(t)$  of Theorem 1 belongs to  $H^s$ , and that the convergence to  $\sum_{j=1}^N R_{c_j, x_j}(t)$  holds in  $H^s$ , for any  $s \geq 1$  (see Proposition 5 of [16]).

We refer to Section 3.1 for a similar existence result for (gKdV) equations with general nonlinearities.

### 1.2. The non linear Schrödinger equations

Now we turn to the case of the non linear Schrödinger equations:

$$(NLS) \quad iu_t + \Delta u + |u|^{p-1}u = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^d, \quad u(t, x) \in \mathbb{C},$$

where  $p > 1$ , for any space dimension  $d \geq 1$ . Concerning the local well-posedness of the Cauchy problem in  $H^1$ , we refer to Ginibre and Velo [10]. Recall that  $H^1$  solutions satisfy the conservation laws

$$\begin{aligned} \int |u|^2(t) &= \int |u_0|^2, \quad \text{Im} \int (\bar{u} \nabla u)(t) = \text{Im} \int \bar{u}_0 \nabla u_0, \\ \frac{1}{2} \int |\nabla u(t)|^2 - \frac{1}{p+1} \int |u|^{p+1}(t) &= \frac{1}{2} \int |\nabla u_0(t)|^2 - \frac{1}{p+1} \int |u_0|^{p+1}. \end{aligned}$$

Consider the radial positive solution  $Q \in H^1(\mathbb{R}^d)$  to

$$(1.5) \quad \Delta Q + Q^p = Q,$$

which is the unique positive solution of this equation up to translations. We refer to [9], [1] and [15] for classical existence and uniqueness results on equation (1.5). Given  $v_0, x_0 \in \mathbb{R}^d$ ,  $\gamma_0 \in \mathbb{R}$  and  $c_0 > 0$ , the function

$$R_{c_0, \gamma_0, v_0, x_0}(t, x) = c_0^{\frac{1}{p-1}} Q(\sqrt{c_0}(x - v_0 t - x_0)) e^{i(\frac{1}{2}v_0 \cdot x - \frac{1}{4}\|v_0\|^2 t + c_0 t + \gamma_0)}$$

is a soliton solution to (NLS), moving on the line  $x = x_0 + v_0 t$ .

We recall the following classical results (for any  $d \geq 1$ ):

- For  $1 < p < 1 + 4/d$ , ( $L^2$  subcritical case) Cazenave and Lions [3] proved that solitons are orbitally stable in  $H^1$ . Multi-solitons (defined in a similar way as for (gKdV)) were constructed in this setting by Martel and Merle [20].
- In the  $L^2$  critical case,  $p = 1 + 4/d$ , solitons are unstable, however multi-solitons were constructed by Merle [23], as a consequence of the construction of special solutions of (NLS) blowing up in finite time at  $N$  prescribed points.
- For  $p \in (1 + \frac{4}{d}, \frac{d+2}{d-2})$  (for  $d = 1, 2$ ,  $p > 1 + \frac{4}{d}$ ): solitons are unstable (see [12]). Recall that  $p = \frac{d+2}{d-2}$  corresponds to the critical  $\dot{H}^1$  case.

We claim the following analogue of Theorem 1 in the context of the  $L^2$  supercritical (NLS) equation.

**Theorem 2** (Multi-solitons for  $L^2$  supercritical (NLS)). *Let  $p \in (1 + \frac{4}{d}, \frac{d+2}{d-2})$  ( $p > 1 + \frac{4}{d}$  for  $d = 1, 2$ ). Let  $c_1, \dots, c_N > 0$ ,  $\gamma_1, \dots, \gamma_N \in \mathbb{R}$ ,  $x_1, \dots, x_N \in \mathbb{R}^d$ , and  $v_1, \dots, v_N \in \mathbb{R}^d$  be such that*

$$\forall k \neq j, \quad v_k \neq v_j.$$

*Then there exist  $T_0 \in \mathbb{R}$ ,  $C, \sigma_0 > 0$ , and a solution  $u \in \mathcal{C}([T_0, \infty), H^1)$  to (NLS) such that*

$$\forall t \in [T_0, \infty), \quad \left\| u(t) - \sum_{j=1}^N R_{c_j, \gamma_j, v_j, x_j}(t) \right\|_{H^1} \leq C e^{-\sigma_0^{3/2} t}.$$

**Remark 2.** The condition on  $p$  means that the problem is  $L^2$  supercritical but  $\dot{H}^1$  subcritical (for  $d \geq 3$ ). In the present paper, we do not treat the  $\dot{H}^1$  critical case – recall that solitons then have only algebraic decay.

The proof of Theorem 2 is completely similar to the one of Theorem 1, see Section 3.2. Note that similarly to the (gKdV) case, we need eigenfunctions for the linearized operator around  $Q$ . For the (NLS) case, see Weinstein [29], Grillakis [11] and Schlag [28].

In Section 1.3, we present an outline of the proof of Theorem 1. A complete proof of Theorem 1 is given in Section 2. Next, extensions of this result to (gKdV) equations with general nonlinearities are presented without proof in Section 3.1. Finally, a sketch of the proof of Theorem 2 is given in Section 3.2. In the Appendix, we gather the proof of two technical lemmas.

### 1.3. Outline of proof of Theorem 1

For simplicity, we consider only positive solitons and pure power nonlinearities for (gKdV). The proof follows a similar initial strategy as in the works of Merle [23] or Martel [16]. Set

$$R_j(t, x) = R_{c_j, x_j}(t, x), \quad R(t, x) = \sum_{j=1}^N R_j(t, x),$$

and consider a sequence  $S_n \rightarrow +\infty$ .

In the subcritical case ([16] and [20]), one considers the sequence  $(u_n)$  of solutions to (gKdV) such that  $u_n(S_n) = R(S_n)$ . The goal is then to obtain backwards uniform estimates on  $u_n(t) - R(t)$  on some time interval  $t \in [T_0, S_n]$ , where  $T_0$  does not depend on  $n$ . From these estimates, one can construct the multi-soliton soliton by compactness arguments. To obtain the uniform estimates, one uses monotonicity properties of local conservation laws and coercivity property of the Hessian of the energy around a soliton:

$$Lv = -v_{xx} - pQ^{p-1}v + v.$$

Indeed, in the subcritical case, it is well-known (see [30]) that  $(Lv, v) \geq \lambda \|v\|_{H^1}^2$  ( $\lambda > 0$ ) provided that  $(v, Q) = (v, Q_x) = 0$ . These two directions are then controlled by modulation with respect to scaling and translation.

In the supercritical case, one cannot obtain uniform estimates in the same way, since the previous property of  $L$  fails. It is known that  $(L\cdot, \cdot)$  is positive definite up to the directions  $Q^{\frac{p+1}{2}}$  and  $Q_x$ ; the direction  $Q_x$  can still be handled using modulation in the translation parameter, but the even direction  $Q^{\frac{p+1}{2}}$  cannot be controlled by the scaling parameter as for the subcritical case (this is of course related to the instable nature of the soliton).

At this point, we need the  $L^2$  eigenfunctions  $Z^\pm$  of the operator  $L\partial_x$ :

$$L(Z_x^\pm) = \pm e_0 Z^\pm, \quad e_0 > 0,$$

constructed by Pego and Weinstein [26]. Following Duyckaerts and Merle [5], we prove that  $(L\cdot, \cdot)$  is positive definite up to the directions  $Z^\pm$  and  $Q_x$  (see Lemma 1 in the present paper). The direction  $Z^-$  being in some sense a stable direction, it does not create any difficulty. For the instable direction  $Z^+$ , we do need an extra parameter, which cannot be controlled by a scaling argument. Thus, instead of considering the final data  $u_n(S_n) = R(S_n)$ , as in [16], we look at solutions  $u_n$  to (gKdV) with final data:

$$u_n(S_n) = R(S_n) + \sum_{j,\pm} \mathbf{b}_{j,n}^\pm Z_j^\pm, \quad \text{where } Z_j^\pm(t, x) = c_j^{\frac{1}{p-1}} Z^\pm(\sqrt{c_j}(x - c_j t - x_j)),$$

and  $\mathbf{b}_n = (\mathbf{b}_{j,n}^\pm)_{j=1,\dots,N;\pm}$  belongs to some small neighborhood of 0 in  $\mathbb{R}^{2N}$ . A topological argument then allows us to select, for all  $n$ , a value of  $\mathbf{b}_n$  so that a uniform estimate of  $\|u_n(t) - R(t)\|_{H^1}$  holds on some interval  $[T_0, S_n]$ .

## 2. Proof of Theorem 1

### 2.1. Preliminary results

Consider the operator

$$Lv = -v_{xx} - pQ^{p-1}v + v.$$

For  $p > 5$ , it is known from the work of Pego and Weinstein [26] that the operator  $\partial_x L$  has two eigenfunctions  $Y^+$  and  $Y^-$  (related by  $Y^-(x) = Y^+(-x)$ ) such that

$$(LY^\pm)_x = \pm e_0 Y^\pm, \quad \text{where } e_0 > 0.$$

In contrast with the (NLS) case (see references in section 3.2), the existence of  $Y^\pm$  is not obtained by variational arguments, but by sharp ODE techniques. Note that [26] provides a complete description of the spectrum of  $\partial_x L$

in  $L^2$  for any  $p > 1$  ; in particular, the existence of such eigenfunctions related to  $\pm e_0$  with  $e_0 > 0$  is proved to be equivalent to super criticality (i.e.  $p > 5$  in the present case).

Next, we observe that  $Z^\pm = LY^\pm$  are eigenfunctions of  $L\partial_x$  (adjoint to  $-\partial_x L$ ). Indeed,

$$L(Z_x^\pm) = \pm e_0 Z^\pm.$$

The functions  $Z^\pm$  are normalized so that  $\|Z^\pm\|_{L^2} = 1$ . Moreover, we recall from [26] (standard ODE arguments) that  $Z^\pm, Y^\pm \in \mathcal{S}(\mathbb{R})$  and have exponential decay, along with their derivatives. Let  $\eta_0 > 0$  such that

$$\forall x \in \mathbb{R}, \quad |Z^+(x)| + |Z^-(x)| + |Z_x^+(x)| + |Z_x^-(x)| \leq C e^{-\eta_0|x|}.$$

Following [5] (concerning the (NLS) case), we claim the following coercivity property of  $L$  (for  $f, g \in L^2$ ,  $(f, g) = \int f g$  denotes the scalar product in  $L^2$ ).

**Lemma 1.** *There exists  $\lambda > 0$  such that*

$$\forall v \in H^1, \quad (Lv, v) \geq \lambda \|v\|_{H^1}^2 - \frac{1}{\lambda} \left( (v, Z^+)^2 + (v, Z^-)^2 + (v, Q_x)^2 \right).$$

**Proof.** The proof is similar to the one of [5, Lemma 5.2]. It is given here for the reader’s convenience.

First we recall the following well-known result.

**Claim.** There exists  $\nu > 0$  such that

$$(2.1) \quad \forall v \in H^1, \quad (Lv, v) \geq \nu \|v\|_{H^1}^2 - \frac{1}{\nu} \left( (v, Q_x)^2 + (v, Q^{\frac{p+1}{2}})^2 \right).$$

Indeed,  $Q_x$  and  $Q^{\frac{p+1}{2}}$  are two eigenfunctions for  $L$ , namely

$$LQ_x = 0 \quad \text{and} \quad LQ^{\frac{p+1}{2}} = \mu_0 Q^{\frac{p+1}{2}}, \quad \text{where } \mu_0 = 1 - \left( \frac{p+1}{2} \right)^2 < 0.$$

The claim then follows from Sturm-Liouville theory.

To prove the Lemma, it suffices to show that

$$(2.2) \quad \text{if } (v, Z^+) = (v, Z^-) = (v, Q_x) = 0 \text{ then } (Lv, v) \geq \lambda \|v\|_{H^1}^2.$$

Let  $v$  satisfy the orthogonality conditions in (2.2) and decompose the functions  $v, Y^\pm$   $L^2$  orthogonally in  $\text{Span}(Q_x, Q^{\frac{p+1}{2}})^\perp$  and  $\text{Span}(Q_x, Q^{\frac{p+1}{2}})$  (the notation  $\perp$  is here related to  $L^2$  orthogonality).

$$v = w + \alpha Q^{\frac{p+1}{2}}, \quad Y^+ = y^+ + \beta Q_x + \gamma Q^{\frac{p+1}{2}}, \quad Y^- = y^- + \delta Q_x + \eta Q^{\frac{p+1}{2}}.$$

By the symmetry  $Y^+(-x) = Y^-(x)$  and uniqueness of the orthogonal decomposition, note that  $\delta = -\beta, \eta = \gamma$  and  $y^+(-x) = y^-(x)$ .

We claim that the functions  $y^+$ ,  $y^-$  are linearly independent. Indeed, decompose into even and odd parts

$$y^+ = y^e + y^o, \quad y^- = y^e - y^o.$$

As  $Y^\pm$ , the functions  $y^\pm$ ,  $y^e$  and  $y^o$  have exponential decay, along with their derivatives.

Let us prove that  $y^e \neq 0$  and  $y^o \neq 0$  ; we observe from  $(LY^+)_x = e_0 Y^+$  that

$$(Ly^e)_x + (Ly^o)_x - \mu_0 \gamma (Q^{\frac{p+1}{2}})_x = e_0 y^e + e_0 y^o + e_0 \beta Q_x + e_0 \gamma Q^{\frac{p+1}{2}}, \text{ i.e.}$$

$$(Ly^e)_x = e_0 (y^o + \beta Q_x) - \mu_0 \gamma (Q^{\frac{p+1}{2}})_x, \quad (Ly^o)_x = e_0 (y^e + \gamma Q^{\frac{p+1}{2}}).$$

If  $y^o = 0$ , then  $y^e = 0$  and  $\gamma = 0$ , hence  $\beta = 0$ , and thus  $Y^+ = Y^- = 0$ , which is a contradiction. Now, if we assume  $y^e = 0$ , by  $(Ly^o)_x = e_0 (y^e + \gamma Q^{\frac{p+1}{2}})$  and  $\int Q^{\frac{p+1}{2}} \neq 0$ , we obtain  $\gamma = 0$ . Thus, from  $0 = (Ly^e)_x = e_0 (y^o + \beta Q_x)$ , we get  $y^o = 0$  and  $\beta = 0$ , so that  $Y^+ = Y^- = 0$ , a contradiction. From the property  $y^e \neq 0$  and  $y^o \neq 0$ , one deduces that  $ay^+ + by^- = 0$  implies  $a = b = 0$ , hence  $y^+$  and  $y^-$  are linearly independent.

We now go back to the proof of coercivity. Note that

$$(LY^\pm, Y^\pm) = \pm e_0^{-1} (LY^\pm, (LY^\pm)_x) = 0.$$

We compute

$$\begin{aligned} 0 &= (v, Z^+) = (v, LY^+) = (Lv, Y^+) = (Lw, y^+) + \alpha \mu_0 \gamma \|Q^{\frac{p+1}{2}}\|_{L^2}^2, \\ 0 &= (v, Z^-) = (v, LY^-) = (Lv, Y^-) = (Lw, y^-) + \alpha \mu_0 \gamma \|Q^{\frac{p+1}{2}}\|_{L^2}^2, \\ 0 &= (LY^+, Y^+) = (Ly^+, y^+) + \gamma^2 \mu_0 \|Q^{\frac{p+1}{2}}\|_{L^2}^2, \\ 0 &= (LY^-, Y^-) = (Ly^-, y^-) + \gamma^2 \mu_0 \|Q^{\frac{p+1}{2}}\|_{L^2}^2. \end{aligned}$$

Hence

(2.3)

$$(Lv, v) = (Lw, w) + \mu_0 \alpha^2 \|Q^{\frac{p+1}{2}}\|_{L^2}^2 = (Lw, w) - \frac{(Lw, y^+)(Lw, y^-)}{\sqrt{(Ly^+, y^+)}\sqrt{(Ly^-, y^-)}}.$$

Consider

$$a = \sup_{\omega \in \text{Span}(y^+, y^-) \setminus \{0\}} \left| \frac{(L\omega, y^+)}{\sqrt{(L\omega, \omega)(Ly^+, y^+)}} \cdot \frac{(L\omega, y^-)}{\sqrt{(L\omega, \omega)(Ly^-, y^-)}} \right|.$$

Recall  $(L \cdot, \cdot)$  is positive definite on  $\text{Span}(Q_x, Q^{\frac{p+1}{2}})^\perp$  ; applying Cauchy-Schwarz inequality to each of the two terms of the product above, we find  $a \leq 1$ . Furthermore, if  $a = 1$ , there exists  $\omega \neq 0$  such that these two Cauchy-Schwarz inequalities are actually equalities, but this is not possible since  $y^+$  and  $y^-$  are independent.



Therefore, we have proved that  $a < 1$ . Using  $L$  orthogonal decomposition on  $\text{Span}(Q_x, Q^{\frac{p+1}{2}})^\perp$ , we also obtain that for all  $\omega \in \text{Span}(Q_x, Q^{\frac{p+1}{2}})^\perp$ ,

$$\left| \frac{(L\omega, y^+)(L\omega, y^-)}{\sqrt{(Ly^+, y^+)}\sqrt{(Ly^-, y^-)}} \right| \leq a(L\omega, \omega).$$

Hence, by (2.3) and next (2.1),

$$(Lv, v) \geq (1 - a)(Lw, w) \geq \nu(1 - a)\|w\|_{H^1}^2 > 0,$$

and so  $(Lw, w) \geq |\mu_0|\alpha^2\|Q^{\frac{p+1}{2}}\|_{L^2}^2$ .

Thus, for  $C = \frac{4}{1-a} \max(\frac{1}{\nu}, \frac{1}{|\mu_0|}\|Q^{\frac{p+1}{2}}\|_{H^1}^2\|Q^{\frac{p+1}{2}}\|_{L^2}^{-2})$  we get

$$\begin{aligned} C(Lv, v) &\geq C(1 - a)(Lw, w) \geq C\frac{1 - a}{2}(Lw, w) + C\frac{1 - a}{2}|\mu_0|\alpha^2\|Q^{\frac{p+1}{2}}\|_{L^2}^2 \\ &\geq 2\|w\|_{H^1}^2 + 2\alpha^2\|Q^{\frac{p+1}{2}}\|_{H^1}^2 \geq \|w + \alpha Q^{\frac{p+1}{2}}\|_{H^1}^2 = \|v\|_{H^1}^2. \end{aligned}$$

■

### 2.2. Main Proposition and proof of Theorem 1

We denote

$$(2.4) \quad \begin{aligned} R_j(t, x) &= c_j^{\frac{1}{p-1}}Q(\sqrt{c_j}(x - c_jt - x_j)), \quad R(t, x) = \sum_{j=1}^N R_j(t, x), \\ Z_j^\pm(t, x) &= c_j^{\frac{1}{p-1}}Z^\pm(\sqrt{c_j}(x - c_jt - x_j)). \end{aligned}$$

Let  $S_n \rightarrow \infty$  be a increasing sequence of time,  $\mathbf{b}_n = (\mathbf{b}_{j,n}^\pm)_{j,\pm} \in \mathbb{R}^{2N}$  be a sequence of parameters to be determined, and let  $u_n$  be the solution to

$$(2.5) \quad \begin{cases} u_{nt} + (u_{nxx} + u_n^p)_x = 0, \\ u_n(S_n) = R(S_n) + \sum_{j \in \{1, \dots, N\}, \pm} \mathbf{b}_{j,n}^\pm Z_j^\pm(S_n). \end{cases}$$

Let

$$(2.6) \quad \sigma_0 = \frac{1}{4} \min \{ \eta_0\sqrt{c_1}, e_0^{2/3}c_1, c_1, c_2 - c_1, \dots, c_N - c_{N-1} \}.$$

**Proposition 1.** *There exist  $n_0 \geq 0$ ,  $T_0 > 0$  and  $C > 0$  (independent of  $n$ ) such that the following holds: For each  $n \geq n_0$ , there exists  $\mathbf{b}_n = (\mathbf{b}_{j,n}^\pm)_{j,\pm} \in \mathbb{R}^{2N}$  with*

$$\left( \sum_{j,\pm} \mathbf{b}_{j,n}^\pm \right)^{1/2} \leq Ce^{-\sigma_0^{3/2}S_n},$$

and such that the solution  $u_n$  to (2.5) is defined on the interval  $[T_0, S_n]$ , and satisfies

$$\forall t \in [T_0, S_n], \quad \|u_n(t) - R(t)\|_{H^1} \leq Ce^{-\sigma_0^{3/2}t}.$$

Assuming this Proposition, we now deduce the proof of Theorem 1. The proof of Proposition 1 is postponed to Section 2.3.

**Proof of Theorem 1 assuming Proposition 1.** It follows closely the proof of Theorem 1 in [16]. We may assume  $n_0 = 0$  in Proposition 1 without loss of generality.

*Step 1: Compactness argument.* From Proposition 1, there exists a sequence  $u_n(t)$  of solutions to (gKdV), defined on  $[T_0, S_n]$  and  $C_0, \sigma_0 > 0$  such that the following uniform estimates hold:

$$(2.7) \quad \forall n \in \mathbb{N}, \forall t \in [T_0, S_n], \quad \|u_n(t) - R(t)\|_{H^1} \leq C_0 e^{-\sigma_0^{3/2} t}.$$

We claim the following compactness result on the sequence  $u_n(T_0)$ .

**Claim.**

$$\limsup_{A \rightarrow \infty} \sup_{n \in \mathbb{N}} \int_{|x| \geq A} u_n^2(T_0, x) dx = 0.$$

**Proof.** Let  $\varepsilon > 0$ ,  $T(\varepsilon) \geq T_0$  be such that  $C_0 e^{-\sigma_0^{3/2} T(\varepsilon)} \leq \sqrt{\varepsilon}$  and  $n$  large enough so that  $S_n \geq T(\varepsilon)$ . Then

$$\int (u_n(T(\varepsilon)) - R(T(\varepsilon)))^2 \leq \varepsilon.$$

Let  $A(\varepsilon)$  be such that  $\int_{|x| \geq A(\varepsilon)} R(T(\varepsilon))^2(x) dx \leq \varepsilon$ ; we get

$$\int_{|x| \geq A(\varepsilon)} u_n^2(T(\varepsilon), x) dx \leq 4\varepsilon.$$

Let  $g(x) \in C^3$  be such that  $g(x) = 0$  if  $x \leq 0$ ,  $g(x) = 1$  if  $x \geq 2$ , and furthermore  $0 \leq g'(x) \leq 1$ ,  $|g'''(x)| \leq 1$ .

Recall that for  $f(x) \in C^3$ , we have (Kato's identity [13])

$$\frac{d}{dt} \int u_n^2 f = -3 \int (u_{nx})^2 f_x + \int u_n^2 f_{xxx} + \frac{2p}{p+1} \int u_n^{p+1} f_x.$$

For  $C(\varepsilon) > 1$  to be determined later, we thus have:

$$\begin{aligned} \frac{d}{dt} \int u_n^2(t, x) g\left(\frac{x - A(\varepsilon)}{C(\varepsilon)}\right) &= -\frac{3}{C(\varepsilon)} \int (u_{nx})^2 g'\left(\frac{x - A(\varepsilon)}{C(\varepsilon)}\right) \\ &+ \frac{1}{C(\varepsilon)^3} \int u_n^2 g'''\left(\frac{x - A(\varepsilon)}{C(\varepsilon)}\right) + \frac{2p}{(p+1)C(\varepsilon)} \int u_n^{p+1} g'\left(\frac{x - A(\varepsilon)}{C(\varepsilon)}\right). \end{aligned}$$

For  $t \geq T_0 \geq 0$ ,  $u_n$  satisfies  $\|u_n(t)\|_{H^1} \leq C_0 + \sum_{j=1}^N \|Q_{c_j}\|_{H^1} \leq C^0$ , and so:

$$\begin{aligned} & \left| \frac{d}{dt} \int u_n^2(t, x) g\left(\frac{x - A(\varepsilon)}{C(\varepsilon)}\right) \right| \\ & \leq \frac{1}{C(\varepsilon)} \left( 3 \int u_{nx}^2(t) + \int u_n^2(t) + \frac{2p}{p+1} \|u_n\|_{L^\infty}^{p-1} \int u_n^2(t) \right) \\ & \leq \frac{1}{C(\varepsilon)} \left( 3C^{02} + \frac{2p}{p+1} 2^{(p-1)/2} C^{0p+1} \right). \end{aligned}$$

Now choose  $C(\varepsilon) = \max \left\{ 1, \frac{T(\varepsilon) - T_0}{\varepsilon} \left( 3C^{02} + \frac{2p}{p+1} 2^{(p-1)/2} C^{0p+1} \right) \right\}$ , so that

$$\left| \frac{d}{dt} \int u_n^2(t, x) g\left(\frac{x - A(\varepsilon)}{C(\varepsilon)}\right) \right| \leq \frac{\varepsilon}{T(\varepsilon) - T_0}.$$

By integration on  $[T_0, T(\varepsilon)]$ :

$$\int_{x \geq 2C(\varepsilon) + A(\varepsilon)} u_n^2(T_0, x) \leq \int u_n^2(T_0, x) g\left(\frac{x - A(\varepsilon)}{C(\varepsilon)}\right) \leq 5\varepsilon.$$

Now considering  $\frac{d}{dt} \int u_n^2(t, x) g\left(\frac{-A(\varepsilon) - x}{C(\varepsilon)}\right)$ , we get in a similar way

$$\int_{x \leq -2C(\varepsilon) - A(\varepsilon)} u_n^2(T_0, x) \leq 5\varepsilon.$$

Therefore, setting  $A_\varepsilon = 2C(\varepsilon/10) + A(\varepsilon/10)$ , we obtain:

$$\forall n \in \mathbb{N}, \quad \int_{|x| \geq A_\varepsilon} u_n^2(T_0, x) \leq \varepsilon. \quad \blacksquare$$

By (2.7), the sequence  $(u_n(T_0))$  is bounded in  $H^1$ , thus we can extract a subsequence (still denoted by  $(u_n)$ ) which converges weakly to  $\varphi_0 \in H^1(\mathbb{R})$ . The previous compactness result ensures that the convergence is strong in  $L^2(\mathbb{R})$ . Indeed, let  $\varepsilon > 0$  and let  $A$  be such that  $\int_{|x| \geq A} \varphi_0^2(x) dx \leq \varepsilon$  and

$$\forall n \in \mathbb{N}, \quad \int_{|x| \geq A} u_n^2(T_0, x) \leq \varepsilon.$$

By the compact embedding  $H^1([-A, A]) \rightarrow L^2([-A, A])$ ,  $\int_{|x| \leq A} |u_n(T_0, x) - \varphi_0(x)|^2 dx \rightarrow 0$  as  $n \rightarrow +\infty$ . We thus derive that

$$\limsup_{n \in \mathbb{N}} \|u_n(T_0) - \varphi_0\|_{L^2(\mathbb{R})}^2 \leq 4\varepsilon.$$

Since this is true for all  $\varepsilon > 0$ ,  $u_n(T_0) \rightarrow \varphi_0$  in  $L^2(\mathbb{R})$  as  $n \rightarrow +\infty$ . By interpolation,  $u_n(T_0)$  converges strongly to  $\varphi_0$  in  $H^s$  for all  $s \in [0, 1)$ .

*Step 2. Construction of the multi-soliton  $u^*$ .* Denote  $u^*(t)$  the solution to

$$\begin{cases} u_t^* + (u_{xx}^* + (u^*)^p)_x = 0, \\ u^*(T_0) = \varphi_0. \end{cases}$$

Due to [14], the Cauchy problem for (gKdV) is locally well-posed in  $H^s$  for  $s \geq 1/2$ : we will work in  $H^{1/2}$  (which is not a critical space) and  $H^1$ . Let  $u^* \in \mathcal{C}([T_0, T^*), H^1)$  be the maximal solution to (gKdV). Recall the blow up alternative: either  $T^* = +\infty$  or  $T^* < \infty$  and then  $\|u^*(t)\|_{H^1} \rightarrow \infty$  as  $t \uparrow T^*$ .

Since the flow is continuous in  $H^{1/2}$ , for any  $t \in [T_0, T^*)$ ,  $u_n(t)$  is defined for  $n$  large enough and  $u_n(t) \rightarrow u^*(t)$  in  $H^{1/2}$  as  $n \rightarrow +\infty$ . By the uniform  $H^1$  bound, we also obtain  $u_n(t) \rightharpoonup u^*(t)$  in  $H^1$ -weak. Hence, using Proposition 1,

$$\forall t \in [T_0, T^*), \quad \|u^*(t) - R(t)\|_{H^1} \leq \liminf_{n \rightarrow \infty} \|u_n(t) - R(t)\|_{H^1} \leq C e^{-\sigma_0^{3/2} t}.$$

In particular, we deduce that

$$\forall t \in [T_0, T^*), \quad \|u^*(t)\|_{H^1} \leq C e^{-\sigma_0^{3/2} t} + \|R(t)\|_{H^1} \leq C + \sum_{j=1}^N \|Q_{c_j}\|_{H^1}.$$

The blow-up alternative implies  $T^* = \infty$ . Hence  $u^* \in \mathcal{C}([T_0, \infty), H^1)$  and moreover  $\|u^*(t) - R(t)\|_{H^1} \leq C e^{-\sigma_0^{3/2} t}$  for all  $t \geq T_0$ . ■

### 2.3. Proof of Proposition 1

The proof proceeds in several steps. For the sake of simplicity, we will drop the index  $n$  for the rest of this section (except for  $S_n$ ). We possibly drop the first terms of the sequence  $S_n$ , so that for all  $n$ ,  $S_n$  is large enough for our purposes.

*Step 1.* Choice of a set of initial data.

**Lemma 2** (Modulation for time independent function). *Let  $0 < c_1 < \dots < c_N$ . There exist  $C, \varepsilon > 0$  such that the following holds. Given  $(\alpha_i)_{i=1, \dots, N}$  such  $\min\{|\alpha_i - \alpha_j| \mid i \neq j\} \geq 1/\varepsilon$ , if  $u(x) \in L^2$  is such that*

$$\left\| u - \sum_{j=1}^N Q_{c_j}(x - \alpha_j) \right\|_{L^2} \leq \varepsilon,$$

*then there exist modulation parameters  $\mathbf{y} = (y_j)_{j=1, \dots, N}$  such that setting*

$$v = u - \sum_{j=1}^N Q_{c_j}(x - \alpha_j - y_j),$$

the following holds

$$(2.8) \quad \|v\|_{L^2} + \sum_{j=1}^N |y_j| \leq C \left\| u - \sum_{j=1}^N Q_{c_j}(x - \alpha_j) \right\|_{L^2},$$

$$(2.9) \quad \text{and } \forall j = 1, \dots, N, \quad \int v(x)(Q_{c_j})_x(x - \alpha_j - y_j)dx = 0.$$

Furthermore,  $u \mapsto (v, \mathbf{y})$  is a smooth diffeomorphism.

*Notation.* For  $\mathbf{b}$  small, from (2.5) and continuity in  $H^1$ ,  $u(t)$  is defined and modulable (in the sense of the previous lemma) for  $t$  close to  $S_n$ . As long as  $u(t)$  is modulable around  $R(t)$ , we denote by  $\mathbf{y}(t) = (y_j(t))_{j=1, \dots, N}$  the modulation parameters, and

$$\tilde{R}_j(t) = R_j(t, x - y_j(t)), \quad \tilde{R}(t) = \sum_{j=1}^N \tilde{R}_j(t), \quad \tilde{Z}_j^\pm(t, x) = Z_j^\pm(t, x - y_j(t)),$$

$$v(t) = u(t) - \tilde{R}(t) \quad \text{so that } \forall j = 1, \dots, N, \quad \int v(t)(\tilde{R}_j)_x(t) = 0,$$

$$\mathbf{a}^\pm(t) = (a_j^\pm(t))_{j=1, \dots, N}, \quad \text{where } a_j^\pm = \int v(t, x) \tilde{Z}_j^\pm(t, x) dx.$$

We also set  $L_j = -\partial_{xx} - p\tilde{R}_j^{p-1}(t) + c_j$ .

We consider  $\mathbb{R}^N$  equipped with the  $\ell^2$  norm. We denote by  $B_{\mathcal{B}}(P, r)$  the closed ball of the Banach space  $\mathcal{B}$ , centered at  $P$  and of radius  $r \geq 0$ . If  $P = 0$ , we simply write  $B_{\mathcal{B}}(r)$ . Finally,  $\mathbb{S}_{\mathbb{R}^N}(r)$  denotes the sphere of radius  $r$  in  $\mathbb{R}^N$ .

In view of Lemma 1, we have to control the functions  $\mathbf{a}^\pm(t)$  on some time interval  $[T_0, S_n]$ . Since  $Z^+$  and  $Z^-$  are not orthogonal and because of the interactions between the various solitons, the values of  $\mathbf{a}^\pm(S_n)$  are not directly related to  $\mathbf{b}$ . The next lemma allows us to establish a one-to-one mapping between the choice of  $\mathbf{b}$  in (2.5) and the suitable constraints  $\mathbf{a}^+(S_n) = \mathbf{a}^+$ ,  $\mathbf{a}^-(S_n) = 0$ , for any choice of  $\mathbf{a}^+$ .

**Lemma 3** (Modulated final data). *There exists  $C > 0$  (independent of  $n$ ) such that for all  $\mathbf{a}^+ \in B_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n})$  there exists a unique  $\mathbf{b}$  with  $\|\mathbf{b}\| \leq C\|\mathbf{a}^+\|$  and such that the modulation  $(v(S_n), \mathbf{y}(S_n))$  of  $u(S_n)$  satisfies*

$$\mathbf{a}^+(S_n) = \mathbf{a}^+ \quad \text{and} \quad \mathbf{a}^-(S_n) = 0.$$

**Proof.** See Appendix. ■

Let  $T_0$  to be determined later in the proof, independent of  $n$ . Let  $\mathbf{a}^+$  to be chosen,  $\mathbf{b}$  be given by Lemma 3 and let  $u$  be the corresponding solution of (2.5). We now define the maximal time interval  $[T(\mathbf{a}^+), S_n]$  on which suitable exponential estimates hold.

**Definition 1.** Let  $T(\mathbf{a}^+)$  be the infimum of  $T \geq T_0$  such that the following properties hold for all  $t \in [T, S_n]$ :

- Closeness to  $R(t)$ :

$$\|u(t) - R(t)\|_{H^1} \leq \varepsilon.$$

In particular, this ensures that  $u(t)$  is modulable around  $R(t)$  in the sense of Lemma 2.

- Estimates on the modulation parameters:

$$\begin{aligned} e^{\sigma_0^{3/2}t}v(t) &\in B_{H^1}(1), & e^{\sigma_0^{3/2}t}\mathbf{y}(t) &\in B_{\mathbb{R}^N}(1), \\ e^{(3/2)\sigma_0^{3/2}t}\mathbf{a}^-(t) &\in B_{\mathbb{R}^N}(1), & e^{(3/2)\sigma_0^{3/2}t}\mathbf{a}^+(t) &\in B_{\mathbb{R}^N}(1). \end{aligned}$$

Observe that Proposition 1 is proved if for all  $n$ , we can find  $\mathbf{a}^+$  such that  $T(\mathbf{a}^+) = T_0$ . The rest of the proof is devoted to prove the existence of such a value of  $\mathbf{a}^+$ .

We claim the following preliminary results on the modulation parameters of  $u(t)$ .

**Claim.**

$$(2.10) \quad v_t + \left( v_{xx} + (v + \tilde{R})^p - \sum_{j=1}^N \tilde{R}_j^p \right)_x - \sum_{j=1}^N \frac{dy_j}{dt} \tilde{R}_{jx} = 0,$$

$$(2.11) \quad \forall t \in [T(\mathbf{a}^+), S_n], \quad \left\| \frac{d\mathbf{y}}{dt}(t) \right\| \leq C\|v(t)\|_{L^2} + Ce^{-2\sigma_0^{3/2}t}.$$

$$(2.12) \quad \forall t \in [T(\mathbf{a}^+), S_n], \quad \forall j, \quad \left| \frac{da_j^\pm}{dt}(t) \pm e_0 c_j^{3/2} a_j^\pm(t) \right| \leq C\|v(t)\|_{H^1}^2 + Ce^{-3\sigma_0^{3/2}t}.$$

**Proof.** The equation of  $v(t)$  is obtained by elementary computations from the equation of  $u(t)$ . Taking the scalar product of this equation with  $\tilde{R}_{jx}$ , we see that  $y_j(t)$  satisfy

$$\frac{dy_j}{dt} \|Q_{c_{jx}}\|_{L^2}^2 = \int \left( v_{xx} + (v + \tilde{R})^p - \sum_{k=1}^N \tilde{R}_k^p \right)_x \tilde{R}_{jx} - \left( c_j + \frac{dy_j}{dt} \right) \int v \tilde{R}_{jxx}.$$

For  $t \geq T_0$  large enough  $\left| \int v \tilde{R}_{jxx} \right| \leq \|Q_{c_{jxx}}\|_{L^2} \|v(t)\|_{L^2} \leq Ce^{-\sigma_0^{3/2}t} \leq \frac{1}{2} \|Q_{c_{jx}}\|_{L^2}^2$ . Using integration by parts to have all the derivatives on  $\tilde{R}_{jx}$  and using Cauchy-Schwarz inequality, we get (2.11).

Now, we prove (2.12). First, note that  $\int \tilde{R}_{jx} \tilde{Z}_j^\pm = 0$  follows from

$$(2.13) \quad \int Q_x Z^\pm = \pm e_0^{-1} \int Q_x L(Z_x^\pm) = \pm e_0^{-1} \int L(Q_x) Z_x^\pm = 0.$$

Using the equation of  $v(t)$  and next the equations of  $Z^\pm$ ,

$$\begin{aligned}
 \frac{da_j^\pm}{dt}(t) &= \int v_t \tilde{Z}_j^\pm + \int v \tilde{Z}_j^\pm_t \\
 &= - \int (v_{xx} + (v + \tilde{R})^p - \sum_k \tilde{R}_k^p)_x \tilde{Z}_j^\pm \\
 &\quad + \sum_k \frac{dy_k}{dt} \int \tilde{R}_{kx} \tilde{Z}_j^\pm - (c_j + \frac{dy_j}{dt}) \int v \tilde{Z}_j^\pm_x \\
 &= - \int (v_{xx} + p\tilde{R}_j^{p-1}v)_x \tilde{Z}_j^\pm - c_j \int v \tilde{Z}_j^\pm_x \\
 &\quad - \int ((v + \tilde{R})^p - \sum_k \tilde{R}_k^p - p\tilde{R}_j^{p-1}v)_x \tilde{Z}_j^\pm \\
 &\quad + \sum_{k \neq j} \frac{dy_k}{dt} \int \tilde{R}_{kx} \tilde{Z}_j^\pm - \frac{dy_j}{dt} \int v \tilde{Z}_j^\pm_x \\
 &= - \int v L_j(\tilde{Z}_j^\pm_x) + \int ((v + \tilde{R})^p - \sum_k \tilde{R}_k^p - p\tilde{R}_j^{p-1}v) \tilde{Z}_j^\pm_x \\
 &\quad + \sum_{k \neq j} \frac{dy_k}{dt} \int \tilde{R}_{kx} \tilde{Z}_j^\pm - \frac{dy_j}{dt} \int v \tilde{Z}_j^\pm_x \\
 &= \mp e_0 c_j^{3/2} a_j^\pm(t) + \int ((v + \tilde{R})^p - \sum_k \tilde{R}_k^p - p\tilde{R}_j^{p-1}v) \tilde{Z}_j^\pm_x \\
 &\quad + \sum_{k \neq j} \frac{dy_k}{dt} \int \tilde{R}_{kx} \tilde{Z}_j^\pm - \frac{dy_j}{dt} \int v \tilde{Z}_j^\pm_x.
 \end{aligned}$$

Using (2.6), for  $k \neq j$ ,

$$\begin{aligned}
 |\tilde{R}_k(t, x)| (|\tilde{Z}_j^\pm(t, x)| + |\tilde{Z}_j^\pm_x(t, x)|) &\leq C e^{-2\sqrt{\sigma_0}(|x-c_k t|+|x-c_j t|)} \\
 (2.14) \qquad \qquad \qquad &\leq C e^{-3\sigma_0^{3/2}t} e^{-\sqrt{\sigma_0}|x-c_j t|}.
 \end{aligned}$$

Hence we have

$$\begin{aligned}
 \left| \int (|v + \tilde{R}|^{p-1}(v + \tilde{R}) - \sum_{k=1}^N \tilde{R}_k^p - p\tilde{R}_j^{p-1}v) \tilde{Z}_j^\pm_x \right| &\leq C \|v(t)\|_{H^1}^2 + C e^{-3\sigma_0^{3/2}t}, \\
 \left| \sum_{k \neq j} \frac{dy_k}{dt} \int \tilde{R}_{kx} \tilde{Z}_j^\pm \right| &\leq C e^{-3\sigma_0^{3/2}t} \|v(t)\|_{H^1} + C e^{-4\sigma_0^{3/2}t} \\
 &\leq C \|v(t)\|_{H^1}^2 + C e^{-4\sigma_0^{3/2}t}.
 \end{aligned}$$

The term  $\frac{dy_j}{dt} \int v \tilde{Z}_j^\pm_x$  is controlled using (2.11). ■

*Step 2.* Conditional stability of  $v$  and  $\mathbf{y}$  under the control of  $\mathbf{a}^\pm$ .

We claim the following improvement of the estimates for  $v(t)$  and  $\mathbf{y}$  on  $[T(\mathbf{a}^+), S_n]$ .

**Lemma 4** (Control of  $v$  and  $\mathbf{y}$ ). *For  $T_0$  large enough (independent of  $n$ ) and for all  $\mathbf{a}^+ \in B_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n})$ , the following holds*

$$(2.15) \quad \forall t \in [T(\mathbf{a}^+), S_n], \quad \|u(t) - R(t)\|_{H^1} \leq Ce^{-\sigma_0^{3/2}t} \leq \varepsilon_0/2,$$

$$(2.16) \quad e^{\sigma_0^{3/2}t}\|v(t)\|_{H^1} \leq 1/2, \quad e^{\sigma_0^{3/2}t}\|\mathbf{y}(t)\| \leq 1/2.$$

The proof of Lemma 4 is postponed to the end of this section. It is very similar to the proofs in the subcritical case (see [16] or [20]).

*Step 3.* Control of  $\mathbf{a}^-(t)$ .

**Lemma 5** (Control of  $\mathbf{a}^-(t)$ ). *For  $T_0$  large enough (independent of  $n$ ) and for all  $\mathbf{a}^+ \in B_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n})$ , the following holds*

$$\forall t \in [T(\mathbf{a}^+), S_n], \quad e^{(3/2)\sigma_0^{3/2}t}\|\mathbf{a}^-(t)\| \leq 1/2.$$

**Proof.** It follows from (2.12), (2.16) and  $a_j^-(S_n) = 0$  that for all  $t \in [T(\mathbf{a}^-), S_n]$ ,

$$|a_j^-(t)| \leq Ce^{e_0c_j^{3/2}t} \int_t^{S_n} e^{-e_0c_j^{3/2}s} \left( e^{-2\sigma_0^{3/2}s} + e^{-3\sigma_0^{3/2}s} \right) ds \leq Ce^{-2\sigma_0^{3/2}t}.$$

Hence, for  $T_0$  large enough,

$$\forall t \in [T(\mathbf{a}^+), S_n], \quad \|\mathbf{a}^-(t)\| \leq Ce^{-2\sigma_0^{3/2}t} \leq \frac{1}{2} e^{-(3/2)\sigma_0^{3/2}t}. \quad \blacksquare$$

*Step 4.* Control of  $\mathbf{a}^+(t)$  by a topological argument.

Finally we turn to the control of  $\mathbf{a}^+(t)$  which will provide us with a suitable value of  $\mathbf{a}^+$ . This is the new key argument of this paper.

**Lemma 6** (Control of  $\mathbf{a}^+(t)$ ). *For  $T_0$  large enough, there exists*

$$\mathbf{a}^+ \in B_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n}) \quad \text{such that} \quad T(\mathbf{a}^+) = T_0.$$

**Proof.** We argue by contradiction.

Assume that for all  $\mathbf{a}^+ \in B_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n})$ , one has  $T(\mathbf{a}^+) > T_0$ . From Lemmas 4 and 5

$$\begin{aligned} u(T(\mathbf{a}^+)) - R(T(\mathbf{a}^+)) &\in B_{H^1}(\varepsilon_0/2), & e^{\sigma_0^{3/2}T(\mathbf{a}^+)}v(T(\mathbf{a}^+)) &\in B_{H^1}(1/2), \\ e^{\sigma_0^{3/2}T(\mathbf{a}^+)}\mathbf{y}(T(\mathbf{a}^+)) &\in B_{\mathbb{R}^N}(1/2), & e^{(3/2)\sigma_0^{3/2}T(\mathbf{a}^+)}\mathbf{a}^-(T(\mathbf{a}^+)) &\in B_{\mathbb{R}^N}(1/2). \end{aligned}$$

Hence by definition of  $T(\mathbf{a}^+)$  and continuity of the flow, one must have

$$(2.17) \quad e^{(3/2)\sigma_0^{3/2}T(\mathbf{a}^+)}\mathbf{a}^+(T(\mathbf{a}^+)) \in \mathbb{S}_{\mathbb{R}^N}(1).$$

Let  $T < T(\mathbf{a}^+)$  be close enough to  $T(\mathbf{a}^+)$  so that the solution  $u(t)$  and its modulation are well-defined on  $[T, S_n]$ . For  $t \in [T, S_n]$ , let

$$(2.18) \quad \mathcal{N}(\mathbf{a}^+(t)) = \mathcal{N}(t) = \left\| e^{(3/2)\sigma_0^{3/2}t}\mathbf{a}^+(t) \right\|^2.$$



Then, by (2.12) and (2.16), we have

$$\begin{aligned} \frac{d\mathcal{N}}{dt}(t) &= e^{3\sigma_0^{3/2}t} \sum_j \left( 3\sigma_0^{3/2} - 2\frac{da_j^+}{dt} \right) a_j^+ \\ &= e^{3\sigma_0^{3/2}t} \sum_j \left( 3\sigma_0^{3/2} - 2e_0c_j^{3/2} \right) |a_j^+|^2 \\ &\quad + O\left( e^{3\sigma_0^{3/2}t} \|\mathbf{a}^+\| (\|v(t)\|_{H^1}^2 + e^{-3\sigma_0^{3/2}t}) \right). \end{aligned}$$

In view of the definition of  $\sigma_0$  (see (2.6)), for all  $j$ ,

$$2e_0c_j^{3/2} - 3\sigma_0^{3/2} \geq e_0c_1^{3/2} \geq 4e_0\sigma_0^{3/2}.$$

For  $t \in [T(\mathbf{a}^+), S_n]$ , due to the bound on  $\|v\|_{H^1}$ , we have

$$e^{3\sigma_0^{3/2}t} \|\mathbf{a}^+\| (\|v(t)\|_{H^1}^2 + e^{-3\sigma_0^{3/2}t}) \leq C e^{-(1/2)\sigma_0^{3/2}t} \sqrt{\mathcal{N}(t)}.$$

Hence we get

$$\frac{d\mathcal{N}}{dt}(t) \leq -4e_0\sigma_0^{3/2}\mathcal{N}(t) + C e^{-(1/2)\sigma_0^{3/2}t} \sqrt{\mathcal{N}(t)}.$$

We consider this estimate at  $t = T(\mathbf{a}^+) \geq T_0$  (so large that  $C e^{-(1/2)\sigma_0^{3/2}t} \leq 2e_0\sigma_0^{3/2}$ ), and using  $\mathcal{N}(T(\mathbf{a}^+)) = 1$ , we get

$$(2.19) \quad \forall \mathbf{a}^+ \in B_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n}), \quad \frac{d\mathcal{N}}{dt}(T(\mathbf{a}^+)) \leq -2e_0\sigma_0^{3/2}.$$

From (2.19), a standard argument says that the map  $\mathbf{a}^+ \mapsto T(\mathbf{a}^+)$  is continuous. Indeed, by (2.19), for all  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $\mathcal{N}(T(\mathbf{a}^+) - \varepsilon) > 1 + \delta$  and, for all  $t \in [T(\mathbf{a}^+) + \varepsilon, S_n]$  (possibly empty),  $\mathcal{N}(t) < 1 - \delta$ . By continuity of the flow of the (gKdV) equation, it follows that there exist  $\eta > 0$  such that for all  $\|\tilde{\mathbf{a}}^+ - \mathbf{a}^+\| \leq \eta$ , the corresponding  $\tilde{\mathbf{a}}^+(t)$  satisfies  $|\mathcal{N}(\tilde{\mathbf{a}}^+(t)) - \mathcal{N}(\mathbf{a}^+(t))| \leq \delta/2$  for all  $t \in [T(\mathbf{a}^+) - \varepsilon, S_n]$ . In particular,  $T(\mathbf{a}^+) - \varepsilon \leq T(\tilde{\mathbf{a}}^+) \leq T(\mathbf{a}^+) + \varepsilon$ .

Now, we consider the continuous map

$$\begin{aligned} \mathcal{M} : \quad B_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n}) &\rightarrow \mathbb{S}_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n}), \\ \mathbf{a}^+ &\mapsto e^{-(3/2)\sigma_0^{3/2}(S_n - T(\mathbf{a}^+))} \mathbf{a}^+(T(\mathbf{a}^+)). \end{aligned}$$

Let  $\mathbf{a}^+ \in \mathbb{S}_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n})$ . From (2.19), it follows that  $T(\mathbf{a}^+) = S_n$  and so  $\mathcal{M}(\mathbf{a}^+) = \mathbf{a}^+$ , which means that  $\mathcal{M}$  restricted to  $\mathbb{S}_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n})$  is the identity. But the existence of such a map  $\mathcal{M}$  contradicts Brouwer's fixed point theorem. In conclusion, there exists  $\mathbf{a}^+ \in B_{\mathbb{R}^N}(e^{-(3/2)\sigma_0^{3/2}S_n})$  such that  $T(\mathbf{a}^+) = T_0$ . ■

The end of this section is devoted to the proof of Lemma 4.

**Proof of Lemma 4.** Let  $\psi$  be defined by

$$\begin{aligned} \psi(x) &= 0 \text{ for } x \leq -1, \quad \psi(x) = 1 \text{ for } x \geq 1, \\ \psi(x) &= \frac{1}{c_0} \int_{-1}^x e^{-\frac{1}{1-y^2}} dy \quad \text{for } x \in (-1, 1), \end{aligned}$$

where  $c_0 = \int_{-1}^1 e^{-\frac{1}{1-y^2}} dy$ . Then  $\psi \in C^\infty(\mathbb{R})$  is non-decreasing and  $0 \leq \psi \leq 1$ . Define  $m_j(t) = \frac{1}{2}((c_j + c_{j+1})t + x_j + x_{j+1})$  for  $j = 1, \dots, N - 1$  and

$$\begin{aligned} \text{for } j = 1, \dots, N - 1, \quad \psi_j(t, x) &= \psi\left(\frac{1}{\sqrt{t}}(x - m_j(t))\right), \quad \psi_N(t) = 1; \\ \text{for } j = 2, \dots, N \quad \phi_j &= \psi_j - \psi_{j-1}, \quad \phi_1 = \psi_1; \\ (2.20) \quad M_j(t) &= \int u^2(t)\phi_j(t), \quad E_j(t) = \int \left(\frac{1}{2}u_x^2 - \frac{1}{p+1}u^{p+1}\right)(t)\phi_j(t). \end{aligned}$$

By the decay properties of  $\tilde{R}_k(t)$ , and the support properties of  $\phi_j$  and its derivatives, we have

$$(2.21) \quad \forall j \neq k, \quad (|\tilde{R}_k(t, x)| + |(\tilde{R}_k)_x(t, x)|)|\phi_j(t, x)| \leq C e^{-3\sigma_0^{3/2}t} e^{-\sqrt{\sigma_0}|x-y_j(t)|},$$

$$(2.22) \quad \forall j, \quad (|\tilde{R}_j(t, x)| + |(\tilde{R}_j)_x(t, x)|)|\phi_j(t, x) - 1| \leq C e^{-3\sigma_0^{3/2}t} e^{-\sqrt{\sigma_0}|x-y_j(t)|},$$

$$(2.23) \quad \forall j, k, \quad |\tilde{R}_k(t, x)|(|\phi_{j_x}(t, x)| + |\phi_{j_{xxx}}(t, x)| + |\phi_{j_t}(t, x)|) \leq \frac{C}{\sqrt{t}} e^{-3\sigma_0^{3/2}t} e^{-\sqrt{\sigma_0}|x-y_k(t)|},$$

$$(2.24) \quad \forall j, \quad |\phi_{j_x}(t, x)| + |\phi_{j_{xxx}}(t, x)| + |\phi_{j_t}(t, x)| \leq \frac{C}{\sqrt{t}}.$$

We begin with some technical claims.

**Claim.**

$$(2.25) \quad \left| \frac{d}{dt} M_j(t) \right| \leq \frac{C}{\sqrt{t}} \|v(t)\|_{H^1}^2 + C e^{-3\sigma_0^{3/2}t},$$

$$(2.26) \quad \left| \frac{d}{dt} \sum_{j=1}^N \left( E_j(t) + \frac{c_j}{2} M_j(t) \right) \right| \leq \frac{C}{\sqrt{t}} \|v(t)\|_{H^1}^2 + C e^{-3\sigma_0^{3/2}t}.$$

**Proof.** By direct computations,

$$\frac{d}{dt} \int u^2 \phi_j = -3 \int u_x^2 \phi_{j_x} + \int u^2 (\phi_{j_{xxx}} + \phi_{j_t}) + \frac{2p}{p+1} \int u^{p+1} \phi_{j_x}.$$

Thus, expanding  $u(t) = \tilde{R}(t) + v(t)$ , and using (2.23) and (2.24), the first two integrals are estimated as desired. For the last term it suffices to observe additionally that  $\|u(t)\|_{L^\infty} \leq C(\|v(t)\|_{H^1} + \|\tilde{R}(t)\|_{H^1}) \leq C$ . This proves (2.25).

Estimate (2.26) is a consequence of (2.25), the conservation of energy and  $\sum_{j=1}^N \phi_j = 1$ .  $\blacksquare$

**Claim.**

$$(2.27) \quad \left| \left( E_j(t) + \frac{c_j}{2} M_j(t) \right) - \left( E(Q_{c_j}) + \frac{c_j}{2} \int Q_{c_j}^2 \right) - \frac{1}{2} H_j(t) \right| \leq C e^{-3\sigma_0^{3/2}t} + C e^{-\sigma_0^{3/2}t} \|v(t)\|_{L^2}^2,$$

where  $H_j(t) = \int (v_x^2(t) - p\tilde{R}_j^{p-1}(t)v^2(t) + c_j v^2(t)) \phi_j(t)$ .

**Proof.** First, we claim

$$(2.28) \quad \left| M_j(t) - \left( \int Q_{c_j}^2 + 2 \int v(t) \tilde{R}_j(t) + \int v^2(t) \phi_j(t) \right) \right| \leq C e^{-3\sigma_0^{3/2}t},$$

$$\left| E_j(t) - E(Q_{c_j}) - \left( \frac{1}{2} \int (v_x^2(t) - p\tilde{R}_j^{p-1}(t)v^2(t)) \phi_j(t) - c_j \int v(t) \tilde{R}_j(t) \right) \right|$$

$$(2.29) \quad \leq C e^{-3\sigma_0^{3/2}t} + C e^{-\sigma_0^{3/2}t} \|v(t)\|_{L^2}^2,$$

Indeed, expanding  $u(t) = v(t) + \sum_k \tilde{R}_k(t)$  in  $M_j(t)$  and  $E_j(t)$ , we get

$$M_j(t) = \int u^2 \phi_j(t) = \int \left( v^2 + 2v\tilde{R} + \left( \sum_{k=1}^N \tilde{R}_k \right)^2 \right) \phi_j(t),$$

$$E_j(t) = \int \left( \frac{1}{2} (v_x^2 + 2v_x \tilde{R}_x + \tilde{R}_x^2) - \frac{1}{p+1} (v + \tilde{R})^{p+1} \right) \phi_j(t)$$

$$= \int \left( \frac{1}{2} (v_x^2 - p\tilde{R}^{p-1}v^2) \right) \phi_j + \int \left( \frac{1}{2} \tilde{R}_x^2 - \frac{1}{p+1} \tilde{R}^{p+1} \right) \phi_j(t)$$

$$- \int v(\tilde{R}_{xx} + \tilde{R}^p) \phi_j - \int \tilde{R}_x v \phi_{j_x}$$

$$+ \frac{1}{p+1} \int \left( -(v + \tilde{R})^{p+1} + \tilde{R}^{p+1} \right) + (p+1)v\tilde{R}^p + \frac{p(p+1)}{2} \tilde{R}^{p-1}v^2 \phi_j(t).$$

Now, estimates (2.21) and (2.22) give (for  $k \neq j$ )

$$\left| \int \tilde{R}_j^2 \phi_j(t) - \int Q_{c_j}^2 \right| + \left| \int \tilde{R}_k^2 \phi_j(t) + \int \left( \frac{1}{2} \tilde{R}_x^2 - \frac{1}{p+1} \tilde{R}^{p+1} \right) \phi_j(t) - E(Q_{c_j}) \right| \leq C e^{-3\sigma_0^{3/2}t}.$$

By  $Q_{xx} + Q^p = Q$ , we have

$$\int v(t)(\tilde{R}_{xx} + \tilde{R}^p)\phi_j = c_j \int v(t)\tilde{R}_j(t) + O(e^{-3\sigma_0^{3/2}t}).$$

Using also (2.23) and for  $k \geq 3$

$$\int |v(t)|^k \phi_j(t) \leq \|v(t)\|_{L^\infty}^{k-2} \int v(t)^2 \phi_j(t) \leq Ce^{-\sigma_0^{3/2}t} \|v\|_{L^2}^2,$$

we obtain (2.28) and (2.29).

Estimate (2.27) is obtained by summing (2.28) and (2.29). Note that in particular that the scalar products  $\int v(t)\tilde{R}_j(t)$  cancel. ■

**Claim.** For  $T_0$  large enough, there exists  $K > 0$  such that for all  $t \in [T_0, S_n]$ ,

$$(2.30) \quad \|v(t)\|_{H^1}^2 \leq K \sum_j H_j(t) + K^2 \sum_j \left( \left( \int v(t)\tilde{Z}_j^+(t) \right)^2 + \left( \int v(t)\tilde{Z}_j^-(t) \right)^2 \right).$$

**Proof.** Estimate (2.30) is a standard consequence of Lemma 1 and  $\int v\tilde{R}_{j_x} = 0$ . See e.g. [21, Lemma 4]. ■

Now, we finish the proof of Lemma 4. Let  $t \in [T(\mathbf{a}^+), S_n]$ . Integrating (2.26) on  $[t, S_n]$ ,

$$\begin{aligned} \left| \sum_{j=1}^N \left\{ \left( E_j(S_n) + \frac{c_j}{2} M_j(S_n) \right) - \left( E_j(t) + \frac{c_j}{2} M_j(t) \right) \right\} \right| &\leq \\ &\leq Ce^{-3\sigma_0^{3/2}t} + C \int_t^{S_n} \|v(s)\|_{H^1}^2 \frac{ds}{\sqrt{s}}. \end{aligned}$$

From (2.27), we get:

$$\begin{aligned} \left| \sum_{j=1}^N (H_j(S_n) - H_j(t)) \right| &\leq \\ &\leq Ce^{-3\sigma_0^{3/2}t} + Ce^{-\sigma_0^{3/2}t} (\|v(t)\|_{L^2}^2 + \|v(S_n)\|_{L^2}^2) + C \int_t^{S_n} \|v(s)\|_{H^1}^2 \frac{ds}{\sqrt{s}}. \end{aligned}$$

Note that from Lemmas 2 and 3, and from the definition of  $T(\mathbf{a}^+)$ ,

$$|H_j(S_n)| \leq C \|v(S_n)\|_{H^1}^2 \leq C \|\mathbf{b}\|^2 \leq Ce^{-3\sigma_0^{3/2}t} \quad \text{and} \quad \|v(t)\|_{H^1}^2 \leq e^{-2\sigma_0^{3/2}t}.$$

By (2.30) and the above estimates

$$\begin{aligned} \|v(t)\|_{H^1}^2 &\leq K \sum_{j=1}^N H_j(t) + K^2 \sum_{j,\pm} a_j^\pm(t)^2 \\ &\leq Ce^{-3\sigma_0^{3/2}t} + C \sum_{j,\pm} a_j^\pm(t)^2 + C \int_t^{S_n} \|v(s)\|_{H^1}^2 \frac{ds}{\sqrt{s}} \\ (2.31) \quad &\leq C_0 e^{-3\sigma_0^{3/2}t} + \frac{C_0}{\sqrt{t}} e^{-2\sigma_0^{3/2}t}. \end{aligned}$$

Hence, for  $T_0$  large enough so that  $C_0 e^{-\sigma_0^{3/2} T_0} \leq 1/8$  and  $C_0/\sqrt{T_0} \leq 1/8$  we get

$$e^{\sigma_0^{3/2} t} \|v(t)\|_{H^1} \leq 1/2.$$

By (2.11) and (2.31),

$$\begin{aligned} \|\mathbf{y}_t(t)\| &\leq C e^{-2\sigma_0^{3/2} t} + C \|v(t)\|_{L^2}, \\ \|\mathbf{y}(t)\| &\leq \|\mathbf{y}(S_n)\| + C \int_t^{S_n} \left( e^{-(3/2)\sigma_0^{3/2} s} + \frac{e^{-\sigma_0^{3/2} s}}{s^{1/4}} \right) ds \\ (2.32) \quad &\leq C e^{-(3/2)\sigma_0^{3/2} t} + \frac{C}{t^{1/4}} e^{-\sigma_0^{3/2} t}, \end{aligned}$$

and we deduce  $e^{\sigma_0^{3/2} t} \|\mathbf{y}(t)\| \leq 1/2$  by possibly taking a larger  $T_0$ . Finally, we have:

$$\begin{aligned} \|u(t) - R(t)\|_{H^1} &\leq \|R(t) - \tilde{R}(t)\|_{H^1} + \|v(t)\|_{H^1} \leq C \|\mathbf{y}(t)\| + \|v(t)\|_{H^1} \\ (2.33) \quad &\leq C e^{-\sigma_0^{3/2} t} \leq \varepsilon_0/2, \end{aligned}$$

by possibly taking a larger  $T_0$ . This concludes the proof of Lemma 4. ■

### 3. Generalizations

#### 3.1. The gKdV equations with general nonlinearities

We now present extensions of Theorem 1 to a more general form of the KdV equation, i.e.

$$(gKdV) \quad u_t + (u_{xx} + f(u))_x = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R}.$$

In order to have both local well-posedness in  $H^1$  from [14] and the existence of eigenvalues for the linearized operator in the unstable case from [26], we assume

$$(3.1) \quad f \text{ is } C^3, \text{ convex on } \mathbb{R}_+, \text{ and } f(0) = f'(0) = 0,$$

but these assumptions can probably be relaxed. Concerning the solitons, we consider velocities  $c_j > 0$  such that

$$\begin{aligned} &\text{a solution } Q_c \text{ of } (Q_c)_{xx} + f(Q_c) = cQ_c \text{ exists for all } c \text{ close to } c_j \\ (3.2) \quad &\text{and } \frac{d}{dc} \int Q_c^2 \Big|_{c=c_j} \neq 0. \end{aligned}$$

Then, combining the proof of Theorem 1 and [16], we claim the following extension of Theorem 1.

**Theorem 3.** *Let  $0 < c_1 < \dots < c_N$  and  $x_1, \dots, x_N \in \mathbb{R}$  be such that for all  $j$ , (3.2) holds. There exist  $T_0 \in \mathbb{R}$ ,  $C, \sigma_0 > 0$ , and a solution  $u \in \mathcal{C}([T_0, \infty), H^1)$  to (gKdV) such that*

$$\forall t \in [T_0, \infty), \quad \left\| u(t) - \sum_{j=1}^N R_{c_j, x_j}(t) \right\|_{H^1} \leq C e^{-\sigma_0^{3/2} t}.$$

**Remark 3.** The critical case  $\frac{d}{dc} \int Q_{c|c=c_j}^2 = 0$  is treated in [16] for the pure power case. We leave open the case where for a general  $f(u)$ ,  $\frac{d}{dc} \int Q_{c|c=c_j}^2 = 0$  for some  $c_j$ , but it probably can be treated by similar techniques.

From the techniques developed in [25], [7] and [8] concerning the (BBM) equation

$$(BBM) \quad (u - u_{xx})_t + (u + u^p)_x = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R},$$

and from the construction of suitable eigenfunctions of the linearized equation by Pego and Weinstein [26] (see page 74), one can also extend the results obtained in this paper to the (BBM) equation for any  $p > 1$ .

### 3.2. The non linear Schrödinger equations

In this section, we sketch the proof of Theorem 2. It is an extension of the proof of Theorem 1 in the present paper and of the main result in [20].

#### 3.2.1. Preliminaries

Let  $v = v_1 + iv_2$ , we define the operator  $\mathcal{L}$  by

$$(3.3) \quad \mathcal{L}v = -L_- v_2 + iL_+ v_1,$$

where the self-adjoint operators  $L_+$  and  $L_-$  are defined by

$$(3.4) \quad L_+ v_1 := -\Delta v_1 + v_1 - pQ^{p-1}v_1, \quad L_- v_2 := -\Delta v_2 + v_2 - Q^{p-1}v_2,$$

From [29], [11] and [28], there exist  $e_0 > 0$ ,  $Y^\pm \in \mathcal{S}(\mathbb{R})$  ( $\bar{Y}^+ = Y^-$ ), normalized so that  $\|Y^\pm\|_{L^2} = 1$  and such that

$$(3.5) \quad \mathcal{L}Y^\pm = \pm e_0 Y^\pm;$$

moreover, for some  $K > 0$ , for any  $v = v_1 + iv_2 \in H^1$  ( $(f, g) = \text{Re} \int f \bar{g}$ )

$$(3.6) \quad \|v\|_{H^1}^2 \leq K(L_+ v_1, v_1) + K(L_- v_2, v_2) + K^2 \left( \int (\nabla Q)v_1 \right)^2 + K^2 \left( \int Qv_2 \right)^2 + K^2 \left( \text{Im} \int Y^+ \bar{v} \right)^2 + K^2 \left( \text{Im} \int Y^- \bar{v} \right)^2.$$

See [5, 6] for the proof of (3.6).

**3.2.2. Proof of Theorem 2 assuming uniform estimates**

We denote

$$(3.7) \quad \begin{aligned} R(t, x) &= \sum_{j=1}^N R_j(t, x) \quad \text{where} \quad R_j(t, x) = R_{c_j, \gamma_j, v_j, x_j}(t, x), \\ Y_j^\pm(t, x) &= c_j^{\frac{1}{p-1}} Y^\pm(\sqrt{c_j}(x - v_j t - x_j)) e^{i(\frac{1}{2}v_j \cdot x - \frac{1}{4}\|v_j\|^2 t + c_j t + \gamma_j)}. \end{aligned}$$

Let  $S_n \rightarrow \infty$  be an increasing sequence of time. We claim the existence of final data giving suitable uniform estimates.

**Proposition 2.** *There exist  $n_0 \geq 0, \sigma_0 > 0, T_0 > 0, C > 0$  (independent of  $n$ ) such that the following holds. For each  $n \geq n_0$ , there exists  $\mathbf{b} = (\mathbf{b}_{j,n}^\pm)_{j,\pm} \in \mathbb{R}^{2N}$  with  $\|\mathbf{b}\| \leq e^{-\sigma_0^{3/2} S_n}$ , and such that the solution  $u_n$  to*

$$(3.8) \quad \begin{cases} iu_{nt} + \Delta u_n + |u_n|^{p-1} u_n = 0, \\ u_n(S_n) = R(S_n) + i \sum_{j \in \{1, \dots, N\}, \pm} \mathbf{b}_{j,n}^\pm Y_j^\pm(S_n) \end{cases}$$

is defined on the interval  $[T_0, S_n]$ , and satisfies

$$\forall t \in [T_0, S_n], \quad \|u_n(t) - R(t)\|_{H^1} \leq C e^{-\sigma_0^{3/2} t}.$$

The proof of Theorem 2 assuming Proposition 2 is completely similar to Section 2.2 in the present paper and to Section 2 in [20], thus it is omitted (note that for this part, as in [20], we use the local  $H^s$  Cauchy theory due to Cazenave and Weissler [4]).

**3.2.3. Proof of the uniform estimates**

We are reduced to prove Proposition 2. We only sketch the proof since it is very similar to Section 2.3 of the present paper combined with Section 3 in [20].

The first step of the proof is to reduce (without loss of generality) to the special case where

$$v_{1,1} < v_{2,1} < \dots < v_{N,1},$$

where  $v_{j,k}$  ( $j \in \{1, \dots, N\}, k \in \{1, \dots, d\}$ ) represents the  $k$ -th component of the velocity vector  $v_j \in \mathbb{R}^d$ . It is a simple observation, based on the invariance by rotation of the (NLS) equation, see [20, Claim 1, page 855].

Next, in the (NLS) case, modulation theory for  $u(t)$  close to  $R(t)$  says that there exist parameters  $\mathbf{y}(t) = (y_1(t), \dots, y_N(t)) \in (\mathbb{R}^d)^N$  and  $\mu(t) =$

$(\mu_1(t), \dots, \mu_N(t)) \in \mathbb{R}^N$  such that

$$\begin{aligned} \tilde{R}_j(t) &= R_j(t, x - y_j(t))e^{i\mu_j(t)}, \quad \tilde{R}(t) = \sum_{j=1}^N \tilde{R}_j(t), \\ \tilde{Y}_j^\pm(t, x) &= Y_j^\pm(t, x - y_j(t))e^{i\mu_j(t)}, \end{aligned}$$

and  $v(t) = u(t) - \tilde{R}(t)$  satisfies

$$\forall j = 1, \dots, N, \quad \operatorname{Re} \int v(t)(\nabla \tilde{R}_j)(t) = \operatorname{Im} \int v(t)\tilde{R}_j(t) = 0.$$

Note that the parameter  $\mu_j(t)$  is used to control the direction  $\operatorname{Im} \int v(t)\tilde{R}_j(t)$ .

In view of (3.6), we are led to set

$$\mathbf{a}^\pm(t) = (a_j^\pm(t))_{j=1, \dots, N}, \quad \text{where } a_j^\pm(t) = \operatorname{Im} \int \tilde{Y}_j^\mp(t, x)\bar{v}(t, x)dx.$$

For given  $\mathbf{a}^+ \in \mathbb{R}^N$ , we define  $\mathbf{b} \in \mathbb{R}^{2N}$  as for the (gKdV) case in Lemma 3.

We define  $T(\mathbf{a}^+)$  as in Definition 1, with the additional requirement  $e^{\sigma_0^{3/2}t}\mu(t) \in B_{\mathbb{R}^N}(1)$ . By standard computations, the following holds on  $[T(\mathbf{a}^+), S_n]$ .

**Claim.** For some  $\sigma_0 > 0$ ,

$$(3.9) \quad \left\| \frac{d\mathbf{y}}{dt}(t) \right\| + \left\| \frac{d\mu}{dt}(t) \right\| \leq C\|v(t)\|_{L^2} + Ce^{-2\sigma_0^{3/2}t},$$

$$(3.10) \quad \left| \frac{da_j^\pm}{dt}(t) \pm e_0 c_j^{3/2} a_j^\pm(t) \right| \leq C\|v(t)\|_{L^2}^2 + Ce^{-3\sigma_0^{3/2}t}.$$

**Proof.** The proof follows from the equation of  $v$

$$\begin{aligned} iv_t + \Delta v + \sum_j \left( |\tilde{R}_j|^{p-1}v + (p-1)|\tilde{R}_j|^{p-2}\operatorname{Re}(\tilde{R}_j v) \right) \\ + O(\|v\|_{H^1}^2 + e^{-4\sigma_0^{3/2}t}) - \sum_j \frac{dy_j}{dt}\tilde{R}_{jx} - \sum_j \frac{d\mu_j}{dt}\tilde{R}_j = 0, \end{aligned}$$

and direct computations using the definition of  $Y^\pm$ . ■

Now we follow exactly the same strategy as in the proof of Theorem 1, by proving analogues of Lemmas 4, 5 and 6.

For the proof of the estimate on  $v(t)$ , we use a functional adapted to the (NLS) equations, as in [20] and [22]:

$$\begin{aligned} \mathcal{G}(t) = \sum_j \left[ \int \left( \frac{1}{2}|\nabla u|^2 - \frac{1}{p+1}|u|^{p+1} \right) \phi_j \right. \\ \left. + \left( c_j + \frac{|v_j|^2}{4} \right) \int |u|^2 \phi_j - v_j \cdot \operatorname{Im} \int \bar{u} \nabla u \phi_j \right], \end{aligned}$$



where

$$\psi_j(t, x) = \psi \left( \frac{1}{\sqrt{t}}(x_1 - m_j(t)) \right), \quad m_j(t) = \frac{1}{2}((v_{j,1} + v_{j+1,1})t + x_{j,1} + x_{j+1,1});$$

$$\phi_1 = \psi_1, \quad \phi_j = \psi_j - \psi_{j-1}.$$

Note that  $\mathcal{G}(t)$  controls the size of  $v(t)$  in  $H^1$  up to  $\mathbf{a}^\pm(t)$  as a consequence of (3.6). As for (gKdV), the following claim allows us to prove the estimate on  $\|v(t)\|_{H^1}$ .

**Claim.**

$$\left| \frac{d\mathcal{G}}{dt}(t) \right| \leq \frac{C}{\sqrt{t}} \|v(t)\|_{H^1}^2 + C e^{-3\sigma_0^{3/2}t}.$$

The estimates of  $\mathbf{a}^\pm(t)$  are exactly the same as in Lemmas 5 and 6, using (3.10).

### A. Appendix

**Proof of Lemma 2.** We use the following notation  $\mathbf{y} = (y_j)_{j=1,\dots,N}$  and

$$R_j(x) = Q_{c_j}(x - \alpha_j), \quad R(x) = \sum_{j=1}^N R_j(x),$$

$$\tilde{R}_j(x) = R_j(x - y_j), \quad \tilde{R}(x) = \sum_{j=1}^N \tilde{R}_j(x).$$

Let  $w = u - R$  small in  $L^2$ . Consider

$$\Phi : L^2 \times \mathbb{R}^N \rightarrow \mathbb{R}^N,$$

$$(w, \mathbf{y}) \mapsto \left( \int (w + R - \tilde{R}) \tilde{R}_{j_x} \right)_{j=1,\dots,N}.$$

Let  $\mathbf{z} = (z_j)_{j=1,\dots,N}$ . By the decay properties of  $\tilde{R}_j$ ,

$$(d_{\mathbf{y}}\Phi(w, \mathbf{y}) \cdot \mathbf{z})_j = \sum_{k=1}^N z_k \int \tilde{R}_{k_x} \tilde{R}_{j_x} - z_j \int (w + R - \tilde{R}) \tilde{R}_{j_{xx}}$$

$$= z_j \|Q_{c_{j_x}}\|_{L^2}^2 + O\left( \sum_{k \neq j} e^{-\sigma_0|\alpha_k - \alpha_j|} |z_k| \right) + O(|z_j| \|w\|_{L^2}) + O(|z_j| \|\mathbf{y}\|).$$

Hence

$$(A.1) \quad d_{\mathbf{y}}\Phi(w, \mathbf{y}) = \text{diag}(\|Q_{c_{j_x}}\|_{L^2}^2) + O\left( \sum_{k \neq j} e^{-\sigma_0|\alpha_k - \alpha_j|} \right) + O(\|w\|_{L^2}) + O(\|\mathbf{y}\|).$$

Therefore, if  $\min\{|\alpha_k - \alpha_j|, i \neq j\}$  is large enough then  $d_{\mathbf{y}}\Phi(0, 0)$  is invertible. Since  $\Phi(0, 0) = 0$ , by the implicit function theorem, it follows that there exists  $\varepsilon > 0$ ,  $\varepsilon \leq \eta$  and a  $C^1$  function  $\phi : B_{L^2}(0, \varepsilon) \rightarrow B_{\mathbb{R}^N}(0, \eta)$  such that  $\Phi(w, \mathbf{y}) = 0$  in  $B_{L^2}(0, \varepsilon) \times \phi(B_{L^2}(0, \varepsilon))$  is equivalent to  $\mathbf{y} = \phi(w)$ . Finally we set  $v = v(w) = w + R - \sum_{j=1}^N R_j(\cdot - \phi(w)_j)$ . ■

**Proof of Lemma 3.** Consider the maps:

$$\begin{aligned} \mathcal{I} : \mathbb{R}^{2N} &\rightarrow H^1 & \Theta : \mathcal{V} &\rightarrow H^1 \times \mathbb{R}^N & \mathcal{S} : H^1 \times \mathbb{R}^N &\rightarrow \mathbb{R}^{2N} \\ \mathbf{b} &\mapsto \sum_{j,\pm} \mathbf{b}_j^\pm Z_j^\pm(S_n) & w &\mapsto (v, \mathbf{y}) & (v, \mathbf{y}) &\mapsto \left( \int v \tilde{Z}_j^\pm \right)_{j,\pm} \end{aligned}$$

where, in the definition of  $\Theta$ ,  $(v, \mathbf{y})$  represents the modulation of  $u = w + R(S_n)$  and  $\mathcal{V} = B_{H^1}(\varepsilon)$  ( $\varepsilon$  being defined in the proof Lemma 2), and in the definition of  $\mathcal{S}$ , we have set  $\tilde{Z}_j^\pm(x) = Z_j^\pm(S_n, x - y_j)$ .

Then  $\mathcal{I}(0) = 0$ ,  $\Theta(0) = (0, 0)$  and  $\mathcal{S}(0, 0) = 0$ . Recall also from Lemma 2 that

$$\|v\|_{L^2} + \|\mathbf{y}\| + \|R_j(S_n) - \tilde{R}_j(S_n)\|_{H^1} \leq C\|w\|_{L^2}.$$

To prove Lemma 3, we claim that  $\Psi = \mathcal{S} \circ \Theta \circ \mathcal{I}$  is a diffeomorphism on a fixed neighbourhood of  $0 \in \mathbb{R}^{2N}$  by computing  $d\Psi = d\mathcal{S} \circ d\Theta \circ d\mathcal{I}$ . Indeed, we claim

**Claim.**

$$d\Psi(\mathbf{b}) = \begin{pmatrix} A & (\int Z^+ Z^-)A \\ (\int Z^+ Z^-)A & A \end{pmatrix} + O(e^{-\sigma_0^{3/2} S_n} + \|\mathbf{b}\|),$$

where  $A = \text{diag}((\|Z_j\|_{L^2}^2)_j) = \text{diag}((c_j^{\frac{5-p}{2(p-1)}})_j)$ .

**Remark 4.** Note that if  $N = 1$  (only one soliton), with e.g.  $c_1 = 1$ , then the map  $\Psi$  is represented by the matrix

$$B = \begin{pmatrix} \int (Z^+)^2 & \int Z^+ Z^- \\ \int Z^+ Z^- & \int (Z^-)^2 \end{pmatrix} = \text{Gramm}(Z^\pm).$$

Indeed, the functions  $Z^\pm$  are orthogonal to  $Q_x$ , so that  $\mathbf{y} = 0$  in this case and  $\Psi$  is linear. Since  $Z^\pm$  are linearly independent (see proof of Lemma 1), the matrix  $B$  is invertible.

The claim means that for the general case  $N \geq 2$ , we obtain a similar behavior around each soliton plus small terms due to the interaction of the various solitons.

**Proof.** We start with the computation of differentials of  $\mathcal{I}$ ,  $\Theta$  and  $\mathcal{S}$ . First,  $\mathcal{I}$  is affine so that  $d\mathcal{I}(\mathbf{b}) = \mathcal{I}$  for all  $\mathbf{b}$ . Second, for  $h \in H^1$ ,  $\mathbf{z} \in \mathbb{R}^N$ ,

$$(d\mathcal{S}(v, \mathbf{y}) \cdot (h, \mathbf{z}))_{j,\pm} = -z_j \int v \tilde{Z}_j^\pm_x + \int h \tilde{Z}_j^\pm.$$

Finally, we consider  $\Theta$ . Let  $\Phi$  and  $\phi$  be defined as in the proof of the Lemma 2 above for  $R(S_n)$ . Then, by (A.1),  $d_{\mathbf{y}}\Phi(w, \mathbf{y})$  is a diagonally dominant matrix and thus it is invertible. Denoting by  $M$  its inverse, it follows from (A.1) that

$$M = \text{diag}((\|Q_{c_{jx}}\|_{L^2}^{-2})_j) + O(\|w\|_{L^2} + \|\mathbf{y}\| + e^{-\sigma_0^{3/2}S_n}).$$

Differentiating  $\Phi(w, \phi(w)) = 0$  with respect to  $w$ , we find  $d\phi = -M \circ d_w\Phi$ . Since  $(d_w\Phi(w, \mathbf{y}).h)_j = \int h\tilde{R}_{jx}(S_n)$  and

$$\Theta(w) = \left( w + R - \sum_j R_j(S_n, \cdot - \phi(w)_j), \phi(w) \right),$$

we obtain

$$\begin{aligned} d\Theta(w).h &= (h - \sum_j \tilde{R}_{jx}(S_n)((M \circ d_w\Phi).h)_j, -M \circ d_w\Phi.h) \\ &= \left( h + \sum_{j=1}^N \|\tilde{R}_{jx}(S_n)\| \frac{\int h\tilde{R}_{jx}(S_n)}{\|Q_{c_{jx}}\|_{L^2}^2}, \left( -\frac{\int h\tilde{R}_{jx}(S_n)}{\|Q_{c_{jx}}\|_{L^2}^2} \right)_{j=1, \dots, N} \right) \\ &\quad + O(\|h\|_{L^2}(e^{-\sigma_0^{3/2}S_n} + \|w\|_{L^2})). \end{aligned}$$

Let  $\tilde{\mathbf{b}} \in \mathbb{R}^{2N}$ . Since  $\mathcal{I}$  is linear, we have

$$d\Psi(\mathbf{b}).\tilde{\mathbf{b}} = d\mathcal{S}(\Theta(\mathcal{I}(\tilde{\mathbf{b}}))).(d\Theta(\mathcal{I}(\tilde{\mathbf{b}})).\mathcal{I}(\tilde{\mathbf{b}})).$$

By the previous computations, we derive

$$\begin{aligned} d\Theta(\mathcal{I}(\tilde{\mathbf{b}})).\mathcal{I}(\tilde{\mathbf{b}}) &= \\ &= \left( \mathcal{I}(\tilde{\mathbf{b}}) + \sum_{j=1}^N \tilde{R}_{jx}(S_n) \frac{\int \mathcal{I}(\tilde{\mathbf{b}})\tilde{R}_{jx}(S_n)}{\|Q_{c_{jx}}\|_{L^2}^2}, \left( -\frac{\int \mathcal{I}(\tilde{\mathbf{b}})\tilde{R}_{jx}(S_n)}{\|Q_{c_{jx}}\|_{L^2}^2} \right)_{j=1, \dots, N} \right) \\ &\quad + O(\|\tilde{\mathbf{b}}\|(e^{-\sigma_0^{3/2}S_n} + \|\tilde{\mathbf{b}}\|)). \end{aligned}$$

Inserting the expression of  $\mathcal{I}(\tilde{\mathbf{b}})$ , using  $\|\mathbf{y}\| \leq C\|\mathbf{b}\|$ ,  $\int Z^\pm Q_x = 0$  and the decay properties of the functions  $Q$  and  $Z$ , we get

$$d\Theta(\mathcal{I}(\tilde{\mathbf{b}})).\mathcal{I}(\tilde{\mathbf{b}}) = (\mathcal{I}(\tilde{\mathbf{b}}), 0) + O(\|\tilde{\mathbf{b}}\|(e^{-\sigma_0^{3/2}S_n} + \|\tilde{\mathbf{b}}\|)).$$

Therefore, using the expression of  $d\mathcal{S}$ , we finally obtain

$$d\Psi(\mathbf{b}) = \text{Gramm}((Z_j^\pm)_{j,\pm}) + O(e^{-\sigma_0^{3/2}S_n} + \|\mathbf{b}\|) = P + O(e^{-\sigma_0^{3/2}S_n} + \|\mathbf{b}\|)$$

where  $\text{Gramm}((Z_j^\pm)_{j,\pm})$  is the Gramm matrix of the family  $(Z_j^\pm)_{j,\pm}$

$$\text{Gramm}((Z_j^\pm)_{j,\pm})_{(j_1,\pm_1),(j_2,\pm_2)} = \int Z_{j_1}^{\pm_1} Z_{j_2}^{\pm_2},$$

and

$$P = \begin{pmatrix} A & (f Z^+ Z^-)A \\ (f Z^+ Z^-)A & A \end{pmatrix},$$

where  $A = \text{diag}((\|Z_j\|_{L^2}^2)_j) = \text{diag}((c_j^{\frac{5-p}{2(p-1)}})_j)$  (recall that  $\|Z^\pm\|_{L^2} = 1$ ). This finishes the proof of the claim. ■

Since  $P$  is invertible ( $Z^+$  and  $Z^-$  are independent, see proof of Lemma 1), we deduce that  $d\Psi$  is invertible on some ball  $B_{\mathbb{R}^{2N}}(\eta)$  ( $\eta > 0$  independent of  $n$  for  $n \geq n_0$  large enough). As a consequence,  $\Psi$  is a diffeomorphism from  $B_{\mathbb{R}^{2N}}(\eta)$  to some neighbourhood  $\mathcal{W}$  of  $0 \in \mathbb{R}^{2N}$ . Let  $\delta > 0$  be such that  $B_{\mathbb{R}^{2N}}(\delta) \subset \mathcal{W}$ . For any  $\mathbf{a}^+ \in B_{\mathbb{R}^N}(\delta)$ , there exist a unique  $\mathbf{b} = \mathbf{b}(\mathbf{a}^+) \in B_{\mathbb{R}^{2N}}(\eta)$  such that  $\Psi(\mathbf{b}(\mathbf{a}^+)) = (\mathbf{a}^+, 0)$  and  $\|\mathbf{b}(\mathbf{a}^+)\| \leq C\|\mathbf{a}^+\|$ . ■

**Acknowledgement.** The authors would like to thank Galina Perelman for useful discussions and comments and the anonymous referee for his careful reading.

## References

- [1] BERESTYCKI, H. AND LIONS, P.-L.: Nonlinear scalar field equations. I. Existence of a ground state. *Arch. Rational Mech. Anal.* **82** (1983), no. 4, 313–345.
- [2] BONA, J. L., SOUGANIDIS, P. E. AND STRAUSS, W. A.: Stability and instability of solitary waves of Korteweg-de Vries type. *Proc. Roy. Soc. London Ser. A* **411** (1987), no. 1841, 395–412.
- [3] CAZENAVE, T. AND LIONS, P.-L.: Orbital stability of standing waves for some nonlinear Schrödinger equations. *Comm. Math. Phys.* **85** (1982), no. 4, 549–561.
- [4] CAZENAVE, T. AND WEISSLER, F.: The Cauchy problem for the critical nonlinear Schrödinger equation in  $H^s$ . *Nonlinear Anal.* **14** (1990), 807–836.
- [5] DUYNCKAERTS, T. AND MERLE, F.: Dynamics of threshold solutions for energy-critical wave equation. *Int. Math. Res. Pap. IMRP* **2007**, no. 4, Art. ID rpn002, 67 pp. (2008).
- [6] DUYNCKAERTS, T. AND ROUDENKO, S.: Threshold solutions for the focusing 3D cubic Schrödinger equation. *Rev. Mat. Iberoam.* **26** (2010), no. 1, 1–56.
- [7] EL DIKA, K.: Asymptotic stability of solitary waves for the Benjamin-Bona-Mahony equation. *Discrete Contin. Dyn. Syst.* **13** (2005), 583–622.
- [8] EL DIKA, K. AND MARTEL, Y.: Stability of  $N$  solitary waves for the generalized BBM equations. *Dyn. Partial Differ. Equ.* **1** (2004), 401–437.

- [9] GIDAS, B., NI, W. M. AND NIRENBERG, L.: Symmetry of positive solutions of nonlinear elliptic equations in  $\mathbb{R}^n$ . In *Mathematical analysis and applications, Part A*, 369–402. Adv. in Math. Suppl. Stud. **7**. Academic Press, New York, 1981.
- [10] GINIBRE, J. AND VELO, G.: On a class of nonlinear Schrödinger equations. I. The Cauchy problem, general case. *J. Funct. Anal.* **32** (1979), no. 1, 1–32.
- [11] GRILLAKIS, M.: Analysis of the linearization around a critical point of an infinite dimensional hamiltonian system. *Comm. Pure Appl. Math.* **43** (1990), no. 3, 299–333.
- [12] GRILLAKIS, M., SHATAH, J. AND STRAUSS, W.: Stability theory of solitary waves in the presence of symmetry. I. *J. Funct. Anal.* **74** (1987), no. 1, 160–197.
- [13] KATO, T.: On the Cauchy problem for the (generalized) Korteweg-de Vries equation. In *Studies in applied mathematics*, 93–128. Adv. Math. Suppl. Stud. **8**. Academic Press, New York, 1983.
- [14] KENIG, C. E., PONCE, G. AND VEGA, L.: Well-posedness and scattering result for the generalized Korteweg-De Vries equation via the contraction principle. *Comm. Pure Appl. Math.* **46** (1993), 527–620.
- [15] KWONG, M. K.: Uniqueness of positive solutions of  $\Delta u - u + u^p = 0$  in  $\mathbb{R}^n$ . *Arch. Rational Mech. Anal.* **105** (1989), no. 3, 243–266.
- [16] MARTEL, Y.: Asymptotic  $N$ -soliton-like solutions of the subcritical and critical generalized Korteweg-de Vries equations. *Amer. J. Math.* **127** (2005), no. 5, 1103–1140.
- [17] MARTEL, Y. AND MERLE, F.: Asymptotic stability of solitons for subcritical generalized KdV equations. *Arch. Ration. Mech. Anal.* **157** (2001), no. 3, 219–254.
- [18] MARTEL, Y. AND MERLE, F.: Instability of solitons for the critical generalized Korteweg-De Vries equation. *Geom. Funct. Anal.* **11** (2001), no. 1, 74–123.
- [19] MARTEL, Y. AND MERLE, F.: Stability of blow-up profile and lower bounds for blow-up rate for the critical generalized KdV equation. *Ann. of Math. (2)* **155** (2002), no. 1, 235–280.
- [20] MARTEL, Y. AND MERLE, F.: Multi solitary waves for nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **23** (2006), 849–864.
- [21] MARTEL, Y., MERLE, F. AND TSAI, T.-P.: Stability and asymptotic stability in the energy space of the sum of  $n$  solitons for subcritical gKdV equations. *Comm. Math. Phys.* **231** (2002), 347–373.
- [22] MARTEL, Y., MERLE, F. AND TSAI, T.-P.: Stability in  $H^1$  of the sum of  $K$  solitary waves for some nonlinear Schrödinger equations. *Duke Math. J.* **133** (2006), no. 3, 405–466.

- [23] MERLE, F.: Construction of solutions with exactly  $k$  blow-up points for the Schrödinger equation with critical nonlinearity. *Comm. Math. Phys.* **129** (1990), no. 2, 223–240.
- [24] MIURA, R. M.: The Korteweg-de Vries equation: a survey of results. *SIAM Rev.* **18** (1976), 412–459.
- [25] MIZUMACHI, T.: Asymptotic stability of solitary wave solutions to the regularized long-wave equation. *J. Differential Equations* **200** (2004), 312–341.
- [26] PEGO, R. L. AND WEINSTEIN, M. I.: Eigenvalues, and instabilities of solitary waves. *Philos. Trans. Roy. Soc. London Ser. A* **340** (1992), no. 1656, 47–94.
- [27] PEGO, R. L. AND WEINSTEIN, M. I.: Asymptotic stability of solitary waves. *Comm. Math. Phys.* **164** (1994), 305–349.
- [28] SCHLAG, W.: Spectral theory and nonlinear partial differential equations: a survey. *Discrete Contin. Dyn. Syst.* **15** (2006), no. 3, 703–723.
- [29] WEINSTEIN, M. I.: Modulational stability of ground states of nonlinear Schrödinger equations. *SIAM J. Math. Anal.* **16** (1985), 472–491.
- [30] WEINSTEIN, M. I.: Lyapunov stability of ground states of nonlinear dispersive evolution equations. *Comm. Pure Appl. Math.* **39** (1986), 51–68.

*Recibido:* 28 de abril de 2009

*Revisado:* 16 de septiembre de 2009

Raphaël Côte  
CNRS and CMLS, UMR 7640  
Ecole Polytechnique  
91128 Palaiseau, France  
cote@math.polytechnique.fr

Yvan Martel  
Mathématiques, UMR 8100  
Univ. Versailles–St. Quentin  
78035 Versailles, France  
yvan.martel@math.uvsq.fr

Frank Merle  
Mathématiques, UMR 8088  
Univ. Cergy-Pontoise  
95302 Cergy-Pontoise, France and IHES  
Frank.Merle@u-cergy.fr