



Erratum: Fitting a C^m -smooth function to data II

Charles Fefferman and Bo'az Klartag

A few small changes have to be made to the proof of Lemma 3 in Section 30, pages 139–140. The definition of P'_t in (52) in page 139 should include a factor of $\lambda^{-\|\gamma_t\|}$. That is, the correct definition should be

$$(52) \quad P'_t = \frac{(\alpha'_t)!}{(\alpha'_t + \gamma_t)!} \lambda^{-\|\gamma_t\|} x^{\gamma_t} \odot_0 \bar{P}_{\alpha'_t}.$$

Furthermore, we need to replace $\lambda^{\|\beta\|}$ by $\lambda^{\|\beta+\gamma_t\|}$ in (54), we need to replace $\lambda^{\|\alpha'_s\|}$ by $\lambda^{\|\alpha'_s+\gamma_s\|}$ in the definition of the matrix A and also in (57), and we need to replace $\lambda^{\|\alpha'_t\|}$ by $\lambda^{\|\alpha'_t+\gamma_t\|}$ in (58) and (59). The rest of the proof stays intact.

References

- [1] FEFFERMAN, C. AND KLARTAG, B.: Fitting a C^m -smooth function to data II. *Rev. Mat. Iberoam.* **25** (2009), no. 1, 49–273.

Received August 8, 2012.

CHARLES FEFFERMAN: Department of Mathematics, Princeton University, Fine Hall, Washington Road, Princeton, New Jersey 08544, USA.

E-mail: cf@math.princeton.edu

BO'AZ KLARTAG: School of Mathematical Sciences, Tel-Aviv University, Tel-Aviv 69978 Israel.

E-mail: klartagb@post.tau.ac.il

Mathematics Subject Classification (2010): 65D05, 65D17.

Keywords: Algorithm, interpolation, approximation, C^m -smoothness.