



Pitt inequalities and restriction theorems for the Fourier transform

Laura De Carli, Dmitry Gorbachev and Sergey Tikhonov

Abstract. We prove new Pitt inequalities for the Fourier transforms with radial and non-radial weights using weighted restriction inequalities for the Fourier transform on the sphere. We also prove new Riemann–Lebesgue estimates and versions of the uncertainty principle for the Fourier transform.

1. Introduction

Weighted inequalities for the Fourier transform provide a natural balance between functional growth and smoothness. On \mathbb{R}^n it is important to determine quantitative comparisons between the relative size of a function and its Fourier transform at infinity. We will let $\widehat{f}(\xi) = \int_{\mathbb{R}^n} e^{ix\xi} f(x) dx$, $\xi \in \mathbb{R}^n$, be the Fourier transform in $L^1(\mathbb{R}^n)$, and $\|\cdot\|_p$ be the standard norm in $L^p(\mathbb{R}^n)$. We consider Pitt type inequalities

$$(1.1) \quad \|u^{1/q} \widehat{f}\|_q \leq C \|v^{1/p} f\|_p, \quad f \in C_0^\infty(\mathbb{R}^n).$$

Here and throughout the paper, u and v are non-negative measurable functions on \mathbb{R}^n , and $1 \leq p, q \leq \infty$ unless otherwise specified. We will use C to denote numeric constants that may change from line to line. We will let $p' = p/(p-1)$ be the conjugate exponent of $1 \leq p \leq \infty$, and we will often let $x = \rho\omega$, with $\omega \in \mathbb{S}^{n-1}$ and $\rho = |x|$. We denote by $|E|$ the Lebesgue measure of a set E and by $\chi_E(x)$ be the characteristic function of E .

In 1983, Heinig [17], Jurkat–Sampson [18] and Muckenhoupt [20], [21] proved:

Theorem 1.1. *Let $n \geq 1$. If the weights u and v satisfy*

$$(1.2) \quad \sup_{s>0} \left(\int_0^s u^*(t) dt \right)^{1/q} \left(\int_0^{1/s} [(1/v)^*(t)]^{1/(p-1)} dt \right)^{1/p'} = C < \infty,$$

Mathematics Subject Classification (2010): Primary 42B10; Secondary 42C20, 46E30.

Keywords: Pitt inequalities, restriction theorems, weights, Riemann–Lebesgue estimate, uncertainty principle.

for $1 < p \leq q < \infty$, where g^* is the non-increasing rearrangement of g , then (1.1) holds.

To formulate necessary conditions for inequality (1.1) to hold, we recall the definition of polar set. If $A \subset \mathbb{R}^n$,

$$A^* = \{\xi \in \mathbb{R}^n : |x\xi| \leq 1, x \in A\}$$

is the polar set of A (see [27], § 4, p. 111). We prove the following.

Theorem 1.2. *Let $n \geq 1$. Suppose that the Pitt inequality (1.1) holds for any $f \in C_0^\infty(\mathbb{R}^n)$ and for $1 < p, q < \infty$.*

(1) *Let $A \subset \mathbb{R}^n$. Then*

$$(1.3) \quad \sup_A \left(\int_{cA^*} u(\xi) d\xi \right)^{1/q} \left(\int_A v^{1-p'}(x) dx \right)^{1/p'} = C < \infty,$$

where $c < \pi/2$ and A^* is the polar set of the set A .

(2) *Let the weights $u(x) = u_0(|x|)$ and $v(x) = v_0(|x|)$ be radial. Then*

$$(1.4) \quad \sup_{s>0} \left(\int_{|x|<c_n/s} u(x) dx \right)^{1/q} \left(\int_{|x|<s} v^{1-p'}(x) dx \right)^{1/p'} = C < \infty,$$

where c_n is any positive number less than $q_{n/2-1}$, the first zero of the Bessel function $J_{n/2-1}(t)$. In particular, $q_{n/2-1} \geq \pi/2$.

(3) *Results of the part (1) also hold if one replaces the sets A and cA^* by a union of their disjoint translations, that is, by the sets $A_1 = \bigcup_{j=1}^{N_1} (A + x_j)$ and $A_2 = \bigcup_{j=1}^{N_2} (cA^* + \xi_j)$ for any x_j and ξ_j .*

Note that in this theorem we do not assume $q \geq p$.

Part (2) of the theorem is known with a smaller constant c ; see the proof of Theorem 3.1 in [17]. Moreover, part (3) generalizes the following necessary condition (see [4], Theorem 3):

$$(1.5) \quad \left(\int_{Q_1} u(\xi) d\xi \right)^{1/q} \left(\int_{Q_2} v^{1-p'}(x) dx \right)^{1/p'} = C < \infty,$$

for all cubes Q_1 and Q_2 such that $|Q_1||Q_2| = 1$.

We should also mention Theorem 2.1 in [19], where a necessary condition similar to (1.5), with u replaced by a measure $d\mu$, was proved.

When $u(x) = u_0(|x|)$ and $v(x) = v_0(|x|)$ are radial, with $u_0(\cdot)$ non-increasing and $v_0(\cdot)$ non-decreasing, then (1.4) is necessary and sufficient for the validity of (1.1) (see [17]). In particular, if $u(x)$ and $v(x)$ are locally integrable power weights, i.e., in the form of $u = |x|^b$ and $v = |x|^a$, with $a, b > -n$, we get that the classical Pitt inequality

$$(1.6) \quad \left(\int_{\mathbb{R}^n} |\widehat{f}(\xi)|^q |\xi|^b d\xi \right)^{1/q} \leq C \left(\int_{\mathbb{R}^n} |f(x)|^p |x|^a dx \right)^{1/p},$$

where $f \in C_0^\infty(\mathbb{R}^n)$, holds if and only if

$$(1.7) \quad \frac{a}{p} + \frac{b}{q} = n \left(1 - \frac{1}{p} - \frac{1}{q} \right),$$

$$(1.8) \quad -n < b \leq 0, \quad \text{and} \quad 0 \leq a < n(p-1);$$

see [22], [24], and [2].

Pitt inequalities with power weights that satisfy less restrictive conditions than those in (1.8) are only valid on special subspaces of $L^p(\mathbb{R}^n)$. We have proved in [11] that if f is a product of a radial function and a spherical harmonics of degree $k \geq 0$, then (1.1) is satisfied with $u = |x|^a$ and $v = |x|^b$ if and only if a and b satisfy (1.7) and

$$(1.8') \quad (n-1) \left(\frac{1}{2} - \frac{1}{p} \right) + \max \left\{ \frac{1}{p'} - \frac{1}{q}, 0 \right\} \leq \frac{b}{p} < \frac{n}{p'} + k,$$

which is less restrictive than the conditions in (1.8) even for $k = 0$.

In this paper we prove L^p - L^q Pitt inequalities for radial and non-radial weights u and v . Our main tools are weighted restriction inequalities for the Fourier transform in \mathbb{R}^n , $n \geq 2$. That is,

$$(1.9) \quad \left(\int_{\mathbb{S}^{n-1}} |\widehat{f}(\omega)|^q U(\omega) d\sigma(\omega) \right)^{1/q} \leq C \left(\int_{\mathbb{R}^n} |f(x)|^p v(x) dx \right)^{1/p},$$

where U and v are non-negative and measurable on \mathbb{S}^{n-1} and \mathbb{R}^n , respectively, and $f \in C_0^\infty(\mathbb{R}^n)$.

We recall several known restriction theorems in Section 2. In Section 3 we present new Pitt inequalities using restriction inequalities. In particular, we prove the following.

Theorem 1.3. *Let $1 \leq p < \frac{2(n+2)}{n+4}$ and $1 \leq q \leq \frac{n-1}{n+1} p'$, with $n \geq 2$. Suppose that $u(x) = u_0(|x|)$ satisfies*

$$(1.10) \quad \int_0^\infty \rho^{n-1-qn/p'} u_0(\rho) d\rho < \infty.$$

Then for every $f \in C_0^\infty(\mathbb{R}^n)$,

$$(1.11) \quad \|u^{1/q} \widehat{f}\|_q \leq C \|f\|_p.$$

Remark 1.4. (i) The proof of Theorem 3.1, of which Theorem 1.3 is a special case, shows that the constant C in (1.11) equals $C' \left(\int_0^\infty \rho^{n-1-qn/p'} u_0(\rho) d\rho \right)^{1/q}$, where C' depends on n, p, q .

(ii) When $u \in L^p(\mathbb{R}^n)$ with $1 \leq p \leq 2$ and $q = 1$, (1.11) is valid also when u is not radial; indeed, by the Hausdorff–Young inequality,

$$\|u \widehat{f}\|_1 \leq \|u\|_p \|\widehat{f}\|_{p'} \leq C \|f\|_p.$$

(iii) Theorem 1.1 and most of the Pitt inequalities in the literature are proved for $1 < p \leq q < \infty$. Theorem 1.3 provides a rather simple sufficient condition for (1.11) that applies either when $p \leq q$ or $p > q$. Note that the known sufficient conditions for (1.11) are usually quite difficult to verify especially in the case $p > q$ (see for example [2]).

Theorem 1.3 applies in cases where Theorem 1.1 does not: in Section 4 we construct a radial weight u for which the inequality (1.2) does not hold, but (1.10) holds for u_0 and therefore (1.11) is valid.

The rest of the paper is organized as follows. In Section 5 we prove necessary conditions for the Pitt inequality (1.1) to hold (Theorem 1.1), necessary conditions for the weighted restriction inequality (1.9) to hold (Proposition 2.2), and sufficient conditions from Section 3. These are the main results of the paper.

In Section 6 we prove new versions of the uncertainty principle for the Fourier transform.

In Section 7 we apply one of our new Pitt-type inequality to get a quantitative version of the Riemann–Lebesgue lemma, which provides an interrelation between the smoothness of a function and the growth properties of the Fourier transforms.

Finally, we would like to mention make the following interesting observation which perhaps is not new: the Pitt inequality (1.1) holds if and only if, for some $s \geq p$, we have $\|u^{1/q}\widehat{f}\|_q \leq C\|w^{-1}\|_{p/(s-p)}^{1/s}\|w^{1/s}v^{1/p}f\|_s$ whenever $w^{-1} \in L^{p/(s-p)}(\mathbb{R}^n)$. In particular, the inequality $\|\widehat{f}\|_{p'} \leq C\|w^{1/s}f\|_s$ holds for every $1 \leq p \leq 2$ whenever $w^{-1} \in L^{p/(s-p)}(\mathbb{R}^n)$, $s \geq p$. We will prove this fact in Section 5.

2. Restriction theorems for the Fourier transform

The Tomas–Stein restriction inequality for the Fourier transform on the unit sphere states that, for every $f \in C_0^\infty(\mathbb{R}^n)$ with $n \geq 2$, for $1 \leq q \leq \frac{n-1}{n+1}p'$ and $1 \leq p \leq \frac{2(n+1)}{n+3}$,

$$(2.1) \quad \left(\int_{\mathbb{S}^{n-1}} |\widehat{f}(\omega)|^q \chi(\omega) d\sigma(\omega) \right)^{1/q} \leq C \left(\int_{\mathbb{R}^n} |f(x)|^p dx \right)^{1/p},$$

where $d\sigma(\omega)$ is the induced Lebesgue measure on \mathbb{S}^{n-1} and $\chi \in C_0^\infty(\mathbb{S}^{n-1})$. The same inequality holds also if $\chi(\omega)d\sigma(\omega)$ is replaced by $d\sigma(\omega)$ ([29], [25], VII, §4).

Thus, if $T(f) = \widehat{f}|_{\mathbb{S}^{n-1}}$ is the restriction operator, T maps $L^p(dx)$ into $L^q(d\sigma)$ boundedly whenever p, q are as in the Tomas–Stein theorem.

Note that (2.1) is trivial when $p = 1$ because

$$\left(\int_{\mathbb{S}^{n-1}} |\widehat{f}(\omega)|^q d\sigma(\omega) \right)^{1/q} \leq \omega_{n-1}^{1/q} \|\widehat{f}\|_\infty \leq \omega_{n-1}^{1/q} \|f\|_1,$$

where $\omega_{n-1} = |\mathbb{S}^{n-1}|$.

The restriction conjecture states that inequality (2.1) is valid for all $1 \leq q \leq \frac{n-1}{n+1} p'$ and $1 \leq p < \frac{2n}{n+1}$. When $n = 2$ the restriction conjecture has been proved by C. Fefferman [13]. When $n \geq 3$, T. Tao [28] has proved that (2.1) is valid for $1 \leq p < \frac{2(n+2)}{n+4}$. Note that $\frac{2(n+2)}{n+4} = \frac{2n}{n+1}$ when $n = 2$.

Weighted versions of the restriction inequality (2.1) in the form of

$$(2.2) \quad \left(\int_{\mathbb{S}^{n-1}} |\widehat{f}(\omega)|^q U(\omega) \, d\sigma(\omega) \right)^{1/q} \leq C \left(\int_{\mathbb{R}^n} |f(x)|^p v(x) \, dx \right)^{1/p}$$

have been proved by several authors. In most of the theorems in the literature, $1 \leq p \leq q \leq \infty$ and $U(\omega)$ is the restriction of a function $\widetilde{U}(x) \in C^\infty(\mathbb{R}^n)$, often with compact support in \mathbb{S}^{n-1} .

Note that, when $v \in L^1_{\text{loc}}(\mathbb{R}^n)$, a necessary condition for the inequality (2.2) to hold is that $U(\omega) \in L^1(\mathbb{S}^{n-1})$. To see this, apply (2.2) to a radial function $f \in C^\infty_0(\mathbb{R}^n)$.

The following duality argument will be used in the proof of the theorems in the next section. The technique is well known, but we state and prove Lemma 2.1 in this paper for the sake of completeness.

Lemma 2.1. *Assume $U(x/|x|) = U(\omega) \in L^1(\mathbb{S}^{n-1})$. Inequality (2.2) is equivalent to*

$$(2.3) \quad \left\| \int_{\mathbb{S}^{n-1}} g(\omega) e^{i\omega y} U^{1/q}(\omega) \, d\sigma(\omega) \right\|_{L^{p'(v^{1-p'} dy)}} \leq C \|g\|_{L^{q'}(\mathbb{S}^{n-1})}.$$

In Section 5 we prove necessary conditions for the weighted restriction inequality (2.2) to hold. To the best of our knowledge these results are new.

Proposition 2.2. *Assume that the inequality (2.2) holds with $U^{1-q'}(\omega) \in L^1(\mathbb{S}^{n-1})$. Then*

$$(2.4) \quad \int_{\mathbb{R}^n} v^{1-p'}(x) |j_{n/2-1}(|x|)|^{p'} \, dx < C,$$

where $j_\nu(t) = \Gamma(\nu + 1)(t/2)^{-\nu} J_\nu(t)$ is the normalized Bessel function.

A special case of (2.4) is in [5], (3.1). In particular, we obtain the following result.

Corollary 2.3. *Assume that the inequality (2.2) holds with $U^{1-q'}(\omega) \in L^1(\mathbb{S}^{n-1})$; assume v is radial and non-negative, and that $v(x) = v_0(|x|)$ satisfies either*

$$(2.5) \quad \int_A v_0^{1-p'}(t - |A|) \, dt \leq C \int_A v_0^{1-p'}(t) \, dt,$$

or

$$(2.6) \quad \int_A v_0^{1-p'}(t + |A|) \, dt \leq C \int_A v_0^{1-p'}(t) \, dt,$$

for all finite intervals A , with a constant C independent of A . Then

$$\int_{\mathbb{R}^n} v^{1-p'}(x) (1 + |x|)^{-p'(n-1)/2} dx < C.$$

Remark 2.4. If $v_0^{1-p'}$ satisfies a doubling condition, that is,

$$\int_{2A} v_0^{1-p'}(t) dt \leq C \int_A v_0^{1-p'}(t) dt,$$

for all intervals A , where $2A$ is the interval twice the length of A and with the mid-point coinciding with that of A , then both (2.5) and (2.6) hold. If v_0 is monotonic, then at least one of the conditions (2.5) and (2.6) hold.

Weighted restriction theorems were intensively studied for piecewise power weights, i.e., in the form of

$$(2.7) \quad v(x) = \begin{cases} |x|^\alpha, & \text{if } |x| \leq 1, \\ |x|^\beta, & \text{if } |x| > 1, \end{cases}$$

see e.g. [5]. The method of the proof of Corollary 2.8 in [8] can be used to prove the following.

Lemma 2.5. *Let $d\mu$ and dv be measures on \mathbb{R}^n , $n \geq 1$, and let $1 \leq p \leq q$ and $s \geq p$. An operator T maps $L^p(d\mu) \rightarrow L^q(dv)$ boundedly if and only if T maps $L^s(w d\mu) \rightarrow L^q(dv)$ boundedly whenever $w^{-1} \in L^{p/(s-p)}(d\mu)$ and*

$$\| \|T\| \|_{L^s(w d\mu) \rightarrow L^q(dv)} \leq C \|w^{-1}\|_{L^{p/(s-p)}(d\mu)}^{1/s}.$$

The proof is in Section 5. If we apply Lemma 2.5 to the restriction operator, with the the Tomas–Stein exponents $s = q = 2$ and $p = \frac{2(n+1)}{n+3}$, we require $w^{-1} \in L^{(n+1)/2}(\mathbb{R}^n)$. This condition, applied to piecewise power weight, allows $\alpha < \frac{2n}{n+1}$ and $\beta > \frac{2n}{n+1}$.

These exponents are not sharp: S. Bloom and G. Sampson have proved in [5] a number of restriction theorems with piecewise power weights, and have obtained, in most cases, sharp conditions on α and β . One of the results in [5], Theorem 5.6, is the following.

Theorem 2.6. *Let $1 < p \leq 2$, $n \geq 2$, $2 \leq q \leq \frac{n-1}{n+1} p'$. Let $v(x)$ given by (2.7). Then (2.2) with $U = 1$ holds if and only if $\alpha < n(p-1)$ and $\beta \geq 0$. Moreover, (2.2) holds with $p = q = 2$ also when $U = 1$ and $v(x)$ is as in (2.7) with $\alpha < n$ and $\beta > 1$.*

We also notice that weighted restriction theorems have been proved for weights in the Campanato–Morrey spaces: for $0 \leq \alpha \leq n/r$ and $r \geq 1$, the Campanato–Morrey space $\mathcal{L}^{\alpha,r}$ is defined as

$$\mathcal{L}^{\alpha,r} = \left\{ f \in L^r_{\text{loc}}(\mathbb{R}^n) : \|f\|_{r,\alpha} = \sup_{\substack{x \in \mathbb{R}^n \\ \rho > 0}} \rho^\alpha \left(\rho^{-n} \int_{|y-x| < \rho} |f(y)|^r dy \right)^{1/r} < \infty \right\}.$$

Note that $\mathcal{L}^{\alpha,n/\alpha} = L^{n/\alpha}(\mathbb{R}^n)$ and $\mathcal{L}^{0,r}(\mathbb{R}^n) = L^\infty(\mathbb{R}^n)$.

A. Ruiz and L. Vega have proved in [23] the following.

Theorem 2.7. *Suppose that $V \in \mathcal{L}^{\alpha,r}$, with $\frac{\alpha}{n} \leq \frac{1}{r} < \frac{2(\alpha-1)}{n-1}$ and $\frac{2n}{n+1} < \alpha \leq n$, $n \geq 2$. Then, the inequality*

$$(2.8) \quad \left(\int_{\mathbb{S}^{n-1}} |\widehat{f}(\omega)|^2 d\sigma(\omega) \right)^{1/2} \leq C \left(\int_{\mathbb{R}^n} |f(x)|^2 V(x) dx \right)^{1/2}$$

holds with $C = C' \|V\|_{\alpha,r}^{1/2}$.

Note that (2.8) holds when $V^{-1} \in L^{(n-1)/2}(\mathbb{R}^n)$. Indeed, the Stein–Tomas restriction inequality (2.1) with $q = 2$ and $p = \bar{p} = \frac{2(n+1)}{n+3}$ and the Hölder inequality imply:

$$\left(\int_{\mathbb{S}^{n-1}} |\widehat{f}(\omega)|^2 d\sigma(\omega) \right)^{1/2} \leq C \|f\|_{\bar{p}} \leq C \|fV^{1/2}\|_2 \|V^{-1/2}\|_{n+1}.$$

Note also that in [23] it is proved that

$$\|\widehat{d\sigma} * f\|_{L^2(V)} \leq C' \|V\|_{\alpha,r} \|f\|_{L^2(V^{-1}\mathbb{R}^n)},$$

but we can use Lemma 2.1 to show that this inequality is equivalent to (2.8). See also [1].

Special cases of the restriction inequality in [23] are in [8] and [9]. F. Chiarenza and A. Ruiz have proved in [9] a version of (2.8) with special doubling weights; S. Chanillo and E. Sawyer have proved in [8], Corollary 2.8, that (2.8) holds when V is in the Fefferman–Phong class F_r , with $r \geq (n - 1)/2$. We recall that $V(x) \in F_r$ if

$$\|v\|_{F_r} = \sup_Q |Q|^{2/n} \left(\frac{1}{|Q|} \int_Q |V(x)|^r \right)^{1/r} < \infty$$

for all cubes $Q \subset \mathbb{R}^n$.

3. New Pitt inequalities

In this section we obtain new Pitt-type inequalities for the Fourier transforms using restriction inequalities from Section 2.

Theorem 3.1. *Assume that the restriction inequality (1.9) holds for some $1 \leq p \leq q \leq \infty$. Let $w(\rho)$ be a measurable function for which $v(\rho x) \leq w(\rho)v(x)$ for a.e. $\rho > 0$ and $x \in \mathbb{R}^n$. Suppose that u is radial, and $u(x) = u_0(|x|)$ satisfies*

$$(3.1) \quad \int_0^\infty \rho^{n-1-qn/p'} u_0(\rho) w^{q/p}(\rho) d\rho < \infty.$$

Then,

$$(3.2) \quad \left(\int_{\mathbb{R}^n} |\widehat{f}(x)|^q U\left(\frac{x}{|x|}\right) u(x) dx \right)^{1/q} \leq C \|v^{1/p} f\|_p.$$

Theorem 1.3 is an easy consequence of Theorem 3.1 (with $U \equiv v \equiv 1$) and the Fefferman–Tao restriction theorem.

In the next section we will show that our theorem can be applied in cases where prior results are not applicable.

Our next result deals with piecewise power weight v defined by (2.7). In order to use Theorem 3.1, we need to find $w(\rho)$ so that $v(\rho x) \leq w(\rho)v(x)$, $\rho > 0$. A straightforward calculation shows that in this case

$$(3.3) \quad w(\rho) \leq w_0(\rho) := \max\{\rho^\alpha, \rho^\beta\}.$$

Using Theorem 3.1 and weighted restriction inequalities from Section 5 of [5], we have:

Corollary 3.2. *Let $1 < p \leq 2$ and $2 \leq q \leq \frac{n-1}{n+1}p'$, with $n \geq 2$. Let v be a piecewise power weight $v(x)$ given by (2.7) with $\alpha < n(p-1)$ and $\beta \geq 0$. Let u be a radial weight that satisfies*

$$\int_0^\infty \rho^{n-1-qn/p'} u_0(\rho) w_0^{q/p}(\rho) d\rho < \infty,$$

where w_0 is given by (3.3). Then, for every $f \in C_0^\infty(\mathbb{R}^n)$,

$$(3.4) \quad \|u^{1/q} \widehat{f}\|_q \leq C \|v^{1/p} f\|_p.$$

Remark 3.3. This corollary is valid for all piecewise power weights v and exponents p, q for which the restriction theorems in [5] hold.

The following result uses weights in a Campanato–Morrey class $\mathcal{L}^{\alpha,r}$ (see Section 2 for a definition).

Corollary 3.4. *Let $V \in \mathcal{L}^{\alpha,r}$, with $\frac{2n}{n+1} < \alpha \leq n$ and $\frac{\alpha}{n} \leq \frac{1}{r} < \frac{2(\alpha-1)}{n-1}$, $n \geq 2$. Assume that there exists a measurable function $w(\rho)$ for which $V(\rho x) \leq w(\rho)V(x)$ for a.e. $\rho > 0$ and $x \in \mathbb{R}^n$, and that $u(x) = u_0(|x|)$ satisfies*

$$(3.5) \quad \int_0^\infty \rho^{-1} u_0(\rho) w(\rho) d\rho < \infty.$$

Then, for every $1 \leq p \leq 2$, the following weighted Hausdorff–Young inequality holds:

$$(3.6) \quad \|u^{2/p'} \widehat{f}\|_{p'} \leq C \|V^{2/p'} f\|_p, \quad f \in C_0^\infty(\mathbb{R}^n).$$

4. Comparison of Theorems 1.1 and 1.3

In this section we give an example of radial weight $u(x) = u_0(|x|)$ that satisfies the conditions of Theorem 1.3 while does not satisfy the conditions (1.2) in Theorem 1.1.

We recall that Theorem 1.3 states that the Pitt inequality

$$(4.1) \quad \|u^{1/q} \widehat{f}\|_q \leq C \|f\|_p$$

holds with $1 \leq q \leq \frac{n-1}{n+1} p'$ and $1 \leq p < \frac{2(n+2)}{n+4}$ whenever $u(x) = u_0(|x|)$ satisfies

$$(4.2) \quad \int_0^\infty \rho^{-a} u_0(\rho) \rho^{n-1} d\rho < \infty, \quad a = \frac{qn}{p'} > 0.$$

On the other hand, when u is radial and $v \equiv 1$, the sufficient condition (1.2) in Theorem 1.1 states that

$$\int_0^s u^*(t) dt \leq C s^{q/p'}.$$

The latter is equivalent to the following condition:

$$(4.3) \quad \sup_E |E|^{-q/p'} \int_E u dx < C,$$

where the supremum is taken over all measurable E , $|E| > 0$.

Let now E_0 be a measurable subset of \mathbb{R}_+ . Consider the radial set $E = \{x \in \mathbb{R}^n : |x| \in E_0\}$. For such set, we can rewrite (4.3) as follows:

$$(4.4) \quad \int_{E_0} u_0(\rho) \rho^{n-1} d\rho \leq C |E|^{q/p'} = C \left(\int_{E_0} \rho^{n-1} d\rho \right)^{q/p'}.$$

Let $A = \cup_{k=1}^\infty A_k$, where $A_k = (k, k + k^{-n-1})$. Set

$$(4.5) \quad u_0(\rho) \rho^{n-1} = \sum_{k=1}^\infty k^n \chi_{A_k}(\rho).$$

Then condition (4.2) holds (and so also the Pitt inequality (4.1)) since

$$\begin{aligned} \int_0^\infty \rho^{-a} u_0(\rho) \rho^{n-1} d\rho &= \sum_{k=1}^\infty k^n \int_k^{k+k^{-n-1}} \rho^{-a} d\rho \\ &\leq \sum_{k=1}^\infty k^n k^{-a} k^{-n-1} = \sum_{k=1}^\infty k^{-1-a} < \infty, \end{aligned}$$

and $a > 0$.

On the other hand, taking $E_N = \{x \in \mathbb{R}^n : |x| \in \cup_{k=1}^N A_k\}$, we get

$$|E_N| = \sum_{k=1}^N \int_k^{k+k^{-n-1}} \rho^{n-1} d\rho \leq \sum_{k=1}^\infty (k+1)^{n-1} k^{-n-1} < C.$$

However,

$$(4.6) \quad \int_{E_N} u(x) dx = \sum_{k=1}^N k^n \int_k^{k+k^{-n-1}} d\rho \asymp \ln N.$$

Therefore, (4.4) (and so also (4.3)), do not hold as $N \rightarrow \infty$.

It is worthwhile to remark that for the radial weights u , the necessary condition (1.4) for the Pitt inequality (4.1) to hold (see Theorem 1.2) can be written as

$$\sup_{s>0} \left(\int_0^s u_0(\rho) \rho^{n-1} d\rho \right)^{1/q} \left(\int_0^{c_n/s} \rho^{n-1} d\rho \right)^{1/p'} < C$$

or, equivalently,

$$(4.7) \quad \sup_{s>0} s^{-a} \int_0^s u_0(\rho) \rho^{n-1} d\rho < C,$$

where $a = qn/p' > 0$. For the weight u given by (4.5) it can be easily checked since

$$\int_0^s u_0(\rho) \rho^{n-1} d\rho \leq \sum_{k=1}^{[s]+1} k^{-1} \leq 1 + \ln(s + 1).$$

This of course implies (4.7) since we only have to consider the case $s \rightarrow \infty$.

5. Proofs of the main results

Proof of Theorem 1.2. Let us assume that Pitt inequality (1.1) holds.

(1) Following [17], consider the function $f = \chi_A v^{1-p'} \in L^p(v)$. For any set $B \subset \mathbb{R}^n$ we get

$$C \|v^{1/p} f\|_p \geq \|u^{1/q} \widehat{f}\|_q \geq \left(\int_B |\widehat{f}(\xi)|^q u(\xi) d\xi \right)^{1/q},$$

where

$$\|v^{1/p} f\|_p = \left(\int_A (v^{1-p'}(x))^p v(x) dx \right)^{1/p} = \left(\int_A v^{1-p'}(x) dx \right)^{1/p} > 0$$

and

$$|\widehat{f}(\xi)| \geq \left| \int_A v^{1-p'}(x) \cos(x\xi) dx \right|, \quad \xi \in B.$$

Let $B = c_n A^*$, where $c_n < \pi/2$ and A^* is polar set of the set A . Then for any $x \in A$ and $\xi \in B$ we have $|x\xi| \leq c_n$ and $\cos(x\xi) \geq \cos c_n > 0$. Therefore,

$$|\widehat{f}(\xi)| \geq \cos c_n \int_A v^{1-p'}(x) dx, \quad \xi \in B.$$

Hence,

$$\begin{aligned} C \left(\int_A v^{1-p'}(x) dx \right)^{1/p} &\geq \left(\int_B |\widehat{f}(\xi)|^q u(\xi) d\xi \right)^{1/q} \\ &\geq \cos c_n \left(\int_A v^{1-p'}(x) dx \right) \left(\int_B u(\xi) d\xi \right)^{1/q}, \end{aligned}$$

or, equivalently,

$$\left(\int_{c_n A^*} u(\xi) d\xi \right)^{1/q} \left(\int_A v^{1-p'}(x) dx \right)^{1/p'} < C.$$

(2) If both weights u and v are radial, then the function $f = \chi_A v^{1-p'}$ and its Fourier transform are also radial. Moreover, taking the ball $A = \{x \in \mathbb{R}^n : |x| \leq s\}$ we get

$$\widehat{f}(\xi) = \omega_{n-1} \int_A v^{1-p'}(x) j_{n/2-1}(|\xi|x) dx.$$

Let $q_{n/2-1}$ be the first zero of the normalized Bessel function $j_{n/2-1}(t)$. It is well known (see e.g. [30], Chapter XV) that $q_{n/2-1} \geq q_{-1/2} = \pi/2$ and $q_{n/2-1} \sim n/2$ for $n \geq 1$. Then $j_{n/2-1}(t) \geq j_{n/2-1}(c_n)$, where $c_n \geq t$ can be taken as follows: $\pi/2 < c_n < q_{n/2-1}$ for $n \geq 2$. The rest of the proof is the same as in (1).

(3) To prove this part, we use ideas similar to [4]. In order to consider translations of the sets A and $c_n A^*$ by the vectors x_0 and ξ_0 correspondingly, it is enough to consider the function $g(x) = f(x - x_0)e^{-ix\xi_0}$ so that $|g(x)| = |f(x - x_0)|$ and $|\widehat{g}(\xi)| = |\widehat{f}(\xi - \xi_0)|$. The integral condition (1.3) easily applies to unions of disjoint translations of A and cA^* . □

Proof of Lemma 2.1. Let $A : L^p(v dx) \rightarrow L^q(\mathbb{S}^{n-1})$ be the operator, initially defined for all $f \in C_0^\infty(\mathbb{R}^n)$, by $Af(\omega) = \widehat{f}(\omega)U^{1/q}(\omega)$. Duality gives

$$\begin{aligned} \|Af\|_{L^q(\mathbb{S}^{n-1})} &= \sup_{\|g\|_{L^{q'}(\mathbb{S}^{n-1})} \leq 1} \left| \int_{\mathbb{S}^{n-1}} Af(\omega) g(\omega) d\sigma(\omega) \right| \\ &= \sup_{\|g\|_{L^{q'}(\mathbb{S}^{n-1})} \leq 1} \left| \int_{\mathbb{R}^n} f(x) A^*g(x) dx \right|, \end{aligned}$$

where

$$(5.1) \quad A^*g(x) = \int_{\mathbb{S}^{n-1}} g(\omega) e^{i\omega x} U^{1/q}(\omega) d\sigma(\omega).$$

By Hölder’s inequality,

$$\int_{\mathbb{R}^n} f(x)A^*g(x) dx \leq \|v^{1/p}f\|_p \|v^{-1/p}A^*g\|_{p'} = \|v^{-1/p}A^*g\|_{p'} \|f\|_{L^p(v dx)}.$$

Therefore, the inequality

$$\|v^{-1/p}A^*g\|_{p'} = \left\| \int_{\mathbb{S}^{n-1}} g(\omega) e^{i\omega x} U^{1/q}(\omega) d\sigma(\omega) \right\|_{L^{p'}(v^{1-p'} dx)} \leq C \|g\|_{L^{q'}(\mathbb{S}^{n-1})}$$

implies (2.2). A similar argument shows that the inequality (2.2), or $\|Af\|_{L^q(\mathbb{S}^{n-1})} \leq C \|f\|_{L^p(v dx)}$, implies

$$(5.2) \quad \|v^{-1/p}A^*g\|_{p'} \leq C \|g\|_{L^{q'}(\mathbb{S}^{n-1})}.$$

□

Proof of Proposition 2.2. Let A and A^* be defined as in Lemma 2.1. Let $g(\omega) = U^{-1/q}(\omega)$. Clearly, $g \in L^{q'}(\mathbb{S}^{n-1})$, and by (5.1)

$$A^*g(x) = \int_{\mathbb{S}^{n-1}} e^{i\omega x} d\sigma(\omega) = \omega_{n-1} j_{n/2-1}(|x|)$$

(see e.g. [16], Appendix B4). From (5.2) it follows that

$$(5.3) \quad \int_{\mathbb{R}^n} v^{1-p'}(x) |j_{n/2-1}(|x|)|^{p'} dx \leq C \left(\int_{\mathbb{S}^{n-1}} U^{1-q'}(\omega) d\sigma(\omega) \right)^{p'/q'}$$

as required. □

Proof of Corollary 2.3. Let $q_k = q_{\nu,k}$ be the positive zeros of the Bessel function $J_\nu(t)$ in nondecreasing order, $k \in \mathbb{N}$, $\nu = n/2 - 1 \geq -1/2$. It is known (see [30], p. 498) that

$$j_\nu(t) = \Gamma(\nu + 1)(t/2)^{-\nu} J_\nu(t) = \prod_{k=1}^{\infty} (1 - t^2/q_k^2),$$

and $|j_\nu(t)| \leq j_\nu(0) = 1$. Moreover,

$$J_\nu(t) = (\pi t/2)^{-1/2} (\cos(t - \nu\pi/2 - \pi/4) + O(t^{-1})), \quad t \rightarrow +\infty,$$

and

$$q_k = \pi(k - 1/2) + \nu\pi/2 + \pi/4 + O(k^{-1}), \quad k \rightarrow \infty$$

(see, e.g., [30], pp. 199, 479, 506). In particular, this gives $|j_\nu(t)| \leq C(1+t)^{-\nu-1/2}$, $t \geq 0$, and $0 < c_\nu \leq q_{k+1} - q_k \leq C_\nu$, $k \geq 0$, where $q_0 = 0$.

Let us now prove the estimate of $|j_\nu(t)|$ from below. Let $q'_k = (q_k + q_{k+1})/2$, $0 < \varepsilon < \min(\pi/2, c_\nu/2)$, $a_0 = 0$, $b_0 = q'_0 + \varepsilon$, $a_k = q'_k - \varepsilon$, $b_k = q'_k + \varepsilon$, $k \geq 1$, and $I_k = [a_k, b_k]$, $k \geq 0$. Then $I_0 \subset [0, q_1]$ and $I_k \subset (q_k, q_{k+1})$, $|I_k| = 2\varepsilon$, $k \geq 1$.

Since for $t = q'_k + u$, $u \in [-\varepsilon, \varepsilon]$, and $k \rightarrow \infty$, we have

$$|\cos(t - \nu\pi/2 - \pi/4)| = |\cos(\pi k + u + O(k^{-1}))| \geq \cos \varepsilon + O(k^{-1}),$$

then $|j_\nu(t)| \geq Ct^{-\nu-1/2}$, $t \in I_k$, for sufficiently large k ($k > k_0$).

For $k \leq k_0$, we have $|j_\nu(t)| \geq \min(|j_\nu(a_k)|, |j_\nu(b_k)|)$, $t \in I_k$. Therefore, the following inequality holds:

$$(5.4) \quad |j_\nu(t)| \geq C(1+t)^{-\nu-1/2}, \quad t \in I := \bigcup_{k=0}^{\infty} I_k.$$

Denote $I'_k = (b_{k-1}, a_k)$, $k \geq 1$, and $I' := \cup_{k=0}^{\infty} I'_k$. We have $|I'_k| = (q_{k+1} - q_{k-1})/2 - 2\varepsilon \leq C_\nu - 2\varepsilon$ and $I \cup I' = \mathbb{R}_+$.

Inequalities (5.3) and (5.4) give

$$\int_{|x| \in I} v^{1-p'}(x) (1 + |x|)^{-p'(n-1)/2} dx < C.$$

Furthermore,

$$\begin{aligned}
 J &:= \omega_{n-1}^{-1} \int_{\mathbb{R}^n} v^{1-p'}(x) (1 + |x|)^{-p'(n-1)/2} dx \\
 &= \int_0^\infty v_0^{1-p'}(t) (1+t)^{-p'(n-1)/2} t^{n-1} dt = \int_{I_0} + \sum_{k=1}^\infty \left(\int_{I_k} + \int_{I'_k} \right).
 \end{aligned}$$

Assume that condition (2.5) holds. Then it is clear that

$$\int_{I'_k} v_0^{1-p'}(t) dt \leq C \int_{I_k} v_0^{1-p'}(t) dt$$

with some constant C . Using this, we get

$$\begin{aligned}
 \int_{I'_k} v_0^{1-p'}(t) (1+t)^{-p'(n-1)/2} t^{n-1} dt &\leq C(1 + b_{k-1})^{-p'(n-1)/2} a_k^{n-1} \int_{I'_k} v_0^{1-p'}(t) dt \\
 &\leq C(1 + b_k)^{-p'(n-1)/2} \int_{I_k} v_0^{1-p'}(t) t^{n-1} dt \leq C \int_{I_k} v_0^{1-p'}(t) (1+t)^{-p'(n-1)/2} t^{n-1} dt,
 \end{aligned}$$

since $b_k = b_{k-1} + |I'_k| + |I_k| \leq b_{k-1} + c \leq Cb_{k-1}$.

Thus,

$$\begin{aligned}
 J &= \int_{I_0} + \sum_{k=1}^\infty \left(\int_{I_k} + \int_{I'_k} \right) v_0^{1-p'}(t) (1+t)^{-p'(n-1)/2} t^{n-1} dt \\
 &\leq C \sum_{k=0}^\infty \int_{I_k} v_0^{1-p'}(t) (1+t)^{-p'(n-1)/2} t^{n-1} dt \\
 &\leq C \int_{|x| \in I} v^{1-p'}(x) (1 + |x|)^{-p'(n-1)/2} dx < C.
 \end{aligned}$$

If the condition (2.6) is satisfied, the proof is similar. □

We prove Lemma 2.5 to make the paper self-contained.

Proof of Lemma 2.5. Assume $s > p$, since the proof in the other case is similar. Let $r = p/(s - p)$. Suppose that $T: L^p(d\mu) \rightarrow L^q(d\nu)$ is bounded. To show that $T: L^s(wd\mu) \rightarrow L^q(d\nu)$ is bounded, we observe that $1/(rs) = (s - p)/(sp) = 1/p - 1/s$. By Hölder’s inequality,

$$\begin{aligned}
 \|Tf\|_{L^q(d\mu)} &\leq C \|f\|_{L^p(d\mu)} = C \|w^{-1/s} w^{1/s} f\|_{L^p(d\mu)} \\
 &\leq C \|w^{-1/s}\|_{L^{rs}(d\mu)} \|w^{1/s} f\|_{L^s(d\mu)} = C \|w^{-1}\|_{L^r(d\mu)}^{1/s} \|w^{1/s} f\|_{L^s(d\mu)},
 \end{aligned}$$

as required.

To prove the other direction we argue as in [8] and as in the proof of Proposition 1.10 in [5]. Observe that

$$\|w^{1/s} f\|_{L^s(d\mu)}^s = \int_{\mathbb{R}^n} w |f(x)|^s d\mu(x) = \int_{\mathbb{R}^n} |f(x)|^p d\mu(x),$$

with $w = |f|^{p-s}$. Since

$$\|w^{-1}\|_{L^r(d\mu)}^{1/s} = \left(\int_{\mathbb{R}^n} |f(x)|^p d\mu(x) \right)^{(s-p)/(sp)} = \|f\|_{L^p(d\mu)}^{1-p/s},$$

we obtain

$$\|Tf\|_{L^q(d\nu)} \leq C \|w^{-1}\|_{L^r(d\mu)}^{1/s} \|f\|_{L^p(d\mu)}^{p/s} = C \|f\|_{L^p(d\mu)}^{1-p/s} \|f\|_{L^p(d\mu)}^{p/s} = C \|f\|_{L^p(d\mu)}. \quad \square$$

Proof of Theorem 3.1. Fix $\rho > 0$ and $f \in C_0^\infty(\mathbb{R}^n)$; let $\delta_\rho\psi(x) = \psi(\rho x)$, and let $g(x) = \rho^{-n}\delta_{\frac{1}{\rho}}f(x)$. We apply (1.9) with g in place of f . Recalling that $\rho^{-n}\widehat{\delta_{1/\rho}f} = \delta_\rho\widehat{f}$, we obtain, by Lemma 2.5 and (1.9) with $d\nu = U d\omega$ and $d\mu = v dx$,

$$\begin{aligned} \left(\int_{\mathbb{S}^{n-1}} |\delta_\rho\widehat{f}(\omega)|^q U(\omega) d\sigma(\omega) \right)^{1/q} &= \left(\int_{\mathbb{S}^{n-1}} |\widehat{g}(\omega)|^q U(\omega) d\sigma(\omega) \right)^{1/q} \\ &\leq C \|g\|_{L^p(v dx)} = C\rho^{-n} \|v^{1/p}\delta_{1/\rho}f\|_p = C\rho^{-n+n/p} \|(\delta_\rho v)^{1/p}f\|_p. \end{aligned}$$

By our assumptions on v we obtain

$$(5.5) \quad \int_{\mathbb{S}^{n-1}} |\widehat{f}(\rho\omega)|^q U(\omega) d\sigma(\omega) \leq C\rho^{-nq/p'} w^{q/p}(\rho) \|v^{1/p}f\|_p^q.$$

We multiply both sides of this inequality by $u_0(\rho)\rho^{n-1}$ and we integrate with respect to ρ . We obtain

$$\begin{aligned} \int_0^\infty \rho^{n-1} \int_{\mathbb{S}^{n-1}} |\widehat{f}(\rho\omega)|^q u_0(\rho) U(\omega) d\sigma(\omega) d\rho \\ \leq C \int_0^\infty \rho^{n-1-nq/p'} u_0(\rho) w^{q/p}(\rho) d\rho \|v^{1/p}f\|_p^q, \end{aligned}$$

which by (3.1) implies $\int_{\mathbb{R}^n} U(x/|x|)u(x)|\widehat{f}(x)|^q dx \leq C\|v^{1/p}f\|_p^q$. □

Proof of Corollary 3.4. When $p = q = 2$, we use Theorem 2.7. The assumptions of Theorem 3.1 are satisfied, and so the following inequality holds:

$$(5.6) \quad \|\widehat{f}\|_{L^2(u dy)} \leq \|f\|_{L^2(V dx)}.$$

To conclude the proof of Corollary 3.4 we use a special case of an interpolation theorem with change of measure proved in [26].

Lemma 5.1. *Let Tf be a linear operator defined in a space of measurable functions that include $L^{p_1}(V_1 dx)$ and $L^{p_2}(V_2 dx)$; assume that*

$$\|Tf\|_{L^{q_1}(u_1 dy)} \leq C \|f\|_{L^{p_1}(V_1 dx)} \quad \text{and} \quad \|Tf\|_{L^{q_2}(u_2 dy)} \leq C \|f\|_{L^{p_2}(V_2 dx)}.$$

Then, for every $0 \leq t \leq 1$,

$$(5.7) \quad \|Tf\|_{L^{q_t}(u_1^t u_2^{1-t} dy)} \leq C \|f\|_{L^{p_t}(V_1^t V_2^{1-t} dx)},$$

where $1/p_t = t/p_1 + (1-t)/p_2$ and $1/q_t = t/q_1 + (1-t)/q_2$.

We apply Lemma 5.1 with $Tf = \widehat{f}$; we interpolate the inequality (5.6) and the $\|\widehat{f}\|_\infty \leq \|f\|_1$; we let $u = u_1$ and $V = V_1$, and $u_2 = V_2 = 1$; we let $1/p_t = t/2 + 1 - t = 1 - t/2$, so that $t = 2(1 - 1/p_t) = 2/p'_t$. Note that $q_t = p'_t$. By (5.7), we have

$$\|\widehat{f}\|_{L^{p'}(u^{2/p'} dy)} \leq \|f\|_{L^p(V^{2/p'} dx)},$$

where we have let $p = p_t$ for simplicity. That concludes the proof of Corollary 3.4. \square

6. Applications to the uncertainty principle

The uncertainty principle is a cornerstone in quantum physics and in Fourier analysis. The simplest formulation of the uncertainty principle in harmonic analysis is *Heisenberg's inequality*, which applies to functions in $L^2(\mathbb{R}^n)$ of norm = 1. It states that the product of the variances of f and \widehat{f} is bounded below by a universal constant, i.e.,

$$\inf_{P \in \mathbb{R}^n} \int_{\mathbb{R}^n} |x - P|^2 |f(x)|^2 dx \inf_{Q \in \mathbb{R}^n} \int_{\mathbb{R}^n} |\xi - Q|^2 |\widehat{f}(\xi)|^2 d\xi \geq \frac{(2\pi)^n n^2}{4}.$$

One of the many consequences of this inequality is that a nonzero function and its Fourier transform cannot both be compactly supported.

The uncertainty principle for L^p functions is also interesting. Inequalities in the form of $\|f\|_2^2 \leq C \|v^{1/p} f\|_p \|w^{1/q} \widehat{f}\|_q$, where v and w are suitable weight functions and $1 \leq p, q \leq \infty$ are discussed in [10]. Power weights are of particular interest: using a standard homogeneity argument, is easy to prove that a necessary condition for the inequality $\|f\|_2^2 \leq C \| |x|^a f \|_p \| |\xi|^b \widehat{f} \|_q$ to hold for all $f \in C_0^\infty(\mathbb{R}^n)$ is that $a + n/p = b + n/q$. See also [14] for a survey on the uncertainty principle.

We prove the following.

Theorem 6.1. *Let u, v be weights for which the Pitt inequality (1.1) holds for some $1 \leq p, q \leq \infty$. Then, for every $f \in C_0^\infty(\mathbb{R}^n)$,*

$$\|f\|_2^2 \leq C \| |u|^{-1/q} |\xi| \widehat{f} \|_{q'} \| |v|^{1/p} |x| f \|_p,$$

where C is independent of f .

Corollary 6.2. *Let $1 \leq p < \frac{2(n+2)}{n+4}$ and $1 \leq q \leq \frac{n-1}{n+1} p'$. Let $s(x) = s_0(|x|)$ be a radial weight that satisfies*

$$(6.1) \quad \int_0^\infty \frac{\rho^{n-1-qn/p'}}{s_0(\rho)} d\rho < \infty.$$

Then,

$$(6.2) \quad \|f\|_2^2 \leq C \| |s_0|^{1/q} (|\xi|) |\xi| \widehat{f} \|_{q'} \| |x| f \|_p, \quad f \in C_0^\infty(\mathbb{R}^n).$$

For example, $s_0(\rho) = \rho^{-m}(1+\rho)^{m+n-nq/p'+\varepsilon}$, with $\varepsilon > 0$, and $m+n-nq/p' > 0$, satisfies (6.1).

Corollary 6.3. *Let*

$$v(x) = \begin{cases} |x|^\alpha, & |x| \leq 1, \\ |x|^\beta, & |x| > 1, \end{cases} \quad \text{and} \quad w_0(\rho) = \max\{\rho^\alpha, \rho^\beta\},$$

Let $1 < p \leq 2$, $2 \leq q \leq \frac{n-1}{n+1} p'$, $\alpha < n(p-1)$, and $\beta \geq 0$. We have

$$(6.3) \quad \|f\|_2^2 \leq C \|s_0^{1/q}(|\xi|) |\widehat{f}| \|_{q'} \| |x| v^{1/p} f \|_p, \quad f \in C_0^\infty(\mathbb{R}^n).$$

provided

$$(6.4) \quad \int_0^\infty \frac{\rho^{n-1-qn/p'} w_0^{q/p}(\rho)}{s_0(\rho)} d\rho < \infty.$$

When $\alpha < n$ and $\beta > 1$, we have

$$(6.5) \quad \|f\|_2^2 \leq C \|s_0^{1/2}(|\xi|) |\widehat{f}| \|_2 \| |x| v^{1/2} f \|_2, \quad f \in C_0^\infty(\mathbb{R}^n),$$

provided

$$\int_0^\infty \frac{w_0(\rho)}{\rho s_0(\rho)} d\rho < \infty.$$

Proof of Theorem 6.1. We use the same idea of the proof of the L^2 Heisenberg principle (see [14]). Let $f \in C_0^\infty(\mathbb{R}^n)$. We denote $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ by (x_1, x') , with $x' \in \mathbb{R}^{n-1}$. We integrate by parts the function $|f(x)|^2 = |f(x_1, x')|^2$ with respect to x_1 . That is,

$$\int_{-\infty}^\infty |f(x_1, x')|^2 dx_1 = x_1 |f(x_1, x')|^2 \Big|_{x_1=-\infty}^\infty - \int_{-\infty}^\infty x_1 \frac{\partial |f(x_1, x')|^2}{\partial x_1} dx_1.$$

A simple calculation shows that

$$\frac{\partial |f(x_1, x')|^2}{\partial x_1} = \frac{\partial}{\partial x_1} (f(x_1, x') \overline{f(x_1, x')}) = 2 \operatorname{Re} \left(\overline{f(x_1, x')} \frac{\partial f(x_1, x')}{\partial x_1} \right).$$

We obtain

$$\int_{-\infty}^\infty |f(x_1, x')|^2 dx_1 = -2 \operatorname{Re} \int_{-\infty}^\infty x_1 \overline{f(x_1, x')} \frac{\partial f(x_1, x')}{\partial x_1} dx_1.$$

We integrate the above identity in x' , to obtain

$$\|f\|_2^2 = -2 \operatorname{Re} \int_{\mathbb{R}^n} x_1 \overline{f(x)} \frac{\partial f(x)}{\partial x_1} dx.$$

We use the identity $\int_{\mathbb{R}^n} f_1 \overline{f_2} dx = (2\pi)^{-n} \int_{\mathbb{R}^n} \widehat{f_1} \overline{\widehat{f_2}} d\xi$, and we recall that the Fourier transform of $\partial f(x)/\partial x_1$ is $-i\xi_1 \widehat{f}(\xi)$. Thus,

$$\begin{aligned} \|f\|_2^2 &= 2(2\pi)^{-n} \operatorname{Re} \left(i \int_{\mathbb{R}^n} \xi_1 \widehat{f}(\xi) \overline{(\widehat{x_1 f})(\xi)} d\xi \right) \\ &= 2(2\pi)^{-n} \operatorname{Re} \left(i \int_{\mathbb{R}^n} (u^{-1/q} \xi_1 \widehat{f}(\xi)) \overline{(u^{1/q} (\widehat{x_1 f})(\xi))} d\xi \right), \end{aligned}$$

and by the Hölder inequality and Theorem 3.1,

$$\begin{aligned} \|f\|_2^2 &\leq C \|u^{-1/q} \widehat{\xi_1 f}\|_{q'} \|u^{1/q} \widehat{x_1 f}\|_q \leq C \|u^{-1/q} \widehat{\xi_1 f}\|_{q'} \|v^{1/p} x_1 f\|_p \\ &\leq C \|u^{-1/q} |\xi| \widehat{f}\|_{q'} \|v^{1/p} |x| f\|_p \end{aligned}$$

as required. □

Proof of Corollary 6.2. Follows from Theorems 1.3 and 6.1, with $v \equiv 1$ and $u_0(\rho) = s_0^{-1}(\rho)$. □

Proof of Corollary 6.3. Follows from Corollary 3.2 and Theorem 6.1, with $u_0(\rho) = s_0^{-1}(\rho)$. □

7. Riemann–Lebesgue estimates via Pitt inequalities

Here we investigate the interrelation between the smoothness of a function and the growth properties of the Fourier transforms. The original result goes back to the Riemann–Lebesgue estimate $|\widehat{f}(\xi)| \rightarrow 0$ as $|\xi| \rightarrow \infty$, where $f \in L^1(\mathbb{R}^n)$, and its quantitative version given by

$$(7.1) \quad |\widehat{f}(\xi)| \leq C \omega_l\left(f, \frac{1}{|\xi|}\right)_1, \quad f \in L^1(\mathbb{R}^n),$$

where the modulus of smoothness $\omega_l(f, \delta)_p$ of a function $f \in L^p(X)$ is defined by

$$(7.2) \quad \omega_l(f, \delta)_p = \sup_{|h| \leq \delta} \|\Delta_h^l f(x)\|_{L^p(\mathbb{R}^n)}, \quad 1 \leq p \leq \infty,$$

and

$$\Delta_h^l f(x) = \Delta_h^{l-1}(\Delta_h f(x)), \quad \Delta_h f(x) = f(x+h) - f(x).$$

Recently this result was extended for L^p -functions [7], [12], [15]. Let us first define the suitable multivariate substitution for the classical modulus of smoothness.

For a locally integrable function f the average on a sphere in \mathbb{R}^n of radius $t > 0$ is given by

$$V_t f(x) := \frac{1}{m_t} \int_{|y-x|=t} f(y) dy, \quad \text{with } V_t 1 = 1, \quad n \geq 2.$$

For $l \in \mathbb{N}$ we define

$$V_{l,t} f(x) := \frac{-2}{\binom{2l}{l}} \sum_{j=1}^l (-1)^j \binom{2l}{l-j} V_{j,t} f(x).$$

and set

$$\Omega_l(f, t)_p = \|f - V_{l,t} f\|_p.$$

In Theorem 2.1 (A), $n \geq 2$ of [15], the following Riemann–Lebesgue type estimate was proved.

Theorem 7.1. *Let $f \in L^p(\mathbb{R}^n)$, $1 < p \leq 2$. Then for $p \leq q \leq p'$ we have $|\xi|^{n(1-1/p-1/q)} \widehat{f}(\xi) \in L^q(\mathbb{R}^n)$, and*

$$\left(\int_{\mathbb{R}^n} [\min(1, t|\xi|)^{2l} |\xi|^{n(1-1/p-1/q)} |\widehat{f}(\xi)|]^q d\xi \right)^{1/q} \leq C \Omega_l(f, t)_p.$$

Note that some partial cases were previously proved in [7], [12]; see also [6]. The essential step in the proof of Theorem 7.1 is the use of Pitt inequalities (1.6) under conditions (1.7) and (1.8) in the case when $b = 0$, that is when the right-hand side of (1.6) is the non-weighted L^p -norm.

Here we refine Theorem 7.1 using inequality (1.11) (see Theorem 1.3).

Theorem 7.2. *Under the assumptions of Theorem 1.3, we have*

$$\left(\int_{\mathbb{R}^n} [\min(1, t|\xi|)^{2l} |\widehat{f}(\xi)|]^q u(\xi) d\xi \right)^{1/q} \leq C \Omega_l(f, t)_p.$$

The proof repeats the proof of Theorem 7.1 with the only modification that one should use the weight $u^{1/q}(\xi)$ in place of $|\xi|^{n(1-1/p-1/q)}$ (see [15], (2.16)) and Theorem 1.3.

8. Other applications

Inequality (1.10) in Theorem 1.3 implies $\int_{\mathbb{R}^n} u(\xi)(1 + |\xi|)^{-qn/p'} d\xi < \infty$. In [3] it is proved that if (1.1) holds for $1 < p \leq q < \infty$, and if $\int_{\mathbb{R}^n} u(\xi)^{1-q'}(1 + |\xi|)^{-M} d\xi < \infty$ for some $M > 0$, then one can prove a Bernstein-type theorem, which characterizes the Fourier transform on weighted Besov spaces. We leave the generalization of the main theorem in [3] to the interested reader.

References

- [1] BARCELÓ, J. A., BENNETT, J. M., CARBERY, A., RUIZ, A. AND VILELA, M. C.: A note on weighted estimates for the Schrödinger operator. *Rev. Mat. Complut.* **21** (2008), no. 2, 481–488.
- [2] BENEDETTO, J. J. AND HENIG H. P.: Weighted Fourier inequalities: new proofs and generalizations. *J. Fourier Anal. Appl.* **9** (2003), no. 1, 1–37.
- [3] BUI, H.-Q.: Bernstein's theorem on weighted Besov spaces. *Forum Math.* **9** (1997), no. 6, 739–750.
- [4] BERNDT, R.: Symmetric conditions for a weighted Fourier transform inequality. *J. Math. Anal. Appl.* **379** (2011), no. 1, 439–443.
- [5] BLOOM, S. AND SAMPSON, G.: Weighted spherical restriction theorem for the Fourier transform. *Illinois J. Math.* **36** (1992), no. 1, 73–101.
- [6] BRAY, W. O.: Growth and integrability of Fourier transforms on Euclidean space. *J. Fourier Anal. Appl.* **20** (2014), no. 6, 1234–1256.

- [7] BRAY, W. O. AND PINSKY, M. A.: Growth properties of Fourier transforms via moduli of continuity. *J. Funct. Anal.* **255** (2008), no. 9, 2265–2285.
- [8] CHANILLO, S. AND SAWYER, E.: Unique continuation for $\Delta+v$ and the C. Fefferman–Phong class. *Trans. Amer. Math. Soc.* **318** (1990), no. 1, 275–300.
- [9] CHIARENZA, F. AND RUIZ, A.: Uniform L_2 -weighted Sobolev inequalities. *Proc. Amer. Math. Soc.* **112** (1991), no. 1, 53–64.
- [10] COWLING, M. G. AND PRICE, J. F.: Bandwidth versus time concentration: the Heisenberg–Pauli–Weyl inequality. *SIAM J. Math. Anal.* **15** (1984) no. 1, 151–165.
- [11] DE CARLI, L., GORBACHEV, D. AND TIKHONOV, S.: Pitt and Boas inequalities for Fourier and Hankel transforms. *J. Math. Anal. Appl.* **408** (2013), no. 2, 762–774.
- [12] DITZIAN, Z.: Smoothness of a function and the growth of its Fourier transform or its Fourier coefficients. *J. Approx. Theory* **162** (2010), no. 5, 980–986.
- [13] FEFFERMAN, C.: Inequalities for strongly singular convolution operators. *Acta Math.* **124** (1970), 9–36.
- [14] FOLLAND, G. B. AND SITARAM, A.: The uncertainty principle: a mathematical survey. *J. Fourier Anal. Appl.* **3** (1997), no. 3, 207–238.
- [15] GORBACHEV, D. AND TIKHONOV, S.: Moduli of smoothness and growth properties of Fourier transforms: two-sided estimates. *J. Approx. Theory* **164** (2012), no. 9, 1283–1312.
- [16] GRAFAKOS, L.: *Classical and modern Fourier analysis*. Pearson/Prentice Hall, NJ, 2004.
- [17] HEINIG, H.: Weighted norm inequalities for classes of operators. *Indiana Univ. Math. J.* **33** (1984), no. 4, 573–582.
- [18] JURKAT, W. B. AND SAMPSON, G.: On rearrangement and weight inequalities for the Fourier transform. *Indiana Univ. Math. J.* **33** (1984), no. 2, 257–270.
- [19] LAKEY, J. D.: Weighted Fourier transform inequalities via mixed norm Hausdorff–Young inequalities. *Canad. J. Math.* **46** (1994), no. 3, 586–601.
- [20] MUCKENHOUPT, B.: A note on two weight function conditions for a Fourier transform norm inequality. *Proc. Am. Math. Soc.* **88** (1983), no. 1, 97–100.
- [21] MUCKENHOUPT, B.: Weighted norm inequalities for the Fourier transform. *Trans. Amer. Math. Soc.* **276** (1983), no. 2, 729–142.
- [22] PITT, H. R.: Theorems on Fourier series and power series. *Duke Math. J.* **3** (1937), no. 4, 747–755.
- [23] RUIZ, A. AND VEGA, L.: Unique continuation for Schrödinger operators with potential in Morrey spaces. *Publ. Mat.* **35** (1991), no. 1, 291–298.
- [24] STEIN, E. M.: Interpolation of linear operators. *Trans. Amer. Math. Soc.* **83** (1956), 482–492.
- [25] STEIN, E. M.: *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*. Princeton Mathematical Series 43, Monographs in Harmonic Analysis III, Princeton University Press, Princeton, NJ, 1993.
- [26] STEIN, E. M. AND WEISS, G.: Interpolation of operators with change of measures. *Trans. Amer. Math. Soc.* **87** (1958), 159–172.
- [27] STEIN, E. M. AND WEISS, G.: *Introduction to Fourier analysis on Euclidean spaces*. Princeton Mathematical Series 32, Princeton University Press, Princeton, NJ, 1971.

- [28] TAO, T.: A sharp bilinear restrictions estimate for paraboloids. *Geom. Funct. Anal.* **13** (2003), no. 6, 1359–1384.
- [29] TOMAS, P. A.: A restriction theorem for the Fourier transform. *Bull. Amer. Math. Soc.* **81** (1975), 477–478.
- [30] WATSON, G. N.: *A treatise on the theory of Bessel functions*. Reprint of the second (1944) edition. Cambridge University Press, Cambridge, 1995.

Received May 29, 2015; revised November 10, 2015.

LAURA DE CARLI: Department of Mathematics, Florida International University, Modesto Maidique Campus, Miami, FL 33199, USA.

E-mail: decarlil@fiu.edu

DMITRY GORBACHEV: Department of Applied Mathematics and Computer Science, Tula State University, 300012 Tula, Russia.

E-mail: dvgmail@mail.ru

SERGEY TIKHONOV: Centre de Recerca Matemàtica, and Universitat Autònoma de Barcelona, Campus de Bellaterra, Edifici C, 08193 Bellaterra (Barcelona), Spain; *and* ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain.

E-mail: stikhonov@crm.cat

D. G. was supported by the RFBR (no. 13-01-00045), the Ministry of Education and Science of the Russian Federation (no. 5414GZ), and by D. Zimin's Dynasty Foundation. S. T. was partially supported by MTM 2014-59174-P, 2014 SGR 289, and by the Alexander von Humboldt Foundation.