



A Nash–Kuiper theorem for $C^{1,1/5-\delta}$ immersions of surfaces in 3 dimensions

Camillo De Lellis, Dominik Inauen and László Székelyhidi Jr.

Abstract. We prove that, given a C^2 Riemannian metric g on the 2-dimensional disk D_2 , any short C^1 immersion of (D_2, g) into \mathbb{R}^3 can be uniformly approximated with $C^{1,\alpha}$ isometric immersions for any $\alpha < 1/5$. This statement improves previous results by Yu. F. Borisov and of a joint paper of the first and third author with S. Conti.

1. Introduction

In this paper we consider isometric immersions of 2-dimensional disks in \mathbb{R}^3 . With $D_r(x_0)$ and $\bar{D}_r(x_0)$ we denote, respectively, the open and closed disks in \mathbb{R}^2 with center x_0 and radius r . When $x_0 = 0$ we write simply D_r , resp. \bar{D}_r . If g is a C^0 Riemannian metric on $\bar{D}_r(x_0)$, an isometric immersion $u: \bar{D}_r(x_0) \rightarrow \mathbb{R}^n$ is a C^1 immersion such that $u^\sharp e = g$, where e denotes the Euclidean metric on \mathbb{R}^n . In other words this means that

$$(1.1) \quad \partial_i u \cdot \partial_j u = g_{ij}.$$

If $0 < \partial_i u \cdot \partial_j u < g_{ij}$ in the sense of quadratic forms, we then call u a short immersion. Our main theorem is the following approximation result which, using a popular terminology, is an “ h -principle” statement, cf. [19], [29], [13].

Theorem 1.1. *Let g be a C^2 metric on \bar{D}_2 and $\bar{u} \in C^1(\bar{D}_2, \mathbb{R}^3)$ a short immersion. For every $\delta > 0$ and $\varepsilon > 0$ there is a $C^{1,1/5-\delta}$ isometric immersion u of (\bar{D}_1, g) in \mathbb{R}^3 such that $\|\bar{u} - u\|_{C^0} < \varepsilon$.*

The well-known ground-breaking result of Nash and Kuiper [23], [22] implies that Theorem 1.1 holds with C^1 replacing $C^{1,1/5-\delta}$. The first extension to the $C^{1,\alpha}$ category was obtained by Yu. F. Borisov: in [3] a version of Theorem 1.1 for $C^{1,1/7-\delta}$ immersions (and embeddings) of 2-dimensional disks with real analytic metrics g

Mathematics Subject Classification (2010): Primary 57N35,58D10; Secondary 53C21.

Keywords: Isometric embedding, convex integration, Nash–Kuiper.

was announced; in fact, more generally, the theorem in [3] applied to $C^{1,\alpha}$ isometric embeddings of n -dimensional balls in \mathbb{R}^{n+1} under the assumption that $\alpha < 1/(1+n+n^2)$. In [4] a detailed proof for the case of 2-dimensional disks and with exponents $\alpha < 1/13$ appeared. In the paper [11] the first and third author jointly with S. Conti gave a detailed and self-contained proof of all the statements contained in [3] for C^2 metrics on n -dimensional balls (with the same Hölder exponents) and analogous generalizations to $C^{1,\beta}$ metrics on compact manifolds without topological restrictions. For the optimal Hölder exponents in the case of rough metrics, which depend on β , the dimension and the topology of the manifold, we refer to [11].

The main contribution of this paper is to be able to raise the optimal Hölder exponent for 2-dimensional disks from $1/7$ to $1/5$, by taking advantage of the theory of conformal maps. The question of the optimal exponent for which an h -principle statement as in Theorem 1.1 can hold is also relevant for rigidity theory, as we will explain below.

It is known that Theorem 1.1 cannot hold for $C^{1,\alpha}$ immersions u when $\alpha > 2/3$ and g has positive Gauss curvature: under these assumptions it was shown by Borisov that $u(\overline{D}_1)$ must be (a portion of) a convex surface. This was the outcome of a series of papers, cf. [1], [2], and an alternative shorter proof has been given in [11]. Borisov's theorem extends the classical rigidity result for the Weyl problem: if (S^2, g) is a compact Riemannian surface with positive Gauss curvature and $u \in C^2$ is an isometric immersion into \mathbb{R}^3 , then u is uniquely determined up to a rigid motion ([9], [20], see also [26] for a thorough discussion).

The technique used to prove approximation results as in Theorem 1.1 follows an iteration scheme called convex integration. The latter was developed by Gromov [18], [19] into a very powerful tool to prove h -principle statements in a wide variety of geometric problems (see also [16], [28]). In general the regularity of solutions obtained via convex integration agrees with the highest derivatives appearing in the equations (see [27]). Thus, an interesting question raised in [19] p. 219 is how one could extend the methods to produce more regular solutions. Essentially the same question, in the case of isometric embeddings, is also mentioned in [31], see Problem 27. In particular, it is tempting to imagine the existence of a threshold α_0 so that:

- the h -principle holds for isometric $C^{1,\alpha}$ immersions of 2-dimensional disks in \mathbb{R}^3 whenever $\alpha < \alpha_0$;
- rigidity holds for $C^{1,\alpha}$ immersions of positively curved 2-dimensional disks in \mathbb{R}^3 whenever $\alpha > \alpha_0$.

Hence a summary of our current knowledge is that, if such a threshold α_0 exists, then it must lie in the interval $[1/5, 2/3]$.

Starting with the work [12], the first and third author pointed out a surprising similarity between the latter question and a long-standing conjecture in the theory of turbulence: in [24] Onsager conjectured the existence of a threshold Hölder regularity discriminating the validity of the conservation of kinetic energy for weak solutions of the incompressible Euler equations. The “rigidity” part of Onsager's

conjecture was established by Eyink and Constantin, E and Titi in the papers [17] and [10]. The paper [14] gave the first proof of the existence of continuous solutions that violate the conservation of total kinetic energy. A series of subsequent works [15], [21], [7], [5], [6], [8] have made a quite substantial progress in settling Onsager’s conjecture.

Theorem 1.1 could be improved in several directions. In particular, with little additional technicalities, which we believe to be of secondary importance, we will also show the following

Theorem 1.2. *Let g be a C^2 metric on $\overline{D_1}$ and $\bar{u} \in C^1(\overline{D_1}, \mathbb{R}^3)$ a short immersion. For every $\delta > 0$ and $\varepsilon > 0$ there is a $C^{1,1/5-\delta}$ isometric immersion u of $(\overline{D_1}, g)$ in \mathbb{R}^3 such that $\|\bar{u} - u\|_{C^0} < \varepsilon$. If in addition \bar{u} is an embedding, then u can be chosen to be an embedding.*

2. Main iteration

Theorem 1.1 is achieved via an iteration, which depends upon several parameters. We start introducing the main ones. The first parameter $\alpha > 0$ is an exponent, which is assumed to be rather small, in fact smaller than a geometric constant:

$$(2.1) \quad 0 < \alpha < \alpha_0.$$

Two further exponents will be called c and b , both assumed to be larger than 1, and a basis a , assumed to be very large. We then define the parameters

$$(2.2) \quad \delta_q := a^{-b^q}, \quad \lambda_q := a^{cb^{q+1}},$$

where q is an arbitrary natural number. The parameter b can in fact be chosen rather close to 1: how much it is allowed to be close to 1 depends on how close is α to 0. The parameter c will be larger but rather close to 5/2, depending on how close are $b - 1$ and α to 0. More precisely, we summarize the conditions which b and c need to satisfy in the following two inequalities:

$$(2.3) \quad \frac{3}{2} > b > \frac{2}{(2 - \alpha)(1 - 2\alpha)}$$

$$(2.4) \quad c > \frac{2(2 - \alpha)b^2 - (3 - 2\alpha)b - 1}{b((2 - \alpha)(1 - 2\alpha)b - 2)} = \frac{((4 - 2\alpha)b + 1)(b - 1)}{b((2 - 5\alpha + 2\alpha^2)b - 2)}.$$

It is moreover convenient to introduce the notation

$$(2.5) \quad g_q := g - \delta_{q+1} e,$$

which simplifies several formulas.

Proposition 2.1. *Fix a metric g as in Theorem 1.1. There is a positive constant α_0 such that for every α as in (2.1) we can choose positive numbers $\sigma_0(\alpha) < 1$ and C_0 with the following property. Assume b and c satisfy (2.3) and (2.4), fix any*

$\bar{C} \geq C_0$ and assume that λ_q and δ_q are defined as in (2.2), where a is sufficiently large depending on α, b, c, g, \bar{C} , namely

$$(2.6) \quad a > a_0(\alpha, b, c, g, \bar{C}).$$

If $q \in \mathbb{N}$ and $u_q : \bar{D}_{1+2^{-q-1}} \rightarrow \mathbb{R}^3$ is an immersion such that

$$(2.7) \quad \|g_q - u_q^\sharp e\|_\alpha \leq \sigma_0 \delta_{q+1}$$

$$(2.8) \quad \|D^2 u_q\|_0 \leq \bar{C} \delta_q^{1/2} \lambda_q,$$

then there is an immersion $u_{q+1} : \bar{D}_{1+2^{-q-2}} \rightarrow \mathbb{R}^3$ such that

$$(2.9) \quad \|g_{q+1} - u_{q+1}^\sharp e\|_0 \leq \frac{\sigma_0}{3} \delta_{q+2} \lambda_{q+1}^{-\alpha}$$

$$(2.10) \quad \|D(g_{q+1} - u_{q+1}^\sharp e)\|_0 \leq \frac{\sigma_0}{3} \delta_{q+2} \lambda_{q+1}^{1-\alpha}$$

$$(2.11) \quad \|u_q - u_{q+1}\|_0 \leq \delta_{q+1}^{1/2} \lambda_{q+1}^{-\gamma}$$

$$(2.12) \quad \|D(u_q - u_{q+1})\|_0 \leq C_0 \delta_{q+1}^{1/2}$$

$$(2.13) \quad \|D^2 u_{q+1}\|_0 \leq \bar{C} \delta_{q+1}^{1/2} \lambda_{q+1},$$

where $\gamma = \gamma(\alpha, b, c) > 0$.

As already mentioned, Proposition 2.1 will be used in an iteration scheme to show Theorem 1.1. The reader will notice that the starting assumption (2.7) does not exactly match the conclusions (2.9)–(2.10). On the other hand, a simple interpolation shows that (2.9) and (2.10) together imply the estimate

$$\|g_{q+1} - u_{q+1}^\sharp e\|_\alpha \leq \sigma_0 \delta_{q+2},$$

which corresponds to (2.7) at the next step of the iteration. It is possible to state a version of Proposition 2.1 where the assumptions and conclusions look more homogeneous, but there would be no real simplification neither in the statement nor in the proof.

Observe that, by our condition upon the parameters, u_q is obviously a short map, because we have

$$u_q^\sharp e \leq g_q + \sigma_0 \delta_{q+1} e = g - (1 - \sigma_0) \delta_{q+1} e < g,$$

where all the inequalities are understood in the sense of quadratic forms. Thus, as a simple corollary we know that

$$(2.14) \quad \|Du_q\|_{C^0} \leq C$$

for some constant C which only depends upon g .

As in the Nash–Kuiper classical theorem, the map u_{q+1} is obtained from the map u_q by adding a certain number of perturbations, each consisting of highly oscillatory functions. As it is clear from the arguments in [11], the threshold Hölder

exponent that can be reached by a Nash–Kuiper type iteration is $1/(1 + 2n_*)$, where n_* is the number of such perturbations. Each perturbation adds, modulo small error terms, a smooth symmetric rank-1 tensor, called “primitive metric”, to $u_q^\sharp e$. The constant n_* is then the smallest number of summands needed to write the metric error $g - u_q^\sharp e$ as a (positive) linear combination of such “primitive metrics”.

We know by the inductive assumption that $(g - u_q^\sharp e)/\|g - u_q^\sharp e\|_0$ is close to e , which implies that n_* can be chosen to be the dimension of the space of symmetric matrices. Thus, if n is the dimension of the manifold, $n_* = n(n + 1)/2$: this explains the threshold $1/(1 + 2n_*) = 1/(1 + n + n^2)$ reached in [11] and claimed originally by Borisov. In particular in dimension 2 the number n_* equals 3 and Borisov’s threshold is $1/7$.

The starting point of this paper is the simple observation that in 2 dimensions we can use a conformal change of coordinates to diagonalize $g - u^\sharp e$ and hence reduce the number n_* from 3 to 2: this justifies the new threshold $1/5$. However, the regularity of the change of coordinates needed to implement this idea deteriorates with q and thus it is not at all clear that the method really improves the regularity of the final map. In fact at first it is not even clear that the new iteration scheme yields any $C^{1,\alpha}$ regularity at all.

In order to overcome this difficulty we obviously need to estimate quite carefully several norms of the conformal change of coordinates, at each step: for this reason we need to keep track of some Hölder norm of $g - u_q^\sharp e$. However, to ensure convergence of the scheme, it does not seem enough to just combine the computations of [11] with the classical estimates on conformal mappings. In particular in order to close the argument we impose a much faster rate of convergence for $g - u_q^\sharp e$: in [11] it was sufficient to choose exponentially decaying δ_q (and exponentially growing λ_q), whereas in this note we take advantage of a double exponential Ansatz. This idea is in fact borrowed from [15], where a scheme with a double exponential decay was used to produce Hölder solutions to the Euler equations.

The rest of the paper is organized as follows.

Section 3 collects the technical preliminary lemmas and propositions which will be used in the proofs of Proposition 2.1 and Theorem 1.1.

The proof of Proposition 2.1 is split into the Sections 4, 5, 6 and 7. Section 4 describes how to reach u_{q+1} from u_q and in particular it gives the precise formulas for the two oscillatory perturbations which we need to add. We will then collect in Section 5 the estimates concerning the first perturbation and in Section 6 the ones concerning the second perturbation. Section 7 will finally conclude the proof of Proposition 2.1.

Section 8 will prove Theorem 1.1 using Proposition 2.1. In fact the proof is not completely straightforward since we have to show the existence of a map u_0 which is C^0 close to the map \bar{u} of Theorem 1.1 and at the same time satisfies the requirements of Proposition 2.1 (with $q = 0$), in order to be able to start the iterative procedure. In Section 9 we give briefly the necessary technical modifications to prove Theorem 1.2.

One key technical point is Proposition 3.4, which addresses rather well-known regularity properties of conformal changes of coordinates. However, it is crucial for us to have an explicit (linear) dependence of certain Hölder norms of the change of coordinates in terms of corresponding norms of the metric. Since we have not been able to find the relevant statements in the literature, we have included a proof of Proposition 3.4 in the Appendix.

3. Preliminaries

3.1. Hölder spaces

In the following, $m \in \mathbb{N}$, $\alpha \in (0, 1)$, and β is a multi-index. Moreover we will always assume that the domain of definition of any map is a disk $\overline{D}_r \subset \mathbb{R}^2$ with radius $r \in [1, 2]$. The maps f can be real-valued, vector-valued, matrix-valued or generally tensor-valued. In all these cases we endow the targets with the standard Euclidean norms, for which we will use the notation $|f(x)|$. We introduce the usual Hölder norms as follows. First of all, the supremum norm is denoted by $\|f\|_0 := \sup |f|$. We define the Hölder seminorms as

$$[f]_m = \max_{|\beta|=m} \|D^\beta f\|_0 \quad \text{and} \quad [f]_{m+\alpha} = \max_{|\beta|=m} \sup_{x \neq y} \frac{|D^\beta f(x) - D^\beta f(y)|}{|x - y|^\alpha}.$$

The Hölder norms are then given by

$$\|f\|_m = \sum_{j=0}^m [f]_j \quad \text{and} \quad \|f\|_{m+\alpha} = \|f\|_m + [f]_{m+\alpha}.$$

We then recall the standard “Leibniz rule” to estimate norms of products

$$(3.1) \quad [fg]_r \leq C([f]_r \|g\|_0 + \|f\|_0 [g]_r) \quad \text{for any } 1 \geq r \geq 0$$

and the usual interpolation inequalities

$$(3.2) \quad [f]_s \leq C \|f\|_0^{1-s/r} [f]_r^{s/r} \quad \text{for all } r \geq s \geq 0.$$

The following version of estimate (3.2), with explicit constant, will be useful at a certain stage:

$$(3.3) \quad \|f\|_\alpha \leq \|f\|_0 + 2\|f\|_0^{1-\alpha} \|Df\|_0^\alpha \quad \text{for all } 0 \leq \alpha \leq 1.$$

We also collect two classical estimates on the Hölder norms of compositions. These are also standard, for instance in applications of the Nash–Moser iteration technique.

Proposition 3.1. *Let $0 \leq \alpha < 1$, $\Psi : \Omega \rightarrow \mathbb{R}$ and $u : \mathbb{R}^n \supset U \rightarrow \Omega$ be two $C^{m,\alpha}$ functions, with $\Omega \subset \mathbb{R}^N$. Then there is a constant C (depending only on α, m, Ω and U) such that*

$$(3.4) \quad [\Psi \circ u]_{m+\alpha} \leq C[u]_{m+\alpha} ([\Psi]_1 + \|u\|_0^{m-1} [\Psi]_m) + C[\Psi]_{m+\alpha} (\|u\|_0^{m-1} [u]_m)^{\frac{m+\alpha}{m}},$$

$$(3.5) \quad [\Psi \circ u]_{m+\alpha} \leq C([u]_{m+\alpha} [\Psi]_1 + [u]_1^{m+\alpha} [\Psi]_{m+\alpha}).$$

Let $f, g : \mathbb{R}^n \supset U \rightarrow \mathbb{R}$ two $C^{m,\alpha}$ functions. Then there is a constant C (depending only on α, m, n and U) such that

$$(3.6) \quad [fg]_{m+\alpha} \leq C(\|f\|_0 [g]_{m+\alpha} + \|g\|_0 [f]_{m+\alpha}).$$

Proof. The chain rule can be written as

$$(3.7) \quad D^m(\Psi \circ u) = \sum_{i=1}^m (D^i \Psi \circ u) \sum_k C_{i,k} (Du)^{k_1} \dots (D^m u)^{k_m},$$

where $C_{i,k}$ are constants and $k = (k_1, \dots, k_m)$ is a multi-index with

$$\sum k_j = i, \quad \sum jk_j = m.$$

The claim then follows by the Leibniz rule (3.1) and a repeated application of the interpolation inequalities (3.2) to (3.7). Statement (3.6) is a straightforward consequence of the usual Leibniz rule, interpolation and the Young inequality. \square

Remark 3.2. Observe that if $\alpha = 0$ we have the estimates

$$(3.8) \quad [\Psi \circ u]_m \leq C[u]_m ([\Psi]_1 + \|u\|_0^{m-1} [\Psi]_m),$$

$$(3.9) \quad [\Psi \circ u]_m \leq C([u]_m [\Psi]_1 + [u]_1^m [\Psi]_m).$$

3.2. Quadratic mollification estimate

We will often use regularizations of maps f by convolution with a standard mollifier $\varphi_\ell(y) := \ell^{-2}\varphi(y/\ell)$, where $\varphi \in C_c^\infty(D_1)$ is assumed to have integral 1 and to be non negative and rotationally symmetric. Since however the domain of f will be D_r (resp. \overline{D}_r), we fix the convention that the convolution $f * \varphi_\ell$ is defined in $D_{r-\ell}$ (resp. $\overline{D}_{r-\ell}$).

Lemma 3.3. *For any $r, s \geq 0$ and $0 < \alpha \leq 1$ we have*

$$(3.10) \quad [f * \varphi_\ell]_{r+s} \leq C \ell^{-s} [f]_r,$$

$$(3.11) \quad [f - f * \varphi_\ell]_r \leq C \ell^2 [f]_{2+r},$$

$$(3.12) \quad \|f - f * \varphi_\ell\|_r \leq C \ell^{2-r} [f]_2, \quad \text{if } 0 \leq r \leq 2,$$

$$(3.13) \quad \|(fg) * \varphi_\ell - (f * \varphi_\ell)(g * \varphi_\ell)\|_r \leq C \ell^{2\alpha-r} \|f\|_\alpha \|g\|_\alpha,$$

where the constants C depend only upon s, r, α and φ .

Proof. Except for (3.12), the other estimates are contained in Lemma 1 of [11]. The additional claim (3.12) can be seen as follows. Recall the estimate

$$\|f - f * \varphi_\ell\|_0 \leq C \ell [f]_1,$$

which can be derived using the mean value theorem and an integration. We combine this estimate with (3.2) and (3.11) to get

$$\begin{aligned} [f - f * \varphi_\ell]_r &\leq C \|f - f * \varphi_\ell\|_0^{1-r} [f - f * \varphi_\ell]_1^r \\ &\leq C (\ell^2 \|D^2 f\|_0)^{1-r} (\ell \|D^2 f\|_0)^r \leq C \ell^{2-r} [f]_2, \end{aligned}$$

whenever $0 \leq r \leq 1$. If however $1 \leq r \leq 2$, we invoke the trivial inequality

$$[f - f * \varphi_\ell]_2 \leq C[f]_2$$

to deduce

$$[f - f * \varphi_\ell]_r \leq C \|\nabla f - \nabla f * \varphi_\ell\|_0^{2-r} [\nabla f - \nabla f * \varphi_\ell]_1^{r-1} \leq C \ell^{2-r} [f]_2,$$

from which the claim follows. □

3.3. Conformal coordinates

A crucial ingredient of our proof is the following proposition on the existence of conformal coordinates for $C^{N,\alpha}$ metrics. Although such existence is a very classical fact, we need an explicit dependence of the norms of the coordinates in terms of the regularity of the metric. Since we have not been able to find a precise reference in the literature, we include a proof in the appendix.

Proposition 3.4. *For any N, α, β with $N \in \mathbb{N}, N \geq 1, 0 < \beta \leq \alpha < 1$ there exist constants $C(N, \alpha, \beta), \sigma_1(N, \alpha, \beta) > 0$ and $\tilde{C}(\alpha)$ such that the following holds. If $1 \leq r \leq 2$ and g is a $C^{N,\alpha}$ metric on \overline{D}_r with*

$$(3.14) \quad \|g - e\|_\alpha \leq \sigma_1$$

then there exists a coordinate change $\Phi: \overline{D}_r \rightarrow \mathbb{R}^2$ and a function $\rho: \overline{D}_r \rightarrow \mathbb{R}^+$ satisfying

$$(3.15) \quad g = \rho^2 (\nabla\Phi_1 \otimes \nabla\Phi_1 + \nabla\Phi_2 \otimes \nabla\Phi_2)$$

and the following estimates:

$$(3.16) \quad \|\rho - 1\|_\alpha + \|D\Phi - \text{Id}\|_\alpha \leq \tilde{C} \|g - e\|_\alpha$$

$$(3.17) \quad \|D^k \rho\|_\beta + \|D^{k+1}\Phi\|_\beta \leq C \|g - e\|_{k+\beta} \quad \forall 1 \leq k \leq N.$$

3.4. Oscillatory functions

The construction of u_{q+1} is based on adding to the map u_q suitable “wrinkles”, namely suitable perturbations. The basic model for this perturbation takes advantage of a pair of real-valued functions with very specific properties, which we will detail here.

Proposition 3.5. *There exists $\delta_\star > 0$ and a function $\Gamma = (\Gamma^t, \Gamma^n) \in C^\infty([0, \delta_\star] \times \mathbb{R}, \mathbb{R}^2)$ with the following properties:*

- (a) $\Gamma(s, \xi) = \Gamma(s, \xi + 2\pi)$ for every s, ξ ;
- (b) $(1 + \partial_\xi \Gamma^t)^2 + (\partial_\xi \Gamma^n)^2 = 1 + s^2$;
- (c) *The following estimates hold:*

$$(3.18) \quad \|\partial_\xi^k \Gamma^n(s, \cdot)\|_0 \leq C(k) s$$

$$(3.19) \quad \|\partial_\xi^k \Gamma^t(s, \cdot)\|_0 \leq C(k) s^2$$

$$(3.20) \quad \|\partial_s \partial_\xi^k \Gamma^t(s, \cdot)\|_0 \leq C(k) s.$$

Proof. Except for (3.19), the remaining claims are contained in Lemma 2 of [11]. The idea is to let Γ have the form

$$\Gamma(s, \xi) := \int_0^\xi (\sqrt{1 + s^2} (\cos(f(s) \sin(\tau)), \sin(f(s) \sin(\tau))) - (1, 0)) d\tau,$$

for an appropriately chosen function f such that (a), (3.18) and (3.20) are fulfilled. (b) is satisfied by construction. The additional statement (3.19) follows from integrating (3.20) in s . □

4. Proof of Proposition 2.1, Part I

4.1. Hierarchy of parameters

A first ingredient in the construction of u_{q+1} is to smooth u_q suitably via a standard mollification. For this we introduce the mollification parameter ℓ , which is rather small: indeed it is defined by the relation

$$(4.1) \quad \ell^{2-\alpha} := \frac{1}{\tilde{C}} \frac{\delta_{q+1}}{\delta_q \lambda_q^2},$$

where \tilde{C} is a constant larger than 1 which depends only upon α, g, σ_0 and \tilde{C} and which will be specified in Section 4.3 below.

The map u_{q+1} will be obtained from (a suitable regularization of) the map u_q in two steps. First we will add an oscillatory perturbation whose frequency is

$$(4.2) \quad \mu := \hat{C} \frac{\delta_{q+1} \lambda_{q+1}^\alpha}{\delta_{q+2} \ell},$$

where the constant \hat{C} , larger than 1, depends only upon α, g , and σ_0 (we specify its choice in Section 7). We will then choose a second perturbation whose frequency is λ_{q+1} .

We next record a few inequalities among the parameters which will be rather useful in simplifying some of our estimates in the remaining sections. Except for the very first inequality in (4.4), which requires a choice of a sufficiently large compared to the constant \hat{C} , all the others are immediate from the restrictions imposed so far on all the various parameters:

$$(4.3) \quad \delta_q \lambda_q^2 \geq 1,$$

$$(4.4) \quad \lambda_{q+1} \geq \mu \geq \ell^{-1} \geq \lambda_q,$$

$$(4.5) \quad \delta_q^{1/2} \lambda_q \leq \delta_q^{1/2} \lambda_q \ell^{-\alpha/2} \leq \delta_{q+1}^{1/2} \ell^{-1} \leq \delta_{q+1}^{1/2} \mu \leq \delta_{q+1}^{1/2} \lambda_{q+1},$$

The first inequality (4.3) follows from $\delta_q \lambda_q^2 = a^{c^2 b^{2q+2} - b^q} \geq a^{b^2 - 1}$ (where we have used $c, b > 1$). Observe that this easily implies $\ell \leq 1$ (recall that δ_{q+2} and \tilde{C}^{-1} are

both smaller than 1), which in turn gives the first inequality in (4.5). Note also that the last inequality in (4.4) is weaker than the second inequality in (4.5):

$$\ell^{-1} \geq \ell^{-1+\alpha/2} \geq \frac{\delta_q^{1/2}}{\delta_{q+1}^{1/2}} \lambda_q \geq \lambda_q.$$

Coming to the second inequality in (4.5), observe that, by the definition of ℓ , this is just the requirement that $\tilde{C} \geq 1$. As for the last two inequalities in (4.5) are equivalent to the first two in (4.4), which will be shown below. Moreover, since $\hat{C} > 1$, $\lambda_{q+1} > 1$ and $\delta_{q+1} \geq \delta_{q+2}$, the second inequality in (4.4) is obvious.

We are therefore left with showing the first inequality in (4.4) which, as already mentioned, needs a sufficiently large a . As it can be readily checked from the definition of μ , such inequality is in fact equivalent to $\delta_{q+2} \lambda_{q+1}^{1-\alpha} \geq \hat{C} \delta_{q+1} \ell^{-1}$. But we record in fact a much stronger inequality, which turns out to be the key relation to conclude the estimates in Proposition 2.1, as it will become apparent in Section 7. More precisely, given any constant \underline{C} which depends upon α, g, σ_0 and \tilde{C} , the following inequality holds provided a is chosen large enough:

$$(4.6) \quad \delta_{q+2}^2 \lambda_{q+1}^{1-2\alpha} \geq \underline{C} \delta_{q+1}^2 \ell^{-1}.$$

In fact such inequality is equivalent to

$$\delta_{q+2}^2 \lambda_{q+1}^{1-2\alpha} \geq \underline{C} \tilde{C}^{1/(2-\alpha)} \delta_{q+1}^{2-1/(2-\alpha)} \delta_q^{1/(2-\alpha)} \lambda_q^{2/(2-\alpha)}.$$

Taking the logarithm in base a , this is equivalent to

$$(c(1 - 2\alpha) - 2)b^{q+2} \geq \left(\frac{1 + 2c}{2 - \alpha} - 2\right)b^{q+1} - \frac{1}{2 - \alpha}b^q + \log_a \underline{C} + \frac{1}{2 - \alpha} \log_a \tilde{C}.$$

The latter follows for a sufficiently large a (depending upon b, c, \tilde{C} and \underline{C}) provided

$$(c(1 - 2\alpha) - 2)b^2 > \left(\frac{1 + 2c}{2 - \alpha} - 2\right)b - \frac{1}{2 - \alpha},$$

which is equivalent to

$$cb((2 - \alpha)(1 - 2\alpha)b - 2) > 2(2 - \alpha)b^2 + (1 - 2(2 - \alpha))b - 1.$$

The latter inequality is however obviously implied by (2.3) and (2.4).

4.2. Constants

In the rest of the paper we will deal with several estimates where we bound norms of various functions using the parameters introduced so far, namely $\delta_q, \lambda_q, \ell, \mu$ and λ_{q+1} . In front of the expressions involving such parameters there will always be some constants, independent of a, b and c . However it is important to distinguish between two types of such constants: the ones which depend only upon α, g and σ_0 will be denoted by C , whereas the ones which depend also upon the \tilde{C} of Proposition 2.1 will be denoted by C^* . Note also that the parameter σ_0 will in fact be chosen as a function of α in Section 4.4. Therefore the constants denoted by C will depend only upon α and g , whereas those denoted by C^* will depend, additionally, also upon \tilde{C} . Moreover, the values of C and C^* may change from line to line.

4.3. Regularization

Having fixed a standard mollifier φ , we then define

$$(4.7) \quad h_q := \frac{g * \varphi_\ell - (u_q * \varphi_\ell)^\sharp e}{\delta_{q+1}} - \frac{\delta_{q+2}}{\delta_{q+1}} e.$$

Observe that

$$(u_q * \varphi_\ell)^\sharp e + \delta_{q+1} h_q = g * \varphi_\ell - \delta_{q+2} e = g_{q+1} + (g * \varphi_\ell - g).$$

So the strategy of the proof will be to perturb $u_q * \varphi_\ell$ to a map u_{q+1} such that

$$u_{q+1}^\sharp e = (u_q * \varphi_\ell)^\sharp e + \delta_{q+1} h_q + E = g_{q+1} + E + (g * \varphi_\ell - g),$$

(cf. (4.26)) where the error term E is suitably small. Before coming to the construction of the map u_{q+1} we deal in this section with the smallness conditions to be imposed on ℓ .

First of all, by choosing \tilde{C} larger than a geometric constant and a sufficiently large (depending upon b and c), we can assume that $\ell \leq 2^{-q-2}$, so that h_q is in fact defined on $\tilde{D}_{1+2^{-q-2}}$. Next, using Lemma 3.3 we can estimate

$$\begin{aligned} \|h_q - e\|_\alpha &\leq \frac{\delta_{q+2}}{\delta_{q+1}} + \frac{1}{\delta_{q+1}} \|g * \varphi_\ell - (u_q * \varphi_\ell)^\sharp e - \delta_{q+1} e\|_\alpha \\ &\leq a^{-(b-1)} + \frac{1}{\delta_{q+1}} (\|(u_q^\sharp e) * \varphi_\ell - (u_q * \varphi_\ell)^\sharp e\|_\alpha + \|(g_q - u_q^\sharp e) * \varphi_\ell\|_\alpha \\ &\quad + \|g - g * \varphi_\ell\|_\alpha) \\ &\leq \sigma_0 + C^* \frac{\ell^{2-\alpha} \delta_q \lambda_q^2}{\delta_{q+1}} + \sigma_0 + \frac{C}{\delta_{q+1}} \|D^2 g\|_0 \ell^{2-\alpha} \\ &\stackrel{(4.3)}{\leq} 2\sigma_0 + C^* \frac{\ell^{2-\alpha} \delta_q \lambda_q^2}{\delta_{q+1}} \leq 3\sigma_0, \end{aligned}$$

where the latter inequality specifies the condition needed on \tilde{C} in (4.1).

Similarly, for $1 \leq k \leq 4$, we can bound

$$(4.8) \quad \begin{aligned} \|D^k h_q\|_0 &\leq \frac{1}{\delta_{q+1}} (\|D^k (g - u_q^\sharp e) * \varphi_\ell\|_0 + \|D^k ((u_q^\sharp e) * \varphi_\ell - (u_q * \varphi_\ell)^\sharp e)\|_0) \\ &\leq C \ell^{\alpha-k} \sigma_0 + C^* \frac{\delta_q \lambda_q^2}{\delta_{q+1}} \ell^{2-k} \leq C \ell^{\alpha-k}, \end{aligned}$$

where we have used (4.1) and Lemma 3.3. Interpolating, for any $0 \leq k \leq 3$ we then get

$$(4.9) \quad \|h_q - e\|_{k+\alpha} \leq C \ell^{-k}.$$

We summarize the conclusions of the previous paragraphs in the following lemma.

Lemma 4.1. *If we choose \tilde{C} sufficiently large, depending upon α, g and \bar{C} , we then have*

$$(4.10) \quad \|h_q - e\|_\alpha \leq 3\sigma_0$$

$$(4.11) \quad \|h_q - e\|_{k+\alpha} \leq C \ell^{-k} \quad \text{for } 1 \leq k \leq 3,$$

where the constant C depends only upon α and g .

4.4. Conformal diffeomorphism

We now wish to apply Proposition 3.4 with $\beta = \alpha > 0$ and $N = 3$. This requires to choose σ_0 such that $3\sigma_0 \leq \sigma_1$, where σ_1 is the constant appearing in (3.14). We thus find maps Φ and ρ such that

$$h_q = \rho^2 (\nabla\Phi_1 \otimes \nabla\Phi_1 + \nabla\Phi_2 \otimes \nabla\Phi_2) .$$

Furthermore, dividing ρ by $\max \rho$, multiplying Φ by $\max \rho$ and using (3.16), we can assume that

$$(4.12) \quad \frac{1}{2} \leq \rho \leq 2, \quad \|D\Phi - \text{Id}\|_0 \leq \frac{1}{2},$$

provided σ_0 is chosen sufficiently small. This exhausts the condition on σ_0 : note that they depend only upon α , since N and β in Proposition 3.4 are fixed to be 3 and α .

Moreover, for any $1 \leq k \leq 3$ we apply (3.17) and (4.11) to estimate

$$(4.13) \quad \|D^k \rho\|_\alpha + \|D^{k+1} \Phi\|_\alpha \leq C \ell^{-k} .$$

4.5. Adding the first primitive metric

We next set $w := u_q * \varphi_\ell$ and we define the following two three-dimensional vectors:

$$(4.14) \quad \tau_1 := Dw(Dw^T Dw)^{-1} \nabla\Phi_1$$

and

$$(4.15) \quad \nu_1 := \frac{\partial_{x_1} w \times \partial_{x_2} w}{|\partial_{x_1} w \times \partial_{x_2} w|} .$$

Observe that ν_1 is in the kernel of Dw^T (or, in other words, $\nu_1(x)$ is a unit normal to the tangent plane $T_{w(x)}(\text{Im}(w))$). Hence it follows easily that τ_1 and ν_1 are orthogonal.

We next normalize these vectors suitably, defining

$$(4.16) \quad t_1 := \frac{\tau_1}{|\tau_1|^2},$$

$$(4.17) \quad n_1 := \frac{\nu_1}{|\tau_1|} .$$

Finally, we define the first perturbation of w , namely the map v given by the formula

$$(4.18) \quad v = w + \frac{1}{\mu} \Gamma^t (\delta_{q+1}^{1/2} |\tau_1| \rho, \mu \Phi_1) t_1 + \frac{1}{\mu} \Gamma^n (\delta_{q+1}^{1/2} |\tau_1| \rho, \mu \Phi_1) n_1,$$

whereas we define

$$(4.19) \quad E_1 := v^\sharp e - (w^\sharp e + \delta_{q+1} \rho^2 \nabla \Phi_1 \otimes \nabla \Phi_1).$$

4.6. Adding the second primitive metric

The map u_{q+1} is then obtained by adding a similar second perturbation to the map v . More precisely we define this time

$$(4.20) \quad \tau_2 := Dv(Dv^T Dv)^{-1} \nabla \Phi_2,$$

$$(4.21) \quad \nu_2 := \frac{\partial_{x_1} v \times \partial_{x_2} v}{|\partial_{x_1} v \times \partial_{x_2} v|},$$

$$(4.22) \quad t_2 := \frac{\tau_2}{|\tau_2|^2},$$

$$(4.23) \quad n_2 := \frac{\nu_2}{|\tau_2|}.$$

The map u_{q+1} is then given by the following formula (analogous to (4.18)):

$$(4.24) \quad u_{q+1} = v + \frac{1}{\lambda_{q+1}} \Gamma^t (\delta_{q+1}^{1/2} |\tau_2| \rho, \lambda_{q+1} \Phi_2) t_2 + \frac{1}{\lambda_{q+1}} \Gamma^n (\delta_{q+1}^{1/2} |\tau_2| \rho, \lambda_{q+1} \Phi_2) n_2.$$

Similarly we define

$$(4.25) \quad E_2 := u_{q+1}^\sharp e - (v^\sharp e + \delta_{q+1} \rho^2 \nabla \Phi_2 \otimes \nabla \Phi_2).$$

Observe that we have the following identity:

$$(4.26) \quad \begin{aligned} E &:= E_1 + E_2 = u_{q+1}^\sharp e - (w^\sharp e + \delta_{q+1} \rho^2 (\nabla \Phi_1 \otimes \nabla \Phi_1 + \nabla \Phi_2 \otimes \nabla \Phi_2)) \\ &= u_{q+1}^\sharp e - w^\sharp e - \delta_{q+1} h_q = u_{q+1}^\sharp e + \delta_{q+2} e - g * \varphi_\ell \\ &= u_{q+1}^\sharp e - g_{q+1} + (g - g * \varphi_\ell). \end{aligned}$$

Hence

$$\begin{aligned} \|g_{q+1} - u_{q+1}^\sharp e\|_0 &\leq \|E\|_0 + \|g - g * \varphi_\ell\|_0, \\ \|D(g_{q+1} - u_{q+1}^\sharp e)\|_0 &\leq \|DE\|_0 + \|D(g - g * \varphi_\ell)\|_0. \end{aligned}$$

For α sufficiently small and a sufficiently big one can achieve

$$(4.27) \quad \|g - g * \varphi_\ell\|_0 \leq C \|D^2 g\|_0 \ell^2 \leq \frac{\sigma_0}{6} \delta_{q+2} \lambda_{q+1}^{-\alpha},$$

$$(4.28) \quad \|D(g - g * \varphi_\ell)\|_0 \leq C \|D^2 g\|_0 \ell \leq \frac{\sigma_0}{6} \delta_{q+2} \lambda_{q+1}^{1-\alpha}.$$

To see this, note that (4.27) is implied by the condition

$$C^\star \frac{\delta_{q+1}}{\delta_q \lambda_q^2} \leq \delta_{q+2} \lambda_{q+1}^{-\alpha},$$

which for $a(\bar{C})$ big enough is guaranteed if

$$b^2 - b + 1 < (2 - \alpha b)cb,$$

or equivalently

$$(4.29) \quad c > \frac{b^2 - b + 1}{b(2 - \alpha b)}.$$

Similarly, (4.28) follows if

$$C^\star \frac{\delta_{q+1}^{1/2}}{\delta_q^{1/2} \lambda_q} \leq \delta_{q+2} \lambda_{q+1}^{1-\alpha},$$

which (for $a(\bar{C})$ big enough) is satisfied whenever

$$(4.30) \quad c > \frac{2b^2 - b + 1}{2b(1 + (1 - \alpha)b)}.$$

Now for any $\alpha > 0, b > 1$ which satisfy the bounds of Proposition 2.1 we have

$$\frac{b^2 - b + 1}{b(2 - \alpha b)} > \frac{2b^2 - b + 1}{2b(1 + (1 - \alpha)b)}.$$

Indeed, since $b < 3/2$ and $\alpha < \alpha_0$, provided α_0 is small enough both denominators in the fractions above are positive. Hence the inequality is equivalent to

$$2b^2 + (\alpha - 4)b + (2 - \alpha) = (b - 1)(\alpha + 2b - 2) > 0,$$

which for $b > 1$ and $\alpha > 0$ is always true. Hence (4.29) implies (4.30).

Next, observe that the left-hand side of (2.4) is larger than

$$g_\alpha(b) = \frac{(4 - 2\alpha)b + 1}{2b},$$

so (2.4) implies $c > g_\alpha(b)$. The bound (4.29) is instead

$$c > h_\alpha(b) = \frac{b^2 - b + 1}{b(2 - \alpha b)}.$$

On the other hand on the interval $[1, 3/2]$, g_α and h_α converge uniformly, as $\alpha \downarrow 0$, to the functions $g_0(b) = 2 + 1/2b$ and $h_0(b) = \frac{b^2 - b + 1}{2b}$. Since on $[1, 3/2]$ g_0 is strictly larger than h_0 , we infer that for α small (2.4) guarantees (4.29). In particular we conclude that for a big enough (2.4) guarantees (4.27) and (4.28).

Thus, the goal of most of the remaining sections is to prove that the desired bounds hold for $\|E\|_0, \|DE\|_0, \|u_{q+1} - u_q\|_0, \|D(u_{q+1} - u_q)\|_0$ and $\|D^2 u_{q+1}\|_0$.

5. Estimates on v and E_1

Our goal in this subsection is to estimate the C^0 norms of $v - u_q$, $D^k v$, E_1 and DE_1 . To this aim we introduce the functions

$$\begin{aligned} A_1^t &:= \partial_\xi \Gamma^t(\delta_{q+1}^{1/2} |\tau_1| \rho, \mu \Phi_1), \\ A_1^n &:= \partial_\xi \Gamma^n(\delta_{q+1}^{1/2} |\tau_1| \rho, \mu \Phi_1), \\ B_1^t &:= \partial_s \Gamma^t(\delta_{q+1}^{1/2} |\tau_1| \rho, \mu \Phi_1), \\ B_1^n &:= \partial_s \Gamma^n(\delta_{q+1}^{1/2} |\tau_1| \rho, \mu \Phi_1), \\ C_1^t &:= \Gamma^t(\delta_{q+1}^{1/2} |\tau_1| \rho, \mu \Phi_1), \\ C_1^n &:= \Gamma^n(\delta_{q+1}^{1/2} |\tau_1| \rho, \mu \Phi_1), \end{aligned}$$

and we decompose the derivative of v as

$$\begin{aligned} Dv &= Dw + \underbrace{A_1^t t_1 \otimes \nabla \Phi_1 + A_1^n n_1 \otimes \nabla \Phi_1}_{=: \mathbf{A}_1} \\ &+ \underbrace{\frac{\delta_{q+1}^{1/2}}{\mu} (B_1^t t_1 + B_1^n n_1) \otimes (\rho \nabla |\tau_1| + |\tau_1| \nabla \rho)}_{=: \mathbf{B}_1} + \underbrace{\frac{1}{\mu} (C_1^t Dt_1 + C_1^n Dn_1)}_{=: \mathbf{C}_1}. \end{aligned}$$

5.1. First technical lemma

In the next lemma we collect the estimates of the C^0 norm of the derivatives of the various quantities introduced above.

Lemma 5.1. *Let \tilde{C} be fixed so that Lemma 4.1 holds and $\hat{C} \geq 1$. If $a \geq a_0(\alpha, g, b, c, \tilde{C})$ for some a_0 sufficiently large, then there are constants C (depending upon α and g but not on \tilde{C}) such that*

$$(5.1) \quad C^{-1} \leq |\tau_1| \leq C$$

and

$$(5.2) \quad \|w - u_q\|_0 \leq C \delta_{q+1}^{1/2} \ell,$$

$$(5.3) \quad \|D(w - u_q)\|_0 \leq C \delta_{q+1}^{1/2},$$

$$(5.4) \quad \|Dw\|_0 \leq C,$$

$$(5.5) \quad \|D^k w\|_0 \leq C \delta_{q+1}^{1/2} \ell^{1-k} \quad \text{for } 2 \leq k \leq 4,$$

$$(5.6) \quad \|D^k \nu_1\|_0 \leq C \delta_{q+1}^{1/2} \ell^{-k} \quad \text{for } 1 \leq k \leq 3,$$

$$(5.7) \quad \|D^k t_1\|_0 + \|D^k \tau_1\|_0 + \|D^k n_1\|_0 \leq C \ell^{-k} \quad \text{for } 0 \leq k \leq 3,$$

$$(5.8) \quad \|D^k A_1^t\|_0 + \|D^k C_1^t\|_0 \leq C \delta_{q+1} \mu^k \quad \text{for } 0 \leq k \leq 3,$$

$$(5.9) \quad \|D^k A_1^n\|_0 + \|D^k B_1^t\|_0 + \|D^k C_1^n\|_0 \leq C \delta_{q+1}^{1/2} \mu^k \quad \text{for } 0 \leq k \leq 3,$$

$$(5.10) \quad \|D^k B_1^n\|_0 \leq C \mu^k \quad \text{for } 0 \leq k \leq 3.$$

Proof. Since $\|D\Phi - \text{Id}\|_0 \leq 1/2$, we obviously have $1/2 \leq |\nabla\Phi_1| \leq 2$. On the other hand, the estimate (4.10) on h_q of the previous section implies

$$g + 5\delta_{q+1} e \geq w^\sharp e \geq g - 5\delta_{q+1} e.$$

If we assume a sufficiently large (depending only upon g, b and c), we conclude $2g \geq w^\sharp e \geq g/2$. Since $w^\sharp e = Dw^T Dw$, this implies that

$$C |\nabla\Phi_1| \geq |\tau_1| \geq C^{-1} |\nabla\Phi_1|$$

for a constant C which depends only upon g , hence (5.1) follows.

Estimates on w . Observe that

$$\begin{aligned} \|w - u_q\|_0 &\leq C \ell^2 \|D^2 u_q\|_0 \leq C^* \ell^2 \delta_q^{1/2} \lambda_q, \\ \|D(w - u_q)\|_0 &\leq C \ell \|D^2 u_q\|_0 \leq C^* \ell \delta_q^{1/2} \lambda_q. \end{aligned}$$

If we choose $a \geq a_0(\alpha, b, c, \bar{C})$ big enough such that $\bar{C} \leq \ell^{-\alpha/2}$, then (5.2) and (5.3) follow with the help of (4.5). Moreover, (5.3) implies (5.4) by (2.14). Finally, (5.5) is a consequence of (3.10), i.e., $\|D^k w\|_0 \leq C \ell^{2-k} \|D^2 u_q\|_0$, and $\bar{C} \leq \ell^{-\alpha/2}$.

Next, observe that $C \geq |\partial_{x_1} w \times \partial_{x_2} w| \geq C^{-1}$ (again due to $2g \geq Dw^T Dw \geq g/2$). Hence (3.8) implies, for $k \geq 1$,

$$\|D^k \nu_1\|_0 \leq C [Dw]_k \|Dw\|_0 \leq C [Dw]_k \leq C \delta_{q+1}^{1/2} \ell^{-k}.$$

Estimates on τ_1, t_1 and n_1 . The C^0 estimates in (5.7) are a trivial consequence of (5.1). Again by Proposition 3.1 we get

$$\begin{aligned} \|D^k \tau_1\|_0 &\leq C \|Dw\|_0 \|D^{k+1} \Phi\|_0 + C \|D\Phi\|_0 (\|D^{k+1} w\|_0 + \|D^2 w\|_0^k) \\ &\leq C \ell^{-k} + C \delta_{q+1}^{1/2} \ell^{-k} \leq C \ell^{-k}. \end{aligned}$$

A second application of Proposition 3.1 (combined with (5.1)) gives the estimates

$$(5.11) \quad \|D^k |\tau_1|\|_0 + \|D^k |\tau_1|^{-1}\|_0 \leq C \ell^{-k}.$$

Combining (5.11) and (5.6), from (3.6) we infer

$$\|D^k n_1\|_0 \leq C \delta_{q+1}^{1/2} \ell^{-k} + C \ell^{-k} \leq C \ell^{-k}.$$

We argue similarly to conclude $\|D^k t_1\|_0 \leq C \ell^{-k}$.

Remaining estimates. The cases $k = 0$ of (5.8), (5.9) and (5.10) are all simple consequences of Proposition 3.5 and $\| |\tau_1| |\rho| \|_0 \leq C$. For the higher derivatives we consider first C_1^t . We introduce the function

$$\Psi(s, \xi) := \delta_{q+1}^{-1} \Gamma^t(\delta_{q+1}^{1/2} s, \xi)$$

and observe that $\|D^i\Psi\|_0 \leq C(i)$ by the estimates in Proposition 3.5 (c). If we introduce the map $U = (|\tau_1|\rho, \mu\Phi_1)$ we can then write

$$\|D^k C_1^t\|_0 = \delta_{q+1} \|D^k(\Psi \circ U)\|_0.$$

On the other hand, observe that

$$\|D^k U\|_0 \leq C \ell^{-k} + C\mu \ell^{1-k} \stackrel{(4.4)}{\leq} C\mu \ell^{1-k}.$$

Hence, using (3.9) we infer

$$\|D^k C_1^t\|_0 \leq C \delta_{q+1} (\mu \ell^{1-k} + \mu^k) \leq C \delta_{q+1} \mu^k.$$

In case of $A_1^t, A_n^1, B_1^t, C_1^n$ and B_1^n we apply the same argument, keeping the map U as defined above, but changing Ψ respectively to

$$\begin{aligned} \Psi(s, \xi) &:= \delta_{q+1}^{-1} \partial_\xi \Gamma^t(\delta_{q+1}^{1/2} s, \xi), \\ \Psi(s, \xi) &:= \delta_{q+1}^{-1/2} \partial_\xi \Gamma^n(\delta_{q+1}^{1/2} s, \xi), \\ \Psi(s, \xi) &:= \delta_{q+1}^{-1/2} \partial_s \Gamma^t(\delta_{q+1}^{1/2} s, \xi), \\ \Psi(s, \xi) &:= \delta_{q+1}^{-1/2} \Gamma^n(\delta_{q+1}^{1/2} s, \xi), \\ \Psi(s, \xi) &:= \partial_s \Gamma^n(\delta_{q+1}^{1/2} s, \xi). \end{aligned}$$

□

5.2. Estimates on $\|v - u_q\|_0, \|D(v - u_q)\|_0$ and $\|D^k v\|_0$

Taking into account Proposition 3.5 we obviously have

$$\|v - w\|_0 \leq C \delta_{q+1}^{1/2} \mu^{-1},$$

whereas, by (5.2),

$$\|u_q - w\|_0 \leq C \delta_{q+1}^{1/2} \ell \leq C \delta_{q+1}^{1/2} \ell^{1-\alpha/2} \leq C \frac{\delta_{q+1}}{\delta_q^{1/2} \lambda_q}.$$

We therefore conclude

$$(5.12) \quad \|u_q - v\|_0 \leq C \delta_{q+1}^{1/2} \mu^{-1} + C \frac{\delta_{q+1}}{\delta_q^{1/2} \lambda_q}.$$

By Lemma 5.1 we easily see that

$$(5.13) \quad \|D(u_q - v)\|_0 \leq C \delta_{q+1}^{1/2}$$

and

$$(5.14) \quad \|D^k v\|_0 \leq C \delta_{q+1}^{1/2} \mu^{k-1} \quad \text{for } k \in \{2, 3\}.$$

Observe also that, by (2.14),

$$(5.15) \quad \|Dv\|_0 \leq C.$$

5.3. Estimates on $\|E_1\|_0$ and $\|DE_1\|_0$

Observe first that due to Proposition 3.5 (b) we have

$$(Dw + \mathbf{A}_1)^T(Dw + \mathbf{A}_1) = w^\sharp e + \delta_{q+1}\rho^2 \nabla\Phi_1 \otimes \nabla\Phi_1.$$

Using the notation $\text{Sym } P$ for the matrix $\frac{1}{2}(P + P^T)$, we can then write

$$E_1 = 2\text{Sym}(Dw^T(\mathbf{B}_1 + \mathbf{C}_1)) + 2\text{Sym}(\mathbf{A}_1^T(\mathbf{B}_1 + \mathbf{C}_1)) + (\mathbf{B}_1 + \mathbf{C}_1)^T(\mathbf{B}_1 + \mathbf{C}_1).$$

We notice that, from Lemma 5.1 and the estimates (4.12) and (4.13) on ρ and Φ , we conclude

$$\begin{aligned} \|\mathbf{A}_1\|_0 + \mu^{-1}\|D\mathbf{A}_1\|_0 &\leq C \delta_{q+1}^{1/2}, \\ \|\mathbf{B}_1\|_0 + \|\mathbf{C}_1\|_0 + \mu^{-1}(\|D\mathbf{B}_1\|_0 + \|D\mathbf{C}_1\|_0) &\leq C \frac{\delta_{q+1}^{1/2}}{\ell\mu}. \end{aligned}$$

It is therefore obvious that, since $\ell\mu \geq 1$,

$$\begin{aligned} \|E_1\|_0 &\leq \|Dw^T\mathbf{B}_1\| + \|Dw^T\mathbf{C}_1\| + C \frac{\delta_{q+1}}{\ell\mu}, \\ \|DE_1\|_0 &\leq \|D(Dw^T\mathbf{B}_1)\|_0 + \|D(Dw^T\mathbf{C}_1)\|_0 + C \delta_{q+1}\ell^{-1}. \end{aligned}$$

We next compute

$$Dw^T\mathbf{B}_1 = \frac{\delta_{q+1}^{1/2}}{\mu} B_1^t(Dw^T t_1) \otimes (\rho\nabla|\tau_1| + |\tau_1|\nabla\rho).$$

Therefore we conclude from Lemma 5.1 that

$$\begin{aligned} \|Dw^T\mathbf{B}_1\|_0 &\leq C \frac{\delta_{q+1}}{\ell\mu}, \\ \|D(Dw^T\mathbf{B}_1)\|_0 &\leq C \delta_{q+1}\ell^{-1}. \end{aligned}$$

Recalling moreover (4.17), we have

$$Dn_1 = \frac{D\nu_1}{|\tau_1|} - n_1 \otimes \frac{\nabla|\tau_1|}{|\tau_1|},$$

and we also conclude that

$$Dw^T\mathbf{C}_1 = \frac{C_1^t}{\mu} Dw^T Dt_1 + \frac{C_1^n}{\mu} Dw^T \frac{D\nu_1}{|\tau_1|}.$$

In particular,

$$\|Dw^T\mathbf{C}_1\|_0 \leq \frac{C\delta_{q+1}}{\mu\ell} + C \frac{\delta_{q+1}^{1/2}}{\mu} \delta_{q+1}^{1/2} \ell^{-1} \leq C \frac{\delta_{q+1}}{\mu\ell}.$$

Similarly we conclude

$$\|D(Dw^T\mathbf{C}_1)\|_0 \leq C \delta_{q+1}\ell^{-1}.$$

Thus we infer

$$\|E_1\|_0 \leq C \frac{\delta_{q+1}}{\ell\mu}, \quad \text{and} \quad \|DE_1\|_0 \leq C \delta_{q+1}\ell^{-1}.$$

6. Estimates on u_{q+1} and E_2

Our goal in this section is to estimate the C^0 norms of $u_{q+1} - v$, Du_{q+1} , D^2u_{q+1} , E_2 and DE_2 . We proceed in the same way as in the previous section and begin by defining the functions

$$\begin{aligned} A_2^t &:= \partial_\xi \Gamma^t(\delta_{q+1}^{1/2}|\tau_2| \rho, \lambda_{q+1}\Phi_2), \\ A_2^n &:= \partial_\xi \Gamma^n(\delta_{q+1}^{1/2}|\tau_2| \rho, \lambda_{q+1}\Phi_2), \\ B_2^t &:= \partial_s \Gamma^t(\delta_{q+1}^{1/2}|\tau_2| \rho, \lambda_{q+1}\Phi_2), \\ B_2^n &:= \partial_s \Gamma^n(\delta_{q+1}^{1/2}|\tau_2| \rho, \lambda_{q+1}\Phi_2), \\ C_2^t &:= \Gamma^t(\delta_{q+1}^{1/2}|\tau_2| \rho, \lambda_{q+1}\Phi_2), \\ C_2^n &:= \Gamma^n(\delta_{q+1}^{1/2}|\tau_2| \rho, \lambda_{q+1}\Phi_2), \end{aligned}$$

and decomposing the derivative of u_{q+1} as

$$\begin{aligned} Du_{q+1} &= Dv + \underbrace{A_2^t t_2 \otimes \nabla \Phi_2 + A_2^n n_2 \otimes \nabla \Phi_2}_{=:A_2} \\ &+ \underbrace{\frac{\delta_{q+1}^{1/2}}{\lambda_{q+1}}(B_2^t t_2 + B_2^n n_2) \otimes (\rho \nabla |\tau_2| + |\tau_2| \nabla \rho)}_{=:B_2} + \underbrace{\frac{1}{\lambda_{q+1}}(C_2^t Dt_2 + C_2^n Dn_2)}_{=:C_2}. \end{aligned}$$

6.1. Second technical lemma

As before, we collect the estimates of the C^0 norm of the derivatives of the various quantities introduced above.

Lemma 6.1. *Assume \tilde{C} is fixed so that Lemma 4.1 holds and $\hat{C} > 1$. If $a \geq a_0(\alpha, g, b, c, \tilde{C}, \hat{C})$ for a sufficiently large a_0 , then there are constants C (depending on α and g but not on \tilde{C}) such that*

$$\begin{aligned} C^{-1} &\leq |\tau_2| \leq C \\ \|D^k \nu_2\|_0 &\leq C \delta_{q+1}^{1/2} \mu^k \quad \text{for } k \in \{1, 2\} \end{aligned}$$

and, for $k \in \{0, 1, 2\}$,

$$\begin{aligned} \|D^k t_2\|_0 + \|D^k \tau_2\|_0 + \|D^k n_2\|_0 &\leq C \ell^{-k} + C \delta_{q+1}^{1/2} \mu^k \\ \|D^k A_2^t\|_0 + \|D^k C_2^t\|_0 &\leq C \delta_{q+1} \lambda_{q+1}^k \\ \|D^k A_2^n\|_0 + \|D^k B_2^t\|_0 + \|D^k C_2^n\|_0 &\leq C \delta_{q+1}^{1/2} \lambda_{q+1}^k \\ \|D^k B_2^n\|_0 &\leq C \lambda_{q+1}^k. \end{aligned}$$

Proof. The arguments are entirely similar to the ones of Lemma 5.1, where we only need to use the estimates (5.13) and (5.14) on $D^k v$ proved in the previous section and the fact that $\lambda_{q+1} \geq \mu$. □

6.2. Estimates on $\|u_{q+1} - v\|_0$, $\|D(u_{q+1} - v)\|_0$ and $\|D^2u_{q+1}\|_0$

The following estimates are straightforward consequences of Lemma 6.1:

$$(6.1) \quad \|u_{q+1} - v\|_0 \leq C \delta_{q+1}^{1/2} \lambda_{q+1}^{-1},$$

$$(6.2) \quad \|Du_{q+1} - Dv\|_0 \leq C \delta_{q+1}^{1/2},$$

$$(6.3) \quad \|D^2u_{q+1}\|_0 \leq C \delta_{q+1}^{1/2} \lambda_{q+1}.$$

6.3. Estimates on $\|E_2\|_0$ and $\|DE_2\|_0$

Arguing as in Section 5.3, we easily see that

$$\|E_2\|_0 \leq C \delta_{q+1} \frac{\mu}{\lambda_{q+1}}, \quad \text{and} \quad \|DE_2\|_0 \leq C \delta_{q+1} \mu.$$

7. Proof of Proposition 2.1, conclusion

Recall that

$$(7.1) \quad \mu := \hat{C} \frac{\delta_{q+1} \lambda_{q+1}^\alpha}{\delta_{q+2} \ell}$$

for an appropriately large constant \hat{C} , depending upon α and g (in particular not on a). It then follows that

$$\|E_1\|_0 + \lambda_{q+1}^{-1} \|DE_1\|_0 \leq \frac{\sigma_0}{12} \delta_{q+2} \lambda_{q+1}^{-\alpha}.$$

Hence, (recall (4.27) and (4.28)) to achieve the estimates (2.9) and (2.10) we need to verify

$$C \delta_{q+1} \frac{\mu}{\lambda_{q+1}} \leq \frac{\sigma_0}{12} \delta_{q+2} \lambda_{q+1}^{-\alpha},$$

which however is implied by (4.6), which is valid provided a is chosen sufficiently large. The three remaining inequalities (2.11), (2.12) and (2.13) are implied by (5.12)–(5.15) and (6.1)–(6.3).

8. Proof of Theorem 1.1

8.1. Step 1

By using the compactness of the domain \bar{D} we may assume without loss of generality that \bar{u} is uniformly strictly short, that is, $g - \bar{u}^\sharp e \geq 2\bar{\delta}$ in \bar{D} for some $\bar{\delta} > 0$. In a first step we will apply the classical Nash–Kuiper argument to obtain a good first approximation.

To this end recall that there exist a finite number¹ of unit vectors $e_i \in \mathbb{R}^2$ and corresponding amplitudes $\phi_i \in C^\infty(\bar{D})$, $i = 1, \dots, N$ such that

$$g - \bar{u}^\sharp e - \bar{\delta} e = \sum_{i=1}^N \phi_i^2 e_i \otimes e_i \quad \text{in } \bar{D}.$$

Define iteratively the smooth mappings $\bar{u}_0 := \bar{u}$, $\bar{u}_1, \dots, \bar{u}_N := \tilde{u}$ by setting, for $i = 1, \dots, N$,

$$\begin{aligned} \tau_i &:= D\bar{u}_{i-1}(D\bar{u}_{i-1}^T D\bar{u}_{i-1})^{-1} e_i, & \nu_i &:= \frac{\partial_{x_1} \bar{u}_{i-1} \times \partial_{x_2} \bar{u}_{i-1}}{|\partial_{x_1} \bar{u}_{i-1} \times \partial_{x_2} \bar{u}_{i-1}|}, \\ t_i &:= \frac{\tau_i}{|\tau_i|^2}, & n_i &:= \frac{\nu_i}{|\tau_i|}. \end{aligned}$$

and

$$(8.1) \quad \bar{u}_i(x) := \bar{u}_{i-1}(x) + \frac{1}{\mu_i} \Gamma^t(\varphi_i |\tau_i|, \mu_i e_i \cdot x) t_i + \frac{1}{\mu_i} \Gamma^n(\varphi_i |\tau_i|, \mu_i e_i \cdot x) n_i.$$

Here the frequencies $1 \leq \mu_1 \leq \mu_2 \leq \dots \leq \mu_N$ will be inductively defined as follows. Let

$$E_i = \bar{u}_i^\sharp e - \bar{u}_{i-1}^\sharp e - \phi_i^2 e_i \otimes e_i$$

so that $\bar{u}_N^\sharp e = g - \bar{\delta} e + \sum_{i=1}^N E_i$. As in Section 5 we can estimate E_1 as

$$\|E_1\|_0 \leq \frac{C(\bar{u})}{\mu_1}, \quad \|E_1\|_1 \leq C(\bar{u}),$$

where $C(\bar{u})$ is a constant depending on \bar{u} . By interpolation we also have

$$\|E_1\|_\alpha \leq \frac{C(\bar{u})}{\mu_1^{1-\alpha}},$$

and moreover $\|\bar{u} - \bar{u}_1\|_0 \leq C\mu_1^{-1}$. Therefore we can choose μ_1 so that

$$\|E_1\|_\alpha \leq \frac{\sigma_1}{2N} \bar{\delta}, \quad \|\bar{u} - \bar{u}_1\|_0 \leq \frac{\varepsilon}{2N}.$$

Continuing, analogously we obtain

$$\|E_2\|_0 \leq \frac{C(\bar{u}, \mu_1)}{\mu_2}, \quad \|E_2\|_1 \leq C(\bar{u}, \mu_1),$$

and hence choose μ_2 so that

$$\|E_2\|_\alpha \leq \frac{\sigma_1}{2N} \bar{\delta}, \quad \|\bar{u}_2 - \bar{u}_1\|_0 \leq \frac{\varepsilon}{2N}.$$

¹Although the number N in this decomposition depends on $\bar{\delta} > 0$, there is a geometric constant N_* such that for any $x \in \bar{D}$ at most N_* of the functions ϕ_i are non-zero. Nevertheless, this information is not required for our purposes.

In a similar manner, we can inductively choose $\mu_i, i = 3, \dots, N$, so that eventually we obtain

$$\|g - \bar{\delta}e - \tilde{u}^\sharp e\|_\alpha \leq \sum_{i=1}^N \|E_i\|_\alpha \leq \frac{\sigma_1}{2} \bar{\delta} \quad \text{and} \quad \|\bar{u} - \tilde{u}\|_0 \leq \frac{\varepsilon}{2}.$$

Remark 8.1. The construction above can be easily adapted to the case when \bar{u} is an embedding, and in this case also \tilde{u} will be an embedding. This is of course well-known and has been proved by Nash and Kuiper. In order to keep our paper self-contained, we nevertheless include here a short proof.

Since the construction of \tilde{u} from \bar{u} involves finite number of steps, it suffices to ensure that at each step \bar{u}_i remains an embedding, i.e., no self-intersections are introduced. To show this, we proceed by induction and assume that \bar{u}_{i-1} is an embedding. By using Proposition 3.5 and the choice of vectors t_i, n_i we can write (8.1) as

$$\bar{u}_i(x) := \bar{u}_{i-1}(x) + \frac{1}{\mu_i} w_i(x, \mu_i x),$$

where $w_i = w_i(x, \xi)$ satisfies

$$\begin{aligned} [D\bar{u}_{i-1}(x) + \partial_\xi w_i(x, \mu_i z)]^T [D\bar{u}_{i-1}(x) + \partial_\xi w_i(x, \mu_i z)] \\ = D\bar{u}_{i-1}(x)^T D\bar{u}_{i-1}(x) + \phi_i^2(x) e_i \otimes e_i. \end{aligned}$$

for any x, z . In particular, since \bar{u}_{i-1} is an immersion, there exists $\omega_1 > 0$ so that

$$(8.2) \quad |(D\bar{u}_{i-1}(x) + \partial_\xi w_i(x, \mu_i z))e| \geq |D\bar{u}_{i-1}(x)e| \geq \omega_1 |e|$$

for any vector e .

Next, let $x, y \in \bar{D}$. By Taylor's theorem and the mean value theorem there exists z on the line segment $[x, y]$ such that

$$\bar{u}_i(x) - \bar{u}_i(y) = D\bar{u}_{i-1}(x)(x - y) + \partial_\xi w_i(x, \mu_i z)(x - y) + \tilde{E},$$

where

$$|\tilde{E}| \leq C \left(|x - y|^2 + \frac{1}{\mu_i} |x - y| \right),$$

and C is a constant depending on the functions $\bar{u}_{i-1}(x)$ and $w_i(x, \xi)$ but not on μ_i . Let $\rho = \omega_1/4C$ and choose $\mu_i > \rho^{-1}$. From (8.2) we deduce that if $|x - y| \leq \rho$, then

$$|\bar{u}_i(x) - \bar{u}_i(y)| \geq \frac{\omega_1}{2} |x - y|.$$

On the other hand, since \bar{u}_{i-1} is assumed to be globally injective and \bar{D} is compact, there exists $\omega_2 > 0$ such that

$$|\bar{u}_{i-1}(x) - \bar{u}_{i-1}(y)| \geq \omega_2 |x - y| \quad \text{for all } |x - y| \geq \rho.$$

Since obviously $\|\bar{u}_i - \bar{u}_{i-1}\|_0 \leq C\mu_i^{-1}$, it follows that for sufficiently large μ_i we will also have

$$|\bar{u}_i(x) - \bar{u}_i(y)| \geq \omega_2 |x - y| \quad \text{for all } |x - y| \geq \rho.$$

In summary, we have shown that, by choosing μ_i sufficiently large, we can ensure that \bar{u}_i is also an embedding.

8.2. Step 2

In step 1 we obtained a good approximation \tilde{u} in the sense that (2.7) from Proposition 2.1 is satisfied. However, although \tilde{u} is smooth, we have no information on the size of the second derivatives $D^2\tilde{u}$. Therefore in this step we obtain a further approximation u_0 , where in addition second derivatives are controlled so that this second approximation can then be used as the starting point of an iteration with Proposition 2.1.

In this step we assume in addition²

$$(8.3) \quad c > \frac{2}{1 - 2\alpha} + \frac{1}{2b}.$$

We show that, no matter how large a is chosen, there is a map u_0 satisfying the assumptions (2.7) and (2.8) of Proposition 2.1, where the constant \bar{C} in the latter estimate is however *independent* of a (because it depends only on g and \tilde{u}). We proceed as in Section 4, except no regularization step is necessary this time. We set

$$h := \frac{g - \tilde{u}^\sharp e}{\bar{\delta}} - \frac{\delta_1}{\bar{\delta}} e$$

and apply Proposition 3.4 to find (C^3) Φ_1, Φ_2 and ρ so that

$$h := \rho^2(\nabla\Phi_1 \otimes \nabla\Phi_1 + \nabla\Phi_2 \otimes \nabla\Phi_2).$$

We then define

$$\tau_1 := D\tilde{u}(D\tilde{u}^T D\tilde{u})^{-1} \nabla\Phi_1, \quad \nu_1 := \frac{\partial_{x_1}\tilde{u} \times \partial_{x_2}\tilde{u}}{|\partial_{x_1}\tilde{u} \times \partial_{x_2}\tilde{u}|},$$

and

$$t_1 := \frac{\tau_1}{|\tau_1|^2}, \quad n_1 := \frac{\nu_1}{|\tau_1|}.$$

Hence we set

$$(8.4) \quad v = \tilde{u} + \frac{1}{\mu} \Gamma^t(\bar{\delta}^{1/2}|\tau_1|, \mu \Phi_1)t_1 + \frac{1}{\mu} \Gamma^n(\bar{\delta}^{1/2}|\tau_1|, \mu \Phi_1)n_1.$$

Then we define

$$\tau_2 := Dv(Dv^T Dv)^{-1} \nabla\Phi_2, \quad \nu_2 := \frac{\partial_{x_1}v \times \partial_{x_2}v}{|\partial_{x_1}v \times \partial_{x_2}v|},$$

²Indeed it could be checked directly that (2.4) implies (8.3) and hence (8.3) is superfluous: however, proceeding as we do we can spare the reader a slightly tedious computation.

and

$$t_2 := \frac{\tau_2}{|\tau_2|^2}, \quad n_2 := \frac{\nu_2}{|\tau_2|}.$$

The map u_0 is finally given by

$$(8.5) \quad u_0 = v + \frac{1}{\lambda} \Gamma^t(\bar{\delta}^{1/2}|\tau_2|\rho, \lambda \Phi_2)t_2 + \frac{1}{\lambda} \Gamma^n(\bar{\delta}^{1/2}|\tau_2|\rho, \lambda \Phi_2)n_2.$$

Again we assume $\lambda \geq \mu \geq 1$. Analogous computations to the ones in Sections 5 and 6 lead to the estimates

$$\begin{aligned} \|g - (u_0^\sharp e + \delta_1 e)\|_\alpha &\leq C \bar{\delta}^{1/2} \mu^{2\alpha-1} + C \bar{\delta} \mu \lambda^{\alpha-1} \\ \|D^2 u_0\|_0 &\leq C \bar{\delta}^{1/2} \lambda, \end{aligned}$$

where the constant C depends only on \tilde{u} and g . We thus set

$$\mu := C_1 \delta_1^{-1/(1-2\alpha)} \quad \text{and} \quad \lambda := C_2 \mu^{1/(1-\alpha)} \delta_1^{-1/(1-\alpha)}.$$

For a sufficiently large choice of C_2 and C_1 we then achieve (2.7) (recall that $\bar{\delta} < 1$).

Clearly

$$\|D^2 u_0\|_0 \leq C_3 \delta_1^{-2/(1-2\alpha)},$$

for a constant C_3 which depends only upon \tilde{u}, g and α . In order to show that (2.8) is satisfied with a constant \bar{C} independent of a , it suffices to show that

$$\delta_1^{-2/(1-2\alpha)} \leq \delta_0^{1/2} \lambda_0.$$

Taking the logarithms in base a the latter inequality is implied by

$$cb \geq \frac{1}{2} + \frac{2}{1-2\alpha} b.$$

8.3. Step 3

Finally we are ready for the iteration based on Proposition 2.1. Fix any α, b and c which satisfy (2.3), (2.4) and (8.3). Then, for any sufficiently large a , we can construct a map u_0 as in the previous step which satisfies $\|\tilde{u} - u_0\|_0 < \varepsilon/2$ and the assumptions of Proposition 2.1, with a constant \bar{C} which does not depend on a . We can apply Proposition 2.1 to generate u_1 . Using (3.3) we conclude

$$\|g_1 - u_1^\sharp e\|_\alpha \leq \|g_1 - u_1^\sharp e\|_0 + 2\|g_1 - u_1^\sharp e\|_0^{1-\alpha} \|D(g_1 - u_1^\sharp e)\|_0^\alpha \leq \sigma_0 \delta_2.$$

Hence u_1 satisfies again the assumptions of Proposition 2.1. More generally, the proposition can be applied inductively to generate a sequence $(u_q)_{q \geq 0}$. Observe that (2.11)–(2.13) imply that

- $(u_q)_{q \geq 0}$ converges uniformly to a map u which (assuming a sufficiently large) satisfies $\|u_0 - u\|_0 < \varepsilon/2$. By assumption on u_0 we therefore have $\|\tilde{u} - u\|_0 < \varepsilon$.

- Interpolating $\|D(u_{q+1} - u_q)\| \leq C \delta_{q+1}^{1/2}$ and

$$\begin{aligned} \|D^2(u_{q+1} - u_q)\|_0 &\leq \|D^2 u_{q+1}\|_0 + \|D^2 u_q\|_0 \leq \bar{C} \delta_{q+1}^{1/2} \lambda_{q+1} + \bar{C} \delta_q^{1/2} \lambda_q \\ &\leq 2 \bar{C} \delta_{q+1}^{1/2} \lambda_{q+1} \end{aligned}$$

shows

$$\|D(u_{q+1} - u_q)\|_\beta \leq C^* \delta_{q+1}^{1/2} \lambda_{q+1}^\beta,$$

for a constant C^* which depends on α, g and \bar{C} . Hence using the definitions (2.2) of δ_q and λ_q we can see that if $\beta < \frac{1}{2bc}$ then $(u_q)_{q \geq 0}$ is a Cauchy sequence on $C^{1,\beta}$.

We next show that, if α is chosen arbitrarily small, bc can be chosen arbitrarily close to $5/2$, which in turn implies that β can be made arbitrarily close to $1/5$. Indeed if we let $\alpha \downarrow 0$, the conditions (2.3), (2.4) and (8.3) become, respectively

$$b > 1, \quad c > \frac{4b^2 - 3b - 1}{2b(b-1)} = 2 + \frac{1}{2b} \quad \text{and} \quad c > 2 + \frac{1}{2b}.$$

This completes the proof.

9. Proof of Theorem 1.2

First of all we notice that, by classical extension theorems, the first statement can be reduced to Theorem 1.1: it suffices to extend both g and \bar{u} smoothly from \bar{D}_1 to \bar{D}_2 . The extended map is not necessarily short for the extended metric, but we can ensure this if we add to the extension of g a tensor of the form $\varphi(|x|)e$, where φ is a rapidly growing C^∞ function which vanishes identically on $[0, 1]$.

Next, observe that the arguments of the Steps 2 and 3 in Section 8, combined with the extension trick outlined above, give in fact the following corollary.

Corollary 9.1. *Let g be a C^2 metric on \bar{D}_1 . Then there are positive constants C_0, \bar{c} and $\bar{\eta}$ with the following properties. Assume that*

- (i) $\underline{u} : \bar{D}_1 \rightarrow \mathbb{R}^3$ is C^∞ ,
- (ii) $\|g - (\underline{u}^\#e + 2\eta e)\|_0 \leq \bar{c} \eta$ for some $\eta \in (0, \bar{\eta})$.

Then for any $\varepsilon > 0$ and $\delta > 0$ there is an isometric map $u \in C^{1,1/5-\delta}(\bar{D}_1)$ such that $\|Du - D\underline{u}\|_0 \leq C_0 \eta^{1/2}$ and $\|u - \underline{u}\|_0 \leq \varepsilon$.

With this corollary at hand we can prove Theorem 1.2 in two easy steps. In the proof we will restrict to the case of embeddings, the case of immersions can be obtained by easy modifications.

Proof of Theorem 1.2 for embeddings. Let g be a C^2 metric on \bar{D}_1 and $\bar{u} \in C^1(\bar{D}_1, \mathbb{R}^3)$ a short embedding. By a simple rescaling and mollification we may assume without loss of generality that \bar{u} is smooth and strictly short. Next,

fix $\omega > 0$ such that $g \geq 16\omega^2e$ and choose $\eta > 0$ such that $\eta \leq \min\{\omega^2, \bar{\eta}\}$ and $C_0\eta^{1/2} \leq \omega$.

As in Step 1 of the proof of Theorem 1.1 (including Remark 8.1), we first construct a smooth embedding \underline{u} with

$$\|\underline{u} - \bar{u}\|_0 < \frac{\varepsilon}{2}$$

and such that

$$\|g - (\underline{u}^\sharp e + 2\eta e)\|_0 \leq \bar{c}\eta.$$

Then the assumptions of Corollary 9.1 are satisfied and we obtain $u \in C^{1,1/5-\delta}(\bar{D}_1)$ with $u^\sharp e = g$ and such that $\|Du - D\underline{u}\|_0 \leq C_0\eta^{1/2}$ and $\|u - \underline{u}\|_0 \leq \varepsilon/2$.

To complete the proof, it remains to show that the map u is an embedding. We again remark that this argument is well-known and is contained in the works of Nash and Kuiper. First of all, since \underline{u} is C^1 , there exists $\rho > 0$ such that $|D\underline{u}(z) - D\underline{u}(y)| \leq \omega$ if $|z - y| \leq \rho$. On the other hand, since \underline{u} is an embedding, then there is $\zeta > 0$ such that $|\underline{u}(z) - \underline{u}(y)| \geq 3\zeta$ if $|z - y| \geq \rho$.

To show global injectivity, we now observe that

$$|u(z) - u(y)| \geq |\underline{u}(z) - \underline{u}(y)| - 2\varepsilon \geq 3\zeta - 2\zeta = \zeta \quad \text{when } |z - y| \geq \rho.$$

On the other hand, if $|z - y| \leq \rho$ we know that

$$|Du(z) - Du(y)| \leq |D\underline{u}(z) - D\underline{u}(y)| + 2\omega \leq 3\omega,$$

and hence, using Taylor’s formula,

$$|u(z) - u(y) - Du(z)(z - y)| \leq 3\omega|z - y|.$$

We therefore can estimate

$$|u(z) - u(y)| \geq |Du(z)(z - y)| - 3\omega|z - y|$$

But $u^\sharp e = g \geq 16\omega^2e$ implies $|Du(z)(z - y)|^2 \geq 16\omega^2|z - y|^2$, which in turn shows $|u(z) - u(y)| \geq \omega|z - y| > 0$.

This completes the proof of Theorem 1.2.

A. Proof of Proposition 3.4

A.1. Beurling and Cauchy transforms

We will need the following two classical integral operators to construct the coordinate transformation of Proposition 3.4. In this section we use the standard notation $z = x + iy$ for complex numbers. Moreover, we recall two standard differential operators $\partial_z = \frac{1}{2}(\partial_x - i\partial_y)$ and $\partial_{\bar{z}} = \frac{1}{2}(\partial_x + i\partial_y)$.

Definition A.1. Suppose $G \subset \mathbb{C}$ is a bounded smooth open set and $f: G \rightarrow \mathbb{C}$ a function. For $z_0 \in \mathbb{C}$ we define the *Cauchy transform*

$$\mathcal{C}_G[f](z_0) := -\frac{1}{\pi} \int_G \frac{f(z)}{z - z_0} dx dy$$

and the *Beurling transform*

$$\mathcal{S}_G[f](z_0) := -\frac{1}{\pi} \int_G \frac{f(z)}{(z - z_0)^2} dx dy.$$

The latter integral must be understood as a Cauchy principal value, in case it exists. As it is easy to check, the Hölder continuity of f is enough to guarantee its existence at every point.

Remark A.2. In the literature the terms Cauchy and Beurling transforms are often used only for the operators $\mathcal{C}_{\mathbb{C}}$ and $\mathcal{S}_{\mathbb{C}}$.

In the book of I. N. Vekua [30] one can find the following very important properties of the operators \mathcal{C}_G and \mathcal{S}_G (cf. Theorem 1.32 in [30]).

Lemma A.3. Let $N \in \mathbb{N}$, $0 < \alpha < 1$, $G \subset \mathbb{C}$ bounded and $f \in C^{N,\alpha}(\overline{G})$. Then we have

- (i) $\mathcal{C}_G[f] \in C^{N+1,\alpha}(\overline{G})$ and $\mathcal{S}_G[f] \in C^{N,\alpha}(\overline{G})$;
- (ii) $\frac{\partial}{\partial \bar{z}} \mathcal{C}_G[f](z) = f(z)$ and $\frac{\partial}{\partial z} \mathcal{C}_G[f](z) = \mathcal{S}_G[f](z) \forall z \in G$;
- (iii) there exists a constant $C_{N,\alpha}$ such that

$$\|\mathcal{S}_G[f]\|_{N+\alpha} \leq \|\mathcal{C}_G[f]\|_{N+1+\alpha} \leq C_{N,\alpha} \|f\|_{N+\alpha}.$$

Property (iii) will be key in order to prove Proposition 3.4. Observe that we can easily find solutions of equations of the type $f_{\bar{z}} = g$ by setting $f = \mathcal{C}_G[g]$. Moreover, we have $\partial_{\bar{z}} \mathcal{S}_G[f] = f_z$, so \mathcal{S}_G links the two operators $\partial_{\bar{z}}$ and ∂_z . To prove regularity and get good estimates we need one more thing, namely that under suitable circumstances the transforms commute with differentiation. This will be the content of Corollary A.7.

Lemma A.4. Let $r > 0$ and $f \in C^1(\overline{D_r})$. Then for any $z_0 \in D_r$ we have the identities

$$\begin{aligned} f(z_0) &= \frac{1}{2\pi i} \int_{\partial D_r} \frac{f(z)}{z - z_0} dz - \frac{1}{\pi} \int_{D_r} \frac{f_{\bar{z}}(z)}{z - z_0} dx dy, \\ \frac{1}{\pi} \int_{D_r} \frac{f(z)}{(z - z_0)^2} dx dy &= \frac{1}{\pi} \int_{D_r} \frac{f_z(z)}{z - z_0} dx dy + \frac{1}{2\pi i} \int_{\partial D_r} \frac{f(z)}{z - z_0} d\bar{z}. \end{aligned}$$

Proof. Take a fixed $z_0 \in D_r$ and look at the differential one-form $\omega = \frac{dz}{z - z_0}$. We can see that

$$d(\omega f) = \frac{f_{\bar{z}}}{z - z_0} d\bar{z} \wedge dz = 2i \frac{f_{\bar{z}}}{z - z_0} dx \wedge dy,$$

hence by Stoke’s theorem we have

$$(A.1) \quad 2i \int_{D_r \setminus D_\varepsilon} \frac{f_{\bar{z}}(z)}{z - z_0} dx dy = \int_{\partial D_r} \frac{f(z)}{z - z_0} dz - \int_{\partial D_\varepsilon} \frac{f(z)}{z - z_0} dz.$$

We can easily compute

$$\lim_{\varepsilon \rightarrow 0} \int_{\partial D_\varepsilon} \frac{f(z)}{z - z_0} dz = 2\pi i f(z_0),$$

and therefore passing to the limit $\varepsilon \rightarrow 0$ in (A.1) yields the first statement; the same reasoning applied to the one-form $\tilde{\omega} = \frac{d\bar{z}}{z - z_0}$ shows the second. \square

Remark A.5. Observe that if we define $\Psi(z_0) = \frac{1}{2\pi i} \int_{\partial D_r} \frac{f(z)}{z - z_0} dz$, then the statements of the previous lemma can be rewritten as

$$\begin{aligned} f(z_0) &= \Psi(z_0) + \mathcal{C}_{D_r}[f_{\bar{z}}](z_0), \\ \mathcal{S}_{D_r}[f](z_0) &= \mathcal{C}_{D_r}[f_z](z_0) - \frac{1}{2\pi i} \int_{\partial D_r} \frac{f(z)}{z - z_0} d\bar{z}. \end{aligned}$$

Remark A.6. It follows from Lemma A.4 that if $f \in C^1_0(\bar{D}_r)$, then

- (i) $\mathcal{C}_{D_r}[f_{\bar{z}}] = f$,
- (ii) $\mathcal{C}_{D_r}[f_z] = \mathcal{S}_{D_r}[f]$.

Combining these two identities with Lemma A.3 we can derive

$$\begin{aligned} (\mathcal{C}_{D_r}[f])_z &= \mathcal{S}_{D_r}[f] = \mathcal{C}_{D_r}[f_z], \\ (\mathcal{C}_{D_r}[f])_{\bar{z}} &= f = \mathcal{C}_{D_r}[f_{\bar{z}}], \end{aligned}$$

and

$$\begin{aligned} (\mathcal{S}_{D_r}[f])_z &= (\mathcal{C}_{D_r}[f_z])_z = \mathcal{S}_{D_r}[f_z], \\ (\mathcal{S}_{D_r}[f])_{\bar{z}} &= (\mathcal{C}_{D_r}[f_z])_{\bar{z}} = \mathcal{C}_{D_r}[(f_z)_{\bar{z}}] = \mathcal{C}_{D_r}[(f_{\bar{z}})_z] = \mathcal{S}_{D_r}[f_{\bar{z}}]. \end{aligned}$$

This shows that for (sufficiently regular) functions with compact support in D_r , the operators \mathcal{C}_{D_r} and \mathcal{S}_{D_r} commute with any linear differential operator \mathcal{D} with constant coefficients. The regularity needed on the function is only linked to the order of the operator \mathcal{D} .

We summarize the latter discussion in the following.

Corollary A.7. *Let $r > 0$ and let \mathcal{D} be a linear differential operator with constant coefficients of order k . Then we have the following identities on $C^k_c(D_r)$:*

- (i) $\mathcal{D} \circ \mathcal{C}_{D_r} = \mathcal{C}_{D_r} \circ \mathcal{D}$ and $\mathcal{D} \circ \mathcal{S}_{D_r} = \mathcal{S}_{D_r} \circ \mathcal{D}$;
- (ii) $\partial_{\bar{z}} \circ \mathcal{C}_{D_r} = \mathcal{C}_{D_r} \circ \partial_{\bar{z}} = \text{Id}$ and $\partial_z \circ \mathcal{C}_{D_r} = \mathcal{C}_{D_r} \circ \partial_z = \mathcal{S}_{D_r}$;
- (iii) $\partial_{\bar{z}} \circ \mathcal{S}_{D_r} = \mathcal{S}_{D_r} \circ \partial_{\bar{z}} = \partial_{\bar{z}}$.

A.2. Beltrami’s equation

With the various properties above established, we take a fundamental step to the proof of Proposition 3.4. As usual we denote by $C_0^{N,\alpha}(\overline{D}_r)$ the closure of $C_c^{N,\alpha}(D_r)$ in the Hölder space $C^{N,\alpha}(\overline{D}_r)$.

Lemma A.8. *Let $r \geq 1$, $N \in \mathbb{N}$, $N \geq 1$, $0 < \beta \leq \alpha < 1$, $\mu, h \in C_0^{N,\alpha}(\overline{D}_r)$. Then there exist constants $C(N, r, \alpha, \beta)$, $c(N, r, \alpha, \beta)$ and $\tilde{C}(\alpha)$ such that if $\|\mu\|_\alpha \leq c$ there exists a solution $\Phi \in C^{N+1,\alpha}(\overline{D}_r)$ to*

$$(A.2) \quad \Phi_{\bar{z}} - \mu \Phi_z = h,$$

with

$$\begin{aligned} \|\Phi\|_{1+\alpha} &\leq \tilde{C} \|h\|_\alpha, \\ \|D^k \Phi\|_{1+\beta} &\leq C (\|D^k h\|_\beta + \|D^k \mu\|_\beta \|h\|_\beta), \end{aligned}$$

for any $1 \leq k \leq N$.

Proof. By a standard approximation argument, it suffices to prove the lemma under the assumption that the supports of μ and h are compactly contained in D_r .

In order to simplify our notation we will use \mathcal{S} and \mathcal{C} in place of \mathcal{S}_{D_r} and \mathcal{C}_{D_r} . We know (thanks to Lemma A.3) that $\mathcal{S} : C^{0,\alpha}(\overline{D}_r) \rightarrow C^{0,\alpha}(\overline{D}_r)$, as well as $\mathcal{C} : C^{0,\alpha}(\overline{D}_r) \rightarrow C^{1,\alpha}(\overline{D}_r)$ and that there exist two constants C_α, C_β (wlog, $C_\alpha, C_\beta > 1$) such that

$$\|\mathcal{S}[f]\|_\alpha \leq \|\mathcal{C}[f]\|_{1+\alpha} \leq C_\alpha \|f\|_\alpha, \quad \text{and} \quad \|\mathcal{S}[f]\|_\beta \leq \|\mathcal{C}[f]\|_{1+\beta} \leq C_\beta \|f\|_\beta.$$

Consider the operator

$$\mathcal{L}_\alpha : C^{0,\alpha}(\overline{D}_r) \rightarrow C^{0,\alpha}(\overline{D}_r), \quad f \mapsto h + \mu \mathcal{S}[f]$$

We have

$$\|\mathcal{L}_\alpha(f_1) - \mathcal{L}_\alpha(f_2)\|_\alpha \leq \|\mu\|_\alpha C_\alpha \|f_1 - f_2\|_\alpha.$$

So, if

$$\|\mu\|_\alpha \leq \frac{1}{2C_\alpha}$$

then \mathcal{L}_α has a unique fixed point $f \in C^{0,\alpha}(\overline{D}_r)$. This means

$$f = h + \mu \mathcal{S}[f],$$

from which we deduce

$$\|f\|_\alpha \leq \frac{\|h\|_\alpha}{1 - \|\mu\|_\alpha C_\alpha} \leq 2\|h\|_\alpha$$

and

$$f = (\text{Id} - \mu \mathcal{S})^{-1} h = \sum_{n \geq 0} (\mu \mathcal{S})^n h =: \sum_{n \geq 0} \omega_n.$$

This shows in particular that f is compactly supported. Using Corollary A.7 one can show by induction that, for any $1 \leq k \leq N$ and any $n \geq 1$,

$$(A.3) \quad \|D^k \omega_n\|_\alpha \leq \tilde{C} C_\alpha (2\tilde{C} C_\alpha \|\mu\|_\alpha)^{n-1} (\|\mu\|_\alpha \|D^k h\|_\alpha + \|D^k \mu\|_\alpha \|h\|_\alpha),$$

where \tilde{C} is the constant in (3.6). Therefore, if we require

$$(A.4) \quad \|\mu\|_\alpha \leq (4\tilde{C}C_\alpha C_\beta (2r)^{\alpha-\beta})^{-1},$$

then the series

$$\sum_{n \geq 0} D^k \omega_n$$

converges uniformly in $C^{0,\alpha}(\overline{D}_r)$ to $D^k f$, hence $f \in C^{N,\alpha}(\overline{D}_r)$. Moreover, by the same argument,

$$(A.5) \quad \begin{aligned} \|D^k f\|_\beta &\leq \tilde{C} C_\beta (\|\mu\|_\beta \|D^k h\|_\beta + \|D^k \mu\|_\beta \|h\|_\beta) \sum_{n \geq 1} (2\tilde{C} C_\beta (2r)^{\alpha-\beta} \|\mu\|_\alpha)^{n-1} \\ &+ \|D^k h\|_\beta \leq C (\|D^k h\|_\beta + \|D^k \mu\|_\beta \|h\|_\beta), \end{aligned}$$

with the help of (A.4), where the constant C depends only on N, r, α and β . Now we define

$$\Phi(z) = \mathcal{C}[f](z), \quad z \in D_r.$$

By property (iii) of Lemma A.3 we have

$$\Phi_{\bar{z}} = f, \quad \Phi_z = \mathcal{S}[f],$$

hence

$$\Phi_{\bar{z}} - \mu \Phi_z = f - \mu \mathcal{S}[f] = (\text{Id} - \mu \mathcal{S})f = h,$$

so the function Φ solves (A.2) and satisfies

$$\|\Phi\|_{1+\alpha} \leq C_\alpha \|f\|_\alpha \leq 2C_\alpha \|h\|_\alpha.$$

Since $D^k \Phi = \mathcal{C}[D^k f]$ by Corollary A.7 we get by recalling (A.5)

$$\|D^k \Phi\|_{1+\beta} \leq C_\beta \|D^k f\|_\beta \leq C (\|D^k h\|_\beta + \|D^k \mu\|_\beta \|h\|_\beta).$$

This shows the claim. □

We immediately get the following.

Corollary A.9. *Let $r \geq 1, N \in \mathbb{N}, N \geq 1, 0 < \beta \leq \alpha < 1, \mu \in C_0^{N,\alpha}(\overline{D}_r)$. Then there exist constants $C(N, r, \alpha, \beta), c(N, r, \alpha, \beta)$ and $\tilde{C}(\alpha)$ such that if $\|\mu\|_\alpha \leq c$ there exists a solution $\Phi \in C^{N+1,\alpha}(\overline{D}_r)$ to the Beltrami equation*

$$\Phi_{\bar{z}} = \mu \Phi_z$$

with

$$\begin{aligned} \|\Phi(z) - z\|_{1+\alpha} &\leq \tilde{C} \|\mu\|_\alpha, \\ \|D^k (\Phi(z) - z)\|_{1+\beta} &\leq C \|D^k \mu\|_\beta, \end{aligned}$$

for any $1 \leq k \leq N$.

Proof. In the Lemma A.8 choose $h = \mu$ to recover a constant c such that if $\|\mu\|_\alpha \leq c$ then we find ϕ solving

$$\phi_{\bar{z}} - \mu \phi_z = \mu.$$

Set $\Phi(z) = z + \phi(z)$. Then obviously

$$\Phi_{\bar{z}} = \mu \Phi_z$$

and using Lemma A.8 we find

$$\|\Phi(z) - z\|_{1+\alpha} = \|\phi\|_{1+\alpha} \leq \bar{C} \|\mu\|_{\alpha},$$

and

$$\|D^k(\Phi(z) - z)\|_{1+\beta} = \|D^k\phi\|_{1+\beta} \leq C \|D^k\mu\|_{\beta}$$

for any $1 \leq k \leq N$, which is what we wanted. □

A.3. Proof of Proposition 3.4

Given the estimates of the previous paragraphs, Proposition 3.4 can be proved following the classical approach, see for instance [25], Addendum 1 to Chapter 9. We report however the argument for the reader's convenience.

With a simple scaling argument we can assume $r = 1$. Let x, y be global coordinates on \bar{D}_1 . Then g takes the form

$$g = \xi dx^2 + 2\zeta dx dy + \omega dy^2,$$

for some functions $\xi, \zeta, \omega \in C^{N,\alpha}(D_1)$. We want to find a function $\Phi: D_1 \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (\Phi_1(x, y), \Phi_2(x, y)) =: (s, t)$ such that in these new coordinates we have

$$g = \rho^2 \circ \Phi^{-1}(s, t) (ds^2 + dt^2),$$

hence

$$g = \rho^2 ((\Phi_{1x}^2 + \Phi_{2x}^2) dx^2 + 2(\Phi_{1x}\Phi_{1y} + \Phi_{2x}\Phi_{2y}) dx dy + (\Phi_{1y}^2 + \Phi_{2y}^2) dy^2),$$

or

$$(A.6) \quad g = \rho^2 (\nabla\Phi_1 \otimes \nabla\Phi_1 + \nabla\Phi_2 \otimes \nabla\Phi_2).$$

A comparison yields

$$\xi\omega - \zeta^2 = \rho^4 (\Phi_{1x}\Phi_{2y} - \Phi_{1y}\Phi_{2x})^2 = \rho^4 J\Phi^2,$$

with $J\Phi = \det \nabla\Phi$. Consequently

$$(A.7) \quad \rho^2 = \frac{\sqrt{\Delta}}{J\Phi},$$

where $\Delta = \xi\zeta - \omega^2$.

It is convenient to switch to complex notation. Consider $z = x + iy$, $\Phi(z) = \Phi_1(x, y) + i\Phi_2(x, y)$. A computation shows that (A.6) is equivalent to the Beltrami equation for Φ :

$$(A.8) \quad \Phi_{\bar{z}}(z) = \mu(z) \Phi_z(z), \quad z \in D_1,$$

with the coefficient

$$(A.9) \quad \mu = \frac{\xi - \omega + 2i\zeta}{\xi + \omega + 2\sqrt{\Delta}}.$$

Now we extend g to a symmetric 2×2 tensor to \mathbb{R}^2 so that

$$\begin{aligned} \|g - e\|_{\alpha; \mathbb{R}^2} &\leq \bar{C}(\alpha) \|g - e\|_{\alpha; \bar{D}_1}, \\ \|g - e\|_{k+\beta; \mathbb{R}^2} &\leq C(N, \alpha, \beta) \|g - e\|_{k+\beta; \bar{D}_1}, \end{aligned}$$

for $1 \leq k \leq N$. In particular note that if σ_1 is chosen sufficiently small, then $g \geq \frac{1}{2}e$ on the whole \mathbb{R}^2 . Repeated applications of (3.4) and (3.6) to the expression (A.9) then yield

$$(A.10) \quad \|\mu\|_{\alpha; \mathbb{R}^2} \leq C \|g - e\|_{\alpha; D_1},$$

$$(A.11) \quad \|\mu\|_{k+\beta; \mathbb{R}^2} \leq C \|g - e\|_{k+\beta; D_1},$$

where the former constant in (A.10) is a universal one and the latter depends only on α, β and N . Hence $\mu \in C^{N, \alpha}(\mathbb{R}^2)$. Next we choose a C^∞ cutoff function η such that

$$\eta(z) = \begin{cases} 1, & \text{if } z \in \bar{D}_1, \\ 0, & \text{if } z \in \mathbb{C} \setminus D_{3/2}. \end{cases}$$

With this define a new function

$$\tilde{\mu} = \eta \mu.$$

By definition we have $\tilde{\mu} \in C_c^{N, \alpha}(D_2)$. Thus, by Corollary A.9, there exist constants C, c and \bar{C} such that if $\|\tilde{\mu}\|_{\alpha; D_2} \leq c$ then there exists $\Phi \in C^{N+1, \alpha}(\bar{D}_2)$ with

$$\Phi_{\bar{z}}(z) = \tilde{\mu}(z) \Phi_z(z), \quad z \in D_2,$$

and

$$(A.12) \quad \begin{aligned} \|\Phi(z) - z\|_{1+\alpha; D_2} &\leq \bar{C} \|\tilde{\mu}\|_{\alpha; D_2}, \\ \|D^k(\Phi(z) - z)\|_{1+\beta; D_2} &\leq C \|\tilde{\mu}\|_{k+\beta; D_2}, \end{aligned}$$

for any $1 \leq k \leq N$. Observe that in particular Φ solves (A.8). Moreover,

$$\|\tilde{\mu}\|_{\alpha; D_2} \leq \|\mu\|_{\alpha; D_2} \|\eta\|_{\alpha; D_2} \leq C \|\mu\|_{\alpha; D_2} \leq C \|g - e\|_{\alpha; D_1},$$

and similarly

$$\|\tilde{\mu}\|_{k+\beta; D_2} \leq C \|\mu\|_{k+\beta; D_2} \leq C \|g - e\|_{k+\beta; D_1},$$

by (A.10) and (A.11). This shows that if $\|g - e\|_{\alpha; D_1} \leq \sigma_1$ with σ_1 small enough, we recover a coordinate change Φ solving (A.8). The estimates for Φ follow immediately. For the estimates of ρ we use the fact that due to (A.12) we have

$$(1 - C \|g - e\|_{\alpha; \bar{D}_1})^2 \leq J\Phi \leq (1 + C \|g - e\|_{\alpha; \bar{D}_1})^2,$$

which together with the expression (A.7), the bounds on Φ , (3.4) and (3.6) imply

$$\|D^k \rho\|_{\beta} \leq C \|g - e\|_{k+\beta; \bar{D}_1}$$

for $1 \leq k \leq N$. This proves the claim.

References

- [1] BORISOV, YU. F.: On the connection between the spatial form of smooth surfaces and their intrinsic geometry. *Vestnik Leningrad. Univ.* **14** (1959), no. 13, 20–26.
- [2] BORISOV, YU. F.: On the question of parallel displacement on a smooth surface and the connection of space forms of smooth surfaces with their intrinsic geometries. *Vestnik Leningrad. Univ.* **15** (1960), no. 19, 127–129.
- [3] BORISOV, YU. F.: $C^{1,\alpha}$ -isometric immersions of Riemannian spaces. *Dokl. Akad. Nauk SSSR* **163** (1965), 11–13.
- [4] BORISOV, YU. F.: Irregular $C^{1,\beta}$ -surfaces with analytic metric. *Sibirsk. Mat. Zh.* **45** (2004), no. 1, 25–61.
- [5] BUCKMASTER, T.: Onsager’s conjecture almost everywhere in time. *Comm. Math. Phys.* **333** (2015), no. 3, 1175–1198.
- [6] BUCKMASTER, T., DE LELLIS, C., ISETT, P. AND SZÉKELYHIDI, JR., L.: Anomalous dissipation for $1/5$ -Hölder Euler flows. *Ann. of Math. (2)* **182** (2015), no. 1, 127–172.
- [7] BUCKMASTER, T., DE LELLIS, C. AND SZÉKELYHIDI, JR., L.: Transporting microstructure and dissipative Euler flows. Preprint, arXiv:1302.2815v3, 2013.
- [8] BUCKMASTER, T., DE LELLIS, C. AND SZÉKELYHIDI, JR., L.: Dissipative Euler flows with Onsager-critical spatial regularity. *Comm. Pure Appl. Math.* **69** (2016), no. 9, 1613–1670.
- [9] COHN-VOSSEN, S.: Zwei Sätze über die Starrheit der Eiflächen. *Nachrichten Göttingen* **1927** (1927), 125–137.
- [10] CONSTANTIN, P., E, W. AND TITI, E. S.: Onsager’s conjecture on the energy conservation for solutions of Euler’s equation. *Comm. Math. Phys.* **165** (1994), no. 1, 207–209.
- [11] CONTI, S., DE LELLIS, C. AND SZÉKELYHIDI, JR., L.: h -principle and rigidity for $C^{1,\alpha}$ isometric embeddings. In *Nonlinear partial differential equations*, 83–116. Abel Sympos. 7, Springer, Heidelberg, 2012.
- [12] DE LELLIS, C. AND SZÉKELYHIDI, JR., L.: The Euler equations as a differential inclusion. *Ann. of Math. (2)* **170** (2009), no. 3, 1417–1436.
- [13] DE LELLIS, C. AND SZÉKELYHIDI, JR., L.: The h -principle and the equations of fluid dynamics. *Bull. Amer. Math. Soc. (N.S.)* **49** (2012), no. 3, 347–375.
- [14] DE LELLIS, C. AND SZÉKELYHIDI, JR., L.: Dissipative continuous Euler flows. *Invent. Math.* **193** (2013), no. 2, 377–407.
- [15] DE LELLIS, C. AND SZÉKELYHIDI, JR., L.: Dissipative Euler flows and Onsager’s conjecture. *J. Eur. Math. Soc. (JEMS)* **16** (2014), no. 7, 1467–1505.
- [16] ELIASHBERG, Y. AND MISHACHEV, N.: *Introduction to the h -principle*. Graduate Studies in Mathematics 48, American Mathematical Society, Providence, RI, 2002.
- [17] EYINK, G. L.: Energy dissipation without viscosity in ideal hydrodynamics. I. Fourier analysis and local energy transfer. *Phys. D* **78** (1994), no. 3-4, 222–240.
- [18] GROMOV, M.: Convex integration of differential relations. I. *Izv. Akad. Nauk U.S.S.R.* **37** (1973), 329–343.
- [19] GROMOV, M.: *Partial differential relations*. Results in Mathematics and Related Areas (3) 9, Springer-Verlag, Berlin, 1986.
- [20] HERGLOTZ, G.: Über die Starrheit der Eiflächen. *Abh. Math. Sem. Hansische Univ.* **15** (1943), 127–129.

- [21] ISETT, P.: *Hölder continuous Euler flows in three dimensions with compact support in time*. Annals of Mathematics Studies 196, Princeton University Press, Princeton, NJ, 2017.
- [22] KUIPER, N.: On C^1 isometric imbeddings. I, II. *Nederl. Akad. Wetensch. Proc. Ser. A.* **58** (1955), 545–556; *Indag. Math.* **17** (1955), 683–689.
- [23] NASH, J.: C^1 isometric imbeddings. *Ann. of Math. (2)* **60** (1954), 383–396.
- [24] ONSAGER, L.: Statistical hydrodynamics. *Nuovo Cimento (9)* **6** (1949), Supplemento, no. 2 (Convegno Internazionale di Meccanica Statistica), 279–287.
- [25] SPIVAK, M.: *A comprehensive introduction to differential geometry. Vol. IV*. Second edition. Publish or Perish, Wilmington, Del., 1979.
- [26] SPIVAK, M.: *A comprehensive introduction to differential geometry. Vol. V*. Second edition. Publish or Perish, Wilmington, Del., 1979.
- [27] SPRING, D.: On the regularity of solutions in convex integration theory. *Invent. Math.* **104** (1991), no. 1, 165–178.
- [28] SPRING, D.: *Convex integration theory. Solutions to the h -principle in geometry and topology*. Monographs in Mathematics 92, Birkhäuser, Basel, 1998.
- [29] SZÉKELYHIDI, JR., L.: From isometric embeddings to turbulence. In *HCDTE lecture notes. Part II. Nonlinear hyperbolic PDEs, dispersive and transport equations*, 63 pp. AIMS Ser. Appl. Math. 7, Am. Inst. Math. Sci. (AIMS), Springfield, MO, 2013.
- [30] VEKUA, I. N.: *Generalized analytic functions*. Pergamon Press, London-Paris-Frankfurt; Addison-Wesley, Reading, Mass., 1962.
- [31] YAU, S.-T.: Open problems in geometry. In *Differential geometry: partial differential equations on manifolds (Los Angeles, CA, 1990)*, 1–28. Proc. Sympos. Pure Math. 54, Part 1, Amer. Math. Soc., Providence, RI, 1993.

Received March 30, 2016; revised September 8, 2016.

CAMILLO DE LELLIS: Institut für Mathematik, Universität Zürich, CH-8057 Zürich, Switzerland.

E-mail: camillo.delellis@math.uzh.ch

DOMINIK INAUEN: Institut für Mathematik, Universität Zürich, CH-8057 Zürich, Switzerland.

E-mail: dominik.inauen@uzh.ch

LÁSZLÓ SZÉKELYHIDI JR.: Institut für Mathematik, Universität Leipzig, D-04109 Leipzig, Germany.

E-mail: laszlo.szekelyhidi@math.uni-leipzig.de