



Numerical evidence towards a positive answer to Morrey's problem

André Guerra and Rita Teixeira da Costa

Abstract. We report on numerical experiments suggesting that rank-one convexity implies quasiconvexity in the planar case. We give a simple heuristic explanation of our findings.

1. Introduction

An important problem in the vectorial calculus of variations is to characterize the integrands $f: \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ for which the functional

$$\mathcal{F}[u] \equiv \int_{\Omega} f(Du(x)) \, dx, \quad \text{where } u: \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m \text{ and } m, n \geq 2,$$

is lower semicontinuous with respect to the weak topology in an appropriate Sobolev space; this is the natural condition for existence of minimizers through the direct method.

In his seminal work [28], Morrey recognized that the weak lower semicontinuity of \mathcal{F} is essentially equivalent to a weak notion of convexity, called quasiconvexity, on f . Despite many efforts in the last five decades, an explicit description of quasiconvex functions remains elusive: for instance, there are fourth-order polynomials whose quasiconvexity has been neither proved nor disproved. Such a description would be relevant not only in the calculus of variations but also in other areas of analysis [17, 22, 27, 39].

Quasiconvexity has been mostly studied in relation with polyconvexity [2] and rank-one convexity; these are respectively stronger and weaker notions that are much easier to tackle [6]. We will focus on the relation between quasiconvexity and rank-one convexity. It is useful to consider certain classes of measures that can be seen as being dual to these notions: gradient Young measures and laminates are, respectively, the probability measures that satisfy Jensen's inequality with respect to quasiconvex and rank-one convex functions; see [31] for more details.

The following remains one of the main open problems in the calculus of variations.

Question 1.1. Are rank-one convex functions quasiconvex? Equivalently, let ν be a compactly supported gradient Young measure in $\mathbb{R}^{m \times n}$; is ν a laminate?

Question 1.1 is usually referred to as Morrey’s problem. It seems that Morrey himself was not sure about what the answer to Question 1.1 should be [28, 29]. A fundamental example of Šverák [37] shows that the answer is negative if $m \geq 3$ and $n \geq 2$, and more recently, Grabovsky [13] obtained a different example when $m = 8$ and $n = 2$. Šverák’s example is a polynomial of degree four; Grabovsky’s example, although analytically more complicated, has the advantage of being 2-homogeneous and invariant under the right-action of $SO(2)$. Question 1.1 remains open in the case of low-dimensional targets, i.e., when $m = 2$ and $n \geq 2$. There is some partial evidence suggesting that the answer might be positive in this case, see e.g. the landmark results of [11, 16, 24, 30]. However, and despite remarkable progress, it is by no means clear that the answer should be positive even in low dimensions.

Since the analytic study of quasiconvexity remains incredibly challenging, it is natural to look for numerical evidence instead. In earlier attempts to do so in [7, 8, 14], the strategy is to fix a rank-one convex function f and look for deformations $u: \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that f does not satisfy Jensen’s inequality with respect to Du . A major shortfall in this approach is that explicit rank-one convex non-polyconvex functions are rare and the available examples are relatively simple and have many symmetries.

In the present note, we take a far more general approach: our candidate rank-one functions are the rank-one envelopes of random functions. This approach enables us to cover a significantly larger portion of the space of rank-one convex functions. To be concrete, in line with the ideas from [37], we fix a Lipschitz deformation $u: \mathbb{T}^n \rightarrow \mathbb{R}^m$ whose gradient has finite image. We check if Jensen’s inequality with respect to Du is falsified by the rank-one convex envelopes of functions of the form

$$f(A) = \begin{cases} g(A), & A \in [Du], \\ 2, & \text{otherwise,} \end{cases}$$

where $[Du]$ denotes the essential range of $Du \in L^\infty(\Omega, \mathbb{R}^{m \times n})$ and $g: [Du] \rightarrow [-1, 1]$ is any function. Note that for a deformation such that $[Du]$ is finite the task of looking for counterexamples is a finite-dimensional problem. As a small technical remark we note that it is important that f only takes finite, although large, values; it is easy to build examples of rank-one convex non-quasiconvex functions if the value $+\infty$ is allowed [3].

We consider random deformations given by the sum of N plane waves for $N \in \{3, 4, 5\}$, see Section 3 for further details. The cases $N = 1, 2$ are not interesting and for $N \geq 6$ we already have that $\#[Du] \geq 64$, so the space of functions $g: [Du] \rightarrow [-1, 1]$ becomes very high-dimensional. Considering such deformations is not very restrictive: in fact, arbitrary deformations can be approximated by sums of plane waves. Moreover, James’s interpretation of Šverák’s example shows that, when $m = 3$, there is already a counterexample for $N = 3$.

Our findings can be summarised as follows. When $m \geq 3$, our approach finds many potential counterexamples, similar to the ones in [37]. When $m = 2$, and despite sampling thousands of different deformations, none were found. This suggests that, when $m = 2$, rank-one convexity and quasiconvexity may be equivalent. We also observe that on average it is easier to check that a given deformation does not yield a counterexample to Question 1.1 as N increases.

We provide a basic heuristic explanation of our findings: for plane wave expansions, the rank-one geometry of the set $[Du]$ is drastically different in the cases $m = 3$ and $m = 2$ and, in the latter, the geometry becomes much richer as N increases. Our considerations are inspired by the very interesting results of Sebestyén–Székelyhidi [36], where the authors tap into this structure to prove that no counterexamples arise when $m = 2$ and $N = 3$, see also [34].

We conclude this introduction by discussing the algorithm we use to look for counterexamples to Question 1.1. By homogenization, the gradient of a Lipschitz deformation $u: \mathbb{T}^n \rightarrow \mathbb{R}^m$ generates a gradient Young measure, which has finite support if $[Du]$ is finite; thus our goal is to determine whether this measure is a laminate. Hence, we are naturally led to consider:

Question 1.2 ([23]). Is there an effective algorithm to decide whether a given probability measure supported on a finite subset of $\mathbb{R}^{m \times n}$ is a laminate?

Deciding whether a given measure is a laminate is difficult, as in principle one has to test Jensen’s inequality with all rank-one convex functions [33]. One possible way of circumventing this issue is to consider just the extremal rank-one convex functions [15]; however, the general structure of these functions remains unclear. A different approach is to use a discretized version of the Kohn–Strang algorithm [25] and in Section 2 we show that this yields a partially satisfying answer to Question 1.2. We rely on the convergence of approximations to the rank-one convex envelope, which were proved in [5, 9, 10, 32], see also [40] for particular examples. We remark that the related problem of calculating the rank-one convex hull of a set still remains poorly understood, see [1] and the references therein.

2. Deciding whether a measure is a laminate

In this section we discuss Question 1.2: throughout, ν is a fixed probability measure with support on a finite set $K \subset \mathbb{R}^{m \times n}$. In this section we use a discretized version of the Kohn–Strang algorithm to show the following.

Proposition 2.1. *Let ν be a probability measure supported in a finite set of points in $\mathbb{R}^{m \times n}$. The problem of deciding whether ν is a laminate is semidecidable, i.e., there is an algorithm which terminates in finite time with a positive answer if ν is a laminate.*

To prove this, we will resort to Pedregal’s theorem [33]: ν is a laminate if and only if

$$(2.1) \quad f^{\text{rc}}(\bar{\nu}) \leq \langle \nu, f \rangle, \quad \bar{\nu} \equiv \langle \nu, \text{id} \rangle$$

for all continuous $f: \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$, where f^{rc} denotes the rank-one convex envelope of f .

Lemma 2.2. *If ν is not a laminate there is $g: K \rightarrow [-1, 1]$ such that, for $0 < \delta < c(K, n, m)$ small enough, the continuous function $f_\delta: \mathbb{R}^{m \times n} \rightarrow [-1, 2]$, defined by*

$$(2.2) \quad f_\delta(A) \equiv \begin{cases} g(A_0) + \frac{2-g(A_0)}{\delta}|A - A_0| & \text{if } |A - A_0| \leq \delta \text{ for some } A_0 \in K, \\ 2 & \text{otherwise,} \end{cases}$$

satisfies $f_\delta^{\text{rc}}(\bar{\nu}) > \langle \nu, f_\delta^{\text{rc}} \rangle$.

Although this is not needed, note that $\lim_{\delta \rightarrow 0} f_\delta = f_0 \equiv g \mathbb{1}_K + 2 \times \mathbb{1}_{\mathbb{R}^{m \times n} \setminus K}$ pointwise.

Proof. Since ν is not a laminate, there is $\tilde{g}: \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ rank-one convex and such that $g(\bar{\nu}) > \langle \nu, g \rangle$; by scaling, we can assume that $\tilde{g}([-2, 2]^{mn}) \subseteq [-1, 1]$. Since \tilde{g} is rank-one convex it is locally Lipschitz and, see [4],

$$\text{Lip}(\tilde{g}, [-1, 1]^{mn}) \leq \min\{m, n\} \text{osc}(\tilde{g}, [-2, 2]^{mn}) \leq 2 \min\{m, n\}.$$

Let us take $g = \tilde{g}|_K$, $\delta \leq \frac{1}{2} \min\{|X_1 - X_2| : X_1, X_2 \in K\}$ so that f_δ is well-defined and, in addition, we require that $\delta < (2 \min\{m, n\})^{-1}$. Thus, for $A \in B_\delta(A_0)$ and $A_0 \in K$,

$$\tilde{g}(A) \leq \tilde{g}(A_0) + 2 \min\{m, n\}|A - A_0| \leq \tilde{g}(A_0) + \frac{1}{\delta}|A - A_0| \leq f_\delta(A).$$

This shows that $f_\delta \geq \tilde{g}$; since g is rank-one convex, also $f_\delta^{\text{rc}} \geq \tilde{g}$ and so

$$f_\delta^{\text{rc}}(\bar{\nu}) \geq \tilde{g}(\bar{\nu}) > \langle \nu, g \rangle = \langle \nu, f_\delta^{\text{rc}} \rangle.$$

The proof is finished. ■

Lemma 2.2 shows that in order to decide whether ν is a laminate one has to explore the finite-dimensional space of functions $g: K \rightarrow [-1, 1]$. In order to compute an approximation of f_δ^{rc} we use a discrete version of the Kohn–Strang algorithm [25].

Algorithm 2.3. We fix $\delta > 0$ small enough so that we do not need to worry about it; thus we drop the subscript δ . By translation invariance we can assume that $\bar{\nu} = 0$. Then:

1. Fix an odd integer L , consider the grid $\mathcal{G}_L \equiv \frac{1}{L}\mathbb{Z}^{mn} \cap [-1, 1]^{mn}$, and choose a finite set of directions \mathcal{D} consisting of rank-one matrices which are in \mathcal{G}_L .
2. Set $f_{L, \mathcal{D}}^{\text{rc}, 0} := f$ and, for $A \in \mathcal{G}_L$,

$$f_{L, \mathcal{D}}^{\text{rc}, i+1}(A) = \min_{X \in \mathcal{D}: A \pm X \in \mathcal{G}_L} \left\{ \frac{f_{L, \mathcal{D}}^{\text{rc}, i}(A + X) + f_{L, \mathcal{D}}^{\text{rc}, i}(A - X)}{2}, f_{L, \mathcal{D}}^{\text{rc}, i}(A) \right\}.$$

We terminate the algorithm if either the maximum difference between iterates stabilizes or $f_{L, \mathcal{D}}^{\text{rc}, i}$ satisfies Jensen’s inequality with respect to ν .

Let $f^{\text{rc}, i}$ be the i -th Kohn–Strang iterate, i.e., $f^{\text{rc}, i}$ is defined inductively by $f^{\text{rc}, 0} = f$ and

$$f^{\text{rc}, i}(A) = \inf \left\{ \lambda f^{\text{rc}, i-1}(X) + (1 - \lambda) f^{\text{rc}, i-1}(Y) : \begin{array}{l} \lambda X + (1 - \lambda)Y = A, \\ \text{rank}(X - Y) = 1 \end{array} \right\},$$

where λ runs over $(0, 1)$. Clearly, for $A \in \mathcal{G}_L$, $f^{\text{rc}, i}(A) \leq f_{L, \mathcal{D}}^{\text{rc}, i}(A) \leq f(A)$, and so if $f_{L, \mathcal{D}}^{\text{rc}, i}$ satisfies Jensen’s inequality with respect to ν , f_δ satisfies (2.1). Conversely, we have:

Proposition 2.3. Let $f_{L, \mathcal{D}}^{\text{rc}} \equiv \lim_{i \rightarrow \infty} f_{L, \mathcal{D}}^{\text{rc}, i}$. Then $f_{L, \mathcal{D}}^{\text{rc}}$ converges uniformly to f_δ^{rc} as $L \rightarrow \infty$ and as the largest angle between any rank-one matrix and its best approximation in \mathcal{D} goes to zero.

For a proof see [32]. Note that we take $0 < \delta < c(m, n, K)$ and that f_δ is continuous, so their results apply. It is clear that combining Lemma 2.2 with Proposition 2.3, we deduce Proposition 2.1.

3. Gradient Young measures versus laminates

In this section we address Question 1.1. Recall that a function $f: \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ is said to be *quasiconvex* if, for all $A \in \mathbb{R}^{m \times n}$,

$$(3.1) \quad f(A) \leq \int_{\mathbb{T}^n} f(A + D\varphi(x)) \, dx \quad \text{for all } \varphi \in C^\infty(\mathbb{T}^n, \mathbb{R}^m).$$

Equivalently, f is quasiconvex if and only if $f(\bar{\nu}) \leq \langle \nu, f \rangle$, where ν is any compactly supported gradient Young measure [20, 31].

We want to test the inequality (3.1) with deformations of the form

$$(3.2) \quad \varphi(x) = \sum_{i=1}^N a_i s(x \cdot n_i + c_i),$$

where $N \in \mathbb{N}$, $a_i \in \mathbb{R}^m$, $n_i \in \mathbb{Z}^n$ are vectors, $c_i \in \mathbb{R}$ are phases and s is the 1-periodic sawtooth function, defined for $t \in [0, 1]$ by $s(t) = t1_{[0, 1/2]}(t) + (1 - t)1_{[1/2, 1]}(t)$. The idea of approximating an arbitrary deformation with a simplified deformation with the form (3.2) is known in the applied harmonic analysis literature as a ridgelet expansion [35]. We remark as a somewhat inconvenient fact that orthonormal ridgelet bases in L^2 , just like Fourier series, are never unconditional bases in L^p for $p \neq 2$, although we do not prove this here.

The advantage of an expansion as in (3.2) is that, with $h \equiv s'$ being the Haar wavelet,

$$D\varphi(x) = \sum_{i=1}^N h(x \cdot n_i + c_i) a_i \otimes n_i;$$

hence the gradient $D\varphi$ takes values in a finite set. In our context, considering plane-wave expansions as in (3.2) is a classical idea, and we are motivated by James’ interpretation of Šverák’s example [31], see also [27], §31, and [34, 36]. Moreover, φ generates a homogeneous gradient Young measure ν , which takes the form

$$(3.3) \quad \nu = \sum_{\varepsilon \in \{-1, 1\}^N} \nu_\varepsilon \delta_{X_\varepsilon},$$

where we define the weights ν_ε and the matrices X_ε as

$$(3.4) \quad \nu_\varepsilon \equiv |\{x \in \mathbb{T}^2 : h(x \cdot n_i + c_i) = \varepsilon_i, i = 1, \dots, N\}|, \quad X_\varepsilon \equiv \sum_{i=1}^N \varepsilon_i a_i \otimes n_i.$$

Note that ν_ε depends on n_i but not on a_i . Furthermore, the measure ν has barycentre zero.

For the sake of conciseness, we introduce the following definition.

Definition 3.1. For $N \in \mathbb{N}$, we say that $f: \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ is *N-wave quasiconvex at zero* if

$$f(0) \leq \sum_{\varepsilon \in \{-1, 1\}^N} \nu_\varepsilon f(X_\varepsilon)$$

for all $(a_i, n_i, c_i) \in \mathbb{R}^m \times \mathbb{Z}^n \times \mathbb{R}$, where ν_ε and X_ε are defined by (3.4). Moreover, f is *N-wave quasiconvex* if, for any $A \in \mathbb{R}^{m \times n}$, the function $f(\cdot - A)$ is *N-wave quasiconvex at zero*.

It seems that variants of this notion were studied in [19] for $N = 3, 4$. By periodicity, in Definition 3.1 we can assume that $c_1 = \dots = c_I = 0$ where $I = \min\{n, N\}$. We have:

Proposition 3.2. *f is quasiconvex if and only if it is N -wave quasiconvex for all N .*

Proof. We prove that if f is N -wave quasiconvex at zero it is quasiconvex at zero, as the converse is clear. We rely on the following standard fact: for $\varphi \in C^\infty(\mathbb{T}^n, \mathbb{R}^m)$ there is a sequence φ_j of the form (3.2) which converges to φ strongly in $W^{1,\infty}(\mathbb{T}^n, \mathbb{R}^m)$. For a quantitative version of this fact when $m = 1$ see e.g. [12], although there the authors take several different functions s_i , for $i = 1, \dots, N$, instead of a fixed sawtooth function; regardless, any s_i can be approximated by scaled and translated copies of s . The general case $m > 1$ follows by straightforward arguments and we omit it.

Let v_j be the gradient Young measure generated by the deformation φ_j ; by assumption,

$$f(0) = f(\bar{v}_j) \leq \langle v_j, f \rangle = \int_{\mathbb{T}^n} f(D\varphi_j) \, dx.$$

Since $\varphi_j \rightarrow \varphi$ in $W^{1,\infty}(\mathbb{T}^n, \mathbb{R}^m)$, we see that $f(0) \leq \int_{\mathbb{T}^n} f(D\varphi) \, dx$. Thus f is quasiconvex at zero. ■

The following theorem gathers several results from the literature.

Theorem 3.3. *N -wave quasiconvexity has the following properties:*

- (a) *1-wave quasiconvexity is equivalent to rank-one convexity;*
- (b) *2-wave quasiconvexity is equivalent to rank-one convexity;*
- (c) *if $m \geq 3$ and $n \geq 2$, then 3-wave quasiconvexity is different from rank-one convexity and is a nonlocal property;*
- (d) *if $m = n = 2$, then 3-wave quasiconvexity is implied by rank-one convexity.*

Proof. (a) follows straightforwardly, (b) follows by Lemma 2.1 in [36], (c) follows from the example in [37] together with an adaptation of the arguments in [26] and (d) is the main result of [36]. ■

4. Counting rank-one connections

The behaviour of gradients of maps changes dramatically from the higher dimensional to the planar case [11, 18, 24]. One of the basic explanations for this difference is that the relative size of the cone

$$\Lambda \equiv \{A \in \mathbb{R}^{m \times 2} : \text{rank } A \leq 1\}$$

is much larger when $m = 2$ than when $m \geq 3$: for instance, it separates the matrix space into two components in the former case.

The previous insight is also relevant towards the goal of understanding the behaviour of the particular deformations of Section 3. In fact, the proof of Theorem 3.3 (d) in [36] also explores the fact that Λ is large: using arguments somewhat in the spirit of [38], the abundance of rank-one connections is used to build complicated laminates supported in the 3-cube $\{X_\varepsilon\}_{\varepsilon \in \{-1,1\}^3}$. In view of Proposition 3.2 it is natural to ponder what can be said for a general $N > 3$.

In this section our goal is to roughly quantify the number of rank-one connections between points in the lamination hull of the N -cube. Our observations are merely heuristic, i.e., we do not provide any proofs, and they are the consequence of analysing thousands of computer-generated random configurations.

For a given choice of matrices X_ε as in (3.4), let us write

$$K_N \equiv \{X_\varepsilon : \varepsilon \in \{-1, 1\}^N\} \subset \mathbb{R}^{m \times 2}, \quad Q_N \equiv [-1, 1]^N \subset \mathbb{R}^N.$$

We can visualise K_N as the vertices of the N -cube Q_N by considering the map $X_\varepsilon \mapsto \varepsilon$. Note, however, that for $N > 2m$ the map $\varepsilon \mapsto X_\varepsilon$ cannot be an embedding.

Let us denote by $K_N^{lc,i}$ the usual i -th lamination convex hull of K_N , see [31] for the definition. Since the edges of the cube correspond to rank-one segments, it is clear that, under the above identification, $K_N^{lc,i}$ contains the i -skeleton of Q_N : for instance, $K_N^{lc,1}$ contains the edges of the cube, $K_N^{lc,2}$ contains the faces, and so on.

We say that X_ε and $X_{\varepsilon'}$ are neighbours if ε and ε' are adjacent vertices in Q_N . Generically, each vertex X_ε is rank-one connected only to its N neighbours and thus $K_N^{lc,1}$ is in fact the 1-skeleton of the N -cube, i.e., it consists of the vertices and the edges $E_N \equiv K_N^{lc,1} \setminus K_N$; note that each edge is an open segment parallel to a rank-one line.

We now want to compare $K_N^{lc,2}$ with the 2-skeleton of the N -cube. We call a rank-one connection *trivial* if it exists in the 2-skeleton of the N -cube. A vertex is trivially connected to the N edges that have that vertex as one of their endpoints. An edge, which we write in the form $\{(\varepsilon_1, \dots, \varepsilon_{i-1}, t, \varepsilon_{i+1}, \dots, \varepsilon_N) : t \in [0, 1]\}$, is trivially connected to the $N - 1$ edges that arise by flipping the sign of one of the ε_j , for $j \neq i$.

Associated to a fixed deformation, we consider two vectors, one with length 2^N and the other with length $N2^{N-1}$. In each of these vectors, the i -th entry represents the number of non-trivial edges to which the i -th vertex, respectively the i -th edge, is rank-one connected. We calculate the mean deviation of each of these vectors. Finally, sampling randomly thousands of deformations, we get approximate values for the average number of connections, see Tables 1 and 2.

	$m = 2$	$m = 3$
$N = 3$	0.95 (0.47)	0 (0)
$N = 4$	4.79 (1.57)	0 (0)
$N = 5$	15.59 (3.65)	0 (0)
$N = 6$	41.70 (8.31)	0 (0)

Table 1. Average number (and average mean deviation) of the number of non-trivial edges to which a vertex is rank-one connected.

Remark 4.1. We would like to make a few points concerning Tables 1 and 2:

- (a) The values obtained should be understood in a probabilistic sense: it is not true that, when $m = 3$, there are never non-trivial connections. In fact, if we randomise vectors $a_i \in (\mathbb{Z} \cap [-L, L])^3, n_i \in (\mathbb{Z} \cap [-L, L])^2$ with L a small number, say $L = 5$, then we find non-trivial rank-one connections in many of the corresponding configurations.

	$m = 2$	$m = 3$
$N = 3$	2.90 (0.63)	0 (0)
$N = 4$	12.50 (2.39)	0 (0)
$N = 5$	36.78 (7.06)	0 (0)
$N = 6$	92.17 (18.28)	0 (0)

Table 2. Average number (and average mean deviation) of the number of non-trivial edges to which an edge is rank-one connected.

- (b) The low average mean deviations in the tables show that the connections are not concentrated in a few vertices or edges; see also Figure 1.
- (c) When $m = 2$, an increase in N also increases the number of connections dramatically. Thus, although the set K_N becomes exponentially more complicated as N increases, the geometry of its rank-one lines also becomes much richer.

Remark 4.2. Rank-one lines are very fragile: even if sometimes rank-one connections exist, they are easily destroyed by small perturbations [21]. It is therefore more appropriate to consider the rank-one convex hull, which is often much larger than the lamination convex hull, albeit it is also much more difficult to calculate.

What we find the most remarkable about Tables 1 and 2 is not the fact that there are almost no rank-one connections when $m = 3$ but rather that there are so many connections when $m = 2$. Thus, in low-dimensions, simple lamination seems to be a viable option to produce very complex gradients. We believe that Tables 1 and 2 can be taken as partial evidence towards a positive answer to Question 1.1 when $m = n = 2$.




Figure 1. Depiction of a “typical” configuration when $N = 3$ and $m = 2$, with an average of 1 vertex-edge connection per vertex and 3 edge-edge connections per edge. Each line denotes the existence of at least one point in the edge which is rank-one connected.

5. Numerical search for counterexamples to Morrey’s problem

In this section we report on numerical experiments which bring together Sections 2 and 3. Our goal was to find numerical evidence towards a resolution of Question 1.1.

We set $n = 2$ and run the following algorithm:

Algorithm 5.1. Fix $L, N \in \mathbb{N}$, with L odd and sufficiently large, and a threshold $\gamma \in [0, 1]$. Set $\mathcal{G}_L \equiv \frac{1}{L}\mathbb{Z}^{2m} \cap [-1, 1]^{2m}$ and choose a finite set of directions \mathcal{D} consisting of rank-one matrices which are in \mathcal{G}_L . Then:

1. Randomly generate a set of directions, $(n_i)_{i=1}^N$ in $([-L, L] \cap \mathbb{Z})^2$. Check that n_i and n_j are linearly independent for $i \neq j$; if not, repeat the previous instruction.
2. For each $(n_i)_{i=1}^N$, randomly generate a set of phases, $(c_i)_{i=1}^N \in \mathbb{R}$. The set $(n_i, c_i)_{i=1}^N$ determines the weights, $(\nu_\varepsilon)_\varepsilon$, at each point in the support of the measure, see (3.4).
3. Randomly generate a set of vectors $(a_i)_{i=1}^N$ in $(\frac{1}{L}\mathbb{Z} \cap [-1, 1])^m$. Check that $a_i \neq 0$ for all i and that the matrices X_ε where the measure is supported, defined in (3.4) in terms of $(a_i, n_i)_{i=1}^N$, are in \mathcal{G}_L ; if not, repeat the previous instruction.

Repeat Step 1 a number M_n of times; for each of those, repeat Step 2 M_c times; and for each $(n_i, c_i)_{i=1}^N$, generate M_a different sets $(a_i)_{i=1}^N$ by 3. We thus obtain $M_v \equiv M_n \times M_c \times M_a$ sets $(a_i, n_i, c_i)_{i=1}^N \in \mathbb{Z}^m \times \mathbb{Z}^2 \times \mathbb{R}$, each of which defines a measure ν supported on \mathcal{G}_L , see (3.3). Then, for each such ν , we execute the following:

4. Randomly generate vectors in $g \in [-1, 1]^{2N}$ and, for each such vector, define a function f as in (2.2).
5. Apply the Kohn–Strang algorithm, as described in Algorithm 2.3, to calculate the approximation $f_{L, \mathcal{D}}^{rc}(0)$ of $f^{rc}(0)$.
6. Check whether $f_{L, \mathcal{D}}^{rc}(0) - \langle \nu, g \rangle > \gamma$. If so, pick another measure of those generated in Steps 1–3 and go back to Step 4. If not, and if this step has not been performed more than M_g times, using the same measure ν , go back to Step 4.

The measure ν is *suspicious* if it seems that Jensen’s inequality fails, i.e., if at least one g generated in Step 4 is such that $f_{L, \mathcal{D}}^{rc}(0) - \langle \nu, g \rangle > 0$. Suspicious measures are further examined:

7. For each suspicious pair (ν, g) , make the changes $(L, \mathcal{D}) \mapsto (L', \mathcal{D}')$, where $L' = 2L - 1$ and $\#\mathcal{D} \leq \#\mathcal{D}'$, and rerun Step 5. Repeat the previous instruction as needed.

Remark 5.1. Note that the the parameter γ ensures that, in Step 4, one keeps looking for g ’s until one finds a sufficiently suspicious measure; we typically took $\gamma = 0.1$ and we note that in Šverak’s example $f^{rc}(0) - \langle \nu, f \rangle \approx 1/4$. In fact, suppose that $0 < f_{L, \mathcal{D}}^{rc}(0) - \langle \nu, g \rangle \ll 1$; when refining the approximation of $f^{rc}(0)$ as in Step 7 it is likely that one finds $f_{L', \mathcal{D}'}(0) - \langle \nu, g \rangle < 0$ and indeed this has often happened in our calculations.

We implemented Steps 1 and 2 of Algorithm 5.1, which determine the weights in the measure (3.3), in Mathematica, as it is well suited to computing ν_ε as given by (3.4). We note that, due to the complexity of this computation, we were unable to apply our algorithm to look for counterexamples with $N \geq 6$. Moreover, for $N = 3$, it follows from

the work of Sebestyén–Székelyhidi [36] that the admissible weights form a line segment in \mathbb{R}^8 , so it is enough to look for counterexamples at the endpoints. Only for $N = 4, 5$ do we, *a priori*, actually require $N_n \times N_c$ to be large in order to have a good sampling of the parameter space.

The bulk of Algorithm 5.1, i.e., Steps 3–7, was implemented in the C programming language. Our implementation is quite fast for $m = 2$: for instance with $L = 25$, $\#\mathcal{D} = 64$ and $M_g = 50$, it typically takes around 3 minutes to perform Steps 4–5, even when Step 5 is performed the maximum number of times. For $m > 2$, the algorithm has a very large computational cost: for instance, with $L = 19$ and $\#\mathcal{D} = 168$, it typically takes around 13 hours to perform Steps 4–5 a number $M_g = 50$ times. We remark that in this case the number of points in the grid is approximately 47×10^6 .

5.1. The case $m = 2$

For $m = 2$ we considered deformations given by sums of N plane waves with $N = 3, 4, 5$.

For $N = 3$, we verified numerically the analytical result of [36]. Using a gridsize of $L = 25$, we selected a total of 210 measures and randomised 50 different functions g , which were rank-one convexified using $\#\mathcal{D} = 64$ rank-one directions, see Table 3. About 5% of the pairs (v, g) were found to be suspicious, though none above the threshold $\gamma = 0.1$. Upon rescaling the grid to $L' = 49$ and increasing the set of rank-one directions to $\#\mathcal{D}' = 256$, all but one of these pairs was shown to satisfy Jensen's inequality; the remaining potential counterexample was ruled out by rescaling the grid again to $L' = 97$ and increasing $\#\mathcal{D}' = 784$.

It is for $N = 4, 5$, where Question 1.1 is open, that our results are most interesting. As the structure of the weights in these cases is unknown, we consider a much larger set of measures, around 1000, in our numerical tests; we have also increased the maximum number of functions g to test to 100×2^N , see Table 3. We have found that, when compared to a run for $N = 3$ with the same L and \mathcal{D} , in the case $N = 4, 5$ there is a drastic decrease in the percentage of suspicious measures initially flagged by Algorithm 5.1: when using a gridsize of $L = 25$ and $\#\mathcal{D} = 64$ rank-one directions, for example, only 0.06% of the pairs (v, g) are found suspicious when $N = 4$ and none are flagged in this way when $N = 5$. From the point of view of our algorithm, Jensen's inequality is clearly easier to verify as N increases, at least within the range of N we test, which could be explained by the increase in size of the 2nd lamination convex hull, c.f. Section 4. None of the pairs (v, g) flagged as suspicious was found to be a counterexample after rescaling the grid to $L' = 49$ and increasing the set of rank-one directions to $\#\mathcal{D}' = 256$. We also tested configurations generated randomly in finer grids, having obtained identical results to the case $L = 25$.

To summarize: after testing thousands of randomly generated measures and hundreds of randomly generated functions, we have not found any counterexamples to Question 1.1.

5.2. The case $m = 3$

For $m = 3$ and $N = 3$, let us consider directions (n_1, n_2, n_3) which are non-degenerate in the sense that, for some choice of phases, there is $\varepsilon \in \{-1, 1\}^3$ with $v_\varepsilon \neq 1/8$. It follows from the example in [37] that, with probability one, any such measure is a counterexample to Question 1.1. Due to the high computational cost of Algorithm 5.1 for $m = 3$, which

N	M_n	M_c	M_a	M_v	M_g
3	7	1	30	210	50
4	7	7	20	980	160
5	7	7	20	980	320

Table 3. Parameter space sampled in numerical experiments with $m = 2$.

severely limits our ability to explore the parameter space, we decided to focus only on $N = 3$ and attempt to recover these analytic results.

With a grid of size $L = 19$ and $\#\mathcal{D} = 168$, over the course of two weeks, we tested around 30 measures corresponding to 3-wave deformations. All but one measure was found to be suspicious and around 90% of the measures were found to be sufficiently suspicious, in the sense that there was one rank-one convexified function for which Jensen’s inequality failed by a margin superior to the threshold of $\gamma = 0.1$. We were unable to verify how many of our candidate counterexamples would survive after rescaling the grid (Step 7 of Algorithm 5.1), as those computations would take around a month per measure. However, these results are in agreement with what is known analytically for $N = 3$, further validating our implementation of Algorithm 5.1.

5.3. The case $m > 3$

It is interesting to consider Grabovsky’s example [13] of a rank-one convex, non quasiconvex function $G: \mathbb{R}^{8 \times 2} \rightarrow \mathbb{R}$. G is quasiconvex at zero, although not at the point $\text{Id}_{\mathbb{H}^2} \equiv e_1 \otimes e_1 + e_5 \otimes e_2$. However, the paper [13] does not give an explicit deformation falsifying the quasiconvexity inequality (3.1); the deformation is only obtained indirectly through the variational principle for the effective tensor in periodic homogenization.

It would be interesting to find an explicit deformation falsifying the quasiconvexity inequality with respect to G and, in particular, to find the smallest value of N for which G is no longer N -wave quasiconvex. In an attempt to do so, we randomly generated deformations according to steps 1–3 of Algorithm 5.1. After testing hundreds of such deformations, and finding no counter-example to Jensen’s inequality, we are led to suppose that G is N -wave quasiconvex for $N \leq 5$. As mentioned above, the case $N \geq 6$ is very computationally demanding.

We also note that there is a curious similarity between the plane-wave expansions of Section 3 and [13], equation (2.18).

Acknowledgements. A. Guerra would like to thank Vladimír Šverák for a very interesting conversation about Morrey’s problem and, in particular, for the suggestion of conducting a numerical experiment roughly along the lines of the one described in Section 5. A. Guerra would also like to thank Jan Kristensen for countless hours of discussion about the problems addressed here. We also thank the anonymous referee for helpful comments.

Funding. André Guerra was sponsored by EP/L015811/1. Rita Teixeira da Costa was sponsored by EP/L016516/1.

References

- [1] Angulo, P., Faraco, D. and García-Gutiérrez, C.: The $2 + 1$ convex hull of a finite set. Preprint 2020, [arXiv:1806.08447](https://arxiv.org/abs/1806.08447).
- [2] Ball, J. M.: Convexity conditions and existence theorems in nonlinear elasticity. *Arch. Rational Mech. Anal.* **63** (1976/77), no. 4, 337–403.
- [3] Ball, J. M.: Sets of gradients with no rank-one connections. *J. Math. Pures Appl.* (9) **69** (1990), no. 3, 241–259.
- [4] Ball, J. M., Kirchheim, B. and Kristensen, J.: Regularity of quasiconvex envelopes. *Calc. Var. Partial Differential Equations* **11** (2000), no. 4, 333–359.
- [5] Bartels, S.: Linear convergence in the approximation of rank-one convex envelopes. *M2AN Math. Model. Numer. Anal.* **38** (2004), no. 5, 811–820.
- [6] Dacorogna, B. *Direct methods in the calculus of variations*. Second edition. Applied Mathematical Sciences 78, Springer, New York, 2008.
- [7] Dacorogna, B., Douchet, J., Gangbo, W. and Rappaz, J.: Some examples of rank one convex functions in dimension two. *Proc. Roy. Soc. Edinburgh Sect. A* **114** (1990), no. 1-2, 135–150.
- [8] Dacorogna, B. and Haeberly, J.-P.: Some numerical methods for the study of the convexity notions arising in the calculus of variations. *RAIRO Modél. Math. Anal. Numér.* **32** (1998), no. 2, 153–175.
- [9] Dolzmann, G. Numerical computation of rank-one convex envelopes. *SIAM J. Numer. Anal.* **36** (1999), no. 5, 1621–1635.
- [10] Dolzmann, G. and Walkington, N. J.: Estimates for numerical approximations of rank one convex envelopes. *Numer. Math.* **85** (2000), no. 4, 647–663.
- [11] Faraco, D. and Székelyhidi, L.: Tartar’s conjecture and localization of the quasiconvex hull in $\mathbb{R}^{2 \times 2}$. *Acta Math.* **200** (2008), no. 2, 279–305.
- [12] Gordon, Y., Maiorov, V., Meyer, M. and Reisner, S.: On the best approximation by ridge functions in the uniform norm. *Constr. Approx.* **18** (2001), no. 1, 61–85.
- [13] Grabovsky, Y.: From microstructure-independent formulas for composite materials to rank-one convex, non-quasiconvex functions. *Arch. Ration. Mech. Anal.* **227** (2018), no. 2, 607–636.
- [14] Gremaud, P.-A.: Numerical optimization and quasiconvexity. *European J. Appl. Math.* **6** (1995), no. 1, 69–82.
- [15] Guerra, A.: Extremal rank-one convex integrands and a conjecture of Šverák. *Calc. Var. Partial Differential Equations* **58** (2019), no. 6, Paper no. 201, 19 pp.
- [16] Harris, T. L. J., Kirchheim, B. and Lin, C.-C.: Two-by-two upper triangular matrices and Morrey’s conjecture. *Calc. Var. Partial Differential Equations* **57** (2018), no. 3, Paper no. 73, 12 pp.
- [17] Iwaniec, T.: Nonlinear Cauchy–Riemann operators in \mathbb{R}^n . *Trans. Am. Math. Soc.* **354** (2002), no. 5, 1961–1995.
- [18] Iwaniec, T., Verchota, G. C. and Vogel, A. L.: The failure of rank-one connections. *Arch. Ration. Mech. Anal.* **163** (2002), no. 2, 125–169.
- [19] Kałamajska, A.: On new geometric conditions for some weakly lower semicontinuous functionals with applications to the rank-one conjecture of Morrey. *Proc. Roy. Soc. Edinburgh Sect. A* **133** (2003), no. 6, 1361–1377.

- [20] Kinderlehrer, D. and Pedregal, P.: Characterizations of Young measures generated by gradients. *Arch. Rational Mech. Anal.* **115** (1991), no. 4, 329–365.
- [21] Kirchheim, B.: *Geometry and rigidity of microstructures*. Diss. habilitation thesis, Universität Leipzig, 2001.
- [22] Kirchheim, B. and Kristensen, J.: On rank one convex functions that are homogeneous of degree one. *Arch. Ration. Mech. Anal.* **221** (2016), no. 1, 527–558.
- [23] Kirchheim, B., Müller, S. and Šverák, V.: Studying nonlinear pde by geometry in matrix space. In *Geometric analysis and nonlinear partial differential equations*, 347–395. Springer, Berlin, 2003.
- [24] Kirchheim, B. and Székelyhidi, L.: On the gradient set of Lipschitz maps. *J. Reine Angew. Math.* **625** (2008), 215–229.
- [25] Kohn, R. V. and Strang, G.: Optimal design and relaxation of variational problems, II. *Comm. Pure Appl. Math.* **39** (1986), no. 2, 139–182.
- [26] Kristensen, J.: On the non-locality of quasiconvexity. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **16** (1999), no. 1, 1–13.
- [27] Milton, G. W.: *The theory of composites*. Cambridge Monographs on Applied and Computational Mathematics 6, Cambridge University Press, Cambridge, 2002.
- [28] Morrey, C. B.: Quasi-convexity and lower semicontinuity of multiple integrals. *Pacific J. Math.* **2** (1952), 25–53.
- [29] Morrey, C. B.: *Multiple integrals in the calculus of variations*. Grundlehren der Mathematischen Wissenschaften 130, Springer, New York, 1966.
- [30] Müller, S.: Rank-one convexity implies quasiconvexity on diagonal matrices. *Internat. Math. Res. Notices* (1999), no. 20, 1087–1095.
- [31] Müller, S.: Variational models for microstructure and phase transitions. In *Calculus of variations and geometric evolution problems (Cetraro, 1996)*, 85–210. Lecture Notes in Math. 1713, Fond. CIME/CIME Found. Subser., Springer, Berlin, 1999.
- [32] Oberman, A. M. and Ruan, Y.: A partial differential equation for the rank one convex envelope. *Arch. Ration. Mech. Anal.* **224** (2017), no. 3, 955–984.
- [33] Pedregal, P.: Laminates and microstructure. *European J. Appl. Math.* **4** (1993), no. 2, 121–149.
- [34] Pedregal, P. and Šverák, V.: A note on quasiconvexity and rank-one convexity for 2×2 matrices. *J. Convex Anal.* **5** (1998), no. 1, 107–117.
- [35] Pinkus, A.: *Ridge functions*. Cambridge Tracts in Mathematics 205, Cambridge University Press, Cambridge, 2015.
- [36] Sebestyén, G. and Székelyhidi Jr., L.: Laminates supported on cubes. *J. Convex Anal.* **24** (2017), no. 4, 1217–1237.
- [37] Šverák, V.: Rank-one convexity does not imply quasiconvexity. *Proc. Roy. Soc. Edinburgh Sect. A* **120** (1992), no. 1-2, 185–189.
- [38] Székelyhidi, L.: Rank-one convex hulls in $\mathbb{R}^{2 \times 2}$. *Calc. Var. Partial Differential Equations* **22** (2005), no. 3, 253–281.
- [39] Tartar, L.: Compensated compactness and applications to partial differential equations. In *Nonlinear analysis and mechanics: Heriot-Watt Symposium, Vol. IV*, 136–212. Res. Notes in Math. 39, Pitman, Boston, Mass.-London, 1979.

- [40] Zhang, K.: Compensated convexity and its applications. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **25** (2008), no. 4, 743–771.

Received May 27, 2020; revised October 12, 2020. Published online June 22, 2021.

André Guerra

University of Oxford, Andrew Wiles Building, Woodstock Rd, Oxford OX2 6GG,
United Kingdom;
guerra@maths.ox.ac.uk

Rita Teixeira da Costa

Department of Pure Mathematics and Mathematical Statistics, University of Cambridge,
Wilberforce Road, Cambridge CB3 0WA, United Kingdom;
rita.t.costa@dpmms.cam.ac.uk