



Minimizing cones for fractional capillarity problems

Serena Dipierro, Francesco Maggi and Enrico Valdinoci

Abstract. We consider a fractional version of Gauß capillarity energy. A suitable extension problem is introduced to derive a boundary monotonicity formula for local minimizers of this fractional capillarity energy. As a consequence, blow-up limits of local minimizers are shown to subsequentially converge to minimizing cones. Finally, we show that in the planar case there is only one possible fractional minimizing cone, the one determined by the fractional version of Young's law.

1. Introduction

In this article we consider local minimizers in the fractional capillarity model introduced in [9], analyze their blow-up limits at boundary points, show, by means of a new monotonicity formula, that these blow-up limits are cones, and give a complete characterization of such cones in the planar case.

In the classical capillarity model of Gauß, see [8], one studies equilibrium configurations of liquid droplets E in a container $\omega \subset \mathbb{R}^n$, $n \geq 2$, by looking at (volume-constrained) local minimizers of the (dimensionally re-normalized) surface tension energy

$$\mathcal{H}^{n-1}(\omega \cap \partial E) + \sigma \mathcal{H}^{n-1}(\partial\omega \cap \partial E),$$

where $\sigma \in (-1, 1)$ is the (constant) *relative adhesion coefficient* determined by the physical properties of the liquid and of the walls of the container. In the model introduced in [9], see (1.1) below, the liquid-air surface energy term $\mathcal{H}^{n-1}(\omega \cap \partial E)$ is replaced by the nonlocal interaction between points $x \in E$ and $y \in \omega \setminus E$; while the liquid-solid surface energy term $\mathcal{H}^{n-1}(\partial\omega \cap \partial E)$ is replaced by the nonlocal interaction between points $x \in E$ and $y \notin \omega$. These nonlocal interactions are measured by the singular fractional kernel $|x - y|^{-(n+s)}$, $s \in (0, 1)$: as $s \rightarrow 1^-$, they are increasingly concentrated, respectively, at points x and y near $\omega \cap \partial E$ and $\partial\omega \cap \partial E$. For this reason, *the fractional capillarity model provides a nonlocal approximation of the Gauß capillarity model in the limit $s \rightarrow 1^-$.*

Mathematics Subject Classification (2020): 76B45, 49Q15, 35R11.

Keywords: Capillarity problems, nonlocal perimeter, extension methods, monotonicity formulae, classification results.

This happens also at the level of the classical equilibrium conditions expressed by the constancy of the mean curvature of $\omega \cap \partial E$ and by the contact angle condition between the liquid-air interface and the walls of the container, valid along $\partial\omega \cap \overline{\omega \cap \partial E}$, and known as *Young's law*. The validity of a *fractional Young law* (see (1.10) below) for sufficiently regular local minimizers of the fractional capillarity energy has been proved in [9], while its precise asymptotics in the limits $s \rightarrow 1^-$ and $s \rightarrow 0^+$ have been presented in [5]. The existence of minimizers in the fractional capillarity model is also addressed in [9]. It is an open problem to understand if these minimizers are regular up to the boundary of the container ω , and thus to confirm the validity of the fractional Young law in a pointwise sense. In this paper we take two important steps in what is a general and well-established strategy for attacking similar questions in geometric variational problems.

Our first result (given in Corollary 1.3) is that blow-up limits of local minimizers sub-sequentially converge to cones (which, in turn, are also local minimizers). This result relies on a new monotonicity formula for the fractional capillarity energy (see Theorem 1.2) and on an equivalence result with a suitable “capillarity adaptation” of the Caffarelli–Silvestre extension problem (given in Proposition 1.1).

Our second result (stated in Theorem 1.4) is a classification theorem for fractional minimizing cones in the half-plane: more precisely, we will show that the only possible fractional minimizing cones in ambient dimension 2 are angular sectors satisfying the fractional version of Young's law.

While the first result about the blow-up limits (as well as the extension theorem and the monotonicity formula used in its proof) is valid in any dimension, the second result about classification of cones is only proved in dimension 2, due to suitable energy estimates that would not be valid in higher dimensions. It is an interesting open problem, which is also open for interior singularities for arbitrary values of $s \in (0, 1)$, to understand if similar rigidity results for minimizing cones are valid in higher dimensions. The other main open problem is that of obtaining a boundary regularity criterion comparable to the one available in the interior [1], and analogous to the ones developed in the classical case to validate Young's law, see [3, 4] and the references therein.

The precise mathematical setting in which we work is the following. Given $s \in (0, 1)$ and two disjoint sets $A, B \subseteq \mathbb{R}^n$, we define the *fractional interaction* between A and B as

$$\mathcal{I}_s(A, B) := \iint_{A \times B} \frac{dx dy}{|x - y|^{n+s}}.$$

Then, given $E \subseteq \omega \subseteq \mathbb{R}^n$ and $\sigma \in (-1, 1)$, we define the *fractional capillarity energy* of E in ω as

$$(1.1) \quad \mathcal{C}_{s,\sigma}(E, \omega) := \mathcal{I}_s(E, E^c \omega) + \sigma \mathcal{I}_s(E, \omega^c).$$

Here above and in the rest of this paper, we use the superscript “ c ” to denote the complementary set in \mathbb{R}^n . Also, given two sets $A, B \subseteq \mathbb{R}^n$ we use the short notation $AB := A \cap B$ (in this way, the notation $E^c \omega$ is short for $(\mathbb{R}^n \setminus E) \cap \omega$). Furthermore, the Lebesgue measure of a set $F \subseteq \mathbb{R}^n$ will be denoted by $|F|$.

We consider the half-space

$$H := \{x = (x_1, \dots, x_n) \in \mathbb{R}^n \text{ such that } x_n > 0\},$$

and, given $R > 0$, we denote by $B_R \subset \mathbb{R}^n$ the n -dimensional Euclidean ball of radius R centered at the origin. In this article, we are interested in local minimizers of the fractional capillarity energy in H . Briefly, we say that $E \subseteq H$ is a *local minimizer in H* if, for every $R > 0$, we have that $\mathcal{J}_s(EB_R, E^c B_R) < +\infty$ and

$$(1.2) \quad \begin{aligned} \mathcal{J}_s(EB_R, E^c H) + \mathcal{J}_s(EB_R^c, E^c B_R H) + \sigma \mathcal{J}_s(EB_R, H^c) \\ \leq \mathcal{J}_s(FB_R, F^c H) + \mathcal{J}_s(FB_R^c, F^c B_R H) + \sigma \mathcal{J}_s(FB_R, H^c), \end{aligned}$$

for every $F \subseteq H$ such that $F \setminus B_R = E \setminus B_R$. In particular, blow-up limits of minimizers in the fractional capillarity problem in bounded domains with smooth boundary are local minimizers in H , see Theorem A.2 in [9].

In order to exploit extension methods (see e.g. [2]), for any $(x, t) \in \mathbb{R}_+^{n+1} := \mathbb{R}^n \times (0, +\infty)$, it is convenient to introduce the fractional Poisson kernel

$$\mathbf{P}_s(x, t) := C_{n,s} \frac{t^s}{(|x|^2 + t^2)^{(n+s)/2}},$$

where $C_{n,s} > 0$ is a normalizing constant (which, from now on, will be omitted) such that

$$\int_{\mathbb{R}^n} \mathbf{P}_s(x, t) dx = 1, \quad \text{for all } t > 0.$$

Given $u \in L^\infty(\mathbb{R}^n)$, we also denote the s -extension of u by

$$\mathbf{E}_u(x, t) := \int_{\mathbb{R}^n} u(y) \mathbf{P}_s(x - y, t) dy, \quad \text{for all } (x, t) \in \mathbb{R}_+^{n+1}.$$

The relevance of this notion of s -extension for our problem lies in the fact that the property of E being a local minimizer in H for the fractional capillarity energy $\mathcal{C}_{s,\sigma}$ is equivalent to the property of \mathbf{E}_u being a local minimizer of a Dirichlet-type energy $\mathcal{F}_{s,\sigma}$ that we are now going to introduce. Indeed, let $X = (x, t) \in \mathbb{R}_+^{n+1}$. As customary, given $E \subseteq \mathbb{R}^n$, we denote by $\chi_E: \mathbb{R}^n \rightarrow \{0, 1\}$ the characteristic function of E . If $u = \chi_E$, we also write $\mathbf{E}_E := \mathbf{E}_{\chi_E}$. In addition, given $\Omega \subseteq \mathbb{R}^{n+1}$ with $\omega := \Omega \cap \{t = 0\}$ and $\Omega^+ := \Omega \cap \mathbb{R}_+^{n+1}$, and $U: \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ with $u(x) := U(x, 0)$, we define the energy

$$(1.3) \quad \mathcal{F}_{s,\sigma}(U, \Omega) := \int_{\Omega^+} t^{1-s} |\nabla U(X)|^2 dX + (\sigma - 1) \iint_{\omega \times H^c} \frac{u(x)}{|x - y|^{n+s}} dx dy.$$

Given $K \subseteq \mathbb{R}^{n+1}$ and $\eta > 0$, we also set

$$(1.4) \quad K_\eta := \{X \in \mathbb{R}^{n+1} \text{ such that } \text{dist}(X, K) < \eta\}.$$

Then, we have the following extension result.

Proposition 1.1. *Let $E \subseteq H$ be such that $\mathcal{J}_s(EB_R, E^c B_R) < +\infty$ for every $R > 0$. The following statements are equivalent:*

- (i) E is a local minimizer in H .

(ii) For all $R > 0$ and all bounded, Lipschitz domains $\Omega \subset \mathbb{R}^{n+1}$ with

$$(1.5) \quad \Omega \cap \{t = 0\} = B_R,$$

we have that

$$(1.6) \quad \mathcal{F}_{s,\sigma}(\mathbf{E}_E, \Omega) \leq \mathcal{F}_{s,\sigma}(U, \Omega)$$

for all $U: \mathbb{R}_+^{n+1} \rightarrow \mathbb{R}$ such that $U(x, 0) = \chi_F(x)$ for all $x \in \mathbb{R}^n$, for some $F \subseteq H$, with $FB_{R-\eta}^c = EB_{R-\eta}^c$, and $U(X) = \mathbf{E}_E(X)$ for all $X \in (\partial\Omega)_\eta \cap \mathbb{R}_+^{n+1}$, for some $\eta \in (0, R)$.

The previous result can be seen as the natural counterpart, in the setting of fractional capillarity problems, of several extension theorems for the fractional Laplacian, for fractional minimal surfaces and, more generally, for nonlocal free boundary problems, see e.g. [1, 2, 6, 10, 13, 14]. Among the many applications of the powerful tool provided by extension results is the possibility of obtaining convenient monotonicity formulae: actually, to the best of our knowledge, all the monotonicity formulae involving nonlocal operators rely on identifying appropriate local extension problems methods.

In the setting considered in this paper, we will exploit Proposition 1.1 to obtain a monotonicity formula that we now describe in detail. We denote by $\mathcal{B}_R \subset \mathbb{R}^{n+1}$ the $(n+1)$ -dimensional Euclidean ball of radius R . For $E \subseteq \omega$ and $r > 0$, we define

$$\Phi_E(r) := r^{s-n} \mathcal{F}_{s,\sigma}(\mathbf{E}_E, \mathcal{B}_r).$$

We observe that the above function is scale invariant, in the sense that

$$(1.7) \quad \Phi_E(r) = \Phi_{E_{r/\rho}}(\rho),$$

where

$$(1.8) \quad E_r := \frac{E}{r} = \left\{ \frac{x}{r}, x \in E \right\}.$$

In this setting, we have the following monotonicity formula.

Theorem 1.2. *Assume that $E \subseteq H$ is a local minimizer for the fractional capillarity energy in H . Then, the function $(0, +\infty) \ni r \mapsto \Phi_E(r)$ is monotone nondecreasing.*

More precisely, for every $r > 0$ we have that

$$(1.9) \quad \Phi'_E(r) \geq r^{s-n} \int_{(\partial\mathcal{B}_r) \cap \{t>0\}} t^{1-s} |\nabla_\nu \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n.$$

Furthermore, we have that Φ_E is constant if and only if E is a cone, i.e., $\tau E = E$ for all $\tau > 0$.

As a consequence of Theorem 1.2, we have that suitable blow-up limits of local minimizers of the fractional capillarity problem are cones:

Corollary 1.3. *Let $\omega \subset \mathbb{R}^n$ be a bounded open set with C^1 -boundary. Let $E \subseteq \omega$ be a minimizer of the capillarity functional in (1.1) among sets of prescribed volume contained in ω .*

Assume that $0 \in \overline{\omega \cap (\partial E)}$. Then for every vanishing sequence r_j there exists (a not relabeled) subsequence and a set $E_0 \subset \mathbb{R}^n$, such that, in the notation of (1.8), we have that $\chi_{E_{r_j}} \rightarrow \chi_{E_0}$ in $L^1_{\text{loc}}(\mathbb{R}^n)$. In addition, E_0 is a cone.

The existence of the minimizers in Corollary 1.3 (and, in fact, of a more general class of minimizers) is warranted by Proposition 1.1 in [9]. As a matter of fact, Corollary 1.3 is also valid for the “almost minimizers”, as introduced in Definition 1.5 of [9], with the same proof that we present here.

In the setting of Corollary 1.3, it is natural to consider locally minimizing cones in H (i.e., sets that are locally minimizing in H and that possess a conical structure). Interestingly, in dimension 2, we can completely characterize locally minimizing cones in H , according to the following result.

Theorem 1.4. *Let $n = 2$ and let E be a locally minimizing cone in $H = \{x_2 > 0\}$. Then, E is made of only one component and, up to a rigid motion, we have that*

$$E = \{x = (x_1, x_2) \in H \text{ such that } x_1 > x_2 \cos \vartheta\},$$

with $\vartheta \in (0, \pi)$ implicitly defined by the formula

$$(1.10) \quad 1 + \sigma = \frac{(\sin \vartheta)^s M(\vartheta, s)}{M(\pi/2, s)},$$

$$\text{where } M(\vartheta, s) := 2 \iint_{(0, \vartheta) \times (0, +\infty)} \frac{r}{(r^2 + 2r \cos t + 1)^{(2+s)/2}} dt dr.$$

Notice that (1.10) expresses the fractional Young law mentioned earlier in this introduction, which, in the limit as $s \rightarrow 1^-$ converges to the contact angle prescription given by the classical Young law. For a detailed asymptotic description of this, see [5].

To prove Theorem 1.4, we use a “translation method” introduced in [11] to prove the regularity of fractional minimizing surfaces in the plane. In our context, however, the cone is going to have a singularity at the origin, hence the notion of “regularity” has to be weakened to a suitable notion of “monotonicity”, taking inspiration by some work in [12].

The rest of this paper is devoted to the proof of the results that we have presented above. More specifically, Section 2 contains some preliminary observations relating the nonlocal surface tension energy introduced in [9] and the nonlocal perimeter functional introduced in [1]. Then, the proof of Proposition 1.1 will be given in Section 3, and the one of Theorem 1.2 in Section 4. Section 5 contains the proof of Corollary 1.3, and Section 6 the one of Theorem 1.4.

2. Capillarity and fractional perimeters

In this section, we point out some useful relations between the capillarity functional given in (1.1) and other fractional energies of geometric type. First of all, we observe that the energy functional in (1.1) can be related to the fractional perimeter introduced in [1]. Indeed, writing, for any given $F, \omega \subseteq \mathbb{R}^n$,

$$\text{Per}_s(F, \omega) := \mathcal{J}_s(F\omega, F^c\omega) + \mathcal{J}_s(F\omega, F^c\omega^c) + \mathcal{J}_s(F\omega^c, F^c\omega),$$

for every $E \subseteq \omega$ we have that

$$\mathcal{C}_{s,\sigma}(E, \omega) = \text{Per}_s(E, \omega) + (\sigma - 1) \mathcal{J}_s(E, \omega^c).$$

It is also useful to define, for all $F \subseteq H$ and all $R > 0$,

$$(2.1) \quad \text{Per}_{s,\sigma}(F, B_R) := \text{Per}_s(F, B_R H) + (\sigma - 1) \mathcal{J}_s(F B_R, H^c).$$

In this setting, we can state the local minimality condition in (1.2) in terms of the fractional perimeter as follows:

Lemma 2.1. *A set $E \subseteq H$ is a local minimizer in H if and only if, for every $R > 0$, we have that $\text{Per}_s(E, B_R H) < +\infty$ and*

$$\text{Per}_{s,\sigma}(E, B_R) \leq \text{Per}_{s,\sigma}(F, B_R)$$

for every $F \subseteq H$ such that $F \setminus B_R = E \setminus B_R$.

Proof. If $F \subseteq H$,

$$\begin{aligned} & \text{Per}_s(F, B_R H) + (\sigma - 1) \mathcal{J}_s(F B_R, H^c) \\ &= \mathcal{J}_s(F B_R H, F^c B_R H) + \mathcal{J}_s(F B_R H, F^c B_R^c H) + \mathcal{J}_s(F B_R H, F^c H^c) \\ & \quad + \mathcal{J}_s(F B_R^c H, F^c B_R H) + \mathcal{J}_s(F H^c, F^c B_R H) + (\sigma - 1) \mathcal{J}_s(F B_R, H^c) \\ &= \mathcal{J}_s(F B_R, F^c B_R H) + \mathcal{J}_s(F B_R, F^c B_R^c H) + \mathcal{J}_s(F B_R, H^c) \\ & \quad + \mathcal{J}_s(F B_R^c, F^c B_R H) + (\sigma - 1) \mathcal{J}_s(F B_R, H^c) \\ &= \mathcal{J}_s(F B_R, F^c H) + \mathcal{J}_s(F B_R^c, F^c B_R H) + \sigma \mathcal{J}_s(F B_R, H^c). \end{aligned}$$

From this, (1.2) and (2.1), the desired result plainly follows. \blacksquare

3. Extension problems and proof of Proposition 1.1

In this section, we analyze the equivalent extension problem stated in Proposition 1.1 and give a proof of it.

Proof of Proposition 1.1. First of all, we observe that, by (1.3) and (2.1), if $V: \mathbb{R}_+^{n+1} \rightarrow \mathbb{R}$ is such that $V(x, 0) = \chi_L(x)$, with $L \subseteq H$, and $\Omega \subset \mathbb{R}^{n+1}$ satisfies (1.5),

$$\begin{aligned} (3.1) \quad & \text{Per}_{s,\sigma}(L, B_R) - \mathcal{F}_{s,\sigma}(V, \Omega) \\ &= \text{Per}_s(L, B_R H) + (\sigma - 1) \mathcal{J}_s(L B_R, H^c) \\ & \quad - \int_{\Omega^+} t^{1-s} |\nabla V(X)|^2 dX - (\sigma - 1) \iint_{B_R \times H^c} \frac{\chi_L(x)}{|x - y|^{n+s}} dx dy \\ &= \text{Per}_s(L, B_R H) - \int_{\Omega^+} t^{1-s} |\nabla V(X)|^2 dX. \end{aligned}$$

We also remark that, if $F \subseteq H$, then

$$\begin{aligned}
& \text{Per}_s(F, B_R) - \text{Per}_s(F, B_R H) \\
&= \mathcal{J}_s(F B_R, F^c) + \mathcal{J}_s(F B_R^c, F^c B_R) - \mathcal{J}_s(F B_R H, F^c) - \mathcal{J}_s(F (B_R H)^c, F^c B_R H) \\
&= \mathcal{J}_s(F B_R H, F^c) + \mathcal{J}_s(F B_R^c, F^c B_R) - \mathcal{J}_s(F B_R H, F^c) - \mathcal{J}_s(F B_R^c, F^c B_R H) \\
&= \mathcal{J}_s(F B_R^c, F^c B_R H) + \mathcal{J}_s(F B_R^c, F^c B_R H^c) - \mathcal{J}_s(F B_R^c, F^c B_R H) \\
(3.2) \quad &= \mathcal{J}_s(F B_R^c, F^c B_R H^c) = \mathcal{J}_s(F B_R^c, B_R H^c).
\end{aligned}$$

We will also exploit Lemma 7.2 of [1], according to which (up to normalizing constants that we omit), given $L, M, \omega \subseteq \mathbb{R}^n$ with $\text{Per}_s(L, \omega), \text{Per}_s(M, \omega) < +\infty$ and $L\tilde{\omega}^c = M\tilde{\omega}^c$, for $\tilde{\omega} \Subset \omega$, then

$$(3.3) \quad \inf_{\Omega^+} \int t^{1-s} (|\nabla V(X)|^2 - |\nabla \mathbf{E}_M(X)|^2) dX = \text{Per}_s(L, \omega) - \text{Per}_s(M, \omega),$$

where the infimum is taken among all bounded Lipschitz domains $\Omega \subseteq \mathbb{R}^{n+1}$ with $\Omega \cap \{t = 0\} \Subset \omega$ and among all functions $V: \mathbb{R}_+^{n+1} \rightarrow \mathbb{R}$ such that $V - \mathbf{E}_M$ is compactly supported in Ω , and $V(x, 0) = \chi_L(x)$.

Now, assume that E is a local minimizer in H , and let R, Ω, η, U and F be as in the assumptions of Proposition 1.1 (ii). In the notation of (1.4), we consider the set

$$\tilde{\Omega} := \left\{ X \in \Omega \text{ such that } \text{dist}(X, \partial\Omega) \geq \frac{\eta}{2} \right\} = \Omega \setminus (\partial\Omega)_{\eta/2}.$$

By the assumptions of Proposition 1.1 (ii), we know that $U - \mathbf{E}_E$ is compactly supported in $\tilde{\Omega}$. Moreover $\tilde{\Omega} \cap \{t = 0\} \Subset \Omega \cap \{t = 0\} = B_R$. Therefore, we can exploit (3.3) with Ω there replaced by $\tilde{\Omega}$ and ω chosen to be B_R , thus obtaining

$$\begin{aligned}
\int_{\tilde{\Omega}^+} t^{1-s} (|\nabla U(X)|^2 - |\nabla \mathbf{E}_E(X)|^2) dX &= \int_{\tilde{\Omega}^+} t^{1-s} (|\nabla U(X)|^2 - |\nabla \mathbf{E}_E(X)|^2) dX \\
&\geq \text{Per}_s(F, B_R) - \text{Per}_s(E, B_R).
\end{aligned}$$

This and (3.1) give that

$$\begin{aligned}
\mathcal{F}_{s,\sigma}(\mathbf{E}_E, \Omega) - \mathcal{F}_{s,\sigma}(U, \Omega) &= \text{Per}_{s,\sigma}(E, B_R) - \text{Per}_s(E, B_R H) + \int_{\tilde{\Omega}^+} t^{1-s} |\nabla \mathbf{E}_E(X)|^2 dX \\
&\quad - \text{Per}_{s,\sigma}(F, B_R) + \text{Per}_s(F, B_R H) - \int_{\tilde{\Omega}^+} t^{1-s} |\nabla U(X)|^2 dX \\
&\leq \text{Per}_{s,\sigma}(E, B_R) - \text{Per}_{s,\sigma}(F, B_R) + \text{Per}_s(F, B_R H) \\
&\quad - \text{Per}_s(E, B_R H) - \text{Per}_s(F, B_R) + \text{Per}_s(E, B_R).
\end{aligned}$$

Consequently, recalling (3.2) and the fact that E and F coincide outside B_R ,

$$\begin{aligned}
\mathcal{F}_{s,\sigma}(\mathbf{E}_E, \Omega) - \mathcal{F}_{s,\sigma}(U, \Omega) \\
&\leq \text{Per}_{s,\sigma}(E, B_R) - \text{Per}_{s,\sigma}(F, B_R) - \mathcal{J}_s(F B_R^c, B_R H^c) + \mathcal{J}_s(E B_R^c, B_R H^c) \\
&= \text{Per}_{s,\sigma}(E, B_R) - \text{Per}_{s,\sigma}(F, B_R).
\end{aligned}$$

The locally minimizing property of E and Lemma 2.1 thereby imply that $\mathcal{F}_{s,\sigma}(\mathbf{E}_E, \Omega) - \mathcal{F}_{s,\sigma}(U, \Omega) \leq 0$, that is (1.6), as desired.

Let us now suppose that, viceversa, the claim in (1.6) holds true. Our objective is now to check that E is a local minimizer. To this end, let $F \subseteq H$ such that $F \setminus B_R = E \setminus B_R$. Also, fixed $\delta > 0$, recalling (3.3), we take a bounded Lipschitz domain $\Omega^{(\delta)} \subseteq \mathbb{R}^{n+1}$ with $\Omega^{(\delta)} \cap \{t = 0\} \Subset B_{R+1}$ and a function $V^{(\delta)}: \mathbb{R}_+^{n+1} \rightarrow \mathbb{R}$ such that $V^{(\delta)} - \mathbf{E}_E$ is compactly supported in $\Omega^{(\delta)}$, and $V^{(\delta)}(x, 0) = \chi_F(x)$, with $\Omega^{(\delta)}$ and $V^{(\delta)}$ attaining the infimum in (3.3) with $\omega := B_{R+1}$ up to an error δ , that is,

$$(3.4) \quad \int_{(\Omega^{(\delta)})^+} t^{1-s} (|\nabla V^{(\delta)}(X)|^2 - |\nabla \mathbf{E}_E(X)|^2) dX - \delta \\ \leq \text{Per}_s(F, B_{R+1}) - \text{Per}_s(E, B_{R+1}).$$

Let

$$\rho' := \sup_{x \in \Omega^{(\delta)} \cap \{t=0\}} |x| \quad \text{and} \quad \rho := \max\{R, \rho'\}.$$

By construction, we have that $\rho' \in [0, R+1)$, and thus $\rho \in [R, R+1)$. Let also $\Omega^{(\delta, \rho)} := \Omega^{(\delta)} \cup \mathcal{B}_\rho$. Then, we have that

$$(3.5) \quad \Omega^{(\delta, \rho)} \cap \{t = 0\} = B_\rho.$$

Furthermore, since $V^{(\delta)} = \mathbf{E}_E$ in $\Omega^{(\delta, \rho)} \setminus \Omega^{(\delta)}$, we have that

$$\int_{(\Omega^{(\delta, \rho)})^+} t^{1-s} (|\nabla V^{(\delta)}(X)|^2 - |\nabla \mathbf{E}_E(X)|^2) dX \\ = \int_{(\Omega^{(\delta)})^+} t^{1-s} (|\nabla V^{(\delta)}(X)|^2 - |\nabla \mathbf{E}_E(X)|^2) dX.$$

Therefore, recalling (3.4),

$$(3.6) \quad \int_{(\Omega^{(\delta, \rho)})^+} t^{1-s} (|\nabla V^{(\delta)}(X)|^2 - |\nabla \mathbf{E}_E(X)|^2) dX - \delta \\ \leq \text{Per}_s(F, B_{R+1}) - \text{Per}_s(E, B_{R+1}).$$

Moreover, in view of (3.5), we are in the position of using (1.6) (with Ω replaced by $\Omega^{(\delta, \rho)}$ and R replaced by ρ). In this way, we find that

$$\mathcal{F}_{s, \sigma}(\mathbf{E}_E, \Omega^{(\delta, \rho)}) \leq \mathcal{F}_{s, \sigma}(V^{(\delta)}, \Omega^{(\delta, \rho)}).$$

Consequently, exploiting (1.3), (3.5) and (3.6),

$$\begin{aligned} & \text{Per}_s(E, B_{R+1}) - \text{Per}_s(F, B_{R+1}) \\ & \leq \int_{(\Omega^{(\delta, \rho)})^+} t^{1-s} (|\nabla \mathbf{E}_E(X)|^2 - |\nabla V^{(\delta)}(X)|^2) dX + \delta \\ & = \mathcal{F}_{s, \sigma}(\mathbf{E}_E, \Omega^{(\delta, \rho)}) - \mathcal{F}_{s, \sigma}(V^{(\delta)}, \Omega^{(\delta, \rho)}) \\ & \quad - (\sigma - 1) \iint_{B_\rho \times H^c} \frac{\chi_E(x)}{|x - y|^{n+s}} dx dy + (\sigma - 1) \iint_{B_\rho \times H^c} \frac{\chi_F(x)}{|x - y|^{n+s}} dx dy + \delta \\ & \leq -(\sigma - 1) \left(\iint_{B_R \times H^c} \frac{\chi_E(x)}{|x - y|^{n+s}} dx dy - \iint_{B_R \times H^c} \frac{\chi_F(x)}{|x - y|^{n+s}} dx dy \right) + \delta \\ & = -(\sigma - 1) (\mathcal{I}_s(EB_R, H^c) - \mathcal{I}_s(FB_R, H^c)) + \delta. \end{aligned}$$

Hence, since

$$\begin{aligned}
 & (\text{Per}_s(E, B_{R+1}) - \text{Per}_s(F, B_{R+1})) - (\text{Per}_s(E, B_R) - \text{Per}_s(F, B_R)) \\
 &= \mathcal{J}_s(EB_{R+1}B_R^c, E^c B_{R+1}^c) + \mathcal{J}_s(EB_R^c, E^c B_{R+1}B_R^c) \\
 &\quad - \mathcal{J}_s(FB_{R+1}B_R^c, F^c B_{R+1}^c) - \mathcal{J}_s(FB_R^c, F^c B_{R+1}B_R^c) \\
 &= 0,
 \end{aligned}$$

we find that

$$\text{Per}_s(E, B_R) - \text{Per}_s(F, B_R) \leq -(\sigma - 1) (\mathcal{J}_s(EB_R, H^c) - \mathcal{J}_s(FB_R, H^c)) + \delta.$$

Then, by (2.1) and (3.2),

$$\begin{aligned}
 & \text{Per}_{s,\sigma}(E, B_R) - \text{Per}_{s,\sigma}(F, B_R) \\
 &= \text{Per}_s(E, B_R H) - \text{Per}_s(F, B_R H) + (\sigma - 1) (\mathcal{J}_s(EB_R, H^c) - \mathcal{J}_s(FB_R, H^c)) \\
 &\leq \delta + \text{Per}_s(E, B_R H) - \text{Per}_s(E, B_R) + \text{Per}_s(F, B_R) - \text{Per}_s(F, B_R H) \\
 &= \delta - \mathcal{J}_s(EB_R^c, B_R H^c) + \mathcal{J}_s(FB_R^c, B_R H^c) = \delta.
 \end{aligned}$$

Sending $\delta \searrow 0$, we thereby conclude that $\text{Per}_{s,\sigma}(E, B_R) \leq \text{Per}_{s,\sigma}(F, B_R)$. This, combined with Lemma 2.1, gives that E is a locally minimizer, as desired. \blacksquare

4. Monotonicity formula and proof of Theorem 1.2

The goal of this section is proving Theorem 1.2.

Proof of Theorem 1.2. Let

$$C_E := \left\{ x \in \mathbb{R}^n \setminus \{0\} \text{ such that } \frac{x}{|x|} \in E \right\}.$$

Given $\varepsilon > 0$, we define

$$E^{(\varepsilon)} := (((1 - \varepsilon)E) \cap B_{1-\varepsilon}) \cup (C_E \cap (B_1 \setminus B_{1-\varepsilon})) \cup (E \setminus B_1),$$

see Figure 1, and

$$U_\varepsilon(X) := \begin{cases} \mathbf{E}_E\left(\frac{X}{1-\varepsilon}\right) & \text{if } X \in \mathcal{B}_{1-\varepsilon}^+, \\ \mathbf{E}_E\left(\frac{X}{|X|}\right) & \text{if } X \in \mathcal{B}_1^+ \setminus \mathcal{B}_{1-\varepsilon}^+, \\ \mathbf{E}_E(X) & \text{if } X \in \mathbb{R}_+^{n+1} \setminus \mathcal{B}_1. \end{cases}$$

We remark that

$$U_\varepsilon(x, 0) = \left\{ \begin{array}{ll} \chi_E\left(\frac{x}{1-\varepsilon}\right) & \text{if } x \in B_{1-\varepsilon}, \\ \chi_E\left(\frac{x}{|x|}\right) & \text{if } x \in B_1 \setminus B_{1-\varepsilon}, \\ \chi_E(x) & \text{if } x \in \mathbb{R}^n \setminus B_1, \end{array} \right\} = \chi_{E^{(\varepsilon)}}(x).$$




Figure 1. The construction used in the proof of Theorem 1.2. The parts of the boundary of $E^{(\epsilon)}$ due to $C_E \cap (B_1 \setminus B_{1-\epsilon})$ are depicted by bold lines.

We also claim that

$$(4.1) \quad E^{(\epsilon)} \subseteq H.$$

Indeed, let $x \in E^{(\epsilon)}$. If $x \in B_{1-\epsilon}$, we have that $x \in (1-\epsilon)E$, and thus $x/(1-\epsilon) \in E$. Since $E \subseteq H$, we deduce that $x_n/(1-\epsilon) \geq 0$, and consequently $x_n \geq 0$, which gives that $x \in H$ in this case.

If instead $x \in B_1 \setminus B_{1-\epsilon}$, we have that $x \in C_E$, and hence $x/|x| \in E$. In this case, since $E \subseteq H$, we find that $x_n/|x| \geq 0$, and again $x \in H$. Finally, if $x \in B_1^c$, we have that $x \in E \subseteq H$, which completes the proof of (4.1).

We also observe that $U_\epsilon = \mathbf{E}_E$ outside \mathcal{B}_1 . Then, in view of (4.1), we can fix $\eta > 0$ and exploit Proposition 1.1 with $\Omega := \mathcal{B}_{1+\eta}$, $R := 1 + \eta$, $U := U_\epsilon$ and $F := E^{(\epsilon)}$. In this way, we conclude that

$$(4.2) \quad \begin{aligned} 0 &\leq \mathcal{F}_{s,\sigma}(U_\epsilon, \mathcal{B}_{1+\eta}) - \mathcal{F}_{s,\sigma}(\mathbf{E}_E, \mathcal{B}_{1+\eta}) \\ &= \int_{\mathcal{B}_{1+\eta}^+} t^{1-s} (|\nabla U_\epsilon(X)|^2 - |\nabla \mathbf{E}_E(X)|^2) dX \\ &\quad + (\sigma - 1) \left(\iint_{\mathcal{B}_{1+\eta} \times H^c} \frac{\chi_{E^{(\epsilon)}}(x)}{|x-y|^{n+s}} dx dy - \iint_{\mathcal{B}_{1+\eta} \times H^c} \frac{\chi_E(x)}{|x-y|^{n+s}} dx dy \right) \\ &= \int_{\mathcal{B}_1^+} t^{1-s} (|\nabla U_\epsilon(X)|^2 - |\nabla \mathbf{E}_E(X)|^2) dX \\ &\quad + (\sigma - 1) (\mathcal{J}_s(B_1 E^{(\epsilon)}, H^c) - \mathcal{J}_s(B_1 E, H^c)). \end{aligned}$$

We set

$$G(r) := r^{s-n} \int_{\mathcal{B}_r^+} t^{1-s} |\nabla \mathbf{E}_E(X)|^2 dX,$$

and, using the change of variable $Y = (y, \theta) := X/(1 - \varepsilon)$, we observe that

$$\begin{aligned}
& \int_{\mathcal{B}_1^+} t^{1-s} |\nabla U_\varepsilon(X)|^2 dX \\
&= \frac{1}{(1-\varepsilon)^2} \int_{\mathcal{B}_{1-\varepsilon}^+} t^{1-s} \left| \nabla \mathbf{E}_E \left(\frac{X}{1-\varepsilon} \right) \right|^2 dX \\
&\quad + \int_{\mathcal{B}_1^+ \setminus \mathcal{B}_{1-\varepsilon}^+} \frac{t^{1-s}}{|X|^2} \left(\left| \nabla \mathbf{E}_E \left(\frac{X}{|X|} \right) \right|^2 - \left| \frac{X}{|X|} \cdot \nabla \mathbf{E}_E \left(\frac{X}{|X|} \right) \right|^2 \right) dX \\
&= (1-\varepsilon)^{n-s} \int_{\mathcal{B}_1^+} \theta^{1-s} |\nabla \mathbf{E}_E(Y)|^2 dY \\
&\quad + \varepsilon \int_{(\partial \mathcal{B}_1) \cap \{t>0\}} t^{1-s} (|\nabla \mathbf{E}_E(X)|^2 - |X \cdot \nabla \mathbf{E}_E(X)|^2) d\mathcal{H}_X^n + o(\varepsilon) \\
&= (1-\varepsilon)^{n-s} G(1) + \varepsilon \int_{(\partial \mathcal{B}_1) \cap \{t>0\}} t^{1-s} |\nabla_\tau \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n + o(\varepsilon),
\end{aligned}$$

where ∇_τ denotes the tangential gradient along $\partial \mathcal{B}_1$.

Similarly,

$$\begin{aligned}
& \int_{\mathcal{B}_1^+} t^{1-s} |\nabla \mathbf{E}_E(X)|^2 dX \\
&= \int_{\mathcal{B}_{1-\varepsilon}^+} t^{1-s} |\nabla \mathbf{E}_E(X)|^2 dX + \varepsilon \int_{(\partial \mathcal{B}_1) \cap \{t>0\}} t^{1-s} |\nabla \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n + o(\varepsilon) \\
&= (1-\varepsilon)^{n-s} G(1-\varepsilon) + \varepsilon \int_{(\partial \mathcal{B}_1) \cap \{t>0\}} t^{1-s} |\nabla \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n + o(\varepsilon),
\end{aligned}$$

and accordingly,

(4.3)

$$\begin{aligned}
& \int_{\mathcal{B}_1^+} t^{1-s} (|\nabla U_\varepsilon(X)|^2 - |\nabla \mathbf{E}_E(X)|^2) dX \\
&= (1-\varepsilon)^{n-s} G(1) - (1-\varepsilon)^{n-s} G(1-\varepsilon) \\
&\quad + \varepsilon \left(\int_{(\partial \mathcal{B}_1) \cap \{t>0\}} t^{1-s} |\nabla_\tau \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n - \int_{(\partial \mathcal{B}_1) \cap \{t>0\}} t^{1-s} |\nabla \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n \right) + o(\varepsilon) \\
&= (1 - (n-s)\varepsilon) (G(1) - G(1-\varepsilon)) - \varepsilon \int_{(\partial \mathcal{B}_1) \cap \{t>0\}} t^{1-s} |\nabla_\nu \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n + o(\varepsilon),
\end{aligned}$$

where ∇_ν denotes the (exterior) normal gradient along $\partial \mathcal{B}_1$.

Furthermore, setting

$$J(r) := r^{s-n} \mathcal{J}_s(B_r E, H^c),$$

using the substitutions $\bar{x} := x/(1 - \varepsilon)$ and $\bar{y} := y/(1 - \varepsilon)$, and noticing that $C_E \cap (\partial B_1) = E \cap (\partial B_1)$, we have that

$$\begin{aligned}
& \mathcal{J}_s(B_1 E^{(\varepsilon)}, H^c) - \mathcal{J}_s(B_1 E, H^c) \\
&= \mathcal{J}_s(B_{1-\varepsilon}((1-\varepsilon)E), H^c) - \mathcal{J}_s(B_{1-\varepsilon}E, H^c) \\
&\quad + \mathcal{J}_s(B_1 B_{1-\varepsilon}^c C_E, H^c) - \mathcal{J}_s(B_1 B_{1-\varepsilon}^c E, H^c) \\
&= \iint_{B_{1-\varepsilon}((1-\varepsilon)E) \times H^c} \frac{dx dy}{|x-y|^{n+s}} - (1-\varepsilon)^{n-s} J(1-\varepsilon) \\
&\quad + \varepsilon \left(\iint_{(\partial B_1) \times H^c} \frac{\chi_{C_E}(x) d\mathcal{H}_x^{n-1} dy}{|x-y|^{n+s}} - \iint_{(\partial B_1) \times H^c} \frac{\chi_E(x) d\mathcal{H}_x^{n-1} dy}{|x-y|^{n+s}} \right) + o(\varepsilon) \\
&= (1-\varepsilon)^{n-s} \iint_{B_1 E \times H^c} \frac{d\bar{x} d\bar{y}}{|\bar{x}-\bar{y}|^{n+s}} - (1-\varepsilon)^{n-s} J(1-\varepsilon) + o(\varepsilon) \\
&= (1-\varepsilon)^{n-s} (J(1) - J(1-\varepsilon)) + o(\varepsilon) \\
&= (1 - (n-s)\varepsilon) (J(1) - J(1-\varepsilon)) + o(\varepsilon).
\end{aligned}$$

Then, plugging this information and (4.3) into (4.2), and noticing that $\Phi_E(r) = G(r) + (\sigma - 1)J(r)$, we conclude that

$$\begin{aligned}
0 &\leq (1 - (n-s)\varepsilon) (G(1) - G(1-\varepsilon)) - \varepsilon \int_{(\partial B_1) \cap \{t>0\}} t^{1-s} |\nabla_v \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n \\
&\quad + (\sigma - 1) (1 - (n-s)\varepsilon) (J(1) - J(1-\varepsilon)) + o(\varepsilon) \\
&= (1 - (n-s)\varepsilon) (\Phi_E(1) - \Phi_E(1-\varepsilon)) \\
&\quad - \varepsilon \int_{(\partial B_1) \cap \{t>0\}} t^{1-s} |\nabla_v \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n + o(\varepsilon) \\
&= \varepsilon \Phi'_E(1) - \varepsilon \int_{(\partial B_1) \cap \{t>0\}} t^{1-s} |\nabla_v \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n + o(\varepsilon).
\end{aligned}$$

Therefore, dividing by ε and sending $\varepsilon \searrow 0$, we see that

$$(4.4) \quad \Phi'_E(1) \geq \int_{(\partial B_1) \cap \{t>0\}} t^{1-s} |\nabla_v \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n.$$

On the other hand, in light of (1.7), we know that

$$(4.5) \quad \Phi_{E_\lambda}(r) = \Phi_{E_{\lambda r/\rho}}(\rho),$$

for all $r, \rho, \lambda > 0$, and thus, choosing $\rho := \lambda r$,

$$\Phi_{E_\lambda}(r) = \Phi_E(\lambda r).$$

As a consequence, taking $\lambda := R$ and $r := 1 + h$, and $\lambda := R$ and $r := 1$, we see that, for all $R > 0$,

$$\Phi'_E(R) = \lim_{h \rightarrow 0} \frac{\Phi_E(R(1+h)) - \Phi_E(R)}{Rh} = \lim_{h \rightarrow 0} \frac{\Phi_{E_R}(1+h) - \Phi_{E_R}(1)}{Rh} = \frac{\Phi'_{E_R}(1)}{R}.$$

Combining this and (4.4) (used here on the set E_R), we obtain that

$$\begin{aligned}\Phi'_E(R) &\geq \frac{1}{R} \int_{(\partial\mathcal{B}_1) \cap \{t>0\}} t^{1-s} |\nabla_\nu \mathbf{E}_{E_R}(X)|^2 d\mathcal{H}_X^n \\ &= R \int_{(\partial\mathcal{B}_1) \cap \{t>0\}} t^{1-s} |\nabla_\nu \mathbf{E}_E(RX)|^2 d\mathcal{H}_X^n \\ &= R^{s-n} \int_{(\partial\mathcal{B}_R) \cap \{t>0\}} t^{1-s} |\nabla_\nu \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n,\end{aligned}$$

that is (1.9), as desired.

Now, if E is a cone, from (1.7) we have that $\Phi_E(r) = \Phi_E(\rho)$ for any $r, \rho > 0$, and therefore Φ_E is constant.

Viceversa, if Φ_E is constant, we deduce from (1.9) that

$$\int_{(\partial\mathcal{B}_r) \cap \{t>0\}} t^{1-s} |\nabla_\nu \mathbf{E}_E(X)|^2 d\mathcal{H}_X^n = 0$$

for all $r > 0$, and therefore $X \cdot \nabla \mathbf{E}_E(X) = 0$ for all $X \in \mathbb{R}_+^{n+1}$. By Euler's formula, this gives that \mathbf{E}_E is homogeneous of degree zero, and consequently, for any $\tau > 0$,

$$\chi_E(\tau x) = \mathbf{E}_E(\tau x, 0) = \mathbf{E}_E(x, 0) = \chi_E(x),$$

and hence E is a cone. ■

5. Homogeneous structure of the blow-up limits and proof of Corollary 1.3

In this section, we analyze the structure of the blow-up limit of local minimizers and we prove Corollary 1.3. To this end, we need the forthcoming auxiliary result which can be seen as the counterpart of Proposition 9.1 in [1] in our setting.

Lemma 5.1. *Let $E \subseteq H$ be a local minimizer in H . Let $E_k \subseteq H$ be a sequence of local minimizers in H and suppose that $E_k \rightarrow E$ in $L^1_{\text{loc}}(\mathbb{R}^n)$ as $k \rightarrow +\infty$.*

Then,

$$\lim_{k \rightarrow +\infty} \Phi_{E_k}(r) = \Phi_E(r) \quad \text{for all } r > 0.$$

Proof. We note that

$$\begin{aligned}(5.1) \quad r^{n-s} \Phi_{E_k}(r) &= \mathcal{F}_{s,\sigma}(\mathbf{E}_{E_k}, \mathcal{B}_r) \\ &= \int_{\mathcal{B}_r^+} t^{1-s} |\nabla \mathbf{E}_{E_k}(X)|^2 dX + (\sigma - 1) \iint_{(\mathcal{B}_r, H) \times H^c} \frac{\chi_{E_k}(x)}{|x - y|^{n+s}} dx dy.\end{aligned}$$

By the dominated convergence theorem, we have that

$$(5.2) \quad \lim_{k \rightarrow +\infty} \iint_{(\mathcal{B}_r, H) \times H^c} \frac{\chi_{E_k}(x)}{|x - y|^{n+s}} dx dy = \iint_{(\mathcal{B}_r, H) \times H^c} \frac{\chi_E(x)}{|x - y|^{n+s}} dx dy.$$

By this and (5.1) we see that, to prove the desired result, it suffices to show that

$$(5.3) \quad \lim_{k \rightarrow +\infty} \int_{\mathcal{B}_r^+} t^{1-s} |\nabla \mathbf{E}_{E_k}(X)|^2 dX = \int_{\mathcal{B}_r^+} t^{1-s} |\nabla \mathbf{E}_E(X)|^2 dX.$$

To this end, we use formula (7.2) in Proposition 7.1 in [1] and we write that, given $r, \delta > 0$,

$$\begin{aligned} \int_{\mathcal{B}_r^+} t^{1-s} |\nabla(\mathbf{E}_{E_k} - \mathbf{E}_E)(X)|^2 dX &= \int_{\mathcal{B}_r^+} t^{1-s} |\nabla \mathbf{E}_{\chi_{E_k} - \chi_E}(X)|^2 dX \\ &\leq C_{r,\delta} \int_{\mathcal{Q}_{r,\delta}} \frac{|\chi_{E_k} - \chi_E(x) - (\chi_{E_k} - \chi_E)(y)|^2}{|x - y|^{n+s}} dx dy, \end{aligned}$$

for some $C_{r,\delta} > 0$, where

$$\mathcal{Q}_{r,\delta} := \mathbb{R}^{2n} \setminus (B_{r+\delta}^c \times B_{r+\delta}^c).$$

Consequently, the claim in (5.3) is established once we show that

$$(5.4) \quad \lim_{k \rightarrow +\infty} \int_{\mathcal{Q}_{r,\delta}} \frac{|\chi_{E_k} - \chi_E(x) - (\chi_{E_k} - \chi_E)(y)|^2}{|x - y|^{n+s}} dx dy = 0.$$

It is convenient to define

$$f_k(x, y) := \frac{\chi_{E_k}(x) - \chi_{E_k}(y)}{|x - y|^{(n+s)/2}} \quad \text{and} \quad f(x, y) := \frac{\chi_E(x) - \chi_E(y)}{|x - y|^{(n+s)/2}}.$$

In this way, claim (5.4) can be written as

$$(5.5) \quad \lim_{k \rightarrow +\infty} \|f_k - f\|_{L^2(\mathcal{Q}_{r,\delta})} = 0.$$

We now use \bowtie as a short notation for $\chi_{\mathcal{Q}_{r,\delta}}(x, y) dx dy / |x - y|^{n+s}$ and set $B := B_{r+\delta}$. We point out that

$$\begin{aligned} (5.6) \quad \frac{\|f_k\|_{L^2(\mathcal{Q}_{r,\delta})}^2}{2} &= \iint_{E_k \times E_k^c} \bowtie = \iint_{(E_k B) \times E_k^c} \bowtie + \iint_{(E_k B^c) \times E_k^c} \bowtie \\ &= \iint_{(E_k B) \times (E_k^c H)} \bowtie + \iint_{(E_k B) \times (E_k^c H^c)} \bowtie \\ &\quad + \iint_{(E_k B^c) \times (E_k^c H)} \bowtie + \iint_{(E_k B^c) \times (E_k^c H^c)} \bowtie \\ &= \mathcal{J}_s(E_k B, E_k^c H) + \mathcal{J}_s(E_k B, E_k^c H^c) \\ &\quad + \mathcal{J}_s(E_k B^c, E_k^c BH) + \mathcal{J}_s(E_k B^c, E_k^c BH^c), \end{aligned}$$

and therefore,

$$\begin{aligned} \frac{\|f_k\|_{L^2(\mathcal{Q}_{r,\delta})}^2}{2} &\leq \mathcal{J}_s(E_k B, E_k^c H) + \mathcal{J}_s(E_k B, E_k^c BH^c) + \mathcal{J}_s(E_k B^c, E_k^c BH) + 2\mathcal{J}_s(B, B^c) \\ &\leq \mathcal{J}_s(E_k B, E_k^c H) + \mathcal{J}_s(E_k B^c, E_k^c BH) + 2\mathcal{J}_s(B, B^c) + \mathcal{J}_s(BH, BH^c) \\ &= \mathcal{J}_s(E_k B, E_k^c H) + \mathcal{J}_s(E_k B^c, E_k^c BH) + C_{r,\delta}, \end{aligned}$$

with $C_{r,\delta}$ independent of k . Hence, using the local minimizing property of E_k in (1.2), and taking $F_k := E_k B^c$,

$$\begin{aligned} & \frac{\|f_k\|_{L^2(\mathcal{Q}_{r,\delta})}^2}{2} \\ & \leq \mathcal{J}_s(F_k B, F_k^c H) + \mathcal{J}_s(F_k B^c, F_k^c B H) + \sigma(\mathcal{J}_s(F_k B, H^c) - \mathcal{J}_s(E_k B, H^c)) + C_{r,\delta} \\ & \leq 0 + \mathcal{J}_s(B^c, B) + \sigma(0 - \mathcal{J}_s(E_k B, H^c)) + C_{r,\delta} \leq 2C_{r,\delta}. \end{aligned}$$

This and Fatou's lemma yield that

$$\|f\|_{L^2(\mathcal{Q}_{r,\delta})}^2 \leq 4C_{r,\delta}.$$

Now we remark that to prove (5.5) it suffices to show that

$$(5.7) \quad \lim_{k \rightarrow +\infty} \|f_k\|_{L^2(\mathcal{Q}_{r,\delta})} = \|f\|_{L^2(\mathcal{Q}_{r,\delta})}.$$

Indeed, suppose that (5.7) holds true and notice that f_k converges to f pointwise. Let $\varphi \in C_0^\infty(\mathcal{Q}_{r,\delta})$ and observe that

$$|f_k(x, y) \varphi(x, y)| \leq \frac{|\varphi(x, y)|}{|x - y|^{(n+s)/2}} \in L^1(\mathcal{Q}_{r,\delta}).$$

Hence, by the dominated convergence theorem,

$$\lim_{k \rightarrow +\infty} \int_{\mathcal{Q}_{r,\delta}} f_k \varphi = \int_{\mathcal{Q}_{r,\delta}} f \varphi.$$

By density, given $\varepsilon > 0$, we can pick $\varphi_\varepsilon \in C_0^\infty(\mathcal{Q}_{r,\delta})$ such that $\|\varphi_\varepsilon - f\|_{L^2(\mathcal{Q}_{r,\delta})} \leq \varepsilon$. In this way, we find that

$$\begin{aligned} \limsup_{k \rightarrow +\infty} \left| \int_{\mathcal{Q}_{r,\delta}} f_k f - \int_{\mathcal{Q}_{r,\delta}} f^2 \right| & \leq \limsup_{k \rightarrow +\infty} \left| \int_{\mathcal{Q}_{r,\delta}} f_k \varphi_\varepsilon - \int_{\mathcal{Q}_{r,\delta}} f^2 \right| + \int_{\mathcal{Q}_{r,\delta}} f_k |f - \varphi_\varepsilon| \\ & \leq \left| \int_{\mathcal{Q}_{r,\delta}} f \varphi_\varepsilon - \int_{\mathcal{Q}_{r,\delta}} f^2 \right| + \limsup_{k \rightarrow +\infty} \|f_k\|_{L^2(\mathcal{Q}_{r,\delta})} \|\varphi_\varepsilon - f\|_{L^2(\mathcal{Q}_{r,\delta})} \\ & \leq \limsup_{k \rightarrow +\infty} (\|f\|_{L^2(\mathcal{Q}_{r,\delta})} + \|f_k\|_{L^2(\mathcal{Q}_{r,\delta})}) \|\varphi_\varepsilon - f\|_{L^2(\mathcal{Q}_{r,\delta})} \leq 4\varepsilon \sqrt{C_{r,\delta}}. \end{aligned}$$

Hence, since ε can be taken arbitrarily small,

$$\lim_{k \rightarrow +\infty} \int_{\mathcal{Q}_{r,\delta}} f_k f = \int_{\mathcal{Q}_{r,\delta}} f^2.$$

As a result, if (5.7) holds true, we obtain that

$$\lim_{k \rightarrow +\infty} \|f_k - f\|_{L^2(\mathcal{Q}_{r,\delta})}^2 = \lim_{k \rightarrow +\infty} \|f_k\|_{L^2(\mathcal{Q}_{r,\delta})}^2 + \|f\|_{L^2(\mathcal{Q}_{r,\delta})}^2 - 2 \int_{\mathcal{Q}_{r,\delta}} f_k f = 0,$$

that is (5.5).

In view of this observation, to complete the proof of Lemma 5.1, we are left with proving (5.7). As a matter of fact, by Fatou's lemma, to prove (5.7) it suffices to check that

$$(5.8) \quad \limsup_{k \rightarrow +\infty} \|f_k\|_{L^2(\mathcal{Q}_{r,\delta})} \leq \|f\|_{L^2(\mathcal{Q}_{r,\delta})},$$

and therefore the remaining part of this proof is devoted to show this inequality. To this end, we let D_k be the symmetric difference of E_k and E , and we define

$$G_k := (EB) \cup (E_k B^c).$$

The local minimizing property of E_k as stated in (1.2) yields that

$$\begin{aligned} & \mathcal{J}_s(E_k B, E_k^c H) + \mathcal{J}_s(E_k B^c, E_k^c BH) + \sigma \mathcal{J}_s(E_k B, H^c) \\ & \leq \mathcal{J}_s(G_k B, G_k^c H) + \mathcal{J}_s(G_k B^c, G_k^c BH) + \sigma \mathcal{J}_s(G_k B, H^c) \\ & = \mathcal{J}_s(EB, G_k^c H) + \mathcal{J}_s(E_k B^c, E^c BH) + \sigma \mathcal{J}_s(EB, H^c) \\ & = \mathcal{J}_s(EB, E^c BH) + \mathcal{J}_s(EB, E_k^c B^c H) + \mathcal{J}_s(E_k B^c, E^c BH) + \sigma \mathcal{J}_s(EB, H^c) \\ & \leq \mathcal{J}_s(EB, E^c BH) + \mathcal{J}_s(EB, E^c B^c H) + \mathcal{J}_s(EB^c, E^c BH) + \sigma \mathcal{J}_s(EB, H^c) \\ & \quad + \mathcal{J}_s(EB, D_k B^c H) + \mathcal{J}_s(D_k B^c, E^c BH) \\ & \leq \mathcal{J}_s(EB, E^c H) + \mathcal{J}_s(EB^c, E^c BH) + \sigma \mathcal{J}_s(EB, H^c) + 2\mathcal{J}_s(B, D_k B^c). \end{aligned}$$

By [1] (see in particular the proof of Theorem 3.3 there), we know that

$$\lim_{k \rightarrow +\infty} \mathcal{J}_s(B, D_k B^c) = 0,$$

and accordingly we can write that

$$\begin{aligned} & \limsup_{k \rightarrow +\infty} \mathcal{J}_s(E_k B, E_k^c H) + \mathcal{J}_s(E_k B^c, E_k^c BH) + \sigma \mathcal{J}_s(E_k B, H^c) \\ & \leq \mathcal{J}_s(EB, E^c H) + \mathcal{J}_s(EB^c, E^c BH) + \sigma \mathcal{J}_s(EB, H^c). \end{aligned}$$

Hence, recalling (5.2),

$$(5.9) \quad \begin{aligned} & \limsup_{k \rightarrow +\infty} \mathcal{J}_s(E_k B, E_k^c H) + \mathcal{J}_s(E_k B^c, E_k^c BH) \\ & \leq \mathcal{J}_s(EB, E^c H) + \mathcal{J}_s(EB^c, E^c BH). \end{aligned}$$

Besides, from (5.6),

$$\begin{aligned} \frac{\|f_k\|_{L^2(\mathcal{Q}_{r,\delta})}^2}{2} & = \mathcal{J}_s(E_k B, E_k^c H) + \mathcal{J}_s(E_k B, E_k^c H^c) \\ & \quad + \mathcal{J}_s(E_k B^c, E_k^c BH) + \mathcal{J}_s(E_k B^c, E_k^c BH^c), \end{aligned}$$

and a similar formula holds true by replacing f_k by f and E_k by E .

In this way, exploiting again the dominated convergence theorem, we deduce that

$$\begin{aligned}
 & \limsup_{k \rightarrow +\infty} \frac{1}{2} \left(\|f_k\|_{L^2(\mathcal{Q}_{r,\delta})}^2 - \|f\|_{L^2(\mathcal{Q}_{r,\delta})}^2 \right) \\
 &= \limsup_{k \rightarrow +\infty} \mathcal{J}_s(E_k B, E_k^c H) + \mathcal{J}_s(E_k B, E_k^c H^c) + \mathcal{J}_s(E_k B^c, E_k^c B H) + \mathcal{J}_s(E_k B^c, E_k^c B H^c) \\
 &\quad - \mathcal{J}_s(EB, E^c H) - \mathcal{J}_s(EB, E^c H^c) - \mathcal{J}_s(EB^c, E^c B H) - \mathcal{J}_s(EB^c, E^c B H^c) \\
 &= \limsup_{k \rightarrow +\infty} \mathcal{J}_s(E_k B, E_k^c H) + \mathcal{J}_s(E_k B^c, E_k^c B H) - \mathcal{J}_s(EB, E^c H) - \mathcal{J}_s(EB^c, E^c B H).
 \end{aligned}$$

From this and (5.9) we obtain (5.8), as desired. \blacksquare

With this preliminary work, we can now complete the proof of Corollary 1.3 by arguing as follows.

Proof of Corollary 1.3. The proof is based on a double blow-up procedure, combined with the monotonicity formula in Theorem 1.2. The advantage of a double blow-up with respect to a single blow-up is that the first blow-up reduces the container ω to a half-space, thus allowing us to use Lemma 5.1 in the second blow-up.

Here are the details of the proof. First of all, we consider the sequence of sets $E_{1/k}$, with $k \in \mathbb{N}$. By Theorem A.2 in [9], up to a subsequence, we know that $\chi_{E_{1/k}}$ converges in $L^1_{\text{loc}}(\mathbb{R}^n)$ to χ_{E^*} as $k \rightarrow +\infty$, for a suitable E^* contained in a half-space H^* , with E^* locally minimizing in H^* . Up to a rigid motion, we can suppose that $H^* = H$.

Now we consider the sequence $E_{1/h}^*$, with $h \in \mathbb{N}$. Using again Theorem A.2 in [9], up to a subsequence, we see that $\chi_{E_{1/h}^*}$ converges as $h \rightarrow +\infty$ in $L^1_{\text{loc}}(\mathbb{R}^n)$ to χ_{E_0} , for a suitable $E_0 \subseteq H$ which is locally minimizing in H . Also, thanks to Lemma 5.1, we have that

$$(5.10) \quad \lim_{h \rightarrow +\infty} \Phi_{E_{1/h}^*}(r) = \Phi_{E_0}(r).$$

Then, Corollary 1.3 will be established once we prove the following claims:

$$(5.11) \quad E_0 \text{ is a cone}$$

and

$$(5.12) \quad \text{there exists an infinitesimal sequence } r_j > 0 \text{ such that } \chi_{E_{r_j}} \text{ converges to } \chi_{E_0} \text{ in } L^1_{\text{loc}}(\mathbb{R}^n) \text{ as } j \rightarrow +\infty.$$

To prove (5.11), we exploit (4.5) with $\lambda := 1/h$ and $\rho := \lambda r$, by writing

$$\Phi_{E_{1/h}^*}(r) = \Phi_{E^*}\left(\frac{r}{h}\right).$$

Hence, in light of (5.10),

$$(5.13) \quad \Phi_{E_0}(r) = \lim_{h \rightarrow +\infty} \Phi_{E_{1/h}^*}(r) = \lim_{h \rightarrow +\infty} \Phi_{E^*}\left(\frac{r}{h}\right) = \lim_{\delta \searrow 0} \Phi_{E^*}(\delta).$$

Notice that the last limit exists, due to the monotonicity of the function proved in Theorem 1.2. Furthermore, the identity in (5.13) says that Φ_{E_0} is constant and then, by Theorem 1.2, E_0 must necessarily be a cone, which proves (5.11).

Now we prove (5.12). For this, let $R > 0$. By the convergence of $E_{1/h}^*$, we know that, given $\varepsilon > 0$, there exists $h_0(R, \varepsilon) \in \mathbb{N}$ such that, for all $h \geq h_0(R, \varepsilon)$,

$$(5.14) \quad \int_{B_R} |\chi_{E_{1/h}^*}(x) - \chi_{E_0}(x)| dx \leq \varepsilon.$$

On the other hand, by the convergence of $E_{1/k}$, there exists $k_0(R, h, \varepsilon) \in \mathbb{N}$ such that, for all $k \geq k_0(R, h, \varepsilon)$,

$$\int_{B_{R/h}} |\chi_{E_{1/k}}(x) - \chi_{E^*}(x)| dx \leq \frac{\varepsilon}{h^n}.$$

Scaling back, and using (5.11), this gives that, for all $k \geq k_0(R, h, \varepsilon)$,

$$\int_{B_R} |\chi_{E_{1/(hk)}}(x) - \chi_{E_{1/h}^*}(x)| dx \leq \varepsilon.$$

Combining this with (5.14), we find that, for all $k \geq k_*(R, \varepsilon) := k_0(R, h_0(R, \varepsilon), \varepsilon)$,

$$\begin{aligned} & \int_{B_R} |\chi_{E_{1/(h_0(R, \varepsilon)k)}}(x) - \chi_{E_0}(x)| dx \\ & \leq \int_{B_R} |\chi_{E_{1/(h_0(R, \varepsilon)k)}}(x) - \chi_{E_{1/h_0(R, \varepsilon)}^*}(x)| dx + \int_{B_R} |\chi_{E_{1/h_0(R, \varepsilon)}^*}(x) - \chi_{E_0}(x)| dx \leq 2\varepsilon. \end{aligned}$$

This establishes (5.12), as desired. \blacksquare

6. Locally minimizing cones in the plane and proof of Theorem 1.4

In this section, we take $n = 2$, and we classify locally minimizing cones, thus proving Theorem 1.4.

Proof of Theorem 1.4. Let $\Psi \in C_0^\infty(\mathcal{B}_{9/10}, [0, 1])$ be a radially decreasing function with $\Psi(X) = 1$ for all $X \in \mathcal{B}_{1/2}$. Given $R > 2$, to be taken as large as we wish in the following, we consider the transformation

$$(6.1) \quad \mathbb{R}^3 \ni X \mapsto Y := X + \Psi\left(\frac{X}{R}\right)e_1,$$

where $e_1 := (1, 0, 0)$. Denoting this map by $Y(X)$ (see Figure 2), we see that it is invertible, and we denote its inverse by $X(Y)$. We also let

$$(6.2) \quad U := \mathbf{E}_E,$$

and

$$U_R^+(Y) := U(X(Y)).$$




Figure 2. Depicting the action of the map Y defined in (6.1) on a set S . Notice that $S \cap \mathcal{B}_{R/2}$ is translated by e_1 , while $S \setminus \mathcal{B}_R$ is left unchanged. Since Ψ is radially decreasing, the slices $S \cap \partial \mathcal{B}_\rho$ corresponding to $\rho \in (1, R)$ are translated by multiples $\lambda(\rho) e_1$ of e_1 , where $\lambda(\rho)$ decreases from $\lambda = 1$ when $\rho = R/2$, to $\lambda = 0$ when $\rho \geq (9/10)R$.

We also denote U_R^- a similar function, in which Ψ is replaced by $-\Psi$. In addition, we set $u(x) := U(x, 0)$, $u_R^+(y) := U_R^+(y, 0)$ and $u_R^-(y) := U_R^-(y, 0)$.

We use coordinates $X = (X_1, X_2, X_3) = (x, t) \in \mathbb{R}^2 \times (0, +\infty)$. We remark that $Y_3(X) = X_3$, hence $X_3(Y) = Y_3$, and accordingly $X_3(y, 0) = 0$. This gives that

$$(6.3) \quad u_R^+(y) = U(X(y, 0)) = U(x(y, 0), 0) = \chi_E(x(y, 0)).$$

Then, in the notation of (1.3), we claim that

$$(6.4) \quad \left| \mathcal{F}_{s,\sigma}(U_R^+, \mathcal{B}_R) + \mathcal{F}_{s,\sigma}(U_R^-, \mathcal{B}_R) - 2\mathcal{F}_{s,\sigma}(U, \mathcal{B}_R) \right| \leq \frac{C}{R^s},$$

for some $C > 0$. To prove this, we let

$$\mathcal{J}_R(U) := \int_{\mathcal{B}_R^+} t^{1-s} |\nabla U(X)|^2 dX \quad \text{and} \quad \mathcal{T}_R(u) := \iint_{\mathcal{B}_R \times H^c} \frac{u(x)}{|x-z|^{2+s}} dx dz.$$

A direct computation (see Lemma 1 in [11]) shows that

$$(6.5) \quad \left| \mathcal{J}_R(U_R^+) + \mathcal{J}_R(U_R^-) - 2\mathcal{J}_R(U) \right| \leq \frac{C}{R^s},$$

for some $C > 0$.

We introduce the following notation: from now on, we denote by \diamond any quantity or bounded function, possibly different from line to line, which changes sign if Ψ is replaced by $-\Psi$. We stress that it is not necessary that \diamond has a sign itself, what matters in this notation is that its pointwise value changes sign if Ψ is replaced by $-\Psi$.

Now, we want to use the change of variable $\tilde{y} := x(y, 0)$ and $\tilde{z} := x(y, 0) - y + z$. In this way, we have that

$$\tilde{y} - \tilde{z} = y - z.$$

We also observe that, if $z \in H^c$, then $\tilde{z}_2 = x_2(y, 0) - y_2 + z_2 = z_2 \leq 0$, and thus $\tilde{z} \in H^c$.

Furthermore, for all $i, j \in \{1, 2, 3\}$,

$$D_{X_i} Y_j(X) = \delta_{ij} + \frac{\delta_{1j}}{R} \partial_i \Psi\left(\frac{X}{R}\right) = \delta_{ij} + \frac{\diamond}{R}.$$

Therefore, we can write that

$$dy dz = \left(1 + \frac{\diamond}{R} + O\left(\frac{1}{R^2}\right)\right) d\tilde{y} d\tilde{z}.$$

We also point out that

$$(6.6) \quad \text{if } y \in B_R, \text{ then } x(y, 0) \in B_R.$$

Indeed, if $|y| \leq 99R/100$, then

$$|x(y, 0)| = \left|y - \Psi\left(\frac{x(y, 0)}{R}\right) e_1\right| \leq \frac{99R}{100} + 1 < R,$$

as long as R is large enough.

If instead $|y| > 99R/100$, it follows that

$$|x(y, 0)| = \left|y - \Psi\left(\frac{x(y, 0)}{R}\right) e_1\right| \geq |y| - 1 > \frac{99R}{100} - 1 > \frac{9R}{10},$$

and consequently $\Psi(x(y, 0)/R) = 0$, whence $x(y, 0) = y$ in this case.

These considerations prove (6.6). Hence, recalling (6.3),

$$\begin{aligned} \mathcal{T}_R(u_R^+) &= \iint_{B_R \times H^c} \frac{u_R^+(y)}{|y-z|^{2+s}} dy dz = \iint_{B_R \times H^c} \frac{\chi_E(x(y, 0))}{|y-z|^{2+s}} dy dz \\ &= \iint_{B_R \times H^c} \frac{\chi_E(\tilde{y})}{|\tilde{y}-\tilde{z}|^{2+s}} \left(1 + \frac{\diamond}{R} + O\left(\frac{1}{R^2}\right)\right) d\tilde{y} d\tilde{z}. \end{aligned}$$

Given our notation related to \diamond , this also says that

$$\mathcal{T}_R(u_R^-) = \iint_{B_R \times H^c} \frac{\chi_E(\tilde{y})}{|\tilde{y}-\tilde{z}|^{2+s}} \left(1 - \frac{\diamond}{R} + O\left(\frac{1}{R^2}\right)\right) d\tilde{y} d\tilde{z}.$$

As a consequence,

$$\begin{aligned} |\mathcal{T}_R(u_R^+) + \mathcal{T}_R(u_R^-) - 2\mathcal{T}_R(u)| &\leq O\left(\frac{1}{R^2}\right) \iint_{B_R \times H^c} \frac{\chi_E(\tilde{y})}{|\tilde{y}-\tilde{z}|^{2+s}} d\tilde{y} d\tilde{z} \\ &\leq O\left(\frac{1}{R^2}\right) \iint_{B_R \times H^c} \frac{d\tilde{y} d\tilde{z}}{|\tilde{y}-\tilde{z}|^{2+s}} \leq O\left(\frac{1}{R^2}\right) \mathcal{I}_s(B_R H, (B_R H)^c) = O\left(\frac{1}{R^s}\right). \end{aligned}$$

From this, (1.3) and (6.5), we obtain (6.4), up to renaming $C > 0$, as desired.

Moreover, from (1.6), we can write that

$$\mathcal{F}_{s,\sigma}(U, \mathcal{B}_R) \leq \mathcal{F}_{s,\sigma}(U_R^-, \mathcal{B}_R).$$

Using this and (6.4), we conclude that

$$(6.7) \quad \begin{aligned} \mathcal{F}_{s,\sigma}(U_R^+, \mathcal{B}_R) - \mathcal{F}_{s,\sigma}(U, \mathcal{B}_R) &\leq \mathcal{F}_{s,\sigma}(U_R^+, \mathcal{B}_R) + \mathcal{F}_{s,\sigma}(U_R^-, \mathcal{B}_R) - 2\mathcal{F}_{s,\sigma}(U, \mathcal{B}_R) \\ &\leq \frac{C}{R^s}. \end{aligned}$$

Now we claim that

$$(6.8) \quad U \text{ is monotone in the direction } e_1, \\ \text{namely either } U(X + \tau e_1) \geq U(X) \text{ or } U(X + \tau e_1) \leq U(X), \text{ for every } \tau > 0.$$

To prove this, we argue by contradiction, supposing that there exist $\bar{X} \in \mathbb{R}_+^3$ and $\bar{\tau}_1, \bar{\tau}_2 > 0$ such that

$$(6.9) \quad U(\bar{X} + \bar{\tau}_1 e_1) > U(\bar{X}) \quad \text{and} \quad U(\bar{X} + \bar{\tau}_2 e_1) < U(\bar{X}).$$

Since E is a cone, we have that U is homogeneous of degree zero, and therefore, letting

$$P := \bar{\tau}_1^{-1} \bar{X} \quad \text{and} \quad Q := \bar{\tau}_2^{-1} \bar{X},$$

we can write (6.9) as

$$(6.10) \quad \begin{aligned} U(P + e_1) &= U(\bar{\tau}_1^{-1} \bar{X} + e_1) = U(\bar{X} + \bar{\tau}_1 e_1) > U(\bar{X}) = U(\bar{\tau}_1^{-1} \bar{X}) = U(P), \\ U(Q + e_1) &= U(\bar{\tau}_2^{-1} \bar{X} + e_1) = U(\bar{X} + \bar{\tau}_2 e_1) < U(\bar{X}) = U(\bar{\tau}_2^{-1} \bar{X}) = U(Q). \end{aligned}$$

We can suppose that

$$(6.11) \quad R/2 > M := 2 + |Q| + |P|,$$

and we set

$$V_R(X) := \min\{U(X), U_R^+(X)\} \quad \text{and} \quad W_R(X) := \max\{U(X), U_R^+(X)\}.$$

We remark that

$$(6.12) \quad \mathcal{F}_{s,\sigma}(V_R, \mathcal{B}_R) + \mathcal{F}_{s,\sigma}(W_R, \mathcal{B}_R) = \mathcal{F}_{s,\sigma}(U, \mathcal{B}_R) + \mathcal{F}_{s,\sigma}(U_R^+, \mathcal{B}_R).$$

In addition, by (1.6),

$$\mathcal{F}_{s,\sigma}(U, \mathcal{B}_R) \leq \mathcal{F}_{s,\sigma}(V_R, \mathcal{B}_R).$$

Combining this and (6.12), we find that

$$(6.13) \quad \mathcal{F}_{s,\sigma}(W_R, \mathcal{B}_R) \leq \mathcal{F}_{s,\sigma}(U_R^+, \mathcal{B}_R).$$

Now, we denote by W_\star the minimizer of $\mathcal{J}_M(W)$ among all the competitors W with $W = W_R$ on $\partial \mathcal{B}_M^+ = ((\partial \mathcal{B}_M) \cap \{t > 0\}) \cup (\mathcal{B}_M \times \{0\})$.

We remark that the minimization of the functional leads to the equation

$$(6.14) \quad \operatorname{div}(t^{1-s} \nabla W_\star) = 0 \quad \text{in } \mathcal{B}_M^+.$$

Also, the same equation is fulfilled by U , in view of (6.2).

We claim that

$$(6.15) \quad W_\star \neq W_R.$$

Indeed, suppose by contradiction that $W_\star = W_R$. Then, since $U \leq W_R = W_\star$, we deduce by the strong maximum principle for the equation in (6.14) (see e.g. Corollary 2.3.10 in [7]) that

$$(6.16) \quad \text{either } U < W_R \quad \text{or } U = W_R \quad \text{in } \mathcal{B}_M^+.$$

On the other hand, by (6.11), we have that

$$Y(P) = P + \Psi\left(\frac{P}{R}\right)e_1 = P + e_1 \quad \text{and} \quad Y(Q) = Q + \Psi\left(\frac{Q}{R}\right)e_1 = Q + e_1.$$

Consequently, by (6.10),

$$\begin{aligned} U_R^+(Y(P)) &= U(P) < U(P + e_1) = U(Y(P)) \\ \text{and} \quad U_R^+(Y(Q)) &= U(Q) > U(Q + e_1) = U(Y(Q)). \end{aligned}$$

Therefore, we see that

$$W_R(Y(P)) = U(Y(P)) \quad \text{and} \quad W_R(Y(Q)) = U_R^+(Y(Q)) > U(Y(Q)),$$

and these observations say that none of the two possibilities in (6.16) can be fulfilled.

This contradiction proves (6.15). Then, from (6.15), we obtain that there exists $\delta_0 > 0$ such that

$$\mathcal{J}_M(W_\star) + \delta_0 \leq \mathcal{J}_M(W_R).$$

We stress that this δ_0 is independent of R , because W_R in \mathcal{B}_M does not depend on R , being

$$W_R(X) = \max\{U(X), U(X - e_1)\} \quad \text{for all } X \in \mathcal{B}_M^+,$$

thanks to (6.11).

Furthermore, if we extend W_\star to be equal to W_R outside \mathcal{B}_M^+ , we have that

$$(6.17) \quad \mathcal{J}_R(W_R) - \mathcal{J}_R(W_\star) = \mathcal{J}_M(W_R) - \mathcal{J}_M(W_\star) \geq \delta_0.$$

Since, by construction $w_\star(x) := W_\star(x, 0) = W_R(x, 0) =: w_R(x)$, we have that $\mathcal{T}_R(w_\star) = \mathcal{T}_R(w_R)$. This and (6.17) give that

$$\mathcal{F}_{s,\sigma}(W_R, \mathcal{B}_R) - \mathcal{F}_{s,\sigma}(W_\star, \mathcal{B}_R) \geq \delta_0.$$

As a consequence, in light of (6.13),

$$(6.18) \quad \mathcal{F}_{s,\sigma}(U_R^+, \mathcal{B}_R) - \mathcal{F}_{s,\sigma}(W_\star, \mathcal{B}_R) \geq \delta_0.$$

On the other hand, using again (1.6),

$$\mathcal{F}_{s,\sigma}(U, \mathcal{B}_R) \leq \mathcal{F}_{s,\sigma}(W_\star, \mathcal{B}_R).$$

Comparing this and (6.18), we see that

$$\mathcal{F}_{s,\sigma}(U_R^+, \mathcal{B}_R) - \widetilde{\mathcal{F}}_{s,\sigma}(U, \mathcal{B}_R) \geq \delta_0.$$

Hence, recalling (6.7),

$$\frac{C}{R^s} \geq \delta_0.$$

We can now send $R \rightarrow +\infty$ and find that $0 \geq \delta_0 > 0$. This contradiction proves the validity of (6.8).

As a consequence of (6.8), we have that u is monotone in the direction e_1 , hence the cone E is made of only one component.

From this and Theorem 1.4 in [9], one also obtains (1.10). ■

Funding. The first author is supported by the Australian Research Council DECRA DE-180100957 “PDEs, free boundaries and applications”. The second author is supported by the NSF Grant DMS 2000034. The third author is supported by the Australian Laureate Fellowship FL190100081 “Minimal surfaces, free boundaries and partial differential equations”. The first and third authors are members of INdAM and AustMS.

References

- [1] Caffarelli, L., Roquejoffre, J.-M. and Savin, O.: Nonlocal minimal surfaces. *Comm. Pure Appl. Math.* **63** (2010), no. 9, 1111–1144.
- [2] Caffarelli, L. and Silvestre, L.: An extension problem related to the fractional Laplacian. *Comm. Partial Differential Equations* **32** (2007), no. 7-9, 1245–1260.
- [3] De Philippis, G. and Maggi, F.: Regularity of free boundaries in anisotropic capillarity problems and the validity of Young’s law. *Arch. Ration. Mech. Anal.* **216** (2015), no. 2, 473–568.
- [4] De Philippis, G. and Maggi, F.: Dimensional estimates for singular sets in geometric variational problems with free boundaries. *J. Reine Angew. Math.* **725** (2017), 217–234.
- [5] Dipierro, S., Maggi, F. and Valdinoci, E.: Asymptotic expansions of the contact angle in nonlocal capillarity problems. *J. Nonlinear Sci.* **27** (2017), 1531–1550.
- [6] Dipierro, S. and Valdinoci, E.: Continuity and density results for a one-phase nonlocal free boundary problem. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **34** (2017), no. 6, 1387–1428.
- [7] Fabes, E. B., Kenig, C. E. and Serapioni, R. P.: The local regularity of solutions of degenerate elliptic equations. *Comm. Partial Differential Equations* **7** (1982), no. 1, 77–116.
- [8] Finn, R.: *Equilibrium capillary surfaces*. Grundlehren der Mathematischen Wissenschaften 284, Springer-Verlag, New York, 1986.
- [9] Maggi, F. and Valdinoci, E.: Capillarity problems with nonlocal surface tension energies. *Comm. Partial Differential Equations* **42** (2017), no. 9, 1403–1446.
- [10] Molčanov, S. A. and Ostrovskiĭ, E.: Symmetric stable processes as traces of degenerate diffusion processes. *Teor. Veroyatnost. i Primenen.* **14** (1969), 127–130.
- [11] Savin, O. and Valdinoci, E.: Regularity of nonlocal minimal cones in dimension 2. *Calc. Var. Partial Differential Equations* **48** (2013), no. 1-2, 33–39.

- [12] Savin, O. and Valdinoci, E.: Some monotonicity results for minimizers in the calculus of variations. *J. Funct. Anal.* **264** (2013), no. 10, 2469–2496.
- [13] Spitzer, F.: Some theorems concerning 2-dimensional Brownian motion. *Trans. Amer. Math. Soc.* **87** (1958), 187–197.
- [14] Stinga, P. R. and Torrea, J. L.: Extension problem and Harnack’s inequality for some fractional operators. *Comm. Partial Differential Equations* **35** (2010), no. 11, 2092–2122.

Received August 14, 2020. Published online June 29, 2021.

Serena Dipierro

Department of Mathematics and Statistics, University of Western Australia, 35 Stirling Highway, Crawley, Perth, WA 6009, Australia;
serena.dipierro@uwa.edu.au

Francesco Maggi

Department of Mathematics, University of Texas at Austin, 2515 Speedway, Stop C1200, Austin, TX 78712-1202, USA;
maggi@math.utexas.edu

Enrico Valdinoci

Department of Mathematics and Statistics, University of Western Australia, 35 Stirling Highway, Crawley, Perth, WA 6009, Australia;
enrico.valdinoci@uwa.edu.au