



On the number of vertices of Newton–Okounkov polygons

Joaquim Roé and Tomasz Szemberg

Abstract. The Newton–Okounkov body of a big divisor D on a smooth surface is a numerical invariant in the form of a convex polygon. We study the geometric significance of the *shape* of Newton–Okounkov polygons of ample divisors, showing that they share several important properties of Newton polygons on toric surfaces. In concrete terms, sides of the polygon are associated to some particular irreducible curves, and their lengths are determined by the intersection numbers of these curves with D .

As a consequence of our description, we determine the numbers k such that D admits some k -gon as a Newton–Okounkov body, elucidating the relationship of these numbers with the Picard number of the surface, which was first hinted at by work of Küronya, Lozovanu and Maclean.

1. Introduction

The Newton–Okounkov body of a line bundle with respect to an admissible flag is defined as follows (see [3, 6, 11]). Let S be a normal projective variety of dimension d (the case we shall deal with in the paper is that of a surface, $d = 2$), let D be a big divisor class on S , and fix a flag

$$Y_\bullet : S = Y_0 \supset Y_1 \supset \cdots \supset Y_d = \{pt\}$$

which is *admissible*, i.e., Y_i is an irreducible subvariety of codimension i , smooth at the point Y_d , for each i . Let g_i be a local equation for Y_i in Y_{i-1} around the point Y_d . Then Y_\bullet determines a rank d valuation v_{Y_\bullet} on the field of rational functions of S , namely $v_{Y_\bullet}(f) = (v_1(f), \dots, v_d(f))$, where v_i are defined recursively setting $f_1 = f$ and

$$\begin{aligned} v_i(f) &= \text{ord}_{Y_i}(f_i), & i &= 1, \dots, d, \\ f_{i+1} &= (f_i/g_i^{v_i(f)})|_{Y_i}, & i &= 1, \dots, d-1. \end{aligned}$$

By trivializing $\mathcal{O}_S(D)$ in an arbitrary neighborhood of Y_d , the valuation v_{Y_\bullet} may be applied to global sections of multiples $\mathcal{O}_S(kD)$, and the Newton–Okounkov body of D with respect to Y_\bullet is the convex body

$$\Delta_{Y_\bullet}(D) = \overline{\left\{ \frac{v_{Y_\bullet}(s)}{k} \mid s \in H^0(S, \mathcal{O}_S(kD)) \right\}}.$$

Whereas the *volume* of the Newton–Okounkov body is well known to be equal to $\text{vol}(D)/(\dim S)!$ for every Y_\bullet , its *shape*, and in particular its dependence on Y_\bullet , is still an intriguing subject.

There are two situations in which the work of Lazarsfeld–Mustață [11] allows to prove that $\Delta_{Y_\bullet}(D)$ is a polytope. If S is a toric variety, D is a torus-invariant divisor, and the flag Y is composed of torus-invariant subvarieties, then [11] proves that $\Delta_{Y_\bullet}(D)$ is (up to the action of $\text{GL}_n(\mathbb{Z})$) just the Newton polytope associated to D in toric geometry (see §3.4 in [4]). If S is a surface, on the other hand, then Küronya–Lozovanu–Maclean [7, 8] used the description of [11] to show that for every D and Y_\bullet , $\Delta_{Y_\bullet}(D)$ is a polygon. A close analysis of their construction reveals that the shape of general Newton–Okounkov polygons (on surfaces) reflects the geometry of the pair (D, Y_\bullet) much like Newton polygons do in the toric case. Indeed, when S, D and Y_\bullet are toric, $\Delta_{Y_\bullet}(D)$ is a polygon with vertices in \mathbb{Z}^2 , with a side corresponding to each prime toric divisor (if D is ample), the selfintersections of these prime divisors determine the slopes of these sides, and their intersection with D equals the lattice length of the corresponding sides. On an arbitrary smooth projective surface, associated to each pair (D, Y_\bullet) , there is a configuration of irreducible curves playing the role of the torus-invariant prime divisors: each side of $\Delta_{Y_\bullet}(D)$ corresponds to one or more of these irreducible curves, their slopes are rational and determined by the intersection matrix of the configuration, and their lengths are determined by the intersection numbers of D with the curves in the configuration. This amounts to Theorem B in [8], which we prefer to state here in the following form (we elaborate on the details in Section 6).

Proposition. *Let S be a normal projective surface, D a big divisor on S , and $Y_\bullet: S \supset C \supset \{p\}$ an admissible flag. Let μ be the maximal real number such that $D - \mu C$ is pseudo-effective, and let $D - \mu C = P + N$ be the Zariski decomposition of $D - \mu C$.*

The intersection matrix of C and of the irreducible components of N determine all possible slopes of sides of $\Delta_{Y_\bullet}(D)$. For a fixed D and C , in the set of all big divisors D' such that the negative part N' of the Zariski decomposition of $D' - \mu_{D'} C$ has the same support as N , and for each possible slope, the length of the corresponding side is a function of the intersection numbers of D' with these curves (where we understand that if this length is zero then no side with that slope exists).

Moreover, the lower sides are related to connected components of N passing through p whereas the upper sides are related to connected components of N intersecting C at other points.

It was observed in [7] that the number of vertices of $\Delta_{Y_\bullet}(D)$ is bounded above by $2\rho(S) + 2$, where $\rho(S)$ denotes the Picard number. We show that the slightly stronger bound $2\rho(S) + 1$ holds and is sharp, i.e., for any given natural number ρ there are surfaces S with $\rho(S) = \rho$, ample divisors D and flags $Y_\bullet: S \supset C \supset \{p\}$ such that $\Delta_{Y_\bullet}(D)$ is a $(2\rho + 1)$ -gon. We also determine, in terms of configurations of negative curves on a given smooth surface S , the numbers k for which there is a flag Y_\bullet such that $\Delta_{Y_\bullet}(D)$ is a k -gon.

The role of Zariski decompositions in the determination of $\Delta_{Y_\bullet}(D)$ provides a strong relationship with Zariski chamber decompositions, and in fact the subdivision of the interval $[0, \mu]$ given by the projections of the sides of the polygon is also induced by the Zariski walls crossed by the ray that starts from D in the direction of $-C$. Somewhat surprisingly

for the authors, the numbers k for which $\Delta_{Y_\bullet}(D)$ can have k sides (or the chambers that can be traversed by a ray emanating from D) are independent of D as long as it is ample.

In order to state our main result, we introduce some new invariants attached to a configuration of negative curves. For an effective divisor $N = C_1 + \dots + C_k$ with negative definite intersection matrix, consider the following two numbers:

- $mc(N)$ denotes the largest number of irreducible components of a connected divisor contained in N .
- $mv(N)$ denotes $k + mc(N) + 4$ if $k < \rho - 1$, and $k + mc(N) + 3$ if $k = \rho - 1$.

Given a smooth projective surface S , let

$$mv(S) = \max\{mv(N) \mid N = C_1 + \dots + C_k \text{ negative definite}\}.$$

Our main result is the following.

Theorem 1.1. *On every smooth projective surface S , and for every big divisor D ,*

$$\max_{Y_\bullet} \{\#\text{vertices}(\Delta_{Y_\bullet}(D))\} \leq mv(S),$$

where the maximum is taken over all admissible flags Y . If D is ample, then for every $3 \leq v \leq mv(S)$ there exists a flag Y_\bullet such that $\Delta_{Y_\bullet}(D)$ has exactly v vertices.

Note that by the Hodge index theorem, $mv(S)$ is defined, and it is bounded above by $2\rho + 1$; we also show that this upper bound is sharp:

Corollary 1.2. *Given a positive integer ρ , there is a projective smooth surface S with Picard number $\rho(S) = \rho$, a divisor D and a flag Y_\bullet such that $\Delta_{Y_\bullet}(D)$ has $2\rho + 1$ vertices.*

The deep analogy of Newton–Okounkov polygons with the Newton polygons in toric geometry is the departure point for our work. In the case of finitely generated valuation semigroup, the connection with toric geometry goes far beyond an analogy and has found important applications via *toric degenerations* (see e.g. [5]). When the pseudo-effective cone of S is finitely generated, only finitely many negative divisors $N = C_1 + \dots + C_k$ exist, with the C_i being generators, and the application of Theorem 1.1 becomes especially straightforward. In these cases, a concrete description of the polygon $\Delta_{Y_\bullet}(D)$ is already available, as a Minkowski sum of triangles and line segments (at least if S is del Pezzo or the flag Y_\bullet is general enough, see [9, 12]).

In the absence of finite generation, the meaning of the toric analogy is far from being well understood, and in [8] it appears implicitly only. In Section 2 we recall the description of Newton–Okounkov polygons uncovered by [11], Section 6.2, and [8] from the point of view outlined above, which we then use to study these polygons, with special emphasis on the number of vertices (or sides) they possess. The analysis of these boundaries, done in Sections 3 and 4, allows to determine the number of lower and upper vertices; Proposition 3.3 is the technical key to all results in this paper. It is worth stressing that, analogously to the boundaries of classical Newton polygons, the lower boundary encodes the local behavior of D near p , whereas the upper boundary encodes the behaviour “at infinity”. In Section 5 we prove Theorem 1.1 and, finally, in Propositions 6.1 and 6.2 we show how to explicitly determine the slopes and lengths (so the whole shape) of Newton–Okounkov polygons from intersection numbers as indicated above. To construct flags Y_\bullet

that give polygons with the desired number of points we use Lemma 5.3, which may be interesting in itself: it shows that the Zariski chambers that can be crossed by a ray starting from an ample class in the Néron–Severi space are independent of the particular class chosen.

2. Newton–Okounkov polygons

In this section we recall the description of Newton–Okounkov polygons on surfaces given by Lazarsfeld–Mustață [11] and Küronya–Lozovanu–Macleán [8]. Given a surface S , we denote $\text{NS}(S)$ its Néron–Severi group (i.e., the group of divisors modulo numerical equivalence), a finitely generated abelian group of rank $\rho(S)$. When needed, we will consider \mathbb{Q} -divisors and \mathbb{R} -divisors with the conventions of [10], and $\text{NS}(S)_{\mathbb{R}}$ will be the space of numerical classes of \mathbb{R} -divisors endowed with the bilinear form given by the intersection product.

Fix a smooth surface S , a big divisor D and an admissible flag $Y_{\bullet}: S \supset C \supset \{p\}$ on S . For every real number t , consider the \mathbb{R} -divisor $D_t = D - tC$ and, if D_t is effective or pseudo-effective, denote its Zariski decomposition

$$D_t = P_t + N_t.$$

Let $\nu = \nu_C(D)$ be the coefficient of C in the negative part N_0 of the Zariski decomposition of D , and let $\mu = \mu_C(D) = \max\{t \in \mathbb{R} \mid D_t \text{ is pseudo-effective}\}$. Note that D_{μ} belongs to the boundary of the pseudo-effective cone, in particular it is not big (as big classes form the interior of the pseudo-effective cone). For every $t \in [\nu, \mu]$, define $\alpha(t) = (N_t \cdot C)_p$, i.e., the local intersection multiplicity of the negative part of D_t and C at p , and $\beta(t) = \alpha(t) + P_t \cdot C$. Lazarsfeld and Mustață showed in Section 6.2 of [11] that $\Delta_{Y_{\bullet}}(D)$ is the region in the plane (t, s) defined by the inequalities $\nu \leq t \leq \mu$, $\alpha(t) \leq s \leq \beta(t)$. Note that α and β are continuous piecewise linear functions in the interval $[\nu, \mu]$, respectively convex and concave.

Observing that N_t increases with t and looking at N_{μ} , Küronya–Lozovanu–Macleán proved in [8] that α is nondecreasing, that the values $t \in (\nu, \mu)$ where α or β fails to be linear are exactly those where D_t crosses walls between Zariski chambers [2], and that there are finitely many such crossed walls, in fact at most as many as components in N_{μ} .

Remark 2.1. Let q_1, \dots, q_r be the intersection points of N_{μ} and C different from p . It follows immediately from the description above that $\beta(t) = D \cdot C - tC^2 - \sum_{i=1}^r (N_t \cdot C)_{q_i}$.

Our approach to understanding the number of vertices in Newton–Okounkov polygons is to further analyze the dependence of N_t on t , and from this derive information on the functions α and β . So, let us briefly recall the proof of polygonality of $\Delta_{Y_{\bullet}}(D)$ due to [11] and [8]. Call C_1, \dots, C_n the irreducible components of N_{μ} , numbered in order of appearance in the support of N_t , that is, denoting

$$t_i = \inf\{t \in [\nu, \mu] \mid C_i \text{ in the support of } N_t\}$$

for each $i \in \{1, \dots, n\}$, one has $t_0 := \nu \leq t_1 \leq \dots \leq t_n < t_{n+1} := \mu$. We can write

$$N_t = a_1(t)C_1 + \dots + a_n(t)C_n,$$

where $a_i(t)$ are (continuous) functions $[v, \mu] \rightarrow \mathbb{R}$. The equations defining the Zariski decomposition $D_t = P_t + N_t$ tell us that, for $t \in [t_{i-1}, t_i]$ and $1 \leq j < i$, $P_t \cdot C_j = 0$, or equivalently $N_t \cdot C_j = D_t \cdot C_j$. Therefore $a_j(t)$ are solutions of the linear system of equations

$$(2.1) \quad \begin{aligned} (a_1(t)C_1 + \cdots + a_{i-1}(t)C_{i-1}) \cdot C_j &= (D - tC) \cdot C_j, & 1 \leq j < i, \\ a_j(t) &= 0, & i \leq j \leq n. \end{aligned}$$

These solutions are unique because the intersection matrix $(C_k \cdot C_j)_{1 \leq k, j < i}$ is nonsingular. Since the independent terms $(D - tC) \cdot C_j$ are affine linear functions of t , so are the solutions $a_j(t) = a_{j0} + a_{j1}t$, i.e., a_j is affine linear on each interval $[t_{i-1}, t_i]$. It follows then that α and β are continuous affine linear on each interval $[t_{i-1}, t_i]$ (which can be degenerate, if $t_i = t_{i-1}$) so $\Delta_{Y_\bullet}(D)$ is a polygon and the first coordinate of every vertex equals one of the t_i , $i \in \{0, \dots, n + 1\}$.

3. Interior vertices

We keep the notations of the previous section, namely $Y_\bullet: S \supset C \supset \{p\}$ is an admissible flag, $D_t = D - tC = P_t + N_t$, and $v = v_C(D)$, $\mu = \mu_C(D)$, α, β, t_i are as above.

The vertices $P = (t_i, s)$ of $\Delta_{Y_\bullet}(D)$ can be classified as *leftmost* (if $t_i = t_0 = v$), *rightmost* (if $t_i = t_{n+1} = \mu$) and *interior* (if $v < t_i < \mu$). A vertex P is also called *upper* if $s = \beta(t_i)$ and *lower* if $s = \alpha(t_i)$. Before proceeding to the determination of the t_i for which $\Delta_{Y_\bullet}(D)$ has upper and lower interior vertices, we recall a result on relative negative parts of Zariski decompositions, essentially due to Zariski:

Lemma 3.1. *Let D be an effective divisor on a smooth surface, let $D = P + N$ be its Zariski decomposition, and let $N = a_1C_1 + \cdots + a_nC_n$, $a_i \in \mathbb{Q}$, be the decomposition into irreducible components. For every subset $I \subset \{1, \dots, n\}$, let $b_i, i \in I$, be the solutions to the system of linear equations*

$$\left(D - \sum_{i \in I} b_i C_i\right) \cdot C_j = 0, \quad j \in I.$$

Then $b_i \leq a_i$ for each $i \in I$.

Proof. First observe that we may assume $I \subsetneq \{1, \dots, n\}$, as otherwise $b_i = a_i$ and there is nothing to prove. The presentation of Zariski decomposition given in [1] in terms of linear algebra will immediately yield that there is $J \subset \{1, \dots, n\}$, $I \subsetneq J$, such that the solutions b'_i to the corresponding system of equations

$$\left(D - \sum_{i \in J} b'_i C_i\right) \cdot C_j = 0, \quad j \in J,$$

satisfy $b_i \leq b'_i$ for each $i \in I$, which applied recursively gives what we need. Indeed, denote $\mathbf{p} = [D - \sum_{i \in I} b_i C_i]$, $\mathbf{v} = [D]$, and $\mathbf{e}_i = [C_i]$, $1 \leq i \leq n$, considered as vectors in $\text{NS}(S)_{\mathbb{R}}$. The space $(\mathbf{e}_i)_{i \in I}$ is, in the language of [1], a special negative definite subspace

of the support space of \mathbf{v} . So, the hypotheses of Lemma 5.3 in [1] are satisfied, and therefore $D' = D - \sum_{i \in I} b_i C_i$ is effective. Since $I \subsetneq \{1, \dots, n\}$, there is at least one curve C_j among C_1, \dots, C_n such that $D' \cdot C_j < 0$ (otherwise the Zariski decomposition of D would not involve all C_1, \dots, C_n); let J be $I \cup \{j \mid D' \cdot C_j < 0\}$. Since the intersection form on $\langle \mathbf{e}_i \rangle_{i \in J}$ is negative definite, there is a unique $\mathbf{n} = \sum_{i \in J} a_i \mathbf{e}_i$ with $\mathbf{n} \cdot \mathbf{e}_i = \mathbf{p} \cdot \mathbf{e}_i \forall i \in J$. Then Lemmas 5.2 and 5.3 in [1] give that \mathbf{n} and $\mathbf{p} - \mathbf{n}$ are effective. The latter effectiveness gives $b_i \leq b'_i$ for each $i \in I$, as wanted. ■

Fix a pair of indices $1 \leq i \leq k \leq n$ such that $t_{i-1} < t_i = \dots = t_k < t_{k+1}$. This means that C_i, \dots, C_k are the components of the negative part of all $N_{t_i+\varepsilon}$ with $\varepsilon > 0$ that are not components of N_{t_i} . Write, for $j = 1, \dots, k$,

$$a_j(t_i + \varepsilon) = \begin{cases} a_{j0} + a_{j1} \varepsilon & \text{if } -1 \ll \varepsilon \leq 0, \\ a_{j0} + a_{j1}^+ \varepsilon & \text{if } 0 < \varepsilon \ll 1, \end{cases}$$

where $a_{j0} = 0$ and $a_{j1} = 0$ for $j \geq i$, and $a_{j1}^+ > 0$ for every $j \leq k$.

Lemma 3.2. *For every $j = 1, \dots, k$, the inequality $a_{j1}^+ \geq a_{j1}$ holds. If $C_j \cdot C_{j'} > 0$ and $a_{j1}^+ > a_{j1}$, then $a_{j'1}^+ > a_{j'1}$.*

Proof. By definition,

$$\left(D_{t_i+\varepsilon} - \sum_{j=1}^k (a_{j1}\varepsilon + a_{j0}) C_j \right) \cdot C_{j'} = 0 \quad \text{for every } \varepsilon \text{ and every } j' < i.$$

Therefore, by Lemma 3.1, for every $0 < \varepsilon \ll 1$ and every j ,

$$a_{j1}^+ \varepsilon + a_{j0} \geq a_{j1} \varepsilon + a_{j0},$$

whence $a_{j1}^+ \geq a_{j1}$.

For the second claim, we only need to take care of the case $j' < i$. Then, we have for every ε ,

$$(3.1) \quad \left(D_{t_i+\varepsilon} - \sum_{j=1}^{i-1} (a_{j1}\varepsilon + a_{j0}) C_j \right) \cdot C_{j'} = 0,$$

$$(3.2) \quad \left(D_{t_i+\varepsilon} - \sum_{j=1}^k (a_{j1}^+\varepsilon + a_{j0}) C_j \right) \cdot C_{j'} = 0.$$

Subtracting both equalities, it results

$$\sum_{j=1}^k (a_{j1}^+ - a_{j1}) C_j \cdot C_{j'} = 0, \quad \text{i.e.,} \quad \sum_{j \neq j'} (a_{j1}^+ - a_{j1}) C_j \cdot C_{j'} = (a_{j'1} - a_{j'1}^+) C_{j'}^2.$$

All terms on the left-hand side of the last equality are nonnegative, and if $C_j \cdot C_{j'} > 0$ and $a_{j1}^+ > a_{j1}$, then at least one of them is positive, so the right-hand side must be positive and $a_{j'1}^+ > a_{j'1}$, as claimed. ■

Proposition 3.3. *Let C_1, \dots, C_n be, as above, the irreducible components of N_μ , and let $t_i = \inf\{t \mid C_i \text{ in } N_t\}$ with $v = t_0 \leq t_1 \leq \dots \leq t_{m+1} = \mu$. Fix some i, k with $t_{i-1} < t_i = \dots = t_k < t_{k+1}$.*

The Newton–Okounkov body $\Delta_{Y_\bullet}(D)$ has an interior lower vertex with first coordinate t_i if and only if, for every $\varepsilon > 0$, there is a connected component of $N_{t_i+\varepsilon}$ that goes through p and contains at least one of the C_i, \dots, C_k .

It has an interior upper vertex with first coordinate t_i if and only if, for every $\varepsilon > 0$, there is a connected component of $N_{t_i+\varepsilon}$ that intersects C at a point different from p and contains at least one of the C_i, \dots, C_k .

Proof. As above, C_1, \dots, C_k are the irreducible components of $N_{t_i+\varepsilon}$ for $0 < \varepsilon \ll 1$, and

$$N_{t_i+\varepsilon} = \sum_{j=1}^k (a_{j1}^+ \varepsilon + a_{j0}) C_j \quad \text{if } 0 < \varepsilon \ll 1.$$

Because of the description above of the lower boundary $\alpha(t)$ of $\Delta_{Y_\bullet}(D)$, it is clear that if no component C_i passes through p then $\alpha(t) = 0$ for all $t \in (v, t_i + \varepsilon)$ for small ε , and there is no lower vertex with first coordinate t_i . So assume some component passes through p , and let $J \subset \{1, \dots, k\}$ be such that $\bigcup_{j \in J} C_j$ is the connected component of $N_{t_i+\varepsilon}$ that contains p . We have

$$\alpha(t_i + \varepsilon) = \begin{cases} \sum_{j \in J} (a_{j1} \varepsilon + a_{j0}) (C_j \cdot C)_p & \text{if } -1 \ll \varepsilon \leq 0, \\ \sum_{j \in J} (a_{j1}^+ \varepsilon + a_{j0}) (C_j \cdot C)_p & \text{if } 0 < \varepsilon \ll 1, \end{cases}$$

so there is a lower vertex with first coordinate t_i if and only if $a_{j1}^+ > a_{j1}$ for some j such that C_j passes through p . By Lemma 3.2, this certainly happens if there is some component C_j , $j \in J$, $j \geq i$. Conversely, if all components C_j , $j \in J$, have $j < i$, the equations involving a_{j1} and a_{j1}^+ in (3.1) and (3.2) are equal, so $a_{j1} = a_{j1}^+$ for all $i \in J$, and there is no interior lower vertex with first coordinate t_i .

The proof of the second claim is entirely analogous, and we will be brief. Because of Remark 2.1, if no component C_j meets C at a point different from p then $\alpha(t) = D \cdot C - t C^2$ for all $t \in (v, t_i + \varepsilon)$ for small ε , and there is no upper vertex with first coordinate t_i . On the other hand, if some component C_j does meet C at a point different from p , by Remark 2.1, there is an upper vertex with first coordinate t_i if and only if $a_{j1}^+ > a_{j1}$ for some j such that C_j meets C at a point different from p , and Lemma 3.2 finishes the proof just as in the case of lower vertices. ■

Corollary 3.4. *The number of interior lower (respectively, upper) vertices of $\Delta_{Y_\bullet}(D)$ is bounded above by the number of irreducible components of the connected component $N(p)$ of N_μ that meets C at p (respectively, the number of irreducible components of N_μ in some $N(q)$ for some $q \in C \setminus \{p\}$).*

4. Rightmost vertices

We keep the notation of the previous section, namely D is a big divisor, C an irreducible curve, p a point on C , $D_t = D - tC$, $\mu = \max\{t \mid D_t \text{ pseudo-effective}\}$, $\nu = \nu_C(D)$, $D_t = P_t + N_t$ the Zariski decomposition for $\nu \leq t \leq \mu$, and the irreducible components of N_μ are C_1, \dots, C_n .

Lemma 4.1. *The subspace*

$$V = \langle [D_\nu], [C_1], \dots, [C_n] \rangle \subset \text{NS}(S)_\mathbb{R}$$

has dimension $n + 1$, and the intersection form restricted to V is nondegenerate with signature $(1, n)$.

Proof. The negative part N_ν of the Zariski decomposition $D_\nu = N_\nu + P_\nu$ (if nonzero) satisfies $N_\nu \leq N_\mu$ and hence is a combination of the C_i . Therefore $[P_\nu] \in V$. Since D is big, P_ν is big and nef, and therefore $P_\nu^2 > 0$. As the intersection matrix of the C_i is negative definite, and $[P_\nu] \in V$, it follows that $\dim V = n + 1$ and there is some class $[P] \in V$ orthogonal to all C_i , which moreover has $P^2 > 0$. ■

Proposition 4.2. *If the numerical equivalence class $[C]$ belongs to the subspace*

$$V = \langle [D], [C_1], \dots, [C_n] \rangle \subset \text{NS}(S)_\mathbb{R},$$

then $\Delta_{Y_\bullet}(D)$ has 1 rightmost vertex. If $[C]$ is ample and does not belong to V , then $\Delta_{Y_\bullet}(D)$ has 2 rightmost vertices.

Proof. By the previous lemma, there is a divisor P orthogonal to all C_i , with $P^2 > 0$ and such that $V = \langle [P], [C_1], \dots, [C_n] \rangle$.

If $[C] \in V$, then $[P_\mu] = [D] - \mu[C] - [N_\mu] \in V$, and since P_μ is orthogonal to all C_i , it must be $[P_\mu] = a[P]$ for some $a \in \mathbb{R}$. If $a \neq 0$, then $P_\mu^2 = a^2 P^2 > 0$ and P_μ would be big, contradicting the definition of μ , so $a = 0$ and $P_\mu = 0$. Hence $\alpha(\mu) - \beta(\mu) = P_\mu \cdot C = 0$, which means that $\Delta_Y(D)$ has a single rightmost vertex.

On the other hand, if $[C] \notin V$, then $[P_\mu] = [D] - \mu[C] - [N_\mu] \notin V$, and in particular $[P_\mu] \neq 0$. If moreover $[C]$ is ample, and so belongs to the interior of the nef cone, then its intersection with every nonzero class on the (dual) pseudo-effective cone is positive. Therefore $\alpha(\mu) - \beta(\mu) = P_\mu \cdot C > 0$, which means that $\Delta_Y(D)$ has two rightmost vertices. ■

Corollary 4.3. *If the negative part N_μ of the Zariski decomposition $D_\mu = P_\mu + N_\mu$ has $\rho(S) - 1$ irreducible components, then the polygon $\Delta_{Y_\bullet}(D)$ has exactly one rightmost vertex.*

Proof. By Lemma 4.1, the subspace

$$V = \langle [D], [C_1], \dots, [C_n] \rangle \subset \text{NS}(S)_\mathbb{R}$$

has dimension equal to $\rho(S) - 1 + 1$ and is therefore equal to the whole Néron–Severi space. The claim then follows from Proposition 4.2. ■

5. Counting vertices

Recall from the introduction that for an effective divisor $N = C_1 + \dots + C_n$ with negative definite intersection matrix, $mc(N)$ denotes the largest number of irreducible components of a connected divisor contained in N ,

$$mv(N) = \begin{cases} n + mc(N) + 4 & \text{if } n < \rho(S) - 1, \\ n + mc(N) + 3 & \text{if } n = \rho(S) - 1, \end{cases}$$

and $mv(S)$ is the maximum of all $mv(N)$ for $N = C_1 + \dots + C_n$ negative definite.

First we prove that $mv(S)$ is an upper bound for the number of vertices of every Newton–Okounkov body on S , and then we give a constructive proof that for every ample divisor class D , every number of vertices allowed by the bound is realized by some flag.

Theorem 5.1. *On every smooth projective surface S , for every big divisor D and every flag Y_\bullet , we have that $\#\text{vertices}(\Delta_{Y_\bullet}(D)) \leq mv(S)$.*

Proof. We shall be more precise, showing that if $v = v_C(D)$, $\mu = \mu_C(D)$ and $N = N_\mu = a_1C_1 + \dots + a_nC_n$ is the negative part of the Zariski decomposition of $D_\mu = D - \mu C$, then the number of vertices is bounded by $mv(N)$. By Proposition 3.3, the number of upper interior vertices is bounded by n , and the number of lower interior vertices is bounded by $mc(N)$; the number of leftmost and rightmost vertices is always at most 2, but if $n = \rho(S) - 1$, then by Corollary 4.3 there is exactly 1 rightmost vertex, and the bound follows. ■

Corollary 5.2. *Let S be a smooth projective algebraic surface, D a big divisor and $Y = \{C, p\}$ an admissible flag on S . The polygon $\Delta_{Y_\bullet}(D)$ has at most $2\rho(S) + 1$ vertices.*

For the construction of flags leading to bodies with the desired number of vertices we shall need the following lemma, which may be of independent interest.

Lemma 5.3. *Let $N = C_1 + \dots + C_k$ be an effective divisor with negative definite intersection matrix (admitting $k = 0$, in which case $N = 0$), and let A be an ample divisor. There is an irreducible curve C whose class is ample, such that for every t with $A - tC$ pseudo-effective, the negative part of its Zariski decomposition is supported on N , and moreover, for every $i = 1, \dots, k$,*

- (1) C intersects C_i in at least two points,
- (2) denoting N_t the negative part of the Zariski decomposition of $A - tC$, $\sup\{t \in \mathbb{Q} \mid C_i \text{ is not contained in } N_t\}$ is a finite positive real number t_i , and
- (3) $t_1 < \dots < t_k$.

Moreover, the numerical class of the curve C can be taken in the span $\langle A, C_1, \dots, C_k \rangle$.

Note that the second condition simply means that the support of N_t is exactly N for t large enough. Observe that the lemma still holds when $N = 0$, as the claims in that case are empty.

Proof. We will prove by induction on k that there are positive rational numbers a_1, \dots, a_k such that $B = A - a_1C_1 - \dots - a_kC_k$ is ample, and that every irreducible curve $C \in |mB|$

satisfies the last two desired properties. Since for every ample class B there is a multiple mB and an irreducible curve $C \in |mB|$ that intersects each C_i in at least two points, we shall be done.

If $k = 1$, choose a positive integer a such that the divisor class $B = A - (1/a)C_1$ is ample. Then for every $C \in |mB|$, $A_{1/m} = A - (1/m)C = (m/a)C_1 = N_{1/m}$, so that

$$0 < \sup\{t \in \mathbb{Q} \mid C_1 \text{ is not contained in } N_t\} < 1/m,$$

and we are done.

Now assume the claim is true for the divisor $C_1 + \dots + C_{k-1}$, and let a_1, \dots, a_{k-1} be positive rational numbers such that $B' = A - a_1C_1 - \dots - a_{k-1}C_{k-1}$ is ample and satisfies the two conditions

- (1) denoting N'_t the negative part of the Zariski decomposition of $A - tB'$, $\sup\{t \in \mathbb{Q} \mid C_i \text{ is not contained in } N'_t\}$ is a finite positive real number t'_i , and
- (2) $t'_1 < \dots < t'_{k-1}$.

Of course, this implies the two analogous conditions for $A - tC$ for every $C \in |mB'|$. Note that for every $t \in [0, 1/m]$, since $A_t = (1 - mt)A + mt(a_1C_1 + \dots + a_{k-1}C_{k-1})$, with $(1 - mt)A$ nef and $a_1C_1 + \dots + a_{k-1}C_{k-1}$ effective, by the extremality properties of the Zariski decomposition it follows that $N_t \leq tm(a_1C_1 + \dots + a_{k-1}C_{k-1})$ (with equality if and only if $t = 1/m$). In particular, all components of N_t are among the C_i .

Choose rational numbers s_i with $0 = s_0 < t'_1 < s_1 < t'_2 < \dots < s_{k-2} < t'_{k-1} < s_{k-1} < 1$. The choices made guarantee that the irreducible components of N_{s_i} are exactly C_1, \dots, C_i , and, since for every $i < j \leq k$ we have $N'_{t'_j} \geq N_{s_i}$ and $P'_{t'_j} \cdot C_j = 0$, it follows that $P'_{s_i} \cdot C_j \geq (t'_j - s_i)B' \cdot C_j > 0$ for all $i < j \leq k$. Therefore, by continuity of the Zariski decomposition (see Proposition 1.14 in [2]), there exist $\varepsilon_i > 0$ such that for all $0 < a_k \leq \varepsilon_i$, the irreducible components of the negative part in the Zariski decomposition of $A - s_i(B' - a_kC_k)$ are also exactly C_1, \dots, C_i . Thus it suffices to choose a rational a_k smaller than $\varepsilon_0, \dots, \varepsilon_{k-1}$ and set $B = A - a_1C_1 - \dots - a_kC_k$, because clearly $N_1 = a_1C_1 + \dots + a_kC_k$ and therefore

$$t_{k-1} < s_{k-1} < \sup\{t \in \mathbb{Q} \mid C_k \text{ is not contained in } N_t\} < 1,$$

completing the induction step.

The class B is by construction a combination of A and the C_i , so the class of C belongs to $\langle A, C_1, \dots, C_k \rangle$ as claimed. ■

Lemma 5.4. *Let $N = C_1 + \dots + C_k$ be a maximal effective divisor with negative definite intersection matrix, i.e., such that there exists no curve C' distinct from C_1, \dots, C_k with $N + C'$ having negative definite intersection matrix, and let A be an ample divisor. If $k < \rho(S) - 1$, there is an irreducible curve C satisfying all properties of Lemma 5.3 but whose numerical class is linearly independent of $\langle A, C_1, \dots, C_k \rangle$.*

Proof. Consider the class B from the proof of Lemma 5.3; we can slightly modify B to obtain a B'' which still satisfies the properties and whose numerical class is independent, as follows. Assuming $t_i = \sup\{t \in \mathbb{Q} \mid C_k \text{ is not contained in } N_t\}$ for $i = 1, \dots, k$ as above, choose rational numbers $s_i \in (t_i, t_{i+1})$ and $s_k \in (t_k, 1)$. Let Z be an irreducible curve

whose numerical class is independent of those of C and the C_i . Then $P_{s_i} \geq (1 - s_i)A$, so $P_{s_i} \cdot Z > 0$, and by continuity of the Zariski decomposition, there exist $\varepsilon_i > 0$ such that for all $|b| \leq \varepsilon_i$ the Zariski decomposition of $A - s_i(B + bZ)$ has exactly the components C_1, \dots, C_i in its negative part. The fact that N is maximal guarantees that for no $t > s_k$ any other negative curve appears in N_t . Then the desired class is $B'' = B + bZ$ for some $|b| \leq \varepsilon_i$ for every i . ■

Theorem 5.5. *On every smooth projective surface S , for every ample divisor A and every integer v , $3 \leq v \leq mv(S)$, there exists a flag Y_\bullet such that $\# \text{vertices}(\Delta_{Y_\bullet}(A)) = v$.*

Proof. Choose an effective divisor $N_{mv} = C_1 + \dots + C_k$ with negative definite intersection matrix, such that $mv(S) = mv(N_{mv})$, and assume that its components have been ordered in such a way that for every $1 \leq i \leq mc(N_{mv})$, the divisor $C_1 + \dots + C_i$ is connected. By the definition of $mv(N)$, it is not restrictive to assume that N_{mv} is maximal, i.e., there exists no curve C' with $N_{mv} + C'$ having negative definite intersection matrix.

If $k < \rho(S) - 1$, for every $i \leq j \leq k$, $mv(C_1 + \dots + C_j) = mv(N_{mv}) - k + j$, and for $0 \leq j \leq i$, $mv(C_1 + \dots + C_j) = mv(N_{mv}) - k - i + 2j$. On the other hand, if $k = \rho(S) - 1$, for every $i \leq j < k$, $mv(C_1 + \dots + C_j) = mv(N_{mv}) - k + j + 1$, and for $0 \leq j \leq i$, $mv(C_1 + \dots + C_j) = mv(N_{mv}) - k - i + 2j + 1$. In any event,

$$\{3, \dots, mv(S)\} \subset \bigcup_{N \leq N_{mv}} \{mv(N) - 1, mv(N) - 2\} \cup \{mv(N_{mv})\}.$$

Therefore, it will be enough to prove that, for every N with negative definite intersection matrix,

- if N is maximal, there is a flag Y_\bullet such that $\Delta_{Y_\bullet}(A)$ has $mv(N)$ vertices;
- if N is nonzero or has less than $\rho(s) - 1$ components, there is a flag Y_\bullet such that $\Delta_{Y_\bullet}(A)$ has $mv(N) - 1$ vertices;
- if N is nonzero and has less than $\rho(s) - 1$ components, there is a flag Y_\bullet such that $\Delta_{Y_\bullet}(A)$ has $mv(N) - 2$ vertices.

In the case of a maximal N with less than $\rho(S) - 1$ components, choose an irreducible curve C satisfying the conditions of Lemma 5.4, and let p be one of the intersection points of C and C_1 (unless $N = 0$ in which case we choose an arbitrary $p \in C$). We claim that $A, Y_\bullet: S \supset C \supset \{p\}$ give a body with $mv(N)$ vertices. On the one hand, since A is ample, $P_0 = A$ and $P_0 \cdot C > 0$, so $v = 0$ and $\Delta_{Y_\bullet}(A)$ has two leftmost vertices. Moreover, Proposition 3.3 ensures that $\Delta_{Y_\bullet}(A)$ has two interior vertices with first coordinate equal to the number t_i given by Lemma 5.4 for $i = 1, \dots, mc(N)$, whereas it only has an upper interior vertex for $mc(N) < i \leq k$. Finally, as the numerical class of C is independent of those of A, C_1, \dots, C_k , by Proposition 4.2, $\Delta_{Y_\bullet}(A)$ has two rightmost vertices. So, the total number of vertices is $mv(N)$.

Now choose C verifying the conditions of Lemma 5.3, so that the class of C belongs to the span $\langle A, C_1, \dots, C_k \rangle$. The shape of $\Delta_{Y_\bullet}(A)$ is as before, but with a single rightmost vertex; if N has $\rho(S) - 1$ components (in particular N is maximal) the total number of vertices is $mv(N)$, otherwise it is $mv(N) - 1$.

Finally, if N is nonzero we can pick p differently, while keeping the same curve C that satisfies the conditions of Lemma 5.3. If $mc(N) = 1$, we let p be a point of C not

on N , and if $mc(N) > 1$ then we take p to be one of the intersection points of C with C_2 . In this way we obtain one lower point less, so if N has $\rho(S) - 1$ components the total number of vertices is $mv(N) - 1$, otherwise it is $mv(N) - 2$. ■

Corollary 5.6. *Given integers v, ρ with $3 \leq v \leq 2\rho + 1$, there exist a smooth projective algebraic surface S with Picard number $\rho(S) = \rho$, a big divisor D and an admissible flag $Y = \{C, p\}$ on S , such that $\Delta_{Y_\bullet}(D)$ has v vertices.*

Proof. For $\rho = 1$ there is nothing to prove, since Theorem 5.1 shows that every big divisor and every admissible flag on a surface with Picard number 1 give rise to a triangular Newton–Okounkov body.

So assume $\rho \geq 2$ and pick a surface S_0 with Picard number 1. Construct S by successively blowing up points $p_1, \dots, p_{\rho-1}$, where $p_1 \in S$ and for $i > 1$, p_i is a point on the exceptional divisor of the previous blowup. Then the exceptional divisor of the composition $S \rightarrow S_0$ is a connected divisor N with $\rho - 1$ components and negative definite intersection matrix, and hence $mv(S) = 2\rho + 1$. ■

Note that this construction can be made starting from $S_0 = \mathbb{P}^2$ and selecting each p_i to lie in the strict transform of a fixed line; in that case the resulting surface S is toric (and the Newton–Okounkov polygons obtained by toric flags have $\rho + 2$ vertices, well short of the $2\rho + 1$ vertices that are attainable with our construction).

6. Slopes and lengths of sides

We keep the notation from previous sections, namely D is a big divisor, C is an irreducible curve, p is a point on C , $v = v_C(D)$, $D_t = D - tC$, $\mu = \max\{t \mid D_t \text{ pseudo-effective}\}$, $D_t = P_t + N_t$ is the Zariski decomposition for $0 \leq t \leq \mu$, C_1, \dots, C_n are the irreducible components in order of appearance, $t_i = \inf\{t \in [v, \mu] \mid C_i \text{ in the support of } N_t\}$, and $N_t = \sum a_i(t)C_i$.

Proposition 6.1. *The slopes of the sides of $\Delta_{Y_\bullet}(D)$ are determined by the intersection numbers $C \cdot C_j$ and $C_i \cdot C_j$ and the local intersection numbers $(C \cdot C_j)_p$.*

Proof. Recall from Section 2 that in the interval $[t_{i-1}, t_i]$, $a_j(t)$ can be written as $a_j(t) = a_{j0} + a_{j1}(t)$ satisfying equations (2.1). By looking at the coefficients of t in (2.1), we see that

$$(6.1) \quad \begin{aligned} a_{11} C_1 \cdot C_j + \dots + a_{i-1,1} C_{i-1} \cdot C_j &= -C \cdot C_j, & 1 \leq j < i, \\ a_{j1} &= 0, & i \leq j \leq n. \end{aligned}$$

Thus the coefficients a_{j1} are determined by the intersection numbers $C \cdot C_j$ and $C_i \cdot C_j$, and in the interval $[t_{i-1}, t_i]$ we have

$$\alpha(t) = (N_t \cdot C)_p = \sum_{j=1}^{i-1} (a_{j0} + a_{j1}t)(C_j \cdot C)_p,$$

i.e., the slope of the corresponding lower side of $\Delta_{Y_\bullet}(D)$ is $\sum a_{j1}(C_j \cdot C)_p$, which is determined by the intersection numbers $C \cdot C_j$ and $C_i \cdot C_j$ and the local intersection numbers $(C \cdot C_j)_p$.

On the other hand,

$$\begin{aligned} \beta(t) &= (N_t \cdot C)_p + P_t \cdot C \\ &= \sum_{j=1}^{i-1} (a_{j0} + a_{j1}t)(C_j \cdot C)_p + D \cdot C - \sum_{j=1}^{i-1} (a_{j0} + a_{j1}t)C_j \cdot C, \end{aligned}$$

i.e., the slope of the corresponding upper side of $\Delta_{Y_\bullet}(D)$ is

$$\sum a_{j1}((C_j \cdot C)_p - C_j \cdot C),$$

which is determined by the intersection numbers $C \cdot C_j$ and $C_i \cdot C_j$ and the local intersection numbers $(C \cdot C_j)_p$. ■

For a fixed flag $Y_\bullet: S \supset C \supset \{p\}$ and a fixed negative definite configuration C_1, \dots, C_n , let $\mathcal{D}_C(C_1, \dots, C_n)$ stand for the set of big divisors D such that the negative components of $D - \mu C$ are exactly the C_i numbered by order of appearance in $D - tC$. Proposition 6.1 shows that the bodies $\Delta_{Y_\bullet}(D)$ for $D \in \mathcal{D}_C(C_1, \dots, C_n)$ share the overall shape (number of vertices and slopes of sides), differing only in the lengths of their sides. These lengths are determined by intersection numbers as follows.

Proposition 6.2. *Fix $Y_\bullet: S \supset C \supset \{p\}$ and C_1, \dots, C_n . For every $D \in \mathcal{D}_C(C_1, \dots, C_n)$, the lengths of the sides of $\Delta_{Y_\bullet}(D)$ are determined by μ and the intersection numbers $D \cdot C_j$, and $D \cdot C$.*

Note that to actually compute the lengths of the sides of $\Delta_{Y_\bullet}(D)$, the intersection numbers appearing in Proposition 6.1 are needed too. The claim in Proposition 6.2 can be rephrased saying that the map

$$\begin{aligned} \mathcal{D}_C(C_1, \dots, C_n) &\rightarrow \mathbb{R}^m \\ D &\mapsto (\text{lengths of sides}) \end{aligned}$$

factors through

$$\begin{aligned} \mathcal{D}_C(C_1, \dots, C_n) &\rightarrow \mathbb{R}^{n+1} \\ D &\mapsto (D \cdot C_1, \dots, D \cdot C_n, D \cdot C). \end{aligned}$$

Proof. The length of the leftmost vertical side is $P_\nu \cdot C = P_0 \cdot C$; since the coefficients of $N_0 = \nu C + a_1 C_1 + \dots + a_k C_k$ (where k is the maximum index with $t_k = \nu$) are determined by

$$\begin{aligned} (6.2) \quad &(D - N_0) \cdot C \geq 0, \quad \text{with equality unless } \nu = 0, \\ (6.3) \quad &(D - N_0) \cdot C_i = 0, \quad i = 1, \dots, k, \end{aligned}$$

it is clear that ν is determined by the claimed intersection numbers. Note that k is also determined by the intersection numbers, as the minimum positive integer such that there is a solution of the form $N_0 = \nu C + a_1 C_1 + \dots + a_k C_k$ to (6.2) and (6.3) satisfying $(D - N_0) \cdot C_j \geq 0$ for all $j > k$.

After Proposition 6.1, the slopes of all sides are determined by the fixed data, thus it is enough to prove that the values t_i are determined by the intersection numbers $D \cdot C_j$, and $D \cdot C$. Let us prove this by induction on i . For $i \leq k$, C_i belongs to N_ν , so $t_i = \nu$, and we already showed that k is determined by the intersection numbers. So assume $t_i > t_{i-1} \geq \nu$ and t_1, \dots, t_{i-1} are determined by $D \cdot C_j$, and $D \cdot C$. Then for $t_{i-1} < t \ll t_{i-1} + 1$,

$$N_t = \sum_{j=1}^{i-1} (a_{j0} + a_{j1}t) C_j,$$

and t_i is the infimum of the t such that

$$D - tC - \sum_{j=1}^{i-1} (a_{j0} + a_{j1}t) C_j$$

intersect some $C_{j'}$, $j' \geq i$, negatively. ■

Remark 6.3. Note that the methods developed in Section 5 provide information on the set $\mathcal{D}_C(C_1, \dots, C_n)$. For example, if $C_1 + \dots + C_n$ is a maximal negative definite configuration and all the t_i are distinct, then this set is an open subset of the big cone.

Acknowledgements. We thank the anonymous referees for carefully reading our manuscript and pointing out an inaccuracy in the original statement and proof of Lemma 5.3. We are grateful to Alex Küronya, Julio Moyano-Fernández and Matthias Nickel for helpful discussions.

Funding. We gratefully acknowledge partial support from the Spanish MinECO Grants No. MTM2016-75980-P and PID2020-116542GB-I00, and by National Science Centre grant 2018/ 30/M/ST1/00148.

References

- [1] Bauer, T., Caibăr, M. and Kennedy, G.: Zariski decomposition: a new (old) chapter of linear algebra. *Amer. Math. Monthly* **119** (2012), no. 1, 25–41.
- [2] Bauer, T., Küronya, A. and Szemberg, T.: Zariski chambers, volumes, and stable base loci. *J. Reine Angew. Math.* **576** (2004), 209–233.
- [3] Boucksom, S.: *Corps d’Okounkov (d’après Okounkov, Lazarsfeld–Mustață et Kaveh–Khovanskii)*. Astérisque no. 361 (2014), Exp. no. 1059, vii, 1–41.
- [4] Fulton, W.: *Introduction to toric varieties*. Annals of Mathematics Studies 131, The William H. Roever Lectures in Geometry, Princeton University Press, Princeton, NJ, 1993.
- [5] Harada, M. and Kaveh, K.: Integrable systems, toric degenerations and Okounkov bodies. *Invent. Math.* **202** (2015), no. 3, 927–985.
- [6] Kaveh, K. and Khovanskii, A. G.: Newton–Okounkov bodies, semigroups of integral points, graded algebras and intersection theory. *Ann. of Math. (2)* **176** (2012), no. 2, 925–978.

- [7] Küronya, A. and Lozovanu, V.: Geometric aspects of Newton–Okounkov bodies. In *Phenomenological approach to algebraic geometry*, pp. 137–212. Banach Center Publications 116, Polish Acad. Sci. Inst. Math., Warsaw, 2018.
- [8] Küronya, A., Lozovanu, V. and Maclean, C.: Convex bodies appearing as Okounkov bodies of divisors. *Adv. Math.* **229** (2012), no. 5, 2622–2639.
- [9] Łuszcz-Świdecka, P. and Schmitz, D.: Minkowski decomposition of Okounkov bodies on surfaces. *J. Algebra* **414** (2014), 159–174.
- [10] Lazarsfeld, R.: *Positivity in algebraic geometry. I. Classical setting: line bundles and linear series*. Ergebnisse der Mathematik und ihrer Grenzgebiete 3 Folge 48, Springer-Verlag, Berlin, 2004.
- [11] Lazarsfeld, R. and Mustață, M.: Convex bodies associated to linear series. *Ann. Sci. Éc. Norm. Supér. (4)* **42** (2009), no. 5, 783–835.
- [12] Pokora, P. and Szemberg, T.: Minkowski bases on algebraic surfaces with rational polyhedral pseudo-effective cone. *Electron. Res. Announc. Math. Sci.* **21** (2014), 126–131.

Received August 2, 2020; revised March 14, 2022. Published online August 5, 2022.

Joaquim Roé

Departament de Matemàtiques, Universitat Autònoma de Barcelona,
08193 Bellaterra (Barcelona), Spain;
jroe@mat.uab.cat

Tomasz Szemberg

Department of Mathematics, Pedagogical University Cracow,
Podchorążych 2, 30-084 Kraków, Poland;
tomasz.szemberg@gmail.com