



Global well-posedness for the defocusing, cubic nonlinear Schrödinger equation with initial data in a critical space

Benjamin Dodson

Abstract. In this note we prove global well-posedness for the defocusing, cubic nonlinear Schrödinger equation with initial data lying in a critical Sobolev space.

1. Introduction

In this note, we discuss the defocusing, cubic, nonlinear Schrödinger equation in three dimensions,

$$(1.1) \quad iu_t + \Delta u = F(u) = |u|^2 u, \quad u(0, x) = u_0 \in \dot{H}^{1/2}(\mathbb{R}^3).$$

Equation (1.1) has a scaling symmetry. For any $\lambda > 0$, if u solves (1.1), then

$$(1.2) \quad u_\lambda(t, x) = \lambda u(\lambda^2 t, \lambda x),$$

also solves (1.1). The initial data $\lambda u_0(\lambda x)$ has $\dot{H}^{1/2}(\mathbb{R}^3)$ norm that is invariant under the scaling (1.2).

The local theory for initial data lying in $\dot{H}^{1/2}(\mathbb{R}^3)$ has been completely worked out, and the scaling symmetry has been shown to control the local well-posedness theory.

Theorem 1.1. *Assume $u_0 \in \dot{H}^{1/2}(\mathbb{R}^3)$, $\|u_0\|_{\dot{H}^{1/2}(\mathbb{R}^3)} \leq A$. Then there exists $\delta = \delta(A)$ such that if $\|e^{it\Delta} u_0\|_{L_{t,x}^5(I \times \mathbb{R}^3)} < \delta$, then there exists a unique solution to (1.1) on $I \times \mathbb{R}^3$ with $u \in C(I; \dot{H}^{1/2}(\mathbb{R}^3))$, and*

$$\|u\|_{L_{t,x}^5(I \times \mathbb{R}^3)} \leq 2\delta.$$

Moreover, if $u_{0,k} \rightarrow u_0$ in $\dot{H}^{1/2}(\mathbb{R}^3)$, then the corresponding solutions $u_k \rightarrow u$ in $C(I; \dot{H}^{1/2}(\mathbb{R}^3))$.

This theorem was proved in [3].

From this, it is straightforward to show that local well-posedness holds for (1.1) for any initial data $u_0 \in \dot{H}^{1/2}(\mathbb{R}^3)$. Indeed, by the dominated convergence principle combined

with Strichartz estimates, for any $u_0 \in \dot{H}^{1/2}(\mathbb{R}^3)$,

$$(1.3) \quad \lim_{T \searrow 0} \|e^{it\Delta} u_0\|_{L^5_{t,x}([-T,T] \times \mathbb{R}^3)} = 0.$$

Since $\delta(A)$ is decreasing as $A \nearrow +\infty$, Strichartz estimates imply that there exists $\delta_0 > 0$ such that if $\|u_0\|_{\dot{H}^{1/2}(\mathbb{R}^3)} < \delta_0$, (1.1) has a global solution that scatters. By scattering, we mean that there exist u_0^+, u_0^- so that

$$\lim_{t \rightarrow +\infty} \|u(t) - e^{it\Delta} u_0^+\|_{\dot{H}^{1/2}} = 0,$$

and

$$\lim_{t \rightarrow -\infty} \|u(t) - e^{it\Delta} u_0^-\|_{\dot{H}^{1/2}} = 0.$$

However, it is important to note that while (1.3) holds for any fixed $u_0 \in \dot{H}^{1/2}(\mathbb{R}^3)$, the convergence is not uniform, even for $\|u_0\|_{\dot{H}^{1/2}(\mathbb{R}^3)} \leq A < \infty$. Thus, one cannot conclude directly from [3] that a uniform bound for $\|u(t)\|_{\dot{H}^{1/2}(\mathbb{R}^3)}$ on the entire time of the existence of the solution to (1.1) implies that the solution is global. This result was instead proved in [9], using concentration compactness methods.

Theorem 1.2. *Suppose that u is a solution of (1.1) with initial data $u_0 \in \dot{H}^{1/2}(\mathbb{R}^3)$ and a maximal interval of existence $I = (T_-, T_+)$. Also assume that $\sup_{t \in (T_-, T_+)} \|u(t)\|_{\dot{H}^{1/2}(\mathbb{R}^3)} = A < \infty$. Then $T_+(u_0) = +\infty$, $T_-(u_0) = -\infty$, and the solution u scatters.*

It is conjectured that (1.1) is globally well-posed and scattering for any $u_0 \in \dot{H}^{1/2}(\mathbb{R}^3)$, without the a priori assumption of a universal bound on the $\dot{H}^{1/2}$ norm of the solution $u(t)$. Partial progress has been made in this direction.

A solution to (1.1) has the conserved quantities mass,

$$M(u(t)) = \int |u(t, x)|^2 dx = M(u(0)),$$

and energy,

$$(1.4) \quad E(u(t)) = \frac{1}{2} \int |\nabla u(t, x)|^2 dx + \frac{1}{4} \int |u(t, x)|^4 dx.$$

This fact implies global well-posedness for (1.1) with $u_0 \in H^1_x(\mathbb{R}^3)$, where $H^1_x(\mathbb{R}^3)$ is the inhomogeneous Sobolev space of order one. In this case, one could also prove bounds on the scattering size directly, using the interaction Morawetz estimate of [5].

Theorem 1.3. *If u is a solution to (1.1), on an interval I , then*

$$(1.5) \quad \|u\|_{L^4_{t,x}(I \times \mathbb{R}^3)} \lesssim \|u\|_{L^\infty_t L^2_x(I \times \mathbb{R}^3)} \|u\|_{L^2_t \dot{H}^{1/2}(I \times \mathbb{R}^3)} \lesssim E(u)^{1/2} M(u)^{3/2}.$$

Interpolating (1.4) and (1.5) then implies

$$(1.6) \quad \|u\|_{L^8_t L^4_x(I \times \mathbb{R}^3)} \lesssim M(u)^{3/4} E(u)^{3/4},$$

with bounds independent of $I \subset \mathbb{R}$. Combining Strichartz estimates and local well-posedness theory, a uniform bound on (1.6) for any $I \subset \mathbb{R}$ directly implies a uniform bound on

$$\|u\|_{L_{t,x}^5(I \times \mathbb{R}^3)}.$$

The argument from [3] implies that proving scattering is equivalent to proving

$$(1.7) \quad \|u\|_{L_{t,x}^5(\mathbb{R} \times \mathbb{R}^3)} < \infty.$$

Indeed, assuming that (1.7) is true, the interval \mathbb{R} may be partitioned into finitely many pieces J_k such that

$$\|u\|_{L_{t,x}^5(J_k \times \mathbb{R}^3)} \leq \delta.$$

Then iterate the argument over the intervals J_k , which proves scattering.

This argument also shows that a solution to (1.1) blowing up at a finite time $T_0 < \infty$ is equivalent to

$$\|u\|_{L_{t,x}^5([0, T_0] \times \mathbb{R}^3)} = \infty.$$

Remark. Prior to [5], [8] and [10] proved scattering using the standard Morawetz estimate. See [12] for more details on Strichartz estimates.

Many have attempted to lower the regularity needed in order to prove global well-posedness. For any $s > 1/2$, the inhomogeneous Sobolev space $H_x^s(\mathbb{R}^3) \subset \dot{H}^{1/2}(\mathbb{R}^3)$. Therefore, if $u_0 \in H_x^s(\mathbb{R}^3)$, then it would be conjectured that the solution to (1.1) with initial data u_0 is global and scatters.

Proving a uniform bound on the $H_x^s(\mathbb{R}^3)$ norm would be enough, since by interpolation this would guarantee a uniform bound on the $\dot{H}_x^{1/2}(\mathbb{R}^3)$ norm. The difficulty is that there does not exist a conserved quantity at regularity s that controls the \dot{H}^s norm for $1/2 < s < 1$.

Instead, [2] used the Fourier truncation method (see also [1] for the cubic problem in two dimensions). Decompose the initial data

$$u_0 = P_{\leq N}u_0 + P_{>N}u_0 = v_0 + w_0.$$

Then $v_0 \in H^1(\mathbb{R}^3)$, and $\|w_0\|_{\dot{H}^{1/2}(\mathbb{R}^3)}$ is small. Thus, (1.1) has a global solution for initial data v_0 or w_0 , call them v and w . Since (1.1) is a nonlinear equation, it is necessary to also estimate the interaction between v and w in the nonlinearity of (1.1). Then, [2] proved global well-posedness for (1.1) with initial data $u_0 \in H_x^s(\mathbb{R}^3)$ when $s > 11/13$. Moreover, [2] proved that the solution is of the form

$$e^{it\Delta}u_0 + v(t), \quad \text{where } v(t) \in H_x^1(\mathbb{R}^3).$$

The results from the Fourier truncation method for (1.1) were improved using the I-method. First, [4] improved the regularity necessary for global well-posedness to $s > 5/6$. Then, [5] improved the necessary regularity to $s > 4/5$. To the author’s best knowledge, the best known regularity result is the result of [11], proving global well-posedness and scattering for regularity $s > 5/7$. For radial initial data, [6] proved global well-posedness and scattering for any $s > 1/2$. This result is almost sharp at high frequencies.

In this paper, we study the cubic nonlinear Schrödinger equation (1.1) with initial data lying in the Sobolev space $W_x^{7/6,11/7}(\mathbb{R}^3)$. That is,

$$\| |\nabla|^{11/7} u_0 \|_{L^{7/6}(\mathbb{R}^3)} < \infty.$$

Remark. This norm is well-defined using the Littlewood–Paley decomposition. See for example [13].

This norm is preserved under the scaling (1.2), and is therefore a critical Sobolev norm. Moreover, $W_x^{7/6,11/7}(\mathbb{R}^3) \subset \dot{H}^{1/2}(\mathbb{R}^3)$, so (1.1) has a local solution for this initial data. We prove global well-posedness for (1.1) with this initial data.

Theorem 1.4. *The cubic nonlinear Schrödinger equation is globally well-posed for initial data $u_0 \in W_x^{7/6,11/7}(\mathbb{R}^3)$.*

The proof of this theorem will heavily utilize dispersive estimates. Interpolating between the fact that $e^{it\Delta}$ is a unitary operator,

$$\| e^{it\Delta} u_0 \|_{L^2(\mathbb{R}^3)} = \| u_0 \|_{L^2(\mathbb{R}^3)},$$

and the dispersive estimate,

$$\| e^{it\Delta} u_0 \|_{L^\infty(\mathbb{R}^3)} \lesssim \frac{1}{t^{3/2}} \| u_0 \|_{L^1(\mathbb{R}^3)},$$

gives the estimate

$$(1.8) \quad \| e^{it\Delta} u_0 \|_{L^7(\mathbb{R}^3)} \lesssim \frac{1}{t^{15/14}} \| u_0 \|_{L^{7/6}(\mathbb{R}^3)}.$$

This implies that the linear solution $e^{it\Delta} u_0$ has very good behavior when $t > 1$, in fact it is integrable in time. We then rescale so that u_0 has a local solution on an interval $[-1, 1]$. We prove that this solution may be decomposed into

$$u(t) = e^{it\Delta} u_0 + v(t) + w(t).$$

In particular,

$$u(1) = e^{i\Delta} u_0 + v(1) + w(1).$$

The term

$$e^{i(t-1)\Delta} e^{i\Delta} u_0 = e^{it\Delta} u_0$$

has good properties when $t > 1$. We can also show that

$$\| \nabla e^{i(t-1)\Delta} v(1) \|_{L^\infty} \lesssim \frac{1}{t^{3/2}},$$

which also has good properties when $t > 1$. Finally, $w(1) \in H_x^1$ and has finite energy. Making a Gronwall argument shows that

$$\| u(t) - e^{it\Delta} u_0 - e^{i(t-1)\Delta} v(1) \|_{\dot{H}^1},$$

is uniformly bounded on $[1, \infty)$. This is enough to give global well-posedness, but not scattering.

This result could be compared to the result in [7] for the nonlinear wave equation. There, the author proved global well-posedness and scattering for the cubic wave equation with initial radial data in the Besov space $B_{1,1}^2 \times B_{1,1}^1$. Here, we do not require radial symmetry, however, we only prove global well-posedness. We are unable to prove scattering at this time due to the lack of a scale invariant conformal symmetry.

We prove a local well-posedness result in section two, and a global result in section three. This argument could be generalized to many intercritical, defocusing nonlinear Schrödinger equations.

2. Local well-posedness

The Sobolev embedding theorem implies that $W_x^{7/6,11/7}(\mathbb{R}^3)$ is embedded into $\dot{H}^{1/2}(\mathbb{R}^3)$. Therefore, (1.1) is locally well-posed, and there exists some $T(u_0) > 0$ such that (1.1) has a solution on $[-T, T]$ and $\|u\|_{L_t^5 L_x^5([-T, T] \times \mathbb{R}^3)} = \epsilon_0$, for some $\epsilon_0(\|u_0\|_{\dot{H}^{1/2}})$ small. After rescaling using (1.2), suppose

$$(2.1) \quad \|u\|_{L_{t,x}^5([-1,1] \times \mathbb{R}^3)} = \epsilon_0.$$

Since (3, 18/5) is an admissible pair, Strichartz estimates imply

$$(2.2) \quad \begin{aligned} & \| |\nabla|^{1/2} u \|_{L_t^\infty L_x^2 \cap L_t^2 L_x^6([-1,1] \times \mathbb{R}^3)} \\ & \lesssim \| |\nabla|^{1/2} u_0 \|_{L_x^2(\mathbb{R}^3)} + \| |\nabla|^{1/2} u \|_{L_t^3 L_x^{18/5}([-1,1] \times \mathbb{R}^3)} \|u\|_{L_{t,x}^5([-1,1] \times \mathbb{R}^3)}. \end{aligned}$$

Therefore,

$$(2.3) \quad \| |\nabla|^{1/2} u \|_{L_t^\infty L_x^2 \cap L_t^2 L_x^6([-1,1] \times \mathbb{R}^3)} \lesssim \|u_0\|_{\dot{H}^{1/2}}.$$

Also, by Duhamel’s principle, for any $t \in [-1, 1]$,

$$(2.4) \quad u(t) = e^{it\Delta} u_0 - i \int_0^t e^{i(t-\tau)\Delta} F(u(\tau)) d\tau = u_l(t) + u_{nl}(t).$$

Remark. Recall from (1.1) that $F(u) = |u|^2 u$.

We begin with a technical lemma. This lemma allows us to make a Littlewood–Paley decomposition of u_{nl} , treat each $P_j u_{nl}$ separately, and then sum up. It also implies that u_{nl} retains all the properties of a solution to the linear Schrödinger equation with initial data in a Besov space.

Remark. In this section, all implicit constants depend on the norm $\|u_0\|_{W^{7/6,11/7}}$.

Remark. Throughout this section we rely very heavily on the bilinear Strichartz estimate

$$\|(e^{it\Delta} P_j u_0)(e^{it\Delta} P_k v_0)\|_{L_{t,x}^2(\mathbb{R} \times \mathbb{R}^3)} \lesssim 2^{-j/2} 2^k \|P_j u_0\|_{L^2} \|P_k v_0\|_{L^2}.$$

See [1] for a proof.

Lemma 2.1. *Let P_j be the customary Littlewood–Paley projection operator. Also suppose that u is a solution to (1.1) satisfying (2.1). Then*

$$(2.5) \quad \sum_j 2^{j/2} \|P_j F(u)\|_{L_t^1 L_x^2([-1,1] \times \mathbb{R}^3)} \lesssim 1.$$

Proof. Decompose the nonlinearity,

$$P_j F(u) = P_j F(P_{\geq j-3}u) + 3P_j((P_{\geq j-3}u)^2(P_{\leq j-3}u)) + 3P_j((P_{j-3 \leq \cdot \leq j+3}u)(P_{\leq j-3}u)^2).$$

By Bernstein’s inequality, and (2.2),

$$(2.6) \quad \begin{aligned} 2^{j/2} \|P_j F(P_{\geq j-3}u)\|_{L_t^1 L_x^2([-1,1] \times \mathbb{R}^3)} \\ \lesssim 2^{j/2} \|P_{\geq j-3}u\|_{L_t^3 L_x^6([-1,1] \times \mathbb{R}^3)}^3 \lesssim 2^{j/2} \left(\sum_{l \geq j-3} 2^{-l/6} \|\nabla\|^{1/6} P_l u \|_{L_t^3 L_x^6} \right)^3. \end{aligned}$$

Next,

$$(2.7) \quad \begin{aligned} 2^{j/2} \|P_j((P_{\geq j-3}u)^2(P_{\leq j-3}u))\|_{L_t^1 L_x^2([-1,1] \times \mathbb{R}^3)} \\ \lesssim 2^{j/2} \left(\sum_{l \geq j-3} 2^{-l/4} \|\nabla\|^{1/4} P_l u \|_{L_t^3 L_x^{36/7}} \right)^2 \|u\|_{L_t^3 L_x^9}. \end{aligned}$$

Finally, by the bilinear Strichartz estimate

$$(2.8) \quad \|(e^{it\Delta} P_j u_0)(e^{it\Delta} P_{l_1} u_0)\|_{L_{t,x}^2(\mathbb{R} \times \mathbb{R}^3)} \lesssim 2^{-j/2} 2^{l_1} \|P_j u_0\|_{L^2} \|P_{l_1} u_0\|_{L^2},$$

combined with the principle of superposition and (2.4),

$$(2.9) \quad \begin{aligned} \|(P_j u)(P_{l_1} u)\|_{L_{t,x}^2} \\ \lesssim 2^{-j/2} 2^{l_1} (\|P_j u_0\|_{L^2} + \|P_j F(u)\|_{L_t^1 L_x^2}) (\|P_{l_1} u_0\|_{L^2} + \|P_{l_1} F(u)\|_{L_t^1 L_x^2}), \end{aligned}$$

and the Sobolev embedding properties of Littlewood–Paley projections,

$$(2.10) \quad \begin{aligned} 2^{j/2} \|(P_{j-3 \leq \cdot \leq j+3}u)(P_{\leq j-3}u)^2\|_{L_t^1 L_x^2([-1,1] \times \mathbb{R}^3)} \\ \lesssim 2^{j/2} \sum_{l_1 \leq j-3} \|(P_{l_1} u)(P_{j-3 \leq \cdot \leq j+3}u)\|_{L_{t,x}^2} \sum_{l_1 \leq l_2 \leq j-3} \|P_{l_2} u\|_{L_t^2 L_x^\infty} \\ \lesssim \|\nabla\|^{1/2} u \|_{L_t^2 L_x^6} \sum_{l_1 \leq j-3} 2^{l_1} (j - l_1) (\|P_{j-3 \leq \cdot \leq j+3}u_0\|_{L^2} \\ + \|P_{j-3 \leq \cdot \leq j+3}F(u)\|_{L_t^1 L_x^2}) \cdot (\|P_{l_1} u_0\|_{L^2} + \|P_{l_1} F(u)\|_{L_t^1 L_x^2}). \end{aligned}$$

By Strichartz estimates, (2.3), Plancherel’s theorem, and the fractional product rule,

$$\begin{aligned} \sum_j 2^j \|P_j u_0\|_{L^2}^2 + \sum_j 2^j \|P_j F(u)\|_{L_t^1 L_x^2([-1,1] \times \mathbb{R}^3)}^2 &\lesssim \|u_0\|_{\dot{H}^{1/2}}^2 + \|\nabla\|^{1/2} F(u) \|_{L_t^1 L_x^2}^2 \\ &\lesssim \|u_0\|_{\dot{H}^{1/2}}^2 + \|\nabla\|^{1/2} u \|_{L_t^3 L_x^{18/5}}^2 \|u\|_{L_t^3 L_x^9}^4 \lesssim 1. \end{aligned}$$

Combining (2.6)–(2.10) with the Cauchy–Schwarz inequality implies

$$(2.11) \quad \sum_j 2^{j/2} \|P_j F(u)\|_{L_t^1 L_x^2([-1,1] \times \mathbb{R}^3)} \lesssim 1,$$

which proves the lemma. ■

Next, decompose u_{nl} in the following manner:

$$u_{nl}(t) = -i \int_0^{(1-\delta)t} e^{i(t-\tau)\Delta} F(u(\tau)) d\tau - i \int_{(1-\delta)t}^t e^{i(t-\tau)\Delta} F(u(\tau)) d\tau = v(t) + w(t),$$

for some $\delta > 0$ sufficiently small, to be specified later.

Lemma 2.2. *For any $t \in [0, 1]$,*

$$(2.12) \quad \|v(t)\|_{L^\infty} \lesssim \frac{1}{\delta^{1/2} t^{1/2}},$$

and

$$(2.13) \quad \|\nabla v(t)\|_{L^\infty} \lesssim \frac{1}{\delta t}.$$

Proof. By the dispersive estimate, since $\|u\|_{L^3} \lesssim \|u\|_{\dot{H}^{1/2}}$ is uniformly bounded on $[0, 1]$,

$$\|v(t)\|_{L^\infty} \lesssim \left\| \int_0^{(1-\delta)t} e^{i(t-\tau)\Delta} F(u) d\tau \right\|_{L^\infty} \lesssim \int_0^{(1-\delta)t} \frac{1}{|t-\tau|^{3/2}} \|u\|_{L^3}^3 d\tau \lesssim \frac{1}{\delta^{1/2} t^{1/2}}.$$

To prove (2.13), observe that by the product rule,

$$\nabla F(u) = 2|u|^2 \nabla u + u^2 \nabla \bar{u}.$$

Interpolating,

$$(2.14) \quad \|\nabla^{1/2} u_l\|_{L^2} \lesssim \|\nabla^{1/2} u_0\|_{L^2} \lesssim 1,$$

with

$$(2.15) \quad t^{15/14} \|\nabla^{11/7} u_l\|_{L^7} \lesssim \|\nabla^{11/7} u_0\|_{L^{7/6}} \lesssim 1,$$

we have

$$(2.16) \quad t^{1/2} \|\nabla u_l\|_{L^3} \lesssim 1.$$

Making a dispersive estimate and using (2.16),

$$\begin{aligned} \left\| \int_0^{(1-\delta)t} e^{i(t-\tau)\Delta} |u|^2 \nabla u_l(\tau) d\tau \right\|_{L^\infty} &\lesssim \int_0^{(1-\delta)t} \frac{1}{|t-\tau|^{3/2}} \|\nabla u_l(\tau)\|_{L^3} \|u\|_{L^3}^2 d\tau \\ &\lesssim \int_0^{(1-\delta)t} \frac{1}{|t-\tau|^{3/2}} \frac{1}{|\tau|^{1/2}} d\tau \lesssim \frac{1}{\delta t}. \end{aligned}$$

The same computation may also be made for $u^2 \nabla \bar{u}_l$.

Next, consider the contribution of $|u|^2 \nabla u_{nl}$. By (2.5), we can, without loss of generality, consider only one P_j Littlewood–Paley multiplier, provided the estimate is uniform in $2^{j/2} \|P_j F(u)\|_{L_t^1 L_x^2}$:

$$|u|^2 (\nabla P_j u_{nl}) = |P_{\leq j} u|^2 (\nabla P_j u_{nl}) + 2\operatorname{Re}((P_{> j} \bar{u})(P_{\leq j} \bar{u})) (\nabla P_j u_{nl}) + |P_{> j} u|^2 (\nabla P_j u_{nl}).$$

Using the bilinear Strichartz estimate in (2.9), as well as (2.11) and the Cauchy–Schwartz inequality,

$$\begin{aligned} (2.17) \quad & \| |u_{\leq j}|^2 (\nabla P_j u_{nl}) \|_{L_t^2 L_x^1([0,1] \times \mathbb{R}^3)} \lesssim \sum_{j_1 \leq j_2 \leq j} \| (P_{j_1} u)(P_{j_2} \nabla u_{nl}) \|_{L_{t,x}^2} \| P_{j_2} u \|_{L_t^\infty L_x^2} \\ & \lesssim \sum_{j_1 \leq j_2 \leq j} 2^{j_1/2} 2^{-j_2/2} 2^{j/2} \| P_j F(u) \|_{L_t^1 L_x^2} (\| |\nabla|^{1/2} P_{j_1} u_0 \|_{L^2} + \| |\nabla|^{1/2} P_{j_1} F(u) \|_{L_t^1 L_x^2}) \\ & \quad \times (\| |\nabla|^{1/2} P_{j_2} u_0 \|_{L^2} + \| |\nabla|^{1/2} P_{j_2} F(u) \|_{L_t^1 L_x^2}) \lesssim 1. \end{aligned}$$

Also, by Bernstein’s inequality and Lemma 2.1,

$$\begin{aligned} & \| |\nabla P_j u_{nl}| |P_{> j} u| (|P_{\leq j} u| + |P_{> j} u|) \|_{L_t^2 L_x^1} \\ & \lesssim \| |\nabla|^{1/2} P_j u_{nl} \|_{L_t^2 L_x^5} \| |\nabla|^{1/2} P_{> j} u \|_{L_t^\infty L_x^2} \| u \|_{L_t^\infty L_x^3} \lesssim 1. \end{aligned}$$

Therefore,

$$\begin{aligned} (2.18) \quad & \left\| \int_0^{(1-\delta)t} e^{i(t-\tau)\Delta} |u|^2 \nabla u_{nl}(\tau) d\tau \right\|_{L^\infty} \lesssim \int_0^{(1-\delta)t} \frac{1}{|t-\tau|^{3/2}} \| |u|^2 \nabla u_{nl} \|_{L^1} d\tau \\ & \lesssim \frac{1}{\delta t} \| |u|^2 \nabla u_{nl} \|_{L_t^2 L_x^1} \lesssim \frac{1}{\delta t}. \end{aligned}$$

The same computation can be also be made for $u^2 \nabla \bar{u}_{nl}$. This completes the proof of Lemma 2.2. ■

Lemma 2.3. For any $t \in [0, 1]$,

$$(2.19) \quad \| |\nabla|^{1/2} w(t) \|_{L^3} \lesssim \frac{1}{\delta^{1/4} t^{1/4}}.$$

Proof. First observe that by interpolation, Bernstein’s inequality, and (2.16),

$$(2.20) \quad \| |\nabla|^{1/2} e^{it\Delta} u_0 \|_{L^3} \lesssim t^{1/4} \| \nabla e^{it\Delta} P_{\geq t^{-1/2}} u_0 \|_{L^3} + t^{-1/4} \| P_{\leq t^{-1/2}} u_0 \|_{\dot{H}^{1/2}} \lesssim t^{-1/4}.$$

Also since $e^{it\Delta}$ is unitary in L^2 , by (2.1) and (2.2),

$$(2.21) \quad \| v(t) \|_{\dot{H}^{1/2}} = \| u_{nl}((1-\delta)t) \|_{\dot{H}^{1/2}} \lesssim \| |\nabla|^{1/2} u \|_{L_t^3 L_x^{18/5}} \| u \|_{L_{t,x}^5}^2 \lesssim \epsilon_0^2,$$

so interpolating (2.12), (2.13), and (2.21),

$$(2.22) \quad \| |\nabla|^{1/2} v \|_{L^3} \lesssim \| |\nabla|^{1/2} v \|_{L^\infty}^{1/3} \| |\nabla|^{1/2} v \|_{L^2}^{2/3} \lesssim \frac{\epsilon_0^{4/3}}{\delta^{1/4} t^{1/4}}.$$

Finally, making a dispersive estimate, for any $t \in [0, 1]$, by (2.20) and (2.22), if $\delta^{1/4} \ll \epsilon_0$,

$$\begin{aligned}
 & \delta^{1/4} t^{1/4} \left\| \int_{(1-\delta)t}^t e^{i(t-\tau)\Delta} |\nabla|^{1/2} F(u) \, d\tau \right\|_{L^3} \\
 (2.23) \quad & \lesssim \delta^{1/4} t^{1/4} \int_{(1-\delta)t}^t \frac{1}{|t-\tau|^{1/2}} \|\nabla|^{1/2} u(\tau)\|_{L^3} \|u(\tau)\|_{L^6}^2 \, d\tau \\
 & \lesssim \left(\sup_{t \in [0,1]} \delta^{1/4} t^{1/4} \|\nabla|^{1/2} u\|_{L^3} \right)^3 \lesssim \epsilon_0^4 + \left(\sup_{t \in [0,1]} \delta^{1/4} t^{1/4} \|\nabla|^{1/2} w\|_{L^3} \right)^3.
 \end{aligned}$$

Thus, absorbing the second term on the right-hand side into the left-hand side of (2.23) proves (2.19):

$$\|\nabla|^{1/2} w(t)\|_{L^3} \lesssim \frac{\epsilon_0^4}{\delta^{1/4} t^{1/4}}. \quad \blacksquare$$

Remark. To make the proofs of Lemmas 2.2 and 2.3 completely rigorous, truncate u_0 in frequency. Then the bounds (2.12), (2.13), and (2.19) all hold on some open subset of $[0, 1]$ that contains 0. Making the bootstrap argument using the proof of Lemma 2.3 gives bounds on all of $[0, 1]$ that do not depend on the frequency truncation of u_0 . Standard perturbation arguments then give the lemmas.

Lemma 2.3 can be strengthened to an estimate on the \dot{H}^1 norm of w .

Lemma 2.4. For any $t \in [0, 1]$,

$$\|\nabla w(t)\|_{L^2} \lesssim \frac{1}{\delta^{1/4} t^{1/4}}.$$

Proof. Once again make use of the bilinear Strichartz estimate. Again by the product rule,

$$\nabla F(u) = 2|u|^2 \nabla u + u^2 \nabla \bar{u}.$$

First, by Strichartz estimates, (2.16), Lemma 2.3, and the Sobolev embedding theorem,

$$\begin{aligned}
 & \left\| \int_{(1-\delta)t}^t e^{i(t-\tau)\Delta} [2|u|^2 \nabla u_l + u^2 \nabla \bar{u}_l] \, d\tau \right\|_{L^2} \lesssim \|2|u|^2 \nabla u_l + u^2 \nabla \bar{u}_l\|_{L_t^2 L_x^{6/5}} \\
 & \lesssim \delta^{1/2} t^{1/2} \|\nabla u_l\|_{L_t^\infty L_x^3((1-\delta)t, t] \times \mathbb{R}^3)} \|u\|_{L_t^\infty L_x^3((1-\delta)t, t] \times \mathbb{R}^3)} \|\nabla|^{1/2} u\|_{L_t^\infty L_x^3((1-\delta)t, t] \times \mathbb{R}^3)} \\
 & \lesssim \frac{\delta^{1/4}}{t^{1/4}}.
 \end{aligned}$$

Next, by (2.19), bilinear Strichartz estimates in (2.9), and the Littlewood–Paley theorem,

$$\begin{aligned}
 & \|2|u_{\leq j}|^2 (\nabla P_j u_{nl}) + (u_{\leq j})^2 (\nabla P_j \bar{u}_{nl})\|_{L_t^2 L_x^{6/5}} \\
 & \lesssim \sum_{k \geq 0} 2^{-k/2} \left\| \left(\sum_{j_1 \leq j} 2^{j_1+k} |P_{j_1+k} u|^2 \right)^{1/2} \left(\sum_{j_1 \leq j} 2^{-j_1} 2^{2j} |P_{j_1} u|^2 |P_j u_{nl}|^2 \right)^{1/2} \right\|_{L_t^2 L_x^{6/5}} \\
 & \lesssim \sum_{k \geq 0} 2^{-k/2} \|\nabla|^{1/2} u(t)\|_{L_t^\infty L_x^3((1-\delta)t, t] \times \mathbb{R}^3)} \\
 & \quad \times \left(\sum_{j_1 \leq j} \|P_{j_1} u_0\|_{\dot{H}^{1/2}}^2 + \|P_{j_1} F(u)\|_{L_t^1 L_x^2}^2 \right)^{1/2} 2^{j/2} \|P_j F(u)\|_{L_t^1 L_x^2} \\
 & \lesssim \frac{1}{\delta^{1/4} t^{1/4}} \|\nabla|^{1/2} P_j F(u)\|_{L_t^1 L_x^2}.
 \end{aligned}$$

Next, by Bernstein’s inequality and (2.19)–(2.21),

$$\begin{aligned} & \|(\nabla P_j u_{nl})|_{u \geq j} \|u\| \|_{L_t^2 L_x^{6/5}} \\ & \lesssim \delta^{1/4} t^{1/4} \| |\nabla|^{1/2} u \|_{L_t^\infty L_x^3([(1-\delta)t, t] \times \mathbb{R}^3)}^2 \| |\nabla|^{1/2} P_j u_{nl} \|_{L_t^4 L_x^3([(1-\delta)t, t] \times \mathbb{R}^3)} \\ & \lesssim \frac{1}{\delta^{1/4} t^{1/4}} \| |\nabla|^{1/2} P_j F(u) \|_{L_t^1 L_x^2([0, 1] \times \mathbb{R}^3)}. \end{aligned}$$

Summing up in j using Lemma 2.1 completes the proof. ■

Remark. The above arguments would work equally well in the time interval $[-1, 0]$.

3. Global well-posedness

We are ready to prove Theorem 1.4. The proof will use conservation of the energy (1.4).

Decompose

$$u(1) = \tilde{v}(1) + w(1),$$

where

$$(3.1) \quad \tilde{v}(1) = u_l(1) + v(1),$$

and $w(1)$ is the w in the previous section. Let $T_0 > 1$ be a time value for which we know that (1.1) has a solution on $[0, T_0)$. By standard local well-posedness arguments and we know that such a T_0 exists. Then on $[1, T_0)$, decompose

$$u(t) = \tilde{v}(t) + w(t),$$

where $\tilde{v}(t)$ is the solution to

$$(3.2) \quad (i \partial_t + \Delta) \tilde{v}(t) = 0, \quad \tilde{v}(1) = \tilde{v}(1, x),$$

and $w(t)$ is the solution to

$$(3.3) \quad (i \partial_t + \Delta) w = |u|^2 u, \quad w(1) = w(1, x).$$

Let $E(t)$ denote the energy of w ,

$$E(t) = \frac{1}{2} \int |\nabla w|^2 + \frac{1}{4} \int |w|^4.$$

First observe that Lemma 2.4 and $\|w(1)\|_{\dot{H}^{1/2}} \lesssim 1$ implies that $E(1) < \infty$. The estimate $\|w(1)\|_{\dot{H}^{1/2}}$ is a consequence of Lemma 2.1 and the definition of w . To prove Theorem 1.4, it suffices to prove that for any $T_0 > 1$ such that (1.1) has a solution on $[0, T_0)$,

$$(3.4) \quad \sup_{t \in [1, T_0)} E(t) < \infty.$$

Indeed, by interpolation and the Sobolev embedding theorem, $E(t) < \infty$ implies that $\|w(t)\|_{L^5} < \infty$. Meanwhile, by (2.14)–(2.16), (2.12), and (2.21), $\|\tilde{v}(t)\|_{L^5}$ is uniformly bounded on \mathbb{R} . Therefore, (3.4) implies

$$\|u\|_{L^{5}_{t,x}([0,T_0] \times \mathbb{R}^3)} < \infty.$$

To estimate the growth of $E(t)$, compute the derivative in time of the energy. By (3.3),

$$\frac{d}{dt} E(t) = -\langle \Delta w, w_t \rangle + \langle |w|^2 w, w_t \rangle = \langle |w|^2 w - |u|^2 u, w_t \rangle,$$

where $\langle \cdot, \cdot \rangle$ is the inner product

$$\langle f, g \rangle = \operatorname{Re} \int f(x) \bar{g}(x) dx.$$

By the product rule,

$$\begin{aligned} \langle w_t, |u|^2 u - |w|^2 w \rangle &= \frac{d}{dt} \langle |w|^2 w, \tilde{v} \rangle + \frac{d}{dt} \langle |\tilde{v}|^2, |w|^2 \rangle \\ (3.5) \quad &+ \frac{1}{2} \frac{d}{dt} \operatorname{Re} \int \bar{w}^2 \tilde{v}^2 + \frac{d}{dt} \langle w, |\tilde{v}|^2 \tilde{v} \rangle - 2 \langle \tilde{v}_t \bar{\tilde{v}}, |w|^2 \rangle \\ &- \langle |w|^2 w, \tilde{v}_t \rangle - \operatorname{Re} \int w^2 \bar{\tilde{v}} \tilde{v}_t - 2 \langle w, |\tilde{v}|^2 \tilde{v}_t \rangle - \langle w, \tilde{v}^2 \bar{\tilde{v}}_t \rangle. \end{aligned}$$

Then define the modified energy,

$$\mathcal{E}(t) = E(t) - \langle |w|^2 w, \tilde{v} \rangle - \langle |\tilde{v}|^2, |w|^2 \rangle - \frac{1}{2} \operatorname{Re} \int w^2 \bar{\tilde{v}}^2 - \langle w, |\tilde{v}|^2 \tilde{v} \rangle.$$

By Hölder’s inequality, and the fact that $\|\tilde{v}\|_{L^4} \lesssim_\delta 1$ for all $t \in [1, \infty)$ (again using (2.14)–(2.16), (2.12), and (2.21)),

$$\langle |w|^2 w, \tilde{v} \rangle + \langle |\tilde{v}|^2, |w|^2 \rangle + \frac{1}{2} \operatorname{Re} \int w^2 \bar{\tilde{v}}^2 + \langle w, |\tilde{v}|^2 \tilde{v} \rangle \lesssim E(t)^{3/4} + E(t)^{1/4}.$$

Therefore, when $E(t)$ is large, $E(t) \sim \mathcal{E}(t)$. Since we are attempting to prove a uniform bound for $E(t)$, it is enough to uniformly bound $\mathcal{E}(t)$.

Also, by (3.5),

$$\frac{d}{dt} \mathcal{E}(t) = -\langle |w|^2 w, \tilde{v}_t \rangle - 2 \langle \tilde{v}_t \bar{\tilde{v}}, |w|^2 \rangle - \operatorname{Re} \int w^2 \bar{\tilde{v}} \tilde{v}_t - 2 \langle w, |\tilde{v}|^2 \tilde{v}_t \rangle - \langle w, \tilde{v}^2 \bar{\tilde{v}}_t \rangle.$$

Since \tilde{v} solves (3.2), $\tilde{v}_t = i \Delta \tilde{v} = i \Delta u_t + i \Delta v$.

Lemma 2.2 implies that for any $t > 1$,

$$(3.6) \quad \|v(t)\|_{L^\infty} + \|\nabla v(t)\|_{L^\infty} = \left\| \int_0^{(1-\delta)t} e^{i(t-\tau)\Delta} \langle \nabla \rangle F(u) d\tau \right\|_{L^\infty} \lesssim \frac{1}{\delta^{3/2} t^{3/2}}.$$

Therefore,

$$\langle |w|^2 w, i \Delta v \rangle = -\langle \nabla (|w|^2 w), i \nabla v \rangle \lesssim \|\nabla v\|_{L^\infty} \|\nabla w\|_{L^2} \|w\|_{L^4}^2 \lesssim_\delta \frac{1}{t^{3/2}} E(t).$$

Remark. Since $\delta > 0$ is fixed, we will ignore it from now on.

Also, by Hölder’s inequality and (1.8),

$$\langle i \Delta(e^{it\Delta} u_0), |w|^2 w \rangle \lesssim \| |\nabla|^{11/7} u_t \|_{L^7} \| \nabla w \|_{L^2}^{3/7} \| w \|_{L^4}^{18/7} \lesssim \frac{1}{t^{15/14}} E(t)^{6/7}.$$

This takes care of the contribution of $\langle \tilde{v}_t, |w|^2 w \rangle$.

Next, integrating by parts,

$$(3.7) \quad 2\langle i(\Delta \tilde{v}) \tilde{v}, |w|^2 \rangle = -2\langle i|\nabla \tilde{v}|^2, |w|^2 \rangle - 2\langle i(\nabla \tilde{v}) \tilde{v}, \nabla |w|^2 \rangle = -2\langle i(\nabla \tilde{v}) \tilde{v}, \nabla |w|^2 \rangle.$$

Then by Hölder’s inequality and (3.6), since $\| \tilde{v} \|_{L^4} \lesssim 1$,

$$\langle i(\nabla v) \tilde{v}, \nabla |w|^2 \rangle \lesssim \| \nabla v \|_{L^\infty} \| \tilde{v} \|_{L^4} \| w \|_{L^4} \| \nabla w \|_{L^2} \lesssim \frac{1}{t^{3/2}} E(t)^{3/4}.$$

Also, by Hölder’s inequality and interpolation,

$$(3.8) \quad \langle i(\nabla u_t)(u_t), \nabla |w|^2 \rangle \lesssim \| \nabla u_t \|_{L^\infty_x} \| u_t \|_{L^4} \| \nabla w \|_{L^2} \| w \|_{L^4} \lesssim \frac{1}{t} \frac{1}{t^{1/8}} E(t)^{3/4}.$$

Finally, by (3.6), and Lemma 2.1, which by the Sobolev embedding theorem and the definition of v implies $\| v \|_{L^3} \lesssim 1$

$$(3.9) \quad \langle i(\nabla u_t)v, \nabla |w|^2 \rangle \lesssim \| \nabla u_t \|_{L^\infty_x} \| v \|_{L^3}^{3/4} \| v \|_{L^\infty}^{1/4} \| \nabla w \|_{L^2} \| w \|_{L^4} \lesssim \frac{1}{t} \frac{1}{t^{3/8}} E(t)^{3/4}.$$

In (3.8) and (3.9) we used:

Lemma 3.1. For any $t \geq 0$,

$$(3.10) \quad \| u_t \|_{L^4} \lesssim \frac{1}{t^{1/8}},$$

and

$$(3.11) \quad \| \nabla u_t \|_{L^\infty} \lesssim \frac{1}{t}.$$

Proof. This is proved by interpolating (2.14)–(2.16). By Bernstein’s inequality, (2.15), (2.16), and the Sobolev embedding theorem,

$$(3.12) \quad \| \nabla P_{\leq t^{-1/2}} u_t \|_{L^\infty} + \| \nabla P_{\geq t^{-1/2}} u_t \|_{L^\infty} \lesssim \frac{1}{t}.$$

Also by the Bernstein inequality and the Sobolev embedding theorem, along with (2.16) and $u_t \in \dot{H}^{1/2}$,

$$(3.13) \quad \| P_{\geq t^{-1/2}} u_t \|_{L^4} + \| P_{\leq t^{-1/2}} u_t \|_{L^4} \lesssim \frac{1}{t^{1/8}}.$$

This proves the lemma. ■

The contribution of $2\text{Re} \int w^2 \bar{v} \bar{v}_t$ may be estimated in a similar manner as the contribution of (3.7), except that there is an additional term to consider,

$$-2\text{Re} \int i w^2 (\nabla \bar{v})^2.$$

Interpolating (3.11) with (2.16),

$$-2\text{Re} \int i w^2 (\nabla \bar{u}_l)^2 \lesssim \|\nabla u_l\|_{L^4}^2 \|w\|_{L^4}^2 \lesssim \frac{1}{t^{5/4}} E(t)^{1/2}.$$

Meanwhile, following (2.17) and using Strichartz estimates,

$$\begin{aligned} & \| |u_{\leq j}|^2 (\nabla P_j u_{nl}) \|_{L_t^1 L_x^{3/2}([0,1] \times \mathbf{R}^3)} \lesssim \sum_{j_1 \leq j_2 \leq j} \| (P_{j_1} u) (P_{j_2} \nabla u_{nl}) \|_{L_{t,x}^2} \| P_{j_2} u \|_{L_t^2 L_x^6} \\ & \lesssim \sum_{j_1 \leq j_2 \leq j} 2^{j_1/2} 2^{-j_2/2} 2^{j/2} \| P_j F(u) \|_{L_t^1 L_x^2} (\| |\nabla|^{1/2} P_{j_1} u_0 \|_{L^2} + \| |\nabla|^{1/2} P_{j_1} F(u) \|_{L_t^1 L_x^2}) \\ & \quad \times (\| |\nabla|^{1/2} P_{j_2} u_0 \|_{L^2} + \| |\nabla|^{1/2} P_{j_2} F(u) \|_{L_t^1 L_x^2}) \lesssim 1. \end{aligned}$$

Plugging this estimate into (2.18) implies that for $t > 1$,

$$\left\| \nabla \int_0^{(1-\delta)} e^{i(t-\tau)\Delta} F(u) \right\|_{L_x^3} \lesssim \frac{1}{t^{1/2}}.$$

Interpolating (3.6) with (3.10),

$$-2\text{Re} \int i w^2 (\nabla \bar{v})^2 \lesssim \|\nabla \bar{v}\|_{L^4}^2 \|w\|_{L^4}^2 \lesssim \frac{1}{t^{3/2}} E(t)^{1/2}.$$

Now treat

$$(3.14) \quad 2\langle w, |\bar{v}|^2 \bar{v}_t \rangle + \langle w, \bar{v}^2 \bar{v}_t \rangle = 2\langle w, |\bar{v}|^2 (i \Delta \bar{v}) \rangle + \langle w, \bar{v}^2 (i \Delta \bar{v}) \rangle.$$

After integrating by parts, by (2.13) and (3.11),

$$\begin{aligned} (3.14) & \lesssim \langle |\nabla \bar{v}|^2, |v| |w| \rangle + \langle |\nabla \bar{v}| |\nabla w|, |v|^2 \rangle \\ & \lesssim \|\nabla \bar{v}\|_{L^4}^2 \|\bar{v}\|_{L^4} \|w\|_{L^4} + \|\nabla w\|_{L^2} \|\nabla \bar{v}\|_{L^\infty} \|\bar{v}\|_{L^4}^2 \lesssim \frac{1}{t^{5/4}} E(t)^{1/4} + \frac{1}{t} E(t)^{1/2} \|\bar{v}(t)\|_{L^4}. \end{aligned}$$

Interpolating (3.6) with $\|v\|_{L^3} \lesssim 1$ implies $\|v\|_{L^4} \lesssim t^{-3/8}$. Meanwhile, (3.10) implies $\|u_l\|_{L^4} \lesssim t^{-1/8}$, so therefore, by (3.1), $\|\bar{v}\|_{L^4} \lesssim 1/t^{1/8}$. Therefore, we have proved

$$(3.15) \quad \frac{d}{dt} \mathcal{E}(t) \lesssim \frac{1}{t^{15/14}} (1 + \mathcal{E}(t)).$$

By Gronwall’s inequality, (3.15) implies a uniform bound on $\mathcal{E}(t)$. This implies a uniform bound on $E(t)$, since $E(t) \sim \mathcal{E}(t)$ when $E(t)$ is large, which proves Theorem 1.4.

Acknowledgements. The author is grateful to Frank Merle and Kenji Nakanishi for helpful conversations regarding the defocusing nonlinear Schrödinger equation. The author is also grateful for the helpful comments of several anonymous referees.

Funding. The author was supported by NSF grants DMS-1500424 and DMS-1764358 during the time of writing this paper.

References

- [1] Bourgain, J.: Refinements of Strichartz' inequality and applications to 2D-NLS with critical nonlinearity. *Internat. Math. Res. Notices* **1998** (1998), no. 5, 253–283.
- [2] Bourgain, J.: Scattering in the energy space and below for 3D NLS. *J. Anal. Math.* **75** (1998), no. 1, 267–297.
- [3] Cazenave, T. and Weissler, F. B.: The Cauchy problem for the critical nonlinear Schrödinger equation in H^s . *Nonlinear Anal.* **14** (1990), no. 10, 807–836.
- [4] Colliander, J., Keel, M., Staffilani, G., Takaoka, H. and Tao, T.: Almost conservation laws and global rough solutions to a Nonlinear Schrödinger equation *Math. Res. Lett.* **9** (2002), no. 5-6, 659–682.
- [5] Colliander, J., Keel, M., Staffilani, G., Takaoka, H. and Tao, T.: Global existence and scattering for rough solutions of a nonlinear Schrödinger equation on \mathbb{R}^3 . *Comm. Pure Appl. Math.* **57** (2004), no. 8, 987–1014.
- [6] Dodson, B.: Global well-posedness and scattering for nonlinear Schrödinger equations with algebraic nonlinearity when $d = 2, 3$ and u_0 is radial. *Camb. J. Math.* **7** (2019), no. 3, 283–318.
- [7] Dodson, B.: Global well-posedness and scattering for the radial, defocusing, cubic wave equation with initial data in a critical Besov space. *Anal. PDE* **12** (2019), no. 4, 1023–1048.
- [8] Ginibre, J. and Velo, G.: On a class of nonlinear Schrödinger equations. II. Scattering theory, general case. *J. Functional Analysis* **32** (1979), no. 1, 33–71.
- [9] Kenig, C. and Merle, F.: Scattering for $\dot{H}^{1/2}$ bounded solutions to the cubic, defocusing NLS in 3 dimensions. *Trans. Amer. Math. Soc.* **362** (2010), no. 4, 1937–1962.
- [10] Lin, J.-E. and Strauss, W. A.: Decay and scattering of solutions of a nonlinear Schrödinger equation. *J. Functional Analysis* **30** (1978), no. 2, 245–263.
- [11] Su, Q.: Global well-posedness and scattering for defocusing, cubic NLS in \mathbb{R}^3 . *Math. Res. Lett.* **19** (2012), no. 2, 431–451.
- [12] Tao, T.: *Nonlinear dispersive equations. Local and global analysis*. CBMS Regional Conference Series in Mathematics 106, American Mathematical Society, Providence, RI, 2006.
- [13] Taylor, M.: *Partial differential equations III. Nonlinear equations*. Second edition. Applied Mathematical Sciences 117, Springer, New York, 2011.

Received October 16, 2020; revised April 15, 2021. Published online July 6, 2021.

Benjamin Dodson

Department of Mathematics, Johns Hopkins University, N. Charles Street, Baltimore,
MD 21218, USA;
bdodson4@jhu.edu