



Mathematics — *On doubly nonlocal fractional elliptic equations*, by GIOVANNI MOLICA BISCI and DUŠAN REPOVŠ, communicated on 13 February 2015.

ABSTRACT. — This work is devoted to the study of the existence of solutions to nonlocal equations involving the fractional Laplacian. These equations have a variational structure and we find a nontrivial solution for them using the Mountain Pass Theorem. To make the nonlinear methods work, some careful analysis of the fractional spaces involved is necessary. In addition, we require rather general assumptions on the local operator. As far as we know, this result is new and represent a fractional version of a classical theorem obtained working with Laplacian equations.

KEY WORDS: Nonlocal problems, fractional equations, Mountain Pass Theorem.

MATHEMATICS SUBJECT CLASSIFICATION: Primary: 49J35, 35S15; Secondary: 47G20, 45G05.

1. INTRODUCTION

As is well-known, nonlocal Laplacian boundary value problems model several physical and biological systems where u describes a process which depends on the average of itself, for example, the population density (see [1, 4, 13, 14, 34]). In the vast literature on this subject, we also refer the reader to some interesting results obtained by Autuori and Pucci in [5, 6, 7] studying Kirchhoff equations by using different approaches.

Very recently, the nonlocal fractional counterpart of Kirchhoff-type problems has been considered (see [12] and [17, 22]). In this order of ideas, we are interested here on the existence of weak solutions for the following (doubly) nonlocal problem:

$$(D_{M,f}) \quad \begin{cases} M\left(\int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy\right) (-\Delta)^s u = f(x, u) & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

where Ω is a bounded domain in $(\mathbb{R}^n, |\cdot|)$ with smooth boundary $\partial\Omega$, $s \in (0, 1)$ is fixed with $s < n/2$ and $(-\Delta)^s$ is the fractional Laplace operator, which (up to normalization factors) can be defined as

$$(-\Delta)^s u(x) := - \int_{\mathbb{R}^n} \frac{u(x+y) + u(x-y) - 2u(x)}{|y|^{n+2s}} dy, \quad x \in \mathbb{R}^n.$$

Further, $f : \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ and $M : [0, +\infty) \rightarrow [0, +\infty)$ are suitable continuous maps.

In our context, problem $(D_{M,f})$ is studied by exploiting classical variational methods. More precisely, we apply the celebrated Mountain Pass Theorem (abbreviated MPT) to this kind of equations motivated by the current literature where the MPT has been intensively applied to find solutions to quasilinear elliptic equations (see [3, 25, 27, 33]).

Technically, this approach is realizable by checking that the associated energy functional satisfies the usual compactness Palais-Smale condition in a suitable variational setting developed by Servadei and Valdinoci (see [30]). Indeed, the nonlocal analysis which we perform in this paper in order to use the Mountain Pass Theorem is quite general and it was successfully exploited for other goals in several recent contributions (see [16, 19, 20, 21, 23, 28, 30, 31, 32] and [15] for various properties on the fractional Sobolev space setting).

This functional analytical context was inspired by (but is not equivalent to) the fractional Sobolev spaces, in order to correctly encode the Dirichlet boundary datum in the variational formulation.

In our context, to avoid some additional technical difficulties due to the presence of the term

$$M\left(\int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy\right),$$

we impose some restrictions on the behavior of the continuous map M .

More precisely, we require that there exists a constant m_0 such that:

$$(C_M^1) \quad 0 < m_0 \leq M(t), \quad \forall t \in [0, +\infty).$$

In addition to the above hypothesis, we assume that:

$$(C_M^2) \quad \text{There exists } t_0 \geq 0 \text{ such that}$$

$$\hat{M}(t) \geq tM(t),$$

$$\text{for every } t \in [t_0, +\infty), \text{ where } \hat{M}(t) := \int_0^t M(s) ds.$$

The above conditions ensure, as proved in Lemma 1, that the potential \hat{M} has a sublinear growth. Under the previous assumptions, by imposing conditions on the nonlinear part f (among others, the Ambrosetti-Rabinowitz relation) we prove the existence of at least one nontrivial weak solution to problem $(D_{M,f})$, see Theorem 2.

This result is related to [30, Theorem 2] where the authors studied a local problem involving a general integro-differential operator of fractional type (see Remark 3) whose prototype is

$$(D_f) \quad \begin{cases} (-\Delta)^s u = f(x, u) & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

We just observe that in our context, in contrast with the cited result, we don't require that

$$\lim_{t \rightarrow 0} \frac{f(x, t)}{t} = 0,$$

uniformly with respect to $x \in \bar{\Omega}$ (see condition (1) in Theorem 2 and Remark 4 below).

Moreover, our main theorem extends to the nonlocal setting a result, already known in the literature for Kirchhoff-type problems obtained by Alves, Corrêa and Ma [2, Theorem 3]. We just point out that M , in the original meaning for Kirchhoff equation, is an increasing function, thus condition (C_M^2) is clearly violated.

We mention, for completeness, that the existence and multiplicity of solutions for elliptic equations in \mathbb{R}^n , driven by a nonlocal integro-differential operator, whose standard prototype is the fractional Laplacian, have been studied by Autuori and Pucci in [9] (this work is related to the results on general quasilinear elliptic problems given in [8]). See also the relevant contributions [10, 24, 26] where Kirchhoff-type problems have been studied by using different methods and approaches.

The plan of the paper is as follows. Section 2 is devoted to our abstract framework and preliminaries. Successively, in Sections 3 we give the main result (see Theorem 2). Finally, a concrete example of an application is presented in the last part of the paper (see Example 1).

2. ABSTRACT FRAMEWORK

This section is devoted to the notations used throughout the paper. We also list some preliminary results which will be useful in the sequel.

Let $H^s(\mathbb{R}^n)$ be the usual fractional Sobolev space endowed with the norm (the so-called *Gagliardo norm*)

$$\|g\|_{H^s(\mathbb{R}^n)} = \|g\|_{L^2(\mathbb{R}^n)} + \left(\int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|g(x) - g(y)|^2}{|x - y|^{n+2s}} dx dy \right)^{1/2}.$$

and defined as the linear space of functions $g \in L^2(\mathbb{R}^n)$ such that

$$\text{the map } (x, y) \mapsto \frac{g(x) - g(y)}{|x - y|^{n/2+s}} \text{ is in } L^2(\mathbb{R}^n \times \mathbb{R}^n, dx dy).$$

Let us consider the subspace $X_0 \subset H^s(\mathbb{R}^n)$ given by

$$X_0 := \{g \in H^s(\mathbb{R}^n) : g = 0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega\}.$$

Of course, the space X_0 is non-empty, since $C_0^2(\Omega) \subseteq X_0$ by [29, Lemma 11] and it depends on the set Ω .

Moreover, by [30, Lemma 6] and the fact that any function $v \in X_0$ is such that $v = 0$ a.e. in $\mathbb{R}^n \setminus \Omega$, we can take in the sequel

$$X_0 \ni v \mapsto \|v\|_{X_0} := \left(\int_{\mathcal{Q}} \frac{|v(x) - v(y)|^2}{|x - y|^{n+2s}} dx dy \right)^{1/2}$$

as norm on X_0 , where $\mathcal{Q} := (\mathbb{R}^n \times \mathbb{R}^n) \setminus (\mathcal{C}\Omega \times \mathcal{C}\Omega)$, and $\mathcal{C}\Omega := \mathbb{R}^n \setminus \Omega$.

Also $(X_0, \|\cdot\|_{X_0})$ is a Hilbert space (for this see [30, Lemma 7]), with the scalar product

$$\langle u, v \rangle_{X_0} := \int_{\mathcal{Q}} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{n+2s}} dx dy.$$

We recall that in [30, Lemma 8] and [32, Lemma 9] the authors proved that the embedding $j : X_0 \hookrightarrow L^v(\mathbb{R}^n)$ is continuous for any $v \in [1, 2^*]$, while it is compact whenever $v \in [1, 2^*)$, where $2^* := 2n/(n - 2s)$ denotes the Sobolev fractional exponent.

Hence, for any $v \in [1, 2^*)$, there exists $c_v > 0$ such that

$$\|v\|_{L^v(\mathbb{R}^n)} \leq c_v \|v\|_{X_0},$$

for every $v \in X_0$.

In the sequel, we will denote by $\lambda_{1,s}$ the first (simple and positive) eigenvalue of the operator $(-\Delta)^s$ with homogeneous Dirichlet boundary data, namely the first eigenvalue of the problem

$$\begin{cases} (-\Delta)^s u = \lambda u & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

that can be characterized as follows

$$\lambda_{1,s} = \min \left\{ \frac{\int_{\mathcal{Q}} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy}{\int_{\Omega} |u(x)|^2 dx} : u \in X_0 \setminus \{0_{X_0}\} \right\}.$$

For the existence and the basic properties of this eigenvalue we refer to [31, Proposition 9 and Appendix A], where a spectral theory for general integro-differential nonlocal operators was developed. Further properties can be also found in [28].

Finally, for the sake of completeness, we recall that a C^1 -functional $J : E \rightarrow \mathbb{R}$, where E is a real Banach space with topological dual E^* , satisfies the *Palais-Smale condition at level* $\mu \in \mathbb{R}$, (abbreviated $(PS)_{\mu}$) when:

$(PS)_{\mu}$ Every sequence $\{u_j\}$ in E such that

$$J(u_j) \rightarrow \mu, \quad \text{and} \quad \|J'(u_j)\|_{E^*} \rightarrow 0,$$

as $j \rightarrow \infty$, possesses a convergent subsequence.

We say that J satisfies the *Palais-Smale condition* (abbreviated (PS)) if $(PS)_\mu$ holds for every $\mu \in \mathbb{R}$.

With the above notation, our main tool is the classical MPT:

THEOREM 1. *Let $(E, \|\cdot\|_E)$ be a real Banach space and let $J \in C^1(E; \mathbb{R})$ be such that $J(0_E) = 0$ and it satisfies the (PS) condition. Suppose that:*

- (I₁) *There exist constants $\rho, \alpha > 0$ such that $J(u) \geq \alpha$ if $\|u\|_E = \rho$.*
- (I₂) *There exists $e \in E$ with $\|e\|_E > \rho$ such that $J(e) \leq 0$.*

Then J possesses a critical value $c \geq \alpha$, which can be characterized as

$$c := \inf_{\gamma \in \Gamma} \max_{u \in \gamma([0,1])} J(u),$$

where

$$\Gamma := \{\gamma \in C([0, 1]; E) : \gamma(0) = 0 \wedge \gamma(1) = e\}.$$

See [27, p. 7; Theorem 2.2].

We cite the monograph [18] as general reference for the variational setting adopted in this paper.

3. THE MAIN RESULT

Our main result is as follows.

THEOREM 2. *Let us assume that $M : [0, +\infty) \rightarrow [0, +\infty)$ is a continuous map such that conditions (C_M^1) and (C_M^2) hold. Further, require that $f : \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function which satisfies the following requirements:*

h₁) *The subcritical growth condition:*

$$|f(x, t)| \leq c(1 + |t|^{q-1}), \quad (\forall x \in \bar{\Omega}, \forall t \in \mathbb{R})$$

where $c > 0$ and $2 < q < 2^*$;

h₂) *The Ambrosetti-Rabinowitz (abbreviated (AR)) condition:*

$$F(x, \xi) := \int_0^\xi f(x, t) dt$$

is θ -superhomogeneous at infinity, i.e. there exists $t^ > 0$ such that*

$$0 < \theta F(x, \xi) \leq f(x, \xi)\xi, \quad (\forall x \in \bar{\Omega}, \forall |\xi| \geq t_*)$$

where $\theta > 2$.

We also assume that

$$(1) \quad \limsup_{t \rightarrow 0} \frac{f(x, t)}{t} \leq \lambda,$$

uniformly for $x \in \bar{\Omega}$, where

$$\lambda < m_0 \lambda_{1,s}.$$

Then the nonlocal problem

$$(D_{M,f}) \quad \begin{cases} M\left(\int_{\Omega} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy\right) (-\Delta)^s u = f(x, u) & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

has at least one nontrivial weak solution.

We recall that a *weak solution* of problem $(D_{M,f})$ is a function $u \in X_0$ such that

$$\begin{aligned} M\left(\int_{\Omega} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy\right) \int_{\Omega} \frac{(u(x) - u(y))(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dx dy \\ = \int_{\Omega} f(x, u(x))\varphi(x) dx, \quad \forall \varphi \in X_0. \end{aligned}$$

3.1. Some remarks on our assumptions

The validity of the next lemma will be crucial in the sequel.

LEMMA 1. *Suppose that conditions (C_M^1) and (C_M^2) hold. Then, there are two positive constants m_1 and m_2 such that*

$$(2) \quad \hat{M}(t) \leq m_1 t + m_2,$$

for every $t \in [0, +\infty)$.

PROOF. Let $t_1 > t_0$, where t_0 appears in hypothesis (C_M^2) . By our assumptions we easily have

$$\frac{M(t)}{\hat{M}(t)} \leq \frac{1}{t},$$

for every $t \in [t_1, +\infty)$. Integrating the above relation, we obtain

$$\int_{t_1}^t \frac{M(s)}{\hat{M}(s)} ds = \log \frac{\hat{M}(t)}{\hat{M}(t_1)} \leq \log \frac{t}{t_1},$$

for every $t \in]t_1, +\infty)$. Thus

$$\hat{M}(t) \leq \frac{\hat{M}(t_1)}{t_1} t,$$

for every $t \in]t_1, +\infty)$. Hence the growth condition (2) holds by taking, for instance, $m_1 := \frac{\hat{M}(t_1)}{t_1}$ and $m_2 := \max_{t \in [0, t_1]} \hat{M}(t)$. The proof is complete.

Owing to conditions (C_M^1) and (C_M^2) , by Lemma 1 one gets the following inequalities:

$$(\hat{C}_M) \quad m_0 \frac{\|u\|_{X_0}^2}{2} \leq \frac{1}{2} \hat{M} \left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy \right) \leq m_1 \frac{\|u\|_{X_0}^2}{2} + \frac{m_2}{2}$$

for every $u \in X_0$.

REMARK 1. We observe that $h_2)$ implies that

$$F(x, \tau \xi) \geq F(x, \xi) \tau^\theta,$$

for every $x \in \bar{\Omega}$, $|\xi| \geq t_*$ and $\tau \geq 1$. Indeed, for $\tau = 1$, clearly the equality holds. Otherwise, fix $|\xi| \geq t_*$ and define $g(x, \tau) := F(x, \tau \xi)$, for every $x \in \bar{\Omega}$ and $\tau \in]1, +\infty)$. By (AR) condition it follows that

$$\frac{g'(x, \tau)}{g(x, \tau)} \geq \frac{\theta}{\tau},$$

for every $x \in \bar{\Omega}$ and $\tau > 1$. By integrating in $]1, \tau]$ we get that

$$\int_1^\tau \frac{g'(x, s)}{g(x, s)} ds = \log \frac{g(x, \tau)}{g(x, 1)} \geq \log \tau^\theta.$$

In conclusion, since for every $x \in \bar{\Omega}$, $|\xi| \geq t_*$ and $\tau > 1$ one has

$$\begin{aligned} F(x, \tau \xi) &=: g(x, \tau) \\ &\geq g(x, 1) \tau^\theta \\ &= F(x, \xi) \tau^\theta, \end{aligned}$$

the claim is verified.

3.2. Proof of Theorem 2

Set

$$\Phi(u) := \frac{1}{2} \hat{M} \left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy \right)$$

for every $u \in X_0$.

Under the assumptions of Theorem 2, we define the C^1 -functional

$$J(u) := \Phi(u) - \int_{\Omega} F(x, u(x)) \, dx, \quad (\forall u \in X_0)$$

where

$$F(x, u(x)) := \int_0^{u(x)} f(x, s) \, ds,$$

whose critical points are the weak solutions of problem $(D_{M,f})$.

In order to prove our result, we apply Theorem 1 to this functional. In the next three lemmas we shall verify the Mountain Pass Theorem conditions.

LEMMA 2. *Every Palais-Smale sequence for the functional J is bounded in X_0 .*

PROOF. Let $\{u_j\} \subset X_0$ be a Palais-Smale sequence i.e.

$$(3) \quad J(u_j) \rightarrow \mu,$$

for $\mu \in \mathbb{R}$ and

$$(4) \quad \|J'(u_j)\|_{X_0^*} = \sup\{|\langle J'(u_j), \varphi \rangle| : \varphi \in X_0, \|\varphi\|_{X_0} = 1\} \rightarrow 0,$$

as $j \rightarrow \infty$.

We argue by contradiction. So, suppose to the contrary that the conclusion were not true. Passing to a subsequence if necessary, we may assume that

$$\|u_j\|_{X_0} \rightarrow +\infty,$$

as $j \rightarrow \infty$.

By conditions (C_M^1) and (C_M^2) , it follows that there exists $j_0 \in \mathbb{N}$ such that

$$\begin{aligned} J(u_j) - \frac{\langle J'(u_j), u_j \rangle}{\theta} &= M(\|u_j\|_{X_0}^2) \left[\frac{\hat{M}(\|u_j\|_{X_0}^2)}{2M(\|u_j\|_{X_0}^2)} - \frac{\|u_j\|_{X_0}^2}{\theta} \right] \\ &\quad + \int_{\Omega} \left[\frac{f(x, u_j(x))u_j(x)}{\theta} - F(x, u_j(x)) \right] dx, \\ &\geq m_0 \left(\frac{1}{2} - \frac{1}{\theta} \right) \|u_j\|_{X_0}^2 \\ &\quad + \int_{\Omega} \left[\frac{f(x, u_j(x))u_j(x)}{\theta} - F(x, u_j(x)) \right] dx, \end{aligned}$$

for every $j \geq j_0$.

Thus

$$\begin{aligned}
 m_0 \left(\frac{\theta - 2}{2\theta} \right) \|u_j\|_{X_0}^2 &\leq J(u_j) - \frac{\langle J'(u_j), u_j \rangle}{\theta} \\
 &\quad - \int_{|u_j(x)| > t_*} \left[\frac{f(x, u_j(x))u_j(x)}{\theta} - F(x, u_j(x)) \right] dx, \\
 &\quad + M \operatorname{meas}(\Omega), \quad \forall j \geq j_0,
 \end{aligned}$$

where “ $\operatorname{meas}(\Omega)$ ” denotes the standard Lebesgue measure of Ω and

$$M := \sup \left\{ \left| \frac{f(x, t)t}{\theta} - F(x, t) \right| : x \in \bar{\Omega}, |t| \leq t_* \right\}.$$

Now, we observe that, the (AR) condition yields

$$\int_{|u_j(x)| > t_*} \left[\frac{f(x, u_j(x))u_j(x)}{\theta} - F(x, u_j(x)) \right] dx \geq 0.$$

So, we deduce that

$$m_0 \left(\frac{\theta - 2}{2\theta} \right) \|u_j\|_{X_0}^2 \leq J(u_j) - \frac{\langle J'(u_j), u_j \rangle}{\theta} + M \operatorname{meas}(\Omega),$$

for every $j \geq j_0$.

Then, for every $j \geq j_0$ one has

$$C \|u_j\|_{X_0}^2 \leq J(u_j) + \theta \|J'(u_j)\|_{X_0^*} \|u_j\|_{X_0} + M \operatorname{meas}(\Omega),$$

where $C := m_0 \left(\frac{\theta - 2}{2\theta} \right) > 0$.

In conclusion, dividing by $\|u_j\|_{X_0}$ and letting $j \rightarrow \infty$, we obtain a contradiction.

The above lemma implies that the C^1 -functional J satisfies the Palais-Smale condition as proved in the next result.

LEMMA 3. *The functional J satisfies the compactness (PS) condition.*

PROOF. Let $\{u_j\} \subset X_0$ be a Palais-Smale sequence. By Lemma 2, the sequence $\{u_j\}$ is necessarily bounded in X_0 . Since X_0 is reflexive, we can extract a subsequence which for simplicity we shall call again $\{u_j\}$, such that $u_j \rightharpoonup u_\infty$ in X_0 . This means that

$$\begin{aligned}
 (5) \quad &\int_{\mathcal{Q}} \frac{(u_j(x) - u_j(y))(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dx dy \\
 &\rightarrow \int_{\mathcal{Q}} \frac{(u_\infty(x) - u_\infty(y))(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dx dy,
 \end{aligned}$$

for any $\varphi \in X_0$, as $j \rightarrow +\infty$.

We will prove that u_j strongly converges to $u_\infty \in X_0$. Exploiting the derivative $J'(u_j)(u_j - u_\infty)$, we obtain

$$(6) \quad \langle a(u_j), u_j - u_\infty \rangle = \langle J'(u_j), u_j - u_\infty \rangle + \int_{\Omega} f(x, u_j(x))(u_j - u_\infty)(x) dx,$$

where we set

$$\begin{aligned} \langle a(u_j), u_j - u_\infty \rangle &:= \left(\int_Q \frac{|u_j(x) - u_j(y)|^2}{|x - y|^{n+2s}} dx dy \right. \\ &\quad \left. - \int_Q \frac{(u_j(x) - u_j(y))(u_\infty(x) - u_\infty(y))}{|x - y|^{n+2s}} dx dy \right) \\ &\quad \times M \left(\int_Q \frac{|u_j(x) - u_j(y)|^2}{|x - y|^{n+2s}} dx dy \right). \end{aligned}$$

Since $\|J'(u_j)\|_{X_0^*} \rightarrow 0$ and the sequence $\{u_j - u_\infty\}$ is bounded in X_0 , taking into account that $|\langle J'(u_j), u_j - u_\infty \rangle| \leq \|J'(u_j)\|_{X_0^*} \|u_j - u_\infty\|_{X_0}$, one gets

$$(7) \quad \langle J'(u_j), u_j - u_\infty \rangle \rightarrow 0,$$

as $j \rightarrow \infty$.

Now observe that, by condition h_1), it follows that

$$\begin{aligned} &\int_{\Omega} |f(x, u_j(x))| |u_j(x) - u_\infty(x)| dx \\ &\leq c \left(\int_{\Omega} |u_j(x) - u_\infty(x)| dx + \int_{\Omega} |u_j(x)|^{q-1} |u_j(x) - u_\infty(x)| dx \right) \\ &\leq c((\text{meas}(\Omega))^{1/q'} + \|u_j\|_{L^q(\Omega)}^{q-1}) \|u_j - u_\infty\|_{L^q(\Omega)}, \end{aligned}$$

where, as usual, q' denotes the conjugate of q .

Since the embedding $X_0 \hookrightarrow L^q(\Omega)$ is compact, clearly $u_j \rightarrow u_\infty$ strongly in $L^q(\Omega)$. So we obtain

$$(8) \quad \int_{\Omega} |f(x, u_j(x))| |u_j(x) - u_\infty(x)| dx \rightarrow 0.$$

By (6), relations (7) and (8) yield

$$(9) \quad \langle a(u_j), u_j - u_\infty \rangle \rightarrow 0,$$

when $j \rightarrow \infty$.

Bearing in mind condition (C_M^1) one obtains

$$(10) \quad 0 < m_0 \leq M \left(\int_Q \frac{|u_j(x) - u_j(y)|^2}{|x - y|^{n+2s}} dx dy \right),$$

for every $j \in \mathbb{N}$.

Hence by (10) and (9) we can write

$$(11) \quad \int_Q \frac{|u_j(x) - u_j(y)|^2}{|x - y|^{n+2s}} dx dy - \int_Q \frac{(u_j(x) - u_j(y))(u_\infty(x) - u_\infty(y))}{|x - y|^{n+2s}} dx dy \rightarrow 0,$$

when $j \rightarrow \infty$.

Thus, by (11) and (5) it follows that

$$\limsup_{j \rightarrow \infty} \int_Q \frac{|u_j(x) - u_j(y)|^2}{|x - y|^{n+2s}} dx dy = \int_Q \frac{|u_\infty(x) - u_\infty(y)|^2}{|x - y|^{n+2s}} dx dy.$$

In conclusion, thanks to [11, Proposition III.30], $u_j \rightarrow u_\infty$ in X_0 . The proof is thus complete.

LEMMA 4. *The functional J has the geometry of the Mountain Pass Theorem. More precisely:*

1. *There exists $r > 0$ such that*

$$\inf_{\|u\|_{X_0} = r} J(u) > 0.$$

2. *For some $u_0 \in X_0$ one has*

$$J(\tau u_0) \rightarrow -\infty,$$

as $\tau \rightarrow +\infty$.

PROOF. 1. We choose $\varepsilon > 0$ small enough, so that it satisfies

$$m_0 > \frac{\lambda + \varepsilon}{\lambda_{1,s}}.$$

By condition (1) there exists $\delta_\varepsilon > 0$ such that

$$\frac{f(x, t)}{t} \leq \lambda + \varepsilon,$$

for every $x \in \bar{\Omega}$ and $|t| \leq \delta_\varepsilon$.

Hence, one has

$$F(x, \xi) \leq \frac{\lambda + \varepsilon}{2} |\xi|^2,$$

for every $|\xi| \leq \delta_\varepsilon$.

As a consequence of the above inequality, using hypotheses h_1), the Sobolev embedding $X_0 \hookrightarrow L^q(\Omega)$ and (\hat{C}_M) , we can write:

$$\begin{aligned} J(u) &\geq \frac{m_0}{2} \|u\|_{X_0}^2 - \int_{|u(x)| \leq \delta_\varepsilon} \frac{\lambda + \varepsilon}{2} |u(x)|^2 dx - C \int_{|u(x)| > \delta_\varepsilon} |u(x)|^q dx \\ &\geq \frac{m_0}{2} \|u\|_{X_0}^2 - \frac{\lambda + \varepsilon}{2\lambda_{1,s}} \|u\|_{X_0}^2 - C \|u\|_{X_0}^q, \end{aligned}$$

for a suitable positive constant C .

Now, set $r := \|u\|_{X_0}^2$ and observe that for $r > 0$ small enough, we have

$$\frac{1}{2} \left(m_0 - \frac{\lambda + \varepsilon}{\lambda_{1,s}} \right) r - Cr^{q/2} > 0,$$

bearing in mind that $q > 2$. Hence

$$\inf_{\|u\|_{X_0}=r} J(u) > 0.$$

2. Let us choose an element $u_0 \in X_0$ such that

$$\text{meas}(\{x \in \Omega : u_0(x) \geq t_*\}) > 0.$$

$F(x, \xi)$ being a θ -superhomogeneous function if $|\xi| \geq t_*$, for $\tau > 1$, we have that

$$\begin{aligned} J(\tau u_0) &\leq \frac{m_1}{2} \|\tau u_0\|_{X_0}^2 + \frac{m_2}{2} - \int_{\Omega} F(x, \tau u_0(x)) dx \\ &\leq m_1 \frac{\|u_0\|_{X_0}^2}{2} \tau^2 - \tau^\theta \int_{|u_0(x)| \geq t_*} F(x, u_0(x)) dx + \frac{m_2}{2} + M \text{meas}(\Omega), \end{aligned}$$

where

$$M := \sup\{|F(x, \xi)| : x \in \bar{\Omega}, |\xi| \leq t_*\}.$$

Thus the (AR) condition implies that

$$J(\tau u_0) \rightarrow -\infty,$$

as $\tau \rightarrow +\infty$. This completes the proof.

4. AN EXAMPLE OF APPLICATION

In this section we present a simple application of our main result.

EXAMPLE 1. Consider the following nonlocal problem:

$$(D_M) \quad \begin{cases} M\left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy\right)(-\Delta)^s u = u^3 + u^4 & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

where

$$M\left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy\right) := 2 + \frac{\sin\left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy\right)}{1 + \left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy\right)^2}.$$

By virtue of Theorem 2, problem (D_M) admits one nontrivial weak solution. Indeed, a direct computation ensures that the continuous function

$$M(t) := 2 + \frac{\sin t}{1 + t^2}, \quad (\forall t \geq 0)$$

satisfies conditions (C_M^1) and (C_M^2) .

Further, the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(t) := t^3 + t^4, \quad (\forall t \in \mathbb{R})$$

satisfies all the hypotheses of Theorems 2.

REMARK 2. We note that in Example 1 condition (C_M^2) is satisfied for every $t \geq t_0$, where t_0 is the unique positive solution of the following real equation

$$\int_0^t \left(2 + \frac{\sin s}{1 + s^2}\right) ds - t\left(2 + \frac{\sin t}{1 + t^2}\right) = 0.$$

REMARK 3. We just observe that Theorem 2 can be proved for a more general class of nonlocal problems of the form

$$\begin{cases} -M\left(\int_{\mathbb{R}^n \times \mathbb{R}^n} |v(x) - v(y)|^2 K(x - y) dx dy\right) \mathcal{L}_K u = f(x, u) & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

where \mathcal{L}_K is defined as follows:

$$\mathcal{L}_K u(x) := \int_{\mathbb{R}^n} (u(x + y) + u(x - y) - 2u(x))K(y) dy, \quad (x \in \mathbb{R}^n)$$

and $K : \mathbb{R}^n \setminus \{0\} \rightarrow (0, +\infty)$ is a function with the properties that:

1. $\gamma K \in L^1(\mathbb{R}^n)$, where $\gamma(x) := \min\{|x|^2, 1\}$;
2. There exists $\beta > 0$ and $s \in (0, 1)$ such that

$$K(x) \geq \beta|x|^{-(n+2s)},$$

for any $x \in \mathbb{R}^n \setminus \{0\}$.

In this case we look for (weak) solutions $u \in X_0$, where

$$X_0 := \{g \in X : g = 0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega\}.$$

Here X denotes the linear space of Lebesgue measurable functions from \mathbb{R}^n to \mathbb{R} such that the restriction to Ω of any function g in X belongs to $L^2(\Omega)$ and

$$((x, y) \mapsto (g(x) - g(y))\sqrt{K(x-y)}) \in L^2((\mathbb{R}^n \times \mathbb{R}^n) \setminus (\mathcal{C}\Omega \times \mathcal{C}\Omega), dx dy).$$

Moreover, hypothesis (1) assumes the form

$$\limsup_{t \rightarrow 0} \frac{f(x, t)}{t} < m_0 \lambda_1,$$

where λ_1 is the first eigenvalue of the problem

$$\begin{cases} -\mathcal{L}_K u = \lambda u & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega. \end{cases}$$

Note that a model for K is given by the singular kernel $K(x) := |x|^{-(n+2s)}$ which gives rise to the fractional Laplace operator.

REMARK 4. The previous remark ensures that our result represents an improvement of [30, Theorems 1 and 2], provided that $M \equiv 1$.

ACKNOWLEDGEMENTS. This paper was written when the first author was visiting professor at the University of Ljubljana in 2014. He expresses his gratitude to the host institution for warm hospitality. The manuscript was realized within the auspices of the GNAMPA Project 2014 titled *Proprietà geometriche ed analitiche per problemi non-locali* and the SRA grants P1-0292-0101 and J1-5435-0101.

REFERENCES

- [1] C. O. ALVES - F. J. S. A. CORRÊA, *On existence of solutions for a class of problem involving a nonlinear operator*, Comm. Appl. Nonlinear Anal. 8 (2001), 43–56.
- [2] C. O. ALVES - F. J. S. A. CORRÊA - T. F. MA, *Positive solutions for a quasilinear elliptic equation of Kirchoff type*, Comput. Math. Appl. 49 (2005), 85–93.
- [3] A. AMBROSETTI - P. RABINOWITZ, *Dual variational methods in critical point theory and applications*, J. Funct. Anal. 14 (1973), 349–381.

- [4] D. ANDRADE - T. F. MA, *An operator equation suggested by a class of stationary problems*, Comm. Appl. Nonlinear Anal. 4 (1997), 65–71.
- [5] G. AUTUORI - P. PUCCI, *Kirchhoff systems with nonlinear source and boundary damping terms*, Commun. Pure Appl. Anal. 9 (2010), 1161–1188.
- [6] G. AUTUORI - P. PUCCI, *Kirchhoff systems with dynamic boundary conditions*, Nonlinear Anal. 73 (2010), 1952–1965.
- [7] G. AUTUORI - P. PUCCI, *Local asymptotic stability for polyharmonic Kirchhoff systems*, Appl. Anal. 90 (2011), 493–514.
- [8] G. AUTUORI - P. PUCCI, *Existence of entire solutions for a class of quasilinear elliptic equations*, NoDEA Nonlinear Differential Equations Appl. 20 (2013), 977–1009.
- [9] G. AUTUORI - P. PUCCI, *Elliptic problems involving the fractional Laplacian in \mathbb{R}^N* , J. Differential Equations 255 (2013), 2340–2362.
- [10] G. AUTUORI - F. COLASUONNO - P. PUCCI, *On the existence of stationary solutions for higher order p -Kirchhoff problems via variational methods*, Commun. Contemp. Math. 16 (2014), 1450002, pages 43.
- [11] H. BRÉZIS, *Analyse fonctionnelle. Théorie et applications*, Masson, Paris (1983).
- [12] L. CAFFARELLI - J. M. ROQUEJOFFRE - O. SAVIN, *Nonlocal minimal surfaces*, Comm. Pure Appl. Math. 63 (2010), 1111–1144.
- [13] M. CHIPOT - B. LOVAT, *Some remarks on non local elliptic and parabolic problems*, Nonlinear Anal. 30 (1997), 4619–4627.
- [14] M. CHIPOT - J. F. RODRIGUES, *On a class of nonlocal nonlinear elliptic problems*, RAIRO Modélisation Math. Anal. Numér. 26 (1992), 447–467.
- [15] E. DI NEZZA - G. PALATUCCI - E. VALDINOCI, *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. Math. 136 (2012), no. 5, 521–573.
- [16] M. FERRARA - G. MOLICA BISCI - B. ZHANG, *Existence of weak solutions for non-local fractional problems via Morse theory*, Discrete and Continuous Dynamical Systems Series B, 19 (2014), 2493–2499.
- [17] A. FISCELLA - E. VALDINOCI, *A critical Kirchhoff type problem involving a non-local operator*, Nonlinear Anal. 94 (2014), 156–170.
- [18] A. KRISTÁLY - V. RĂDULESCU - Cs. VARGA, *Variational Principles in Mathematical Physics, Geometry, and Economics: Qualitative Analysis of Nonlinear Equations and Unilateral Problems*, Encyclopedia of Mathematics and its Applications, No. 136, Cambridge University Press, Cambridge, 2010.
- [19] G. MOLICA BISCI, *Fractional equations with bounded primitive*, Appl. Math. Lett. 27 (2014), 53–58.
- [20] G. MOLICA BISCI, *Sequences of weak solutions for fractional equations*, Math. Res. Lett. 21 (2014), 1–13.
- [21] G. MOLICA BISCI - B. A. PANSERA, *Three weak solutions for nonlocal fractional equations*, Adv. Nonlinear Stud. 14 (2014), 591–601.
- [22] G. MOLICA BISCI - D. REPOVŠ, *Fractional nonlocal problems involving nonlinearities with bounded primitive*, J. Math. Anal. Appl. 420 (2014), 167–176.
- [23] G. MOLICA BISCI - R. SERVADEI, *A bifurcation result for nonlocal fractional equations*, to appear in Anal. Appl.; DOI:10.1142/S0219530514500067.
- [24] P. PUCCI - Q. ZHANG, *Existence of entire solutions for a class of variable exponent elliptic equations*, J. Differential Equations 257 (2014), 1529–1566.
- [25] P. PUCCI - V. RĂDULESCU, *The impact of the mountain pass theory in nonlinear analysis: a mathematical survey*, Boll. Unione Mat. Ital. Series IX, No. 3, (2010) 543–584.

- [26] P. PUCCI - S. SALDI, *Critical stationary Kirchhoff equations in \mathbb{R}^n involving nonlocal operators*, to appear in Rev. Mat. Iberoam., pages 23.
- [27] P. H. RABINOWITZ, *Minimax methods in critical point theory with applications to differential equations*, CBMS Reg. Conf. Ser. Math., 65, American Mathematical Society, Providence, RI (1986).
- [28] R. SERVADEI, *The Yamabe equation in a non-local setting*, to appear in Adv. Nonlinear Anal. 3 (2013).
- [29] R. SERVADEI - E. VALDINOCI, *Lewy-Stampacchia type estimates for variational inequalities driven by nonlocal operators*, to appear in Rev. Mat. Iberoam. 29 (2013).
- [30] R. SERVADEI - E. VALDINOCI, *Mountain Pass solutions for non-local elliptic operators*, J. Math. Anal. Appl. 389 (2012), 887–898.
- [31] R. SERVADEI - E. VALDINOCI, *Variational methods for non-local operators of elliptic type*, Discrete Contin. Dyn. Syst. 33, 5 (2013), 2105–2137.
- [32] R. SERVADEI - E. VALDINOCI, *The Brézis-Nirenberg result for the fractional Laplacian*, to appear in Trans. Amer. Math. Soc.
- [33] M. STRUWE, *Variational methods, Applications to nonlinear partial differential equations and Hamiltonian systems*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3, Springer Verlag, Berlin–Heidelberg (1990).
- [34] C. F. VASCONCELOS, *On a nonlinear stationary problem in unbounded domains*, Rev. Mat. Univ. Complut. Madrid 5 (1992), 309–318.

Received 18 November 2014,
and in revised form 22 January 2015.

Giovanni Molica Bisci
Dipartimento P.A.U.
Università degli Studi Mediterranea di Reggio Calabria
Salita Melissari, 89100 Reggio Calabria, Italy
gmolica@unirc.it

Dušan Repovš
Faculty of Education and Faculty of Mathematics and Physics
University of Ljubljana, POB 2964, Slovenia 1001
dusan.repovs@guest.arnes.si