



**Partial Differential Equations.** – *Some of my favorite open problems*, by HAÏM BREZIS, communicated on 10 November 2022.

*To the memory of Antonio Ambrosetti, a dearly missed friend,  
and master of Nonlinear Analysis.*

ABSTRACT. – This is a selection of the main open problems I raised throughout my career and that have resisted so far. This is not an exhaustive list, but striking questions fairly easy to state; some were raised 40 years ago, others quite recently.

KEYWORDS. – Critical exponent, radial symmetry, harmonic maps, degree theory, extremal solutions, Bourgain–Brezis estimates, relative isoperimetric inequality.

2020 MATHEMATICS SUBJECT CLASSIFICATION. – Primary 35B06; Secondary 35A16, 35B33.

### 1. AN ELLIPTIC EQUATION INVOLVING THE CRITICAL EXPONENT IN 3D

Let  $\Omega$  be the unit ball in  $\mathbb{R}^3$ . Consider the equation

$$(1.1) \quad \begin{cases} -\Delta u = u^5 + \lambda u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where the unknown  $u : \Omega \rightarrow \mathbb{R}$  is a smooth function and  $\lambda \in \mathbb{R}$  is a parameter.

A natural question is whether (1.1) admits a non-trivial solution,  $u \neq 0$ . Note that the exponent 5 corresponds to the critical Sobolev exponent  $(N + 2)/(N - 2)$  when  $N = 3$ , which produces notorious difficulties. The answer, which depends on  $\lambda$ , is known for a large class of  $\lambda$ 's; however, for one interval of  $\lambda$ 's the answer has remained undecided over the past forty years. Let  $\lambda_1 = \pi^2$  be the first eigenvalue of  $-\Delta$  on  $\Omega$  with zero Dirichlet condition.

OPEN PROBLEM 1.1 (Implicit in [30]). Assume that

$$(1.2) \quad 0 < \lambda \leq \lambda_1/4.$$

Does there exist a solution  $u \neq 0$  of (1.1)?

Several comments are in order:

- (a) The answer is not known even if (1.2) is replaced by a sub-interval, e.g.  $0 < \lambda < \delta$  with  $\delta$  small.
- (b) When  $\lambda \leq 0$ , the only solution of (1.1) is  $u \equiv 0$ ; this is a celebrated result of Pohozaev (1965).
- (c) When

$$\lambda_1/4 < \lambda < \lambda_1,$$

there exists a radial solution  $u > 0$  in  $\Omega$  of (1.1). This is a central result from Brezis–Nirenberg [30, Theorem 1.2] (see also [11, 38]).

- (d) When

$$0 < \lambda \leq \lambda_1/4,$$

any radial solution  $u$  of (1.1) must be  $u \equiv 0$  (see [30, proof of Lemma 1.4]). In particular (via Gidas–Ni–Nirenberg) there exists no solution  $u > 0$  in  $\Omega$  of (1.1). Therefore, if (1.4) holds and a solution  $u \not\equiv 0$  of (1.1) exists, it must be *non-radial and sign-changing*.

- (e) When

$$\lambda \geq \lambda_1,$$

there exist sign-changing solutions of (1.1) – but no solution  $u > 0$  of (1.1), (see [37]). In the bifurcation diagram, branches of solutions emanate from the eigenvalues associated with non-radial sign-changing eigenfunctions. It would be interesting to decide whether such branches “reach” the interval  $(0, \lambda_1/4)$ ; they might instead admit, e.g., vertical asymptotes at values of  $\lambda \geq \lambda_1/4$ .

## 2. QUESTIONS OF UNIQUENESS AND RADIAL SYMMETRY ARISING FROM THE GINZBURG–LANDAU (GL) SYSTEM

Let  $\Omega$  be the unit disc in  $\mathbb{R}^2$ . Consider the system

$$(2.1) \quad \begin{cases} -\Delta u = \frac{1}{\varepsilon^2} u(1 - |u|^2) & \text{in } \Omega, \\ u(x) = x & \text{on } \partial\Omega, \end{cases}$$

where  $\varepsilon > 0$  is a given parameter and the unknown  $u$  maps  $\Omega$  into  $\mathbb{R}^2$ .

It is easy to check that (2.1) admits a solution  $u$  of the form

$$(2.2) \quad u(x) = \frac{x}{|x|} f_\varepsilon(|x|),$$

where  $f_\varepsilon : [0, 1] \rightarrow \mathbb{R}$  satisfies the ordinary differential equation (ODE)

$$(2.3) \quad \begin{cases} -f'' - \frac{1}{r}f' + \frac{1}{r^2}f = \frac{1}{\varepsilon^2}f(1-f^2) & \text{in } (0, 1), \\ f(0) = 0 \quad \text{and} \quad f(1) = 1. \end{cases}$$

In fact, (2.3) admits a unique solution (see [3, Appendix II]). We will denote by  $U_\varepsilon(x)$  the solution of (2.1) given by (2.2)–(2.3), and we call it the radially symmetric (or just the radial) solution of (2.1). A long-standing open problem is whether  $U_\varepsilon(x)$  is also the unique solution of (2.1).

**OPEN PROBLEM 2.1** ([3, Problem 10 in Chapter XI]). Is the radial solution  $U_\varepsilon$  the only solution of (2.1)?

A positive answer would, in particular, imply that solutions of some specific nonlinear *systems* of PDEs inherit the radial symmetry of the data – a property reminiscent of the celebrated Gidas–Ni–Nirenberg result relative to *positive* solutions of some *scalar* PDEs.

Note that (2.1) has a variational structure: the solutions of (2.1) are the critical points of the GL energy,

$$E_\varepsilon(u) := \frac{1}{2} \int_\Omega |\nabla u|^2 + \frac{1}{4\varepsilon^2} \int_\Omega (|u|^2 - 1)^2,$$

subject to the boundary condition  $u \in H_g^1(\Omega; \mathbb{R}^2)$ , where  $g(x) = x$  on  $\partial\Omega$ .

The answer to Open Problem 2.1 is known to be positive in two “opposite” cases.

(a) When  $\varepsilon$  is sufficiently *large*; more precisely  $\varepsilon \geq 1/\sqrt{\lambda_1}$ , where  $\lambda_1$  is the first eigenvalue of  $-\Delta$  under zero Dirichlet condition. Indeed, it is easy to check that  $E_\varepsilon$  is strictly convex when  $\varepsilon \geq 1/\sqrt{\lambda_1}$  and its unique minimizer is also its unique critical point.

(b) When  $\varepsilon$  is sufficiently *small*:  $\varepsilon < \varepsilon_0$  for some appropriate  $\varepsilon_0$ . This result is due to Pacard–Rivière [56]. Their proof is highly non-trivial and fills a significant part of the monograph [56]; it would be interesting to find a simpler proof.

The intermediate range  $\varepsilon_0 \leq \varepsilon < 1/\sqrt{\lambda_1}$  is totally open. An easier question still unresolved is the following.

**OPEN PROBLEM 2.2.** Is  $U_\varepsilon$  a minimizer of  $E_\varepsilon$  on  $H_g^1(\Omega; \mathbb{R}^2)$  for any  $\varepsilon > 0$ ?

Note that a positive answer to Open Problem 2.1 implies a positive answer to Open Problem 2.2, since any minimizer of  $E_\varepsilon$  on  $H_g^1(\Omega; \mathbb{R}^2)$  is a solution of (2.1), and by uniqueness it would coincide with  $U_\varepsilon$ .

The next result provides substantial evidence that the answer to Open Problem 2.2 is positive.

**THEOREM 2.1** (Mironescu [50], see also Lieb–Loss [49]). *For every  $\varepsilon > 0$ ,  $U_\varepsilon$  is a local minimizer of  $E_\varepsilon$ ; moreover  $D^2 E_\varepsilon(U_\varepsilon)$  is positive definite.*

Following Brezis [13, Open Problem 6], one may ask similar questions when  $\Omega$  is the unit ball in  $\mathbb{R}^N$ ,  $N \geq 3$ , and  $u : \Omega \rightarrow \mathbb{R}^2$ ; the counterpart of (2.3) is

$$\begin{cases} -f'' - \frac{(N-1)}{r} f' + \frac{N-1}{r^2} f = \frac{1}{\varepsilon^2} f(1 - f^2) & \text{in } (0, 1), \\ f(0) = 0 \quad \text{and} \quad f(1) = 1. \end{cases}$$

Ignat–Nguyen [42] established the analog of Theorem 2.1 in any dimension  $N \geq 3$ , while Ignat–Nguyen–Slastikov–Zarnescu [43] proved that in dimension  $N \geq 7$ ,  $U_\varepsilon$  is a *global* minimizer of  $E_\varepsilon$  on  $H_g^1(\Omega; \mathbb{R}^N)$  for any  $\varepsilon > 0$ ; in fact  $U_\varepsilon$  is the unique global minimizer of  $E_\varepsilon$ .

One can also raise identical questions for the  $p$ -GL energy

$$E_{\varepsilon,p}(u) := \frac{1}{p} \int_{\Omega} |\nabla u|^p + \frac{1}{4\varepsilon^2} \int_{\Omega} (|u|^2 - 1)^2,$$

where  $p > 1$ ,  $\Omega$  is the unit disc in  $\mathbb{R}^2$ , and  $u \in W_g^{1,p}(\Omega; \mathbb{R}^2)$ , with  $g(x) = x$  on  $\partial\Omega$ .

A somewhat related direction concerns the “asymptotic radial symmetry” as  $\varepsilon \rightarrow 0$ ; it is more specifically illustrated as follows.

**OPEN PROBLEM 2.3** ([26, Open Problem 30]). Assume  $p > 2$  and let  $u_\varepsilon$  be any minimizer (resp., critical point) of  $E_{\varepsilon,p}$  on  $W_g^{1,p}(\Omega, \mathbb{R}^2)$ . Does  $u_\varepsilon(x) \rightarrow x/|x|$  in  $\Omega \setminus \{0\}$  as  $\varepsilon \rightarrow 0$ ?

Note that the answer to the same problem when  $p = 2$  is positive (see [3] for minimizers and [56] for critical points). When  $p < 2$ , the answer is positive for minimizers (this is an immediate consequence of [26, Theorem 13.6]) and is open for general critical points.

Entire solutions of the GL equation are also of interest. Consider the system

$$(2.4) \quad -\Delta u = u(1 - |u|^2) \quad \text{on } \mathbb{R}^2,$$

and the condition at infinity

$$(2.5) \quad \lim_{|x| \rightarrow \infty} |u(x)| = 1,$$

where  $u$  is a smooth function from  $\mathbb{R}^2$  into  $\mathbb{R}^2 \simeq \mathbb{C}$ . Here the parameter  $\varepsilon$  is irrelevant since it can be scaled out.

Property (2.5) allows to define the degree of  $u$  at infinity

$$\deg(u, \infty) := \deg \left( \frac{u(Rx)}{|u(Rx)|}; x \in \mathbb{S}^1 \right) \quad \text{for } R \text{ sufficiently large.}$$

Given any  $q \in \mathbb{Z}$ ,  $q \neq 0$ , there exists a distinguished solution  $u$  of (2.4)-(2.5) given in polar coordinates by

$$u(r, \theta) = e^{iq\theta} g_q(r),$$

where  $g_q : [0, \infty) \rightarrow [0, 1)$  satisfies the ODE

$$(2.6) \quad \begin{cases} -g'' - \frac{1}{r}g' + \frac{q^2}{r^2}g = g(1 - g^2) & \text{on } (0, \infty), \\ g(0) = 0 & \text{and } \lim_{r \rightarrow \infty} g(r) = 1. \end{cases}$$

In fact, (2.6) admits a unique solution denoted by  $g_q$  (see [3, Appendix III]). When  $q \in \mathbb{Z}$ ,  $q \neq 0$ , we set  $V_q(r, \theta) := e^{iq\theta} g_q(r)$ . We also set  $V_0 \equiv 1$ . Note that

$$\deg(V_q, \infty) = q \quad \forall q \in \mathbb{Z}.$$

A long-standing open problem is whether the functions  $V_q$ ,  $q \in \mathbb{Z}$ , are the *only* solutions of (2.4)-(2.5); more precisely:

**OPEN PROBLEM 2.4** ([3, Problem 14], [13, 25]). Let  $u$  be any solution of (2.4)-(2.5). Does  $u$  coincide with  $V_q$  modulo rotation and translation, where  $q = \deg(u, \infty)$ ? I.e., is  $u(x) = \alpha V_q(x - x_0)$  for some  $\alpha \in \mathbb{C}$ ,  $|\alpha| = 1$ , and  $x_0 \in \mathbb{R}^2$ ?

Two partial results are known so far:

**THEOREM 2.2** ([25]). Assume that  $u$  is a solution of (2.4)-(2.5) such that

$$\deg(u, \infty) = 0,$$

and which satisfies in addition

$$(2.7) \quad \int_{\mathbb{R}^2} (|u|^2 - 1)^2 < \infty.$$

Then  $u = V_0$  modulo rotation and translation; i.e.,  $u$  is a constant of modulus 1.

**THEOREM 2.3** ([51]). Assume that  $u$  is a solution of (2.4)-(2.5) such that

$$\deg(u, \infty) = \pm 1,$$

and which satisfies in addition (2.7).

Then  $u = V_{\pm 1}$  modulo rotation and translation.

Addressing Open Problem 2.4 when  $|\deg(u, \infty)| \geq 2$ , Ovchinnikov and Sigal [55] have devised a strategy to construct non-radial solutions of (2.4)-(2.5), thereby providing a negative answer to Open Problem 2.4. However, their proposed construction has been criticized (see Esposito [39] and Kurzke [47]), and the problem remains open.

In another direction, we point out that it is not known whether the conclusions of Theorem 2.2 and Theorem 2.3 remain true if one removes assumption (2.7). That is more generally the following.

OPEN PROBLEM 2.5 ([25, Problem 2], [13, Open Problem 2]). Assume that  $u$  satisfies (2.4) and (2.5). Does (2.7) hold?

Finally, we mention that property (2.7) appears quite naturally in connection with solutions of (2.4). In particular, the functions  $V_q, q \in \mathbb{Z}$ , satisfy (2.7); more precisely (see, e.g., [60]), for any  $q \in \mathbb{Z}$ ,

$$|V_q(x)| = 1 - \frac{q^2}{2|x|^2} + o\left(\frac{1}{|x|^2}\right) \quad \text{as } |x| \rightarrow \infty.$$

Also, one can show (see [25, 60]) that any solution of (2.4) satisfying (2.7) enjoys the following properties:

- (a)  $\lim_{|x| \rightarrow \infty} |u(x)| = 1$ , so that  $q = \deg(u, \infty)$  is well defined,
- (b)  $|u(x)| = 1 - \frac{q^2}{2|x|^2} + o\left(\frac{1}{|x|^2}\right)$  as  $|x| \rightarrow \infty$ ,
- (c)  $\int_{\mathbb{R}^2} (|u|^2 - 1)^2 = 2\pi q^2$ ,
- (d)  $\lim_{|x| \rightarrow \infty} |u(x) - \alpha V_q(x)| = 0$  for some  $\alpha \in \mathbb{C}, |\alpha| = 1$ .

### 3. HARMONIC MAPS FROM THE DISC TO $\mathbb{S}^2$

Let  $\Omega$  be the unit disc in  $\mathbb{R}^2$ . A harmonic map to  $\mathbb{S}^2$  is a smooth map  $u : \Omega \rightarrow \mathbb{R}^3$  satisfying

$$(3.1) \quad -\Delta u_i = u_i |\nabla u|^2, \quad i = 1, 2, 3, \text{ in } \Omega,$$

$$(3.2) \quad |u(x)| = 1 \quad \text{in } \Omega.$$

Given a smooth map  $g : \partial\Omega \rightarrow \mathbb{S}^2$ , we add the boundary condition

$$(3.3) \quad u = g \quad \text{on } \partial\Omega.$$

One could also define a concept of weak harmonic map, i.e., a map  $u \in H_g^1(\Omega; \mathbb{S}^2)$  satisfying (3.1) in the sense of distributions. A celebrated result of F. Hélein (1990) asserts that weak harmonic maps are smooth.

The solutions of (3.1)–(3.3) correspond to critical points of the energy

$$E(u) = \int_{\Omega} |\nabla u|^2 \quad \text{on } \mathcal{E} = H_g^1(\Omega; \mathbb{S}^2).$$

It is easy to produce a solution of (3.1)–(3.3) by minimizing  $E$  on  $\mathcal{E}$ . Denote by  $\underline{u}$  such a minimizer (it need not be unique). A natural question is whether there are other solutions. In fact, one might expect that problem (3.1)–(3.3) admits infinitely many solutions. The reason is that  $\mathcal{E}$  has infinitely many connected components classified by a topological degree. More precisely, given  $u \in \mathcal{E}$ , let  $v : \mathbb{S}^2 \rightarrow \mathbb{S}^2$  be defined by

$$v(x, y, z) := \begin{cases} u(x, y) & \text{if } (x, y, z) \in \mathbb{S}^2 \text{ and } z \geq 0, \\ \underline{u}(x, y) & \text{if } (x, y, z) \in \mathbb{S}^2 \text{ and } z \leq 0. \end{cases}$$

Note that  $v \in H^1(\mathbb{S}^2; \mathbb{S}^2)$  since  $u = \underline{u}$  on  $\partial\Omega$ . Hence  $\deg v$  is well defined (see [31]) and in fact

$$\deg v = \frac{1}{4\pi} \int_{\Omega} u \cdot u_x \wedge u_y - \frac{1}{4\pi} \int_{\Omega} \underline{u} \cdot \underline{u}_x \wedge \underline{u}_y,$$

by Kronecker's formula. One may thus decompose  $\mathcal{E}$  into its connected components

$$\mathcal{E} = \bigcup_{k \in \mathbb{Z}} \mathcal{E}_k, \quad \text{where } \mathcal{E}_k = \{u \in \mathcal{E}; \deg v = k\}.$$

It is tempting to minimize  $E$  in each class  $\mathcal{E}_k$ . However,  $\text{Inf}_{\mathcal{E}_k} E$  need not be achieved (except when  $k = 0$  since  $\underline{u} \in \mathcal{E}_0$ ), the reason being that the degree is *not* continuous under weak convergence in  $H^1$ . Thus, if  $(u_n)$  is a minimizing sequence in  $\mathcal{E}_k$  and  $u_n \rightharpoonup u_\infty$  weakly in  $H^1$ , the limit  $u_\infty$  might “jump” to another class  $\mathcal{E}_\ell$ ,  $\ell \neq k$ , and will not be a minimizer of  $E$  in  $\mathcal{E}_k$ . This scenario can really occur. For example, if  $g \equiv C$  is a constant, a result of Lemaire [48] asserts that  $u \equiv C$  is the only solution of (3.1)–(3.3). As a consequence,  $\text{Inf}_{\mathcal{E}_k} E$  is achieved only when  $k = 0$ .

Here is a general result in the positive direction.

**THEOREM 3.1** ([22, 44]). *Assume  $g \not\equiv C$ , then  $\text{Inf}_{\mathcal{E}_k} E$  is achieved at least in one of the classes  $\mathcal{E}_{+1}$  or  $\mathcal{E}_{-1}$ . As a consequence, problem (3.1)–(3.3) admits at least two solutions (including  $\underline{u}$ ).*

Little is known concerning the existence of additional solutions, even when  $g$  has a simple form (see, however, works by Jie Qing, A. Soyeur, Morgan Pierre, L. Oswald, and G. Paulik). Consider the boundary condition

$$(3.4) \quad g(x, y) = (Rx, Ry, \sqrt{1 - R^2}) \quad \text{for } (x, y) \in \partial\Omega,$$

with  $0 < R < 1$ . In this case, one can write down two explicit solutions of (3.1)–(3.3):

$$\begin{aligned}\underline{u}(x, y) &= \frac{2\lambda}{\lambda^2 + r^2}(x, y, \lambda) + (0, 0, -1) \quad \text{for } (x, y) \in \Omega, \\ \bar{u}(x, y) &= \frac{2\mu}{\mu^2 + r^2}(x, y, -\mu) + (0, 0, 1) \quad \text{for } (x, y) \in \Omega,\end{aligned}$$

where  $r^2 = x^2 + y^2$ ,  $\lambda = \frac{1}{R} + \sqrt{\frac{1}{R^2} - 1}$ , and  $\mu = \frac{1}{R} - \sqrt{\frac{1}{R^2} - 1}$ .

It is not difficult to check that  $\underline{u}$  is a minimizer of  $E$  in  $\mathcal{E}$  and that  $\bar{u}$  is a minimizer of  $E$  in  $\mathcal{E}_{-1}$ .

More precisely,  $\underline{u}$  is the unique minimizer of  $E$  in  $\mathcal{E}_0$  and  $\bar{u}$  is the unique minimizer of  $E$  in  $\mathcal{E}_{-1}$ . Moreover,  $\text{Inf}_{\mathcal{E}_k} E$  is not achieved when  $k \neq 0$  and  $k \neq -1$  (see [22, Theorem 2]). This does not exclude the possible existence of other solutions of (3.1)–(3.3) as follows.

**OPEN PROBLEM 3.1** ([22]). Assume that  $g$  is given by (3.4). Are there other solutions of (3.1)–(3.3) besides  $\underline{u}$  and  $\bar{u}$ ?

Either way, the answer to Open Problem 3.1 would be illuminating. A negative answer might possibly shed some light on the important question whether solutions of specific nonlinear systems inherit the symmetry of the data – assuming the first step in the proof establishes that any solution is radially symmetric. A positive answer (more than two solutions) might involve the development of new techniques for finding non-minimizing critical points in variational problems with lack of compactness.

#### 4. CONTINUOUS HARMONIC MAPS FROM $B^3$ TO $\mathbb{S}^2$

Let  $\Omega = B^3$  be the unit ball in  $\mathbb{R}^3$ . A (weak) harmonic map to  $\mathbb{S}^2$  is a map  $u \in H^1(\Omega; \mathbb{R}^3)$  satisfying

$$(4.1) \quad -\Delta u_i = u_i |\nabla u|^2 \quad i = 1, 2, 3, \text{ in } \Omega,$$

$$(4.2) \quad |u(x)| = 1 \quad \text{in } \Omega.$$

Given a smooth map  $g : \partial\Omega \simeq \mathbb{S}^2 \rightarrow \mathbb{S}^2$ , we add the boundary condition

$$(4.3) \quad u = g \quad \text{on } \partial\Omega.$$

Solutions of (4.1)–(4.3) correspond to critical points of the energy

$$E(u) = \int_{\Omega} |\nabla u|^2 \quad \text{on } H_g^1(\Omega; \mathbb{S}^2).$$

Note that  $H_g^1(\Omega; \mathbb{S}^2)$  is always non-empty since  $u(x) = g(x/|x|) \in H_g^1(\Omega; \mathbb{S}^2)$ .



It is, therefore, easy to produce solutions of (4.1)–(4.3), e.g. by considering minimizers of the problem

$$(4.4) \quad \text{Min} \{E(u); u \in H_g^1(\Omega; \mathbb{S}^2)\}.$$

In contrast with the 2D case (see Section 3), weak harmonic maps need not be smooth – and not even continuous. The optimal regularity result for minimizers is known from the works of Schoen–Uhlenbeck [59] and Brezis–Coron–Lieb [23]: any minimizer  $u$  of (4.4) is smooth in  $\bar{\Omega}$  except at a finite number of points  $(a_i)$  in  $\Omega$ , and near each  $a_i$ ,  $u$  behaves like  $\pm(x - a_i)/|x - a_i|$  modulo a rotation.

When  $\deg g \neq 0$ ,  $\{u \in C(\bar{\Omega}; \mathbb{S}^2); u = g \text{ on } \partial\Omega\} = \emptyset$ , and thus singularities are unavoidable. Since we will be concerned with the existence of continuous harmonic maps satisfying (4.3), we assume throughout this section that

$$(4.5) \quad \deg g = 0.$$

Here is a long-standing open problem originally posed by R. Schoen in the mid-1980's.

**OPEN PROBLEM 4.1** ([40, 41], [15, Open Problem 3]). Assume that (4.5) holds. Does there exist a continuous harmonic map satisfying (4.1)–(4.3)?

Even in the absence of a topological obstruction (i.e., when (4.5) holds), minimizers in (4.4) can still have singularities, and therefore will not provide a solution to Open Problem 4.1. This is a consequence of a remarkable gap phenomenon discovered by Hardt–Lin [40] (see also [12]): there exist smooth maps  $g : \partial\Omega \rightarrow \mathbb{S}^2$  satisfying (4.5) and such that

$$(4.6) \quad \text{Min}\{E(u); u \in H_g^1(\Omega; \mathbb{S}^2)\} < \text{Inf}\{E(u); u \in H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega})\}.$$

In order to solve Open Problem 4.1, it is tempting to tackle the following problem.

**OPEN PROBLEM 4.2** ([40, 41]). Is the

$$(4.7) \quad \text{Inf}\{E(u); u \in H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega})\}$$

achieved?

Clearly, a positive answer to Open Problem 4.2 would provide a solution to Open Problem 4.1. But, in principle, it might happen that the answer to Open Problem 4.2 is negative while the answer to Problem 4.1 is positive. (Can this scenario occur?)

A natural strategy to solve Open Problem 4.2 has been developed by Bethuel–Brezis–Coron [2] via the concept of *relaxed energy* defined as follows.

Fix  $u \in H_g^1(\Omega; \mathbb{S}^2)$ . In general, there exist no sequence  $(u_n)$  in  $H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega})$  such that  $u_n \rightarrow u$  strongly in  $H^1$ . (This is, e.g., a consequence of the gap phenomenon (4.6).) However, there always exists a sequence  $(u_n)$  in  $H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega})$  such that  $u_n \rightharpoonup u$  weakly in  $H^1$  (see [1]). Set

$$(4.8) \quad R(u) := \text{Inf} \left\{ \liminf_{n \rightarrow \infty} E(u_n); u_n \in H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega}) \text{ and } u_n \rightharpoonup u \text{ weakly in } H^1 \right\},$$

where the first Inf is taken over all sequences  $(u_n)$  as above. ( $R$  stands for relaxed).

The functional  $R$  is well defined on  $H_g^1(\Omega; \mathbb{S}^2)$  and it is weakly lower semi-continuous. Therefore,

$$(4.9) \quad \text{Min} \{ R(u); u \in H_g^1(\Omega; \mathbb{S}^2) \} \text{ is achieved.}$$

We claim that if the Inf in (4.7) is achieved, say by some  $\underline{u} \in H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega})$ , then  $\underline{u}$  is also a minimizer in (4.9).

Indeed, we clearly have

$$\begin{aligned} E(v) &\leq R(v) \quad \forall v \in H_g^1(\Omega; \mathbb{S}^2), \\ R(v) &\leq E(v) \quad \forall v \in H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega}) \end{aligned}$$

(just take  $u_n \equiv v$  in (4.8)).

Therefore,

$$(4.10) \quad R(v) = E(v) \quad \forall v \in H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega}).$$

We also have

$$(4.11) \quad \text{Inf}_{v \in H_g^1} R(v) = \text{Inf}_{v \in H_g^1 \cap C} E(v).$$

Indeed, it is clear that

$$(4.12) \quad \text{Inf}_{v \in H_g^1} R(v) \leq \text{Inf}_{v \in H_g^1 \cap C} R(v) = \text{Inf}_{v \in H_g^1 \cap C} E(v) \quad \text{by (4.10).}$$

On the other hand, given any  $w \in H_g^1(\Omega; \mathbb{S}^2)$ , there exists (by definition of  $R$ ) a sequence  $(w_n)$  in  $H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega})$  such that  $w_n \rightharpoonup w$  weakly in  $H^1$  and

$$(4.13) \quad E(w_n) \rightarrow R(w).$$

Thus

$$(4.14) \quad \text{Inf}_{v \in H_g^1 \cap C} E(v) \leq E(w_n) \quad \forall n.$$

Passing to the limit in (4.14) using (4.13) gives

$$\text{Inf}_{v \in H_g^1 \cap C} E(v) \leq R(w) \quad \forall w \in H_g^1(\Omega; \mathbb{S}^2),$$

so that

$$(4.15) \quad \inf_{v \in H_g^1 \cap C} E(v) \leq \inf_{v \in H_g^1} R(v).$$

Combining (4.12) and (4.15) yields (4.11).

We may now return to the above claim concerning (4.7). Assume that  $\inf_{v \in H_g^1 \cap C} E(v)$  is achieved, say by some  $\underline{u} \in H_g^1 \cap C$ , then by (4.10) and (4.11)

$$R(\underline{u}) = E(\underline{u}) = \inf_{v \in H_g^1 \cap C} E(v) = \inf_{v \in H_g^1} R(v)$$

and hence  $\underline{u}$  is a minimizer for the problem (4.9).

Therefore, prospective solutions of Open Problem 4.2 are to be found among the minimizers of (4.9). This leads us to the following problem.

**OPEN PROBLEM 4.3 ([2]).** Are the minimizers of (4.9) continuous on  $\bar{\Omega}$ ? (Is it true for at least one of the minimizers?)

If the answer to Open Problem 4.3 is positive and  $\underline{u} \in H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega})$  is such a minimizer, then  $\underline{u}$  satisfies, by (4.10),

$$E(\underline{u}) = R(\underline{u}) = \inf_{v \in H_g^1} R(v) = \inf_{v \in H_g^1 \cap C} E(v),$$

and, therefore, we have solved Open Problem 4.2 since  $\underline{u}$  is a minimizer for Open Problem 4.2.

In tackling Open Problem 4.3, we have at our disposal explicit representation formulas for  $R$ . We first need some notations. Given  $u \in H^1(\Omega; \mathbb{S}^2)$ , consider the  $D$ -field (introduced in [23])

$$D(u) = \left( \det \left( u, \frac{\partial u}{\partial x_2}, \frac{\partial u}{\partial x_3} \right), \det \left( \frac{\partial u}{\partial x_1}, u, \frac{\partial u}{\partial x_3} \right), \det \left( \frac{\partial u}{\partial x_1}, \frac{\partial u}{\partial x_2}, u \right) \right),$$

so that  $D(u) \in L^1(\Omega; \mathbb{R}^3)$ , and one can define the distribution

$$Ju := \frac{1}{3} \operatorname{div} D(u) \in \mathcal{D}'(\Omega; \mathbb{R}).$$

$J$  stands for Jacobian because  $Ju$  coincides with the usual Jacobian when  $u \in C^2(\Omega; \mathbb{R}^3)$ . Since  $u$  takes its values in  $\mathbb{S}^2$ , it follows that  $Ju = 0$  in the region where  $u$  is smooth. As we are going to see below (in (4.18)),  $Ju$  carries important information about the location of the topological singularities of  $u$ .

**THEOREM 4.1 ([2]).** For every  $u \in H_g^1(\Omega; \mathbb{S}^2)$

$$(4.16) \quad R(u) = \int_{\Omega} |\nabla u|^2 + S(u),$$

where

$$(4.17) \quad S(u) := 2 \operatorname{Sup} \left\{ \int_{\Omega} D(u) \cdot \nabla \zeta - \int_{\partial\Omega} (\operatorname{Jac} g) \zeta; \zeta \in W^{1,\infty}(\Omega; \mathbb{R}), \|\nabla \zeta\|_{L^\infty} \leq 1 \right\},$$

and  $\operatorname{Jac} g$  denotes the Jacobian determinant of  $g : \partial\Omega \simeq \mathbb{S}^2 \rightarrow \mathbb{S}^2$ .

As a consequence of (4.16) and (4.17), we see that  $R$  is continuous for the strong topology of  $H^1$  (it is even locally Lipschitz). Thus, it is helpful to know the value of  $R$  on a dense subset of  $H_g^1(\Omega; \mathbb{S}^2)$  for the strong topology, in particular on the class

$$\mathcal{C} = \{u \in H_g^1(\Omega; \mathbb{S}^2); u \text{ is continuous on } \bar{\Omega} \text{ except on a finite set } (a_i), 1 \leq i \leq k \text{ in } \Omega\}$$

which is dense in  $H_g^1(\Omega; \mathbb{S}^2)$  (by a classical result of Bethuel).

When  $u \in \mathcal{C}$ , we have (see [23]) the important formula

$$Ju = \frac{4\pi}{3} \sum_{i=1}^k \operatorname{deg}(u, a_i) \delta_{a_i},$$

where  $\operatorname{deg}(u, a_i)$  is the degree of  $u$  restricted to a small ball centered at  $a_i$ . Relabeling the points  $(a_i)$  as  $P_i$  and  $N_i$ ,  $1 \leq i \leq \ell$ , including multiplicities, we may write

$$(4.18) \quad Ju = \frac{4\pi}{3} \sum_{i=1}^{\ell} (\delta_{P_i} - \delta_{N_i}).$$

(Here we use the fact that  $\sum_i \operatorname{deg}(u, a_i) = \operatorname{deg} g = 0$ .) Inserting (4.18) in (4.17) yields (after integration by parts)

$$S(u) = 8\pi \operatorname{Sup} \left\{ \sum_{i=1}^{\ell} [\zeta(P_i) - \zeta(N_i)]; \zeta \in W^{1,\infty} \text{ and } \|\nabla \zeta\|_{L^\infty} \leq 1 \right\},$$

which implies (see [23]) that

$$(4.19) \quad S(u) = 8\pi \operatorname{Min}_{\sigma} \sum_{i=1}^{\ell} |P_i - N_{\sigma(i)}|,$$

where the  $\operatorname{Min}_{\sigma}$  is taken over all permutations  $\sigma$  of the integers  $\{1, \dots, \ell\}$ . This formula is closely connected to optimal transport as explained in [19, 26].

Combining (4.16) and (4.19), we derive a remarkable explicit formula for  $R$  when  $u \in \mathcal{C}$ :

$$R(u) = \int_{\Omega} |\nabla u|^2 + 8\pi \operatorname{Min}_{\sigma} \sum_{i=1}^{\ell} |P_i - N_{\sigma(i)}|.$$

In fact, there is a similar formula, just slightly more complicated for a general  $u \in H_g^1(\Omega; \mathbb{S}^2)$ . Namely, one can show (see [26]) that, for every  $u \in H_g^1(\Omega; \mathbb{S}^2)$ , there exist sequences  $(P_i)$  and  $(N_i)$  such that  $\sum_{i=1}^{\infty} |P_i - N_i| < \infty$ , and

$$Ju = \frac{4\pi}{3} \sum_{i=1}^{\infty} (\delta_{P_i} - \delta_{N_i}).$$

Moreover,

$$S(u) = 8\pi \operatorname{Inf} \left\{ \sum_{i=1}^{\infty} |\tilde{P}_i - \tilde{N}_i|; \sum (\delta_{\tilde{P}_i} - \delta_{\tilde{N}_i}) = \sum (\delta_{P_i} - \delta_{N_i}) \right\}.$$

The relaxed energy consists, therefore, of the usual energy  $\int_{\Omega} |\nabla u|^2$  plus an additional term involving the “interaction of singularities” – a quantity which may possibly be of physical interest.

Some partial regularity results concerning the minimizers of the relaxed energy have been obtained by Giaquinta–Modica–Soucek, Hardt–Lin–Poon and others, but the answer to Open Problem 4.3 remains elusive. An easier question still unresolved is the following.

**OPEN PROBLEM 4.4.** Let  $\underline{u}$  be a minimizer of (4.9). Is it true that  $S(\underline{u}) = 0$ , i.e.,  $J\underline{u} = 0$ ? or equivalently that there exists a sequence  $(u_n)$  in  $H_g^1(\Omega; \mathbb{S}^2) \cap C(\bar{\Omega})$  such that  $u_n$  converges to  $\underline{u}$  strongly in  $H^1$ ?

## 5. DEGREE, VMO, $W^{1/p,p}$ , AND FOURIER

Recall that

$$\begin{aligned} & \text{BMO}(\mathbb{S}^1) \\ & := \left\{ f \in L^1(\mathbb{S}^1; \mathbb{C}); |f|_{\text{BMO}} := \sup_I \frac{1}{|I|^2} \int_I \int_I |f(x) - f(y)| dx dy < \infty \right\}, \end{aligned}$$

where  $\sup_I$  is taken over all arcs of circle in  $\mathbb{S}^1$ ,

$$\text{VMO}(\mathbb{S}^1) := \left\{ f \in \text{BMO}(\mathbb{S}^1); \lim_{|I| \rightarrow 0} \frac{1}{|I|^2} \int_I \int_I |f(x) - f(y)| dx dy = 0 \right\}.$$

Obviously,  $L^\infty \subset \text{BMO}$  and  $C \subset \text{VMO}$ ; moreover, VMO is the closure of  $C$  in BMO.

For any  $1 < p < \infty$ ,

$$\begin{aligned} & W^{1/p,p}(\mathbb{S}^1) \\ & := \left\{ f \in L^1(\mathbb{S}^1; \mathbb{C}); |f|_{W^{1/p,p}}^p := \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \frac{|f(x) - f(y)|^p}{|x - y|^2} dx dy < \infty \right\}. \end{aligned}$$

As usual,  $H^{1/2} = W^{1/2,2}$ . It follows easily from Hölder that

$$W^{1/p,p}(\mathbb{S}^1) \subset \text{VMO}(\mathbb{S}^1) \quad \forall p \in (1, \infty).$$

Clearly, the classes  $W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$  increase with  $p$  on  $(1, \infty)$ ; indeed if  $p > q$ ,

$$|f(x) - f(y)|^p \leq 2^{p-q} |f(x) - f(y)|^q.$$

Brezis–Nirenberg [31] have established that degree theory persists in  $\text{VMO}(\mathbb{S}^1; \mathbb{S}^1)$ ; in particular, the new degree coincides with the classical degree on  $C(\mathbb{S}^1; \mathbb{S}^1)$ , and if  $f_n, f \in \text{VMO}$  satisfy  $f_n \rightarrow f$  in  $\text{BMO}$ , then  $\deg f_n \rightarrow \deg f$ .

The starting point in this section is the following estimate for the degree.

**THEOREM 5.1** ([8, Corollary 0.5]). *For every  $1 < p < \infty$ , there exists a constant  $C_p$  (depending only on  $p$ ) such that*

$$(5.1) \quad |\deg f| \leq C_p |f|_{W^{1/p,p}}^p \quad \forall f \in W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1).$$

Note that estimate (5.1) “deteriorates” as  $p \searrow 1$  since  $|f|_{W^{1/p,p}} \rightarrow \infty$  as  $p \searrow 1$  unless  $f$  is a constant (see [14] and the beginning of Section 9). Therefore, it is tempting to monitor the behavior of the constant  $C_p$  as  $p \searrow 1$ . A reasonable conjecture is the following.

**OPEN PROBLEM 5.1** ([17, Remark 7]). Does there exist a (universal) constant  $c$  such that, for every  $1 < p \leq 2$ ,

$$(5.2) \quad |\deg f| \leq c(p-1) \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \frac{|f(x) - f(y)|^p}{|x - y|^2} \quad \forall f \in W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)?$$

There is strong evidence in support of (5.2) as  $p \searrow 1$ . Indeed, we have (as a consequence of [28, Proposition 1]), when  $f$  is smooth,

$$\lim_{p \searrow 1} (p-1) \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \frac{|f(x) - f(y)|^p}{|x - y|^2} \simeq \int_{\mathbb{S}^1} |\dot{f}|,$$

while

$$|\deg f| \leq \frac{1}{2\pi} \int_{\mathbb{S}^1} |\dot{f}| \quad \text{by the Cauchy formula.}$$

A far-reaching extension of (5.1) is the following striking estimate for the degree.

**THEOREM 5.2.** *There exists a constant  $C$  such that for every  $f \in C(\mathbb{S}^1; \mathbb{S}^1)$*

$$(5.3) \quad |\deg f| \leq C \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \frac{dx dy}{|x - y|^2} \Big|_{[|f(x) - f(y)| \geq \sqrt{3}]}$$

A weaker version of (5.3) where  $\sqrt{3}$  is replaced by a small constant  $\delta_0 > 0$  was originally announced in [8] and proved in [6, Theorem 4]. Subsequently, Bourgain–Brezis–Nguyen [9] pushed the estimate up to  $\delta_0 = \sqrt{2}$ . Finally, Nguyen [53] established that (5.3) holds and that  $\sqrt{3}$  is optimal (see also [26, Section 12.5]); the current proof is quite involved and it is natural to raise the following problem.

**OPEN PROBLEM 5.2** ([26, Open Problem 18]). Is there a simpler, more geometric proof of Theorem 5.2? What is the best constant  $C$  in (5.3)? Is it achieved?

Given  $f \in C(\mathbb{S}^1; \mathbb{S}^1)$  and  $\delta > 0$ , set

$$I_\delta(f) := \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \frac{dx dy}{|x - y|^2} \Big|_{[|f(x) - f(y)| \geq \delta]}$$

As a consequence of (5.3), we know that  $\forall f$ ,

$$(5.4) \quad |\deg f| \leq C I_\delta(f) \quad \forall \delta \leq \sqrt{3}.$$

Note that this estimate deteriorates as  $\delta \searrow 0$  since  $I_\delta(f) \nearrow \infty$  as  $\delta \searrow 0$  (unless  $f$  is a constant). Therefore, it is natural to try to improve (5.4) by replacing  $C$  with a constant  $C_\delta$  which tends to 0 as  $\delta \searrow 0$ . A reasonable conjecture is the following.

**OPEN PROBLEM 5.3** ([17, Open Problem 3]). Does there exist a (universal) constant  $c$  such that

$$(5.5) \quad |\deg f| \leq c \delta I_\delta(f) \quad \forall f \in C(\mathbb{S}^1; \mathbb{S}^1) \quad \forall \delta \leq \sqrt{3}?$$

There is strong evidence in support of (5.5) as  $\delta \rightarrow 0$ . Indeed, when  $f$  is smooth, we have  $\lim_{\delta \rightarrow 0} \delta I_\delta(f) \simeq \int_{\mathbb{S}^1} |\dot{f}|$  (see [18, Theorem 3.1] and [29, Proposition 1]), while  $|\deg f| \leq \frac{1}{2\pi} \int_{\mathbb{S}^1} |\dot{f}|$  by the Cauchy formula.

An  $\mathbb{S}^N$ -version of (5.5) was established by Nguyen [54] for any  $N \geq 2$ , but, surprisingly, the case  $N = 1$  remains elusive!

Nonlocal energies have become popular in recent years and it is of interest to study the “least amount of  $W^{1/p,p}$ -energy” necessary to produce a map  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  of prescribed degree  $d$  (see [17, Remark 5]). More precisely, given  $1 < p < \infty$  and  $d \in \mathbb{Z}$ , set

$$(5.6) \quad m_{p,d} := \text{Inf} \left\{ |f|_{W^{1/p,p}}^p; f \in W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1), \deg f = d \right\}.$$

From Theorem 5.1 above and [26, Theorem 12.9], we know that, for every  $1 < p < \infty$ , there exist two positive constants  $\underline{c}_p$  and  $\bar{c}_p$  such that

$$\underline{c}_p |d| \leq m_{p,d} \leq \bar{c}_p |d| \quad \forall d \in \mathbb{Z}.$$

This suggests the following.

OPEN PROBLEM 5.4 ([26, Open Problems 22 and 23]). Let  $1 < p < \infty$ . Is it true that

$$(5.7) \quad m_{p,d} = |d|m_{p,1} \quad \forall d \in \mathbb{Z}?$$

Is the Inf in (5.6) achieved?

The answer to both questions is positive for  $p = 2$  (see [26, Theorems 12.9 and 12.10]). Assuming that (5.7) does not hold, is there an explicit formula for

$$\lim_{d \rightarrow +\infty} \frac{m_{p,d}}{d} = \inf_{d > 0} \frac{m_{p,d}}{d}?$$

Recall that the Fourier coefficients of a function  $f \in L^2(\mathbb{S}^1; \mathbb{C})$  are given, for every  $n \in \mathbb{Z}$ , by

$$a_n = a_n(f) := \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\theta}) e^{-in\theta} d\theta = \frac{1}{2\pi} \int_{\mathbb{S}^1} f(z) \bar{z}^n d\ell,$$

so that

$$f(e^{i\theta}) = \sum_{n \in \mathbb{Z}} a_n e^{in\theta} \quad \text{in } L^2(0, 2\pi).$$

An elementary computation (see [17, Lemma 5] or [26, Lemma 12.5]) yields, for every  $f \in H^{1/2}(\mathbb{S}^1)$ ,

$$|f|_{H^{1/2}(\mathbb{S}^1)}^2 = \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \frac{|f(x) - f(y)|^2}{|x - y|^2} dx dy = 4\pi^2 \sum_{n \in \mathbb{Z}} |n| |a_n|^2.$$

The following striking formula connecting the degree of a map  $f \in H^{1/2}(\mathbb{S}^1; \mathbb{S}^1)$  and its Fourier coefficients was brought to light by Brezis [17, Theorem 4] (see also [26, Theorem 12.6]).

THEOREM 5.3. For every  $f \in H^{1/2}(\mathbb{S}^1; \mathbb{S}^1)$ , we have

$$(5.8) \quad \deg f = \sum_{n \in \mathbb{Z}} n |a_n(f)|^2.$$

Equality (5.8) implies, in particular, that if  $f, g \in H^{1/2}(\mathbb{S}^1; \mathbb{S}^1)$  satisfy  $|a_n(f)| = |a_n(g)| \forall n \in \mathbb{Z}$ , then  $\deg f = \deg g$ . This formula has become the starting point of a challenging direction of research labeled ‘‘Can one hear the degree?’’ in [17]. More precisely,

$$(5.9) \quad \left\{ \begin{array}{l} \text{Given two maps } f, g \in \text{VMO}(\mathbb{S}^1; \mathbb{S}^1) \text{ such that } |a_n(f)| = |a_n(g)| \\ \text{for all } n \in \mathbb{Z}, \text{ can one conclude that } \deg f = \deg g? \end{array} \right.$$

The answer to question (5.9) turns out to be negative in general.



**THEOREM 5.4** ([10]). *There exist two functions  $f, g \in C(\mathbb{S}^1; \mathbb{S}^1)$  such that  $|a_n(f)| = |a_n(g)| \forall n \in \mathbb{Z}$ , and  $\deg f \neq \deg g$ .*

The construction of Bourgain–Kozma in [10] is quite elaborate and it would be desirable to find a simpler one.

On the other hand, the answer is positive in some classes bigger than  $H^{1/2}(\mathbb{S}^1; \mathbb{S}^1)$  as follows.

**THEOREM 5.5** ([17, Corollary 2], [26, Corollary 12.3], and [45]). *Assume that  $f, g \in W^{1/3,3}(\mathbb{S}^1; \mathbb{S}^1)$  satisfy  $|a_n(f)| = |a_n(g)| \forall n \in \mathbb{Z}$ , then  $\deg f = \deg g$ . In particular, the conclusion holds if  $f, g \in C^{0,\alpha}(\mathbb{S}^1; \mathbb{S}^1)$  with  $\alpha > 1/3$ .*

Note that the assertion in Theorem 5.5 is far from obvious since  $W^{1/3,3}(\mathbb{S}^1; \mathbb{S}^1)$  is strictly bigger than  $H^{1/2}(\mathbb{S}^1; \mathbb{S}^1)$  and thus the series  $\sum_{n \in \mathbb{Z}} |n| |a_n(f)|^2$  can be divergent for a general  $f \in W^{1/3,3}(\mathbb{S}^1; \mathbb{S}^1)$ . Note also that there is a wide “gap” between the positive result in Theorem 5.5 and the counterexample by Bourgain–Kozma in Theorem 5.4. It is not known whether  $W^{1/3,3}$  is the sharp borderline.

**OPEN PROBLEM 5.5.** What happens to Theorem 5.5 when  $f, g \in W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$  with  $p > 3$  (resp.,  $f, g \in C^{0,\alpha}(\mathbb{S}^1; \mathbb{S}^1)$  with  $\alpha \leq 1/3$ )? The problem is open even under the stronger assumption that  $f \in W^{1/p,p}$  with  $2 < p \leq 3$  and  $g \in W^{1/q,q}$  with  $q > 3$  (resp.,  $f \in C^{0,\alpha}$  with  $\alpha > 1/3$  and  $g \in C^{0,\beta}$  with  $\beta \leq 1/3$ ).

Theorem 5.5 is an immediate consequence of the following summation formula.

**THEOREM 5.6** ([17, Theorem 6], [26, Theorem 12.7], and [45]). *For every  $f \in W^{1/3,3}(\mathbb{S}^1; \mathbb{S}^1)$ , we have*

$$(5.10) \quad \deg f = \lim_{\varepsilon \searrow 0} \frac{\sum_{n \in \mathbb{Z}} n |a_n(f)|^2 \frac{\sin n\varepsilon}{n\varepsilon}}{n \neq 0}.$$

*In particular, (5.10) holds if  $f \in C^{0,\alpha}(\mathbb{S}^1; \mathbb{S}^1)$  with  $\alpha > 1/3$ .*

It is not known whether different summation formulas might produce improvements of Theorem 5.5, e.g., assuming that  $f$  belongs to a larger class  $W^{1/p,p}$  with  $p > 1/3$ , or  $C^{0,\alpha}$  with  $\alpha \leq 1/3$ . In fact, it is an open problem whether one can capture the degree of any map  $f \in \text{VMO}(\mathbb{S}^1; \mathbb{S}^1)$  via a summation process involving only  $|a_n(f)|$ ,  $n \in \mathbb{Z}$ . More precisely, by a summation process we mean a family  $(\sigma_{n,\varepsilon})$ ,  $n \in \mathbb{Z}$ ,  $0 < \varepsilon < 1$ , satisfying

$$(5.11) \quad \forall \varepsilon \in (0, 1), \quad \sup_{n \in \mathbb{Z}} |n| |\sigma_{n,\varepsilon}| < \infty,$$

$$(5.12) \quad \forall n \in \mathbb{Z}, \quad \lim_{\varepsilon \searrow 0} \sigma_{n,\varepsilon} = 1.$$

Note that if (5.11)-(5.12) hold, then  $\sum_{n \in \mathbb{Z}} n |a_n(f)|^2 \sigma_{n,\varepsilon}$  is well defined  $\forall \varepsilon \in (0, 1)$  and the question of interest is whether

$$(5.13) \quad \lim_{\varepsilon \searrow 0} \sum_{n \in \mathbb{Z}} n |a_n(f)|^2 \sigma_{n,\varepsilon} = \deg f?$$

OPEN PROBLEM 5.6. Given any  $p > 3$  (resp.,  $\alpha \leq 1/3$ ), does there exist a summation process  $(\sigma_{n,\varepsilon})$ , depending only on  $p$  (resp.,  $\alpha$ ) such that (5.13) holds for every  $f \in W^{1/p,p}$  (resp.,  $f \in C^{0,\alpha}$ )?

In the same vein one can raise the following problem.

OPEN PROBLEM 5.7. Given any  $f \in C(\mathbb{S}^1; \mathbb{S}^1)$  (resp.,  $f \in \text{VMO}(\mathbb{S}^1; \mathbb{S}^1)$ ), does there exist a summation process  $(\sigma_{n,\varepsilon})$  (depending on  $f$ ) such that (5.13) holds?

We call attention to the following assertion.

COROLLARY 5.1. *Given any summation process  $(\sigma_{n,\varepsilon})$ , there exists some  $f \in C(\mathbb{S}^1; \mathbb{S}^1)$  (depending on  $\sigma_{n,\varepsilon}$ ) such that  $\sum_{n \in \mathbb{Z}} n |a_n(f)|^2 \sigma_{n,\varepsilon}$  does not converge to  $\deg f$ .*

Corollary 5.1 is an immediate consequence (by contradiction) of Theorem 5.4. For special summation processes, explicit  $f$ 's satisfying the conclusion of Corollary 5.1 have been constructed by Korevaar [46] and Kahane [45]; these are

- (a)  $\sigma_{n,\varepsilon} := \begin{cases} 1 & \text{if } |n| \leq [1/\varepsilon] \text{ (the integer part of } 1/\varepsilon), \\ 0 & \text{if } |n| > [1/\varepsilon], \end{cases}$
- (b)  $\sigma_{n,\varepsilon} := (1 - \varepsilon)^{|n|}$ ,
- (c)  $\sigma_{n,\varepsilon} := \frac{\sin n\varepsilon}{n\varepsilon}$  if  $n \neq 0$  and  $\sigma_{0,\varepsilon} = 1$ .

A stronger version of Corollary 5.1 is open:

OPEN PROBLEM 5.8. Does there exist some  $f \in C(\mathbb{S}^1; \mathbb{S}^1)$  (resp.,  $f \in \text{VMO}(\mathbb{S}^1; \mathbb{S}^1)$ ) such that for any summation process  $(\sigma_{n,\varepsilon})$ ,  $\sum_{n \in \mathbb{Z}} n |a_n(f)|^2 \sigma_{n,\varepsilon}$  does not converge to  $\deg f$ ?

Note that a negative answer to Open Problem 5.8 amounts to a positive answer to Open Problem 5.7.

An interesting direction of research concerns the distance between the homotopy classes of  $W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$ ,  $1 < p < \infty$ , which are given by

$$\mathcal{E}_d = \{f \in W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1); \deg f = d\}, \quad d \in \mathbb{Z}.$$

There are two natural notions of distance:

$$\text{dist}_{W^{1/p,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = \text{Inf}_{f \in \mathcal{E}_{d_1}} \text{Inf}_{g \in \mathcal{E}_{d_2}} \|f - g\|_{W^{1/p,p}}$$

and

$$\text{Dist}_{W^{1/p,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = \text{Sup}_{f \in \mathcal{E}_{d_1}} \text{Inf}_{g \in \mathcal{E}_{d_2}} |f - g|_{W^{1/p,p}}.$$

It turns out (see [27, 31]) that

$$\text{dist}_{W^{1/p,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = 0, \quad \forall d_1, d_2 \in \mathbb{Z}.$$

On the other hand,  $\text{Dist}_{W^{1/p,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2})$  has an interesting interpretation.

Given  $f \in \mathcal{E}_{d_1}$ ,  $\text{Inf}_{g \in \mathcal{E}_{d_2}} |f - g|_{W^{1/p,p}}$  represents the least amount of energy required to pass from the given  $f \in \mathcal{E}_{d_1}$  to a configuration in  $\mathcal{E}_{d_2}$ , and  $\text{Dist}_{W^{1/p,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2})$  is the “highest price” one may have to pay as  $f$  runs in  $\mathcal{E}_{d_1}$ . A remarkable result of Shafrir asserts that this quantity depends only on  $|d_1 - d_2|$ .

**THEOREM 5.7** ([61]). *We have*

$$(5.14) \quad \text{Dist}_{W^{1/p,p}}^p(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = m_{p,|d_1-d_2|}, \quad \forall p \in (1, \infty), \forall d_1, d_2 \in \mathbb{Z},$$

where  $m_{p,d}$  is defined in (5.6).

In particular (when  $p = 2$ ), (5.14) becomes

$$\text{Dist}_{H^{1/2}}^2(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = 4\pi^2 |d_1 - d_2|.$$

A similar conclusion when  $\mathbb{S}^1$  is replaced by  $\mathbb{S}^N$ , with  $N \geq 2$  or just  $N = 2$ , is widely open. Consider, e.g.,  $H^1(\mathbb{S}^2; \mathbb{S}^2)$  and its homotopy classes

$$\mathcal{E}_d = \{f \in H^1(\mathbb{S}^2; \mathbb{S}^2); \deg f = d\}, \quad d \in \mathbb{Z}.$$

The quantity of interest is

$$\text{Dist}_{H^1}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = \text{Sup}_{f \in \mathcal{E}_{d_1}} \text{Inf}_{g \in \mathcal{E}_{d_2}} \|\nabla(f - g)\|_{L^2}.$$

**OPEN PROBLEM 5.9** ([27]). Is it true that

$$\text{Dist}_{H^1}^2(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = 8\pi |d_1 - d_2| \quad \forall d_1, d_2 \in \mathbb{Z}?$$

The inequality  $\leq$  is known (see [27] and the references therein), but the reverse inequality ( $\geq$ ) has been established only when  $d_2 > d_1 \geq 0$  (see [27, Proposition 7.3]). Even a much easier problem is open:

$$\text{is it true that } \text{Dist}_{H^1}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = \text{Dist}_{H^1}(\mathcal{E}_{d_2}, \mathcal{E}_{d_1}) \quad \forall d_1, d_2 \in \mathbb{Z}?$$

In a totally different (but somewhat related) direction, one may ask whether a version of the Brouwer fixed point theorem holds for VMO maps.

**OPEN PROBLEM 5.10.** Let  $B$  be the closed unit ball in  $\mathbb{R}^N$  and let  $f$  be a VMO map from  $B$  into itself. Is it true that for every  $\varepsilon > 0$ , the set

$$\{x \in B; |f(x) - x| < \varepsilon\}$$

has positive measure?

## 6. UNBOUNDED EXTREMAL SOLUTIONS

Consider the nonlinear elliptic equation

$$(6.1) \quad \begin{cases} -\Delta u = \lambda f(u) & \text{in } \Omega \subset \mathbb{R}^N, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega$  is a smooth bounded domain in  $\mathbb{R}^N$ ,  $\lambda > 0$  is a constant, and  $f : [0, +\infty) \rightarrow (0, +\infty)$  is a smooth function satisfying

$$(6.2) \quad f(0) > 0, \quad f \text{ is increasing and convex,}$$

and

$$(6.3) \quad \lim_{t \rightarrow \infty} \frac{f(t)}{t} = +\infty.$$

Some basic general results concerning problem (6.1) are summarized in the following (see [16, 34] and the references therein).

**THEOREM 6.1.** *There exists a constant  $\lambda^* \in (0, +\infty)$  such that the following hold.*

- (a) *For every  $\lambda \in (0, \lambda^*)$ , problem (6.1) admits a minimal smooth solution denoted by  $\underline{u}(\lambda)$ ; moreover  $\underline{u}(\lambda)$  increases with  $\lambda$  (for every  $x \in \Omega$ ).*
- (b) *For  $\lambda > \lambda^*$ , there is no solution of (6.1).*
- (c)  *$u^* = \lim_{\lambda \uparrow \lambda^*} \underline{u}(\lambda)$  is a weak solution of (6.1) in the sense that  $u^* \in L^1(\Omega)$ ,  $f(u^*)\delta \in L^1(\Omega)$ , where  $\delta(x) = \text{dist}(x, \partial\Omega)$  and*

$$(6.4) \quad - \int_{\Omega} u^* \Delta \zeta = \lambda^* \int_{\Omega} f(u^*) \zeta \quad \forall \zeta \in C^2(\bar{\Omega}), \zeta = 0 \text{ on } \partial\Omega.$$

It was originally established by Nedev [52] that  $u^* \in L^\infty(\Omega)$  when  $N \leq 3$ . The question whether the same conclusion holds when  $4 \leq N \leq 9$  for every  $f$  satisfying (6.2)-(6.3) was a long-standing open problem raised by Brezis [16]. It was recently solved in a splendid piece of work by Cabré–Figalli–Ros–Oton–Serra.

**THEOREM 6.2** ([35]). *Assume (6.2)-(6.3) and*

$$N \leq 9,$$

*then  $u^* \in L^\infty(\Omega)$ .*

Assumption  $N \leq 9$  is optimal since a celebrated result by Joseph–Lundgren (1973) provides an explicit solution of (6.4) when  $N \geq 10$ ,  $\Omega = B_1$  is the unit ball in  $\mathbb{R}^N$ , and  $f(u) = e^u$ . Namely,  $\lambda^* = 2(N - 2)$  and  $u^*(x) = \log(1/|x|^2)$ , so that  $u^* \notin L^\infty(\Omega)$ .

This completes the case  $N \leq 9$ . On the other hand, many interesting questions remain open when  $N \geq 10$ . Here are some of them.

OPEN PROBLEM 6.1 ([16, 34]). Assume that  $\Omega$  is a bounded smooth convex set in  $\mathbb{R}^N$ ,  $N \geq 10$ . Let  $f(u) = e^u$ . Is  $u^*$  unbounded? If the answer is negative for some domains  $\Omega$ , can one find other functions  $f$  satisfying (6.2)-(6.3) (possibly depending on  $\Omega$ ) such that  $u^* \notin L^\infty(\Omega)$ ?

OPEN PROBLEM 6.2 ([16, 34]). Assume  $u^* \notin L^\infty(\Omega)$ . What can be said about the blow-up set of  $u^*$ ? Does it consist of a single point when  $\Omega$  is convex?

## 7. ESTIMATES À LA BOURGAIN–BREZIS

The starting point in the following (non-trivial) estimate for the phase of  $\mathbb{S}^1$ -valued maps. For simplicity, we work on  $\mathbb{T}^N$ ,  $N \geq 2$ .

THEOREM 7.1 ([4, Corollary 1]). *Let  $\varphi : \mathbb{T}^N \rightarrow \mathbb{R}$  be a smooth function, and set  $u := e^{i\varphi}$ . Then*

$$(7.1) \quad \left\| \varphi - \int \varphi \right\|_{L^{N/(N-1)}} \leq C(|u|_{H^{1/2}} + |u|_{H^{1/2}}^2).$$

A basic ingredient in the proof of Theorem 7.1 is the following.

THEOREM 7.2 ([4, Theorem 2]). *Given any  $f \in L^N(\mathbb{T}^N)$  such that  $\int f = 0$ , there exists  $Y \in W^{1,N} \cap L^\infty(\mathbb{T}^N; \mathbb{R}^N)$  satisfying*

$$\operatorname{div} Y = f \quad \text{on } \mathbb{T}^N$$

and

$$\|Y\|_{W^{1,N}} + \|Y\|_{L^\infty} \leq C_N \|f\|_{L^N}.$$

It is easy to check that  $W^{1,N} \subset H^{1/2}$  and, therefore,  $Y \in H^{1/2} \cap L^\infty$  with the corresponding weaker estimate

$$(7.2) \quad \|Y\|_{H^{1/2}} + \|Y\|_{L^\infty} \leq C_N \|f\|_{L^N}.$$

The assertion of Theorem 7.2 is equivalent via duality to the estimate

$$\left\| \psi - \int \psi \right\|_{L^{N/(N-1)}} \leq C_N \|\nabla \psi\|_{W^{-1,N/(N-1)} + L^1} \quad \forall \psi,$$

while the weaker assertion (7.2) corresponds to the weaker estimate

$$(7.3) \quad \left\| \psi - \int \psi \right\|_{L^{N/(N-1)}} \leq C_N \|\nabla \psi\|_{H^{-1/2} + L^1}.$$

Theorem 7.1 can be deduced easily from estimate (7.2). Indeed, we have

$$(7.4) \quad \nabla \varphi = -i\bar{u}\nabla u.$$

Multiplying (7.4) by  $Y$  and applying (7.2) yields

$$\begin{aligned} \left| \int \varphi f \right| &= \left| \int \varphi \operatorname{div} Y \right| = \left| \int \nabla \varphi \cdot Y \right| = \left| \int \bar{u} \nabla u \cdot Y \right| \\ &\leq C |u|_{H^{1/2}} |\bar{u} Y|_{H^{1/2}} \leq C |u|_{H^{1/2}} (|u|_{H^{1/2}} \|Y\|_{L^\infty} + |Y|_{H^{1/2}}) \\ &\leq C |u|_{H^{1/2}} (1 + |u|_{H^{1/2}}) \|f\|_{L^N}, \end{aligned}$$

which implies (7.1).

It turns out that there is a substantial improvement of Theorem 7.1.

**THEOREM 7.3** ([4, Theorem 4], [7, Theorem 3], and [26, Theorem 9.7]). *With the same notations as in Theorem 7.1, we have*

$$(7.5) \quad \left\| \varphi - \int \varphi \right\|_{H^{1/2} + W^{1,1}} \leq C_N (|u|_{H^{1/2}} + |u|_{H^{1/2}}^2).$$

[Note that  $H^{1/2} \subset L^{N/(N-1)}$  and  $W^{1,1} \subset L^{N/(N-1)}$ .]

If we try to establish (7.5) by the same method as above we would need to invoke a stronger version of (7.3); namely

$$(7.6) \quad \left\| \psi - \int \psi \right\|_{H^{1/2} + W^{1,1}} \leq C \|\nabla \psi\|_{H^{-1/2} + L^1} \quad \forall \psi.$$

However, such an estimate is still undecided.

**OPEN PROBLEM 7.1** ([4]). Does (7.6) hold?

## 8. REGULARITY OF MINIMIZERS FOR FUNCTIONALS INVOLVING THE TOTAL VARIATION

Let  $\Omega$  be a smooth bounded domain in  $\mathbb{R}^N$ ,  $N \geq 2$ . Assume (for simplicity) that  $f$  is a smooth function on  $\bar{\Omega}$  and consider the functional

$$\Phi(u) := \int_{\Omega} |\nabla u| + \frac{1}{2} \int_{\Omega} |u - f|^2$$

defined for  $u \in \operatorname{BV}(\Omega) \cap L^2(\Omega)$ . This functional has been extensively used, e.g., in image processing following the classical work of Rudin–Osher–Fatemi.

Standard functional analysis yields the existence and uniqueness of a minimizer denoted by  $U \in \text{BV}(\Omega) \cap L^2(\Omega)$  for the problem

$$(8.1) \quad \text{Min}_{u \in \text{BV} \cap L^2} \Phi(u).$$

So far the best regularity result is the following.

**THEOREM 8.1** ([36, 58]). *The minimizer  $U$  satisfies  $\nabla U \in L^\infty(\Omega)$ , but in general  $\nabla U$  is not continuous.*

This still leaves room for improvement.

**OPEN PROBLEM 8.1** ([20, Open Problem 1]). Does  $\nabla U$  belong to  $\text{BV}(\Omega)$ ?

When  $N = 1$ , the answer to Open Problem 8.1 is positive (see [20, Theorem 1]). The proof is based on a transformation relating the solution of (8.1) to the derivative of the solution of an obstacle problem. One may then apply a result of [24] (valid in all dimensions  $N$ ). Unfortunately, this transformation seems to be restricted to  $N = 1$ .

## 9. CHARACTERIZATIONS OF CONSTANT FUNCTIONS AND BEYOND

Let  $\Omega$  be a smooth bounded connected domain in  $\mathbb{R}^N$ ,  $N \geq 1$ . It is known (see [14], [26, Corollary 6.4]) that any measurable function  $u : \Omega \rightarrow \mathbb{R}$  satisfying

$$\int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{p+N}} dx dy < \infty,$$

for some  $1 \leq p < \infty$ , must be a constant (i.e., there exists a constant  $c$  such that  $u(x) = c$  a.e.). As was pointed out in [14], this fact is an immediate consequence of the Bourgain–Brezis–Mironescu (BBM) formula [5]; alternative direct elementary proofs are presented in [26].

A significant extension of this result goes as follows. Given  $\lambda > 0$ ,  $p \geq 1$ , and a measurable function  $u : \Omega \rightarrow \mathbb{R}$ , set

$$\Phi_{\lambda,p}(u) := \text{meas} \left\{ (x, y) \in \Omega \times \Omega; \frac{|u(x) - u(y)|^p}{|x - y|^{p+N}} \geq \lambda \right\}.$$

**THEOREM 9.1.** *Assume that a measurable function  $u$  satisfies*

$$\lim_{\lambda \rightarrow \infty} \lambda \Phi_{\lambda,p}(u) = 0 \quad \text{for some } p \geq 1,$$

*then  $u$  must be a constant.*

Theorem 9.1 is due to Brezis–Van Schaftingen–Yung [33] when  $p > 1$  (with  $\Omega = \mathbb{R}^N$  but the proof is unchanged when  $\Omega$  is a ball) and to Poliakovsky [57, Corollary 1.1]

when  $p = 1$ . Actually, Poliakovsky derives the result from a deeper assertion (see below); as a consequence, the proof of Theorem 9.1 when  $p = 1$  is quite intricate and it would be desirable to find a more direct and elementary proof, possibly in the spirit of [33].

Here is a natural question related to Theorem 9.1.

OPEN PROBLEM 9.1 ([33, Open Problem 1]). Assume that  $u$  satisfies

$$\liminf_{\lambda \rightarrow \infty} \lambda \Phi_{\lambda,p}(u) = 0 \quad \text{for some } p \geq 1.$$

Can one conclude that  $u$  is a constant?

A far-reaching version of this question is the following.

OPEN PROBLEM 9.2 ([32, Section 7.2]). Given any  $1 < p < \infty$ , is there a constant  $C = C(p, N, \Omega)$  such that, for all measurable functions  $u$ ,

$$(9.1) \quad \|\nabla u\|_{L^p}^p \leq C \liminf_{\lambda \rightarrow \infty} \lambda \Phi_{\lambda,p}(u)?$$

in the sense that  $\|\nabla u\|_{L^p} = \infty$  if  $u \notin W^{1,p}(\Omega)$ .

Similarly for  $p = 1$ , is there a constant  $C = C(N, \Omega)$  such that, for all measurable functions  $u$ ,

$$(9.2) \quad \|\nabla u\|_M \leq C \liminf_{\lambda \rightarrow \infty} \lambda \Phi_{\lambda,1}(u)?$$

where  $\|\nabla u\|_M$  denotes the total mass of  $\nabla u$  if  $u \in \text{BV}(\Omega)$ , and  $\|\nabla u\|_M = \infty$  if  $u \notin \text{BV}(\Omega)$ .

Poliakovsky [57] gave a positive answer to Open Problem 9.2 under the stronger assumption that  $\liminf$  is replaced by  $\limsup$  in (9.1) and (9.2).

Along the same lines one may even try to go one step further.

OPEN PROBLEM 9.3 ([32, Section 7.3]). Does the family of functionals  $\lambda \Phi_{\lambda,p}$  converge as  $\lambda \rightarrow \infty$ , in the sense of  $\Gamma$ -convergence, to the functional  $\Psi$  defined, when  $p > 1$ , by

$$\Psi(u) := \begin{cases} c_p \|\nabla u\|_{L^p}^p & \text{if } u \in W^{1,p}(\Omega), \\ +\infty & \text{if } u \notin W^{1,p}(\Omega), \end{cases}$$

and when  $p = 1$  by

$$\Psi(u) := \begin{cases} c_1 \|\nabla u\|_M & \text{if } u \in \text{BV}(\Omega), \\ +\infty & \text{if } u \notin \text{BV}(\Omega), \end{cases}$$

for some positive constants  $c_p$  and  $c_1$ ?

The three problems above are already interesting when  $N = 1$ !



## 10. A SHARP RELATIVE ISOPERIMETRIC INEQUALITY FOR THE CUBE

Let  $Q$  be the unit cube in  $\mathbb{R}^N$ ,  $N \geq 2$ . Given a measurable set  $S \subset Q$ , denote by  $\mathbf{1}_S$  the characteristic function of  $S$ , by  $|S| = \|\mathbf{1}_S\|_{L^1}$  the volume of  $S$ , and by  $P(S)$  the relative perimeter of  $S$ , i.e., taking into account only the part of the boundary of  $S$  inside  $Q$ ; in other words  $P(S)$  is the total mass of the measure  $\nabla \mathbf{1}_S$  (possibly infinite if  $S$  is not rectifiable). Consider the function  $f_N(t)$  defined for  $0 \leq t \leq 1$  by

$$(10.1) \quad f_N(t) := \inf\{P(S); S \text{ is a measurable subset of } Q \text{ such that } |S| = t\}.$$

Clearly,  $f_N(t) = f_N(1-t) \forall t \in [0, 1]$ ; just replace  $S$  by  $Q \setminus S$  in (10.1). An explicit formula for  $f_N(t)$  is known when  $N = 2$ .

THEOREM 10.1 ([21]). *We have*

$$(10.2) \quad f_2(t) = \begin{cases} (\pi t)^{1/2} & \text{if } 0 \leq t \leq \frac{1}{\pi}, \\ 1 & \text{if } \frac{1}{\pi} \leq t \leq \frac{1}{2}. \end{cases}$$

OPEN PROBLEM 10.1 ([21]). Is there a formula similar to (10.2) for  $f_N(t)$  when  $N \geq 3$ ? In particular, is it true that  $f_N(t) \equiv 1$  near  $t = 1/2$ ?

## REFERENCES

- [1] F. BETHUEL, [A characterization of maps in  \$H^1\(B^3, S^2\)\$  which can be approximated by smooth maps](#). *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **7** (1990), no. 4, 269–286. Zbl [0708.58004](#) MR [1067776](#)
- [2] F. BETHUEL – H. BREZIS – J.-M. CORON, [Relaxed energies for harmonic maps](#). In *Variational methods (Paris, 1988)*, pp. 37–52, Progr. Nonlinear Differential Equations Appl. 4, Birkhäuser, Boston, MA, 1990. Zbl [0793.58011](#) MR [1205144](#)
- [3] F. BETHUEL – H. BREZIS – F. HÉLEIN, [Ginzburg–Landau vortices](#). Progr. Nonlinear Differential Equations Appl. 13, Birkhäuser, Boston, MA, 1994. Zbl [0802.35142](#) MR [1269538](#)
- [4] J. BOURGAIN – H. BREZIS, [On the equation  \$\operatorname{div} Y = f\$  and application to control of phases](#). *J. Amer. Math. Soc.* **16** (2003), no. 2, 393–426. Zbl [1075.35006](#) MR [1949165](#)
- [5] J. BOURGAIN – H. BREZIS – P. MIRONESCU, [Another look at Sobolev spaces](#). In *Optimal control and partial differential equations*, pp. 439–455, IOS, Amsterdam, 2001. Zbl [1103.46310](#) MR [3586796](#)
- [6] J. BOURGAIN – H. BREZIS – P. MIRONESCU, [Complements to the paper: Lifting, degree, and distributional Jacobian, revisited](#). 2004, <https://hal.sorbonne-universite.fr/LJLL/hal-00747668> visited on 16 November 2023.

- [7] J. BOURGAIN – H. BREZIS – P. MIRONESCU,  $H^{1/2}$  maps with values into the circle: minimal connections, lifting, and the Ginzburg–Landau equation. *Publ. Math. Inst. Hautes Études Sci.* (2004), no. 99, 1–115. Zbl [1051.49030](#) MR [2075883](#)
- [8] J. BOURGAIN – H. BREZIS – P. MIRONESCU, Lifting, degree, and distributional Jacobian revisited. *Comm. Pure Appl. Math.* **58** (2005), 529–551. Zbl [1077.46023](#) MR [2119868](#)
- [9] J. BOURGAIN – H. BREZIS – H.-M. NGUYEN, A new estimate for the topological degree. *C. R. Math. Acad. Sci. Paris* **340** (2005), no. 11, 787–791. Zbl [1071.55002](#) MR [2139888](#)
- [10] J. BOURGAIN – G. KOZMA, One cannot hear the winding number. *J. Eur. Math. Soc. (JEMS)* **9** (2007), no. 4, 637–658. Zbl [1134.42301](#) MR [2341826](#)
- [11] H. BREZIS, Elliptic equations with limiting Sobolev exponents—the impact of topology. *Comm. Pure Appl. Math.* **39** (1986), S17–S39. Zbl [0601.35043](#) MR [861481](#)
- [12] H. BREZIS,  $S^k$ -valued maps with singularities. In *Topics in calculus of variations (Montecatini Terme, 1987)*, pp. 1–30, Lecture Notes in Math. 1365, Springer, Berlin, 1989. Zbl [0684.49015](#) MR [994017](#)
- [13] H. BREZIS, Symmetry in nonlinear PDE’s. In *Differential equations: La Pietra 1996 (Florence)*, pp. 1–12, Proc. Sympos. Pure Math. 65, American Mathematical Society, Providence, RI, 1999. Zbl [0927.35038](#) MR [1662746](#)
- [14] H. BREZIS, How to recognize constant functions. A connection with Sobolev spaces. *Uspekhi Mat. Nauk* **57** (2002), no. 4(346), 59–74; translation in *Russian Math. Surveys* **57** (2002), no. 4, 693–708. Zbl [1072.46020](#) MR [1942116](#)
- [15] H. BREZIS, The interplay between analysis and topology in some nonlinear PDE problems. *Bull. Amer. Math. Soc. (N.S.)* **40** (2003), no. 2, 179–201. Zbl [1161.35354](#) MR [1962295](#)
- [16] H. BREZIS, Is there failure of the inverse function theorem? In *Morse theory, minimax theory and their applications to nonlinear differential equations*, pp. 23–33, New Stud. Adv. Math. 1, International Press, Somerville, MA, 2003. Zbl [1200.35144](#) MR [2056500](#)
- [17] H. BREZIS, New questions related to the topological degree. In *The unity of mathematics*, pp. 137–154, Progr. Math. 244, Birkhäuser, Boston, MA, 2006. Zbl [1109.58017](#) MR [2181804](#)
- [18] H. BREZIS, New approximations of the total variation and filters in imaging. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **26** (2015), no. 2, 223–240. Zbl [1325.26036](#) MR [3341570](#)
- [19] H. BREZIS, Remarks on the Monge–Kantorovich problem in the discrete setting. *C. R. Math. Acad. Sci. Paris* **356** (2018), no. 2, 207–213. Zbl [1384.49037](#) MR [3758722](#)
- [20] H. BREZIS, Remarks on some minimization problems associated with BV norms. *Discrete Contin. Dyn. Syst.* **39** (2019), no. 12, 7013–7029. Zbl [1428.26017](#) MR [4026179](#)
- [21] H. BREZIS – A. BRUCKSTEIN, A sharp relative isoperimetric inequality for the square. *C. R. Math. Acad. Sci. Paris* **359** (2021), 1191–1199. MR [4346201](#)

- [22] H. BREZIS – J.-M. CORON, [Large solutions for harmonic maps in two dimensions](#). *Comm. Math. Phys.* **92** (1983), no. 2, 203–215. Zbl [0532.58006](#) MR [728866](#)
- [23] H. BREZIS – J.-M. CORON – E. H. LIEB, [Harmonic maps with defects](#). *Comm. Math. Phys.* **107** (1986), no. 4, 649–705. Zbl [0608.58016](#) MR [868739](#)
- [24] H. BREZIS – D. KINDERLEHRER, [The smoothness of solutions to nonlinear variational inequalities](#). *Indiana Univ. Math. J.* **23** (1973/74), 831–844. Zbl [0278.49011](#) MR [361436](#)
- [25] H. BREZIS – F. MERLE – T. RIVIÈRE, [Quantization effects for  \$-\Delta u = u\(1 - |u|^2\)\$  in  \$\mathbf{R}^2\$](#) . *Arch. Rational Mech. Anal.* **126** (1994), no. 1, 35–58. Zbl [0809.35019](#) MR [1268048](#)
- [26] H. BREZIS – P. MIRONESCU, [Sobolev maps to the circle—from the perspective of analysis, geometry, and topology](#). *Progr. Nonlinear Differential Equations Appl.* 96, Birkhäuser, New York, 2021. Zbl [1501.46001](#) MR [4390036](#)
- [27] H. BREZIS – P. MIRONESCU – I. SHAFRIR, [Distances between homotopy classes of  \$W^{s,p}\(\mathbb{S}^N; \mathbb{S}^N\)\$](#) . *ESAIM Control Optim. Calc. Var.* **22** (2016), no. 4, 1204–1235. Zbl [1371.46027](#) MR [3570500](#)
- [28] H. BREZIS – H.-M. NGUYEN, [Two subtle convex nonlocal approximations of the BV-norm](#). *Nonlinear Anal.* **137** (2016), 222–245. Zbl [1344.49017](#) MR [3485124](#)
- [29] H. BREZIS – H.-M. NGUYEN, [Non-local functionals related to the total variation and connections with image processing](#). *Ann. PDE* **4** (2018), no. 1, article no. 9. Zbl [1398.49042](#) MR [3749763](#)
- [30] H. BREZIS – L. NIRENBERG, [Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents](#). *Comm. Pure Appl. Math.* **36** (1983), no. 4, 437–477. Zbl [0541.35029](#) MR [709644](#)
- [31] H. BREZIS – L. NIRENBERG, [Degree theory and BMO. I. Compact manifolds without boundaries](#). *Selecta Math. (N.S.)* **1** (1995), no. 2, 197–263. Zbl [0852.58010](#) MR [1354598](#)
- [32] H. BREZIS – A. SEEGER – J. VAN SCHAFTINGEN – P.-L. YUNG, [Families of functionals representing Sobolev norms](#). 2021, *Analysis & PDE* (to appear); arXiv:[2109.02930](#).
- [33] H. BREZIS – J. VAN SCHAFTINGEN – P.-L. YUNG, [Going to Lorentz when fractional Sobolev, Gagliardo and Nirenberg estimates fail](#). *Calc. Var. Partial Differential Equations* **60** (2021), no. 4, article no. 129. Zbl [1471.35008](#) MR [4279388](#)
- [34] H. BREZIS – J. L. VÁZQUEZ, [Blow-up solutions of some nonlinear elliptic problems](#). *Rev. Mat. Univ. Complut. Madrid* **10** (1997), no. 2, 443–469. Zbl [0894.35038](#) MR [1605678](#)
- [35] X. CABRÉ – A. FIGALLI – X. ROS-OTON – J. SERRA, [Stable solutions to semilinear elliptic equations are smooth up to dimension 9](#). *Acta Math.* **224** (2020), no. 2, 187–252. Zbl [1467.35172](#) MR [4117051](#)
- [36] V. CASELLES – A. CHAMBOLLE – M. NOVAGA, [Regularity for solutions of the total variation denoising problem](#). *Rev. Mat. Iberoam.* **27** (2011), no. 1, 233–252. Zbl [1228.94005](#) MR [2815736](#)

- [37] M. COMTE, [Solutions of elliptic equations with critical Sobolev exponent in dimension three](#). *Nonlinear Anal.* **17** (1991), no. 5, 445–455. Zbl 0754.35040 MR 1124117
- [38] O. DRUET, [Elliptic equations with critical Sobolev exponents in dimension 3](#). *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **19** (2002), no. 2, 125–142. Zbl 1011.35060 MR 1902741
- [39] P. ESPOSITO, [Some remarks concerning symmetry-breaking for the Ginzburg–Landau equation](#). *J. Funct. Anal.* **265** (2013), no. 10, 2189–2203. Zbl 1283.35133 MR 3091811
- [40] R. HARDT – F.-H. LIN, [A remark on  \$H^1\$  mappings](#). *Manuscripta Math.* **56** (1986), no. 1, 1–10. Zbl 0618.58015 MR 846982
- [41] R. M. HARDT, [Singularities of harmonic maps](#). *Bull. Amer. Math. Soc. (N.S.)* **34** (1997), no. 1, 15–34. Zbl 0871.58026 MR 1397098
- [42] R. IGNAT – L. NGUYEN, [Local minimality of  \$\mathbb{R}^N\$ -valued and  \$\mathbb{S}^N\$ -valued Ginzburg–Landau vortex solutions in the unit ball  \$B^N\$](#) . 2021, arXiv:2111.07669.
- [43] R. IGNAT – L. NGUYEN – V. SLASTIKOV – A. ZARNESCU, [Uniqueness of degree-one Ginzburg–Landau vortex in the unit ball in dimensions  \$N \geq 7\$](#) . *C. R. Math. Acad. Sci. Paris* **356** (2018), no. 9, 922–926. Zbl 1398.49003 MR 3849078
- [44] J. JOST, [The Dirichlet problem for harmonic maps from a surface with boundary onto a 2-sphere with nonconstant boundary values](#). *J. Differential Geom.* **19** (1984), no. 2, 393–401. Zbl 0551.58012 MR 755231
- [45] J.-P. KAHANE, [Sur l'équation fonctionnelle  \$\int\_{\mathbb{T}} \(\psi\(t+s\) - \psi\(s\)\)^3 ds = \sin t\$](#) . *C. R. Math. Acad. Sci. Paris* **341** (2005), no. 3, 141–145. Zbl 1085.45006 MR 2158834
- [46] J. KOREVAAR, [On a question of Brézis and Nirenberg concerning the degree of circle maps](#). *Selecta Math. (N.S.)* **5** (1999), no. 1, 107–122. Zbl 0938.55005 MR 1694896
- [47] M. KURZKE, [The correlation coefficient for vortices in the plane](#). *J. Math. Anal. Appl.* **479** (2019), no. 1, 333–336. Zbl 1457.76049 MR 3987037
- [48] L. LEMAIRE, [Applications harmoniques de surfaces riemanniennes](#). *J. Differential Geometry* **13** (1978), no. 1, 51–78. Zbl 0388.58003 MR 520601
- [49] E. H. LIEB – M. LOSS, [Symmetry of the Ginzburg–Landau minimizer in a disc](#). *Math. Res. Lett.* **1** (1994), no. 6, 701–715. Zbl 0842.49014 MR 1306015
- [50] P. MIRONESCU, [On the stability of radial solutions of the Ginzburg–Landau equation](#). *J. Funct. Anal.* **130** (1995), no. 2, 334–344. Zbl 0839.35011 MR 1335384
- [51] P. MIRONESCU, [Les minimiseurs locaux pour l'équation de Ginzburg–Landau sont à symétrie radiale](#). *C. R. Acad. Sci. Paris Sér. I Math.* **323** (1996), no. 6, 593–598. Zbl 0858.35038 MR 1411048
- [52] G. NEDEV, [Regularity of the extremal solution of semilinear elliptic equations](#). *C. R. Acad. Sci. Paris Sér. I Math.* **330** (2000), no. 11, 997–1002. Zbl 0955.35029 MR 1779693
- [53] H.-M. NGUYEN, [Optimal constant in a new estimate for the degree](#). *J. Anal. Math.* **101** (2007), 367–395. Zbl 1147.47046 MR 2346551

- [54] H.-M. NGUYEN, [A refined estimate for the topological degree](#). *C. R. Math. Acad. Sci. Paris* **355** (2017), no. 10, 1046–1049. Zbl [1422.55004](#) MR [3716480](#)
- [55] Y. N. OVCHINNIKOV – I. M. SIGAL, [Symmetry-breaking solutions of the Ginzburg–Landau equation](#). *Zh. Èksper. Teoret. Fiz.* **126** (2004), no. 5, 1249–1266. MR [2108700](#)
- [56] F. PACARD – T. RIVIÈRE, [Linear and nonlinear aspects of vortices](#). Progr. Nonlinear Differential Equations Appl. 39, Birkhäuser, Boston, MA, 2000. Zbl [0948.35003](#) MR [1763040](#)
- [57] A. POLIAKOVSKY, [Some remarks on a formula for Sobolev norms due to Brezis, Van Schaftingen and Yung](#). *J. Funct. Anal.* **282** (2022), no. 3, article no. 109312. Zbl [1479.35029](#) MR [4339013](#)
- [58] A. PORRETTA, [On the regularity of the total variation minimizers](#). *Commun. Contemp. Math.* **23** (2021), no. 1, article no. 1950082. Zbl [1450.35096](#) MR [4169243](#)
- [59] R. SCHOEN – K. UHLENBECK, [A regularity theory for harmonic maps](#). *J. Differential Geometry* **17** (1982), no. 2, 307–335. Zbl [0521.58021](#) MR [664498](#)
- [60] I. SHAFRIR, [Remarks on solutions of  \$-\Delta u = \(1 - |u|^2\)u\$  in  \$\mathbf{R}^2\$](#) . *C. R. Acad. Sci. Paris Sér. I Math.* **318** (1994), no. 4, 327–331. Zbl [0806.35030](#) MR [1267609](#)
- [61] I. SHAFRIR, [On the distance between homotopy classes in  \$W^{1/p,p}\(\mathbb{S}^1, \mathbb{S}^1\)\$](#) . *Confluentes Math.* **10** (2018), no. 1, 125–136. Zbl [1493.46058](#) MR [3869013](#)

---

Received 26 August 2022,  
and in revised form 6 October 2022

Haïm Brezis  
Department of Mathematics, Rutgers University,  
Hill Center, Busch Campus, 110 Frelinghuysen Rd, Piscataway, NJ 08854, USA  
[hbrezis@gmail.com](mailto:hbrezis@gmail.com)