



Mathematical Physics. – *Propagation of chaos for topological interactions by a coupling technique*, by PIERRE DEGOND, MARIO PULVIRENTI and STEFANO ROSSI, communicated on 21 April 2023.

ABSTRACT. – We consider a system of particles that interact through a jump process. The jump intensities are functions of the proximity rank of the particles, a type of interaction referred to as topological in the literature. Such interactions have been shown relevant for the modelling of bird flocks. We show that, in the large number of particles limit and under minimal smoothness assumptions on the data, the model converges to a kinetic equation which was derived in earlier works both formally and rigorously under more stringent regularity assumptions. The proof relies on the coupling method which assigns to the particle and limiting processes a joint process posed on the cartesian product of the two configuration spaces of the former processes. By appropriate estimates in a suitable Wasserstein metric, we show that the distance between the two processes tends to zero as the number of particles tends to infinity, with an error typical of the law of large numbers.

KEYWORDS. – Propagation of chaos, rank-based interaction, coupling method.

2020 MATHEMATICS SUBJECT CLASSIFICATION. – Primary 35Q92; Secondary 70K55, 92D50.

1. INTRODUCTION

Systems of self-propelled agents undergoing local interactions are ubiquitous in nature, from migrating cells [16] to locust swarms [2] and fish schools [18]. They form intriguing patterns such as coherent motion, travelling bands, oscillations, etc., encompassed in the generic term of collective dynamics (see a review in [24]). Most models of collective dynamics are based on mean-field interactions (such as the Cucker–Smale [12] or Vicsek [23] models) or binary contact interactions [4]. However, a third type of interaction has been suggested following observations of bird flocks [1, 9] and referred to as “topological interaction”. In this kind of interaction, the strength of the interaction of an agent with another one is a function of the proximity rank of the latter with respect to the former. The seminal paper [1] has been followed by a number of papers studying various aspects of this phenomenon; see, e.g., [7, 8, 15, 20, 21].

Mathematically, flocking of systems of topologically interacting particles has been investigated in [19, 22, 26]. In [17], in addition to studying flocking, the author proposes kinetic and fluid models derived from mean-field topological interactions. The present

work is strongly aligned with [5, 6, 13] where kinetic models are derived for topological interaction models based on jump processes. More precisely, [13] proves propagation of chaos and provides a rigorous proof of the model formally derived in [5]. The proof of [13] makes the limiting assumption that the interaction strength is an analytic function of the normalized rank (a concept precisely defined below) and is based on the BBGKY hierarchy. In the present work, we propose an alternative proof of the result of [13] based on the coupling method. The advantage of the coupling method over the BBGKY hierarchy is that it only requires the interaction strength to be Lipschitz continuous, a much more general and natural assumption than that of [13]. On the other hand, [6] formally derives a kinetic model for a more singular interaction. The mathematical validity of this formal result is still open. The literature on propagation of chaos and derivation of kinetic models from particle ones is huge, and it is difficult to provide a fair account of all relevant contributions in a short introduction. We refer the interested reader to the reviews [10, 11] which provide a fairly detailed description of the subject.

The outline of this paper is as follows. In Section 2, we present the model and provide a formal derivation of the macroscopic model. We then state the theorem and comment on it in view of the previous results. Section 3 is devoted to the proof.

2. PRESENTATION OF THE MODEL AND MAIN RESULTS

We recall the model and notations introduced in [5, 13] and state our result. We study an N -particle system in \mathbb{R}^d , $d = 1, 2, 3, \dots$ (or in \mathbb{T}^d the d -dimensional torus). Each particle, say particle i , has a position x_i and velocity v_i . The configuration of the system is denoted by

$$Z_N = \{z_i\}_{i=1}^N = \{(x_i, v_i)\}_{i=1}^N = (X_N, V_N).$$

Given the particle i , we order the remaining particles j_1, j_2, \dots, j_{N-1} according to their distance from i , namely, by the following relation:

$$|x_i - x_{j_h}| \leq |x_i - x_{j_{h+1}}|, \quad h = 1, 2, \dots, N-1.$$

The rank $R(i, k)$ of particle $k = j_h$ (with respect to i) is h . Note that if $B_r(x)$ denotes the closed ball of center $x \in \mathbb{R}^d$ and radius $r > 0$, we have

$$R(i, k) = \sum_{\substack{1 \leq h \leq N \\ h \neq i}} \mathcal{X}_{B_{|x_i - x_k|}(x_i)}(x_h),$$

where \mathcal{X}_A is the characteristic function of the set A .

Given a non-increasing Lipschitz continuous function

$$K : [0, 1] \rightarrow \mathbb{R}^+ \quad \text{s.t.} \quad \int_0^1 K(r) \, dr = 1,$$

we introduce the transition probabilities

$$(2.1) \quad \pi_{i,j}^N = \frac{K(r(i, j))}{\sum_{s=1}^N K\left(\frac{s}{N-1}\right)},$$

where $r(i, j)$ is the normalized rank:

$$r(i, j) = \frac{R(i, j)}{N-1} \in \left\{ \frac{1}{N-1}, \frac{2}{N-1}, \dots \right\}.$$

Thanks to the normalization in (2.1), we have that $\sum_j \pi_{i,j}^N = 1$. We can also rewrite $\pi_{i,j}^N$ as

$$(2.2) \quad \pi_{i,j}^N = \alpha_N K(r(i, j)),$$

where

$$(2.3) \quad \alpha_N = \frac{1}{(N-1)(1 - e_K(N))}$$

and $e_K(N)$ is the error given by the Riemann sums

$$(2.4) \quad e_K(N) = \int_0^1 K(x) \, dx - \frac{1}{N-1} \sum_s K\left(\frac{s}{N-1}\right).$$

We are now in position to introduce a stochastic process describing alignment via a topological interaction. The particles go freely: $x_i + v_i t$. At some random time dictated by a Poisson process of intensity N , choose a particle (say i) with probability $\frac{1}{N}$ and a partner particle, say j , with probability $\pi_{i,j}$. Then, perform the transition

$$(v_i, v_j) \rightarrow (v_j, v_j).$$

After that, the system goes freely with the new velocities and so on.

The process is described by the following Markov generator given, for any $\Phi \in C_b^1(\mathbb{R}^{2dN})$, by

$$(2.5) \quad L_N \Phi(X_N, V_N) = \sum_{i=1}^N v_i \cdot \nabla_{x_i} \Phi(X_N, V_N) + \sum_{i=1}^N \sum_{\substack{1 \leq j \leq N \\ i \neq j}} \pi_{i,j}^N [\Phi(X_N, V_N^i(v_j)) - \Phi(X_N, V_N)],$$

where

$$V_N^i(v_j) = (v_1, \dots, v_{i-1}, v_j, v_{i+1}, \dots, v_N) \quad \text{if } V_N = (v_1, \dots, v_{i-1}, v_i, v_{i+1}, \dots, v_N).$$

Note that $\pi_{i,j}^N$ depends not only on N but also on the whole spatial configuration X_N . Therefore, the law of the process $\mathcal{W}^N(Z_N; t) dZ_N$ is driven by the following evolution equation:

$$(2.6) \quad \begin{aligned} \partial_t \int \mathcal{W}^N(Z_N; t) \Phi(Z_N) dZ_N \\ = \sum_{i=1}^N \int \mathcal{W}^N(Z_N; t) v_i \cdot \nabla_{x_i} \Phi(Z_N) dZ_N \\ + \sum_{i=1}^N \sum_{\substack{1 \leq j \leq N \\ i \neq j}} \int \mathcal{W}^N(Z_N; t) \pi_{i,j}^N [\Phi(X_N, V_N^i(v_j)) - \Phi(X_N, V_N)] dZ_N, \end{aligned}$$

for any test function Φ . Here, $\mathcal{W}^N(Z_N; t)$ is the density with respect to the Lebesgue measure.

We assume that the initial measure $\mathcal{W}^N(Z_N; 0) dZ_N$ factorizes; namely, $\mathcal{W}^N(0) = f_0^{\otimes N}$, where f_0 is the initial datum for the limiting kinetic equation we are going to establish. Note also that $\mathcal{W}^N(Z_N; t)$, for $t \geq 0$, is symmetric in the exchange of particles.

The strong form of equation (2.6) is

$$\left(\partial_t + \sum_{i=1}^N v_i \cdot \nabla_{x_i} \right) \mathcal{W}^N(t) = -N \mathcal{W}^N(t) + \mathcal{L}_N \mathcal{W}^N(t)$$

where

$$\mathcal{L}_N \mathcal{W}^N(X_N, V_N; t) = \sum_{i=1}^N \sum_{\substack{1 \leq j \leq N \\ i \neq j}} \delta(v_i - v_j) \int \pi_{i,j}^N \mathcal{W}^N(X_N, V_N^{(i)}(u); t) du.$$

2.1. Heuristic derivation

We now want to derive the kinetic equation we expect to be valid in the limit $N \rightarrow \infty$. Setting $\Phi(Z_N) = \varphi(z_1)$ in (2.6), we obtain

$$(2.7) \quad \begin{aligned} \partial_t \int f_1^N(z_1) \varphi(z_1) dz_1 \\ = \int f_1^N(z_1) v_1 \cdot \nabla_{x_1} \varphi(z_1) dz_1 - \int f_1^N(z_1) \varphi(z_1) dz_1 \\ + \sum_{j \neq 1} \int \mathcal{W}^N(Z_N; t) \pi_{1,j}^N \varphi(x_1, v_j) dZ_N. \end{aligned}$$

Here, f_1^N denotes the one-particle marginal of the measure \mathcal{W}^N . We recall that the s -particle marginals are defined by

$$(2.8) \quad f_s^N(Z_s) = \int \mathcal{W}^N(Z_s, z_{s+1}, \dots, z_N) dz_{s+1}, \dots, dz_N, \quad s = 1, 2, \dots, N,$$

and are the distribution of the first s particles (or of any group of s tagged particles).

In order to describe the system in terms of a single kinetic equation, we expect that chaos propagates. Actually, since \mathcal{W}^N is initially factorizing, although the dynamics creates correlations, we hope that, due to the weakness of the interaction, factorization still holds approximately also at any positive time t ; namely,

$$f_s^N \approx f_1^{\otimes s}.$$

In this case, the law of large numbers does hold; that is,

$$\frac{1}{N} \sum_j \delta(z - z_j) \approx f_1^N(z, t)$$

for \mathcal{W}^N -almost all $Z_N = \{z_1, \dots, z_N\}$. Then,

$$\begin{aligned} \pi_{i,j}^N &\approx \frac{1}{N-1} K \left(\frac{1}{N-1} \sum_k \mathcal{X}_{B_{|x_i-x_j|}(x_i)}(x_k) \right) \\ &\approx \frac{1}{N-1} K(M_{\rho_1^N}(B_{|x_1-x_2|}(x_1))) \end{aligned}$$

where

$$(2.9) \quad M_{\rho_1^N}(B_R(x)) = \int_{B_R(x)} \rho_1^N(y) dy,$$

and $\rho_1^N(x) = \int dv f_1^N(x, v)$ is the spatial density. Motivated by this remark, from now on, we use the following notation:

$$M_{X_N}(B_{|x_i-x_j|}(x_i)) = r(i, j) = \frac{1}{N-1} \sum_k \mathcal{X}_{B_{|x_i-x_j|}(x_i)}(x_k).$$

Here, M stands for ‘‘mass’’ and the notation introduced is justified by the law of large numbers.

In conclusion, we expect that, by (2.7), in the limit $N \rightarrow \infty$, $f_1^N \rightarrow f$ and $f_2^N \rightarrow f^{\otimes 2}$, where f solves

$$\begin{aligned} \partial_t \int f(z_1) \varphi(z_1) dz_1 &= \int f(z_1) v_1 \cdot \nabla_{x_1} \varphi(z_1) dz_1 - \int f(z_1) \varphi(z_1) dz_1 \\ &\quad + \int f(z_1) f(z_2) \varphi(x_1, v_2) K(M_\rho(B_{|x_1-x_2|}(x_1))) dz_1 dz_2 \end{aligned}$$

and $\rho(x, t) = \int f(x, v, t) dv$. This is the weak form of the equation

$$(2.10) \quad (\partial_t + v \cdot \nabla_x) f(x, v, t) \\ = -f(x, v, t) + \rho(x, t) \int K(M_\rho(B_{|x-y|}(x))) f(y, v, t) dy.$$

We remark that existence and uniqueness of global solutions in $L^1(\mathbb{R}^{2d})$ for the kinetic equation (2.10) can be proved by using a standard Banach fixed-point argument.

Once f is known, we can construct the one-particle nonlinear process given by the generator

$$L_1^{(1)} \phi(x, v) \\ = (v \cdot \nabla_x - 1) \phi(x, v) + \int f(y, w) \phi(x, w) K(M_\rho(B_{|x-y|}(x))) dy dw.$$

We also introduce the N -particle process given by N independent copies of the above process. Its generator is

$$(2.11) \quad L_N^{(1)} \Phi(Z_N) = V_N \cdot \nabla_{X_N} \Phi(Z_N) \\ + \sum_{i=1}^N \left[\int \Phi(X_N, V_N^i(w_i)) K(M_\rho(B_{|x_i-y_i|}(x_i))) f(y_i, w_i) dy_i dw_i - \Phi(X_N, V_N) \right].$$

2.2. Motivations and main result

This work aims to prove propagation of chaos for the N -particle process described by (2.5). Propagation of chaos consists in preparing a system of N particles with initial configurations i.i.d with a given law f_0 , showing that, considering any group of fixed s particles between the N ones, this independence (chaos) is also recovered for future times for the fixed s -group when $N \rightarrow \infty$. This is expressed mathematically by saying that the s -particle marginal $f_s^N(t)$ introduced in (2.8) approximates $f^{\otimes s}(t)$ for positive times, where $f(t)$ is the solution with initial datum f_0 of the limit equation (2.10).

As mentioned in the introduction, the propagation of chaos result for (2.5) was already obtained in [13] using hierarchical techniques. Indeed, the BBGKY hierarchies are a powerful approach but their structure is such that the equation for the s -marginal depends on the contributions given by different integral terms each of which involving only a single $(s+r)$ -marginal for $r = 1, 2, \dots$. In this case, the non-binary nature of the topological interaction does not allow deriving this hierarchical structure unless the interaction function K is real analytic and therefore expandable in series, which is exactly the assumption made in [13].

The reason for this work is to provide a different derivation of the limit kinetic equation, using the classic probabilistic coupling technique. In general, given two stochastic processes X and Y , a coupling is a realization of a new process on a product probability space that has as marginal distributions those of X and Y . This approach brings a more natural and general proof, avoiding the analyticity assumption on K .

THEOREM 1. *Let $f \in C([0, T]; L^1(\mathbb{R}^{2d}))$ be the solution of the limit equation (2.10) with initial datum $f_0 \in L^1(\mathbb{R}^{2d})$. Assume that the interaction function K is Lipschitz-continuous and consider the N -particle dynamics such that*

$$\mathcal{W}_N(0) = f_0^{\otimes N}.$$

If f_s^N denotes the s -marginal as defined in (2.8), for $t \in [0, T]$ and $s \in \{1, \dots, N\}$, it holds that

$$(2.12) \quad \|f_s^N(t) - f^{\otimes s}(t)\|_{L^1(\mathbb{R}^{2ds})} \leq s \frac{e^{C_K T}}{\sqrt{N-1}},$$

where C_K is a constant depending only on the Lipschitz constant of K .

The topological character of the interaction brings us naturally to work with norms of strong type and in particular with the L_1 /total variation distance (see also [3] where a distance similar to the total variation has been used to prove the validity of the mean-field limit for a deterministic Cucker–Smale model with topological interactions introduced in [17]).

Indeed, given two measures ρ_1 and ρ_2 , from (2.9), we have

$$|M_{\rho_1}(B_r(x)) - M_{\rho_2}(B_r(x))| \leq \|\rho_1 - \rho_2\|_{TV}$$

where, given (X, \mathcal{A}) a measurable space and two measures μ and ν over X , the total variation distance is defined as

$$\|\mu - \nu\|_{TV} = \sup_{A \in \mathcal{A}} |\mu(A) - \nu(A)|.$$

In the present work, we use the equivalence between the L^1 distance and the total variation for regular measures and the characterization of the total variation distance given by the Wasserstein distance

$$\|\mu - \nu\|_{TV} = \inf_{\pi \in \mathcal{C}(\mu, \nu)} \int_{X \times X} d(x, y) d\pi(x, y),$$

where $\mathcal{C}(\mu, \nu)$ is the set of all couplings, i.e., measures on the product space with marginals, respectively, μ and ν in the first and second variables, and $d(a, b) = 1 - \delta_{a,b}$ is the discrete distance (see [25]).

3. PROOF OF THE RESULT

3.1. Coupling and strategy of the proof

We introduce, as a coupling between (2.5) and (2.11), the process $t \rightarrow (Z_N(t); \Sigma_N(t))$ on the product space $\mathbb{R}^{2dN} \times \mathbb{R}^{2dN}$, where $\Sigma_N(t) = (Y_N(t), W_N(t))$. The generator of the new process is

$$Q_N = Q_0 + \tilde{Q}_N,$$

where

$$(3.1) \quad Q_0 \Phi(Z_N; \Sigma_N) = (V_N \cdot \nabla_{X_N} + W_N \cdot \nabla_{Y_N}) \Phi(Z_N; \Sigma_N)$$

is the free-stream operator, while

$$(3.2a) \quad \tilde{Q}_N \Phi(Z_N; \Sigma_N) = \sum_{i=1}^N \sum_{j \neq i} \lambda_{i,j} [\Phi(X_N, V_N^i(v_j); Y_N, W_N^{(i)}(w_j)) - \Phi(Z_N; \Sigma_N)]$$

$$(3.2b) \quad + \sum_{i=1}^N \sum_{j \neq i} [\pi_{i,j}^N(X_N) - \lambda_{i,j}] [\Phi(X_N, V_N^i(v_j); \Sigma_N) - \Phi(Z_N; \Sigma_N)]$$

$$(3.2c) \quad + \sum_{i=1}^N \sum_{j \neq i} [\pi^\rho(y_i, y_j) - \lambda_{i,j}] [\Phi(Z_N; Y_N, W_N^{(i)}(w_j)) - \Phi(Z_N; \Sigma_N)]$$

$$(3.2d) \quad + \sum_{i=1}^N \int du \mathcal{E}_i^N(u) [\Phi(Z_N; Y_N, W_N^{(i)}(u)) - \Phi(Z_N; \Sigma_N)]$$

tends to penalize the discrepancies that can occur over time between Z_N and Σ_N .

Indeed, in (3.2a), the process jumps jointly on both variables with a rate given by

$$(3.3) \quad \lambda_{i,j}(X_N; y_i, y_j) := \min \{ \pi_{i,j}^N(X_N), \pi^\rho(y_i, y_j) \},$$

where

$$(3.4) \quad \pi^\rho(y_i, y_j) := \alpha_N K(M_\rho(B_{|y_i - y_j|}(y_i))).$$

In (3.2b) and (3.2c), the jumps occur only for one of the pair, with a transition probability given by the error between $\lambda_{i,j}$ and π^N or π^ρ . Finally, in (3.2d),

$$\mathcal{E}_i^N(u) = \int K(M_\rho(B_{|y_i - y|}(y_i))) f(y, u) dy - \sum_{j \neq i} \pi^\rho(y_i, y_j) \delta(u - w_j)$$

is the last error due to the approximation of the limit kinetic equation by the N -particle

dynamics with transition probabilities given by π^ρ and will be treated using the law of large numbers.

We remark that since $\int K(x) dx = 1$, formally, we have

$$\begin{aligned} \int K(M_\rho(B_{|x-y|}(x)))\rho(y) dy &= \int_0^{+\infty} dr K(M_\rho(B_r(x))) \int_{|x-y|=r} \rho(y) d\mathcal{H}^{n-1}(dy) \\ &= \int_0^{+\infty} dr K(M_\rho(B_r(x))) \frac{d}{dr} [M_\rho(B_r(x))] \\ &= \int K(x) dx = 1. \end{aligned}$$

This is generally true for $\rho \in L^1(\mathbb{R}^d)$ and it is a consequence of the coarea formula (see [14, Thm. 3.12, p. 140]).

From the previous formula, it follows that Q_N is a coupling of the two previously described processes; i.e., we recover, considering test functions depending only on Z_N and Σ_N , respectively, the two processes as the two marginals.

We want to prove that f and f_1^N (defined as in (2.8)) agree asymptotically in the limit $N \rightarrow +\infty$. To do this, we consider $R^N(t) = R^N(Z_N, \Sigma_N; t)$ the law at time t for the coupled process. As initial distribution at time 0, we assume

$$(3.5) \quad R^N(0) = f_0^{\otimes N}(Z_N)\delta(Z_N - \Sigma_N).$$

Let $D_N(t)$ be the average fraction of particles having different positions or velocities, i.e., using the symmetry of the law,

$$(3.6) \quad D_N(t) = \int dR^N(t) \frac{1}{N} \sum_{i=1}^N d(z_i, \sigma_i) = \int dR^N(t) d(z_1, \sigma_1),$$

where $z_i = (x_i, v_i)$, $\sigma_i = (y_i, w_i)$, and $d(a, b) = 1 - \delta_{a,b}$ is the discrete distance.

The aim is to show that $D_N(t) \rightarrow 0$. This means the following: initially, the coupled system has all the pairs of particles overlapping. The dynamics creates discrepancies and the average number of separated pairs is exactly D_N which is also the total variation distance ($L^1(x, v)$ in our case) between f_1^N and f .

Notice that the convergence of the s -marginals f_s^N toward $f^{\otimes s}$ claimed in (2.12) is easily recovered by the fact that

$$\begin{aligned} \|f_s^N(t) - f^{\otimes s}(t)\|_{TV} &\leq \int (Z_s, \Sigma_s) dR^N(Z_N, \Sigma_N; t) \\ &\leq \sum_{i=1}^s \int d(z_i, \sigma_i) dR^N(Z_N, \Sigma_N; t) = sD_N(t) \end{aligned}$$

where $\delta(a, b)$ denotes the discrete distance on the space $\mathbb{R}^{2ds} \times \mathbb{R}^{2ds}$.

3.2. Convergence estimates

Let S_t^N be the semigroup defined by the free-stream generator Q_0 in (3.1). To estimate $D_N(t)$, we apply the Duhamel formula in (3.6), and we get

$$(3.7) \quad \int dR^N(t)d(z_1, \sigma_1) = \int dR^N(0)d(S_t^N(z_1, \sigma_1)) \\ + \int_0^t d\tau \int dR^N(\tau) \tilde{Q}_N d(S_{t-\tau}^N(z_1, \sigma_1)),$$

where \tilde{Q}_N is defined in (3.2).

The first term in (3.7) is negligible: indeed, from (3.5), we have

$$\int dR^N(0)d(S_t^N(z_1, \sigma_1)) = \int df_0^{\otimes N}(Z_N)d(S_t^N(z_1, z_1)) \equiv 0.$$

Concerning the second term in (3.7), we define

$$\bar{z}_1 = (x_1 + v_1(t - \tau), v_1), \quad \bar{z}_1^{(j)} = (x_1 + v_1(t - \tau), v_j),$$

and $\bar{X}_N = (x_1 + v_1(t - \tau), \dots, x_N + v_N(t - \tau))$, similarly, for $\bar{\sigma}$, $\bar{\sigma}^{(j)}$, and \bar{Y}_N .

By (3.2), we get

$$\int dR^N(\tau) \tilde{Q}_N d(S_{t-\tau}^N(z_1, \sigma_1)) = A_1(\tau) + A_2(\tau) + A_3(\tau),$$

where

$$A_1(\tau) = \sum_{j \neq 1} \int dR^N(\tau) \lambda_{1,j}(\bar{X}_N; \bar{y}_1, \bar{y}_j) [d(\bar{z}_1^{(j)}; \bar{\sigma}_1^{(j)}) - d(\bar{z}_1; \bar{\sigma}_1)]$$

is due to the term of the generator \tilde{Q}_N where the velocities of the particles jump simultaneously,

$$A_2(\tau) = \sum_{j \neq 1} \int dR^N(\tau) (\pi_{1,j}^N(\bar{X}_N) - \lambda_{1,j}) [d(\bar{z}_1^{(j)}; \bar{\sigma}_1) - d(\bar{z}_1; \bar{\sigma}_1)] \\ + \sum_{j \neq 1} \int dR^N(\tau) (\pi^\rho(\bar{y}_1, \bar{y}_j) - \lambda_{1,j}) [d(\bar{z}_1; \bar{\sigma}_1^{(j)}) - d(\bar{z}_1; \bar{\sigma}_1)]$$

is due to the terms of the generator where only one of the two coupled processes jump, and

$$A_3(\tau) = \int dR^N(\tau) \int du \bar{\mathcal{E}}_1^N(u) [d(\bar{z}_1; \bar{\sigma}_1^{(u)}) - d(\bar{z}_1; \bar{\sigma}_1)]$$

is due to the remainder term. Here, $\bar{\mathcal{E}}_1^N(u)$ is $\mathcal{E}_1^N(u)$ evaluated along the moving frame of the free transport.

Here, we have used that $d(z_1, \sigma_1)$ depends only on the configurations of the first particle; hence, the only non-zero contribution in the sum over i is given for $i = 1$.

Concerning $A_1(\tau)$, it follows from (2.3) and (2.4) that

$$|e_K(N)| \leq \frac{\text{Lip}(K)}{N-1}$$

and that, for $N > 2\text{Lip}(K) + 1$,

$$\alpha_N \leq \frac{4e^{\frac{\text{Lip}(K)}{N-1}}}{N-1},$$

using the inequality $1/(1-x) \leq 4e^x$ for $x \in (0, 1/2)$. Therefore, from (3.3), we get

$$\lambda_{1,j} \leq \alpha_N \|K\|_\infty \leq \frac{4\sqrt{e}\text{Lip}(K)}{N-1}.$$

By the symmetry of R^N and denoting $C_K := 8\sqrt{e}\text{Lip}(K)$,

$$(3.8) \quad A_1(\tau) \leq \frac{C_K}{2(N-1)} \sum_{j \neq 1} \int dR^N(\tau) [d(z_j, \sigma_j) + d(z_1, \sigma_1)] \leq C_K D_N(\tau)$$

since $d(\bar{z}_1^{(j)}; \bar{\sigma}_1^{(j)}) \leq d(z_j, \sigma_j) + d(z_1, \sigma_1)$. Indeed, the right-hand side is vanishing iff $z_1 = \sigma_1$ and $z_j = \sigma_j$, and, in this case, also the left-hand side is clearly vanishing.

We now give a bound on $A_2(\tau)$. Since $\lambda_{1,j}$ is the minimum between $\pi_{1,j}^N$ and $\pi_{i,j}^\rho$, we have

$$(3.9) \quad |A_2(\tau)| \leq \sum_{j \neq 1} \int dR^N(\tau) |\pi_{1,j}^N(\bar{X}_N) - \pi_{1,j}^\rho(\bar{y}_1, \bar{y}_j)|.$$

From (2.2) and (3.4),

$$|\pi_{1,j}^N(\bar{X}_N) - \pi_{1,j}^\rho(\bar{y}_1, \bar{y}_j)| \leq \alpha_N \text{Lip}(K) |M_{\bar{X}_N}(\bar{B}_{1,j}^x) - M_\rho(\bar{B}_{1,j}^y)|,$$

where we are using the shorthand notation

$$\bar{B}_{1,j}^x = B_{|\bar{x}_1 - \bar{x}_j|}(\bar{x}_1) \quad \text{and} \quad \bar{B}_{1,j}^y = B_{|\bar{y}_1 - \bar{y}_j|}(\bar{y}_1).$$

By the triangular inequality,

$$\begin{aligned} |M_{\bar{X}_N}(\bar{B}_{1,j}^x) - M_\rho(\bar{B}_{1,j}^y)| &\leq |M_{\bar{X}_N}(\bar{B}_{1,j}^x) - M_{\bar{X}_N}(\bar{B}_{1,j}^y)| \\ &\quad + |M_{\bar{X}_N}(\bar{B}_{1,j}^y) - M_{\bar{Y}_N}(\bar{B}_{1,j}^y)| \\ &\quad + |M_{\bar{Y}_N}(\bar{B}_{1,j}^y) - M_\rho(\bar{B}_{1,j}^y)|. \end{aligned}$$

Hence, we divide the estimate (3.9), respectively, in three terms:

$$|A_2(\tau)| \leq T_1(\tau) + T_2(\tau) + T_3(\tau).$$

In $T_1(\tau)$, we are considering particles with spatial configuration given by X_N , and we want to estimate the discrepancy of the configuration over two different balls $\bar{B}_{1,j}^x$ and $\bar{B}_{1,j}^y$. Since $\bar{B}_{1,j}^x = \bar{B}_{1,j}^y$ iff $z_1 = \sigma_1$ and $z_j = \sigma_j$, using that $M_{\bar{X}_N} \in [0, 1]$, we have

$$|M_{\bar{X}_N}(\bar{B}_{1,j}^x) - M_{\bar{X}_N}(\bar{B}_{1,j}^y)| \leq d(z_1, \sigma_1) + d(z_j, \sigma_j).$$

Therefore, by the symmetry of R^N ,

$$T_1(\tau) \leq \alpha_N \text{Lip}(K) \sum_{j \neq 1} \int dR^N(\tau) [d(z_1, \sigma_1) + d(z_j, \sigma_j)] \leq C_K D_N(\tau).$$

Regarding $T_2(\tau)$, we are considering the discrepancy of two different configurations over the same ball $\bar{B}_{1,j}^y$. Since

$$|M_{\bar{X}_N}(\bar{B}_{1,j}^y) - M_{\bar{Y}_N}(\bar{B}_{1,j}^y)| \leq \frac{1}{N} \sum_{i=1}^N d(z_i, \sigma_i),$$

using again the symmetry of the law, we get

$$T_2(\tau) \leq \alpha_N \text{Lip}(K) \sum_{j \neq 1} \int dR^N(\tau) d(z_1, \sigma_1) \leq C_K D_N(\tau).$$

The last estimate on $T_3(\tau)$ is a consequence of the law of large numbers. After a change of variable, using the symmetry of the law R^N and the fact that this last term depends only on the Y_N configuration, we have that

$$T_3(\tau) = \alpha_N \text{Lip}(K) \sum_{j \neq 1} \int d\rho^{\otimes N}(\tau) |M_{Y_N}(B_{1,j}^y) - M_\rho(B_{1,j}^y)|,$$

where $B_{1,j}^y = B_{|y_1 - y_j|}(y_1)$. By Cauchy-Schwartz,

$$\begin{aligned} & \left| \int d\rho^{\otimes N}(\tau) |M_{Y_N}(B_{1,j}^y) - M_\rho(B_{1,j}^y)| \right|^2 \\ & \leq \int d\rho^{\otimes N}(\tau) \left| \frac{1}{N-1} \sum_{h \neq 1} [\mathcal{X}_{B_{1,j}^y}(y_h) - M_\rho(B_{1,j}^y)] \right|^2 \\ & \leq \sum_{h_1, h_2 \neq 1} \int \frac{d\rho^{\otimes N}(\tau)}{(N-1)^2} [\mathcal{X}_{B_{1,j}^y}(y_{h_1}) - M_\rho(B_{1,j}^y)] [\mathcal{X}_{B_{1,j}^y}(y_{h_2}) - M_\rho(B_{1,j}^y)]. \end{aligned}$$

Thanks to the independence of the limit process, we get that the only non-zero contributions are given when $h_1 = h_2$, and this happens only for $N - 1$ terms. Hence,

$$T_3(\tau) \leq \frac{C_K}{\sqrt{N-1}}.$$

Collecting the estimates on T_1 , T_2 , and T_3 , we obtain that

$$(3.10) \quad A_2(\tau) \leq C_K \left(D_N(\tau) + \frac{1}{\sqrt{N-1}} \right).$$

We conclude the proof estimating $A_3(\tau)$. Since this term depends only on the independent Y_N configuration,

$$\begin{aligned} |A_3(\tau)| &\leq \int \frac{df^{\otimes N}(\tau)}{N-1} \sum_{j \neq 1} \left| \int K(M_\rho(B_{|\bar{y}_1-y|}(\bar{y}_1))) d\rho(y) - K(M_\rho(\bar{B}_{1,j}^y)) \right| \\ &\quad + \frac{1}{N-1} \sum_{j \neq 1} \int df^{\otimes N}(\tau) (1 - (N-1)\alpha_N) K(M_\rho(\bar{B}_{1,j}^y)), \end{aligned}$$

where we added and subtracted the term $\sum_j K(M_\rho(\bar{B}_{1,j}^y))/(N-1)$.

Applying again the law of large numbers on the first term, estimating the second term thanks to (2.3), and using that

$$1 - (N-1)\alpha_N = \frac{e_K(N)}{1 - e_K(N)} \leq \frac{C_K}{N-1},$$

we arrive at

$$(3.11) \quad |A_3(\tau)| \leq \frac{C_K}{\sqrt{N-1}}.$$

Collecting the estimates in (3.8), (3.10), and (3.11) and using Gronwall's lemma, we conclude the proof of the theorem.

ACKNOWLEDGMENTS. – P. Degond holds a visiting professor association with the Department of Mathematics, Imperial College London, UK.

REFERENCES

- [1] M. BALLERINI ET AL., [Interaction ruling animal collective behavior depends on topological rather than metric distance: Evidence from a field study](#). *Proc. Natl. Acad. Sci. USA* **105** (2008), 1232–1237.
- [2] S. BAZAZI ET AL., [Collective motion and cannibalism in locust migratory bands](#). *Curr. Biol.* **18** (2008), 735–739.

- [3] D. BENEDETTO – E. CAGLIOTI – S. ROSSI, [Mean-field limit for particle systems with topological interactions](#). *Math. Mech. Complex Syst.* **9** (2021), no. 4, 423–440. Zbl 1499.35585 MR 4393188
- [4] E. BERTIN – M. DROZ – G. GRÉGOIRE, [Boltzmann and hydrodynamic description for self-propelled particles](#). *Phys. Rev. E* **74** (2006), no. 2, article no. 022101.
- [5] A. BLANCHET – P. DEGOND, [Topological interactions in a Boltzmann-type framework](#). *J. Stat. Phys.* **163** (2016), no. 1, 41–60. Zbl 1352.92182 MR 3472093
- [6] A. BLANCHET – P. DEGOND, [Kinetic models for topological nearest-neighbor interactions](#). *J. Stat. Phys.* **169** (2017), no. 5, 929–950. Zbl 1380.82032 MR 3719633
- [7] N. W. BODE – D. W. FRANKS – A. J. WOOD, [Limited interactions in flocks: relating model simulations to empirical data](#). *J. R. Soc. Interface* **8** (2011), 301–304.
- [8] M. CAMPERI – A. CAVAGNA – I. GIARDINA – G. PARISI – E. SILVESTRI, [Spatially balanced topological interaction grants optimal cohesion in flocking models](#). *Interface Focus* **2** (2012), 715–725.
- [9] A. CAVAGNA ET AL., [From empirical data to inter-individual interactions: unveiling the rules of collective animal behavior](#). *Math. Models Methods Appl. Sci.* **20** (2010), 1491–1510. Zbl 1197.92053 MR 3090590
- [10] L.-P. CHAINTRON – A. DIEZ, [Propagation of chaos: A review of models, methods and applications. I. Models and methods](#). *Kinet. Relat. Models* **15** (2022), no. 6, 895–1015. Zbl 1496.82016 MR 4489768
- [11] L.-P. CHAINTRON – A. DIEZ, [Propagation of chaos: A review of models, methods and applications. II. Applications](#). *Kinet. Relat. Models* **15** (2022), no. 6, 1017–1173. Zbl 1496.82017 MR 4489769
- [12] F. CUCKER – S. SMALE, [Emergent behavior in flocks](#). *IEEE Trans. Automat. Control* **52** (2007), no. 5, 852–862. Zbl 1366.91116 MR 2324245
- [13] P. DEGOND – M. PULVIRENTI, [Propagation of chaos for topological interactions](#). *Ann. Appl. Probab.* **29** (2019), no. 4, 2594–2612. Zbl 1456.60254 MR 3984258
- [14] L. C. EVANS – R. F. GARIÉPY, [Measure theory and fine properties of functions](#). Revised edn., Textb. Math., CRC Press, Boca Raton, FL, 2015. Zbl 1310.28001 MR 3409135
- [15] F. GINELLI – H. CHATÉ, [Relevance of metric-free interactions in flocking phenomena](#). *Phys Rev Lett* **105** (2010), article no. 168103.
- [16] R. GINIŪNAITĖ – R. E. BAKER – P. M. KULESA – P. K. MAINI, [Modelling collective cell migration: neural crest as a model paradigm](#). *J. Math. Biol.* **80** (2020), no. 1-2, 481–504. Zbl 1432.92016 MR 4062827
- [17] J. HASKOVEC, [Flocking dynamics and mean-field limit in the Cucker–Smale-type model with topological interactions](#). *Phys. D* **261** (2013), 42–51. Zbl 1310.92062 MR 3144007

- [18] U. LOPEZ – J. GAUTRAIS – I. D. COUZIN – G. THERAULAZ, [From behavioural analyses to models of collective motion in fish schools](#). *Interface Focus* **2** (2012), 693–707.
- [19] S. MARTIN, [Multi-agent flocking under topological interactions](#). *Systems Control Lett.* **69** (2014), 53–61. Zbl [1288.93005](#) MR [3212821](#)
- [20] T. NIZATO – H. MURAKAMI – Y. P. GUNJI, [Emergence of the scale-invariant proportion in a flock from the metric-topological interaction](#). *Biosystems* **119** (2014), 62–68.
- [21] Y. SHANG – R. BOUFFANAIS, [Consensus reaching in swarms ruled by a hybrid metric-topological distance](#). *Eur. Phys. J. B* **87** (2014), no. 12, article no. 294. MR [3284898](#)
- [22] Y. SHANG – R. BOUFFANAIS, [Influence of the number of topologically interacting neighbors on swarm dynamics](#). *Sci. Rep.* **4** (2014), article no. 4184.
- [23] T. VICSEK – A. CZIRÓK – E. BEN-JACOB – I. COHEN – O. SHOCHET, [Novel type of phase transition in a system of self-driven particles](#). *Phys. Rev. Lett.* **75** (1995), no. 6, 1226–1229. MR [3363421](#)
- [24] T. VICSEK – A. ZAFEIRIS, [Collective motion](#). *Phys. Rep.* **517** (2012), 71–140.
- [25] C. VILLANI, [Topics in optimal transportation](#). Grad. Stud. Math. 58, American Mathematical Society, Providence, RI, 2003. Zbl [1106.90001](#) MR [1964483](#)
- [26] L. WANG – G. CHEN, [Synchronization of multi-agent systems with metric-topological interactions](#). *Chaos* **26** (2016), no. 9, article no. 094809. Zbl [1382.93005](#) MR [3518988](#)

Received 17 December 2022,
and in revised form 27 March 2023

Pierre Degond
Institut de Mathématiques de Toulouse (UMR 5219), Université de Toulouse, CNRS UPS
31062 Toulouse Cedex 9, France;
pierre.degond@math.univ-toulouse.fr

Mario Pulvirenti
Dipartimento di Matematica, Università di Roma ‘La Sapienza’
Piazzale Aldo Moro 2, 00185 Roma, Italy;
pulviren@mat.uniroma1.it

Stefano Rossi
Dipartimento di Matematica, Università di Roma ‘La Sapienza’
Piazzale Aldo Moro 2, 00185 Roma, Italy;
stef.rossi@uniroma1.it