

Binormal Deformation and Bimicrolocalization

By

Kiyoshi TAKEUCHI*

Contents

- §1. Introduction
- §2. Geometry
 - 2.1. Preliminary results on normal bundles
 - 2.2. Binormal deformation
 - 2.3. Binormal cones
- §3. Bispecialization Functor
 - 3.1. Functorial properties of the functor v_{NM}
 - 3.2. Structure of the functor v_{NM}
- §4. Bimicrolocalization Functors
 - 4.1. Functors $v\mu_{NM}$ and μ_{NM}
 - 4.2. Functorial properties of $v\mu_{NM}$ and μ_{NM}
 - 4.3. Various generalizations of the functor μ_{hom}
- §5. Applications
 - 5.1. Second microfunctions associated to a submanifold
 - 5.2. Second microfunctions associated to a foliation
 - 5.3. Second microdifferential operators

§1. Introduction

The specialization functor is an important tool in algebraic geometry as well as in algebraic analysis. In the real case, it associates to a sheaf F on a real manifold X and to a submanifold M of X , a sheaf (i.e. an object of the derived category of sheaves) $v_M(F)$ on the normal bundle $T_M X$ which describes the “boundary values” of F along M . Its Fourier transform is the sheaf $\mu_M(F)$ of Sato’s microlocalization of F along M .

Communicated by M. Kashiwara, March 29, 1995.

1991 Mathematics Subject Classification(s): 14F05, 32C38, 35A27, 58G07.

* Department of Mathematics, Hiroshima University, 1–3–1, Kagamiyama, Higashi-Hiroshima-shi, 739, Hiroshima, Japan.

Now consider a triplet of real manifolds $N \subset M \subset X$. In order to analyze the sheaf $v_M(F)$ along the submanifold $N \times_M T_M X$ of $T_M X$, a natural idea is to introduce the second specialization $v_{(N \times_M T_M X)}(v_M(F))$. Taking the Fourier transform of this object, we get the theory of the second microlocalization introduced by M. Kashiwara [4] in 1972 and developed by many authors, in particular by Kashiwara-Kawai [7] and Laurent [15]. However, the sheaf one obtains by this method is very “big” and the direct image of this sheaf on the bundle $N \times_M T_M^* X$ is not the restriction to $N \times_M T_M^* X$ of the sheaf $\mu_M(F)$. This leads to many difficulties.

In this paper, we propose a new approach to the second specialization by considering the inverse image \tilde{M} of M in \tilde{X}_N , the normal deformation of X along N and defining the binormal deformation of X along (M, N) as the deformation of \tilde{X}_N along \tilde{M} . Then, by paraphrasing the construction of the functor $v_M(\cdot)$, we construct the functor $v_{NM}(\cdot)$ which associates to a sheaf F on X a sheaf on $T_N M \times_M T_M X$. By taking the Fourier transform of this sheaf, we obtain the sheaf $v\mu_{NM}(F)$ on $T_N M \times_M T_M^* X$ and the sheaf $\mu_{NM}(F)$ on $T_N^* M \times_M T_M^* X$. We study the functorial properties of these new sheaves and we show that they enjoy the desired properties. In particular the direct image of $\mu_{NM}(F)$ on $N \times_M T_M^* X$ coincides with the restriction of $\mu_M(F)$ to this set.

It would then be possible to develop a whole theory of the functor $\mu_{NM}(\cdot)$, in the lines of Sato-Kawai-Kashiwara [19] or Kashiwara-Schapira [11], but this will not be done here. Let us only mention that the author ([24] and [25]) has applied this theory to the study of boundary value problems in higher codimension.

Finally, notice that the idea of second normal deformation already appeared in Sabbah [18] (this idea has been developed by Bost [1] to obtain a proof of Atiyah-Singer theorem) and a construction of a sheaf of second microfunctions, following the lines of [19], based on a geometrical idea very closed to ours, had been proposed by Kataoka-Tose in [13]. We thank N. Tose to have drawn our attention on this (unpublished) paper. Another construction (based on integral formulas) of the same sheaf of second microfunctions was performed later in [14]. Also notice that Sjöstrand [23] introduced a different language and with different methods, various spaces of second microfunctions. On the other hand, “simultaneous specialization” is a slightly different geometrical construction introduced by Delort [2].

The main results of this paper were announced in [22]. The idea of bispecialization is essentially due to P. Schapira. The author and he tried to realize his idea to construct a framework of higher codimensional boundary value problems treated in [24] and [25]. In this process, the relation with the theory of second microlocal analysis was found and developed by the author. The present article gives complete proofs of this theory including the

developments after the introduction of bispecialization. The author would like to express his profound gratitude to Prof. Schapira for giving him many valuable suggestions and technical aids throughout this work. This theory were not realized without his constant encouragements. He is also very grateful to Prof. Honda and Mr. Watanabe for careful reading of some parts of this manuscript.

§2. Geometry

2.1. Preliminary results on normal bundles

Let $X \supset M \supset N$ be a sequence of C^∞ -manifolds. Let us denote it by (X, M, N) and call it a triplet of manifolds. We sometimes identify M with the zero-section of the normal bundle $T_M X$, and N is considered as a submanifold of $T_M X$. We recall some elementary geometrical results on normal bundles.

Proposition 2.1. *Let (X, M, N) be a triplet of manifolds. There are canonical isomorphisms of normal bundles:*

$$(2.1) \quad \begin{array}{ccc} & T_{(T_N M)}(T_N X) & \\ & \uparrow \mathbf{A} & \\ T_N M \times_M T_M X & \xrightarrow{\mathbf{B}} & T_N(T_M X), \\ & \uparrow \mathbf{C} & \\ & T_{(N \times_M T_M X)}(T_M X). & \end{array}$$

where N is considered as a submanifold of the zero section M of $T_M X$ and $T_N M$ is imbedded into $T_N X$ by the tangent map associated to the inclusion $(M, N) \rightarrow (X, N)$.

Proof. We will construct the canonical morphisms **A**, **B** and **C**. Then, using local coordinate systems, it will be obvious that these morphisms are isomorphisms. For this purpose, the simple lemma below is crucial.

Lemma 2.2. *Let $\tau: E \rightarrow Z, (x; \xi) \mapsto x$ be a vector bundle over a C^∞ -manifold Z .*

(i) *Let $\left(x, \xi; x^* \frac{\partial}{\partial x} + \xi^* \frac{\partial}{\partial \xi}\right)$ be a linear coordinate system on TE . Then, there exists a canonical injective morphism of vector bundles over Z :*

$$(2.2) \quad \psi_\tau: E \longrightarrow Z \times_E TE, \quad (x, \xi) \longmapsto \left(x, 0; \xi \frac{\partial}{\partial \xi}\right).$$

(ii) Let E_1 and E_2 be two copies of E and $((x, \xi_1), (x, \xi_2))$ be a linear coordinate system on $E_1 \times_Z E_2$. Then there exists a canonical injective morphism of vector bundles over E_1 :

$$(2.3) \quad \phi_\tau: E_1 \times_Z E_2 \longrightarrow TE_1, \quad ((x, \xi_1), (x, \xi_2)) \longmapsto \left(x, \xi_1; \xi_2 \frac{\partial}{\partial \xi_1} \right).$$

Proof.

(i) First consider the exact sequence:

$$0 \longrightarrow TZ \longrightarrow Z \times_E TE \longrightarrow T_Z E \longrightarrow 0.$$

This sequence splits since the projection $E \rightarrow Z$ gives the morphism $TE \rightarrow E \times_Z TZ$, hence the morphism: $Z \times_E TE \rightarrow TZ$. Therefore we get the injection:

$$\psi_\tau: E \simeq T_Z E \longrightarrow Z \times_E TE.$$

(ii) Applying (i) to the vector bundle $E_1 \times_Z E_2 \rightarrow E_1$, we get the injective morphism:

$$E_1 \times_Z E_2 \longrightarrow E_1 \times_{(E_1 \times_Z E_2)} T(E_1 \times_Z E_2) \longrightarrow T(E_1 \times_Z E_2).$$

Next consider the tangent map: $T(E_1 \times_Z E_2) \rightarrow TE_1$ associated to the addition map $E_1 \times_Z E_2 \rightarrow E_1$. The composition of these two morphisms is the desired one:

$$\phi_\tau: E_1 \times_Z E_2 \longrightarrow TE_1.$$

■

Applying Lemma 2.2 (i) to $\tau: T_M X \rightarrow M$, we have an injective morphism of vector bundles over M :

$$(2.4) \quad \psi_\tau: T_M X \longrightarrow M \times_{(T_M X)} T(T_M X).$$

Restricting it to N , we get a chain of morphisms on N :

$$(2.5) \quad N \times_M T_M X \xrightarrow{\psi_\tau|_N} N \times_{(T_M X)} T(T_M X) \longrightarrow T_N(T_M X).$$

The composition of the above morphisms gives a morphism:

$$(2.6) \quad N \times_M T_M X \longrightarrow T_N(T_M X).$$

On the other hand, we have a tangent map associated to the zero section embedding $(M, N) \rightarrow (T_M X, N)$, that is:

$$(2.7) \quad T_N M \longrightarrow T_N(T_M X).$$

By adding the morphisms (2.6) and (2.7), we get the morphism **B**:

$$(2.8) \quad \mathbf{B}: T_N M \times_M T_M X \longrightarrow T_N(T_M X).$$

To construct the morphism **A**, first notice the commutative diagram:

$$(2.9) \quad \begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & TN & \longrightarrow & TN & \longrightarrow & 0 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & N \times_M TM & \longrightarrow & N \times_X TX & \longrightarrow & N \times_M T_M X \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & T_N M & \longrightarrow & T_N X & \longrightarrow & N \times_M T_M X \longrightarrow 0, \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

where all columns and the first and the second rows are exact. Hence, by virtue of the nine Lemma, the third row is also exact. That is, we have a canonical isomorphism of vector bundles over N :

$$(2.10) \quad T_N X / T_N M \xrightarrow{\sim} N \times_M T_M X,$$

in which the left hand side is a quotient bundle over N .

If we apply Lemma 2.2 (ii) to $\tau_1: T_N M \rightarrow N$ and $\tau_2: T_N X \rightarrow N$, we get:

$$(2.11) \quad \phi_{\tau_1}: T_N M \times_N T_N M \longrightarrow T(T_N M)$$

and

$$(2.12) \quad \phi_{\tau_2}: T_N X \times_N T_N X \longrightarrow T(T_N X).$$

Since ϕ_{τ_2} is a morphism of vector bundles over $T_N X$, it can be restricted to $T_N M \subset T_N X$, and we denote it by $\phi_{\tau_2}|_{T_N M}$. Now we have a commutative diagram:

$$(2.13) \quad \begin{array}{ccc} T_N M \times_N T_N M & \xrightarrow{\phi_{\tau_1}} & T(T_N M) \\ \mathbf{D} \downarrow & & \downarrow \mathbf{E} \\ T_N M \times_N T_N X & \xrightarrow{\phi_{\tau_2}|_{T_N M}} & T_N M \times_{(T_N X)} T(T_N X), \end{array}$$

where the morphism **D** is induced by the injection $T_N M \rightarrow T_N X$, and the morphism **E** is the tangent map associated to the inclusion $T_N M \hookrightarrow T_N X$. The

commutativity is easily checked by fixing a local coordinate system. Therefore, we can construct a morphism between the quotient bundle of the morphism **D** and that of **E**:

$$(2.14) \quad T_N M \times_N (T_N X / T_N M) \longrightarrow T_{(T_N M)}(T_N X).$$

The left hand side is isomorphic to $T_N M \times_M T_M X$ by (2.10), and we have thus constructed the morphism **A**.

The morphism **C** is the tangent map induced by the projection:

$$(T_M X, N \times_M T_M X) \longrightarrow (M, N),$$

that is,

$$(2.15) \quad T_{(N \times_M T_M X)}(T_M X) \longrightarrow (N \times_M T_M X) \times_N T_N M \simeq T_N M \times_M T_M X.$$

This completes the proof of Proposition 2.1. ■

2.2. Binormal deformation

Let (X, M, N) be a triplet of manifolds. First, recall the construction of the normal deformation of X along N in [11]. We shall denote it by \tilde{X}_N and let $t_1 \in \mathbb{R}$ be the deformation parameter. Then we have the commutative diagram below:

$$(2.16) \quad \begin{array}{ccc} T_N X & \xrightarrow{s_N} & \tilde{X}_N \xleftarrow{j_N} \Omega_N = \{t_1 > 0\} \\ \tau_N \downarrow & & p_N \downarrow \swarrow \tilde{p}_N \\ N & \xrightarrow{i_N} & X. \end{array}$$

Set $\hat{\Omega}_N := \{(x, t_1) \in \tilde{X}_N; t_1 \neq 0\}$ and consider:

$$(2.17) \quad \tilde{M} := \overline{(p_N|_{\hat{\Omega}_N})^{-1} M} \subset \tilde{X}_N,$$

the closure of $(p_N|_{\hat{\Omega}_N})^{-1} M$ in \tilde{X}_N . Then \tilde{M} is an intrinsically defined smooth submanifold of \tilde{X}_N . If we choose a local coordinate system $x = (x', x'', x''')$ of X such that

$$(2.18) \quad \begin{cases} M = \{x' = 0\}, \\ N = \{x' = 0, x'' = 0\}, \end{cases}$$

then $\tilde{M} = \{x' = 0\}$ in \tilde{X}_N . Now we shall define the binormal deformation \tilde{X}_{NM} of X along (M, N) as the normal deformation of \tilde{X}_N along \tilde{M} .

Definition 2.3. One sets:

$$(2.19) \quad \tilde{X}_{NM} := (\tilde{X}_N)_{\tilde{M}}.$$

Define the morphism $p: \tilde{X}_{NM} \rightarrow X$ as the composition of the natural morphisms

$p_{\tilde{M}}: \tilde{X}_{NM} \rightarrow \tilde{X}_N$ and $p_N: \tilde{X}_N \rightarrow X$. Using the local coordinate system in (2.18), this morphism p is described as follows:

$$(2.20) \quad (x', x'', x''', t_1, t_2) \longmapsto (t_2 x', x'', x''', t_1) \longmapsto (t_1 t_2 x', t_1 x'', x'''),$$

where t_2 denotes the deformation parameter of $\tilde{X}_{NM} \rightarrow \tilde{X}_N$. Since $\tilde{M} \cap \{t_1 = 0\} \simeq T_N M \subset T_N X$ in \tilde{X}_N , we have an isomorphism:

$$(2.21) \quad \tilde{X}_{NM} \cap \{t_1 = 0\} \simeq (\tilde{X}_N \cap \{t_1 = 0\})_{(\tilde{M} \cap \{t_1 = 0\})} \simeq (T_N X)_{(T_N M)}$$

and

$$(2.22) \quad \tilde{X}_{NM} \cap \{t_1 = t_2 = 0\} \simeq T_{(T_N M)}(T_N X).$$

By virtue of Proposition 2.1, we can identify it with $T_N M \times_M T_M X$, and there exists a commutative diagram below:

$$(2.23) \quad \begin{array}{ccc} T_N M \times_M T_M X & \xrightarrow{s} \tilde{X}_{NM} & \xleftarrow{j} \Omega := \{t_1 > 0, t_2 > 0\} \\ \tau \downarrow & & \downarrow p \quad \swarrow \tilde{p} \\ N & \longrightarrow & X. \end{array}$$

Remark 2.4. (i) \tilde{M} is isomorphic to the normal deformation \tilde{M}_N of M along N .

(ii) We have another construction of \tilde{X}_{NM} starting from the normal deformation $p_M: \tilde{X}_M \rightarrow X$ with the deformation parameter t_2 . Setting:

$$\tilde{N} := \overline{(p_M|_{\{t_2 \neq 0\}})^{-1} N} \subset \tilde{X}_M,$$

the binormal deformation \tilde{X}_{NM} of X along (M, N) can be defined by:

$$(2.24) \quad \tilde{X}_{NM} := (\tilde{X}_M)_{\tilde{N}}.$$

Of course, it is also possible to construct \tilde{X}_{NM} directly by gluing local charts as in [11].

2.3. Binormal cones

There is a $(\mathbb{R}^+)^2$ -action

$$(2.25) \quad \tilde{X}_{NM} \times (\mathbb{R}^+)^2 \longrightarrow \tilde{X}_{NM}$$

on \tilde{X}_{NM} which is described in the local coordinate system $(x', x'', x''', t_1, t_2)$ associated to the one in (2.18) of X by:

$$(2.26) \quad ((x', x'', x''', t_1, t_2), (c, d)) \longmapsto \left(cdx', cx'', x''', \frac{t_1}{c}, \frac{t_2}{d} \right).$$

It is easily checked that this definition is intrinsic.

Definition 2.5. Let S be a subset of X . We define the binormal cone to S along (M, N) by:

$$(2.27) \quad C_{NM}(S) := (T_N M \times_M T_M X) \cap \overline{\tilde{p}^{-1}(S)}.$$

This is a closed subset of $T_N M \times_M T_M X$.

In this section, we will investigate the properties of $C_{NM}(S)$. The problems are always local, and we shall restrict ourselves to the following situation:

$$(2.28) \quad \begin{cases} X = X'_{x'} \times X''_{x''} \times X'''_{x'''} \\ M = \{0\} \times X''_{x''} \times X'''_{x'''} \\ N = \{0\} \times \{0\} \times X'''_{x'''} \end{cases}$$

where $X'_{x'} = \mathbb{R}^{n'}$, $X''_{x''} = \mathbb{R}^{n''}$ and $X'''_{x'''} = \mathbb{R}^{n'''}$ for some n' , n'' and n''' . Then there exists an associated coordinate system $\left(v' \frac{\partial}{\partial x'}, v'' \frac{\partial}{\partial x''}, x'''\right)$ of

$$(2.29) \quad T_N M \times_M T_M X \simeq \mathbb{R}^{n'}_{v'} \times \mathbb{R}^{n''}_{v''} \times \mathbb{R}^{n'''}_{x'''}$$

Proposition 2.6. Let S be a subset of X .

(i) $C_{NM}(S)$ is a closed biconic subset of $T_N M \times_M T_M X$.

(ii) $(v'_0, v''_0, x'''_0) \in C_{NM}(S) \Leftrightarrow$ There exists a sequence $\{(x'_n, x''_n, x'''_n), c_n, d_n\}$ in $S \times (\mathbb{R}^+)^2$ with $x_n = (x'_n, x''_n, x'''_n)$ such that

$$(2.30) \quad \begin{cases} x_n \longrightarrow (0, 0, x'''_0), \\ c_n d_n x'_n \longrightarrow v'_0, \\ c_n x''_n \longrightarrow v''_0, \\ d_n \longrightarrow +\infty. \end{cases}$$

Proof.

(i) Since the restriction of the $(\mathbb{R}^+)^2$ -action (2.25) to

$$\tilde{X}_{NM} \cap \{t_1 = t_2 = 0\} = T_N M \times_M T_M X$$

coincides with the biconic structure of $T_N M \times_M T_M X$, the set $C_{NM}(S)$ is biconic.

(ii) (A) First assume $(v'_0, v''_0, x'''_0) \in C_{NM}(S)$. Then there exists a sequence $\{(x'_n, x''_n, x'''_n, t_n, s_n)\}$ in $\tilde{p}^{-1}(S)$ which satisfies:

$$(x'_n, x''_n, x'''_n, t_n, s_n) \longrightarrow (v'_0, v''_0, x'''_0, 0, 0).$$

We can easily see that the sequence $\left\{(t_n s_n x'_n, t_n x''_n, x'''_n), \frac{1}{t_n}, \frac{1}{s_n}\right\}$ in $S \times (\mathbb{R}^+)^2$ satisfies the condition (2.30).

(B) Conversely, assume the condition (2.30).

(a) The case when $\{c_n\}$ is not bounded.

Taking a subsequence if necessary, we may assume $c_n \rightarrow +\infty$. Then, the sequence in $\tilde{p}^{-1}(S)$:

$$\left\{ (c_n d_n x'_n, c_n x''_n, x'''_n), \frac{1}{c_n}, \frac{1}{d_n} \right\}$$

converges to $(v'_0, v''_0, x'''_0, 0, 0) \in T_N M \times_M T_M X$. Thus, we have $(v'_0, v''_0, x'''_0) \in C_{NM}(S)$.

(b) The case when $\{c_n\}$ and $\{c_n d_n\}$ are bounded. Because $x_n = (x'_n, x''_n, x'''_n) \rightarrow (0, 0, x'''_0)$,

$$\begin{cases} c_n d_n x'_n \rightarrow 0 \\ c_n x''_n \rightarrow 0 \end{cases} \quad \text{and} \quad \begin{cases} v'_0 = 0 \\ v''_0 = 0. \end{cases}$$

Moreover, taking a subsequence, we may assume

$$|x'_n| < \frac{1}{4^n} \quad \text{and} \quad |x''_n| < \frac{1}{2^n} \quad \text{for } \forall n \in \mathbb{N}.$$

If we define the sequences $\{\varepsilon_n\}$ and $\{\delta_n\}$ by:

$$\varepsilon_n = \frac{n}{2^n}, \quad \delta_n = \frac{1}{2^n},$$

then we have:

$$\left(\left(\frac{x'_n}{\varepsilon_n \delta_n}, \frac{x''_n}{\varepsilon_n}, x'''_n \right), \varepsilon_n, \delta_n \right) \rightarrow ((0, 0, x'''_0), 0, 0) \in C_{NM}(S).$$

(c) The case when $\{c_n\}$ is bounded and $\{c_n d_n\}$ is not bounded. Taking a subsequence, it may be assumed:

$$\begin{cases} |x''_n| < \frac{1}{n^2} \\ \left| \frac{1}{c_n d_n} \right| < \frac{1}{n^2}, \end{cases}$$

because $x''_n \rightarrow 0$. If we set $\varepsilon_n := \frac{1}{n}$, we have:

$$(2.31) \quad \left(c_n d_n x'_n, \frac{x''_n}{\varepsilon_n}, x'''_n, \varepsilon_n, \frac{1}{\varepsilon_n c_n d_n} \right) \in \tilde{p}^{-1}(S)$$

and it converges to the point:

$$(2.32) \quad (v'_0, v''_0, x'''_0, 0, 0) = (v'_0, 0, x'''_0, 0, 0) \in T_N M \times_M T_M X.$$

Therefore this completes the proof of Proposition 2.6. ■

There is another interpretation of $C_{NM}(S)$.

Proposition 2.7. *Let $p_0 = (v'_0, v''_0, 0) \in T_N M \times_M T_M X$ and set:*

$$\Gamma_\varepsilon := \{(x', x'', x''') \in X; |x'| < \varepsilon |x''|\}.$$

(i) *Assume $v'_0 \neq 0, v''_0 \neq 0$. Then:*

$$(2.33) \quad p_0 \notin C_{NM}(S)$$

$$\iff \begin{cases} \text{there exist an open neighborhood } U \text{ of } 0, \\ \text{open convex cones } \Gamma' \subset X_{x'} \text{ and } \Gamma'' \subset X_{x''}, \\ \text{which satisfy } v'_0 \in \Gamma' \text{ and } v''_0 \in \Gamma'' \text{ respectively,} \\ \text{and } \varepsilon > 0, \text{ such that:} \\ S \cap \{U \cap (\Gamma' \times \Gamma'' \times X_{x'''}') \cap \Gamma_\varepsilon\} = \emptyset, \end{cases}$$

(ii) *Assume $v'_0 = 0, v''_0 \neq 0$. Then:*

$$(2.34) \quad p_0 = (0, v''_0, 0) \notin C_{NM}(S)$$

$$\iff \begin{cases} \text{there exist an open neighborhood } U \text{ of } 0, \\ \text{an open convex cone } \Gamma'' \subset X_{x''} \text{ which satisfy} \\ v''_0 \in \Gamma'', \text{ and } \varepsilon > 0, \text{ such that:} \\ S \cap \{U \cap (X_{x'}' \times \Gamma'' \times X_{x'''}') \cap \Gamma_\varepsilon\} = \emptyset. \end{cases}$$

(iii) *Assume $v'_0 \neq 0, v''_0 = 0$. Then:*

$$(2.35) \quad p_0 = (v'_0, 0, 0) \notin C_{NM}(S)$$

$$\iff \begin{cases} \text{there exist an open neighborhood } U \text{ of } 0 \text{ and} \\ \text{an open convex cone } \Gamma' \subset X_{x'} \text{ which satisfies} \\ v'_0 \in \Gamma', \text{ such that:} \\ S \cap \{U \cap (\Gamma' \times X_{x''}'' \times X_{x'''}')\} = \emptyset. \end{cases}$$

(iv) *Finally assume $v'_0 = 0, v''_0 = 0$. Then:*

$$(2.36) \quad p_0 = (0, 0, 0) \notin C_{NM}(S)$$

$$\iff \begin{cases} \text{there exists an open neighborhood } U \text{ of } 0, \\ \text{such that:} \\ S \cap U = \emptyset. \end{cases}$$

Proof. The parts (\Leftarrow) are always easy, so we only prove the parts (\Rightarrow). Throughout the proof, we use the results of the preceding proposition

as a definition of $C_{NM}(S)$.

(i) Let $\{U_n\}_{n \in \mathbb{N}}$ be a decreasing sequence of open neighborhoods of 0 in X such that:

$$\bigcap_{n \in \mathbb{N}} U_n = \{0\}.$$

Let $\{\Gamma'_n\}_{n \in \mathbb{N}}$ and $\{\Gamma''_n\}_{n \in \mathbb{N}}$ be decreasing sequences of open convex cones of $X'_{x'}$ and $X''_{x''}$ respectively which satisfy:

$$\bigcap_{n \in \mathbb{N}} \Gamma'_n = \{c \cdot v'_0; c \in \mathbb{R}^+\}, \quad \bigcap_{n \in \mathbb{N}} \Gamma''_n = \{c \cdot v''_0; c \in \mathbb{R}^+\}.$$

We assume: for every $n \in \mathbb{N}$ there exists:

$$x_n = (x'_n, x''_n, x'''_n) \in S \cap \{U_n \cap (\Gamma'_n \times \Gamma''_n \times X'''_n) \cap \Gamma_{\frac{1}{n}}\}.$$

Then, we have:

$$\begin{cases} x_n \longrightarrow 0, x'_n \in \Gamma'_n, x''_n \in \Gamma''_n \quad \text{and} \\ |x'_n| < \frac{1}{n} |x''_n|, \quad \text{for every } n \in \mathbb{N}. \end{cases}$$

Since the “directions” of x'_n and x''_n tend to v'_0 and v''_0 respectively, if we set

$$(2.37) \quad d_n := \frac{|v'_0|}{|v''_0|} \cdot \frac{|x''_n|}{|x'_n|} > \frac{|v'_0|}{|v''_0|} \cdot n > 0,$$

the “direction” of the vector $\begin{pmatrix} d_n x'_n \\ x''_n \end{pmatrix}$ tends to the vector $\begin{pmatrix} v'_0 \\ v''_0 \end{pmatrix}$, that is, there exists a sequence $c_n > 0$, such that:

$$c_n \begin{pmatrix} d_n x'_n \\ x''_n \end{pmatrix} \longrightarrow \begin{pmatrix} v'_0 \\ v''_0 \end{pmatrix}.$$

Moreover we have $d_n \rightarrow +\infty$ by (2.37). This contradicts the assumption that $(v'_0, v''_0, 0) \notin C_{NM}(S)$.

(ii) Assume as in the proof of (i): for every $n \in \mathbb{N}$, there exists:

$$x_n \in S \cap \{U_n \cap (X'_{x'} \times \Gamma''_n \times X'''_n) \cap \Gamma_{\frac{1}{n}}\}.$$

First of all, since the “direction” of x''_n tends to v''_0 , there exists a sequence $c_n > 0$, such that:

$$c_n x''_n \longrightarrow v''_0 \neq 0.$$

In particular, the sequence $\{c_n x_n''\}$ is bounded, and

$$|c_n x_n'| < \frac{1}{n} |c_n x_n''| < \frac{c}{n} \text{ for some } c > 0.$$

Now set $d_n := \sqrt{n} > 0$. Then:

$$\begin{cases} d_n \longrightarrow +\infty \\ c_n d_n x_n' \longrightarrow 0 \\ c_n x_n'' \longrightarrow v_0'' \neq 0. \end{cases}$$

This is a contradiction.

(iii) Assume: for every $n \in \mathbb{N}$, there exists:

$$x_n \in \mathcal{S} \cap \{U_n \cap (\Gamma_n' \times X_{x_n}'' \times X_{x_n}''')\}.$$

Then, extracting a subsequence, we may assume:

$$|x_n'| < \frac{1}{n}, \text{ for every } n \in \mathbb{N}.$$

Since the “direction” of x_n' tends to v_0' , by setting

$$d_n := \frac{|v_0'|}{|x_n'|} > |v_0'| \cdot n > 0,$$

we have:

$$d_n x_n' \longrightarrow v_0'.$$

Now set $c_n \equiv 1$ for every $n \in \mathbb{N}$, then we have a convergence:

$$\begin{cases} d_n \longrightarrow +\infty \\ c_n d_n x_n' \longrightarrow v_0' \neq 0 \\ c_n x_n'' \longrightarrow 0. \end{cases}$$

This is a contradiction.

(iv) Assume: there exists a sequence $x_n \in \mathcal{S}$, such that $x_n \rightarrow 0$. Extracting a subsequence, we may assume that:

$$|x_n'| < \frac{1}{n}, \text{ for every } n \in \mathbb{N}.$$

Set $c_n \equiv 1$, $d_n := \sqrt{n}$. Then:

$$\begin{cases} d_n \longrightarrow +\infty \\ c_n d_n x_n' \longrightarrow 0 \\ c_n x_n'' \longrightarrow 0. \end{cases}$$

This completes the proof of Proposition 2.7. ■

In terms of the normal cones $C_N(S)$ and $C_M(S)$ of [11], we get:

Corollary 2.8.

$$(2.38) \quad C_{NM}(S) \cap T_N M = C_N(S) \cap T_N M,$$

$$(2.39) \quad C_{NM}(S) \cap N \times_M T_M X = C_M(S) \cap N \times_M T_M X.$$

The following two propositions are essential tools for the study in the next section.

Proposition 2.9. *Let V be a biconic open subset of $T_N M \times_M T_M X$.*

(i) *Let W be an open neighborhood of V in \tilde{X}_{NM} , and set $U = \tilde{p}(W \cap \Omega)$. Then $V \cap C_{NM}(X \setminus U) = \emptyset$.*

(ii) *Let $U \subset X$ be an open set such that $V \cap C_{NM}(X \setminus U) = \emptyset$. Then $\tilde{p}^{-1}(U) \sqcup V$ is an open neighborhood of V in $\Omega \sqcup (T_N M \times_M T_M X)$.*

Proof. The proof is almost the same as that of Proposition 4.1.3 of [11], and we shall not repeat it. ■

The next proposition is analogous to Proposition 4.1.4 of [11].

Proposition 2.10. *Assume (2.28) and (2.29):*

$$T_N M \times_M T_M X \simeq \mathbb{R}_v^n \times \mathbb{R}_v'' \times \mathbb{R}_x'''.$$

Let $V = V' \times V'' \times V''' \subset T_N M \times_M T_M X$ be an open subset of the product-type with $V' \subsetneq \mathbb{R}_v^n$ and $V'' \subsetneq \mathbb{R}_v''$ are conic open subsets and $V''' \subset \mathbb{R}_x'''$ is an open subset. Then, the family of open neighborhoods W of V in \tilde{X}_{NM} which satisfy the following condition (2.40) forms a fundamental neighborhood system of V .

$$(2.40) \quad R\Gamma(F_x, \mathbb{C}_{F_x}) \simeq \mathbb{C} \text{ if } F_x \neq \emptyset \text{ for every } x \in X,$$

where $F_x := \tilde{p}^{-1}(x) \cap (W \cap \Omega)$.

Proof. Set $\dot{\Omega} := \Omega \cap \{x' \neq 0, x'' \neq 0\}$ and $S := \Omega \cap \{|x'| = 1, |x''| = 1\}$. Then $\dot{\Omega}$ is closed under the $(\mathbb{R}^+)^2$ -action (2.25) on \tilde{X}_{NM} and

$$S \simeq \dot{\Omega} / (\mathbb{R}^+)^2.$$

For a point $p_0 = (x', x'', x''', t_1, t_2) \in S$ which satisfies $|x'| = |x''| = 1$ and $t_1, t_2 > 0$, the orbit of the $(\mathbb{R}^+)^2$ -action through p_0 can be expressed as follows:

$$O_{p_0} := \left\{ \left(cdx', cx'', x''', \frac{t_1}{c}, \frac{t_2}{d} \right); (c, d) \in (\mathbb{R}^+)^2 \right\}$$

$$\simeq (\mathbb{R}^+)^2.$$

We identify O_{p_0} with $(\mathbb{R}^+)_{(c,d)}^2$ and the original point $p_0 \in S$ with the point $(c, d) = (1, 1)$ of $(\mathbb{R}^+)^2$. Now let us take an arbitrary open neighborhood W of V . One may assume $W \cap \Omega \subset \dot{\Omega}$, because $\{0\} \notin V'$ and $\{0\} \notin V''$. For $p_0 \in S \cap W$, $O_{p_0} \cap W$ corresponds to an open subset of $(\mathbb{R}^+)^2$ containing the point $(1, 1)$, and set:

$$\tilde{W}_{p_0} := \{ \text{maximal star-shaped open subset of } (\mathbb{R}^+)^2 \\ \text{centered at the point } (1, 1) \text{ in } O_{p_0} \cap W \}.$$

Define a subset \tilde{W} of $W \cap \Omega$ by:

$$\tilde{W} := \bigcup_{p_0 \in S \cap W} \tilde{W}_{p_0}.$$

Claim 1. \tilde{W} is an open subset of $W \cap \Omega$.

Proof. Take an arbitrary point $q_0 \in \tilde{W}_{p_0}$ ($p_0 \in S \cap W$), and an open subset U_{q_0} of $O_{p_0} \simeq (\mathbb{R}^+)^2$, such that:

$$[p_0, q_0] \subset U_{q_0} \subset \subset \tilde{W}_{p_0} \subset O_{p_0},$$

where $[p_0, q_0]$ denotes the line segment connecting the points p_0 and q_0 in $O_{p_0} \simeq (\mathbb{R}^+)^2$.

Now consider the continuous map:

$$\phi: S \times (\mathbb{R}^+)^2 \longrightarrow \tilde{X}_{NM}$$

defined by

$$((x', x'', x''', t_1, t_2), (c, d)) \longmapsto \left(cdx', cx'', x''', \frac{t_1}{c}, \frac{t_2}{d} \right).$$

Then we have $\phi(\{p_0\} \times \bar{U}_{q_0}) \subset W$. Since $\phi(\{p_0\} \times \bar{U}_{q_0})$ is compact in W , there exists an open neighborhood V_{p_0} of p_0 in S such that:

$$\begin{cases} p \in V_{p_0} \subset S \cap W, \text{ and} \\ \phi(V_{p_0} \times \bar{U}_{q_0}) \subset W. \end{cases}$$

Therefore ϕ induces a diffeomorphism between $V_{p_0} \times U_{q_0}$ and $\phi(V_{p_0} \times U_{q_0})$, and $\phi(V_{p_0} \times U_{q_0})$ is an open subset of W which contains q_0 . By the definition of \tilde{W} , we see that there exists an open neighborhood U of q_0 in \tilde{X}_{NM} , such that $U \subset \tilde{W}$. This completes the proof of Claim 1.

Claim 2. For every $x_0 = (x'_0, x''_0, x'''_0) \in V$ (hence $x'_0 \neq 0$ and $x''_0 \neq 0$), there exists an open neighborhood U_{x_0} of $(x_0, 0, 0)$ in \tilde{X}_{NM} such that:

$$U_{x_0} \cap \Omega \subset \tilde{W}.$$

Proof. Let us take an open neighborhood D_{x_0} of x_0 in V such that:

$$D_{x_0} = D_{x'_0} \times D_{x''_0} \times D_{x'''_0},$$

where $D_{x'_0} \subset \subset V'$, $D_{x''_0} \subset \subset V''$ and $D_{x'''_0} \subset \subset V'''$ are open balls centered at x'_0 , x''_0 and x'''_0 respectively. We only have to find $\delta > 0$ such that:

$$D_{x_0} \times \{0 < t_1 < \delta\} \times \{0 < t_2 < \delta\} \subset \tilde{W}.$$

First let us define K_S as the image of $\bar{D}_{x_0} = \bar{D}_{x'_0} \times \bar{D}_{x''_0} \times \bar{D}_{x'''_0}$ by the natural continuous mapping:

$$\phi_S: (\mathbb{R}^{n'} \setminus \{0\}) \times (\mathbb{R}^{n''} \setminus \{0\}) \times \mathbb{R}^{n'''} \longrightarrow S^{n'-1} \times S^{n''-1} \times \mathbb{R}^{n'''}$$

defined by

$$(x', x'', x''') \longmapsto \left(\frac{x'}{|x'|}, \frac{x''}{|x''|}, x''' \right),$$

and define $\hat{K}_{(\mathbb{R}^+)^2}$ as the image of $\bar{D}_{x'_0} \times \bar{D}_{x''_0}$ by the continuous map:

$$\phi_{(\mathbb{R}^+)^2}: (\mathbb{R}^{n'} \setminus \{0\}) \times (\mathbb{R}^{n''} \setminus \{0\}) \longrightarrow (\mathbb{R}^+)^2$$

defined by

$$(x', x'') \longmapsto \left(|x''|, \frac{|x'|}{|x''|} \right).$$

Next take an arbitrary compact star-shaped subset $K_{(\mathbb{R}^+)^2}$ in $(\mathbb{R}^+)^2$ centered at $(1, 1)$, such that:

$$\{(1, 1)\} \cup \hat{K}_{(\mathbb{R}^+)^2} \subset K_{(\mathbb{R}^+)^2},$$

and consider the continuous map:

$$\psi: (S^{n'-1} \times S^{n''-1} \times \mathbb{R}^{n'''}) \times (\mathbb{R}^+)^2 \longrightarrow \mathbb{R}^{n'} \times \mathbb{R}^{n''} \times \mathbb{R}^{n'''}$$

defined by

$$((y', y'', y'''), (c, d)) \longmapsto (cdy', cy'', y''').$$

Then $K := \psi(K_S \times K_{(\mathbb{R}^+)^2})$ is compact by definition, and it satisfies:

$$D_{x_0} \subset K \subset \subset V.$$

Since W is an open neighborhood of V in \tilde{X}_{NM} , there exists $\varepsilon_1 > 0$ such that:

$$K \times \{0 < t_1 < \varepsilon_1\} \times \{0 < t_2 < \varepsilon_1\} \subset W.$$

By virtue of the compactness of $K_{(\mathbb{R}^+)^2}$ in $(\mathbb{R}^+)^2$, there exists $\varepsilon_2 > 0$ such that:

$$\left\{ \left(cdy', cy'', y''', \frac{t_1}{c}, \frac{t_2}{d} \right); (y', y'', y''') \in K_S, (c, d) \in K_{(\mathbb{R}^+)^2}, 0 < t_1, t_2 < \varepsilon_2 \right\} \subset W.$$

Now we will complete the proof of Claim 2. For $(x', x'', x''') \in D_{x_0}$, take the points:

$$\left(\frac{x'}{|x'|}, \frac{x''}{|x''|}, x''' \right) \in K_S \quad \text{and} \quad \left(|x''|, \frac{|x'|}{|x''|} \right) \in \hat{K}_{(\mathbb{R}^+)^2} \subset K_{(\mathbb{R}^+)^2}.$$

The star-shapedness of $K_{(\mathbb{R}^+)^2}$ allows us to joint the points (1, 1) and $\left(|x''|, \frac{|x'|}{|x''|} \right)$ in $K_{(\mathbb{R}^+)^2}$ by a line segment, and we have:

$$\left(x', x'', x''', \frac{t_1}{|x''|}, \frac{t_2}{|x'|/|x''|} \right) \in \tilde{W},$$

for $0 < t_1, t_2 < \varepsilon_2$. Since we may assume there exists $c > 0$ such that:

$$|x''| < c \quad \text{and} \quad \frac{|x'|}{|x''|} < c \quad \text{for every } x \in D_{x_0},$$

we have by setting $\delta := \frac{\varepsilon_2}{c}$:

$$D_{x_0} \times \{0 < t_1 < \delta\} \times \{0 < t_2 < \delta\} \subset \tilde{W}.$$

Thus the proof of Claim 2 is also complete. Now we will finish the proof of the proposition. By Claim 2, we find an open neighborhood U_{x_0} of $(x_0, 0, 0)$ in \tilde{X}_{NM} for every $x_0 \in V$, such that:

$$U_{x_0} \subset W \quad \text{and} \quad U_{x_0} \cap \Omega \subset \tilde{W}.$$

Set:

$$\hat{W} := \tilde{W} \cup \left(\bigcup_{x_0 \in V} U_{x_0} \right).$$

Then it is an open neighborhood of V in \tilde{X}_{NM} and satisfies the condition:

$$\begin{cases} \hat{W} \subset W \\ \hat{W} \cap \Omega = \tilde{W}. \end{cases}$$

Therefore for every $x \in X$, $F_x := \tilde{p}^{-1}(x) \cap (\hat{W} \cap \Omega)$ is a star-shaped open subset

of $(\mathbb{R}^+)^2$ or an empty set. ■

§3. Bispecialization Functor

3.1. Functorial properties of the functor v_{NM}

In this section, we shall define the functor v_{NM} of bispecialization along (M, N) , and investigate its functorial properties. From now on, we will frequently use the terminology of derived categories. Let A be a commutative ring with finite global dimension. We denote by $\mathbf{D}^b(X)$ the derived category of complexes of sheaves of A_X -module with bounded cohomologies on a manifold X .

Definition 3.1. Let $F \in \mathbf{D}^b(X)$. One sets:

$$(3.1) \quad v_{NM}(F) := s^{-1}Rj_*\tilde{p}^{-1}F,$$

and says that $v_{NM}(F)$ is the bispecialization of F along (M, N) .

As we see in the next proposition, the functor v_{NM} is a generalization of the usual one v_M .

Proposition 3.2. *Let $M = N$ and identify $T_M M \times_M T_M X$ with $T_M X$. Then for every $F \in \mathbf{D}^b(X)$, we have a canonical isomorphism:*

$$(3.2) \quad v_M(F) \xrightarrow{\sim} v_{MM}(F).$$

Proof. First consider the natural morphism $q: \tilde{X}_{MM} \rightarrow \tilde{X}_M$ given by $(t_1, t_2) \mapsto t_1 t_2$ and the commutative diagram:

$$(3.3) \quad \begin{array}{ccccccc} T_M M \times_M T_M X & \xrightarrow{s} & \tilde{X}_{MM} & \xleftarrow{j} & \Omega & \xrightarrow{\tilde{p}} & X \\ \sim \downarrow & & q \downarrow & & q|_{\Omega} \downarrow & & id_X \downarrow \\ T_M X & \xrightarrow{s_M} & \tilde{X}_M & \xleftarrow{j_M} & \Omega_M & \xrightarrow{\tilde{p}_M} & X, \end{array}$$

where Ω_M is the open subset of \tilde{X}_M defined by $\Omega_M := \{t > 0\}$. Then we can construct the morphism:

$$v_M(F) \longrightarrow v_{MM}(F)$$

by the same way as in the proof of Proposition 4.2.5 of [11]. The stalks of $v_{MM}(F)$ coincide with those of $v_M(F)$ because we can easily obtain the results corresponding to Proposition 4.1.3 and Proposition 4.1.4 of [11] for the case $\tilde{X}_{MM} \rightarrow X$. This completes the proof. ■

Let (X, M, N) and (X', M', N') be triplets of manifolds, and f be a morphism from X' to X such that

$$(3.4) \quad \begin{cases} f(M') \subset M \\ f(N') \subset N. \end{cases}$$

For the sake of simplicity, we shall often write:

$$(3.5) \quad f: (X', M', N') \longrightarrow (X, M, N).$$

If we take a local coordinate system (x', x'', x''') (resp. (y', y'', y''')) of (X, M, N) (resp. (X', M', N')) and set

$$(3.6) \quad f(y', y'', y''') = (\phi'(y), \phi''(y), \phi'''(y)),$$

then we can define a morphism \tilde{f}' from $\tilde{X}'_{N'M'}$ to \tilde{X}_{NM} as follows:

$$(3.7) \quad \begin{cases} x' = \frac{1}{t_1 t_2} \phi'(t_1 t_2 y', t_1 y'', y''') \\ x'' = \frac{1}{t_1} \phi''(t_1 t_2 y', t_1 y'', y''') \\ x''' = \phi'''(t_1 t_2 y', t_1 y'', y''') \\ t_1 = t_1 \\ t_2 = t_2. \end{cases}$$

Restricting it to $\{t_1 = t_2 = 0\} = T_{N'}M' \times_M T_M X'$, we get a morphism Tf :

$$(3.8) \quad \begin{cases} v' = \frac{\partial \phi'}{\partial x'}(0, 0, y''') \cdot u' \\ v'' = \frac{\partial \phi''}{\partial x''}(0, 0, y''') \cdot u'' \\ x''' = \phi'''(0, 0, y'''), \end{cases}$$

where $\left(v' \frac{\partial}{\partial x'}, v'' \frac{\partial}{\partial x''}, x'''\right)$ (resp. $\left(u' \frac{\partial}{\partial y'}, u'' \frac{\partial}{\partial y''}, y'''\right)$) is the associated

coordinate system of $T_N M \times_M T_M X$ (resp. $T_{N'} M' \times_{M'} T_{M'} X'$). Thus, we get a commutative diagram below:

$$(3.9) \quad \begin{array}{ccccccc} T_{N'}M' \times_{M'} T_{M'}X' & \xrightarrow{sx'} & \tilde{X}'_{N'M'} & \xleftarrow{jx'} & \Omega_{X'} & \xrightarrow{\tilde{p}x'} & X' \\ Tf \downarrow & & \tilde{f}' \downarrow & & \tilde{f} \downarrow & & f \downarrow \\ T_N M \times_M T_M X & \xrightarrow{sx} & \tilde{X}_{NM} & \xleftarrow{jx} & \Omega_X & \xrightarrow{\tilde{p}x} & X, \end{array}$$

in which all squares are Cartesian.

By paraphrasing the arguments of [11], one obtains functorial properties of the functor v_{NM} .

Proposition 3.3. *Let $G \in \mathbf{D}^b(X')$.*

(i) *There exists a commutative diagram:*

$$(3.10) \quad \begin{array}{ccc} R(Tf)_! v_{N'M'}(G) & \longrightarrow & v_{NM}(Rf_! G) \\ \downarrow & & \downarrow \\ R(Tf)_* v_{N'M'}(G) & \longleftarrow & v_{NM}(Rf_* G). \end{array}$$

(ii) *Assume $f: X' \rightarrow X$ is a closed embedding, $f^{-1}(M) = M'$, $f^{-1}(N) = N'$, and $f: X' \rightarrow X$ and $f|_{M'}: M' \rightarrow M$ are clean w.r.t. M and N respectively. Then all these morphisms are isomorphisms.*

Proposition 3.4. *Let $F \in \mathbf{D}^b(X)$.*

(i) *There are canonical morphisms:*

$$(3.11) \quad \begin{cases} \alpha: (Tf)^{-1} v_{NM}(F) \longrightarrow v_{N'M'}(f^{-1}F), \\ \beta: v_{N'M'}(f^!F) \longrightarrow (Tf)^! v_{NM}(F), \end{cases}$$

such that the following diagram commutes:

$$(3.12) \quad \begin{array}{ccc} (Tf)^! A_{T_N M \times_M T_M X} \otimes^L (Tf)^{-1} v_{NM}(F) & \xrightarrow{\alpha} & v_{N'M'}(f^! A_X \otimes^L f^{-1}F) \\ \downarrow & & \downarrow \\ (Tf)^! v_{NM}(F) & \xleftarrow{\beta} & v_{N'M'}(f^!F). \end{array}$$

(ii) *If $f: X' \rightarrow X$, $f|_{M'}: M' \rightarrow M$ and $f|_{N'}: N' \rightarrow N$ are all smooth, then all these morphisms are isomorphisms.*

Proposition 3.5. *Let $F \in \mathbf{D}^b(X)$ and $G \in \mathbf{D}^b(X')$. Then we have a canonical morphism:*

$$(3.13) \quad v_{NM}(F) \boxtimes^L v_{N'M'}(G) \longrightarrow v_{(N \times N')(M \times M')}(F \boxtimes^L G)$$

in $\mathbf{D}^b(T_{(N \times N')(M \times M')} \times_{(M \times M')} T_{(M \times M')}(X \times X'))$.

Proposition 3.6. *Let $F, G \in \mathbf{D}^b(X)$. Then we have a canonical morphism:*

$$(3.14) \quad v_{NM}(F) \otimes^L v_{NM}(G) \longrightarrow v_{NM}(F \otimes^L G).$$

As a particular case, consider the morphism $f = \text{id}_X: (X, M, N) \rightarrow (X, M, M)$ and the commutative diagram:

$$(3.15) \quad \begin{array}{ccccccc} T_N M \times_M T_M X & \xrightarrow{s} & \tilde{X}_{NM} & \xleftarrow{j} & \Omega & \xrightarrow{\tilde{p}} & X \\ \swarrow & Tf \downarrow & \tilde{f} \downarrow & & \tilde{f} \downarrow & & f \downarrow \\ N \times_M T_M X & \longrightarrow & T_M X & \xrightarrow{s'} & \tilde{X}_{MM} & \xleftarrow{j'} & \Omega' & \xrightarrow{\tilde{p}'} & X \end{array}$$

in which all squares are Cartesian and Ω' is the open subset of \tilde{X}_{MM} defined by $\Omega' := \{t_1 > 0, t_2 > 0\}$. Then by Proposition 3.4, we get a natural morphism:

$$(3.16) \quad \alpha_0: (Tf)^{-1} v_{MM}(F) \longrightarrow v_{NM}(F)$$

Next, consider the following commutative diagram:

$$(3.17) \quad \begin{array}{ccccccc} T_N M \times_M T_M X & \xrightarrow{s} & \tilde{X}_{NM} & \xleftarrow{j} & \Omega & \xrightarrow{\tilde{p}} & X \\ \swarrow & \tau' \downarrow & p_{\tilde{M}} \downarrow & & \tilde{p}_{\tilde{M}} \downarrow & & id_X \downarrow \\ T_N M & \longrightarrow & T_N X & \xrightarrow{s_N} & \tilde{X}_N & \xleftarrow{j_N} & \Omega_N & \xrightarrow{\tilde{p}_N} & X, \end{array}$$

where Ω_N is the open subset of \tilde{X}_N defined by $\Omega_N := \{t_1 > 0\}$. Then we can construct a canonical morphism:

$$(3.18) \quad \beta_0: \tau'^{-1} v_N(F) \longrightarrow v_{NM}(F)$$

by the same way as in the proof of Proposition 4.2.5 of [11].

Theorem 3.7. *The morphisms $\alpha := \alpha_0|_{N \times_M T_M X}$ and $\beta := \beta_0|_{T_{NM}}$ are isomorphisms. In view of Proposition 3.2, for $F \in \mathbf{D}^b(X)$ we have natural isomorphisms:*

$$(3.19) \quad \alpha: v_M(F)|_{N \times_M T_M X} \xrightarrow{\sim} v_{NM}(F)|_{N \times_M T_M X},$$

$$(3.20) \quad \beta: v_N(F)|_{T_{NM}} \xrightarrow{\sim} v_{NM}(F)|_{T_{NM}}.$$

Proof. First let us show that α is an isomorphism, i.e.

$$\alpha: \tilde{f}'^{-1} Rj'_* \tilde{p}'^{-1} F|_{N \times_M T_M X} \xrightarrow{\sim} Rj_* \tilde{f}^{-1} \tilde{p}^{-1} F|_{N \times_M T_M X}.$$

For $p_0 = (x'_0, 0, 0, 0, 0) \in N \times_M T_M X \subset \tilde{X}_{NM}$ and for every $j \in \mathbb{Z}$:

$$\begin{aligned} H^j[Rj_*\tilde{f}^{-1}\tilde{p}'^{-1}F]_{p_0} &\simeq \varinjlim_{\varepsilon>0} H^j(U_\varepsilon; \tilde{f}^{-1}\tilde{p}'^{-1}F) \\ &\simeq \varinjlim_{\varepsilon>0} H^j(\tilde{f}(U_\varepsilon); \tilde{p}'^{-1}F), \end{aligned}$$

where

$$\begin{aligned} U_\varepsilon := \{x'; |x' - x'_0| < \varepsilon\} \times \{|x''| < \varepsilon\} \times \{|x'''| < \varepsilon\} \\ \times \{0 < t_1 < \varepsilon, 0 < t_2 < \varepsilon\} \end{aligned}$$

and we have used the fact that \tilde{f} is a diffeomorphism between Ω and Ω' in the second equality. Since \tilde{f} is explicitly described by:

$$(x', x'', x''', t_1, t_2) \longmapsto (x', t_1x'', x''', t_1, t_2)$$

by using a local coordinate system,

$$\begin{aligned} \tilde{f}(U_\varepsilon) = \{(x', x'', x''', t_1, t_2) \in \tilde{X}_{MM}; |x' - x'_0| < \varepsilon, |x''| < \varepsilon t_1, \\ |x'''| < \varepsilon, 0 < t_1 < \varepsilon, 0 < t_2 < \varepsilon\} \subset \Omega'. \end{aligned}$$

Now set

$$\begin{aligned} V_\varepsilon := \{(x', x'', x''', t_1, t_2) \in \tilde{X}_{MM}; |x' - x'_0| < \varepsilon, |x''| < \varepsilon^2, \\ |x'''| < \varepsilon, 0 < t_1 < \varepsilon, 0 < t_2 < \varepsilon\} \subset \Omega'. \end{aligned}$$

Then $V_\varepsilon \supset \tilde{f}(U_\varepsilon)$. On the other hand, we have:

$$H^j|\tilde{f}^{-1}Rj'_*\tilde{p}'^{-1}F]_{p_0} \simeq H^j[Rj'_*\tilde{p}'^{-1}F]_{\tilde{f}'(p_0)} \simeq \varinjlim_{\varepsilon>0} H^j(V_\varepsilon; \tilde{p}'^{-1}F)$$

for every $j \in \mathbb{Z}$, because $\tilde{f}'(p_0) = (x'_0, 0, 0, 0, 0)$ in \tilde{X}_{MM} . Thus it remains to show the isomorphism:

$$R\Gamma(V_\varepsilon; \tilde{p}'^{-1}F) \xrightarrow{\sim} R\Gamma(\tilde{f}(U_\varepsilon); \tilde{p}'^{-1}F)$$

for every $\varepsilon > 0$. For $x_1 = (x'_1, x''_1, x'''_1) \in X$, the inverse image $\tilde{p}'^{-1}(x_1)$ in $\Omega' \subset \tilde{X}_{MM}$ is expressed in the following way:

$$\tilde{p}'^{-1}(x_1) = \left\{ \left(\frac{x'_1}{t_1 t_2}, x''_1, x'''_1, t_1, t_2 \right); (t_1, t_2) \in (\mathbb{R}^+)^2 \right\}.$$

We can find that $\tilde{p}'^{-1}(x_1) \cap V_\varepsilon$ and $\tilde{p}'^{-1}(x_1) \cap \tilde{f}(U_\varepsilon)$ are isomorphic to open rectangles in $(\mathbb{R}^+)^2$ if they are not empty. Moreover we have:

$$\tilde{p}'^{-1}(x_1) \cap V_\varepsilon \neq \emptyset \iff \tilde{p}'^{-1}(x_1) \cap \tilde{f}(U_\varepsilon) \neq \emptyset.$$

Therefore, the required isomorphism is deduced from a slight modification of Corollary 3.7.3 of [11], and the morphism α is an isomorphism.

Next, we shall show that β is an isomorphism, i.e.

$$\beta: p_{\tilde{M}}^{-1} Rj_{N*} \tilde{p}_N^{-1} F|_{T_N M} \xrightarrow{\sim} Rj_{*} \tilde{p}_{\tilde{M}}^{-1} \tilde{p}_N^{-1} F|_{T_N M}.$$

For $q_0 = (0, x''_0, 0, 0) \in T_N M \subset \tilde{X}_{NM}$ and for every $j \in \mathbb{Z}$:

$$H^j[Rj_{*} \tilde{p}_{\tilde{M}}^{-1} \tilde{p}_N^{-1} F]_{q_0} \simeq \varinjlim_{\varepsilon > 0} H^j(W_\varepsilon; \tilde{p}_{\tilde{M}}^{-1} \tilde{p}_N^{-1} F),$$

where

$$\begin{aligned} W_\varepsilon := & \{ |x'| < \varepsilon \} \times \{ x''; |x'' - x''_0| < \varepsilon \} \times \{ |x'''| < \varepsilon \} \\ & \times \{ 0 < t_1 < \varepsilon, 0 < t_2 < \varepsilon \} \subset \Omega \subset \tilde{X}_{NM}. \end{aligned}$$

Now we have:

$$\begin{aligned} \tilde{p}_{\tilde{M}}(W_\varepsilon) = & \{ |x'| < \varepsilon^2 \} \times \{ x''; |x'' - x''_0| < \varepsilon \} \times \{ |x'''| < \varepsilon \} \\ & \times \{ 0 < t_1 < \varepsilon \} \subset \Omega_N \subset \tilde{X}_N, \end{aligned}$$

because $\tilde{p}_{\tilde{M}}: \Omega \rightarrow \Omega_N$ is explicitly described by:

$$(x', x'', x''', t_1, t_2) \longmapsto (t_2 x', x'', x''', t_1)$$

in a local coordinate system. Since $\tilde{p}_{\tilde{M}}|_{W_\varepsilon} = p_{\tilde{M}}|_{W_\varepsilon}: W_\varepsilon \rightarrow p_{\tilde{M}}(W_\varepsilon)$ is a topological submersion with contractible fibers, we have an isomorphism:

$$\text{id} \xrightarrow{\sim} R(p_{\tilde{M}}|_{W_\varepsilon})_*(p_{\tilde{M}}|_{W_\varepsilon})^{-1}$$

in $\mathbf{D}^b(p_{\tilde{M}}(W_\varepsilon))$ by Proposition 3.3.9 of [11]. Therefore, for every $j \in \mathbb{Z}$:

$$H^j(W_\varepsilon; \tilde{p}_{\tilde{M}}^{-1} \tilde{p}_N^{-1} F) \xleftarrow{\sim} H^j(p_{\tilde{M}}(W_\varepsilon); \tilde{p}_N^{-1} F).$$

On the other hand,

$$\begin{aligned} H^j[p_{\tilde{M}}^{-1} Rj_{N*} \tilde{p}_N^{-1} F]_{q_0} & \simeq H^j(Rj_{N*} \tilde{p}_N^{-1} F)_{p_{\tilde{M}}(q_0)} \\ & \simeq \varinjlim_{\varepsilon > 0} H^j(p_{\tilde{M}}(W_\varepsilon); \tilde{p}_N^{-1} F). \end{aligned}$$

because $p_{\tilde{M}}(q_0) = (0, x''_0, 0, 0)$ in \tilde{X}_N . This completes the proof. ■

3.2. Structure of the functor v_{NM}

In this section, we shall describe the global sections of $v_{NM}(F)$ on product type biconic open subsets of $T_N M \times_M T_M X$.

Assume:

$$(3.21) \quad \begin{cases} X = X'_{x'} \times X''_{x''} \times X'''_{x'''} \\ M = \{0\} \times X''_{x''} \times X'''_{x'''}, \\ N = \{0\} \times \{0\} \times X'''_{x'''} \end{cases}$$

where X' , X'' and X''' are real finite dimensional vector spaces. Then we have:

$$(3.22) \quad T_N M \times_M T_M X = X'_{v'} \times X''_{v''} \times X'''_{v'''}$$

Theorem 3.8. *Let $F \in \mathbf{D}^b(X)$. Then:*

(i) $v_{NM}(F) \in \mathbf{D}^b_{\mathbb{R}^+ \times \mathbb{R}^+}(T_N M \times_M T_M X)$, that is, it is a biconic object on $T_N M \times_M T_M X$, and

$$(3.23) \quad \text{supp } v_{NM}(F) \subset C_{NM}(\text{supp } F).$$

(ii) *Let $V \subset T_N M \times_M T_M X$ be a biconic open subset of the product type:*

$$(3.24) \quad V = V' \times V'' \times V''',$$

that is, $V' \subset X'_{v'}$ and $V'' \subset X''_{v''}$ are conic open subsets and $V''' \subset X'''_{v'''}$ is an open set. Then for every $j \in \mathbb{Z}$:

$$(3.25) \quad H^j(V; v_{NM}(F)) \simeq \varinjlim_{C_{NM}(X \setminus U) \cap V \neq \emptyset} H^j(U; F).$$

Proof.

(i) Obvious. (ii) **Case 1.** Assume $V' \subsetneq X'_{v'}$ and $V'' \subsetneq X''_{v''}$. Then the proof is almost similar to that of Theorem 4.2.3 of [11]. (Use Proposition 2.9 and Proposition 2.10. See Remark 3.3.10 of [11].)

Case 2. Assume $V' = X'_{v'}$, i.e.

$$(3.26) \quad V = X'_{v'} \times V'' \times V'''$$

Since $C_{NM}(S)$ is a biconic closed subset of $T_N M \times_M T_M X$ for every $S \subset X$, we have:

$$(3.27) \quad \begin{aligned} C_{NM}(X \setminus U) \cap V &= \emptyset \\ \iff C_{NM}(X \setminus U) \cap (\{0\} \times V'' \times V''') &= \emptyset \\ \iff C_N(X \setminus U) \cap (\{0\} \times V'' \times V''') &= \emptyset \end{aligned}$$

for every open subset $U \subset X$, where we identified $\{0\} \times V'' \times V''' \subset T_N M$ with a subset of $T_N X$ and we have used Corollary 2.38. For such $U \subset X$, we can construct a morphism

$$(3.28) \quad A: R\Gamma(U; F) \longrightarrow R\Gamma(V; v_{NM}(F))$$

by Proposition 2.9 (ii). (See the formula (4.2.2) of [11]). Since $v_{NM}(F)$ is a biconic object of $\mathbf{D}^b(T_N M \times_M T_M X)$,

$$(3.29) \quad R\Gamma(V; v_{NM}(F)) \xrightarrow{\sim} R\Gamma(\{0\} \times V'' \times V'''; v_{NM}(F)|_{T_{NM}}),$$

and there is a commutative diagram:

$$(3.30) \quad \begin{array}{ccc} R\Gamma(U; F) & \xrightarrow{\mathbf{A}} & R\Gamma(V; v_{NM}(F)) \\ \mathbf{B} \downarrow & & \downarrow \sim \\ R\Gamma(\{0\} \times V'' \times V'''; v_N(F)|_{T_{NM}}) & \xrightarrow{\mathbf{C}} & R\Gamma(\{0\} \times V'' \times V'''; v_{NM}(F)|_{T_{NM}}). \end{array}$$

The morphism \mathbf{C} is an isomorphism by Theorem 3.7, and it follows from Corollary 3.7.3 of [11] that:

$$(3.31) \quad H^j(\{0\} \times V'' \times V'''; v_N(F)|_{T_{NM}}) \xleftarrow{\sim} \varinjlim_{C_N(X \setminus U) \cap (\{0\} \times V'' \times V''') = \emptyset} H^j(U; F)$$

for every $j \in \mathbb{Z}$. Thus, by virtue of (3.27) and the commutativity of the diagram (3.30), we get:

$$(3.32) \quad H^j(V; v_{NM}(F)) \xleftarrow{\sim} \varinjlim_{C_{NM}(X \setminus U) \cap V = \emptyset} H^j(U; F)$$

for every $j \in \mathbb{Z}$.

Case 3. Assume $V'' = X''_{v''}$, i.e.

$$(3.33) \quad V = V' \times X''_{v''} \times V'''.$$

If we use (3.19):

$$v_{NM}(F)|_{N \times_M T_M X} \xleftarrow{\sim} v_M(F)|_{N \times_M T_M X},$$

Corollary 2.38 and Proposition 2.9 (ii), the proof proceeds similarly to that of Case 2. ■

Now, we consider $T_N M$ and $N \times_M T_M X$ as zero-sections of $T_N M \times_M T_M X$. We want to know the structure of $R\Gamma_{(T_N M)}(v_{NM}(F))$ and $R\Gamma_{(N \times_M T_M X)}(v_{NM}(F))$. Consider the diagram (3.17). Then we have canonical morphisms:

$$(3.34) \quad \begin{array}{ccc} \tau'^{-1} v_N(R\Gamma_M(F)) & \longrightarrow & v_{NM}(R\Gamma_M(F)) \\ & \xleftarrow{\sim} & R\Gamma_{(T_N M)}(v_{NM}(R\Gamma_M(F))) \\ & & \longrightarrow R\Gamma_{(T_N M)}(v_{NM}(F)), \end{array}$$

where the first arrow is obtained by (3.18) and the second isomorphism is a consequence of:

$$(3.35) \quad \text{supp } v_{NM}(R\Gamma_M(F)) \subset T_N M.$$

Theorem 3.9. *Let $F \in \mathbf{D}^b(X)$. Then the restriction of the morphism (3.34) to $T_N M \subset T_N M \times_M T_M X$:*

$$(3.36) \quad v_N(R\Gamma_M(F))|_{T_N M} \longrightarrow R\Gamma_{(T_N M)}(v_{NM}(F))|_{T_N M}$$

is an isomorphism.

Proof. Notice:

$$(3.37) \quad T_N M \times_M T_M X \setminus T_N M \simeq T_N M \times_M \dot{T}_M X,$$

and denote this set by A . Then we have a sequence of morphisms:

$$(3.38) \quad \begin{array}{c} \tau'^{-1}v_N(R\Gamma_{(X \setminus M)}(F)) \longrightarrow v_{NM}(R\Gamma_{(X \setminus M)}(F)) \\ \qquad \qquad \qquad \qquad \qquad \qquad \longrightarrow R\Gamma_A(v_{NM}(R\Gamma_{(X \setminus M)}(F))) \\ \qquad \qquad \qquad \qquad \qquad \qquad \longleftarrow \sim R\Gamma_A(v_{NM}(F)), \end{array}$$

where the last isomorphism follows from the formula $R\Gamma_A(v_{NM}(R\Gamma_M(F))) \simeq 0$ (cf. (3.35)). Now we have a morphism of distinguished triangles

$$(3.39) \quad \begin{array}{ccccc} \tau'^{-1}v_N(R\Gamma_M F) & \longrightarrow & \tau'^{-1}v_N(F) & \longrightarrow & \tau'^{-1}v_N(R\Gamma_{X \setminus M}(F)) & \xrightarrow{+1} \\ \downarrow & & \downarrow & & \downarrow & \\ R\Gamma_{(T_N M)}(v_{NM}(F)) & \longrightarrow & v_{NM}(F) & \longrightarrow & R\Gamma_A(v_{NM}(F)) & \xrightarrow{+1}, \end{array}$$

and we know by Theorem 3.7:

$$v_N(F)|_{T_N M} \xrightarrow{\sim} v_{NM}(F)|_{T_N M}.$$

Hence, it is enough to show

$$(3.40) \quad v_N(R\Gamma_{(X \setminus M)}(F))|_{T_N M} \xrightarrow{\sim} R\Gamma_A(v_{NM}(F))|_{T_N M}.$$

For this purpose, suppose the condition (3.21) and (3.22), and choose:

$$p_0 = (0, x''_0, 0) \in T_N M \subset T_N M \times_M T_M X.$$

Since $v_{NM}(F)$ is conic w.r.t. the v' -variables,

$$(3.41) \quad H^j[R\Gamma_{(T_N M \times_M T_M X \setminus T_N M)}(v_{NM}(F))]_{p_0} \simeq \varinjlim_{\varepsilon > 0} H^j(U_\varepsilon; \tilde{p}^{-1}F),$$

where

$$(3.42) \quad \begin{aligned} U_\varepsilon := & \{1 - \varepsilon < |x'| < 1 + \varepsilon\} \times \{|x'' - x''_0| < \varepsilon\} \times \{|x'''| < \varepsilon\} \\ & \times \{0 < t_1 < \varepsilon, 0 < t_2 < \varepsilon\} \subset \Omega \subset \tilde{X}_{NM}. \end{aligned}$$

Now recall $p_{\tilde{M}}: \tilde{X}_{NM} \rightarrow \tilde{X}_N$ is described by

$$(x', x'', x''', t_1, t_2) \longmapsto (t_2 x', x'', x''', t_1),$$

and $p_{\tilde{M}}|_{U_\varepsilon} = \tilde{p}_{\tilde{M}}|_{U_\varepsilon}: U_\varepsilon \rightarrow p_{\tilde{M}}(U_\varepsilon)$ is a smooth morphism with contractible fibers. Thus by Proposition 3.3.9 of [11], we have:

$$(3.43) \quad R\Gamma(p_{\tilde{M}}(U_\varepsilon); \tilde{p}_N^{-1}F) \xrightarrow{\sim} R\Gamma(p_{\tilde{M}}(U_\varepsilon); R(p_{\tilde{M}}|_{U_\varepsilon})_*(p_{\tilde{M}}|_{U_\varepsilon})^{-1}\tilde{p}_N^{-1}F) \\ \simeq R\Gamma(U_\varepsilon; \tilde{p}^{-1}F).$$

On the other hand,

$$(3.44) \quad p_{\tilde{M}}(U_\varepsilon) = \{0 < |x'| < \varepsilon(1 + \varepsilon)\} \times \{|x'' - x_0''| < \varepsilon\} \\ \times \{|x'''| < \varepsilon\} \times \{0 < t_1 < \varepsilon\} \subset \Omega_N \subset \tilde{X}_N,$$

and it follows from $\tilde{p}_N^{-1}R\Gamma_{(X \setminus M)}(F) \simeq R\Gamma_{(\Omega_N \setminus \tilde{M})}(\tilde{p}_N^{-1}F)$ that:

$$(3.45) \quad H^j[v_N(R\Gamma_{(X \setminus M)}(F))]_{p_0} \simeq \varinjlim_{\varepsilon > 0} H^j(p_{\tilde{M}}(U_\varepsilon); \tilde{p}_N^{-1}F).$$

This completes the proof of Theorem 3.9. ■

Let:

$$(3.46) \quad \iota: N \times_M T_M X \hookrightarrow T_N M \times_M T_M X$$

be the zero-section embedding and

$$(3.47) \quad \rho: T_N X \longrightarrow N \times_M T_M X$$

be the canonical projection. Then:

Theorem 3.10. *There exists a canonical isomorphism for $F \in \mathbf{D}^b(X)$:*

$$(3.48) \quad R\rho_! v_N(F) \xrightarrow{\sim} \iota^! v_{NM}(F).$$

Proof. Consider the following commutative diagram:

$$(3.49) \quad \begin{array}{ccccccc} T_N X & \xrightarrow{s''} & \tilde{X}_{NM} & \xleftarrow{j''} & \Omega'' & \xrightarrow{\tilde{p}''} & X \\ \swarrow \rho & Tg \downarrow & \tilde{g}' \downarrow & & \tilde{g} \downarrow & & \downarrow g = id_X \\ N \times_M T_M X & \xrightarrow{\iota} & T_N M \times_M T_M X & \xrightarrow{s} & \tilde{X}_{NM} & \xleftarrow{j} & \Omega & \xrightarrow{\tilde{p}} & X. \end{array}$$

This is a particular case of the diagram (3.9) for the morphism

$$g = id_X: (X, N, N) \longrightarrow (X, M, N)$$

(Note that $\tilde{g}: \Omega'' \simeq \Omega$ is an isomorphism). By virtue of Proposition 3.3 (i), we have a canonical morphism:

$$(3.50) \quad R(Tg)_! v_{NN}(F) \longrightarrow v_{NM}(F).$$

Hence there is a sequence of canonical morphisms:

$$(3.51) \quad \begin{aligned} R\rho_! v_{NN}(F) &\longrightarrow i^! R i_! R\rho_! v_{NN}(F) \simeq i^! R(Tg)_! v_{NN}(F) \\ &\longrightarrow i^! v_{NM}(F). \end{aligned}$$

Since $v_{NN}(F)$ coincides with $v_N(F)$ by Proposition 3.2, we have constructed a canonical morphism:

$$(3.52) \quad R\rho_! v_N(F) \longrightarrow i^! v_{NM}(F).$$

To verify that it is in fact an isomorphism, it is enough to show the following lemma. ■

Lemma 3.11. *Assume (3.21) and (3.22), and let $\left(v' \frac{\partial}{\partial x'}, v'' \frac{\partial}{\partial x''}, x'''\right)$ be a coordinate system of $T_N M \times_M T_M X$. Then, for $p_0 = (v'_0, 0, 0) \in N \times_M T_M X$ and for every $j \in \mathbb{Z}$,*

$$(3.53) \quad H^j[R\Gamma_{(N \times_M T_M X)}(v_{NM}(F))]_{p_0} \xleftarrow{\sim} \varinjlim_{Z, U} H^j_{Z \cap U}(U; F),$$

where

$$Z = \{|x'| \geq \varepsilon |x''|\}, \quad U = (U' \times X'' \times X''') \cap \{|x| < \varepsilon\},$$

for open cones U' of $X'_{x'}$, such that $v'_0 \in U'$ and $\varepsilon > 0$.

Proof. Fix an open cone $U' \subset X'_{x'}$, such that $v'_0 \in U'$ and $\varepsilon > 0$, and set:

$$(3.54) \quad \begin{cases} U := (U' \times X'' \times X''') \cap \{|x| < \varepsilon\}, \\ \tilde{U} := U \cap \{|x'| < \varepsilon |x''|\}, \\ V := U' \times X''_{v''} \times \{|x'''| < \varepsilon\}, \\ \tilde{V} := U' \times (X''_{v''} - \{0\}) \times \{|x'''| < \varepsilon\}, \end{cases}$$

where V and \tilde{V} are biconic open cones of $T_N M \times_M T_M X$ of the product-type. Then by Proposition 2.7,

$$(3.55) \quad C_{NM}(X \setminus U) \cap V = \emptyset \quad \text{and} \quad C_{NM}(X \setminus \tilde{U}) \cap \tilde{V} = \emptyset.$$

Applying Proposition 2.9 (ii), we can construct a natural morphism:

$$(3.56) \quad R\Gamma(U; F) \longrightarrow R\Gamma(U; R\tilde{p}_* \tilde{p}^{-1} F)$$

$$\begin{aligned} &\simeq R\Gamma(\tilde{p}^{-1}(U); \tilde{p}^{-1}F) \\ &\longrightarrow R\Gamma(V; s^{-1}Rj_*\tilde{p}^{-1}F) \\ &\simeq R\Gamma(V; v_{NM}(F)) \end{aligned}$$

and similarly we have:

$$(3.57) \quad R\Gamma(\tilde{U}; F) \longrightarrow R\Gamma(\tilde{V}; v_{NM}(F)).$$

Notice that

$$(3.58) \quad \overline{\tilde{p}^{-1}(U \setminus \tilde{U})} \cap V \subset (V \setminus \tilde{V}) = U' \times \{0\} \times \{|x''| < \varepsilon\}$$

and there exists a natural morphism:

$$(3.59) \quad R\Gamma_{(U \setminus \tilde{U})}(U; F) \longrightarrow R\Gamma_{(V \setminus \tilde{V})}(V; F).$$

Now we have a morphism of distinguished triangles:

$$(3.60) \quad \begin{array}{ccccc} R\Gamma_{(U \setminus \tilde{U})}(U; F) & \longrightarrow & R\Gamma(U; F) & \longrightarrow & R\Gamma(\tilde{U}; F) \xrightarrow{+1} \\ \downarrow & & \downarrow & & \downarrow \\ R\Gamma_{(V \setminus \tilde{V})}(V; v_{NM}(F)) & \longrightarrow & R\Gamma(V; v_{NM}(F)) & \longrightarrow & R\Gamma(\tilde{V}; v_{NM}(F)) \xrightarrow{+1}. \end{array}$$

If we shrink U, \tilde{U}, V and \tilde{V} by replacing U' and $\varepsilon > 0$, we obtain:

$$(3.61) \quad \begin{cases} \varinjlim_U H^j(U; F) \xrightarrow{\sim} \varinjlim_V H^j(V; v_{NM}(F)) \\ \varinjlim_{\tilde{U}} H^j(\tilde{U}; F) \xrightarrow{\sim} \varinjlim_{\tilde{V}} H^j(\tilde{V}; v_{NM}(F)), \end{cases}$$

by Theorem 3.8 (ii) and Proposition 2.7. Applying the five lemma to (3.60), we have:

$$(3.62) \quad \begin{aligned} \varinjlim_{U, \tilde{U}} H^j_{(U \setminus \tilde{U})}(U; F) &\xrightarrow{\sim} \varinjlim_{V, \tilde{V}} H^j_{(V \setminus \tilde{V})}(V; v_{NM}(F)) \\ &\simeq H^j[R\Gamma_{(N \times_M T_M X)}(v_{NM}(F))]_{p_0}. \end{aligned}$$

Hence the proof is complete. ■

The method of the proof of the above Lemma allows us to obtain the following result:

Proposition 3.12. *Assume (3.21) and (3.22), and set:*

$$(3.63) \quad T_N M \times_M T_M^* X = X'_{\xi'} \times X''_{v''} \times X'''_{x'''}.$$

Then for $p_0 = (\zeta'_0, v''_0, 0) \in \dot{T}_N M \times_M \dot{T}_M^* X$ and every $j \in \mathbb{Z}$, we have:

$$(3.64) \quad \varinjlim_A H^j [R\Gamma_A(v_{NM}(F))]_{(0, v''_0, 0)} \xleftarrow{\sim} \varinjlim_{Z, U} H^j_{Z \cap U}(U; F),$$

where $A = A' \times X'' \times X'''$, A' ranges through the family of closed convex cones of X' such that $(A' - \{0\}) \subset \{x' \in X' \mid \langle x', \xi'_0 \rangle > 0\}$,

$$(3.65) \quad \begin{cases} Z = A' \times X'' \times X''', \\ U = (X' \times U'' \times X''') \cap \{|x'| < \varepsilon |x''|, |x| < \varepsilon\}, \varepsilon > 0, \end{cases}$$

and U'' ranges through the family of open cones of X'' such that $v''_0 \in U''$.

Proof. Similar to that of Lemma 3.11, and we shall omit it. ■

Denote by τ_M and τ_N the projection:

$$(3.66) \quad T_N M \xleftarrow{\tau_M} T_N M \times_M T_M X \xrightarrow{\tau_N} N \times_M T_M X,$$

and let us summarize the results of Proposition 3.2, Theorem 3.7, Theorem 3.9 and Theorem 3.10.

Theorem 3.13. *Let $F \in \mathbf{D}^b(X)$. Then there are canonical isomorphisms:*

$$(3.67) \quad R\tau_{N*} v_{NM}(F) \xleftarrow{\sim} v_M(F)|_{N \times_M T_M X},$$

$$(3.68) \quad R\tau_{M*} v_{NM}(F) \xleftarrow{\sim} v_N(F)|_{T_N M},$$

$$(3.69) \quad R\tau_{N!} v_{NM}(F) \xleftarrow{\sim} R\rho_! v_N(F),$$

$$(3.70) \quad R\tau_{M!} v_{NM}(F) \xleftarrow{\sim} v_N R\Gamma_M(F),$$

where ρ denotes the projection $T_N X \rightarrow N \times_M T_M X$.

§4. Bimicrolocalization Functors

4.1. Functors $v\mu_{NM}$ and μ_{NM}

Let

$$\mathcal{F}_1: \mathbf{D}^b(T_N M \times_M T_M X) \longrightarrow \mathbf{D}^b(T_N M \times_M T_M^* X)$$

and

$$\mathcal{F}_2: \mathbf{D}^b(T_N M \times_M T_M^* X) \longrightarrow \mathbf{D}^b(T_N^* M \times_M T_M^* X)$$

be Fourier-Sato transformations on $T_M X$ and $T_N M$ respectively. We shall use the projections:

$$(4.1) \quad T_N M \xleftarrow{\pi_M} T_N M \times_M T_M^* X \xrightarrow{\tau_N} N \times_M T_M^* X,$$

and

$$(4.2) \quad T_N^* M \xleftarrow{\pi_M} T_N^* M \times_M T_M^* X \xrightarrow{\pi_N} N \times_M T_M^* X,$$

Definition 4.1. For $F \in \mathbf{D}^b(X)$, set:

$$(4.3) \quad v\mu_{NM}(F) := \mathcal{F}_1(v_{NM}(F)) \in \mathbf{D}^b(T_N M \times_M T_M^* X),$$

$$(4.4) \quad \mu_{NM}(F) := \mathcal{F}_2(v\mu_{NM}(F)) \in \mathbf{D}^b(T_N^* M \times_M T_M^* X).$$

We leave the construction of the functor $\mu_{v_{NM}(F)}$ to the reader.

Theorem 4.2. Let $F \in \mathbf{D}^b(X)$. Then:

(i) $v\mu_{NM}(F)$ is a biconic object of $\mathbf{D}^b(T_N M \times_M T_M^* X)$.

(ii) In the situation of (3.21) and (3.22), let $\left(\xi' dx', v'' \frac{\partial}{\partial x''}, x'''\right)$ be the coordinates of $T_N M \times_M T_M^* X$. Let $p_0 = \left(\xi'_0 dx', v''_0 \frac{\partial}{\partial x''}, 0\right) \in \dot{T}_N M \times_M \dot{T}_M^* X$. Then for every $j \in \mathbb{Z}$:

$$(4.5) \quad H^j[v\mu_{NM}(F)]_{p_0} = \varinjlim_{Z, U} H^j_{Z \cap U}(U; F),$$

where the inductive limit is taken over the family of subsets Z and U of X such that: $Z = \gamma' \times X''_{x'} \times X'''_{x''}$, γ' is a closed cone of X' , such that

$$(4.6) \quad (\gamma' \setminus \{0\}) \subset \{x' \in X'; \langle x', \xi'_0 \rangle > 0\},$$

$$U = (X'_{x'} \times U'' \times X'''_{x''}) \cap \{|x'| < \varepsilon |x''|, |x| < \varepsilon\}, \quad \varepsilon > 0,$$

and U'' is an open cone of $X''_{x'}$ such that $v''_0 \in U''$.

(iii) We have canonical isomorphisms:

$$(4.7) \quad R\tau_{N^*} v\mu_{NM}(F) \xrightarrow{\sim} \mu_M(F)|_{N \times_M T_M^* X},$$

$$(4.8) \quad R\pi_{M^*} v\mu_{NM}(F) \xrightarrow{\sim} v_N R\Gamma_M(F),$$

$$(4.9) \quad R\tau_{N!} v\mu_{NM}(F) \xrightarrow{\sim} \mu_N(F)|_{N \times_M T_M^* X},$$

$$(4.10) \quad R\pi_{M!} \nu \mu_{NM}(F) \xleftarrow{\sim} \nu_N(F)|_{T_{NM}} \otimes \omega_{M \setminus X}.$$

Proof. (i) Obvious.

(ii) This is nothing but Proposition 3.12.

(iii) Using Proposition 3.7.13 and Proposition 3.7.14 of [11], these formulas are immediate consequences of Theorem 3.13. ■

Theorem 4.3. *Let $F \in \mathbf{D}^b(X)$. Then:*

(i) $\mu_{NM}(F)$ is a biconic object of $\mathbf{D}^b(T_N^*M \times_M T_M^*X)$.

(ii) In the situation of (3.21) and (3.22), let $(\xi' dx', \xi'' dx'', x''')$ be the coordinates of $T_N^*M \times_M T_M^*X$. Let $q_0 = (\xi'_0 dx', \xi''_0 dx'', 0) \in \dot{T}_N^*M \times_M \dot{T}_M^*X$. Then for every $j \in \mathbb{Z}$:

$$(4.11) \quad H^j[\mu_{NM}(F)]_{q_0} = \varinjlim_{Z,U} H^j_{Z \cap U}(U; F),$$

where the inductive limit is taken over the family of subsets Z and U of X such that: $Z = \{(\{0\} \times \gamma'' + \gamma) \times X'''_x, \gamma'' \subset X''_x, \text{ and } \gamma \subset X'_x \times X''_x\}$ are closed cones such that

$$(4.12) \quad (\gamma'' \setminus \{0\}) \subset \{x''; \langle x'', \xi''_0 \rangle > 0\},$$

$$(4.13) \quad (\gamma \setminus \{0\}) \subset \{(x', x''); \langle x', \xi'_0 \rangle > 0\},$$

and $U = \{x \in X; |x| < \varepsilon\}$ for some $\varepsilon > 0$.

(iii) We have canonical isomorphisms:

$$(4.14) \quad R\pi_{N*} \mu_{NM}(F) \xleftarrow{\sim} \mu_N(F)|_{N \times_M T_M^*X},$$

$$(4.15) \quad R\pi_{M*} \mu_{NM}(F) \xleftarrow{\sim} \mu_N R\Gamma_M(F),$$

$$(4.16) \quad R\pi_{N!} \mu_{NM}(F) \xleftarrow{\sim} \mu_M(F)|_{N \times_M T_M^*X} \otimes \omega_{N/M},$$

$$(4.17) \quad R\pi_{M!} \mu_{NM}(F) \xleftarrow{\sim} R\theta_! \mu_N(F),$$

where θ denotes the projection $T_N^*X \rightarrow T_N^*M$.

Proof. (i) is obvious.

(iii) follows from Theorem 4.2 (iii) by using Proposition 3.7.13 and Proposition 3.7.14 of [11]. It remains to show (ii).

(ii) Assume $x' = (x'_1, \dots, x'_n)$, $x'' = (x''_1, \dots, x''_n)$, $\xi'_0 dx' = dx'_1$, and $\xi''_0 dx'' = dx''_1$. Now set for $\varepsilon > 0$:

$$(4.18) \quad \begin{cases} U_\varepsilon := \{(x', x'', x''') \in X; |x'| < \varepsilon^2, |x''| < \varepsilon, |x'''| < \varepsilon\}, \\ \hat{U}_\varepsilon := U_\varepsilon \cap \{(x', x'', x''') \in X; |x'| < \varepsilon |x''|, x'_1 < \varepsilon |(x''_2, \dots, x''_{n'})|\} \\ \hat{Z}_\varepsilon := \{(x', x'', x''') \in X; x'_1 \geq \varepsilon |(x''_2, \dots, x''_{n'})|\}. \end{cases}$$

Since $T_N M \times_M T_M X = \mathbb{R}_v^{n'} \times \mathbb{R}_{v''}^{n''} \times \mathbb{R}_{x'''}^{n'''}$, with the notations $v' = (v'_1, \dots, v'_{n'})$ and $v'' = (v''_1, \dots, v''_{n''})$, we can define following subsets of $T_N M \times_M T_M X$ for $\varepsilon > 0$:

$$(4.19) \quad \begin{cases} V_\varepsilon := \{(v', v'', x'''); |x'''| < \varepsilon\}, \\ \hat{V}_\varepsilon := \{(v', v'', x'''); |x'''| < \varepsilon, v''_1 < \varepsilon |(v''_2, \dots, v''_{n''})|\}, \\ \hat{A}_\varepsilon := \{(v', v'', x'''); v'_1 \geq \varepsilon |(v''_2, \dots, v''_{n''})|\}. \end{cases}$$

Then, by the techniques used in the proof of Lemma 3.11, we have for every $j \in \mathbb{Z}$:

$$(4.20) \quad \varinjlim_{\varepsilon > 0} H_{(\hat{A}_\varepsilon \cap V_\varepsilon)}^j(V_\varepsilon; v_{NM}(F)) \xleftarrow{\sim} \varinjlim_{\varepsilon > 0} H_{(\hat{Z}_\varepsilon \cap U_\varepsilon)}^j(U_\varepsilon; F),$$

$$(4.21) \quad \varinjlim_{\varepsilon > 0} H_{(\hat{A}_\varepsilon \cap \hat{V}_\varepsilon)}^j(\hat{V}_\varepsilon; v_{NM}(F)) \xleftarrow{\sim} \varinjlim_{\varepsilon > 0} H_{(\hat{Z}_\varepsilon \cap \hat{U}_\varepsilon)}^j(\hat{U}_\varepsilon; F).$$

Now again, set for every $\varepsilon > 0$:

$$(4.22) \quad \begin{cases} Z_\varepsilon := (\hat{Z}_\varepsilon \cap U_\varepsilon) \setminus \hat{U}_\varepsilon, \\ A_\varepsilon := (\hat{A}_\varepsilon \cap V_\varepsilon) \setminus \hat{V}_\varepsilon. \end{cases}$$

Then one has

$$(4.23) \quad C_{NM}(Z_\varepsilon) \cap V_\varepsilon \subset A_\varepsilon$$

by Proposition 2.6 (ii), and there exist a canonical morphism:

$$(4.24) \quad R\Gamma_{Z_\varepsilon}(U_\varepsilon; F) \longrightarrow R\Gamma_{A_\varepsilon}(V_\varepsilon; v_{NM}(F))$$

and a morphism between distinguished triangles:

$$(4.25) \quad \begin{array}{ccccc} R\Gamma_{Z_\varepsilon}(U_\varepsilon; F) & \longrightarrow & R\Gamma_{(\hat{Z}_\varepsilon \cap U_\varepsilon)}(U_\varepsilon; F) & \longrightarrow & R\Gamma_{(\hat{Z}_\varepsilon \cap \hat{U}_\varepsilon)}(\hat{U}_\varepsilon; F) & \xrightarrow{+1} \\ \downarrow & & \downarrow & & \downarrow & \\ R\Gamma_{A_\varepsilon}(V_\varepsilon; v_{NM}(F)) & \longrightarrow & R\Gamma_{(\hat{A}_\varepsilon \cap V_\varepsilon)}(V_\varepsilon; v_{NM}(F)) & \longrightarrow & R\Gamma_{(\hat{A}_\varepsilon \cap \hat{V}_\varepsilon)}(\hat{V}_\varepsilon; v_{NM}(F)) & \xrightarrow{+1}. \end{array}$$

By virtue of (4.20), (4.21) and the five Lemma, we have for every $j \in \mathbb{Z}$:

$$(4.26) \quad \varinjlim_{\varepsilon > 0} H_{A_\varepsilon}^j(V_\varepsilon; v_{NM}(F)) \xleftarrow{\sim} \varinjlim_{\varepsilon > 0} H_{Z_\varepsilon}^j(U_\varepsilon; F),$$

where the left hand side coincides with $H^j[\mu_{NM}(F)]_{q_0}$ ($q_0 = (dx'_1, dx''_1, 0)$).
Now notice:

(4.27)

$$Z_\varepsilon = U_\varepsilon \cap \{(x', x'', x'''); x'_1 \geq \varepsilon |(x'_2, \dots, x'_n)|\} \\ \cap [\{(x', x'', x'''); x'_1 \geq \varepsilon |(x''_2, \dots, x''_n)|\} \cup \{(x', x'', x'''); |x'| \geq \varepsilon |x''|\}],$$

and it is a direct calculation that the right hand side of (4.26) coincides with the right hand side of (4.11). This completes the proof of the theorem. ■

4.2. Functorial properties of $v\mu_{NM}$ and μ_{NM}

Let $f: (X', M', N') \rightarrow (X, M, N)$ be a morphism of triplets of manifolds. The morphism:

$$Tf: T_{N'}M' \times_{M'} T_{M'}X' \longrightarrow T_N M \times_M T_M X$$

induced by f is decomposed in the following way:

$$(4.28) \quad T_{N'}M' \times_{M'} T_{M'}X' \xrightarrow{f_1} T_{N'}M' \times_{M'} (M' \times_M T_M X) \\ \simeq T_{N'}M' \times_{M'} T_M X \\ \xrightarrow{f_2} (N' \times_N T_N M) \times_M T_M X \\ \xrightarrow{f_3} T_N M \times_M T_M X.$$

If we consider these morphisms w.r.t. the second vector bundle structure, f_1 is a morphism of vector bundles over $T_{N'}M'$ and f_2 and f_3 are base changes of vector bundles. Taking the dual of the above sequence of morphisms, we have:

$$(4.29) \quad T_{N'}M' \times_{M'} T_{M'}^* X' \xleftarrow{\tau_{f_1}} T_{N'}M' \times_{M'} T_M^* X \\ \xrightarrow{f_{2\pi}} T_N M \times_M (N' \times_M T_M^* X) \\ \xrightarrow{f_{3\pi}} T_N M \times_M T_M^* X.$$

Next let us consider them w.r.t. the first vector bundle structure. Then f_1 and $f_{3\pi}$ are base changes of vector bundles and $f_{2\pi}$ is a morphism of vector bundles over $N' \times_M T_M^* X$. Taking the dual again, we have:

$$(4.30) \quad T_{N'}^* M' \times_{M'} T_{M'}^* M' \xleftarrow{\tau_{f_{1\pi}}} T_{N'}^* M' \times_{M'} T_M^* X$$

$$\begin{aligned} & \xleftarrow{f_{2\pi}^!} T_N^* M \times_M (N' \times_M T_M^* X) \\ & \xrightarrow{f_{3\pi\pi}} T_N^* M \times_M T_M^* X. \end{aligned}$$

The following propositions are very similar to the corresponding results in [11], and the proofs proceed completely similarly. Hence we shall omit them.

Proposition 4.4. *Let $F \in \mathbf{D}^b(X)$, $G \in \mathbf{D}^b(X')$. Then there are commutative diagrams below:*

$$(4.31) \quad \begin{array}{ccc} R(f_{3\pi} \circ f_{2\pi})_! f_1^{-1} v\mu_{N'M'}(G) & \longrightarrow & v\mu_{NM}(Rf_! G) \\ \downarrow & & \downarrow \\ R(f_{3\pi} \circ f_{2\pi})_* [f_1^! v\mu_{N'M'}(G) \otimes (f_1^! A_{T_N^* M' \times_M T_M^* X'})^{\otimes -1}] & \longleftarrow & v\mu_{NM}(Rf_* G) \end{array}$$

and

$$(4.32) \quad \begin{array}{ccc} Rf_{1!} [(f_{3\pi} \circ f_{2\pi})^! A_{T_N^* M' \times_M T_M^* X} \otimes (f_{3\pi} \circ f_{2\pi})^{-1} v\mu_{NM}(F)] & \longrightarrow & v\mu_{N'M'}(f^! A_X \otimes f^{-1} F) \\ \downarrow & & \downarrow \\ Rf_{1*} (f_{3\pi} \circ f_{2\pi})^! v\mu_{NM}(F) & \longleftarrow & v\mu_{N'M'}(f^! F). \end{array}$$

Proposition 4.5. *Let $F \in \mathbf{D}^b(X)$, $G \in \mathbf{D}^b(X')$. Then there are commutative diagrams below:*

$$(4.33) \quad \begin{array}{ccc} Rf_{3\pi\pi!} (f_{1\pi} \circ f_{2\pi})^{-1} \mu_{N'M'}(G) & \longrightarrow & \mu_{NM}(Rf_! G) \\ \downarrow & & \downarrow \\ Rf_{3\pi\pi*} [(f_{1\pi} \circ f_{2\pi})^! \mu_{N'M'}(G) \otimes \{(f_{1\pi} \circ f_{2\pi})^! A_{T_N^* M' \times_M T_M^* X'}\}^{\otimes -1}] & \longleftarrow & \mu_{NM}(Rf_* G) \end{array}$$

and

$$(4.34) \quad \begin{array}{ccc} R(f_{1\pi} \circ f_{2\pi})_! [f_{3\pi\pi}^! A_{T_N^* M' \times_M T_M^* X} \otimes f_{3\pi\pi}^{-1} \mu_{NM}(F)] & \longrightarrow & \mu_{N'M'}(f^! A_X \otimes f^{-1} F) \\ \downarrow & & \downarrow \\ R(f_{1\pi} \circ f_{2\pi})_* f_{3\pi\pi}^! \mu_{NM}(F) & \longleftarrow & \mu_{N'M'}(f^! F). \end{array}$$

Proposition 4.6. *Let $F \in \mathbf{D}^b(X)$, $G \in \mathbf{D}^b(X')$. Then, we have canonical morphisms:*

$$(4.35) \quad v\mu_{NM}(F) \boxtimes^L v\mu_{N'M'}(G) \longrightarrow v\mu_{(N \times N')(M \times M')}(F \boxtimes^L G)$$

and

$$(4.36) \quad \mu_{NM}(F) \boxtimes^L \mu_{N'M'}(G) \longrightarrow \mu_{(N \times N')(M \times M')}(F \boxtimes^L G).$$

Now let us consider the natural morphisms given by the addition of covectors:

$$(4.37) \quad \gamma: (T_N M \times_M T_M^* X) \times_{T_N M} (T_N M \times_M T_M^* X) \longrightarrow T_N M \times_M T_M^* X$$

and

$$(4.38) \quad \theta: (T_N^* M \times_M T_M^* X) \times_N (T_N^* M \times_M T_M^* X) \longrightarrow T_N^* M \times_M T_M^* X.$$

Then we have:

Proposition 4.7. *Let $F, G \in \mathbf{D}^b(X)$. There are canonical morphisms of “product”:*

$$(4.39) \quad R\gamma_! [\nu\mu_{NM}(F) \boxtimes_{T_N M}^L \nu\mu_{NM}(G)] \longrightarrow \nu\mu_{NM}(F \otimes^L G) \otimes \omega_{M/X}$$

and

$$(4.40) \quad R\theta_! [\mu_{NM}(F) \boxtimes_N^L \mu_{NM}(G)] \longrightarrow \mu_{NM}(F \otimes^L G) \otimes \omega_{N/X}.$$

4.3. Various generalizations of the functor μ_{hom}

Using the diagonal embedding, the functor of Sato’s microlocalization allows one to define the bifunctor $\mu_{hom}(\cdot, \cdot)$ ([11]). Here we shall proceed similarly, using the functor of second microlocalization, in various geometrical situations. The results in this section were obtained during some discussions with P. Schapira. In particular, the idea to work in the relative setting is due to him.

First, consider a smooth morphism of manifolds $f: X \rightarrow S$. We identify $X \times_S T^*S$ to an involutive submanifold of T^*X that we denote by V :

$$V := X \times_S T^*S,$$

and one defines as usual the relative cotangent bundle $T^*(X/S)$ by the exact sequence of vector bundles:

$$0 \longrightarrow X \times_S T^*S \longrightarrow T^*X \longrightarrow T^*(X/S) \longrightarrow 0.$$

Set $L := X \times_S X$ and consider the triplet of manifolds:

$$X \simeq \Delta_X \hookrightarrow L := X \times_S X \hookrightarrow X \times X.$$

The first projection from $T^*(X \times X)$ to T^*X defines the isomorphisms:

$$\begin{aligned} T_{\Delta_X}^*(X \times X) &\simeq T^*X, \\ \Delta_X \times_L T_L^*(X \times X) &\simeq X \times_S T^*S. \end{aligned}$$

Moreover, the natural morphism $T_{\Delta_X}^*(X \times X) \rightarrow T_{\Delta_X}^*(X \times_S X)$ being zero on $X \times_S T^*S$, it defines the morphism:

$$T^*(X/S) \longrightarrow T_{\Delta_X}^*(X \times_S X) = T_{\Delta_X}^* L$$

and one checks immediately that this is an isomorphism. Summarizing, we have got a natural isomorphism:

$$\begin{aligned} T_{\Delta_X}^* L \times_L T_L^*(X \times X) &\simeq T^*(X/S) \times_S T^*S \\ &\simeq T^*(X/S) \times_X V. \end{aligned}$$

The canonical projection $T^*X \rightarrow X$ and the Hamiltonian isomorphism define a sequence of morphisms:

$$\begin{aligned} V \times_X T^*X &\hookrightarrow V \times_{(T^*M)} T^*(T^*X) \\ &\simeq V \times_{(T^*X)} T(T^*X) \longrightarrow T_V(T^*X). \end{aligned}$$

The composition of these morphisms sends $V \times_X (X \times_S T^*S)$ to the zero-section of $T_V(T^*X)$ and we obtain a morphism:

$$V \times_X T^*(X/S) \longrightarrow T_V(T^*X)$$

and one immediately checks, using local coordinates, that this morphism is an isomorphism.

Denote as usual by q_1 and q_2 the first and second projections from $X \times X$ to X .

Definition 4.8. Let F and G belong to $\mathbf{D}^b(X)$. We set:

$$\mu\mathit{hom}_f(G, F) = \mu_{\Delta_{X \times L}} \mathcal{R}\mathit{Hom}(q_2^{-1}G, q_1^!F).$$

Hence $\mu\mathit{hom}_f(G, F)$ is a biconic object on $T_{\Delta_X}^* L \times_L T_L^*(X \times X) \simeq T^*(X/S) \times_X V \simeq T_V(T^*X)$. When restricting to V , we find:

$$\mu\mathit{hom}_f(G, F)|_V \simeq \mu\mathit{hom}(G, F)|_V$$

and when restricting to $T^*(X/S)$, we find a sheaf which coincides with the “relative $\mu\mathit{hom}$ -sheaf” of Ishimura [3].

It is possible to extend the previous construction to the case where f is no more smooth by decomposing f by the graph embedding, $f = p \circ s$, with $s: X \hookrightarrow X \times S$, $p: X \times S \rightarrow S$, and setting:

$$\mu\mathit{hom}_f(G, F) = \mu\mathit{hom}_p(s_*G, s_*F).$$

Of course, one should prove that if f is smooth, one recovers the previous definition. This is left to the reader. (Notice that a similar trick already appears in [21].)

Next, consider the case of a closed embedding: $f: N \hookrightarrow X$. Set

$V := N \times_X T^*X$, and consider the triplet of manifolds:

$$N \hookrightarrow X \simeq \Delta_X \hookrightarrow X \times X.$$

The projection $(T^*X, V) \rightarrow (X, N)$ defines $T_V(T^*X) \rightarrow T_N X$. Combining with the projection $T_V(T^*X) \rightarrow V$, we find the isomorphism:

$$T_V(T^*X) \simeq T_N X \times_X V \simeq T_N X \times_X T^*X.$$

Definition 4.9. Let F and G belong to $\mathbf{D}^b(X)$. We set:

$$\mu\mathit{hom}_f(G, F) = v\mu_{N\Delta_X} R\mathcal{H}om(q_2^{-1}G, q_1^!F).$$

Hence, $\mu\mathit{hom}_f(G, F)$ is a biconic object on $T_N X \times_X T^*X \simeq T_V(T^*X)$. When restricting to V , we find:

$$\mu\mathit{hom}_f(G, F)|_V \simeq \mu\mathit{hom}(G, F)|_V,$$

and when restricting to $T_N X$, we find

$$\mu\mathit{hom}_f(G, F)|_{T_N X} \simeq v_N R\mathcal{H}om(G, F).$$

Here again, one could release the hypothesis that f is an embedding by decomposing f as $f = p \circ s$, with $s: N \hookrightarrow N \times X$, $p: N \times X \rightarrow X$, and setting:

$$\mu\mathit{hom}_f(G, F) = \mu\mathit{hom}_s(p^{-1}G, p^{-1}F).$$

Remark 4.10. Let us consider a geometrical construction which often appears when dealing with involutive submanifolds. Let $f: Y \rightarrow X$ be a smooth morphism, let $g: S \rightarrow Y$ be a closed embedding, let U be an open subset of T^*Y and assume the intersection of $W := U \cap (S \times_Y T^*Y)$ and $Y \times_X T^*X$ is clean and f_π induces an isomorphism:

$$V' := U \cap (S \times_Y T^*Y) \cap (Y \times_X T^*X) \simeq V,$$

where V is an involutive submanifold of T^*X . Denote by ρ and ϖ the natural morphisms from $T_{V'}(Y \times_X T^*X)$ to $T_W T^*Y$ and $T_V T^*X$ respectively. Let F and G be two objects of $\mathbf{D}^b(X)$. Then it would be interesting to construct an object $\mu\mathit{hom}_V(G, F)$ such that:

$$\varpi^{-1} \mu\mathit{hom}_V(G, F) \simeq \rho^{-1} \mu\mathit{hom}_g(f^{-1}G, f^{-1}F).$$

§5. Applications

5.1. Second microfunctions associated to a submanifold

Let M be a real analytic manifold of dimension n and N be a submanifold of M of codimension d . We denote by \mathcal{O}_X the sheaf of holomorphic functions on a complexification X of M . Let us apply our theory to the triplet (X, M, N) .

Definition 5.1. We set:

$$(5.1) \quad \begin{cases} \hat{\mathcal{C}}_{NM} := \nu\mu_{NM}(\mathcal{O}_X) \otimes \text{or}_M[n], \\ \mathcal{C}_{NM} := \mu_{NM}(\mathcal{O}_X) \otimes \text{or}_N[n]. \end{cases}$$

These are objects of the derived categories $\mathbf{D}^b(T_N M \times_M T_M^* X)$ and $\mathbf{D}^b(T_N^* M \times_M T_M^* X)$ respectively.

Assume $M = \mathbb{R}^n$, $N = \{0\} \times \mathbb{R}_x^{n-d}$ and $\Omega = \Omega' \times \mathbb{R}_x^{n-d} \subset M$ for an open convex proper cone Ω' in \mathbb{R}_x^d , such that $(1, 0, \dots, 0) \in \Omega'$, and take the associated coordinate system $\left(\sqrt{-1}\eta dx, v' \frac{\partial}{\partial x'}, x''\right)$ of $T_N M \times_M T_M^* X$. Then we know by [24] and [25] that the complex $\hat{\mathcal{C}}_{NM}$ coincides with the inductive limit of the complex $\mathcal{C}_{\Omega|X}$ due to [20] in the following sense.

Theorem 5.2. *The complex $\hat{\mathcal{C}}_{NM}$ is concentrated in degree 0, and for $p_0 := \left(\frac{\partial}{\partial x_1}, x'_0\right) \in T_N M$ we have an isomorphism:*

$$(5.2) \quad \varinjlim_{(1,0,\dots,0) \in \Omega'} \mathcal{C}_{\Omega|X}|_{\{x'_0\} \times_M T_M^* X} \simeq \hat{\mathcal{C}}_{NM}|_{\{p_0\} \times_M T_M^* X}.$$

Now by virtue of Theorem 4.2 (iii), we can construct an infinitesimal version of the boundary value morphism. In fact consider the morphism:

$$(5.3) \quad \hat{\mathcal{C}}_{NM} \longrightarrow \tau^! R\tau_!(\hat{\mathcal{C}}_{NM}) \simeq \tau^{-1}(\mathcal{C}_{N|X}|_{N \times_M T_M^* X})[d],$$

where $\tau: T_N M \times_M T_M^* X \rightarrow N \times_M T_M^* X$ denotes the canonical projection and $\mathcal{C}_{N|X}$ is the sheaf introduced in [19]. On account of the above theorem, we have constructed a morphism:

$$(5.4) \quad \varinjlim_{(1,0,\dots,0) \in \Omega'} \mathcal{C}_{\Omega|X} \longrightarrow \mathcal{C}_{N|X}[d]$$

on $N \times_M T_M^* X$. Furthermore, let \mathcal{M} be a coherent \mathcal{D}_X -module for which N is non-characteristic. Then [24][25] shows the injectivity of the microlocal boundary value morphism:

$$(5.5) \quad H^0 R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \hat{\mathcal{C}}_{NM}) \longrightarrow \tau^{-1} H^0 R\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{C}_{N|X}[d])|_{N \times_M T_M^* X}.$$

Remark 5.3. It gives the uniqueness of microlocal boundary value problems and generalizes the result of Schapira [20] to higher codimensions. It can also be considered as a microlocal extension of Oaku's theorem in [16]. However we have not yet shown the injectivity by using the framework developed in this paper.

Finally let us consider the complex \mathcal{C}_{NM} on $T_N^* M \times_M T_M^* X$. We can

easily show by Theorem 4.3 (ii) and Kashiwara’s abstract edge of the wedge theorem that it is concentrated in degree 0. Let π be the canonical projection $T_N^*M \times_M T_M^*X \rightarrow N \times_M T_M^*X$, and apply Sato’s distinguished triangle $R\pi_! \rightarrow R\pi_* \rightarrow R\tilde{\pi}_* \rightarrow_{+1}$ to the sheaf \mathcal{C}_{NM} . Then by Theorem 4.3 (iii) we get a distinguished triangle:

$$(5.6) \quad \mathcal{C}_M|_{N \times_M T_M^*X} \longrightarrow \mathcal{C}_{N|X}|_{N \times_M T_M^*X}[d] \longrightarrow R\tilde{\pi}_* \mathcal{C}_{NM}[d] \longrightarrow_{+1},$$

where \mathcal{C}_M denotes the sheaf of Sato’s microfunctions. It microlocalizes the distinguished triangle:

$$(5.7) \quad \mathcal{B}_M|_N \longrightarrow \Gamma_N \mathcal{B}_M|_N[d] \longrightarrow R\tilde{\pi}_{N*} \mu_N \mathcal{B}_M[d] \longrightarrow_{+1}$$

considered in Kashiwara-Kawai [6] by setting $\tilde{\pi}_N: \dot{T}_N^*M \rightarrow N$, and we believe that the sheaf \mathcal{C}_{NM} is a useful object when studying higher codimensional boundary value problems.

5.2. Second microfunctions associated to a foliation

Let M be a real analytic manifold of dimension n and X be its complexification as in the previous section. We assume the manifold M is endowed with an analytic foliation by leaves of dimension d and denote by L the union of their complexification in X . Hence, locally there exist a smooth surjective real analytic morphism $g: M \rightarrow M'$ and its complexification $g_{\mathbb{C}}: X \rightarrow X'$, and $L = g_{\mathbb{C}}^{-1}(M')$. We shall apply our theory to the triplet (X, L, M) . It follows from Theorem 4.3 (ii) and Kashiwara’s abstract edge of the wedge theorem that the complex $\mu_{ML}(\mathcal{O}_X)[n]$ on $T_M^*L \times_L T_L^*X$ is concentrated in degree 0.

Definition 5.4. We set:

$$(5.8) \quad \mathcal{C}_{ML} := \mu_{ML}(\mathcal{O}_X) \otimes_{\text{or}_M} [n]$$

and call it the sheaf of second microfunctions along L .

Now let π_M denote the projection $T_M^*L \times_L T_L^*X \rightarrow M \times_L T_L^*X$. Set $A := M \times_L T_L^*X \subset T_M^*X$ and denote by π the composition of π_M and the projection $A \rightarrow M$. We denote by $\mathcal{C}\mathcal{O}$ the sheaf of microfunctions with holomorphic parameters defined by:

$$(5.9) \quad \mathcal{C}\mathcal{O} := \mu_L(\mathcal{O}_X) \otimes_{\text{or}_L} [n - d].$$

Applying Sato’s distinguished triangle $R\pi_{M!} \rightarrow R\pi_{M*} \rightarrow R\tilde{\pi}_{M*} \rightarrow_{+1}$ to the sheaf \mathcal{C}_{ML} , we get:

Theorem 5.5. *There exists an exact sequence on $A = M \times_L T_L^*X$:*

$$(5.10) \quad 0 \longrightarrow \mathcal{C}\mathcal{O}|_A \longrightarrow \mathcal{C}_M|_A \longrightarrow \dot{\pi}_{M*} \mathcal{C}_{ML} \longrightarrow 0.$$

Hence we have functorially constructed a sheaf which decomposes the singularities of the microfunctions on A . Notice that the sheaf of second microfunctions of Kashiwara [8], that is $\mu_A(\mu_L(\mathcal{O}_X))[n]$ is different from ours. Also notice that the same sheaf as \mathcal{C}_{ML} was already obtained by Kataoka-Tose [13] making use of their second comonoidal transformation. Refer also to the paper of Kataoka-Tose-Okada [14]. However these authors did not construct it including the zero sections and their formulation was not functorial. In particular, it does not allow them to make use of algebraic machinery of \mathcal{D}_X -modules. An attempt to extend our construction to the third microlocalization was recently made by Watanabe [26].

Definition 5.6. We denote by sp_L the natural morphisms:

$$(5.11) \quad \mathcal{B}_M \xrightarrow{\sim} \pi_* \mathcal{C}_{ML}, \mathcal{C}_M|_A \xrightarrow{\sim} \pi_{M*} \mathcal{C}_{ML}.$$

If u is a hyperfunction on M (resp. a microfunction on A), we denote by $\text{ss}_L(u)$ the support of $\text{sp}_L(u)$ in $T_M^*L \times_L T_L^*X$.

From now on, we shall explain how the basic operations on second microfunctions can easily be deduced from the morphisms constructed in Section 4. Let $f: (Y, H, N) \rightarrow (X, L, M)$ be a morphism of triplets of manifolds. Here we assume that $f|_Y: Y \rightarrow X$ is a complexification of the real analytic morphism $f|_N: N \rightarrow M$, H is defined similarly to L and $f(H) \subset L$. Then we have canonical morphisms associated to f :

$$(5.12) \quad T_N^*H \times_H T_H^*Y \xleftarrow[\rho_f]{} (N \times_M T_M^*L) \times_L T_L^*X \xrightarrow[\varpi_f]{} T_M^*L \times_L T_L^*X.$$

The following theorem is a direct consequence of Proposition 4.5.

Theorem 5.7. *There exist canonical morphisms of “restriction” and “integration” of second microfunctions:*

$$(5.13) \quad \begin{cases} \rho_{f_!} \varpi_f^{-1} \mathcal{C}_{ML} \longrightarrow \mathcal{C}_{NH} \text{ and} \\ \varpi_{f_!} \rho_f^{-1} (\mathcal{C}_{NH} \otimes_{\mathcal{A}_N} \mathcal{V}_N) \longrightarrow \mathcal{C}_{ML} \otimes_{\mathcal{A}_M} \mathcal{V}_M, \end{cases}$$

where \mathcal{A} (resp. \mathcal{V}) denotes the sheaf of analytic functions (resp. analytic densities) on a real analytic manifold.

For example, assume N is a submanifold of M which is transversal to the leaves of the foliation of M , Y a complexification of N in X and $H = Y \cap L$. Set $A' := N \times_H T_H^*Y$. Then $A' \simeq N \times_M A$ and we have canonical morphisms:

$$(5.14) \quad T_N^*H \times_N A' \xleftarrow[\rho]{} (N \times_M T_M^*L) \times_M A \xrightarrow[\varpi]{} T_M^*L \times_M A,$$

where ρ is smooth and ϖ is a closed embedding. Hence we find that if $u \in \mathcal{C}_M|_A$ and $ss_L(u)$ is proper with respect to ρ , then $u|_N \in \mathcal{C}_N|_A$ is well-defined. It implies the possibility of defining a local restriction of microfunctions under a suitable condition on the second singular spectrum. It seems to be a new fact, because until now the restriction of microfunctions were always considered to be global.

Remark 5.8. The operations in Theorem 5.7 generalizes those obtained by Okada-Tose [17] for compactly supported hyperfunctions in a non functorial way. They defined them by means of the FBI transformation of [23]. Our construction ensures that these operations are coordinate invariant and well-defined for microfunctions.

Finally let us consider the “product” of second microfunctions. Let (X, L, M) be the triplet as before and take a coordinate system $(x, \sqrt{-1}\eta'dx', \sqrt{-1}\eta''dx'')$ of $T_M^*L \times_L T_L^*X$ in which $(x, \sqrt{-1}\eta'dx')$ belongs to T_M^*L . In this case, the morphism θ in (4.38) is the morphism:

$$(T_M^*L \times_L T_L^*X) \times_M (T_M^*L \times_L T_L^*X) \longrightarrow T_M^*L \times_L T_L^*X$$

which is described by

$$(5.15) \quad \begin{aligned} & ((x, \sqrt{-1}\eta'_1 dx', \sqrt{-1}\eta''_1 dx''), (x, \sqrt{-1}\eta'_2 dx', \sqrt{-1}\eta''_2 dx'')) \\ & \longmapsto (x, \sqrt{-1}(\eta'_1 + \eta'_2) dx', \sqrt{-1}(\eta''_1 + \eta''_2) dx''). \end{aligned}$$

Here we sometimes consider M as a submanifold of $T_M^*L \times_L T_L^*X$ by identifying it with the intersection of two zero sections, that is, $T_M^*L \cap (M \times_L T_L^*X)$. In general, we cannot make a product of two hyperfunctions u_1 and u_2 . In order to give a criterion which assures the possibility of the product operation, we prepare the next lemma.

Lemma 5.9. *Let S be a closed subset of $(T_M^*L \times_L T_L^*X) \times_M (T_M^*L \times_L T_L^*X)$ which is conic with respect to the four conic structures. Then the following two conditions are equivalent each other.*

- (i) θ is a proper mapping over S .
- (ii) θ satisfies:

$$(5.16) \quad \begin{cases} S \cap \theta^{-1}(T_M^*L) \subset (T_M^*L) \times_M (T_M^*L) \quad \text{and} \\ S \cap \theta^{-1}(M \times_L T_L^*X) \subset (M \times_L T_L^*X) \times_M (M \times_L T_L^*X). \end{cases}$$

Proof. It follows from the condition (ii) that:

$$(5.17) \quad S \cap \theta^{-1}(M) \subset M \times_M M \simeq M.$$

Hence θ is proper over S , because θ is a linear mapping w.r.t. the fibers. Next

assume the condition (i) and for example take a point:

$$(5.18) \quad p_0 := ((x, \sqrt{-1}\eta'_1 dx', \sqrt{-1}\eta''_1 dx''), (x, \sqrt{-1}\eta'_2 dx', \sqrt{-1}\eta''_2 dx''))$$

in $S \cap \theta^{-1}(M \times_L T_L^* X)$, that is, $\eta'_1 + \eta'_2 = 0$. Suppose p_0 does not belong to $(M \times_L T_L^* X) \times_M (M \times_L T_L^* X)$, then $\eta'_1 = -\eta'_2 \neq 0$. Since S is conic w.r.t. η''_1 and η''_2 variables, we get a point:

$$(5.19) \quad q_0 := ((x, \sqrt{-1}\eta'_1 dx', \sqrt{-1}0 dx''), (x, \sqrt{-1}\eta'_2 dx', \sqrt{-1}0 dx''))$$

which belongs to $[S \cap \theta^{-1}(M)] \setminus [M \times_M M]$. It contradicts the assumption (i). \blacksquare

Now we can give a new condition which ensures the existence of the product of two hyperfunctions.

Theorem 5.10. *Let u_1 and u_2 be hyperfunctions on M . Assume every point:*

$$(5.20) \quad ((x, \sqrt{-1}\eta'_1 dx', \sqrt{-1}\eta''_1 dx''), (x, \sqrt{-1}\eta'_2 dx', \sqrt{-1}\eta''_2 dx''))$$

in $ss_L(u_1) \times_M ss_L(u_2)$ satisfies the following conditions:

$$(5.21) \quad \begin{cases} \eta'_1 + \eta'_2 = 0 \implies \eta'_1 = \eta'_2 = 0 & \text{and} \\ \eta''_1 + \eta''_2 = 0 \implies \eta''_1 = \eta''_2 = 0. \end{cases}$$

Then there exists a “product” $u_1 \cdot u_2$ of u_1 and u_2 , and

$$(5.22) \quad ss_L(u_1 \cdot u_2) \subset \theta[ss_L(u_1) \times_M ss_L(u_2)].$$

Proof. It is an immediate consequence of Proposition 4.7 and Lemma 5.9. \blacksquare

5.3. Second microdifferential operators

In this section, we shall explain some ideas in order to develop the theory of second microdifferential operators which act on our microfunctions. Notice that the class of operators of Laurent [15] seems to be too wide to operate on the sheaf \mathcal{C}_{ML} . First, let (X, Y, Z) be a triplet of complex manifolds and assume Z is of complex codimension d in X . Then we will show the result below in a forthcoming paper.

Proposition 5.11. *The complex $\mu_{ZY}(\mathcal{O}_X)[d]$ of sheaves on $T_Z^* Y \times_Y T_Y^* X$ is concentrated in degree 0, and we define the sheaf $\mathcal{C}_{ZY|X}^{\mathbb{R}}$ of second holomorphic microfunctions by:*

$$(5.23) \quad \mathcal{C}_{ZY|X}^{\mathbb{R}} := \mu_{ZY}(\mathcal{O}_X)[d].$$

It is possible to write down the various functorial properties of second holomorphic microfunctions as in [19], but we shall not develop it here. Let

$X = X' \times X''$ be a product of two complex manifolds. We denote by $\Delta_X = \Delta_{X'} \times \Delta_{X''}$ the diagonal set of $X \times X$ and set $\hat{L} := X' \times X' \times \Delta_{X''}$ in $X \times X$. Let us apply our functor to the triplet $(X \times X, \hat{L}, \Delta_X)$ of complex manifolds.

Definition 5.12. We define the sheaf $\mathcal{E}_{\hat{X}L}^{\mathbb{R}}$ of second microdifferential operators on

$$(5.24) \quad T_{\Delta_X}^* \hat{L} \times_{\hat{L}} T_{\hat{L}}^*(X \times X) \simeq T^*X' \times T^*X''$$

by $\mathcal{E}_{\hat{X}L}^{\mathbb{R}} := \mu_{\Delta_X \hat{L}}(\mathcal{O}_{X \times X}^{(0,n)})[n]$, where $n = \dim_{\mathbb{C}} X$ and $\mathcal{O}_{X \times X}^{(0,n)}$ denotes the sheaf of holomorphic differential forms of maximal degree w.r.t. the second factor.

We can easily verify that the sheaf $\mathcal{E}_{\hat{X}L}^{\mathbb{R}}$ is endowed with a ring structure by the results of Kashiwara-Schapira [9]. Moreover the restriction of the sheaf $\mathcal{E}_{\hat{X}L}^{\mathbb{R}}$ to the zero section

$$(5.25) \quad \Delta_X \times_{\hat{L}} T_{\hat{L}}^*(X \times X) \simeq X' \times T^*X''$$

coincides with the restriction of the sheaf $\mathcal{E}_X^{\mathbb{R}}$ of holomorphic microdifferential operators to it. This is an explicit advantage of our theory. Now let $M = M' \times M''$ be a product of two real analytic manifolds, $X = X' \times X''$ a complexification and $L := X' \times M''$ a partial complexification of $M = M' \times M''$ in X . We shall consider the action of the Ring $\mathcal{E}_{\hat{X}L}^{\mathbb{R}}$ on the sheaf \mathcal{C}_{ML} of second microfunctions along L . For this purpose, we perform a so-called S-K-K realification. To begin with, define the sheaf $\mathcal{C}_{(M \times M)(L \times L)}$ of second microfunctions on

$$(5.26) \quad T_{(M \times M)(L \times L)}^* \times_{(L \times L)} T_{(L \times L)}^*(X \times X) \simeq T^*(M' \times M') \times T^*(M'' \times M'')$$

by:

$$(5.27) \quad \mathcal{C}_{(M \times M)(L \times L)} := \mu_{(M \times M)(L \times L)}(\mathcal{O}_{X \times X}) \otimes \text{or}_{M \times M}[2n].$$

Then we have:

Theorem 5.13. *There exists a canonical morphism of “realification”:*

$$(5.28) \quad \mathcal{E}_{\hat{X}L}^{\mathbb{R}}|_{T_{M'}^*X' \times T_{M''}^*X''} \longrightarrow \Gamma_E \mathcal{C}_{(M \times M)(L \times L)} \otimes_{\mathcal{A}_M} \mathcal{V}_M,$$

where we set:

$$(5.29) \quad \begin{aligned} E &:= T_{\Delta_{M'}}^*(M' \times M') \times T_{\Delta_{M''}}^*(M'' \times M'') \\ &\simeq T^*M' \times T^*M'' \simeq T_{M'}^*X' \times T_{M''}^*X''. \end{aligned}$$

Proof. The intersection of the triplets $(X \times X, \hat{L}, \Delta_X)$ and $(X \times X, L \times L, M \times M)$ is equal to

$$(5.30) \quad (X \times X, \hat{L} \cap (L \times L), \Delta_X \cap (M \times M)) = (X \times X, L', \Delta_M)$$

by setting $L' := X' \times X' \times \Delta_{M''}$. Therefore there exist morphisms of triplets of manifolds:

$$(5.31) \quad \begin{cases} f := \text{id}_{X \times X}: (X \times X, L', \Delta_M) \longrightarrow (X \times X, \hat{L}, \Delta_X), \\ g := \text{id}_{X \times X}: (X \times X, L', \Delta_M) \longrightarrow (X \times X, L \times L, M \times M). \end{cases}$$

By virtue of the results in Section 4.2, we have associated injective morphisms:

$$(5.32) \quad \begin{aligned} T_{\Delta_M}^* L' \times_{L'} T_{L'}^*(X \times X) &\xleftarrow{\rho_f} \Delta_M \times_{\Delta_X} [T_{\Delta_X}^* \hat{L} \times_{\hat{L}} T_{\hat{L}}^*(X \times X)] \\ &\longrightarrow T_{\Delta_X}^* \hat{L} \times_{\hat{L}} T_{\hat{L}}^*(X \times X) \\ &\quad \varpi_f \end{aligned}$$

and

$$(5.33) \quad \begin{aligned} T_{\Delta_M}^* L' \times_{L'} T_{L'}^*(X \times X) &\xleftarrow{\rho_g} \Delta_M \times_{(M \times M)} [T_{(M \times M)}^*(L \times L) \times_{(L \times L)} T_{(L \times L)}^*(X \times X)] \\ &\longrightarrow T_{(M \times M)}^*(L \times L) \times_{(L \times L)} T_{(L \times L)}^*(X \times X). \end{aligned}$$

The next lemma is a simple corollary of Proposition 4.5.

Lemma 5.14. *Let $F \in \mathbf{D}^b(X \times X)$. Then there exist canonical morphisms:*

$$(5.34) \quad \begin{cases} \varpi_f^{-1} \mu_{\Delta_X \hat{L}}(F) \longrightarrow \rho_f^1 \mu_{\Delta_M L'}(F) \otimes \text{or}_M[n], \\ \rho_g^{-1} \mu_{\Delta_M L'}(F) \longrightarrow \varpi_g^1 \mu_{(M \times M)(L \times L)}(F). \end{cases}$$

Now let us set:

$$(5.35) \quad \begin{aligned} E &:= \Delta_M \times_{\Delta_X} [T_{\Delta_X}^* \hat{L} \times_{\hat{L}} T_{\hat{L}}^*(X \times X)] \\ &\cap \Delta_M \times_{(M \times M)} [T_{(M \times M)}^*(L \times L) \times_{(L \times L)} T_{(L \times L)}^*(X \times X)] \\ &\simeq T_{\Delta_{M'}}^*(M' \times M') \times T_{\Delta_{M''}}^*(M'' \times M'') \end{aligned}$$

in $T_{\Delta_M}^* L' \times_{L'} T_{L'}^*(X \times X)$, and consider the Cartesian square below:

$$(5.36) \quad \begin{array}{ccc} E & \xrightarrow{i_f} & \Delta_M \times_{(M \times M)} [T_{(M \times M)}^*(L \times L) \times_{(L \times L)} T_{(L \times L)}^*(X \times X)] \\ \downarrow i_g & & \downarrow \rho_g \\ \Delta_M \times_{\Delta_X} [T_{\Delta_X}^* \hat{L} \times_{\hat{L}} T_{\hat{L}}^*(X \times X)] & \xrightarrow{\rho_f} & T_{\Delta_M}^* L' \times_{L'} T_{L'}^*(X \times X) \end{array}$$

in which all morphisms are injective. Therefore we get a chain of morphisms:

$$\begin{aligned}
 \mu_{\Delta_X} \hat{\mathcal{L}}(\mathcal{O}_{X \times X})|_E[n] &= \iota_g^{-1} \varpi_f^{-1} \mu_{\Delta_X} \hat{\mathcal{L}}(\mathcal{O}_{X \times X})[n] \\
 &\rightarrow \iota_g^{-1} \rho_f^! \mu_{\Delta_M L}(\mathcal{O}_{X \times X}) \otimes \text{or}_M[2n] \\
 (5.37) \quad &\rightarrow \iota_f^! \rho_g^{-1} \mu_{\Delta_M L}(\mathcal{O}_{X \times X}) \otimes \text{or}_M[2n] \\
 &\rightarrow \iota_f^! \varpi_g^! \mu_{(M \times M)(L \times L)}(\mathcal{O}_{X \times X}) \otimes \text{or}_M[2n] \\
 &\simeq R\Gamma_E \mu_{(M \times M)(L \times L)}(\mathcal{O}_{X \times X}) \otimes \text{or}_M[2n]
 \end{aligned}$$

due to the above lemma, and the proof of the theorem is complete. ■

Theorem 5.13 asserts that our second microdifferential operators are transformed to second microlocal operators, i.e. elements of $\Gamma_E \mathcal{C}_{(M \times M)(L \times L)} \otimes_{\mathcal{S}_M} \mathcal{V}_M$. The action of second microlocal operators on the sheaf \mathcal{C}_{ML} can be easily obtained by making use of Theorem 5.7 and Theorem 5.10 along the same lines as in Sato-Kawai-Kashiwara [19], and we shall omit precise arguments here.

References

- [1] Bost, J-B., Deformation to the normal cone and index theorem, *Exposé au congrès satellite ECM*, Université Paris Nord, 30 juin, 1992.
- [2] Delort, J-M., Microlocalisation simultanée et problème de Cauchy ramifié, *Compositio Math.*, to appear.
- [3] Ishimura, R., Le foncteur μhom relatif et l'action de \mathcal{E}^R relatif sur les fonctions holomorphes, *Mem. Fac. Sci. Kyushu Univ.*, Ser A46 (1992), 69–77.
- [4] Kashiwara, M., Deuxième microlocalisation et théorème du “watermelon”, *Talks at Nice*, unpublished, 1972.
- [5] ———, Systems of microdifferential equations, *Progr. Math.* 34, Birkhäuser, Boston, 1983.
- [6] Kashiwara, M. and Kawai, T., On the boundary value problem for elliptic systems of linear partial differential equations I-II, *Proc. Japan. Acad.*, 48 (1971), 712–715; *ibid.*, 49 (1972), 164–168.
- [7] ———, Second microlocalization and asymptotic expansions, *LN in Physics* 126, Proc. Les Houches, D. Iagolnitzer editor, Springer-Verlag, 79 (1979), 21–76.
- [8] Kashiwara, M. and Laurent, Y., Théorèmes d’annulation et deuxième microlocalisation, *prépublication d’Orsay*, 1983.
- [9] Kashiwara, M. and Schapira, P., Micro-hyperbolic systems, *Acta Math.*, 142 (1979), 1–55.
- [10] ———, Microlocal study of sheaves, *Astérisque*, 128 (1985).
- [11] ———, *Sheaves on manifolds*, Grundlehren der Math. Wiss. 292, Springer-Verlag, 1990.
- [12] Kataoka, K., On the theory of Radon transformations of hyperfunctions, *J. Fac. Sci. Univ. Tokyo*, 28 (1981), 331–413.
- [13] Kataoka, K. and Tose, N., Some remarks in 2nd microlocalization (in Japanese), *RIMS Kôkyu-roku, Kyoto Univ.*, 660 (1988), 52–63.
- [14] Kataoka, K., Tose, N. and Okada, Y., Decomposition of second microlocal analytic singularities, *D-Modules and Microlocal Geometry*, M. Kashiwara, T. Monteiro-Fernandès and P. Schapira editors., De Gruyter, (1993), 163–171.
- [15] Laurent, Y., *Théorie de la deuxième microlocalisation dans le domaine complexe*, Progress in Math., Birkhäuser, 1985.

- [16] Oaku, T., Higher-codimensional boundary value problem and F -mild hyperfunctions, *Algebraic Analysis, II*, (Papers dedicated to Prof. Sato), M. Kashiwara and T. Kawai editors., Academic Press, (1988), 571–586.
- [17] Okada, Y. and Tose, N., FBI-transformation and microlocalization-equivalence of the second analytic wave front sets and the second singular spectrum, *J. Math. Pures Appl.*, **70.4** (1991), 427–455.
- [18] Sabbah, C., Quelques remarques sur la géométrie des cônes conormaux, *Astérisque*, **130** (1985), 161–192.
- [19] Sato, M., Kawai, T. and Kashiwara, M., Hyperfunctions and pseudodifferential equations, *LN in Math*, Springer-Verlag, **287** (1973), 265–529.
- [20] Schapira, P., Microfunctions for boundary value problems, *Algebraic Analysis, II*, (Papers dedicated to Prof. Sato), M. Kashiwara and T. Kawai editors., Academic Press, (1988) 809–819.
- [21] Schapira, P. and Schneiders, J-P., Index theorem for elliptic pairs, *Astérisque*, **224** (1994).
- [22] Schapira, P. and Takeuchi, K., Déformation binormale et bispécialisation, *C.R. Acad. Sci.*, **319**, Série I (1994), 707–712.
- [23] Sjöstrand, J., Singularités analytiques microlocales, *Astérisque*, **95** (1982).
- [24] Takeuchi, K., Étude microlocale des problèmes aux limites en codimension supérieure, *C.R. Acad. Sci.*, **320**, Série I (1995), 441–443.
- [25] ———, Microlocal boundary value problem in higher codimensions, to appear in *Bull. Soc. Math. France*.
- [26] Watanabe, S., On third microlocal analysis, *Master thesis presented to University of Chiba.*, 1995.