

Quantum Toroidal Algebras and Their Vertex Representations

By

Yoshihisa SAITO*

Abstract

We construct the vertex representations of the quantum toroidal algebras $U_q(\widehat{\mathfrak{sl}}_{n+1,tor})$. In the classical case the vertex representations are not irreducible. However in the quantum case they are irreducible.

§1. Introduction

The classical toroidal algebras have been studied by many authors [MEY], [S], [Y], etc. Here “classical” means $q = 1$. The definition of the quantum toroidal algebras is given in [GKV]. They gave a geometric realization of the quantum toroidal algebras without any results on their representation theory. Recently Varagnolo and Vasserot [VV] proved Schur-type duality between representations of the quantum toroidal algebras and the double affine Hecke algebra introduced by Cherednik [C]. This is an analogue of the duality between the quantum affine algebras and the affine Hecke algebras given by Chari and Pressly [CP]. In [VV] only the representations of “trivial central charge” was studied. It is known that there are two subalgebras $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ and $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ of $U_q(\widehat{\mathfrak{sl}}_{n+1,tor})$ such that there are surjective algebra homomorphisms $U_q(\widehat{\mathfrak{sl}}_{n+1}) \rightarrow U_q^{(i)}(\widehat{\mathfrak{sl}}_{n+1})$ for $i=1,2$. In this paper we say that M has a level $(0,0)$ instead of the trivial central charge. The first 0 means that M has a level 0 as a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module and the second 0 means that M has a level 0 as a $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ -module. This notation is an analogue of level 0 representations of the affine quantum algebras.

In this paper we try to consider an analogue of “integrable representations”

Communicated by T. Miwa, June 12, 1997

1991 Mathematics Subject Classification(s): 17B37, 81R50

*Research Institute for Mathematical Sciences, Kyoto University, Japan.

Current Address: Department of Mathematics, University of Hiroshima, Higashi-Hiroshima, Hiroshima, 739-8526, Japan.

The author is supported by the JSPS Research Fellowships for Young Scientists

in the toroidal case. Let us recall the integrability of quantum Kac-Moody modules. Let $U_q(\mathfrak{g})$ be a quantum Kac-Moody algebra and V a $U_q(\mathfrak{g})$ -module. We say V is integrable if V has a weight space decomposition and locally nilpotent actions of the Chevalley generators of $U_q(\mathfrak{g})$. Therefore the definition of “integrability” needs Chevalley-type generators. The toroidal algebras is defined through Drinfeld type of generators and its Chevalley-type generators are not known. Therefore we are not able to define the integrability at this moment. However, in the affine case, Frenkel-Jing [FJ] realized the integrable representations with level 1 by the vertex representations. Thus if there are “vertex representations” of quantum toroidal algebras, they must be interesting example of the integrable representations still not defined. In the $q = 1$ case, vertex representations of the toroidal algebras have been already considered by Moody-Eswara Rao-Yokonuma [MEY]. In this paper we construct the q -analogue of the representations defined by them for $\mathfrak{g} = \mathfrak{sl}_{n+1}$ with level (1,0) and (1,1). Therefore we give a new class of the representations of the quantum toroidal algebras. In the $q = 1$ case, the Fock modules are not irreducible over the Heisenberg algebra and the vertex representations are not irreducible over the toroidal algebra. In the quantum case, it is not the case: the Fock modules are irreducible and also the vertex representations with level (1,0) are irreducible.

The algebra $U_q(\mathfrak{g}_{tor})$ has infinitely many generators satisfying infinitely many relations (See §2). It is preferable that $U_q(\mathfrak{g}_{tor})$ is written by finitely many generators with finitely many relations. According to [GKV] there are finitely many generators of $U_q(\mathfrak{sl}_{2,tor})$ but the relations among these generators are highly non-trivial. In this paper we give an explicit form of finitely many generators of $U_q(\mathfrak{sl}_{2,tor})$ and closed relations of them (See §4). They coincide with the generators by Vasserot [V].

§2. Definition of Quantum Toroidal Algebras

2.1. Notations. Let \mathfrak{g} be a complex semisimple Lie algebra of type A_n and $\hat{\mathfrak{g}}$ an affine Kac-Moody Lie algebra of type $A_n^{(1)}$. We denote their Cartan subalgebras by \mathfrak{h} and $\hat{\mathfrak{h}}$ respectively. We denote by $\bar{\alpha}_1, \dots, \bar{\alpha}_n$ the simple roots of \mathfrak{g} , by $\bar{h}_1, \dots, \bar{h}_n$ the simple coroots of \mathfrak{g} , by $\bar{\Lambda}_1, \dots, \bar{\Lambda}_n$ the fundamental weights of \mathfrak{g} , by $\alpha_0, \dots, \alpha_n$ the simple roots of $\hat{\mathfrak{g}}$, by h_0, \dots, h_n the simple coroots of $\hat{\mathfrak{g}}$ and $\Lambda_0, \dots, \Lambda_n$ the fundamental weights of $\hat{\mathfrak{g}}$. Let $\bar{Q} = \bigoplus_{i=1}^n \mathbb{Z}\bar{\alpha}_i$ be the root lattice of \mathfrak{g} , $\bar{P} = \bigoplus_{i=1}^n \mathbb{Z}\bar{\Lambda}_i$ the weight lattice of \mathfrak{g} , $Q = \bigoplus_{i=0}^n \mathbb{Z}\alpha_i$ the root lattice of $\hat{\mathfrak{g}}$ and $P = \bigoplus_{i=0}^n \mathbb{Z}\Lambda_i \oplus \mathbb{Z}\delta$ the weight lattice of $\hat{\mathfrak{g}}$. Here δ is the null root.

We denote the pairing of \mathfrak{h} and \mathfrak{h}^* (resp. $\hat{\mathfrak{h}}$ and $\hat{\mathfrak{h}}^*$) by \langle, \rangle . The invariant bilinear form on P is given by $\langle \alpha_i | \alpha_j \rangle = -\delta_{ij-1} + 2\delta_{ij} - \delta_{ij+1}$ and $\langle \delta | \delta \rangle = 0$. The projection form P to \bar{P} is given by $\bar{\Lambda}_i = \Lambda_i - \Lambda_0$ and $\bar{\delta} = 0$.

2.2. We will give the definition of the quantum toroidal algebra $U_q(\mathfrak{g}_{tor})$.

Definition 2.2.1. Let $M = (m_{ij})_{0 \leq i, j \leq n}$ be a skew-symmetric $(n+1) \times (n+1)$ -matrix with integral coefficients and let κ be an element of $\mathbb{Q}(q)^*$. $U_q(\mathfrak{g}_{tor})$ is an associated algebra over $\mathbb{Q}(q)$ with generators:

$$E_{i,k}, F_{i,k}, H_{i,l}, K_i^\pm, q^{\pm \frac{1}{2}c}, q^{\pm d_1}, q^{\pm d_2},$$

for $k \in \mathbb{Z}, l \in \mathbb{Z} \setminus \{0\}$ and $i = 0, 1, \dots, n$.

We introduce $K_{i,k}^\pm$ as the Fourier components of the following generating functions:

$$K_i^+(z) = \sum_{k \geq 0} K_{i,k}^+ z^{-k} = K_i^+ \exp\left((q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}\right),$$

$$K_i^-(z) = \sum_{k \leq 0} K_{i,k}^- z^{-k} = K_i^- \exp\left(- (q - q^{-1}) \sum_{k \geq 1} H_{i,-k} z^k\right).$$

The defining relations of $U_q(\mathfrak{g}_{tor})$ are then written as follows:

$$(2.2.1) \quad q^{\pm \frac{1}{2}c} \text{ are central,}$$

$$(2.2.2) \quad K_i^+ K_i^- = K_i^- K_i^+ = 1.$$

$$(2.2.3) \quad [K_i^\pm, K_j^\pm] = 0,$$

$$(2.2.4) \quad [K_i^\pm, H_{j,l}] = 0,$$

$$(2.2.5) \quad [H_{i,k}, H_{j,l}] = \delta_{k+l,0} \frac{1}{k} [k \langle h_i, \alpha_j \rangle] \frac{q^{kc} - q^{-kc}}{q - q^{-1}} \kappa^{-km_i},$$

$$(2.2.6) \quad [q^{\pm d_i}, K_j^\pm] = 0,$$

$$(2.2.7) \quad q^{d_1} H_{j,l} q^{-d_1} = q^l H_{j,l},$$

$$(2.2.8) \quad [q^{\pm d_2}, H_{j,l}] = 0,$$

$$(2.2.9) \quad q^{d_1} E_{j,k} q^{-d_1} = q^k E_{j,k},$$

$$q^{d_1} F_{j,k} q^{-d_1} = q^k F_{j,k},$$

$$(2.2.10) \quad q^{d_2} E_{j,k} q^{-d_2} = q^{\delta_{j0}} E_{j,k},$$

$$q^{d_2} F_{j,k} q^{-d_2} = q^{-\delta_{j0}} F_{j,k},$$

$$(2.2.11) \quad K_i^+ E_{j,k} K_i^- = q^{\langle h_i, \alpha_j \rangle} E_{j,k},$$

$$K_i^+ F_{j,k} K_i^- = q^{-\langle h_i, \alpha_j \rangle} F_{j,k},$$

$$(2.2.12) \quad [H_{i,k}, E_{j,l}] = \frac{1}{k} [k \langle h_i, \alpha_j \rangle] q^{-\frac{1}{2} |k|c} \kappa^{-km_i} E_{j,k+l},$$

$$(2.2.13) \quad [H_{i,k}, F_{j,l}] = -\frac{1}{k} [k \langle h_i, \alpha_j \rangle] q^{\frac{1}{2}|k|c} \kappa^{-kmu} F_{j,k+l},$$

$$\kappa^{mu} E_{i,k+1} E_{j,l} - q^{\langle h_i, \alpha_j \rangle} \kappa^{mu} E_{j,l} E_{i,k+1} = q^{\langle h_i, \alpha_j \rangle} E_{i,k} E_{j,l+1} - E_{j,l+1} E_{i,k},$$

$$\kappa^{mu} F_{i,k+1} F_{j,l} - q^{-\langle h_i, \alpha_j \rangle} \kappa^{mu} F_{j,l} F_{i,k+1} = q^{-\langle h_i, \alpha_j \rangle} F_{i,k} F_{j,l+1} - F_{j,l+1} F_{i,k},$$

$$(2.2.14) \quad [E_{i,k}, F_{j,l}] = \delta_{i,j} \frac{1}{q - q^{-1}} \{ q^{\frac{1}{2}(k-l)c} K_{i,k+l}^+ - q^{\frac{1}{2}(l-k)c} K_{i,k+l}^- \},$$

$$(2.2.15) \quad \sum_{\sigma \in \mathfrak{S}_m} \sum_{r=0}^m (-1)^r \begin{bmatrix} m \\ r \end{bmatrix} E_{i,k_{\sigma(1)}} \dots E_{i,k_{\sigma(r)}} E_{j,l} E_{i,k_{\sigma(r+1)}} \dots E_{i,k_{\sigma(m)}} = 0,$$

$$\sum_{\sigma \in \mathfrak{S}_m} \sum_{r=0}^m (-1)^r \begin{bmatrix} r \\ m \end{bmatrix} F_{i,k_{\sigma(1)}} \dots F_{i,k_{\sigma(r)}} F_{j,l} F_{i,k_{\sigma(r+1)}} \dots F_{i,k_{\sigma(m)}} = 0,$$

for $i \neq j$,

where $m = 1 - \langle h_i, \alpha_j \rangle$.

In these relations we denote $[k] = \frac{q^k - q^{-k}}{q - q^{-1}}$, $[n]! = \prod_{k=1}^n [k]$, $\begin{bmatrix} m \\ r \end{bmatrix} = \frac{[m]!}{[r]! [m-r]!}$.

2.3. Let $U'_q(\mathfrak{g}_{tor})$ be the subalgebra of $U_q(\mathfrak{g}_{tor})$ generated by $E_{i,k}, F_{i,k}, K_i^\pm, H_{i,l}, q^{\pm \frac{1}{2}c}$. Let $U_q^{\phi'}(\mathfrak{g}_{tor})$ (resp. $U_q^{\phi}(\mathfrak{g}_{tor})$) be the subalgebra generated by $U'_q(\mathfrak{g}_{tor})$ and $q^{\pm d_1}$ (resp. $q^{\pm d_2}$). Let $U_q^{(1)'}(\widehat{\mathfrak{sl}}_{n+1})$ be the subalgebra generated by $E_{i,k}, F_{i,k}, K_i^\pm, H_{i,l}, q^{\pm \frac{1}{2}c}$ ($1 \leq i \leq n, k \in \mathbb{Z}, l \in \mathbb{Z} \setminus \{0\}$) and $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ the subalgebra generated by $U_q^{(1)'}(\widehat{\mathfrak{sl}}_{n+1})$ and $q^{\pm d_1}$. Let $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ the subalgebra generated by $E_{i,0}, F_{i,0}, K_i^\pm$ ($0 \leq i \leq n$) and $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ the subalgebra generated by $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ and $q^{\pm d_2}$. By the definition it is clear that there are surjective homomorphisms $U_q(\widehat{\mathfrak{sl}}_{n+1}) \rightarrow U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ and $U'_q(\widehat{\mathfrak{sl}}_{n+1}) \rightarrow U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$.

The following are straightforward.

Lemma 2.3.1. For $1 \leq i \leq n$ let $\overline{E_{i,k}} = E_{i,k} \kappa^{\sum_{j=1}^{i-1} m_j - l}$, $\overline{F_{i,k}} = F_{i,k} \kappa^{\sum_{j=1}^{i-1} m_j - l}$, $\overline{H_{i,l}} = H_{i,l} \kappa^{\sum_{j=1}^{i-1} m_j - l}$. Then the relations between $\overline{E_{i,k}}, \overline{F_{i,k}}, \overline{H_{i,l}}$ and K_i^\pm are precisely the relations of Drinfeld generators of $U_q(\widehat{\mathfrak{sl}}_{n+1})$. That is, there are surjective homomorphisms $U_q(\widehat{\mathfrak{sl}}_{n+1}) \rightarrow U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ and $U'_q(\widehat{\mathfrak{sl}}_{n+1}) \rightarrow U_q^{(1)'}(\widehat{\mathfrak{sl}}_{n+1})$.

Lemma 2.3.2. Let $K_{\mathfrak{F}}^\pm = \prod_{i=0}^n K_i^\pm$. Then $K_{\mathfrak{F}}^\pm$ are central elements of $U_q(\mathfrak{g}_{tor})$.

Note that $q^{\pm c}$ is the central elements of $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ and $K_{\mathfrak{F}}^\pm$ the central

elements of $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$.

§3. Vertex Representations

3.1. Heisenberg algebras. In this section we shall give the vertex representations of $U_q(\mathfrak{g}_{tor})$. We assume $c=1$.

Consider a $\mathbf{Q}(q)$ -algebra S_n generated by $H_{i,l}$ ($0 \leq i \leq n, l \in \mathbf{Z} \setminus \{0\}$) satisfying:

$$(3.1.1) \quad [H_{i,k}, H_{j,l}] = \delta_{k+l,0} \frac{1}{k} [k \langle h_i, \alpha_j \rangle] \frac{q^k - q^{-k}}{q - q^{-1}} \kappa^{-km_i}.$$

We call S_n the Heisenberg algebra.

Let S_n^+ (resp. S_n^-) be the subalgebra of S_n generated by $H_{i,l}$ ($0 \leq i \leq n, l > 0$) (resp. $0 \leq i \leq n, l < 0$).

We introduce the Fock space

$$\mathcal{F}_n = S_n v_0$$

with the defining relations:

$$(3.1.2) \quad H_{i,l} v_0 = 0, \quad \text{for } l > 0,$$

$$(3.1.3) \quad q^{\frac{1}{2} \epsilon} v_0 = q^{\frac{1}{2}} v_0.$$

Note that \mathcal{F}_n is a free S_n^- -module of rank 1.

Let \mathbf{F} be a field of characteristic zero and let \mathfrak{a} be an associative \mathbf{F} -algebra generated by x_p, y_p ($p \in \mathbf{Z}_{>0}$), z and its inverse z^{-1} with the following relations:

$$[x_p, z] = [y_p, z] = 0,$$

$$[x_p, x_r] = [y_p, y_r] = 0,$$

$$[x_p, y_r] = \delta_{pr} z.$$

Let \mathfrak{a}^+ (resp. \mathfrak{a}^-) be the subalgebra of \mathfrak{a} generated by x_p (resp. y_p). We set $\mathfrak{b} = \mathfrak{a}^+ \otimes_{\mathbf{F}} \mathbf{F}[z, z^{-1}]$. This is a maximal abelian subalgebra of \mathfrak{a} . Fix a nonzero scalar $\lambda \in \mathbf{F}^\times$. Let \mathbf{F}_λ be the one-dimensional space \mathbf{F} viewed as a \mathfrak{b} -module by:

$$z \cdot 1 = \lambda, \quad \mathfrak{a}^+ \cdot 1 = 0.$$

Let $F(\lambda)$ be the induced \mathfrak{a} -module

$$F(\lambda) = \text{Ind}_{\mathfrak{b}}^{\mathfrak{a}} \mathbf{F}_\lambda = \mathfrak{a} \otimes_{\mathfrak{b}} \mathbf{F}_\lambda.$$

By the defining relations of \mathfrak{a} we obtain an \mathbf{F} -linear isomorphism

$$F(\lambda) \cong \mathfrak{a}^-.$$

Since \mathfrak{a}^- is abelian we may regard it as the algebra of polynomials in the variables y_1, y_2, \dots . Then we see that z acts on $\mathfrak{a}^- \cong \mathbb{F}[y_1, y_2, \dots]$ by the multiplication of λ , x_p acts by $\lambda \frac{\partial}{\partial y_p}$. By this realization we immediately have the following lemma.

Lemma 3.1.1. *$F(\lambda)$ is an irreducible \mathfrak{a} -module.*

Fix an skew-symmetric $(n+1) \times (n+1)$ -matrix with integral coefficients $M = (m_{ij})_{0 \leq i, j \leq n}$. We say that $\kappa \in \mathbb{Q}(q)^*$ is generic with respect to M if for any $k \in \mathbb{Z}_{>0}$ the matrix $([k \langle h_i, \alpha_j \rangle] \kappa^{-km_{ij}})$ is invertible.

Note that if $n = 1$ any κ is not generic with respect to any M . Since the matrix $([k \langle h_i, \alpha_j \rangle])_{0 \leq i, j \leq n}$ is invertible for $n > 1$, there exists a generic κ for $n > 1$.

Lemma 3.1.2. (1) *For a fixed M , we assume that $\kappa \in \mathbb{Q}(q)^*$ is generic with respect to M (in particular $n > 1$). Then \mathcal{F}_n is an irreducible S_n -module.*
 (2) *\mathcal{F}_1 is not irreducible.*

Proof. (1) Set $G(k) = (g(k)_{ij}) ([k \langle h_i, \alpha_j \rangle] \kappa^{-km_{ij}})$. Since κ is generic with respect to M there exists its inverse $G(k)^{-1} = (g(k)^{ij})$ for any k . Note that by the definition $\sum_{0 \leq s \leq n} g(k)^{is} g(k)_{sj} = \delta_{ij}$.

We set

$$\tilde{H}_{i,k} = \begin{cases} \sum_{0 \leq s \leq n} \frac{k}{[k]} g(k)^{is} H_{s,k}, & \text{for } k > 0, \\ H_{i,k}, & \text{for } k < 0. \end{cases}$$

Then we have

$$[\tilde{H}_{i,k}, \tilde{H}_{j,l}] = [\tilde{H}_{i,-k}, \tilde{H}_{j,-l}] = 0, \\ [\tilde{H}_{i,k}, \tilde{H}_{j,-l}] = \delta_{k,l} \delta_{ij},$$

for $k, l > 0$. Since all $G(k)$ are regular, $\tilde{H}_{i,k} (0 \leq i \leq n, k > 0)$ generate S_n^+ .

We shall use Lemma 3.1.1. Put $\mathbb{F} = \mathbb{Q}(q)$, $\mathfrak{a} = S_n$, $\mathfrak{a}^\pm = S_n^\pm$, $\lambda = 1$, $x_p = \tilde{H}_{i,k} (k > 0)$, $y_r = \tilde{H}_{j,l} (l < 0)$ where $p = (k-1)(n+1) + i + 1$ and $r = (-l-1)(n+1) + j + 1$.

Then it is clear that $F(1) = \mathcal{F}_n$. By Lemma 3.1.1 we conclude that \mathcal{F}_n is an irreducible S_n -module.

(2) It is easy to see that $\kappa^{-km_{0i}} H_{0,-k} + H_{1,-k}$ is a central element of S_1 for each $k \in \mathbb{Z}_{>0}$. Therefore \mathcal{F}_1 has infinitely many singular vectors. \square

3.2. Construction of level $(1,0)$ modules. In this subsection we assume $\kappa = 1$. Note that $1 \in \mathbb{Q}(q)$ is generic with respect to any M with $n > 1$.

Note that one can rewrite $\bar{P} = \bigoplus_{i=2}^n \mathbf{Z} \bar{\alpha}_i \oplus \mathbf{Z} \bar{\Lambda}_n$. We introduce a twisted version of the group algebra $\mathbf{Q}(q)[\bar{P}]$ by $\mathbf{Z}/2\mathbf{Z}$. We denote it by $\mathbf{Q}(q)[\bar{P}]$. This is the $\mathbf{Q}(q)$ -algebra generated by symbols $e^{\bar{\alpha}_2}, e^{\bar{\alpha}_3}, \dots, e^{\bar{\alpha}_n}, e^{\bar{\Lambda}_n}$ which satisfy the following relations:

$$(3.2.1) \quad e^{\bar{\alpha}_i} e^{\bar{\alpha}_j} = (-1)^{\langle \bar{h}_i, \bar{\alpha}_j \rangle} e^{\bar{\alpha}_j} e^{\bar{\alpha}_i}$$

$$(3.2.2) \quad e^{\bar{\alpha}_i} e^{\bar{\Lambda}_n} = (-1)^{\delta_{in}} e^{\bar{\Lambda}_n} e^{\bar{\alpha}_i}.$$

For $\bar{\alpha} = \sum_{i=2}^n m_i \bar{\alpha}_i + m_{n+1} \bar{\Lambda}_n$ we denote $e^{\bar{\alpha}} = (e^{\bar{\alpha}_2})^{m_2} (e^{\bar{\alpha}_3})^{m_3} \dots (e^{\bar{\alpha}_n})^{m_n} (e^{\bar{\Lambda}_n})^{m_{n+1}}$. For example $e^{\bar{\alpha}_1} = e^{-2\bar{\alpha}_2} e^{-3\bar{\alpha}_3} \dots e^{-n\bar{\alpha}_n} e^{(n+1)\bar{\Lambda}_n}$, $e^{\bar{\Lambda}_1} = e^{-\bar{\alpha}_{i+1}} e^{-2\bar{\alpha}_{i+2}} \dots e^{-(n-i)\bar{\alpha}_n} e^{(n-i+1)\bar{\Lambda}_n}$ where $\bar{\Lambda}_i$ is the i -th fundamental weight. We denote $\bar{\alpha}_0 = -\sum_{i=1}^n \bar{\alpha}_i$ and $\bar{h}_0 = -\sum_{i=1}^n \bar{h}_i$.

Note that $\langle h_i, \alpha_j \rangle = \langle \bar{h}_i, \bar{\alpha}_j \rangle$ for $0 \leq i, j \leq n$.

We denote by $\mathbf{Q}(q)\{\bar{Q}\}$ the subalgebra of $\mathbf{Q}(q)[\bar{P}]$ generated by $e^{\bar{\alpha}_i}$ ($1 \leq i \leq n$).

Set

$$W(p)_n = \mathcal{F}_n \otimes \mathbf{Q}(q)\{\bar{Q}\} e^{\bar{\Lambda}_p} \quad \text{for } 1 \leq p \leq n$$

and

$$W(0)_n = \mathcal{F}_n \otimes \mathbf{Q}(q)\{\bar{Q}\}.$$

We define the operators $H_{i,1} e^{\bar{\alpha}} (\bar{\alpha} \in \bar{Q})$, $\partial_{\bar{\alpha}_i}$ on $W(p)_n$ for $i=0,1, \dots, n$ as follows: for $v \otimes e^{\bar{\beta}} = H_{i_1, -k_1} \dots H_{i_N, -k_N} v_0 \otimes e^{\bar{\beta}} \in W(p)_n$,

$$H_{i,1} (v \otimes e^{\bar{\beta}}) = (H_{i,1} v) \otimes e^{\bar{\beta}},$$

$$e^{\bar{\alpha}} (v \otimes e^{\bar{\beta}}) = v \otimes e^{\bar{\alpha} + \bar{\beta}},$$

$$\partial_{\bar{\alpha}_i} (v \otimes e^{\bar{\beta}}) = \langle \bar{h}_i, \bar{\beta} \rangle v \otimes e^{\bar{\beta}},$$

$$d(v \otimes e^{\bar{\beta}}) = \left(-\sum_{s=1}^N k_s - \frac{(\bar{\beta}|\bar{\beta})}{2} + \frac{(\bar{\Lambda}_p|\bar{\Lambda}_p)}{2} \right) v \otimes e^{\bar{\beta}}.$$

We have the following lemma.

Lemma 3.2.1. *As operators on $W(p)_n$,*

$$e^{\bar{\alpha}_i} e^{\bar{\alpha}_j} = (-1)^{\langle \bar{h}_i, \bar{\alpha}_j \rangle} e^{\bar{\alpha}_j} e^{\bar{\alpha}_i},$$

$$e^{\bar{\alpha}_i} q^{\partial_{\bar{\alpha}_i}} = q^{\langle \bar{h}_i, \bar{\alpha}_i \rangle} q^{\partial_{\bar{\alpha}_i}} e^{\bar{\alpha}_i},$$

$$e^{\bar{\alpha}_i} z^{\partial_{\bar{\alpha}_i}} = z^{\langle \bar{h}_i, \bar{\alpha}_i \rangle} z^{\partial_{\bar{\alpha}_i}} e^{\bar{\alpha}_i}$$

for $0 \leq i, j \leq n$.

We introduce the following generating functions:

$$E_i(z) = \sum_{k \in \mathbb{Z}} E_{i,k} z^{-k},$$

$$F_i(z) = \sum_{k \in \mathbb{Z}} F_{i,k} z^{-k}.$$

Proposition 3.2.2. *Let $c = 1$ and $\kappa = 1$. Then for each p and n , the following action gives a $U_q^{\phi'}(\mathfrak{g}_{tor})$ -module structure on $W(p)_n$:*

$$q^{\frac{1}{2}c} \mapsto q^{\frac{1}{2}},$$

$$q^{d_1} \mapsto q^d,$$

$$E_i(z) \mapsto \exp\left(\sum_{k \geq 1} \frac{H_{i,-k}}{[k]} (q^{-1/2}z)^k\right) \exp\left(\sum_{k \geq 1} -\frac{H_{i,k}}{[k]} (q^{1/2}z)^{-k}\right) e^{\bar{\alpha}_i z^{\bar{\alpha}_i + 1}},$$

$$F_i(z) \mapsto \exp\left(\sum_{k \geq 1} -\frac{H_{i,-k}}{[k]} (q^{1/2}z)^k\right) \exp\left(\sum_{k \geq 1} \frac{H_{i,k}}{[k]} (q^{-1/2}z)^{-k}\right) e^{-\bar{\alpha}_i z^{-\bar{\alpha}_i + 1}},$$

$$K_i^+(z) \mapsto \exp\left((q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}\right) q^{\bar{\alpha}_i},$$

$$K_i^-(z) \mapsto \exp\left(- (q - q^{-1}) \sum_{k \geq 1} H_{i,-k} z^k\right) q^{-\bar{\alpha}_i}$$

for $0 \leq i \leq n$.

The proof will be given in Appendix.

We have immediately the following lemma.

Lemma 3.2.3. *The $U_q^{\phi'}(\mathfrak{g}_{tor})$ -module $W(p)_n$ is cyclic:*

$$W(p)_n = U_q^{\phi'}(\mathfrak{g}_{tor}) (v_0 \otimes e^{A_p}).$$

Theorem 3.2.4. *If $n > 1$ then $W(p)_n$ is irreducible for any p .*

Proof. Since \mathcal{F}_n is irreducible with respect to the action of S_n , it is enough to show that for any non-zero $v = v_0 \otimes \sum_{\bar{\alpha} \in \bar{Q}} a_{\bar{\alpha}} e^{\bar{\alpha}} e^{\bar{A}_1}$ ($a_{\bar{\alpha}} \in \mathbb{Q}(q)$) there exists $X \in U_q^{\phi'}(\mathfrak{g}_{tor})$ such that $Xv = v_0 \otimes e^{\bar{A}_p}$. Let \bar{S}_n be the subalgebra of S_n generated by $H_{i,l}$ ($1 \leq i \leq n, l \in \mathbb{Z} \setminus \{0\}$) and $q^{\frac{1}{2}c}$. Let $\bar{\mathcal{F}}_n$ be the \bar{S}_n -submodule of \mathcal{F}_n generated by $v_0 \otimes e^{\bar{A}_1}$, and let $\overline{W(p)}_n = \bar{\mathcal{F}}_n \otimes \mathbb{Q}(q) \{ \bar{Q} \} e^{\bar{A}_p}$. As already known $\overline{W(p)}_n$ is an irreducible $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module. It is obvious that $v \in \overline{W(p)}_n$. Therefore there exists $X \in U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1}) \subset U_q^{\phi'}(\mathfrak{g}_{tor})$ such that $Xv = v_0 \otimes e^{\bar{A}_p}$. \square

Remark 3.2.5. Since \mathcal{F}_1 is not irreducible as an S_1 -module $W(p)_1$ is not

irreducible.

Remark 3.2.6. Since $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ and $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ are subalgebras of $U_q^{(2)'}(\mathfrak{g}_{tor})$ we can regard $W(p)_n$ as a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module or as a $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ -module. As a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module, $W(p)_n$ is a level 1 module. On the other hand it is a level 0 module as a $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ -module.

3.3. On the structure of level (1,0) modules. In this subsection we will study the level 1 $U_q(\widehat{\mathfrak{sl}}_{n+1})$ -module structure of $W(p)_n$.

Let M be a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module. We can regard M as a $U_q(\widehat{\mathfrak{sl}}_{n+1})$ -module via a surjective homomorphism $U_q(\widehat{\mathfrak{sl}}_{n+1}) \rightarrow U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$. As a $U_q(\widehat{\mathfrak{sl}}_{n+1})$ -module, we denote the character of M by ch_M .

Let $L(\Lambda_p)$ be the irreducible highest weight $U_q(\widehat{\mathfrak{sl}}_{n+1})$ -module with highest weight Λ_p . Note that the following identity holds:

$$(3.3.1) \quad ch_{L(\Lambda_p)} = \frac{e^{\Lambda_p} \sum_{\alpha \in \widehat{\mathcal{Q}}} e^{\alpha - (\frac{1}{2}(\alpha|\alpha) - (\alpha|\overline{\Lambda_p}))\delta}}{\varphi(e^{-\delta})^n}.$$

Here $\varphi(x) = \prod_{k>0} (1-x^k)$.

We denote δ by the null root of $U_q(\widehat{\mathfrak{sl}}_{n+1})$.

By the definition of $W(p)_n$ and (3.2.1) it is immediate to see the following proposition.

Proposition 3.3.1. *As a $U_q(\widehat{\mathfrak{sl}}_{n+1})$ -module, we have*

$$\begin{aligned} ch_{W(p)_n} &= \frac{e^{\Lambda_p} \sum_{\alpha \in \widehat{\mathcal{Q}}} e^{\alpha - (\frac{1}{2}(\alpha|\alpha) - (\alpha|\overline{\Lambda_p}))\delta}}{\varphi(e^{-\delta})^{n+1}} \\ &= \frac{ch_{L(\Lambda_p)}}{\varphi(e^{-\delta})}. \end{aligned}$$

Lemma 3.3.2. *For each $l \in \mathbb{Z} \setminus \{0\}$ there exist $\widehat{H}_l = \sum_{i=0}^n \alpha_{1,i} H_{1,i}$ ($\alpha_{1,i} \in \mathbb{Q}(q)$) such that*

$$[\widehat{H}_l, H_{j,k}] = 0$$

for any $1 \leq j \leq n$ and $k \in \mathbb{Z} \setminus \{0\}$. Moreover such \widehat{H}_l is unique up to scalar.

Proof. Note that $\kappa=1$. The rank of $n \times (n+1)$ -matrix $([l\langle h_i, \alpha_j \rangle])_{1 \leq i \leq n, 0 \leq j \leq n}$ is equal to n . The lemma follows from this fact immediately. \square

By the definition of \widehat{H}_l we have

$$(3.3.2) \quad [\widehat{H}_k, \widehat{H}_l] = \delta_{k+l,0} \gamma_k.$$

where $\gamma_k \in \mathbb{Q}(q)$. We fix a normalization of \widehat{H}_l by putting $\gamma_k = 1$ for all k .

Let \widehat{S}_n be the subalgebra of S_n generated by \widehat{H}_l . By the definition, \widehat{S}_n acts on $W(p)_n$.

The following two lemmas are easy to see.

Lemma 3.3.3. For $l > 0$, $\widehat{H}_l(v_0 \otimes e^{\bar{p}}) = 0$.

Lemma 3.3.4. The action of $U'_q(\widehat{\mathfrak{sl}}_{n+1})$ on $W(p)_n$ commutes with the action of \widehat{S}_n .

Let \widehat{S}_n^- be the subalgebra of \widehat{S}_n generated by $\widehat{H}_l (l < 0)$.

Proposition 3.3.5. As $U_q(\widehat{\mathfrak{sl}}_{n+1})$ -module

$$W(p)_n \cong L(\Lambda_p)^{\oplus \infty}.$$

Proof. Set $\deg(\widehat{H}_k) = k$. Let $M_k = M_k(\widehat{H}_{-1}, \widehat{H}_{-2}, \dots)$ be a monomial of degree k in variables $\widehat{H}_{-1}, \widehat{H}_{-2}, \dots$. Then by the above two lemmas, $M_k v_0 \otimes e^{\bar{\Lambda}_p}$ is a singular vector of $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module $W(p)_n$. Let W_{M_k} be the $U_q(\widehat{\mathfrak{sl}}_{n+1})$ -submodule which is generated by $M_k v_0 \otimes e^{\bar{\Lambda}_p}$. Then by the definition of the action of $U_q(\widehat{\mathfrak{sl}}_{n+1})$ on $W(p)_n$, we have

$$W_{M_k} \cong L(\Lambda_p - k\delta) \cong L(\Lambda_p).$$

The vectors $\{M_k v_0 \otimes e^{\bar{\Lambda}_p}\}$ are linearly independent. The number of the monomials of degree k is equal to the k -th partition number $p(k)$. Therefore there is a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -submodule W of $W(p)_n$ which is isomorphic to $\bigoplus_{k \geq 0} L(\Lambda_p - k\delta)^{\oplus p(k)}$. By Proposition 3.3.1 it coincides with $W(p)_n$. This completes proof. \square

By Lemma 3.3.4 and the proof of Proposition 3.3.5, the following corollary follows immediately.

Corollary 3.3.6. As $U'_q(\widehat{\mathfrak{sl}}_{n+1}) \otimes \widehat{S}_n^-$ -module $W(p)_n$ is isomorphic to $L(\Lambda_p) \otimes \widehat{S}_n^-$.

3.4. Construction of level $(1, 1)$ modules. We introduce a twisted version of the group algebra $\mathbb{Q}(q)[Q]$ by $\mathbb{Z} \setminus 2\mathbb{Z}$. We denote it by $\mathbb{Q}(q)\{Q\}$. This is the $\mathbb{Q}(q)$ -algebra generated by symbols $e^{\alpha_0}, e^{\alpha_1}, \dots, e^{\alpha_n}$ which satisfy the following relations:

$$(3.4.1) \quad e^{\alpha_i} e^{\alpha_j} = (-1)^{\langle h_i, \alpha_j \rangle} e^{\alpha_j} e^{\alpha_i}.$$

Similarly to §3.2, we denote $e^\alpha = (e^{\alpha_0})^{m_0} (e^{\alpha_1})^{m_1} \dots (e^{\alpha_n})^{m_n}$ for $\alpha = \sum_{i=0}^n m_i \alpha_i \in Q$.

Let

$$V(p)_n = \mathcal{F}_n \otimes \mathbf{Q}(q) \{Q\} e^{\Lambda_p}.$$

Here we regard e^{Λ_p} only a symbol indexed by p .

We define the operators $H_{i,l}$ ($0 \leq i \leq n, l \neq 0$), e^α ($\alpha \in Q$), ∂_{α_i} and $z^{H_{i,0}}$ ($0 \leq i \leq n$) on $V(p)_n$ as follows:

$$\text{for } v \otimes e^\beta e^{\Lambda_p} = H_{i_1, -k_1} \dots H_{i_N, -k_N} v_0 \otimes e^\beta e^{\Lambda_p} \in V(p)_n (\beta = \sum_{k=0}^n m_k \alpha_k \in Q),$$

$$H_{i,l}(v \otimes e^\beta e^{\Lambda_p}) = (H_{i,l}v) \otimes e^\beta e^{\Lambda_p},$$

$$e^\alpha(v \otimes e^\beta e^{\Lambda_p}) = v \otimes (e^\alpha e^\beta) e^{\Lambda_p},$$

$$\partial_{\alpha_i}(v \otimes e^\beta e^{\Lambda_p}) = \langle h_{i,\beta} + \Lambda_p \rangle v \otimes e^\beta e^{\Lambda_p},$$

$$z^{H_{i,0}}(v \otimes e^\beta e^{\Lambda_p}) = z^{\langle h_{i,\beta} + \Lambda_p \rangle} \kappa^{\frac{1}{2} \sum_{i=0}^n \langle h_i, m_i \alpha_i \rangle m_{i_1}} v \otimes e^\beta e^{\Lambda_p},$$

$$d_1(v \otimes e^\beta e^{\Lambda_p}) = \left(-\sum_{s=1}^N k_s - \frac{(\beta|\beta)}{2} - (\beta|\Lambda_p) \right) v \otimes e^\beta e^{\Lambda_p},$$

$$d_2(v \otimes e^\beta e^{\Lambda_p}) = m_0(v \otimes e^\beta e^{\Lambda_p}).$$

The following lemma is easy.

Lemma 3.4.1. *As operators on $V(p)_n$,*

$$e^{\alpha_i} e^{\alpha_j} = (-1)^{\langle h_i, \alpha_j \rangle} e^{\alpha_j} e^{\alpha_i}$$

$$q^{\partial_{\alpha_i}} e^{\alpha_j} = q^{\langle h_i, \alpha_j \rangle} e^{\alpha_j} q^{\partial_{\alpha_i}}$$

$$z^{H_{i,0}} e^{\alpha_j} = z^{\langle h_i, \alpha_j \rangle} \kappa^{\frac{1}{2} \langle h_i, \alpha_j \rangle m_{i_1}} e^{\alpha_j} z^{H_{i,0}}.$$

Proposition 3.4.2. *Assume $c = 1$ then for each p and n , the following action gives a $U_q(\mathfrak{g}_{tor})$ -module structure on $V(p)_n$:*

$$q^{\frac{1}{2}c} \mapsto q^{\frac{1}{2}},$$

$$q^{d_1} \mapsto q^{d_1},$$

$$q^{d_2} \mapsto q^{d_2},$$

$$E_i(z) \mapsto \exp \left(\sum_{k \geq 1} \frac{H_{i,-k}}{[k]} (q^{-1/2}z)^k \right) \exp \left(\sum_{k \geq 1} -\frac{H_{i,k}}{[k]} (q^{1/2}z)^{-k} \right) e^{\alpha_i} z^{H_{i,0}+1},$$

$$F_i(z) \mapsto \exp \left(\sum_{k \geq 1} -\frac{H_{i,-k}}{[k]} (q^{1/2}z)^k \right) \exp \left(\sum_{k \geq 1} \frac{H_{i,k}}{[k]} (q^{-1/2}z)^{-k} \right) e^{-\alpha_i} z^{-H_{i,0}+1},$$

$$K_i^+(z) \mapsto \exp \left((q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k} \right) q^{\partial_{\alpha_i}}.$$

$$K_i^-(z) \mapsto \exp\left(- (q-q^{-1}) \sum_{k \geq 1} H_{1,-k} z^k\right) q^{-\partial a_i}$$

for $0 \leq i \leq n$.

The proof will be given in Appendix.

It is easy to see the following lemma.

Lemma 3.4.3. $V(p)_n$ is a cyclic $U_q(\mathfrak{g}_{tor})$ -module: $V(p)_n = U_q(\mathfrak{g}_{tor})(v_0 \otimes e^{A_p})$.

Lemma 3.4.4. $V(p)_n$ has level 1 as a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module and as a $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ -module.

Proof. It is clear that $V(p)_n$ is a level 1 $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module. The center of $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ is $\prod_{k=0}^n K_k$. By the definition it acts as the scalar q on $V(p)_n$. \square

§4. On $U_q(\mathfrak{sl}_{2,tor})$

4.1. In this section we assume that $\mathfrak{g} = \mathfrak{sl}_2$. We shall try to find finitely many generators of $U_q(\mathfrak{sl}_{2,tor})$.

Let

$$\begin{aligned} E_i &= E_{i,0}, F_i = F_{i,0}, q^{\pm h_i} = K_i^{\pm}, \text{ for } i=0,1, \\ E_{-1} &= F_{0,1}, K_0^-, F_{-1} = K_0^+ E_{0,-1}, q^{\pm h_{-1}} = q^{\pm c} K_0^{\mp}. \end{aligned}$$

Proposition 4.1.1. $U_q(\mathfrak{sl}_{2,tor})$ is generated by $E_i, F_i, q^{\pm h_i}$ ($i = -1, 0, 1$), $q^{\pm \frac{1}{2}c}$, $q^{\pm d_1}, q^{\pm d_2}$.

Proof. Let \mathcal{A} be the subalgebra of $U_q(\mathfrak{sl}_{2,tor})$ generated by $E_i, F_i, q^{\pm h_i}$ ($i = -1, 0, 1$), $q^{\pm \frac{1}{2}c}$, $q^{\pm d_1}, q^{\pm d_2}$. By the definition we have $E_{0,-1} = q^{-h_0} F_{-1}$ and $F_{0,1} = E_{-1} q^{h_0}$. Since

$$\begin{aligned} [E_{0,0}, F_{0,1}] &= \frac{1}{q - q^{-1}} q^{-\frac{1}{2}c} K_{0,1}^+ \\ &= q^{-\frac{1}{2}c} q^{h_0} H_{0,1} \end{aligned}$$

and

$$\begin{aligned} [E_{0,-1}, F_{0,0}] &= -\frac{1}{q - q^{-1}} q^{\frac{1}{2}c} K_{0,-1}^- \\ &= q^{\frac{1}{2}c} q^{-h_0} H_{0,-1}. \end{aligned}$$

We deduce $H_{0,1}$, and $H_{0,-1} \in \mathcal{A}$. We recall (2.2.7)

$$[H_{t,k}, E_{j,l}] = \frac{1}{k} [k \langle h_t, \alpha_j \rangle] q^{-\frac{1}{2}|k|c} \kappa^{-km_u} E_{j,k+l},$$

$$[H_{t,k}, F_{j,l}] = -\frac{1}{k} [k \langle h_t, \alpha_j \rangle] q^{\frac{1}{2}|k|c} \kappa^{-km_u} F_{j,k+l}.$$

By these formulas we have $E_{t,k}, F_{t,k} \in \mathcal{A}$ for $i=0,1, k \in \mathbf{Z}$ in inductively.

On the other hand we know

$$\begin{aligned} [E_{0,1}, F_{0,1}] &= \frac{1}{q-q^{-1}} q^{-\frac{1}{2}c} K_{0,2}^+ \\ &= \frac{1}{2} (q-q^{-1}) H_{0,1}^2 + H_{0,2}. \end{aligned}$$

Therefore we get $H_{0,2} \in \mathcal{A}$. Similarly we have $H_{t,i} \in \mathcal{A}$ for any i,l .

This completes the proof. \square

Lemma 4.1.2. *The following relations hold in $U_q(\mathfrak{sl}_{2,tor})$:*

$$(4.1.1) \quad [q^{\pm h_i}, q^{\pm h_j}] = 0,$$

$$(4.1.2) \quad [q^{\pm d_i}, q^{\pm h_j}] = 0,$$

$$(4.1.3) \quad q^{d_1} E_j q^{-d_1} = q^{\delta_{j,-1}} E_j,$$

$$q^{d_1} F_j q^{-d_1} = q^{-\delta_{j,-1}} F_j,$$

$$(4.1.4) \quad q^{d_2} E_j q^{-d_2} = q^{-1+\delta_{j,1}} E_j,$$

$$q^{d_2} F_j q^{-d_2} = q^{1-\delta_{j,1}} F_j,$$

$$(4.1.5) \quad q^{h_i} E_j q^{-h_i} = q^{a_{ij}} E_j,$$

$$q^{h_i} F_j q^{-h_i} = q^{-a_{ij}} F_j,$$

where

$$(a_{ij})_{-1 \leq i, j \leq 1} = \begin{pmatrix} 2 & -2 & 2 \\ -2 & 2 & -2 \\ 2 & -2 & 2 \end{pmatrix}.$$

$$(4.1.6) \quad [E_i, F_j] = \delta_{ij} \frac{q^{h_i} - q^{-h_i}}{q - q^{-1}}, \quad \text{for } |i-j| \leq 1,$$

$$(4.1.7) \quad E_{-1}^3 F_1 - q^{-2} [3] E_{-1}^2 F_1 E_{-1} + q^{-4} [3] E_{-1} F_1 E_{-1}^2 - q^{-6} F_1 E_{-1}^3 = 0,$$

$$(4.1.8) \quad E_1^3 F_{-1} - q^2 [3] E_1^2 F_{-1} E_1 + q^4 [3] E_1 F_{-1} E_1^2 - q^6 F_{-1} E_1^3 = 0,$$

$$(4.1.9) \quad F_{-1}^3 E_1 - q^{-2} [3] F_{-1}^2 E_1 F_{-1} + q^{-4} [3] F_{-1} E_1 F_{-1}^2 - q^{-6} E_1 F_{-1}^3 = 0,$$

$$(4.1.10) \quad F_i^3 E_{-1} - q^2 [3] F_i^2 E_{-1} F_1 + q^4 [3] F_1 E_{-1} F_i^2 - q^6 E_{-1} F_i^3 = 0,$$

$$(4.1.11) \quad E_i^3 E_j - [3] E_i^2 E_j E_i + [3] E_i E_j E_i^2 - E_j E_i^3 = 0, \quad \text{for } |i-j|=1,$$

$$(4.1.12) \quad F_i^3 F_j - [3] F_i^2 F_j F_i + [3] F_i F_j F_i^2 - F_j F_i^3 = 0, \quad \text{for } |i-j|=1,$$

$$(4.1.13) \quad E_{-1} E_1 - q^2 E_1 E_{-1} = 0,$$

$$(4.1.14) \quad F_{-1} F_1 - q^2 F_1 F_{-1} = 0.$$

Proof. By the definition of $U_q(\mathfrak{g}_{tor})$ and E_i, F_i and q^h , it is easy to check these relations. \square

Let \mathcal{U} be an associative algebra over $\mathbb{Q}(q)$ generated by $E_i, F_i, q^{\pm h_i}$ ($i = -1, 0, 1$), $q^{\pm \frac{1}{2}c}, q^{\pm d_1}, q^{\pm d_2}$ with relations (4.1.1) – (4.1.14). Then we have,

Corollary 4.1.3. *There is a canonical surjective algebra homomorphism $\Psi: \mathcal{U} \rightarrow U_q(\mathfrak{sl}_{2,tor})$.*

Remark 4.1.4. Ψ has a highly nontrivial kernel. It is important to determine it. For example the following formulas holds in $U_q(\mathfrak{sl}_{2,tor})$:

$$\kappa^{m_{01}} E_{0,0} E_{1,-1} - q^{-2} \kappa^{m_{01}} E_{1,-1} E_{0,0} = q^{-2} E_{0,-1} E_{1,0} - E_{1,0} E_{0,-1},$$

$$E_{0,-1} = q^{-h_0} F_{-1}$$

and

$$E_{1,-1} = \frac{\kappa^{-m_{01}}}{[-2]} [F_{-1} F_0 - q^{-2} F_0 F_{-1}, E_1].$$

Therefore we have

$$\begin{aligned} X &= \frac{1}{[-2]} E_0 [F_{-1} F_0 - q^{-2} F_0 F_{-1}, E_1] - \frac{q^{-2}}{[-2]} [F_{-1} F_0 - q^{-2} F_0 F_{-1}, E_1] E_0 \\ &\quad - q^{-2} q^{-h_0} F_{-1} E_1 - E_1 q^{-h_0} F_{-1} \\ &= 0 \end{aligned}$$

in $U_q(\mathfrak{sl}_{2,tor})$. Thus $X \in \text{Ker } \Psi$. But, as an element of \mathcal{U} , X is not equal to 0.

4.2. Let

$$E_{0*} = F_{1,1} K_1^-, \quad F_{0*} = K_1^+ E_{1,-1}, \quad q^{\pm h_{0*}} = q^{\pm c} K_1^{\mp}.$$

Proposition 4.2.1. *The subalgebra generated by E_i, F_i , and q^{h_i} for $i = 0, 1$, $E_{0*}, F_{0*}, q^{h_{0*}}, q^{\pm c}, q^{\pm d_1}, q^{\pm d_2}$, is equal to $U_q(\mathfrak{sl}_{2,tor})$. That is, they are generators of*

$U_q(\widehat{\mathfrak{sl}}_{2,tor})$. Moreover these generators satisfy relations similar to the ones in Lemma 4.1.2.

Proof. This proposition is proved in the same way as Proposition 4.1.1 and Lemma 4.1.2. \square

We have immediately the following lemma.

Lemma 4.2.2. *Let $U_q^{(1)}$ be the subalgebras generated by E_i, F_i, q^{h_i} for $i=1,0^*$ and $q^{\pm(d_1+d_2)}$, $U_q^{(2)}$ the subalgebras generated by E_i, F_i, q^{h_i} for $i=0, 1$ and $q^{\pm(d_1+d_2)}$, $U_q^{(3)}$ the subalgebras generated by $E_i, F_i, q^{\pm h_i}$ for $i=0^*, -1$ and $q^{\pm(d_1+d_2)}$, and $U_q^{(4)}$ the subalgebras generated by $E_i, F_i, q^{\pm h_i}$ for $i=0, -1$ and $q^{\pm(d_1+d_2)}$. Then $U_q^{(i)}$ ($i=1,2,3,4$) are isomorphic to $U_q(\widehat{\mathfrak{sl}}_2)$.*

Those four algebras are schematically visualized by Fig. 1.




Fig. 1.

Let $U_q(\widehat{\mathfrak{sl}}_2)_{(i)}$ ($i=-1,0,1,0^*$) be the subalgebra of $U_q(\widehat{\mathfrak{sl}}_{2,tor})$ generated by $E_i, F_i, q^{\pm h_i}$. All $U_q(\widehat{\mathfrak{sl}}_2)_{(i)}$ ($i \in \{-1,0,1,0^*\}$) are isomorphic to $U_q(\widehat{\mathfrak{sl}}_2)$. The upper left circle in Fig. 1 means $U_q(\widehat{\mathfrak{sl}}_2)_{(0)}$, the upper right one means $U_q(\widehat{\mathfrak{sl}}_2)_{(1)}$, the lower left one means $U_q(\widehat{\mathfrak{sl}}_2)_{(-1)}$ and the lower right one means $U_q(\widehat{\mathfrak{sl}}_2)_{(0^*)}$. The diagram

$$\circ' \iff \circ' \quad (i, j \in \{-1,0,1,0^*\})$$

means the algebra generated by $U_q(\widehat{\mathfrak{sl}}_2)_{(i)}$ and $U_q(\widehat{\mathfrak{sl}}_2)_{(j)}$ is isomorphic to $U_q(\widehat{\mathfrak{sl}}_2)$. For example $\circ^0 \iff \circ^1$ means the algebra generated by $U_q(\widehat{\mathfrak{sl}}_2)_{(0)}$ and $U_q(\widehat{\mathfrak{sl}}_2)_{(-1)}$ which we call $U_q^{(2)}$ is isomorphic to $U_q(\widehat{\mathfrak{sl}}_2)$. The meaning of the diagram

$$\circ' \implies \circ'$$

is as follows: In the algebra generated by $U_q(\widehat{\mathfrak{sl}}_2)_{(i)}$ and $U_q(\widehat{\mathfrak{sl}}_2)_{(j)}$, the following relations hold

$$(4.2.1) \quad q^{h_i} E_j q^{-h_i} = q^2 E_j, \quad q^{h_i} E_i q^{-h_i} = q^2 E_i,$$

$$(4.2.2) \quad q^{h_i} F_j q^{-h_i} = q^{-2} F_j, \quad q^{h_j} F_i q^{-h_j} = q^{-2} F_i,$$

$$(4.2.3) \quad E_i^3 F_j - q^{-2} [3] E_i^2 F_j E_i + q^{-4} [3] E_i F_j E_i^2 - q^{-6} F_j E_i^3 = 0,$$

$$E_j^3 F_i - q^2 [3] E_j^2 F_i E_j + q^4 [3] E_j F_i E_j^2 - q^6 F_i E_j^3 = 0,$$

$$(4.2.4) \quad F_i^3 E_j - q^{-2} [3] F_i^2 E_j F_i + q^{-4} [3] F_i E_j F_i^2 - q^{-6} E_j F_i^3 = 0,$$

$$F_j^3 E_i - q^2 [3] F_j^2 E_i F_j + q^4 [3] F_j E_i F_j^2 - q^6 E_i F_j^3 = 0,$$

$$(4.2.5) \quad E_i E_j - q^2 E_j E_i = 0,$$

$$(4.2.6) \quad F_i F_j - q^2 F_j F_i = 0.$$

Appendix A.

A.1. Proof of Proposition 3.2.2 and 3.4.2. For the proof we rewrite the defining relation $U_q(\mathfrak{g}_{\text{tor}})$ generating function level.

$$(A.1.1) \quad q^{\pm \frac{1}{2}c} \text{ are central.}$$

$$(A.1.2) \quad K_i^+ K_i^- = K_i^- K_i^+ = 1,$$

$$(A.1.3) \quad K_i^\pm(z) K_j^\pm(w) = K_j^\pm(w) K_i^\pm(z)$$

$$(A.1.4)$$

$$\theta_{-\langle h_i, \alpha \rangle} \left(q^{-c} \kappa^{-m_u \frac{z}{w}} \right) K_i^-(z) K_j^+(w) = \theta_{-\langle h_i, \alpha \rangle} \left(q^c \kappa^{-m_u \frac{z}{w}} \right) K_j^+(w) K_i^-(z)$$

$$(A.1.5) \quad q^{d_1} K_i^\pm(z) q^{-d_1} = K_i^\pm(q^{-1}z),$$

$$(A.1.6) \quad [q^{d_2}, K_i^\pm(z)] = 0,$$

$$(A.1.7) \quad q^{d_1} E_j(z) q^{-d_1} = E_j(q^{-1}z),$$

$$q^{d_1} F_j(z) q^{-d_1} = F_j(q^{-1}z),$$

$$(A.1.8) \quad q^{d_2} E_j(z) q^{-d_2} = q^{\delta_{j0}} E_j(z),$$

$$q^{d_2} F_j(z) q^{-d_2} = q^{-\delta_{j0}} F_j(z),$$

$$(A.1.9) \quad K_i^+(z) E_j(w) = \theta_{-\langle h_i, \alpha \rangle} \left(q^{-\frac{1}{2}c} \kappa^{-m_u \frac{w}{z}} \right) E_j(w) K_i^+(z)$$

$$K_i^-(z) E_j(w) = \theta_{\langle h_i, \alpha \rangle} \left(q^{-\frac{1}{2}c} \kappa^{m_u \frac{z}{w}} \right) E_j(w) K_i^-(z)$$

$$K_i^+(z) F_j(w) = \theta_{\langle h_i, \alpha \rangle} \left(q^{\frac{1}{2}c} \kappa^{-m_u \frac{w}{z}} \right) F_j(w) K_i^+(z)$$

$$K_i^-(z)F_j(w) = \theta_{-\langle h_i, \alpha_j \rangle} \left(q^{\frac{1}{2}c} \kappa^{m_i} \frac{z}{w} \right) F_j(w) K_i^-(z)$$

(A.1.10)

$$[E_i(z), F_j(w)] = \delta_{i,j} \frac{1}{q-q^{-1}} \left\{ \delta \left(q^c \frac{w}{z} \right) K_i^+ \left(q^{\frac{1}{2}c} w \right) - \delta \left(q^c \frac{z}{w} \right) K_i^- \left(q^{\frac{1}{2}c} z \right) \right\}$$

$$(A.1.11) \quad (\kappa^{m_i} z - q^{\langle h_i, \alpha_j \rangle} w) E_i(z) E_j(w) = (q^{\langle h_i, \alpha_j \rangle} \kappa^{m_i} z - w) E_j(w) E_i(z)$$

$$(\kappa^{m_i} z - q^{-\langle h_i, \alpha_j \rangle} w) F_i(z) F_j(w) = (q^{-\langle h_i, \alpha_j \rangle} \kappa^{m_i} z - w) F_j(w) F_i(z)$$

(A.1.12)

$$\sum_{\sigma \in \mathfrak{S}_m} \sum_{r=0}^m (-1)^r \begin{bmatrix} m \\ r \end{bmatrix}_q E_i(z_{\sigma(1)}) \cdots E_i(z_{\sigma(r)}) E_j(w) E_i(z_{\sigma(r+1)}) \cdots E_i(z_{\sigma(m)}) = 0$$

$$\sum_{\sigma \in \mathfrak{S}_m} \sum_{r=0}^m (-1)^r \begin{bmatrix} m \\ r \end{bmatrix}_q F_i(z_{\sigma(1)}) \cdots F_i(z_{\sigma(r)}) F_j(w) F_i(z_{\sigma(r+1)}) \cdots F_i(z_{\sigma(m)})$$

where $i \neq j$ and $m = 1 - \langle h_i, \alpha_j \rangle$.

In these formulas we denote $\theta_m(z) = \frac{zq^{m-1}}{z-q^m}$ for $m \in \mathbb{Z}$, $\delta(z) = \sum_{k \in \mathbb{Z}} z^k$.

If Proposition 3.4.2 holds, then, from Lemma 3.2.1 and 3.4.1, we have Proposition 3.2.2 by putting $h_i \mapsto \bar{h}_i$, $\alpha_i \mapsto \bar{\alpha}_i$, $\kappa \mapsto 1$ and $z^{H_{i,0}} \mapsto z^{\bar{0}_i}$. Therefore it is enough to show Proposition 3.4.2.

The relations (A.1.1), (A.1.2) and (A.1.3) are trivial. (A.1.4) is just the commutation relations of Heisenberg algebra S_n . Therefore, by the definition of $V(p)_n$, it is clear that (A.1.4) holds. The relations (A.1.5), (A.1.6) immediately follows from the definition of d_1 and d_2 .

Let us show (A.1.7) and (A.1.8). Take $v \otimes e^\beta c^{A_p} \in V(p)_n$ where $\beta = \sum_{k=0}^n m_k \alpha_k \in Q$. Then we have

$$q^{d_1} e^{\pm \alpha_j} z^{\pm H_{j,0}+1} q^{-d_1} (v \otimes e^\beta e^{A_p})$$

$$= q^{\mp \langle \alpha_j, \beta + A_p \rangle - 1} z^{\pm \langle h_j, \beta + A_p \rangle + 1} \kappa^{\frac{1}{2} \sum_{k=0}^n \langle h_j, m_k \alpha_k \rangle m_{j,k}} v \otimes e^{\pm \alpha_j} e^\beta e^{A_p}$$

$$= (q^{-1} z)^{\pm \langle h_j, \beta + A_p \rangle + 1} \kappa^{\frac{1}{2} \sum_{k=0}^n \langle h_j, m_k \alpha_k \rangle m_{j,k}} v \otimes e^{\pm \alpha_j} e^\beta e^{A_p}$$

$$= e^{\pm \alpha_j} (q^{-1} z)^{\pm H_{j,0}+1} (v \otimes e^\beta e^{A_p}).$$

Therefore we have

$$q^{d_1} E_j(z) q^{-d_1} = q^{d_1} \exp \left(\sum_{k \geq 1} \frac{H_{j,-k}}{[k]} q^{-\frac{1}{2}k} z^k \right) \exp \left(- \sum_{k \geq 1} \frac{H_{j,k}}{[k]} q^{-\frac{1}{2}k} z^{-k} \right) e^{\alpha_j} z^{H_{j,0}+1} q^{-d_1}$$

$$= \exp \left(\sum_{k \geq 1} \frac{H_{j,-k}}{[k]} q^{-\frac{1}{2}k} (q^{-1} z)^k \right) \exp \left(- \sum_{k \geq 1} \frac{H_{j,k}}{[k]} q^{-\frac{1}{2}k} (q^{-1} z)^{-k} \right) e^{\alpha_j} (q^{-1} z)^{H_{j,0}+1}$$

$$=E_j(q^{-1}z).$$

Similarly we have $q^{d_1}F_j(z)q^{-d_1}=F_j(q^{-1}z)$.

It is clear that

$$(A.1.13) \quad q^{d_2}e^{\pm\alpha_j}z^{\pm H_{j,0}+1}q^{-d_2}(v\otimes e^\beta e^{A_p})=q^{\pm\delta_{j,0}}e^{\pm\alpha_j}z^{\pm H_{j,0}+1}(v\otimes e^\beta e^{A_p}).$$

From (A.1.13) and the fact that q^{d_2} commutes with $H_{j,k}$, we have (A.1.8).

We shall show (A.1.9). We denote

$$\begin{aligned} E_i^+(z) &= \exp\left(\sum_{k\geq 1} \frac{H_{i,-k}}{[k]} q^{-\frac{1}{2}k} z^k\right), \\ E_i^-(z) &= \exp\left(-\sum_{k\geq 1} \frac{H_{i,k}}{[k]} q^{-\frac{1}{2}k} z^{-k}\right), \\ F_i^+(z) &= \exp\left(-\sum_{k\geq 1} \frac{H_{i,-k}}{[k]} q^{\frac{1}{2}k} z^k\right), \\ F_i^-(z) &= \exp\left(\sum_{k\geq 1} \frac{H_{i,k}}{[k]} q^{\frac{1}{2}k} w^{-k}\right). \end{aligned}$$

Let us proof

$$K_i^+(z)E_j(w)=\theta_{-\langle h_i,\alpha_j\rangle}\left(q^{-\frac{1}{2}}\kappa^{m-u}\frac{w}{z}\right)E_j(w)K_i^+(z).$$

We have

$$\begin{aligned} \left[(q-q^{-1})\sum_{k\geq 1} H_{i,k}z^{-k}, \sum_{l\geq 1} \frac{H_{j,-l}}{[l]} q^{-\frac{1}{2}l}w^l\right] &= \sum_{k,l} (q-q^{-1}) \frac{1}{[l]} [H_{i,k}, H_{j,-l}] z^{-k} q^{-\frac{1}{2}l}w^l \\ &= \sum_k (q-q^{-1}) \frac{[k\langle h_i,\alpha_j\rangle]}{k} \kappa^{-km_u} q^{-\frac{1}{2}k} \left(\frac{w}{z}\right)^k \\ &= \sum_k \frac{1}{k} (q^{k(\langle h_i,\alpha_j\rangle-\frac{1}{2})} - q^{k(-\langle h_i,\alpha_j\rangle-\frac{1}{2})}) \kappa^{-km_u} \left(\frac{w}{z}\right)^k \\ &= \log \frac{1 - q^{-\langle h_i,\alpha_j\rangle-\frac{1}{2}} \kappa^{-m_u} \frac{w}{z}}{1 - q^{\langle h_i,\alpha_j\rangle-\frac{1}{2}} \kappa^{-m_u} \frac{w}{z}} \end{aligned}$$

and

$$\left[(q-q^{-1})\sum_{k\geq 1} H_{i,k}z^{-k}, -\sum_{l\geq 1} \frac{H_{j,l}}{[l]} q^{-\frac{1}{2}l}w^{-l}\right]=0.$$

We recall Campbell-Hausdorff formula: let A and B be noncommutative operators and $C=[A, B]$. If $[C, A]=[C, B]=0$ then we have $e^A e^B = e^C e^B e^A$.

By Campbell-Hausdorff formula we get

$$\begin{aligned} & \exp\left((q-q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}\right) E_j^+(w) E_j^-(w) \\ &= \frac{1 - q^{-\langle h_i, \alpha_j \rangle - \frac{1}{2} \kappa^{-m_{ij}} \frac{w}{z}}}{1 - q^{\langle h_i, \alpha_j \rangle - \frac{1}{2} \kappa^{-m_{ij}} \frac{w}{z}}} E_j^+(w) E_j^-(w) \exp\left((q-q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}\right). \end{aligned}$$

On the other hand, by Lemma 3.4.1 we have

$$q^{\partial_{\alpha_i}} e^{\alpha_j} w^{H_{j,0}} = q^{\langle h_i, \alpha_j \rangle} e^{\alpha_j} w^{H_{j,0}} q^{\partial_{\alpha_i}}.$$

Thus we get

$$\begin{aligned} & K_i^+(z) E_j(w) \\ &= \exp\left((q-q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}\right) E_j^+(w) E_j^-(w) q^{\partial_{\alpha_i}} e^{\alpha_j} w^{H_{j,0}+1} \\ &= \frac{1 - q^{-\langle h_i, \alpha_j \rangle - \frac{1}{2} \kappa^{-m_{ij}} \frac{w}{z}}}{1 - q^{\langle h_i, \alpha_j \rangle - \frac{1}{2} \kappa^{-m_{ij}} \frac{w}{z}}} E_j(w) K_i^+(z) \\ &= \theta_{-\langle h_i, \alpha_j \rangle} \left(q^{-\frac{1}{2} \kappa^{-m_{ij}} \frac{w}{z}} \right) E_j(w) K_i^+(z). \end{aligned}$$

The other formulas in (A.1.9) can be checked by similar arguments.

Let us show (A.1.10). We have

$$\begin{aligned} & \left[- \sum_{k \geq 1} \frac{H_{i,k}}{[k]} q^{-\frac{1}{2} k} z^{-k}, - \sum_{k \geq 1} \frac{H_{j,-k}}{[k]} q^{\frac{1}{2} k} w^k \right] \\ &= \sum_{k \geq 1} \frac{1}{k [k]} [k \langle h_i, \alpha_j \rangle] \kappa^{-m_{ij}} \left(\frac{w}{z} \right)^k \\ &= \begin{cases} \log \frac{1}{(1 - q \frac{w}{z}) (1 - q^{-1} \frac{w}{z})}, & \langle h_i, \alpha_j \rangle = 2 \quad (i=j), \\ \log(1 - q \kappa^{-m_{ij}} \frac{w}{z}) (1 - q^{-1} \kappa^{-m_{ij}} \frac{w}{z}), & \langle h_i, \alpha_j \rangle = -2, \\ \log(1 - \kappa^{-m_{ij}} \frac{w}{z}), & \langle h_i, \alpha_j \rangle = -1, \\ 0, & \langle h_i, \alpha_j \rangle = 0. \end{cases} \end{aligned}$$

For example we will show in the case of $\langle h_i, \alpha_j \rangle = -2$. By Campbell-Hausdorff formula we get

$$E_i^\pm(z) F_j^\pm(w) = F_j^\pm(w) E_i^\pm(z)$$

and

$$E_i^-(z)F_j^+(w) = \left(1 - q\kappa^{-m_i}\frac{w}{z}\right) \left(1 - q^{-1}\kappa^{-m_i}\frac{w}{z}\right) F_j^+(w)E_i^-(z).$$

On the other hand, by Lemma 3.4.1, we have

$$z^{H_{i,0}}e^{-\alpha_i} = z^2\kappa^{m_i}e^{-\alpha_i}z^{H_{i,0}}.$$

Therefore we get

$$\begin{aligned} E_i(z)F_j(w) &= E_i^+(z)E_i^-(z)e^{\alpha_i z^{H_{i,0}+1}}F_j^+(w)F_j^-(w)e^{-\alpha_i w^{-H_{i,0}+1}} \\ &= (z\kappa^{\frac{1}{2}m_i} - qw\kappa^{-\frac{1}{2}m_i})(z\kappa^{\frac{1}{2}m_i} - q^{-1}w\kappa^{-\frac{1}{2}m_i})E_i^+(z)F_j^+(w)E_i^-(z)F_j^-(w) \\ &\quad \times e^{\alpha_i e^{-\alpha_i} z^{H_{i,0}+1} w^{-H_{i,0}+1}}. \end{aligned}$$

By a similar argument we have

$$\begin{aligned} F_j(w)E_i(z) &= (w\kappa^{-\frac{1}{2}m_i} - qz\kappa^{\frac{1}{2}m_i})(w\kappa^{-\frac{1}{2}m_i} - q^{-1}z\kappa^{\frac{1}{2}m_i})E_i^+(z)F_j^+(w)E_i^-(z)F_j^-(w) \\ &\quad \times e^{\alpha_i e^{-\alpha_i} z^{H_{i,0}+1} w^{-H_{i,0}+1}}. \end{aligned}$$

Therefore we get

$$[E_i(z), F_j(w)] = 0.$$

Similarly one can check the other formulas.

We will show (A.1.11). We have

$$\begin{aligned} &\left[-\sum_{k \geq 1} \frac{H_{i,k}}{[k]} q^{-\frac{1}{2}k} z^{-k}, \sum_{k \geq 1} \frac{H_{j,-k}}{[k]} q^{-\frac{1}{2}k} w^k \right] \\ &= \sum_{k \geq 1} \frac{1}{k[k]} [k \langle h_i, \alpha_j \rangle] \kappa^{-m_i} \left(\frac{w}{z}\right)^k \\ &= \begin{cases} \log\left(1 - \frac{w}{z}\right) \left(1 - q^{-2} \frac{w}{z}\right), & i=j, \\ \log \frac{1}{\left(1 - \kappa^{-m_i} \frac{w}{z}\right) \left(1 - q^{-2} \kappa^{-m_i} \frac{w}{z}\right)}, & \langle h_i, \alpha_j \rangle = -2, \\ \log \frac{1}{1 - q^{-1} \kappa^{-m_i} \frac{w}{z}}, & \langle h_i, \alpha_j \rangle = -1, \\ 0, & \langle h_i, \alpha_j \rangle = 0. \end{cases} \end{aligned}$$

For example let us show in the case of $\langle h_i, \alpha_j \rangle = -2$. If $\langle h_i, \alpha_j \rangle \neq -2$ one can show the formula by a similar argument. By Lemma 3.4.1 we get

$$(z\kappa^{m_i} - q^{\langle h_i, \alpha_j \rangle} w)E_i(z)E_j(w)$$

$$\begin{aligned}
 &= (z\kappa^{m_u} - q^{-2}w) E_i^+(z) E_i^-(z) e^{\alpha_i} z^{H_{i,0}+1} E_j^+(w) E_j^-(w) e^{\alpha_j} w^{H_{j,0}+1} \\
 &= \frac{z\kappa^{m_u} - q^{-2}w}{z^2 \kappa^{m_u} \left(1 - \kappa^{-m_u} \frac{w}{z}\right) \left(1 - q^{-2} \kappa^{-m_u} \frac{w}{z}\right)} E_i^+(z) E_i^+(w) E_i^-(z) E_j^-(w) \\
 &\quad \times e^{\alpha_i} e^{\alpha_j} z^{H_{i,0}+1} w^{H_{j,0}+1} \\
 &= \frac{1}{(z - \kappa^{-m_u} w)} E_i^+(z) E_i^+(w) E_j^-(z) E_j^-(w) e^{\alpha_i} e^{\alpha_j} z^{H_{i,0}+1} w^{H_{j,0}+1}.
 \end{aligned}$$

On the other hand we have

$$\begin{aligned}
 &(q^{\langle h_i, \alpha_j \rangle} \kappa^{m_u} z - w) E_j(w) E_i(z) \\
 &= \frac{q^{-2} \kappa^{m_u} z - w}{w^2 \kappa^{-m_u} \left(1 - \kappa^{m_u} \frac{z}{w}\right) \left(1 - q^{-2} \kappa^{m_u} \frac{z}{w}\right)} E_j^+(w) E_i^+(z) E_j^-(w) E_i^-(z) \\
 &\quad \times e^{\alpha_i} e^{\alpha_j} w^{H_{j,0}+1} z^{H_{i,0}+1} \\
 &= \frac{1}{(z - \kappa^{-m_u} w)} E_i^+(z) E_j^+(w) E_i^-(z) E_j^-(w) e^{\alpha_i} e^{\alpha_j} z^{H_{i,0}+1} w^{H_{j,0}+1}.
 \end{aligned}$$

Thus we have $(z\kappa^{m_u} - q^{\langle h_i, \alpha_j \rangle} w) E_i(z) E_j(w) = (q^{\langle h_i, \alpha_j \rangle} \kappa^{m_u} z - w) E_j(w) E_i(z)$.

The formula $(z\kappa^{m_u} - q^{-\langle h_i, \alpha_j \rangle} w) F_i(z) F_j(w) = (q^{-\langle h_i, \alpha_j \rangle} \kappa^{m_u} z - w) F_j(w) F_i(z)$ is proved similarly.

Let us prove (A.1.12). Assume that $\langle h_i, \alpha_j \rangle = -2$. This is the most complicated case. The other cases can be proved similarly.

We have following formulas:

$$\begin{aligned}
 &E_i(z_1) E_i(z_2) E_i(z_3) E_j(w) \\
 &= \frac{(z_1 - z_2) (z_1 - q^{-2} z_2) (z_1 - z_3) (z_1 - q^{-2} z_3) (z_2 - z_3) (z_2 - q^{-2} z_3)}{(\kappa^{\frac{1}{2} m_u} z_1 - \kappa^{-\frac{1}{2} m_u} w) (\kappa^{\frac{1}{2} m_u} z_1 - q^{-2} \kappa^{-\frac{1}{2} m_u} w) (\kappa^{\frac{1}{2} m_u} z_2 - \kappa^{-\frac{1}{2} m_u} w)} \\
 &\quad \times \frac{1}{(\kappa^{\frac{1}{2} m_u} z_2 - q^{-2} \kappa^{-\frac{1}{2} m_u} w) (\kappa^{\frac{1}{2} m_u} z_3 - \kappa^{-\frac{1}{2} m_u} w) (\kappa^{\frac{1}{2} m_u} z_3 - q^{-2} \kappa^{-\frac{1}{2} m_u} w)} \\
 &\quad \times E_i^+(z_1) E_i^+(z_2) E_i^+(z_3) E_j^+(w) E_i^-(z_1) E_i^-(z_2) E_i^-(z_3) E_j^-(w) \\
 &\quad \times e^{3\alpha_i} e^{\alpha_j} z_1^{H_{i,0}+1} z_2^{H_{i,0}+1} z_3^{H_{i,0}+1} w^{H_{j,0}+1},
 \end{aligned}$$

$$\begin{aligned}
 &E_i(z_1) E_i(z_2) E_j(w) E_i(z_3) \\
 &= \frac{(z_1 - z_2) (z_1 - q^{-2} z_2) (z_1 - z_3) (z_1 - q^{-2} z_3) (z_2 - z_3) (z_2 - q^{-2} z_3)}{(\kappa^{\frac{1}{2} m_u} z_1 - \kappa^{-\frac{1}{2} m_u} w) (\kappa^{\frac{1}{2} m_u} z_1 - q^{-2} \kappa^{-\frac{1}{2} m_u} w) (\kappa^{\frac{1}{2} m_u} z_2 - \kappa^{-\frac{1}{2} m_u} w)}
 \end{aligned}$$

$$\begin{aligned}
 & \times \frac{1}{(\kappa^{\frac{1}{2}m_\nu} z_2 - q^{-2} \kappa^{-\frac{1}{2}m_\nu} w) (\kappa^{-\frac{1}{2}m_\nu} w - \kappa^{\frac{1}{2}m_\nu} z_3) (\kappa^{-\frac{1}{2}m_\nu} w - q^{-2} \kappa^{\frac{1}{2}m_\nu} z_3)} \\
 & \times E_i^+(z_1) E_i^+(z_2) E_i^+(z_3) E_j^+(w) E_i^-(z_1) E_i^-(z_2) E_i^-(z_3) E_j^-(w) \\
 & \times e^{3\alpha_i} e^{\alpha_j} z_1^{H_{i,0}+1} z_2^{H_{i,0}+1} z_3^{H_{i,0}+1} w^{H_{j,0}+1}, \\
 E_i(z_1) E_j(w) E_i(z_2) E_i(z_3) \\
 & = \frac{(z_1 - z_2) (z_1 - q^{-2} z_2) (z_1 - z_3) (z_1 - q^{-2} z_3) (z_2 - z_3) (z_2 - q^{-2} z_3)}{(\kappa^{\frac{1}{2}m_\nu} z_1 - \kappa^{-\frac{1}{2}m_\nu} w) (\kappa^{\frac{1}{2}m_\nu} z_1 - q^{-2} \kappa^{-\frac{1}{2}m_\nu} w) (\kappa^{-\frac{1}{2}m_\nu} w - \kappa^{\frac{1}{2}m_\nu} z_2)} \\
 & \times \frac{1}{(\kappa^{-\frac{1}{2}m_\nu} w - q^{-2} \kappa^{\frac{1}{2}m_\nu} z_2) (\kappa^{-\frac{1}{2}m_\nu} w - \kappa^{\frac{1}{2}m_\nu} z_3) (\kappa^{-\frac{1}{2}m_\nu} w - q^{-2} \kappa^{\frac{1}{2}m_\nu} z_3)} \\
 & \times E_i^+(z_1) E_i^+(z_2) E_i^+(z_3) E_j^+(w) E_i^-(z_1) E_i^-(z_2) E_i^-(z_3) E_j^-(w) \\
 & \times e^{3\alpha_i} e^{\alpha_j} z_1^{H_{i,0}+1} z_2^{H_{i,0}+1} z_3^{H_{i,0}+1} w^{H_{j,0}+1}, \\
 E_j(w) E_i(z_1) E_i(z_2) E_i(z_3) \\
 & = \frac{(z_1 - z_2) (z_1 - q^{-2} z_2) (z_1 - z_3) (z_1 - q^{-2} z_3) (z_2 - z_3) (z_2 - q^{-2} z_3)}{(\kappa^{-\frac{1}{2}m_\nu} w - \kappa^{\frac{1}{2}m_\nu} z_1) (\kappa^{-\frac{1}{2}m_\nu} w - q^{-2} \kappa^{\frac{1}{2}m_\nu} z_1) (\kappa^{-\frac{1}{2}m_\nu} w - \kappa^{\frac{1}{2}m_\nu} z_2)} \\
 & \times \frac{1}{(\kappa^{-\frac{1}{2}m_\nu} w - q^{-2} \kappa^{\frac{1}{2}m_\nu} z_2) (\kappa^{-\frac{1}{2}m_\nu} w - \kappa^{\frac{1}{2}m_\nu} z_3) (\kappa^{-\frac{1}{2}m_\nu} w - q^{-2} \kappa^{\frac{1}{2}m_\nu} z_3)} \\
 & \times E_i^+(z_1) E_i^+(z_2) E_i^+(z_3) E_j^+(w) E_i^-(z_1) E_i^-(z_2) E_i^-(z_3) E_j^-(w) \\
 & \times e^{3\alpha_i} e^{\alpha_j} z_1^{H_{i,0}+1} z_2^{H_{i,0}+1} z_3^{H_{i,0}+1} w^{H_{j,0}+1}.
 \end{aligned}$$

Therefore it is enough to show that

(A.1.14)

$$\begin{aligned}
 & \sum_{\sigma \in \mathfrak{S}_3} \frac{\prod_{i < j} (z_{\sigma(i)} - z_{\sigma(j)}) (z_{\sigma(i)} - q^{-2} z_{\sigma(j)})}{\prod_{i=1}^3 (\kappa^{\frac{1}{2}m_\nu} z_{\sigma(i)} - \kappa^{-\frac{1}{2}m_\nu} w)} \\
 & \times \left\{ \frac{1}{(\kappa^{\frac{1}{2}m_\nu} z_{\sigma(1)} - q^{-2} \kappa^{-\frac{1}{2}m_\nu} w) (\kappa^{\frac{1}{2}m_\nu} z_{\sigma(2)} - q^{-2} \kappa^{-\frac{1}{2}m_\nu} w) (\kappa^{\frac{1}{2}m_\nu} z_{\sigma(3)} - q^{-2} \kappa^{-\frac{1}{2}m_\nu} w)} \right. \\
 & + \frac{q^2 + 1 + q^{-2}}{(\kappa^{\frac{1}{2}m_\nu} z_{\sigma(1)} - q^{-2} \kappa^{-\frac{1}{2}m_\nu} w) (\kappa^{\frac{1}{2}m_\nu} z_{\sigma(2)} - q^{-2} \kappa^{-\frac{1}{2}m_\nu} w) (\kappa^{-\frac{1}{2}m_\nu} w - q^{-2} \kappa^{\frac{1}{2}m_\nu} z_{\sigma(3)})} \\
 & \left. + \frac{q^2 + 1 + q^{-2}}{(\kappa^{\frac{1}{2}m_\nu} z_{\sigma(1)} - q^{-2} \kappa^{-\frac{1}{2}m_\nu} w) (\kappa^{-\frac{1}{2}m_\nu} w - q^{-2} \kappa^{\frac{1}{2}m_\nu} z_{\sigma(2)}) (\kappa^{-\frac{1}{2}m_\nu} w - q^{-2} \kappa^{\frac{1}{2}m_\nu} z_{\sigma(3)})} \right)
 \end{aligned}$$

$$+ \left. \frac{1}{(\kappa^{-\frac{1}{2}m\nu}w - q^{-2}\kappa^{\frac{1}{2}m\nu}z_{\sigma(1)}) (\kappa^{-\frac{1}{2}m\nu}w - q^{-2}\kappa^{\frac{1}{2}m\nu}z_{\sigma(2)}) (\kappa^{-\frac{1}{2}m\nu}w - q^{-2}\kappa^{\frac{1}{2}m\nu}z_{\sigma(3)})} \right\} = 0.$$

This identity is proved by a direct calculation.

Thus the proposition is proved.

Acknowledgment

The author would like to thank Masaki Kashiwara, Tetsuji Miwa and Michio Jimbo for their encouragement and valuable discussions. He also thanks Kenji Iohara, Norio Suzuki, Eric Vasserot and Hiroshi Yamada for stimulating discussions.

References

- [Be] Beck, J., Braid group action and quantum affine algebras, *Comm. Math. Phys.*, **165** (1994), 555-568.
- [B] Berman, S., On generators and relations for certain involutory subalgebras Kac-Moody Lie algebras, *Comm. Alg.*, **17**(12) (1989) 3165-3185.
- [BM] Berman, S. and Moody, R., Lie algebras graded by finite root systems and the intersection matrix algebras of Slodowy, *Inv. Math.*, **108**(1992), 323-347.
- [C] Cherednik, I., Double-affine Hecke algebras, Knizhnik-Zamolodchikov equations and Macdonald's operators, *Intern. Math. Research Notices*, **9**(1992), 171-180.
- [CP] Chari, V. and Pressly, A., Quantum affine algebras and affine Hecke algebras, *Pacific J. Math.*, **174**(1996), 295-326.
- [D1] Drinfeld, V., Quantum groups, *Proc. ICM Berkeley*, **1**(1986), 789-820.
- [D2] ———, A new realization of Yangians and quantized affine algebras, *Sov. Math. Dokl.*, **36** (1988), 212-216.
- [EM] Eswara Rao, S. and Moody, R. V., Vertex representations for N -toroidal Lie algebras and a Generalization of the Virasoro algebra, *Comm. Math. Phys.*, **159**(1994), 239-264.
- [FJ] Frenkel, I. and Jing, N., Vertex representations of quantum affine algebras, *Proc. Natl. Acad. Sci. USA*, **85** (1988), 9373-9377.
- [FK] Frenkel, I. and Kac, V., Basic Representations of affine Lie algebras and dual Resonance models, *Inv. Math.*, **62**(1980), 23-66.
- [GKV] Ginzburg, V., Kapranov, M. and Vasserot, E., Langlands reciprocity for algebraic surface, *Math. Res. Lett.*, **2**(1995), 147-160.
- [K] Koyama, Y., Staggered polarization of vertex model with $U_q(\widehat{\mathfrak{sl}}(n))$ -symmetry, *Comm. Math. Phys.*, **164** (1994), 277-291.
- [MEY] Moody, R. V., Eswara Rao, S. and Yokomura, T., Toroidal Lie algebras and vertex representations, *Geom. Dedicata*, **35**(1990), 283-307.
- [S] Slodowy, S., Beyond Kac-Moody algebras, and inside, *Canadian Math. Soc. Conference Proceedings*, **5**(1986), 361-371.
- [V] Vasserot, E., Private communication.
- [VV] Varagnolo, M. and Vasserot, E., Schur duality in the toroidal setting, *Comm. Math. Phys.*, **182**(1996), 469-484.
- [Y] Yamada, H., Extended affine Lie algebra and their vertex representations, *Publ. RIMS. Kyoto Univ.*, **25** (1989), 587-603.

