

On the Representation of Orthogonally Additive Polynomials in ℓ_p

By

Alberto IBORT*, Pablo LINARES** and José G. LLAVONA**

Abstract

We present a new proof of a Sundaresan's result which shows that the space of orthogonally additive polynomials $\mathcal{P}_o({}^k\ell_p)$ is isometrically isomorphic to $\ell_{p/p-k}$ if $k < p < \infty$ and to ℓ_∞ if $1 \leq p \leq k$.

§1. Introduction

A continuous scalar-valued map P in a Banach space is called a k -homogeneous polynomial if there exists a continuous k -linear form ϕ on X such that $P(x) = \phi(x, \dots, x)$. The space of k -homogeneous polynomials on X will be denoted by $\mathcal{P}({}^kX)$. It is a Banach space with the norm

$$\|P\| = \sup_{\|x\| \leq 1} |P(x)|$$

We will denote by $\widehat{\otimes}_{\pi,k} X$ and by $\widehat{\otimes}_{\pi,s,k} X$ the completed k -fold projective tensor product and the completed k -fold projective symmetric tensor product

Communicated by H. Okamoto. Received January 18, 2008.

2000 Mathematics Subject Classification(s): Primary 46G25; Secondary 46B42, 46M05.

Key words: orthogonally additive polynomials, tensor diagonal.

The first author was supported in part by Project MTM 2004-07090-C03. The second author was partially supported by the "Programa de formación del profesorado universitario del MEC". The second and third author were supported in part by Project MTM 2006-03531.

*Departamento de Matemáticas, Universidad Carlos III de Madrid, Avda. de la Universidad 30, 28911 Leganés, Spain.

e-mail: alberto@math.uc3m.es

**Departamento de Análisis Matemático, Facultad de Matemáticas, Universidad Complutense de Madrid, 28040 Madrid, Spain.

e-mail: plinares@mat.ucm.es and JL.Llavona@mat.ucm.es

respectively, where π denotes the projective norm. For $X = \ell_p$, the tensor diagonal is defined to be the closed subspace of $\widehat{\otimes}_{\pi,s,k} X$ generated by

$$e_n \otimes \overbrace{\cdots}^k \otimes e_n.$$

It will be denoted by $D_{k,p}$.

It will be needed the well known result that the dual of the space $\widehat{\otimes}_{\pi,s,k} X$ is isometrically isomorphic to the space $\mathcal{P}({}^k X)$. For notations and results about homogeneous polynomials, the reader is referred to [4] or [6], and for the theory of tensor products to [8].

We are interested in the subspace of $\mathcal{P}({}^k X)$ consisting of all orthogonally additive polynomials. Recall that if X is a Banach lattice, a k -homogeneous polynomial P is said to be orthogonally additive if $P(x + y) = P(x) + P(y)$ whenever x and y are orthogonal or disjoint elements of X (that is $|x| \wedge |y| = 0$). We will consider $X = \ell_p$ as a Banach lattice with its natural order given by $x = (x_n) \leq y = (y_n)$ if $x_n \leq y_n$. See [5] for more information about the theory of Banach lattices.

We give here a new proof of a result of Sundaresan (see [9]). This result has been recently generalized to all Banach lattices by Benyamini, Lasalle and Llavona in [2]. There are also independent proofs for the case of $X = C(K)$, see [3] and [7]. The reason to present this new proof is that, in our opinion, it is much simpler in the sense that there is no need of hard tools such as the representation of orthogonally additive functionals using Caratheodory functions, as in [9], or the Kakutani representation theorem for Banach lattices used in [2]. This new proof also makes more evident the underlying ideas.

§2. The Tensor Diagonal

In this section we generalize Example 2.23 in [8] to give a description of the tensor diagonal in $\widehat{\otimes}_{\pi,s,k} \ell_p$. The main technique was a Rademacher averaging ([8], Lemma 2.22). Its generalization requires the k -Rademacher generalized functions introduced by Aron and Globevnik in [1]:

Definition 2.1 [1]. Fix $k \in \mathbb{N}$, $k \geq 2$ and let $\alpha_1 = 1, \alpha_2, \dots, \alpha_k$ denote the k^{th} roots of unity. Let $r_1 : [0, 1] \rightarrow \mathbb{C}$ be the step function taking the value α_j on $(j - 1/k, j/k)$ for $j = 1, \dots, k$. Assuming that r_{n-1} has been defined, define r_n in the following way: fix any of the k^{n-1} sub-intervals I of $[0, 1]$ used in the definition of r_{n-1} . Divide I into k equal intervals I_1, \dots, I_k and set $r_n(t) = \alpha_j$ if $t \in I_j$.

We will also need the following

Lemma 2.2 [1]. *For each $k = 2, 3, \dots$ the associated functions r_n satisfy the following properties:*

- $|r_n(t)| = 1$ for all $n \in \mathbb{N}$ and all $t \in [0, 1]$.
- For any choice of n_1, \dots, n_k

$$\int_0^1 r_{n_1}(t) \cdots r_{n_k}(t) dt = \begin{cases} 1 & \text{if } n_1 = \cdots = n_k \\ 0 & \text{otherwise.} \end{cases}$$

The next lemma is a generalization of Lemma 2.22 in [8]:

Lemma 2.3 Rademacher Averaging. *Let X_1, \dots, X_k vector spaces and let $x_{1,1}, \dots, x_{1,n} \in X_1, \dots$, and $x_{k,1}, \dots, x_{k,n} \in X_k$. Then*

$$\sum_{i=1}^n x_{1,i} \otimes \cdots \otimes x_{k,i} = \int_0^1 \left(\sum_{i=1}^n r_i(t)x_{1,i} \right) \otimes \cdots \otimes \left(\sum_{i=1}^n r_i(t)x_{k,i} \right) dt.$$

Proof. Just expand the integral and use the second property of Lemma 2.2. □

Theorem 2.4. *Let $1 \leq p < \infty$. The tensor diagonal $D_{k,p}$ in $\widehat{\otimes}_{\pi,s,k} \ell_p$ is isometrically isomorphic to $\ell_{p/k}$ if $k < p < \infty$ and to ℓ_1 if $1 \leq p \leq k$.*

Proof.

1. $k < p < \infty$.

Let $u = \sum_{i=1}^n a_i e_i \otimes \cdots \otimes e_i \in D_{k,p}$. Using Lemma 2.3, we write

$$u = \int_0^1 \left(\sum_{i=1}^n \text{sign}(a_i) |a_i|^{1/k} r_i(t) e_i \right) \otimes \cdots \otimes \left(\sum_{i=1}^n |a_i|^{1/k} r_i(t) e_i \right) dt$$

and, like in [8] (pages 34–35) we get:

$$\begin{aligned} \pi(u) &\leq \sup_{0 \leq t \leq 1} \left\| \sum_{i=1}^n \text{sign}(a_i) |a_i|^{1/k} r_i(t) e_i \right\|_p \cdots \left\| \sum_{i=1}^n |a_i|^{1/k} r_i(t) e_i \right\|_p = \\ &= \left(\sum_{i=1}^n |a_i|^{p/k} \right)^{k/p} = \|(a_i)\|_{p/k} \end{aligned}$$

To prove the identity, define a k -linear form on ℓ_p by $B(x_1, \dots, x_k) = \sum b_i x_{1,i} \cdots x_{k,i}$ where $x_j = (x_{j,n})$ and $b_i = \text{sign}(a_i) |a_i|^{p/k-1}$. Using Hölder's inequality, it is easy to see that $\|B\| \leq (\sum_{i=1}^n |a_i|^{p/k})^{1-p/k}$ and then

$$\sum_{i=1}^n |a_i|^{p/k} = |\langle u, B \rangle| \leq \pi(u) \left(\sum_{i=1}^n |a_i|^{p/k} \right)^{1-p/k}.$$

Hence $\|(a_i)\|_{p/k} \leq \pi(u)$ and therefore $D_{k,p}$ is isometrically isomorphic to $\ell_{p/k}$.

2. $1 \leq p \leq k$.

Let be $u = \sum_{i=0}^n a_i e_i \otimes \cdots \otimes e_i \in D_{k,p}$, then $\pi(u) \leq \sum_{i=0}^n |a_i|$. Reciprocally, define $B(x_1, \dots, x_k) = \sum_{i=0}^n \text{sign}(a_i) x_{1,i} \cdots x_{k,i}$, we have that $|B(x_1, \dots, x_k)| \leq \|x_1\|_k \cdots \|x_k\|_k \leq \|x_1\|_p \cdots \|x_k\|_p$ and so $\|B\| \leq 1$. Then

$$\pi(u) \geq \langle u, B \rangle = \sum_{i=1}^{\infty} |a_i|$$

and we are done. \square

Remark 1. Definition 2.1 gives the classical Rademacher functions for the case $k = 2$ and these results are those in [8], (Example 2.23, page 34).

§3. The Main Result

Our proof is based in the fact that the orthogonally additive polynomials are isometrically isomorphic to the dual of the tensor diagonal $D_{k,p}$. We need a previous lemma:

Lemma 3.1. *Let $1 \leq p < \infty$. The dual of the tensor diagonal $D_{k,p}^*$ is isometrically isomorphic to ℓ_∞ if $1 \leq p \leq k$ and to $\ell_{p/p-k}$ if $k < p < \infty$ in the sense that for every $F \in D_{k,p}^*$, $(F(e_i \otimes \cdots \otimes e_i))$ is in ℓ_∞ for the first case and in $\ell_{p/p-k}$ for the second.*

Proof. The proof is standard, observe that $\ell_{p/p-k}$ is the dual of $\ell_{p/k}$ and carry on the same proof of $\ell_q^* = \ell_{q'}$ for $1/q + 1/q' = 1$ with the identification of the projective norm shown in Theorem 2.4. \square

We are now ready to prove the theorem:

Theorem 3.2. *Let $1 \leq p < \infty$. The space of orthogonally additive, k -homogeneous polynomials $\mathcal{P}_o({}^k\ell_p)$ is isometrically isomorphic to ℓ_∞ for $1 \leq p \leq k$ and to $\ell_{p/p-k}$ for $k < p < \infty$.*

Proof. The proof consists on showing that $\mathcal{P}_o({}^k\ell_p)$ is isometrically isomorphic to $D_{k,p}^*$. We will suppose that $k < p < \infty$. The other case is analogous.

Let $F \in D_{k,p}^*$, the correspondence is established by associating F to the polynomial $P(x) = \tilde{F}(x \otimes \cdots \otimes x)$ where $\tilde{F} \in (\widehat{\otimes}_{\pi,k,s} \ell_p)^*$ is defined by

$$\tilde{F}(e_{n_1} \otimes \cdots \otimes e_{n_k}) = \begin{cases} F(e_{n_1} \otimes \cdots \otimes e_{n_k}) & \text{if } n_1 = \cdots = n_k \\ 0 & \text{otherwise} \end{cases}$$

To see that \tilde{F} is well defined let $x = \sum x_n e_n \in \ell_p$, then

$$\begin{aligned} |\tilde{F}(x \otimes \cdots \otimes x)| &\leq \sum |x_{n_1} \cdots x_{n_k} \tilde{F}(e_{n_1} \otimes \cdots \otimes e_{n_k})| = \sum |x_n^k F(e_n \otimes \cdots \otimes e_n)| \\ &\leq \|x_n^k\|_{p/k} \|F(e_n \otimes \cdots \otimes e_n)\|_{p/p-k} \leq \|F\| \|x\|_p^k = \\ &= \|F\| \pi(x \otimes \cdots \otimes x) \end{aligned}$$

hence \tilde{F} is continuous and $\|\tilde{F}\| \leq \|F\|$, then we have the equality since \tilde{F} was an extension of F .

To see that $P(x) = \tilde{F}(x \otimes \cdots \otimes x)$ is orthogonally additive, note that is enough to check that the k -linear symmetric form ϕ associated to P , verifies that $\phi(e_{n_1}, \dots, e_{n_k})$ is zero whenever at least two of its entries will be different and this is true by definition of \tilde{F} .

Finally as the correspondence between polynomials and the dual of the symmetric tensor product is an isometric isomorphism, $\|P\| = \|\tilde{F}\| = \|F\|$ which completes the proof. \square

As it has been shown, if $P(x) = \phi(x, \dots, x)$ is an orthogonally additive polynomial on ℓ_p , the essential information of P is contained in the sequence $(\phi(e_n, \dots, e_n))$. There is another possible proof of Theorem 3.2 using a result by Zalduendo (see Corollary 1 of [10]):

Lemma 3.3 [10]. *Let $k < p$ and let ϕ a continuous k -linear form on ℓ_p . Then*

$$(\phi(e_n, \dots, e_n)) \in \ell_{p/p-k}.$$

From this result, it can be shown as well that the space of orthogonally additive polynomials is isometrically isomorphic to $\ell_{p/p-k}$ if $k < p$. We don't repeat the proof since the ideas are essentially the same, however the proof presented here is self-contained.

Remark 2. This method will also be valid for $1 \leq p \leq k$ since in this case, trivially $(\phi(e_n, \dots, e_n)) \in \ell_\infty$. It is shown in [10] that this is the best characterization.

References

- [1] R. M. Aron and J. Globevnik, *Analytic functions on c_0* , Rev. Mat. Univ. Complut. Madrid **2** (1989), suppl., 27–33.
- [2] Y. Benyamini, S. Lassalle and J. G. Llavona, *Homogeneous orthogonally additive polynomials on Banach lattices*, Bull. London Math. Soc. **38** (2006), no. 3, 459–469.
- [3] D. Carando, S. Lassalle and I. Zalduendo, *Orthogonally additive polynomials over $C(K)$ are measures—a short proof*, Integral Equations Operator Theory **56** (2006), no. 4, 597–602.
- [4] S. Dineen, *Complex analysis on infinite-dimensional spaces*, Springer Monographs in Mathematics, Springer, London, 1999.
- [5] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces. II*, Springer, Berlin, 1979.
- [6] J. Mujica, *Complex analysis in Banach spaces*, North-Holland, Math. Studies, 120, Amsterdam, 1986.
- [7] D. Pérez-García and I. Villanueva, *Orthogonally additive polynomials on spaces of continuous functions*, J. Math. Anal. Appl. **306** (2005), no. 1, 97–105.
- [8] R. A. Ryan, *Introduction to tensor products of Banach spaces*, Springer, London, 2002.
- [9] K. Sundaresan, *Geometry of spaces of homogeneous polynomials on Banach lattices*, in Applied geometry and discrete Mathematics, 571–586, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., 4, Amer. Math. Soc., Providence, RI, 1991.
- [10] I. Zalduendo, *An estimate for multilinear forms on ℓ_p spaces*, Proc. Roy. Irish Acad. Sect. A **93** (1993), no. 1, 137–142.